1   **Well-resolved phylogeny reveals multiple origins of keel flowers during the**
2   **diversification of papilionoid legumes**
3

4   Liming Cai[1], Domingos Cardoso[2,3], Lydia G. Tressel[1], Chaehee Lee[1,4], Bikash Shrestha[1,5], In-
5   Su Choi[6,7], Haroldo C. de Lima[3], Luciano P. de Queiroz[8], Tracey A. Ruhlman[1], Robert K.
6   Jansen[1], and Martin F. Wojciechowski[6]
7
8

[1] Department of Integrative Biology, University of Texas at Austin, Austin, Texas, USA
[2] Instituto de Biologia, Universidade Federal da Bahia, Salvador, Bahia, Brazil
[3] Instituto de Pesquisas Jardim Botânico do Rio de Janeiro, Rio de Janeiro, Brazil
[4] Department of Plant Sciences, University of California Davis, Davis, California, USA
[5] DOE Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, CA, USA
[6] School of Life Sciences, Arizona State University, Tempe, Arizona, USA
[7] Department of Biological Sciences and Biotechnology, Hannam University, Daejeon, Korea
[8] Universidade Estadual de Feira de Santana, Feira de Santana, Bahia, Brazil
17
18

Authors for correspondence:
Liming Cai
Email: cail@ufl.edu
Domingos Cardoso
Email: cardosobot@gmail.com
Martin F. Wojciechowski
Email: mfwojciechowski@asu.edu
26
27

Summary

● The butterfly-shaped keel flowers are among the most successful pollination syndromes

in angiosperms. These flowers steer the mechanical interaction with bees and thus are

hypothesized to accelerate pollinator-driven diversification. The highly labile evolution of

keel flowers in Papilionoideae (Fabaceae) provides a suitable system to test this hypothesis

but a robust phylogeny is lacking.

● Using 1,456 low-copy nuclear loci, we confidently resolve the early divergence history of

Papilionoideae. Constrained by this backbone phylogeny, we generated a time tree for

3,326 Fabales to evaluate the tempo and mode of diversification within a state-dependent

evolutionary framework.

● The first keel flowers emerged around 59.0 Ma in Papilionoideae, predating the earliest

fossil by 3–4 million years. The Miocene diversification of Papilionoideae coincided with

rapid evolution of keel flowers. At least seven independent origins and twenty-seven losses

of keel flowers were identified in Papilionoideae, Polygalaceae, and Cercidoideae during

this time. However, the state-dependent diversification model was not favored.

● Lack of radiation associated with keel flowers suggests that diversification within

Papilionoideae was not solely driven by pollinator-mediated selection, but instead an

outcome of the synergistic effects of multiple innovations including nitrogen fixing and

chemical defense as well as dispersal to the subtropical and temperate regions.

Keywords: Fabales, floral evolution, pollination-driven speciation, coalescent method,

phylogenomics, SSE model

INTRODUCTION

Biodiversity is unevenly distributed across the Tree of Life and elucidating the underlying processes is a fundamental task in biological research (Vargas & Zardoya, 2014). In plants, many speciation mechanisms have been proposed including adaptation to pollinators and arms race with herbivores (Givnish, 2010; Van der Niet *et al.*, 2014; Johnson & Schiestl, 2016; Maron *et al.*, 2019). There is also a growing consensus that these factors may act in synergy to promote diversification (Donoghue & Sanderson, 2015). Legumes (Fabaceae) are among the most ecologically and evolutionarily successful plant families on Earth. With approximately 23,000 species and 800 genera, legumes have defined biomes from tropical savannas to temperate sagebrush steppes (Schrire *et al.*, 2005; Lewis *et al.*, 2005; LPWG, 2017; Legume Data Portal, https://www.legumedata.org/). This massive diversity has been hypothesized to be an outcome of a suite of key innovations including nitrogen fixation symbiotically with soil bacteria (Sprent, 2001; Werner *et al.*, 2015; Doyle, 2016; Faria *et al.*, 2022), direct chemical defense with alkaloids and canavanine (Bell, 1981; Bisby *et al.*, 1994; Kursar *et al.*, 2009; Wink, 2013), indirect defense via ant domatia and extrafloral nectaries (Marazzi & Sanderson, 2010; Chomicki & Renner, 2015; Marazzi *et al.*, 2019), and an extraordinary range of floral architectures to enhance pollination interaction (Arroyo, 1981; Polhill *et al.*, 1981; Pennington *et al.*, 2000; Cardoso *et al.*, 2012, 2013b).

Compared to nitrogen fixation and defense chemicals, the macroevolution of floral architecture has received relatively less attention in Fabaceae. Members of Fabaceae subfamilies often exhibit lineage-specific floral traits tailored to particular pollinators, such as the showy stamens in Mimosoideae for bees, the night-blooming Caesalpinioideae for moths and bats, and the distinctive butterfly-like keel flowers in Papilionoideae, which are closely linked to bee pollination (Arroyo, 1981). Keel flowers are bilateral flowers with two showy wings and a lower keel protecting the reproductive parts (Westerkamp, 1997). They are most commonly found in Papilionoideae and Polygalaceae, but also independently evolved several times in other angiosperm families such as Ranunculaceae, Geraniaceae, and Commelinaceae (Westerkamp, 1997). These repeated evolutions exhibit functional convergence but divergent developmental pathways (Bello *et al.,* 2012). The bilateral symmetry and fused stamens can effectively restrict the direction of entry by pollinators

and aid with precise pollen placement (Neal *et al.*, 1998). Furthermore, many

Papilionoideae flowers exhibit primary and secondary pollination mechanisms such as

valvular, pump, explosive, and brush mechanisms to prevent pollen waste and deposit

pollen on particular bee body parts (Etcheverry *et al.*, 2012; Uluer *et al.*, 2022). Therefore,

the evolution of keel flowers is considered to be an outcome of pollinator-mediated

selection (Grant, 1949; Stebbins, 1970; Rosas-Guerrero *et al.*, 2014; Stewart *et al.*, 2022).

There is mounting evidence from other empirical systems that pollinator-mediated

selection can be a powerful driver of diversification (e.g., the bat-pollinated *Burmeistera*;

the beak of the galea in *Pedicularis*; Ree, 2005; Muchhala, 2006), but this has yet to be

tested in legumes.

To rigorously assess trait-driven diversification, a system with high trait variability is

essential for robust comparative hypothesis testing (Ricklefs & Renner, 1994, 2000;

Donoghue & Sanderson, 2015). The remarkable diversity in floral symmetry and

architecture in the Papilionoideae offers an ideal model for such investigations. Although

keel flowers are considered an essential feature and a hallmark of Papilionoideae (Polhill *et*

al., 1981), there is considerable variation in floral symmetry in early diverging

Papilionoideae clades. For example, members from the tribes Sophoreae and Swartzieae

are frequently actinomorphic, with five petals undifferentiated or even lost to various

degrees (Pennington *et al.*, 2000; Cardoso *et al.*, 2012, 2013b). These genera are typically

species-poor and more than 80% have no more than ten extant species based on records

from the Plants of the World Online (POWO; https://powo.science.kew.org). In contrast,

the keel flower is highly constrained among the large non-protein-amino-acid accumulating

clade (NPAAA; ca. 10,200 species), which is mostly known for economically important

crops like soybeans, peas, lentils, and clovers (Choi *et al.*, 2022). These general

observations have prompted our speculation that the evolution of keel flowers may be

associated with increased diversification rates in Papilionoideae.

Testing these macroevolutionary hypotheses requires a robust and densely sampled

phylogeny. Over the past three decades, a general phylogenetic framework for Fabaceae

has been developed thanks to a tremendous network of global collaboration (LPWG, 2013,

2017). Six distinct subfamilies are recognized by the Legume Phylogeny Working Group

(LPWG), with Papilionoideae as the most species-rich subfamily comprising 14,000 species

in 503 genera (Lewis *et al.*, 2005; LPWG, 2017). However, the LPWG phylogeny and

classification is largely based on two plastid genes *matK* and *rbcL*, which have obvious

limitations when dealing with rapid radiations or complex evolutionary scenarios such as

hybridization (Barrett *et al.*, 2014; Morales-Briones *et al.*, 2021). In Papilionoideae, studies

based on *matK*, *rbcL*, and complete plastomes have consistently defined 22 well-supported

clades but the relationships among these clades remain poorly resolved (Wojciechowski *et*

al., 2004; McMahon & Sanderson, 2006; Cardoso *et al.*, 2012, 2013b; Zhang *et al.*, 2020;

Choi *et al.*, 2022). Most significantly, Swartzieae and the ADA clade have been individually

proposed as sister to all remaining Papilionoideae. Given that Swartzieae contains mostly

non-keel flowers with frequent petal losses whereas the ADA clade contains largely keel

flowers (Torke & Schaal, 2008; Cardoso *et al.*, 2015), their placement will have a strong

bearing on the timing and evolutionary trajectory of keel flowers. Furthermore, many of

the 22 clades are characterized by tropical trees that are scarce in distribution and

underrepresented in previous studies. These include species-poor clades such as the

African *Amphimas*, the Amazonian *Aldina,* and the North American *Dermatophyllum*

(Cardoso *et al.*, 2013). Along these lines, recent advances in legume phylogeny using multi-

locus nuclear datasets have promising applications in resolving complex evolutionary

histories (Koenen *et al.*, 2021; Zhao *et al.*, 2021) and inspired us to apply this technique to

address the recalcitrant relationships within Papilionoideae.

In this study, we generated a phylogenomic dataset of 1,456 low-copy genes from 287

species with a focus on recalcitrant clades in Papilionoideae. Using both coalescent and

concatenation methods, we inferred a robust species tree and consistently resolved the

early divergence history of Papilionoideae. This nuclear phylogeny was subsequently used

as the backbone to reconstruct a 3,326-species time tree of Fabales to explore the evolution

of keel flowers and to test whether it has promoted diversification in Fabales.

MATERIALS AND METHODS

Taxon sampling

Our taxon sampling included 279 species broadly distributed across Papilionoideae, plus

one Cercidoideae, one Detarioideae, one Dialioideae, and five Caesalpinioideae species as

outgroups (Tables S1–2). The dataset comprised an aggregation of 47 newly generated

transcriptomes and 32 low-pass genomic sequences (Table S1) as well as published

sequences from 208 species from Zhao *et al.* (2021) to broaden taxon sampling (Table S2).

Newly generated transcriptomes and genomic sequences were mostly from

morphologically and evolutionarily representative genera from the early diverging clades

of Papilionoideae (Cardoso *et al.*, 2012; Choi *et al.*, 2022). Collectively the dataset

represented 19 of the 22 major clades consistently recognized in the subfamily (Cardoso *et*

al., 2012; Cardoso *et al.*, 2013; Choi *et al.*, 2022). The three clades we did not include are

Amphimas, *Aldina*, and the recently described *Cabari* (Gregório *et al.*, 2024). All newly

generated sequences were derived from plant materials collected in the field or grown in

the University of Texas at Austin greenhouses (Table S1). Voucher specimens were

deposited in the herbaria of the Rio de Janeiro Botanic Garden (RB), State University at

Feira de Santana (HUEFS), and the Billie L. Turner Plant Resources Center (TEX-LL). For

DNA and RNA extraction, the methods described previously by Lee *et al.* (2021) and Zhang

et al. (2013) were employed, respectively.

RNA sequencing and transcriptome assembly

Total RNA samples were shipped to GENEWIZ (South Plainfield, NJ, USA) for ribosomal

RNA (rRNA) depletion, library preparation, and sequencing of approximately 25 million

paired-end reads on an Illumina HiSeq platform (Illumina, San Diego, CA). We accessed the

quality of raw 150 bp paired-end reads using FastQC v.0.11.5

(www.bioinformatics.babraham.ac.uk/projects/fastqc) and removed filtered adapters and

low-quality reads using TrimGalore v.0.5.0

(https://github.com/FelixKrueger/TrimGalore). The rRNA was removed from the merged

trimmed reads using SortMeRNA v.2.1 (Kopylova *et al.*, 2012) with eight rRNA databases

for prokaryotes, archaea, and eukaryotes. Clean RNA-Seq reads were assembled *de novo*

using Trinity v.2.8.4 (Grabherr *et al.*, 2011), and the assembled transcripts were subjected

to the assessment of completeness and redundancy using BUSCO v.4.1.0 (Benchmarking

Universal Single-Copy Orthologs; Simão *et al.*, 2015) with the eudicots_odb10.2019-11-20

database. The number of raw reads and BUSCO evaluation results are provided in Table S1.

All newly generated reads are available on GenBank Sequence Read Archive under

Bioproject PRJNA1120003 (https://www.ncbi.nlm.nih.gov/sra/PRJNA1120003).

Low-copy nuclear dataset assembly

We used the 1,559 low-copy nuclear loci identified by Zhao *et al.* (2021) for our

phylogenetic reconstruction. To recover these loci in our transcriptomes, we used the

profile hidden Markov models implemented in HMMER v3.3.2 (Johnson *et al.,* 2010) to

identify orthologs. Briefly, a DNA profile database was built for each locus using the

function 'hmmbuild' based on the DNA alignments from Zhao *et al.* (2021). These profiles

were then searched against the target transcriptomes with an e-value threshold of 1e-50

using 'hmmsearch'. We then extracted the best hit from each species as the candidate

ortholog using a custom Python script (all scripts available on GitHub repository

https://github.com/lmcai/Papilionoideae_phylogenomics).

To recover the same set of loci from our genome sequence data, we developed a new

function of the PhyloHerb pipeline (Cai *et al.*, 2022) to automatically extract nuclear loci

from low-pass genome sequencing data. Briefly, PhyloHerb accomplishes this in three steps

— mapping, assembly, and scaffolding. First, it mapped reads to the reference sequence

using the very-sensitive mode of Bowtie v2.3.4.1 (Langmead & Salzberg, 2012). Mapped

reads were then assembled into contigs using SPAdes v3.8.0 under the default setting

(Bankevich *et al.*, 2012). The contigs were scaffolded into full-length loci based on syntenic

information compared to the reference using BLAST with an e-value threshold of 1e-40.

These functions were integrated as the '-m assemb' and '-m ortho -nuc' functions in

PhyloHerb v1.1. For details, see the GitHub page

(https://github.com/lmcai/PhyloHerb/#iv-retrieve-low-copy-nuclear-genes). Finally, for

quality control purposes, we also estimated the nuclear genome coverage of genome

sequences using Jellyfish v2.3.0 with a kmer size of 21 (Marçais & Kingsford, 2011).

Sequence alignment, dataset cleaning, and phylogenetic inference

Sequence alignment of each locus was implemented in MAFFT v7.245 (Katoh & Standley,

2013) under the default settings. The aligned sequences were subjected to three rounds of

stringent cleaning to remove excessive missing data and potential paralogs. First, we used

the '-missing' function in PhyloHerb to remove sequences with >70% ambiguous sites

including gaps in each locus. Second, a phylogeny-based ortholog identification method

from Yang and Smith (2014) was used to remove potential paralogs. To accomplish this, a

maximum likelihood (ML) phylogeny was inferred for each gene using the DNA alignments

from the previous cleaning step. The best substitution model was automatically

determined by ModelFinder implemented in IQ-TREE v.2.2.2.6 (Minh *et al.*, 2020).

Branches that were more than fifteen times the mean branch length were pruned from the

resulting trees using a custom Python script 'prune_long_branches.py' (available on

GitHub). Then, a phylogeny-based ortholog identification approach using the Python script

'prune_paralogs_RT.py' (Yang & Smith 2014) was applied to further remove potential

contaminants of paralogs. Third, all DNA sequences were realigned using the MAFFT E-INS-

i algorithm (--genafpair --maxiterate 1000) after removing spurious sequences identified in

the previous steps. Sites with more than 70% missing data were removed from the

alignment using the '-gt 0.7' flag in trimAL. We also found that the published alignments

from Zhao *et al.* (2021) were site-trimmed, thus creating systematic biases for phylogenetic

inference (e.g., shared gaps for sequences from Zhao *et al.* (2021) compared to species

sampled by us). To accommodate for this, we used a custom Python script to mask all sites

in our sequences that were present in >60% of species but absent in >90% of species from

Zhao *et al.* (2021). Loci with less than 199 (70%) taxa were removed from downstream

phylogenomic assessments. In the end, 1,456 out of the 1,559 loci passed our screening

criteria (Fig. S1).

To estimate a species tree using the multispecies-coalescent model, we first inferred

individual gene trees for each of the 1,456 loci in IQ-TREE. The best substitution model was

automatically determined by ModelFinder implemented in IQ-TREE and branch support

was evaluated by 1000 ultrafast bootstrap replications. The species tree was inferred from

1,456 gene trees using ASTRAL-III v5.7.8 (Zhang *et al.*, 2017) under the default settings.

Meanwhile, to further mitigate the impact of artificially resolved gene tree branches with

low support values, we used newick-utils v1.6 (Junier & Zdobnov, 2010) to collapse gene

tree branches with ultrafast bootstrap support values less than 70. We then inferred a

species tree with these gene trees containing polytomies in ASTRAL-III.

To infer the species tree using the concatenation methods, we first further subsampled the

loci based on length and missing data. Any loci with alignments shorter than 900 base pairs

(bp; 33% loci removed) or less than 227 species (80% of the total species) were excluded

from the concatenation analysis, which resulted in 672 loci. The concatenated supermatrix

was generated by the '-m conc' function in PhyloHerb, along with a gene delineation file

input into PartitionFinder 2. The best partition scheme for ML analyses employing the

general time reversible (GTR) + gamma model was selected by PartitionFinder 2 using the

heuristic search algorithm 'rcluster' (Lanfear *et al.*, 2017). We then inferred an ML

phylogeny of the 287 species using IQ-TREE with 100 traditional nonparametric bootstrap

replications (BP) under the GTR + gamma nucleotide substitution model and linked but

variable evolution rates for each partition with the '-p' flag.

Topology test

Several conflicting relationships emerged from the comparison between concatenation and

coalescent analyses. To explore the origin of these conflicts and identify the best supported

phylogenetic scenarios, we conducted topology tests on six focal groups defined in Table

S3. By alternating the position of the focal group and fixing the rest of the tree, we inferred

an optimum phylogeny for competing scenarios in IQ-TREE. We then conducted the

approximately unbiased (AU), Kishino-Hasegawa (KH), and Shimodaira-Hasegawa (SH)

tests with 10,000 replicates as implemented in IQ-TREE. To compare the goodness-of-fit of

alternative species trees to gene trees under the coalescent model, we conducted a

pairwise likelihood ratio test (Liu *et al.*, 2019). We similarly optimized the branch length of

fixed topologies with alternative positions of the focal group in MP-EST v3.0 (Liu *et al.*,

2010). We used the test2.sptree function from the R package Phybase v2.0 (Liu & Yu, 2010)

to conduct the likelihood ratio test described in Liu *et al.* (2019). The likelihood ratio test

statistic is given by $\tau = 2(L_{\text{Tree2}} - L_{\text{Tree1}})$, in which L_{Tree1} and L_{Tree2} are the log-likelihoods of

the null and alternative hypotheses, respectively. The statistical significance (*p*-value) of

the comparison is evaluated by generating a null distribution of τ via bootstrap sampling

gene trees 100 times. Finally, to diagnose the causes of conflicting results, we examined the

gene (gCF) and site concordance factors (sCF), as well as gene-wise likelihood scores

implemented in IQ-TREE.

Bayesian divergence time estimation

To infer a Bayesian time tree, we subsampled the nuclear dataset to select the most clock-

like genes using SortaDate (Smith *et al.*, 2018). Total tree length, root-to-tip variance, and

topological congruence with the species tree were used as key indicators for phylogenetic

informativeness and stable molecular rate. We selected the 30 best genes (Table S4) for

dating to facilitate a reasonable running time of the Bayesian analysis, an approach

comparable to a recent study of Fabaceae divergence time (Koenen *et al.*, 2021). We

carefully selected eleven fossils to calibrate our molecular dating analyses, including four

fossils from early diverging eudicot lineages and seven well-recognized fossils within

Fabales (Table S5). In particular, the earliest fossil fruits and leaflets of Fabaceae from the

Early Paleocene (63.5 Ma) of the Denver Basin in Colorado, USA, were used to constrain the

stem group Fabaceae (Lyson *et al.*, 2019). Many recent fossils within Fabaceae were not

included because they provide redundant constraints compared to the eleven fossils we

selected. We used BEAST v.2.2.1 (Bouckaert *et al.*, 2014) to estimate divergence times

under the GTR substitution model, an uncorrelated relaxed molecular clock model, and a

birth-death tree prior. The topology was fixed to reflect the optimum species tree inferred

from the ASTRAL analysis. Fossil calibrations were set as exponential priors with the offset

minimum ages aligned with the fossils described in Table S5. The root node, for which we

used a normal prior at 128.6 Ma with a standard deviation of 1.0, truncated to minimum

and maximum ages of 113 Ma (the Aptian–Albian boundary) and 136 Ma (the oldest known

crown angiosperm fossil (Magallón *et al.*, 2015). We ran analyses under the uncorrelated

lognormal model for 100 million generations and confirmed convergence with Tracer

v1.7.1 (Rambaut *et al.*, 2018). The first 30% of the generations were discarded as burn-in

before summarizing median branch lengths and substitution rates with TreeAnnotator

from the BEAST package.

Time tree inference for a 3,326-species *matK* dataset of Fabales

To evaluate the macroevolutionary trends of floral diversification in Papilionoideae under a broader taxonomic context, we generated a plastid *matK* gene dataset that included 3,106 species from 794 genera in Fabaceae (92% genus-level completeness), plus 220 additional species from all other families in Fabales and outgroups. Sequences from Choi *et al.* (2022), Legume Phylogeny Working Group (LPWG 2017), and Gregório *et al.* (2024) were consolidated to generate the most comprehensive *matK* database for Fabaceae, and combined with *matK* sequences from Polygalaceae, Surianaceae, and Quillajaceae available on GenBank, which served as outgroups in Fabales (Table S6).

As a first step of data curation, we updated the nomenclature to reflect the most recent effort from POWO, Flora e Funga do Brasil (https://floradobrasil.jbrj.gov.br/consulta; BFG 2022), and taxonomic studies from individual clades (e.g., Hughes *et al.*, 2022). Second, we employed multiple rounds of phylogeny-guided cleaning for the *matK* sequence alignment. Initial alignment was conducted with the translation align option of MAFFT v.7.490 implemented in Geneious Prime 2023.1.2 (https://www.geneious.com/) and was manually inspected thereafter. For outgroup species, a separate alignment was generated using the L-INS-i algorithm in MAFFT to accommodate higher sequence divergence across families. These two alignments were then merged using the profile-profile alignment algorithm implemented in ClustalX (Thompson *et al.*, 2003). A draft ML phylogeny was inferred under the GTR+I+G model with 1000 ultrafast bootstrap replications in IQ-TREE and we filtered the *matK* matrix accordingly to remove misidentified and rogue taxa.

To infer a robust *matK* phylogeny for divergence time estimation, we used our well-supported nuclear phylogeny to constrain the topology and node ages of major clades in the *matK* dataset. This is necessary because the slow substitution rate of *matK* may result in contentious relationships and the lack of sequence divergence among close relatives may impact ancestral state estimation and divergence time estimation. As a result, relationships *between* the following clades were constrained: the ADA, Swartzieae, Cladrastis, Vataireoid, Exostyleae, Dalbergioid s.l., Genistoid s.l., Andira, Baphieae, and NPAAA. Relationships *within* each clade and *between* all subfamilies and families remained as-is. With this

partially fixed topology, the ML phylogeny was inferred under the best fitting model in IQ-
TREE.

Thirty secondary calibration points from the Bayesian time tree (Table S7) were used to
infer the *matK* gene time tree under the penalized likelihood method implemented in
TreePL v1.0 (Smith & O'Meara, 2012). These calibration points were imposed as both
maximum and minimum age constraints to fix node ages. We then primed and cross-
validated our data to establish the best smoothing parameter (smooth = 10) for our
dataset. Finally, we conducted a thorough search to infer the best time tree.

Testing state-dependent diversification analysis
We performed a Bayesian analysis of macroevolutionary mixtures using the program
BAMM v2.5.0 (Rabosky, 2014) to detect shifts in diversification rates. Reversible-jump
Markov chain Monte Carlo was run for 20 million generations and sampled every 1,000
generations. Posterior probabilities of rate shifts were summarized with the R package
BAMMtools v.2.1.6 (Rabosky *et al.*, 2014) and visualized using the scripts from (Padfield *et*
al., 2024).

To explore whether the evolution of keel flowers drives pollinator-associated
diversification in clades like Papilionoideae (Fabaceae) and Polygaleae (Polygalaceae), we
estimated the divergence time history of Fabales under the state-dependent speciation and
extinction (SSE) models. First, we coded the floral type of all species as either keel or non-
keel flowers (Table S6). Here, keel flowers are defined as bilaterally symmetrical,
specialized flowers with petals highly differentiated into a standard, two wings, and a keel
composed of two partially to fully fused petals enclosing the reproductive organs.
Connation among floral components, including stamens and keel petals, is common. To
infer the sampling ratio, we used the powoGenera function from the R package expowo
(Zuanny *et al.*, 2024) to obtain the number of species for each genus based on the Plants of
the World Online (https://powo.science.kew.org) database. The total species number for
the keel flower clades and non-keel flower clades was enumerated to calculate the
sampling ratio for these two groups. Finally, we inferred the diversification rate under the

state-dependent and state-independent models using the R package hisse v2.1.8 (Beaulieu

& O'Meara, 2016). We applied our dataset to the Binary State Speciation and Extinction

model (BiSSE), the Hidden State Speciation and Extinction model (HiSSE), as well as the

state-independent models CID-2 and CID-4 that have equal parameter complexity

compared to BiSSE and HiSSE, respectively (Maddison *et al.*, 2007; Beaulieu & O'Meara,

2016). We used the function 'hisse' followed by 'MarginReconHiSSE' to infer the speciation

and extinction rates as well as ancestral states under the HiSSE model.

RESULTS

An efficient phylogenetic data assembly pipeline based on a curated set of loci

BUSCO assessments for the assembled transcriptomes showed a high level of

completeness, with the percentage of complete orthologous genes ranging from 70.3% to

91.1% (Table S1). The complete multi-loci dataset included 1,456 genes with a mean length

of 1,003 bp after trimming (Data S1). These loci on average contained 253 (85%) species

and the concatenated DNA dataset consisted of 941,354 sites with a gene and character

occupancy of 91.5% and 83.0%, respectively. None of our newly sequenced species

exhibited exceptional levels of missing data (<30% missing; Table S1), but three species

added from Zhao *et al.* (2021), which included *Pisum sativum* L., *Hesperothamnus*

pentaphyllus (Harms) Harms, and *Euchresta japonica* Hook. f. ex Regel, contained the

highest level of missing data (69–85% missing; Table S2).

Our comprehensive, genus-level taxon sampling represents multiple data collection

strategies, including high-quality reference genomes, RNA-seq, and low-pass genome

sequencing (i.e., genome skimming). Among these three approaches, our *in-silico* extraction

of phylogenetic markers from genomic sequencing data using PhyloHerb represents a cost-

effective approach to construct a multi-locus phylogeny, eliminating the need for target

enrichment library preparation. Genomic sequencing of the 32 Papilionoideae taxa yielded

25–30 million 150 bp paired-end reads and 1–15× nuclear coverage according to k-mer

distribution (Table S8; Data S2). In the final DNA matrix, the level of missing data in these

32 species was not statistically different from transcriptomic data (Welch two sample t-test

p-value = 0.08). Even for species with less than 2× nuclear genome coverage, at least 1,075

(74%) of the targeted loci were recovered (Table S8). Computationally, PhyloHerb is also

highly accessible. Extracting ~1,500 loci from each species on average requires 903.2s CPU

time and 2.9 GB peak memory including assembly and ortholog identification, which can be

completed on most laptops.

Multi-locus nuclear dataset resolves contentious relationships in Papilionoideae

To accommodate for incomplete lineage sorting, we used both coalescent and

concatenation methods for phylogenetic inference. The coalescent-based phylogeny

estimated by ASTRAL was largely congruent with the concatenation-based phylogeny

inferred by IQ-TREE and 92.6% and 89.1% of the nodes received maximum support (100

BP; 1.0 posterior probability, PP), respectively (Fig. 1).

Within Papilionoideae, the monophyly of these 19 clades was fully supported by both

coalescent and concatenation analyses (Fig. 1; Fig. S2). Relationships among these clades

were confidently resolved including those of recalcitrant lineages such as *Dermatophyllum,*

Sesbania, and the Andira clade (Fig. 1). Among the earliest-diverging lineages in

Papilionoideae, the ADA, Swartzieae and Cladrastis clades formed successive sister groups

to the remaining Papilionoideae (i.e., the 50-kb inversion clade, Fig. 1). Within the 50-kb

inversion clade, the Vataireoid and Exostyleae clades together formed a well-supported

clade (100 BP, 1.0 PP) sister to all other groups. The monophyly of the Vataireoid and

Exostyleae clades was strongly favored by both the AU test using the concatenated matrix

and the likelihood ratio test based on the multispecies coalescent (MSC) model. The

alternative topology suggested by Zhao *et al.* (2021), where these two groups were not

monophyletic, was rejected (AU test p-value < 0.01; MSC likelihood ratio test p-value <0.01;

Table S3).

One major conflict in our results is the relative placement of the Dalbergioid s.l. (1,580

spp.) and Genistoid s.l. (2,100 spp.) clades within the 50-kb inversion clade. A

monophyletic Dalbergioid s.l. + Genistoid s.l. clade was recovered but poorly supported by

the concatenation method (41 BP). On the other hand, the coalescent method recovered a

closer relationship between Dalbergioid s.l. and the Andira, Baphieae, and NPAAA clades

423
Figure 1 Phylogeny of Papilionoideae (continued on the next page).

with high confidence (0.98 PP; Fig. S2). The AU test on these two alternative topologies did not provide definitive conclusions (*p*-value >0.05; Table S3) and the gene-wise likelihood supporting each topology was approximately the same (Fig. S3). The MSC likelihood ratio test, however, rejected the monophyletic Dalbergioid s.l. + Genistoid s.l. clade (*p*-value = 0.03). Within the Genistoid s.l. clade, the genus *Ormosia* Jacks was strongly supported as sister to all remaining lineages of the clade (100 BP; 1.0 PP), while *Dermatophyllum* was confidently placed as an early-diverging lineage sister to Brongniartieae (88 BP; 1.0 PP). The African monotypic genus *Haplormosia* Harms was nested within the genistoids (98 BP; 0.90 PP). Within the Dalbergioid s.l. clade, all previously identified major subclades were maximally supported. The Amorpheae appeared as sister to all remaining dalbergioids, and the *Adesmia*, *Dalbergia*, and *Pterocarpus* subclades were phylogenetically successive sisters (100 BP; >=0.99 PP).

The Andira, Baphieae, and NPAAA clades were each monophyletic and they formed a grade comprising the remaining Papilionoideae. Within the NPAAA clade, the genus *Hypocalyptus* Thunb. was the earliest diverging lineage (97 BP; 0.94 PP), while the Mirbelioid, Hologalegina, and Indigofereae + Millettioid-Phaseoloid clades formed successive sisters to Hypocalypteae. Within mirbelioids, both coalescent and concatenation methods placed lineages from tribe Bossiaeeae as the earliest diverging clade sister to a poorly supported clade composed mostly of tribe Mirbelieae (<58 BP; <0.83 PP). Such low support could be

attributed to the two short internal branches (1.3–1.7e-3 in mutation units; 2.0–3.9e-2 in

coalescent units) separating Bossiaeeae and two non-monophyletic groups within

Mirbelieae — (*Daviesia* Sm. + *Viminaria* Sm.) and *Gompholobium* Sm. These rapid radiations

have rendered little phylogenetic information to reconstruct this section of the phylogeny

— the gene concordant factor was 16.3% for the branch separating Bossiaeeae and

Mirbelieae, while 14.0% of the genes supported the other two alternative quartet

topologies at this node; similarly, 16.0% of the genes supported the paraphyly of *Daviesia* +

Viminaria and *Gompholobium*, but one of the alternative topologies was supported by

17.4% of the genes (Fig. 1; Data S3). Results from the site concordance factor largely

reflected a similar trend where each of the three quartet topologies was supported by 33%

of the sites (Data S3).

The Millettioid-Phaseoloid clade within the larger NPAAA clade harbors more uncertainty.

The three genera *Aganope* Miq., *Austrosteenisia* R. Geesink and *Dalbergiella* Baker f. formed

a well-supported basal grade within the millettioids (100 BP; 1.0 PP). The concatenation

method confidently placed (100 BP) subtribe Clitoriinae—including *Centrosema* (DC.)

Benth., *Periandra* Mart. ex Benth. and *Clitoria* L.—as sister to *Dalbergiella*. In contrast, the

coalescent method placed Clitoriinae with a clade consisting of members from the

polyphyletic Phaseoleae, Desmodieae, and Psoraleeae with low confidence (0.31 PP). A

clade comprising Abreae, Diocleae, Millettieae formed the next diverging group in the

millettioids. The remaining millettioids included tribes Desmodieae, Psoraleeae, and the

polyphyletic Phaseoleae. In both coalescent and concatenation analyses, *Platycyamus*

regnellii Benth. was the earliest diverging lineage (100 BP; 1.0 PP), but conflicts exist in

terms of the placement of the *Butea* clade (genera *Spatholobus* Hassk., *Butea* Roxb. ex

Willd., *Hardenbergia* Benth., *Kennedia* Vent.), the Apios clade (genera *Mucuna* Adans. and

Apios Fabr.), and *Shuteria* Wight & Arn. Both coalescent and concatenation methods

strongly supported the *Butea* clade to be the earliest diverging among the three (100 BP;

1.0 PP), while the Apios clade and *Shuteria* were recovered as monophyletic (99 BP) in the

concatenation analysis but paraphyletic (0.89 PP) in the coalescent analysis.

Evolution of keel flowers in Fabales

The robust nuclear phylogeny was used to constrain the uncertain relationships among

major clades in the *matK*-based phylogeny. This broadly sampled phylogeny of Fabales

included 3,326 species — 3,106 Fabaceae (15.9% species diversity), 1 Quillajaceae (100%),

7 Surianaceae (55.6%), 206 Polygalaceae (15.4%), and 6 additional eudicots for dating

purposes. With fossil constraints, the time tree inferred by BEAST supported a Late

Cretaceous origin of crown group Fabales at 78.0 Ma (95% HPD 71.9–88.3 Ma; Fig. S4), and

an early Paleocene origin of crown group Fabaceae and Polygalaceae at 66.0 Ma (95% HPD

64.7–66.8 Ma) and 56.1 Ma (95% HPD 55.8–56.6 Ma), respectively. Within Fabaceae, we

inferred an explosive diversification leading to all six subfamilies within 7 million years

after the origin of the family. Crown group Papilionoideae was inferred to originate at 60.9

Ma (95% HPD 59.7–62.1 Ma; Fig. S4).

Keel flowers, which are mostly known in Papilionoideae and tribe Polygaleae

(Polygalaceae), have been independently lost and gained multiple times. Ancestral

character reconstruction suggested at least seven independent origins of keel flowers in

tribe Polygaleae (Polygalaceae), *Cercis* L. (Cercidoideae), *Angylocalyx* Taub. (ADA clade,

Papilionoideae), tribe Dipterygeae (ADA clade, Papilionoideae), the MRCA of *Petaladenium*

Ducke and *Dussia* Krug & Urb. ex Taub. (ADA clade, Papilionoideae), *Myrospermum* Jacq.

(ADA clade, Papilionoideae), and the MRCA of the Cladrastis clade and the 50-kb inversion

clade (Papilionoideae). In addition, there were two independent losses of keel flowers in

Polygalaceae and twenty-five in Papilionoideae (summarized in Table S9; Fig. S5). These

numbers, especially the losses, were strongly dependent on the phylogeny. We inferred the

ancestral floral architecture of Papilionoideae to be non-keel-like, with the first keel

flowers emerging in the common ancestor of the Cladrastis and the 50-kb inversion clades

around 59.0 Ma (95% HPD 57.9–60.2 Ma). This is more recent than the origination of keel

flowers in crown group Polygalaceae, which included *Xanthophyllum* at 62.8 Ma (95% HPD

59.6–64.9 Ma), though the exact order cannot be established with confidence due to

overlapping confidence intervals.

State-dependent diversification

The BAMM analysis inferred 18 rate shifts in Fabales and 11 of these shifts are nested

within Papilionoideae (Fig. 2a, c). The diversification rate increased gradually over time in

Fabales and nearly all rate shifts represented a significant acceleration in diversification

rates rather than a decrease (gray lines in Fig. 2b). With few exceptions, 9 out of the 11 rate

shifts in Papilionoideae took place during the Miocene, which marked a time of global

cooling and aridification (Herbert *et al.*, 2016).

Although most rate shifts took place in the keel-flowered Papilionoideae, our

diversification rate analysis did not support the SSE models. Among the four state-

dependent (BiSSE and HiSSE) and state-independent models (CID-2 and CID-4), CID-4 was

significantly better than the second-best HiSSE model with a ΔAIC of 168.772. However,

diversification parameters from both BiSSE and HiSSE models suggested an orders-of-

magnitude higher net diversification rate in keel-flowered clades than non-keel-flowered

clades. Specifically, HiSSE suggested the net diversification rate to be 0.13 in keel-flowered

clades and 5.5e-4 in non-keel-flowered clades; BiSSE suggested it to be 0.084 in keel-

flowered clades and 8.1e-04 in non-keel-flowered clades.

DISCUSSION

Insights into the early divergence relationships within papilionoids

Previous phylogenetic studies focusing on the early divergences of papilionoid legumes

have largely utilized plastid markers (Wojciechowski *et al.*, 2004; Lavin *et al.*, 2005;

McMahon & Sanderson, 2006; Cardoso *et al.*, 2012, 2013b; Zhang *et al.*, 2020; Choi *et al.*,

2022). These studies have consistently recognized well-supported clades but struggled in

resolving inter-clade relationships. Based on 1,456 low-copy nuclear genes, we show that

the ADA, Swartzieae, and Cladrastis clades are successive sister groups to the 50-kb

inversion clade (Fig. 1). This result is congruent with the recent nuclear phylogeny from

Zhao *et al.* (2021) but differs from most plastid-based phylogenies where the Swartzieae

clade was placed as sister to all other Papilionoideae (Cardoso *et al.*, 2012, 2013b; Zhang *et*

al., 2020; Choi *et al.*, 2022).

Figure 2. Rate heterogeneity in the diversification of Fabales. (a) Time-calibrated tree of
Fabales species based on plastid *matK* gene sequences. Branch colors show net
diversification rates inferred from BAMM and circles at nodes refer to clades with
significant rate shifts. Purple lines in the outer ring indicate the presence of keel flowers.
(b) Rate-through-time plot showing increased net diversification over time. The black line
represents the average across the whole tree; the gray lines represent the rate through
time on parts of the tree where significant rate shifts were identified. Shaded regions
represent 95% confidence intervals. (c) Posterior distribution of the number of rate shifts
inferred using BAMM. (d) Variation in tip-specific diversification rates for keel-flowered
and non-keel-flowered lineages. All rates are relative and only allow us to look at relative
differences between biome preferences and parts of the phylogeny. The photo by Janet
Davis shows a *Bombus griseocollis* bee on a *Baptisia australis* flower.

The 50-kb inversion clade constitutes 95% of the remaining species diversity in

Papilionoideae. Our well-resolved nuclear phylogeny provides several important insights

into relationships within this clade that were inconclusive in previous plastid sequence-

based studies. First, the Exostyleae and Vataireoid clades form a monophyletic group sister

to the remaining members of the 50-kb inversion clade (100 BP; 1.0 PP). Their close

affinity is morphologically supported by the denticulate, serrate, or occasionally spinescent

leaf margin, which is highly unusual outside the NPAAA clade in Papilionoideae (Mansano

et al., 2004; Cardoso *et al.*, 2012, 2013a,b). Second, although conflicting topologies are

obtained when placing the Genistoid s.l. clade, statistically significant support is only

achieved for the coalescent analysis (Table S3). Thus, based on the ASTRAL tree, the

Genistoid s.l. clade is sister to a clade comprised of the Dalbergioid s.l., Andira, Baphieae,

and NPAAA clades (PP = 0.98). Notably, the Dalbergioid s.l., Andira, and Baphieae clades

form successive sister groups to the large NPAAA clade — a relationship that has not been

observed in any plastid sequence-based study (Cardoso *et al.*, 2012, 2013b; Choi *et al.*,

2022).

Within the Genistoid s.l. clade, the Ormosieae, Brongniartieae, and Leptolobieae form

successive sister groups to all remaining species. Particularly, the genus *Dermatophyllum,*

comprised of four species, is a well-supported sister lineage to tribe Brongniartieae (88 BP;

1.0 PP). In previous studies, *Dermatophyllum* often formed a polytomy with the Dalbergioid

s.l., Andira, and Genistoid s.l. clades (Wojciechowski et al., 2004; Cardoso et al., 2012,

Cardoso et al., 2013). The close affinity between *Dermatophyllum* and Brongniartieae is

supported by their shared niche preference and the ability to synthesize a class of highly

distinct alkaloids. The preferred habitat of *Dermatophyllum* in warm temperate woodlands,

chaparral, and deserts of southern United States to central Mexico overlaps with many

Mexican Brongniartieae especially in the genus *Brongniartia* (Pennington *et al.*, 2005; Ross

& Crisp, 2005). The presence of quinolizidine alkaloids in *Dermatophyllum* further supports

its placement within the Genistoid s.l. clade (Cardoso *et al.*, 2013; Choi *et al.*, 2022). This

class of alkaloids is absent in all other legumes and only sparsely distributed in eight other

vascular plant families (Bisby *et al.*, 1994; Kite & Pennington, 2003; Lee *et al.*, 2013; Wink,

2013; Cely-Veloza *et al.*, 2023). Therefore, our new result provides compelling evidence for

a single origin of alkaloid biosynthesis in papilionoids and a chemical synapomorphy for

the Genistoid s.l. clade.

In addition to *Dermatophyllum*, the close relationship between the African monotypic genus

Haplormosia and the American-Australian Brongniartieae clade proposed by (Cardoso *et*

al., 2017) is also challenged by our new result. Although several reproductive and

vegetative features are shared between these two lineages — including unifoliolate leaves;

single-seeded, tardily dehiscent fruits with thick, woody valves; and stamens with distinctly

dimorphic anthers (Cardoso *et al.*, 2017) — the distinct biogeographic distribution makes

the proposed relationship contentious. Our results instead place *Haplormosia* within a

group of largely Old-World lineages, including *Camoensia*, Crotalarieae, Genisteae,

Podalyrieae, and Sophoreae (98 BP; 0.90 PP). Morphologically, species from Crotalarieae,

Podalyrieae, and Genistieae also exhibit unifoliolate leaves and dimorphic anthers that are

alternately basifixed and dorsifixed (Schutte-Vlok & van Wyk, 2011; Le Roux & Van Wyk,

2012; du Preez *et al.*, 2021).

Finally, the NPAAA clade contains a whopping ca. 10,200 species that account for nearly

70% of species level diversity in papilionoids. It is sister to the predominantly African

Baphieae clade with ca. 60 species (100 BP; 1.0 PP). The NPAAA clade comprises two large,

strongly-supported subclades, the Hologalegina (Robinioid+IR-lacking clade) and

Indigofereae + Millettioid-Phaseoloid clades, in agreement with all previous molecular

phylogenetic studies (Wojciechowski *et al.*, 2004; Cardoso *et al.*, 2013b; LPWG, 2017; Zhao

et al., 2021). Within Hologalegina (ca. 5,300 spp.), the pantropical genus *Sesbania* was

placed within the Robinioid clade. In plastid-based studies, the relationship between

Sesbania, and tribes Loteae and Robinieae is highly unstable with all combinations of

relationships recovered (Wojciechowski *et al.*, 2004; Farruggia *et al.*, 2018; Zhang *et al.*,

2020; Choi *et al.*, 2022). Our results confirm *Sesbania* as sister to the warm temperate,

primarily northern hemisphere Loteae (100 BP; 1.0 PP), and together they form a sister

group to Robinieae (100 BP; 1.0 PP). The paraphyletic relationship among *Sesbania,* Loteae,

and Robinieae proposed by Zhao *et al.* (2021) is strongly rejected by our AU test (*p*-value <

0.01) and MSC likelihood ratio test (*p*–value < 0.01). Relationships within the IR-lacking

clade (IRLC) align with recent studies using plastid and nuclear sequences (Choi *et al.*,

2022; Zhao *et al.*, 2021).

Relationships within the primarily subtropical to pantropical Indigofereae + Millettioid-

Phaseoloid clade (ca. 4,050 spp.) are consistent with earlier findings (Hu *et al.*, 2000;

Wojciechowski *et al.*, 2004; Stefanović *et al.*, 2009; LPWG, 2017; Oyebanji *et al.*, 2020; Zhao

et al., 2021; Choi *et al.*, 2022). In the Millettieae s.l. and Phaseoleae s.l. (Schrire, 2005a,b),

the "core Millettieae" (Hu *et al.*, 2002) and the three traditional tribes Abreae, Desmodieae,

and Psoraleeae (Lewis *et al.*, 2005) are each well supported monophyletic lineages. Tribes

Millettieae and Phaseoleae are polyphyletic with unresolved generic-level relationships.

Loss and gain of keel flowers during the diversification of the papilionoids

A suite of reproductive, nutritional, and defense traits has contributed to the assemblage of

present legume biodiversity (Lewis *et al.*, 2005). Here we focused on the pollinator-driven

diversification in Fabales, which has been widely invoked to explain the radiation of many

angiosperms (Sargent, 2004; Kay & Sargent, 2009; Givnish *et al.*, 2015). Bilaterally

symmetrical flowers such as a keel flower are strongly associated with bees by restricting

the entry to and the precise placement of pollen to discourage gene flow between incipient

species (Neal *et al.*, 1998). Despite being a stereotyped search image for legumes, there is

great lability in floral form and symmetry during the early evolution history of papilionoids

(Fig. 2). This provides an ideal comparative framework to test whether the evolution of

keel flowers drives diversification (Ricklefs & Renner, 1994, 2000).

Our well-resolved nuclear gene and densely-sampled plastid *matK* gene phylogenies

provide the essential evolutionary framework to address this question. Crown group

Fabaceae was estimated to originate at the K-Pg boundary at 66.0 Ma (95% HPD 64.7–66.8

Ma), which is in line with previous molecular-based estimates and fossil records (Lavin *et*

al., 2005; Bruneau *et al.*, 2008; Cannon *et al.*, 2015; Koenen *et al.*, 2021; Zhao *et al.*, 2021).

The ancestral floral architecture of Papilionoideae is estimated to be non-keel-like, with the

first keel flowers in Fabaceae emerging in the common ancestor of the Cladrastis and the

50-kb inversion clades at 59.0 Ma (95% HPD 57.9–60.2 Ma). This estimation predates the

earliest papilionoid-like flower fossil *Barnebyanthus buchananensis* by 3 million years

(Herendeen & Crepet, 1992) and closely succeeded the evolution of keel flowers in crown

group Polygalaceae at 62.8 Ma (95% HPD 59.6–64.9 Ma). However, these estimates

contradict the results of a recent study which concluded that papilionoid keel flowers

evolved 22.03 to 10.32 Ma before they occurred in Polygalaceae (Uluer *et al.*, 2022).

Such a discrepancy comes mainly from the ancestral state reconstruction analysis of keel

flowers themselves, but also from minor differences in divergence time estimation due to

taxon sampling and methodology. Unlike the crown group Papilionoideae origination

proposed by Uluer *et al.* (2022), our results suggest a more dynamic evolutionary history of

keel flowers with five independent origins that postdate crown group Papilionoideae

(Table S9; Fig. S5). Four of these events are nested within the ADA clade, taking place in the

Miocene of tropical Africa (e.g., *Angylocalyx*) and America (e.g., *Myrospermum*). The two

oldest keel flower lineages originated during the Selandian age of the Paleocene in

Papilionoideae and Polygalaceae, though the exact order cannot be confidently established.

However, the short interval between these two old events and multiple more recent origins

of keel flowers in the ADA clades may all result from the same mutualistic process where

younger lineages benefitted from and contributed to the strong relationship with

pollinators already established by older keel lineages (Tucker, 2002; Bello *et al.*, 2012;

Uluer *et al.*, 2022).

In addition to gains, perhaps what is more intriguing is the numerous losses of keel flowers.

We identified more than two dozen independent losses in Papilionoideae alone (Table S9).

Only one out of these 27 losses is nested within the NPAAA clade in genus *Erythrina* L.,

even though the NPAAA clade accounts for more than 70% of the Fabaceae species. These

results suggest that ancestral Fabaceae likely took a generalist pollination strategy,

allowing floral symmetry to be more labile. This is exemplified in multiple early branching

papilionoids with lost, reduced, or poorly differentiated petals. These modifications are

accompanied by expansion or shifts of pollinators to bats, passerine birds, hummingbirds,

and even beetles (e.g., Arroyo, 1981; Pennington *et al.*, 2000; Lewis *et al.*, 2003). In

contrast, derived lineages may have experienced stronger stabilizing selection to maintain

specialized floral architecture for coevolved pollinators. The NPAAA clade has diversified

largely during the Miocene under global cooling, aridification, and expansion of grasslands

(Estep *et al.*, 2014; Herbert *et al.*, 2016). The IRLC, Indigofereae, and Robinioid clades are

thus strongly associated with herbaceous growth habit in arid to semi-arid temperate

regions (Wojciechowski *et al.*, 2004; Schrire *et al.*, 2005, 2009). Adaptation to these

habitats may cast additional geographical, environmental, and developmental constraints

to limit the breadth of viable pollinators, making it advantageous to maintain ancestral

states rather than evolve new ones.

Finally, although the HiSSE model supports a higher net diversification rate in keel-

flowered lineages, the best-supported model was one where the diversification rate was

independent of floral architecture. The orders-of-magnitude difference in net

diversification rate estimated by the HiSSE model is driven by handful keel-flowered

lineages (Fig. 2d). Not all keel-flowered Fabaceae radiated extensively (e.g., *Cercis*), and

Swartzia (Swartzieae) diversified rapidly without keel flowers. Along these lines, Afkhami

et al. (2018) also reported a lack of correlation between the nodulation ability and net

diversification rate despite that nodulating legume genera contain three times more

species than non-nodulating ones. The lack of support for SSE models suggests that

individual traits including keel flowers and nodulation are not the sole driver of

diversification, but instead are one of several diversification-promoting traits within a

broader confluence of factors that have additively contributed to lineage diversification

(Donoghue & Sanderson, 2015). Besides innovations of key traits, recurrent biome shifts

and rapid colonization of novel ecological niches also played a significant role in promoting

diversification in Fabaceae. Examples include the rapid radiation of the primarily

herbaceous IRLC clade in temperate regions (Wojciechowski *et al.*, 2004), the dominance of

mimosoids in seasonally dry forests (Pennington *et al.*, 2009; Ringelberg *et al.*, 2023), and

the remarkable diversification of Indigofereae in tropical fire-prone savannas (Schrire *et*

al., 2009). In summary, while pollinator interactions have played a role in the

diversification of papilionoid legumes, a combination of intrinsic factors (e.g., chemical

defenses and nodulation) and extrinsic factors (e.g., biome shifts and ecological

specialization) were synergistically involved to drive the remarkable diversification of
Papilionoideae.
722

CONCLUSIONS
Drawing insights from phylogenomics, divergence time estimation, and comparative
methods, we present novel insights into the early divergence history and floral evolution of
papilionoid legumes. With 1,456 low-copy nuclear loci, we confidently resolved the
recalcitrant relationships in Papilionoideae and inferred a Paleocene origin of keel flowers
at 59 Ma, predating the earliest fossil record by 3–4 million years. We identified at least
seven independent origins and 27 losses of keel flowers across Fabales, with losses
predominantly taking place in early diverging papilionoid lineages. Ancestral Fabaceae
thus likely displayed a more generalist pollination strategy, allowing for greater floral
flexibility over time. Although these independent gains and losses coincide with the
Miocene diversification of Papilionoideae, our results do not support floral architecture as
the sole driver of diversification in Fabaceae. Instead, a confluence of intrinsic and extrinsic
factors including nitrogen fixation, chemical defenses, and recurrent dispersal into new
biomes such as subtropical and temperate regions, likely contributed synergistically to the
diversification of papilionoids.
738

ACKNOWLEDGEMENTS
This work was supported by grants from the National Science Foundation to MFW (DEB-
1853010) and RKJ, TAR (DEB-1853024), the Texas Ecological Laboratory Program
(EcoLab) to RKJ, TAR, I-SC, and LC, the Sidney F. and Doris Blake Professorship in
Systematic Botany to RKJ, the Stengl Wyer Fellowship to LC, the CNPq (Research
Productivity Fellowship no. 314187/2021-9), and FAPERJ (Programa Jovem Cientista do
Nosso Estado - 2022, grant no. 200.153/2023) to DC, the National Research Foundation of
Korea (NRF) grant funded by the Korea government (MSIT) (RS-2024-00341022) to I-SC,
and funds from the College of Liberal Arts and Sciences at Arizona State University to MFW.
We thank the TEX-LL, HUEFS, and RB herbaria for voucher deposition, the Desert Legume
Program at the University of Arizona for seeds; Alessandra Schnadelbach, Henrique Batalha
Filho, and Paula Ristow for permitting us to use their labs at UFBA; and George Yatskievych

(TEX/LL) for arranging a formal Material Transfer Agreement (Decree number 8772)

under the SisGen Cadastro RDC6BE9, which facilitated research activities between the

institutions.

SUPPLEMENTARY INFORMATION

Supplementary figures

Figure S1 Histogram of the number of species per loci. The vertical line represents the

30% missing data threshold to select the 1,456 loci in this study.

Figure S2 Topological congruences and conflicts of Papilionoideae phylogeny inferred

from IQTREE (left) and ASTRAL (right). Tangled grey lines indicate conflicting

relationships. Branch supports represent nonparametric bootstrap replicates values from

IQ-TREE and local posterior probability from ASTRAL, respectively. Only branches that are

not maximally supported are labeled.

Figure S3 Quantification of phylogenetic signal from individual genes. The two alternative

placements of the focal clades are illustrated with red or blue, with the red topology being

favored by the concatenation analysis. Genes are ordered in descending order based on

their support for the red topology. Differences in the log-likelihood scores (ΔlogL) between

these two topologies are calculated for individual genes (left) or the concatenated matrix

(right). Panels A-E represent five contentious relationships within Papilionoideae defined

in Table S3.

Figure S4 Bayesian inference of divergence time in Fabaceae based on 30 top clock-like

loci. Divergence time is shown in millions of years ago (Ma) and the blue bars represent

95% credibility intervals. Eleven fossil calibration points including a constraint for the root

age are marked by numbered stars. See Table S5 for a full list of fossils and references.

Figure S5 Ancestral character reconstruction of keel flowers in Fabales inferred under the

marginal reconstruction algorithm in HiSSE. An ultrametric tree of 3,326 Fabales species is

inferred from matK sequences with branch lengths drawn in proportion to divergence

time. Solid and hollow branches indicate the presence and absence of keel flowers, respectively. Events of keel flower gains and losses are marked by orange and magenta circles on the branch, respectively. See Table S9 for age estimations of these events.

Supplementary Tables

Table S1 Voucher information and gene assembly information of 79 newly sequenced species in this study. Species name, lineage affiliation, assembly quality, SRA accession number, and voucher information including collection locality are provided.

Table S2 Species name, lineage affiliation, and number of sampled genes from 208 species from Zhao et al. (2021).

Table S3 Testing alternative phylogenetic placement of Papilionoideae clades under the maximum likelihood framework for concatenated DNA sequences.

Table S4 List of top 30 clock-like genes with quantified root-to-tip branch length variation, total tree length, and topological similarity to the species tree measured by bipartition score.

Table S5 Fossil calibration points for divergence time estimation.

Table S6 Time tree, Genbank ID, and keel flower trait status for the 3,326 Fabales species from the *matK* dataset. Numbers on the phylogeny are divergence time estimation for each node based on treePL.

Table S7 Secondary calibration points for the treePL analysis. All node ages in million years ago (Ma) are derived from the BEAST time tree and are fixed in treePL. Nodes are defined by the most recent common ancestor of two species indicated in the spreadsheet.

Table S8 Nuclear genome sequencing coverage of 32 whole genome sequencing datasets

estimated based on 21-mer frequency distribution. See Table S1 for the species name of

each ID.

Table S9 Summary of independent origins and losses of keel flower in Fabales. Binary

ancestral state reconstruction is inferred under the marginal reconstruction algorithm in

HiSSE. The timing of these events is determined by the crown group age of involved clades.

The precise age of events involving a single species cannot be determined using available

data. Therefore the age of these events is represented by a range from present to their stem

group age.

Supplementary Data

Data S1 DNA alignments and maximum likelihood phylogenies inferred for the 1,456 low-

copy nuclear genes used in this study. The alignments are inferred using the MAFFT-linsi

algorithm. The gene trees are reconstructed using IQTREE and branch support is evaluated

with 1000 ultrafast bootstrap replication. Species names and their corresponding ID are

provided in sp_ID.csv. The list of 672 loci used to generate the concatenated matrix for

species tree estimation is provided in G672_len900_sp227.list.

Data S2 Distribution of 21-mer frequency in 32 species with genome skimming data. The

21-mer frequency is inferred from Jellyfish v2.3.0 to estimate sequencing coverage.

Data S3 Gene and site concordance factors of the Papilionoideae phylogeny. The species

tree and concatenated alignment of the 672-gene dataset were used as inputs in IQ-TREE to

calculate these concordance factors.

COMPETING INTERESTS

None declared.

AUTHOR CONTRIBUTIONS

Conception and experimental design, LC, DC, I-SC, RKJ, TAR, MFW. Acquisition of funds, RKJ,

TAR, MFW. Field collections, I-SC, DC, HCL, LPQ, RKJ, TAR, MFW. Nucleic acid isolation, I-SC,

CL, TAR. RNA isolations, TAR. Transcriptome and genome assembly, CL, LT, BS, LC. DNA

alignments and phylogenetic analyses, LC, I-SC, DC, MFW. Data interpretation, LC, I-SC, DC,

MFW. Production of figures and tables, LC, DC. Writing the initial draft of the manuscript,

LC, DC, MFW, TAR, RKJ. All authors revised and approved the final version of the

manuscript.

DATA AVAILABILITY

The raw sequence data generated and analyzed during this study are available in the NCBI

GenBank repository under Bioproject PRJNA1120003. The alignment data supporting the

findings of this study are available as supplementary data and have been deposited in the

Figshare Digital Repository (DOI: 10.6084/m9.figshare.27135984). The code used for data

processing and analysis is openly available on GitHub at

https://github.com/lmcai/Papilionoideae_phylogenomics.

REFERENCES

Afkhami ME, Luke Mahler D, Burns JH, Weber MG, Wojciechowski MF, Sprent J,
Strauss SY. **2018**. Symbioses with nitrogen-fixing bacteria: nodulation and phylogenetic
data across legume genera. *Ecology* **99**: 502–502.

Arroyo MTK. **1981**. Breeding systems and pollination biology. In: R. M. Polhill and P. H.
Raven, Ed., Advances in Legume Systematics, Part 2. Richmond, UK: Royal Botanic Garden,
Kew, 723–769.

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,
Nikolenko SI, Pham S, Prjibelski AD, *et al.* **2012**. SPAdes: a new genome assembly
algorithm and its applications to single-cell sequencing. *Journal of Computational Biology*
19: 455–477.

Barrett CF, Specht CD, Leebens-Mack J, Stevenson DW, Zomlefer WB, Davis JI. **2014**.
Resolving ancient radiations: can complete plastid gene sets elucidate deep relationships
among the tropical gingers (Zingiberales)? *Annals of Botany* **113**: 119.

Beaulieu JM, O'Meara BC. **2016**. Detecting hidden diversification shifts in models of trait-
dependent speciation and extinction. *Systematic Biology* **65**: 583–601.

Bell EA. **1981**. Non-protein amino acids in the Leguminosae. In: Advances in Legume Systematics, Part 2, eds Polhill R. M., Raven P. H. Richmond, UK: Royal Botanic Gardens, Kew, 489–499.

Bello MA, Rudall PJ, Hawkins JA. **2012**. Combined phylogenetic analyses reveal interfamilial relationships and patterns of floral evolution in the eudicot order Fabales. *Cladistics* **28**: 393–421.

Bisby FA, Buckingham J, harborne JB. **1994**. *Phytochemical Dictionary of the Leguminosae*. London: Chapman and Hall.

Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, Suchard MA, Rambaut A, Drummond AJ. **2014**. BEAST 2: a software platform for Bayesian evolutionary analysis. *PLOS Computational Biology* **10**: e1003537.

Bruneau A, Mercure M, Lewis GP, Herendeen PS. **2008**. Phylogenetic patterns and diversification in the caesalpinioid legumes. *Botany* **86**: 697–718.

Cai L, Zhang H, Davis CC. **2022**. PhyloHerb: a high-throughput phylogenomic pipeline for processing genome skimming data. *Applications in Plant Sciences* **10**: e11475.

Cannon SB, McKain MR, Harkess A, Nelson MN, Dash S, Deyholos MK, Peng Y, Joyce B, Stewart CN, Rolf M. **2015**. Multiple polyploidy events in the early radiation of nodulating and nonnodulating legumes. *Molecular Biology and Evolution* **32**: 193–210.

Cardoso D, Harris DJ, Wieringa JJ, São-Mateus WMB, Batalha-Filho H, Torke BM, Prenner G, Queiroz LP. **2017**. A molecular-dated phylogeny and biogeography of the monotypic legume genus *Haplormosia*, a missing African branch of the otherwise American-Australian Brongniartieae clade. *Molecular Phylogenetics and Evolution* **107**: 431–442.

Cardoso D, Paganucci Queiroz L, Cavalcante Lima H, Suganuma E, van den Berg C, Lavin M. **2013a**. A molecular phylogeny of the vataireoid legumes underscores floral evolvability that is general to many early-branching papilionoid lineages. *American Journal of Botany* **100**: 403–421.

Cardoso D, Pennington RT, Queiroz LP, Boatwright JS, Van Wyk B-E, Wojciechowski MF, Lavin M. **2013b**. Reconstructing the deep-branching relationships of the papilionoid legumes. *South African Journal of Botany* **89**: 58–75.

Cardoso D, Queiroz LP, Pennington RT, Lima HC, Fonty É, Wojciechowski MF, Lavin M. **2012**. Revisiting the phylogeny of papilionoid legumes: new insights from comprehensively sampled early-branching lineages. *American Journal of Botany* **99**: 1991–2013.

Cardoso D, São-Mateus WMB, Cruz DT, Zartman CE, Komura DL, Kite G, Prenner G, Wieringa JJ, Clark A, Lewis G, *et al.* **2015**. Filling in the gaps of the papilionoid legume

phylogeny: The enigmatic Amazonian genus *Petaladenium* is a new branch of the early-diverging Amburaneae clade. *Molecular Phylogenetics and Evolution* **84**: 112–124.

Cely-Veloza W, Kato MJ, Coy-Barrera E. **2023**. Quinolizidine-type alkaloids: chemodiversity, occurrence, and bioactivity. *ACS Omega* **8**: 27862–27893.

Choi I-S, Cardoso D, Queiroz LP, Lima HC, Lee C, Ruhlman TA, Jansen RK, Wojciechowski MF. **2022**. Highly resolved papilionoid legume phylogeny based on plastid phylogenomics. *Frontiers in Plant Science* **13**.

Chomicki G, Renner SS. **2015**. Phylogenetics and molecular clocks reveal the repeated evolution of ant-plants after the late Miocene in Africa and the early Miocene in Australasia and the Neotropics. *New Phytologist* **207**: 411–424.

Donoghue MJ, Sanderson MJ. **2015**. Confluence, synnovation, and depauperons in plant diversification. *New Phytologist* **207**: 260–274.

Doyle JJ. **2016**. Chasing unicorns: nodulation origins and the paradox of novelty. *American Journal of Botany* **103**: 1865–1868.

Estep MC, McKain MR, Diaz DV, Zhong J, Hodge JG, Hodkinson TR, Layton DJ, Malcomber ST, Pasquet R, Kellogg EA. **2014**. Allopolyploidy, diversification, and the Miocene grassland expansion. *Proceedings of the National Academy of Sciences of the United States of America* **111**: 15149–15154.

Etcheverry AV, Alemán MM, Figueroa-Fleming T, López-Spahr D, Gómez CA, Yáñez C, Figueroa-Castro DM, Ortega-Baes P. **2012**. Pollen:ovule ratio and its relationship with other floral traits in Papilionoideae (Leguminosae): an evaluation with Argentine species. *Plant Biology* **14**: 171–178.

Faria SM, Ringelberg JJ, Gross E, Koenen EJM, Cardoso D, Ametsitsi GKD, Akomatey J, Maluk M, Tak N, Gehlot HS, *et al.* 2022. The innovation of the symbiosome has enhanced the evolutionary stability of nitrogen fixation in legumes. *New Phytologist* **235**: 2365–2377.

Farruggia FT, Lavin M, Wojciechowski MF. **2018**. Phylogenetic systematics and biogeography of the pantropical genus *Sesbania* (Leguminosae). *Systematic Botany* **43**: 414–429.

Givnish TJ. **2010**. Ecology of plant speciation. *Taxon* **59**: 1326–1366.

Givnish TJ, Spalink D, Ames M, Lyon SP, Hunter SJ, Zuluaga A, Iles WJ, Clements MA, Arroyo MT, Leebens-Mack J. **2015**. Orchid phylogenomics and multiple drivers of their extraordinary diversification. *Proceedings of the Royal Society B: Biological Sciences* **282**: 20151553.

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
Raychowdhury R, Zeng Q. **2011**. Trinity: reconstructing a full-length transcriptome
without a genome from RNA-Seq data. *Nature Biotechnology* **29**: 644.

Grant V. **1949**. Pollination systems as isolating mechanisms in angiosperms. *Evolution* **3**:
82–97.

Gregório BS, Carvalho CS, Ramos G, Rocha L, Stirton CH, Lima HC, Zartman CE, Lewis
GP, Torke BM, Snak C, et al. 2024. A molecular phylogeny of the early-branching
Genistoid lineages of papilionoid legumes reveals a new Amazonian genus segregated from
Clathrotropis. *Botanical Journal of the Linnean Society* **205**: 1–14.

Herbert TD, Lawrence KT, Tzanova A, Peterson LC, Caballero-Gill R, Kelly CS. **2016**.
Late Miocene global cooling and the rise of modern ecosystems. *Nature Geoscience* **9**: 843–
847.

Herendeen PS, Crepet WL. **1992**. Paleobotanical and biogeographic history of the
legumes (Leguminosae), an important component in Cenozoic and modern tropical
terrestrial ecosystems. *The Paleontological Society Special Publications* **6**: 127–127.

Hu J-M, Lavin M, Wojciechowski MF, Sanderson MJ. **2000**. Phylogenetic systematics of
the tribe Millettieae (Leguminosae) based on chloroplast *trnK/matK* sequences and its
implications for evolutionary patterns in Papilionoideae. *American Journal of Botany* **87**:
418–430.

Hughes CE, Queiroz LP, Lewis GP. 2022. Advances in Legume Systematics 14.
Classification of Caesalpinioideae Part 1: new generic delimitations. *PhytoKeys* **205**: 1–2.

Johnson SD, Schiestl FP. **2016**. *Floral Mimicry*. Oxford, New York: Oxford University Press.

Johnson LS, Eddy SR, Portugaly E. **2010**. Hidden Markov model speed heuristic and
iterative HMM search procedure. *BMC Bioinformatics* **11**: 431.

Junier T, Zdobnov EM. **2010**. The Newick utilities: high-throughput phylogenetic tree
processing in the UNIX shell. *Bioinformatics* **26**: 1669–1670.

Katoh K, Standley DM. **2013**. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular Biology and Evolution* **30**: 772–780.

Kay KM, Sargent RD. 2009. The role of animal pollination in plant speciation: integrating
ecology, geography, and genetics. *Annual Review of Ecology, Evolution, and Systematics* **40**:
637–656.

Kite GC, Pennington RT. **2003**. Quinolizidine alkaloid status of *Styphnolobium* and
Cladrastis (Leguminosae). *Biochemical Systematics and Ecology* **31**: 1409–1416.

Koenen EJM, Ojeda DI, Bakker FT, Wieringa JJ, Kidner C, Hardy OJ, Pennington RT,
Herendeen PS, Bruneau A, Hughes CE. **2021**. The origin of the legumes is a complex
paleopolyploid phylogenomic tangle closely associated with the cretaceous–paleogene (K–
Pg) mass extinction event. *Systematic Biology* **70**: 508–526.

Kopylova E, Noé L, Touzet H. **2012**. SortMeRNA: fast and accurate filtering of ribosomal
RNAs in metatranscriptomic data. *Bioinformatics* **28**: 3211–3217.

Kursar TA, Dexter KG, Lokvam J, Pennington RT, Richardson JE, Weber MG,
Murakami ET, Drake C, McGregor R, Coley PD. **2009**. The evolution of antiherbivore
defenses and their contribution to species coexistence in the tropical tree genus *Inga*.
Proceedings of the National Academy of Sciences of the United States of America **106**: 18073–
18078.

Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B. **2017**. PartitionFinder 2: new
methods for selecting partitioned models of evolution for molecular and morphological
phylogenetic analyses. *Molecular Biology and Evolution* **34**: 772–773.

Langmead B, Salzberg SL. **2012**. Fast gapped-read alignment with Bowtie 2. *Nature*
Methods **9**: 357–359.

Lavin M, Herendeen PS, Wojciechowski MF. **2005**. Evolutionary rates analysis of
Leguminosae implicates a rapid diversification of lineages during the tertiary. *Systematic*
Biology **54**: 575–594.

Le Roux MM, Van Wyk BE. **2012**. The systematic value of flower structure in *Crotalaria*
and related genera of the tribe Crotalarieae (Fabaceae). *Flora* **207**: 414–426.

Lee C, Choi I-S, Cardoso D, Lima HC, Queiroz LP, Wojciechowski MF, Jansen RK,
Ruhlman TA. 2021. The chicken or the egg? Plastome evolution and an independent loss
of the inverted repeat in papilionoid legumes. *The Plant Journal* **107**: 861–875.

Lee ST, Cook D, Molyneux RJ, Davis TZ, Gardner DR. **2013**. Alkaloid profiles of
Dermatophyllum arizonicum, *Dermatophyllum gypsophilum*, *Dermatophyllum secundiflorum*,
Styphnolobium affine, and *Styphnolobium japonicum* previously classified as *Sophora*
species. *Biochemical Systematics and Ecology* **49**: 87–93.

Lewis GP, Knudsen JT, Klitgaard BB, Pennington RT. **2003**. The floral scent of
Cyathostegia mathewsii (Leguminosae, Papilionoideae) and preliminary observations on
reproductive biology. *Biochemical Systematics and Ecology* **31**: 951–962.

Lewis G, Schrire B, Mackinder B, Lock M. **2005**. *Legumes of the World*. Richmond, UK:
Royal Botanic Gardens, Kew.

Liu L, Anderson C, Pearl D, Edwards SV. **2019**. Modern phylogenomics: building
phylogenetic trees using the multispecies coalescent model. In: Evolutionary Genomics.
Springer, 211–239.

Liu L, Yu L. **2010**. Phybase: an R package for species tree analysis. *Bioinformatics* **26**: 962–
963.

Liu L, Yu L, Edwards SV. **2010**. A maximum pseudo-likelihood approach for estimating
species trees under the coalescent model. *BMC Evolutionary Biology* **10**: 302.

LPWG. **2013**. Legume phylogeny and classification in the 21st century: progress, prospects
and lessons for other species–rich clades. *Taxon* **62**: 217–248.

LPWG. **2017**. A new subfamily classification of the Leguminosae based on a taxonomically
comprehensive phylogeny. *Taxon* **66**: 44–77.

**Lyson TR, Miller IM, Bercovici AD, Weissenburger K, Fuentes AJ, Clyde WC, Hagadorn
JW, Butrim MJ, Johnson KR, Fleming RF**, ***et al.*** **2019**. Exceptional continental record of
biotic recovery after the Cretaceous–Paleogene mass extinction. *Science* **366**: 977–983.

Maddison WP, Midford PE, Otto SP. **2007**. Estimating a binary character's effect on
speciation and extinction. *Systematic Biology* **56**: 701–710.

Magallón S, Gómez-Acevedo S, Sánchez-Reyes LL, Hernández-Hernández T. **2015**. A
metacalibrated time-tree documents the early rise of flowering plant phylogenetic
diversity. *New Phytologist* **207**: 437–453.

Mansano VF, Bittrich V, Azevedo Tozzi AMG, Souza AP. **2004**. Composition of the
Lecointea clade (Leguminosae, Papilionoideae, Swartzieae), a re-evaluation based on
combined evidence from morphology and molecular data. *Taxon* **53**: 1007–1018.

Marazzi B, Gonzalez AM, Delgado-Salinas A, Luckow MA, Ringelberg JJ, Hughes CE.
2019. Extrafloral nectaries in Leguminosae: phylogenetic distribution, morphological
diversity and evolution. *Australian Systematic Botany* **32**: 409–458.

Marazzi B, Sanderson MJ. **2010**. Large-scale patterns of diversification in the widespread
legume genus Senna and the evolutionary role of extrafloral nectaries. *Evolution* **64**: 3570–
3592.

Marçais G, Kingsford C. **2011**. A fast, lock-free approach for efficient parallel counting of
occurrences of k-mers. *Bioinformatics* **27**: 764–770.

Maron JL, Agrawal AA, Schemske DW. **2019**. Plant–herbivore coevolution and plant
speciation. *Ecology* **100**: e02704.

McMahon MM, Sanderson MJ. **2006**. Phylogenetic supermatrix analysis of GenBank
sequences from 2228 papilionoid legumes. *Systematic Biology* **55**: 818–836.

**Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A,
Lanfear R**. **2020**. IQ-TREE 2: New models and efficient methods for phylogenetic inference
in the genomic era. *Molecular Biology and Evolution* **37**: 1530–1534.

Morales-Briones DF, Kadereit G, Tefarikis DT, Moore MJ, Smith SA, Brockington SF,
Timoneda A, Yim WC, Cushman JC, Yang Y. **2021**. Disentangling sources of gene tree
discordance in phylogenomic data sets: testing ancient hybridizations in Amaranthaceae s.l.
Systematic Biology **70**: 219–235.

Muchhala N. **2006**. The pollination biology of *Burmeistera* (Campanulaceae): specialization
and syndromes. *American Journal of Botany* **93**: 1081–1089.

Neal PR, Dafni A, Giurfa M. **1998**. Floral symmetry and its role in plant-pollinator
systems: terminology, distribution, and hypotheses. *Annual Review of Ecology, Evolution,*
and Systematics **29**: 345–373.

Oyebanji O, Zhang R, Chen S-Y, Yi T-S. **2020**. New insights into the plastome evolution of
the Millettioid/Phaseoloid Clade (Papilionoideae, Leguminosae). *Frontiers in Plant Science*
11.

Padfield D, Kay S, Vos R, Quince C, Vos M. **2024**. Macroevolutionary dynamics in micro-
organisms: generalists give rise to specialists across biomes in the ubiquitous bacterial
phylum Myxococcota. *Molecular Biology and Evolution* **41**: msae088.

Pennington R, Klitgaard B, Ireland H, Lavin M. **2000**. New insights into floral evolution
of basal Papilionoideae from molecular phylogenies. In: P.S. Herendeen and A. Bruneau
(Eds). Advances in Legume Systematics 9. Royal Botanic Gardens, Kew. In: 233–248.

Pennington RT, Lavin M, Oliveira-Filho A. **2009**. Woody plant diversity, evolution, and
ecology in the tropics: perspectives from seasonally dry tropical forests. *Annual Review of*
Ecology, Evolution, and Systematics **40**: 437–457.

Pennington RT, Stirton CH, Schrire BD. **2005**. Tribe Sophoreae. In: Lewis, G., Schrire, B.,
Mackinder, B., Lock,M. (Eds.), Legumes of the World. In: Richmond, UK: Royal Botanic
Gardens, Kew, 227–249.

Polhill RM, Raven PH, Stirton CH. **1981**. Evolution and systematics of the Leguminosae.
In: Polhill, R.M. and Raven, P.H., Eds., Advances in Legume Systematics, Part I. Richmond,
UK: Royal Botanic Garden, Kew, 1–26.

Preez B, Dreyer LL, Stirton CH, Muasya AM. **2021**. A monograph of the genus *Polhillia*
(Genisteae: Fabaceae). *South African Journal of Botany* **138**: 156–183.

Rabosky DL. **2014**. Automatic detection of key innovations, rate shifts, and diversity-
dependence on phylogenetic trees. *PLOS ONE* **9**: e89543.

Rabosky DL, Grundler M, Anderson C, Title P, Shi JJ, Brown JW, Huang H, Larson JG.
2014. BAMM tools: an R package for the analysis of evolutionary dynamics on phylogenetic
trees. *Methods in Ecology and Evolution* **5**: 701–707.

Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. **2018**. posterior summarization
in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* **67**: 901–904.

Ree RH. **2005**. Phylogeny and the evolution of floral diversity in *Pedicularis*
(Orobanchaceae). *International Journal of Plant Sciences* **166**: 595–613.

Ricklefs RE, Renner SS. **1994**. Species richness within families of flowering plants.
Evolution **48**: 1619–1636.

Ricklefs RE, Renner SS. **2000**. Evolutionary flexibility and flowering plant familial
diversity: a comment on Dodd, Silvertown, and Chase. *Evolution* **54**: 1061–1065.

Ringelberg JJ, Koenen EJM, Sauter B, Aebli A, Rando JG, Iganci JR, Queiroz LP, Murphy
DJ, Gaudeul M, Bruneau A, et al. 2023. Precipitation is the main axis of tropical plant
phylogenetic turnover across space and time. *Science Advances* **9**: eade4954.

Rosas-Guerrero V, Aguilar R, Martén-Rodríguez S, Ashworth L, Lopezaraiza-Mikel M,
Bastida JM, Quesada M. **2014**. A quantitative review of pollination syndromes: do floral
traits predict effective pollinators? *Ecology Letters* **17**: 388–400.

Ross JH, Crisp MD. **2005**. Brongniartieae. In: Lewis, G., Schrire, B., Mackinder, B., Lock, M.
(Eds.), Legumes of the World. In: Richmond, UK: Royal Botanic Gardens, Kew, 52–259.

Sargent RD. **2004**. Floral symmetry affects speciation rates in angiosperms. *Proceedings of*
the Royal Society of London. Series B: Biological Sciences **271**: 603–608.

Schrire BD. **2005a**. Tribe Millettieae. In: Lewis, G., Schrire, B., Mackinder, B., Lock,M. (Eds.),
Legumes of the World. In: Richmond, UK: Royal Botanic Gardens, Kew, 367–387.

Schrire BD. **2005b**. Tribe Phaseoleae. In: Lewis, G., Schrire, B., Mackinder, B., Lock,M.
(Eds.), Legumes of the World. In: Richmond, UK: Royal Botanic Gardens, Kew, 393–431.

Schrire BD, Lavin M, Barker NP, Forest F. **2009**. Phylogeny of the tribe Indigofereae
(Leguminosae–Papilionoideae): geographically structured more in succulent-rich and
temperate settings than in grass-rich environments. *American Journal of Botany* **96**: 816–
852.

Schrire B, Lavin M, Lewis G, Friis I, Balslev H. **2005**. Global distribution patterns of the
Leguminosae: insights from recent phylogenies: insights from recent phylogenies.
Biologiske Skrifter **55**: 375–422.

Schutte-Vlok AL, van Wyk B-E. **2011**. A taxonomic revision of *Podalyria* (Fabaceae).
Systematic Botany **36**: 631–660.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. **2015**. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics **31**: 3210–3212.

Smith SA, Brown JW, Walker JF. **2018**. So many genes, so little time: a practical approach
to divergence-time estimation in the genomic era. *PLOS ONE* **13**: e0197433.

Smith SA, O'Meara BC. **2012**. treePL: divergence time estimation using penalized
likelihood for large phylogenies. *Bioinformatics* **28**: 2689–2690.

Sprent JI. **2001**. *Nodulation in legumes*. London: Royal Botanic Gardens, Kew.

Stebbins GL. **1970**. Adaptive radiation of reproductive characteristics in angiosperms, I:
Pollination mechanisms. *Annual Review of Ecology and Systematics* **1**: 307–326.

Stefanović S, Pfeil BE, Palmer JD, Doyle JJ. **2009**. Relationships among Phaseoloid
legumes based on sequences from eight chloroplast regions. *Systematic Botany* **34**: 115–
128.

Stewart AB, Diller C, Dudash MR, Fenster CB. **2022**. Pollination-precision hypothesis:
support from native honey bees and nectar bats. *New Phytologist* **235**: 1629–1640.

Thompson JD, Gibson TJ, Higgins DG. **2003**. Multiple sequence alignment using ClustalW
and ClustalX. *Current Protocols in Bioinformatics*: **2.3**: 2.3.1–2.3.22.

Torke BM, Schaal BA. **2008**. Molecular phylogenetics of the species-rich neotropical genus
Swartzia (Leguminosae, Papilionoideae) and related genera of the swartzioid clade.
American Journal of Botany **95**: 215–228.

Tucker SC. **2002**. Floral ontogeny of Cercis (Leguminosae: Caesalpinioideae: Cercideae):
Does it show convergence with Papilionoids? *International Journal of Plant Sciences* **163**:
75–87.

Uluer DA, Forest F, Armbruster S, Hawkins JA. **2022**. Reconstructing an historical
pollination syndrome: keel flowers. *BMC Ecology and Evolution* **22**: 45.

Van der Niet T, Peakall R, Johnson SD. **2014**. Pollinator-driven ecological speciation in
plants: new evidence and future perspectives. *Annals of Botany* **113**: 199–212.

Vargas P, Zardoya R. **2014**. *The Tree of Life: Evolution and Classification of Living
Organisms*. Sunderland MA: Sinauer Associates.

Werner GDA, Cornwell WK, Cornelissen JHC, Kiers ET. **2015**. Evolutionary signals of
symbiotic persistence in the legume–rhizobia mutualism. *Proceedings of the National
Academy of Sciences of the United States of America* **112**: 10262–10269.

Westerkamp C. **1997**. Keel blossoms: bee flowers with adaptations against bees. *Flora*
192: 125–132.

Wink M. **2013**. Evolution of secondary metabolites in legumes (Fabaceae). *South African
Journal of Botany* **89**: 164–175.

Wojciechowski MF, Lavin M, Sanderson MJ. **2004**. A phylogeny of legumes
(Leguminosae) based on analysis of the plastid *matK* gene resolves many well-supported
subclades within the family. *American Journal of Botany* **91**: 1846–1862.

Yang Y, Smith SA. **2014**. Orthology inference in nonmodel organisms using transcriptomes
and low-coverage genomes: improving accuracy and matrix occupancy for phylogenomics.
Molecular Biology and Evolution **31**: 3081–3092.

Zhang J, Ruhlman TA, Mower JP, Jansen RK. **2013**. Comparative analyses of two
Geraniaceae transcriptomes using next-generation sequencing. *BMC Plant Biology* **13**: 228.

Zhang C, Rabiee M, Sayyari E, Mirarab S. 2018. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. *BMC Bioinformatics* **19**: 153.

Zhang R, Wang Y-H, Jin J-J, Stull GW, Bruneau A, Cardoso D, Queiroz LP, Moore MJ,
Zhang SD, Chen SY. **2020**. Exploration of plastid phylogenomic conflict yields new insights
into the deep relationships of Leguminosae. *Systematic Biology* **69**: 613–622.

Zhao Y, Zhang R, Jiang K-W, Qi J, Hu Y, Guo J, Zhu R, Zhang T, Egan AN, Yi T-S, *et al.*
2021. Nuclear phylotranscriptomics and phylogenomics support numerous
polyploidization events and hypotheses for the evolution of rhizobial nitrogen-fixing
symbiosis in Fabaceae. *Molecular Plant* **14**: 748–773.

Zuanny DC, Vilela B, Moonlight PW, Särkinen TE, Cardoso D. **2024**. expowo: an R
package for mining global plant diversity and distribution data. *Applications in Plant*
Sciences: e11609.

1166

(a)

CAESALPINIOIDEAE

DETARIOIDEAE

CERC.

POLYGALACEAE

PAPILIONOIDEAE

Floral architecture
(tip points)

non-keel

keel

Net diversification
(branch colours)

Fast

Slow

125 Ma   100   75   50   25

| Cretaceous | Pal | Eocene | Oligo | Miocene | Pli |

(b)

Net diversification rate

0   50   100

Relative time

(c)

Probability

0.20

0.15

0.10

0.05

0.00

15   20   25   30

Number of shifts

(d)

Net diversification rate

2.0

1.5

1.0

0.5

0.0

non-keel   keel

Floral architecture