The utility of single-cell RNA sequencing data in predicting plant metabolic pathway genes

Jingwei Ma[1,2‡], Liting Zou[1‡], Zhenglin Wang[1‡], Xiaoxue Wang[1,2], Xinyi Zuo[1,3], Fei Wang[1,3], Zhiqing Wang[1,4], Zhimei Li[1], Lin Li[2]*, Peipei Wang[1]*

[1] Kunpeng Institute of Modern Agriculture at Foshan, Agricultural Genomics Institute at Shenzhen, Chinese Academy of Agricultural Sciences, Shenzhen, Guangdong 518124, China

[2] College of Life Science, Shanxi Normal University, Taiyuan 030031, China

[3] College of Horticulture, Northwest A&F University, Yangling, Shaanxi 712100, China

[4] College of Plant Protection, South China Agricultural University, Guangzhou, Guangdong 510642, China

Summary

● It is an ever challenging task to make genome-wide predictions for plant metabolic pathway genes (MPGs) encoding enzymes that catalyze the biosynthesis of plant natural products.

● Here, starting from 1,130 benchmark MPGs that have experimental evidence in *Arabidopsis thaliana*, we investigate the utilities of single-cell RNA sequencing (scRNA-seq) data—a recently arisen omics data that has been used in several other fields—in predicting MPGs using four machine learning (ML) algorithms that support multi-label tasks.

● Compared with traditional bulk RNA-seq data, scRNA-seq data lead to different and tighter co-expression networks among MPGs within metabolic classes, but relatively lower prediction accuracy of MPGs into classes. Splitting the scRNA-seq data into tissue-specific subsets can improve the gene co-expression network tightness and prediction accuracy of MPGs for some classes. Expression features from the same tissue types in bulk RNA-seq and scRNA-seq data have different contributions to the prediction of MPGs into classes. Models built using the ensemble algorithm AutoGluon outperforms those using other three classical ML algorithms.

28 ● Our results demonstrate the usefulness and characteristics of scRNA-seq data in
29   predicting MPGs into metabolic classes, and propose that more effort is needed in the
30   future to improve the model prediction performance.

32 **Keywords:** *Arabidopsis thaliana*; single-cell RNA sequencing data; bulk RNA sequencing data;
33 plant metabolic pathway genes; machine learning

Introduction

As sessile organisms, plants protect themselves from herbivores and stresses mainly via
producing plant natural products (PNPs) (Al-Khayri *et al.*, 2023). Some PNPs, such as
anthocyanin (Appelhagen *et al.*, 2018), paclitaxel (Yang *et al.*, 2020), salicylic acid (Ahmadi *et*
al., 2016) and pyrethroids (Elshafie *et al.*, 2023), have also been widely used in human nutrition,
medicine, cosmetics, agriculture and other fields (Tiwari & Rana, 2015; Fang *et al.*, 2019; Jucá
et al., 2020; Bhambhani *et al.*, 2021). The content of these natural products in plants is generally
quite low, and the extraction of them is usually inefficient and environmentally unfriendly
(Zhang *et al.*, 2023). Thus, synthesizing PNPs in engineered microbial cells or plant cells has
become a hotspot in synthetic biology, which requires the elucidation of metabolic pathway
genes (MPGs) that encode enzymes catalyzing the biosynthesis of these PNPs. However, only
about 0.1% of metabolic pathways of the estimated ~1 million metabolites synthesized by plants
have been revealed (Schläpfer *et al.*, 2017). Before the genome era, the identification of plant
MPGs relied heavily on traditional genetic and/or biochemical approaches, which are time- and
labor-consuming. For example, the identification of the genes participating in the biosynthesis
pathway of avenacin spanned about 22 years, from the isolation of saponin-deficient (*sad*)
mutants (Papadopoulou *et al.*, 1999) to the identification of the 12-gene cluster (Li *et al.*, 2021).

With the development of omics technologies, more and more metabolic pathways have been
deciphered by researchers via approaches such as genome-wide association studies (GWAS)
(Shang *et al.*, 2014; Dong *et al.*, 2015) and association analysis between transcriptome and
metabolome (Magnard *et al.*, 2015; Tohge *et al.*, 2020). Although these approaches have
improved the efficiency and accuracy of MPG identification, they generally target one or a few
pathways with relatively low throughput, but require large-scale genotyping or RNA-sequencing,
in company with metabolomic profiling.

The development of computational tools, such as those for identifying plant metabolic gene

clusters (Schläpfer *et al.*, 2017; Kautsar *et al.*, 2017; Töpfer *et al.*, 2017) or annotating enzyme

genes (Schläpfer *et al.*, 2017), and machine learning (ML) approaches utilizing one or multiple

types of omics data (Moore *et al.*, 2019; Wang *et al.*, 2021), has made it possible to predict plant

MPGs on a larger scale. In one of our previous studies, we investigated the utility of

transcriptome data in predicting MPGs in tomato using ML approaches, based on the hypothesis

that genes in the same metabolic pathways may be transcribed similarly to cooperatively

synthesize the target metabolites (Wang *et al.*, 2021). Although the ML approaches

outperformed naive methods that soly compared expression similarities among genes in tomato,

the global prediction accuracy was still not good enough (the median optimal F1 was 0.4 for 85

pathways) even though exhaustive combinations of the ways how transcriptome data were used

and types of ML algorithms were evaluated. We reasoned that the failure in predicting tomato

MPGs using transcriptome data may be partially due to the potential shortcomings of the

traditional bulk RNA sequencing (bulk RNA-seq) data used in the study: the RNA sequencing is

generally processed for a mixed pool of multiple types of cells within the sampled tissues.

Considering that the biosynthesis of many PNPs are cell type-specific (Sun *et al.*, 2022), and that

single-cell RNA sequencing (scRNA-seq) data have been used to explore the spatial organization

of PNPs and the transcripts of contributing MPGs (Sun *et al.*, 2022; Yu *et al.*, 2023), we

hypothesize that the prediction of plant MPGs may be potentially improved by using scRNA-seq

data compared with bulk RNA-seq data.

Here, we compare the utilities of scRNA-seq vs. bulk RNA-seq data in predicting MPGs in

Arabidopsis thaliana, for which there are the most comprehensive MPGs with experimental

evidence (hereafter referred to as benchmark MPGs) and scRNA-seq data among other plant

species. We first compare the degrees of co-expression levels for benchmark MPGs within the

same metabolic classes using these two types of RNA-seq data. Next, we establish ML models

for predicting MPGs into metabolic classes using these two data, employing four ML algorithms

that support multi-class and multi-label classification. Finally, we apply our models to enzyme

genes that lack experimental evidence, and compare the predicted metabolic classes of these

genes with the annotated classes of them in the Plant Metabolic Network (PMN) database.

Results

Different co-expression modules of benchmark metabolic pathway genes based on two types of RNA-seq data

Bulk RNA-seq and scRNA-seq data of tissues collected from Arabidopsis wild-type plants grown under normal conditions or continuous light were collected, since the majority of public scRNA-seq data in Arabidopsis were from these tissues, resulting in a total of 13 scRNA-seq datasets comprising 146 cell types and 5 bulk RNA-seq datasets encompassing 40 tissue types (**Table S1–S4**, see **Methods**). We obtained the metabolic pathway annotations for Arabidopsis genes from the PMN database (Schläpfer *et al.*, 2017), and only retained 1,130 benchmark MPGs belonging to 635 metabolic pathways (**Table S5**, **Methods**). Considering that the majority of these 635 metabolic pathways (75.32%) only contained ≤5 benchmark MPGs (**Fig. 1A**), we grouped them into 13 metabolic classes according to a previous paper (Chae *et al.*, 2014) (**Table S5**) to facilitate the following analysis. This classification resulted in an average number of MPGs at 121 across 13 classes (a MPG can belong to multiple classes). Additionally, 34 genes from 37 pathways that do not belong to any of the 13 classes were assigned to the category "Others" (**Fig. 1B**).

To compare how well genes within different metabolic classes can be distinguished using bulk RNA-seq and scRNA-seq data, we established co-expression networks for benchmark MPGs using the weighted gene co-expression network analysis (WGCNA) (**Methods**). These 1,130 MPGs were clustered into 14 co-expression modules by both data: $M1_{bulk}$–$M14_{bulk}$ (**Fig. 1C, Table S6**) and $M1_{sc}$–$M14_{sc}$ (**Fig. 1E**, **Table S7**). In general, all the 14 M_{bulk} modules were overrepresented by MPGs from 1~3 metabolic classes (**Fig. 1D**), whereas 5 M_{sc} modules were not overrepresented by MPGs from any classes (**Fig. 1F**). On the other hand, MPGs from the amines and polyamines (AP) and redox classes were not overrepresented in any of the M_{bulk} modules, but overrepresented in $M8_{sc}$ and $M2_{sc}$, respectively. Similarly, MPGs from the detoxification and intermediate metabolites (IM) classes were not overrepresented in any of the M_{sc} modules, but in $M5_{bulk}$ and $M8_{bulk}$/$M13_{bulk}$, respectively.

In addition, the gene components differed for M_{bulk} and M_{sc} modules, even those that were overrepresented by MPGs from the same metabolic classes. For example, $M14_{bulk}$ (containing 20 hormone genes) and $M9_{sc}$ (15) were overrepresented by MPGs from the hormone class, and

MPGs in M14$_{bulk}$ and M9$_{sc}$ were preferentially expressed in flowers/siliques and flowers/embryo-
related cell types, respectively (**Fig. 1C,E, Table S3,4,6,7**). However, there was no overlap
between hormone genes from M14$_{bulk}$ and M9$_{sc}$ (**Fig. S1A**). MPGs from specialized metabolism
(SM) were enriched in M7$_{bulk}$ and M11$_{bulk}$ (containing 50 and 28 SM genes, respectively), and
M4$_{sc}$ and M7$_{sc}$ (each containing 16 SM genes, **Fig. 1D,F**). MPGs in M7$_{bulk}$, M4$_{sc}$ or M7$_{sc}$ were
preferentially expressed in root-related tissues or cell types, whereas MPGs in M11$_{bulk}$ tended to
be expressed in broad types of tissues (**Fig. 1C,E**). There were 10 and 6 SM genes shared
between M7$_{bulk}$ and M4$_{sc}$, and between M7$_{bulk}$ and M7$_{sc}$, respectively, but each 1 SM gene shared
between M11$_{bulk}$ and M4$_{sc}$, and between M11$_{bulk}$ and M7$_{sc}$ (**Fig. S1B**). Taken together, these
results highlight the usefulness of both types of RNA-seq data in distinguishing MPGs from
different metabolic classes, while underscoring the differences in co-expression networks among
MPGs revealed by these two types of data.

Clustering coefficient of gene expression within 13 metabolic classes based on two types of
RNA-seq data

To better assess the tightness of the gene co-expression networks for individual metabolic
classes, we calculated the clustering coefficients (*C* values) for MPGs within 13 metabolic
classes using these two types of RNA-seq data (see **Methods**). The *C* value is normally used to
measure the degree to which nodes in a graph tend to cluster together (Onnela *et al.*, 2005), and
has been widely used in several real-world networks, particularly in social networks (Opsahl &
Panzarasa, 2009); the higher the *C* value, the tighter the network. To illustrate how well the *C*
value reflects the tightness of co-expression network within metabolic classes (**Table S8**), we
further calculated the *z*-score of the observed *C* value in 10,000 background *C* values where the
metabolic class annotations of all the 1,130 MPGs were randomly shuffled (**Table S9**, see
Methods). A *z*-score>1.645 (with a corresponding *p*-value<0.05) indicates that MPGs in a class
are significantly more tightly co-expressed than random gene groups (upper inset in **Fig. 2A**),
whereas a negative *z*-score indicates not tightly co-expressed (lower inset in **Fig. 2A**). Only the
energy metabolism (EM) and cofactors classes had *z*-scores >1.645 both when two types of
RNA-seq data were used (**Fig. 2A**); another two (IM and redox) and one (detoxification) class
had *z*-scores >1.645 only when scRNA-seq or bulk RNA-seq data were used, respectively. All
the remaining eight classes had *z*-scores <1.645 regardless of the types of RNA-seq data used.

Due to the fact that some PNPs may be synthesized in specific tissues or cell types (Daloso
et al., 2023), we hypothesize that co-expression signals of MPGs synthesizing these natural
products within tissues/cell types may be diluted by combining RNA-seq data from all the
samples. To test this, we split the scRNA-seq data into four subsets based on the tissue types:
root, shoot, leaf and flower (the embryo subset was not considered due to the small number of
cell types [n=9]); the bulk RNA-seq data were not split further because it contained only 40
samples. When the four scRNA-seq subsets were used, more metabolic classes had z-
scores >1.645 (**Fig. 2B**, **Fig. S3,S4**, **Table S10**). For example, detoxification had z-scores >1.645
when the shoot, leaf and flower subsets were used; SM had z-scores >1.645 when root, shoot and
flower subsets were used; the carbohydrates and hormones classes had z-scores >1.645 when
shoot and flower subsets were used. In contrast, IM had z-scores <1.645 when four subsets were
used, compared with the z-score≈5 when the entire scRNA-seq data were used. However, AP
had z-scores <1.645 regardless of which types of RNA-seq data or which subsets of scRNA-seq
data were used. These results are consistent with our previous findings for bulk RNA-seq data
(Wang *et al.*, 2021) that subsets of transcriptome data provided different co-expression patterns
for plant MPGs, and highlight the need to explore different RNA-seq data when examining co-
expression tightness among MPGs for different metabolic classes.

Prediction performance of machine learning models built using two types of RNA-seq data
Thus far, we showed that the benchmark MPGs within most metabolic classes can be
distinguished from others using at least one type of RNA-seq data or one subset of scRNA-seq
data. Next, we asked to what degree benchmark MPGs can be correctly predicted into metabolic
classes. Considering that some MPGs belong to multiple metabolic classes (**Table S5**), we built
ML models using four algorithms that support multi-class and multi-label classification tasks: K-
nearest neighbors (KNN, Guo *et al.*, 2003), Random Forest (RF, Breiman, 2001), eXtreme
Gradient Boosting (XGBoost, Chen & Guestrin, 2016), and AutoGluon-Tabular (AutoGluon,
Erickson *et al.*, 2020). KNN makes predictions of classes for a given gene based on proximity of
this gene with genes in the training set; RF builds a number of classification decision trees using
a number of sub-samples of the dataset and then aggregates the trees into a single ensemble
model; XGBoost is an optimized distributed gradient boosting library that provides parallel tree
boosting; AutoGluon ensembles multiple ML algorithms and stacks them in multiple layers.

Twenty percent of benchmark MPGs within each class were first held out as the test set, and the
remaining 80% MPGs were used to train the models using a five-fold cross-validation scheme
(see **Methods**). F1 scores on the cross-validation ($F1_{CV}$) and test ($F1_{test}$) sets for models with the
optimal combination of parameters were both reported, where the F1 score was the harmonic
mean of precision and recall of the prediction (see **Methods**).

When measuring F1 scores for MPGs across all the 14 classes (13 metabolic classes plus the
"Others" class), we found that models built using bulk RNA-seq data outperformed scRNA-seq-
based models in terms of both $F1_{CV}$ and $F1_{test}$ scores, regardless of which algorithm was used
(**Fig. 3A**). Although having the highest prediction accuracy, AutoGluon models had potential
issues of overfitting, as $F1_{CV}$ scores for AutoGluon models were much higher than the
corresponding $F1_{test}$ scores. RF models had the lowest prediction accuracy, with both $F1_{CV}$ and
$F1_{test}$ <0.14. Although F1 scores for our models were mostly < 0.3, they were significantly higher
than those for 1,000 random guesses where the metabolic classes of all the benchmark MPGs
were randomly shuffled (the corresponding z-scores >1.645, **Fig. 3B, Fig. S5, Table S11,12**),
suggesting the usefulness of these two types of RNA-seq data in predicting MPGs into metabolic
classes.

Considering that MPGs within different classes behaved differently in the co-expression
analysis as shown in the above sections, we also calculated F1 scores for MPGs within each class.
Consistent with the overall F1 scores, bulk RNA-seq data led to significantly higher $F1_{CV}$ and
$F1_{test}$ scores for more classes than scRNA-seq data regardless of which algorithms were used
(**Fig. 3C,D**, **Fig. S6A,B**), and the F1 scores for individual classes were more similar between
KNN, RF and XGBoost models than with AutoGluon models (**Fig. 3E**, **Fig. S6E,F**, only the
comparison between $F1_{CV}$ scores were shown). In addition, we found that the prediction
performance for a class was positively correlated with the number of MPGs within the class both
when bulk RNA-seq and scRNA-seq data were used (**Fig. 3F**, only results of $F1_{CV}$ for XGBoost
models were shown as examples). This result suggests that with the increase of knowledge about
plant MPGs in the future, the prediction accuracy for these classes is expected to be improved.
When examining the z-scores of F1 scores, which were supposed to better reflect the prediction
performance of models built with different factors than F1 scores *per se*, we found that most of
these 13 classes had z-scores >1.645 for both $F1_{CV}$ and $F1_{test}$ regardless of which type of RNA-
seq data and which algorithms were used (**Fig. 3G,H**, **Fig. S6C,D**, **Fig. S7–14**, **Table S13–20**).

In the above section, we showed that tissue-specific scRNA-seq data led to higher C values
for some classes (**Fig. 2B**) than the entire scRNA-seq data. We next asked whether the tissue-
specific scRNA-seq data can also lead to better prediction for MPGs. $F1_{CV}$ scores of three, nine,
four and seven classes can be improved by some of the tissue-specific scRNA-seq data for KNN,
RF, XGBoost and AutoGluon models, respectively, while $F1_{test}$ scores of seven, six, three and
four classes were improved for KNN, RF, XGBoost and AutoGluon models, respectively (**Fig.**
S15). These results further highlight the need to explore different scRNA-seq data when
predicting MPGs into different metabolic classes.

Furthermore, we investigated whether combining the two types of RNA-seq data would
improve the prediction accuracy for MPGs. For the KNN models, the combined data led to
significantly higher $F1_{CV}$ and $F1_{test}$ for four and two classes, respectively, both than when bulk
RNA-seq or scRNA-seq data were used (**Fig. S16A**). Particularly, the average $F1_{CV}$ and $F1_{test}$
scores of 10 replicate runs for inorganic nutrients (IN) were improved from 0.034/0.028 (bulk
RNA-seq/scRNA-seq) and 0/0 to 0.169 and 0.136, respectively. However, for RF, XGBoost and
AutoGluon models, only $F1_{test}$ scores of one (cofactors), two (amino acids [AA] and AP), and
two (redox and detoxification) classes were significantly improved, respectively, when combined
data were used compared with two individual types of RNA-seq data (**Fig. S16B–D**). These
results suggest that, although bulk RNA-seq and scRNA-seq data led to different gene co-
expression networks and different prediction performance for MPGs within classes, combining
these two data does not necessarily leverage the strengths of both data and lead to better
prediction performance than individual data in most cases.

Important features revealed by interpreting the predictive models

To evaluate how the gene expression data from two types of RNA-seq data contribute to the
prediction of MPGs into metabolic classes, we examined the feature importance for models built
using these two data (see **Methods**, feature importance is not applicable for KNN models). We
found that when the same RNA-seq data were used, models built using RF and AutoGluon
tended to have more similar feature importance than with XGBoost models (**Fig. 4A, Fig. S17,18,**
Table S21–26). In addition, the correlations of feature importance between different algorithms
were higher in bulk RNA-seq-based models than in those based on scRNA-seq data (**Fig. 4A**).
Together with the above findings that bulk RNA-seq-based models tended to have better

prediction of MPGs for more metabolic classes than scRNA-seq-based models, this result
suggests that bulk RNA-seq features may be more stable to reflect the co-expression correlations
among MPGs than scRNA-seq features.

On the other hand, when the same algorithm was used, features from the same tissue types
between bulk RNA-seq and scRNA-seq data tended to have different importance ranks. For
example, in RF models, bulk RNA-seq features from different types of tissues were generally
ranked evenly according to the feature importance (**left panel in Fig. 4B**, **Fig. S17A, Table S21**),
while the scRNA-seq features from flowers tended to have higher importance and rank relatively
higher than features from other tissues (**right panel in Fig. 4B**, **Fig. S18A, Table S22**). This
result is consistent with the findings that RF models built using scRNA-seq data from flowers
tended to have the highest or relatively higher prediction accuracy for most classes (**Fig. S15B**),
and that the number of MPGs that were specifically expressed in flowers in the scRNA-seq data
was the highest among others (**Fig. 1E**). Similarly, for AutoGluon models, scRNA-seq features
from flowers and embryos tended to be more important than those from other tissues (**Fig. S18C,**
Table S26), whereas bulk RNA-seq features from different tissues ranked more evenly (**Fig.**
S17C, Table S25). AutoGluon models built using scRNA-seq data from flowers and embryos
had better prediction accuracies for detoxification and IN than models based on data from other
tissues (**Fig. S15D**). In contrast, for XGBoost models, both the bulk RNA-seq and scRNA-seq
features from different tissues ranked relatively evenly (**Fig. S17B**, **Fig. S18B, Table S23,24**),
and much fewer metabolic classes had improved prediction accuracy when cell type-specific
scRNA-seq data were used compared with RF and AutoGluon models (**Fig. S15**). These results
suggest that RF and AutoGluon models may outperform XGBoost models in distinguishing the
most important predictors.

Beside the advantages of having higher prediction accuracy and being potentially better at
distinguishing important predictors, the AutoGluon models also allow us to examine the
importance of features to the prediction of individual classes (**Table S25,26**). For example, gene
expression in the cell culture tissues at an early stage ("Cell_culture_early" in **Fig. 4A**) and gene
expression in central cells of female gametophytes ("Central_cell", **Fig. 4A**) were consistently
important for models built using three algorithms, and these two features were most important
for predicting MPGs into the carbohydrates (**Fig. S19B, Table S25**) and EM (**Fig. S19A, Table**
S26) classes. To understand why these features were more important for predicting MPGs into

the corresponding metabolic classes than others, more analysis is needed to be conducted in the
future.

Prediction of metabolic classes for enzyme genes without experimental evidence

To evaluate how our models performed for new genes, we applied our models to 9,441
enzyme genes that have not been assigned to any metabolic pathways supported by experimental
evidence (hereafter referred to as unknown genes). Only AutoGluon and XGBoost models (the
last model from 10 replicate runs) were used due to the relatively better performance compared
with models built using the other two algorithms. Using AutoGluon models, 8,361 (88.6%) and
8,322 (88.1%) unknown genes were predicted into at least one of the 13 metabolic classes when
bulk RNA-seq and scRNA-seq data were used (**Fig. 5A**), whereas 4,623 (49.0%) and 3,425
(36.3%) unknown genes were predicted by bulk RNA-seq-based and scRNA-seq-based
XGBoost models, respectively(**Fig. S20A**). The number of unknown genes that were predicted
into each metabolic class was positively correlated with the number of benchmark MPGs within
the class (**Fig. 5A, Fig. S20A**). This result suggests two potential scenarios that are not mutually
exclusive: 1) the current pre-knowledge about MPGs are relatively proportionally distributed
among metabolic classes; 2) the data structure of predicted instances are influenced by the data
structure of the training instances.

Next, we examined whether the predicted metabolic classes for unknown genes are
consistent between models built using two types of RNA-seq data and with the annotation from
the PMN database (**Table S5, S27–30**). Using AutoGluon models, 2,051 unknown genes were
predicted into exactly the same metabolic classes using bulk RNA-seq and scRNA-seq data, and
119 out of them were annotated in the same classes by PMN (strict criteria, upper panel in **Fig.**
5B); the numbers were 2,281 and 251 for a loose criteria where a gene had at least one common
class between different datasets (lower panel in **Fig. 5B**). In comparison, using XGBoost models,
692/33 and 1,003/121 unknown genes shared common class memberships between two
predictions/among three datasets, when using strict or loose criteria, respectively (**Fig. S20B**),
much fewer than those using AutoGluon models. These results further indicate that the ensemble
algorithm AutoGluon outperforms XGBoost (and also the other two algorithms) in predicting
MPGs into metabolic classes using transcriptomic data.

Considering that a benchmark MPG can be annotated with more than one class, and the
multiple class annotations for a gene were not randomly distributed (e.g., MPGs in detoxification
tended to be also annotated as in redox, compared with other classes, **Fig. 5C**), we further
investigated whether the mismatches of classes between PMN annotations and model predictions
have a similar bias among classes (**Fig. 5D,E, Fig. S20C,D**). For some cases the mismatches
resembled the multi-label annotations. For example, unknown genes that were annotated as
within SM by PMN tended to be predicted as within the class hormones by the bulk RNA-seq-
based AutoGluon model (**Fig. 5D**) and both XGBoost models built using bulk RNA-seq and
scRNA-seq data (**Fig. S20C,D**). However, the global mismatch patterns did not necessarily
repeat the multi-label annotation patterns. For example, benchmark MPGs in EM tended to also
belong to carbohydrates and AA (**Fig. 5C**), whereas unknown genes in EM from PMN
annotations tended to be predicted into cofactors and hormones by bulk RNA-seq-based
AutoGluon model (**Fig. 5D**), fatty acids and lipids by scRNA-seq-based AutoGluon model (**Fig.
5E**), and SM by both XGBoost models (**Fig. S20C,D**). This discrepancy can be mainly due to the
unbalanced numbers of benchmark MPGs from different classes used to train the models, and the
far-from-perfect prediction accuracy of our models built using transcriptomic data. The potential
low quality of PMN annotations for these unknown genes can be another rationale that should
not be excluded at the current stage.

Discussion

Models built using two types of RNA-seq data had significantly higher prediction accuracy
for most metabolic classes than random guesses, suggesting the usefulness of both RNA-seq data
in predicting MPGs into metabolic classes. Generally, bulk RNA-seq-based models led to higher
performance in predicting MPGs into classes than scRNA-seq-based models, opposed to our
assumption that scRNA-seq data may provide more detailed expression profiles for genes than
bulk RNA-seq data. There can be three potential reasons for this result. Firstly, the sequencing
and analytical techniques for single-cell transcriptome at present still have multiple limitations,
such as the low capture efficiency for transcripts within cells, especially for genes with low
expression, resulting in inherently sparse and noisy nature of the scRNA-seq data (Kharchenko,
2021); the analysis of scRNA-seq data is complex and challenging, and requires prior knowledge
about marker genes (Ke *et al.*, 2022). With the development and improvement of sequencing and

analytical techniques, scRNA-seq data is expected to show more promise in multiple fields,

including predicting plant MPGs.

Secondly, the biosynthesis of a PNP may occur sequentially in a series of specific cells. For

example, the transcription of 20 genes responsible for the biosynthesis of monoterpenoid indole

alkaloids occur in three types of cells (namely, internal phloem-associated parenchyma cells,

epidermal cells, and laticifer cells) in the *Catharanthus roseus* leaves, and the intermediates

produced during the biosynthesis process are transported between different cell types (Sun *et al.*,

2022). If this phenomenon is common for PNPs, bulk RNA-seq data that capture gene expression

profiles for the entire tissues rather than individual cell types are supposed to be superior to

scRNA-seq data in predicting MPGs. Bioinformatic approaches that take into account the

temporal or spatial frameshift in expression when measuring the expression correlation between

genes could potentially make better use of the scRNA-seq data in predicting MPGs.

Thirdly, our predictive models used the expression profiles of genes in bulk RNA-seq and

scRNA-seq data, rather than the expression correlations among genes (e.g., C values of genes).

In our clustering coefficient analysis, scRNA-seq data led to higher C values (z-scores > 1.645)

for more classes than bulk RNA-seq data (**Fig. 2A**), and for even more classes when tissue-

specific scRNA-seq data were used (**Fig. 2B**). Models built using C values of genes may

improve the prediction performance when scRNA-seq data were used. However, a major

challenge for using C values of genes as predictors in the models is that the C value of a gene is

calculated by considering all the neighbors that are directly connected with the gene. It is a tricky

problem to calculate the C values for genes in the validation and test subsets with neighbors from

the training subset while controlling the risks of data leakage during the calculation.

The finding that prediction performance for a given class was influenced by the number of

MPGs within the class regardless of the types of RNA-seq data used (**Fig. 3F**) suggests an urgent

need to expand our knowledge about plant MPGs to improve the prediction accuracy, which in

turn needs assistance from other approaches (such as ML approaches in our study) beyond

conventional forward genetics. One potential solution for this seemingly "endless loop" is to

experimentally evaluate part of the preliminary predictions (error-prone) and feed the validated

knowledge back to the ML models to boost the prediction performance. Another potential

solution would be the development of ML algorithms which can handle with giant amount of

features (e.g., expression and epigenetic profiles of genes across a large number of tissues, cell

types, stages, environments, and so on) while learning from a limited numbers of instances (i.e.,
genes). The AutoGluon algorithm has shown great performance in prediction tasks such as
OpenML AutoML and Kaggle Benchmark (Erickson *et al.*, 2020). Although outperformed
models built using other three classical ML algorithms in predicting MPGs into metabolic
classes in our study, AutoGluon models had the risk of overfitting. This indicates that for some
prediction tasks in biology with limited numbers of instances (e.g., plant MPGs), the algorithms
for ensemble learning (e.g., AutoGluon) that show great promise in other fields (e.g., image
recognition) should be used with caution.

In our study, due to the limitation of publicly released scRNA-seq data, we only focused on
scRNA-seq and bulk RNA-seq data from wild-type tissues that were grown under normal
conditions. A number of PNPs, especially specialized metabolites, are only synthesized in
specific tissues or cell types, or under specific conditions, such as biotic and abiotic stresses. For
example, acyl sugars are specifically synthesized in the glandular trichomes of certain plants
(Schilmiller *et al.*, 2008); the rupture of a cassava leaf mesophyll cell caused by herbivore attack
or animal chewing leads to the production of cyanide which has the bitter taste for mammals
(McMahon *et al.*, 1995). Our findings showed that tissue-specific scRNA-seq data led to better
prediction for some classes than the entire scRNA-seq data we collected. More scRNA-seq data
for samples collected under different conditions are needed to improve the prediction accuracy
for MPGs responsible for the biosynthesis of the PNPs that are specific to the corresponding
conditions (Wang *et al.*, 2021).

In summary, the prediction of MPGs is a complex and challenging task. More efforts should
be put in the future to expand our knowledge about the contributing genes underlying PNPs with
higher efficiency and to accelerate the development of synthetic biology.

Materials and Methods

RNA-seq data collection and preprocessing

A total of 13 scRNA-seq datasets of *A. thaliana* (**Table S1**) were downloaded from the
PlantscRNAdb (http://ibi.zju.edu.cn/plantscrnadb/, accessed on Mar 20th, 2023) and Gene
Expression Omnibus (GEO, https://www.ncbi.nlm.nih.gov/geo/, accessed on March 20th, 2023)
databases, or from the publication (Song *et al.*, 2020). These scRNA-seq datasets contained
samples taken from root, shoot, leaf, flower and embryo tissues of wild-type plants grown under

normal conditions or continuous lights. Five bulk RNA-seq datasets that contained samples from
the same types of tissues of wild-type plants grown under normal conditions or continuous lights
(**Table S2**) were downloaded from the Expression Atlas database
(https://www.ebi.ac.uk/gxa/sc/home) and GEO (accessed on Mar 20th, 2023) databases, or from
the publication (Mergner *et al.*, 2020).

For the preprocessing of scRNA-seq data, genes expressed in <3 cells and cells with <200
genes expressed were first filtered out from the following analysis. Missing values in the
resulting expression matrix were filled with zeros. The Seurat (v4.3.0) package (Hao *et al.*, 2021)
in R (v4.2.0) was used to preprocess the scRNA-seq data. Specifically, the "RunPCA" function
was used for dimensionality reduction; "NormalizeData" was used to normalize gene expression
levels among different cells, where read count of each gene was divided by the total counts
within individual cells, then multiplied by a scale factor of 10,000 and log1p-transformed;
"FindClusters" was used to cluster cell types; "UMAP" was used to visualize the clusters.
Marker genes that were specifically expressed in certain cell types were obtained from the
PlantscRNAdb and were used to annotate the corresponding cell types. A total of 146 cell types
were annotated and gene expression levels were averaged among cells with the same type (**Table
S3**).

For the preprocessing of bulk RNA-seq data, the gene expression levels (transcripts per
million, TPM) were first averaged across biological replicates, and then were normalized using
the "normalize.quantiles" function in R among different samples, where the expression values
within individual samples were adjusted to ensure a consistent distribution and to eliminate
technical variations between samples. The normalized expression values were further log1p-
transformed to enable comparison with those in scRNA-seq data. (**Table S4**).

Metabolic pathway gene annotation collection

We first downloaded the metabolic pathway gene annotation from the PMN database
(https://plantcyc.org/, accessed on Sep 9th, 2022, PlantCyc v15.1.0, Schläpfer *et al.*, 2017), which
contained 3,508 MPGs annotated in 635 metabolic pathways. To obtain benchmark MPGs, we
wrote a python script to crawl the PMN webpages of individual MPGs (accessed on Dec 19th,
2022) and kept MPGs with experimental evidence, such as those inferred from direct assay,
mutant phenotype, etc. This resulted in a total of 1,131 benchmark MPGs, which belong to 636

419 metabolic pathways (**Table S5**). The gene *AT5G23960* and its corresponding pathway β-
420 caryophyllene biosynthesis (PWY-6275) was removed from the following analysis since it
421 lacked expression information in the scRNA-seq data. Additional potential enzyme genes of *A.*
422 *thaliana* were obtained from The Arabidopsis Information Resource (TAIR) databases
423 (https://www.arabidopsis.org/), or annotated using the Ensemble Enzyme Prediction Pipeline
424 (E2P2, v3.1) software. In summary, a total of 10,572 potential enzyme-encoding genes were
425 annotated, including 1,131 benchmark MPGs (**Table S5**).

426 **Weighted gene co-expression network analyses**

427  Gene co-expression network analysis was conducted using the WGCNA (v1.72.1, Zhang &
428 Horvath, 2005) package in R. During the network analysis, the matrix of Pearson correlation
429 coefficient (PCC) values for gene pairs was transformed into an adjacency matrix to achieve a
430 scale-free network structure, with the parameter 'networkType' being set to 'signed', where the
431 PCC values were processed using the equation:

432
$$a_{i,j} = \left(\frac{1+cor(i,j)}{2}\right)^{\beta} \qquad (1)$$

433 where, $a_{i,j}$ represents the weight of the edge between two genes *i* and *j*; *cor*(*i, j*) represents the
434 PCC between the two genes; *β* is the best soft threshold power estimated by the software
435 WGCNA using the function "pickSoftThreshold", which was estimated at 12 for both bulk
436 RNA-seq and scRNA-seq data. The "cutreeDynamic" function was employed to divide the
437 network into 14 co-expression modules for both RNA-seq data, using parameters as
438 minModuleSize=30, mergeCutHeight=0.1 for both data, while deepSplit=3 and 4 for scRNA-seq
439 and bulk RNA-seq data, respectively.

440 **Clustering coefficient calculation**

441  To measure the tightness of gene co-expression within metabolic classes, we calculated the
442 clustering coefficient (*C* value) for each class. The gene expression values in two RNA-seq data
443 first plus one and were log-transformed with base 10, and then processed through the WGCNA
444 pipeline to obtain the gene co-expression networks. In the networks, the nodes were benchmark
445 MPGs, edges between nodes were co-expression relationships between MPGs with the thickness
446 of edges (i.e., the strength of the connection) being the transformed PCC values. The networks

were next processed using the NetworkX (v3.1, Hagberg *et al.*, 2008) package in Python to

calculate the local C values for individual genes, with the equation (Onnela *et al.*, 2005):

$$c_u = \frac{1}{deg(u)(deg(u)-1)} \sum_{vw} (\widehat{w}_{uv} \widehat{w}_{uw} \widehat{w}_{vw})^{1/3} \qquad (2)$$

where, for a node (i.e., a gene) u with two adjacent nodes v and w as an example, c_u represents

the local C value of the node u; $deg(u)$ is the degree of the node u, representing the number of

neighboring nodes connected to the node u; w_{uv}, w_{uw}, and w_{vw} represent the weights of the

edges between nodes u, v, and w; the weights are normalized by being divided by the maximum

weight in the network, e.g., $\widehat{w}_{uv} = w_{uv}/max(w)$; $(\widehat{w}_{uv} \widehat{w}_{uw} \widehat{w}_{vw})^{1/3}$ indicates the geometric

mean of the normalized weights between nodes u, v, and w; $\sum_{vw}$ represents the summation

over all pairs of neighboring nodes (i.e., v,w) that are directly connected to node u. The local C

value ranges from 0 to 1. The average local C value of all genes within a metabolic class (the

global C value, hereafter referred to as C value for short) was used to represent the compactness

of the class in terms of gene expression (**Table S8**).

Since the C values can be influenced by the number of nodes within a network (Kaiser,

2008), to understand what C values would indicate a tight network and to make the C values

among different metabolic classes comparable, we also estimated the C values for 10,000

random simulations (**Table S9,10**) where the class memberships of all the benchmark MPGs

were randomly shuffled to simulate the out-of-order relationships among these MPGs. Then a z-

score of the observed C values in the distribution of 10,000 background C values were calculated

using the equation:

$$z = \frac{x-\mu}{\sigma} \qquad (3)$$

where, z represents the z-score value, x represents the observed C value, μ represents the mean of

10,000 background C values, and σ represents the standard deviation of 10,000 background C

values.

Machine learning model building

Four algorithms were used to build ML models: KNeighborsClassifier (K-Nearest

Neighbors, KNN), RandomForestClassifier (RF), and XGBClassifier (eXtreme Gradient

Boosting, XGBoost) from scikit-learn (v1.2.2, implemented in Python v3.9.16, Pedregosa *et al.*,

2011), and AutoGluon-Tabular (AutoGluon, Erickson *et al.*, 2020) from the AutoGluon library

(v0.8.2, Python). Out of the benchmark MPGs within each class, 20% were used as the test set to
evaluate the performance of the models. The remaining 80% of benchmark MPGs within each
class were further split into five folds; MPGs in four folds were used to train the model and
MPGs in the remaining fifth fold were used to validate the model performance; this training-
validation step was conducted five times to make sure that MPGs in all five folds were used as in
the validation subset once. Caution should be taken to avoid that genes that belong to multiple
classes are held out as in the test set for one class but remain as in the training set for another
class, and the same for the training/validation splits.

Hyperparameter tuning was conducted for KNN, RF and XGBoost models (this step is
automatically embedded within AutoGluon) to select the optimal combination of parameters
using GridSearch. Specifically, the hyperparameter spaces for KNN models were {3, 4, 5, 6, 7}
for the 'n_neighbors' (*k* values) and {'uniform', 'distance'} for the 'weights'; for RF models: {3, 5,
10} for the 'max_depth', {0.1, 0.25, 0.5, 0.75, 'sqrt', 'log2', None} for the 'max_features' and {10,
50, 100, 500} for the 'n_estimators'; for XGBoost modes: {3, 5, 10} for 'max_depth', {0.01, 0.1,
0.5, 1} for 'learning_rate', {10, 50, 100, 500} for 'n_estimators' and {0.1, 5} for 'gamma'. The
best combination of parameters for each model was determined based on the F1 scores on the
cross-validation set (F1$_{CV}$), where F1 was calculated with the equations:

493
$$F1 = \frac{2 \times precision \times recall}{precision + recall} \tag{4}$$

494
$$precision = \frac{TP}{TP + FP} \tag{5}$$

495
$$recall = \frac{TP}{TP + FN} \tag{6}$$

where TP (true positive) represents the number of benchmark MPGs that are correctly predicted
into a class; FP (false positive) represents the number of benchmark MPGs that are mis-predicted
into a class; FN (false negative) represents the number of benchmark MPGs that are not
predicted into the correct class.

Finally, the feature importance was obtained for RF ("gini"), XGBoost ("weight"), and
AutoGluon ("permutation-based") models (**Table S21–26**), and the feature importance for all the
classes were averaged for AutoGluon models to make them comparable to those for RF and
XGBoost models.

Acknowledgments

We thank Xinhao Fan, Qiaowei Li, Kangli Li, Ruipu Chen, Zhuobiao Wang, for their helpful suggestions in preprocessing the scRNA-seq data. This work was supported by the Scientific Research Foundation for Principle Investigator, Kunpeng Institute of Modern Agriculture at Foshan (KIMAQD2022003 to PW) and National Natural Science Foundation of China (32370241 to PW).

Author contributions

PW conceived and designed this study. ZW collected the metabolic pathway gene annotations; LZ and JM collected and preprocessed the bulk RNA-seq and scRNA-seq data, respectively. ZW wrote the machine learning pipelines. JM conducted all the other analyses with help from LZ, XW and ZW. JM and PW wrote the manuscript with inputs from all the other authors. All authors read and approved the final manuscript.

Data availability

All the data used in this study are provided in the supplementary materials, and all the scripts used are available on Github at:

https://github.com/peipeiwang6/Manuscript/tree/main/2024_scRNA_in_pathway_prediction.

Figure legends

Fig. 1. Different expression patterns for genes within metabolic classes revealed using bulk RNA-seq and single-cell RNA-seq (scRNA-seq) data. (**A**) A pie plot showing the percentages of 635 metabolic pathways that have a certain number of benchmark metabolic pathway genes (MPGs): blue, 1 MPG; purple, 2; cyan, 3; orange, 4; green, 5; red, $\geq$ 6 MPGs. (**B**) A bar chart showing the numbers of MPGs belonging to 13 metabolic classes and the "others" category that contains all the other pathways not belonging to any classes. (**C,E**) Heatmaps showing the expression patterns of 1,130 MPGs, which were grouped into 14 co-expression modules (M1$_{bulk}$–M14$_{bulk}$ for **C**, M1$_{sc}$–M14$_{sc}$ for **E**) inferred by weighted gene co-expression network analysis using the bulk RNA-seq (**C**) and scRNA-seq (**E**) data. Color scale in the heatmaps: the

standardized expression values (transcripts per kilobase million) using z-score normalization; red: high expression, blue: low expression; all values > 3 were assigned a value of 3. Tissue types that bulk RNA-seq samples or scRNA-seq cell types belong to are indicated above the heatmaps with colors: dark green, root; green, shoot; olive, leaf; orange, flower; pink, embryo; light pink, others. (**D,F**) Heatmaps showing enrichment of genes from 13 different metabolic classes (columns) within 14 co-expression modules (rows) when bulk RNA-seq (**D**) and scRNA-seq (**F**) data were used. The ordering of metabolic classes is the same as in (**B**). Color scale: odds ratio from Fisher's exact test; all values > 4 were assigned a value of 4. Asterisk(s) indicate(s) significance levels of the Fisher's exact test: *, p-value<0.05; **, p-value<0.01; ***, p-value<0.001; only the over-representative cases (odds ratio > 1) are indicated with the significance levels. SM: specialized metabolism; Carb: carbohydrates; FAL: fatty acids and lipids; AA: amino acids; Cof: cofactors; Horm: hormones; EM: energy metabolism; NT: nucleotides; IN: inorganic nutrients; DTX: detoxification; AP: amines and polyamines; IM: intermediate metabolism.

Fig. 2. Clustering coefficients (*C* values) for genes within 13 metabolic classes calculated using two types of RNA-seq data. (**A**) The z-scores of C values for genes within 13 metabolic classes compared with background values, when bulk RNA-seq (blue) and scRNA-seq (orange) data were used. The dashed horizontal line indicates a z-score of 1.645, corresponding to a p-value of 0.05. Light gray background: metabolic classes with significantly higher C values than background values when both types of RNA-seq data were used; light orange and light blue backgrounds: classes with significantly higher C values only when scRNA-seq or bulk RNA-seq data were used, respectively. The two inset density plots show the distribution of 10,000 background C values for genes in the energy metabolism (EM, upper) and specialized metabolism (SM, lower) classes as examples when scRNA-seq data were used. Red solid vertical line: the observed C value for genes within EM or SM class; black dashed vertical line: the average C value of 10,000 background values. (**B**) z-scores of C values for genes within 13 metabolic classes when scRNA-seq data from four different types of tissues (root, shoot, leaf, and flower) were used.

Fig. 3. Prediction accuracy of genes in metabolic classes for models built using two types of RNA-seq data and different algorithms. (**A**) Overall $F1_{CV}$ (left panel) and $F1_{test}$ (right) scores for all 13 metabolic classes when bulk RNA-seq (blue) and scRNA-seq (orange) data were used (n=10 replicate runs). KNN, RF, XGBoost and AutoGluon: models built using K-Nearest Neighbors, Random Forest, eXtreme Gradient Boosting and AutoGluon-Tabular, respectively. (**B**) z-scores of $F1_{CV}$ (left panel) and $F1_{test}$ (right) for models built using two types of RNA-seq data and different algorithms. A z-score > 1.645 (dashed line, p-value=0.05) indicates that the model's performance is significantly better than random guess. (**C,D**) $F1_{CV}$ (left panel) and $F1_{test}$ (right) scores for individual metabolic classes for XGBoost (**C**) and AutoGluon (**D**) models. **E:** Correlation of $F1_{CV}$ between XGBoost (x-axis) and KNN models (y-axis) built using bulk RNA-seq (left panel) and scRNA-seq (right) data. (**F**) Correlation between the number of MPGs in a class and the $F1_{CV}$ for the class in XGBoost models, when bulk RNA-seq (left panel) and scRNA-seq (right) data were used. Error bars indicate the standard deviation of results from 10 replicate runs. PCC: Pearson correlation coefficient. (**G,H**) z-score values of $F1_{CV}$ (left panel) and $F1_{test}$ (right) scores for 13 metabolic classes compared with values of random guess for 1,000 times for XGBoost (**G**) and 100 times for AutoGluon (**H**) models. Asterisk(s) indicate(s) significant levels for the two-sided Wilcoxon rank-sum test: *, p-value < 0.05; **, p-value < 0.01; ***: p-value < 0.001. EM: energy metabolism; Cof: cofactors; IM: intermediate metabolism; DTX: detoxification; AA: amino acids; NT: nucleotides; AP: amines and polyamines; Carb: carbohydrates; FAL: fatty acids and lipids; IN: inorganic nutrients; SM: specialized metabolism; Horm: hormones.

Fig. 4. Feature importance for RF, XGBoost and AutoGluon models. (**A**) Correlation of importance for bulk RNA-seq (upper panels) and scRNA-seq (lower) features between RF, XGBoost and AutoGluon models. PCC: Pearson correlation coefficient. (**B**) Importance of bulk RNA-seq (left panel) and scRNA-seq (right) features in RF models. For the details of feature names, see **Fig. S17,18**. Different colors indicate features from different tissue types: dark green, root; green, shoot; olive, leaf; orange, flower; pink, embryo; light pink, others.

Fig. 5. Prediction of metabolic classes for unknown genes by AutoGluon models. (**A**) Count of unknown genes that were predicted into 13 metabolic classes by AutoGluon models built

using bulk RNA-seq (upper panel) and scRNA-seq (lower) data. (**B**) Venn diagrams showing the

overlap of unknown genes that were predicted into metabolic classes by AutoGluon models built

using bulk RNA-seq (blue) and scRNA-seq (orange) data, and those that were annotated as

within metabolic classes in the PMN database (purple). In the upper and lower panels,

overlapping regions show the number of genes with exactly the same classes or at least one

common class between datasets, respectively. (**C**) A heatmap showing the numbers of

benchmark MPGs that were annotated within a certain metabolic class (diagonal cells) and

within two different classes (non-diagonal cells). For example, from 242 benchmark MPGs that

were annotated as in SM, 31 MPGs were also annotated as in the hormone class. (**D,E**)

Heatmaps showing the numbers of unknown genes that were predicted as in a class (x-axis) by

XGBoost models built using bulk RNA-seq (**D**) and scRNA-seq (**E**) data, and were also

annotated as within the same or other classes (y-axis) by PMN database. Values in the column

"Not-assigned" indicate the numbers of unknown genes that were predicted into neither any of

the 13 metabolic classes nor the "Others" category. Cell colors in (**C,D,E**) indicate the numbers

of genes. EM: energy metabolism; Cof: cofactors; IM: intermediate metabolism; DTX:

detoxification; AA: amino acids; NT: nucleotides; AP: amines and polyamines; Carb:

carbohydrates; FAL: fatty acids and lipids; IN: inorganic nutrients; SM: specialized metabolism;

Horm: hormones.

Supplementary figure legends

Fig. S1. Shared benchmark MPGs within co-expression modules inferred by bulk RNA-seq

and scRNA-seq data. (**A**) A venn diagram showing the numbers of benchmark MPGs shared by

$M3_{bulk}$, $M4_{bulk}$, $M14_{bulk}$, $M7_{sc}$, and $M9_{sc}$, for which MPGs from the hormone class were

overrepresented. (**B**) A venn diagram showing the numbers of benchmark MPGs shared by

$M7_{bulk}$, $M11_{bulk}$, $M4_{sc}$, and $M7_{sc}$, for which MPGs from the specialized metabolism class were

overrepresented.

Fig. S2. Clustering coefficients (*C* values) for MPGs within each metabolic class calculated

using bulk RNA-seq (A) and scRNA-seq (B) data. The density plot shows the distribution of

background *C* values for MPGs within each metabolic class for 10,000 random simulations

where the metabolic class annotations for MPGs were randomly shuffled. The red solid line

denotes the observed *C* value for the co-expression network of genes within each class; the black dashed line indicates the average *C* value of 10,000 random simulations. The *z* represents the *z*-score of the observed *C* value in the background distribution; *p* represents the corresponding *p*-value of the *z*-score.

Fig. S3. *C* values for MPGs within each metabolic class calculated using scRNA-seq data of roots (A) and shoots (B). The density plot shows the distribution of background *C* values. The density plot shows the distribution of background *C* values for MPGs within each metabolic class for 10,000 random simulations where the metabolic class annotations for MPGs were randomly shuffled. The red solid line denotes the observed *C* value for the co-expression network of genes within each class; the black dashed line indicates the average *C* value of 10,000 random simulations. The *z* represents the *z*-score of the observed *C* value in the background distribution; *p* represents the corresponding *p*-value of the *z*-score.

Fig. S4. *C* values for MPGs within each metabolic class calculated using scRNA-seq data of leaves (A) and flowers (B). The density plot shows the distribution of background *C* values for MPGs within each metabolic class for 10,000 random simulations where the metabolic class annotations for MPGs were randomly shuffled. The red solid line denotes the observed *C* value for the co-expression network of genes within each class; the black dashed line indicates the average *C* value of 10,000 random simulations. The *z* represents the *z*-score of the observed *C* value in the background distribution; *p* represents the corresponding *p*-value of the *z*-score.

Fig. S5. Overall F1 scores for KNN, RF, XGBoost, and AutoGluon models when bulk RNA-seq (A) and scRNA-seq (B) data were used. The density plot shows the distribution of overall F1 scores for models where the labels of benchmark MPGs in the training set were randomly shuffled (random guesses). Red solid line: the average $F1_{CV}$ or $F1_{test}$ score for models built using the true labels of MPGs; pink rectangle area: the range of F1 scores of 10 replicate runs; black dashed line: the average F1 score of 1,000 random guesses for KNN, RF, and XGBoost models and 100 random guesses for AutoGluon models. *z*: *z*-score of the average F1 score of 10 replicate runs in the background distribution; *p*: the corresponding *p*-value of the *z*-score.

Fig. S6. Prediction accuracy of MPGs within individual metabolic classes for KNN and RF

models. **(A,B)** $F1_{CV}$ (left panel) and $F1_{test}$ (right) scores for individual metabolic classes for KNN

(A) and RF **(B)** models when bulk RNA-seq (blue) and scRNA-seq (orange) data were used.

Error bars indicate the standard deviation of results from 10 replicate runs. Asterisk(s) indicate(s)

significant levels for the two-sided Wilcoxon rank-sum test: *, p-value < 0.05; **, p-value < 0.01;

***: p-value < 0.001. **(C,D)** z-scores of $F1_{CV}$ (left panel) and $F1_{test}$ (right) scores for 13 metabolic

classes compared with values of random guess for 1,000 times for KNN **(C)** and RF **(D)** models.

A z-score above the dashed line (z-score=1.645, p-value=0.05) indicates that the model's

performance is significantly better than random guess. **(E,F)** The correlation of $F1_{CV}$ between

models built using XGBoost (x-axis) and RF (y-axis in **E**), and between models built using

XGBoost and AutoGluon (y-axis in **F**), when bulk RNA-seq (left panel) and scRNA-seq (right)

data were used. PCC: Pearson correlation coefficient. EM: energy metabolism; Cof: cofactors;

IM: intermediate metabolism; DTX: detoxification; AA: amino acids; NT: nucleotides; AP:

amines and polyamines; Carb: carbohydrates; FAL: fatty acids and lipids; IN: inorganic nutrients;

SM: specialized metabolism; Horm: hormones.

Fig. S7. $F1_{CV}$ (A) and $F1_{test}$ (B) scores of individual classes for KNN models built using bulk

RNA-seq data. The density plot shows the distribution of F1 scores for models where the labels

of benchmark MPGs in the training set were randomly shuffled (random guesses). Red solid line:

the average F1 score for models built using the true labels of MPGs; pink rectangle area: the

range of F1 scores of 10 replicate runs; black dashed line: the average F1 score of 1,000 random

guesses. z: z-score of the average F1 score of 10 replicate runs in the background distribution; p:

the corresponding p-value of the z-score.

Fig. S8. $F1_{CV}$ (A) and $F1_{test}$ (B) scores of individual classes for KNN models built using

scRNA-seq data. The density plot shows the distribution of F1 scores for models where the

labels of benchmark MPGs in the training set were randomly shuffled (random guesses). Red

solid line: the average F1 score for models built using the true labels of MPGs; pink rectangle

area: the range of F1 scores of 10 replicate runs; black dashed line: the average F1 score of 1,000

random guesses. z: z-score of the average F1 score of 10 replicate runs in the background

distribution; p: the corresponding p-value of the z-score.

Fig. S9. F1$_{CV}$ (A) and F1$_{test}$ (B) scores of individual classes for RF models built using bulk RNA-seq data. The density plot shows the distribution of F1 scores for models where the labels of benchmark MPGs in the training set were randomly shuffled (random guesses). Red solid line: the average F1 score of models built using the true labels of MPGs; pink rectangle area: the range of F1 scores of 10 replicate runs; black dashed line: the average F1 score of 1,000 random guesses. z: z-score of the average F1 score of 10 replicate runs in the background distribution; p: the corresponding p-value of the z-score.

Fig. S10. F1$_{CV}$ (A) and F1$_{test}$ (B) scores of individual classes for RF models built using scRNA-seq data. The density plot shows the distribution of F1 scores for models where the labels of benchmark MPGs in the training set were randomly shuffled (random guesses). Red solid line: the average F1 score of models built using the true labels of MPGs; pink rectangle area: the range of F1 scores of 10 replicate runs; black dashed line: the average F1 score of 1,000 random guesses. z: z-score of the average F1 score of 10 replicate runs in the background distribution; p: the corresponding p-value of the z-score.

Fig. S11. F1$_{CV}$ (A) and F1$_{test}$ (B) scores of individual classes for XGBoost models built using bulk RNA-seq data. The density plot shows the distribution of F1 scores for models where the labels of benchmark MPGs in the training set were randomly shuffled (random guesses). Red solid line: the average F1 score of models built using the true labels of MPGs; pink rectangle area: the range of F1 scores of 10 replicate runs; black dashed line: the average F1 score of 1,000 random guesses. z: z-score of the average F1 score of 10 replicate runs in the background distribution; p: the corresponding p-value of the z-score.

Fig. S12. F1$_{CV}$ (A) and F1$_{test}$ (B) scores of individual classes for XGBoost models built using scRNA-seq data. The density plot shows the distribution of F1 scores for models where the labels of benchmark MPGs in the training set were randomly shuffled (random guesses). Red solid line: the average F1 score of models built using the true labels of MPGs; pink rectangle area: the range of F1 scores of 10 replicate runs; black dashed line: the average F1 score of 1,000

random guesses. z: z-score of the average F1 score of 10 replicate runs in the background
distribution; p: the corresponding p-value of the z-score.

Fig. S13. F1$_{CV}$ (A) and F1$_{test}$ (B) scores of individual classes for AutoGluon models built
using bulk RNA-seq data. The density plot shows the distribution of F1 scores for models
where the labels of benchmark MPGs in the training set were randomly shuffled (random
guesses). Red solid line: the average F1 score of models built using the true labels of MPGs; pink
rectangle area: the range of F1 scores of 10 replicate runs; black dashed line: the average F1
score of 100 random guesses. z: z-score of the average F1 score of 10 replicate runs in the
background distribution; p: the corresponding p-value of the z-score.

Fig. S14. F1$_{CV}$ (A) and F1$_{test}$ (B) scores of individual classes for AutoGluon models built
using scRNA-seq data. The density plot shows the distribution of F1 scores for models where
the labels of benchmark MPGs in the training set were randomly shuffled (random guesses). Red
solid line: the average F1 score of models built using the true labels of MPGs; pink rectangle
area: the range of F1 scores of 10 replicate runs; black dashed line: the average F1 score of 100
random guesses. z: z-score of the average F1 score of 10 replicate runs in the background
distribution; p: the corresponding p-value of the z-score.

Fig. S15. Performance of models built with four algorithms using the entire and subsets of
scRNA-seq data from five different tissues. KNN (**A**), RF (**B**), XGBoost (**C**), and AutoGluon
(**D**) models were built using the K-Nearest Neighbors, Random Forest, eXtreme Gradient
Boosting, and AutoGluon-Tabular algorithms, respectively. Left panel: F1$_{CV}$; right panel: F1$_{test}$.
Models built using different scRNA-seq datasets are indicated using different colors: dark blue,
the entire scRNA-seq data; dark green, scRNA-seq data from root; green, those from shoot; olive,
leaf; orange, flower; pink, embryo. Error bars indicate the standard deviation of results from 10
replicate runs. Asterisk(s) indicate(s) significant levels for the two-sided Wilcoxon rank-sum test,
showing that the models' performance using tissue subset data is significantly higher than when
using the entire scRNA-seq data: *, p-value < 0.05; **, p-value < 0.01; ***: p-value < 0.001. EM:
energy metabolism; Cof: cofactors; IM: intermediate metabolism; DTX: detoxification; AA:

amino acids; NT: nucleotides; AP: amines and polyamines; Carb: carbohydrates; FAL: fatty
acids and lipids; IN: inorganic nutrients; SM: specialized metabolism; Horm: hormones.

Fig. S16. Performance of models built with four algorithms using bulk RNA-seq (blue),
scRNA-seq data (orange) and the combined data (gray). KNN (**A**), RF (**B**), XGBoost (**C**),
and AutoGluon (**D**) models were built using K-Nearest Neighbors, Random Forest, eXtreme
Gradient Boosting, and AutoGluon-Tabular algorithms, respectively. Left panel: $F1_{CV}$; right
panel: $F1_{test}$. Error bars indicate the standard deviation of results from 10 replicate runs.
Asterisk(s) indicate(s) significant levels for the two-sided Wilcoxon rank-sum test: *, p-value <
0.05; **, p-value < 0.01; ***: p-value < 0.001. The asterisks are only marked for the classes for
which the combined data-based models significantly outperformed both models built using bulk
RNA-seq and scRNA-seq data. EM: energy metabolism; Cof: cofactors; IM: intermediate
metabolism; DTX: detoxification; AA: amino acids; NT: nucleotides; AP: amines and
polyamines; Carb: carbohydrates; FAL: fatty acids and lipids; IN: inorganic nutrients; SM:
specialized metabolism; Horm: hormones.

Fig. S17. Global feature importance for RF (A), XGBoost (B) and AutoGluon (C) models
built using bulk RNA-seq data. Bar chart shows the average importance of bulk RNA-seq
features for 10 replicate runs; bar colors indicate different tissue types to which the features
belong: dark green, root; green, shoot; olive, leaf; orange, flower; pink, embryo; light pink,
others.

Fig. S18. Global feature importance for RF (A), XGBoost (B) and AutoGluon (C) models
built using scRNA-seq data. The bar chart shows the average importance of scRNA-seq
features for 10 replicate runs; bar colors indicate different tissue types to which the features
belong: dark green, root; green, shoot; olive, leaf; orange, flower; pink, embryo; light pink,
others.

Fig. S19. Feature importance for individual metabolic classes for AutoGluon models when
scRNA-seq (A) and bulk RNA-seq (B) data were used. Colors within the heatmaps indicate
the average importance of features for 10 replicate runs; the background colors of the texts

indicate different tissue types to which the features belong: dark green, root; green, shoot; olive, leaf; orange, flower; pink, embryo; light pink, others.

Fig. S20. Prediction of metabolic classes for unknown genes by XGBoost models. (**A**) Count of unknown genes that were predicted into 13 metabolic classes by XGBoost models built using bulk RNA-seq (upper panel) and scRNA-seq data (lower). (**B**) Venn diagrams showing the overlap of unknown genes that were predicted into metabolic classes by XGBoost models built using bulk RNA-seq (blue) and scRNA-seq data (orange), and those that were annotated as within metabolic classes in the PMN database (purple). In the upper and lower panels, overlapping regions show the number of genes with exactly the same classes or at least one common class between datasets, respectively. (**C,D**) Heatmaps showing the numbers of unknown genes that were predicted as in a class (x-axis) by XGBoost models built using bulk RNA-seq (**C**) and scRNA-seq data (**D**), and were also annotated as within the same or other classes (y-axis) by PMN database. Values in the column "Not-assigned" indicate the numbers of unknown genes that were predicted into neither any of the 13 metabolic classes nor the "Others" category. Cell colors in (**C,D**) indicate the numbers of genes. EM: energy metabolism; Cof: cofactors; IM: intermediate metabolism; DTX: detoxification; AA: amino acids; NT: nucleotides; AP: amines and polyamines; Carb: carbohydrates; FAL: fatty acids and lipids; IN: inorganic nutrients; SM: specialized metabolism; Horm: hormones.

Supplementary tables

Table S1. Information of 13 single-cell RNA sequencing (scRNA-seq) datasets.

Table S2. Information of five bulk RNA-seq datasets.

Table S3. Gene expression matrix of scRNA-seq data.

Table S4. Gene expression matrix of bulk RNA-seq data.

Table S5. 10,572 enzyme-encoding genes obtained from the PMN, TAIR databases, or annotated with EC numbers using the E2P2 software.

Table S6. 14 gene co-expression modules determined by WGCNA using bulk RNA-seq data.

Table S7. 14 gene co-expression modules determined by WGCNA using scRNA-seq data.

Table S8. Clustering coefficients of metabolic pathway genes within each class based on two types of RNA-seq data and four tissue-specific scRNA-seq data.

Table S9. Clustering coefficients for 10,000 random simulations calculated using two types of
RNA-seq data.

Table S10. Clustering coefficients for 10,000 random simulations calculated using four tissue-
specific scRNA-seq data.

Table S11. Overall $F1_{CV}$ and $F1_{test}$ scores for 1,000 random simulations of KNN, RF, and
XGBoost models and for 100 random simulations of AutoGluon model when bulk RNA-seq data
were used.

Table S12. Overall $F1_{CV}$ and $F1_{test}$ scores for 1,000 random simulations of KNN, RF, and
XGBoost models and for 100 random simulations of AutoGluon model when scRNA-seq data
were used.

Table S13. $F1_{CV}$ and $F1_{test}$ scores for 1,000 random simulations of KNN models when bulk
RNA-seq data were used.

Table S14. $F1_{CV}$ and $F1_{test}$ scores for 1,000 random simulations of KNN models when scRNA-
seq data were used.

Table S15. $F1_{CV}$ and $F1_{test}$ scores for 1,000 random simulations of RF models when bulk RNA-
seq data were used.

Table S16. $F1_{CV}$ and $F1_{test}$ scores for 1,000 random simulations of RF models when scRNA-seq
data were used.

Table S17. $F1_{CV}$ and $F1_{test}$ scores for 1,000 random simulations of XGBoost models when bulk
RNA-seq data were used.

Table S18. $F1_{CV}$ and $F1_{test}$ scores for 1,000 random simulations of XGBoost models when
scRNA-seq data were used.

Table S19. $F1_{CV}$ and $F1_{test}$ scores for 100 random simulations of AutoGluon models when bulk
RNA-seq data were used.

Table S20. $F1_{CV}$ and $F1_{test}$ scores for 100 random simulations of AutoGluon models when
scRNA-seq data were used.

Table S21. Global feature importance of 40 bulk RNA-seq features for RF models.

Table S22. Global feature importance of 146 scRNA-seq features for RF models.

Table S23. Global feature importance of 40 bulk RNA-seq features for XGBoost models.

Table S24. Global feature importance of 146 scRNA-seq features for XGBoost models.

Table S25. Feature importance for individual metabolic classes of 40 bulk RNA-seq features for
AutoGluon models.

Table S26. Feature importance for individual metabolic classes of 146 scRNA-seq features for
AutoGluon models.

Table S27. Prediction of 9,441 unknown genes into individual metabolic classes by the
AutoGluon model when bulk RNA-seq data were used.

Table S28. Prediction of 9,441 unknown genes into individual metabolic classes by the
AutoGluon model when scRNA-seq data were used.

Table S29. Prediction of 9,441 unknown genes into individual metabolic classes by the
XGBoost model when bulk RNA-seq data were used.

Table S30. Prediction of 9,441 unknown genes into individual metabolic classes by the
XGBoost model when scRNA-seq data were used.

References

Ahmadi MK, Fang L, Moscatello N, Pfeifer BA. **2016**. *E. coli* metabolic engineering for gram scale
production of a plant-based anti-inflammatory agent. *Metabolic Engineering* **38**: 382–388.

Al-Khayri JM, Rashmi R, Toppo V, Chole PB, Banadka A, Sudheer WN, Nagella P, Shehata WF,
Al-Mssallem MQ, Alessa FM, Almaghasla MI, Rezk AA-S. **2023**. Plant secondary metabolites:
the weapons for biotic stress management. *Metabolites* **13**: 716.

Appelhagen I, Wulff-Vester AK, Wendell M, Hvoslef-Eide A-K, Russell J, Oertel A, Martens S,
Mock H-P, Martin C, Matros A. **2018**. Colour bio-factories: towards scale-up production of
anthocyanins in plant cell cultures. *Metabolic Engineering* **48**: 218–232.

Bhambhani S, Kondhare KR, Giri AP. **2021**. Diversity in chemical structures and biological properties
of plant alkaloids. *Molecules* **26**: 3374.

Breiman L. **2001**. Random Forests. *Machine Learning* **45**: 5–32.

Chae L, Kim T, Nilo-Poyanco R, Rhee SY. **2014**. Genomic signatures of specialized metabolism in
plants. *Science* **344**: 510–513.

Chen T, Guestrin C. **2016**. XGBoost: a scalable tree boosting system. In: Proceedings of the 22nd ACM
SIGKDD International Conference on Knowledge Discovery and Data Mining. San Francisco
California USA: ACM, 785–794.

Daloso DDM, Morais EG, Oliveira E Silva KF, Williams TCR. **2023**. Cell-type-specific metabolism in
plants. *The Plant Journal* **114**: 1093–1114.

Dong X, Gao Y, Chen W, Wang W, Gong L, Liu X, Luo J. **2015**. Spatiotemporal distribution of
phenolamides and the genetics of natural variation of hydroxycinnamoyl spermidine in rice.
Molecular Plant **8**: 111–121.

Elshafie HS, Camele I, Mohamed AA. **2023**. A comprehensive review on the biological, agricultural
and pharmaceutical properties of secondary metabolites based-plant origin. *International Journal*
of Molecular Sciences **24**: 3266.

Erickson N, Mueller J, Shirkov A, Zhang H, Larroy P, Li M, Smola A. **2020**. AutoGluon-tabular:
robust and accurate AutoML for structured data. https://doi.org/10.48550/arXiv.2003.06505.

Fang C, Fernie AR, Luo J. **2019**. Exploring the diversity of plant metabolism. *Trends in Plant Science*
24: 83–98.

Guo G, Wang H, Bell D, Bi Y, Greer K. **2003**. KNN model-based approach in classification. In: On The
Move to Meaningful Internet Systems: CoopIS DOA and ODBASE. Berlin, Germany: Springer,
986–996.

Hagberg A, Swart PJ, Schult DA. **2008**. Exploring network structure, dynamics, and function using
networkx. In: Proceedings of the 7th Python in Science Conference. Los Alamos National
Laboratory (LANL), Los Alamos, NM (United States), 11–15.

Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, Lee MJ, Wilk AJ, Darby C,
Zager M, Hoffman P, Stoeckius M, Papalexi E, Mimitou EP, Jain J, Srivastava A, Stuart T,
Fleming LM, Yeung B, Rogers AJ, McElrath JM, Blish CA, Gottardo R, Smibert P, Satija
R. **2021**. Integrated analysis of multimodal single-cell data. *Cell* **184**: 3573–3587.

Jucá MM, Cysne Filho FMS, De Almeida JC, Mesquita DDS, Barriga JRDM, Dias KCF, Barbosa
TM, Vasconcelos LC, Leal LKAM, Ribeiro JE, Vasconcelos SMM. **2020**. Flavonoids:
biological activities and therapeutic potential. *Natural Product Research* **34**: 692–705.

Kaiser M. **2008**. Mean clustering coefficients: the role of isolated nodes and leafs on clustering measures
for small-world networks. *New Journal of Physics* **10**: 083042.

Kautsar SA, Suarez Duran HG, Blin K, Osbourn A, Medema MH. **2017**. plantiSMASH: automated
identification, annotation and expression analysis of plant biosynthetic gene clusters. *Nucleic*
Acids Research **45**: W55–W63.

Ke M, Elshenawy B, Sheldon H, Arora A, Buffa FM. **2022**. Single cell RNA-sequencing: a powerful
yet still challenging technology to study cellular heterogeneity. *BioEssays* **44**: 2200084.

Kharchenko PV. **2021**. The triumphs and limitations of computational methods for scRNA-seq. *Nature*
Methods **18**: 723–732.

Li Y, Leveau A, Zhao Q, Feng Q, Lu H, Miao J, Xue Z, Martin AC, Wegel E, Wang J, Orme A, Rey
M-D, Karafiátová M, Vrána J, Steuernagel B, Joynson R, Owen C, Reed J, Louveau T,

Stephenson MJ, Zhang L, Huang X, Huang T, Fan D, Zhou C, Tian Q, Li W, Lu Y, Chen J, Zhao Y, Lu Y, Zhu C, Liu Z, Polturak G, Casson R, Hill L, Moore G, Melton R, Hall N, Wulff BBH, Doležel J, Langdon T, Han B, Osbourn A. 2021. Subtelomeric assembly of a multi-gene pathway for antimicrobial defense compounds in cereals. *Nature Communications* **12**: 2563.

Magnard J-L, Roccia A, Caissard J-C, Vergne P, Sun P, Hecquet R, Dubois A, Hibrand-Saint Oyant L, Jullien F, Nicolè F, Raymond O, Huguet S, Baltenweck R, Meyer S, Claudel P, Jeauffre J, Rohmer M, Foucher F, Hugueney P, Bendahmane M, Baudino S. 2015. Biosynthesis of monoterpene scent compounds in roses. *Science* **349**: 81–83.

McMahon JM, White WL, Sayre RT. 1995. Cyanogenesis in cassava (*Manihot esculenta* crantz). *Journal of Experimental Botany* **46**: 731–741.

Mergner J, Frejno M, List M, Papacek M, Chen X, Chaudhary A, Samaras P, Richter S, Shikata H, Messerer M, Lang D, Altmann S, Cyprys P, Zolg DP, Mathieson T, Bantscheff M, Hazarika RR, Schmidt T, Dawid C, Dunkel A, Hofmann T, Sprunck S, Falter-Braun P, Johannes F, Mayer KFX, Jürgens G, Wilhelm M, Baumbach J, Grill E, Schneitz K, Schwechheimer C, Kuster B. 2020. Mass-spectrometry-based draft of the arabidopsis proteome. *Nature* **579**: 409–414.

Moore BM, Wang P, Fan P, Leong B, Schenck CA, Lloyd JP, Lehti-Shiu MD, Last RL, Pichersky E, Shiu S-H. 2019. Robust predictions of specialized metabolism genes through machine learning. *Proceedings of the National Academy of Sciences, USA* **116**: 2344–2353.

Onnela J-P, Saramäki J, Kertész J, Kaski K. 2005. Intensity and coherence of motifs in weighted complex networks. *Physical Review E, Statistical, Nonlinear, and Soft Matter Physics* **71**: 065103.

Opsahl T, Panzarasa P. 2009. Clustering in weighted networks. *Social Networks* **31**: 155–163.

Papadopoulou K, Melton RE, Leggett M, Daniels MJ, Osbourn AE. 1999. Compromised disease resistance in saponin-deficient plants. *Proceedings of the National Academy of Sciences, USA* **96**: 12923–12928.

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M, Prettenhofer P, Weiss R, Dubourg V, Vanderplas J, Passos A, Cournapeau D. 2011. Scikit-learn: machine learning in python. *Journal of Machine Learning Research* **12**: 2825–2830.

Schilmiller AL, Last RL, Pichersky E. 2008. Harnessing plant trichome biochemistry for the production of useful compounds. *The Plant Journal* **54**: 702–711.

Schläpfer P, Zhang P, Wang C, Kim T, Banf M, Chae L, Dreher K, Chavali AK, Nilo-Poyanco R, Bernard T, Kahn D, Rhee SY. 2017. Genome-wide prediction of metabolic enzymes, pathways, and gene clusters in plants. *Plant Physiology* **173**: 2041–2059.

Shang Y, Ma Y, Zhou Y, Zhang H, Duan L, Chen H, Zeng J, Zhou Q, Wang S, Gu W, Liu M, Ren J,
Gu X, Zhang S, Wang Y, Yasukawa K, Bouwmeester HJ, Qi X, Zhang Z, Lucas WJ, Huang
S. **2014**. Biosynthesis, regulation, and domestication of bitterness in cucumber. *Science* **346**:
1084–1088.

Song Q, Ando A, Jiang N, Ikeda Y, Chen ZJ. **2020**. Single-cell RNA-seq analysis reveals ploidy-
dependent and cell-specific transcriptome changes in *Arabidopsis* female gametophytes. *Genome*
Biology **21**: 178.

Sun S, Shen X, Li Y, Li Y, Wang S, Li R, Zhang H, Shen G, Guo B, Wei J, Xu J, St-Pierre B, Chen
S, Sun C. **2022**. Single-cell RNA sequencing provides a high-resolution roadmap for
understanding the multicellular compartment of specialized metabolism. *Nature Plants* **9**:
179–190.

Tiwari R, Rana CS. **2015**. Plant secondary metabolites: a review. *International Journal of Engineering*
Research and General Science **3**: 661–670.

Tohge T, Scossa F, Wendenburg R, Frasse P, Balbo I, Watanabe M, Alseekh S, Jadhav SS, Delfin
JC, Lohse M, Giavalisco P, Usadel B, Zhang Y, Luo J, Bouzayen M, Fernie AR. **2020**.
Exploiting natural variation in tomato to define pathway structure and metabolic regulation of
fruit polyphenolics in the lycopersicum complex. *Molecular Plant* **13**: 1027–1046.

Töpfer N, Fuchs L-M, Aharoni A. **2017**. The PhytoClust tool for metabolic gene clusters discovery in
plant genomes. *Nucleic Acids Research* **45**: 7049–7063.

Wang P, Moore BM, Uygun S, Lehti-Shiu MD, Barry CS, Shiu S. **2021**. Optimising the use of gene
expression data to predict plant metabolic pathway memberships. *New Phytologist* **231**: 475–489.

Yang Y-H, Mao J-W, Tan X-L. **2020**. Research progress on the source, production, and anti-cancer
mechanisms of paclitaxel. *Chinese Journal of Natural Medicines* **18**: 890–897.

Yu C, Hou K, Zhang H, Liang X, Chen C, Wang Z, Wu Q, Chen G, He J, Bai E, Li X, Du T, Wang
Y, Wang M, Feng S, Wang H, Shen C. **2023**. Integrated mass spectrometry imaging and
single-cell transcriptome atlas strategies provide novel insights into taxoid biosynthesis and
transport in *Taxus mairei* stems. *The Plant Journal* **115**: 1243–1260.

Zhang B, Horvath S. **2005**. A general framework for weighted gene co-expression network analysis.
Statistical Applications in Genetics and Molecular Biology **4**: 1–45.

Zhang S, Ye T, Liu Y, Hou G, Wang Q, Zhao F, Li F, Meng Q. **2023**. Research advances in clinical
applications, anticancer mechanism, total chemical synthesis, semi-synthesis and biosynthesis of
paclitaxel. *Molecules* **28**: 7517.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5