Cerebral Small Vessel Disease genetic determinant TRIM47 controls brain homeostasis *via* the NRF2 antioxidant system

Authors: Cécile Duplàa[1], Valentin Delobel[1], Romain Boulestreau[1], Sébastien Rubin[1], Juliette Vaurs[1], Béatrice Jaspard-Vinassa[1], Muriel Busson[1], Cloé Combrouze[1], Carole Proust[3], Jean-Luc Morel[2], Bruno Bontempi[2], Aniket Mishra[3], Stéphanie Debette[3], Thierry Couffinhal[1,#] and Claire Peghaire[1,#,*].

[1] Univ. Bordeaux, INSERM, Biologie des maladies cardiovasculaires, U1034, F-33600 Pessac, France

[2] CNRS, IMN, UMR 5293, University Bordeaux, F-33000, Bordeaux, France

[3] Univ. Bordeaux, INSERM, Bordeaux Population Health, U1219, F-33000 Bordeaux, France

[#] These authors contributed equally to this work

[*] email of corresponding author: claire.peghaire@u-bordeaux.fr

Abstract

Cerebral small vessel disease (SVD) is a leading cause of strokes and a significant contributor to dementia, yet the precise mechanisms underlying its pathogenesis remain elusive. In a recent whole-exome association study involving population cohorts with SVD, we identified a variant on *TRIM47* locus. The ubiquitin ligase *TRIM47* is highly expressed in brain endothelial cells (ECs), suggesting its potential role in blood brain barrier (BBB) integrity. Here, we show that TRIM47 regulates brain ECs resilience and adaptive responses to oxidative stress by binding to KEAP1, stabilizing NRF2 protein levels and promoting NRF2 antioxidant signaling pathway. *In vivo*, *Trim47*-deficient mice exhibit downregulation of NRF2 target genes, BBB dysfunction, astrogliosis and cognitive impairments. Endothelial-specific deletion of *Trim47* recapitulates this vascular dementia phenotype, emphasizing the critical role of endothelial TRIM47 in protecting brain function. Treatment with the NRF2 activator

tert-butylhydroquinone normalized BBB integrity and cognitive function in *Trim47*-deficient mice, highlighting the role of TRIM47 in driving brain homeostasis through NRF2 pathway activation. This work indicates that the loss of the protective TRIM47/NRF2 axis may increase the susceptibility to developing human SVD and that targeting the TRIM47/NRF2 axis could offer novel therapeutic strategies for vascular dementia and related neurovascular diseases.

Introduction

Cerebral small vessel disease (SVD) encompasses a range of pathological processes that damage arterioles, capillaries or venules supplying brain tissue. It is the most common, chronic and progressive vascular disease. It is a leading cause of strokes (25%), a major contributor to cognitive decline (45%) and a common feature of vascular dementia and Alzheimer disease. SVD is highly prevalent in the population and is often detectable on brain imaging before clinical symptoms manifest[1,2]. In fact, this disease covers a spectrum of pathological conditions that impair the brain microvasculature leading to diverse brain damages, as evidenced by magnetic resonance key features such as white matter hyperintensities, microbleeds, infarcts or enlarged perivascular spaces[3].

Despite the prevalence of SVD and its significant impact on public health, particularly with aging populations, the molecular pathways driving its etiology remain poorly understood. Various risk factors, including hypertension and aging, are associated with SVD, but no established therapeutic strategies currently exist for its prevention or treatment. Current prevention efforts mainly focus on lowering blood pressure and reducing known cardiovascular risk factors (beta blockers, calcium channel blockers, statins, ACE inhibitors, aspirin…). However, these interventions are insufficient for stopping the progression of SVD and improving cognitive function, especially in older adults[4,5]. Thus, understanding the

precise molecular and cellular mechanisms underlying SVD pathophysiology is an absolute prerequisite for identifying novel targets, developing new effective therapeutic strategies and possibly accelerating precision medicine.

To identify novel potential genetic determinants of SVD, we recently performed a genome-wide association study and whole-exome sequencing on population-based cohorts of extreme MRI-defined SVD[6]. This study identified a probably damaging common missense variant in the Tripartite Motif 47 (*TRIM47*) locus and highlighted an inverse relation between TRIM47 expression in the brain blood vessels and extensive SVD severity. Moreover, transcriptomic data showing an enrichment of TRIM47 expression in brain blood vessels and endothelial cells (ECs) as well as preliminary functional evaluation suggest a putative role of TRIM47 in human brain ECs[6]. In the present study, we wanted to understand the link between TRIM47 and SVD.

TRIM47 belongs to a subfamily of E3 ubiquitin ligases involved in many cellular and physiological processes including cell proliferation, apoptosis, innate immunity and autophagy. The E3 ubiquitin ligases regulate protein degradation, cell trafficking and are also involved in DNA repair[7,8]. TRIM47 was first identified as being overexpressed in astrocytoma[9] and its role in cancers has been studied extensively. As an oncogene, TRIM47 promotes tumorigenesis, and is overexpressed in multiple cancers (breast[10], ovarian[11], colorectal[12], glioma[13,14], renal[15], gastric cancers[16]). It is also used as a prognostic biomarker in the cancer field[16–19].

Regarding its vascular role, previous works demonstrated that TRIM47 regulate cellular functions associated with angiogenesis *in vitro*[13], with its expression being upregulated under various pathological conditions, including inflammation, hypoxia, and oxidative stress

in human umbilical vein endothelial cells[20]. More recently, few studies have further suggested a potential role for TRIM47 in vascular biology during pathological conditions. *In vivo*, deletion of *Trim47* alleviates inflammation in a mouse model of LPS-induced sepsis[20], while overexpression of *Trim47* markedly accelerates cerebral ischemia injury though promoting apoptosis and inflammation in rats[21].

In this study, we identified the precise role of *Trim47* in brain physiology, demonstrating its major contribution to brain endothelial cell homeostasis, blood brain barrier integrity and cognitive function, through the regulation of the antioxidant Nrf2 pathway. Our data highlight the therapeutic potential of targeting the TRIM47/NRF2 protective axis in patients with SVD, as well as in broader contexts such as vascular dementia and Alzheimer's disease.

Results

TRIM47 is an important activator of the antioxidant NRF2 pathway in human brain endothelial cells

Our group, along with others, has recently reported that TRIM47 is not only expressed by various types of ECs, but is also particularly enriched in brain vessels and plays a crucial role in regulating ECs functions *in vitro*[6,13,20]. To identify the specific pathways regulated by TRIM47 in brain ECs, we performed bulk RNA-sequencing on human brain microvascular endothelial cells (HBMEC) treated with either control siRNA (siCtl) or *TRIM47*-targeting siRNA (si*TRIM47*) for 72 h. Transcriptomic data analysis revealed that 208 genes were downregulated while 214 genes were upregulated following *TRIM47* depletion (Fig. 1a). Pathway enrichment analysis revealed that the genes regulated by TRIM47, particularly those downregulated in *TRIM47*-deficient cells, were associated with the nuclear erythroid

2-related factor 2 (NRF2) pathway and oxidative stress (Fig. 1b). Interestingly, *TRIM47* depletion did not affect the mRNA levels of the transcription factor *NFE2L2* (NRF2) and *KEAP1*, a regulator of NRF2, but it significantly downregulated key NRF2 target genes including *HMOX1*, *NQO1* and *GCLM* (Fig. 1c). Further experiments showed that inhibition of TRIM47 expression in HBMEC by siRNA treatment reduced *NQO1* and *HMOX1* (HO1) mRNA levels over 24, 48 and 72 h (Supplementary Fig. 1a-c) with a corresponding decrease in HO1 protein levels (Fig. 1d).

Additionally, and as reported[20] under oxidative stress conditions induced by H_2O_2, *TRIM47* expression was upregulated but *NFE2L2* mRNA levels were not affected (Supplementary Fig. 1d-e). Crucially, TRIM47 was required to fully induce *HMOX1* and *NQO1* mRNA levels in both basal and oxidative stress conditions in HBMEC, mirroring the effects of *NRF2* siRNA (Fig. 1e-f). In line with these data, H_2O_2 treatment activated the HO1 promoter (HO1-4.5kb); *TRIM47* depletion caused a significant decrease in HO1 promoter activity in both baseline and oxidative stress conditions (Fig. 1g). Together, these data showed that TRIM47 is a key and novel activator of the NRF2 antioxidant pathway in human brain ECs.

TRIM47 regulates NRF2 stability *via* its interaction with KEAP1 in brain endothelial cells

To uncover the precise mechanism by which TRIM47 regulates the NRF2 pathway, we investigated the potential interaction between TRIM47 and the transcription factor NRF2 in brain ECs using *in vitro* models. *TRIM47* overexpression (Supplementary Fig. 2a) led to a modest increase in *NFE2L2* (NRF2) expression (Supplementary Fig. 2b) and to a significant increase in the expression of NRF2 target genes *NQO1* and *HMOX1* (Fig. 2a-b). This induction was abolished by *NRF2* siRNA treatment showing that TRIM47's effect on these genes is dependent on NRF2 (Fig. 2a-b).

Given that NRF2 is the master regulator of antioxidant defense gene expression and directly drives the expression of HO1 and NQO1 by binding to antioxidant response elements (ARE), on their promoters (reviewed in[22]), we sought to determine whether TRIM47 requires these NRF2 binding sites to increase HO1 expression. A luciferase assay in HeLa cells revealed that *TRIM47* overexpression activates the HO1 promoter (Fig. 2c). Mutations in the two ARE sites on the HO1 promoter (HO1-4.5kb 2 Mut, double mutant construct)[23,24] blocked HO1 promoter transactivation, underscoring the necessity of NRF2 binding to ARE sites for HO1 induction by TRIM47-mediated HO1 induction (Fig. 2c). As a control, a triple mutant construct with additional mutation in the E box of the HO1 promoter (HO1-4.5kb 3 Mut) was used which, as expected, fully blocked HO1 promoter transactivation (Fig. 2c).

The activity of NRF2, both under basal conditions and in response to stress, is tightly controlled. NRF2 protein levels are finely regulated by Kelch-like ECH-associated protein 1 (KEAP1), a master negative regulator that binds to NRF2, leading to its ubiquitination and subsequent degradation by the proteasome (reviewed in[22]). To investigate whether TRIM47 modulates NRF2 pathway through the formation of a regulatory complex, we performed co-immunoprecipitation assay (Co-IP) assays. Our data revealed TRIM47 interacts with KEAP1 in HeLa cells overexpressing both TRIM47-myc and Keap1-Flag (Fig. 2d), but not with NRF2 itself (data not shown). Importantly, although TRIM47 depletion in HBMEC did not affect KEAP1 protein levels, Western blot analysis, with and without a proteasome inhibitor treatment (MG132), showed a significant decrease in NRF2 protein (Fig. 2e). This suggests that TRIM47 may prevent NRF2 protein degradation by binding to KEAP1. ECs were then treated with an NRF2 pathway activator, namely the tert-butylhydroquinone (tBHQ), which promotes NRF2 protein stability by destabilizing the KEAP1-NRF2 complex. This NRF2 pathway activator did not affect the mRNA levels *of TRIM47* (Supplementary Fig. 2d) and

NFE2L2 (Supplementary Fig. 2e). Moreover and importantly, tBHQ was able to partly rescue the effect of *TRIM47* siRNA on *NQO1* (Fig. 2f) and *HMOX1* (Fig. 2g) expression, indicating that TRIM47 acts upstream of NRF2. As expected, tBHQ could not rescue the effects of *NRF2* siRNA on these target genes (Fig. 2f and g). These findings highlight TRIM47 as a key modulator of NRF2 stability through its interaction with KEAP1.

In summary, these data indicate that under basal conditions when TRIM47 is expressed in ECs, it binds to KEAP1 in the cytoplasm, thereby reducing NRF2 protein degradation and enhancing its stability. NRF2 can then accumulate in the cytoplasm and translocate to the nucleus to induce the transcription of its antioxidant target genes. Conversely, when TRIM47 is repressed, KEAP1 is free to bind to NRF2, leading to its ubiquitination and degradation by the proteasome, which decreases the NRF2-mediated antioxidant pathway in ECs. The NRF2 pathway can be reactivated by the use of the NRF2 activator tBHQ as depicted in the model (Fig. 2h).

TRIM47 protects from oxidative stress both *in vitro* and *in vivo*

Since TRIM47 promotes the antioxidant NRF2 pathway, we hypothesized that loss of TRIM47 in brain ECs might lead to a higher susceptibility to stress. Indeed, depletion of *TRIM47* in HBMEC led to a significant increase in oxidative stress as detected by a CellRox green dye. Additionally, *TRIM47* siRNA worsened the effect of two oxidative stress inducers, tert-butyl hydroperoxide (TBHP) and Angiotensin II (Supplementary Fig. 3a and b). Notably, activation of the NRF2 pathway by stabilizing NRF2 protein with the tBHQ compound was able to dampen the oxidative stress induced by *TRIM47* depletion under both basal and stress conditions (Fig. 3a-b). This indicates that TRIM47 contributes to antioxidant defenses in HBMEC, at least in part by activating the NRF2 pathway.

To explore the antioxidant potential of TRIM47 *in vivo*, we used mice deleted for the *Trim47* gene across all tissues and cell types (knock out (KO) mice also denoted *Trim47-/-* mice) (Supplementary Fig. 4a). *Trim47-/-* mice were viable and fertile (Supplementary Fig. 4b), developed normally, showing no significant difference in weight compared to their littermate control littermates (*Trim47+/+* mice) (Supplementary Fig. 4c-e) and displaying no obvious macroscopic phenotype. The efficacy of *Trim47* deletion was confirmed at the mRNA level by qPCR on brain lysates and isolated brain ECs from *Trim47-/-* and *Trim47+/+* mice (Fig. 3c-d and Supplementary Fig. 4f).

To challenge these KO mice, we employed a pathological model of oxidative stress induced by hypoxia, namely the oxygen-induced retinopathy (OIR) model, which is characterized by increased reactive oxygen species production[25,26]. qPCR screening of *Nfe2l2* and its target genes on brain tissues from *Trim47+/+* control mice exposed to either room air or the OIR model, confirmed that this pathological model activates the NRF2 protective antioxidant system in the central nervous system (Fig. 3c). Notably, the deletion of *Trim47* resulted in a decreased in the *Nrf2* pathway under both baseline and pathological (OIR) conditions (Fig. 3c). A similar reduction in Nrf2 pathway was also observed in brain ECs isolated from *Trim47+/+ and Trim47-/-* mice at postnatal day 16, following exposure to the OIR model (Fig. 3d). Additionally, histological analysis of the retinas at P16 indicated that the loss of *Trim47* was associated with an increase in avascular areas (Fig. 3e-f), suggesting that *Trim47* plays a protective role in this model.

Together, these results demonstrate that TRIM47 provides antioxidant properties to brain ECs *in vitro* and protects from oxidative stress in a mouse model of OIR, partly by promoting the Nrf2 pathway.

Trim47 deletion leads to cognitive impairments

Our recent analysis of extensive human genomic datasets has identified TRIM47 as a novel biological target linked to SVD[6]. To investigate the impact of *Trim47* deletion on cognitive functions and brain physiology, we submitted adult *Trim47* -/- mice and their control littermates to two complementary behavioral tests assessing spatial recognition memory in the Y-maze and spatial discrimination in the Morris water maze. Contrary to controls, male *Trim47* -/- mice (Fig. 4a-b) failed to recognize the unexplored (previously inaccessible) arm during the test phase of the Y-maze procedure, indicating an inability to form short-term spatial recognition memory. Adult *Trim47* -/- mice were also impaired in the cognitively-challenging water maze paradigm where they were unable to learn the spatial location of the hidden platform. Accordingly, latency and distance swum to the platform decreased over the 4 training days in control but not *Trim47* -/- mice (Fig. 4c-e). Potential confounds such as locomotor deficits were absent in the *Trim47* -/- whose swim speed was comparable to that of controls (Fig. 4f). Cognitive impairments were similar in female cohorts, suggesting that the phenotype observed in *Trim47*-deficient mice is not gender-specific (Fig. 4g-l).

Interestingly, the sucrose preference test did not reveal any sign of anhedonia, a key symptom of depression in humans, suggesting that *Trim47*-/- mice develop a purely cognitive impairment without any apparent anxiety-like symptoms (Supplementary Fig. 4g).

Trim47 deletion triggers BBB dysfunction associated with decreased Nrf2 pathway *in vivo*

The enrichment of *Trim47* expression in brain ECs[6] pointed towards its potential role in blood brain barrier (BBB) development and/or function. Notably, Glut1 immunostaining analysis in brain sections from adult *Trim47*-/- and control mice showed no differences in

vascular density in the cortical regions and hippocampus compared to control mice (Supplementary Fig. 5a-b). These data prove that *Trim47* is not essential for the development of brain blood vessels.

To determine whether the cognitive impairments observed in *Trim47* KO mice were linked to compromised BBB function, we assessed BBB integrity by injecting two different tracers. Quantification of cadaverine (Fig. 5a) and dextran (Fig. 5b) extravasation, expressed as the permeability index ratio, revealed a significantly increase in BBB leakiness in adult *Trim47-/-* mice compared to control mice. Importantly, tracer permeability in the kidney was not affected, suggesting that *Trim47* specifically acts on brain vasculature which is characterized by a specialized continuous endothelium (Fig. 5a-b). Immunostaining of brain sections with fibrinogen and podocalyxin antibodies, to visualize blood vessels, revealed the presence of fibrinogen, a blood component, outside the vessels in the hippocampus and cortical regions of *Trim47-/-* mice. In contrast, fibrinogen was confined within the vessels in control mice, confirming a localized increase in BBB permeability (in some capillaries and larger vessels) in *Trim47-/-* mice (Fig. 5c and Supplementary Fig. 5c).

To identify the cause of BBB defects, we analyzed the expression of key components of tight and adherent junctions which are essential for BBB integrity, in adult *Trim47-/-* and *Trim47+/+* mice. qPCR analysis of brain ECs (Fig. 5d) showed a mild but significant decrease in the expression of *Cldn5* (Claudin5), *Cldn11* (Claudin11), and *Ocln* (Occludin) in *Trim47-/-* mice compared to control mice (Fig. 5d). Additionally, there was a trend towards downregulation of *Cdh5* (VE-cadherin) in both brain ECs (Fig. 5d) and in whole brain lysates (Supplementary Fig. 5e). However, the mRNA expression level of *Tjp1* (ZO1) was not affected in *Trim47-/-* mice compared to their littermate controls (Fig. 5d). Immunofluorescence

staining of brain sections confirmed the reduced protein levels of Claudin5 in the hippocampus and cortical regions of *Trim47-/-* mice (Fig. 5e and Supplementary Fig. 5d). These data indicated that the increased BBB permeability observed in *Trim47-/-* mice is likely due to the decreased expression of specific tight/adherent junction components in brain ECs.

qPCR analysis confirmed the downregulation of the Nrf2 pathway in adult *Trim47-*deficient mice both in brain EC (Fig. 5f) and the whole brain (Supplementary Fig. 5e). This finding aligns with our *in vitro* and *in vivo* data from the hypoxia-induced oxidative stress model (OIR) at p16. Blood vessels are particularly vulnerable to localized oxidative stress, and reduced antioxidant protection in ECs may further contribute to BBB impairment. Our results demonstrate that the loss of BBB integrity in *Trim47-/-* mice is associated with decreased expression of tight junction proteins. Nrf2 has been reported as a positive transcriptional regulator of Claudin5 expression in ECs[27], suggesting that the decreased Nrf2 pathway in *Trim47*-deficient brain ECs could aggravate damage through the downregulation of Claudin5, thereby directly contributing to BBB dysfunction.

BBB impairment-induced by *Trim47* deletion leads to brain lesions *in vivo*

Since BBB disruption can lead to the leakage and accumulation of neurotoxic material or blood components (such as fibrinogen) in the brain parenchyma, we next wanted to determine whether BBB impairment in *Trim47-/-* mice resulted in additional brain defects. Astrocytes, which are in close contact with ECs, are highly plastic and can be directly affected by BBB dysfunction and changes in their local environment[28,29]. Supporting this idea, GFAP staining indicated a significant increase in astrocyte activation in the hippocampus, particularly in the dentate gyrus region of *Trim47-/-* mice where blood vessel leakiness was

observed (Fig. 5g).

As astrocytes reactivity can be modulated by neuroinflammation[28], we next assessed the expression of adhesion molecules and cytokines in brain ECs and in whole brain lysates. qPCR analysis revealed no evidence of a pro-inflammatory endothelial state or a global pro-inflammatory profile in *Trim47-/-* mice, as shown by a reduction in *Icam1* and *Vcam1* expression (Fig. 5d and Supplementary Fig. 5e). Additionally, we analyzed Cd11b mRNA expression levels (Supplementary Fig. 6a) and performed immunofluorescence staining with Iba1 (Supplementary Fig. 6b) and Cd68 (Supplementary Fig. 6c) antibodies in *Trim47+/+* and *Trim47-/-* mouse brain samples. No significant activation of microglia and perivascular macrophages or immune cells recruitment was observed following *Trim47* deletion *in vivo*.

Since *Trim47* mutant mice show cognitive impairments, we hypothesized that the loss of *Trim47* may affect the number of neurons. However, histological analyses of coronal brain sections revealed no evidence of detectable neuronal loss in *Trim47-/-* mice compared to controls, as shown by NeuN immunostaining in the hippocampus and cortical regions (Supplementary Fig. 6d-e). GWAS have identified the *TRIM47* locus as a top genetic risk factor for the formation of white matter hyperintensities (WMH) in humans[30]. To evaluate the impact of *Trim47* deletion on WM, we performed electron microscopy on brain tissues. The measurement of g-ratio[31] in the corpus callosum region did not reveal significant axonal demyelination in *Trim47-/-* mice compared with *Trim47+/+* mice (Fig. 5h). In line with this, myelin basic protein (MBP) staining did not show any significant reduction in myelination in the hippocampus of 10-months-old *Trim47-/-* mice (Fig. 5i), suggesting that *Trim47*-deficient mice do not display major detectable WM lesions at this stage.

Endothelial-specific deletion of *Trim47* is sufficient to recapitulate a vascular-dementia phenotype

To demonstrate that the phenotype observed in *Trim47-/-* mice was primarily due to the loss of endothelial *Trim47*, we generated mice deleted for *Trim47* in ECs (*Trim47*[iEC-KO]) and compared them with littermate controls (*Trim47*[iEC-WT]) using an inducible CDH5-Cre line (Supplementary Fig. 7a-d). The efficiency of *Trim47* deletion was confirmed on lung ECs isolated from *Trim47*[iEC-KO] *versus Trim47*[iEC-WT] adult mice (Fig. 6a). Behavioral tests conducted on adult mice revealed that male *Trim47*[iEC-KO] mice displayed impaired performance in the Y-maze (Fig. 6b-c) and water maze procedures (Fig. 6d-g) similar to the deficits observed in *Trim47-/-* mice. These finding confirm that the loss of endothelial *Trim47* is sufficient to induce cognitive impairment.

Importantly, the endothelial-specific deletion of *Trim47* also led to impaired BBB function, as shown by an increased BBB permeability index following 3kDa dextran injection (Fig. 6h) and a decreased expression of key adherent and tight junction components in brain ECs isolated from these mice (Fig. 6i). Finally, qPCR analysis also highlighted the downregulation of critical Nrf2 target genes, including *Hmox1* and *Slc7a11*, in brain ECs from *Trim47*[iEC-KO] compared to *Trim47*[iEC-WT] controls (Fig. 6j).

These results demonstrate that the loss of *Trim47* specifically in ECs recapitulates the vascular dementia phenotype observed in the global KO mouse, underscoring the crucial role of endothelial Trim47 in maintaining brain homeostasis.

Activation of the NRF2 pathway prevents brain defects and cognitive impairment observed in *Trim47*-deficient mice

Since we showed that TRIM47 is a key regulator of the NRF2 antioxidant pathway both *in vitro* and *in vivo*, we hypothesized that *Trim47* could promote brain homeostasis by activating the Nrf2 protective pathway in adult mice. To test this, we performed experiments on male *Trim47+/+* and *Trim47-/-* mice, feeding them a diet containing the Nrf2 activator tBHQ (1% w/w) (Fig. 7a). As previously reported[32,33], the tBHQ diet was well tolerated by the mice and did not induce any weight loss (Supplementary Fig. 8a). qPCR analysis confirmed that the tBHQ diet effectively prevented the impairment of the antioxidant pathway in *Trim47-/-* mice, both in brain ECs (Fig. 7b) and whole brain tissue (Supplementary Fig. 8b).

Notably, the expression of tight junction components in brain ECs was restored to normal levels in *Trim47-/-* mice treated with the Nrf2 activator compared to control mice (Fig. 7b). The diet with the antioxidant molecule prevented the increased BBB permeability (Fig. 7c), proving that antioxidant therapy successfully restored BBB integrity and function in *Trim47*-deficient mice.

In line with its known anti-inflammatory properties[34,35], tBHQ did not normalize the mRNA levels of the adhesion molecules *Icam1* and *Vcam1* in brain ECs isolated from *Trim47 -/-* mice (Supplementary Fig. 8c). Interestingly, the Nrf2 activator diet prevented astrocyte activation in *Trim47*-deficient mice, as shown by GFAP staining in the hippocampus (particularly in the dentate gyrus) (Fig. 7d).

Finally, we assessed whether increasing the Nrf2 pathway could also restore the cognitive function of *Trim47-/-* adult mice. Behavioral tests revealed that reactivation of the Nrf2 pathway *via* a tBHQ diet was sufficient to normalize the performances of the *Trim47-/-*

males in both the Y-maze (Fig. 7e-f) and water maze tests (Fig. 7g-h and Supplementary Fig. 8d-e), indicating a reversal of cognitive impairments.

Together, these data demonstrate that enhancing the Nrf2 pathway was sufficient to prevent BBB leakage, brain defects and cognitive impairment in *Trim47-/-* mice. This highlights the important role of *Trim47* in maintaining brain physiology and cognitive function, primarily through the activation of the Nrf2 antioxidant pathway.

Discussion

In this study, we report that the SVD genetic determinant TRIM47 activates the NRF2 pathway *in vitro* and *in vivo*. TRIM47 controls brain ECs resilience to oxidative stress by promoting NRF2 protein stability. *In vivo*, global deletion or loss of endothelial *Trim47* result in a spontaneous mild vascular dementia phenotype characterized by increased BBB permeability, brain lesions and cognitive dysfunction. Activation of the NRF2 pathway using tBHQ treatment prevents these defects, showing that the protective TRIM47/NRF2 axis is required to ensure cerebral homeostasis.

Despite progress in understanding the cause of SVD, specific signaling pathways leading to its development are still poorly understood. Current primary prevention strategies focusing on managing blood pressure and reducing risk factors are insufficient for slowing down the progression of SVD, which ultimately leads to cognitive decline. The urgent need for mechanism-based therapies is clear and could dramatically improve outcomes in stroke and dementia prevention[3–5]. Deciphering the molecular mechanisms underlying SVD etiology is essential for identifying novel targetable pathways. Our recent analyses of extensive human genomic datasets identified TRIM47 as a potential genetic determinant of SVD[6].

In the present study, we aim to investigate how this novel candidate selected from human data could contribute to brain pathophysiology using preclinical models. We found that adult mice (males and females) deleted for *Trim47* display spatial memory deficits associated with increased BBB permeability (cadaverin, dextran, fibrinogen leakages) and decreased expression of tight junction proteins (Claudin5, Occludin) in cortical and hippocampal regions, suggesting that this pathology is driven by early blood vessels' damages. In line with this concept, endothelial-specific deletion of *Trim47* was able to fully mimic this phenotype, showing that the loss of *Trim47* in brain ECs is primarily responsible for the development of the disease. To our knowledge, only eNOS hemi-deficiency which has been reported as a model of cerebral chronic hypoperfusion, also causes spontaneous mild cognitive impairment associated with BBB leakage in aged mice[36,37].

Additionally, we describe increased GFAP-positive astrocytes in *Trim47* deleted mice specifically in the dentate gyrus and in cortical regions where blood vessel's leakiness was detected. This astrogliosis in *Trim47*-deficient mice is presumably due to the localized extravasation of plasmatic protein such as fibrinogen. Supporting this hypothesis, previous elegant works from other groups have demonstrated the deleterious role of fibrinogen deposits into the CNS and its numerous neuropathological effects including astrocytes and microglia activation[38–40]. Notably, *Trim47-/-* mice do not show activation of microglia or clear sign of neuroinflammation. These data concur with previous studies reporting the role of *Trim47* in activating NF-kB signalling pathway and thus playing a pro-inflammatory role in different pathological models[20,21]. Together, these data point toward a dual role of TRIM47, which can activate the protective NRF2 system or promote the pro-inflammatory NF-kB pathway depending on the context and the model. Finally, while *Trim47-/-* mice exhibit localized BBB leakage and early damages of blood vessels, no major white matter lesion was

detectable at this stage, suggesting that challenging these mice with an additional pathological condition could be considered to reveal potential demyelination in this model.

Our *in vitro* data identified TRIM47 as a novel and important partner of the Nrf2 antioxidant system. TRIM47 acts upstream of NRF2 and activates this pathway by inducing NRF2 protein stability, potentially through its binding to its inhibitor KEAP1 in brain ECs. *In vivo*, our results confirmed TRIM47 regulation of the NRF2 system at postnatal age, in baseline and oxidative stress conditions, but also in adult mice. Crucially, we demonstrated that activation of the Nrf2 pathway with tBHQ is sufficient to prevent the BBB defects and the vascular dementia phenotype observed in *Trim47* mutant mice, proving that endothelial TRIM47/NRF2 axis is a pivotal antioxidant mechanism actively promoting brain homeostasis.

In fact, cerebral vasculature is one of the main targets of localized oxidative stress processes. Reactive oxygen species and oxidative stress have been identified as factors with a potentially strong impact on the etiology of cerebrovascular diseases[41]. If previously underestimated, age-associated vascular oxidative stress[42], impairment of NRF2 signaling pathway[43] and BBB disruption[44,45] are hypothesized to play a potential role in the pathogenesis of vascular dementia[46,47]. Nrf2 is a transcription factor that controls the expression of an array of antioxidant and detoxification enzymes[48] and has been shown recently to regulate the expression of tight junction proteins including Claudin5, even if the exact mechanism is not fully understood[27]. Importantly, Nrf2 deficiency exacerbates obesity-[49] or hypoperfusion-induced[50] BBB disruption and cognitive decline in mice. Conversely, Nrf2 activation with the antioxidant sulforaphane improves BBB function after cerebral ischemic stroke[51]. These data support our *in vivo* findings showing that the loss of *Trim47* induces Nrf2 pathway impairment, triggering primary blood vessels' insults, and leading ultimately to cognitive deficits. Together, our data demonstrate that *Trim47*-deficient mice display some

key features of human SVD making this tool an innovative preclinical model of vascular dementia that could be used by the scientific community in the vascular biology and neurosciences fields to better understand the pathophysiology of SVD. This could be a helpful model for testing the efficacy of new therapeutical strategies developed to tackle SVD or Alzheimer disease.

This work fully reveals the role of TRIM47 in brain physiology and the important contribution of TRM47/NRF2 pathway to SVD pathophysiology. Our findings suggest that patients with *TRIM47* mutation may be more susceptible to developing SVD by losing their NRF2 antioxidant protection, highlighting NRF2 targeting as a viable therapeutic strategy for managing vascular cognitive impairment. This work will hopefully pave the way to accelerate the use of interventions with NRF2 activators for preventing or treating SVD and more broadly vascular dementia.

Methods

Cell culture and treatments

Human brain microvascular endothelial cells (HBMEC) were purchased from SicenCell (#1000) and cultured in Endothelial Cell Growth Medium-2 media (EGM-2) (Lonza). HBMEC were transfected with lentivirus or siRNA in EGM-2 media (Lonza). HeLa (ATCC CCL-2) and Hek293 (ATCC CRL-1573) were cultured in DMEM (Dulbecco's Modified Eagle Medium, Thermo Fisher Scientific) supplemented with 10% FBS (fetal bovine serum) and 1% penicillin-streptomycin (Thermo Fisher Scientific). Cells were transfected with a pool of two siRNA against TRIM47 (siTRIM47#1 and #2, denoted as siTRIM47 in the text) (15nM for each TRIM47 siRNA, thus siRNAs at a final concentration of 30 nM) or with siRNA against NRF2 (15 nM) using INTERFERin® transfection reagent (Polyplus). Negative control siRNA

(Eurogentec) was used, which is denoted as siCont. Sequences of the siRNA are listed in **Supplementary Table 1**. Plasmids were transfected using jetPRIME (Polyplus) following the manufacturer's protocol. Lentiviruses were transduced at a multiplicity of infection of 30.

For some experiments, cells were stimulated with H2O2 (Sigma), TBHP (Luperox®, Sigma) or Angiotensin II (ANGII, Sigma, A9525) to induce oxidative stress conditions. In some instances, HBMEC were treated with the proteasome inhibitor MG-132 for 4 h (5 µM, M7449, Sigma) to inhibit protein degradation. For activation of the NRF2 pathway, cells were pre-treated with tert-butylhydroquinone (tBHQ) (Sigma, 112941) or dimethyl fumarate (DMF) (Santa Cruz, sc-239774).

Plasmids and lentiviruses

The plasmid pCMV6 TRIM47 (Myc-DDK-tagged) was purchased from Origene (RC218521). It contains the full sequence of human TRIM47 (NM_033452) which was then subcloned into the lentiviral vector pRRLsin-MND-MCS-WPRE. Preparations of lentivirus encoding for human TRIM47 were produced at the Bordeaux University lentivirus platform (TBMCore).

Flag-Keap1 was a gift from Qing Zhong (Addgene plasmid #28023; http://n2t.net/addgene:28023; RRID:Addgene_28023). pCDNA3-Myc3-Nrf2 was a gift from Yue Xiong (Addgene plasmid #21555; http://n2t.net/addgene:21555; RRID:Addgene_21555).

pHOGL3/4.5 and pHOGL3/4.5 Triple mutant plasmids were a gift from Prof. Anupam Agarwal (University of Alabama at Birmingham, US). The pHOGL3/4.5 double mutant was obtained from the pHOGL3/4.5 triple mutant in which mutation C>T was created into the mutated EBox by a rapid-site-directed-mutagenesis using the following non-overlapping primers (F: cacgtgacccgccgagcata; R: ggccaggcggaacagc) and Platinum™SuperFi™II DNA Polymerase

(Invitrogen). pGL3 Firefly Luciferase basic empty vector (Promega) was used as a control. pGL4.74 [hRluc/TK] vector] (Promega) Renilla luciferase vector was used as an internal control.

Immunoblotting analysis and co-immunoprecipitation

Mouse brains were collected; cortices were dissected and then frozen in liquid nitrogen. HBMEC or mouse cortices were lysed in RIPA buffer supplemented with protease and phosphatase inhibitors using a Tissue-Lyser (Qiagen). All protein lysates were then centrifuged for 15 min at $16000 \times g$ at 4 °C and supernatants were isolated. Proteins were loaded in 7 or 12% acrylamide gels and then transferred on a polyvinylidene difluoride (PVDF Immobilon membrane, Merck) membrane and incubated in PBS 0.1% Tween supplemented with 5% milk for 1 h to block non-specific binding. Immunoblots were next labeled with the primary antibodies listed in **Supplementary Table 2**. Primary antibodies were detected using fluorescently labeled secondary antibodies: anti-rabbit IgG Alexa Fluor™ 750 and anti-mouse IgG Alexa Fluor™ 700 (Fisher Scientific). Detection of fluorescence intensity was performed using an Odyssey imaging system (Li-COR, ScienceTec) and Odyssey ver.3 software. Fluorescence intensity was quantified by densitometry using Fiji-ImageJ software to measure protein levels and normalized against loading controls. See the source data file for the uncropped immunoblots.

For immunoprecipitation experiments, HEK293 were co-transfected with KEAP1-Flag (0.125 µg.ml−1) and TRIM47-myc-Flag (0.125 µg. ml−1). Proteins lysates were incubated with primary antibody at +4 °C, followed by incubation with Pierce™ Protein A/G Magnetic Agarose Beads (Thermo Fisher Scientific, 78609). Immunoprecipitation was carried out using

primaries antibodies at 2 µg for 500 µg of protein lysates. The primary antibodies used are listed in **Supplementary Table2**.

Reporter luciferase assay

HeLa were transfected with control or TRIM47 siRNA for 24h or transfected with either 0.1 µg of TRIM47 plasmid or pcDNA3.1 empty vector and with HO1 luciferase reporter plasmids or pGl3 vector for 24 h. Luciferase activity was measured using the Dual-Luciferase® Reporter Assay System (Promega) and a Spark® Multimode Microplate Reader (Tecan). Luciferase reporter activity was normalized to the internal Renilla luciferase control and is expressed relative to the pGL3 empty vector.

Real-time polymerase chain reaction

Total RNA from mouse tissues (brain and brain endothelial cells) and human cells (HBMEC and HeLa) was isolated by using the Direct-zol™ RNA MicroPrep kit (R2062, Zymo Research) and reverse transcribed into cDNA using the M-MLV Reverse Transcriptase (Promega). Quantitative real-time PCR was performed using the GoTaq qPCR master mix (Promega) on a QuantStudio™ 3 qPCR System (Thermo Fisher Scientific). Gene expression values were normalized to cyclophilin (PPIA) expression (human cells), *Pecam1* (mouse brain endothelial cells) or *18s* (mouse brain). See **Supplementary Tables 3 and 4** for the list of oligonucleotides.

Bulk RNA-sequencing on HBMEC

HBMEC were transfected with control *vs* TRIM47 siRNA for 72h. Total RNAs were isolated as described above. RNA-seq analysis was conducted on three biological replicates. Library preparation and sequencing were performed at the transcriptomic platform of the Paris

Brain Institute (ICM). mRNA library preparation was performed based on the manufacturer's recommendations (KAPA mRNA HyperPrep Kit [ROCHE]). Pooled library preparations were sequenced on NextSeq® 500 whole genome sequencing (Illumina®), corresponding to 2x30 million reads per sample after demultiplexing. The quality of raw data was evaluated with FastQC. Poor-quality sequences were trimmed or removed with Trimmomatic software to retain only good quality paired reads. Star v2.5.3a 53 was used to align reads on the human GRCh37/hg19 reference genome using standard options. Quantification of gene was done using RNA-Seq by Expectation-Maximization (RSEM) v 1.2.28, prior to normalization using the edgeR Bioconductor software package. Data analysis and visualization of the results was done with the Data Analysis Core facility from ICM and their user interface (QuBy). Differential analysis was conducted with the generalized linear model framework likelihood ratio test from edgeR using a log2FC threshold set up to 1. Multiple hypothesis adjusted p-values were calculated using the Benjamini-Hochberg procedure to control for the false discovery rate (FDR). Genes with low counts (CPM<5) were excluded from the analysis. Genes with an adjusted p-value (FDR) <0.05 were considered as differentially expressed genes (DEG). Gene set enrichment analysis was performed using the Molecular Signatures Database (MSigDB) for the Wikipathways. Volcano plots and dotplots with enrichment scores (NES) were directly downloaded from QuBy.

Oxidative stress measurement

HBMEC seeded on 24 well plates were transfected with control or TRIM47 siRNA and then replated on Nunc™ Lab-Tek™ II Chamber Slide™ (Thermo fisher Scientific) after 24h. Cells were then pre-treated with DMSO, tBHQ (10 µM) or DMF (10 µM) for 24h and then oxidative stress was induced by TBHP (Luperox, 150 µM, 1h) or with angiotensin II (750nM, 2h).

CellROX® Green reagent (C10444, Invitrogen) for oxidative stress detection and Hoechst for identifying nuclei (Thermo Fisher Scientific) were added directly to the live cells in complete EGM2 medium. After 30 minutes incubation, cells were washed 3 times in PBS and imaged with a fluorescent microscope (Observer Z.1, Zeiss). Oxidative stress was measured by the quantification of green fluorescence in cytoplasm using the "Measure nuclear and cytoplasmic intensities tool" from ImageJ macro.

Mice and breeding

Animal experiments were performed in accordance with the guidelines from Directive 2010/63/EU of the European Parliament on the protection of animals used for testing and research and approved by the local Animal Care and Use Committee of the Bordeaux University CEEA50 (IACUC protocol #36971). General surgical procedures were conducted in mice according to ARRIVE guidelines (https://www.nc3rs.org.uk/arrive-guidelines). All animals used were retained on a C57BL/6 background and both male and female mice were used for experiments.

Trim47 full KO mice (C57BL/6N-Trim47em1(IMPC)Bay/Mmmh) were purchased from the national public repository system for mutant mice Mutant Mouse Resource and Research Center. The inducible endothelial-specific *Trim47* knockout mouse model (*Trim47*iEC-KO) was generated by breeding *Trim47* $^{fl/fl}$ mice (strain ID: T005408, GemPharmtech) with Cdh5(PAC)-CreERT2 mice[52]. Endothelial deletion of *Trim47* was induced in adult mice, by tamoxifen injection (5648, Sigma) (five injections of 0.5 mg daily). All experiments were conducted using littermate controls denoted in the text as *Trim47* +/+ for the full KO mice and as *Trim47*iEC-WT for the endothelial-specific deleted mice line.

For the activation of the Nrf2 pathway, adult *Trim47 -/-* and *Trim47 +/+* males were kept on a diet containing tBHQ (Sigma, 1% w/w). The food pellets were supplemented with tBHQ by the manufacturer Safe. Body weight and food intake were monitored regularly. Behavioral tests were performed after a 1 month-diet and tissues were collected for histology, BBB permeability assessment or transcriptomic analysis after 2 months.

In vivo model of oxidative stress - oxygen-induced retinopathy model

The oxygen-induced retinopathy model (OIR) was carried out as described previously[53,54]. Mice pups and their nursing mother were exposed to 75% oxygen for 5 days from age P8 to age P13 in a hyperoxic chamber (ProOx Model 110, BioSpherix). At P13, they were returned to normal room air. Pups were euthanized at P16 to assess the degree of vascular regression (avascular area) and to evaluate retinal revascularization with pathological vessels (NV tufts). Only the pups weighing >4.5 g at P16 were included in the study since low weight impact the outcome and severity of the OIR model[55]. Mouse eyes at P16 postnatal stage were dissected and stained with isolectin-B4 in PBS supplemented with 3% bovine serum albumin, 1% donkey serum and 0.5% Triton. Retinal vasculatures were imaged with a fluorescent microscope (Observer Z.1, Zeiss). Avascular retinal areas were quantified with Zen blue software (version 3.5, Zeiss), normalized to the total retinal area and expressed as a percentage compared to littermate controls.

Behavioral tests

Y-maze

The Y-maze apparatus consisted of three arms (8 × 30 × 15 cm) separated by an angle of 120°. The recognition memory procedure took advantage of the innate tendency of rodents

to explore novel environments. It consisted of four trials performed over 2 consecutive days. During the exploration (encoding) trial, mice were allowed to explore only two arms (starting arm and familiar arm) of the maze for 5 minutes, with the third arm being blocked (novel arm). The mice were given two exploration trials the first day with an inter-trial interval of 3 h and one additional trial the second day to reinforce encoding. Three hours after the last exploration trial, mice underwent a recognition trial. Each mouse was put back in the starting arm of the maze, with free access to all three arms for 5 min. Time spent by mice in each arm and their trajectories were recorded and analyzed using an automated videotracking system (EthoVisionXT 16, Noldus). Discriminating the novel arm from the two familiar arms was used as an index of spatial recognition memory. Memory performance was expressed as the percentage of time spent in novel arm calculated as follows: (time spent in novel arm/time spent in all three arms) × 100.

Morris water maze

This test used a circular pool (180 cm in diameter) located in a dimly lit room decorated with various distal cues. The pool contained opaque water (22 ± 1 °C) and was subdivided into four virtual quadrants (A, B, C and D). A circular escape platform (10 cm in diameter) was submerged 1 cm beneath the water surface. The hidden platform remained in the same fixed position (quadrant D) during training. For each training trial, mice could swim for a maximum of 60 seconds to locate the hidden platform and were allowed to stay on it for 15 seconds upon finding it. Mice unable to locate the platform were gently guided to it. Mice were trained over 4 consecutive days and given 2 daily trials separated by an inter-trial interval of 3 h. The performance of each mouse (latency to reach the platform, distance

swum, swim speed and trajectory) was acquired and analyzed using an automated videotracking system using (EthoVisionXT 16, Noldus).

Sucrose preference test

To assess anhedonia and anxiety, we performed a sucrose preference test on adult mice. The mice were previously accustomed to having 2 bottles in their cage, one week before the test. Mice were then offered two 50-ml bottles filled with water and 1% sucrose for 4 days. The 1% sucrose solution was then replaced by a 3% sucrose solution which was again replaced 4 days later by a 9% sucrose solution. Each day, water and sucrose amounts were monitored by weighing the water bottles, and the positions of the bottles in the cages were counterbalanced to avoid spatial preference. Sucrose preference was determined as the percentage of sucrose intake normalized to the total liquid intake.

Endothelial cells isolation

Brain endothelial cells

Mouse brains collected from 2 adult mice were pooled and transferred into a petri dish containing cold HBSS without calcium/magnesium. Cerebellum and brain stem were then removed, leaving cerebral hemispheres intact. Each cerebral hemisphere was then cut in 4 sagittal slides before being enzymatically and mechanically digested using a gentleMACS™ Dissociator (#130-093-235, Miltenyi Biotec, 37C_ABDK_01 program) and the Adult Brain Dissociation kit (#130-107-677, Miltenyi Biotec) following the manufacturer. Brain lysates were filtered through a 70 μm MACS Smart Strainer (#130-198-462, Miltenyi Biotec). Incubation of the filtered suspension with Myelin Removal Beads II for 15 minutes (#130-096-733, Miltenyi Biotec) allowed depletion of myelin. ECs enrichment was performed by

incubating the suspension with CD31 Microbeads for 30 minutes (#130-097-418, Miltenyi Biotec). The positive selection of ECs was done using LS column (#130-042-201, Miltenyi Biotec) placed on a magnetic separator (MACS® MultiStand and QuadroMACS™ Separator, Miltenyi Biotec). Endothelial cells were centrifuged (10 minutes, 500 g) and cell pellet was resuspended in 300 µl of TRI Reagent® (MRC gene) and then processed for RNA isolation.

Lung endothelial cells

Mouse lungs were collected from adult *Trim47* or *Trim47* -/- mice, minced with a scalpel and then digested using a gentleMACS™ (37C_m_LDK_1 program) and the Lung Dissociation kit (P) (Miltenyi Biotec, Cat#130-095-927). Lung lysates were filtered through a 70 µm MACS Smart Strainer. Endothelial cell enrichment was performed by incubating the suspension with CD31 Microbeads for 15 minutes (#130-097-418, Miltenyi Biotec). Endothelial cells were centrifuged (10 minutes, 500 g) and cell pellet was resuspended in TRI Reagent® and then processed for RNA isolation.

Blood brain barrier permeability assessment *in vivo*

To assess BBB permeability, fluorescent tracers were injected intravenously (retro-orbital) in adult (5-6 months) mice and left to circulate for 20 min as described previously[56]. Cadaverine (0.95 kDa) conjugated to Alexa Fluor-555 (A30677, Invitrogen) was injected at a concentration of 100 µg Cadaverine/20 g of mice and lysine-fixable 3 kDa dextran conjugated to tetramethylrhodamine (D3308, Invitrogen) at a concentration of 250 µg dextran/20 g. To assess tracers leak, a 300 µL blood sample was collected by cardiac puncture and centrifuged at 10 000 g, 10 min at 4 °C for serum preparation. Animals were then perfused in the left ventricle with warmed PBS. One hemi-brain (free of olfactory lobes and cerebellum) and kidney were then collected, and their weight measured before being

snap-frozen in liquid nitrogen. Tissues were homogenized in PBS using a Tissue Lyser (Qiagen) and centrifuged at 15,000 g, 20 min, and 4 °C. Fluorescence measurement (RFUs) of 50 μL of diluted serum, brain and kidney supernatants was done using a Spark® Multimode Microplate Reader (Tecan). A Permeability Index (PI) was calculated as follows: PI (mL/g) = (Tissue RFUs/g tissue weight)/(Serum RFUs/mL serum) for each animal. The PI of each animal was divided with the mean PI of the *Trim47 +/+* group and data were represented as a ratio of PI. For additional histological analysis, the contralateral hemi-brain was directly OCT-embedded and preserved for immunohistochemistry.

Immunofluorescence analysis of brain tissues

10 μm cryosections (sagittal) were prepared using a cryostat, then fixed with 4% PFA for 10 min at room temperature, blocked with blocking solution (PBS1X, 1 % BSA, 0.5 % Triton) for 1 h and incubated for 1.5 h at room temperature with primary antibodies diluted in buffer containing PBS1X, 0.5 % BSA and 0.25 % Triton X. For other histological analysis, brains were collected and placed in 4% PFA overnight at 4 °C. Brains were then washed 2 times 1h with PBS1X and embedded in 4% agarose. 50 μm coronal sections were prepared using a Leica VT 1000S vibratome and placed in storage solution (Glycerol 30%, Ethylene glycol 30%, PBS-1X 40%) -20 °C. Brain coronal sections were blocked with PBS 1X, 10% donkey serum, 0.05% Triton)) for 4h prior to incubation with primary antibodies for 48h and then with secondary antibodies (1/400 dilution). After 3x 30 min wash with PBS1X, sections were mounted using mounting medium (Fluoromount G with DAPI, #00-4958-02, Thermo Fisher Scientific). Sections were imaged with a fluorescent microscope (Observer Z.1, Zeiss) and with a confocal microscope (LSM 900, Zeiss). Images were analyzed with Zen blue (version 3.5, Zeiss) and ImageJ (version 2.3.0, NIH).

Transmission electron microscopy

Animals were perfused in the left ventricle with 0,9% NaCl and 20 IU/mL heparin, then 4% PFA with 2.5% glutaraldéhyde in 0.1M Phosphate Buffer pH7.4, and 4% PFA in 0.1M Phosphate Buffer pH7.4. Brains were collected and 150 µm coronal sections were prepared using a vibratome and placed in 4% PFA overnight at 4 °C. Samples were post-fixed with 1% osmium tetroxide and dehydrated with graded ethanol series. A final dehydration step is performed with acetone and embedded in Epon. Ultrathin sections (70 nm) were cut by using an ultramicrotome (Leica EM UC6, Leica Microsystems) and observed under a transmission electron microscope (MET 120kV HT7650, HITACHI, Japan).

Statistical analysis

All *in vitro* results presented in this study are representative of at least three independent experiments. '*n*' represents the number of biological replicates unless otherwise stated. All *in vivo* experiments were done on littermates with similar body weights per condition. Males and females were analyzed separately for behavioral tests. Data are shown as the mean ± standard error of the mean (s.e.m.) or standard deviation (s.d) as stated in the text. Statistical analysis was performed using GraphPad Prism 8 software. A two-sided unpaired *t-test* was performed for statistical analysis of two groups for *in vitro* data. A two-sided Mann–Whitney test was performed for the analysis of two groups for *in vivo* results. One-way ANOVA followed by Bonferroni's multiple comparisons test was performed for statistical analysis between 3 or more groups. Two-way ANOVA with multiple comparisons test was performed to evaluate mouse performances (2 or 3 groups) and progression over time during the Water maze. Differences were considered statistically significant with a P value < 0.05.

Data availability Statement

All other data supporting the findings of this work are available within the paper and its Supplementary Information file. Any other data are available from the corresponding author upon reasonable request.

Acknowledgment

We thank Professor Anupam Agarwal (The University of Alabama at Birmingham, School of Medicine, United States) for providing the HO1 promoter luciferase constructs. We thank Philippe Alzieu, Sylvain Grolleau and Maxime David for their technical support and Christelle Boulle for administrative assistance. This work benefited from equipment and services from the Bordeaux Imaging Center (BIC) platform and the iGenSeq (RNA sequencing) and iCONICS (RNAseq analysis) core facilities at the ICM (Institut du Cerveau et de la Moelle épinière, PARIS, France).

Funding

This project is supported by a grant overseen by the French National Research Agency (ANR) as part of the Investment for the Future Programme ANR-18-RHUS-0002 (SHIVA Project) and by the Precision and Global Vascular Brain Health Institute (VBHI) funded by the France 2030 IHU3 initiative.

Author contribution

C.D. contributed to study design, performed experiments, analyzed and conceptualized results, V.D., R.B., S.R, M.B., C.C. and C.Proust performed *in vivo* experiments, JV and BJV contributed to experimental design and performed *in vitro* experiments and histology, J.L.M and B.B provided equipment for behavioral tests, analyzed data for behavior assessment,

A.M. and S.D contributed to scientific discussion, T.C. designed the study, conceptualized results, and wrote the manuscript, C.Peghaire designed the study, performed experiments, analyzed and conceptualized results and wrote the manuscript

Corresponding author

Correspondence to Claire Peghaire: claire.peghaire@u-bordeaux.fr

Competing interests

The authors declare no competing interests.

References

(1) Nation, D. A.; Sweeney, M. D.; Montagne, A.; Sagare, A. P.; D'Orazio, L. M.; Pachicano, M.; Sepehrband, F.; Nelson, A. R.; Buennagel, D. P.; Harrington, M. G.; Benzinger, T. L. S.; Fagan, A. M.; Ringman, J. M.; Schneider, L. S.; Morris, J. C.; Chui, H. C.; Law, M.; Toga, A. W.; Zlokovic, B. V. Blood-Brain Barrier Breakdown Is an Early Biomarker of Human Cognitive Dysfunction. *Nat. Med.* **2019**, *25* (2), 270–276. https://doi.org/10.1038/s41591-018-0297-y.

(2) Barisano, G.; Montagne, A.; Kisler, K.; Schneider, J. A.; Wardlaw, J. M.; Zlokovic, B. V. Blood-Brain Barrier Link to Human Cognitive Impairment and Alzheimer's Disease. *Nat. Cardiovasc. Res.* **2022**, *1* (2), 108–115. https://doi.org/10.1038/s44161-021-00014-4.

(3) Wardlaw, J. M.; Smith, C.; Dichgans, M. Small Vessel Disease: Mechanisms and Clinical Implications. *Lancet Neurol.* **2019**, *18* (7), 684–696. https://doi.org/10.1016/S1474-4422(19)30079-1.

(4) Gao, Y.; Li, D.; Lin, J.; Thomas, A. M.; Miao, J.; Chen, D.; Li, S.; Chu, C. Cerebral Small Vessel Disease: Pathological Mechanisms and Potential Therapeutic Targets. *Front. Aging Neurosci.* **2022**, *14*.

(5) Markus, H. S.; Erik de Leeuw, F. Cerebral Small Vessel Disease: Recent Advances and Future Directions. *Int. J. Stroke* **2023**, *18* (1), 4–14. https://doi.org/10.1177/17474930221144911.

(6) Mishra, A.; Duplaà, C.; Vojinovic, D.; Suzuki, H.; Sargurupremraj, M.; Zilhão, N. R.; Li, S.; Bartz, T. M.; Jian, X.; Zhao, W.; Hofer, E.; Wittfeld, K.; Harris, S. E.; van der Auwera-Palitschka, S.; Luciano, M.; Bis, J. C.; Adams, H. H. H.; Satizabal, C. L.; Gottesman, R. F.; Gampawar, P. G.; Bülow, R.; Weiss, S.; Yu, M.; Bastin, M. E.; Lopez, O. L.; Vernooij, M. W.; Beiser, A. S.; Völker, U.; Kacprowski, T.; Soumare, A.; Smith, J. A.; Knopman, D. S.; Morris, Z.; Zhu, Y.; Rotter, J. I.; Dufouil, C.; Valdés Hernández, M.; Muñoz Maniega, S.; Lathrop, M.; Boerwinkle, E.; Schmidt, R.; Ihara, M.; Mazoyer, B.; Yang, Q.; Joutel, A.; Tournier-Lasserve, E.; Launer, L. J.; Deary, I. J.; Mosley, T. H.; Amouyel, P.; DeCarli, C. S.; Psaty, B. M.; Tzourio, C.; Kardia, S. L. R.; Grabe, H. J.; Teumer, A.; van Duijn, C. M.; Schmidt, H.; Wardlaw, J. M.; Ikram, M. A.; Fornage, M.; Gudnason, V.; Seshadri, S.; Matthews, P. M.; Longstreth, W. T.; Couffinhal, T.; Debette, S. Gene-Mapping Study of Extremes of Cerebral Small Vessel Disease Reveals TRIM47 as a Strong Candidate. *Brain* **2022**, *145* (6), 1992–2007. https://doi.org/10.1093/brain/awab432.

(7) Hatakeyama, S. TRIM Family Proteins: Roles in Autophagy, Immunity, and Carcinogenesis. *Trends Biochem. Sci.* **2017**, *42* (4), 297–311. https://doi.org/10.1016/j.tibs.2017.01.002.

(8)   Borlepawar, A.; Frey, N.; Rangrez, A. Y. A Systematic View on E3 Ligase Ring TRIMmers with a Focus on Cardiac Function and Disease. *Trends Cardiovasc. Med.* **2019**, *29* (1), 1–8. https://doi.org/10.1016/j.tcm.2018.05.007.

(9)   Vandeputte, D. A. A.; Meije, C. B.; van Dartel, M.; Leenstra, S.; IJlst-Keizers, H.; Das, P. K.; Troost, D.; Bosch, D. A.; Baas, F.; Hulsebos, T. J. M. GOA, a Novel Gene Encoding a Ring Finger B-Box Coiled-Coil Protein, Is Overexpressed in Astrocytoma. *Biochem. Biophys. Res. Commun.* **2001**, *286* (3), 574–579. https://doi.org/10.1006/bbrc.2001.5431.

(10) Azuma, K.; Ikeda, K.; Suzuki, T.; Aogi, K.; Horie-Inoue, K.; Inoue, S. TRIM47 Activates NF-κB Signaling via PKC-ε/PKD3 Stabilization and Contributes to Endocrine Therapy Resistance in Breast Cancer. *Proc. Natl. Acad. Sci. U. S. A.* **2021**, *118* (35), e2100784118. https://doi.org/10.1073/pnas.2100784118.

(11) Wang, X.; Fu, Y.; Xing, Y. TRIM47 Promotes Ovarian Cancer Cell Proliferation, Migration, and Invasion by Activating STAT3 Signaling. *Clinics* **2022**, *77*, 100122. https://doi.org/10.1016/j.clinsp.2022.100122.

(12) Liang, Q.; Tang, C.; Tang, M.; Zhang, Q.; Gao, Y.; Ge, Z. TRIM47 Is Up-Regulated in Colorectal Cancer, Promoting Ubiquitination and Degradation of SMAD4. *J. Exp. Clin. Cancer Res.* **2019**, *38* (1), 159. https://doi.org/10.1186/s13046-019-1143-x.

(13) Wang, Z.; Li, Z.; Han, C.; Cheng, Y.; Wang, K. TRIM47 Promotes Glioma Angiogenesis by Suppressing Smad4. *Vitro Cell. Dev. Biol. - Anim.* **2022**, *58* (9), 771–779. https://doi.org/10.1007/s11626-022-00722-6.

(14) Chen, L.; Li, M.; Li, Q.; Xu, M.; Zhong, W. Knockdown of TRIM47 Inhibits Glioma Cell Proliferation, Migration and Invasion through the Inactivation of Wnt/β-Catenin Pathway. *Mol. Cell. Probes* **2020**, *53*, 101623. https://doi.org/10.1016/j.mcp.2020.101623.

(15) Chen, J.; Xu, D.; Cao, J.; Zuo, L.; Han, Z.; Tian, Y.; Chu, C.; Zhou, W.; Pan, X.; Cui, X. TRIM47 Promotes Malignant Progression of Renal Cell Carcinoma by Degrading P53 through Ubiquitination. *Cancer Cell Int.* **2021**, *21* (1), 129. https://doi.org/10.1186/s12935-021-01831-0.

(16) Xia, Y.; Wei, Z.; Huang, W.; Wei, X.; He, Y. Trim47 Overexpression Correlates with Poor Prognosis in Gastric Cancer. *Neoplasma* **2021**, *68* (02), 307–316. https://doi.org/10.4149/neo_2020_200708N706.

(17) Han, Y.; Tian, H.; Chen, P.; Lin, Q. TRIM47 Overexpression Is a Poor Prognostic Factor and Contributes to Carcinogenesis in Non-Small Cell Lung Carcinoma. *Oncotarget* **2017**, *8* (14), 22730–22740. https://doi.org/10.18632/oncotarget.15188.

(18) Fujimura, T.; Inoue, S.; Urano, T.; Takayama, K.; Yamada, Y.; Ikeda, K.; Obinata, D.; Ashikari, D.; Takahashi, S.; Homma, Y. Increased Expression of Tripartite Motif (TRIM) 47 Is a Negative Prognostic Predictor in Human Prostate Cancer. *Clin. Genitourin. Cancer* **2016**, *14* (4), 298–303. https://doi.org/10.1016/j.clgc.2016.01.011.

(19) Dai, W.; Wang, J.; Wang, Z.; Xiao, Y.; Li, J.; Hong, L.; Pei, M.; Zhang, J.; Yang, P.; Wu, X.; Tang, W.; Jiang, X.; Jiang, P.; Xiang, L.; Li, A.; Lin, J.; Liu, S.; Wang, J. Comprehensive Analysis of the Prognostic Values of the TRIM Family in Hepatocellular Carcinoma. *Front. Oncol.* **2021**, *11*, 767644. https://doi.org/10.3389/fonc.2021.767644.

(20) Qian, Y.; Wang, Z.; Lin, H.; Lei, T.; Zhou, Z.; Huang, W.; Wu, X.; Zuo, L.; Wu, J.; Liu, Y.; Wang, L.-F.; Guan, X.-H.; Deng, K.-Y.; Fu, M.; Xin, H.-B. TRIM47 Is a Novel Endothelial Activation Factor That Aggravates Lipopolysaccharide-Induced Acute Lung Injury in Mice via K63-Linked Ubiquitination of TRAF2. *Signal Transduct. Target. Ther.* **2022**, *7* (1), 148. https://doi.org/10.1038/s41392-022-00953-9.

(21) Hao, M.-Q.; Xie, L.-J.; Leng, W.; Xue, R.-W. Trim47 Is a Critical Regulator of Cerebral Ischemia-Reperfusion Injury through Regulating Apoptosis and Inflammation. *Biochem. Biophys. Res. Commun.* **2019**, *515* (4), 651–657. https://doi.org/10.1016/j.bbrc.2019.05.065.

(22) Hayes, J. D.; Dinkova-Kostova, A. T. The Nrf2 Regulatory Network Provides an Interface between Redox and Intermediary Metabolism. *Trends Biochem. Sci.* **2014**, *39* (4), 199–218. https://doi.org/10.1016/j.tibs.2014.02.002.

(23) Liby, K.; Hock, T.; Yore, M. M.; Suh, N.; Place, A. E.; Risingsong, R.; Williams, C. R.; Royce, D. B.; Honda, T.; Honda, Y.; Gribble, G. W.; Hill-Kapturczak, N.; Agarwal, A.; Sporn, M. B. The Synthetic Triterpenoids, CDDO and CDDO-Imidazolide, Are Potent Inducers of Heme Oxygenase-1 and Nrf2/ARE Signaling. *Cancer Res.* **2005**, *65* (11), 4789–4798. https://doi.org/10.1158/0008-5472.CAN-04-4539.

(24) Deshane, J.; Kim, J.; Bolisetty, S.; Hock, T. D.; Hill-Kapturczak, N.; Agarwal, A. Sp1 Regulates Chromatin Looping between an Intronic Enhancer and Distal Promoter of the Human Heme Oxygenase-1 Gene in Renal Cells. *J. Biol. Chem.* **2010**, *285* (22), 16476–16486. https://doi.org/10.1074/jbc.M109.058586.

(25) Li, S.-Y.; Fu, Z. J.; Lo, A. C. Y. Hypoxia-Induced Oxidative Stress in Ischemic Retinopathy. *Oxid. Med. Cell. Longev.* **2012**, *2012*, 426769. https://doi.org/10.1155/2012/426769.

(26) Deliyanti, D.; Lee, J. Y.; Petratos, S.; Meyer, C. J.; Ward, K. W.; Wilkinson-Berka, J. L.; de Haan, J. B. A Potent Nrf2 Activator, Dh404, Bolsters Antioxidant Capacity in Glial Cells and Attenuates Ischaemic Retinopathy. *Clin. Sci.* **2016**, *130* (15), 1375–1387. https://doi.org/10.1042/CS20160068.

(27) Mao, L.; Yang, T.; Li, X.; Lei, X.; Sun, Y.; Zhao, Y.; Zhang, W.; Gao, Y.; Sun, B.; Zhang, F. Protective Effects of Sulforaphane in Experimental Vascular Cognitive Impairment: Contribution of the Nrf2 Pathway. *J. Cereb. Blood Flow Metab. Off. J. Int. Soc. Cereb. Blood Flow Metab.* **2019**, *39* (2), 352–366. https://doi.org/10.1177/0271678X18764083.

(28) Giovannoni, F.; Quintana, F. J. The Role of Astrocytes in CNS Inflammation. *Trends Immunol.* **2020**, *41* (9), 805–819. https://doi.org/10.1016/j.it.2020.07.007.

(29) Patani, R.; Hardingham, G. E.; Liddelow, S. A. Functional Roles of Reactive Astrocytes in Neuroinflammation and Neurodegeneration. *Nat. Rev. Neurol.* **2023**, *19* (7), 395–409. https://doi.org/10.1038/s41582-023-00822-1.

(30) Yeung, S. H.-S.; Lee, R. H.-S.; Cheng, G. W.-Y.; Ma, I. W.-T.; Kofler, J.; Kent, C.; Ma, F.; Herrup, K.; Fornage, M.; Arai, K.; Tse, K.-H. White Matter Hyperintensity Genetic Risk Factor TRIM47 Regulates Autophagy in Brain Endothelial Cells. bioRxiv December 19, 2023, p 2023.12.18.566359. https://doi.org/10.1101/2023.12.18.566359.

(31) Wang, Y.; Sun, B.; Shibata, B.; Guo, F. Transmission Electron Microscopic Analysis of Myelination in the Murine Central Nervous System. *STAR Protoc.* **2022**, *3* (2), 101304. https://doi.org/10.1016/j.xpro.2022.101304.

(32) Akhter, H.; Katre, A.; Li, L.; Liu, X.; Liu, R.-M. Therapeutic Potential and Anti-Amyloidosis Mechanisms of Tert-Butylhydroquinone for Alzheimer's Disease. *J. Alzheimers Dis. JAD* **2011**, *26* (4), 767–778. https://doi.org/10.3233/JAD-2011-110512.

(33) Guan, Y.; Tan, Y.; Liu, W.; Yang, J.; Wang, D.; Pan, D.; Sun, Y.; Zheng, C. NF-E2-Related Factor 2 Suppresses Intestinal Fibrosis by Inhibiting Reactive Oxygen Species-Dependent TGF-B1/SMADs Pathway. *Dig. Dis. Sci.* **2018**, *63* (2), 366–380. https://doi.org/10.1007/s10620-017-4710-z.

(34) Bian, H.; Yan, F.; Li, W.; Tu, W.; Ji, X. Tert-Butylhydroquinone Prevents Neuroinflammation and Relieves Depression via Regulation of NLRP3 Signaling in Mice. *Int. Immunopharmacol.* **2022**, *107*, 108723. https://doi.org/10.1016/j.intimp.2022.108723.

(35) Zhang, Z.-W.; Liang, J.; Yan, J.-X.; Ye, Y.-C.; Wang, J.-J.; Chen, C.; Sun, H.-T.; Chen, F.; Tu, Y.; Li, X.-H. TBHQ Improved Neurological Recovery after Traumatic Brain Injury by Inhibiting the Overactivation of Astrocytes. *Brain Res.* **2020**, *1739*, 146818. https://doi.org/10.1016/j.brainres.2020.146818.

(36) Tan, X.-L.; Xue, Y.-Q.; Ma, T.; Wang, X.; Li, J. J.; Lan, L.; Malik, K. U.; McDonald, M. P.; Dopico, A. M.; Liao, F.-F. Partial eNOS Deficiency Causes Spontaneous Thrombotic Cerebral Infarction, Amyloid Angiopathy and Cognitive Impairment. *Mol. Neurodegener.* **2015**, *10*, 24. https://doi.org/10.1186/s13024-015-0020-0.

(37) Chen, X.; Chen, L.; Lin, G.; Wang, Z.; Kodali, M. C.; Li, M.; Chen, H.; Lebovitz, S. G.; Ortyl, T. C.; Li, L.; Ismael, S.; Singh, P.; Malik, K. U.; Ishrat, T.; Zhou, F.-M.; Zheng, W.; Liao, F.-F. White Matter Damage as a Consequence of Vascular Dysfunction in a Spontaneous Mouse Model of Chronic

Mild Chronic Hypoperfusion with eNOS Deficiency. *Mol. Psychiatry* **2022**, *27* (11), 4754–4769. https://doi.org/10.1038/s41380-022-01701-9.

(38) Ryu, J. K.; Rafalski, V. A.; Meyer-Franke, A.; Adams, R. A.; Poda, S. B.; Rios Coronado, P. E.; Pedersen, L. Ø.; Menon, V.; Baeten, K. M.; Sikorski, S. L.; Bedard, C.; Hanspers, K.; Bardehle, S.; Mendiola, A. S.; Davalos, D.; Machado, M. R.; Chan, J. P.; Plastira, I.; Petersen, M. A.; Pfaff, S. J.; Ang, K. K.; Hallenbeck, K. K.; Syme, C.; Hakozaki, H.; Ellisman, M. H.; Swanson, R. A.; Zamvil, S. S.; Arkin, M. R.; Zorn, S. H.; Pico, A. R.; Mucke, L.; Freedman, S. B.; Stavenhagen, J. B.; Nelson, R. B.; Akassoglou, K. Fibrin-Targeting Immunotherapy Protects against Neuroinflammation and Neurodegeneration. *Nat. Immunol.* **2018**, *19* (11), 1212–1223. https://doi.org/10.1038/s41590-018-0232-x.

(39) Petersen, M. A.; Ryu, J. K.; Akassoglou, K. Fibrinogen in Neurological Diseases: Mechanisms, Imaging and Therapeutics. *Nat. Rev. Neurosci.* **2018**, *19* (5), 283–301. https://doi.org/10.1038/nrn.2018.13.

(40) Dean, T.; Mendiola, A. S.; Yan, Z.; Meza-Acevedo, R.; Cabriga, B.; Akassoglou, K.; Ryu, J. K. Fibrin Promotes Oxidative Stress and Neuronal Loss in Traumatic Brain Injury via Innate Immune Activation. *J. Neuroinflammation* **2024**, *21* (1), 94. https://doi.org/10.1186/s12974-024-03092-w.

(41) De Silva, T. M.; Miller, A. A. Cerebral Small Vessel Disease: Targeting Oxidative Stress as a Novel Therapeutic Strategy? *Front. Pharmacol.* **2016**, *7*, 61. https://doi.org/10.3389/fphar.2016.00061.

(42) Song, K.; Li, Y.; Zhang, H.; An, N.; Wei, Y.; Wang, L.; Tian, C.; Yuan, M.; Sun, Y.; Xing, Y.; Gao, Y. Oxidative Stress-Mediated Blood-Brain Barrier (BBB) Disruption in Neurological Diseases. *Oxid. Med. Cell. Longev.* **2020**, *2020*, 1–27. https://doi.org/10.1155/2020/4356386.

(43) Gureev, A. P.; Khorolskaya, V. G.; Sadovnikova, I. S.; Shaforostova, E. A.; Cherednichenko, V. R.; Burakova, I. Y.; Plotnikov, E. Y.; Popov, V. N. Age-Related Decline in Nrf2/ARE Signaling Is Associated with the Mitochondrial DNA Damage and Cognitive Impairments. *Int. J. Mol. Sci.* **2022**, *23* (23), 15197. https://doi.org/10.3390/ijms232315197.

(44) Montagne, A.; Barnes, S. R.; Sweeney, M. D.; Halliday, M. R.; Sagare, A. P.; Zhao, Z.; Toga, A. W.; Jacobs, R. E.; Liu, C. Y.; Amezcua, L.; Harrington, M. G.; Chui, H. C.; Law, M.; Zlokovic, B. V. Blood-Brain Barrier Breakdown in the Aging Human Hippocampus. *Neuron* **2015**, *85* (2), 296–302. https://doi.org/10.1016/j.neuron.2014.12.032.

(45) Bennett, H. C.; Zhang, Q.; Wu, Y.; Manjila, S. B.; Chon, U.; Shin, D.; Vanselow, D. J.; Pi, H.-J.; Drew, P. J.; Kim, Y. Aging Drives Cerebrovascular Network Remodeling and Functional Changes in the Mouse Brain. *Nat. Commun.* **2024**, *15* (1), 6398. https://doi.org/10.1038/s41467-024-50559-8.

(46) George, M.; Tharakan, M.; Culberson, J.; Reddy, A. P.; Reddy, P. H. Role of Nrf2 in Aging, Alzheimer's and Other Neurodegenerative Diseases. *Ageing Res. Rev.* **2022**, *82*, 101756. https://doi.org/10.1016/j.arr.2022.101756.

(47) Yang, T.; Zhang, F. Targeting Transcription Factor Nrf2 (Nuclear Factor Erythroid 2-Related Factor 2) for the Intervention of Vascular Cognitive Impairment and Dementia. *Arterioscler. Thromb. Vasc. Biol.* **2020**. https://doi.org/10.1161/ATVBAHA.120.314804.

(48) Zhang, H.; Davies, K. J. A.; Forman, H. J. Oxidative Stress Response and Nrf2 Signaling in Aging. *Free Radic. Biol. Med.* **2015**, *88*, 314–336. https://doi.org/10.1016/j.freeradbiomed.2015.05.036.

(49) Tarantini, S.; Valcarcel-Ares, M. N.; Yabluchanskiy, A.; Tucsek, Z.; Hertelendy, P.; Kiss, T.; Gautam, T.; Zhang, X. A.; Sonntag, W. E.; de Cabo, R.; Farkas, E.; Elliott, M. H.; Kinter, M. T.; Deak, F.; Ungvari, Z.; Csiszar, A. Nrf2 Deficiency Exacerbates Obesity-Induced Oxidative Stress, Neurovascular Dysfunction, Blood–Brain Barrier Disruption, Neuroinflammation, Amyloidogenic Gene Expression, and Cognitive Decline in Mice, Mimicking the Aging Phenotype. *J. Gerontol. Ser. A* **2018**, *73* (7), 853–863. https://doi.org/10.1093/gerona/glx177.

(50) Sigfridsson, E.; Marangoni, M.; Hardingham, G. E.; Horsburgh, K.; Fowler, J. H. Deficiency of Nrf2 Exacerbates White Matter Damage and Microglia/Macrophage Levels in a Mouse Model of Vascular Cognitive Impairment. *J. Neuroinflammation* **2020**, *17* (1), 367. https://doi.org/10.1186/s12974-020-02038-2.

(51) Fan, W.; Chen, H.; Li, M.; Fan, X.; Jiang, F.; Xu, C.; Wang, Y.; Wei, W.; Song, J.; Zhong, D.; Li, G. NRF2 Activation Ameliorates Blood–Brain Barrier Injury after Cerebral Ischemic Stroke by

Regulating Ferroptosis and Inflammation. *Sci. Rep.* **2024**, *14* (1), 5300. https://doi.org/10.1038/s41598-024-53836-0.

(52) Sörensen, I.; Adams, R. H.; Gossler, A. DLL1-Mediated Notch Activation Regulates Endothelial Identity in Mouse Fetal Arteries. *Blood* **2009**, *113* (22), 5680–5688. https://doi.org/10.1182/blood-2008-08-174508.

(53) Smith, L. E.; Wesolowski, E.; McLellan, A.; Kostyk, S. K.; D'Amato, R.; Sullivan, R.; D'Amore, P. A. Oxygen-Induced Retinopathy in the Mouse. *Invest. Ophthalmol. Vis. Sci.* **1994**, *35* (1), 101–111.

(54) Bats, M.-L.; Bougaran, P.; Peghaire, C.; Gueniot, F.; Abelanet, A.; Chan, H.; Séguy, C.; Jeanningros, S.; Jaspard-Vinassa, B.; Couffinhal, T.; Duplàa, C.; Dufourcq, P. Therapies Targeting Frizzled-7/β-Catenin Pathway Prevent the Development of Pathological Angiogenesis in an Ischemic Retinopathy Model. *FASEB J.* **2020**, *34* (1), 1288–1303. https://doi.org/10.1096/fj.201901886R.

(55) Stahl, A.; Chen, J.; Sapieha, P.; Seaward, M. R.; Krah, N. M.; Dennison, R. J.; Favazza, T.; Bucher, F.; Löfqvist, C.; Ong, H.; Hellström, A.; Chemtob, S.; Akula, J. D.; Smith, L. E. H. Postnatal Weight Gain Modifies Severity and Functional Outcome of Oxygen-Induced Proliferative Retinopathy. *Am. J. Pathol.* **2010**, *177* (6), 2715–2723. https://doi.org/10.2353/ajpath.2010.100526.

(56) Devraj, K.; Guérit, S.; Macas, J.; Reiss, Y. An In Vivo Blood-Brain Barrier Permeability Assay in Mice Using Fluorescently Labeled Tracers. *J. Vis. Exp. JoVE* **2018**, No. 132, 57038. https://doi.org/10.3791/57038.

Figure legends

Figure 1 TRIM47 is an activator of the antioxidant NRF2 pathway in human brain endothelial cells

a. RNA-sequencing was performed on HBMEC treated with control or TRIM47 siRNA for 72h. Volcano plot showing log2 fold change (FC) *vs* -log10 FDR of differentially expressed genes (DEG) in response to TRIM47 knockdown. Significant downregulated (log2FC<-1, in blue) and upregulated (log2FC>1, in red) genes were identified by edgeR analysis (FDR<0.05 and CPM >=5). **b.** Pathway enrichment analysis of DEG. Gene set enrichment analysis (Wikipathway) revealed a significant (p-adj=0.025) downregulation of genes associated with the NRF2 signalling pathway. Dotplot depicted the normalized enrichment scores (NES) for the top 10 pathways. The dots are colored by the adjusted p-value and their size is proportional with the size of the gene-set. **c.** Heatmap showing the expression of selected genes related to the NRF2 pathway in HBMEC treated with TRIM47 siRNA *vs* control. Data are expressed as log2FC. *FDR<0.05. **d.** Immunoblot and quantification of WB for TRIM47 and HO1 expression in control and TRIM47-deficient HBMEC treated with DMSO or MG132 for 5h. Data were normalized to α-tubulin (n=3 or 4 experiments). **** P<0.0001, Student's t-test. **e-f.** qPCR analysis of (**e**) *HMOX1* and (**f**) *NQO1* gene expression in control, TRIM47 or NRF2 siRNA-treated HBMEC for 72h and treated with either PBS or H2O2 (200 uM for 3h). Data were normalized to *cyclophilin* (n=3 experiments). * P<0.05; ** P<0.01; *** P<0.001, One-way ANOVA, # P<0.05; ## P<0.01, Student's t-test. **G.** HO1 promoter luciferase reporter assay. HeLa were co-transfected with HO1 promoter-luciferase construct (HO1-4.5bp) or a pGL3 empty vector with control or siRNA targeting TRIM47. Cells were treated with either PBS or H2O2 (200 uM for 1h). Luciferase activity was measured in cell lysates. Values represent the fold change in relative luciferase activity over the HO1-4.5kb + siControl condition and normalized to Renilla (n=4 experiments). **** P<0.0001 One-way ANOVA; #### P<0.0001 compared to pGL3 + siControl condition, Student's t-test. All graphical data are mean ± s.d.

Figure 2 TRIM47 cooperates with NRF2 in driving antioxidant target genes

a-b. qPCR analysis of (**a**) *NQO1* and (**b**) *HMOX1* gene expression in siCont or siNRF2 treated HBMEC and co-transduced with a control or TRIM47 lentivirus for 24h. Data were normalized to *cyclophilin* (n=3 experiments). * P<0.05; ** P<0.01; *** P<0.001; **** P<0.0001, One-way ANOVA. **c.** HO1 promoter luciferase reporter assay. TRIM47 cDNA expression plasmid (TRIM47) or empty expression plasmid (pcDNA) were co-transfected with HO1 promoter-luciferase constructs (HO1-4.5bp, HO1-4.5bp with 2 or 3 mutations) or a pGL3 empty vector in HeLa, and luciferase activity was measured. Values represent the fold change in relative luciferase activity over pGL3 vector (n=4 experiments). * P<0.05; ** P<0.01; **** P<0.0001, One-way ANOVA; ### P<0.001: compared to pGL3 + Lenti control condition, Student's t-test. **d.** TRIM47 and KEAP1 interaction was assessed by Co-IP assay in whole cell lysates from HEK293. Cells were transfected with TRIM47-myc pcDNA, KEAP1-Flag or both expression plasmids. Lysates were immunoprecipitated with myc antibody and immuno-blotted for myc (TRIM47) and KEAP1. **e.** Immunoblot and quantification of WB for NRF2 and KEAP1 in control and TRIM47-deficient HBMEC treated with DMSO or MG132 for 5h (n=4 experiments). Data were normalized to α-tubulin.* P<0.05; ** P<0.01, Student's t-test. **f-g.** qPCR analysis of (**f**) *NQO1* and (**g**) *HMOX1* gene expression in HBMEC treated with control, TRIM47 or NRF2 siRNA for 72h and with tBHQ (10μM) for 24h (n=3 experiments). ** P<0.01; *** P<0.001; **** P<0.0001, One-way ANOVA; ##, P<0.01 Student's t-test. All graphical data are mean ± s.e.m. **h.** Model of TRIM47 activation of the antioxidant NRF2 pathway in endothelial cells. 1. In basal conditions, TRIM47 binds to KEAP1 in the cytoplasm and thus alleviates KEAP1-dependent NRF2 degradation and increases NRF2 protein stabilization that can translocate to the nucleus to drive the expression of NRF2-target genes (*HMOX1*, *NQO1*) promoting cellular antioxidant protection. 2. In the absence of TRIM47 (TRIM47 siRNA condition), KEAP1 strongly binds to NRF2 leading to its ubiquitination and degradation by proteosomal complex, thus leading to the decreased of the protective NRF2 pathway.

Figure 3 TRIM47 protects from oxidative stress *in vitro* and *in vivo*

a-b. Representative immunofluorescence image (**a**) and quantification (**b**) of CellRox dye (green, marker of oxidative stress) in HBMEC transfected with siCont or si*TRIM47* for 24 h, then pre-treated with DMSO or tBHQ (10 μM for 24 h) and treated with PBS or tert-butyl hydroperoxide (TBHP, 200 μM for 1 h); nuclei are identified by Hoechst (blue). Scale bar 100 μm. Quantification represents the mean percentage of cytoplasmic area of Cellrox dye per field (n=4 wells from 3 independent experiments). **** $P<0.0001$, One-way ANOVA; #### $P<0.001$: comparison with siCont + THBP without drug treatment. **c.** Heatmaps of selected genes link to the Nrf2 pathway. qPCR screening performed on whole brain lysates isolated from P16 *Trim47+/+* and *Trim47-/-* mice exposed to normoxia condition or oxygen-induced retinopathy (OIR) model and. Data were normalized to *18S* (n=5-7 replicates per genotype and condition). # $P<0.01$, +/+ OIR *vs* +/+ 21%O_2, * $P<0.0001$, -/- OIR *vs* +/+ OIR. One-way ANOVA. **d**. qPCR in brain ECs isolated from P16 *Trim47+/+* and *Trim47-/-* mice exposed to OIR model. Data were normalized to *Pecam1* and expressed as fold change (n=3 replicates/ genotype; 2 mice/replicate).**e.** Representative images of isolectin B4 (IB4, green or black) staining of postnatal day 16 retina from *Trim47+/+* and *Trim47-/-* mice following OIR model characterized by oxidative stress. Right panel highlights, in blue, the vascular area of the retina after OIR. Scale bar 500 μm. **f.** Quantification represents the percentage of avascular area (μm^2) in *Trim47+/+* and *Trim47-/-* mice at P16 following OIR (n=9 mice per genotype). * $P<0.05$; Mann-Whitney test. All graphical data are mean ± s.d.

Figure 4 *Trim47* deletion leads to cognitive impairments

a-b. Y-maze test was performed on males (8-9 months; *Trim47+/+*: n=9; *Trim47-/-:* n=10). Data revealed recognition memory in *Trim47-/-* mice compared to *Trim47+/+* mice. (**a**) Y-maze data are presented as a ratio between the time spent in novel arm and the cumulative time spent in the novel and the familiar arms. **$p<0.01$, -/- *vs* +/+ mice, Mann-Whitney test. (**b**) Representative heatmap images of the pathes adopted in the Y-maze test. Mice started in position a; novel arm is arm c. **c-f.**

Water maze test was performed in adults male (7-8 months) *Trim47+/+* (n=11) *and Trim47-/-* (n=11)

mice and revealed lower performances. *Trim47 -/-* mice failed to learn the spatial position of the

hidden platform (**c**) Graph shows the latency to reach the hidden platform (expressed in seconds)

over 4 training days. **p<0.01 *-/- vs +/+* mice, Two-way ANOVA with repeated measures. (**d**)

Representative heatmap images of the pathes adopted by the mice during the first and last training

days. The orange circle shows the position where the platform was located; the white cross indicates

where the mice started the test. (**e**) Graph shows the distance to reach platform (expressed in cm) on

day 4. *p<0.05, *-/- vs +/+* mice, Mann-Whitney test. (**f**) Data represent the velocity expressed as cm

per second of each mouse on day 4 of water maze training. **g-h.** Y-maze was also performed in

females (8-10 months; *Trim47+/+*: n=14; *Trim47-/-*: n=13). . *p<0.05, *-/- vs +/+* mice. **i-l.** Water maze

test was repeated using females (8-9 months, *Trim47+/+:* n=10; *Trim47-/-*: n=11). Female *Trim47-/-*

exhibited cognitive impairments comparable to males. *p<0.05, *-/- vs +/+* mice. All graphical data are

mean ± s.d.

Figure 5 *Trim47* deletion leads to brain defects associated with Nrf2 pathway impairment

a-c. Blood brain barrier permeability assessment was done on *Trim47+/+* and *Trim47-/-* mice (5-7

months, males). Quantification of (**a**) cadaverine and (**b**) dextran contents in brain and kidney. Data

indicate the permeability index ratio (n=5 *Trim47+/+* and n=5 *Trim47-/-*). ** P<0.01, Mann-Whitney

test. (**c**) Representative confocal images and quantification of fibrinogen leakage (green) in brain

cryosections from *Trim47+/+* and *Trim47-/-* mice. Tissues are co-stained for podocalyxin (blood

vessels, pink). Scale bars 20 µm. White arrows highlight fibrinogen extravasation in *Trim47-/-* mice.

Top graph shows permeability in hippocampus (n=3 *Trim47+/+*, n=5 *Trim47-/-*) and bottom one in

cortex (n=5 *Trim47+/+*, n=5 *Trim47-/-*). * P<0.05; *** P<0.001, Mann-Whitney. **d.** qPCR analysis of

genes linked to junction and inflammation in brain ECs isolated from *Trim47+/+* and *Trim47-/-* mice

(7-10 months). Data normalized to *Pecam1* (n=6 replicates/genotype; 2 mice/replicate, except for

Il1b, *Il6* and *Plvap*: n=3/genotype). ND: not detectable; * P<0.05; ** P<0.01, Mann-Whitney. **e.**

Representative images and quantification of claudin5 (green) in brain cryosections from *Trim47+/+* and *Trim47-/-* mice (6-7 months); nuclei identified by DAPI (blue). Scale bars 20 µm. Top graph shows claudin5 expression in hippocampus (n=4 *Trim47+/+*, n=5 *Trim47-/-*) and bottom one in cortex (n=5 per genotype). * P<0.05; ** P<0.01, Mann-Whitney. **f.** qPCR for Nrf2 target genes in brain EC from *Trim47+/+* and *Trim47-/-* mice (7-10 months). Data normalized to *Pecam1* (n=6 replicates/genotype; * P<0.05; ** P<0.01, Mann-Whitney. **g**. Representative images and quantification of GFAP (green) in thick brain coronal sections (hippocampus) from *Trim47+/+* and *Trim47-/-* mice (10 months) (n=5 *Trim47+/+* and n=5 *Trim47-/-*). Scale bars 200 or 100 µm. * P<0.05, Mann-Whitney. **h.** Electron microscopy was performed on brain thin sections to analyze white matter lesions in corpus callosum. G-ratio (ratio of the inner axonal diameter to the total outer diameter, including the myelin sheath) was measured as a readout of axonal myelination in *Trim47+/+* and *Trim47-/-* mice (8 months) (n=3 per genotype). Scale bars 2 µm. i. Representative image of MBP (white) staining in thick brain coronal sections (hippocampus) from *Trim47+/+* and *Trim47-/-* mice (8 months). Scale bars 100 µm. All graphical data are mean ± s.d.

Figure 6 *Trim47* deletion in endothelial cells is sufficient to mimic the phenotype observed in *Trim47* full KO mice

a. qPCR analysis for *Trim47* expression was done on lung ECs isolated from *Trim47^iEC-WT^* and *Trim47^iEC-KO^* males (5 months). Data normalized to *Pecam1* (n=6 *Trim47 ^iEC-WT^* and n=9 *Trim47 ^iEC-KO^*); *** P<0.001, Mann-Whitney. **b-c.** Y-maze test was performed on males (3-5 months; *Trim47^iEC-WT^:* n=14; *Trim47 ^iEC-KO^:* n=13). (**b**) Y-maze data are presented as a ratio between the time spent in novel arm and the cumulative time spent in the novel and the familiar arms. *p<0.05, Mann-Whitney test. (**c**) Representative heatmap images of the path for the Y-maze test. Mice started in position a; novel arm is arm c. **d-g.** Water maze test was performed on males (3-5 months; *Trim47 ^iEC-WT^:* n=12; *Trim47^iEC-KO^:* n=13). (**d**) Graph shows the latency to reach the hidden platform (expressed in seconds) for the 4 training days. **p<0.01, *iEC-WT vs iEC-KO* mice, Two-way ANOVA with repeated measures.

(**e**) Representative heatmap images of the pathes for the first and last training days. (**f**) Graph shows the distance to reach platform (expressed in cm) on day 4. *p<0.05, *iEC-WT vs iEC-KO* mice, Mann-Whitney test. (**g**) Data represent the velocity expressed as cm per second (swimming speed) of each mouse on day 4 during water maze training. **h.** Assessment of BBB permeability was done on *Trim47^{iEC-WT}* and *Trim47^{iEC-KO}* mice (4-6 months, males). Quantification of dextran contents in brain cortex and kidney, 20 minutes after I.V injection of fluorescent tracer. Data indicate the permeability index ratio (normalization to fluorescence detected in serum) (n=5 *Trim47^{iEC-WT}* and n=5 *Trim47^{iEC-KO}*). * P<0.01, Mann-Whitney test. **i-j.** qPCR for (**i**) genes linked to junctions and inflammation and (**j**) Nrf2 targets in brain EC isolated from *Trim47^{iEC-WT}* and *Trim47^{iEC-KO}* mice (4-6 months). Data normalized to *Pecam1* (n=4-6 replicates/genotype with 2 mice/replicate; * P<0.05; ** P<0.01, ND: not detectable, Mann-Whitney. All graphical data are mean ± s.d.

Figure 7 Activation of NRF2 pathway prevents cognitive impairment and brain defects observed in *Trim47 -/- mice*

a. Cartoon depicting the experimental therapeutical approach chosen in mouse**.** 6-8 months *Trim47+/+ and Trim47-/-* males were placed on a pellet diet containing the NRF2 pathway activator tert-butylhydroquinone (tBHQ, 1%W/W) for 1 month. Behavioral tests were then performed on these mice and on untreated *Trim47 -/-* males before brain harvesting for qPCR analysis, BBB function and histology. **b**. qPCR for Nrf2 pathway performed on brain EC from *Trim47+/+ and Trim47-/-* males (8-10 months) with tBHQ diet (n=5 replicates/genotype; 2 mice/replicate). Mann-Whitney. ND: not detectable. **c**. BBB permeability assessment done on *Trim47+/+* and *Trim47-/-* males with tBHQ (8-9 months). Quantification of dextran in brain cortex and kidney (n=5 *Trim47+/+* + tBHQ and n=5 *Trim47-/-* +tBHQ). Mann-Whitney. **d.** Representative image and quantification of GFAP expression (green) in brain sections (coronal) from *Trim47+/+* and *Trim47-/-* adult mice (8-10 months) after tBHQ diet (n=3 per genotype). Scale bars 100 μm. Mann-Whitney. **e-f.** Y-maze test was performed on adult mice after a 1 month diet with tBHQ. (**e**) Data are presented as a ratio between

the time spent in novel arm and the cumulative time spent in the novel and familiar arm (n=12 *Trim47+/+* + tBHQ, n=13 *Trim47-/-* + tBHQ and n=5 *Trim47-/-* without diet). *p<0.05; **p<0.01, One-way ANOVA. (**f**) Representative heatmap images of the pathes for the Y-maze test. **g-h.** Water maze test performed on adult mice after tBHQ diet. (**g**) Latency to reach platform (expressed in seconds) is shown for the 4 training days (n=14 *Trim47 +/+* + tBHQ, n=13 *Trim47 -/-* + tBHQ, n= *Trim47 -/-*). *p<0.05, *Trim47 -/- vs Trim47 +/+* + tBHQ. Two-way ANOVA with repeated measures. (**h**) Representative heatmap of the pathes of *Trim47 +/+ and Trim47 +/+* treated with tBHQ. All graphical data are mean ± s.d.

Figures

TRIM47 , NRF2 pathway and SVD

Title:

Cerebral Small Vessel Disease genetic determinant TRIM47 controls brain homeostasis via the NRF2 antioxidant system

Figure 1

Figure 2

Figure 3

a

siCont | siTRIM47

PBS
- DMSO: CellRox / CellRox/Hoechst
- tBHQ
100 µm

TBHP
- DMSO
- tBHQ

b

Oxydative stress

- siCont
- siTRIM47

PBS+DMSO

PBS+tBHQ

TBHP+DMSO

TBHP+tBHQ ####

****

****

Cytoplasmic area of CellRox-%

c

Expression profile on P16 whole brain

Relative gene expression
fold change (normallized to 18S)

0.5 1.0 1.5 2.0 2.5

Genes: Tubb3, S100b, Pdgfrb, Trim47, Pecam1, Cdh5, Nfe2l2, Keap1, Hmox1, Nqo1, Gclm, Me1

21%O2: +/+, -/- (ND for Trim47)

OIR: +/+ (# for Keap1, Hmox1, Nqo1, Gclm, Me1), -/- (ND for Trim47; * for Hmox1, Nqo1, Gclm, Me1)

d

Expression profile on P16 brain ECs

Relative gene expression
fold change (normalized to Pecam1)

0.5 1.5 2.5

Genes: Trim47, Cdh5, Nfe2l2, Keap1, Hmox1, Nqo1, Gclm, Me1

OIR: +/+, -/- (ND for Trim47)

e

Trim47 +/+
IB4
Avascular area IB4
500 µm

Trim47 -/-
IB4
Avascular area IB4
500 µm

f

- Trim47 +/+ n=9
- Trim47 -/- n=9

*

% of avascular area

Avascular area - P16

Figure 4

Figure 5

Figure 6

Figure 7

a

Trim47 +/+ vs Trim47 -/-
(6-8 months)

Weeks: 0 — 4 — 5 — 6 — 8

Diet with tBHQ (1%w/w)

Water maze | Y maze | Brain tissue analysis

b

Expression profile on Brain EC
+ tBHQ diet

Trim47 — ND
Slc2a1
Mfsd2a
Abcb1a

Nrf2 pathway:
Nfe2l2
Keap1
Hmox1
Nqo1
Gclm
Me1
Slc7a11

Junctions:
Cdh5
Cldn5
Cldn11
Ocln
Tjp1

+/+   -/-

Relative gene expression fold change (normalized to Pecam1)

c

Dextran 3kDa

- Trim47 WT + tBHQ n=5
- Trim47 KO + tBHQ n=5

Dextran 3kDa-Permeability index ratio

Brain | Kidney

d

tBHQ diet

Hippocampus | Dentate gyrus

Trim47 +/+: GFAP/DAPI, CA1, DG, GFAP, 200 μm, 100 μm

Trim47 -/-: CA1, DG

- Trim47 +/+ + tBHQ
- Trim47 -/- + tBHQ

Area of GFAP staining (μm2) per field

NS

Hippocampus

e

- Trim47 +/+ + tBHQ n=12
- Trim47 -/- + tBHQ n=13
- Trim47 -/- n=5

NS | * | **

T novel/(T novel+T old)

Y maze

f

tBHQ diet

Trim47 +/+ | Trim47 -/-

c, b, Novel arm, a

g

- Trim47 +/+ + tBHQ n=14
- Trim47 -/- + tBHQ n=13
- Trim47 -/- n=5

Latency to plateform (s)

Day 1 | Day 2 | Day 3 | Day 4 (**)

Training days

h

tBHQ diet

Trim47 +/+ | Trim47 -/-

Day 1

Day 4

Start: ✕   Plateform: ●