Torsion-Induced Traumatic Optic Neuropathy (TITON): A Physiologically Relevant Animal Model of Traumatic Optic Neuropathy

Running Title: **Torsion-Induced Traumatic Optic Neuropathy (TITON)**

Annie K. Ryan,[1] Brooke I. Asemota,[2] Tyler Heisler-Taylor,[3] Claire Mello,[3] Luis Rodriguez,[4,5] William E. Sponsel,[6,7] Julie Racine,[8] Tonia S. Rex,[9] Randolph D. Glickman,[10] Matthew A. Reilly[1,3,*]

Affiliations

1. Department of Biomedical Engineering, The Ohio State University, Columbus, OH
2. Dept of Pediatrics, Pediatric Neurology, Duke Children's Hospital, Durham, NC
3. Department of Ophthalmology and Visual Science, The Ohio State University, Columbus, OH
4. Department of Bioengineering, The University of Texas at Dallas, Richardson, TX
5. Department of Physical Medicine & Rehabilitation, UT Southwestern Medical Center, Dallas, TX
6. WESMD Professional Associates, San Antonio, TX
7. Rosenberg School of Optometry, University of the Incarnate Word, San Antonio, TX
8. Department of Ophthalmology, Nationwide Children's Hospital, Columbus, OH
9. Vanderbilt Eye Institute, Vanderbilt University Medical Center, Nashville, TN
10. Department of Ophthalmology, UT Health-San Antonio, San Antonio, TX

To Whom Correspondence Should Be Addressed:
Matthew A. Reilly, Ph.D.
3008 Mars G. Fontana Laboratories
140 W. 19th Ave
Columbus, OH 43210
reilly.196@osu.edu

Keywords: electroretinogram; traumatic brain injury; traumatic optic neuropathy; animal model; models of injury; ocular trauma

Abstract:

Traumatic optic neuropathy (TON) is a common cause of irreversible blindness following head injury. TON is characterized by axon damage in the optic nerve followed by retinal ganglion cell death in the days and weeks following injury. At present, no therapeutic or surgical approach has been found to offer any benefit beyond observation alone. This is due in part to the lack of translational animal models suitable for understanding mechanisms and evaluating candidate treatments. In this study, we developed a rat model of TON in which the eye is rapidly rotated, inflicting mechanical stress on the optic nerve and leading to significant visual deficits. These functional deficits were thoroughly characterized up to one week after injury using electrophysiology and immunohistochemistry. The photopic negative response (PhNR) of the light adapted full field electroretinogram (LA ffERG) was significantly altered following injury. This correlated with increased biomarkers of retinal stress, axon disruption, and ganglion cell death. Together, this evidence suggests the utility of our model for mimicking clinically relevant TON and that the PhNR may be an early diagnostic for TON. We also found indirect evidence that ketamine, which was used for anesthesia, may ameliorate TON. Future studies will utilize this animal model for evaluation of candidate treatments.

1. Introduction:

Traumatic optic neuropathy (TON) is closely associated with traumatic brain injury (TBI), and results in irreversible blindness (1). TON occurs in up to 5% of all closed head injuries and 2.5% of maxillofacial and mid-face traumas (2–4). This results in upwards of 75,000 people per year developing irreversible vision loss as a result of TON. Ocular injuries are the 4th most common battlefield injury and, in the US, make up around 2.5 million emergency department visits per year (5,6).

Intervention relies primarily on high dose corticosteroid administration and/or surgical decompression; however, both have been observed to have limited success (2,3,7,8). In addition, treatment with both corticosteroids and surgical decompression have been called into question due to the increasing evidence of a higher risk of morbidity when administered in TBI patients (3,8–10). In this study, 21% of TBI participants who were administered corticosteroids died at 2 weeks whereas in the placebo group only 18% died within this timeframe (10). The authors concluded, corticosteroids should not be utilized to treat TBI patients (10). Due to the common coexistence of TBI and TON, corticosteroid administration has been minimized, resulting in an urgent need for therapeutic intervention before vision loss occurs. The first step towards this goal is development of a physiologically-relevant animal model in which to determine the cellular injury pathway and find a diagnostic criterion capable of locating visual dysfunction before significant cell loss occurs.

In general, TON is characterized by a loss of vision associated with retinal ganglion (RGC) death and axonal loss of the optic nerve; however, the exact mechanism by which the visual system is affected is not fully understood (1,11,12). Translationally relevant animal models are limited with most studies relying on optic nerve crush or transection. These models focus on a direct injury inducement, but do not address the more common indirect causes of injury (13). One indirect TON small animal model utilizes air blast trauma to develop a repeatable injury (14). This model serves to mimic the improvised explosive device blast many military personnel experience; however, this model does not mimic the torsional shearing that is sometimes observed in other forms of coup-contrecoup TBI. In our model we aim to develop a torsional indirect traumatic optic neuropathy (TITON) event.

Our injury mechanism more closely aligns with closed head TBIs as it induces a torsional shearing event similar to what the optic nerves experience during the coup-countercoup motion of the brain during the acceleration and deceleration event of TBI (8,15,16). After coup-contrecoup motion, deformational injuries to the optic nerve such as bending, shearing, and compression have been reported (15,16). As the optic nerve experiences biomechanical stress, the retina can experience injury events as well. RGCs are the cell type on the anterior portion of the retina, with axons that extend posteriorly and bundle together to form the optic nerve. Thus, since the optic nerve is largely comprised of RGC axons, this injury event may translate to the retina. The direct method in which vision loss occurs and which cell types are initially afflicted is debated.

Determination of the cells in the retina and optic nerve that are injured is crucial to
determining diagnostic and treatment protocols. We therefore employed light adapted
full field electroretinograms (LA ffERGs) to find early diagnostic protocols and to
determine the cell types afflicted after TITON.

In this study we developed a translationally relevant small animal model of TON via a
torsional rotational injury event, similar to closed head TBI. This TITON model was
thoroughly characterized using ffERG. Based on these observations, we also propose
the LA ffERG as a means of early diagnostic screening for TON.

2. Materials and Methods:

108   The overall experimental design was comprised of two phases. First, a pilot study was
109   conducted to estimate the ocular rotation required to achieve a low level of irreversible
110   functional loss. Second, a more extensive characterization of the low-level injury was
111   undertaken to evaluate potential diagnostic biomarkers.

2.1 Animals:
For the pilot study, 27 adult female Sprague Dawley rats (350-400g; Charles River
Laboratories International, Inc., Wilmington, MA) were utilized to estimate amplitude,
velocity, and degree of rotation required to achieve a low-level, irreversible injury. For
the current study, 40 ~200g male Sprague Dawley rats were used (Charles River). All
experiments were conducted in accordance with the ARVO Statement for the Use of
Animals in Ophthalmic and Visual Research and institutionally approved protocols.

2.2 Injury Induction:
A robot was constructed (**Fig. 1**) to allow the application of rapid, coaxial rotation to the
eyes of a rat. The foundation of the instrument was a rat stereotaxic (Mouse Specific
Stereotaxic Base; Leica Microsystems, Inc., Richmond, IL) that allowed complete
immobilization of the rat's head and accurate positioning of the actuator relative to the
superior-inferior axis of rotation of the left eye. The actuator (BLM-N23; Galil Motion
Control, Rocklin, CA) was fitted with a quadrature encoder (4000 counts/revolution)
yielding an angular resolution of 0.09 degrees. The actuator was controlled using a
programmable motion controller (DMC-30012; Galil Motion Control) that allowed
prescription of maximum angular displacement, velocity, and acceleration.

Figure 1: Schematic (A) and photographs (B, C) of the TITON apparatus. (A) The
instrument allows independent variation of the amplitude, rate, and direction of rotation
to either eye of the rat via a custom user interface. (B) A motor was rigidly mounted on a
stereotaxic unit to allow precise positioning of the motor relative to the eye's axis of
rotation while immobilizing the animal's head to ensure accurate rotation. (C) Rat in
stereotaxic with fixation hook attached to left eye.

135   Prior to induction of the traumatic event, an initial calibration of the robot
apparatus was conducted using the control software (GalilTools; Galil Motion Control) at

the desired parametric rate and amplitude of rotation to ensure proper connectivity of
the motor and ensure that the eye and rig were able to achieve the desired range of
motion. A scleral fixation hook (47-2650 Guthrie Fixation Hook; Sklar Instruments, West
Chester, PA) was used to rigidly connect the axis to the temporal extraocular
musculature (Fig. 1C). For the pilot study the left eyes of animals were utilized while in
the current study the right eye was subjected to rotations up to 47 degrees at angular
velocities up to 3320 degrees/s. Peak amplitude and velocity were then recorded.

2.3 Identification of Injury Thresholds
Flash visual evoked potentials (fVEPs) were recorded before, immediately after injury,
and again seven days post injury to quantify the extent of injury. The animal was
anesthetized throughout visual testing and the trauma event with ketamine (80 mg/kg)
and xylazine (8 mg/kg), which was administered intraperitoneally. Following anesthesia,
a thin layer of optical ointment (neomycin and polymyxin B sulfates, and bacitracin zinc
ophthalmic ointment; Akorn, Inc., Lake Forest, IL) was applied to the eye to prevent
drying and infection.

fVEPs utilized a ~10μs flash was presented by an external stimulator (Grass PS22D
Photic Stimulator, Grass RPS107 Regulated Power Supply; Grass Instrument Co.,
Quincy, MA). These light flashes induced fVEPs in the corresponding visual cortex of
the brain. The fVEPs were sampled as an average of 64 signal data points and were
acquired using sub-dermal electrodes (Low Profile Needle Electrode for EEG; Chalgren
Enterprises, Inc., Gilroy, CA; TDS 2014C Oscilloscope, Tektronix Inc., Beaverton, OR;
IGMEB-NUM25 Mini Electrode Board; Grass Inst. Co., Quincy, MA; 12B-8-23
Neurodata Acquisition System; Grass Inst. Co., Quincy, MA) that were placed to
maximize the signal-to-noise ratio.(17) Three data sets were collected as follows. One
minute of dark adaptation preceded fVEP followed by one minute of dark adaptation
prior to collection of a control waveform recorded in an identical way except that the eye
was not exposed to the flash stimulus to ensure the recorded activity was a direct result
of the external stimulus. Three such data sets were recorded at each time point
immediately before, immediately after, and one week after trauma.

After the fVEPs had been taken following the traumatic event, the rat was given 2.1
mg/kg yohimbine (Yobine®; Lloyd Laboratories, Shenandoah, IA) to reverse the initial
anesthesia. One week after the initial session, the rat was anesthetized as above and
another set of fVEPs were acquired using the previously mentioned protocol.

Changes in fVEP amplitude and latency at N1 and P1 were normalized against baseline
readings to account for inter-animal variability. These metrics were computed for fVEPs
immediately after injury (D0) and seven days after injury (D7).

Negative controls were placed in the apparatus and removed without applying any
ocular rotation. Positive controls were subjected to saccade-level ocular rotation. An
example of the angular position and velocity from one experiment is shown in **Fig. 2 A
& B, respectively**. Signal – control fVEP waveforms were used to measure the

difference in amplitude and latency from N1 to P1 (**Fig. 2C**). A threshold-based statistical model was used to estimate the angular rotation required to achieve a functional deficit immediately after injury as well as 7 days after injury.

Figure 2: Example of the rapid rotation applied to the specified (A) angular displacement (45 degrees) and (B) peak velocity (3000 degrees/second) to the left eye of a rat. (C) Sample fVEP before and immediately after ocular rotation shown in A and B. Overlaid is a schematic data extraction procedure for an fVEP waveform. The waveform itself was generated by collecting and averaging numerous signals following flash exposure, then subtracting an averaged signal collected in the same way without the flash.

2.4 Functional Characterization of the Injury

Prior to testing, rats were anesthetized with an IP injection of a mixture of ketamine (West-Ward; Eatontown, NJ) and xylazine (Akorn Inc.; Lake Forest, IL) (90mg/kgketamine and 10mg/kg xylazine). Once the pedal withdrawal reflex was absent, the eyes were dilated with 1% Tropicamide Ophthalmic Solution (Akorn Inc.). In preparation for functional characterization, one platinum subdermal needle electrode (Natus Manufacturing Limited; Gort, CO. Galway Ireland) was placed in the tail, another subdermal needle electrode (Natus) was placed in the center of the head, one 6mm gold surface cup electrode (Natus) was placed in the mouth and finally, after the eyes were numbed with 0.5% Tetracaine Hydrochloride Ophthalmic Solution (Bausch & Lomb Inc.; Tampa, FL) and lubricated with Systane gel (Alcon Laboratories Inc.; Fort Worth, Texas) or GentleTears gel (Alcon), the Celeris electrodes (Diagnosys LLC; Lowell, MA) were placed perpendicularly on the surface of each eye slightly touching the cornea. To record the full field electroretinograms and the flash visual evoked potentials, the Celeris system (Diagnosys LLC; Lowell, MA) was used. During the entire testing period, the internal temperature of the rats was kept at 37^{o}C.

LA ffERGs, oscillatory potentials (OPs) and fVEPs were obtained against a background of 30 cd/m^2 using a white flash (flash intensity: 200 cd.s/m^2) with an interstimulus interval (ISI) of 1000ms. An average of 50 sweeps were collected. The light-adapted flicker ERGs were also recorded. A white flash (flash intensity: 200 cd.s.m^2) was used. A total of 200 sweeps were collected with a frequency of 20 Hz. Finally, the photopic negative response (PhNR) were also recorded and used a white stimulus (flash intensity: 200 cd.s/m^2) against a green background (40 cd/m^2). An average of 100 sweeps were collected.

A-waves were measured from baseline to the first negative trough. The b-waves were measured from the a-wave peak to the highest peak of the response. Flicker ERG and OPs were measured from preceding trough to peak. OPs were calculated utilizing the sum of the OPs (SOP) and only OPs that occur prior to the peak of the b-wave were utilized (SOP= OP1-2 + OP3 + OP4 + OP5). PhNR amplitudes were measured from baseline to the trough following the b-wave. Peak times were measured utilizing the time of the flash onset to respective peaks.

Visual electrophysiology testing was performed at baseline (no more than 7 days before
injury event) and at 1 and 7 days (D1 and D7, respectively) after injury. At the end of the
testing period, the animals were injected with an IP dose of Antipamizole (Modern
Veterinary Therapeutics LLC; Miami, FL) (2mg/kg) to reverse the effect of xylazine and
ease the recovery of the animals.

2.5 Histology and Immunohistochemistry
Animals were humanely euthanized immediately following D1 or D7 ERG
measurements using established and approved IACUC protocol as described below.
Animals underwent CO_2 asphyxiation followed by a lateral incision of the atlanto-
occipital joint in preparation of tissue collection. Briefly, the skull was removed. The
brain was reflected rostrally to expose the optic nerves. The nerves were then
separated from the brain, and the brain was removed. The nerves and eyes were then
dissected en bloc for immunohistochemistry analysis. For immunohistochemistry, rats
were perfused with phosphate-buffered saline (PBS) followed by 4% paraformaldehyde,
and the eye and optic nerve were dissected and incubated overnight at 4°C in 4%
paraformaldehyde. The optic nerves and chiasm were dissected away from the brain
and cryo-preserved in 20% sucrose in PBS for 2 hrs at 4°C followed by 30% sucrose in
PBS overnight at 4°C. Tissues were embedded in OCT (Fisher Scientific, Waltham, MA)
and longitudinally sectioned on a cryostat at 10 μm thickness. For
immunohistochemistry, sections were incubated in PBS to remove the OCT and then
incubated in 1:20 normal donkey serum in PBS plus Triton-X-100 (PBT) at room
temperature for 2 hrs. Sections were then incubated with specific antibodies as follows:
anti-β-tubulin (1:1000; MAB5564; Millipore, Burlington, MA) and anti-glial fibrillary acidic
protein (GFAP; 1:50; Z0334; DAKO, Santa Clara, CA) in PBT overnight at 4°C, rinsed
with PBS, and incubated in donkey anti-mouse Alexa 488 and donkey anti-rabbit Alexa
594 (1:200; Fisher Scientific) in PBT overnight at 4°C. Finally, the sections were rinsed,
mounted in Vectashield plus DAPI (Vector laboratories, Burlingame CA). Tissue
samples were imaged on a wide-field fluorescence microscope (Nikon Eclipse, Melville,
NY) or confocal microscope (Nikon AXR) using consistent settings.

A terminal deoxynucleotidyl transferase (dUTP) nick end labeling (TUNEL) assay was
performed to assay cell death. Eyes with optic nerves attached were assayed. A 28-
gauge needle was utilized at the temporal limbus of the eye to allow better perfusion of
4% PFA to the ocular interior. Eyes and nerves were then placed in 4% PFA and gently
agitated for 2 hours before 2x 15-minute PBS wash. The samples were then transferred
to 30% sucrose and incubated overnight at 4°C, then embedded in OCT (Fisher
Scientific, Waltham, MA) and longitudinally sectioned on a cryostat at 10 μm thickness.
Slides were then heated for 10 minutes on low heat. Once at room temperature, slides
were soaked in an ABT solution for 2x 15min cycles with gentle shaking. Slides were
then washed 2x with PBS for 10mins. TUNEL staining was completed following
manufacturer's instructions (TMR red kit, cat#: 12156792910, Roche) and with
previously reported methods (18). After TUNEL kit application slides were washed 2x in
PBS for 5 minutes. Hoechst nuclear stain was then applied to the slides for 10 minutes

in a humidity chamber at room temperature. Slides were washed in PBS for 5 minutes
before coverslips were mounted using 50% glycerol. Samples were then imaged with a
Nikon Eclipse Ts2R fluorescent microscope with a panda sCMOS camera with NIS-
Elements AR software (18). TUNEL positive (TUNEL+) cells counts were determined
through MATLAB code quantification.

3. Results:

3.1 Injury Threshold Estimation

Optic neuropraxia, defined as a reversible functional deficit in fVEP, was observed at
insults with experimental extents of rotation between 20.5° - 38.1° (corresponding
angular velocities 1475 - 3320°/s), which exceeded the maximal physiological saccadic
parameters of 10° at 500°/s (**Fig. 3A,C**). Immediately after injury, fVEP amplitude
decreased with faster peak times   in a dose-dependent trend above 20.5° (effects of -
0.0221 and -0.0090 mV/°; $p < 0.0001$ and 0.0011, respectively).

Figure 3: Normalized functional changes in fVEP amplitude (A, B) and latency (C, D) as
a function of the angle of rotation Θ. Left: Both fVEP amplitude (A) and latency (C) were
negatively correlated with the amplitude of ocular rotation above 20.5 degrees. Right:
One week after ocular rotation, the fVEP amplitude increased with amplitude of ocular
rotation above 38.1 degrees (B) while both fVEP amplitude and latency had higher
variability. Subscripts 1 and 7 refer to the after-injury timepoint (i.e. either day 1 or day
7).

TON, defined as statistically significant changes from baseline measured one week
after injury, was induced by rotations of larger than 38.1° with peak velocity greater than
3320°/s (**Fig. 3B,D**). Specifically, the final fVEP amplitude increased significantly
relative to baseline above this threshold (0.1257 per degree; $p = 0.0005$), while the
variance in both amplitude and latency increased ($p = 0.0313$ and 0.0317, respectively).
The magnitude of the latency change was highly variable and did not exhibit any
particular trend with respect to the angle of rotation (0.0113 per degree; $p = 0.8168$).

3.2 Functional Characterization of Injury

Significant alterations to the amplitude of the PhNR were observed following injury
(**Fig.4F**). The right, injured, eye (OD), showed a significant PhNR amplitude alteration
between D1 and D7 ($p = 0.0002$), with D1 amplitudes being decreased as compared to
elevated D7 amplitudes. Also of note, OD showed a decrease in amplitude at D1
($p = 0.0225$) as compared to OS. OD D7 PhNR amplitudes were found to significantly
overshoot OD Baseline amplitudes ($p = 0.0225$). In the injured group PhNR amplitude
significant changes were depicted for the following: OD D1 vs OS D7 ($p < 0.0001$), OD
Baseline vs OS D7 ($p = 0.0012$), OS Baseline vs OS D7 ($p = 0.0008$), OS Baseline vs OD
D7 ($p = 0.0094$).

Figure 4: Average ERG amplitudes between injured and experimental groups. ffERG a-
wave amplitudes with significant amplitude differences noted between OD Baseline and
OD D7 ($p = 0.0051$), OS Baseline and OD D7 ($p = 0.0010$), and OS Baseline and OS D7

(p=0.0294) (A). ffERG amplitudes of OD D1 compared to OS D7 were determined to be
statistically significant (B). No statistically relevant differences were noted with regard to
FVEP amplitudes (C). SOP amplitudes showed significant alteration between OD D1
and OD D7 (p=0.0251) (D). Flicker amplitudes depicted significant changes between
OS Baseline and OD D7 (p=0.0125) (E). Significant differences were noted for the
injured animal PhNR amplitudes between OD D1 and OD D7 (p=0.0002), OD D1 and
OS D7 (p<0.0001), OD D1 and OS D1 (p=0.0225), OS Baseline and OS D7 (p=0.0008),
OS Baseline and OD D7 (p=0.0094), OD Baseline and OS D7 (p=0.0012), and OD
Baseline and OD D7 (p=0.0129) (F).

A significant rise in amplitude was detected between OD D7 and OD Baseline a-wave
amplitude (p=0.0051). Additionally, we observed a-wave amplitude statistical
significances between OS Baseline and OD D7 (p=0.0010) and OS Baseline and OS
D7 (p=0.0294). The b-wave showed one statistically significant amplitude increase
between OD D1 and OS D7 (p=0.0153). The SOPs had one detectable significant
alteration between the injured OD at D1 as compared to the right eye sham control at
D7 (p=0.0251). The flicker waveform depicted one statistically significant rise in of
amplitude between OS Baseline and OD D7 (p=0.0125). We did not observe any
statistically significant alterations in the fVEP waveforms between eyes or between
groups. Additionally, we did not observe any significant changes to peak latency for any
of the electrophysiological waveforms.

3.3 Histology and Imaging:
3.3.1   Immunofluorescent Staining
Fluorescent labeling of the retina in injured eyes revealed increased GFAP labeling at
D7 (**Fig. 5C,H**). This suggests the presence of stressed conditions within the retina.
Increased GFAP labeling was observed to extend from the ganglion cell layer into the
photoreceptor layer, which indicated Müller cells were becoming hypertrophic. GFAP
labeling in injured D1 retinas appeared to be thickening at the RGC layer, but retinas did
not appear to have increased Müller glial extensions (**Fig. 5A,F**). DAPI and β-tubulin did
not show a significant alteration between injured eye, contralateral eye, or sham eyes.

Figure 5: Representative epifluorescence retinal images labeled with the nuclear
marker, DAPI, and immunolabeled for the axon marker, β-tubulin (green), and the glial
marker, GFAP (red) (A-E). GFAP marked channels (F-J). Injured eye D1 (A,F);
contralateral eye (B,G); injured eye D7 (C,H); contralateral eye (D, I); sham (E,J). GFAP
appears to be thicker at the retinal ganglion cell (RGC) layer of D1 injured retinas (A,F).
The injured eye at D7 was observed to develop increased hypertrophic activity of the
Müller glial cells as observed by the increased GFAP presence and the extensions of
the Müller glial cells from the RGC bodies through the retina to the photoreceptors (C,
H). These extensions were not observed in the contralateral eye at D1 (B,G) or D7 (D,
I). Nor were they represented in the sham (E,J), or the injured eye at D1 (A,F).

D7 optic nerve immunohistochemical analysis found decreased labeling intensity β-
tubulin in injured nerves relative to sham (**Fig. 6D**). Changes were observed bilaterally,

though not symmetrically, suggesting that the biomechanical mechanism of injury
employed must transmit stresses through the chiasm. Biomechanical stress
transmission through the chiasm was therefore investigated and directly observed in a
cadaver animal following craniotomy.

Figure 6: Representative epifluorescence micrographs of longitudinal sections of optic
nerves labeled with the nuclear marker, DAPI, and immunolabeled for the axon marker,
β-tubulin (green), and the glial marker, GFAP (red): uninjured (A); uninjured, no primary
antibody (B); left (C) and right (D) optic nerves of D7 injured animal; left (E) and right (F)
optic nerves of D1 injured animals. In the control (A), β-tubulin is highly punctate with
marginally higher intensity near the surface of the nerve; GFAP is most prevalent in
localized gaps between β-tubulin and slightly brighter near the surface. (C) In the
contralateral ON, β-tubulin has a lower intensity throughout and is disrupted across a
diagonal portion of the ON (yellow arrow); GFAP also has a significantly lower intensity
across the ON. (D) In the ipsilateral eye, β-tubulin labeling was lower intensity and more
diffuse in most of the ON but with large, high-intensity puncta at left; larger bundles of
GFAP were colocalized with axial columns of DAPI, but also showed some radial
bundling. β-tubulin presence in the injured eye at D1 (F) appears to have a lower
intensity as compared to the contralateral eye (E).

3.3.2 TUNEL
TUNEL+ cells were detected in the retina and optic nerve of injured animals in both OD
and OS (**Fig. 7**). Generally, retinas showed similar counts of TUNEL+ cells in both eyes
at each timepoint. Both OD and OS retinas appeared to feature a rise in TUNEL+ cells
by D7 (**Fig. 7 A-D,I**). Optic nerve TUNEL staining showed an increased amount of
TUNEL+ cells detected in OD samples at both D1 and D7 as compared to the
contralateral eye; however, there were no statistically significant differences detected
(**Fig. 7 E-H,J**).

Figure 7: Representative TUNEL micrographs of retina (A-D) and optic nerves (E-H).
Some TUNEL positive (TUNEL+) cells were detected 24 hours after injury (D1) with an
increase in TUNEL+ cells in both retina and optic nerve occurring by 7 days after injury
(D7) (I,J). TUNEL+ cell counts were higher in OD optic nerves as compared to OS optic
nerves at both D1 and D7 (J).

4. Discussion:

Based on our findings and clinical considerations, we believe that the PhNR of the LA
ffERG may be the most beneficial test for early diagnosis of TON since we observed
statistically significant alterations in amplitudes between the injured eye and
contralateral eye 24 hours after injury inducement (D1), and we also observed a
significant alteration in PhNR amplitude between D1 and D7 (**Fig. 4**) (8,19). Our retinal
immunofluorescent staining supports these findings with our observations of increased
GFAP staining at D1, via the thickening of Müller glial cell bodies in the RGC layer of
the retina at D1 and then the increased hypertrophic activity of the Müller glial cells

observed in the retinas of animals at D7 (**Fig.5**). Epifluorescence indicated significant
changes in both β-tubulin and GFAP labeling within the injured optic nerve (**Fig. 6**).
Reduced β-tubulin labeling suggests a decrease in the number of microtubules within
each axon and/or fewer axons. Furthermore, we observed an increase in TUNEL+ cells
in injured retinas and optic nerves, supporting our model of torsion-induced TON.
Combined with diminished PhNRs, these findings support the hypothesis that our model
is causing a loss of activity in the ganglion cells, which is indicative optic neuropathy.

Together, these findings provide novel insights into the secondary neurodegenerative
mechanisms leading to vision loss from TON. The developed animal model represents
an excellent platform for studying TON, developing diagnostics, and evaluating
candidate treatments for safety and efficacy. This model has tremendous potential for
evaluation of both pre- and post-traumatic therapies.

ffERGs are a series of visual stimuli-electrophysiologic response waveforms which
serve to analyze the functional output of the various cell types within the visual system.
Light adapted ERGs require no dark adaptation time, which makes them relatively faster
for real world injury applications. The different cell types within the retina can be
observed through the inducement of various light stimulating protocols. Generally, the a-
wave of the LA ffERG consists of the photoreceptor response. The a-wave is the first
waveform of the ffERG and arises due to the hyperpolarization of the photoreceptor
membranes during phototransduction (20–23). The b-wave is the positive component of
the ffERG signal and reflects the depolarization of the bipolar cells (20–23).The OPs are
observed in the ffERG as the oscillations on the ascending limb of the b-wave, and in
rodents the OPs can also be on the descending limb as well (22–24). These signals are
generated by the inhibitory effect of the amacrine cells. The amacrine cells release γ-
aminobutyric acid (GABA) and glycine, which act as inhibitory neurotransmitters, and
create the oscillatory features observed (22,23). The PhNR is the negative deflection
observed after the b-wave, and is representative of retinal ganglion cell (RGC) and
amacrine cell activity (25–27). The VEP is a waveform which can assess the post-retinal
function by characterizing the electrical output of the visual cortex (28). The light-
adapted flicker ERG at 20Hz is known to isolate the cone function in rat retinas (29).
With analysis of each of these waveforms we can determine which cell types are
afflicted after injury and find a diagnostic criterion for TON.

During our ffERG analysis we noticed the sham animals were displaying rising signals
similar to our left (contralateral) eye (OS) recordings. Repetitive ketamine dosing may
have impacted our inner retinal cells as the inner retinal ffERG waveforms such as the
PhNR, b-wave, and SOPs displayed increasing signals over time. This phenomenon
has been reported in other literature and we believe ketamine has been acting on these
inner retinal cells via the mechanisms outlined in **Figure 8** (30–34). Briefly, after TBI,
glutamate levels rise, and NMDA receptor function increases, and ketamine has been
reported to block both glutamate and NMDA receptors. The highest concentration of
NMDA receptors are found in the inner retinal cells, which are also the cell types

dominating the PhNR, b-wave, and SOPs (33). Due to this, we believe the back-to-back
ketamine doses at injury and then 24 hours later for D1 ERG may be depressing the
inner retinal ERG dependent waveforms. This would be why our sham animals are
following the same trend of our contralateral controls. Our study design inherently
controls for this by ensuring all animals have the same number and timing of anesthetic
events. In addition, we normalized the electroretinogram data against baseline
measurements, but there were not any significant changes between the non-normalized
and normalized datasets, indicating changes observed were significant to the model
and not the result of miscellaneous factors.

Figure 8: Potential flow chart of the injury cascade of events following TBI/TON
inducement. Importantly, the injury event increases the activity of the NMDA receptor
and elevates glutamate levels. Both of these factors are then attenuated by ketamine,
which can lead to electrophysiological changes.

The observation of ketamine's effect after repetitive dosing poses an interesting theory
as to whether it can be used as a neuroprotective therapeutic, especially since it can
target inner retinal cells via the suppression of the NMDA receptor. One rodent study
investigating neuroprotection after light induced damage reported ketamine/xylazine as
a potential pretreatment neuroprotectant (32). In the future we plan to elucidate
therapeutic strategies for retinal cell preservation. Alterations of the PhNR amplitude
indicate inner retinal cell damage, so therapeutics capable of preservation of this area of
the retina may prove to be the most beneficial in minimizing cell and vision loss after
injury. Other potential therapeutic strategies could target the anti-inflammatory aspect of
TON. The inflammatory response that happens in the days following the primary injury
may be just as harmful if not more damaging than the initial injury event. Being able to
suppress this inflammatory response may help to prevent the subsequent damage
characteristic of TON. Potential therapeutics for this type of therapeutic intervention
include ibudilast, TUDCA, cyclosporine, erythropoietin, dopamine, and vitamins such as
C and E (1,35–41).

This study has potential limitations. One such limitation was the utilization of only one
animal sex for each portion of the study: animals were either all female or all male but
did not include subjects from both sexes for each sub experiment. Due to timing the
main experiment utilized male rodents as their hormonal levels did not change
throughout the month, resulting in more consistent ffERGs. In future work, both sexes of
animals should be evaluated for diagnostic and therapeutic intervention strategies.
Generally, men are reported to have the highest incidence of TBI; however, women
suffering TBI have a greater risk of death as compared to male counterparts (42). This
indicates the biomechanical mechanisms governing TBI and potentially TON may differ
with regard to sex. Another limitation to this study is we utilized LA ffERGs, which target
more cone receptors and rodents are known to be a rod dominated species (43). We
may have observed larger deficits in ffERG amplitudes had we utilized dark adapted
ffERGs. LA ffERGs were chosen as they are quicker to conduct in a clinical setting

since they avoid dark adaptation time. Finally, the mechanism of TON induction used in this study was designed to provide a repeatable mechanical stimulus rather than mimic what might be considered a more typical TON-inducing injury. This decision was taken to maximize experimental precision and thereby reduce the number of animals required to achieve statistically significant results when evaluating candidate therapeutics. In reality, TON may be caused by many unique insults; however, the end result of RGC death and irreversible visual dysfunction is thought to be quite similar to what we observed in our model.

5. Conclusions

We have shown that our torsion-induced TON animal model can produce a diagnosable functional change in ffERGs via analysis of the PhNR. This functional change correlates with stressed conditions in the retina as evidenced by our tissue analysis. This is the first step in understanding how the retina and optic nerve are afflicted after TON insult. Future studies will determine therapeutic intervention strategies capable of electrophysiological function in the retinal neuronal circuits. Through our therapeutic intervention we hope to elucidate the cellular mechanisms responsible for the characteristic visual system deterioration of TON. By selecting therapeutics which target different suspected pathways of secondary neurodegeneration we hope to determine how TON progresses and at which time points therapeutic intervention would be the most efficacious.(1)

6. Transparency, Rigor, and Reproducibility:

All anesthetic events were controlled by utilizing the same dosing schedule and quantities between animals and groups. In addition, ffERG amplitude differences were controlled by comparing animals of the same sex and by collecting all visual electrophysiological data during the same time of day. This set of controls negates sex-based differences and then also ensures circadian rhythm effects have been minimized. All animals used were withing the same weight range, minimizing animal-to-animal differences. Both sham animals and contralateral eyes (OS) were used as controls. Optic nerves cross with partial decussation at the chiasm, but nerve fibers never coalesce to form new, unique fibers. We hypothesized there may be an intrachiasmal transference between the nerves due to our injury event. As such, we utilized sham animals as an additional control parameter. The injury device is controlled and allows for a higher rate of reproducibility as the motorized system executes the torsional motion. This motion is preprogrammed in terms of degrees, velocity, and torque. This allows the user to ensure the parameters of the injury event are the same between animals.

7. Acknowledgements:

This work was partially funded by US Department of Defense Vision Research Program Awards W81XWH-15-1-0074 and W81XWH-22-1-0989. The opinions or assertions contained herein are the private views of the authors and are not to be construed as

official or as reflecting the views of the Department of the Army or the Department of Defense. This research grant was supported in part by the Ohio Affiliate of Prevent Blindness Young Investigator Student Fellowship Award for Females Scholars in Vision Research. Services were performed at the OSU Vision Sciences Research Core Program under P30EY032857. We would like to thank The Ohio State University laboratory and animal resources (ULAR), Dr. Stacey Meeker, Dr. Colleen Cebulla. In addition, we would like to thank Reilly undergraduate lab members: Stephanie Small, Michelle Mosko, Emma Lally, Vidhya Kannan, Sam Duckworth, and Eve Howard. We would also like to thank Dr. Wade Rich and Elizabeth Urbanski.

8. Authorship confirmation/contribution statement:

AKR contributed to data collection, analysis, and drafted the manuscript. BIA contributed to experimental design, data collection, data analysis, and manuscript editing. THT and CM contributed to data collection and analysis. LR contributed to data collection and manuscript editing.TR contributed to data collection, analysis, and edited the manuscript. WES contributed to experimental conceptualization, experimental design, and manuscript editing. RDG contributed to experimental design, data collection, and manuscript editing. MAR contributed to experimental conceptualization, experimental design, and manuscript editing. All authors contributed to the literature review and discussions and approved the submitted version.

9. Author(s') Disclosure Statements:

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

10. Funding Statement:

This work was supported by the US Department of Defense Vision Research Program Awards W81XWH-15-1-0074 and W81XWH-22-1-0989. This research grant was supported in part by the Ohio Affiliate of Prevent Blindness Young Investigator Student Fellowship Award for Females Scholars in Vision Research.

11. References:

1. Ryan AK, Rich W, Reilly MA. Oxidative stress in the brain and retina after traumatic injury. Front Neurosci [Internet]. 2023;17. Available from: https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2023.1021152

2. Chan JW. Optic Nerve Disorders: Diagnosis and Management [Internet]. 1st ed. New York, NY: Springer; 2008. 284 p. Available from: https://doi.org/10.1007/978-0-387-68979-1

3. Sarkies N. Traumatic optic neuropathy. Eye. 2004 Nov 1;18(11):1122–5.

4. Prevention C for DC and. Report to congress: Traumatic brain injury in the United
States. 2016; Available from:
https://www.cdc.gov/traumaticbraininjury/pubs/tbi_report_to_congress.html

5. Scott R. The injured eye. Philos Trans R Soc B Biol Sci. 2011 Jan
27;366(1562):251–60.

6. Sherwood D, Sponsel WE, Lund BJ, Gray W, Watson R, Groth SL, et al. Anatomical
Manifestations of Primary Blast Ocular Trauma Observed in a Postmortem Porcine
Model. Invest Ophthalmol Vis Sci. 2014;55(2):1124–32.

7. Steinsapir KD, Goldberg RA. Traumatic optic neuropathy. Surv Ophthalmol. 1994
Jun;38(6):487–518.

8. Blanch RJ, Joseph IJ, Cockerham K. Traumatic optic neuropathy management: a
systematic review. Eye. 2024 Aug 1;38(12):2312–8.

9. Steinsapir KD, Goldberg RA. Traumatic optic neuropathy: an evolving
understanding. (1879-1891 (Electronic)).

10. Ian Roberts, Yates D, Sandercock P, Farrell B, Wasserberg J, Lomas G, et al. Effect
of intravenous corticosteroids on death within 14 days in 10008 adults with clinically
significant head injury (MRC CRASH trial): randomised placebo-controlled trial.
Lancet. 2004 Oct;364(9442):1321–8.

11. Tse BC, Dvoriantchikova G, Tao W, Gallo RA, Lee JY, Pappas S, et al. Tumor
Necrosis Factor Inhibition in the Acute Management of Traumatic Optic Neuropathy.
Invest Ophthalmol Vis Sci. 2018 Jun 1;59(7):2905–12.

12. Khan RS, Ross AG, Aravand P, Dine K, Selzer EB, Shindler KS. RGC and Vision
Loss From Traumatic Optic Neuropathy Induced by Repetitive Closed Head Trauma
Is Dependent on Timing and Force of Impact. Transl Vis Sci Technol. 2021
Jan;10(1):8.

13. Burke EG, Cansler SM, Evanson NK. Indirect traumatic optic neuropathy: modeling
optic nerve injury in the context of closed head trauma. Neural Regen Res [Internet].
2019;14(4). Available from:
https://journals.lww.com/nrronline/fulltext/2019/14040/indirect_traumatic_optic_neur
opathy__modeling.9.aspx

14. Bricker-Anthony C, Hines-Beard J, Rex TS. Molecular changes and vision loss in a
mouse model of closed-globe blast trauma. Invest Ophthalmol Vis Sci. 2014 Jul
3;55(8):4853–62.

15. Li Y, Singman E, McCulley T, Wu C, Daphalapurkar N. The Biomechanics of Indirect
Traumatic Optic Neuropathy Using a Computational Head Model With a Biofidelic
Orbit. Front Neurol. 2020;11:346.

16. Galgano M, Toshkezi G, Qiu X, Russell T, Chin L, Zhao LR. Traumatic Brain Injury:
Current Treatment Strategies and Future Endeavors. Cell Transplant. 2017
Jul;26(7):1118–30.

17. You Y, Klistorner A, Thie J, Graham SL. Improving reproducibility of VEP recording
in rats: electrodes, stimulus source and peak analysis. Doc Ophthalmol Adv
Ophthalmol. 2011 Oct;123(2):109–19.

18. Heisler-Taylor T, Kim B, Reese AY, Hamadmad S, Kusibati R, Fischer AJ, et al. A
new multichannel method quantitating TUNEL in detached photoreceptor nuclei.
Exp Eye Res. 2018 Nov;176:121–9.

19. Dhillon A, Ahmad MSZ, Breeze J, Blanch RJ. Prolonged deployed hospital care in
the management of military eye injuries. Eye Lond Engl. 2020 Nov;34(11):2106–11.

20. Perlman I. University of Utah Health Sciences Center. In: Webvision: The
Organization of the Retina and Visual System [Internet] [Internet]. Salt Lake City,
UT: University of Utah Health Sciences Center; 2001. Available from:
https://www.ncbi.nlm.nih.gov/books/NBK11554/

21. Reichel E, Klein K. Retinal Electrophysiology. In: Ophthalmology [Internet]. 5th ed.
Elsevier; 2019. Available from: https://www.clinicalkey.com/#!/content/book/3-s2.0-
B9780323528191002875

22. Audo I, Robson AG, Holder GE, Moore AT. The negative ERG: clinical phenotypes
and disease mechanisms of inner retinal dysfunction. Surv Ophthalmol. 2008
Feb;53(1):16–40.

23. Mallery R, Mackay D, Prasad S. Ocular Functional and Structural Investigations. In:
Bradley and Daroff's Neurology in Clinical Practice [Internet]. 8th ed. Elsevier; 2022.
p. 601–13. Available from: https://www.clinicalkey.com/#!/content/book/3-s2.0-
B9780323642613000437?scrollTo=%23hl0000474

24. Rosolen SG, Rigaudière F, LeGargasson JF, Chalier C, Rufiange M, Racine J, et al.
Comparing the photopic ERG i-wave in different species. Vet Ophthalmol. 2004
Jun;7(3):189–92.

25. Li B, Barnes GE, Holt WF. The decline of the photopic negative response (PhNR) in
the rat after optic nerve transection. Doc Ophthalmol Adv Ophthalmol. 2005
Jul;111(1):23–31.

26. Jnawali A, Lin X, Patel NB, Frishman LJ, Ostrin LA. Retinal ganglion cell ablation in
guinea pigs. Exp Eye Res. 2021 Jan;202:108339.

27. Miura G, Wang MH, Ivers KM, Frishman LJ. Retinal pathway origins of the pattern
ERG of the mouse. Exp Eye Res. 2009 Jun 15;89(1):49–62.

28. Ridder WH, Nusinowitz S. The visual evoked potential in the mouse—Origins and
response characteristics. Vision Res. 2006;46(6):902–13.

29. Goto Y, Yasuda T, Tobimatsu S, Kato M. 20-Hz flicker stimulus can isolate the cone
function in rat retina. Ophthalmic Res. 1998;30(6):368–73.

30. Choh V, Gurdita A, Tan B, Feng Y, Bizheva K, McCulloch DL, et al. Isoflurane and
ketamine:xylazine differentially affect intraocular pressure-associated scotopic
threshold responses in Sprague-Dawley rats. Doc Ophthalmol Adv Ophthalmol.
2017 Oct;135(2):121–32.

31. Prando S, Carneiro C de G, Otsuki DA, Sapienza MT. Effects of ketamine/xylazine
and isoflurane on rat brain glucose metabolism measured by (18) F-
fluorodeoxyglucose-positron emission tomography. Eur J Neurosci. 2019
Jan;49(1):51–61.

32. Arango-Gonzalez B, Schatz A, Bolz S, Eslava-Schmalbach J, Willmann G, Zhour A,
et al. Correction: Effects of Combined Ketamine/Xylazine Anesthesia on Light
Induced Retinal Degeneration in Rats. PLOS ONE. 2012 Aug
6;7(8):10.1371/annotation/d52a610e-6a56-4158-90bf-45e37f053567.

33. Shen Y, Liu XL, Yang XL. N-methyl-D-aspartate receptors in the retina. Mol
Neurobiol. 2006 Dec;34(3):163–79.

34. Deng Y, Thompson BM, Gao X, Hall ED. Temporal relationship of peroxynitrite-
induced oxidative damage, calpain-mediated cytoskeletal degradation and
neurodegeneration after traumatic brain injury. Exp Neurol. 2007 May;205(1):154–
65.

35. Bond WS, Rex TS. Evidence That Erythropoietin Modulates Neuroinflammation
through Differential Action on Neurons, Astrocytes, and Microglia. Front Immunol.
2014;5:523.

36. Daruich A, Picard E, Boatright JH, Behar-Cohen F. Review: The bile acids urso- and
tauroursodeoxycholic acid as neuroprotective therapies in retinal disease. Mol Vis.
2019;25:610–24.

37. Cueva Vargas JL, Belforte N, Di Polo A. The glial cell modulator ibudilast attenuates
neuroinflammation and enhances retinal ganglion cell viability in glaucoma through
protein kinase A signaling. Neurobiol Dis. 2016 Sep;93:156–71.

38. Bernardo-Colón A, Vest V, Clark A, Cooper ML, Calkins DJ, Harrison FE, et al.
Antioxidants prevent inflammation and preserve the optic projection and visual
function in experimental neurotrauma. Cell Death Dis. 2018 Oct 26;9(11):1097.

39. DeJulius CR, Bernardo-Colón A, Naguib S, Backstrom JR, Kavanaugh T, Gupta
MK, et al. Microsphere antioxidant and sustained erythropoietin-R76E release

functions cooperate to reduce traumatic optic neuropathy. J Control Release Off J
Control Release Soc. 2021 Jan 10;329:762–73.

40. Lan YL, Li S, Lou JC, Ma XC, Zhang B. The potential roles of dopamine in traumatic
brain injury: a preclinical and clinical update. Am J Transl Res. 2019;11(5):2616–31.

41. Lendvai-Emmert D, Magyar-Sumegi ZD, Hegedus E, Szarka N, Fazekas B, Amrein
K, et al. Mild traumatic brain injury-induced persistent blood–brain barrier disruption
is prevented by cyclosporine A treatment in hypertension. Front Neurol [Internet].
2023;14. Available from:
https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2023.1252796

42. Munivenkatappa A, Agrawal A, Shukla DP, Kumaraswamy D, Devi BI. Traumatic
brain injury: Does gender influence outcomes? Int J Crit Illn Inj Sci. 2016
Jun;6(2):70–3.

43. Galindo-Romero C, Norte-Muñoz M, Gallego-Ortega A, Rodríguez-Ramírez KT,
Lucas-Ruiz F, González-Riquelme MJ, et al. The retina of the lab rat: focus on
retinal ganglion cells and photoreceptors. Front Neuroanat [Internet]. 2022;16.
Available from:
https://www.frontiersin.org/journals/neuroanatomy/articles/10.3389/fnana.2022.9948
90

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A

Microcontroller

Motor

PC

Rat

B

C

Figure 3