Endogenous tagging of Munc13-1 with a SNAP tag as a tool for monitoring synapse nanoarchitecture

Maria Kowald[1,‡], Sylvestre P. J. T. Bachollet[1,‡], Fritz Benseler[2], Maria Steinecker[3], Sofia Kaushik[1], Tolga Soykan[1], Sun Siqi[1], Ramona Birke[1], Dragana Ilic[1], Nils Brose[2], Hanna Hörnberg[3], Martin Lehmann[1], Silvio O. Rizzoli[4,5,6], Johannes Broichhagen[1,*], Noa Lipstein[1,2*]

[1]   *Leibniz-Forschungsinstitut für Molekulare Pharmakologie (FMP), Berlin, Germany*

[2]   *Max Planck Institute for Multidisciplinary Sciences, Göttingen, Germany*

[3]   *Max Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), Berlin, Germany*

[4]   *Department of Neuro- and Sensory Physiology, University Medical Center Göttingen, Germany*

[5]   *Excellence Cluster Multiscale Bioimaging (MBExC), Göttingen, Germany*

[6]   *Center for Biostructural Imaging of Neurodegeneration (BIN), University Medical Center Göttingen, Göttingen, Germany*

KEYWORDS Active zone, Munc13-1, Unc13a, Synapse, SNAP tag, SBG-SiR-d12.

ABSTRACT: Synaptic function is governed by highly regulated protein machineries, whose abundance and spatial localization change continually. Studies to determine the dynamic changes in synaptic proteins nanoarchitecture typically rely on immunolabeling, or on the expression of fluorescent proteins. The former employs chemical fluorophores and signal amplification, but requires fixation. The latter disturbs the cells minimally, but uses suboptimal fluorophores. Self-labeling tags have been introduced to combine the advantages of these approaches, and here we introduce a knock-in mouse line where the essential synaptic protein Munc13-1 is endogenously fused to the self-labeling SNAP tag. We demonstrate efficient Munc13-1-SNAP labeling in fixed neurons and brain sections by various SNAP dyes, as well as by a novel bright and far-red compound, SBG-SiR-d12, which we introduce here. SBG-SiR-d12 is designed as a membrane impermeable dye, but we repurposed it to stain cytosolic Munc13-1-SNAP in permeabilized, fixed cells and tissue, where we find it outperforms other dyes as evaluated by both conventional and super-resolution microscopy. Finally, we show that Munc13-1-SNAP can also be monitored by live cell imaging. We conclude that the Munc13-1[SNAP] mouse line is a useful tool for the analysis of Munc13-1 nanoarchitectural dynamics in synapses, with a potential for wide adoption.

INTRODUCTION

Dynamic changes in protein copy numbers, complex composition, and nanoscale organization are often targets in the regulation of cellular activity. To reliably monitor proteins in time and space, efficient labeling strategies have been developed. A key development has been the introduction of fluorescent protein tags (e.g. green fluorescent protein, GFP[1]). However, and despite continuous improvements, the photophysical properties of fluorescent proteins often fall short compared to chemical dyes in terms of brightness and stability[2, 3]. To address this limitation, self-labeling tags were created - engineered protein tags that can be genetically encoded and covalently bind bioorthogonal synthetic probes. Belonging to this group, the SNAP tag is an engineered O^6-alkylguanine-DNA alkyltransferase that binds O^6-benzylguanine (BG) derivatives in a covalent, non-reversible manner[4-6]. As such, proteins fused to a SNAP tag can be visualized in live cells or in fixed tissue by the addition of a BG-fluorophore conjugate. This mode of protein labeling is advantageous as an alternative to genetic conjugation with fluorescent protein tags, because it enables the flexible attachment of bright and stable dyes to the protein of interest, in the living cell, and at a time of choice. The small tag size (~19 kDa) is optimal for super-resolution imaging, where it is desired to have a fluorophore in close proximity to the protein of interest[7, 8]. Moreover, flexible use of chemical dyes can enable pulse-chase labeling[9-12] or signal multiplexing[13]. The toolkit for SNAP tag labeling is rapidly expanding: BG derivatives carrying an array of fluorophores are available, and several chemical modifications of the BG-dye conjugates have been introduced to modify their properties, for example to make the conjugate membrane impermeable for extracellular labeling[10] or to boost labeling kinetics[14].

Here, we opted to leverage the advantages of self-labeling tags and develop novel tools to monitor synapses. Munc13-1 (protein unc-13 homolog A, encoded by the *Unc13a* gene) is a presynaptic protein with a central function in the preparation of synaptic vesicles (SVs) for fast exocytosis[15, 16]. Munc13-1 is expressed in the majority of neuronal subtypes in the central and peripheral nervous system, as well as in some neurosecretory cell types including chromaffin cells or insulin-releasing beta-pancreatic cells[17, 18]. In neurons, it resides in the active zone, a protein-dense compartment at the presynaptic membrane where SVs undergo fusion, and is absolutely essential for synaptic transmission. At the molecular level, Munc13-1 catalyzes the formation of SNARE complexes, that link the SV membrane with the presynaptic plasma membrane, thus making SVs fusion competent[19, 20]. Munc13-1 function sets multiple synapse properties, including synaptic strength, the release probability of SVs, the rate of SV replenishment after depletion, and synaptic plasticity[16, 20-29]. In humans, genetic variations in the *UNC13A* gene are associated with a neurodevelopmental syndrome[30], and non-coding intronic variants are amongst the strongest genetic risk factors for the neurodegenerative conditions amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD)[31, 32].

Alongside changes in its function, the arrangement of Munc13-1 molecules at the active zone has been deemed critical for shaping synaptic transmission properties. In *D. melanogaster* neuromuscular junction synapses, the Munc13-1 ortholog Unc13a forms rings ~70 nm around voltage-gated Ca^{2+} channels, contributing to the strong temporal coupling of synaptic transmission and Ca^{2+} influx triggered by an action potential[33, 34]. Upon synapse silencing, Unc13a expression is altered, with results pointing to an increase in the expression levels of Unc13a[35] and/or compaction of the already-available Unc13a[36], both associated with a homeostatic increase in synaptic strength. In mammalian synapses, Munc13-1 is arranged in nanoclusters and the number of nanoclusters is positively correlated with the strength of glutamate release[37]. CryoEM analysis and models based on the crystal structure of Munc13-1 promote the view that Munc13-1 forms hexameric rings surrounding one synaptic vesicle, acting in a cooperative manner to drive fusion[38, 39]. These rings have not been resolved in synapses yet, and, in general, tools are still lacking to visualize dynamic changes of Munc13-1 nanoarchitecture in mammalian synapses.

Here, we present a novel CRISPR/Cas9 knock-in mouse line, where we inserted a SNAP tag cassette at the endogenous *Unc13a* gene sequence to generate a Munc13-1 variant that is C-terminally fused to a SNAP tag. We validate the Unc13a[SNAP] mouse line, and interrogate neurons via confocal, stimulated emission-depletion (STED) microscopy, and live cell imaging. Due to the complex nature of cultured neurons and brain tissue used, and because Munc13-1 is a protein with a moderate to low expression level[40], we evaluated the performance of several SNAP tag substrates for efficient labeling. We tested the bright and stable far-red SNAP dyes BG-JF$_{646}$[41] and BG-SiR-d12[42], and, in addition, their membrane impermeable variants, SBG-JF$_{646}$[10] and SBG-SiR-d12, the synthesis and characterization of which we present here. Munc13-1 is a cytosolic protein and should not be labeled by membrane impermeable dyes, but given that some of our experiments were dealing with fixed and permeabilized preparations, we hypothesized that the charge originally installed to prevent membrane permeability may be useful in reducing background levels by repulsion. We report here that SBG-SiR-d12 has superior performance in labeling fixed cultured neurons and brain slices in terms of brightness and specificity, and conclude that repurposing membrane-impermeable dyes for staining in permeabilized preparations could be a promising approach to enhance labeling quality in complex samples, in cases where membrane permeability is irrelevant. We also conclude that the novel Unc13a[SNAP] mouse line is useful for detection of synapses and active zones at multiple scales, and thus may be useful to characterize their rapid dynamics in live and fixed samples, *in vivo* or in cultured neurons, for monitoring the reorganization of Munc13-1 in response to plasticity-inducing stimuli, and for live-imaging of Munc13-1 nanoclusters.

RESULTS

Generation and validation of an Unc13a-SNAP knock-in mouse line

We opted to generate a mouse model where endogenous Munc13-1 is C-terminally fused with a SNAP tag, with the two protein modules separated by a short and flexible, 9 amino acid long linker (sequence: (GGS)$_3$; Figure 1A). Supporting our design strategy, we relied on a previously-generated knock-in mouse line where an enhanced yellow fluorescent protein was added C-terminally to the Unc13a sequence[43]. A structural model of Munc13-1-SNAP generated by the recently-developed AlphaFold3[44] (Figure 1B) predicts three important features, i.e. 1) the addition of the SNAP tag likely does not change the Munc13-1 structure, 2) the SNAP tag likely does not exhibit protein-protein interactions with Munc13-1, and, importantly, 3) the C-terminal C2C domain likely remains structured and accessible to protein-protein or protein-lipid interactions, both of which are critical for Munc13-1 function in SV priming and thus for setting the strength of neurotransmission[45-47].

The Unc13a[WT] allele (Figure 1A) was targeted by site-directed CRISPR-Cas9 mutagenesis. The correct integration of the SNAP tag sequence was validated by long-range location PCRs (see Materials and Methods and the Supporting Information section for more information). In subsequent PCR analysis in genomic DNA from wild-type (WT), heterozygous (Unc13a[WT/SNAP]), and homozygous (Unc13a[SNAP/SNAP]) knock-in mice, we were able to amplify a DNA fragment spanning the last *Unc13a* exon and the SNAP tag cassette sequence (Unc13a[SNAP] allele), confirming correct integration and enabling routine mouse genotyping (Figure 1C and Supporting Information section).

The resulting homozygous Unc13a[SNAP] mice (Unc13a[em1(SNAP)Bros]) were viable and fertile, and exhibited no observable changes in survival, breeding performance, or cage behavior. Monitoring the well-being of the mice up to 8 months of age, we did not observe burden inflicted by the genetic modification. Because Munc13-1 loss results in perinatal lethality shortly after birth[23], we conclude that the Munc13-1-SNAP protein fusion is functional.

Figure 1: Validation of the Unc13a^{SNAP} mouse line. (A) Scheme of the 3' region of the *Unc13a* gene in WT and in targeted, Unc13a^{SNAP} mice, including the DNA sequences at the site of SNAP tag insertion. Black arrows illustrate the location of the genotyping oligonucleotides, a black line illustrates the PCR fragment. The linker, stop codon, and genotyping oligonucleotides are not drawn to scale. **(B)** An illustration of the Munc13-1 domain structure in the Unc13a^{SNAP} mice (up) and AlphaFold3 structural prediction of the Munc13-1-SNAP protein (down). **(C)** Genotyping PCR results for the indicated genotypes. **(D)** A Western blot analysis of Munc13-1 (upper blot) and the SNAP tag (lower blot) in brain synaptosome homogenates from WT, heterozygous and homozygous Unc13a^{SNAP} mice. **(E)** Quantification of Munc13-1 levels from samples as in **(D)**. **(F)** A Western blot analysis of RIM1, Syntaxin 1 and Synapsin 1 in brain homogenates from WT, heterozygous and homozygous Unc13a^{SNAP} mice.

Next, we evaluated the expression levels of Munc13-1 by Western blot analysis of a crude synaptosomal fraction (P2) from WT, heterozygous, and homozygous mouse brains. We found a non-significant, mild reduction in the expression levels of Munc13-1-SNAP in comparison to the Munc13-1 WT protein (Figure 1D, 1E). Interestingly, in samples from heterozygous mice the expression of the WT Munc13-1 protein appeared higher than that of the tagged Munc13-1 (Figure 1D, middle lane). The expression of the SNAP tag cassette was confirmed in samples from heterozygous and homozygous Unc13a^{SNAP} mice, but not in the corresponding WT littermates (Figure 1D). We did not find evidence for the emergence of a truncated Munc13-1 protein in samples from heterozygous and homozygous Unc13a^{SNAP} mice, indicating the stability of the tagged protein variant (Supporting Information, Figure S1). Finally, we found no change in the expression level of the major Munc13-1 interacting proteins syntaxin 1A/B and RIM1 (Regulating synaptic membrane exocytosis protein 1), as well as in the levels of Synapsin 1 in synaptosomal fractions from Unc13a$^{SNAP/SNAP}$ mouse brains and WT littermate samples (Figure 1F). We conclude that the genetic integration of a SNAP tag at the C-terminus of Munc13-1 does not lead to overt changes in synaptic composition.

Next, we performed immunocytochemical analysis in hippocampal neuronal cultures from littermate WT and Unc13a$^{SNAP/SNAP}$ mouse brains (Figure 2A, 2B), and immunolabeled the neurons with 1) an antibody to label the Munc13-1 protein, 2) an antibody against Bassoon, a presynaptic active zone marker, and 3) an antibody against MAP2 to label the dendritic extensions of the neuron. We quantified the expression levels of Munc13-1 in WT and in Unc13a$^{SNAP/SNAP}$ synapses based on the Munc13-1 antibody signal intensity (Figure 2C), and found no changes between samples. We also quantified the intensity of the signal arising from the labeling of Bassoon, and found no differences (Figure 2D), indicating that the expression of Munc13-1-SNAP does not change Bassoon expression. To evaluate whether Munc13-1-SNAP is correctly localized in the AZ, we determined the colocalization coefficient between the Bassoon and Munc13-1 signals (Figure 2E). We found a high degree of colocalization between the two signals, with a minor but significant reduction in the degree of signal colocalization in samples from Unc13a$^{SNAP/SNAP}$ synapses. However, since the correlation coefficient values remain relatively similar (WT, 0.55 ± 0.003, n=5284 synapses; Unc13a$^{SNAP/SNAP}$, 0.52 ± 0.0025, n=7186 synapses) we conclude that Munc13-1-SNAP is comparably-expressed and largely localized properly in the AZ.

Figure 2: Validation of Munc13-1 localization in hippocampal neurons from Unc13a$^{SNAP/SNAP}$ mice. Example images of cultured hippocampal neurons from WT **(A)** and Unc13a$^{SNAP/SNAP}$ mouse brains **(B)**, immunolabeled with antibodies against Munc13-1, the active zone marker Bassoon, and MAP2 (Scale bar: 20 μm). **(C)** Quantification of the fluorescence signal arising from antibody labeling of Munc13-1 and **(D)** of the fluorescence signal arising from antibody labeling of Bassoon in neurons from WT and Unc13a$^{SNAP/SNAP}$ mice from images as in A and B. **(E)** The colocalization of Munc13-1 and Bassoon signals evaluated by calculating the Pearson's correlation coefficient.

Labeling of Munc13-1-SNAP in fixed samples using BG-dye compounds, and the development of SBG-SiR-d12

Following the validation of the knock-in mouse line, we next tested the efficiency of Munc13-1 labeling *via* the SNAP tag in fixed and cultured primary neurons. With the aim of performing super-resolution STED imaging in mind, we opted for bright, stable and far-red dyes. Silicon Rhodamine (SiR), and the next generation fluorophores JF_{646} and SiR-d12, were developed to exhibit boosted brightness without loss of resolution[41, 42]. Conjugated to BG (i.e. $BG-JF_{646}$ and BG-SiR-d12), these dyes can be used for SNAP tag labeling. To improve the specificity of staining, sulfonated BG (SBG) substrates have been created[10] and benchmarked in complex tissue[48]. The SBG moiety was originally designed for extracellular protein labeling, as it renders the conjugates membrane-impermeable. Thus, SBG-dye conjugates are not anticipated to label Munc13-1-SNAP, which is a cytosolic protein. Nonetheless, with the knowledge that the SBG moiety renders the conjugate less lipophilic, thus improving water solubility over prolonged periods of time[10], we hypothesized that the charged SBG-dye conjugate will reduce background levels through surface repulsion and generate less non-specific deposits. Cytosolic proteins could still be labeled, if the sample has been fixed and permeabilized prior to the SNAP dye application. In addition to the already described $SBG-JF_{646}$, we report here the synthesis and validation of a novel membrane impermeable SNAP dye, SBG-SiR-d12 (see methods and Supporting Information section). We were able to obtain and validate SBG-SiR-d12 against a cohesive palette of far-red dyes for SNAP tag labeling (Figure 3A). Moreover, we could show that in live HEK293 cells transfected with a construct encoding for the fusion protein SNAP-TM-HTP (extracellular SNAP, separated by a transmembrane domain from an intracellular HaloTag[49]), SBG-SiR-d12 acts as a non-membrane permeable dye (Figure S2). Intracellular pools were successfully labeled using $CA-JF_{519}$ as a control.

Next, we used primary neuronal cultures from WT and Unc13a$^{SNAP/SNAP}$ mice, and tested $BG-JF_{646}$, $SBG-JF_{646}$, BG-SiR-d12 and SBG-SiR-d12 for their efficiency in labeling Munc13-1-SNAP. We immunolabeled fixed and permeabilized neurons with antibodies against Munc13-1, Bassoon, and MAP2, and applied one of the BG-dye conjugates mentioned above (Figure 3B, 3C). Neurons from Unc13a$^{SNAP/SNAP}$ mice were efficiently-labeled by all four SNAP-targeted dyes (Figure 3C, left panel 'SNAP'). In WT, negative control neurons, we did not observe unspecific labeling, except in the case of $SBG-JF_{646}$, where substantial background was observed (Figure 3B, left panel 'SNAP'). We analyzed the signal intensity arising from the SNAP tag labeling (Figure 3D), and found that the fluorescent signal was strongest for $SBG-JF_{646}$ and SBG-SiR-d12 in knock-in synapses. Importantly, the best signal to background ratio, which we define here as the fold-change in the median value of the signal intensity between Unc13a$^{SNAP/SNAP}$ synapses and WT synapses for each SNAP dye, was superior for SBG-SiR-d12 ($BG-JF_{646}$, 3.5-fold; $SBG-JF_{646}$, 2.2-fold; BG-SiR-d12, 9.8-fold; SBG-SiR-d12, 10.1-fold). To demonstrate specificity, we assessed the colocalization between two signals arising from labeling of the same antigen: the signal for Munc13-1 arising from the SNAP tag labeling, and the signal of Munc13-1 arising from the antibody labeling. We found, as

expected, a high degree of co-localization for all dyes tested (Figure 3E). We conclude that the SNAP tag conjugated to Munc13-1 is functional and enables specific and bright labeling of Munc13-1 in fixed tissue at confocal resolution. SBG-SiR-d12 has risen as the dye of choice to label the SNAP tag in permeabilized, fixed samples.

Figure 3: Development of SBG-SiR-d12 and validation of the SNAP tag functionality in cultured, fixed hippocampal neurons. (A) Chemical structures of silicone rhodamine variants for SNAP tag labeling, including SBG-SiR-d12 developed here. (B-C) Example images of fixed cultured hippocampal neurons from WT (B) and Unc13a$^{SNAP/SNAP}$ (C) mouse brains immunolabeled with the SNAP tag compounds $BG-JF_{646}$, $SBG-JF_{646}$, BG-SiR-d12, SBG-SiR-d12, as well as with antibodies against Munc13-1, Bassoon, and MAP2 (Scale bar: 20 µm). (D) Quantification of the fluorescence signal arising from Munc13-1-SNAP labeling by $BG-JF_{646}$, $SBG-JF_{646}$, BG-SiR-d12, SBG-SiR-d12, in neurons from WT and Unc13a$^{SNAP/SNAP}$ mice. (E) Colocalization of signals arising from simultaneous Munc13-1 antibody labeling and SNAP tag labeling using the Person's correlation coefficient.

Next, we tested whether the Unc13a^{SNAP} mice can be useful in imaging presynapses in slices of fixed brain tissue, which are typically difficult to label due to a high degree of non-specific binding of antibodies and reduced antibody permeability

(Figure 4, Figure S3). Sagittal sections from Unc13a^{WT} and Unc13a$^{SNAP/SNAP}$ mice were labeled with either BG-JF$_{646}$, SBG-JF$_{646}$, BG-SiR-d12 and SBG-SiR-d12, in parallel to immuno-labeling with an antibody against the presynaptic marker synaptophysin, and a nuclear stain (DAPI). We obtained excellent labeling with SBG-SiR-d12 in regions of high synapse density and where Munc13-1 has been shown to be enriched in a Munc13-1-YFP knock-in mouse line[43], for example in the cortex, in the hippocampal *Stratum Orients* and *Stratum Radium*, and in the molecular layer of the cerebellum formation. This signal colocalized well with antibody-stained synaptophysin signal, and little background was evident in WT control sections. The signal to background ratio of intensity measured between slices obtained from Unc13a$^{SNAP/SNAP}$ and WT mice was 2.4 for SBG-SiR-d12, and lower for other dyes tested (BG-JF$_{646}$, 1.8-fold; SBG-JF$_{646}$, 1.3-fold; BG-SiR-d12, 0.96-fold; Figure S3). We conclude that fixed tissue from Unc13a^{SNAP} mouse brains can be used for the rapid labeling of presynaptic terminals in cultured neurons and in brain tissue.

Figure 4: Labeling of presynaptic terminal in brain sections of Unc13a$^{SNAP/SNAP}$ mice. Sagittal brain sections from WT (**A**) and Unc13a$^{SNAP/SNAP}$ (**B**) mice were stained with 1 µM SBG-SiR-d12, an antibody against synaptophysin (green; to stain synapses), and with DAPI (cyan), to stain cell nuclei. Example images in the region of the hippocampus (middle) and of the cerebellum (down).

Super-resolution imaging of Munc13-1-SNAP

Next, we were interested to detect Munc13-1-SNAP by super-resolution imaging, to enable the localization of Munc13-1 beyond the diffraction limit[8], which is essential for monitoring presynaptic nanoarchitecture. We used stimulated emission depletion (STED) nanoscopy and imaged synapses from WT and Unc13a$^{SNAP/SNAP}$ hippocampal neurons in culture, stained with antibodies against MAP2 (dendrites), Munc13-1, Bassoon, and one of the SNAP-targeting dyes described above. Example images of three synapses per compound are presented in Figure 5A and 5B. Using antibody staining (Munc13-1 antibody), we identified Munc13-1 puncta that were localized on one edge of the active zone protein Bassoon, and were in part concentrated in nanoclusters. Imaging of Munc13-1 via the SNAP tag resulted in weaker signals compared to the antibody staining, which is expected due to the lack of signal amplification by primary and secondary antibodies. The SNAP tag signals, however, were well-colocalized with the signals arising from the Munc13-1 antibody labeling (Figure 5D). As in confocal

imaging, we found a high degree of background when labeling WT samples with SBG-JF$_{646}$, and the best signal to background ratio when using SBG-SiR-d12 (Figure 5C; BG-JF$_{646}$, 4.6-fold; SBG-JF$_{646}$, 2.3-fold; BG-SiR-d12, 7.4-fold; SBG-SiR-d12, 9.1-fold). In summary, these data highlight that endogenous Munc13-1 labeled via the SNAP tag can be imaged at super-resolution, with a lower signal intensity in comparison to antibody staining, but with a minimal fluorophore-protein distance.

Figure 5: STED microscopy of Munc13-1-SNAP. (A) Example images of synapses from WT and **(B)** Unc13a$^{SNAP/SNAP}$ neurons labeled with the SNAP tag compounds BG-JF$_{646}$, SBG-JF$_{646}$, BG-SiR-d12, SBG-SiR-d12, as well as with antibodies against Munc13-1 and Bassoon, and imaged using STED microscopy. Three synapses are illustrated per condition (Scale bar: 0.5 µm) **(C)** Quantification of the fluorescence signal arising from Munc13-1-SNAP labeling by BG-JF$_{646}$, SBG-JF$_{646}$, BG-SiR-d12, SBG-SiR-d12, in neurons from WT and Unc13a$^{SNAP/SNAP}$ mice. **(D)** Colocalization of signals arising from Munc13-1 antibody labeling and SNAP tag labeling using the Pearson's correlation coefficient.

Live labeling and imaging of Munc13-1-SNAP

Finally, we tested whether we could utilize the Unc13a^{SNAP} mice to monitor Munc13-1 using live imaging in neuronal cultures. We incubated WT and Unc13a$^{SNAP/SNAP}$ neurons with 0.5 µM BG-SiR-d12 for 2 h, briefly washed the neurons, and immediately subjected them to live cell imaging on a Stimulated Emission Depletion (STED) microscope with 640 nm Excitation and 775 nm depletion at 37 °C. In neuronal processes of WT neurons, we observed a significant level of background staining. Nonetheless, bright and discrete puncta were readily visible along processes of Unc13a$^{SNAP/SNAP}$ neurons, which is reminiscent to synaptic labeling (Figure 6). We conclude that

Munc13-1-SNAP labeling and imaging is possible in living neurons.

Figure 6: Live imaging of cultured neurons from Unc13a$^{SNAP/SNAP}$ mice. Live imaging of neuronal processes following incubation with 0.5 μM BG-SiR-d12 for 120 minutes. After staining, the coverslips were imaged using a Nikon Eclipse TI research microscope with a STEDYCON system. Four example images are shown for WT neurons (left) and Unc13a$^{SNAP/SNAP}$ neurons (right).

DISCUSSION

Changes in synaptic function have been repeatedly linked to changes in the organization and composition of the neurotransmitter release machinery. At the presynaptic compartment, the proximity (coupling) of release sites to voltage-gated Ca^{2+} channels determine the strength and probability of neurotransmitter release and can be dynamically modulated following plasticity-inducing triggers[50-53]. Plasticity also evokes changes in the size of the active zone[54], and in the abundance of key synaptic proteins[33, 35, 55, 56]. Munc13-1 is a central presynaptic protein that is absolutely essential for neurotransmitter release, and changes in Munc13-1 nanoarchitecture are emerging as critical for setting neurotransmitter release properties[37, 57]. However, tools are still lacking to monitor such nanoarchitectural changes for endogenous Munc13-1 *in vivo* and *in vitro*.

We used CRISPR-Cas9 to knock-in the SNAP tag cassette in the last exon and prior to the stop codon of the *Unc13a* gene. The successful targeting in mouse embryos enabled us to establish a mouse line constitutively expressing Munc13-1-SNAP at endogenous expression levels. In hippocampal neuronal cultures of this mouse, Munc13-1-SNAP is expressed at WT levels, localizes well to the active zone, and can be labeled via SNAP tag ligands to enable confocal and STED imaging in fixed neuronal cultures. We also demonstrate that labeling can be achieved in brain slices, and in living neurons. With these, we validate the Unc13a^{SNAP} mice for future studies of synapse and active zone nanoarchitecture. This mouse line complements already-available tools for synapse nanoscopy, in particular a PSD95-HaloTag line that has proven as highly useful for analyzing postsynaptic density at super-resolution[58], but also for imaging synapses in brain slices[58].

Key advantages of self-labeling tags include great flexibility in dye selection and a large and consistently growing toolbox of chemicals available. Importantly, reagents do not only differ in the wavelength of the fluorophore attached to them, but also in their chemical and biological properties, e.g. they undergo uptake to different extents, and have variable solubilities. Nonetheless, a large fraction of reported investigations using self-labeling tags rely on the application of one derivative only. We show here by testing several BG derivatives simultaneously that the performance of different BG-fluorophore compounds is assay-dependent and needs to be carefully controlled[2, 59]. To enhance the labeling specificity of our samples, we chose to synthesize SBG-SiR-d12. While a priori, using a sulfonated and impermeable dye seems counterintuitive, we reasoned that enhanced solubility is beneficial in the context of fixed and permeabilized samples. Indeed, SBG-SiR-d12 outperformed all other conjugates in signal strength, and had the highest signal to background ratio in fixed cell (10-fold) and in complex brain tissue samples (2.4-fold). SBG-SiR-d12 leverages an outstanding performance for instance, in comparison to our recently developed probe Alexa488-C3 to image beta-secretase (signal to background 1.7-fold[60]). Its enhanced performance with respect to JF dyes could be attributed to the higher fluorogenicity and a resulting higher degree of non-fluorescent spirolactones of JF dyes in fixed preparations. We propose SNAP labeling in fixed cell cultures and slices with SBG-SiR-d12 as an alternative to membrane-permeable SNAP dyes. The protocol presented here joins protocols for SNAP tag labeling in the live setting, e.g. in immortalized cell lines[2], *Drosophila* brain[61, 62], in mice for neural identification and ablation[63], for receptor localization including optical manipulation[64], and in pancreatic islets[65].

Undoubtedly, further development of imaging configurations and SNAP tag reagents, as well as testing of additional labeling protocols will expand the range of experimentation possible using the Unc13a^{SNAP} mice. Because it is directly fused to Munc13-1, the SNAP tag offers the unique advantage that it carries the dye in close proximity to the molecule, reducing to minimum the aberrations, i.e. the distances between fluorophores and the actual positions of the biological molecule. This is immensely important for precise protein localization as the resolution of imaging is improving to the single nanometer scale[66-69], making the POI-fluorophore distance generated by primary and secondary antibody labeling unacceptable. The genetic fusion via CRISPR/Cas9 ensures the visualization of Munc13-1 at endogenous expression patterns and levels, independent of constrains related to antibody species, and in the context of neuronal networks in either fixed or live preparations. These key features will surely make the Unc13a^{SNAP} mice useful in the discovery of synaptic nanoarchitectural principles in the future.

MATERIALS AND METHODS

General. BG-JF$_{549}$, SBG-JF$_{549}$, and BG-SiR-d12 have been synthesized and used previously in our laboratories[10, 42].

Chemical Synthesis. SBG-SiR-d12 was synthesized as follows: In an Eppendorf tube 1.2 mg (2.0 μmol, 1.0 equiv.) of SiR-d12-COOH[42] was dissolved in 100 μL/mg DMF and 8.0 equiv. of DIPEA. Upon addition of 2.3 equiv. TSTU (from a 41 mg/mL stock in DMSO) the reaction mixture was vortexed and allowed to incubate for 10 min, before 1.2 equiv. of SBG-NH$_2$ was added. The mixture was vortexed again and allowed to

incubate for 60 min before it was quenched by addition of 20 equiv. of acetic acid and 25 vol% of water. C18 RP-HPLC (MeCN:H$_2$O+0.1% TFA = 10:90 to 90:10 over 45 minutes) provided the desired compound, SBG-SiR-d12, which was obtained as a blue powder after lyophilization in 42% yield (0.84 μmol) and aliquoted to 5 nmol to be stored at –20 °C.

^{1}H NMR (600 MHz, MeOD-d$_4$): δ [ppm] = 9.29–9.21 (m, 1H), 8.23–8.16 (m, 1H), 8.12–8.08 (m, 1H), 7.71 (s, 1H), 7.50 (d, J = 7.9 Hz, 2H), 7.39 (d, J = 7.9 Hz, 2H), 7.20 (s, 2H), 6.86 (d, J = 9.1 Hz, 2H), 6.70 (dd, J = 9.3, 3.0 Hz, 2H), 5.59 (s, 2H), 4.59–4.55 (m, 2H), 3.54 (t, J = 6.5 Hz, 2H), 3.45–3.41 (m, 1H), 3.20–3.18 (m, 1H), 3.12 (t, J = 7.0 Hz, 2H), 2.90 (t, J = 6.5 Hz, 2H), 2.70 (t, J = 7.0 Hz, 2H), 0.63 (s, 3H), 0.57 (s, 3H).

HRMS (ESI): calc. for C$_{45}$H$_{38}$D$_{12}$N$_9$O$_8$SSi [M+H]$^+$: 916.4020, found: 916.4165.

Animal study approval. The generation and use of the Unc13a-SNAP (Unc13a$^{em1(SNAP)Bros}$) knockin mice were approved by the responsible local government organization (Niedersächsisches Landesamt für Verbraucherschutz und Lebensmittelsicherheit; 33.9-42502-04-13/1359 and 33.19- 33.19-42502-04-20/3589). All the experiments performed in Berlin complied with European law and the state of Berlin animal welfare body (LAGeSo).

Generation of an Unc13a-SNAP knock-in mouse line. Superovulated C57BL6/J females were mated with C57BL6/J males and fertilized eggs collected. In-house prepared CRISPR reagents (hCas9_mRNA, sgRNAs, preformed Cas9_sgRNA RNP complexes, and the dsDNA used as a repair template (HDR fragment), were microinjected into the pronucleus and the cytoplasm of zygotes at the pronuclear stage using an Eppendorf Femtojet. Importantly, all nucleotide-based CRISPR-Cas9 reagents (sgRNAs and hCAS9_mRNA) were used as RNA molecules and were not plasmid-encoded, reducing the probability of off-target effects, due to the short live of RNA-based reagents[70, 71]. The sgRNAs targeting the region around the Munc13-1 STOP-codon were selected using the guide RNA selection tool CRISPOR[72, 73]. The correct site-specific insertion of the HDR fragment was confirmed by two localization PCR with primers upstream and downstream of the HDR sequence, followed by direct sequencing of the obtained PCR products. The sequences of the RNA and DNA fragments used to generate and validate the mice are available in the Supporting Information section. We term the line as Unc13a$^{em1(SNAP)Bros}$. For questions about usage of the line, please contact Nils Brose and Noa Lipstein.

Synaptosome preparation and Western blot analysis. A crude synaptosomal fraction (P2) was prepared as described in[74] from mice age 7-11 weeks. The sample was solubilized to a final concentration of 2 mg/ml in 50 mM Tris/HCl pH 8, 150 mM NaCl, 1 mM CaCl$_2$, 1 mM EGTA, 1% NP-40, 0.2 mM phenylmethylsulfonyl fluoride, 1 mg/ml aprotinin, and 0.5 mg/ml leupeptin, stirred on ice for at least 15 min at 4 °C, centrifuged at 4 °C, 20,000g for 5 minutes to remove insoluble material, mixed with Laemmli sample buffer and boiled for 10 minutes at 99 °C. 5 μg of each sample were loaded on a 4-12% Bis-Tris gel, and the proteins were separated and transferred to a nitrocellulose membrane. The membrane was washed three times with ddH$_2$O, stained with Pierce™ Reversible Protein Stain Kit

for Nitrocellulose Membranes (Thermo Scientific 24580) and scanned. Then, the membrane was blocked for 60 minutes with blocking buffer (5% (*w/v*) low fat milk in Tris buffered Saline (TBS) buffer supplemented with 0.1% Tween20 (TBS-T)), and blotted with the following antibodies diluted in blocking buffer for 2 hours at RT with gentle shaking: Rabbit anti Munc13-1 (Synaptic Systems 126 103, diluted 1:1000), Rabbit anti SNAP (New England Biolabs, P9310S, diluted 1:500), Mouse anti synapsin 1 (Synaptic Systems 106 011, diluted 1:1000), Mouse anti Syntaxin 1A/B (Synaptic Systems 110 011, diluted 1:2000), Rabbit anti Rim1 (Synaptic Systems 140 003, diluted 1:1000). The membrane was washed three times for 10 minutes each with TBS-T before blotted for 1 hour at RT with secondary antibodies conjugated with horseradish peroxidase (HRP) diluted in blocking buffer: Goat-anti-Rabbit-HRP (Jackson immune 111-035-114, diluted 1:5000) and Goat-anti-Mouse-HRP (Jackson immuno 115-035-146, diluted 1:5000). Before developing the chemiluminescence signal using the standard HRP signal amplification system, the membrane was washed three times for 10 minutes each in TBST buffer and once shortly in TBS buffer.

HEK293T cultures. HEK293T cells were cultured in growth medium (DMEM, Glutamax, 4.5 g Glucose, 10% FCS, 1% PS; Invitrogen) at 37 °C and 5% CO$_2$. 30,000 cells per well were seeded on 8-well μL slides (Ibidi) previously coated with 0.25 mg ml^{-1} poly-L-lysine (Aldrich, mol wt 70 000–150 000). The next day, 400 ng DNA was transfected using 0.8 μL Jet Prime reagent in 40 μL Jet Prime buffer (VWR) per well. Medium was exchanged against antibiotic-free media before the transfection mix was pipetted on the cells. After 4 hours incubation at 37 °C and 5% CO$_2$, medium was exchanged against growth media. After 24 hours cells were stained and imaged. All dyes were used at a concentration of 500 nM. 5 μM Hoechst 33342 was used to stain DNA. Staining was done in growth medium at 37 °C, 5% CO$_2$ for 30 minutes. Afterwards cells were washed once in growth media and imaged live in cell imaging buffer (Invitrogen).

Neuronal cultures. Primary hippocampal neurons were cultured as previously described [75]. Briefly, hippocampi were dissected from Unc13a$^{SNAP/SNAP}$, Unc13a$^{WT/SNAP}$, and WT/WT P0-P1 littermate mice, and were incubated at 37 °C gently shaking in a solution containing 0.2 mg/ml L-cysteine, 1 mM CaCl$_2$, 0.5 mM ethylenediaminetetraacetic acid (EDTA) and 25 units/ml of papain (Worthington Biochemicals), pH 8 in Dulbecco's modified eagle medium. After 45 minutes, the solution was replaced by a prewarmed solution containing 2.5 mg/ml Bovine Serum Albumin, 2.5 mg/ml trypsin inhibitor, 1% Fetal Bovine Serum, heat inactivated in Dulbecco's modified eagle medium, and the hippocampi were incubated for 15 minutes at 37 °C gently shaking. The solution was replaced by neuronal culture medium (Neurobasal™-A medium supplemented with 2% B-27™ Plus Supplement, 1% GlutaMAX™ supplement, and 0.2% Penicillin-Streptomycin), and the hippocampi were gently triturated to produce a cell suspension that was plated on PLL-coated coverslips for 2-3 weeks at 37 °C and 5% CO$_2$. The culture contains primarily neurons and few astrocytes. Neurons were used for experiments at day-in-vitro 14-17.

Immunocytochemical staining protocols. Experiments were made in neuronal cultures at days-in-vitro 14-17. Neurons

were washed twice with PBS, and fixed by adding cold 1% paraformaldehyde for 10 minutes on ice. Neurons were again washed and leftover paraformaldehyde was quenched by adding 50 mM cold glycine for 10 minutes, followed by two additional washes with PBS. The cells were either stored in PBS at 4 °C or immediately used for immunostaining. The cell membrane was permeabilized with cold 0.25% Triton-X-100 in PBS for 10 minutes and rinsed once in pure PBS. To block nonspecific binding sites, the cells were incubated in cold 0.3% NGS in PBS for 20 minutes. All steps were performed with the multi-well plate stored on ice to prevent protein degradation.

To label Munc13-1 using the SNAP tag, 250 nmol BG-dye conjugates were diluted in blocking solution and applied to the neurons for 30 minutes at room temperature (RT) in the dark. Subsequently, the cells were extensively washed up to eight times in blocking solution and the following antibodies were applied in blocking buffer for immunolabeling: Chicken polyclonal MAP2 (Novus Biologicals NB300-213, diluted 1:1000), Rabbit polyclonal Munc13-1 (Synaptic Systems AB_126103, diluted 1:500), and Guinea Pig polyclonal Bassoon (Synaptic Systems AB_141004, diluted 1:500). Incubation was performed for 1 h at RT in the dark. To remove unbound antibodies, the cultures were washed three times with blocking solution before applying the following secondary antibodies conjugated with fluorophores: Anti-Chicken-405, Anti-Guinea Pig-488 and Anti-Rabbit-594 (all diluted 1:500). The neurons were incubated with the secondary antibodies for 30 minutes at RT in the dark and washed eight times in PBS to remove unbound products. The coverslips on which the neurons were fixed were mounted with ProLong™ Gold Antifade Mountant (Invitrogen™ P36934) on microscope glass slides and stored in the dark at RT for 48 hours until dry before imaging.

Brain sections and immunohistochemistry. Animals were transcardially perfused with fixative (4% paraformaldehyde in 1X PBS, pH 7.4). Brains were post-fixed overnight at 4 °C, incubated in 30% sucrose in 1X PBS for a minimum of 48 h, and snap frozen with 2-methybutan on dry ice. Tissues were sectioned at 30 μm on a cryostat. Floating sections were kept in 30% ethylene glycol and 30% glycerol in 1X PBS at -20 °C. For immunostaining, sections were washed three times in 1X PBS before incubation with blocking solution containing 10% normal donkey serum (NDS) in TBS-T (0.05% Triton X-100 in 1X TBS). The slices were incubated with one of the SNAP tag dyes (1 μM) and anti-synaptophysin1 (Synaptic Systems AB_101002, diluted 1:200,) at 4 °C overnight in TBS-T containing 2% NDS. Sections were washed three times in 1X TBS-T, followed by incubation for 2 h at room temperature in the dark with secondary antibody (Alexa Fluor 488-AffiniPure Donkey Anti-Rabbit IgG, Jackson Research 711-545-152, diluted 1:1000,) and DAPI (Sigma MBD0015-1ML, diluted 1:10000). The sections were washed three times in 1X TBS-T before mounted onto microscope slides with ProLong Glass antifade (Invitrogen™P36982).

Confocal and STED Imaging. Confocal and STED microscopy on fixed neuronal samples was performed using a Leica SP8 TCS STED FALCON (Leica Microsystems). Confocal images were collected using a time gated Hybrid detector (0.5–6 ns). Images of 1024x1024 pixel had a pixel size of 130 nm for confocal imaging and 20 nm for STED imaging. For imaging far red SNAP dyes, we used the following settings: excitation wavelength 646 nm, emission wavelength 656–751 nm. Brain sections were imaged on a Nikon Spinning Disk Field Scanning Confocal System using 10X objective (Plan Apo λD, NA 0.45), acquired in the "Large Image" mode of NIS 5.4 Software and stitched with 20% overlap between adjacent frames. For imaging DAPI, Synaptophysin and SNAP dyes 405, 488 and 640 nm excitation lasers and 440/40, 525/50, 708/75 emission filters were used, respectively. CA-JF$_{519}$ and SBG-SiR-d12 in live transfected HEK293Ts cells was imaged on a TIE Nikon epifluorescence microscope equipped with a pE4000 (cool LED), Penta Cube (AHF 66–615), 60 oil NA 1.49 (Apo TIRF Nikon) and a sCMOS camera (Prime 95B, Photometrics) operated by NIS Elements (Nikon). For excitation the following LED wavelengths were used: Hoechst: 405 nm; JF$_{519}$: 480 nm; SiR-d12: 635 nm.

Live Imaging. Live confocal imaging was done on a STEDYCON system (Abberior Instruments GmbH, Göttingen), mounted on a Nikon Eclipse TI research microscope, equipped with a Plan APO Lambda 100X/1.45 NA oil objective (Nikon) and controlled by NIS Elements (Nikon). 24 h prior to live-cell imaging, an incubation chamber surrounding the microscope set-up was set to 37 °C. In order to provide stable focus during imaging the Perfect Focus System (Nikon) was used. Imaging was performed in a medium containing 140 mM NaCl, 2.4 mM KCl, 10 mM HEPES, 10 mM glucose, 2 mM CaCl$_2$, and 2 mM MgCl$_2$ (320 mOsm/l) at 37 °C. Excitation was done with a 640 nm diode laser and Emission was detected with a single counting avalanche photodiode (650-700 nm). A pixel size of 20 nm by 20 nm and a line accumulation of 2 was used.

Image data analysis. Confocal and STED images (Figures 3 and 5) were analyzed using routines generated in Matlab (The Mathworks Inc., Natick MA, USA; version R2022b). The Bassoon images were subjected to an automated thresholding procedure, to identify signals above background (using a threshold equal to the mean intensity value plus the standard deviation, calculated for the entire image). The resulting regions of interest (ROIs) were processed by an erosion procedure, using a kernel of 4 pixels, to remove noise events, and only retain synapse-like signals. The remaining ROIs were then automatically dilated, to include the surrounding synaptic areas, and the average intensity values were measured and reported, for all channels. At the same time, Pearson's correlation coefficients (across all pixels in a ROI) were determined, for each ROI, between the relevant channels (SNAP tag labeling, Munc13 antibody immunostaining).

Statistics. Statistical analysis was conducted in Prism10. Experimental groups were compared against each other using the Kruskal–Wallis test, and the Dunn's test for multiple comparisons was used to perform pairwise comparisons between each independent group.

Data Availability. The data that support the findings of this study are available upon request from the corresponding author.

ASSOCIATED CONTENT

Supporting Information

Supplemental Figures 1-3, Generation of an Unc13a^{Snap} knock-in mouse mutant using CRISPR/Cas9 gene editing, Location PCR and genotyping, general chemistry, SBG-SiR-d12 analysis (PDF)

AUTHOR INFORMATION

Corresponding Authors

Johannes Broichhagen − *Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Berlin 13125, Germany; orchid.org/0000-0003-3084-6595; Email:* broichhagen@fmp-berlin.de

Noa Lipstein − *Max Planck Institute for Multidisciplinary Sciences, Göttingen 37077, Germany; Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Berlin 13125, Germany; orchid.org/0000-0003-3084-6595; Email:* lipstein@fmp-berlin.de

Authors

Maria Kowald − *Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Berlin 13125, Germany; orchid.org/0009-0001-4769-4834; orchid.org/0009-0001-4769-4834*

Sylvestre P. J. T. Bachollet − *Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Berlin 13125, Germany; orchid.org/0000-0002-1249-5296*

Fritz Benseler − *Max Planck Institute for Multidisciplinary Sciences, Göttingen 37077, Germany; orchid.org/0009-0006-3846-9447*

Maria Steinecker − *Max Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), Berlin 13125, Germany*

Sofia Kaushik − *Department of Pharmacology and Cell Biology, Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Berlin 13125, Germany; orcid.org/0000-0002-5757-764X*

Tolga Soykan − *Department of Pharmacology and Cell Biology, Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Berlin 13125, Germany; orcid.org/*0000-0002-1324-9601

Siqi Sun − *Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Berlin 13125, Germany; orchid.org/0009-0002-8291-3358*

Ramona Birke − *Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Berlin 13125, Germany*

Dragana Ilic − *Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Berlin 13125, Germany; orchid.org/0000-0002-3725-7964*

Nils Brose − *Max Planck Institute for Multidisciplinary Sciences, Göttingen 37077, Germany*

Hanna Hörnberg − *Max Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), Berlin 13125, Germany; orcid.org/*0000-0001-9142-1932

Martin Lehmann − *Department of Pharmacology and Cell Biology, Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Berlin 13125, Germany; orcid.org/0000-0002-8370-6353*

Silvio O. Rizzoli − *Department of Neuro- and Sensory Physiology, University Medical Center, Göttingen 37073, Germany; Excellence Cluster Multiscale Bioimaging (MBExC), Göttingen 37073, Germany; Center for Biostructural Imaging of Neurodegeneration (BIN), University Medical Center Göttingen, Göttingen 37073, Germany; orcid.org/0000-0002-1667-7839*

Author Contributions

Maria Kowald: Conceptualization, Formal analysis, Investigation, Methodology, visualization, Writing − Review and Editing; **Sylvestre Bachollet**: Conceptualization, Formal analysis, Investigation, Methodology; **Fritz Benseler**: Conceptualization, Formal analysis, Investigation, Methodology, Supervision, Writing − Review and Editing; **Maria Steinecker**: Investigation; **Sofia Kaushik**: Investigation, Methodology; **Tolga Soykan**: Investigation, Methodology, formal analysis; **Sun Siqi**: Formal analysis, Investigation, Writing − Review and Editing; **Ramona Birke**: Investigation; **Dragana Ilic**: Formal analysis, Investigation, Writing − Review and Editing; **Nils Brose**: Conceptualization, Funding acquisition, Supervision, Writing − Review and Editing; **Hanna Hörnberg**: Methodology, Supervision, Writing − Review and Editing; **Martin Lehmann**: Investigation, Methodology, Writing − Review and Editing; **Silvio O. Rizzoli**: Formal analysis, Funding acquisition, visualization, Writing − Review and Editing; **Johannes Broichhagen**: Conceptualization, Formal analysis, Funding acquisition, Investigation, Methodology, Supervision, visualization, Writing − Review and Editing; **Noa Lipstein**: Conceptualization, Formal analysis, Funding acquisition, Investigation, Methodology, Supervision, visualization, Writing − Original draft. All authors have given approval to the final version of the manuscript. / ‡These authors contributed equally.

Funding Sources

This work was supported by the German Research Foundation Excellence Strategy EXC-2049-390688087 (to N. Lipstein and H. Hörnberg); CRC 1286 "Quantitative Synaptology" project A11 (to N. Lipstein), project A03 (to Silvio O. Rizzoli), and A09 (to N. Brose), the TargetALS foundation (Industry-led collaborative consortia: 'Correcting Aberrant Splicing of UNC13A as a Therapeutic Approach for ALS and FTD' to N. Lipstein), and the Einstein Center for Neurosciences (to S. Sun). This project has received funding from the European Union's Horizon Europe Framework Programme (deuterON, grant agreement no. 101042046 to J. Broichhagen), and from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation, to J. Broichhagen (A Chemical Biology Approach for all-in-one cryoCLEM probes; CC RFA).

Notes

Conflict of interest: Noa Lipstein is a scientific advisory board member of TRACE Neuroscience Inc. Johannes Broichhagen receives licensing revenue from Celtarys Research, which is unrelated to this project.

ACKNOWLEDGMENT

The authors would like to thank Christiane Harenberg and Dayana Warnecke from the AGCTLab (MPI for Multidisciplinary Sciences) for expert technical assistance and genotyping. We thank the animal facility of the Max Planck Institute of Multidisciplinary Sciences City Campus for their excellent work towards generating the mouse model, the animal facility of the Max-Delbruck Center

Campus Berlin Buch for mouse husbandry, and Kerstin Steinhagen for technical assistance.

ABBREVIATIONS

Munc13-1, protein unc-13 homolog A; SV, synaptic vesicles; BG, O^6-benzylguanine; CA, chloroalkane; ALS, amyotrophic lateral sclerosis; FTD, frontotemporal dementia; SiR, silicon rhodamine; WT, wild type

REFERENCES

(1) Rodriguez, E. A.; Campbell, R. E.; Lin, J. Y.; Lin, M. Z.; Miyawaki, A.; Palmer, A. E.; Shu, X.; Zhang, J.; Tsien, R. Y. The Growing and Glowing Toolbox of Fluorescent and Photoactive Proteins. *Trends Biochem Sci* **2017**, *42* (2), 111-129. DOI: 10.1016/j.tibs.2016.09.010 From NLM Medline.

(2) Erdmann, R. S.; Baguley, S. W.; Richens, J. H.; Wissner, R. F.; Xi, Z.; Allgeyer, E. S.; Zhong, S.; Thompson, A. D.; Lowe, N.; Butler, R.; et al. Labeling Strategies Matter for Super-Resolution Microscopy: A Comparison between HaloTags and SNAP-tags. *Cell Chem Biol* **2019**, *26* (4), 584-592 e586. DOI: 10.1016/j.chembiol.2019.01.003 From NLM Medline.

(3) Zhang, D.; Chen, Z.; Du, Z.; Bao, B.; Su, N.; Chen, X.; Ge, Y.; Lin, Q.; Yang, L.; Hua, Y.; et al. Design of a palette of SNAP-tag mimics of fluorescent proteins and their use as cell reporters. *Cell Discov* **2023**, *9* (1), 56. DOI: 10.1038/s41421-023-00546-y From NLM PubMed-not-MEDLINE.

(4) Sun, X.; Zhang, A.; Baker, B.; Sun, L.; Howard, A.; Buswell, J.; Maurel, D.; Masharina, A.; Johnsson, K.; Noren, C. J.; et al. Development of SNAP-tag fluorogenic probes for wash-free fluorescence imaging. *Chembiochem* **2011**, *12* (14), 2217-2226. DOI: 10.1002/cbic.201100173 From NLM Medline.

(5) Los, G. V.; Encell, L. P.; McDougall, M. G.; Hartzell, D. D.; Karassina, N.; Zimprich, C.; Wood, M. G.; Learish, R.; Ohana, R. F.; Urh, M.; et al. HaloTag: a novel protein labeling technology for cell imaging and protein analysis. *ACS Chem Biol* **2008**, *3* (6), 373-382. DOI: 10.1021/cb800025k From NLM Medline.

(6) Gautier, A.; Juillerat, A.; Heinis, C.; Correa, I. R., Jr.; Kindermann, M.; Beaufils, F.; Johnsson, K. An engineered protein tag for multiprotein labeling in living cells. *Chem Biol* **2008**, *15* (2), 128-136. DOI: 10.1016/j.chembiol.2008.01.007 From NLM Medline.

(7) Masch, J. M.; Steffens, H.; Fischer, J.; Engelhardt, J.; Hubrich, J.; Keller-Findeisen, J.; D'Este, E.; Urban, N. T.; Grant, S. G. N.; Sahl, S. J.; et al. Robust nanoscopy of a synaptic protein in living mice by organic-fluorophore labeling. *Proc Natl Acad Sci U S A* **2018**, *115* (34), E8047-E8056. DOI: 10.1073/pnas.1807104115 From NLM Medline.

(8) Sahl, S. J.; Hell, S. W.; Jakobs, S. Fluorescence nanoscopy in cell biology. *Nat Rev Mol Cell Biol* **2017**, *18* (11), 685-701. DOI: 10.1038/nrm.2017.71 From NLM Medline.

(9) Bodor, D. L.; Rodriguez, M. G.; Moreno, N.; Jansen, L. E. Analysis of protein turnover by quantitative SNAP-based pulse-chase imaging. *Curr Protoc Cell Biol* **2012**, *Chapter 8*, Unit8 8. DOI: 10.1002/0471143030.cb0808s55 From NLM Medline.

(10) Poc, P.; Gutzeit, V. A.; Ast, J.; Lee, J.; Jones, B. J.; D'Este, E.; Mathes, B.; Lehmann, M.; Hodson, D. J.; Levitz, J.; Broichhagen, J. Interrogating surface versus intracellular transmembrane receptor populations using cell-impermeable SNAP-tag substrates. *Chem Sci* **2020**, *11* (30), 7871-7883. DOI: 10.1039/d0sc02794d From NLM PubMed-not-MEDLINE.

(11) Neukam, M.; Sala, P.; Brunner, A. D.; Ganss, K.; Palladini, A.; Grzybek, M.; Topcheva, O.; Vasiljevic, J.; Broichhagen, J.; Johnsson, K.; et al. Purification of time-resolved insulin granules reveals proteomic and lipidomic changes during granule aging. *Cell Rep* **2024**, *43* (3), 113836. DOI: 10.1016/j.celrep.2024.113836 From NLM Medline.

(12) Yim, W. W.; Yamamoto, H.; Mizushima, N. A pulse-chasable reporter processing assay for mammalian autophagic flux with Halo-Tag. *Elife* **2022**, *11*. DOI: 10.7554/eLife.78923 From NLM Medline.

(13) Frei, M. S.; Tarnawski, M.; Roberti, M. J.; Koch, B.; Hiblot, J.; Johnsson, K. Engineered HaloTag variants for fluorescence lifetime multiplexing. *Nat Methods* **2022**, *19* (1), 65-70. DOI: 10.1038/s41592-021-01341-x From NLM Medline.

(14) Kühn, S.; Nasufovic, V.; Wilhelm, J.; Kompa, J.; de Lange, E. M. F.; Lin, Y.-H.; Egoldt, C.; Fischer, J.; Lennoi, A.; Tarnawski, M.; et al. SNAP-tag2: faster and brighter protein labeling. *bioRxiv* **2024**, 2024.2008.2028.610127. DOI: 10.1101/2024.08.28.610127.

(15) Brose, N.; Hofmann, K.; Hata, Y.; Sudhof, T. C. Mammalian homologues of Caenorhabditis elegans unc-13 gene define novel family of C2-domain proteins. *J Biol Chem* **1995**, *270* (42), 25273-25280.

(16) Dittman, J. S. Unc13: a multifunctional synaptic marvel. *Curr Opin Neurobiol* **2019**, *57*, 17-25. DOI: 10.1016/j.conb.2018.12.011.

(17) Augustin, I.; Betz, A.; Herrmann, C.; Jo, T.; Brose, N. Differential expression of two novel Munc13 proteins in rat brain. *Biochem J* **1999**, *337 (Pt 3)*, 363-371.

(18) Man, K. N.; Imig, C.; Walter, A. M.; Pinheiro, P. S.; Stevens, D. R.; Rettig, J.; Sorensen, J. B.; Cooper, B. H.; Brose, N.; Wojcik, S. M. Identification of a Munc13-sensitive step in chromaffin cell large dense-core vesicle exocytosis. *Elife* **2015**, *4*. DOI: 10.7554/eLife.10635.

(19) Brunger, A. T.; Choi, U. B.; Lai, Y.; Leitz, J.; Zhou, Q. Molecular Mechanisms of Fast Neurotransmitter Release. *Annu Rev Biophys* **2018**, *47*, 469-497. DOI: 10.1146/annurev-biophys-070816-034117.

(20) Imig, C.; Min, S. W.; Krinner, S.; Arancillo, M.; Rosenmund, C.; Sudhof, T. C.; Rhee, J.; Brose, N.; Cooper, B. H. The morphological and molecular nature of synaptic vesicle priming at presynaptic active zones. *Neuron* **2014**, *84* (2), 416-431. DOI: 10.1016/j.neuron.2014.10.009.

(21) Lipstein, N.; Chang, S.; Lin, K. H.; Lopez-Murcia, F. J.; Neher, E.; Taschenberger, H.; Brose, N. Munc13-1 is a Ca(2+)-phospholipid-dependent vesicle priming hub that shapes synaptic short-term plasticity and enables sustained neurotransmission. *Neuron* **2021**, *109* (24), 3980-4000 e3987. DOI: 10.1016/j.neuron.2021.09.054 From NLM Medline.

(22) Lipstein, N.; Sakaba, T.; Cooper, B. H.; Lin, K. H.; Strenzke, N.; Ashery, U.; Rhee, J. S.; Taschenberger, H.; Neher, E.; Brose, N. Dynamic control of synaptic vesicle replenishment and short-term plasticity by Ca(2+)-calmodulin-Munc13-1 signaling. *Neuron* **2013**, *79* (1), 82-96. DOI: 10.1016/j.neuron.2013.05.011.

(23) Augustin, I.; Rosenmund, C.; Sudhof, T. C.; Brose, N. Munc13-1 is essential for fusion competence of glutamatergic synaptic vesicles. *Nature* **1999**, *400* (6743), 457-461. DOI: 10.1038/22768.

(24) Basu, J.; Betz, A.; Brose, N.; Rosenmund, C. Munc13-1 C1 domain activation lowers the energy barrier for synaptic vesicle fusion. *J Neurosci* **2007**, *27* (5), 1200-1210. DOI: 10.1523/JNEUROSCI.4908-06.2007.

(25) Junge, H. J.; Rhee, J. S.; Jahn, O.; Varoqueaux, F.; Spiess, J.; Waxham, M. N.; Rosenmund, C.; Brose, N. Calmodulin and Munc13 form a Ca2+ sensor/effector complex that controls short-term synaptic plasticity. *Cell* **2004**, *118* (3), 389-401. DOI: 10.1016/j.cell.2004.06.029.

(26) Rhee, J. S.; Betz, A.; Pyott, S.; Reim, K.; Varoqueaux, F.; Augustin, I.; Hesse, D.; Sudhof, T. C.; Takahashi, M.; Rosenmund, C.; Brose, N. Beta phorbol ester- and diacylglycerol-induced augmentation of transmitter release is mediated by Munc13s and not by PKCs. *Cell* **2002**, *108* (1), 121-133.

(27) Rosenmund, C.; Sigler, A.; Augustin, I.; Reim, K.; Brose, N.; Rhee, J. S. Differential control of vesicle priming and short-term plasticity by Munc13 isoforms. *Neuron* **2002**, *33* (3), 411-424.

(28) Shin, O. H.; Lu, J.; Rhee, J. S.; Tomchick, D. R.; Pang, Z. P.; Wojcik, S. M.; Camacho-Perez, M.; Brose, N.; Machius, M.; Rizo, J.; et al. Munc13 C2B domain is an activity-dependent Ca2+ regulator of synaptic exocytosis. *Nat Struct Mol Biol* **2010**, *17* (3), 280-288. DOI: 10.1038/nsmb.1758.

(29) Varoqueaux, F.; Sigler, A.; Rhee, J. S.; Brose, N.; Enk, C.; Reim, K.; Rosenmund, C. Total arrest of spontaneous and evoked

synaptic transmission but normal synaptogenesis in the absence of Munc13-mediated vesicle priming. *Proc Natl Acad Sci U S A* **2002**, *99* (13), 9037-9042. DOI: 10.1073/pnas.122623799.

(30) Lipstein, N.; Verhoeven-Duif, N. M.; Michelassi, F. E.; Calloway, N.; van Hasselt, P. M.; Pienkowska, K.; van Haaften, G.; van Haelst, M. M.; van Empelen, R.; Cuppen, I.; et al. Synaptic UNC13A protein variant causes increased neurotransmission and dyskinetic movement disorder. *J Clin Invest* **2017**, *127* (3), 1005-1018. DOI: 10.1172/JCI90259.

(31) Akiyama, T.; Koike, Y.; Petrucelli, L.; Gitler, A. D. Cracking the cryptic code in amyotrophic lateral sclerosis and frontotemporal dementia: Towards therapeutic targets and biomarkers. *Clin Transl Med* **2022**, *12* (5), e818. DOI: 10.1002/ctm2.818 From NLM Medline.

(32) van Rheenen, W.; van der Spek, R. A. A.; Bakker, M. K.; van Vugt, J.; Hop, P. J.; Zwamborn, R. A. J.; de Klein, N.; Westra, H. J.; Bakker, O. B.; Deelen, P.; et al. Common and rare variant association analyses in amyotrophic lateral sclerosis identify 15 risk loci with distinct genetic architectures and neuron-specific biology. *Nat Genet* **2021**, *53* (12), 1636-1648. DOI: 10.1038/s41588-021-00973-1 From NLM Medline.

(33) Bohme, M. A.; Beis, C.; Reddy-Alla, S.; Reynolds, E.; Mampell, M. M.; Grasskamp, A. T.; Lutzkendorf, J.; Bergeron, D. D.; Driller, J. H.; Babikir, H.; et al. Active zone scaffolds differentially accumulate Unc13 isoforms to tune Ca(2+) channel-vesicle coupling. *Nat Neurosci* **2016**, *19* (10), 1311-1320. DOI: 10.1038/nn.4364.

(34) Reddy-Alla, S.; Bohme, M. A.; Reynolds, E.; Beis, C.; Grasskamp, A. T.; Mampell, M. M.; Maglione, M.; Jusyte, M.; Rey, U.; Babikir, H.; et al. Stable Positioning of Unc13 Restricts Synaptic Vesicle Fusion to Defined Release Sites to Promote Synchronous Neurotransmission. *Neuron* **2017**, *95* (6), 1350-1364 e1312. DOI: 10.1016/j.neuron.2017.08.016.

(35) Bohme, M. A.; McCarthy, A. W.; Grasskamp, A. T.; Beuschel, C. B.; Goel, P.; Jusyte, M.; Laber, D.; Huang, S.; Rey, U.; Petzoldt, A. G.; et al. Rapid active zone remodeling consolidates presynaptic potentiation. *Nat Commun* **2019**, *10* (1), 1085. DOI: 10.1038/s41467-019-08977-6.

(36) Mrestani, A.; Pauli, M.; Kollmannsberger, P.; Repp, F.; Kittel, R. J.; Eilers, J.; Doose, S.; Sauer, M.; Siren, A. L.; Heckmann, M.; Paul, M. M. Active zone compaction correlates with presynaptic homeostatic potentiation. *Cell Rep* **2021**, *37* (1), 109770. DOI: 10.1016/j.celrep.2021.109770 From NLM Medline.

(37) Sakamoto, H.; Ariyoshi, T.; Kimpara, N.; Sugao, K.; Taiko, I.; Takikawa, K.; Asanuma, D.; Namiki, S.; Hirose, K. Synaptic weight set by Munc13-1 supramolecular assemblies. *Nat Neurosci* **2018**, *21* (1), 41-49. DOI: 10.1038/s41593-017-0041-9.

(38) Li, F.; Grushin, K.; Coleman, J.; Pincet, F.; Rothman, J. E. Diacylglycerol-dependent hexamers of the SNARE-assembling chaperone Munc13-1 cooperatively bind vesicles. *Proc Natl Acad Sci U S A* **2023**, *120* (44), e2306086120. DOI: 10.1073/pnas.2306086120 From NLM Medline.

(39) Grushin, K.; Kalyana Sundaram, R. V.; Sindelar, C. V.; Rothman, J. E. Munc13 structural transitions and oligomers that may choreograph successive stages in vesicle priming for neurotransmitter release. *Proc Natl Acad Sci U S A* **2022**, *119* (7). DOI: 10.1073/pnas.2121259119 From NLM Medline.

(40) Wilhelm, B. G.; Mandad, S.; Truckenbrodt, S.; Krohnert, K.; Schafer, C.; Rammner, B.; Koo, S. J.; Classen, G. A.; Krauss, M.; Haucke, V.; et al. Composition of isolated synaptic boutons reveals the amounts of vesicle trafficking proteins. *Science* **2014**, *344* (6187), 1023-1028. DOI: 10.1126/science.1252884.

(41) Grimm, J. B.; English, B. P.; Chen, J.; Slaughter, J. P.; Zhang, Z.; Revyakin, A.; Patel, R.; Macklin, J. J.; Normanno, D.; Singer, R. H.; et al. A general method to improve fluorophores for live-cell and single-molecule microscopy. *Nat Methods* **2015**, *12* (3), 244-250, 243 p following 250. DOI: 10.1038/nmeth.3256 From NLM Medline.

(42) Rossmann, K.; Akkaya, K. C.; Poc, P.; Charbonnier, C.; Eichhorst, J.; Gonschior, H.; Valavalkar, A.; Wendler, N.; Cordes, T.; Dietzek-Ivansic, B.; et al. N-Methyl deuterated rhodamines for protein labelling in sensitive fluorescence microscopy. *Chem Sci* **2022**, *13* (29), 8605-8617. DOI: 10.1039/d1sc06466e From NLM PubMed-not-MEDLINE.

(43) Kalla, S.; Stern, M.; Basu, J.; Varoqueaux, F.; Reim, K.; Rosenmund, C.; Ziv, N. E.; Brose, N. Molecular dynamics of a presynaptic active zone protein studied in Munc13-1-enhanced yellow fluorescent protein knock-in mutant mice. *J Neurosci* **2006**, *26* (50), 13054-13066. DOI: 10.1523/JNEUROSCI.4330-06.2006.

(44) Abramson, J.; Adler, J.; Dunger, J.; Evans, R.; Green, T.; Pritzel, A.; Ronneberger, O.; Willmore, L.; Ballard, A. J.; Bambrick, J.; et al. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **2024**, *630* (8016), 493-500. DOI: 10.1038/s41586-024-07487-w From NLM Medline.

(45) Padmanarayana, M.; Liu, H.; Michelassi, F.; Li, L.; Betensky, D.; Dominguez, M. J.; Sutton, R. B.; Hu, Z.; Dittman, J. S. A unique C2 domain at the C terminus of Munc13 promotes synaptic vesicle priming. *Proc Natl Acad Sci U S A* **2021**, *118* (11). DOI: 10.1073/pnas.2016276118 From NLM Medline.

(46) Stevens, D. R.; Wu, Z. X.; Matti, U.; Junge, H. J.; Schirra, C.; Becherer, U.; Wojcik, S. M.; Brose, N.; Rettig, J. Identification of the minimal protein domain required for priming activity of Munc13-1. *Curr Biol* **2005**, *15* (24), 2243-2248. DOI: 10.1016/j.cub.2005.10.055.

(47) Quade, B.; Camacho, M.; Zhao, X.; Orlando, M.; Trimbuch, T.; Xu, J.; Li, W.; Nicastro, D.; Rosenmund, C.; Rizo, J. Membrane bridging by Munc13-1 is crucial for neurotransmitter release. *Elife* **2019**, *8*. DOI: 10.7554/eLife.42806.

(48) Ast, J.; Nasteska, D.; Fine, N. H. F.; Nieves, D. J.; Koszegi, Z.; Lanoiselee, Y.; Cuozzo, F.; Viloria, K.; Bacon, A.; Luu, N. T.; et al. Revealing the tissue-level complexity of endogenous glucagon-like peptide-1 receptor expression and signaling. *Nat Commun* **2023**, *14* (1), 301. DOI: 10.1038/s41467-022-35716-1 From NLM Medline.

(49) Birke, R.; Ast, J.; Roosen, D. A.; Lee, J.; Rossmann, K.; Huhn, C.; Mathes, B.; Lisurek, M.; Bushiri, D.; Sun, H.; et al. Sulfonated red and far-red rhodamines to visualize SNAP- and Halo-tagged cell surface proteins. *Org Biomol Chem* **2022**, *20* (30), 5967-5980. DOI: 10.1039/d1ob02216d From NLM Medline.

(50) Chen, Z.; Das, B.; Nakamura, Y.; DiGregorio, D. A.; Young, S. M., Jr. Ca2+ channel to synaptic vesicle distance accounts for the readily releasable pool kinetics at a functionally mature auditory synapse. *J Neurosci* **2015**, *35* (5), 2083-2100. DOI: 10.1523/JNEUROSCI.2753-14.2015.

(51) Nakamura, Y.; Harada, H.; Kamasawa, N.; Matsui, K.; Rothman, J. S.; Shigemoto, R.; Silver, R. A.; DiGregorio, D. A.; Takahashi, T. Nanoscale distribution of presynaptic Ca(2+) channels and its impact on vesicular release during development. *Neuron* **2015**, *85* (1), 145-158. DOI: 10.1016/j.neuron.2014.11.019.

(52) Rebola, N.; Reva, M.; Kirizs, T.; Szoboszlay, M.; Lorincz, A.; Moneron, G.; Nusser, Z.; DiGregorio, D. A. Distinct Nanoscale Calcium Channel and Synaptic Vesicle Topographies Contribute to the Diversity of Synaptic Function. *Neuron* **2019**, *104* (4), 693-710 e699. DOI: 10.1016/j.neuron.2019.08.014.

(53) Keller, D.; Babai, N.; Kochubey, O.; Han, Y.; Markram, H.; Schurmann, F.; Schneggenburger, R. An Exclusion Zone for Ca2+ Channels around Docked Vesicles Explains Release Control by Multiple Channels at a CNS Synapse. *PLoS Comput Biol* **2015**, *11* (5), e1004253. DOI: 10.1371/journal.pcbi.1004253.

(54) Bell, M. E.; Bourne, J. N.; Chirillo, M. A.; Mendenhall, J. M.; Kuwajima, M.; Harris, K. M. Dynamics of nascent and active zone ultrastructure as synapses enlarge during long-term potentiation in mature hippocampus. *J Comp Neurol* **2014**, *522* (17), 3861-3884. DOI: 10.1002/cne.23646 From NLM Medline.

(55) Goel, P.; Dufour Bergeron, D.; Bohme, M. A.; Nunnelly, L.; Lehmann, M.; Buser, C.; Walter, A. M.; Sigrist, S. J.; Dickman, D. Homeostatic scaling of active zone scaffolds maintains global synaptic strength. *J Cell Biol* **2019**, *218* (5), 1706-1724. DOI: 10.1083/jcb.201807165.

(56) Dorrbaum, A. R.; Alvarez-Castelao, B.; Nassim-Assir, B.; Langer, J. D.; Schuman, E. M. Proteome dynamics during homeostatic scaling in cultured neurons. *Elife* **2020**, *9*. DOI: 10.7554/eLife.52939 From NLM Medline.

(57) Fukaya, R.; Hirai, H.; Sakamoto, H.; Hashimotodani, Y.; Hirose, K.; Sakaba, T. Increased vesicle fusion competence underlies long-term potentiation at hippocampal mossy fiber synapses. *Sci Adv* **2023**, *9* (8), eadd3616. DOI: 10.1126/sciadv.add3616 From NLM Medline.

(58) Wegner, W.; Mott, A. C.; Grant, S. G. N.; Steffens, H.; Willig, K. I. In vivo STED microscopy visualizes PSD95 sub-structures and morphological changes over several hours in the mouse visual cortex. *Sci Rep* **2018**, *8* (1), 219. DOI: 10.1038/s41598-017-18640-z From NLM Medline.

(59) Bosch, P. J.; Correa, I. R., Jr.; Sonntag, M. H.; Ibach, J.; Brunsveld, L.; Kanger, J. S.; Subramaniam, V. Evaluation of fluorophores to label SNAP-tag fused proteins for multicolor single-molecule tracking microscopy in live cells. *Biophys J* **2014**, *107* (4), 803-814. DOI: 10.1016/j.bpj.2014.06.040 From NLM Medline.

(60) Stockinger, F.; Poc, P.; Mohwald, A.; Karch, S.; Hafner, S.; Alzheimer, C.; Sandoz, G.; Huth, T.; Broichhagen, J. Multicolor, Cell-Impermeable, and High Affinity BACE1 Inhibitor Probes Enable Superior Endogenous Staining and Imaging of Single Molecules. *J Med Chem* **2024**, *67* (12), 10152-10167. DOI: 10.1021/acs.jmedchem.4c00339 From NLM Medline.

(61) Kohl, J.; Ng, J.; Cachero, S.; Ciabatti, E.; Dolan, M. J.; Sutcliffe, B.; Tozer, A.; Ruehle, S.; Krueger, D.; Frechter, S.; et al. Ultrafast tissue staining with chemical tags. *Proc Natl Acad Sci U S A* **2014**, *111* (36), E3805-3814. DOI: 10.1073/pnas.1411087111 From NLM Medline.

(62) Sutcliffe, B.; Ng, J.; Auer, T. O.; Pasche, M.; Benton, R.; Jefferis, G. S.; Cachero, S. Second-Generation Drosophila Chemical Tags: Sensitivity, Versatility, and Speed. *Genetics* **2017**, *205* (4), 1399-1408. DOI: 10.1534/genetics.116.199281 From NLM Medline.

(63) Yang, G.; de Castro Reis, F.; Sundukova, M.; Pimpinella, S.; Asaro, A.; Castaldi, L.; Batti, L.; Bilbao, D.; Reymond, L.; Johnsson, K.; Heppenstall, P. A. Genetic targeting of chemical indicators in vivo. *Nat Methods* **2015**, *12* (2), 137-139. DOI: 10.1038/nmeth.3207 From NLM Medline.

(64) Gutzeit, V. A.; Acosta-Ruiz, A.; Munguba, H.; Hafner, S.; Landra-Willm, A.; Mathes, B.; Mony, J.; Yarotski, D.; Borjesson, K.; Liston, C.; et al. A fine-tuned azobenzene for enhanced photopharmacology in vivo. *Cell Chem Biol* **2021**, *28* (11), 1648-1663 e1616. DOI: 10.1016/j.chembiol.2021.02.020 From NLM Medline.

(65) Ast, J.; Arvaniti, A.; Fine, N. H. F.; Nasteska, D.; Ashford, F. B.; Stamataki, Z.; Koszegi, Z.; Bacon, A.; Jones, B. J.; Lucey, M. A.; et al. Super-resolution microscopy compatible fluorescent probes reveal endogenous glucagon-like peptide-1 receptor distribution and dynamics. *Nat Commun* **2020**, *11* (1), 467. DOI: 10.1038/s41467-020-14309-w From NLM Medline.

(66) Shaib, A. H.; Chouaib, A. A.; Chowdhury, R.; Mihaylov, D.; Zhang, C.; Imani, V.; Georgiev, S. V.; Mougios, N.; Monga, M.; Reshetniak, S.; et al. Visualizing proteins by expansion microscopy. *bioRxiv* **2023**, 2022.2008.2003.502284. DOI: 10.1101/2022.08.03.502284.

(67) Schmidt, R.; Weihs, T.; Wurm, C. A.; Jansen, I.; Rehman, J.; Sahl, S. J.; Hell, S. W. MINFLUX nanometer-scale 3D imaging and microsecond-range tracking on a common fluorescence microscope. *Nat Commun* **2021**, *12* (1), 1478. DOI: 10.1038/s41467-021-21652-z From NLM Medline.

(68) Truckenbrodt, S.; Maidorn, M.; Crzan, D.; Wildhagen, H.; Kabatas, S.; Rizzoli, S. O. X10 expansion microscopy enables 25-nm resolution on conventional microscopes. *EMBO Rep* **2018**, *19* (9). DOI: 10.15252/embr.201845836.

(69) Chen, F.; Tillberg, P. W.; Boyden, E. S. Optical imaging. Expansion microscopy. *Science* **2015**, *347* (6221), 543-548. DOI: 10.1126/science.1260088 From NLM Medline.

(70) Doench, J. G.; Fusi, N.; Sullender, M.; Hegde, M.; Vaimberg, E. W.; Donovan, K. F.; Smith, I.; Tothova, Z.; Wilen, C.; Orchard, R.; et al. Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. *Nat Biotechnol* **2016**, *34* (2), 184-191. DOI: 10.1038/nbt.3437 From NLM Medline.

(71) Tycko, J.; Wainberg, M.; Marinov, G. K.; Ursu, O.; Hess, G. T.; Ego, B. K.; Aradhana; Li, A.; Truong, A.; Trevino, A. E.; et al. Mitigation of off-target toxicity in CRISPR-Cas9 screens for essential non-coding elements. *Nat Commun* **2019**, *10* (1), 4063. DOI: 10.1038/s41467-019-11955-7 From NLM Medline.

(72) Haeussler, M.; Schonig, K.; Eckert, H.; Eschstruth, A.; Mianne, J.; Renaud, J. B.; Schneider-Maunoury, S.; Shkumatava, A.; Teboul, L.; Kent, J.; et al. Evaluation of off-target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. *Genome Biol* **2016**, *17* (1), 148. DOI: 10.1186/s13059-016-1012-2 From NLM Medline.

(73) Concordet, J. P.; Haeussler, M. CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. *Nucleic Acids Res* **2018**, *46* (W1), W242-W245. DOI: 10.1093/nar/gky354 From NLM Medline.

(74) Biesemann, C.; Gronborg, M.; Luquet, E.; Wichert, S. P.; Bernard, V.; Bungers, S. R.; Cooper, B.; Varoqueaux, F.; Li, L.; Byrne, J. A.; et al. Proteomic screening of glutamatergic mouse brain synaptosomes isolated by fluorescence activated sorting. *EMBO J* **2014**, *33* (2), 157-170. DOI: 10.1002/embj.201386120.

(75) Burgalossi, A.; Jung, S.; Man, K. N.; Nair, R.; Jockusch, W. J.; Wojcik, S. M.; Brose, N.; Rhee, J. S. Analysis of neurotransmitter release mechanisms by photolysis of caged Ca(2)(+) in an autaptic neuron culture system. *Nat Protoc* **2012**, *7* (7), 1351-1365. DOI: 10.1038/nprot.2012.074.