Multiple neural pathways to successful visual short-term memory across the adult lifespan

Michelle G. Jansen[1], Alireza Salami[2, 3, 4, 5], Fernando Ruiz Martínez[1], Daniel J. Mitchell[6], Joukje M. Oosterman[1], Cam-CAN[7], Linda Geerligs[1*]

[1]Donders Institute for Brain, Cognition, and Behaviour, Radboud University, Nijmegen, the Netherlands

[2]Department of Integrative Medical Biology, Umeå University, Umeå, Sweden

[3]Umea Center for Functional Brain Imaging, Umeå University, Umeå, Sweden

[4]Wallenberg Centre for Molecular Medicine, Umeå University, Umeå, Sweden

[5]Aging Research Center, Karolinska Institutet & Stockholm University, Stockholm, Sweden

[6]MRC Cognition and Brain Sciences Unit, University of Cambridge, Cambridge, United Kingdom

[7]Cambridge Centre for Ageing and Neuroscience (Cam-CAN), University of Cambridge and MRC Cognition and Brain Sciences Unit, University of Cambridge, United Kingdom

*Corresponding author: linda.geerligs@donders.ru.nl

Abstract

Cognitive task performance can be supported through multiple neural pathways, a concept referred to as brain degeneracy. We used a novel approach to consider brain degeneracy during a visual short-term memory task (VSTM) across the adult lifespan in the Cam-CAN study (n = 113, 23-87 years old). Our main goal was to identify subgroups of participants whose VSTM performance was characterized by distinct brain activation patterns. First, we identified seven brain modules that responded similarly to the VSTM task and resembled previously identified functional networks (adjusted mutual information [aMI] = 0.45). Subsequently, latent profile analysis revealed four distinct subgroups of participants. Each subgroup was characterized by different recruitment patterns of these brain modules, predominantly in the frontal control module (FCM), visual module (VM), and default mode module (DMM). Subgroups did not differ in demographics or task performance. However, associations between brain activity and performance varied across subgroups, particularly in the FCM, suggesting that individuals may use different cognitive operations to perform the VSTM task. Further analyses revealed group differences in white matter integrity, mostly in the uncinate fasciculus, suggested that individual differences in structural brain properties may shape the different brain activation patterns. Altogether, our study contributes to our understanding of how multiple neural pathways could underlie cognitive performance.

Keywords: brain degeneracy; many-to-one mapping; fMRI; visual short-term memory; lifespan; white matter integrity

1 Introduction

Over the past decades, research sought to elucidate how the brain's functional and structural architecture gives rise to cognitive functioning (Finn et al., 2023; Genon et al., 2022; Park & Friston, 2013; Poldrack, 2006). Nevertheless, this relationship is far from straightforward. Not only do brain measures typically account for only a small portion of the variance in cognitive functioning (Hedden et al., 2016), but there are also several contradictory findings in terms of which neural mechanisms are adaptive or maladaptive for cognitive performance. For example, age-related increases in prefrontal activation have been linked to both superior and inferior performance in older adults (Grady, 2012; Nyberg et al., 2020). The complex nature of brain-cognition associations is further emphasized by previous studies showing that the extent to which brain structure or function relates to cognitive performance varies substantially across individuals (Lindenberger, 2014; Lövdén et al., 2018; Patel et al., 2022; Salami et al., 2018).

To gain a better understanding of how cognitive functions are supported throughout the lifespan, we may need to revisit the common assumption that most individuals exhibit similar patterns of neural activity to perform a certain cognitive task (one-to-one mapping) (Seghier & Price, 2018; Westlin et al., 2023). Conversely, brain degeneracy suggests that a specific task could be executed through multiple neural pathways (many-to-one mapping) (Edelman & Gally, 2001; Noppeney et al., 2004; Price & Friston, 2002). Degeneracy is evident in many neural systems, including those supporting cognitive functions (Kelso, 2012; Noppeney et al., 2004). In line with this, previous studies have identified subgroups of individuals showing unique neural activity patterns while performing the same task (Cerliani et al., 2017; Kherif et al., 2009; Seghier et al., 2008). Interestingly, this neural variability was associated with the use of different strategies to complete the task (Kherif et al., 2009; Noppeney et al., 2006), despite similar levels of task performance (Fischer-Baum et al., 2018; Noppeney et al., 2006). This illustrates how multiple neural routes can lead to successful cognitive performance. However, if we focus on similarities across individuals these possibilities are overlooked, emphasizing the need to understand brain degeneracy and what it might mean for how we study the associations between brain and cognition.

Building on the concept of many-to-one mapping, it has been suggested that secondary activation repertoires may be prompted when the primary circuit is unable to sustain successful task completion (Mason et al., 2015; Noppeney et al., 2004; Price & Friston, 2002). For instance, studies using transcranial magnetic stimulation showed that younger adults adapt to temporary functional lesions by recruiting alternative neural circuits (Lee et al., 2003; Pascual-Leone et al.,

2005). Furthermore, differential recruitment of brain regions during task performance has been

related to functional recovery after acquired brain injury (Nudo, 2013). In the context of normal

ageing, alterations in brain activity have been linked to the degradation of major white matter

tracts (Burzynska et al., 2013; Lucas et al., 2018), and localized grey matter loss (Kalpouzos et

al., 2012; Salami et al., 2012; Salami et al., 2014). Although this involves functional deterioration,

older adults are able to compensate for impaired brain function by recruiting additional brain

regions, particularly in the frontal lobes (Cabeza et al., 2002; Grady, 2012; Park & Reuter-Lorenz,

2009). In contrast, structural preservation in older individuals (i.e., brain maintenance; Nyberg

et al., 2012) has been related to neural activation patterns more closely resembling those

typically observed in younger adults (Cabeza et al., 2018; Nyberg et al., 2012). Together, these

findings stress that it is important to consider interindividual variability in brain structure to

understand why different neural systems may be recruited during task performance.

In the present study, we aimed to investigate brain degeneracy across the adult lifespan in the

context of visual short-term memory (VSTM), using data from the Cam-CAN project (n = 122, 23-

87 years old). VSTM is particularly interesting in the context of degeneracy, given that high

performance can be achieved via different strategies (Ritakallio et al., 2024). VSTM refers to our

ability to maintain visual information and is crucial for many other cognitive functions (Luck &

Vogel, 2013; Pearson & Keogh, 2019). However, the precise neural mechanisms underlying

successful VSTM performance are not fully understood, partially due to inconclusive findings in

prior research that did not consider the possibility of different brain activation patterns (Pearson

and Keogh, 2019). Our primary aim was therefore to identify subgroups of participants whose

VSTM performance was characterized by distinct brain activation patterns. We first identified

modules of brain regions that showed similar responses across participants during the VSTM

task. Next, we applied latent profile analysis to identify sub-groups of participants that

demonstrated differential recruitment patterns of the previously identified brain modules. To

better understand the variability in brain activity during the VSTM task, we characterized the

most significant differences in brain activity across subgroups.

Furthermore, we sought to elucidate factors contributing to the observed subgroup differences

in brain activity patterns. Specifically, we investigated the extent to which these subgroups

differed in demographic variables (i.e., age, sex, educational attainment) and task performance.

We were particularly interested in the role of age, as this factor was previously related to both

performance and neural correlates of the VSTM task (Lugtmeijer et al., 2023; Mitchell & Cusack,

2018; Morcom & Henson, 2018; Sander et al., 2012). For example, if age-related differences in

brain function were the largest source of inter-individual variability in brain activation patterns

during the VSTM task, we expected to identify subgroups that differed in their age and task performance. Conversely, if age-invariant inter-individual differences were most prominent, we expected to see participant subgroups that were roughly age-matched. Additionally, we investigated whether structural brain characteristics, such as grey matter volumes and white matter integrity, would explain variations in brain activity among the subgroups, as reflected by subgroup differences in these measures.

Lastly, we assessed associations between brain activation profiles and task performance within each subgroup. For example, increased activity in brain module X might have been beneficial for performance for subgroup Y but hindered this for subgroup Z. Such unique associations would not only provide further evidence for the presence of different neural routes to facilitate successful VSTM performance, but also potentially suggest cognitive strategies that were used to succeed.

2 Methods

2.1 Participants

This study included data from 122 participants (59 female, 48.4%) who were aged 23–87 (mean age 53.9, SD = 18.1) from the healthy, population-derived cohort tested in Stage III of the Cam-CAN project (Shafto et al., 2014; Taylor et al., 2017).

A complete overview of the study procedures and inclusion criteria have been described previously (Shafto et al., 2014; Taylor et al., 2017). Briefly, in Stage I, participants engaged in home-based interviews where demographic information, cognitive measures, and other measures of health (e.g., mental and physical health) were collected. In Stage II, participants engaged in three testing sessions, including cognitive and behavioural assessments, structural MRI (e.g., T1-weighted imaging and diffusion-weighted imaging), functional MRI (e.g., resting-state), and MEG recordings. During Stage III, among other things, structural MRI (e.g., T1-weighted images) and task-related functional MRI (fMRI) data during a VSTM task were acquired. In this manuscript, we use the demographic information from Stage I, the diffusion-weighted images from Stage II, and the T1-weighted images and VSTM fMRI paradigm from Stage III. Participants were native English-speakers with normal or corrected-to-normal vision and hearing and had no neurological disorders. Additional exclusion criteria were history of drug or alcohol abuse, and poor hearing or vision.

The study was conducted in accordance with ethical approval obtained from the Cambridgeshire
2 (now East of England - Cambridge Central) Research Ethics Committee. All participants gave
written informed consent prior to participation.

We excluded a total of 9 participants from our analyses due to poor VSTM task performance (*n*
= 8) or data quality issues (*n* = 1*)*, leaving a final sample of 113 participants (see below for more
details).

2.2 Data acquisition

2.2.1 Demographic information

Demographic information, including age, sex, and educational attainment, was acquired during
home-based interviews during Stage I of the Cam-CAN project. Educational attainment in this
study was based on the highest level of completed education (i.e., categorized in 1 = no
education above the age of 16 years old; 2 = GCSEs or equivalent; 3 = A levels or equivalent; 4 =
university levels or equivalent).

2.2.2 MRI data

MRI data was acquired on a 3T Siemens TIM Trio System scanner at the MRC Cognition Brain
and Sciences Unit, Cambridge, UK. During each of three VSTM task runs, functional data were
acquired using a multi-echo, T2*-weighted EPI sequence (TR = 2000 milliseconds, TE = 12
milliseconds, 25 milliseconds, 38 milliseconds, flip angle = 78 degrees, 32 axial slices of thickness
of 3.7 mm with an interslice gap of 20%, FOV = 294mm × 294 mm, voxel-size = 3 mm × 3 mm ×
3.48 mm). Each volume had 34 axial slices (acquired in descending order), slice thickness of 2.9
mm, with an interslice gap of 20%. The multiple echoes were combined by computing their
average weighted by their estimated T2* contrast. The number of scans acquired varied per run,
depending on reaction time, between 294 and 349.

A high-resolution 3D T1-weighted structural image was acquired using a Magnetization Prepared
Rapid Gradient Echo (MPRAGE) sequence, with the following parameters: TR = 2250 ms; TE =
2.99 ms; TI = 900 ms; flip angle = 9 degrees; FOV = 256 × 240 × 192 mm; voxel size = 1 mm
isotropic; GRAPPA acceleration factor = 2. This T1 image was missing for two of the participants
in this sample, however for these participants T1 images were acquired during the scanning
session that took place during Stage II of Cam-CAN, approximately 1-3 years earlier.

Diffusion-weighted images (DWIs) were acquired with a twice-refocused spin-echo sequence
during Stage II of Cam-CAN. Thirty diffusion gradient directions were obtained for each of two

b-values: 1000 and 2000 s/mm2, plus three images acquired with a b-value of 0. These

parameters were optimized for estimation of the diffusion kurtosis tensor and associated scalar

metrics (Shafto et al., 2014). Other DWI parameters were TR = 9100, TE = 104 ms, FOV = 192

mm × 192 mm, voxel size 2 mm isotropic, 66 axial slices, with acquisition time 10 minutes and 2

seconds.

2.2.3 Visual short-term memory (VSTM) task

During fMRI scanning, participants performed a VSTM paradigm (see **Figure 1)**, based on Emrich

et al. (2013). A detailed description of the task is provided in Lugtmeijer et al. (2023).

Briefly, three displays containing different coloured moving dots (red, yellow and blue) were

presented sequentially on each trial. Each trial began with a grey fixation dot for 5 seconds.

Subsequently, the fixation dot brightened for 2 seconds and was followed by the coloured dot

displays. Each coloured dot display was shown for 500ms, with a 250ms black screen in between.

As a manipulation of the number of items in memory, trials could contain one, two or three

arrays of dots moving in a single direction (load 1, 2, and 3, respectively).

Participants were instructed to remember the direction of motion for dot displays in which all

dots moved in a single direction, and to ignore any dot displays that rotated around the fixation

location (the encoding phase). As three displays were shown in each trail, the memory load was

manipulated by varying the number of rotating dots in relation to the number of moving dot

displays. After the third coloured dot display, there was a blank display of 8 seconds (the

maintenance phase), followed by a probe display. The probe display consisted of a coloured

circle that indicated which of the three dot displays (red, yellow, or blue) the participant should

recall. Participants had up to 5 seconds to adjust a pointer on the circle until it matched the

direction of motion of the probed dot display (the response phase).

Figure 1: Adapted from Morcom and Henson (2018). An example of the VSTM task. Trials began
with a fixation dot for 7 seconds. On each trial, three moving dot displays were then presented
in red, yellow, and blue for 500 ms each (250 ms in between). After the last display, the screen
was blank for an 8 second maintenance period and then a probe display was presented. A
coloured circle indicated which dot display participants should recall (red, yellow, or blue).
Participants had up to 5 seconds to adjust a pointer until its angle matched the motion direction
of the probed display.

2.3 VSTM performance

Several factors can limit VSTM performance. For example, memory imprecision (e.g.,
uncertainty between 80 or 85 degrees), random guessing (e.g., when forgetting an item
entirely), and reporting an incorrect item (mis-binding, e.g., reporting the angle of the red dots
instead of the blue dots). To distinguish these different sources of error, we applied a three-
component mixture model (Bays et al., 2009). A complete description of the application of the
mixture model to the VSTM task is provided in previous work (Mitchell et al., 2018; Lugtmeijer
et al., 2023).

For our subsequent analyses, we used the estimates of a memory item's precision (the inverse
of the standard deviation in degrees) and the number of items (directions) that were stored in
memory (K). Eight participants in our sample showed very poor performance for memory of a
single item (>30 degrees deviation on average). This suggests that they may not have understood
the task instructions and were therefore excluded from further analyses. Since there was a
ceiling effect for the number of items in memory at load 1 (mean = 0.96, SD = 0.06) and to some
extent at load 2 (mean = 1.76, SD = 0.24), we decided to focus on performance at load 3 (mean

= 2.40, SD = 0.46). The distribution of scores for the number of items in memory at each load is

shown in **Supplementary Figure 1**.

2.4 MRI pre-processing and first level analyses

All MRI data were pre-processed using SPM 12 (http://www.fil.ion.ucl.ac.uk/spm) and

"automatic analysis" software (Cusack et al., 2015). The T1 and T2 weighted structural images

were rigid body registered with a MNI template, bias corrected and segmented into six tissue

classes using the newer segment protocol in SPM 12 (Ashburner & Friston, 2005). Non-linear

transformation using DARTEL normalization was performed to match a grey matter template

created from the Cam-CAN stage III sample. The functional images were spatially realigned and

interpolated to correct for differences in acquisition times, and co-registered to the structural

image using rigid body transformations. Both grey matter volume images and functional images

were transformed to MNI space using the warps and affine transformation estimated from the

structural images. For grey matter volume images, this was done while preserving the total

amount of signal from each region in the images. The functional images were resliced to 3x3x3

mm voxels, while grey matter volume images were resliced to 1.5x1.5x1.5 mm voxels and both

were smoothed with an 8 mm full-width half-maximum (FWHM) Gaussian kernel. For the two

participants with missing T1-weighted scans, normalization and co-registration were performed

based on the T1 images collected during stage II of the Cam-CAN project. One participant was

excluded from further analyses due to data quality issues.

For each participant and voxel, we fit a general linear model (GLM) with three phases of each

trial per run: encoding, maintenance, and response. The encoding phase was modelled as an

epoch of 2 seconds starting from the beginning of the first dot display movement. The

maintenance phase was modelled as an epoch of 8 seconds (i.e., the duration of the blank

display that was shown after all probes were displayed). The response phase was modelled as

an epoch of the time between the probe onset and response. These three components were

each split into three regressors: one for each load. Six additional regressors were added per run,

representing the motion parameters estimated in the realignment stage. The means of each run

were also modelled separately. The main contrast of interest in this study was the difference

between the highest and the lowest memory load during the maintenance period (load 3 vs.

load 1). This contrast allows us to equate the amount of visual information that was presented

just before the maintenance period as well as the motor response that was required just after,

while maximizing the difference in memory load.

2.5 Identifying modules with similar inter-individual variability

To make the analyses more computationally efficient in later stages, we reduced the dimensionality of the data by grouping voxels into a set of small regions of interest (ROI). Specifically, we used the 748 (out of 840) regions from the Craddock atlas (2012) that had sufficient coverage in our recent paper (Geerligs et al., 2015), allowing us to use the network labels defined in our previous study. To select ROIs that might be involved in task-performance in a subset of participants, we first extracted the average t-values for the contrast of interest (i.e., load 3 vs. load 1 during the maintenance period) across all voxels in each ROI for each participant. To further reduce the dimensionality of the data, we only included ROIs that showed increased or reduced activation in load 3 versus load 1 in at least 10% of the participants at a liberal significance threshold (T > 1.96, p < 0.05). This resulted in the final inclusion of 418 ROIs.

For each included ROI, we extracted the average beta-values. Subsequently, we calculated the correlation values between ROIs for load 3 vs. load 1 across participants (i.e., ROI-by-ROI correlation matrix). A high correlation coefficient between two ROIs suggests that these regions show similar inter-individual variability in load-dependent modulation of the BOLD response, and therefore may contribute to the same brain module.

In the next step, we used the Brain Connectivity Toolbox to identify groups of ROIs, or brain modules, that showed similar brain activity patterns across participants (Rubinov & Sporns, 2010). Specifically, we applied consensus partitioning based on the Louvain modularity algorithm (Bassett et al., 2013; Lancichinetti & Fortunato, 2012). The ROI-by-ROI correlation matrix was used to create an initial partition of the ROIs into distinct modules with the Louvain modularization algorithm (Blondel et al., 2008), and was further refined using an optimization procedure (Sun et al., 2009). This partitioning procedure was repeated 500 times, and all the repetitions were assembled into an ROI-by-ROI agreement matrix, where each number indicates the proportion of repetitions in which a pair of ROIs were assigned to the same module. The ROI-by-ROI agreement matrix was then used as an input for consensus partitioning, where a permuted version of the agreement matrix was used for thresholding. This was repeated until the algorithm converged into a matrix of ones and zeros, indicating that a single partition was reached.

The complete partitioning procedure was performed for multiple values of the resolution parameter γ, ranging from 1 to 1.5 with steps of 0.05. The γ parameter is a penalty term that biases the size of the communities to be found. Smaller values of γ will result in larger modules,

while higher values will result in smaller modules. As an γ value of 1.5 already resulted in 49
modules, we did not explore a wider range of this parameter space.

Next, we selected the partition with the highest resemblance to a previously defined set of age-
representative networks (Geerligs et al., 2015b), as measured with the adjusted mutual
information (aMI), as our final set of brain modules. The optimum we identified was at γ = 1.25,
resulting in 13 distinct modules (aMI = 0.45). However, as many modules consisted of only one
ROI or a few ROIs, we only included modules with at least 15 ROIs in our analyses. This resulted
in the exclusion of 6 brain modules, comprising of 28 ROIs in total. The final set of 7 brain
modules was used for further analyses (see "Results").

Within each brain module and for each participant, we computed the average brain activity for
load 3 vs. load 1. However, we observed strong correlations between the average brain activity
values across all brain modules (ranging from r=0.39 to 0.77). This is consistent with prior
research which demonstrated that overall network responsivity, rather than module-specific
activation patterns, were related to age and task performance (Samu et al., 2017). To be able to
look beyond this general factor in our subsequent analyses, we computed the residual activity
within each brain module after regressing out the mean activity across the 418 ROIs. To be able
to retain information about the engagement of each network during the task, the mean activity
across all subjects was added back after computing the residuals. The extent to which the mean
activity across these ROIs explained the variance within each brain module varied between 0.61-
0.75, as indicated by the adjusted R^2 values.

2.6 Brain activity profiles

To identify latent subgroups of participants characterized by different patterns of brain activity,
we applied latent profile analysis (LPA) using the Mclust package in R (Scrucca et al., 2016).
Briefly, LPA is a data-driven Gaussian mixture modelling method to identify hidden subgroups in
a population (Vermunt & Magidson, 2002). A Gaussian mixture model assumes a multivariate
Gaussian distribution for each subgroup. Therefore, every observation (here, participant) will
have a specific probability of belonging to a specific subgroup, as specified by a probability
density function determined by the average, the variance, and the covariance of each of the
subgroup distributions.

We applied LPA using both the residual activity in each brain module and the overall responsivity
across all networks, specifying between 1 and 5 subgroups. This allowed us to look at module-
specific variations in responsivity in addition to the overall responsivity across brain modules.
We restricted the model space to models that do not allow for covariance between modules

within subgroups of participants (i.e., models with a spherical or diagonal distribution of the
within-group covariance matrix). This way, our model assumes that any correlations between
modules must be caused by participants with different brain activity profiles (similar to Lövdén
et al., 2018). The Bayesian Information Criterion (BIC), Integrated Completed Likelihood (ICL)
and bootstrapped Likelihood Ratio Test were used to select the most optimal model and class
solution (i.e., the number of subgroups).

2.7 Structural brain characteristics

To investigate subgroup differences in grey matter, we extracted grey matter volumes for each
of the 418 ROIs that were included in our main analysis and subsequently averaged the grey
matter volumes within each module. The two participants with the missing T1-scan at Stage III
were excluded from this analysis, resulting in a sample of 111 participants for the grey matter
data.

To investigate subgroup differences in white matter microstructure, we used the diffusion data
to estimate mean kurtosis. Mean kurtosis is influenced by changes in organelles, cell membranes
and the ratio of intracellular to extracellular water compartments (Falangola et al., 2008). It
provides a sensitive metric to determine how aging affects the complexity of white matter
microstructure, and typically decreases in late-life (Benitez et al., 2018).

Diffusion data were pre-processed and analysed as described in a previous paper (Geerligs et
al., 2018). Briefly, the data were co-registered, normalized to MNI space and smoothed with a
1-mm FWHM Gaussian kernel to reduce residual interpolation errors. Subsequently, we
calculated the average mean kurtosis values for 18 tract-based regions of interest defined by
the Johns Hopkins University white matter tractography atlas (Hua et al., 2008). These included
the uncinate fasciculus, superior longitudinal fasciculus, inferior fronto-occipital fasciculus,
anterior thalamic radiations, forceps minor, forceps major, cerebrospinal tract, the inferior
longitudinal fasciculus, ventral cingulate gyrus, and the dorsal cingulate gyrus. All tracts were
split into their left and right hemisphere sections, except for the forceps minor and forceps
major.

Participants with severe head motion during the DWI scan, as revealed by an automated striping
detection method on the diffusion data, were excluded (Neto Henriques and Correia, 2015). In
addition, for each tract, participants with mean kurtosis values of 3 SDs above or below the
group average were excluded. This resulted in a final sample of 96 participants for all tracts,
except for the inferior longitudinal fasciculus (n = 95), superior longitudinal fasciculus (n = 95),
cerebrospinal tract (n = 94), and ventral cingulate gyrus (n = 86).

2.8 Statistical tests

All statistical tests were performed using R (version 4.3.1).

Given that overall responsivity (i.e., average activity across all included ROIs) explained most of the variability in brain module activity, we first explored how this factor related to age and task performance using Pearson correlations. To investigate whether associations between overall responsivity and task performance were age-independent, we additionally calculated the partial correlations between these two variables while accounting for age.

Next, we performed several between-group comparisons to better understand the diverse brain activity profiles associates with the VSTM task that were identified with the latent profile analysis. To assess which brain modules showed the most pronounced differences across subgroups we statistically compared brain module activity between groups. We also tested whether subgroups differed in demographic characteristics (age, sex, educational attainment), VSTM task performance (the number and precision of items in memory at load 3), grey matter volumes (i.e., average grey matter volumes in each of the brain modules), and white matter microstructure (i.e., average mean kurtosis in each of the 18 tracts). These comparisons were performed to understand which factors contributed to the differences in brain activity profiles.

We applied a two-step approach to determine between-group differences. First, Welch's ANOVA was used to determine if there were differences between subgroups in the variable of interest. This was used instead of the regular Fisher's one-way ANOVA because of unequal sizes and variances between groups. Second, each significant ANOVA was followed by pairwise Welch's t-tests between groups (i.e., using non-pooled standard deviations to consider the unequal variances between groups). Chi-squared tests were used in case of categorical data (i.e., comparing sex or educational attainment across groups). As educational attainment had less than < 5 observations in certain categories, this variable was split into a dichotomous category prior to performing the comparisons (university level vs. other educational categories), however for descriptive purposes all categories are reported.

Analyses on grey matter volumes were corrected for intracranial volume. All analyses were repeated while additionally correcting for the effects of age, except for the comparisons on demographic characteristics. Specifically, we regressed the covariate(s) on the variable of interest and used the residuals of that regression for the ANOVA and subsequent t-tests.

To further investigate the extent to which different neural routes facilitated successful VSTM performance, we investigated the associations between brain module activity and task

performance within each subgroup. If certain associations are present in one subgroup but not in another, this could suggest that these subgroups relied on different cognitive operations to successfully complete the task. To test this, we calculated Pearson correlations between brain module activity and task performance within each subgroup, adjusting for the effects of age (i.e., partial correlations). The difference between coefficients were statistically compared using Fisher's z-tests.

We used false discovery rate (FDR) corrections (Benjamini & Hochberg, 1995) to adjust for multiple comparisons across different brain modules, white matter tracts or pairwise tests. In addition to reporting (FDR-corrected) p-values, we computed Bayes Factors (BF) for the alternative hypothesis using the *BayesFactor* package (https://cran.r-project.org/web/packages/BayesFactor/index.html). Effects were considered as strongly supported in case of both FDR-adjusted $p < 0.05$ and $BF > 3$ (Buades-Rotger et al., 2023; Driessen et al., 2018). In addition, we calculated Bayesian R^2 to evaluate the fit of each model (Gelman et al., 2019), using the *performance* package (https://easystats.github.io/performance/index.html).

3 Results

3.1 Brain modules with similar brain activity patterns across participants

We identified seven distinct modules of brain regions with similar patterns of inter-individual variability in terms of their brain activity while performing the VSTM task (**Figure 2**). The brain activity metric utilized to define these modules (and used in all subsequent analyses) is based on load 3 vs. load 1 during the maintenance period.

The modules included a frontal control module (FCM), which overlapped largely with the traditional fronto-parietal control network. This module encompasses the lateral prefrontal cortex and the anterior cingulate but did not include parietal areas. A distinct temporo-parietal module (TPM) included parietal areas that typically fall within the default mode and fronto-parietal control network, predominantly in the left hemisphere, but also included temporal areas. The cerebellum and thalamus module (CTM) covered the cerebellum and thalamus, and the visual module (VM) included the occipital cortex. Another module overlapped with the dorsal attention network, covering the frontal eye fields and the intraparietal sulcus, which we refer to as the dorsal attention module (DAM). The motor module (MM) covered the bilateral precentral gyrus but also extended into the right posterior temporal cortex. Lastly, one module

largely overlapped with the default mode network, covering the medial prefrontal cortex,
precuneus, and temporal areas, which we refer to as the default mode module (DMM).

Figure 2: An illustration of the seven brain modules we identified.

3.2 Overall responsivity increases with age and accounts for module-specific brain activity

As previously described, the average activity across all included ROIs explained a substantial
proportion of the variance in brain activity within each module (R^2 = 0.61-0.75).

We explored how the overall responsiveness related to age and task performance. Average
activity across all brain modules increased with age (r = 0.37, 95% CI = [0.19; 0.52], uncorrected
p < 0.001, BF = 338.96). However, we did not observe significant associations between overall
activity and memory precision (r = 0.04, uncorrected p = 0.688, BF = 0.15) or number of items in
memory (r = -0.07, uncorrected p = 0.460, BF = 0.18).

3.3 Latent profile analysis identifies subgroups with distinct activation signature across
modules

To look beyond overall responsivity, this factor was regressed out of the seven identified brain
networks. To account for all variance in brain activity, the LPA grouped participants based on
the residual brain activity within each brain network, in addition to the overall responsivity
across all ROIs. The BIC and ICL values suggested that the most optimal fit was acquired with a
model of four latent subgroups with equal shape, variable volume and assumed diagonal
covariance matrix (n = 35, n = 24, n = 42, and n = 12 in groups 1-4, respectively). The distributions
of BIC and ICL values for the different latent profile models are shown in **Supplementary Figure**
2 and 3. A more detailed description of the decision on the number of classes, as well as the
corresponding fit indices, is provided in **Supplementary Methods.**

An overview of the brain module activation profiles across subgroups is displayed in **Figure 3.** To
investigate which networks were the strongest drivers of the subgroup differences, we

statistically compared the brain activity profiles across subgroups **(Table 1).** It should be noted

that the LPA was designed to produce subgroups with distinct brain activity profiles. Therefore,

these significance values are only meaningful in relation to each other as a comparison of the

relative contributions of different modules and subgroups, but not in isolation. A complete

overview of the between-group statistics for the brain module activity is depicted in

Supplementary Table 1.

The most substantial differences in residual brain activity were observed for the FCM, VM, DAM,

and DMM, as reflected by the corresponding Bayes Factors and values of Bayesian R^2 (all ≥ 0.26,

indicating substantial effect size; Cohen, 2013). Focusing on four most influential modules,

subgroup 1 is characterized by less FCM activity compared to subgroups 3 and 4, the lowest

DAM activity across all groups, and the least DMM deactivity. Subgroup 2 demonstrates the least

FCM activity but the highest VM activity across all subgroups. In contrast, group 3 shows higher

FCM activity than subgroups 1 and 2, but lower VM activation than subgroup 2. Subgroup 4

shows a similar pattern as subgroup 3 with even higher FCM and lower VM activity. We observed

no significant differences for the TPM and overall responsivity of brain activity to the task load

manipulation. These results remained consistent after applying age corrections.

In summary, while subgroup 2 exhibited the lowest frontal activation paired with higher visual

activation, subgroups 3 and 4 displayed the opposite pattern. In contrast, subgroup 1 was

characterized by having the least DAM activity and DMM deactivation. This may suggest that

different neural mechanisms were used to complete the VSTM task.

Figure 3: *For each of the four identified subgroups this figure shows the mean-regressed average*

brain activity in each of the seven identified brain networks as well as the overall responsivity

across all ROIs (Resp). The most influential brain modules were the FCM, VM, DAM, and DMM.

FCM = frontal control module; TPM = temporal parietal module; CTM = cerebellum/thalamus

module; VM = visual module; DAM = dorsal attention module; MM = motor module; DMM =

default mode module.

Table 1. *Comparing residual network activity across subgroups*

	F	Bayesian R^2	t 2>1	t 3>1	t 4>1	t 3>2	t 4>2	t 4>3
FCM	54.29**	0.51†	-5.37**	4.82**	3.43**	12.45**	4.98**	2.00
TPM	1.05	0.00						
CTM	10.92**	0.15	4.42**	5.33**	-0.03	-0.45	-1.42	-1.32
VM	33.93**	0.46†	7.48**	-0.89	-2.54*	-9.43**	-6.22**	-2.23
DAM	20.00**	0.28†	5.69**	7.01**	3.06*	0.30	1.11	1.03
MM	9.45**	0.15	-0.05	-4.97**	-1.91	-3.29*	-1.75	-0.17
DMM	42.56**	0.47†	-8.99**	-10.49**	-4.32*	-0.37	-1.31	-1.21
Resp	0.77	0.00						

479 *FDR-corrected p-value < 0.05 & BR > 3;

480 **=FDR-corrected p-value < 0.001 & BR > 3.

481 †$R^2 \geq 0.26$.

3.4 Subgroups showed no differences in demographics or VSTM task performance

Subgroup characteristics are displayed in **Table 2.** No significant between-group differences were found for age, sex, education (all $p > 0.05$, BF < 3). We found a trend level age difference between the subgroups ($F_{3,40.8}$=2.79, $p = 0.053$, BF = 0.43), with subgroup 4 tended to be younger than the other participant groups (subgroup 1: $t_{23.6}$ = 2.83, $p = 0.009$, BF = 3.70; subgroup 2: $t_{27.7}$ = 2.29, $p = 0.030$, BF = 1.74; subgroup 3: t_{18} = 2.01, $p = 0.059$, BF = 1.49). VSTM performance did not significantly differ across subgroups (precision and number of items in memory at load 3, all $p > 0.05$, BF < 3). These results were unaffected by age corrections. A complete overview of the between-group statistics for the subgroup characteristics can be found in **Supplementary Table 2**.

Table 2. Subgroup characteristics.

	All participants	Subgroup 1	Subgroup 2	Subgroup 3	Subgroup 4
N	113	35	24	42	12
Age, mean (SD)	53 (18.0)	57 (19.1)	55 (19.8)	51 (15.6)	41 (15.4)
Sex					
Male, n (%)	59 (52%)	14 (40%)	15 (63%)	24 (57%)	6 (50%)
Female, n (%)	54 (48%)	21 (60%)	9 (38%)	18 (43%)	6 (50%)
Highest education					
University, n (%)	82 (73%)	21 (60%)	19 (79%)	32 (76%)	10 (83%)
A' levels, n (%)	17 (15%)	6 (17%)	3 (13%)	7 (17%)	1 (8.3%)
GCSE grade, n (%)	9 (8%)	4 (11%)	1 (4.2%)	3 (7.1%)	1 (8.3%)
None > 16, n (%)	5 (4.4%)	4 (11%)	1 (4.2%)	0 (0%)	0 (0%)
Number of items in memory (load 3), mean (SD)	2.40 (0.46)	2.35 (0.50)	2.38 (0.47)	2.43 (0.47)	2.49 (0.34)
Memory precision (load 3), mean (SD)	0.12 (0.05)	0.10 (0.04)	0.12 (0.04)	0.12 (0.05)	0.13 (0.04)

3.5 Subgroups revealed distinct associations between brain activity and VSTM task performance

Next, we investigated the association between brain activity and task performance within each subgroup, while correcting for age. Since this concerned within-group analyses, we excluded subgroup 4 because of its small number of participants (*n* = 12). To limit the number of comparisons, we focused on the brain modules with the most pronounced differences between the subgroups: the FCM, VM, DAM, and DMM. A complete overview of the associations between brain module activity and performance on the VSTM task for each of the subgroups is displayed in **Supplementary Table 3.**

For the FCM, we found that the subgroups 1 and 2 showed negative associations between FCM

activity and memory precision (subgroup 1: r = −0.37, 95% CI = [−0.62; −0.04], uncorrected p =

0.030, BF = 2.36; subgroup 2: r = −0.41, 95% CI = [−0.70; 0.01], uncorrected p = 0.045, BF = 1.93,

see **Figure 5A**). However, this association was absent in subgroup 3 (r = 0.11, p = 0.465, BF =

0.30). When we compared the slopes between subgroups 1 and 3, we found that they were

significantly different (z = -2.11, 95% CI = [-0.87; -0.03], uncorrected p-value = 0.035), the same

was true for the comparison between subgroups 2 and 3 (z = -2.05, 95% CI = [-0.94; -0.02],

uncorrected p-value = 0.040). Notably, subgroups 1 and 2 were characterized by having

relatively low FCM recruitment than subgroup 3 and 4. Therefore, these results suggest that the

low recruitment of the FCM within subgroups 1 and 2 may have been beneficial to their memory

precision. No association was observed between FCM activity and number of items in memory

in any of the subgroups.

For the VM, no significant associations with items in memory or memory precision were

observed.

In the DAM, we observed that subgroup 1 demonstrated a positive association between DAM

activity and memory precision (r = 0.42, 95% CI = [0.10; 0.66], uncorrected p = 0.012, BF = 4.97,

see **Figure 5B).** In subgroup 2, this association tended in a similar direction; however, this was

not significant (r = 0.30, uncorrected p = 0.15). The slope from subgroup 1 significantly differed

from subgroup 3 (z = 2.14, 95% CI = [0.04; 0.86], uncorrected p = 0.032). Subgroup 1 was

characterized by having the least DAM recruitment. This suggests that for subgroup 1, their low

DAM activity may have negatively impacted their memory precision. There were no significant

associations between DAM activity and number of items in memory.

In the DMM, no associations were observed between DMM activity and memory precision. A

negative association was observed between reduced DMM deactivation and the number of

items in memory in subgroup 1 (r = -0.38, 95% CI = [-0.63; -0.05], uncorrected p = 0.026, BF =

2.65), however no associations were found in subgroup 2 (r = -0.11, p = 0.597, BF = 0.35) or

subgroup 3 (r = 0.01, p = 0.942, BF = 0.23; see **Figure 5C**). The slopes did not significantly differ

between groups. These results suggest that the increased activity (or reduced suppression) of

the DMM in subgroup 1 may have hindered the number of items participants in that subgroup

could recall.

Taken together, we observed several unique associations between brain module activity and

VSTM performance that were subgroup-specific, even though task performance did not differ

across groups. This further supports the notion that these distinct brain activity profiles reflect

different yet equally effective pathways to perform the task. It should be noted that after
applying FDR-corrections for multiple testing across the three subgroups and four networks,
none of these associations remained significant, and therefore should be interpreted with
caution.

Figure 4: Associations between brain module activity in the frontal control module (FCM), visual module (VM), dorsal attention module (DAM), and default mode module (DMM) and performance on the visual short-term memory task. All analyses were corrected for age. **A.** *Subgroups 1 and 2 showed negative associations between FCM activity and memory precision (uncorrected p = 0.030, uncorrected p = 0.045).* **B.** *Subgroup 1 demonstrated a positive association between DAM activity and memory precision (uncorrected p = 0.012).* **C.** *Subgroup 1 demonstrated a negative association between reduced DMM suppression and the number of items in memory (uncorrected p = 0.026).*

3.6 Subgroups differ in white matter integrity

Next, we investigated whether structural brain characteristics differed between groups. First, we compared grey matter volumes within each of the modules between groups while correcting for total intracranial volume. Grey matter volume in the TPM significantly differed across subgroups ($F_{3,39.84}$ = 5.67, FDR-adjusted p-value = 0.017, BF = 5.10), however this effect disappeared after additionally correcting for age (unadjusted p = 0.048; BF = 0.72).

We also investigated whether the different subgroups showed significant differences in white matter microstructure, as measured by mean kurtosis (MK). We found significant differences between subgroups for four white matter tracts (**see Figure 5**). These included the left inferior frontal occipital fasciculus (IFOF; $F_{3,37.8}$ = 4.73, FDR-adjusted p = 0.030, BF = 5.04), right IFOF

559   (F$_{3,36.10}$ = 4.08, FDR-adjusted p-value = 0.049, BF = 3.49), left inferior longitudinal fasciculus

560   (F$_{3,35.09}$ = 6.71, FDR-adjusted p-value = 0.011, BF = 28.10) and the left uncinate fasciculus (F$_{3,35.43}$

561   = 6.60, FDR-adjusted p-value = 0.011, BF = 16.93).

Figure 5: The upper part of the figure shows the average mean kurtosis values in each subgroup for the white matter tracts that displayed significant between-group differences (FDR-corrected p-value < 0.05 and Bayes Factor > 3). The bottom part visualizes the white matter tracts in a glass brain, generated with MRIcroGL (https://www.nitrc.org/projects/mricrogl). Only the effects in the left inferior longitudinal fasciculus and left uncinate fasciculus survived additional age corrections.

Only the effects in the left inferior longitudinal fasciculus and the left uncinate fasciculus

remained significant after correcting for age (F$_{3,34.8}$ = 5.42, FDR-adjusted p-value = 0.033, BF =

5.81; F$_{3,35.4}$ = 6.64, FDR-corrected p-value = 0.020, BF = 17.32, respectively). The left inferior

fasciculus showed lower MK in subgroup 2 relatively to subgroup 3 (t$_{53.56}$ = 3.71, FDR-correct p =

0.003, BF = 30.24) and subgroup 4 ($t_{13.37}$ = 2.65, unadjusted p = 0.019, BF = 11.23), however the

latter effect did not survive FDR-corrections.

The left uncinate fasciculus showed lower mean kurtosis in subgroup 2 relatively to subgroup 1

($t_{45.50}$ = 2.58, FDR-corrected p-value = 0.039, BF = 3.49), subgroup 3 ($t_{50.46}$ = 4.37, FDR-corrected

p-value < 0.001, BF > 100), and subgroup 4 ($t_{15.49}$ = 2.60, FDR-corrected p-value = 0.039, BF =

6.62). So, subgroup 2, which is the subgroup with the least FCM activity but the highest VM

activity across all subgroups, specifically shows lower MK in the left uncinate fasciculus and the

left inferior longitudinal fasciculus. Notably, the low recruitment of the FCM within this subgroup

appeared to be beneficial to their memory precision.

A complete overview of the between-group statistics for grey matter volumes and MK is

provided in **Supplementary Tables 4 and 5.**

4 Discussion

While we often assume that similar patterns of neural activity are engaged to complete a

cognitive task across individuals, accumulating evidence suggests that this is not necessarily the

case, even in the absence of brain injury or disease (Noppeney et al., 2004; Price & Friston, 2002;

Seghier & Price, 2018). Instead of discarding inter-individual variability in task-related functional

activity, we leveraged it in our analyses to investigate brain degeneracy or many-to-one mapping

in the context of VSTM performance. We identified four participant subgroups with distinct

brain activity profiles, particularly within the FCM, VM, DAM, and DMM. Despite substantial

differences in brain activity, these subgroups showed no difference in task performance or age,

suggesting that these profiles may reflect different yet equally effective pathways to perform

the task. This was further supported by the small, albeit distinct associations between task

performance and brain modular activity across subgroups. Individual differences in brain

structure may partly account for differences in brain activity, as subgroups showed age-

independent differences in white matter integrity for the left inferior longitudinal fasciculus and

the left uncinate fasciculus. Here, we first discuss our findings in relation to many-to-one

mapping. Subsequently, we describe the modules and subgroups that we identified in more

detail as well as the role of global versus local inter-individual variability in brain activity. Finally,

we highlight the factors that contributed to subgroup variations in brain activity.

Degeneracy suggests that there can be multiple neural pathways leading to the same

behavioural outcome (Edelman & Gally, 2001; Mason et al., 2015; Price & Friston, 2002; Westlin

et al., 2023). We identified four subgroups with marked differences in recruitment of various brain modules during the VSTM task, after accounting for differences in overall network activity. These results add to previous studies that revealed subgroups with distinct profiles of task-related functional activity (Cerliani et al., 2017; Fischer-Baum et al., 2018; Kherif et al., 2009; Noppeney et al., 2006; Noppeney et al., 2005; Seghier et al., 2008; Seghier & Price, 2016). Similarly to previous findings (Fischer-Baum et al., 2018; Noppeney et al., 2006), these subgroups demonstrated similar levels of performance. These findings illustrate how different patterns of brain activity can lead to the same functional outcome, providing support for the notion of brain degeneracy. Additionally, we observed several unique associations between brain modular activity and task performance. This finding suggests that there is not a single route to successful cognitive performance, and the optimal route may vary from person to person. This finding highlights the individual-specific nature of brain-cognition relationship, where diverse neural mechanisms can yield similar outcomes (Noppeney et al., 2006; Seghier & Price, 2018).

As a first step toward identifying subgroups, we reduced the dimensionality of our data, identifying seven brain modules that showed similar between-participant variability in their activity patterns during the maintenance period of the VSTM task (load 3 vs. load 1). Strikingly, these modules show strong correspondence to brain networks that are typically identified using functional connectivity analyses, such as the frontoparietal control network and default mode network (Power et al., 2011; Yeo et al., 2011). This happens even though functional connectivity is typically based on temporal correlations in activity within participants while we investigated correlations in activity between participants, suggesting that there is a shared underlying source. These results corroborate the idea that there is a stable intrinsic network architecture that is largely uniform across individuals but also shows reliable individual differences (Finn et al., 2015; Fox & Raichle, 2007; Gratton et al., 2018). This intrinsic architecture manifests itself both in how networks are functionally connected within a single person (within-subject) and in the differences in network activity among individuals (between-subjects). In line with this, previous studies demonstrated that functional connectivity can be used to predict task-activation, and that individual differences in connectivity are related to variability in task-related activation (Cole et al., 2014; Cole et al., 2021; Finn et al., 2015; Tavor et al., 2016).

We observed that the brain activity across the seven brain modules was highly correlated, corroborating previous findings where the global recruitment of task-related functional activity explained most inter-individual variability, both in relation to performance and age (Chen et al., 2022; Mitchell et al., 2023; Samu et al., 2017). Consistent with prior studies, the average activity

across all modules explained 60-75% of the variability in activity observed within each brain module. To be able to look beyond this general factor, we used module-specific activity for our subsequent analysis, where the average activity within each module was corrected for the average activity across all modules. Results from the latent profile analysis suggests that there is meaningful inter-individual variability beyond this general factor, by revealing subgroups with differences in white matter integrity and unique associations between brain activity and performance. These findings illustrate how individual differences in task-related brain activation arise from both overall responsivity to the task (Samu et al., 2017), and the specific recruitment patterns of brain regions, highlighting the importance of individual-specific factors in shaping task-related brain activity (Cole et al., 2021; Finn et al., 2015; Gratton et al., 2018).

Of the seven modules we identified, four showed the most significant differences in activity across subgroups: the FCM, VM, DAM, and DMM. These brain modules largely overlap with the frontoparietal network, visual network, dorsal attention network, and default mode network, respectively (Power et al., 2011; Yeo et al., 2011). With increasing task load, activity typically increases in the frontoparietal and dorsal attention networks, whereas the default mode network becomes more suppressed (Huang et al., 2016; Nagel et al., 2009; Salami et al., 2019; Salami et al., 2018; Zuo et al., 2019). This reflects a shift away from internally focused processes to task-oriented activity (Fox et al., 2005; Seeley et al., 2007; Smith et al., 2009). Notably, this pattern was weaker for subgroup 1, suggesting that this subgroup might have been less attentive during the maintenance phase (Anticevic et al., 2012; Berry et al., 2017). Surprisingly, this subgroup showed task performance comparable to the other subgroups, despite the positive association between DAM activity and memory precision, and the negative association between DMM activity and the number of items in memory. One possibility is that VSTM performance was sustained in this subgroup through increased engagement during the encoding or retrieval phase (Linke et al., 2011; Magen, 2017). Stimulus information might also have been preserved in a passive state, where maintenance is facilitated by neural mechanisms not characterized by increased activity (i.e., activity-silent mechanisms), such as short-term synaptic plasticity (Lewis-Peacock et al., 2014; Stokes, 2015).

Successful VSTM performance is further facilitated through interactions between the frontoparietal and visual networks (D'Esposito & Postle, 2015; Eriksson et al., 2015; Scimeca et al., 2018). The sensory recruitment hypothesis of working memory suggests that the same brain regions responsible for encoding the stimulus are also essential for maintaining this information (D'Esposito, 2007; D'Esposito & Postle, 2015). In line with this, in the context of VSTM, visual regions have been shown to maintain detailed sensory representations of the stimuli (Emrich et

al., 2013; Postle, 2015; Serences, 2016). The frontoparietal network is thought to exert top-down control over these areas and support the maintenance of more abstract representations of the stimuli, such as verbal descriptions (Christophel et al., 2017; Gayet et al., 2018; Scimeca et al., 2018; Serences, 2016). Interestingly, subgroup 2 demonstrated the highest VM activity and lowest FCM activity. The reduced FCM recruitment in this subgroup was beneficial to VSTM performance, as reflected by the negative associations between FCM activity and memory precision in these participants. In contrast, subgroups 3 and 4 demonstrated relatively high FCM activity and lower VM activity. The use of different strategies to complete the task is an important source of variability for task-related brain activity (Miller et al., 2012; Sanfratello et al., 2014). Consistently, previous studies that identified subgroups with distinct brain activity profiles during task engagement attributed this variability to the adoption of different strategies (Kherif et al., 2009). It is possible that subgroup 2 might have relied more on visual representations, while subgroup 3 and 4 maintained stimulus information through more abstract forms, thereby relying more on the FCM. This would also align with the two most commonly reported strategies to perform VSTM tasks: maintaining visual representations of the stimuli (e.g., picturing the red dots moving to the left corner) and translating salient stimuli features into verbal representations (e.g., mentally repeating "the red dots moved to the left corner") (Berger & Gaunitz, 1979; Pearson & Keogh, 2019).

Subgroup 2 also showed decreased white matter integrity in the inferior longitudinal fasciculus and the uncinate fasciculus, whereas this was preserved in subgroups 3 and 4. The inferior longitudinal fasciculus is involved in modulating visual information and semantic-lexical processing (Herbet et al., 2018; Shekari & Nozari, 2023). The uncinate fasciculus is associated with associative learning and semantic retrieval (Olson et al., 2015; Von Der Heide et al., 2013). This suggests that the use of a visually oriented strategy in subgroup 2 may be linked to the reduced microstructural integrity in two white matter tracts that play a role in creating and retrieving verbal representations (Duffau et al., 2013; Herbet et al., 2018; Shekari & Nozari, 2023). However, we did not find any direct association between brain activity and performance in subgroup 3, which makes it challenging to interpret how this brain activity profile precisely facilitated task performance in this group.

The age-independent differences in white matter integrity across subgroups were consistent with our expectation that structural brain characteristics would contribute to the variability in brain activity among the subgroups (or vice versa). This also aligns with previous research linking white matter microstructure to the recruitment of different brain regions during task performance (Burzynska et al., 2013; Burzynska et al., 2015). We did not observe any differences

in grey matter across subgroups, suggesting that white matter integrity may be relatively more important to explain inter-individual variability in brain activity in our sample (Miller et al., 2012). Typically, white matter degradation is associated with declines in cognitive performance, including working memory (Bennett & Madden, 2014; Oschwald et al., 2020). Our results suggest that this is not necessarily the case when different neural pathways can be used to perform a task (Madden et al., 2009), corresponding to the notion of brain degeneracy (Noppeney et al., 2004; Price & Friston, 2002). It should be noted that the differences in white matter integrity between subgroups were relatively small, emphasizing the need to further investigate these associations in future studies. Nevertheless, our findings underscore that individual differences in structural brain characteristics are an important source of variability in brain activity and cognitive performance (Genon et al., 2022; Kanai & Rees, 2011; Miller et al., 2012; Seghier & Price, 2018).

Although prior research has demonstrated strong age-related differences in brain activity and performance during VSTM tasks, including the one used here (Lugtmeijer et al., 2023; Morcom & Henson, 2018; Sander et al., 2012), we did not observe any differences in age or performance between the subgroups. This suggests that other factors beyond age or task performance were the primary drivers behind the distinct brain activation patterns observed across subgroups (Finn et al., 2015; Gratton et al., 2018). These factors may have included the extent to which participants were engaged throughout the task (Linke et al., 2011; Magen, 2017; Rose, 2020), the strategies adopted to complete the task (Kherif et al., 2009; Miller et al., 2012), and variations in white matter integrity (Burzynska et al., 2013; Burzynska et al., 2015), as discussed above. Nonetheless, age was positively correlated with the global recruitment of brain activity, consistent with prior research showing that that variability in overall responsivity was primarily due to age-differences (Mitchell et al., 2023; Samu et al., 2017). Therefore, the lack of age differences between subgroups might also be explained by our approach of clustering participants based on the residual brain activity in each module (i.e., after regressing out the overall responsiveness). In contrast, task performance was not associated with overall responsivity, suggesting that this factor is not the most substantial driver of inter-individual variability in brain activity. Therefore, studies that are interested in explaining performance metrics would benefit from distinguishing individuals based on performance itself rather than brain activity (e.g., Lövden et al., 2018; Salami et al., 2018).

There are several limitations that should be considered when interpreting these findings. First, we explored how the brain activity differences across subgroups may be associated with different cognitive processes. Although the use of different strategies has been recognized an

important source of task-related functional activity (Kherif et al., 2009; Miller et al., 2012; Noppeney et al., 2006; Sanfratello et al., 2014), we note that this interpretation remains speculative as we do not have any information about which strategies participants used when they performed the task. Therefore, the evidence here should be taken as proof of principle, showing that when we use a different approach to look at task-related brain activity, interesting dimensions of inter-individual variability are revealed that may otherwise remain hidden (Cerliani et al., 2017; Fischer-Baum et al., 2018; Kherif et al., 2009; Noppeney et al., 2006; Seghier et al., 2008; Seghier & Price, 2016; Seghier & Price, 2018). Second, our dataset consists of a relatively small number of participants, which may have limited our ability to observe within-group associations with performance. Lastly, although we have discussed the role of several factors in explaining the variability in brain activity between subgroups, this does not rule out the possibility that other variables that were not considered here could also contribute, such as other early-life factors, genetic factors or brain development (Walhovd et al., 2023). We hope that future studies focusing on this subject will include larger samples and a more extensive characterization of individual-specific factors that could further explain this variability.

Taken together, the novel approach introduced here allowed us to uncover distinct subgroups with marked differences in neural activity during the VSTM task, associations between brain module activity and VSTM performance, and white matter microstructural properties. However, these subgroups did not significantly differ in their demographic characteristics or VSTM performance. The identification of distinct subgroups illustrates how multiple neural pathways may underlie successful VSTM performance. This further highlights the importance of considering brain degeneracy to understand variability in brain-cognition associations (Edelman & Gally, 2001; Mason et al., 2015; Price & Friston, 2002; Westlin et al., 2023). Recognizing this possibility provides us with a more comprehensive understanding of the different ways in which our brains support cognitive functions and how this may vary from person to person. Consequently, these insights may help to unravel new ways in which structure and function interact, and how complementary routes can lead to the same cognitive outcomes.

Data and code availability

Data availability is provided via the Cam-CAN data portal (https://camcan-archive.mrc-cbu.cam.ac.uk/dataaccess/). The scripts and analysis pipeline are made available via Github (https://github.com/michellegjansen/neural_pathways_VSTM).

Author contributions

Michelle G. Jansen: Conceptualization, methodology, software, investigation, visualization, validation, data curation, formal analysis, writing – original draft. Alireza Salami: Conceptualization, methodology, writing – review & editing. Fernando-Ruiz Martínez: Methodology, software, investigation, data curation, formal analysis. Daniel J. Mitchell: Methodology, data curation, formal analysis, writing – review & editing. Joukje M. Oosterman: Supervision, writing – review & editing. Linda Geerligs: Conceptualization, methodology, software, investigation, visualization, validation, data curation, formal analysis, supervision, writing – original draft.

Funding

The Cam-CAN study received funding from the UK Biotechnology and Biological Sciences Research Council (BB/H008217/1), the UK Medical Research Council Cognition & Brain Sciences Unit (CBU), University of Cambridge, UK.

Alireza Salami is supported by the Swedish Research Council (grant number 2016-01936), Riksbankens Jubileumsfond (grant number P20-0515), Knut and Alice Wallenberg Foundation (Wallenberg Fellow grant to A.S.), and StratNeuro grant at Karolinska Institutet. Daniel J. Mitchell is supported by the Medical Research Council Intramural Program MC_UU_00030/7. Linda Geerligs supported by a Vidi grant from the Dutch Research Council (NWO, VI.Vidi.201.150).

Declaration of Competing Interests

The authors declare no competing interests.

Acknowledgments

We thank the Cam-CAN respondents and their primary care teams in Cambridge for their participation in this study, and colleagues at the MRC Cognition and Brain Sciences Unit MEG and MRI facilities for their assistance. Further information about the Cam-CAN corporate authorship membership can be found at http://www.cam-can.com/publications/Cam-CAN_Corporate_Author.html (list #15).

References

Anticevic, A., Cole, M. W., Murray, J. D., Corlett, P. R., Wang, X. J., & Krystal, J. H. (2012). The role of default network deactivation in cognition and disease. *Trends Cogn Sci*, *16*(12), 584-592. https://doi.org/10.1016/j.tics.2012.10.008

Ashburner, J., & Friston, K. J. (2005). Unified segmentation. *Neuroimage*, *26*(3), 839-851. https://doi.org/10.1016/j.neuroimage.2005.02.018

Bassett, D. S., Porter, M. A., Wymbs, N. F., Grafton, S. T., Carlson, J. M., & Mucha, P. J. (2013). Robust detection of dynamic community structure in networks. *Chaos*, *23*(1), 013142. https://doi.org/10.1063/1.4790830

Bays, P. M., Catalao, R. F., & Husain, M. (2009). The precision of visual working memory is set by allocation of a shared resource. *Journal of vision*, *9*(10), 7-7.

Benitez, A., Jensen, J. H., Falangola, M. F., Nietert, P. J., & Helpern, J. A. (2018). Modeling white matter tract integrity in aging with diffusional kurtosis imaging. *Neurobiology of Aging*, *70*, 265-275. https://doi.org/https://doi.org/10.1016/j.neurobiolaging.2018.07.006

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1), 289-300. https://doi.org/https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Bennett, I. J., & Madden, D. J. (2014). Disconnected aging: cerebral white matter integrity and age-related differences in cognition. *Neuroscience*, *276*, 187-205. https://doi.org/10.1016/j.neuroscience.2013.11.026

Berger, G. H., & Gaunitz, S. C. B. (1979). Self-rated imagery and encoding strategies in visual memory. *British Journal of Psychology*, *70*(1), 21-24. https://doi.org/https://doi.org/10.1111/j.2044-8295.1979.tb02137.x

Berry, A. S., Sarter, M., & Lustig, C. (2017). Distinct Frontoparietal Networks Underlying Attentional Effort and Cognitive Control. *J Cogn Neurosci*, *29*(7), 1212-1225. https://doi.org/10.1162/jocn_a_01112

Buades-Rotger, M., Smeijers, D., Gallardo-Pujol, D., Krämer, U. M., & Brazil, I. A. (2023). Aggressive and psychopathic traits are linked to the acquisition of stable but imprecise hostile expectations. *Translational Psychiatry*, *13*(1), 197. https://doi.org/10.1038/s41398-023-02497-0

Burzynska, A. Z., Garrett, D. D., Preuschhof, C., Nagel, I. E., Li, S.-C., Bäckman, L., Heekeren, H. R.,
& Lindenberger, U. (2013). A Scaffold for Efficiency in the Human Brain. *The Journal of*
Neuroscience, *33*(43), 17150-17159. https://doi.org/10.1523/jneurosci.1426-13.2013

Burzynska, A. Z., Wong, C. N., Voss, M. W., Cooke, G. E., McAuley, E., & Kramer, A. F. (2015).
White Matter Integrity Supports BOLD Signal Variability and Cognitive Performance in
the Aging Human Brain. *PLOS ONE*, *10*(4), e0120315.
https://doi.org/10.1371/journal.pone.0120315

Cabeza, R., Albert, M., Belleville, S., Craik, F. I. M., Duarte, A., Grady, C. L., Lindenberger, U.,
Nyberg, L., Park, D. C., Reuter-Lorenz, P. A., Rugg, M. D., Steffener, J., & Rajah, M. N.
(2018). Maintenance, reserve and compensation: the cognitive neuroscience of healthy
ageing. *Nat Rev Neurosci*, *19*(11), 701-710. https://doi.org/10.1038/s41583-018-0068-2

Cabeza, R., Anderson, N. D., Locantore, J. K., & McIntosh, A. R. (2002). Aging Gracefully:
Compensatory Brain Activity in High-Performing Older Adults. *Neuroimage*, *17*(3), 1394-
1402. https://doi.org/https://doi.org/10.1006/nimg.2002.1280

Cerliani, L., Thomas, R. M., Aquino, D., Contarino, V., & Bizzi, A. (2017). Disentangling subgroups
of participants recruiting shared as well as different brain regions for the execution of
the verb generation task: A data-driven fMRI study. *Cortex*, *86*, 247-259.
https://doi.org/https://doi.org/10.1016/j.cortex.2016.11.017

Chen, X., Rundle, M. M., Kennedy, K. M., Moore, W., & Park, D. C. (2022). Functional activation
features of memory in successful agers across the adult lifespan. *Neuroimage*, *257*,
119276. https://doi.org/https://doi.org/10.1016/j.neuroimage.2022.119276

Christophel, T. B., Klink, P. C., Spitzer, B., Roelfsema, P. R., & Haynes, J.-D. (2017). The Distributed
Nature of Working Memory. *Trends in Cognitive Sciences*, *21*(2), 111-124.
https://doi.org/https://doi.org/10.1016/j.tics.2016.12.007

Cohen, J. (2013). *Statistical power analysis for the behavioral sciences*. Academic press.

Cole, M. W., Bassett, D. S., Power, J. D., Braver, T. S., & Petersen, S. E. (2014). Intrinsic and task-
evoked network architectures of the human brain. *Neuron*, *83*(1), 238-251.
https://doi.org/10.1016/j.neuron.2014.05.014

Cole, M. W., Ito, T., Cocuzza, C., & Sanchez-Romero, R. (2021). The Functional Relevance of Task-
State Functional Connectivity. *J Neurosci*, *41*(12), 2684-2702.
https://doi.org/10.1523/jneurosci.1713-20.2021

Cusack, R., Vicente-Grabovetsky, A., Mitchell, D. J., Wild, C. J., Auer, T., Linke, A. C., & Peelle, J.
E. (2015). Automatic analysis (aa): efficient neuroimaging workflows and parallel
processing using Matlab and XML [Technology Report]. *Frontiers in Neuroinformatics*, *8*.
https://doi.org/10.3389/fninf.2014.00090

D'Esposito, M. (2007). From cognitive to neural models of working memory. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *362*(1481), 761-772.
https://doi.org/doi:10.1098/rstb.2007.2086

D'Esposito, M., & Postle, B. R. (2015). The Cognitive Neuroscience of Working Memory. *Annual*
Review of Psychology, *66*(Volume 66, 2015), 115-142.
https://doi.org/https://doi.org/10.1146/annurev-psych-010814-015031

Driessen, J. M. A., Fanti, K. A., Glennon, J. C., Neumann, C. S., Baskin-Sommers, A. R., & Brazil, I.
A. (2018). A comparison of latent profiles in antisocial male offenders. *Journal of*
Criminal Justice, *57*, 47-55.
https://doi.org/https://doi.org/10.1016/j.jcrimjus.2018.04.001

Duffau, H., Herbet, G., & Moritz-Gasser, S. (2013). Toward a pluri-component, multimodal, and
dynamic organization of the ventral semantic stream in humans: lessons from
stimulation mapping in awake patients [Opinion]. *Frontiers in Systems Neuroscience*, *7*.
https://doi.org/10.3389/fnsys.2013.00044

Edelman, G. M., & Gally, J. A. (2001). Degeneracy and complexity in biological systems.
Proceedings of the National Academy of Sciences, *98*(24), 13763-13768.

Emrich, S. M., Riggall, A. C., LaRocque, J. J., & Postle, B. R. (2013). Distributed Patterns of Activity
in Sensory Cortex Reflect the Precision of Multiple Items Maintained in Visual Short-
Term Memory. *The Journal of Neuroscience*, *33*(15), 6516-6523.
https://doi.org/10.1523/jneurosci.5732-12.2013

Eriksson, J., Vogel, Edward K., Lansner, A., Bergström, F., & Nyberg, L. (2015). Neurocognitive
Architecture of Working Memory. *Neuron*, *88*(1), 33-46.
https://doi.org/https://doi.org/10.1016/j.neuron.2015.09.020

Finn, E. S., Poldrack, R. A., & Shine, J. M. (2023). Functional neuroimaging as a catalyst for
integrated neuroscience. *Nature*, *623*(7986), 263-273. https://doi.org/10.1038/s41586-
023-06670-9

Finn, E. S., Shen, X., Scheinost, D., Rosenberg, M. D., Huang, J., Chun, M. M., Papademetris, X.,
& Constable, R. T. (2015). Functional connectome fingerprinting: identifying individuals
using patterns of brain connectivity. *Nature Neuroscience*, *18*(11), 1664-1671.
https://doi.org/10.1038/nn.4135

Fischer-Baum, S., Kook, J. H., Lee, Y., Ramos-Nuñez, A., & Vannucci, M. (2018). Individual
Differences in the Neural and Cognitive Mechanisms of Single Word Reading. *Front Hum*
Neurosci, *12*, 271. https://doi.org/10.3389/fnhum.2018.00271

Fox, M. D., & Raichle, M. E. (2007). Spontaneous fluctuations in brain activity observed with
functional magnetic resonance imaging. *Nat Rev Neurosci*, *8*(9), 700-711.
https://doi.org/10.1038/nrn2201

Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E. (2005).
The human brain is intrinsically organized into dynamic, anticorrelated functional
networks. *Proc Natl Acad Sci U S A*, *102*(27), 9673-9678.
https://doi.org/10.1073/pnas.0504136102

Gayet, S., Paffen, C. L. E., & Van der Stigchel, S. (2018). Visual Working Memory Storage Recruits
Sensory Processing Areas. *Trends in Cognitive Sciences*, *22*(3), 189-190.
https://doi.org/10.1016/j.tics.2017.09.011

Geerligs, L., Rubinov, M., Cam, C., & Henson, R. N. (2015). State and Trait Components of
Functional Connectivity: Individual Differences Vary with Mental State. *J Neurosci*,
35(41), 13949-13961. https://doi.org/10.1523/jneurosci.1324-15.2015

Gelman, A., Goodrich, B., Gabry, J., & Vehtari, A. (2019). R-squared for Bayesian Regression
Models. *The American Statistician*, *73*(3), 307-309.
https://doi.org/10.1080/00031305.2018.1549100

Genon, S., Eickhoff, S. B., & Kharabian, S. (2022). Linking interindividual variability in brain
structure to behaviour. *Nature Reviews Neuroscience*, *23*(5), 307-318.
https://doi.org/10.1038/s41583-022-00584-7

Grady, C. (2012). The cognitive neuroscience of ageing. *Nature Reviews Neuroscience*, *13*(7),
491-505. https://doi.org/10.1038/nrn3256

Gratton, C., Laumann, T. O., Nielsen, A. N., Greene, D. J., Gordon, E. M., Gilmore, A. W., Nelson,
S. M., Coalson, R. S., Snyder, A. Z., Schlaggar, B. L., Dosenbach, N. U. F., & Petersen, S. E.
(2018). Functional Brain Networks Are Dominated by Stable Group and Individual

Factors, Not Cognitive or Daily Variation. *Neuron*, *98*(2), 439-452.e435.
https://doi.org/10.1016/j.neuron.2018.03.035

Hedden, T., Schultz, A. P., Rieckmann, A., Mormino, E. C., Johnson, K. A., Sperling, R. A., &
Buckner, R. L. (2016). Multiple Brain Markers are Linked to Age-Related Variation in
Cognition. *Cereb Cortex*, *26*(4), 1388-1400. https://doi.org/10.1093/cercor/bhu238

Herbet, G., Zemmoura, I., & Duffau, H. (2018). Functional Anatomy of the Inferior Longitudinal
Fasciculus: From Historical Reports to Current Hypotheses. *Front Neuroanat*, *12*, 77.
https://doi.org/10.3389/fnana.2018.00077

Huang, A. S., Klein, D. N., & Leung, H.-C. (2016). Load-related brain activation predicts spatial
working memory performance in youth aged 9–12 and is associated with executive
function at earlier ages. *Developmental Cognitive Neuroscience*, *17*, 1-9.
https://doi.org/https://doi.org/10.1016/j.dcn.2015.10.007

Kalpouzos, G., Persson, J., & Nyberg, L. (2012). Local brain atrophy accounts for functional
activity differences in normal aging. *Neurobiol Aging*, *33*(3), 623.e621-623.e613.
https://doi.org/10.1016/j.neurobiolaging.2011.02.021

Kanai, R., & Rees, G. (2011). The structural basis of inter-individual differences in human
behaviour and cognition. *Nature Reviews Neuroscience*, *12*(4), 231-242.
https://doi.org/10.1038/nrn3000

Kelso, J. A. (2012). Multistability and metastability: understanding dynamic coordination in the
brain. *Philos Trans R Soc Lond B Biol Sci*, *367*(1591), 906-918.
https://doi.org/10.1098/rstb.2011.0351

Kherif, F., Josse, G., Seghier, M. L., & Price, C. J. (2009). The main sources of intersubject
variability in neuronal activation for reading aloud. *J Cogn Neurosci*, *21*(4), 654-668.
https://doi.org/10.1162/jocn.2009.21084

Lancichinetti, A., & Fortunato, S. (2012). Consensus clustering in complex networks. *Scientific
Reports*, *2*(1), 336. https://doi.org/10.1038/srep00336

Lee, L., Siebner, H. R., Rowe, J. B., Rizzo, V., Rothwell, J. C., Frackowiak, R. S., & Friston, K. J.
(2003). Acute remapping within the motor system induced by low-frequency repetitive
transcranial magnetic stimulation. *Journal of Neuroscience*, *23*(12), 5308-5318.

Lewis-Peacock, J. A., Drysdale, A. T., & Postle, B. R. (2014). Neural Evidence for the Flexible
Control of Mental Representations. *Cerebral Cortex*, *25*(10), 3303-3313.
https://doi.org/10.1093/cercor/bhu130

Lindenberger, U. (2014). Human cognitive aging: Corriger la fortune? *Science*, *346*(6209), 572-
578. https://doi.org/doi:10.1126/science.1254403

Linke, A. C., Vicente-Grabovetsky, A., Mitchell, D. J., & Cusack, R. (2011). Encoding strategy
accounts for individual differences in change detection measures of VSTM.
Neuropsychologia, *49*(6), 1476-1486.
https://doi.org/https://doi.org/10.1016/j.neuropsychologia.2010.11.034

Lövdén, M., Karalija, N., Andersson, M., Wåhlin, A., Axelsson, J., Köhncke, Y., Jonasson, L. S.,
Rieckman, A., Papenberg, G., Garrett, D. D., Guitart-Masip, M., Salami, A., Riklund, K.,
Bäckman, L., Nyberg, L., & Lindenberger, U. (2018). Latent-Profile Analysis Reveals
Behavioral and Brain Correlates of Dopamine-Cognition Associations. *Cereb Cortex*,
28(11), 3894-3907. https://doi.org/10.1093/cercor/bhx253

Lucas, M., Wagshul, M. E., Izzetoglu, M., & Holtzer, R. (2018). Moderating Effect of White Matter
Integrity on Brain Activation During Dual-Task Walking in Older Adults. *The Journals of
Gerontology: Series A*, *74*(4), 435-441. https://doi.org/10.1093/gerona/gly131

Luck, S. J., & Vogel, E. K. (2013). Visual working memory capacity: from psychophysics and
neurobiology to individual differences. *Trends Cogn Sci*, *17*(8), 391-400.
https://doi.org/10.1016/j.tics.2013.06.006

Lugtmeijer, S., Geerligs, L., Tsvetanov, K. A., Mitchell, D. J., Cam, C. A. N., & Campbell, K. L. (2023).
Lifespan differences in visual short-term memory load-modulated functional
connectivity. *Neuroimage*, *270*, 119982.
https://doi.org/https://doi.org/10.1016/j.neuroimage.2023.119982

Madden, D. J., Bennett, I. J., & Song, A. W. (2009). Cerebral White Matter Integrity and Cognitive
Aging: Contributions from Diffusion Tensor Imaging. *Neuropsychology Review*, *19*(4),
415-435. https://doi.org/10.1007/s11065-009-9113-2

Magen, H. (2017). The role of central attention in retrieval from visual short-term memory.
Psychonomic Bulletin & Review, *24*(2), 423-430. https://doi.org/10.3758/s13423-016-
1111-9

Mason, P. H., Winter, B., & Grignolio, A. (2015). Hidden in plain view: degeneracy in complex
systems. *Biosystems*, *128*, 1-8.

Miller, M. B., Donovan, C.-L., Bennett, C. M., Aminoff, E. M., & Mayer, R. E. (2012). Individual
differences in cognitive style and strategy predict similarities in the patterns of brain
activity between individuals. *Neuroimage*, *59*(1), 83-93.
https://doi.org/https://doi.org/10.1016/j.neuroimage.2011.05.060

Mitchell, D. J., & Cusack, R. (2018). Visual short-term memory through the lifespan: Preserved
benefits of context and metacognition. *Psychol Aging*, *33*(5), 841-854.
https://doi.org/10.1037/pag0000265

Mitchell, D. J., Mousley, A. L. S., Shafto, M. A., Cam-CAN, & Duncan, J. (2023). Neural
Contributions to Reduced Fluid Intelligence across the Adult Lifespan. *The Journal of*
Neuroscience, *43*(2), 293-307. https://doi.org/10.1523/jneurosci.0148-22.2022

Morcom, A. M., & Henson, R. N. A. (2018). Increased Prefrontal Activity with Aging Reflects
Nonspecific Neural Responses Rather than Compensation. *J Neurosci*, *38*(33), 7303-
7313. https://doi.org/10.1523/jneurosci.1701-17.2018

Nagel, I. E., Preuschhof, C., Li, S.-C., Nyberg, L., Bäckman, L., Lindenberger, U., & Heekeren, H. R.
(2009). Performance level modulates adult age differences in brain activation during
spatial working memory. *Proceedings of the National Academy of Sciences*, *106*(52),
22552-22557. https://doi.org/doi:10.1073/pnas.0908238106

Noppeney, U., Friston, K. J., & Price, C. J. (2004). Degenerate neuronal systems sustaining
cognitive functions. *J Anat*, *205*(6), 433-442. https://doi.org/10.1111/j.0021-
8782.2004.00343.x

Noppeney, U., Penny, W. D., Price, C. J., Flandin, G., & Friston, K. J. (2006). Identification of
degenerate neuronal systems based on intersubject variability. *Neuroimage*, *30*(3), 885-
890. https://doi.org/10.1016/j.neuroimage.2005.10.010

Noppeney, U., Price, C. J., Penny, W. D., & Friston, K. J. (2005). Two Distinct Neural Mechanisms
for Category-selective Responses. *Cerebral Cortex*, *16*(3), 437-445.
https://doi.org/10.1093/cercor/bhi123

Nudo, R. (2013). Recovery after brain injury: mechanisms and principles [Review]. *Frontiers in*
Human Neuroscience, *7*. https://doi.org/10.3389/fnhum.2013.00887

Nyberg, L., Boraxbekk, C.-J., Sörman, D. E., Hansson, P., Herlitz, A., Kauppi, K., Ljungberg, J. K., Lövheim, H., Lundquist, A., Adolfsson, A. N., Oudin, A., Pudas, S., Rönnlund, M., Stiernstedt, M., Sundström, A., & Adolfsson, R. (2020). Biological and environmental predictors of heterogeneity in neurocognitive ageing: Evidence from Betula and other longitudinal studies. *Ageing Research Reviews*, *64*, 101184. https://doi.org/https://doi.org/10.1016/j.arr.2020.101184

Nyberg, L., Lövdén, M., Riklund, K., Lindenberger, U., & Bäckman, L. (2012). Memory aging and brain maintenance. *Trends Cogn Sci*, *16*(5), 292-305. https://doi.org/10.1016/j.tics.2012.04.005

Olson, I. R., Von Der Heide, R. J., Alm, K. H., & Vyas, G. (2015). Development of the uncinate fasciculus: Implications for theory and developmental disorders. *Dev Cogn Neurosci*, *14*, 50-61. https://doi.org/10.1016/j.dcn.2015.06.003

Oschwald, J., Guye, S., Liem, F., Rast, P., Willis, S., Röcke, C., Jäncke, L., Martin, M., & Mérillat, S. (2020). Brain structure and cognitive ability in healthy aging: a review on longitudinal correlated change. *Reviews in the Neurosciences*, *31*(1), 1-57. https://doi.org/doi:10.1515/revneuro-2018-0096

Park, D. C., & Reuter-Lorenz, P. (2009). The Adaptive Brain: Aging and Neurocognitive Scaffolding. *Annual Review of Psychology*, *60*(Volume 60, 2009), 173-196. https://doi.org/https://doi.org/10.1146/annurev.psych.59.103006.093656

Park, H.-J., & Friston, K. (2013). Structural and Functional Brain Networks: From Connections to Cognition. *Science*, *342*(6158), 1238411. https://doi.org/doi:10.1126/science.1238411

Pascual-Leone, A., Amedi, A., Fregni, F., & Merabet, L. B. (2005). The plastic human brain cortex. *Annual Review of Neuroscience*, *28*(1), 377-401. https://doi.org/10.1146/annurev.neuro.27.070203.144216

Patel, R., Mackay, C. E., Jansen, M. G., Devenyi, G. A., O'Donoghue, M. C., Kivimaki, M., Singh-Manoux, A., Zsoldos, E., Ebmeier, K. P., Chakravarty, M. M., & Suri, S. (2022). Inter- and intra-individual variation in brain structural-cognition relationships in aging. *Neuroimage*, *257*, 119254. https://doi.org/10.1016/j.neuroimage.2022.119254

Pearson, J., & Keogh, R. (2019). Redefining Visual Working Memory: A Cognitive-Strategy, Brain-Region Approach. *Current Directions in Psychological Science*, *28*(3), 266-273. https://doi.org/10.1177/0963721419835210

Poldrack, R. A. (2006). Can cognitive processes be inferred from neuroimaging data? *Trends in*
Cognitive Sciences, *10*(2), 59-63.
https://doi.org/https://doi.org/10.1016/j.tics.2005.12.004

Postle, B. R. (2015). The cognitive neuroscience of visual short-term memory. *Current Opinion in*
Behavioral Sciences, *1*, 40-46.
https://doi.org/https://doi.org/10.1016/j.cobeha.2014.08.004

Power, J. D., Cohen, A. L., Nelson, S. M., Wig, G. S., Barnes, K. A., Church, J. A., Vogel, A. C.,
Laumann, T. O., Miezin, F. M., Schlaggar, B. L., & Petersen, S. E. (2011). Functional
network organization of the human brain. *Neuron*, *72*(4), 665-678.
https://doi.org/10.1016/j.neuron.2011.09.006

Price, C. J., & Friston, K. J. (2002). Degeneracy and cognitive anatomy. *Trends in Cognitive*
Sciences, *6*(10), 416-421.

Ritakallio, L., Fellman, D., Salmi, J., Jylkkä, J., & Laine, M. (2024). Self-reported strategy use in
working memory tasks. *Scientific Reports*, *14*(1), 4893. https://doi.org/10.1038/s41598-
024-54160-3

Rose, N. S. (2020). The Dynamic-Processing Model of Working Memory. *Current Directions in*
Psychological Science, *29*(4), 378-387. https://doi.org/10.1177/0963721420922185

Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity: uses and
interpretations. *Neuroimage*, *52*(3), 1059-1069.
https://doi.org/10.1016/j.neuroimage.2009.10.003

Salami, A., Eriksson, J., & Nyberg, L. (2012). Opposing effects of aging on large-scale brain
systems for memory encoding and cognitive control. *J Neurosci*, *32*(31), 10749-10757.
https://doi.org/10.1523/jneurosci.0278-12.2012

Salami, A., Garrett, D. D., Wåhlin, A., Rieckmann, A., Papenberg, G., Karalija, N., Jonasson, L.,
Andersson, M., Axelsson, J., Johansson, J., Riklund, K., Lövdén, M., Lindenberger, U.,
Bäckman, L., & Nyberg, L. (2019). Dopamine $D_{2/3}$ Binding Potential
Modulates Neural Signatures of Working Memory in a Load-Dependent Fashion. *The*
Journal of Neuroscience, *39*(3), 537-547. https://doi.org/10.1523/jneurosci.1493-
18.2018

Salami, A., Rieckmann, A., Fischer, H., & Bäckman, L. (2014). A multivariate analysis of age-
related differences in functional networks supporting conflict resolution. *Neuroimage*,
86, 150-163. https://doi.org/10.1016/j.neuroimage.2013.08.002

Salami, A., Rieckmann, A., Karalija, N., Avelar-Pereira, B., Andersson, M., Wåhlin, A., Papenberg,
G., Garrett, D. D., Riklund, K., Lövdén, M., Lindenberger, U., Bäckman, L., & Nyberg, L.
(2018). Neurocognitive Profiles of Older Adults with Working-Memory Dysfunction.
Cerebral Cortex, *28*(7), 2525-2539. https://doi.org/10.1093/cercor/bhy062

Samu, D., Campbell, K. L., Tsvetanov, K. A., Shafto, M. A., & Tyler, L. K. (2017). Preserved cognitive
functions with age are determined by domain-dependent shifts in network responsivity.
Nature communications, *8*(1), 14743.

Sander, M. C., Lindenberger, U., & Werkle-Bergner, M. (2012). Lifespan age differences in
working memory: A two-component framework. *Neuroscience & Biobehavioral*
Reviews, *36*(9), 2007-2033.
https://doi.org/https://doi.org/10.1016/j.neubiorev.2012.06.004

Sanfratello, L., Caprihan, A., Stephen, J. M., Knoefel, J. E., Adair, J. C., Qualls, C., Lundy, S. L., &
Aine, C. J. (2014). Same task, different strategies: How brain networks can be influenced
by memory strategy. *Human Brain Mapping*, *35*(10), 5127-5140.
https://doi.org/https://doi.org/10.1002/hbm.22538

Scimeca, J. M., Kiyonaga, A., & D'Esposito, M. (2018). Reaffirming the Sensory Recruitment
Account of Working Memory. *Trends in Cognitive Sciences*, *22*(3), 190-192.
https://doi.org/10.1016/j.tics.2017.12.007

Scrucca, L., Fop, M., Murphy, T. B., & Raftery, A. E. (2016). mclust 5: Clustering, Classification and
Density Estimation Using Gaussian Finite Mixture Models. *R j*, *8*(1), 289-317.

Seeley, W. W., Menon, V., Schatzberg, A. F., Keller, J., Glover, G. H., Kenna, H., Reiss, A. L., &
Greicius, M. D. (2007). Dissociable Intrinsic Connectivity Networks for Salience
Processing and Executive Control. *The Journal of Neuroscience*, *27*(9), 2349-2356.
https://doi.org/10.1523/jneurosci.5587-06.2007

Seghier, M. L., Lee, H. L., Schofield, T., Ellis, C. L., & Price, C. J. (2008). Inter-subject variability in
the use of two different neuronal networks for reading aloud familiar words.
Neuroimage, *42*(3), 1226-1236.
https://doi.org/https://doi.org/10.1016/j.neuroimage.2008.05.029

Seghier, M. L., & Price, C. J. (2016). Visualising inter-subject variability in fMRI using threshold-
weighted overlap maps. *Sci Rep*, *6*, 20170. https://doi.org/10.1038/srep20170

Seghier, M. L., & Price, C. J. (2018). Interpreting and Utilising Intersubject Variability in Brain
Function. *Trends in Cognitive Sciences*, *22*(6), 517-530.
https://doi.org/https://doi.org/10.1016/j.tics.2018.03.003

Serences, J. T. (2016). Neural mechanisms of information storage in visual short-term memory.
Vision Res, *128*, 53-67. https://doi.org/10.1016/j.visres.2016.09.010

Shafto, M. A., Tyler, L. K., Dixon, M., Taylor, J. R., Rowe, J. B., Cusack, R., Calder, A. J., Marslen-
Wilson, W. D., Duncan, J., Dalgleish, T., Henson, R. N., Brayne, C., & Matthews, F. E.
(2014). The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) study protocol:
a cross-sectional, lifespan, multidisciplinary examination of healthy cognitive ageing.
BMC Neurol, *14*, 204. https://doi.org/10.1186/s12883-014-0204-1

Shekari, E., & Nozari, N. (2023). A narrative review of the anatomy and function of the white
matter tracts in language production and comprehension. *Front Hum Neurosci*, *17*,
1139292. https://doi.org/10.3389/fnhum.2023.1139292

Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., Filippini, N., Watkins,
K. E., Toro, R., Laird, A. R., & Beckmann, C. F. (2009). Correspondence of the brain's
functional architecture during activation and rest. *Proc Natl Acad Sci U S A*, *106*(31),
13040-13045. https://doi.org/10.1073/pnas.0905267106

Stokes, M. G. (2015). 'Activity-silent' working memory in prefrontal cortex: a
dynamic coding framework. *Trends in Cognitive Sciences*, *19*(7), 394-405.
https://doi.org/10.1016/j.tics.2015.05.004

Tavor, I., Parker Jones, O., Mars, R. B., Smith, S. M., Behrens, T. E., & Jbabdi, S. (2016). Task-free
MRI predicts individual differences in brain activity during task performance. *Science*,
352(6282), 216-220. https://doi.org/10.1126/science.aad8127

Taylor, J. R., Williams, N., Cusack, R., Auer, T., Shafto, M. A., Dixon, M., Tyler, L. K., Cam, C., &
Henson, R. N. (2017). The Cambridge Centre for Ageing and Neuroscience (Cam-CAN)
data repository: Structural and functional MRI, MEG, and cognitive data from a cross-
sectional adult lifespan sample. *Neuroimage*, *144*(Pt B), 262-269.
https://doi.org/10.1016/j.neuroimage.2015.09.018

Vermunt, J. K., & Magidson, J. (2002). Latent class cluster analysis. *Applied latent class analysis*,
11(89-106), 60.

Von Der Heide, R. J., Skipper, L. M., Klobusicky, E., & Olson, I. R. (2013). Dissecting the uncinate
fasciculus: disorders, controversies and a hypothesis. *Brain*, *136*(Pt 6), 1692-1707.
https://doi.org/10.1093/brain/awt094

Walhovd, K. B., Lövden, M., & Fjell, A. M. (2023). Timing of lifespan influences on brain and
cognition. *Trends in Cognitive Sciences*, *27*(10), 901-915.
https://doi.org/10.1016/j.tics.2023.07.001

Westlin, C., Theriault, J. E., Katsumi, Y., Nieto-Castanon, A., Kucyi, A., Ruf, S. F., Brown, S. M.,
Pavel, M., Erdogmus, D., & Brooks, D. H. (2023). Improving the study of brain-behavior
relationships by revisiting basic assumptions. *Trends in Cognitive Sciences*.

Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman, J.
L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fischl, B., Liu, H., & Buckner, R. L. (2011). The
organization of the human cerebral cortex estimated by intrinsic functional connectivity.
J Neurophysiol, *106*(3), 1125-1165. https://doi.org/10.1152/jn.00338.2011

Zuo, N., Salami, A., Yang, Y., Yang, Z., Sui, J., & Jiang, T. (2019). Activation-based association
profiles differentiate network roles across cognitive loads. *Human Brain Mapping*, *40*(9),
2800-2812. https://doi.org/https://doi.org/10.1002/hbm.24561

1154