**Revisiting the paradigm of anhematophagy in male mosquitoes**

Jovana Bozic[1,2,3¶,] Renuka E. Joseph[1,2,3¶,] Tse-Yu Chen[1,2,3], Margaret V. Becker[4], Rachel S. Krizek[2,3,5], Amber Holley[6], Renee L.M.N. Ali[4], Miguel Á. Miranda[7,8], Carlos Barceló[7], Mikel A. González[9], Maureen Laroche[10,11,12], Rafael Gutiérrez-López[13], Renke Lühken[14], Douglas E. Norris[4], Perran A. Ross[15], Joshua B. Benoit[16], and Jason L. Rasgon[1,2,3,5*]

1. Department of Entomology, The Pennsylvania State University, University Park, PA, USA
2. The Center for Infectious Disease Dynamics, The Pennsylvania State University, University Park, PA, USA
3. The Huck Institutes of the Life Sciences, The Pennsylvania State University, University Park, PA, USA
4. The W. Harry Feinstone Department of Molecular Microbiology and Immunology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD, USA
5. Department of Biochemistry and Molecular Biology, The Pennsylvania State University, University Park, PA, USA
6. Department of Pathology, The University of Texas Medical Branch, Galveston, TX, USA
7. Applied Zoology and Animal Conservation Group, University of the Balearic Islands, Palma, Spain
8. Instituto de Investigaciones Agroambientales y de Economía del Agua (INAGEA), University of the Balearic Islands, Palma, Spain
9. Departamento de Soluciones ambientales y Entomología. Grupo SASTI, Spain
10. Department of Microbiology & Immunology, School of Medicine, The University of Texas Medical Branch, Galveston, TX, USA
11. Department of Population Health & Health Disparities, School of Public and Population Health, The University of Texas Medical Branch, Galveston, TX, USA
12. Clima, Latin American Center of Excellence for Climate Change and Health, Universidad Peruana Cayetano Heredia (UPCH), Lima-Peru.
13. Centro Nacional de Microbiología, Instituto de Salud Carlos III (ISCIII), CIBERINFEC. Madrid, Spain.
14. Bernhard Nocht Institut for Tropical Medicine, Hamburg, Germany
15. School of BioSciences, Bio21 Institute, The University of Melbourne, Parkville, VIC 3010, Australia
16. Department of Biological Sciences, University of Cincinnati, Cincinnati, OH, USA

¶ These authors contributed equally to this work

* email for correspondence: jlr54@psu.edu

**Abstract**

Most female mosquitoes are reproductively obligate feeders on vertebrate blood to
obtain nutrients required for egg production (driving transmission of vector-borne
pathogens in the process), and which rely on plant sugars for their non-reproductive
energy requirements. Male mosquitoes, on the other hand, are thought to rely
exclusively on plant sugars for their energetic needs; indeed, this dichotomy is one of
the central paradigms of medical entomology. Here, we provide multiple lines of
evidence challenging this assumption. We show that when reared under
dehydration/starvation conditions, male mosquitoes will readily take blood from a
membrane feeder while CRISPR mutants with impaired humidity detection do not
increase their bloodfeeding rates. For multiple mosquito species, dehydrated males are
attracted to human hosts and attempt to probe. We also observed several instances
where dehydrated probing males were able to successfully pierce vertebrate host skin
and acquire blood from a vertebrate host. When fed a blood meal containing West Nile
virus, male *Cx. tarsalis* mosquitoes became infected with and orally transmitted the
pathogen at rates and titers equivalent to females. Finally, we collected wild, putatively
blood-fed male mosquitoes from Texas, USA and Mallorca, Spain, and identified
vertebrate DNA in these samples (canine and human, respectively). These data
suggest that under specific circumstances male mosquitoes may be able to probe
and/or ingest blood and transmit pathogens to vertebrate hosts. These results are a
paradigm shift in our understanding of male mosquito biology and suggest they may be
more directly involved in pathogen transmission cycles than previously recognized.

**Key words:** *Culex tarsalis*, *Culex quinquefasciatus*, *Aedes aegypti*, *Aedes notoscriptus*,
*Anopheles stephensi*, mosquito, bloodfeeding, sugar feeding, dehydration, vector-borne
disease

**Introduction**

Female mosquitoes (except those which are autogenous) are reproductively obligate bloodfeeders, feeding on vertebrate blood to obtain nutrients required for egg production, and relying on plant sugars for their non-reproductive energy requirements. In contrast, male mosquitoes, on the other hand, are thought to rely exclusively on plant sugars for their energetic needs [1]. Indeed, this difference is one of the central tenants of medical entomology; female mosquitoes bloodfeed, males do not. Female bloodfeeding facilitates transmission of blood-borne pathogens, such as viruses or parasites, between vertebrate hosts, which is why the majority of mosquito research is performed on females rather than males; even when males are studied, it is usually within the context of how they affect females (mating behavior and fertility, pathogen transmission modulation, etc.) [2-5].

Female mosquitoes evolved to feed on blood, and this adaptation is reflected in the biology of their midgut, where transcripts related to blood digestion are enriched in the female compared to the male [6]; one would expect that male mosquitoes should not be attracted to blood as a nutrition source as they are thought to lack the proper physiology to digest and process it. However, there is one interesting report in the literature where male mosquitoes were attracted to and fed on blood. Nikbakhtzadeh and colleagues [7] documented bloodfeeding behavior in a laboratory colony of the mosquito *Culex quinquefasciatus.* When presented with defibrinated sheep blood on a cotton pledget (and to a much less efficient extent, a Parafilm membrane), male mosquitoes took a bloodmeal. However, in this study blood was toxic to male mosquitoes, which died in a dose-dependent manner when blood was mixed with sugar [7], consistent with physiological adaptations to sugar vs bloodfeeding in males vs. females [6]. Interestingly, males did not show a preference for sugar compared to blood in a dual-choice assay [7], and the reason they fed on blood at all, particularly as it was toxic, remains an open question. As this is (to our knowledge) the only published observation of male mosquito bloodfeeding behavior, it is difficult to speculate. However, there are a multitude of observations that male mosquitoes are attracted to human host odors and

this behavior is suppressed by mosquito repellents [8], which includes species from
arguably the three most important mosquito genera that act as pathogen vectors to
humans (*Anopheles*, *Culex*, and *Aedes*).

Here, we present studies on male bloodfeeding behavior in the mosquitoes *Cx. tarsalis,*
*Cx. quinquefasciatus, Ae. aegypti, Ae. notoscriptus,* and *An. stephensi*. *Cx. tarsalis* is
one of the major West Nile virus (WNV) vectors in North America, where it is widely
distributed across the western United States [9]. It is genetically diverse, generally feeds
on birds in the wild, and can be facultatively autogenous [9-10]. After becoming infected
with WNV during a bloodfeeding event, it can also transmit the virus vertically to
offspring at relatively high rates [11-12]. *Ae. aegypti* is one of the major invasive
arbovirus vectors in the world [13]. *An. stephensi* is a major vector of malaria in Asia,
and has been recently expanded its range into Africa [14]. *Cx. quinquefasciatus* is a
major arbovirus vector with a cosmopolitan distribution [15], while *Ae. notoscriptus* is a
vector of Ross River virus in Australia [16]. We opportunistically observed *Cx. tarsalis*
and *Ae. aegypti* males taking blood during unrelated laboratory studies, and undertook
experiments to document and understand the behavior in multiple species. We found
that when dehydrated, male mosquitoes will predictably take blood from a membrane
feeder, determined the physiological mechanisms driving male bloodfeeding behavior,
demonstrated that some males were capable of obtaining a bloodmeal from a
vertebrate host through biting, and present results of experiments examining potential
for male mosquitoes to be involved in pathogen transmission cycles. We also collected
bloodfed male mosquitoes in the wild and identified the hosts that they fed upon. These
results are a paradigm shift in our understanding of male mosquito biology and suggest
they may be more directly involved in pathogen transmission cycles than previously
recognized.

**Methods**
Human and animal subjects approvals: Human landing experiments with *Cx. tarsalis*
used the senior author (JLR) under PSU IRB Exempt Protocol STUDY00024284.
Human landing experiments with *Aedes* mosquitoes and *Cx. quinquefasciatus* used

author PAR under University of Melbourne Human Ethics committee approval 0723847.

Experiments with mice were conducted under JHU IACUC protocol MO21H10.

<u>Mosquitoes</u>: *Cx. tarsalis* (KNWR strain), *Cx. quinquefasciatus* (sourced from

Queensland, Australia), *Ae. aegypti* (Liverpool, IR93a mutant, or Cairns strains), *Ae.*

*notoscriptus* (sourced from Queensland, Australia), and *An. stephensi* (Liston strain)

were maintained at 25°C, 16:8 h light:dark diurnal cycle with 80% relative humidity, with

10% sucrose solution provided at all times through a cotton wick. For general rearing,

mosquitoes were provided with expired anonymous human blood through a water-

jacketed glass membrane (Parafilm) feeder or a Hemotek feeder for egg development.

<u>Male dehydration</u>: To stimulate bloodfeeding, male mosquitoes were held at 25°C, 75%

RH without sugar or water for 24 hours [17].

<u>Survival analysis</u>: Bloodfed male *Cx. tarsalis* mosquitoes were isolated and placed into

cup cages, held at the previously described standard insectary conditions, and provided

with a cotton ball soaked in 10% sucrose solution. Control males were non-bloodfed

and were maintained under the same conditions. Dead mosquitoes were counted every

day and removed from the cages. Significant differences in survival between mosquito

groups was determined with Kaplan-Meier analysis using GraphPad Prism version

9.0.4.

<u>Dehydration and bloodfeeding behavior in male *Cx. tarsalis*</u>: *Cx. tarsalis* males were

reared conventionally (80% RH, with free access to 10% sucrose solution in water), or

under dehydrating conditions as described above, then were offered a bloodmeal

through a membrane feeder for 30 minutes. The number of fed and unfed mosquitoes at

the end of the feeding period were counted. Data were analyzed by Fishers Exact test.

<u>Ionotropic receptor 93a (Ir93a) mutant mosquito assays:</u> We obtained an *Ae. aegypti*

line that was a CRISPR knockout mutant for the Ir93a gene, which inhibits their ability to

sense humidity [18]. The mutation was introgressed into the wild-type Liverpool

background for comparison with Liverpool controls, and both lines reared as described above. For experiments, at 5-6 days post-emergence, males of each strain were transferred to 10 x 10 x 10 cm cages and deprived of sucrose and water (or held at normal conditions as controls) for 24 hours before being offered an anonymous human bloodmeal (Innovative Research) using an artificial feeding system (Hemotek). Bloodfeeding rates for each genotype and condition were recorded. Data were analyzed by Fishers Exact tests and confidence intervals calculated from the binomial distribution.

Landing and probing experiments: Cages of 50 male mosquitoes (*Cx. tarsalis*, *Cx. quinquefasciatus*, *Ae. notoscriptus,* or wild-type *Ae. aegypti*) were reared under standard or dehydrating insectary conditions and were allowed to probe on the hand of JLR (*Cx. tarsalis*) or PAR (*Ae. aegypti, Ae. notoscriptus,* and *Cx. quinquefasciatus*) for five minute trials. Mosquito landings (defined as a mosquito alighting on the volunteer hand for any period of time) and probing behavior (defined as exploring and probing with mouthparts) were counted during the 5 minute interval. Each experiment was repeated 6 times. Data were analyzed by Mann-Whitney U test.

Host bloodfeeding by male mosquitoes:

*Cx. tarsalis –* The senior author (JLR) had an unrelated small (3mm) scratch on their hand obtained from a pet cat a day earlier. A sterile razor blade was used to pick the scab off the scratch allowing a minimal amount of blood to be exposed. The wounded hand was placed in a cage of 20 dehydrated male *Cx. tarsalis* mosquitoes and their behavior recorded.

*An. stephensi –* Mosquitoes were reared at 27°C (± 1°C) with 75% ± 5% relative humidity and a 12-h day–night cycle by Johns Hopkins Malaria Research Institute Insectary Core facility. Male mosquitoes were transferred to cups (40-50 per cup) and held under dehydrating conditions for 24h. Prior to the feed the number of surviving mosquitoes was counted. Mosquitoes were allowed to feed on a sedated 8 week old female Swiss Webster mouse for 15 mins. The mouse was placed belly down on the top of the cup. No wound was created, and the fur was not shaved. Cups were

immediately frozen after the attempted feed and mosquitoes screened under a
microscope for the presence of a blood meal.

WNV feeds: Dehydrated *Cx. tarsalis* males and female controls were allowed access to
an infectious blood meal consisting of a 1:1 mix of anonymous human blood spiked with
$5.0 \times 10^7$ FFU/ml (focus-forming units/ml) of WN02-1956 (GenBank: AY590222). A
subset of male and females were processed immediately after feeding ("day zero") to
check for virus viability. Mosquito virus infection and transmission assays were
performed at 7 and 14 days post-blood feeding. Fully engorged mosquitoes were sorted
from non-fed ones for analysis. Mosquitoes were anesthetized with triethylamine
(Sigma, St. Louis, MO), legs/wings from each mosquito were removed and placed
separately in a 2-mL tube filled with 0.5 mL mosquito diluent (MD: 20% heat-inactivated
fetal bovine serum (FBS) in Dulbecco's phosphate-buffered saline, 50 µg/mL
penicillin/streptomycin, 50 ug/mL gentamicin, and 2.5 µg/mL fungizone, with a sterile 2.0
mm stainless steel bead (Next Advance, Inc. Innovative Lab Products for the Life
Sciences). The proboscis of each mosquito was positioned in a tapered capillary tube
containing approximately 10 µL of a 1:1 solution of 50% sucrose and FBS to induce
salivation. After 30 min, the tube contents were expelled into 0.3 mL MD, and bodies
were placed individually into a 2-mL tube filled with 0.5 mL MD and a stainless steel
bead as described above. Mosquito bodies and legs/wings were homogenized for 30
sec with TissueLyser (QIAGEN, Hilden, Germany) at 24 cycles/sec, followed by
centrifugation for 1 min. Mosquito bodies, legs/wings, and salivary secretion samples
were tested for live, infectious WNV using focus-forming assays (FFAs; see below).

WNV FFAs: WNV titers were quantified by FFA, which detects live, infectious virus.
C6/36 cells were seeded into 96-well plates at a density of $1 \times 10^5$ cells/well and
incubated overnight at 28°C in complete RPMI medium without $CO_2$. The next day,
medium was removed from the wells. Samples from male and female bodies or
legs/wings were serially diluted in a serum-free RPMI medium; saliva samples were
undiluted. 30 µL of each sample was added in duplicate to the prepared C6/36 cells.

Cells were incubated for 1 hour at 28°C without $CO_2$, after which the inoculum was
removed. 100 µL of RPMI containing 0.8% methylcellulose was added to limit viral
spread. Infected cells were incubated for 48 hours at 28°C without $CO_2$. At 48 hours
post-infection, infected C6/36 cells were fixed with 50 µL 4% formaldehyde for 30
minutes at room temperature (RT). Cells were washed, permeabilized with 0.2% triton-
X, and blocked with 3% BSA. 30 µL monoclonal flavivirus antibody (Clone D1-4G2-4-15,
BEI-resources, NR-50327) was added and incubated overnight at 4°C. After washing,
30µl of fluorescent secondary antibody (1:1000 dilution; Goat anti-Mouse IgG (H+L)
Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 488, Invitrogen/Thermo
Fischer, A-11029) was added and incubated overnight at 4°C. Cells were maintained in
100 µL PBS to prevent drying. WNV foci were imaged using a FITC filter on an Olympus
BX41 microscope with a UPlanFI 4x objective and counted. Infection rate (IR) was
defined as the proportion of mosquitoes exposed to virus that had WNV-positive bodies.
Dissemination rate (DR) was defined as the proportion of mosquitoes with WNV-positive
bodies that had WNV-positive legs/wings. Transmission rate (TR) was defined as the
proportion of mosquitoes with WNV-positive legs/wings that had WNV-positive saliva.
IR, DR, and TR were analyzed with Fisher's exact tests. Viral titers were analyzed using
Mann-Whitney U tests.

<u>Wild mosquito collection, DNA extraction and molecular detection and identification of</u>
<u>vertebrate DNA:</u> Mosquitoes were collected using BG-Sentinel traps (Biogents) [19-20]
on Galveston Island, in South Texas, August 12-13, 2024, and in Mallorca, Spain, July
9, 2020. Fresh mosquitoes were sexed, morphologically identified the same day and
placed at -20°C until processed. For Texas samples, after identification, whole male
mosquitoes were individually incubated with 20µL of proteinase K and 180µL of G2
buffer (Qiagen) at 56°C for one hour then were homogenized using tungsten beads and
a TissueLyser (Qiagen) for one minute at 30 cycles/sec. After a quick spin to pellet
debris, DNA was extracted from the lysate using a Qiagen EZ1 automated DNA
extraction instrument and reagents from the associated EZ1&2 DNA Tissue Kit. DNA
samples were screened for vertebrate DNA using vertebrate-specific qPCR assay
targeting a fragment of the 16s rRNA on a CFX Duet PCR system (Bio-Rad) [21].

Human blood obtained from the University of Texas Medical Branch Blood Bank was
used as a positive control and PCR mix without DNA was used as a negative control.
Samples were analyzed in duplicate and the entire assay repeated twice. Samples
positive for vertebrate DNA had a 690 bp sequence of the COI gene amplified by
primers ModRepCOIF and ModRepCOIR [22], the amplicons cloned using the NEB
PCR cloning kit (New England Biolabs, Ipswich, MA), sequenced, and identity
confirmed by BLAST analysis. For Mallorca samples, whole engorged specimens were
placed into a 2 mL tube, along with 20 2.0 mm zirconia beads (BioSpec Products,
Bartlesville, USA) and 1 mL of cell culture medium (high-glucose Dulbecco's modified
Eagle's medium; Sigma-Aldrich, St. Louis, MO, USA) and homogenized with a
TissueLyser (Qiagen, Hilden, Germany) for 2 minutes at 50 oscillations per second.
Following clarification through centrifugation for 1 minute at 8000 rpm and 4°C, the
resulting suspension was transferred to a new safe-lock tube. DNA was extracted from
200 µL of the homogenate using the KingFisher™ Flex Magnetic Particle Processor
with the MagMAX™ Pathogen Ribonucleic Acid/DNA Kit (both Thermo Fisher Scientific,
Waltham, MA, USA). A primer set targeting the 16S rRNA gene was used following
previously published protocols [23]. All amplicons were directly sequenced (LGC
Genomics, Berlin, Germany).

Conceptual replication: As we are aware that the results presented here are
controversial, we endeavored to have multiple independent laboratories conduct the
experiments to test the generality of the observations. Work with *Cx. tarsalis* was
performed at Penn State; work with mutant *Ae. aegypti* was conducted at the University
of Cincinnati; work with *Ae. notoscriptus, Cx. quinquefasciatus,* and wild-type *Ae.*
*aegypti* was conducted at the University of Melbourne; work with *An. stephensi* was
conducted at the Johns Hopkins Bloomberg School of Public Health.

**Results**

Bloodfeeding is not toxic to *Cx. tarsalis* males: While we were bloodfeeding during an
experiment related to relative humidity (e.g. [24]), we noted incidentally that in addition

to females, male mosquitoes were probing the membrane and were taking blood
(Figure 1A,B). As this was a spontaneous occurrence, the total number of males in the
cage was not recorded but was on the order of 50-70 based on standard rearing
practices in our lab. Out of this total, we isolated seven blood-engorged males. These
males were placed into a survival cup and a survival experiment conducted, comparing
their survival to 10 non-bloodfed males from the same initial cage. Although
Nikbakhtzadeh et al. [7] demonstrated that blood was highly toxic to male *Cx.*
*quinquefasciatus* in laboratory studies, we did not observe any acute toxicity to blood in
male *Cx. tarsalis*; indeed, survival in bloodfed males was marginally (although not
statistically) higher than non-bloodfed males (Figure 1C).

*Cx. tarsalis* male bloodfeeding is driven by dehydration: As we previously demonstrated
that dehydration stimulates elevated bloodfeeding behavior in females [17, 24-25], we
tested the hypothesis that dehydration was driving bloodfeeding behavior in males.
Cages of male mosquitoes were reared under conventional insectary conditions or
under dehydrating conditions [17]. No conventionally reared male mosquito (N = 64)
took a bloodmeal from the membrane feeder, while 44/163 dehydrated males took a
bloodmeal ($P < 0.00001$).

Male mosquito bloodfeeding behavior is dependent on their ability to sense humidity:
Mosquitoes sense humidity through ionotropic receptor Ir93a, by which they locate
oviposition sites, and CRISPR Ir93a knock-out mutants are impaired in this behavior
[18]. We noted in further experiments that male *Ae. aegypti* mosquitoes would also take
blood from a membrane feeder, so we used an available Ir93a *Ae. aegypti* KO mutant
for these assays [18]. When reared under standard insectary conditions, bloodfeeding
rates did not differ statistically between wild-type and mutant mosquitoes. However,
when reared under dehydrating conditions, bloodfeeding rates for the mutant did not
increase, while wild-type mosquitoes had significantly elevated bloodfeeding behavior
($P = 0.0284$) (Figure 2).

<u>Dehydrated male mosquitoes will probe the hand of a human volunteer:</u> The hand of a
human volunteer was exposed to cages of conventionally reared or dehydrated male
mosquitoes (*Cx. tarsalis, Cx. quinquefasciatus, Ae. aegypti, and Ae. notoscriptus*).
Conventionally reared males showed little interest in the host, with infrequent landings
that lasted less than 5 seconds. In all trials, only one conventionally reared mosquito
(*Ae. aegypti*) demonstrated probing behavior. In contrast, for all tested species
dehydrated males landed significantly more often on the hand of the volunteer, most
landings lasted until the end of the time period, and probing behavior was observed in
the majority of landings (Figures 3-4, Supplementary Videos 1-9). One dehydrated *Cx.*
*tarsalis* male mosquito (out of 6 separate trials) was able to pierce the skin of the
volunteer at the base of the wrist, although it was unable to reach the capillaries and
acquire a bloodmeal (Figure 4, Supplementary Video 10). The bite resulted in a mild
immunogenic reaction that disappeared after approximately 10 minutes (Figure 4G,H).
To confirm that only males were in the cage, after the study was concluded the entire
cage was killed by freezing and every mosquito visually examined for the presence of a
female or a gynandromorph; only males were identified.
364
<u>Dehydrated male mosquitoes will take blood from vertebrate hosts:</u> Dehydrated male
mosquitoes show keen interest in probing a human host, but in these experiments were
not able to acquire blood, even from the single observed "successful" probing attempt.
We hypothesized that if blood was made more accessible, male mosquitoes would take
a bloodmeal. The senior author serendipitously had a small scratch on their hand
(acquired from a pet cat a day earlier). The scab was peeled back using a sterile razor
blade, exposing a small amount of blood. The volunteer placed their hand in a cage of
20 dehydrated *Cx. tarsalis* male mosquitoes. Males were attracted to the wound, and
wound probing behavior was observed by 5 males (Supplementary Video 11). One
male out of the 5 that probed fed and took a bloodmeal from the wound (Figure 5A-C).
At the conclusion of the experiment, the fed male was dissected to confirm the presence
of blood in the gut (Figure 5D).
377

Further experiments were conducted examining the ability for male *An. stephensi*
mosquitoes to take blood from a mouse host (which has thinner skin compared to a
human). *An. stephensi* males were more sensitive to the dehydration procedure
compared to *Aedes* or *Culex*, and out of the 2 replicate experiments most did not
survive the dehydration procedure (Rep 1: 34/39 dead; Rep 2: 46/50 dead). However,
out of the 9 surviving mosquitoes across both replicates, one mosquito (in replicate 1)
was able to bite the mouse and acquire a bloodmeal (Figure 5E). These data prove that
male mosquitoes do have the ability to pierce the skin of some vertebrates and can
canulate capillaries to acquire blood.

Male *Cx. tarsalis* mosquitoes are competent vectors for West Nile virus: Since we
determined that male mosquitoes will probe vertebrate hosts, can pierce skin, and
ingest blood, we asked the question: can male mosquitoes become infected with and
transmit pathogens? We offered dehydrated male *Cx. tarsalis* mosquitoes an artificial
bloodmeal spiked with WNV through a membrane feeder and assayed their virus vector
competence at day 7 and day 14 post-feeding. Female *Cx. tarsalis* were exposed to
virus at the same time as a control. We found that both female and male *Cx. tarsalis*
were able to become infected with, disseminate, and orally transmit virus; males
transmitted at both day 7 and 14, while females only had detectable virus in their saliva
at day 14. After adjusting for multiple comparisons, infection rates (IR), dissemination
rates (DR), and transmission rates (TR) did not differ statistically between males and
females at either timepoint (Table 1).

We quantitated all viral titers using an infectious virus assay. First, a subsample of
males and females were assayed immediately after feeding ("day zero") to confirm virus
viability. All fed males and females had detectable live infectious virus in their bodies,
although females had statistically higher viral titers ($P = 0.005$), likely because they
could physically ingest a larger volume of blood. At day 7 post-exposure, viral titers
were not statistically different between males and females in the bodies, the legs/wings,
or the saliva (Figure 6A). At day 14 post-exposure, females had higher viral titers in their
bodies ($P = 0.001$) and legs/wings ($P = 0.0083$) compared to males, suggesting either

greater viral replication rates, or simply more tissue available for virus replication due to

larger size. However, viral titers in saliva between males and females did not differ

statistically (Figure 6B).

<u>Vertebrate DNA and blood can be detected in wild-caught male mosquitoes:</u> While BG-

Sentinel traps are designed to collect female mosquitoes, they do also occasionally

catch males [e.g. 20-21]. During collections in Texas, we collected a total of 15 male

mosquitoes (ten *Ae. albopictus*, one *Ae. sollicitans,* and four *Cx. quinquefasciatus*).

Vertebrate DNA was not detected in *Ae. albopictus* or *Ae. sollicitans.* However, three of

the four collected *Cx. quinquefasciatus* males had detectable vertebrate DNA (Ct values

ranging from 15.55 to 18.36) (Supplementary Table 1). When homogenized during

extraction, the supernatants from these positive males were reddish-brown, possibly

suggesting the presence of blood in the specimen. We sequenced the vertebrate COI

sequences in these samples; all 3 hit canine (best hits - *Canis lupis familaris*, or

domestic dog) (Supplementary file 1). In collections in Mallorca, Spain, we collected a

male *Cx. pipiens* biotype *pipiens* mosquito with what appeared to be partially digested

blood in its midgut (Figure 7). We sequenced the vertebrate mitochondrial 16S RNA

gene sequences in this sample, which suggested the mosquito had fed on a human

host (Supplementary file 1).

**Discussion**

Previous work showed that in the laboratory, blood was toxic to male *Cx.*

*quinquefasciatus* mosquitoes [7], suggesting that in this species male bloodfeeding

seems to be a maladaptive trait. In our study, we demonstrate that males of multiple

species can tolerate bloodfeeding, and that male bloodfeeding behavior is driven by

water homeostasis during dehydration conditions. When mosquitoes cannot sense

humidity due to Ir93a mutagenesis, dehydration does not increase blood seeking

behavior. These results are consistent with the role of dehydration on bloodfeeding

behavior in female mosquitoes, where dehydration can stimulate females to increase

their bloodfeeding rates as well [17, 24-25] and thus may reflect an adaptive trait where

mosquitoes (female or male) can maximize their water intake during drought or periods

of low relative humidity if other sources (nectar or free water) are not available.

The mouthparts of male mosquitoes are thought to be physically incapable of

penetrating vertebrate skin; however, our data suggest that this paradigm may be

incomplete; in our experiments male mosquito mouthparts were proven adequate to

pierce a Parafilm membrane, as well as vertebrate (human and mouse) skin.

Dehydrated males were significantly attracted to and actively probed the hand of a

human host, and one individual was even able to penetrate the outer epidermis, leading

to a transitory immune reaction (Supplementary Video 10 and Figure 4G). As the saliva

of males differs from that of females, lacking various proteins needed for

immunomodulation and bloodmeal acquisition [26], and it is likely that very little saliva

was transferred compared to the bite of a female mosquito, it is not surprising that the

host immune reaction was mild and rapidly resolving. When allowed access to a wound,

dehydrated *Cx. tarsalis* male mosquitoes readily probed the wound and one took a

bloodmeal. As this experiment was facilitated by the fortuitous presence of a pre-

existing wound on the hand of the senior author, it could not be deliberately repeated

(as we were not allowed to make a deliberate wound due to IRB concerns). Further

experiments using *An. stephensi* and a mouse model demonstrated that male

mosquitoes can and will bite vertebrate hosts and can canulate capillaries to acquire

blood. In aggregate, these experiments suggest that male mosquitoes do have the

ability to take blood from vertebrate hosts under specific circumstances and that

although not necessary, feeding may be facilitated by the presence of host wounds.

According to fossil evidence, male mosquitoes are thought to once have had the ability

to feed on vertebrate blood, and to have lost this ability over evolutionary time [27]. It is

possible that the neural circuitry regulating host seeking and bloodfeeding behavior may

still be conserved among male mosquitoes, or alternatively that this is simply a unique

response to dehydration conditions to allow replenishment of water reserves, as a lack

of humidity detection observed in Ir93a mutants muted the blood feeding phenotype.

However, more in-depth functional neural studies are required to fully investigate this

phenomenon.

Interestingly our data demonstrate that, if male *Cx. tarsalis* orally acquire a WNV
infection, they are competent vectors and transmit the virus at similar rates and titers
compared to females. In our experiments we explicitly used an assay that quantified
live, infectious viral particles rather than quantitative PCR to rule out results that might
be due to carryover of non-infectious viral RNA. Our results suggest that male *Cx.*
*tarsalis* retain the receptors necessary for viral infection on their midgut, salivary glands,
and other body tissues. Intriguingly, there is a recent report of wild-caught male *Cx.*
*pipiens* mosquitoes that were infected with WNV from accidently feeding on infected
blood droplets during a laboratory assay [28]. *Cx. tarsalis* males (as well as *Cx. pipiens*
and other *Culex* mosquitoes) can also be infected with WNV due to vertical
transmission from infected mothers [11, 29-30], and it is possible that the ability for
males to become infected with virus orally is a consequence of their retaining receptors
necessary for vertical infection.

Finally, there is the question "is bloodfeeding behavior by male mosquitoes
epidemiologically significant?"  It is already known that male mosquitoes can be
indirectly important for vector-borne disease transmission dynamics. For example,
mating can affect key physiological parameters in females related to pathogen infection
and transmission [2]. More directly, in some species including *Cx. tarsalis* and WNV,
male mosquitoes can be infected with arboviruses by vertical transmission from infected
mothers [11, 29-30]. Infected males can also transmit some viruses venereally to
females during mating where they can be transmitted to vertebrate hosts during feeding
[29-30]. Consistent collection of males using host-derived attractants suggest that males
are commonly found to move toward hosts [8], increasing the potential of male feeding
on host-derived fluids under specific conditions (dry periods with a lack of water and/or
sugar resources). In field collections, we were able to detect vertebrate DNA/blood in
wild-caught *Culex* males in two separate geographic locations, suggesting that male
mosquitoes do encounter vertebrate hosts for bloodfeeding, either through wounds or
possibly by direct biting. In total, our study suggests the possibility of edge cases where
male mosquitoes could be more directly implicated in pathogen transmission, where

males undergoing dehydrating or starvation conditions (for example, during drought) acquire virus through vertical transmission from infected mothers or by feeding on an infected vertebrate host, then transmit to a naïve host through feeding on an open wound or by probing the skin, as mosquitoes often transmit the bulk of virus when probing skin prior to actually taking a bloodmeal [31].

We must emphasize that while compelling, the ability for male mosquitoes to bloodfeed or transmit pathogens in nature have not been widely studied (we suspect that researchers have not rigorously examined these phenomena as the dogma in the field is that males do not bloodfed). However, our data suggest that the canonical role of males as non-bloodfeeders needs to be re-examined and their contribution to pathogen transmission explicitly quantified (even if minor), particularly in light of recent vector-borne disease control strategies that rely on the mass release of male mosquitoes into natural populations [32-35].

**References**

1.  Clements AN. The Biology of Mosquitoes, Volume 1. CABI Publishers; 1992.

2.  Dahalan FA, Churcher TS, Windbichler N, Lawniczak MKN. The male mosquito contribution towards malaria transmission: Mating influences the *Anopheles* female midgut transcriptome and increases female susceptibility to human malaria parasites. PLoS Pathog. 2019; 15:e1008063.

3.  Marcenac P, Shaw WR, Kakani EG, Mitchell SN, South A, Werling K, Marrogi E, Abernathy DG, Yerbanga RS, Dabiré RK, Diabaté A, Lefèvre T, Catteruccia F. A mating-induced reproductive gene promotes *Anopheles* tolerance to *Plasmodium falciparum* infection. PLoS Pathog. 2020;16:e1008908.

4.  Carraretto D, Soresinetti L, Rossi I, Malacrida AR, Gasperi G, Gomulski LM. Behavioural responses of male *Aedes albopictus* to different volatile chemical compounds. Insects. 2022;13:290.

5.  Peng D, Kakani EG, Mameli E, Vidoudez C, Mitchell SN, Merrihew GE, MacCoss MJ, Adams K, Rinvee TA, Shaw WR, Catteruccia F. A male steroid controls female sexual behaviour in the malaria mosquito. Nature. 2022;608:93.

6.  Warr E, Aguilar R, Dong Y, Mahairaki V, Dimopoulos G. Spatial and sex-specific dissection of the *Anopheles gambiae* midgut transcriptome. BMC Genomics. 2007;8:37.

7.  Nikbakhtzadeh MR, Buss GK, Leal WS. Toxic effect of blood feeding in male mosquitoes. Front Physiol. 2016;7:4.

8.  Paris V, Hardy C, Hoffmann AA, Ross PA. How often are male mosquitoes attracted to humans? R Soc Open Sci. 2023 Oct 25;10(10):230921.

9.  Venkatesan M, Rasgon JL. Population genetic data suggest a role for mosquito-mediated dispersal of West Nile virus across the western United States. Mol Ecol. 2010;19:157.

10.  Provost-Javier KN, Chen S, Rasgon JL. Vitellogenin gene expression in autogenous *Culex tarsalis*. Insect Mol Biol. 2010;19:423.

11. Goddard LB, Roth AE, Reisen WK, Scott TW. Vertical transmission of West Nile Virus by three California *Culex* (Diptera: Culicidae) species. J Med Entomol. 2003;40:743.

12. Nelms BM, Fechter-Leggett E, Carroll BD, Macedo P, Kluh S, Reisen WK. Experimental and natural vertical transmission of West Nile virus by California *Culex* (Diptera: Culicidae) mosquitoes. J Med Entomol. 2013;50:371.

13. Brady OJ, Hay SI. The Global Expansion of Dengue: How *Aedes aegypti* Mosquitoes Enabled the First Pandemic Arbovirus. Annu Rev Entomol. 2020 Jan 7;65:191-208.

14. Acosta AL, Castro MC, Laporta GZ, Conn JE, Sallum MAM. Future global distribution and climatic suitability of *Anopheles stephensi*. Sci Rep. 2025 Jul 1;15(1):22268.\

15. Farajollahi A, Fonseca DM, Kramer LD, Marm Kilpatrick A. "Bird biting" mosquitoes and human disease: a review of the role of *Culex pipiens* complex mosquitoes in epidemiology. Infect Genet Evol. 2011 Oct;11(7):1577-85.

16. Watson TM, Kay BH. Vector competence of *Aedes notoscriptus* (Diptera: Culicidae) for Ross River virus in Queensland, Australia. J Med Entomol. 1998 Mar;35(2):104-6.

17. Hagan RW, Didion EM, Rosselot AE, Holmes CJ, Siler SC, Rosendale AJ, Hendershot JM, Elliot KSB, Jennings EC, Nine GA, Perez PL, Rizlallah AE, Watanabe M, Romick-Rosendale LE, Xiao Y, Rasgon JL, Benoit JB. Dehydration prompts increased activity and blood feeding by mosquitoes. Sci Rep. 2018;8:6804.

18. Laursen WJ, Budelli G, Tang R, Chang EC, Busby R, Shankar S, Gerber R, Greppi C, Albuquerque R, Garrity PA. Humidity sensors that alert mosquitoes to nearby hosts and egg-laying sites. Neuron. 2023 Mar 15;111(6):874-887.e8.

19. Velo E, Balestrino F, Kadriaj P, Carvalho DO, Dicko A, Bellini R, Puggioli A, Petrić D, Michaelakis A, Schaffner F, Almenar D, Pajovic I, Beqirllari A, Ali M, Sino G, Rogozi E, Jani V, Nikolla A, Porja T, Goga T, Fălcuă E, Kavran M, Pudar D, Mikov O, Ivanova-Aleksandrova N, Cvetkovikj A, Akıner MM, Mikovic R, Tafaj L, Bino S, Bouyer J, Mamai

W. A Mark-release-recapture study to estimate field performance of imported radio-sterilized male *Aedes albopictus* in Albania. Front Bioeng Biotechnol. 2022 Aug 16;10:833698.

20. Staunton KM, Goi J, Townsend M, Ritchie SA, Crawford JE, Snoad N, Karl S, Burkot TR. Effect of BG-lures on the male *Aedes* (Diptera: Culicidae) sound trap capture rates. J Med Entomol. 2021 Nov 9;58(6):2425-2431.

21. Taniguchi K, Akutsu T, Watanabe K, Ogawa Y, Imaizumi K. A vertebrate-specific qPCR assay as an endogenous internal control for robust species identification. Forensic Sci Int Genet. 2022 Jan;56:102628.

22. Reeves LE, Gillett-Kaufman JL, Kawahara AY, Kaufman PE. Barcoding blood meals: New vertebrate-specific primer sets for assigning taxonomic identities to host DNA from mosquito blood meals. PLoS Negl Trop Dis. 2018 Aug 30;12(8):e0006767.

23. Kitano T, Umetsu K, Tian W, Osawa M. Two universal primer sets for species identification among vertebrates. Int J Legal Med. 2007 Sep;121(5):423-7.

24. Manzano-Alvarez J, Terradas G, Holmes CJ, Benoit JB, Rasgon JL. Dehydration stress and Mayaro virus vector competence in *Aedes aegypti*. J Virol. 2023 Dec 21;97(12):e0069523.

25. Holmes CJ, Chakraborty S, Ajayi OM, Unran MR, Frigard RA, Stacey CL, Susanto EE, Chen SC, Rasgon JL, DeGennaro MJ, Xiao Y, Benoit JB. Multiple blood feeding bouts in mosquitoes allow for prolonged survival and are predicted to increase viral transmission during dry periods. iScience. 2025 Feb;28:111760.

26. Lu S, Martin-Martin I, Ribeiro JM, Calvo E. A deeper insight into the sialome of male and female *Ochlerotatus triseriatus* mosquitoes. Insect Biochem Mol Biol. 2022 Aug;147:103800.

27. Azar D, Nel A, Huang D, Engel MS. The earliest fossil mosquito. Curr Biol. 2023 Dec 4;33(23):5240-5246.e2.

28. Linthout C, Martins AD, de Wit M, Delecroix C, Abbo SR, Pijlman GP, Koenraadt CJM. The potential role of the Asian bush mosquito *Aedes japonicus* as spillover vector for West Nile virus in the Netherlands. Parasit Vectors. 2024 Jun 17;17(1):262.

29. Schopen S, Labuda M, Beaty B. Vertical and venereal transmission of California group viruses by *Aedes triseriatus* and *Culiseta inornata* mosquitoes. Acta Virol. 1991:35:37.

30. Patrican LA, DeFoliart GR. *Aedes triseriatus* and La Crosse virus: similar venereal infection rates in females given the first bloodmeal immediately before mating or several days after mating. Am J Trop Med Hyg. 1987 May;36(3):648-52

31. Visser I, Koenraadt CJM, Koopmans MPG, Rockx B. The significance of mosquito
saliva in arbovirus transmission and pathogenesis in the vertebrate host. One Health.
2023 Feb 12;16:100506.

32. Bansal S, Lim JT, Chong CS, Dickens B, Ng Y, Deng L, Lee C, Tan LY, Kakani EG,
Yoong Y, Du Yu D, Chain G, Ma P, Sim S, Ng LC, Tan CH. Effectiveness of *Wolbachia*-
mediated sterility coupled with sterile insect technique to suppress adult *Aedes aegypti*
populations in Singapore: a synthetic control study. Lancet Planet Health. 2024
Sep;8(9):e617-e628.

33. Carvalho DO, McKemey AR, Garziera L, Lacroix R, Donnelly CA, Alphey L,
Malavasi A, Capurro ML. Suppression of a field population of *Aedes aegypti* in Brazil by
sustained release of transgenic male mosquitoes. PLoS Negl Trop Dis. 2015 Jul
2;9(7):e0003864.

34. Apte RA, Smidler AL, Pai JJ, Chow ML, Chen S, Mondal A, Sánchez C HM,
Antoshechkin I, Marshall JM, Akbari OS. Eliminating malaria vectors with precision-
guided sterile males. Proc Natl Acad Sci U S A. 2024 Jul 2;121(27):e2312456121.

35. Rasgon JL. Precision-guided tools for malaria control. Proc Natl Acad Sci U S A.
2024 Aug 6;121(32):e2411587121.

**Declarations**

Ethics approval and consent to participate: All experiments with a human volunteer

used JLR (PSU IRB Exempt Protocol STUDY00024284) or PAR (University of

Melbourne Human Ethics committee approval 0723847). Mouse experiments were

performed under Johns Hopkins University IACUC protocol MO21H10.

Acknowledgements: This research was supported by NIH/NIAID grant R01AI150251,

USDA Hatch Project 4769, a grant with the Pennsylvania Department of Health using

Tobacco Settlement Funds, and funds from the Dorothy Foehr Huck and J. Lloyd Huck

endowment to JLR, and NIH/NIAID grant R01AI148551 to JBB and JLR. RLMNA, MVB,

DEN were supported in part by the Bloomberg Philanthropies; MVB was additionally

supported by NIH/NIAID training grant 2T32AI138953-06A1. Mosquito collections in

Spain were funded by the *Aedes* Invasive Mosquitoes AIM COST Action CA17108

funded by COST (European Cooperation in Science and Technology). PAR was

supported by an Australian Research Council Discovery Early Career Researcher

Award (DE230100067) funded by the Australian Government. RL was funded by the Federal Ministry of Research, Technology and Space of Germany (Grant Number 01KI2022). We thank Ms. Francine McCullogh for laboratory assistance, Ms. Amelia Romo and Ms. Heather Engler for assistance with mosquito rearing, Mabel Tao and Cindy Kimura for assistance with preparing mice for mosquito feeds, Apeksha Warusawithana for assistance with probing experiments, and Paul Garrity for providing Ir93a mutant mosquitoes used in this study. Finally, we thank Jiji the cat for his participation in this research.

Authors' contributions: JB, REJ, RSK, AH, ML, PAR, RL, T-YC, MAM, CB, MAG, RG-L, RLMNA, MVB, DEN, JBB, and JLR conducted the research; JBB contributed materials and reagents; JB, REJ, PAR and JLR analyzed the data; JB, REJ, PAR, JBB, and JLR wrote the manuscript.

**Figure legends**

**Figure 1. Male *Cx. tarsalis* bloodfeeding behavior and survival**. A) mosquitoes congregating at and bloodfeeding from a paraffin membrane. Arrow points to male orienting toward the membrane. B) Blood-engorged male mosquito. Unengorged male can be seen in-frame. C) Survival curve of bloodfed vs. non-bloodfed male *Cx. tarsalis* mosquitoes. No significant difference was observed between treatments.

**Figure 2. CRISPR deletion of Ir93a ablates male mosquito bloodfeeding behavior under dehydration conditions.** When reared conventionally, both wild-type and mutant *Ae. aegypti* exhibit baseline levels of bloodfeeding behavior. When reared under dehydration conditions, wild-type males significantly increase bloodfeeding behavior but humidity-insensitive mutant mosquitoes do not. Confidence intervals were calculated from the binomial distribution. WT = wild-type.

**Figure 3. Host probing behavior of dehydrated male mosquitoes**. Top: Landing responses for dehydrated vs. conventionally reared ("standard") male mosquitoes. Bottom: Probing behavior for dehydrated vs. conventionally reared ("standard") male mosquitoes. ** = $P < 0.01$

**Figure 4.** Landing and probing behavior of dehydrated mosquitoes. A) Conventionally reared mosquitoes (*Cx. tarsalis*) ignore a human host, while B) dehydrated mosquitoes actively land on a human host. Dehydrated mosquitoes C) *Cx. tarsalis* D) *Ae. aegypti* E) *Ae. notoscriptus* F) *Cx. quinquefasciatus* actively probe after landing. G) Bite reaction 2 minutes post-probing (arrow) H) Immune reaction resolved by 10 minutes post-probing. See Supplementary videos 1-9 for complete behavioral responses of each mosquito species and Supplementary video 10 for biting behavior.

**Figure 5. Male mosquito bloodfeeding.** A) Male *Cx. tarsalis* feeding on an open wound. B) Close-up of *Cx. tarsalis* feeding behavior. C) Blood is observable in the male *Cx. tarsalis* mosquito gut. D) Blood in the male *Cx. tarsalis* mosquito gut was confirmed by dissection. E) Male *An. stephensi* mosquito with visible blood in the midgut after feeding on a mouse. See Supplementary Video 11 for complete *Cx. tarsalis* behavioral response.

**Figure 6. West Nile virus vector competence for male and female *Cx. tarsalis*.** A) 7 days post-exposure. B) 14 days post-exposure. Red = bodies (infection); blue = legs/wings (dissemination); black = saliva (transmission). Males = closed circles, females = open circles. Zero values had 0.01 added purely for log-scale plotting purposes ($10^{-2}$ = uninfected); analysis was performed on untransformed data. ** = $P < 0.01$.

**Figure 7. Wild bloodfed *Cx. quinquefasciatus* mosquito.** Top: Male mosquito was collected incidentally during collections in Mallorca, Spain. What appears to be partially digested blood is visible in the midgut. Bottom: Closeup of mosquito genitalia to confirm sex of the specimen.

**Supplemental legends**

**Supplementary Video 1.** Probing behavior of dehydrated male *Cx. tarsalis* mosquito on the thumb of a human volunteer.

**Supplementary Video 2.** Probing behavior of dehydrated male *Cx. tarsalis* mosquito on the index finger of a human volunteer.

**Supplementary Video 3.** Probing behavior of dehydrated male *Ae. aegypti* mosquito on the thumb of a human volunteer.

**Supplementary Video 4.** Probing behavior of dehydrated male *Ae. aegypti* mosquito on the index finger of a human volunteer.

**Supplementary Video 5.** Probing behavior of dehydrated male *Ae. notoscriptus* mosquito on the index finger of a human volunteer.

**Supplementary Video 6.** Probing behavior of dehydrated male *Ae. notoscriptus*
mosquitoes on the index finger of a human volunteer.

**Supplementary Video 7.** Probing behavior of dehydrated male *Cx. quinquefasciatus*
mosquito on the hand of a human volunteer.

**Supplementary Video 8.** Probing behavior of dehydrated male *Cx. quinquefasciatus*
mosquito on the hand of a human volunteer.

**Supplementary Video 9.** Attraction and landing behavior of dehydrated male *Ae.*
*aegypti* mosquitoes on the hand of a human volunteer.

**Supplementary Video 10.** Probing behavior of dehydrated *Cx. tarsalis* male mosquito
on the wrist of a human volunteer. This mosquito succeeded in penetrating the outer
epidermis.

**Supplementary Video 11.** Male *Cx. tarsalis* feeding from a wound on a human
volunteer.

**Supplementary Table 1.** Ct values for vertebrate DNA detected in male mosquitoes
collected from Galveston, TX.
806
807    **Supplementary File 1.** Vertebrate 16S and COI sequences obtained from field-
808    collected mosquitoes, with associated BLAST statistics.
809
810

**Figure 1**

**Figure 2**

**Figure 3**

**Figure 4**

**Figure 5**

**Figure 6**

**Figure 7**

Table 1. WNV Infection rate (IR), dissemination rate (DR) and transmission rate (TR) of dehydrated males and females at 7 and 14 days post-infection. No comparisons were statistically significant after correcting for multiple comparisons.

|  | N | IR | DR | TR |
|---|---|---|---|---|
| Day 7 |  |  |  |  |
| Males | 43 | 0.53 | 0.70 | 0.56 |
| Females | 10 | 0.50 | 0.60 | 0.00 |
| Day 14 |  |  |  |  |
| Males | 14 | 0.50 | 0.86 | 0.50 |
| Females | 12 | 0.92 | 0.91 | 0.30 |