Revisiting the paradigm of anhematophagy in male mosquitoes

Jovana Bozic[1,2,3¶], Renuka E. Joseph[1,2,3¶], Rachel S. Krizek[2,3,4], Amber Holley[5],

Maureen Laroche[6,7,8], Joshua B. Benoit[9], and Jason L. Rasgon[1,2,3,4]*

[1] Department of Entomology, The Pennsylvania State University, University Park, PA,

USA

[2] The Center for Infectious Disease Dynamics, The Pennsylvania State University,

University Park, PA, USA

[3] The Huck Institutes of the Life Sciences, The Pennsylvania State University, University

Park, PA, USA

[4] Department of Biochemistry and Molecular Biology, The Pennsylvania State

University, University Park, PA, USA

[5] Department of Pathology, The University of Texas Medical Branch, Galveston, TX,

USA

[6] Department of Microbiology & Immunology, The University of Texas Medical Branch,

Galveston, TX, USA

[7] Department of Global Health, The University of Texas Medical Branch, Galveston, TX,

USA

[8] Clima, Latin American Center of Excellence for Climate Change and Health,

Universidad Peruana Cayetano Heredia (UPCH), Lima-Peru.

[9] Department of Biological Sciences, University of Cincinnati, Cincinnati, OH, USA

[¶] These authors contributed equally to this work

* email for correspondence: jlr54@psu.edu

Abstract

Female mosquitoes are reproductively obligate bloodfeeders which feed on vertebrate blood to obtain nutrients required for egg production (driving transmission of vector-borne pathogens in the process), and which rely on plant sugars for their non-reproductive energy requirements. Male mosquitoes, on the other hand, are thought to rely exclusively on plant sugars for their energetic needs; indeed, this dichotomy is one of the central tenets of medical entomology. Here, we show that male *Culex tarsalis* and *Aedes aegypti* mosquitoes will readily take blood from a membrane feeder when reared under dehydration conditions with no toxic effects. Mosquitoes with impaired humidity detection do not increase their bloodfeeding rates when dehydrated compared to wild-type controls. While conventionally reared males ignore a human host, dehydrated males are attracted to and attempt to probe, with some success, although they cannot access host capillaries. However, they will take blood from a vertebrate host wound. When fed a blood meal containing West Nile virus, male mosquitoes can become infected with and orally transmit the pathogen at rates and titers equivalent to females. Finally, vertebrate DNA, likely from blood, was detected in wild-caught specimens of male *Culex quinquefasciatus* mosquitoes from Texas. These data suggest that under some circumstances male mosquitoes may be able to probe and/or ingest blood and transmit pathogens to vertebrate hosts, and that their role in maintaining pathogen transmission cycles should be re-examined.

Key words: *Culex tarsalis*, *Aedes aegypti*, mosquito, bloodfeeding, sugar feeding, vector-borne disease

Introduction

Female mosquitoes (except those which are autogenous) are reproductively obligate bloodfeeders, feeding on vertebrate blood to obtain nutrients required for egg production, and relying on plant sugars for their non-reproductive energy requirements. Male mosquitoes, on the other hand, rely exclusively on plant sugars for their energetic needs [1]. Indeed, this difference is one of the central tenants of medical entomology; female mosquitoes bloodfeed, males do not. Female bloodfeeding allows transmission of blood-borne pathogens, such as viruses or parasites, between vertebrate hosts, which is why the majority of mosquito research is performed on females rather than males; even when males are studied, it is usually within the context of how they affect females (mating behavior and fertility, pathogen transmission modulation, etc.) [2-5].

Female mosquitoes are adapted to feed on blood, and this adaptation is reflected in the biology of their midgut, where transcripts related to blood digestion are enriched in the female compared to the male [6]; one would expect that male mosquitoes should not be attracted to blood as a nutrition source as they are thought to lack the proper physiology to digest and process it. However, there is one interesting report in the literature where male mosquitoes were attracted to and fed on blood. Nikbakhtzadeh and colleagues [7] documented bloodfeeding behavior in a laboratory colony of the mosquito *Culex quinquefasciatus.* When presented with defibrinated sheep blood on a cotton pledget (and to a much less efficient extent, a Parafilm membrane), male mosquitoes took a bloodmeal. However, blood was toxic to male mosquitoes, which died in a dose-dependent manner when blood was mixed with sugar [7], consistent with physiological adaptations to sugar vs bloodfeeding in males vs. females [6]. Interestingly, males did not show a preference for sugar compared to blood in a dual-choice assay [7], and the reason they fed on blood at all, particularly as it was toxic, remains an open question. As this is (to our knowledge) the only observation of male mosquito bloodfeeding behavior, it is difficult to speculate. However, there are a multitude of observations that male mosquitoes are attracted to human host odors and this behavior is suppressed by

93 mosquito repellants [8], which includes species from arguably the three most important
94 mosquito genera that act as disease vectors to humans (*Anopheles*, *Culex*, and *Aedes*).
95

Here, we present studies on male bloodfeeding behavior in the mosquitoes *Cx. tarsalis*
and Ae. aegypti. *Cx. tarsalis* is one of the major West Nile virus (WNV) vectors in North
America, where it is widely distributed across the Western United States [9]. It is
genetically diverse, generally feeds on birds in the wild, and can be facultatively
autogenous [9-10]. After becoming infected with WNV during a bloodfeeding event, it
can also transmit the virus vertically to offspring at relatively high rates [11-12]. *Ae.*
aegypti is one of the major invasive arbovirus vectors in the world [13]. We
opportunistically observed *Cx. tarsalis* and *Ae. aegypti* males taking blood during
unrelated laboratory studies, and undertook experiments to document and understand
the behavior. We found that when dehydrated, male *Cx. tarsalis* and *Ae. aegypti* will
predictably take human blood from a membrane feeder, determined the mechanism
driving male bloodfeeding behavior, and present results of experiments examining
potential for male mosquitoes to be involved in pathogen transmission cycles. These
results are a paradigm shift in our understanding of male mosquito biology and suggest
they may be more directly involved in pathogen transmission cycles than previously
recognized.
112

113 **Methods**
114 Human subjects: All experiments with a human volunteer used the senior author (JLR)
115 under PSU IRB Exempt Protocol STUDY00024284.
116

117 Mosquitoes: *Cx. tarsalis* strain (KNWR) and *Aedes aegypti* (Liverpool) were maintained
118 at 25°C, 16:8 h light:dark diurnal cycle with 80% relative humidity, with 10% sucrose
119 solution provided at all times through a cotton wick. For general rearing, mosquitoes
120 were provided with expired anonymous human blood (BioIVT) through a water-jacketed
121 glass membrane (Parafilm) feeder or a Hemotek feeder for egg development.
122

<u>Male dehydration:</u> To stimulate bloodfeeding, male mosquitoes were held at 25°C, 75%
RH without sugar or water for 24 hours [14].

<u>Survival analysis:</u> Bloodfed male mosquitoes were isolated and placed into cup cages,
held at the previously described standard insectary conditions, and provided with a
cotton ball soaked in 10% sucrose solution. Control males were non-bloodfed and were
maintained under the same conditions. Dead mosquitoes were counted every day and
removed from the cages. Significant differences in survival between mosquito groups
was determined with Kaplan-Meier analysis using GraphPad Prism version 9.0.4.

<u>Dehydration and bloodfeeding behavior in male *Cx. tarsalis*:</u> *Cx. tarsalis* males were
reared conventionally (80% RH, with free access to 10% sucrose solution in water), or
under dehydrating conditions (75% RH, 25°C, with no access to water or sugar for 24
hours) [14], then were offered a bloodmeal through a membrane feeder for 30 minutes.
The number of fed and unfed mosquitoes at the end of the feeding period were counted.
Data were analyzed by Fishers Exact test.

<u>Ionotropic receptor 93a (Ir93a) mutant mosquito assays:</u> CRISPR protocols have not yet
been developed for *Cx. tarsalis*, but we noted during experiments that males of the
species *Ae. aegypti* (where CRISPR mutagenesis is routine) would also take blood from
a membrane feeder, so we obtained an *Ae. aegypti* line that was a CRISPR knockout
mutant for the Ir93a gene, which inhibits its ability to sense humidity [15]. The mutation
was introgressed into the wild-type Liverpool background for comparison with Liverpool
controls, and both lines reared as described above. For experiments, at 5-6 days post-
emergence, males of each strain were transferred to 10 x 10 x 10 cm cages and
deprived of sucrose and water (or held at normal conditions as controls) for 24 hours
before being offered an anonymous human bloodmeal using an artificial feeding system
(Hemotek). Bloodfeeding rates for each genotype and condition were recorded. Data
were analyzed by Fishers Exact tests and confidence intervals calculated from the
binomial distribution.

Landing and probing experiments: Cages of 50 *Cx. tarsalis* males (reared under standard or dehydrating insectary conditions) were allowed to probe on the hand of the senior author for five minute trials. Mosquito landings (defined as a mosquito alighting on the volunteer hand for any period of time) and probing behavior (defined as exploring and probing with mouthparts) were counted during the 5 minute interval. The experiment was repeated 6 times. Data were analyzed by Mann-Whitney U test.

Host bloodfeeding by male mosquitoes: The senior author had an unrelated small (3mm) scratch on their hand from obtained from a pet cat a day earlier. A sterile razor blade was used to pick the scab off the scratch allowing a minimal amount of blood to be exposed. The wounded hand was placed in a cage of 20 dehydrated male *Cx. tarsalis* mosquitoes and their behavior recorded.

WNV feeds: Dehydrated *Cx. tarsalis* males and female controls were allowed access to an infectious blood meal consisting of a 1:1 mix of anonymous human blood (BioIVT) and 5.0×10^{7} FFU/ml (focus-forming units/ml) of WN02-1956 (GenBank: AY590222). A subset of male and females were processed immediately after feeding ("day zero") to check for virus viability. Mosquito virus infection and transmission assays were performed at 7 and 14 days post-blood feeding. Fully engorged mosquitoes were sorted from non-fed ones for analysis. Mosquitoes were anesthetized with triethylamine (Sigma, St. Louis, MO), legs/wings from each mosquito were removed and placed separately in a 2-mL tube filled with 0.5 mL mosquito diluent (MD: 20% heat-inactivated fetal bovine serum (FBS) in Dulbecco's phosphate-buffered saline, 50 µg/mL penicillin/streptomycin, 50 ug/mL gentamicin, and 2.5 µg/mL fungizone, with a sterile 2.0 mm stainless steel bead (Next Advance, Inc. Innovative Lab Products for the Life Sciences). The proboscis of each mosquito was positioned in a tapered capillary tube containing approximately 10 µL of a 1:1 solution of 50% sucrose and FBS to induce salivation. After 30 min, the tube contents were expelled into 0.3 mL MD, and bodies were placed individually into a 2-mL tube filled with 0.5 mL MD and a stainless steel bead as described above. Mosquito bodies and legs/wings were homogenized for 30 sec with TissueLyser (QIAGEN, Hilden, Germany) at 24 cycles/sec, followed by

centrifugation for 1 min. Mosquito bodies, legs/wings, and salivary secretion samples
were tested for live, infectious WNV using focus-forming assays (FFAs; see below).

WNV FFAs: WNV titers were quantified by FFA, which detects live, infectious virus.
C6/36 cells were seeded into 96-well plates at a density of 1×10^5 cells/well and
incubated overnight at 28°C in complete RPMI medium without CO_2. The next day,
medium was removed from the wells. Samples from male and female bodies or
legs/wings were serially diluted in a serum-free RPMI medium; saliva samples were
undiluted. 30 µl of each sample was added in duplicate to the prepared C6/36 cells.
Cells were incubated for 1 hour at 28°C without CO_2, after which the inoculum was
removed. 100 µl of RPMI containing 0.8% methylcellulose was added to limit viral
spread. Infected cells were incubated for 48 hours at 28°C without CO_2. At 48 hours
post-infection, infected C6/36 cells were fixed with 50 µl 4% formaldehyde for 30
minutes at room temperature (RT). Cells were washed, permeabilized with 0.2% triton-
X, and blocked with 3% BSA. 30 µl monoclonal flavivirus antibody (Clone D1-4G2-4-15,
BEI-resources, NR-50327) was added and incubated overnight at 4°C. After washing,
30µl of fluorescent secondary antibody (1:1000 dilution; Goat anti-Mouse IgG (H+L)
Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 488, Invitrogen/Thermo
Fischer, A-11029) was added and incubated overnight at 4°C. Cells were maintained in
100 µl PBS to prevent drying. West Nile virus foci were imaged using a FITC filter on an
Olympus BX41 microscope with a UPlanFl 4x objective and counted. Infection rate (IR)
was defined as the proportion of mosquitoes exposed to virus that had WNV-positive
bodies. Dissemination rate (DR) was defined as the proportion of mosquitoes with
WNV-positive bodies that had WNV-positive legs/wings. Transmission rate (TR) was
defined as the proportion of mosquitoes with WNV-positive legs/wings that had WNV-
positive saliva. IR, DR, and TR were analyzed with Fisher's exact tests. Viral titers were
analyzed using Mann-Whitney U tests.

Wild mosquito collection, DNA extraction and molecular detection of vertebrate DNA:
Mosquitoes were collected using BG-Sentinel traps (Biogents) on Galveston Island, in
South Texas, August 12-13, 2024. Traps were placed in a residential area bordered by

marshes frequently visited by wildlife (e.g., coyotes, birds, racoons, opossums, etc.).
Traps were placed at 6pm and collected the next morning at 6am. Fresh mosquitoes
were sexed, morphologically identified the same day and placed at -20°C until
processed. After identification, whole male mosquitoes were individually incubated with
20µL of proteinase K and 180µL of G2 buffer (Qiagen) at 56°C for one hour then were
homogenized using tungsten beads and a TissueLyser (Qiagen) for one minute at 30
cycles/sec. After a quick spin to pellet debris, DNA was extracted from the lysate using
a Qiagen EZ1 automated DNA extraction instrument and reagents from the associated
EZ1&2 DNA Tissue Kit. DNA samples were screened for vertebrate DNA using
vertebrate-specific qPCR assay targeting a fragment of the 16s rRNA on a CFX Duet
PCR system (Bio-Rad) [16]. Human blood obtained from the University of Texas
Medical Branch Blood Bank was used as a positive control and PCR mix without DNA
was used as a negative control. Samples were analyzed in duplicate and the entire
assay repeated twice.
230
231

Results
Bloodfeeding is not toxic to *Cx. tarsalis* males: While we were bloodfeeding during an
experiment related to relative humidity (e.g. [17]), we noted incidentally that in addition
to females, male mosquitoes were probing the membrane and were taking blood
(Figure 1A,B). As this was a spontaneous occurrence, the total number of males in the
cage was not recorded but was on the order of 50-70 based on standard rearing
practices in our lab. Out of this total, we isolated seven blood-engorged males. These
males were placed into a survival cup and a survival experiment conducted, comparing
their survival to 10 non-bloodfed males from the same initial cage. Although
Nikbakhtzadeh et al. [7] demonstrated than blood was highly toxic to male *Cx.*
quinquefasciatus in laboratory studies, we did not observe any acute toxicity to blood in
male *Cx. tarsalis*; indeed, survival in bloodfed males was marginally (although not
statistically) higher than non-bloodfed (Figure 1C).
245

Cx. tarsalis male bloodfeeding is driven by dehydration: As we previously demonstrated
that dehydration stimulates elevated bloodfeeding behavior in females [14, 17], we
tested the hypothesis that dehydration was driving bloodfeeding behavior in males.
Cages of male mosquitoes were reared under conventional insectary conditions or
under dehydrating conditions [14]. No conventionally reared male mosquito (N = 64)
took a bloodmeal from the membrane feeder, while 44/163 dehydrated males took a
bloodmeal ($P < 0.00001$).

Male mosquito bloodfeeding behavior is dependent on their ability to sense humidity:
Ae. aegypti and _Anopheles gambiae_ mosquitoes sense humidity through ionotropic
receptor Ir93a, by which they locate oviposition sites, and CRISPR Ir93a knock-out
mutants are impaired in this behavior [15]. We noted anecdotally that in our lab, male
Ae. aegypti mosquitoes would also take blood from a membrane feeder, and as
CRISPR protocols have not yet been developed for _Cx. tarsalis_, we used an Ir93a _Ae._
aegypti KO mutant for these assays [15]. When reared under standard insectary
conditions, bloodfeeding rates did not differ statistically between wild-type and mutant
mosquitoes. However, when reared under dehydrating conditions, bloodfeeding rates
for the mutant did not increase, while wild-type mosquitoes had significantly elevated
bloodfeeding behavior ($P = 0.0284$) (Figure 2).

Dehydrated male mosquitoes will probe the hand of a human volunteer: The hand of a
human volunteer was exposed to cages of conventionally reared or dehydrated male
Cx. tarsalis. Conventionally reared males showed little interest in the host, with
infrequent landings that lasted less than 5 seconds. None demonstrated probing
behavior. In contrast, dehydrated males landed significantly more often on the hand of
the volunteer ($P = 0.0065$), most landings lasted until the end of the time period, and
probing behavior was observed in the majority of landings ($P = 0.0022$) (Figure 3,
Supplementary Videos 1 and 2). One dehydrated male mosquito (out of 6 separate
trials) was able to lightly pierce the skin of the volunteer at the base of the wrist,
although it was unable to reach the capillaries and acquire a bloodmeal (Supplementary
Video 3). The bite resulted in a mild immunogenic reaction that disappeared after

approximately 10 minutes (Supplementary Figure 1). To confirm that only males were in
the cage, after the study was concluded the entire cage was killed by freezing and every
mosquito visually examined for the presence of a female or a gynandromorph; only
males were identified. While this is only a single observation and thus definitive
conclusions cannot be drawn, to our knowledge, this is the first documented case of a
male mosquito biting a vertebrate host.

Dehydrated male mosquitoes will take blood from a vertebrate host wound: Dehydrated
male *Cx. tarsalis* show keen interest in probing a human host, but in these experiments
were not able to acquire blood, even from the single observed "successful" probing
attempt. We hypothesized that if blood was made more accessible, male mosquitoes
would take a bloodmeal. The senior author serendipitously had a small scratch on their
hand (acquired from a pet cat a day earlier). The scab was peeled back using a sterile
razor blade, exposing a small amount of blood. The volunteer placed their hand in a
cage of 20 dehydrated male mosquitoes. Males were attracted to the wound, and
wound probing behavior was observed by 5 males (see Supplementary Video 4 for
example). One male out of the 5 that probed fed and took a bloodmeal from the wound
(Supplementary Video 5, Figure 4). At the conclusion of the experiment, the fed male
was dissected to confirm the presence of blood in the gut (Figure 4).

Male *Cx. tarsalis* mosquitoes are competent vectors for West Nile virus: Since we
determined that male *Cx. tarsalis* will probe a human hand or ingest blood from a
wound, allowing ingestion of a blood meal from a vertebrate host, we asked the
question: can male mosquitoes become infected with and transmit arboviruses? We
offered dehydrated male mosquitoes a bloodmeal spiked with WNV and assayed their
vector competence at day 7 and day 14 post-exposure. Female *Cx. tarsalis* were
exposed to virus at the same time as a control. We found that both female and male *Cx.*
tarsalis were able to become infected with, disseminate, and orally transmit virus; males
transmitted at both day 7 and 14, while females only had detectable virus in their saliva
at day 14. After adjusting for multiple comparisons, infection rates (IR), dissemination

rates (DR), and transmission rates (TR) did not differ statistically between males and
females at either timepoint (Table 1).

We quantitated all viral titers using an infectious virus assay. First, a subsample of
males and females were assayed immediately after feeding ("day zero") to confirm virus
viability. All fed males and females had detectable live infectious virus in their bodies,
although females had statistically higher viral titers (P = 0.005), likely because they
could physically ingest a larger volume of blood. At day 7 post-exposure, viral titers
were not statistically different between males and females in the bodies, the legs/wings,
or the saliva (Figure 5A). At day 14 post-exposure, females had higher viral titers in their
bodies (P = 0.001) and legs/wings (P = 0.0083) compared to males, suggesting either
greater viral replication rates, or simply more tissue available for virus replication due to
the larger size of the females. However, viral titers in saliva between males and females
were statistically similar (Figure 5B).

<u>Vertebrate DNA can be detected in wild-caught male mosquitoes:</u> While BG-Sentinel
traps are designed to collect female mosquitoes, they do also catch males [e.g. 18-19].
We collected a total of 15 male mosquitoes (ten *Ae. albopictus*, one *Ae. sollicitans,* and
four *Cx. quinquefasciatus*). Vertebrate DNA was not detected in *Ae. albopictus* or *Ae.*
sollicitans. However, three of the four collected *Cx. quinquefasciatus* males had
detectable vertebrate DNA (Ct values ranging from 15.55 to 18.36) (Supplementary
Table 1). When homogenized during extraction, the supernatants from these positive
males were reddish-brown, possibly suggesting the presence of blood in the specimen.

Discussion
Previous work showed that in the laboratory, blood was toxic to male *Cx.*
quinquefasciatus mosquitoes [7], suggesting that in this species male bloodfeeding
seems to be a maladaptive trait, perhaps a laboratory artifact. In our study, we
demonstrate that males of other species (*Cx. tarsalis* and *Ae. aegypti*) can tolerate
bloodfeeding, and that male bloodfeeding behavior is driven by water homeostasis
during dehydration conditions. When mosquitoes cannot sense humidity due to Ir93a

mutagenesis, dehydration does not increase blood seeking behavior. These results are
consistent with the role of dehydration on bloodfeeding behavior in female mosquitoes,
where dehydration can stimulate females to increase their bloodfeeding rates as well
[14, 17, 20] and thus may reflect an adaptive trait where mosquitoes (female or male)
can maximize their water intake during drought or periods of low relative humidity if
other sources (nectar or free water) are not available.
344
The mouthparts of male mosquitoes are thought to be physically incapable of
penetrating vertebrate skin; however, in our experiments they were proven adequate to
pierce a Parafilm membrane. Dehydrated *Cx. tarsalis* males were significantly attracted
to and actively probed the hand of a human host, and one individual was even able to
slightly penetrate the outer epidermis, leading to a transitory immune reaction
(Supplementary Video 3 and Supplementary Figure 1). As the saliva of males differs
from that of females, lacking various proteins needed for immunomodulation and
bloodmeal acquisition [21], and it is likely that very little saliva was transferred compared
to the bite of a female mosquito, it is not surprising that the host immune reaction was
mild and rapidly resolving. When allowed access to a wound, dehydrated male
mosquitoes readily probed the wound and one took a bloodmeal. As this experiment
was facilitated by the fortuitous presence of a pre-existing wound on the hand of the
senior author, it could not be deliberately repeated (as we were not allowed to make a
deliberate wound due to IRB concerns). Still, it does suggest that male mosquitoes have
the ability to take blood under specific rare circumstances that require dry periods and,
likely, a host with a wound. According to fossil evidence, male mosquitoes are thought
to once have had the ability to feed on vertebrate blood, and to have lost this ability over
evolutionary time [22]. It is possible that the neural circuitry regulating host seeking and
bloodfeeding behavior may still be conserved among male mosquitoes, or alternatively
that this is simply a unique response to dehydration conditions in the lab.
365
Interestingly our data demonstrate that, if male *Cx. tarsalis* orally acquire a WNV
infection, they are competent vectors and transmit the virus at similar rates and titers
compared to females. In our experiments we explicitly used an assay that quantified

live, infectious viral particles rather than quantitative PCR to rule out results that might
be due to carryover of non-infectious viral RNA. Our results suggest that male *Cx.*
tarsalis retain the receptors necessary for viral infection on their midgut, salivary glands,
and other body tissues. Interestingly, there is a recent report of wild male *Cx. pipiens*
mosquitoes that were artificially infected with WNV from accidently feeding on infected
blood droplets during a laboratory assay [23]. *Cx. tarsalis* males (as well as *Cx. pipiens*
and other *Culex* mosquitoes) can be infected with WNV due to vertical transmission
from infected mothers [11, 24-25], and it is possible that the ability for males to become
infected with virus orally is a consequence of their retaining receptors necessary for
vertical infection.
379
Finally, there is the question "is bloodfeeding behavior by male mosquitoes
epidemiologically significant"? It is already known that male mosquitoes can be
indirectly important for vector-borne disease transmission dynamics. For example,
mating can affect key physiological parameters in females related to pathogen infection
and transmission [2]. More directly, in some species, including *Cx. tarsalis* and WNV,
male mosquitoes can be infected with arboviruses by vertical transmission from infected
mothers [11, 24-25]. Infected males can also transmit some viruses venereally to
females during mating where they can be transmitted to vertebrate hosts during feeding
[24-25]. Consistent collection of males using host-derived attractants suggest that males
are commonly found to move toward hosts [8], increasing the potential of male feeding
on host-derived fluids under specific conditions (dry periods with a lack of sugar and
water resources). In field collections, we were able to detect vertebrate DNA in wild-
caught *Cx. quinquefasciatus* males, and the color of the homogenates for these
specimens indicated that blood may have been present, suggesting that male
mosquitoes do encounter vertebrate hosts and acquire DNA, possibly by probing and,
potentially, wound feeding. In total, our study suggests the rare possibility of edge cases
where male mosquitoes could be more directly implicated in virus transmission, where
males undergoing dehydrating conditions (for example, during drought) acquire virus
through vertical transmission from infected mothers or by feeding on an open wound of
an infected vertebrate host, then transmit to a naïve host through feeding on an open

wound or by probing the skin, as mosquitoes often transmit the bulk of virus when
probing skin prior to actually taking a bloodmeal [26].

We must emphasize that while compelling, the ability for male mosquitoes to bloodfeed
or transmit pathogens in nature have not been widely studied (we suspect that
researchers have not rigorously examined these phenomena). However, our data
suggest that the canonical role of males as non-bloodfeeders needs to be re-examined
and their contribution to pathogen transmission explicitly quantified, particularly in light
of recent vector-borne disease control strategies that rely on the mass release of male
mosquitoes into natural populations [27-30].

References

1.   Clements AN. The Biology of Mosquitoes, Volume 1. CABI Publishers; 1992.

2.   Dahalan FA, Churcher TS, Windbichler N, Lawniczak MKN. The male mosquito contribution towards malaria transmission: Mating influences the *Anopheles* female midgut transcriptome and increases female susceptibility to human malaria parasites. PLoS Pathog. 2019; 15:e1008063.

3.   Marcenac P, Shaw WR, Kakani EG, Mitchell SN, South A, Werling K, Marrogi E, Abernathy DG, Yerbanga RS, Dabiré RK, Diabaté A, Lefèvre T, Catteruccia F. A mating-induced reproductive gene promotes *Anopheles* tolerance to Plasmodium falciparum infection. PLoS Pathog. 2020;16:e1008908.

4.   Carraretto D, Soresinetti L, Rossi I, Malacrida AR, Gasperi G, Gomulski LM. Behavioural Responses of male *Aedes albopictus* to different volatile chemical compounds. Insects. 2022;13:290.

5.   Peng D, Kakani EG, Mameli E, Vidoudez C, Mitchell SN, Merrihew GE, MacCoss MJ, Adams K, Rinvee TA, Shaw WR, Catteruccia F. A male steroid controls female sexual behaviour in the malaria mosquito. Nature. 2022;608:93.

6.   Warr E, Aguilar R, Dong Y, Mahairaki V, Dimopoulos G. Spatial and sex-specific dissection of the *Anopheles gambiae* midgut transcriptome. BMC Genomics. 2007;8:37.

7.   Nikbakhtzadeh MR, Buss GK, Leal WS. Toxic effect of blood feeding in male mosquitoes. Front Physiol. 2016;7:4.

8.   Paris V, Hardy C, Hoffmann AA, Ross PA. How often are male mosquitoes attracted to humans? R Soc Open Sci. 2023 Oct 25;10(10):230921.

9.   Venkatesan M, Rasgon JL. Population genetic data suggest a role for mosquito-mediated dispersal of West Nile virus across the western United States. Mol Ecol. 2010;19:157.

10.   Provost-Javier KN, Chen S, Rasgon JL. Vitellogenin gene expression in autogenous *Culex tarsalis*. Insect Mol Biol. 2010;19:423.

473   11. Goddard LB, Roth AE, Reisen WK, Scott TW. Vertical transmission of West Nile
474   Virus by three California *Culex* (Diptera: Culicidae) species. J Med Entomol.
475   2003;40:743.
476
477   12. Nelms BM, Fechter-Leggett E, Carroll BD, Macedo P, Kluh S, Reisen WK.
478   Experimental and natural vertical transmission of West Nile virus by California *Culex*
479   (Diptera: Culicidae) mosquitoes. J Med Entomol. 2013;50:371.
480
481   13. Brady OJ, Hay SI. The Global Expansion of Dengue: How Aedes aegypti
482   Mosquitoes Enabled the First Pandemic Arbovirus. Annu Rev Entomol. 2020 Jan
483   7;65:191-208.
484
485   14. Hagan RW, Didion EM, Rosselot AE, Holmes CJ, Siler SC, Rosendale AJ,
486   Hendershot JM, Elliot KSB, Jennings EC, Nine GA, Perez PL, Rizlallah AE, Watanabe
487   M, Romick-Rosendale LE, Xiao Y, Rasgon JL, Benoit JB. Dehydration prompts
488   increased activity and blood feeding by mosquitoes. Sci Rep. 2018;8:6804.
489
490   15. Laursen WJ, Budelli G, Tang R, Chang EC, Busby R, Shankar S, Gerber R, Greppi
491   C, Albuquerque R, Garrity PA. Humidity sensors that alert mosquitoes to nearby hosts
492   ad egg-laying sites. Neuron. 2023 Mar 15;111(6):874-887.e8.
493
494   16. Taniguchi K, Akutsu T, Watanabe K, Ogawa Y, Imaizumi K. A vertebrate-specific
495   qPCR assay as an endogenous internal control for robust species identification.
496   Forensic Sci Int Genet. 2022 Jan;56:102628.
497
17. Velo E, Balestrino F, Kadriaj P, Carvalho DO, Dicko A, Bellini R, Puggioli A, Petrić
D, Michaelakis A, Schaffner F, Almenar D, Pajovic I, Beqirllari A, Ali M, Sino G, Rogozi
E, Jani V, Nikolla A, Porja T, Goga T, Fălcuă E, Kavran M, Pudar D, Mikov O, Ivanova-
Aleksandrova N, Cvetkovikj A, Akıner MM, Mikovic R, Tafaj L, Bino S, Bouyer J, Mamai
W. A Mark-Release-Recapture Study to Estimate Field Performance of Imported Radio-
Sterilized Male Aedes albopictus in Albania. Front Bioeng Biotechnol. 2022 Aug
16;10:833698.
505
506   18. Staunton KM, Goi J, Townsend M, Ritchie SA, Crawford JE, Snoad N, Karl S,
507   Burkot TR. Effect of BG-Lures on the Male Aedes (Diptera: Culicidae) Sound Trap
508   Capture Rates. J Med Entomol. 2021 Nov 9;58(6):2425-2431.
509
510   19. Manzano-Alvarez J, Terradas G, Holmes CJ, Benoit JB, Rasgon JL. Dehydration
511   stress and Mayaro virus vector competence in *Aedes aegypti*. J Virol. 2023 Dec
512   21;97(12):e0069523.
513
514   20. Holmes CJ, Chakraborty S, Ajayi OM, Unran MR, Frigard RA, Stacey CL, Susanto
515   EE, Chen SC, Rasgon JL, DeGennaro MJ, Xiao Y, Benoit JB. Multiple bouts of blood
516   feeding in mosquitoes allow prolonged survival and are predicted to increase viral
517   transmission during drought. bioRxiv [Preprint]. 2024 Jun 22:2024.05.28.595907.
518

21. Lu S, Martin-Martin I, Ribeiro JM, Calvo E. A deeper insight into the sialome of male and female *Ochlerotatus triseriatus* mosquitoes. Insect Biochem Mol Biol. 2022 Aug;147:103800.

22. Azar D, Nel A, Huang D, Engel MS. The earliest fossil mosquito. Curr Biol. 2023 Dec 4;33(23):5240-5246.e2.

23. Linthout C, Martins AD, de Wit M, Delecroix C, Abbo SR, Pijlman GP, Koenraadt CJM. The potential role of the Asian bush mosquito Aedes japonicus as spillover vector for West Nile virus in the Netherlands. Parasit Vectors. 2024 Jun 17;17(1):262.

24. Schopen S, Labuda M, Beaty B. Vertical and venereal transmission of California group viruses by *Aedes triseriatus* and *Culiseta inornata* mosquitoes. Acta Virol. 1991:35:37.

25. Patrican LA, DeFoliart GR. *Aedes triseriatus* and La Crosse virus: similar venereal infection rates in females given the first bloodmeal immediately before mating or several days after mating. Am J Trop Med Hyg. 1987 May;36(3):648-52

26. Visser I, Koenraadt CJM, Koopmans MPG, Rockx B. The significance of mosquito saliva in arbovirus transmission and pathogenesis in the vertebrate host. One Health. 2023 Feb 12;16:100506.

27. Bansal S, Lim JT, Chong CS, Dickens B, Ng Y, Deng L, Lee C, Tan LY, Kakani EG, Yoong Y, Du Yu D, Chain G, Ma P, Sim S, Ng LC, Tan CH. Effectiveness of *Wolbachia*-mediated sterility coupled with sterile insect technique to suppress adult *Aedes aegypti* populations in Singapore: a synthetic control study. Lancet Planet Health. 2024 Sep;8(9):e617-e628.

28. Carvalho DO, McKemey AR, Garziera L, Lacroix R, Donnelly CA, Alphey L, Malavasi A, Capurro ML. Suppression of a Field Population of *Aedes aegypti* in Brazil by Sustained Release of Transgenic Male Mosquitoes. PLoS Negl Trop Dis. 2015 Jul 2;9(7):e0003864.

29. Apte RA, Smidler AL, Pai JJ, Chow ML, Chen S, Mondal A, Sánchez C HM, Antoshechkin I, Marshall JM, Akbari OS. Eliminating malaria vectors with precision-guided sterile males. Proc Natl Acad Sci U S A. 2024 Jul 2;121(27):e2312456121.

30. Rasgon JL. Precision-guided tools for malaria control. Proc Natl Acad Sci U S A. 2024 Aug 6;121(32):e2411587121.

Declarations

Ethics approval and consent to participate: All experiments with a human volunteer
used the senior author (JLR) under PSU IRB Exempt Protocol STUDY00024284.

Acknowledgements: This research was supported by NIH/NIAID grant R01AI150251,
USDA Hatch Project 4769, a grant with the Pennsylvania Department of Health using
Tobacco Settlement Funds, and funds from the Dorothy Foehr Huck and J. Lloyd Huck
endowment to JLR, and NIH/NIAID grant R01AI148551 to JBB and JLR. We thank Ms.
Amelia Romo and Ms. Heather Engler for assistance with mosquito rearing and Paul
Garrity for providing the transgenic mosquitoes used in this study (Ir93a).

Authors' contributions: JB, REJ, RSK, AH, ML, JBB, and JLR conducted the research,
JBB contributed materials and reagents, JLR analyzed the data, JB, REJ, JBB, and JLR
wrote the manuscript.

Figure legends

Figure 1. Male *Cx. tarsalis* bloodfeeding behavior and survival. A) mosquitoes congregating at and bloodfeeding from a paraffin membrane. Arrow points to male orienting toward the membrane. B) Blood-engorged male mosquito. Un-engorged male can be seen in-frame. C) Survival curve of bloodfed vs. non-bloodfed male *Cx. tarsalis* mosquitoes. No significant difference was observed between treatments.

Figure 2. CRISPR deletion of Ir93a ablates male mosquito bloodfeeding behavior under dehydration conditions. When reared conventionally, both wild-type and mutant *Ae. aegypti* exhibit baseline levels of bloodfeeding behavior. When reared under dehydration conditions, wild-type males significantly increase bloodfeeding behavior but humidity-insensitive mutant mosquitoes do not. Confidence intervals were calculated from the binomial distribution. WT = wild-type.

Figure 3. Host probing behavior of dehydrated male mosquitoes. A) Control cage of conventionally reared male *Cx. tarsalis*. Mosquitoes ignore the human host. B) Cage of dehydrated male *Cx. tarsalis*. Dehydrated males land on and probe the human host. C, D) Stills from Supplementary Videos 1 and 2 showing dehydrated male mosquito probing behavior. See videos for complete behavioral responses. E) Landing responses for dehydrated vs. conventionally reared ("standard") male *Cx. tarsalis*. F) Probing behavior for dehydrated vs. conventionally reared ("standard") male *Cx. tarsalis*. Error bars = SEM. ** = $P < 0.01$

624 **Figure 4. Male *Cx. tarsalis* taking a human bloodmeal from a wound.** A) Male *Cx.*
625 *tarsalis* feeding on an open wound. B) Close-up of feeding behavior. C) Blood is
626 observable in the male mosquito gut. D) Blood in the male mosquito gut was confirmed
627 by dissection. See Supplementary Video 3 for complete behavioral response.
628
Figure 5. West Nile virus vector competence for male and female *Cx. tarsalis*. A) 7
days post-exposure. B) 14 days post-exposure. Red = bodies (infection); blue =
legs/wings (dissemination); black = saliva (transmission). Males = closed circles,
females = open circles. Zero values had 0.01 added purely for log-scale plotting
purposes (10^{-2} = uninfected); analysis was performed on untransformed data. ** = $P <$
0.01.
635
636

Table 1. WNV Infection rate (IR), dissemination rate (DR) and transmission rate (TR) of dehydrated males and females at 7 and 14 days post-infection. No comparisons were statistically significant after correcting for multiple comparisons.

	N	IR	DR	TR
Day 7				
Males	43	0.53	0.70	0.56
Females	10	0.50	0.60	0.00
Day 14				
Males	14	0.50	0.86	0.50
Females	12	0.92	0.91	0.30

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5