1 **Nonequilibrium polysome dynamics promote chromosome segregation and its**

2 **coupling to cell growth in *Escherichia coli***

Alexandros Papagiannakis[1,2,3], Qiwei Yu[5], Sander K. Govers[1,&], Wei-Hsiang Lin[1,2,#], Ned S. Wingreen[5,6],

and Christine Jacobs-Wagner[1,2,3,4]

[1]Howard Hughes Medical Institute, Stanford University, Stanford, CA 94305, USA

[2]Sarafan Chemistry, Engineering, and Medicine for Human Health Institute, Stanford University, Stanford,

CA 94305, USA

[3]Department of Biology, Stanford University, Stanford, CA 94305, USA

[4]Department of Microbiology and Immunology, School of Medicine, Stanford, CA 94305, USA

[5]Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton NJ 08544, USA

[6]Department of Molecular Biology, Princeton University, Princeton NJ 08544, USA

[&]Current address: Department of Biology, KU Leuven, Leuven, Belgium

[#]Current address: Institute of Molecular Biology, Academia Sinica, Taipei, Taiwan

*Corresponding author: email: jacobs-wagner@stanford.edu

19 **ABSTRACT**

Chromosome segregation is essential for cellular proliferation. Unlike eukaryotes, bacteria lack

cytoskeleton-based machinery to segregate their chromosomal DNA (nucleoid). The bacterial ParABS

system segregates the duplicated chromosomal regions near the origin of replication. However, this

function does not explain how bacterial cells partition the rest (bulk) of the chromosomal material.

Furthermore, some bacteria, including *Escherichia coli*, lack a ParABS system. Yet, *E. coli* faithfully

segregates nucleoids across various growth rates. Here, we provide theoretical and experimental evidence

that polysome production during chromosomal gene expression helps compact, split, segregate, and

position nucleoids in *E. coli* through out-of-equilibrium dynamics and polysome exclusion from the DNA

meshwork, inherently coupling these processes to biomass growth across nutritional conditions. Halting

chromosomal gene expression and thus polysome production immediately stops sister nucleoid migration

while ensuing polysome depletion gradually reverses nucleoid segregation. Redirecting gene expression

away from the chromosome and toward plasmids causes ectopic polysome accumulations that are

sufficient to drive aberrant nucleoid dynamics. Cell width enlargement suggest that the proximity of the DNA

to the membrane along the radial axis is important to limit the exchange of polysomes across DNA-free

regions, ensuring nucleoid segregation along the cell length. Our findings suggest a self-organizing mechanism for coupling nucleoid segregation to cell growth.

INTRODUCTION

All cells must segregate their replicated chromosomes to propagate genetic information to daughter cells at division. Remarkably, this essential process is far less understood in bacteria than eukaryotes. Eukaryotic cells use the mitotic spindle, a sophisticated cytoskeleton-based machine, for chromosome segregation. No equivalent structure has been identified in bacteria, which fold their chromosomal material into a membrane-less organelle called the nucleoid. Many bacterial species use a ParABS system to segregate a specific DNA region proximal to the chromosomal origin of replication (*oriC*) (Figge et al., 2003; Ireton et al., 1994; Jalal and Le, 2020; Lim et al., 2014; Livny et al., 2007; Mohl and Gober, 1997). However, this initial step does not explain how the terminal regions of the chromosome separates (Harju et al., 2024) or how sister nucleoids move away from each other. Moreover, the chromosomally encoded *parA* and/or *parB* gene are often not essential for cell viability in various bacteria (Bartosik et al., 2009; Charaka and Misra, 2012; Donczew et al., 2016; Donovan et al., 2010; Du et al., 2016; Ireton et al., 1994; Jakimowicz et al., 2007a, 2007b; Jecz et al., 2015; Kadoya et al., 2011; Kawalek et al., 2020; Kim et al., 2000; Lagage et al., 2016; Lee and Grossman, 2006; Lewis et al., 2002; Li, 2019; Li et al., 2015; Minnen et al., 2011; Santi and McKinney, 2015; Takacs et al., 2022; Yamaichi et al., 2007; Yu et al., 2010). In fact, some bacteria do not even encode a ParABS system for *oriC* partitioning (Livny et al., 2007), with *Escherichia coli* being a prime example. Yet, *E. coli* segregates its duplicated nucleoids faithfully.

Several non-dedicated cellular processes have been proposed to play a role. For example, DNA replication may facilitate chromosome partitioning though a DNA polymerase-dependent extrusion mechanism (Lemon and Grossman, 2000). Simulations of coarse-grained polymer models have also suggested that the conformational entropy of confined circular DNA chains could provide an unmixing force that separates two mixed DNA polymers (Jun and Mulder, 2006; Jun and Wright, 2010; Pande et al., 2023). However, recent modeling work has argued that entropic forces generated from excluded volume interactions between partially replicated chromosomes inhibit (rather than promote) their segregation (Harju et al., 2024). Mechanical stress between overlapping sister nucleoids (Fisher et al., 2013) and DNA loop extrusion (Harju et al., 2024) have also been proposed to contribute to chromosome segregation. *E. coli* mutants in which the *mukB* gene required for DNA loop extrusion is deleted exhibit moderate nucleoid segregation defects linked to the organization and condensation of the nucleoid (Danilova et al., 2007; Mäkelä et al., 2021; Sawitzke and Austin, 2000). However, even in these mutants, most cells successfully segregate and partition their nucleoids between the daughter cells. Furthermore, topoisomerase I mutations or *seqA* deletion suppress the chromosome segregation defects associated with the *mukB* deletion (Danilova et al., 2007; Sawitzke and Austin, 2000; Weitao et al., 1999), suggesting that other mechanisms exist.

Furthermore, while these various proposed mechanisms likely contribute to the splitting of the nucleoid into
two objects, they cannot explain how separated sister nucleoids move away from each other. In addition,
the dominant mechanism underlying the nucleoid diffusional bias must somehow be coordinated with
growth rate. In the presence of a poor-quality carbon source, *E. coli* initiates nucleoid segregation late in
the cell division cycle. As the quality of the carbon source improves and the growth rate increases, nucleoid
segregation occurs at an increasingly earlier stage of the cell division cycle until it initiates in the preceding
cycle to accommodate faster division times (Bates and Kleckner, 2005; Govers et al., 2024; Tiruvadi-
Krishnan et al., 2022). Jacob et al. (1963) offered an elegant solution to couple chromosome segregation
to cell growth by hypothesizing that duplicated chromosomes migrate apart through membrane attachment
and localized growth of the membrane between the attachment points. Drug experiments have indeed
suggested that the chromosomal DNA is attached to the cytoplasmic membrane along the radial axis
through a process known as transertion (Bakshi et al., 2015; Binenbaum et al., 1999; Kaval et al., 2023;
Kim et al., 2024; Matsumoto et al., 2015; Spahn et al., 2023; Woldringh, 2002), which occurs when
transcription, translation, and insertion of membrane proteins occur at the same time. However, for the
nucleoids to migrate directionally, the 1963 model requires localized growth of the cytoplasmic membrane
between the splitting nucleoids. To our knowledge, there is no evidence of such localized expansion of the
inner membrane. Even if the connection between the DNA and cytoplasmic membrane extends to the
peptidoglycan cell wall, *E. coli* cells elongate via a dispersed mode of peptidoglycan growth along the cell
length up to the final stage of the division cycle (Cooper and Hsieh, 1988; Navarro et al., 2022; Wientjes
and Nanninga, 1989; Woldringh et al., 1987). In addition, experiments with *B. subtilis* cells stripped of their
cell walls indicate that nucleoid segregation does not require cell wall attachment (Wu et al., 2020). Finally,
the segregation of chromosomal loci is faster than the rate of cell elongation, ruling out cell elongation and
nucleoid tethering to the membrane or cell wall as a major driver of chromosome segregation (Kuwada et
al., 2013).
95
We hypothesized that the mechanism(s) driving nucleoid segregation and its coupling to growth rate may
be linked to the bacterial cellular organization. In contrast to eukaryotes, bacteria do not insulate their DNA
in membrane-bound compartments. As a result, the chromosomal meshwork can freely interact with the
macromolecules present in the cytoplasm, which exhibits a high level of polydispersity and crowding
(McGuffee and Elcock, 2010; Zimmerman and Trach, 1991). Experiments and modeling have suggested
that macromolecular crowders exert compaction forces on the compressible nucleoids through steric (i.e.,
excluded-volume) interactions (Mondal et al., 2011; Pelletier et al., 2012; Wu et al., 2019b; Yang et al.,
2020; Zhang et al., 2009). Monosomes and polysomes (hereafter collectively referred to as "polysomes" for
simplicity), which consist of mRNAs loaded with one or more translating ribosomes, constitute a sizeable
and abundant cytoplasmic crowder (McGuffee and Elcock, 2010). The proteome fraction allocated to
ribosomes increases with growth rate across nutrient conditions (Chure and Cremer, 2023; Dai et al., 2017;
Dourado and Lercher, 2020; Hu et al., 2020; Molenaar et al., 2009; Scott et al., 2014; Si et al., 2017). The

proportion of ribosomes engaged in translation also increases with improving nutrient quality, attaining 70-
80% under fast growth (Dai et al., 2017; Forchhammer and Lindahl, 1971; Mohapatra and Weisshaar, 2018;
Sanamrad et al., 2014; Varricchio and Monier, 1971). These polysomes form structures (Brandt et al., 2009)
of comparable or larger size than the average ~50 nm mesh size of the nucleoid (Xiang et al., 2021).
Consistent with a steric clash between polysomes and the nucleoid meshwork, fluorescently-labeled
ribosomes accumulate in DNA-free regions such as the cell poles and between segregated nucleoids
(Azam et al., 2000; Bakshi et al., 2015, 2012; Chai et al., 2014; Gray et al., 2019; Lewis et al., 2000;
Mohapatra and Weisshaar, 2018; Robinow and Kellenberger, 1994; Sanamrad et al., 2014; Xiang et al.,
2021). Interestingly, theoretical work suggests that excluded-volume effects alone may be sufficient for the
DNA to spontaneously phase separate from polysomes and compact into its observed nucleoid form
(Mondal et al., 2011). Furthermore, a recent nonequilibrium statistical physics model proposes that nucleoid
positioning and segregation could potentially be explained by the steric interaction (i.e., repulsion) between
polysomes and DNA and the nonequilibrium effects associated with mRNA synthesis and degradation
(Miangolarra et al., 2021). In this theoretical model, polysome accumulation in the middle of the nucleoid
emerges due to polysomes born in the middle of the nucleoid taking longer to diffuse out of the nucleoid
than those born closer to the nucleoid edges. Polysome accumulation beyond a certain concentration
threshold drives phase separation between the DNA and polysomes at the mid-nucleoid location, resulting
in nucleoid splitting. However, we lack experimental validation of this model. Furthermore, the coupling
between nucleoid segregation and growth rate has not been addressed either experimentally or
theoretically.

Here, we quantitatively characterize the temporal and spatial dynamics of nucleoids and polysomes in *E.*
coli under different conditions and perturbations. In this work, we refer to nucleoid segregation as a series
of events observable by microscopy (Figure 1A): (i) the initiation of nucleoid splitting, marked by the
depletion of a DNA marker near the mid position of the nucleoid, (ii) the end of nucleoid splitting, which
results in the generation of two separable nucleoid objects, and (iii) the migration of the sister nucleoids
away from each other, marked by the increasing distance between the centroid of the nucleoids. This
sequence of events can be initiated in the preceding cell division cycle under nutrient-rich (fast-growth)
conditions. Our experimental findings, combined with modeling, provide evidence that out-of-equilibrium
and asymmetric polysome rearrangements in the cell result in both DNA compaction and the choreography
of the nucleoid segregation cycle during growth. The dual involvement of polysomes in protein synthesis
and nucleoid dynamics ensures self-regulating coordination between cell growth and chromosome
segregation across a wide range of growth rates, even under conditions where *E. coli* divides faster than it
can replicate its chromosome.

RESULTS

The rate of polysome accumulation correlates with the rate of nucleoid segregation across cells.

To carefully examine the dynamics between nucleoids and polysomes with high temporal resolution (every 1 or 3 min, Figure 1 – figure supplement 1A-D), we ran time-lapse microfluidic experiments using an *E. coli* strain carrying the 50S ribosomal protein RplA fused to GFP and the nucleoid-associated protein HupA tagged with mCherry (Xiang et al., 2021). Cells were grown at 30°C in M9 buffer supplemented with glucose, casamino acids and thiamine (M9gluCAAT). While the ribosome signal displayed the expected polar and inter-nucleoid enrichments, the accrual of ribosome signal in the middle of the cells was particularly strong compared to the polar regions, as shown by representative individual cells (Figure 1B and Movie S1) and average kymographs (Figure 1C, n = 4122 cell division cycles). Cell constriction contributed to the apparent depletion of ribosomal signal from the mid-cell region at the end of the cell division cycle (Figure 1B-C and Figure 1 – figure supplement 2). In predivisional cells, the ribosomal signal accumulated in the middle of the segregated nucleoids near the ¼ and ¾ cell positions, as shown in average one-dimensional (1D) and two-dimensional (2D) fluorescence profiles (n = 1907 predivisional cells, Figure 1D). Drug treatment and single-ribosome tracking experiments have demonstrated that ribosomal accumulations inside *E. coli* correspond to polysomes whereas free ribosomal (or ribosomal subunits) are homogeneously distributed (Bakshi et al., 2012a; Gray et al., 2019; Lewis et al., 2000; Linnik et al., 2024; Sanamrad et al., 2014; Xiang et al., 2021). This is in agreement both with our own results obtained using rifampicin (Figure 1 – figure supplement 3), a transcriptional inhibitor that causes polysome depletion over time (Blundell and Wild, 1971; Campbell et al., 2001; Hartmann et al., 1967), and with the polysome classification in cryo-electron tomograms of *E. coli* sections (Xiang et al., 2021). Since ribosomal enrichments consist of polysomes, we will therefore refer to these enrichments as polysome accumulations hereafter.

Under our relatively nutrient-rich growth conditions (M9gluCAAT), nucleoid splitting occurred early in the division cycle and occasionally initiated in the segregated nucleoids from the preceding division cycle (Movie S1 and Figure 1B). Given the variability in nucleoid segregation timing across cells (Figure 1 – figure supplement 4A), we developed a computational method to track nucleoid dynamics independently of the cell division cycle (see Methods and Figure 1 – figure supplement 4B-C). Specifically, we focused on the nucleoid cycle—defined as the period between the ends of two nucleoid splitting events (Figure 1 – figure supplement 4D)—instead of the division cycle. By tracking the accumulation of RplA-GFP and the depletion of HupA-mCherry in the middle of nucleoids, we found that polysome accumulation and nucleoid splitting correlated in time (Figure 1E). Furthermore, the rate of polysome accumulation at mid-nucleoid ($\Delta RplA_{mid\text{-}nuc}/\Delta T$) correlated with the rate of DNA depletion in the same region ($\Delta HupA_{mid\text{-}cell}/\Delta T$) at the single-cell level (ρ = -0.52, Figure 1F). This indicates that cells that accumulated polysomes faster also split their nucleoids faster. Importantly, the fitted linear regression had an intercept close to zero for both axes (Figure 1F), indicating that when the rate of polysome accumulation approached zero, so did the rate of nucleoid splitting.

To examine what happens when sister nucleoids move away from each other, we divided the time between
the end of nucleoid splitting and cell division into four bins. We found that right after nucleoid splitting (0-
25% bin), most (~70%) cells continued to accumulate polysomes at mid-cell, i.e., between the sister
nucleoids (Figure 1G). For these cells, the rate of polysome accumulation ($\Delta RplA_{mid-cell}/\Delta T$) positively
correlated with the rate of nucleoid migration ($\Delta Distance_{nuc}/\Delta T$) ($\rho = 0.47$, Figure 1H). Thus, the faster that
cells accumulated polysomes at mid-cell, the faster the sister nucleoids migrated apart (and vice versa).

Cell elongation may also contribute to sister nucleoid migration near the end of the division cycle.
As noted in the Introduction, previous work has shown that the rate of chromosomal loci is faster than that
of cell elongation, indicating that cell elongation is not the predominant process driving chromosome
segregation (Kuwada et al., 2013). This interpretation is consistent with our rate measurements of nucleoid
segregation and cell elongation (Figure 1H). However, several observations suggest that cell elongation
may play a complementary role to polysome accumulations in nucleoid segregation, particularly near the
end of the division cycle. First, we noted that there was a substantial percentage (~30%) of cells with
decreasing RplA-GFP signal at mid-cell right after nucleoid splitting ($\Delta RplA_{mid-cell}/\Delta T \leq 0$, Figure 1H).
Interestingly, in these cells, nucleoid migration did not stop; instead, its average rate was similar to the
average rate of cell elongation ($\Delta Length_{cell}/\Delta T$) (Figure 1H). In fact, when the cell elongation rate was
subtracted from the nucleoid migration rate for each single cell (Figure 1 – figure supplement 5A), the
positive correlation between polysome accumulation and nucleoid migration rates remained ($\rho = 0.47$), but
now the average rate of nucleoid migration was near zero in cells with no polysome accumulation ($\Delta RplA_{mid-}$
$_{cell}/\Delta T \leq 0$).

This finding may suggest a mixed contribution between polysome accumulation and cell elongation to
nucleoid migration. This was interesting considering that polysomes became less enriched at mid-cell but
more enriched in the middle of sister nucleoids in predivisional cells (Figure 1B-D). This change
corresponded to a redistribution of polysome enrichments, as the average ribosome concentration
remained constant during the cell division cycle (Figure 1 – figure supplement 1E). This spatial change in
polysome enrichments was accompanied by a steady decline in the percentage of cells that continued to
accumulate polysomes at mid-cell between the end of nucleoid splitting and cell division (Figure 1G).
Concurrent with this decline, the migration rate of sister nucleoids became less correlated with the rate of
polysome accumulation and more correlated with the rate of cell elongation (Figure 1 – figure supplement
5B). To examine the relative correlation of polysome accumulation and cell elongation with nucleoid
migration over time, we used a linear mixed-effects model to analyze each relative nucleoid migration
interval (see Methods). The coefficients of the fitted mixed linear regressions suggest the following
hypothesis: Early during nucleoid migration (0-25% between the completion of nucleoid splitting and cell
division), polysome accumulation contributes most to the measured variance in the displacement of the

sister nucleoids (Figure 1I). This contribution progressively decreases over time while that of cell elongation
increases (Figure 1I). Such a switch in relative contribution would be consistent with the spatiotemporal
dynamics of polysome accumulation and cell wall synthesis (see Discussion).
221
Polysome accumulation at mid-cell correlates with the relative timing of nucleoid segregation
across nutrient conditions and growth rates.
If polysome production plays a role in nucleoid segregation, it predicts that the timing and amount of
polysome accumulation at mid-cell will correlate with the timing of nucleoid segregation across nutrient
conditions. To test this prediction, we analyzed images of fluorescently labeled ribosomes and nucleoids in
cells grown under 30 different carbon source conditions (Table S1) that vary the doubling times (~40 min
to ~4 h) and average cell areas (~1.9 to ~3 μm^2) of *E. coli*. This dataset included both previously published
(Gray et al., 2019) and new microscopy snapshots from our laboratory. Demographs generated from these
images revealed that the polysome accumulation at mid-cell was reproducible across all conditions and
strains, irrespective of the ribosomal subunit protein (RplA or RpsB) or the fluorescent tag (msfGFP, mEos2
or GFP) used to mark ribosomes (Figure 2A and Figure 2 – figure supplement 1). Since these profiles were
generated from snapshot images, they confirmed that the mid-cell polysome accumulation observed in the
time-lapse microfluidic experiments (Figure 1) was not caused by a photobleaching effect.
235
For population snapshots, nucleoids were typically imaged using DAPI (rather than a fluorescent fusion to
HupA), indicating that polysome accumulation at mid-nucleoid was independent of the DNA labeling
method. Importantly, and consistent with our prediction, the richer the growth conditions (i.e., the larger the
average cell area), the earlier the polysome accumulation and the nucleoid splitting occurred in the division
cycle based on relative cell lengths (Figure 2A and Figure 2 – figure supplement 1). In addition, the
polysome accumulation at mid-cell was more pronounced in nutrient-rich media (e.g., M9malaCAAT)
compared to nutrient-poor ones (e.g., M9mann) where the nucleoid segregated later in the division cycle
(Figure 2A and Figure 2 – figure supplement 1).
244
To quantify these phenotypes across strains and nutrient conditions, we extracted and correlated
population-level polysome and nucleoid statistics. We found a strong correlation (Spearman correlation ρ
= -0.85) between the amplitude of the average ribosomal signal accumulation and the average nucleoid
signal depletion at mid-cell (Figure 2B). Since the average cell area (colormap in Figure 2B) correlated with
the growth rate of the population (Schaechter et al., 1958), this plot also confirmed that faster growing
populations displayed stronger polysome accumulation and greater DNA depletion at mid-cell on average
(Figure 2B). This was also observed across cells within a population under the same nutritional condition
(ρ = -0.84, Figure 2 – figure supplement 2).
253

254 Both the average proteome fraction dedicated to ribosomes and the average fraction of ribosomes engaged
255 in translation are known to correlate with growth rate across nutrient conditions (Chure and Cremer, 2023;
256 Dai et al., 2017; Dourado and Lercher, 2020; Hu et al., 2020; Molenaar et al., 2009; Scott et al., 2014; Si et
257 al., 2017). Thus, a role for polysome production in nucleoid segregation may provide a mechanistic link
258 between growth rate and the relative timing of nucleoid splitting.
259
Spatial polysome asymmetry correlates with nucleoid positioning.
A surprising result was the apparent higher polysome accumulation at mid-cell relative to the cell pole
regions (Figures 1B-C and 2A). This was not artificially created by the smaller cytoplasmic volumes at the
cell poles or constriction sites due to membrane curvature (Figure 2 – figure supplement 3). It was also not
caused by photobleaching artefact during the timelapse microscopy since it was also observed from
snapshot images, particularly under nutrient-rich conditions (Figure 2 – figure supplement 1 and Figure 2 –
figure supplement 3). The uneven distribution of polysomes suggested limited diffusion-driven equilibration
of polysome concentration between the DNA-free regions. At division, such a disequilibrium could lead to
a higher concentration of polysomes at the new cell pole relative to the old pole in daughter cells through
inheritance. To examine this possibility, we went back to the microfluidic experiments in which we traced
cell lineages from mother to daughter cells (Figure 1 – figure supplement 1A), determined the pole identity
(new vs. old) of each tracked cell (Figure 1 – figure supplement 4B), and compared the polysome
accumulations between the new and old poles (see Methods and Figure 3 – figure supplement 1). Old
mother cells (located at the end of the microfluidic channels) and their daughters were excluded from our
analysis to avoid cell aging effects (Chao et al., 2024; Coquel et al., 2013; Koleva and Hellweger, 2015;
Łapińska et al., 2019; Lindner et al., 2008; Proenca et al., 2019). We found that newborn cells had more
polysomes at the new pole compared to the old one on average (Figure 3A), consistent with the notion that
polysomes do not rapidly equilibrate in concentration between DNA-free regions through diffusion.
278
Across cells, the polysome distribution asymmetry between cell poles negatively correlated with the position
of the nucleoid in newborn daughter cells (Figure 3B, cells always oriented with the new pole to the right).
The x and y intercepts of a fitted linear regression to the data were not zero, indicating that a nucleoid
positioned precisely at the cell center did not equate with an even distribution of polysomes between poles.
Instead, newborn cells with centrally located nucleoids tended to have more polysomes at the new pole
compared to the old pole (example #1 in Figure 3B-C) whereas cells born with symmetric polysome
enrichments between poles tended to display an off-center nucleoid closer to the new pole (example #2 in
Figure 3B-C).
287
288 ***Spatial polysome asymmetry correlates with asymmetric nucleoid compaction in newborn cells.***
289 Construction of average 2D cell projections of the RplA-GFP signal in cells sorted based on their relative
290 timing to cell division confirmed the polysome asymmetry between poles in newborn cells (Figure 3D and

Movie S2). Strikingly, the corresponding 2D projections of the HupA-mCherry signal revealed another spatial asymmetry, this time, in the average DNA mass distribution along the nucleoid length (axial asymmetry). The average HupA-mCherry signal concentration was higher toward the new pole right after birth or, correspondingly, toward the middle of the cell (future new pole) in the period prior to division (Figure 3D and Movie S2). Quantification of this axial asymmetry in newborn cells revealed that the HupA-mCherry concentration is ~20% higher toward the new pole on average (Figure 3E), suggesting that the DNA density is uneven along the nucleoid length. This nucleoid mass asymmetry emerged late in the nucleoid migration cycle, typically before cell division such that it was inherited by newborn cells (Figure 3D).

To examine if features of polysome accumulations (e.g., position, amplitude, or fraction of cell length covered) correlate with the asymmetric nucleoid density and its variability among cells, we used a linear mixed-effects model (see Methods). We found that in newborn cells, the positions of polysome enrichments in the old-pole and mid-cell regions significantly correlated with the HupA-mCherry density asymmetry (Figure 3 – figure supplement 2). Guided by this finding, we hypothesized that the positions of the accumulating polysomes along the cell length determine the space available for the chromosome to occupy and thereby dictate the DNA density distribution along the nucleoid. To examine this hypothesis, we combined the correlated polysome accumulation characteristics into a compound statistic that describes the relative polysome-free space between two cell halves (see Methods). Compared to other polysome statistics, the relative availability of cell space depleted of polysomes (RplA-GFP signal) correlated most strongly with the asymmetric distribution of DNA (HupA-mCherry signal) in newborn cells (Figure 3F). In other words, the DNA concentration was higher in cell regions with more space available between polysome enrichments. In most (~60%) cells, the distance between polysome enrichments, and thus the DNA concentration along the nucleoid, was greater between mid-cell and the new pole (i.e., x and y values >1, example #1 in Figure 3F-G). The opposite pattern was true for a small fraction (~14%) of cells, where the larger distance between polysome enrichments was located between mid-cell and the old pole (x and y values < 1, example #2 in Figure 3F-G). Thus, polysome asymmetry correlates with asymmetric nucleoid compaction.

A minimal reaction-diffusion model generates experimentally observed cellular asymmetries and growth rate-dependent nucleoid segregation.

Our single-cell correlative studies were consistent with exclusion between polysomes and DNA contributing to nucleoid compaction and segregation. However, it remained unclear whether the same mechanism could also explain the growth rate-dependent trends and cellular asymmetries that we observed (Figures 2 and 3). Therefore, we built a minimal reaction-diffusion model (see Methods) to describe the dynamics of polysomes and DNA during the cell cycle based on previous work (Miangolarra et al., 2021). Our model takes into account two ingredients important for nucleoid segregation: effective repulsion between DNA and polysomes from steric effects (described by the Cahn-Hilliard theory) and the nonequilibrium processes of

polysome synthesis and degradation (described by linear reaction kinetics). The model is based on realistic
parameters of polysome diffusion, production, and degradation (see Methods). We assume polysome
production (i.e., mRNA synthesis and ribosome loading) to be uniform within the nucleoid and polysome
degradation (i.e., mRNA decay) to be uniform across the entire cell (from pole to pole). These assumptions
consider the most trivial scenarios (see Discussion for other scenarios). We also assume that the cell grows
exponentially and that the nucleoid expands in size proportionally to the cell during growth, which has been
experimentally verified (Campos et al., 2014; Govers et al., 2024; Gray et al., 2019). Since *E. coli* grows
along its long axis and polysomes do not readily diffuse around the nucleoid to equilibrate (Figure 3A-D and
Figure 2 – figure supplement 3), we reduced the problem to one dimension, the cell length. Importantly, the
system is driven out of equilibrium by the continuous production and degradation of polysomes. It operates
at a nonequilibrium steady state even at fixed cell length.

First, for simplicity and illustrative purposes, we considered the case of a non-growing virtual cell with the
nucleoid initially spread throughout most of the cell to show that the repulsion between polysomes and DNA
is sufficient to both demix (phase separate) these two macromolecules and compact the nucleoid when the
simulation reaches steady state (Figure 4A). The compaction force originates from the nucleoid/polysome
steric repulsion (which drives the phase separation). The resulting higher concentration of polysomes on
each side of the nucleoid produces a difference in osmotic pressure that condenses the nucleoid. This
compaction force is consistent with drug experiments. Depletion of polysomes through inhibition of
transcription with rifampicin leads to nucleoid expansion whereas stabilization of polysomes through
inhibition of ribosome translocation with chloramphenicol results in greater nucleoid compaction (Bakshi et
al., 2014; Cabrera et al., 2009; Farrar et al., 2025; Spahn et al., 2023, 2018, 2018; Stracy et al., 2015; Xiang
et al., 2021). In the case of chloramphenicol, fusion of nucleoids has been reported (Bakshi et al., 2014;
Spahn et al., 2023). This phenomenon is expected to occur for nucleoids in close proximity at the time of
chloramphenicol treatment. These nucleoids may touch through diffusion (thermal fluctuation) and fuse to
reduce their interaction with the polysomes and minimize their conformational energy. Well-separated
nucleoids typically did not fuse. Since division could still occur during chloramphenicol treatment, the lack
of fusion between well-separated condensed nucleoids was more evident in filamenting cells inhibited for
cell division by cephalexin (Figure 4 – figure supplement 1).

Next, we tested whether adding cell growth to our model recapitulates the experimental observations.
Simulations showed that polysomes accumulate in the middle of the nucleoid during growth (Figure 4B and
Movie S3, left panel). This is followed by division of the nucleoid into two entities, which then move apart
from each other as polysomes accumulated between them. Thus, the model provides a minimal mechanism
for nucleoid segregation: At any given point, polysomes that form in the middle of the nucleoid have a lower
probability of escaping the nucleoid through diffusion compared to polysomes born at the edge of the
nucleoid (Figure 4C). Consequently, the polysome concentration rises monotonically towards the center of

the nucleoid, with its level increasing with nucleoid length (quadratically in the quasi-steady-state limit).

Once the mid-cell polysome concentration reaches a threshold (the spinodal concentration), phase

separation occurs spontaneously (i.e. via spinodal decomposition), creating a new polysome-rich phase

that splits the nucleoid in two. Compared to regions near the poles, this new phase has a higher polysome

concentration and therefore a higher osmotic pressure. This pressure difference results in a net poleward

force on the sister nucleoids that drives their migration toward the poles (Figure 4B and Movie S3, left

panel). Therefore, both phase separation (due to the steric repulsion described above) and nonequilibrium

polysome production and degradation (which creates the initial accumulation of polysomes around midcell)

are essential ingredients for nucleoid segregation.

We wondered whether our simple model could also explain the correlation between growth rate and the

relative timing of nucleoid segregation (Govers et al., 2024). Therefore, we performed simulations for

different growth rates, matching the cell and nucleoid length at birth with population-level measurements

(Figure 4 – figure supplement 2). To initialize each simulation in a realistic fashion, we used the last

timepoint (i.e., half of the predivisional cell) of the previous simulation as initial conditions, capturing the

new/old pole identity as well as any cellular asymmetries inherited between generations. The relative timing

of nucleoid splitting was measured as nucleoid depletion at mid-cell. We found that our model successfully

captures the negative trend between the growth rate and the relative timing of nucleoid splitting. Simulated

cells that grew faster also split their nucleoids earlier from birth to division, agreeing with population-level

data (Figure 4D and Movie S3). The same simulations also reproduced the relatively constant cell length

at which nucleoid splitting occurs across different growth rates (Figure 4 – figure supplement 3), which was

previously discovered in population-level measurements (Govers et al., 2024). These phenomenological

principles are expressions of the link between the absolute nucleoid length and the rate of polysome

accumulation in the middle of the nucleoid, which is explained by our mechanistic model.

To examine a potential origin for the asymmetries in polysome distribution and nucleoid compaction that

we observed (Figure 3), we examined the effect of the nucleoid diffusion coefficient D_n, which is a model

parameter that describes how fast the nucleoid relaxes towards its equilibrium configuration. Large D_n

represents the quasi-steady-state limit where the nucleoid relaxation time scale is much shorter than the

cell doubling time, and the nucleoid always assumes its (symmetric) equilibrium state. Conversely, small

D_n (i.e., slower relaxation time) can lead to asymmetric concentration profiles that will be inherited by the

daughter cells. Consistent with this expectation, we found that during relatively fast growth (~40 min

doubling time), a lower nucleoid diffusion coefficient results in a larger deviation from the equilibrium

concentration profiles for the nucleoid and polysomes (dotted curves vs. solid curves, Figure 4E). In fact,

at diffusion coefficients below 0.005 μm^2/s, the model (Figure 4E) reproduced the experimentally observed

asymmetries, including the nucleoid position offset towards the new pole at birth, the higher polysome

concentration at the new pole compared to the old one, and the asymmetric nucleoid compaction (Figure

3). Our minimal model thus suggests that the material properties of the nucleoid (e.g., stiffness) may
contribute to the observed nucleoid and polysome asymmetries in *E. coli* (see Discussion).
404
Polysomes accumulate at mid-nucleoid in DNA regions inaccessible to freely diffusing particles of
similar sizes.
In the model, the early polysome accumulation in the middle of nucleoid is caused by the nonequilibrium
processes of polysomes being born within the nucleoid while being degraded uniformly across the cell (due
to mRNA turnover). It predicts that the early mid-nucleoid enrichment of polysome signal observed in our
experiments is the product of such nonequilibrium processes rather than of polysomes simply diffusing into
undetected DNA-free space. If this is correct, freely diffusing objects of similar sizes to polysomes should
accumulate at mid-cell after polysomes accumulate there, i.e., after free DNA-space has been generated
through nucleoid splitting. To test this expectation, we compared the average distribution of RplA-msfGFP
with that of freely diffusing mCherry-labeled μNS particles from snapshot images of DAPI-stained cells
(Figure 5A). These μNS particles consist of a fragment of a mammalian reovirus protein (Broering et al.,
2005, 2002) that self-assembles into a particle, typically one per cell, when produced orthogonally in *E. coli*
(Parry et al., 2014). They have sizes between 50 and 200 nm (Parry et al., 2014; Xiang et al., 2021) similar
to polysomes (Brandt et al., 2009; Slayter et al., 1968) and are therefore largely excluded by the nucleoid
mesh (Xiang et al., 2021). After sorting cells by length into four bins, the positions of mCherry-labeled μNS
particles from approximately 2580 cells per bin were superimposed using the relative cellular coordinates
to construct particle density maps. Cells were randomly oriented in this analysis (meaning that asymmetries
between poles cannot be observed), as the pole identity cannot be assigned from snapshot images.
423
We found that in the shortest (i.e., newborn) cells, RplA-msfGFP-labeled polysomes had already
accumulated within the nucleoid (Figure 5B, bin 1). In contrast, mCherry-μNS particles were restricted to
the cell poles and were not able to access the mid-cell region until after nucleoid splitting was clearly visible
(Figure 5B, bins 2-4). This was also shown in the corresponding 1D average concentration and particle
density profiles (Figure 5C). These observations support the notion that the early mid-nucleoid
accumulation of RplA-msfGFP is caused by nonequilibrium effects associated with polysome synthesis and
degradation rather than polysome diffusion into DNA-free space.
431
Arrest of polysome production immediately stops nucleoid segregation while polysome depletion
gradually reverses it.
Our correlative analyses and model (Figures 1-4) support the hypothesis that the interactions and ensuing
exclusion between polysomes and nucleoid promote nucleoid segregation and macromolecular
asymmetries along the cell length. To probe causality, we used two tests. The first one aimed to disrupt the
proposed mechanism using rifampicin. Rifampicin treatment is known to homogenize ribosome distribution
and expand the nucleoid over time through polysome depletion (Bakshi et al., 2014; Dworsky and

Schaechter, 1973; Koch and Gross, 1979; Pettijohn and Hecht, 1974; Xiang et al., 2021). However, our
hypothesis predicts a faster effect on nucleoid segregation. Blocking transcription should instantly reduce
the rate of polysome production to zero, causing an immediate arrest of nucleoid segregation. Gradual
depletion of the existing polysomes due to mRNA decay should then cause, on a slower time scale, a
dissipation of the phase separation between DNA and ribosomes.
444
To test these predictions, we subjected cells growing in M9gluCAAT in microfluidic channels to two rounds
of rifampicin treatment (Figure 6A and Movie S4). Rifampicin resulted in growth rate inhibition (Figure 6A)
and changes in the nucleoid area and nucleoid-to-cell area (NC) ratio (Figure 6 – figure supplement 1), as
previously described (Bakshi et al., 2014; Dworsky and Schaechter, 1973; Koch and Gross, 1979; Pettijohn
and Hecht, 1974; Xiang et al., 2021). In cells that completed their division cycle before antibiotic addition
(squares and fitted blue curve in Figure 6B), the distance between the intensity peak of each sister nucleoid
increased monotonically between birth and division, displaying the dynamics of normal, unperturbed
nucleoid segregation. Cells born 22 min to 12 min before the treatment (circles in Figure 6B) experienced
the same nucleoid segregation dynamics up to the time of rifampicin addition. Exposure to rifampicin led to
the near-immediate arrest of nucleoid segregation (Figure 6B), consistent with our prediction.
455
Interestingly, the distance between the sister nucleoids remained the same for close to 30 min into
rifampicin treatment, after which it started to decrease (Figure 6B). Average 1D and 2D cell projections of
the scaled (divided by the whole cell average) RplA-GFP and HupA-mCherry concentration suggest that
the time delay between the arrest in nucleoid segregation and its reversal is likely due to the compressible
nature of the nucleoid, which has been demonstrated in vitro (Pelletier et al., 2012). As the polysomes in
the middle of the cells started to visibly deplete (> 6 min after rifampicin addition), the DNA signal expanded
to fill the emerging available space without affecting the distance between the peaks of the sister nucleoids
(white crosses, Figure 6C). This is consistent with the removal of a compaction force exerted by the
accumulating polysomes on the soft nucleoid. About 30 min after rifampicin addition and further polysome
depletion, the peak signals of the sister nucleoids (white crosses, Figure 6C) started migrating closer to
each other. Eventually, after 1 h of rifampicin treatment, when the RplA-GFP fluorescence was
homogeneous, the two nucleoid objects fused into one (Figure 6C), consistent with the dissipation of phase
separation.
469
These results support the notion that in untreated cells, polysome accumulation effectively exerts a force
on the compressible nucleoid, which translates into its observed compaction and translocation (hence,
segregation). Gradual polysome depletion through rifampicin treatment progressively decreased this effect,
reversing the process. This reversal became obvious when we plotted the correlation between the relative
polysome accumulation and nucleoid depletion at mid-cell for two cell lineages that experienced rifampicin
at different times after birth (Figure 6D). Irrespective of their birth time (12 or 3 min before the addition of

rifampicin), the negative correlation between the two variables was reversed ~9 min after the addition of
the antibiotic, following the same path as for the untreated cells but in the opposite direction (Figure 6D).
Polysome depletion during rifampicin treatment also resulted in correlated loss of asymmetric nucleoid
compaction in newborn cells (Figure 6E). Altogether, these results support the notion that the asymmetric
accumulation of polysomes results in an anisotropic force that asymmetrically compacts and segregates
nucleoids.

Ectopic polysome production redirects nucleoid dynamics.
Our second approach to test causality experimentally was to redirect polysome production away from the
chromosome to achieve polysome accumulation at an ectopic site. To achieve polysome accumulation at
ectopic sites in experiments, we overexpressed a useless protein (mTagBFP2) for the cell from a T7
promoter on a multi-copy pET28 plasmid (Figure 7A). The resulting CJW7798 strain also carried the
ribosome (RplA-msfGFP) and DNA (HupA-mCherry) markers. We reasoned that high expression of BFP2
from the plasmid would slow polysome production within the nucleoid and create polysome accumulations
at ectopic cellular locations through recruitment of ribosomes to plasmid transcripts. This, in turn, should
affect nucleoid dynamics according to our model.

Plasmid expression of mTagBFP2 was induced by addition of 100 µM IPTG and expression of a
chromosomally encoded T7 RNA polymerase (Figure 7B). The gradual increase of mTagBFP2
fluorescence in the cells was associated with a concomitant decrease in cell growth rate (Figure 7C),
consistent with reduced gene expression from the chromosome. We verified by flow cytometry that
induction did not block DNA replication. IPTG-induced cells displayed a similar scaling relationship between
the intensity of DNA (labeled with DRAQ5) and cell size (side scatter area) than uninduced cells (Figure 7
– figure supplement 1).

We found that induced cells displayed various patterns of polysome accumulations (Movie S5), presumably
due to stochastic clustering of plasmids in DNA-free regions as previously reported for multi-copy plasmids
devoid of DNA partitioning genes (Hsu and Chang, 2019; Reyes-Lamothe et al., 2014; Yao et al., 2007).
Importantly, the ectopic accumulations of polysomes had a drastic effect on nucleoid dynamics. In some
cells, polysomes accumulated at one pole instead of the middle of the nucleoid, preventing the nucleoid
from splitting (Figure 7D). Expansion of the polysome accumulation at a pole effectively pushed the nucleoid
toward the opposite pole of the cell. In other cells, polysome accumulation occurred between sister
nucleoids, but did not relocate to the segregated nucleoids at the ¼ and ¾ cell positions. Rather, polysome
accumulation persisted and expanded between the sister nucleoids, effectively further pushing them apart
(Figure 7E). Both of these phenotypes were reproduced by our model when we intentionally caused an
accumulation of polysomes, either at a pole or between segregated nucleoids (Figure 7 – figure supplement
2). In our experiments we also observed filamenting cells with correlated polysome and nucleoid dynamics

that changed in time (Figure 7F), resulting in transient events of nucleoid fusion, splitting, or changes in

migration direction depending on where polysomes accumulated. Movie S5 shows additional examples of

such dependency.

Upon induction, we observed the appearance of diffuse mTagBFP2 fluorescence (Figure 7B). We also

observed the formation of inclusion bodies (bright phase contrast) that typically remained at a pole or

sometimes moved along the edge of a growing polysome accumulation (Figure 7F). After a long of period

of IPTG induction (> 8 h), polysome accumulation eventually decreased, leading to nucleoid decompaction

(Figure 7 – figure supplement 3).

This experiment effectively decoupled polysome accumulation from cell growth. By redirecting a substantial

fraction of chromosome gene expression to a single plasmid-encoded gene, we reduced the rate of cell

growth but still created a large accumulation of polysomes at an ectopic location. This ectopic polysome

accumulation was sufficient to affect nucleoid dynamics in a correlated fashion. Altogether, these results

support the notion that ectopic polysome accumulation drives nucleoid dynamics.

Cell width enlargement leads to nucleoid splitting along the incorrect cell axis and to the fusion of

polysome accumulations from distinct DNA-free regions.

A previous study on *Bacillus subtilis* L-forms suggests that the width of the cell is also important for nucleoid

segregation (Wu et al., 2020). In that study, L-forms, which are spherical cells stripped of their cell wall,

were squeezed into narrow microfluidic channels of similar width than the diameter of walled rod-shaped

cells. Growth in these channels resulted into elongated cells with improved efficiency in nucleoid

segregation (Wu et al., 2020). In normal (walled) cells, the nucleoid is kept close to the cytoplasmic

membrane across the cell width, likely due to transertion (co-transcriptional translation and translocation of

membrane and secreted proteins) (Bakshi et al., 2014; Spahn et al., 2023; Youngren et al., 2014). We

reasoned that this geometric constraint reproduces some important aspects of our simple model. The

nucleoid proximity to the cell membrane along the cell width ensures that the nucleoid grows exclusively

along one dimension, the cell length. Furthermore, the nucleoid attachment to the membrane along this

radial cell axis may act as a diffusion barrier and limit polysome exchange between distinct DNA-free

regions.

To test these ideas, we first treated cultures with A22 to inhibit cell width control through the inactivation of

MreB (Bean et al., 2009; Iwai et al., 2002). This resulted in cells with a polysome phase at the cell center

surrounded by a nucleoid phase around the cell periphery (Figure 8 – figure supplement 1, top). The

peripheral nucleoid localization is likely due to membrane attachment through transertion, as previously

shown (Spahn et al., 2023). To further increase the cell width (> 2.5-fold), we exposed cells to the cell

division inhibitor cephalexin in addition to A22 (Figure 8A and Figure 8 – figure supplement 1, bottom). As

a control, we showed that cephalexin treatment alone resulted in filamentous cells (of constant cell width)
with multiple nucleoids separated by polysome accumulations (Figure 8B), consistent with previous reports
(Chai et al., 2014; Gray et al., 2019; Thappeta et al., 2024). In cells treated with both drugs, we observed
two types of subcellular rearrangements. In smaller cells, which started with a single nucleoid, drug
treatment resulted in a single large polysome accumulation at the cell center, with the nucleoid displaying
a toroidal shape at the cell periphery (Figure 8C and Movie S6). In longer cells with two segregated
nucleoids, the nucleoids expanded and often aberrantly segregated along the cell width concomitant with
polysome accumulation at the site of nucleoid splitting (white arrowheads, Figure 8D). Consequently, a
polysome "bridge" was formed between the polysome accumulations flanking the nucleoid. These
polysome bridges resulted in a characteristic cross-like polysome pattern, marking the two axes
(longitudinal and radial) of nucleoid segregation (Figure 8E). Cell width enlargement led to coalescence of
polysome accumulations toward the cell center and fusion of nucleoids around the cell periphery (Figure
8C-E and Figure 8 – figure supplement 1, bottom). As a result, the nucleoids and polysome accumulations
decreased in number while increasing in size in cephalexin/A22-treated cells compared to cephalexin-
treated cells with the same cell area distribution but normal cell width (Figure 8F). These results suggest a
critical role for cell width regulation in limiting the diffusion of polysomes around the nucleoid, thereby
promoting nucleoid segregation specifically along the cell length.
567

DISCUSSION
The flow of genetic information intrinsically couples nucleoid segregation to cell growth.
This study provides experimental and theoretical evidence (Figures 1-8) that polysome production within
the nucleoid—an inherent product of chromosomal gene expression—contributes to nucleoid segregation
and positioning in *E. coli* cells. This may also be true in other bacteria, as reduced or abrogated transcription
via gene deletion or antibiotic treatment causes chromosome segregation defects in *Streptococcus*
pneumoniae (Kjos and Veening, 2014) and *Bacillus subtilis* (Dworkin and Losick, 2002).
575

An appealing feature of this proposed model is that polysomes inherently integrate the rate of nucleoid
segregation with that of gene expression and cell growth. The concentration of polysomes and their rate of
accumulation in the cell are a direct reflection of transcriptional and translational activities in the cell
(Balakrishnan et al., 2022). The higher the concentration of polysomes, the faster the growth rate becomes.
Thus, in our proposed model, an increase in polysome concentration not only leads to more protein
synthesis and faster cell growth but also results in faster nucleoid segregation. The reverse is true for a
decrease in polysome concentration, inherently coupling these processes without the help of a dedicated
regulatory system. Such coupling was observed across isogenic cells with variable growth rates in the same
nutrient condition (Figure 2 – figure supplement 2) as well as across nutrient conditions that led to a wide
range of growth rates (Figure 2 and Figure 2 – figure supplement 1).
586

Directional nucleoid splitting requires DNA/polysome exclusion and cell width control.
A key aspect of our proposed mechanism for nucleoid segregation is that polysomes form within nucleoids
due to chromosomal gene expression, which, together with polysome turnover due to mRNA decay, creates
an out-of-equilibrium system (Figure 4) (Miangolarra et al., 2021). Redirecting polysome formation to
plasmid gene expression leads to ectopic polysome accumulations that is sufficient to alter nucleoid
dynamics (Figure 7 and Movie S5).
593
In normal cells, the effective force that segregates nucleoids appears to be linked to the propensity of the
chromosomal meshwork and polysomes to separate from each other. Mutual exclusion is, at least in part,
caused by the steric repulsion between the two macromolecules. While ribosomes or ribosomal subunits
freely diffuse across the cell unobstructed by the presence of the nucleoid (Bakshi et al., 2012; Sanamrad
et al., 2014), the larger polysomes are impeded by the chromosomal mesh based on size considerations
alone (Xiang et al., 2021). Modeling studies have suggested that such steric hindrance between large
crowders (polysomes) and a polymeric meshwork (chromosome) can result in phase separation and
polymer compaction (Bakshi et al., 2014; Castellana et al., 2016; Miangolarra et al., 2021; Mondal et al.,
2011; Wu et al., 2019b). Theoretically, mRNAs alone are large enough to phase separate from DNA, though
to a lesser degree than polysomes (Miangolarra et al., 2021). It is also possible that phase separation
between nucleoids and polysomes (or mRNAs) involves non-steric interactions such as electrostatic
repulsion between the negatively charged DNA and RNA (mRNA and rRNA), as previously hypothesized
(Joyeux, 2015). Such steric and non-steric interactions may contribute to the effective poor solvent quality
of the polysome-rich cytoplasm for the chromosome (Xiang et al., 2021). Future research will be necessary
to elucidate the precise nature of the interaction between chromosomes and mRNAs, whether individually
or in complex with ribosomes.
610
We found that the width of the cell controls the exclusion dynamics between polysomes and nucleoids by
directing nucleoid growth and segregation along a single cellular dimension, cell length (Figure 8). The
close proximity of the nucleoid to the membrane presumably due to transertion (Bakshi et al., 2014;
Rabinovitch et al., 2003; Spahn et al., 2023) effectively limits the diffusion and fusion of polysome
accumulations from distinct DNA-free regions, which leads to alternating enrichments of polysomes and
DNA along the length of elongating cells (Figure 8B). Limited polysome diffusion around the nucleoids is
consistent with the observed differences in ribosome concentrations between DNA-free regions (Figure 3A-
B, and D). Otherwise, we would expect polysome enrichments on each side of the nucleoids to rapidly
equilibrate in concentration. Diffusion limitation around nucleoids is consistent with our previous report that
polysomes diffuse much faster over short distances (within DNA-free domains) than long distances (across
DNA-free domains) (Gray et al., 2019). Control of cell width is vital to uphold this constraint and promote
that nucleoid grows along a single cell axis. This explains the drastic improvement in nucleoid segregation
of wall-less cells (L-forms) when compressed to normal width (Wu et al., 2020). It highlights the importance

of cell width regulation and suggests that nucleoid segregation may have imposed an evolutionary

constraint on cell width control.

Nucleoid segregation likely involves multiple factors.

The model shows that the most trivial case of uniform production of polysomes (i.e., uniform mRNA

synthesis) within the nucleoid is sufficient to cause an enrichment of polysomes at mid-nucleoid (Figure 4)

(Miangolarra et al., 2021). Inside cells, this polysome enrichment at mid-nucleoid may be enhanced by a

bias in mRNA synthesis across the nucleoid. For instance, the chromosomal region close to the origin of

replication has been shown to be more highly expressed per gene copy than other regions on the

chromosome (Scholz et al., 2019) and this region is located in the middle of the nucleoid prior to DNA

replication (Bates and Kleckner, 2005; Cass et al., 2016; Fisher et al., 2013; Kuwada et al., 2013, 2013;

Mäkelä et al., 2021; Sadhir and Murray, 2023; Wang et al., 2006). In addition, this highly expressed

chromosomal region is the first one to replicate, which should lead to further local increase in mRNA

expression due to a doubling in gene dosage (Pountain et al., 2022). We did not consider such localized

mRNA synthesis in our model. However, if we did, it would only help the mechanism that we proposed by

increasing the polysome built-up at mid-nucleoid.

Other factors are likely involved in nucleoid segregation. In fact, our data revives the largely abandoned

sixty-year-old hypothesis by Jacob et al. (1963) that cell growth separates the sister nucleoids through their

potential attachment to the cell wall, but with two notable differences. First, the contribution of cell growth

to nucleoid splitting would be minor relative to the polysome contribution (Figure 1I and Figure 1 – figure

supplement 5). Second, cell growth would contribute predominantly near the end of the division cycle

(Figure 1I). This late timing would be attractive for two reasons. First, it corresponds to the time when

polysomes stop accumulating between the separated sister nucleoids and polysome accumulations

emerge at the middle of these nucleoids (i.e., at the ¼ and ¾ cell positions) to start the next round of

segregation (Figure 1D and Movie S2). Second, this is also when *E. coli* switches its cell wall growth pattern

from dispersed along the cell body to zonal and divisome-dependent at mid-cell (Cooper and Hsieh, 1988;

Gray et al., 2015; Navarro et al., 2022; Wientjes and Nanninga, 1989; Woldringh et al., 1987). Indeed, zonal

cell growth between the sister nucleoids was a key assumption of the 1963 model (Jacob et al., 1963).

What is not entirely clear is how the DNA would be attached to the peptidoglycan cell wall. Transertion links

the DNA to the cytoplasmic membrane. Perhaps the coupling between transcription, translation, and

membrane insertion extends to peptidoglycan binding.

Beyond cell elongation, thermodynamic demixing and other cellular processes such as DNA replication,

loop extrusion, supercoiling, and preferential loading of DNA remodeling complexes are also likely to be

important for robust chromosome segregation and organization (Danilova et al., 2007; Harju et al., 2024;

Hofmann et al., 2019; Holmes and Cozzarelli, 2000; Lemon and Grossman, 2000; Mäkelä et al., 2021;

661 Minnen et al., 2011; Sawitzke and Austin, 2000; Weitao et al., 1999; Wu et al., 2019a; Youngren et al.,

662 2014).

664 We note that in our time-lapse experiments, the accumulation of polysome signal appeared to slightly

665 precede the depletion of DNA signal that marked the initiation of nucleoid splitting (Figure 1E). Polysome

666 enrichment in the middle of unconstricted nucleoids was also occasionally observed in snapshot images of

667 cells growing on glycerol, a slow growth condition that results in a single nucleoid segregation event late

668 during the cell division cycle (Figure 1 – figure supplement 6). This is not seen in our model in which

669 polysome accumulation and nucleoid splitting occur at the same time (Figure 4B and Movie S3). This small

670 discrepancy may reflect a limitation of our experimental or modeling approach. For example, it is possible

671 that the point spread function of our fluorescent DNA marker slightly delays the moment at which we can

672 detect signal depletion at mid-nucleoid and thereby the initiation of nucleoid splitting. Alternatively, the small

673 difference in timing may be associated with a model simplification. In our model, the nucleoid is effectively

674 a solution of DNA fragments. In reality, the nucleoid consists of a circular polymer, crosslinked by nucleoid-

675 associated proteins. These DNA crosslinks may cause a small resistance that marginally delays the

676 initiation of nucleoid splitting relative to the polysome enrichment at mid-nucleoid.

678 **E. coli *is an asymmetric organism.***

679 The spatial macromolecular asymmetries uncovered in our study contrast with the common perception of

680 *E. coli* as a symmetric organism. In our relatively rich growth medium (M9gluCAAT), the distribution of

681 polysomes at the new pole in newborn cells was, on average, higher than at the old pole through inheritance

682 of the large mid-cell accumulation of polysomes from their mother cells (Figure 3A and D). We also

683 observed an asymmetric distribution in DNA density within nucleoids, which correlated with the availability

684 of polysome-free space along the cell length and width (Figure 3E-G). This asymmetry emerged before cell

685 division (Figure 3D and Movie S2). Simulations of our reaction-diffusion model suggest that slower

686 diffusing/relaxing nucleoids are more likely to reproduce these nucleoid position and compaction

687 asymmetries during the finite course of the cell division cycle (Figure 4E). A reduction in the apparent

688 nucleoid diffusivity has been linked to nucleoid-associated proteins, which bridge and thus stiffen the DNA

689 polymer (Subramanian and Murray, 2023). The physiological significance for these dynamic cellular

690 asymmetries is not clear at this time, though it is conceivable that a difference in DNA compaction within

691 the nucleoid may affect gene expression. Regardless, our study illustrates how spatial and temporal

692 asymmetries in the cytoplasm can emerge from the interactions between two of the most important

693 macromolecules in the cell.

697   **METHODS**
698
699   **Strains and constructs**
700   Strains and plasmids used for this study are listed in Tables S2 and S3, respectively, while the sequences
701   of the oligonucleotides used to make constructs can be found in Table S4.
702
To measure the concentration and spatial heterogeneity of ribosomes inside *E. coli*, we used strains in
which the RplA 50S ribosome subunit protein (strains CJW7323, CJW6768, CJW7020 and CJW7651) or
the RpsB 30S ribosome subunit protein (strains CJW6769 and CJW7021) are fused with mEos2 (strains
CJW6768 and CJW6769), msfGFP (strains CJW7020, CJW7021, CJW7651, and CJW7798) or GFP
(CJW7323) (Gray et al., 2019; Xiang et al., 2021). Nucleoid characteristics were measured using strains in
which HupA, a nucleoid-associated protein, is fused with mCherry (strains CJW7323, CJW6723 and
CJW7798) (Xiang et al., 2021). Alternatively, DAPI was used to stain the DNA (strains CJW6768,
CJW6769, CJW7020, CJW7021, and CJW7651).
711
The mCherry-µNS (CJW7651) particles were chromosomally expressed from the native Lac promoter after
induction with 150 µM IPTG (for 3 h for agarose pad experiments, or continuously for microfluidics
experiments). Strain CJW7651 was constructed as follows. The *gfp* coding sequence in the pER12
(pBAD322A-gfp-µNS) plasmid (kind gift from Dr. A. Janakiraman, City College of New York) was swapped
with the mCherry-coding sequence using megaprimer whole plasmid (MEGAWHOP) cloning (Bryksin and
Matsumura, 2010) and the primer pair ER12-MCR-fwd/ER12-MCR-rev2 to generate plasmid pER12-
mCherry. The mCherry-µNS coding sequence and the *rrnB* transcriptional terminator were amplified from
the pER12-mCherry plasmid (primer pair µNSmCherry fwd/rev) and assembled with the *frt* site flanked with
a kanamycin resistance cassette (amplified from pKD13 (Datsenko and Wanner, 2000) using the primer
pair FRT_KanR fwd/rev) and the ColE1 origin of replication (PCR amplified from the pBAD22A plasmid
(Guzman et al., 1995) using the primer pair ColE1 fwd/rev) using Gibson DNA assembly (Gibson et al.,
2009) to form the pAPG1 plasmid. The pAPG1 plasmid also included two 50 bp sites homologous to the
attB region of the *E. coli* chromosome, introduced as overhangs in the primers used for Gibson DNA
assembly. The *attB* homologous regions allowed for the integration of the arabinose inducible mCherry-
µNS expression cassette into the respective site, though this was not used in this study. The pAPG1
plasmid was verified by sequencing using the pAPG1 seq1-5 primers. The mCherry-µNS coding sequence
was then PCR amplified from the pAPG1 plasmid using primer pair lacZYA_redµNS fwd/rev, which includes
50-bp overhangs homologous to the region upstream and downstream of the *lacZYA* operon, and
integrated downstream of the *lac* promoter in the MG1655 strain (Guyer et al., 1981; Jensen, 1993) using
lambda red recombination and the pKD46 plasmid (Datsenko and Wanner, 2000) for the construction of
the CJW7144 strain. Correct insertion of the mCherry-µNS coding sequence to substitute the *lacZYA*
operon coding sequences was confirmed by colony PCR using primer pairs LacI_fwd/CynX_rev and
LacI_fwd/mCherry_rev. Transduction with a P1 phage lysate of the CJW7144 strain was used to transfer

the mCherry-µNS expression cassette into the CJW7020 strain using kanamycin as a selection marker. As
a result, the CJW7145 strain was constructed, which was verified using colony PCR and the primer pairs
LacI_fwd/CynX_rev and LacI_fwd/mCherry_rev, KanR_fwd/CynX_rev. The CJW7145 strain was then
transformed using the pCP20 plasmid (Datsenko and Wanner, 2000), which encodes the FLP recombinase
(Cherepanov and Wackernagel, 1995) to remove the kanamycin resistance cassette, yielding the CJW7651
strain.
741
The CJW7798 strain was derived from the MG1655 (DE3) strain (Tseng et al., 2010a), which carries the
T7 RNA polymerase-encoding gene under *lacUV5* control. Transduction with a P1 phage lysate of the
CJW5158 strain (Gray et al., 2019) was used to transfer the gene encoding the HupA-mCherry fusion into
the MG1655 (DE3) strain using kanamycin as a selection marker. The pCP20 plasmid (Datsenko and
Wanner, 2000), which encodes the FLP recombinase (Cherepanov and Wackernagel, 1995), was used to
remove the kanamycin resistance cassette, yielding the CJW7466 strain. Transduction with a P1 phage
lysate of the CJW7019 strain, a kanamycin-resistant intermediate of strain CJW7020 (Gray et al., 2019),
was used to transfer the gene encoding the RplA-msfGFP fusion into the CJW7466 strain using kanamycin
as a selection marker. The resulting strain (CJW7766) was grown in the presence of kanamycin since the
cells tend to lose msfGFP fluorescence in the absence of the antibiotic. The sequence of the CJW7766
strain was confirmed by whole genome sequencing.
753
The pET28:mTagBFP2 plasmid variant was derived from the pET28:GFP plasmid (Shis and Bennett,
2013), which was a gift from Mathew Bennett (Addgene plasmid # 60733 ; http://n2t.net/addgene:60733;
RRID:Addgene_60733). First, the GFP coding sequence was substituted with the mTagBFP2 coding
sequence from the pBAD-mTagBFP2 plasmid (Subach et al., 2011), a gift from Vladislav Verkhusha
(plasmid # 34632; http://n2t.net/addgene:34632; RRID:Addgene_34632). Specifically, the mTagBFP2
coding sequence was amplified using the primer pair mTagBFP2 fwd/rev and assembled (Gibson DNA
assembly) with the pET28 backbone, which was amplified in two pieces using the pET28_one fwd/rev and
the pET28_two fwd/rev primer pairs to derive the pET28:mTagBFP2 plasmid. Then, the kanamycin
resistance cassette that was originally present in the pET28 backbone was substituted with the
chloramphenicol resistance cassette from the pSB3C5-proA-B0032-E0051 plasmid (Davis et al., 2011)
which was a gift from Joseph Davis and Robert Sauer (Addgene plasmid # 107244;
http://n2t.net/addgene:107244; RRID:Addgene_107244). Specifically, the backbone of the
pET28:mTagBFP2 plasmid excluding the kanamycin resistance cassette was amplified using the
pET28mTagBFP2 fwd/rev primers and assembled (Gibson DNA assembly) with the coding sequence of
the chloramphenicol resistance cassette that was amplified using the cmR fwd/rev primer pair. As a result,
the pET28:mTagBFP2-CmR plasmid was created. The proper assembly of the pET28:mTagBFP2 and
pET28:mTagBFP2-CmR plasmids were confirmed by sequencing.
771

772   The pET28:mTagBFP2-CmR plasmid was introduced into CJW7766 by electroporation to create the
773   CJW7798 strain, which was used to redirect the ribosomes away from the nucleoid onto the plasmid-
774   expressed mTagBFP2 mRNAs after inducing the expression of T7 RNA polymerase with IPTG.
775
Growth conditions
Strains CJW6769, CJW7020 and CJW7021 used in Figure 2 and Figure 2 – figure supplement 1 were
grown as previously described (Gray et al., 2019) using a basic M9 medium formulation without trace
elements and supplemented with 0.2% (w/v) carbon source and when specified with 0.1% (w/v) casamino
acids (CAA) and 1 µg/mL thiamine (T). The strain CJW6768 (used in Figure 2 and and Figure 2 – figure
supplement 1) was grown using the same medium formulation. The abbreviations of the medium growth
conditions presented in Figure 2A and Figure 2 – figure supplement 1 are defined in Table S1. For the rest
of our experiments on agarose pads or in a microfluidic device, strains CJW7323, CJW7651 and 7798 were
grown in M9 salts (final concentrations: 33.7 mM Na_2HPO_4, 22 mM KH_2PO_4, 8.55 mM NaCl, 9.35 mM
NH_4Cl, 1 mM $MgSO_4$, 0.3 mM $CaCl_2$) supplemented with trace elements (Fe, Zn, Cu, Co, B, Mn), 0.1%
(w/v) thiamine, and, when specified, 0.4% (w/v) casamino acids. The pH of the 10x M9 salts was adjusted
to 7.2 with NaOH. The trace elements were added at a final concentration of 13.4 mM ethylene-diamine-
tetra-acetic-acid, 3.1 mM of $FeCl_3$-$6H_2O$, 0.62 mM of $ZnCl_2$, 76 µM of $CuCl_2$-$2H_2O$, 42 µM of $CoCl_2$-$2H_2O$,
162 µM of HBO_3-$2H_2O$, 8.1 µM of $MnCl_2$-$4H_2O$.
790
To achieve steady-state exponential growth prior to imaging, a stationary phase liquid culture in the
appropriate growth medium was diluted at least 10,000 times and grown to an optical density at 600 nm
(OD_{600}) between 0.1 and 0.3. These cells were either loaded in a mother-machine-type microfluidics device
(Lin and Jacobs-Wagner, 2022; Wang et al., 2010) where they were grown under the constant flow of
medium (~0.5 µL/sec), or spotted on 1% agarose pads prepared with the same growth medium. In the
microfluidic device, cells were grown for at least 3 h prior to image acquisition. All precultures and
experiments were performed at 30ºC.
798
799   The mCherry-µNS particles (CJW7651 strain) were imaged on agarose pads after inducing exponentially
800   growing cells with 150 µM IPTG for 3 h (Figure 4E-G). Before spotting on the 1% agarose pad, the induced
801   cells were stained with DAPI (1 µg/mL) for 5 min. In both experiments, the cells were growing in
802   M9gluCAAT.
803
804   To redirect ribosomes to plasmid-expressed, T7 promoter-driven mTagBFP2 mRNAs, exponentially
805   growing CJW7798 cells grown in M9glyCAAT to an OD ~ 0.2 were spotted on a 1% agarose pad containing
806   M9glyCAAT and 100 µM IPTG, the latter to induce the expression of the T7 RNA polymerase.
807
808
809

Rifampicin treatment

The antibiotic rifampicin (see Table S5) was used to block transcription, deplete mRNAs, and release the mRNA-bound ribosomes (polysomes). Rifampicin treatment was performed either in batch culture (Figure 1 – figure supplement 2) or in microfluidics (Figure 6 and Movie S4).

For the microfluidic experiment, two rounds of rifampicin treatment were performed by switching between M9gluCAAT containing antibiotic (100µg/mL) and antibiotic-free medium at 30ºC. For the medium switches, solenoid valves were used in a custom-built pressurized perfusion system, achieving fast (within 1 min) changes in the cellular environment. The first switch to rifampicin occurred 2 h after normal growth in antibiotic-free medium (Figure 6A and Movie S4). Rifampicin treatment lasted 2 h until the system switched back to antibiotic-free medium, where cells were left to recover for 8 h. Twelve hours into the experiment, the system switched again to rifampicin-containing medium.

A22 and cephalexin treatment

Cephalexin (50 µg/mL) and A22 (4 µg/mL) (see Table S5) were added to exponentially growing CJW7323 cell cultures (OD_{600} ~ 0.1) in M9gluCAAT just prior to spotting on a 1% agarose pad, which was made with the same growth medium and contained the same antibiotic concentrations. The radial expansion of the cells and the polysome and nucleoid signals were tracked over time in a time-lapse experiment. The selected concentration of A22 has previously been shown to not affect cell growth (Takacs et al., 2010). All precultures and time-lapse imaging experiments were performed at 30ºC.

Chloramphenicol and cephalexin treatment

Exponentially growing CJW7323 cells (OD_{600} ~ 0.1) in M9gluCAAT were pre-treated with cephalexin (50 µg/mL) for 1 h in the shaking flask. Then the cells were spotted on a 1% agarose pad, which was made with the same growth medium and contained cephalexin (50 µg/mL) and chloramphenicol (75 µg/mL). All precultures and time-lapse imaging experiments were performed at 30ºC.

Ectopic polysome accumulation experiment

Exponentially growing CJW7798 cells in M9glyCAAT supplemented with kanamycin (50 µg/mL) and chloramphenicol (35 µg/mL) were washed one time with M9glyCAAT minimal medium lacking antibiotics but supplemented with 100 µM IPTG (see Table S5). The washed cells were spotted on a 1% agarose pad prepared with M9glyCAAT also supplemented with IPTG (100 µM) for time-lapse imaging. All pre-cultures and time-lapse imaging experiments were performed at 30ºC. Since chloramphenicol affects protein synthesis and thus polysome formation, the agarose pads lacked chloramphenicol, resulting in a fraction that loss the plasmids based on the absence of blue fluorescence. Therefore, only cells that expressed blue fluorescence (and thus carried the pET28:mTagBFP2-CmR plasmid) were analyzed.

DNA quantification using flow cytometry

Exponentially growing CJW7798 cells (OD$_{600}$ ~1) in M9glyCAAT supplemented with kanamycin (50 μg/mL) and chloramphenicol (35 μg/mL) were washed with M9glyCAAT minimal medium lacking antibiotics. The washed cells were used to start two cultures in M9glyCAAT without antibiotics. In one culture, the expression of the mTagBFP2 was induced with 100 μM ITPG. In the second culture, the cells were not supplemented with IPTG (i.e., no induction). After 200 min, the cells were fixed with 70% cold ethanol for 30 min, followed with stained with 5μM DRAQ5 (Invitrogen^{TM} eBioscience^{TM} Cat. 65-0880-92) for 30 min at 37ºC as previously described (Silva et al., 2010). The stained cells were washed with 0.1 Tris, 2mM MgCl$_2$ buffer (pH 7.4) and then used for flow cytometry. The Attune^{TM} CytPix flow cytometer (Invitrogen^{TM}) was used to quantify the DRAQ5 fluorescence in the red laser 2 (RL2) channel (637 nm excitation, 690DLP beamsplitter, 720/30 emission, 440 voltage) and the cell size in the side scatter (SSC) channel (320 voltage). A low threshold of 0.1x10^3 was set for the forward scatter and 0.3x10^3 for the side scatter signal. The flow was set at 12.5 μL/min for a total of 10^5 counted events per sample. The stained cells were diluted appropriately (usually 100-fold) to ensure less than 10^3 events per second. For each biological replicate, the DNA content (RL2-Area) and the cell size (SSC-Area) were quantified. Lower thresholds in the RL2-Area and the SSC-Area channels were applied to exclude very small events without fluorescence, as well as a polygon gate in the SSC-Area vs. SSC-Height statistics to exclude events with more than one cells. These gates were first applied on the data from the uninduced sample and then replicated on the data from the induced sample for each biological replicate. As a result, the exact same thresholds were applied between the induced and uninduced samples per biological replicate for fair comparison. The analysis of the flow cytometry data was performed using a custom Python package (*flowio_to_pandas*), which is based on the FlowIO flow cytometry standard (FCS) file parser (White et al., 2021) and which allows for interactive gating.

Microscopy

Strains CJW6769, CJW7020, and CJW7021 used in Figure 2A-B and Figure 2 – figure supplement 1, were imaged on agarose pads using the same microscopy set-up and optical configurations as previously described (Gray et al., 2019). Snapshots of the CJW6768 strain (used in Figure 2 and Figure 2 – figure supplement 1) were taken using a Nikon Ti-E microscope equipped with a 100x Plan Apo 1.45NA Ph3 oil objective, a Hamamatsu Orca-Flash4.0 V2 CMOS camera (16-bit, Slow Scan sensor mode) and a Lumencor Spectra X LED (Light Emitting Diode) engine. The 395/25nm LED was used to excite DAPI and the 470/24nm LED was used to excite GFP. DAPI fluorescence was acquired using an ET350/50x (excitation filter), RT400lp (dichroic mirror), ET460/50m (emission filter) filter cube from Chroma. The ET470/40x (excitation filter), T495lpxr (dichroic mirror), ET525/50m (emission filter) configuration from Chroma was used for GFP fluorescence acquisition. The microscope was controlled using the NIS Elements software by Nikon and the ND acquisition module.

For the time-lapse observation of cells (strain CJW7323) treated with chloramphenicol, A22 and/or cephalexin, images were taken using a Nikon Ti-E microscope, equipped with a 100x Plan Apo 1.45NA Ph3 oil objective, a Hamamatsu Orca-Flash4.0 V2 CMOS camera (16-bit, Slow Scan sensor mode), and a Sola solid state white light source (Lumencor), in a temperature-controlled enclosure (Okolabs). The AT470/40x excitation filter, combined with a T495LPXR beam-splitter and an ET525/50m emission filter, was used for GFP. For mCherry visualization, the ET560/40x excitation filter, combined with a T585lp beam-splitter and a ET630/75m emission filter, was used. A neutral density filter (32x) was applied in the excitation path to reduce phototoxicity and photobleaching. Images were taken every 2.5 min in the brightfield channel (phase contrast) and every 5 min in the fluorescence channels (RplA-GFP and HupA-mCherry).

For the rest of the brightfield and epi-fluorescence wide-field microscopy experiments (strains CJW7323, CJW7651, and CJW7798), snapshots or time-lapse images were taken using a Nikon Ti2-E inverted microscope, equipped with a 100x Plan Apo 1.45NA Ph3 oil objective, a Photometrics Prime BSI back-illuminated sCMOS camera (2048x2048 pixels sensor with a pixel size of 6.5 μm), and a Lumencor Spectra III LED (Light Emitting Diode) engine, in a temperature-controlled enclosure (Okolabs). The HDR 16-bit sensor mode was used to acquire images. The microscope was controlled using the NIS Elements software by Nikon, and the JOBS or the ND acquisition module were used to acquire snapshots or time-lapse images. The Perfect Focus System (by Nikon) was used to maintain focus in microfluidics experiments. The auto-focus function in NIS Elements was used to locate the optimal z-position in agarose pad experiments (snapshots or time-lapse). For the DAPI, BFP, GFP, or mCherry channels, a polychroic mirror (FF-409/493/596-Di02 by Shemrock) combined with a triple-pass emitter (FF-1-432/523/702-25 by Shemrock) was used. For DAPI, BFP, and GFP imaging, additional emission filters were applied in the optical path (FF01-432-36 by Shemrock, FF01-432-36 by Shemrock and ET525/50M by Chroma, respectively). DAPI, BFP, GFP, and mCherry were excited using a 390/22nm, 390/22nm, 475/28nm and 575/25nm LED, respectively.

For the microfluidic experiments (strain CJW7323), the excitation light intensity was reduced to 20% for the 1-min interval and 30% for the 3-min interval imaging of the RplA-GFP and HupA-mCherry. An exposure time of 120-ms was used for both markers. In all time-lapse experiments, a neutral density filter (absorptive ND filter, OD:1.3 / 5% transmission, NE13B by Thorlabs) was also applied to reduce the LED power and minimize phototoxicity and photobleaching. The LED powers (factored by the neutral density filters) were calibrated using a microscope slide power meter with an 18x18 mm sensor size (S170C by Thorlabs). The light intensity was measured for each excitation wavelength (475/28nm and 575/25nm) at the end of the objective (without immersion oil) for different LED intensities (% of maximum intensity). From the generated calibration curves, the RplA-GFP excitation light power was estimated to be 524 μW and 629 μW for the 1-

min and 3-min fluorescence interval imaging, respectively. HupA-mCherry was excited with 109 μW light
power for the 1-min and 191 μW for the 3-min fluorescence interval imaging.

The type N immersion oil (by Nikon) was used in all experiments except for the time-lapse observation of
cephalexin, A22, A22+cephalexin, and chloramphenicol- and cephalexin-treated cells where the type F
immersion oil (by Nikon) was used. All imaging (time-lapse and snapshots) was done at 30ºC.

Analysis software and code availability
Cropping and alignment of the microfluidic channels were performed with MATLAB (www.mathworks.com)
using a previously published pipeline from our lab (Lin and Jacobs-Wagner, 2022). Supervised classification
and curation of the Oufti cell meshes was also implemented in MATLAB (Campos et al., 2018; Govers et
al., 2024; Gray et al., 2019). For the T7 experiments, segmentation and tracking of the CJW7798 cells was
performed using the Omnipose deep neural network architecture (Cutler et al., 2022), using a previously
trained model (Thappeta et al., 2024) and the SuperSegger MATLAB-based package (Stylianidou et al.,
2016). The remaining analyses were performed using Python 3.9 (www.python.org), that included the
numpy (Harris et al., 2020), *scipy* (Virtanen et al., 2020), *pandas* (McKinney, 2010), *scikit-mage* (Van Der
Walt et al., 2014), *scikit-learn* (Pedregosa et al., 2012)*, shapely* (Gillies, Sean et al., 2023), *statsmodels*
(Seabold and Perktold, 2010) and *pytorch* (Paszke et al., 2019) libraries. The *matplotlib* (Hunter, 2007) and
seaborn (Waskom, 2021) libraries were used for plotting. The analysis pipeline and functions (summarized
in Table S6) are available in the Jacobs-Wagner lab GitHub repository
(http://www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025) and the GitHub
repository of Alexandros Papagiannakis (https://github.com/alexSysBio).

Cell segmentation and tracking
For the CJW6768, CJW6769, CJW7020, and CJW7021 strains used in Figure 2 and Figure 2 – figure
supplement 1, Oufti (Paintdakhi et al., 2016) was used to draw cell meshes on the phase contrast snapshots
and a MATLAB-based support vector machine model was used to remove the badly segmented cells as
previously described (Campos et al., 2018; Govers et al., 2024; Gray et al., 2019).

The Omnipose deep neural network architecture (Cutler et al., 2022) with a previously trained model
(Thappeta et al., 2024) was used to segment the CJW7798 cells during T7 RNA polymerase induction on
an agarose pad. The segmented cells were then tracked using SuperSegger (Stylianidou et al., 2016). A
custom class (*omnipose_to_python_ghv.py*) was developed to transfer the SuperSegger segmentation and
tracking data into Python for post-processing.

Due to the unusual and variable morphology of the A22/cephalexin-treated cells, a custom Python class
was developed for their segmentation and tracking. The *Otsu_phase_segmentation_ghv.py* class

segments the cells by applying an Otsu-threshold (Otsu, 1979) on the inverted phase images, followed by

binary dilation (*scikit-image.morphology.binary_dilation* Python function) hole filling

(*scipy.ndimage.binary_fill_holes* Python function), and binary closing (*scikit-*

image.morphology.binary_closing Python function). The segmentation masks were tracked between

subsequent timepoints, using cell distance as well as cell area constraints, and linked into cell growth

trajectories. Cell morphology criteria such as the maximum pixel distance from the medial axis and the

medial axis sinuosity were used to remove bad segmentations. The remaining segmentation masks were

manually curated.

For the other experiments on agarose pads, a neural network (Wiktor et al., 2021; Zhou et al., 2020) with a

previously designed and trained U-net architecture (Mäkelä et al., 2024) was used to segment single cells

based on phase contrast snapshots. The generated segmentation masks were further processed by

watershed separation, filling the holes within masks and removing unusually small masks. Finally, a

graphical interface was used to manually remove the badly curated cells (less than 5% of the segmented

cell population).

In contrast to the Oufti software (Paintdakhi et al., 2016) that generated sub-pixel meshes around the cell

boundaries, the other applied segmentation algorithms returned pixel-based cell masks. To deal with this

discrepancy, the sub-pixel cell meshes from Oufti were converted into pixel-based cell masks by collecting

the pixels within the circumscribed single cell area in Python (*oufti_snapshots_GrayGovers* class, included

in the *snapshots_analysis_OUFTI_GrayGovers* Python script). After this conversion, the same Phyton-

based functions were applied for the analysis of the cell fluorescence and morphology statistics regardless

of the segmentation method used.

To segment cells growing in the microfluidic device, a custom library of image analysis functions

(*mother_machine_segmentation* class, included in the *microfluidics_segmentation_ghv* Python script) was

developed in Python. This algorithm was applied on the cropped, aligned, background-subtracted, and

inverted phase contrast images that were produced using a previously published pipeline (Lin and Jacobs-

Wagner, 2022) in MATLAB. Cell segmentation was performed in three steps. First, all the cells within the

microfluidics channel, which were brighter than the microfluidic channel background in the inverted and

background-corrected phase-contrast channel, were segmented using an Otsu intensity threshold (Otsu,

1979). This crude thresholding step, which separated the cell (brighter: 1) from the background (darker: 0)

pixels, yielded at least one masked label for the entire row of stacked cells in each microfluidic channel. A

watershed segmentation was then applied to define the boundaries between individual cells and split the

Otsu-based binary mask(s). The watershed algorithm was guided by the number of cells and their relative

positions in the microfluidic channel, which were determined using an adaptive filter combined with the local

decrease of the inverted phase intensity between the poles of adjacent cells. Finally, a graphical interface
was used to curate the cell masks by manually splitting or merging cell labels.
995

After segmentation, the curated single-cell masks were tracked over time and linked into trajectories from
birth to division using the centroid distance and the relative area difference between cells at consecutive
time-points (*mother_machine_tracking* class, included in the *microfluidics_segmentation_ghv* Python
script, or the *fluorescence_analysis* class and its depending functions in the *Time_lapse_on_agarose_pads*
Python package). Examples of cell segmentation and tracking in microfluidics are shown in Movie S1 and
Figure 1 – figure supplement 1A. Regardless of the image acquisition time intervals for the fluorescence
channels, phase-contrast images were acquired every minute, which allowed for accurate cell tracking.
1003

Fluorescence background correction was different between microfluidic and agarose-pad experiments. In
microfluidics, the average background was measured within two 8-pixel (0.528 µm) wide areas, one on the
left side and one the right side of the channel, at least 10 pixels (0.66 µm) away from the channel boundaries
from top to bottom. The estimated background was then subtracted along the channel length. The
background correction was integrated in the *get_fluorescence_image* function, located in the
*microfluidics_analysis_functions_*ghv Python script.
1010

In agarose-pad experiments, an Otsu threshold (Otsu, 1979) was used on the inverted phase-contrast
image to segment all cells. A binary dilation was then performed on these crude cell masks before
estimating the local average fluorescence of the unmasked pixels (pixels outside the dilated cell masks).
The locally-averaged background was then used to fill in the cell areas and reconstruct the background of
the entire field of view in the absence of cells. The smoothed (Gaussian smoothing) reconstructed
background fluorescence was subtracted from each fluorescence image. This background correction
pipeline (*back_sub* function) is integrated in both the *unet_snapshots* class and the
oufti_snapshots_GoversGray class. For the time-lapse imaging of A22-treated CJW7323 cells (Figure 7)
and CJW7798 cells during induction of T7 RNA polymerase expression (Figure 6), the fluorescence
background was not subtracted since we did not need to quantify the raw pixel values. The normalized
polysome and nucleoid fluorescence normalized by the average whole cell fluorescence is shown instead,
marking the relative macromolecular rearrangements.
1023

1024 **mCherry-µNS particle localization and tracking**
1025 For the localization of the diffraction-limited mCherry-µNS particles (Figure 5), a set of custom functions
1026 were developed in Python and implemented in the *particle_positions_snapshots* class, included in the
1027 *snapshots_analysis_UNET_ghv* Python script.
1028

The fluorescent particles were segmented in the background corrected mCherry-µNS fluorescence images
using a Laplace of Gaussian (LoG) and an adaptive filter combined. The particle masks that had an area
larger than a specified threshold (80 pixels for the microfluidics and 90 pixels for the agarose-pad
experiments), which usually included two particles from the same or adjacent cells, were further processed
by applying a relative fluorescence threshold (90[th] percentile of masked pixel intensity) to find the local
fluorescence maxima and separate the two objects. Additional minimum area and aspect ratio constraints
were applied. To accurately estimate the particle position with sub-pixel resolution, a 2D Gaussian function
(eq. 1) with rotation (eq. 2) was fitted (least square method: *scipy.optimize.leastsq*) to an area of 7x7 pixels
centered at the centroid of the particle mask:

1038
$$g(x,y) = A \exp\left(-\frac{1}{2}\left(\left(\frac{x_0 - x_{rot}}{\sigma_x}\right)^2 + \left(\frac{y_0 - y_{rot}}{\sigma_y}\right)^2\right)\right) \quad \text{[eq. 1]}$$

1039
1040 and
1041
$$x_{rot} = x\cos(\theta) - y\sin(\theta)$$
1042
$$y_{rot} = x\sin(\theta) + y\cos(\theta) \qquad \text{[eq. 2]}$$
1043
1044 where *x* and *y* are the coordinates in the 2D imaging plane, *A* is the amplitude of the fitted Gaussian, *σ* is
1045 the standard deviation in each dimension, and *θ* is the rotation angle in radians, which was used to rotate
1046 the Gaussian distribution (x_{rot}, y_{rot}) around the particle center or the Gaussian mean (x_0, y_0).

1047

Two-dimensional cell mapping and alignment
Ribosome and nucleoid fluorescence statistics, as well as particle positions within cells, were mapped in
relative 2D cellular coordinates from pole to pole and across the cell width. Such mapping allowed us to
project the fluorescence or position statistics in 1 or 2D for specific cell length ranges and cell division cycle
intervals. For this analysis, it was necessary to draw the medial axis for each of the segmented cells, which
was obtained differently in microfluidic and agarose-pad images.

1054

In the microfluidic experiments, where all the cells were stacked in a vertically oriented channel, the medial
axis was drawn by fitting a second-degree polynomial to the most distant coordinates from the cell
boundaries, excluding the cell caps at the poles where the medial axis was linearly extrapolated. The length
of the cells, which was parallel to the length of the microfluidic channel, was used as the independent
variable for the fitting. This medial axis estimation is implemented via the *all_medial_axis* function in the
microfluidics_analysis_functions_ghv Python script.

1061

However, in the agarose-pad experiments, cells were randomly oriented in the imaging plane. Thus, it was
impossible to use one of the coordinates (x or y) as the independent variable as this would bias our medial
axis estimation toward the same coordinate for cells that were not diagonally oriented. Instead, we
developed an algorithm that scanned through the middle of the cells with a fixed sub-pixel step and
directionality constraints to identify nodes at the most distant locations from the cell boundaries. These

ordered (from pole to pole) and numbered nodes were used to fit the x and y coordinates of the medial axis
separately, using the number of the node as an independent variable and its position coordinates as
dependent variables. The degree of the fitted polynomials (d) scaled linearly with the total number of nodes
(N) based on the relation $d = 0.1N - 5$. Similar to the microfluidic experiments, the medial axis was linearly
extrapolated at the cell caps. This medial axis estimation is implemented using the *get_medial_axis* function
in both the *unet_snapshots* and *oufti_snapshots_GoversGray* classes located in the
snapshots_analysis_UNET_ghv and *snapshots_analysis_OUFTI_GrayGovers_ghv* Python scripts
respectively. The medial axis estimation function is also provided as a separate function in the
Bivariate_medial_axis_estimation.py Python script.
1076
The medial axis of each cell, with a resolution of 0.1 pixel, was used to map the cell pixels and particle
positions along the cell length and width. The projection of each pixel or particle position on the medial axis,
defined as the most proximal node on the central line, was used to determine its cell length coordinate. The
distance between the pixel or particle position and its medial axis projection was used to determine its
absolute cell width coordinate. The sign of the cross product $\overrightarrow{PL} \times \overrightarrow{PC}$ (plus or minus), where L was the
location of the particle, P was its projection on the medial axis, and C was the center of mass of the cell
mask, was used to determine the position of particles or cell pixels on the sagittal plane.
1084

Cell polarity and ages
In time-lapse experiments, the old and new poles of the cells were determined as follows. The pole of the
daughter cell that was closer to the mid-cell position of its predivisional mother cell, where cell constriction
occurs, was defined as the new pole. The medial axis coordinates were adjusted based on this polarity. As
a result, the relative cell length extended from -1 to 1, with -1 corresponding to the old pole, 1 to the new
pole, and zero to the cell center (see Figure 3 – figure supplement 1B-D). The implementation of this method
is included in the *get_polarity* function in the *microfluidics_analysis_functions_ghv* Python script.
1092

Cell lineages and ages relative to the oldest mother cells (first lineage) at the closed end of the microfluidic
channel were determined using the orientation of the cell poles. The daughter cells that had the same
polarity as the oldest mother cells were assigned the same age and lineage as their mothers. The daughter
cells with the opposite polarity, which inherited the new pole of their mother cells, were considered younger
and belonged to the second lineage. The daughter cells of mother cells belonging to the second lineage
were classified as lineages three and four depending on their polarity. Cells from the third lineage had
opposite polarity and those from the fourth lineage had the same polarity as their second lineage mother
cells. Finally, mother cells from the third lineage, which inherited the new pole of their second-lineage
mother cells, divided to yield the fifth and sixth cell lineages. The fifth lineage had the same polarity as its
third-lineage mother and its sister lineage (sixth) had the opposite polarity and inherited the new pole from
its third-lineage mother cell. The implementation of this method is included in the *get_ages* function in the

*microfluidics_analysis_functions_*ghv Python script. For the analysis of the data presented in Figure 3, only

the third, fourth, fifth and sixth lineages were considered to avoid age-related effects at the cell poles (Chao

et al., 2024; Coquel et al., 2013; Koleva and Hellweger, 2015; Łapińska et al., 2019; Lindner et al., 2008;

Proenca et al., 2019).

1108

1109 **Construction of intensity profiles, 2D cell projections, demographs, and kymographs**

1110 With the ribosome and nucleoid fluorescence pixels as well as the µNS particle positions inside the cells

1111 having been mapped, the fluorescence and particle statistics from pole to pole (medial axis projection) and

1112 across the cell width (distance from the medial axis) were plotted for different cell division cycle intervals or

1113 cell length ranges.

1114

An intensity profile represents the change in a fluorescence statistic along the cell from one pole to the

other. Such a fluorescence statistic can be the average fluorescence intensity (as in Figures 1D, 3G, 5C,

and Figure 3- figure supplement 1B), which corresponds to the concentration of the reported protein per

cell area (sum of pixels divided by the number of pixels). In other analysis (as in Figure 6C and Figure 1 –

figure supplement 6C), the protein concentration along the cell length was divided by the whole cell

average. This scaled statistic is expressed as a percentage change relative to average concentration (mean

%). A value above 100 indicates increase above the average, whereas a value below 100 corresponds to

a decrease below the average concentration. This scaled statistic is insensitive to the RplA and HupA

concentration variability between cells, or to the maturation of the GFP or mCherry fluorophores. As a result,

the scaled concentrations showcase rearrangements of the tagged proteins and do not imply changes in

protein synthesis relative to cell growth.

1126

Intensity profiles were plotted for single cells (Figure 3 – figure supplement 1B and Figure 1 – figure

supplement 6A) or populations (as in Figures 1D, 3G, 5C, 6C, 6E). In the second instance, the intensity

profiles represent the average 1D projection of the fluorescence statistic for a specific cell division cycle

interval, cell length range, growth rate range, polysome and nucleoid asymmetry group, or time range (e.g.,

during antibiotic treatment). To generate a single-cell intensity profile, the medial axis length and its

projected pixels were binned, and the average value of the selected fluorescence statistic was calculated

per bin. If the number of bins was higher than the length of the medial axis in pixels, the missing values

were filled in by linear interpolation. To avoid artefacts from the cell boundaries, the intensity profile was

estimated for a box with a width of six pixels (~0.4 µm) centered on the medial axis (i.e., three pixels on

each side of the medial axis). The intensity profile was occasionally smoothed by a moving average of a

specified cell-length window, without excluding the bins at the edges which were not smoothed.

Alternatively, a univariate spline (*scipy.interpolate.UnivariateSpline*) was fitted to the cell-length-binned data

(as in Figure 5C). The univariate splines were fitted to the binned fluorescence intensity along the cell

length. The number of cell length bins (N_{bin}) linearly scaled with the cell length in µm (l_{cell}) using the first-

order relationship: $N_{bin} = \frac{100}{4.3} l_{cell}$ (only the whole part of the decimal was considered). Similar to the

intensity profile where a fluorescence statistic was plotted along the cell length, the density profiles

(histograms of the particle positions along the cell) for the mCherry-µNS (Figure 5C) particles along the cell

were also plotted. Note that the intensity profiles shown in Figures 1D, 3G, 5C, 6C, 6E, Figure 1 – figure

supplement 1D, and Figure 2 – figure supplement 3D are averages of many segmented cells or cell division

cycles. In these averaged profiles, the variability in the timing of nucleoid segregation across cells

attenuates the appearance of polysome accumulation. Furthermore, the point spread function of fluorescent

ribosome markers causes an underestimation of the magnitude of the polysome accumulations in epi-

fluorescence micrographs (Bakshi et al., 2012b).

1150

The intensity profiles were also used to construct demographs from agarose-pad experiments (Figure 2A,

Figure 2 – figure supplement 1, and Figure 1 – figure supplement 6B) where cells were sorted by cell length

from the shorter newborn to the longer predivisional cells. Intensity profiles were also used to construct

ensemble kymographs from microfluidic experiments where the cells were sorted by their relative position

in the cell division cycle (Figure 1C and Figure 2 – figure supplement 2B) or single-cell kymographs from

agarose pad time-lapse measurements (Figure 7D-F). Specifically, the average intensity profile was

estimated for each cell length or cell division cycle bin, and the average intensity profiles were stacked from

birth to division (left to right) oriented according to the cell polarity (top to bottom). In the ensemble

kymographs, the fluorescence intensities were projected along the relative cell length (cell length %),

whereas in the single-cell kymographs the length of the projection linearly scales with the cell length. A

Gaussian smoothing (*skimage.filters.gaussian*) was applied to smooth the demographs and ensemble

kymographs.

1163

In addition to the intensity or density profiles for a fluorescence statistic or the density of the µNS particles

along the medial axis, 2D projections of the ribosome or nucleoid signal distribution (Figures 1D, 3D, 3G,

5B, 6C, 6E, and Figure 2 – figure supplement 2A) as well as the µNS particle positions (Figure 5B) were

also constructed. These 2D intensity or density maps were constructed by binning the cell pixels or particle

positions not only by cell length, but also by cell width. The average fluorescence statistic or the particle

density was then shown per bin, including data from many single cells within a specified cell division cycle

or cell length range. A Gaussian smoothing (*skimage.filters.gaussian*) was applied to smooth the 2D

projections.

1172

Extraction of population-level statistics of polysomes and nucleoids

To quantify the correlation between polysome accumulation and nucleoid segregation at the population

level (Figure 2 and Figure 2 – figure supplement 1), the average polysome accumulation at mid-cell was

extracted from cell snapshot images. This statistic was obtained by averaging the scaled (divided by the

whole cell average concentration) RplA-GFP and HupA-mCherry intensity profiles of all the cells in the

population. Therefore, this statistic describes the average "behavior" of the population under a specific
growth condition. We reasoned that if, in a specific nutrient condition, nucleoid segregation happens earlier
during the division cycle, then a higher fraction of the population will have segregated nucleoids. If so, the
scaled HupA-mCherry concentration should exhibit a stronger depletion at mid-cell on average. This
depletion was measured as the relative concentration difference between the HupA-mCherry peaks, which
corresponded to the two lobes of the constricting nucleoid, and the trough at mid-cell, which corresponds
to the point of nucleoid splitting. Similarly, the average polysome accumulation was measured as the RplA-
GFP relative concentration difference between the mid-cell accumulation (peak) and the polysome depleted
regions near the quarter cell positions, or at the centers of the segregating sister nucleoids (troughs). The
same statistics were used to quantify the relative polysome accumulation at mid-cell and the relative
nucleoid depletion in the same region between cell division cycles with different growth rates in the same
nutrient condition (Figure 2 – figure supplement 2).

1190

Nucleoid segregation cycle tracking
In this work, the nucleoid segregation cycle was defined as the period from the end of a nucleoid splitting
event (i.e., when a segregating nucleoid is segmented by local thresholding (custom Python function:
LoG_adaptive_image_filter.py) as a separate object from its sister nucleoid) until the end of the splitting of
the same nucleoid object (Figure 1 – figure supplement 2D), usually after cell division but sometimes in the
same cell division cycle (Figure 1 – figure supplement 2A).

1197

To track the nucleoid objects through the nucleoid segregation cycle from the mother cells to their
daughters, we used the cell polarity, i.e., based on pole identity (Figure 1 – figure supplement 4B). The
nucleoids of daughter cells that had opposite polarity to their mother cells were inherited from the new pole
of the mother cells (groups 1 and -1 in Figure 1 – figure supplement 4B-C). The nucleoids in the daughter
cells that had the same polarity as their mother cells were inherited from the old pole of the mother cells
(groups 2 and -2 in Figure 1 – figure supplement 4B-C). During a complete nucleoid segregation cycle, the
nucleoid was tracked from the quarter position of the mother cell to the center of its daughter cells (Figure
1 – figure supplement 4C). This method is implemented in the *track_nucleoids* function, included in the
microfluidics_analysis_functions_ghv Python script.

1207

1208   To quantify the polysome accumulation and nucleoid splitting during the nucleoid segregation cycle (Figure
1209   1E-F), we measured the RplA-GFP and HupA-mCherry concentration within the cell length region of 2.5
1210   pixels adjacent to the center of the tracked nucleoid object (5 pixels or 0.33 µm in total).

1211

1212

1213

Quantification and segmentation of polysomes and nucleoids segmentation in cells treated with cephalexin or cephalexin + A22

The polysome accumulations and nucleoids objects were also segmented in A22 and cephalexin-treated cells using an adaptive filter (custom Python function: *LoG_adaptive_image_filter.py*) on the masked cell fluorescence. The number of the detected polysome accumulations or nucleoid objects corresponds to the number of the segmented labels in the RplA-GFP or HupA-mCherry channel respectively. Similarly, the average polysome accumulation or nucleoid area corresponds to the average area of all segmented labels per channel. For a fair comparison between cells treated with A22 + cephalexin and cells treated with cephalexin alone, cells were randomly sampled without substitution from 12 cell area bins between 10 and 13 μm^2 (bin width = 0.25 μm^2) such that the compared populations in Figure 7F had the same cell area distributions (same number of cells per cell area bin).

Quantification of the distance between fusing or non-fusing nucleoids in cephalexin- and chloramphenicol-treated cells

The distance between nucleoids corresponds to the cell length difference between the peaks of the longitudinal HupA-mCherry signal projection (Figure 4 – figure supplement 1A-B). To calculate these distances, the HupA-mCherry longitudinal profile was first smoothed using a fourth-roder univariate regression (*scipy.interpolate.UnivariateSpline*). The optimal smoothing factor (10^3) was determined using the L-curve method (Nasehi Tehrani et al., 2012) where the x-axis corresponds to the value of the smoothing factor and the y-axis to the variance in detected nucleoid number during cell growth. The average variance was considered across the tracked cell trajectories. The smoothing factor at the elbow of the L-curve is the optimal point between overfitting, where the noise in the HupA-mCherry signal is wrongly assigned to nucleoid objects, and underfitting, where two nucleoids in proximity may be wrongly identified as one. After smoothing, the peaks of the HupA-mCherry projection were detected using the *scipy.signal.find_peaks* function, applying a lower threshold of 125 arbitrary units. The number of the detected peaks is equal to the nucleoid number per cell, whereas the distance between the peaks is equal to the distance between the nucleoid centroids. A nucleoid fusion event is indicated by the reduction in nucleoid number between consecutive frames in a cell trajectory.

The tracked cell trajectories were classified into those that fuse their nucleoids and those that do not. The first class includes cell trajectories that contain a nucleoid fusion event and end with less detected nucleoids objects compared to their beginning. The second class includes cell trajectories that do not contain any nucleoid fusion event and end with an equal number or more detected nucleoids compared to their beginning. All classified trajectories start at time-point 0 min, which corresponds to the first time-point of the experiment, or the first microscopy image taken after spotting the cells on the chloramphenicol-containing agarose pad. Only trajectories with more than one nucleoid at timepoint 0 min are considered. This first time-point also corresponds to when the minimum distance between adjacent nucleoids is quantified for

1251   each cell trajectory (plotted in Figure 4 – figure supplement 1C). The nucleoid segregation variability shown

1252   in Figure 4 – figure supplement 1B, is consistent with previous observations (Spahn et al., 2014).

1253

Linear mixed-effects models to determine the relative contribution of polysome accumulation and

cell elongation to the migration of the sister nucleoids

To estimate the relative contribution of the polysome accumulation at mid-cell and that of cell elongation to

the migration of the separated sister nucleoids (first seen in Figure 1H), linear mixed-effects regressions

were fitted to the scaled single cell data shown in Figure 1 – figure supplement 5B (*statsmodel* package in

Python, *statsmodels.formula.api.mixedlm* function). One model was fitted for each relative time interval

between the end of nucleoid splitting and cell division, describing the rate of distance increase between the

separated sister nucleoids (σ, response variable) as a function of the polysome concentration increase rate

(ρ), the rate of cell elongation (λ), their interaction ($\rho\lambda$), and noise (ε) (*statsmodels.formula.api.ols*) as

follows:

1264
1265
$$\sigma = \beta_0 + \beta_1\rho + \beta_2\lambda + \beta_3(\rho\lambda) + \varepsilon \qquad \text{[eq. 3]}$$
1266

The single cell data were scaled by subtracting the population mean and then dividing by the standard

deviation prior to model fitting for each of the four relative time intervals (in Figure 1 – figure supplement

5B). The coefficients of the linear regression model (shown in Figure 1I) provide a measure of the relative

effect of the independent variables on the dependent variable. The cell elongation (λ) had a statistically

significant effect to the response variable (σ) Prob(>|Z|) < 10^{-9}) across all relative time intervals. The

polysome accumulation coefficients (ρ) had a statistically significant effect to the response variable (σ)

(Prob(>|Z|) < 10^{-14}) for the first three relative time intervals (0-25%, 25-50% and 50-75%) and a marginally

significant one Prob(>|Z|) = 0.02) for the last quartile (75-100%). The interaction term ($\rho\lambda$) had a marginally

significant effect (Prob(>|Z|) = 0.002) for the first quartile (0-25%) yet with a very small coefficient of 0.06

and did not present any significance (Prob(>|Z|) > 0.05) for the remaining time intervals (25-50%, 50-75%

and 75-100%). Hence, it is not shown in Figure 1I. The most significant effect on the migration of the sister

nucleoids (σ) was presented by the rate of polysome accumulation at mid-cell (ρ), 0-25% from the end of

nucleoid splitting to cell division, with a coefficient of 0.46 and Prob(>|Z|) < 10^{-84}.

1280
1281
Calculation of rates

The rates of the polysome accumulation ($\frac{\Delta RplA_{mid-nuc\ conc.}}{\Delta T}$) or nucleoid depletion ($\frac{\Delta HupA_{mid-nuc\ conc.}}{\Delta T}$) in the

middle of the nucleoid (2.5 pixels adjacent the nucleoid center) (Figure 1F), correspond to the slope of a

linear regression (Python fitting function: *numpy.polyfit*) fitted to the change of the mid-nucleoid RplA-GFP

or HupA-mCherry concentration over time, 40-90% into the nucleoid segregation cycle (as in Figure 1E).

This nucleoid cycle interval corresponds to the average time from the initiation of nucleoid splitting to just

before its completion. Choosing the appropriate range to correlate the two statistics was important since

polysomes appeared to accumulate in the middle of the nucleoid before the onset of nucleoid splitting
(Figures 1E and Figure 1 – figure supplement 6D).
1291
The rate of RplA-GFP concentration increase at mid-cell ($\frac{\Delta RplA_{mid-cell\ conc.}}{\Delta T}$) was calculated (Figure 1H) within
a region covering 10% of the total cell length at the cell center. The same results were obtained when the
rates were calculated within a cell region covering 2.5 pixels adjacent to the cell center. The rate of nucleoid
migration ($\frac{\Delta Distance_{nuc}}{\Delta T}$) corresponds to the slope of a linear regression that describes the natural log-
transformed distance increase between the separated sister nucleoids over time (Figure 1H). Similarly, the
rate of cell elongation ($\frac{\Delta Length_{cell}}{\Delta T}$) was calculated considering the natural log-transformed cell length
increase over time (Figure 1H). The rate of RplA-GFP concentration increase at mid-cell ($\frac{\Delta RplA_{mid-cell\ conc.}}{\Delta T}$),
the rate of nucleoid migration ($\frac{\Delta Distance_{nuc}}{\Delta T}$), and the rate of cell elongation ($\frac{\Delta Length_{cell}}{\Delta T}$) were calculated within
four relative time intervals from the end of nucleoid splitting to cell division (Figure 1G and Figure 1 – figure
supplement 5B). Only single nucleoid migration intervals with more than four timepoints were considered
to ensure a reliable fitting. The small fraction (~8%) of cells that were born with two separately detected
nucleoid objects, which indicates that the end of nucleoid splitting occurred in the previous cell division
cycle, were removed from the analysis. The cell division cycles with a negative cell elongation rate (<1%)
were also excluded from the analysis.
1306
Reaction-diffusion model
Our theoretical model is an extension of previous work (Miangolarra et al., 2021), which showed that steric
effects between the DNA and the transcription-translation machinery (including polysomes) are sufficient
to drive nucleoid segregation. To simply describe these interactions, we employed the Flory-Huggins theory
for regular solutions (Flory, 1942; Huggins, 1941) and modeled the non-dimensionalized free-energy
density f as a function of the local volume fractions of the nucleoid (n) and polysomes (p):
1313
1314 $$f(n,p) = \frac{p}{v_p}\ln p + \frac{n}{v_n}\ln n + (1-p-n)\ln(1-p-n) + \chi_p p(1-p-n) + \chi_n n(1-p-n) + \chi_{np}np -$$
1315 $$\kappa_p \nabla p \nabla(1-p-n) - \kappa_n \nabla n \nabla(1-p-n) - \kappa_{np}\nabla n \nabla p. \qquad \text{[eq. 4]}$$
1316
1317   Here, $(1-n-p)$ is the volume fraction of the cytoplasm not occupied by nucleoid or polysomes, whereas
1318 v_n and v_p are proportional to the molecular volumes of the elementary translational degrees of freedom
1319   (i.e., of the nucleoid element and single polysome, respectively). χ_p, χ_n and χ_{np} are the Flory-Huggins
1320   interaction parameters, and κ_p, κ_n and κ_{np} are the interfacial tensions. We take $\kappa_{ij} = \lambda^2 \chi_{ij}$, with λ being a
1321   characteristic interface width.
1322

1323   The time evolution of the volume fractions is described by a combination of Cahn-Hilliard theory (Cahn and

1324   Hilliard, 1958) and the reactions kinetics of the polysomes. We assume that polysomes are produced inside

1325   the nucleoid at a constant rate k_1 and degraded uniformly along the cell at a rate k_{-1}:

1326

1327
$$\frac{\partial p}{\partial t} = \nabla(M_p \nabla \mu_p) + k_1 n - k_{-1} p, \quad \text{[eq. 5]}$$

1328
$$\frac{\partial n}{\partial t} = \nabla(M_n \nabla \mu_n), \quad \text{[eq. 6]}$$

1329

1330   where $\mu_p = \frac{\delta F}{\delta p}$ and $\mu_n = \frac{\delta F}{\delta n}$ are the local chemical potentials of polysomes and nucleoid elements

1331   respectively, with $F = \int f[n(x), p(x)] dx$ representing the total free energy. $M_p = v_p D_p p$ and $M_n = v_n D_n n$

1332   are the mobility coefficients of the polysomes and the nucleoid, respectively, which depend on the diffusion

1333   coefficients of the polysomes (D_p) and the nucleoid (D_n). This recovers the Fick law of diffusion in the

1334   noninteracting limit.

1335

Since *E. coli* grows by elongation while maintaining its cylindrical shape and cell width, we reduce the

problem to one dimension, with x representing the position along the long axis of the cell. We assume that

the cell length grows exponentially such that the length at time t is $L(t) = L(0)e^{\gamma t}$, with γ being the growth

rate and $L(0)$ the initial cell length at birth. Exponential growth dilutes the existing polysomes with rate γ.

On the other hand, we assume that the nucleoid is not diluted since DNA replication occurs continuously,

and the nucleoid length is known to grow in proportion to cell length (Gray et al., 2019). Letting $\tilde{x} = x/L(t)$

be the relative position along the cell's long axis, the time evolution is given by:

1343

1344
$$\frac{\partial p}{\partial t} = \partial_{\tilde{x}}(\tilde{M}_p \partial_{\tilde{x}} \mu_p) + k_1 n - (k_{-1} + \gamma)p, \quad \text{[eq. 7]}$$

1345
$$\frac{\partial n}{\partial t} = \partial_{\tilde{x}}(\tilde{M}_n \partial_{\tilde{x}} \mu_n), \quad \text{[eq. 8]}$$

1346

1347   where $\partial_{\tilde{x}} = \frac{\partial}{\partial \tilde{x}} = L(t)\frac{\partial}{\partial x}$, and $\tilde{M}_{n,p} = M_{n,p} L(t)^{-2}$. The chemical potentials are:

1348

1349
$$\mu_p = v_p^{-1} \ln p - \ln(1 - p - n) + \chi_p(1 - 2p - n) + (\chi_{np} - \chi_n)n$$

1350
$$-\chi_p \tilde{\lambda}^2 \partial_{\tilde{x}}^2 (2p + n) + (\chi_{np} - \chi_n)\tilde{\lambda}^2 \partial_{\tilde{x}}^2 n, \quad \text{[eq. 9]}$$

1351

1352
$$\mu_n = v_n^{-1} \ln n - \ln(1 - p - n) + \chi_n(1 - 2n - p) + (\chi_{np} - \chi_p)p$$

1353
$$-\chi_n \tilde{\lambda}^2 \partial_{\tilde{x}}^2 (2n + p) + (\chi_{np} - \chi_p)\tilde{\lambda}^2 \partial_{\tilde{x}}^2 p, \quad \text{[eq. 10]}$$

1354

1355   where $\tilde{\lambda} = \lambda/L(t)$.

1356

We take the non-dimensionalized Flory-Huggins interaction parameters to be $\chi_p = 0.2$, $\chi_n = 0.4$ and $\chi_{np} = 1.2$. The χ_{np} parameter captures the steric repulsion between the nucleoid and the polysomes (Miangolarra et al., 2021). The diffusion coefficient of the polysome was set to $D_p = 0.023 \mu m^2/s$, which lies within the same order of magnitude as was previously measured by single-molecule tracking (Bakshi et al., 2012b). For simplicity, we adopt an average description of all polysomes with an average diffusion coefficient. Considering multiple polysome species does not change the physical picture. To illustrate, we considered an extension of our model, which contains three polysome species, each with a different diffusion coefficient (DP = 0.018, 0.023, or 0.028 µm2/s), reflecting that polysomes with more ribosomes will have a lower diffusion coefficient. Simulation of this model reveals that the different polysome species have essentially the same concentration distribution (Figure 4 — figure supplement 4), suggesting that the average description in our minimal model is sufficient for our purposes.

Different diffusion coefficients were tested for the nucleoid ($D_n = 0.0005, 0.001$ or 0.005 µm^2/s) to capture different levels of nucleoid stiffness and hence response time to the local changes in polysome concentration. The polysome degradation rate was set to $k_{-1} = 0.003 s^{-1}$, which corresponds to a half-life of around 5 min (Bernstein et al., 2004). We take the polysome production rate to be $k_1 = k_{1,0} \left(1 + \frac{\gamma}{k_{-1}}\right)$, with $k_{1,0} = 0.002 s^{-1}$, to match the observation that the nucleoid length is proportional to cell length (Gray et al., 2019). The initial cell length $L(0)$ matched the cell length at birth, which has been previously shown to increase exponentially with the growth rate (Govers et al., 2024). Hence, each of the simulated growth rates ($\gamma = 0.25, 0.36, 0.46, 0.57, 0.67, 0.78, 0.88, 0.99, 1.09, 1.2$ h^{-1}) was matched to a cell length at birth (Figure 4 – figure supplement 2) by fitting a linear regression ($L(0) = l_0 e^{\gamma/\gamma_0}$, with fitted parameters $l_0 = 2.00$ µm and $\gamma_0 = 2.81$ h^{-1}) to the experimental data (Govers et al., 2024). Other parameters are chosen as follows: $\lambda = 0.03$ µm, $v_n = 10$, $v_p = 5$.

The dynamic equations (eqs. 7-8) were solved numerically in a fixed 1D domain $\tilde{x} \in (0,1)$, with no-flux boundary conditions. Space is discretized into $N = 128$ grid points, and time is discretized to steps of Δt. Time stepping was implemented with an implicit-explicit scheme as previously demonstrated (Mao et al., 2020, 2019). The simulations were initialized from the (symmetric) steady-state solution of the system at the initial cell length $L(0)$ (Figure 4B, dashed curves in Figure 4E, and Movie S3), or using the (asymmetric) macromolecular distribution of the predivisional cell of the former simulation as the initial condition (Figure 4D, solid curves in Figure 4E), which allowed us to capture cell polarity (new versus old pole). The cell length $L(t)$ was updated and recorded at each time step during the simulation. The nucleoid splitting time (shown in Figure 4D) was determined by thresholding the relative decrease of the nucleoid concentration at the cell center. A depletion threshold of 30% was used to mark the event of nucleoid splitting.

The *p* field in our model describes the distribution of all polysomes, which could include one, two, or multiple
ribosomes (in this paper the term "polysomes" refers to both monosomes and true polysomes). For
simplicity, we adopt an average description of all polysomes with an average diffusion coefficient and
interaction parameters, which is sufficient for capturing the fundamental mechanism underlying nucleoid
segregation. The model can be extended to include multiple polysome species: $p = \sum_i p_i$, with each species'
evolution following Eq. 5 with a different diffusion coefficient. Figure 4 – figure supplement 4 shows an
example containing 3 polysome species, with diffusion coefficients $D_p = 0.018, 0.023, 0.028 \ \mu m^2/s$.

1399

1400 Our model can also be extended to consider ectopic polysome production, which introduces an additional
1401 polysome production term:

1402
$$\frac{\partial p}{\partial t} = \nabla\big(M_p \nabla \mu_p\big) + k_1' n - k_{-1} p + k_{\text{ect}}(x) \ , \ \ [\text{eq. 5*}]$$

where $k_{\text{ect}}(x)$ describes ectopic polysome production from plasmids, with x being the distance from one
pole measured in μm. k_1' represents a reduced polysome production rate in the nucleoid. Figure 7 – figure
supplement 2 shows two examples with details as follows. For ectopic production near the poles, we use
$k_{\text{ect}}(x) = 0.8 \ k_1 \ \Theta(x) \ \Theta(0.8 - x)$ and $k_1' = 0.5 k_1$, where $\Theta(x)$ is the Heaviside step function; this restricts
ectopic production to within $0.8 \ \mu m$ from one pole. For ectopic production between sister nucleoids, we use
$k_{\text{ect}}(x) = 0.5 \ k_1 \ \Theta\left(0.4 - \left|\frac{L}{2} - x\right|\right)$ and $k_1' = 0.5 k_1$, where L is the cell length; this restricts ectopic production
to within $0.4 \ \mu m$ from mid-cell. These ectopic production rates are for illustrative purposes only and the
qualitative result does not depend sensitively on parameter values.

1411

1412 The code for the model (including scripts to reproduce all simulation figures) is available at
1413 https://github.com/qiweiyuu/polysome.

1414

Polysome and nucleoid analysis based on fitted Gaussian functions
Quantification of polysome and nucleoid asymmetries within cells along the cell division cycle was achieved
by fitting Gaussian functions (Python least squares optimization: *scipy.optimize.curve_fit*) to the RplA-GFP
or the HupA-mCherry intensity profiles of cells growing in M9gluCAAT in the microfluidic device. Three
Gaussian functions were fitted to the polysome signal concentration early in the cell division cycle. Two of
them captured the nucleoid-excluded polysomes accumulating at the poles and the third captured the
polysome accumulation in the middle of nucleoid (Figure 3 – figure supplement 1A-B):

1422

1423
$$Poly(l) = A_{old} \exp\left(-\frac{1}{2}\left(\frac{l_{old}-l}{\sigma_{old}}\right)^2\right) + A_{mid} \exp\left(-\frac{1}{2}\left(\frac{l_{mid}-l}{\sigma_{mid}}\right)^2\right) + A_{new} \exp\left(-\frac{1}{2}\left(\frac{l_{new}-l}{\sigma_{new}}\right)^2\right) \ \ [\text{eq. 11}],$$

1424
1425 where *A_old, A_new* and *A_mid* are the amplitudes, *l_old, l_new* and *l_mid* are the means, and *σ_old, σ_new* and *σ_mid* are the
1426 standard deviations of the respective fitted functions along the relative cell length (*l*).

1427

1428 Two Gaussian functions were fitted to the concentration of constricting nucleoids (Figure 3 – figure
1429 supplement 1A-B):

$$Nuc(l) = A_{old} \exp\left(-\frac{1}{2}\left(\frac{l_{old}-l}{\sigma_{old}}\right)^2\right) + A_{new} \exp\left(-\frac{1}{2}\left(\frac{l_{new}-l}{\sigma_{new}}\right)^2\right) \quad \text{[eq. 12]}$$

All Gaussian functions were fitted over the cell background fluorescence, which corresponds to the baseline

fluorescence within the masked cell boundaries and along the relative cell length coordinates from the old

pole (-1) via the cell center (0) to the new pole (1). The Gaussian functions were fitted either to single

newborn cells (0-2.5% into the cell division cycle) as shown in Figure 3 – figure supplement 1A-B and as

applied in Figures 3A and 3E, or to the less noisy average intensity profile of the segmented cells within the

0-10% range of their cell division cycle, as shown in Figure 3 – figure supplement 1C-D and used in Figures

3B and 3F. The areas of the Gaussian functions fitted to the polysomes (used in Figure 3A-B) were

estimated using the function:

$$Area = A\,\sigma\,\sqrt{2\pi} \quad \text{[eq. 13]},$$

1444 where A is the amplitude and σ *is* the standard deviation.

1446 The Gaussian areas were used to quantify the polysome asymmetries between the poles:

$$Poly_{asym} = log_{10}\left(\frac{Area_{new}}{Area_{old}}\right) \quad \text{[eq. 14]},$$

1451 where $Area_{new}$ and $Area_{old}$ are the areas of the fitted Gaussians to the nucleoid-excluded polysomes

1452 accumulating in the new and the old pole, respectively. The statistic in eq. 14 corresponds to the x-axis in

1453 Figure 3B. It is a positive value when the new pole contains more polysomes than the old one, and a

1454 negative value when the old pole contains more polysomes than the new one.

1456 The nucleoid Gaussian statistics were used to quantify the nucleoid position asymmetry defined as:

$$Nuc_{pos} = \frac{l_{new}+l_{old}}{2}\frac{l_{cell}}{2} \quad \text{[eq. 15]},$$

1461 and the nucleoid compaction asymmetry was calculated as:

$$Nuc_{comp} = \frac{A_{new}}{A_{old}} \quad \text{[eq. 16]},$$

where l_{old} and l_{new} are the means of the nucleoid Gaussians in relative cell coordinates (as in eq. 12) and

l_{cell} is the cell length in μm. The nucleoid position was first estimated in relative cell coordinates from the old

pole (-1) via the cell center (0) to the new pole (1) and then multiplied by half the cell length to convert to

absolute cell length units. A_{old} and A_{new} are the amplitudes of the fitted nucleoid Gaussians (as in eq. 12) in

arbitrary fluorescence units. The nucleoid position corresponds to the mid-point between the two Gaussian

means (eq. 15) and is a positive number when the nucleoid is shifted toward the new pole and a negative
number when it is positioned closer to the old pole (y-axis in Figure 3B).

1472

1473   The nucleoid compaction asymmetry was calculated as the relative nucleoid Gaussian amplitude (eq. 16).
1474   It had a positive value greater than 1 when the nucleoid concentration was higher toward the new pole and
1475   had a positive value smaller than 1 if the nucleoid was more concentrated toward the old pole (y-axis in
1476   Figure 3F).

1477

1478   To determine which polysome Gaussian statistics and related asymmetries may contribute to nucleoid
1479   compaction asymmetry (eq. 16), a linear mixed-effects model was used (*statsmodel* package in Python,
1480   *statsmodels.formula.api.mixedlm* function). The polysome Gaussian statistics were used as independent
1481   variables and the nucleoid compaction asymmetry as a dependent variable:

1482

1483

1484 $$Nuc_{comp} = \beta_0 + \beta_1 l_{old} + \beta_2 l_{mid} + \beta_3 l_{new} + \beta_4 \sigma_{old} + \beta_5 \sigma_{mid}$$
1485 $$+\beta_6 \sigma_{new} + \beta_7 A_{old} + \beta_8 A_{mid} + \beta_9 A_{new} + \varepsilon \qquad \text{[eq. 17]},$$

1486

where β_0 is the constant (intercept), β_n *(n=1,2…,9)* are the coefficients for each of the polysome Gaussian
statistics (same as in eq. 4 and plotted in Figure 3 – figure supplement 2A-B), and ε is the error. The entire
dataset was treated as a single group. Before fitting the linear mixed-effects model, all data were normalized
by subtracting the average and dividing the resulting value by their standard deviation (Figure 3 – figure
supplement 2B). The relative cell length positions of the polysomes at the old pole (l_{old}) and in the middle
of the nucleoid (l_{mid}) were found to significantly influence the asymmetric nucleoid compaction (Figure 3 –
figure supplement 2C). We reasoned that the available spaces between the polysome positions (Figure 3
– figure supplement 2A) may influence the compaction of the nucleoid and devised a new statistic that
compares the available polysome-free space toward the new pole to that of toward the old pole:

1496
1497 $$Poly_{space} = \frac{l_{new} - l_{mid}}{l_{mid} - l_{old}} \qquad \text{[eq. 18]},$$

1498
1499   where l_{new}, l_{mid} and l_{old} are the polysome Gaussian means (as in eq. 11), which represent the positions of
1500   the polysomes at the new pole, mid-cell region, and old pole, respectively, expressed as relative cell
1501   coordinates from the old pole (-1) via the cell center (0) to the new pole (1). The fraction above (eq. 18) has
1502   a value greater than 1 when the polysome-free space is greater toward the new pole and a value smaller
1503   than 1 when the polysome-free space is greater toward the old pole (x-axis in Figure 3F).

1504

Linear regressions
Two types of linear regressions were used in this work. In Figure 2B, a first-degree polynomial was fitted
using the ordinary least squares method and the numpy library in Python (*numpy.polyfit*), always using the

x-axis as the independent variable and the *y*-axis as the dependent variable. In Figures 1F and 3B, a

principal component regression was fitted using the scikit-learn library in Python

(*sklearn.decomposition.PCA*). A principal component analysis was first applied on the two-dimensional z-

transformed data to find the linear regressor that explained most of the variance (the first principal

component). This linear regressor was then rescaled to the original 2D plane. The principal component

regression eliminates the dependent variable bias associated with a traditional univariate fit. This bias is

introduced when calculating the prediction error (sum of squares) only along the dependent variable axis

during the least squares optimization. The elimination of this univariate bias is particularly important for the

fitted regressions in Figures 1F and 3B since their parameters were used to estimate the rate of nucleoid

splitting in the absence of polysome accumulation in the middle (y-intercept in Figure 1F), the position of

the nucleoid with equal polysomes at the poles (y-intercept in Figure 3B), or the polar asymmetry required

for a centered nucleoid (x-intercept in Figure 3B).

1520

1521 **Kernel density estimations**

1522 One-dimensional (Figures 3A, E, and Figure 1- figure supplement 4C) or two-dimensional (Figures 1F, 1H,

1523 Figure 1 – figure supplement 5A-B, and Figure 3 – figure supplement 2B) Gaussian kernel density

1524 estimations were fitted using the *scipy.stats.gaussian_kde* method. The bandwidth was determined using

1525 Scott's rule (Scott, 1992).

1526

1527 **Correlation coefficient calculations**

1528 All Spearman correlation coefficients (referred to as Spearman ρ) were estimated using the *scipy.stats*

1529 Python library and the *spearmanr* function. Only those Spearman correlation coefficients with a p-value

1530 below 0.05 are shown. Otherwise, the correlation is annotated as not-significant (NS) (as in Figure 1 –

1531 figure supplement 5B).

1532

1533 **ADDITIONAL RESOURCES**

1534 This study did not generate any additional resources.

1535

ACKNOWLEDGMENTS

We are grateful to Dr. A Janakiraman (City College of New York) for the pER12 (pBAD322A-gfp-µNS)

plasmid, Dr K. Prather (Massachusetts Institute of Technology) for the MG1655 (DE3) strain, and the

members of the Jacobs-Wagner laboratory for fruitful discussions and critical reading of the manuscript.

This work was supported in part by the National Institutes of Health (R01 GM082938 to N.S.W.) and by the

Netherlands Organization for Scientific Research (research program Rubicon Science 2018-1, project

number 019.181EN.018 to A.P). C.J.W. is an investigator of the Howard Hughes Medical Institute.

1543

1544

AUTHOR CONTRIBUTIONS

Conceptualization, A.P. and C.J.-W.; methodology, A.P.; software, A.P. and Q.Y.; formal analysis, A.P.; investigation, A.P., Q.Y, and S.K.G.; resources, A.P., Q.Y., and W.-H.L.; data curation, A.P.; writing – original draft, A.P. and C.J.-W.; writing – review and editing, A.P., C.J.-W , N. W., Q.Y., S.K.G., and W.-H.L; visualization, A.P. and C.J.-W.; supervision, N.W. and C.J.-W.; project administration, C.J.-W.; funding acquisition, A.P., N.W. and C.J.-W.

DECLARATION OF INTERESTS

The authors declare no competing interests.

DATA AVAILABILITY

The microscopy images and dataframes used on this study are available on BioStudies and the BioImage Archive (accession # S-BIAD-1658).

REFERENCES

Bakshi, S., Choi, H., Mondal, J., Weisshaar, J.C., 2014. Time-dependent effects of transcription- and translation-halting drugs on the spatial distributions of the *E scherichia coli* chromosome and ribosomes: Time-dependent drug effects on *E. coli* chromosome spatial distribution. Mol. Microbiol. 94, 871–887. https://doi.org/10.1111/mmi.12805

Bakshi, S., Choi, H., Weisshaar, J.C., 2015. The spatial biology of transcription and translation in rapidly growing Escherichia coli. Front. Microbiol. 6. https://doi.org/10.3389/fmicb.2015.00636

Bakshi, S., Siryaporn, A., Goulian, M., Weisshaar, J.C., 2012a. Superresolution imaging of ribosomes and RNA polymerase in live Escherichia coli cells. Mol. Microbiol. 85, 21–38. https://doi.org/10.1111/j.1365-2958.2012.08081.x

Bakshi, S., Siryaporn, A., Goulian, M., Weisshaar, J.C., 2012b. Superresolution imaging of ribosomes and RNA polymerase in live Escherichia coli cells: Superresolution imaging of ribosome and RNAP. Mol. Microbiol. 85, 21–38. https://doi.org/10.1111/j.1365-2958.2012.08081.x

Balakrishnan, R., Mori, M., Segota, I., Zhang, Z., Aebersold, R., Ludwig, C., Hwa, T., 2022. Principles of gene regulation quantitatively connect DNA to RNA and proteins in bacteria. Science 378, eabk2066. https://doi.org/10.1126/science.abk2066

Bartosik, A.A., Mierzejewska, J., Thomas, C.M., Jagura-Burdzy, G., 2009. ParB deficiency in Pseudomonas aeruginosa destabilizes the partner protein ParA and affects a variety of physiological parameters. Microbiology 155, 1080–1092. https://doi.org/10.1099/mic.0.024661-0

Bates, D., Kleckner, N., 2005. Chromosome and Replisome Dynamics in E. coli: Loss of Sister Cohesion Triggers Global Chromosome Movement and Mediates Chromosome Segregation. Cell 121, 899–911. https://doi.org/10.1016/j.cell.2005.04.013

Bean, G.J., Flickinger, S.T., Westler, W.M., McCully, M.E., Sept, D., Weibel, D.B., Amann, K.J., 2009. A22 Disrupts the Bacterial Actin Cytoskeleton by Directly Binding and Inducing a Low-Affinity State in MreB. Biochemistry 48, 4852–4857. https://doi.org/10.1021/bi900014d

Bernstein, J.A., Lin, P.-H., Cohen, S.N., Lin-Chao, S., 2004. Global analysis of *Escherichia coli*
RNA degradosome function using DNA microarrays. Proc. Natl. Acad. Sci. 101, 2758–
2763. https://doi.org/10.1073/pnas.0308747101
Binenbaum, Z., Parola, A.H., Zaritsky, A., Fishov, I., 1999. Transcription- and translation-
dependent changes in membrane dynamics in bacteria: testing the transertion model for
domain formation. Mol. Microbiol. 32, 1173–1182. https://doi.org/10.1046/j.1365-
2958.1999.01426.x
Blundell, M.R., Wild, D.G., 1971. Altered ribosomes after inhibition of *Escherichia coli* by
rifampicin. Biochem. J. 121, 391–398. https://doi.org/10.1042/bj1210391
Brandt, F., Etchells, S.A., Ortiz, J.O., Elcock, A.H., Hartl, F.U., Baumeister, W., 2009. The
Native 3D Organization of Bacterial Polysomes. Cell 136, 261–271.
https://doi.org/10.1016/j.cell.2008.11.016
Broering, T.J., Arnold, M.M., Miller, C.L., Hurt, J.A., Joyce, P.L., Nibert, M.L., 2005. Carboxyl-
Proximal Regions of Reovirus Nonstructural Protein μNS Necessary and Sufficient for
Forming Factory-Like Inclusions. J. Virol. 79, 6194–6206.
https://doi.org/10.1128/JVI.79.10.6194-6206.2005
Broering, T.J., Parker, J.S.L., Joyce, P.L., Kim, J., Nibert, M.L., 2002. Mammalian Reovirus
Nonstructural Protein μNS Forms Large Inclusions and Colocalizes with Reovirus
Microtubule-Associated Protein μ2 in Transfected Cells. J. Virol. 76, 8285–8297.
https://doi.org/10.1128/JVI.76.16.8285-8297.2002
Bryksin, A.V., Matsumura, I., 2010. Overlap extension PCR cloning: a simple and reliable way
to create recombinant plasmids. BioTechniques 48, 463–465.
https://doi.org/10.2144/000113418
Cabrera, J.E., Cagliero, C., Quan, S., Squires, C.L., Jin, D.J., 2009. Active Transcription of
rRNA Operons Condenses the Nucleoid in *Escherichia coli* : Examining the Effect of
Transcription on Nucleoid Structure in the Absence of Transertion. J. Bacteriol. 191,
4180–4185. https://doi.org/10.1128/JB.01707-08
Cahn, J.W., Hilliard, J.E., 1958. Free Energy of a Nonuniform System. I. Interfacial Free Energy.
J. Chem. Phys. 28, 258–267. https://doi.org/10.1063/1.1744102
Campbell, E.A., Korzheva, N., Mustaev, A., Murakami, K., Nair, S., Goldfarb, A., Darst, S.A.,
2001. Structural Mechanism for Rifampicin Inhibition of Bacterial RNA Polymerase.
Cell 104, 901–912. https://doi.org/10.1016/S0092-8674(01)00286-0
Campos, M., Govers, S.K., Irnov, I., Dobihal, G.S., Cornet, F., Jacobs-Wagner, C., 2018.
Genomewide phenotypic analysis of growth, cell morphogenesis, and cell cycle events in
Escherichia coli. Mol. Syst. Biol. 14, e7573. https://doi.org/10.15252/msb.20177573
Campos, M., Surovtsev, I.V., Kato, S., Paintdakhi, A., Beltran, B., Ebmeier, S.E., Jacobs-
Wagner, C., 2014. A constant size extension drives bacterial cell size homeostasis. Cell
159, 1433–1446. https://doi.org/10.1016/j.cell.2014.11.022
Cass, J.A., Kuwada, N.J., Traxler, B., Wiggins, P.A., 2016. Escherichia coli Chromosomal Loci
Segregate from Midcell with Universal Dynamics. Biophys. J. 110, 2597–2609.
https://doi.org/10.1016/j.bpj.2016.04.046
Castellana, M., Hsin-Jung Li, S., Wingreen, N.S., 2016. Spatial organization of bacterial
transcription and translation. Proc. Natl. Acad. Sci. 113, 9286–9291.
https://doi.org/10.1073/pnas.1604995113

Chai, Q., Singh, B., Peisker, K., Metzendorf, N., Ge, X., Dasgupta, S., Sanyal, S., 2014. Organization of Ribosomes and Nucleoids in Escherichia coli Cells during Growth and in Quiescence. J. Biol. Chem. 289, 11342–11352. https://doi.org/10.1074/jbc.M114.557348

Chao, L., Chan, C.K., Shi, C., Rang, U.C., 2024. Spatial and temporal distribution of ribosomes in single cells reveals aging differences between old and new daughters of Escherichia coli. eLife 12, RP89543. https://doi.org/10.7554/eLife.89543.3

Charaka, V.K., Misra, H.S., 2012. Functional Characterization of the Role of the Chromosome I Partitioning System in Genome Segregation in Deinococcus radiodurans. J. Bacteriol. 194, 5739–5748. https://doi.org/10.1128/JB.00610-12

Cherepanov, P.P., Wackernagel, W., 1995. Gene disruption in Escherichia coli: TcR and KmR cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance determinant. Gene 158, 9–14. https://doi.org/10.1016/0378-1119(95)00193-A

Chure, G., Cremer, J., 2023. An optimal regulation of fluxes dictates microbial growth in and out of steady state. eLife 12, e84878. https://doi.org/10.7554/eLife.84878

Cooper, S., Hsieh, M.L., 1988. The Rate and Topography of Cell Wall Synthesis during the Division Cycle of Escherichia coli Using N-Acetylglucosamine as a Peptidoglycan Label. Microbiology 134, 1717–1721. https://doi.org/10.1099/00221287-134-6-1717

Coquel, A.-S., Jacob, J.-P., Primet, M., Demarez, A., Dimiccoli, M., Julou, T., Moisan, L., Lindner, A.B., Berry, H., 2013. Localization of Protein Aggregation in Escherichia coli Is Governed by Diffusion and Nucleoid Macromolecular Crowding Effect. PLoS Comput. Biol. 9, e1003038. https://doi.org/10.1371/journal.pcbi.1003038

Cutler, K.J., Stringer, C., Lo, T.W., Rappez, L., Stroustrup, N., Brook Peterson, S., Wiggins, P.A., Mougous, J.D., 2022. Omnipose: a high-precision morphology-independent solution for bacterial cell segmentation. Nat. Methods 19, 1438–1448. https://doi.org/10.1038/s41592-022-01639-4

Dai, X., Zhu, M., Warren, M., Balakrishnan, R., Patsalo, V., Okano, H., Williamson, J.R., Fredrick, K., Wang, Y.-P., Hwa, T., 2017. Reduction of translating ribosomes enables Escherichia coli to maintain elongation rates during slow growth. Nat. Microbiol. 2, 16231. https://doi.org/10.1038/nmicrobiol.2016.231

Danilova, O., Reyes-Lamothe, R., Pinskaya, M., Sherratt, D., Possoz, C., 2007. MukB colocalizes with the oriC region and is required for organization of the two Escherichia coli chromosome arms into separate cell halves. Mol. Microbiol. 65, 1485–1492. https://doi.org/10.1111/j.1365-2958.2007.05881.x

Datsenko, K.A., Wanner, B.L., 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. Proc. Natl. Acad. Sci. 97, 6640–6645. https://doi.org/10.1073/pnas.120163297

Davis, J.H., Rubin, A.J., Sauer, R.T., 2011. Design, construction and characterization of a set of insulated bacterial promoters. Nucleic Acids Res. 39, 1131–1141. https://doi.org/10.1093/nar/gkq810

Donczew, M., Mackiewicz, P., Wróbel, A., Flärdh, K., Zakrzewska-Czerwińska, J., Jakimowicz, D., 2016. ParA and ParB coordinate chromosome segregation with cell elongation and division during *Streptomyces* sporulation. Open Biol. 6, 150263. https://doi.org/10.1098/rsob.150263

Donovan, C., Schwaiger, A., Krämer, R., Bramkamp, M., 2010. Subcellular Localization and Characterization of the ParAB System from *Corynebacterium glutamicum*. J. Bacteriol. 192, 3441–3451. https://doi.org/10.1128/JB.00214-10

Dourado, H., Lercher, M.J., 2020. An analytical theory of balanced cellular growth. Nat. Commun. 11, 1226. https://doi.org/10.1038/s41467-020-14751-w

Du, W.-L., Dubarry, N., Passot, F.M., Kamgoué, A., Murray, H., Lane, D., Pasta, F., 2016. Orderly Replication and Segregation of the Four Replicons of Burkholderia cenocepacia J2315. PLOS Genet. 12, e1006172. https://doi.org/10.1371/journal.pgen.1006172

Dworkin, J., Losick, R., 2002. Does RNA polymerase help drive chromosome segregation in bacteria? Proc. Natl. Acad. Sci. 99, 14089–14094. https://doi.org/10.1073/pnas.182539899

Dworsky, P., Schaechter, M., 1973. Effect of Rifampin on the Structure and Membrane Attachment of the Nucleoid of *Escherichia coli*. J. Bacteriol. 116, 1364–1374. https://doi.org/10.1128/jb.116.3.1364-1374.1973

Farrar, A., Turner, P., El Sayyed, H., Feehily, C., Chatzimichail, S., Ta, S., Crook, D., Andersson, M., Oakley, S., Barrett, L., Nellåker, C., Stoesser, N., Kapanidis, A., 2025. Ribosome phenotypes for rapid classification of antibiotic-susceptible and resistant strains of Escherichia coli. Commun. Biol. 8, 319. https://doi.org/10.1038/s42003-025-07740-6

Figge, R.M., Easter, J., Gober, J.W., 2003. Productive interaction between the chromosome partitioning proteins, ParA and ParB, is required for the progression of the cell cycle in Caulobacter crescentus: Dominant-negative ParB alleles. Mol. Microbiol. 47, 1225–1237. https://doi.org/10.1046/j.1365-2958.2003.03367.x

Fisher, J.K., Bourniquel, A., Witz, G., Weiner, B., Prentiss, M., Kleckner, N., 2013. Four-Dimensional Imaging of E. coli Nucleoid Organization and Dynamics in Living Cells. Cell 153, 882–895. https://doi.org/10.1016/j.cell.2013.04.006

Flory, P.J., 1942. Thermodynamics of High Polymer Solutions. J. Chem. Phys. 10, 51–61. https://doi.org/10.1063/1.1723621

Forchhammer, J., Lindahl, L., 1971. Growth rate of polypeptide chains as a function of the cell growth rate in a mutant of Escherichia coli 15. J. Mol. Biol. 55, 563–568. https://doi.org/10.1016/0022-2836(71)90337-8

Gibson, D.G., Young, L., Chuang, R.-Y., Venter, J.C., Hutchison, C.A., Smith, H.O., 2009. Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat. Methods 6, 343–345. https://doi.org/10.1038/nmeth.1318

Gillies, Sean, van der Wel, Casper, Van den Bossche, Joris, Taves, Mike W., Arnott, Joshua, Ward, Brendan C., Others, 2023. Shapely. https://doi.org/10.5281/ZENODO.5597138

Govers, S.K., Campos, M., Tyagi, B., Laloux, G., Jacobs-Wagner, C., 2024. Apparent simplicity and emergent robustness in the control of the Escherichia coli cell cycle. Cell Syst. 15, 19-36.e5. https://doi.org/10.1016/j.cels.2023.12.001

Gray, A.N., Egan, A.J., Van'T Veer, I.L., Verheul, J., Colavin, A., Koumoutsi, A., Biboy, J., Altelaar, A.F.M., Damen, M.J., Huang, K.C., Simorre, J.-P., Breukink, E., Den Blaauwen, T., Typas, A., Gross, C.A., Vollmer, W., 2015. Coordination of peptidoglycan synthesis and outer membrane constriction during Escherichia coli cell division. eLife 4, e07118. https://doi.org/10.7554/eLife.07118

Gray, W.T., Govers, S.K., Xiang, Y., Parry, B.R., Campos, M., Kim, S., Jacobs-Wagner, C., 2019. Nucleoid Size Scaling and Intracellular Organization of Translation across Bacteria. Cell 177, 1632-1648.e20. https://doi.org/10.1016/j.cell.2019.05.017

Guyer, M.S., Reed, R.R., Steitz, J.A., Low, K.B., 1981. Identification of a Sex-factor-affinity
Site in E. coli as. Cold Spring Harb. Symp. Quant. Biol. 45, 135–140.
https://doi.org/10.1101/SQB.1981.045.01.022
Guzman, L.M., Belin, D., Carson, M.J., Beckwith, J., 1995. Tight regulation, modulation, and
high-level expression by vectors containing the arabinose PBAD promoter. J. Bacteriol.
177, 4121–4130. https://doi.org/10.1128/jb.177.14.4121-4130.1995
Harju, J., Van Teeseling, M.C.F., Broedersz, C.P., 2024. Loop-extruders alter bacterial
chromosome topology to direct entropic forces for segregation. Nat. Commun. 15, 4618.
https://doi.org/10.1038/s41467-024-49039-w
Harris, C.R., Millman, K.J., Van Der Walt, S.J., Gommers, R., Virtanen, P., Cournapeau, D.,
Wieser, E., Taylor, J., Berg, S., Smith, N.J., Kern, R., Picus, M., Hoyer, S., Van
Kerkwijk, M.H., Brett, M., Haldane, A., Del Río, J.F., Wiebe, M., Peterson, P., Gérard-
Marchant, P., Sheppard, K., Reddy, T., Weckesser, W., Abbasi, H., Gohlke, C., Oliphant,
T.E., 2020. Array programming with NumPy. Nature 585, 357–362.
https://doi.org/10.1038/s41586-020-2649-2
Hartmann, G., Honikel, K.O., Knüsel, F., Nüesch, J., 1967. The specific inhibition of the DNA-
directed RNA synthesis by rifamycin. Biochim. Biophys. Acta BBA - Nucleic Acids
Protein Synth. 145, 843–844. https://doi.org/10.1016/0005-2787(67)90147-5
Hofmann, A., Mäkelä, J., Sherratt, D.J., Heermann, D., Murray, S.M., 2019. Self-organised
segregation of bacterial chromosomal origins. eLife 8, e46564.
https://doi.org/10.7554/eLife.46564
Holmes, V.F., Cozzarelli, N.R., 2000. Closing the ring: Links between SMC proteins and
chromosome partitioning, condensation, and supercoiling. Proc. Natl. Acad. Sci. 97,
1322–1324. https://doi.org/10.1073/pnas.040576797
Hsu, T.-M., Chang, Y.-R., 2019. High-Copy-Number Plasmid Segregation—Single-Molecule
Dynamics in Single Cells. Biophys. J. 116, 772–780.
https://doi.org/10.1016/j.bpj.2019.01.019
Hu, X.-P., Dourado, H., Schubert, P., Lercher, M.J., 2020. The protein translation machinery is
expressed for maximal efficiency in Escherichia coli. Nat. Commun. 11, 5260.
https://doi.org/10.1038/s41467-020-18948-x
Huggins, M.L., 1941. Solutions of Long Chain Compounds. J. Chem. Phys. 9, 440–440.
https://doi.org/10.1063/1.1750930
Hunter, J.D., 2007. Matplotlib: A 2D Graphics Environment. Comput. Sci. Eng. 9, 90–95.
https://doi.org/10.1109/MCSE.2007.55
Ireton, K., Gunther, N.W., Grossman, A.D., 1994. spo0J is required for normal chromosome
segregation as well as the initiation of sporulation in Bacillus subtilis. J. Bacteriol. 176,
5320–5329. https://doi.org/10.1128/jb.176.17.5320-5329.1994
Iwai, N., Nagai, K., Wachi, M., 2002. Novel S-Benzylisothiourea Compound That Induces
Spherical Cells in Escherichia coli Probably by Acting on a Rod-shape-determining
Protein(s) Other Than Penicillin-binding Protein 2. Biosci. Biotechnol. Biochem. 66,
2658–2662. https://doi.org/10.1271/bbb.66.2658
Jacob, F., Brenner, S., Cuzin, F., 1963. On the Regulation of DNA Replication in Bacteria. Cold
Spring Harb. Symp. Quant. Biol. 28, 329–348.
https://doi.org/10.1101/SQB.1963.028.01.048
Jakimowicz, D., Brzostek, A., Rumijowska-Galewicz, A., Żydek, P., Dołzbłasz, A., Smulczyk-
Krawczyszyn, A., Zimniak, T., Wojtasz, Ł., Zawilak-Pawlik, A., Kois, A., Dziadek, J.,

Zakrzewska-Czerwińska, J., 2007a. Characterization of the mycobacterial chromosome
segregation protein ParB and identification of its target in Mycobacterium smegmatis.
Microbiology 153, 4050–4060. https://doi.org/10.1099/mic.0.2007/011619-0
Jakimowicz, D., Żydek, P., Kois, A., Zakrzewska-Czerwińska, J., Chater, K.F., 2007b.
Alignment of multiple chromosomes along helical ParA scaffolding in sporulating
Streptomyces hyphae. Mol. Microbiol. 65, 625–641. https://doi.org/10.1111/j.1365-
2958.2007.05815.x
Jalal, A.S.B., Le, T.B.K., 2020. Bacterial chromosome segregation by the ParABS system. Open
Biol. 10, 200097. https://doi.org/10.1098/rsob.200097
Jecz, P., Bartosik, A.A., Glabski, K., Jagura-Burdzy, G., 2015. A Single parS Sequence from the
Cluster of Four Sites Closest to oriC Is Necessary and Sufficient for Proper Chromosome
Segregation in Pseudomonas aeruginosa. PLOS ONE 10, e0120867.
https://doi.org/10.1371/journal.pone.0120867
Jensen, K.F., 1993. The Escherichia coli K-12 "wild types" W3110 and MG1655 have an rph
frameshift mutation that leads to pyrimidine starvation due to low pyrE expression levels.
J. Bacteriol. 175, 3401–3407. https://doi.org/10.1128/jb.175.11.3401-3407.1993
Jun, S., Mulder, B., 2006. Entropy-driven spatial organization of highly confined polymers:
Lessons for the bacterial chromosome. Proc. Natl. Acad. Sci. 103, 12388–12393.
https://doi.org/10.1073/pnas.0605305103
Jun, S., Wright, A., 2010. Entropy as the driver of chromosome segregation. Nat. Rev.
Microbiol. 8, 600–607. https://doi.org/10.1038/nrmicro2391
Kadoya, R., Baek, J.H., Sarker, A., Chattoraj, D.K., 2011. Participation of Chromosome
Segregation Protein ParAI of *Vibrio cholerae* in Chromosome Replication. J. Bacteriol.
193, 1504–1514. https://doi.org/10.1128/JB.01067-10
Kaval, K.G., Chimalapati, S., Siegel, S.D., Garcia, N., Jaishankar, J., Dalia, A.B., Orth, K., 2023.
Membrane-localized expression, production and assembly of Vibrio parahaemolyticus
T3SS2 provides evidence for transertion. Nat. Commun. 14, 1178.
https://doi.org/10.1038/s41467-023-36762-z
Kawalek, A., Wawrzyniak, P., Bartosik, A.A., Jagura-Burdzy, G., 2020. Rules and Exceptions:
The Role of Chromosomal ParB in DNA Segregation and Other Cellular Processes.
Microorganisms 8, 105. https://doi.org/10.3390/microorganisms8010105
Kim, H.-J., Calcutt, M.J., Schmidt, F.J., Chater, K.F., 2000. Partitioning of the Linear
Chromosome during Sporulation of *Streptomyces coelicolor* A3(2) Involves an *oriC* -
Linked *parAB* Locus. J. Bacteriol. 182, 1313–1320.
https://doi.org/10.1128/JB.182.5.1313-1320.2000
Kim, Seunghyeon, Wang, Y.-H., Hassan, A., Kim, Sangjin, 2024. Re-defining how mRNA
degradation is coordinated with transcription and translation in bacteria.
https://doi.org/10.1101/2024.04.18.588412
Kjos, M., Veening, J.-W., 2014. Tracking of chromosome dynamics in live *S treptococcus*
pneumoniae reveals that transcription promotes chromosome segregation: Transcription
promotes chromosome segregation. Mol. Microbiol. 91, 1088–1105.
https://doi.org/10.1111/mmi.12517
Koch, A.L., Gross, G.H., 1979. Growth Conditions and Rifampin Susceptibility. Antimicrob.
Agents Chemother. 15, 220–228. https://doi.org/10.1128/AAC.15.2.220

Koleva, K.Z., Hellweger, F.L., 2015. From protein damage to cell aging to population fitness in
E. coli: Insights from a multi-level agent-based model. Ecol. Model. 301, 62–71.
https://doi.org/10.1016/j.ecolmodel.2015.01.024
Kuwada, N.J., Cheveralls, K.C., Traxler, B., Wiggins, P.A., 2013. Mapping the driving forces of
chromosome structure and segregation in Escherichia coli. Nucleic Acids Res. 41, 7370–
7377. https://doi.org/10.1093/nar/gkt468
Lagage, V., Boccard, F., Vallet-Gely, I., 2016. Regional Control of Chromosome Segregation in
Pseudomonas aeruginosa. PLOS Genet. 12, e1006428.
https://doi.org/10.1371/journal.pgen.1006428
Łapińska, U., Glover, G., Capilla-Lasheras, P., Young, A.J., Pagliara, S., 2019. Bacterial ageing
in the absence of external stressors. Philos. Trans. R. Soc. B Biol. Sci. 374, 20180442.
https://doi.org/10.1098/rstb.2018.0442
Lee, P.S., Grossman, A.D., 2006. The chromosome partitioning proteins Soj (ParA) and Spo0J
(ParB) contribute to accurate chromosome partitioning, separation of replicated sister
origins, and regulation of replication initiation in Bacillus subtilis. Mol. Microbiol. 60,
853–869. https://doi.org/10.1111/j.1365-2958.2006.05140.x
Lemon, K.P., Grossman, A.D., 2000. Movement of Replicating DNA through a Stationary
Replisome. Mol. Cell 6, 1321–1330. https://doi.org/10.1016/S1097-2765(00)00130-1
Lewis, P.J., Thaker, S.D., Errington, J., 2000. Compartmentalization of transcription and
translation in Bacillus subtilis. EMBO J. 19, 710–718.
https://doi.org/10.1093/emboj/19.4.710
Lewis, R.A., Bignell, C.R., Zeng, W., Jones, A.C., Thomas, C.M., 2002. Chromosome loss from
par mutants of Pseudomonas putida depends on growth medium and phase of growth.
Microbiology 148, 537–548. https://doi.org/10.1099/00221287-148-2-537
Li, H., 2019. Random Chromosome Partitioning in the Polyploid Bacterium *Thermus*
thermophilus HB27. G3 GenesGenomesGenetics 9, 1249–1261.
https://doi.org/10.1534/g3.119.400086
Li, H., Angelov, A., Pham, V.T.T., Leis, B., Liebl, W., 2015. Characterization of chromosomal
and megaplasmid partitioning loci in Thermus thermophilus HB27. BMC Genomics 16,
317. https://doi.org/10.1186/s12864-015-1523-3
Lim, H.C., Surovtsev, I.V., Beltran, B.G., Huang, F., Bewersdorf, J., Jacobs-Wagner, C., 2014.
Evidence for a DNA-relay mechanism in ParABS-mediated chromosome segregation.
eLife 3, e02758. https://doi.org/10.7554/eLife.02758
Lin, W.-H., Jacobs-Wagner, C., 2022. Connecting single-cell ATP dynamics to overflow
metabolism, cell growth, and the cell cycle in Escherichia coli. Curr. Biol. 32, 3911-
3924.e4. https://doi.org/10.1016/j.cub.2022.07.035
Lindner, A.B., Madden, R., Demarez, A., Stewart, E.J., Taddei, F., 2008. Asymmetric
segregation of protein aggregates is associated with cellular aging and rejuvenation. Proc.
Natl. Acad. Sci. 105, 3076–3081. https://doi.org/10.1073/pnas.0708931105
Linnik, D., Maslov, I., Punter, C.M., Poolman, B., 2024. Dynamic structure of E. coli cytoplasm:
supramolecular complexes and cell aging impact spatial distribution and mobility of
proteins. Commun. Biol. 7, 508. https://doi.org/10.1038/s42003-024-06216-3
Livny, J., Yamaichi, Y., Waldor, M.K., 2007. Distribution of Centromere-Like *parS* Sites in
Bacteria: Insights from Comparative Genomics. J. Bacteriol. 189, 8693–8703.
https://doi.org/10.1128/JB.01239-07

Mäkelä, J., Papagiannakis, A., Lin, W.-H., Lanz, M.C., Glenn, S., Swaffer, M., Marinov, G.K., Skotheim, J.M., Jacobs-Wagner, C., 2024. Genome concentration limits cell growth and modulates proteome composition in *Escherichia coli*. https://doi.org/10.1101/2024.02.26.582181

Mäkelä, J., Uphoff, S., Sherratt, D.J., 2021. Nonrandom segregation of sister chromosomes by *Escherichia coli* MukBEF. Proc. Natl. Acad. Sci. 118, e2022078118. https://doi.org/10.1073/pnas.2022078118

Mao, S., Chakraverti-Wuerthwein, M.S., Gaudio, H., Košmrlj, A., 2020. Designing the Morphology of Separated Phases in Multicomponent Liquid Mixtures. Phys. Rev. Lett. 125, 218003. https://doi.org/10.1103/PhysRevLett.125.218003

Mao, S., Kuldinow, D., Haataja, M.P., Košmrlj, A., 2019. Phase behavior and morphology of multicomponent liquid mixtures. Soft Matter 15, 1297–1311. https://doi.org/10.1039/C8SM02045K

Matsumoto, K., Hara, H., Fishov, I., Mileykovskaya, E., Norris, V., 2015. The membrane: transertion as an organizing principle in membrane heterogeneity. Front. Microbiol. 6. https://doi.org/10.3389/fmicb.2015.00572

McGuffee, S.R., Elcock, A.H., 2010. Diffusion, Crowding & Protein Stability in a Dynamic Molecular Model of the Bacterial Cytoplasm. PLoS Comput. Biol. 6, e1000694. https://doi.org/10.1371/journal.pcbi.1000694

McKinney, W., 2010. Data Structures for Statistical Computing in Python. Presented at the Python in Science Conference, Austin, Texas, pp. 56–61. https://doi.org/10.25080/Majora-92bf1922-00a

Miangolarra, A.M., Li, S.H.-J., Joanny, J.-F., Wingreen, N.S., Castellana, M., 2021. Steric interactions and out-of-equilibrium processes control the internal organization of bacteria. Proc. Natl. Acad. Sci. 118, e2106014118. https://doi.org/10.1073/pnas.2106014118

Minnen, A., Attaiech, L., Thon, M., Gruber, S., Veening, J.-W., 2011. SMC is recruited to oriC by ParB and promotes chromosome segregation in Streptococcus pneumoniae: Chromosome segregation in S. pneumoniae. Mol. Microbiol. 81, 676–688. https://doi.org/10.1111/j.1365-2958.2011.07722.x

Mohapatra, S., Weisshaar, J.C., 2018. Functional mapping of the *E. coli* translational machinery using single-molecule tracking. Mol. Microbiol. 110, 262–282. https://doi.org/10.1111/mmi.14103

Mohl, D.A., Gober, J.W., 1997. Cell Cycle–Dependent Polar Localization of Chromosome Partitioning Proteins in Caulobacter crescentus. Cell 88, 675–684. https://doi.org/10.1016/S0092-8674(00)81910-8

Molenaar, D., Van Berlo, R., De Ridder, D., Teusink, B., 2009. Shifts in growth strategies reflect tradeoffs in cellular economics. Mol. Syst. Biol. 5, 323. https://doi.org/10.1038/msb.2009.82

Mondal, J., Bratton, B.P., Li, Y., Yethiraj, A., Weisshaar, J.C., 2011. Entropy-Based Mechanism of Ribosome-Nucleoid Segregation in E. coli Cells. Biophys. J. 100, 2605–2613. https://doi.org/10.1016/j.bpj.2011.04.030

Nasehi Tehrani, J., McEwan, A., Jin, C., Van Schaik, A., 2012. L1 regularization method in electrical impedance tomography by using the L1-curve (Pareto frontier curve). Appl. Math. Model. 36, 1095–1105. https://doi.org/10.1016/j.apm.2011.07.055

Navarro, P.P., Vettiger, A., Ananda, V.Y., Llopis, P.M., Allolio, C., Bernhardt, T.G., Chao, L.H., 2022. Cell wall synthesis and remodelling dynamics determine division site architecture

and cell shape in Escherichia coli. Nat. Microbiol. 7, 1621–1634.
https://doi.org/10.1038/s41564-022-01210-z

Otsu, N., 1979. A Threshold Selection Method from Gray-Level Histograms. IEEE Trans. Syst. Man Cybern. 9, 62–66. https://doi.org/10.1109/TSMC.1979.4310076

Paintdakhi, A., Parry, B., Campos, M., Irnov, I., Elf, J., Surovtsev, I., Jacobs-Wagner, C., 2016. Oufti: an integrated software package for high-accuracy, high-throughput quantitative microscopy analysis: Oufti: image analysis software. Mol. Microbiol. 99, 767–777. https://doi.org/10.1111/mmi.13264

Pande, S., Mitra, D., Chatterji, A., 2023. Entropy mediated organization of E.coli chromosome in fast growth conditions. https://doi.org/10.48550/ARXIV.2304.02275

Parry, B.R., Surovtsev, I.V., Cabeen, M.T., O'Hern, C.S., Dufresne, E.R., Jacobs-Wagner, C., 2014. The Bacterial Cytoplasm Has Glass-like Properties and Is Fluidized by Metabolic Activity. Cell 156, 183–194. https://doi.org/10.1016/j.cell.2013.11.028

Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z., Gimelshein, N., Antiga, L., Desmaison, A., Köpf, A., Yang, E., DeVito, Z., Raison, M., Tejani, A., Chilamkurthy, S., Steiner, B., Fang, L., Bai, J., Chintala, S., 2019. PyTorch: An Imperative Style, High-Performance Deep Learning Library. https://doi.org/10.48550/ARXIV.1912.01703

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Müller, A., Nothman, J., Louppe, G., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., Duchesnay, É., 2012. Scikit-learn: Machine Learning in Python. https://doi.org/10.48550/ARXIV.1201.0490

Pelletier, J., Halvorsen, K., Ha, B.-Y., Paparcone, R., Sandler, S.J., Woldringh, C.L., Wong, W.P., Jun, S., 2012. Physical manipulation of the *Escherichia coli* chromosome reveals its soft nature. Proc. Natl. Acad. Sci. 109. https://doi.org/10.1073/pnas.1208689109

Pettijohn, D.E., Hecht, R., 1974. RNA Molecules Bound to the Folded Bacterial Genome Stabilize DNA Folds and Segregate Domains of Supercoiling. Cold Spring Harb. Symp. Quant. Biol. 38, 31–41. https://doi.org/10.1101/SQB.1974.038.01.006

Pountain, A.W., Jiang, P., Yao, T., Homaee, E., Guan, Y., Podkowik, M., Shopsin, B., Torres, V.J., Golding, I., Yanai, I., 2022. Transcription-replication interactions reveal principles of bacterial genome regulation (preprint). Systems Biology. https://doi.org/10.1101/2022.10.22.513359

Proenca, A.M., Rang, C.U., Qiu, A., Shi, C., Chao, L., 2019. Cell aging preserves cellular immortality in the presence of lethal levels of damage. PLOS Biol. 17, e3000266. https://doi.org/10.1371/journal.pbio.3000266

Rabinovitch, A., Zaritsky, A., Feingold, M., 2003. DNA–membrane interactions can localize bacterial cell center. J. Theor. Biol. 225, 493–496. https://doi.org/10.1016/S0022-5193(03)00292-3

Reyes-Lamothe, R., Tran, T., Meas, D., Lee, L., Li, A.M., Sherratt, D.J., Tolmasky, M.E., 2014. High-copy bacterial plasmids diffuse in the nucleoid-free space, replicate stochastically and are randomly partitioned at cell division. Nucleic Acids Res. 42, 1042–1051. https://doi.org/10.1093/nar/gkt918

Sadhir, I., Murray, S.M., 2023. Mid-cell migration of the chromosomal terminus is coupled to origin segregation in Escherichia coli. Nat. Commun. 14, 7489. https://doi.org/10.1038/s41467-023-43351-7

Sanamrad, A., Persson, F., Lundius, E.G., Fange, D., Gynnå, A.H., Elf, J., 2014. Single-particle tracking reveals that free ribosomal subunits are not excluded from the *Escherichia coli* nucleoid. Proc. Natl. Acad. Sci. 111, 11413–11418. https://doi.org/10.1073/pnas.1411558111

Santi, I., McKinney, J.D., 2015. Chromosome Organization and Replisome Dynamics in Mycobacterium smegmatis. mBio 6, e01999-14. https://doi.org/10.1128/mBio.01999-14

Sawitzke, J.A., Austin, S., 2000. Suppression of chromosome segregation defects of *Escherichia coli muk* mutants by mutations in topoisomerase I. Proc. Natl. Acad. Sci. 97, 1671–1676. https://doi.org/10.1073/pnas.030528397

Schaechter, M., MaalOe, O., Kjeldgaard, N.O., 1958. Dependency on Medium and Temperature of Cell Size and Chemical Composition during Balanced Growth of Salmonella typhimurium. J. Gen. Microbiol. 19, 592–606. https://doi.org/10.1099/00221287-19-3-592

Scholz, S.A., Diao, R., Wolfe, M.B., Fivenson, E.M., Lin, X.N., Freddolino, P.L., 2019. High-Resolution Mapping of the Escherichia coli Chromosome Reveals Positions of High and Low Transcription. Cell Syst. 8, 212-225.e9. https://doi.org/10.1016/j.cels.2019.02.004

Scott, D.W., 1992. Multivariate Density Estimation: Theory, Practice, and Visualization, 1st ed, Wiley Series in Probability and Statistics. Wiley. https://doi.org/10.1002/9780470316849

Scott, M., Klumpp, S., Mateescu, E.M., Hwa, T., 2014. Emergence of robust growth laws from optimal regulation of ribosome synthesis. Mol. Syst. Biol. 10, 747. https://doi.org/10.15252/msb.20145379

Seabold, S., Perktold, J., 2010. Statsmodels: Econometric and Statistical Modeling with Python. Presented at the Python in Science Conference, Austin, Texas, pp. 92–96. https://doi.org/10.25080/Majora-92bf1922-011

Shis, D.L., Bennett, M.R., 2013. Library of synthetic transcriptional AND gates built with split T7 RNA polymerase mutants. Proc. Natl. Acad. Sci. 110, 5028–5033. https://doi.org/10.1073/pnas.1220157110

Si, F., Li, D., Cox, S.E., Sauls, J.T., Azizi, O., Sou, C., Schwartz, A.B., Erickstad, M.J., Jun, Y., Li, X., Jun, S., 2017. Invariance of Initiation Mass and Predictability of Cell Size in Escherichia coli. Curr. Biol. 27, 1278–1287. https://doi.org/10.1016/j.cub.2017.03.022

Silva, F., Lourenço, O., Pina-Vaz, C., Rodrigues, A.G., Queiroz, J.A., Domingues, F.C., 2010. The Use of DRAQ5 to Monitor Intracellular DNA in Escherichia coli by Flow Cytometry. J. Fluoresc. 20, 907–914. https://doi.org/10.1007/s10895-010-0636-y

Slayter, H., Kiho, Y., Hall, C.E., Rich, A., 1968. AN ELECTRON MICROSCOPIC STUDY OF LARGE BACTERIAL POLYRIBOSOMES. J. Cell Biol. 37, 583–590. https://doi.org/10.1083/jcb.37.2.583

Spahn, C., Endesfelder, U., Heilemann, M., 2014. Super-resolution imaging of Escherichia coli nucleoids reveals highly structured and asymmetric segregation during fast growth. J. Struct. Biol. 185, 243–249. https://doi.org/10.1016/j.jsb.2014.01.007

Spahn, C., Middlemiss, S., Gómez-de-Mariscal, E., Henriques, R., Bode, H.B., Holden, S., Heilemann, M., 2023. Transertion and cell geometry organize the *Escherichia coli* nucleoid during rapid growth. https://doi.org/10.1101/2023.10.16.562172

Spahn, C.K., Glaesmann, M., Grimm, J.B., Ayala, A.X., Lavis, L.D., Heilemann, M., 2018. A toolbox for multiplexed super-resolution imaging of the E. coli nucleoid and membrane using novel PAINT labels. Sci. Rep. 8, 14768. https://doi.org/10.1038/s41598-018-33052-3

Stracy, M., Lesterlin, C., Garza De Leon, F., Uphoff, S., Zawadzki, P., Kapanidis, A.N., 2015.
Live-cell superresolution microscopy reveals the organization of RNA polymerase in the
bacterial nucleoid. Proc. Natl. Acad. Sci. 112. https://doi.org/10.1073/pnas.1507592112
Stylianidou, S., Brennan, C., Nissen, S.B., Kuwada, N.J., Wiggins, P.A., 2016. *SuperSegger* :
robust image segmentation, analysis and lineage tracking of bacterial cells. Mol.
Microbiol. 102, 690–700. https://doi.org/10.1111/mmi.13486
Subach, O.M., Cranfill, P.J., Davidson, M.W., Verkhusha, V.V., 2011. An Enhanced Monomeric
Blue Fluorescent Protein with the High Chemical Stability of the Chromophore. PLoS
ONE 6, e28674. https://doi.org/10.1371/journal.pone.0028674
Subramanian, S., Murray, S.M., 2023. Subdiffusive movement of chromosomal loci in bacteria
explained by DNA bridging. Phys. Rev. Res. 5, 023034.
https://doi.org/10.1103/PhysRevResearch.5.023034
Takacs, C.N., Wachter, J., Xiang, Y., Ren, Z., Karaboja, X., Scott, M., Stoner, M.R., Irnov, I.,
Jannetty, N., Rosa, P.A., Wang, X., Jacobs-Wagner, C., 2022. Polyploidy, regular
patterning of genome copies, and unusual control of DNA partitioning in the Lyme
disease spirochete. Nat. Commun. 13, 7173. https://doi.org/10.1038/s41467-022-34876-4
Thappeta, Y., Cañas-Duarte, S.J., Kallem, T., Fragasso, A., Xiang, Y., Gray, W., Lee, C.,
Cegelski, L., Jacobs-Wagner, C., 2024. Glycogen phase separation drives
macromolecular rearrangement and asymmetric division in *E. coli*.
https://doi.org/10.1101/2024.04.19.590186
Tiruvadi-Krishnan, S., Männik, Jaana, Kar, P., Lin, J., Amir, A., Männik, Jaan, 2022. Coupling
between DNA replication, segregation, and the onset of constriction in Escherichia coli.
Cell Rep. 38, 110539. https://doi.org/10.1016/j.celrep.2022.110539
Tseng, H.-C., Harwell, C.L., Martin, C.H., Prather, K.L., 2010a. Biosynthesis of chiral 3-
hydroxyvalerate from single propionate-unrelated carbon sources in metabolically
engineered E. coli. Microb. Cell Factories 9, 96. https://doi.org/10.1186/1475-2859-9-96
Tseng, H.-C., Harwell, C.L., Martin, C.H., Prather, K.L., 2010b. Biosynthesis of chiral 3-
hydroxyvalerate from single propionate-unrelated carbon sources in metabolically
engineered E. coli. Microb. Cell Factories 9, 96. https://doi.org/10.1186/1475-2859-9-96
Van Der Walt, S., Schönberger, J.L., Nunez-Iglesias, J., Boulogne, F., Warner, J.D., Yager, N.,
Gouillart, E., Yu, T., 2014. scikit-image: image processing in Python. PeerJ 2, e453.
https://doi.org/10.7717/peerj.453
Varricchio, F., Monier, R., 1971. Ribosome Patterns in *Escherichia coli* Growing at Various
Rates. J. Bacteriol. 108, 105–110. https://doi.org/10.1128/jb.108.1.105-110.1971
Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski,
E., Peterson, P., Weckesser, W., Bright, J., Van Der Walt, S.J., Brett, M., Wilson, J.,
Millman, K.J., Mayorov, N., Nelson, A.R.J., Jones, E., Kern, R., Larson, E., Carey, C.J.,
Polat, İ., Feng, Y., Moore, E.W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R.,
Henriksen, I., Quintero, E.A., Harris, C.R., Archibald, A.M., Ribeiro, A.H., Pedregosa,
F., Van Mulbregt, P., SciPy 1.0 Contributors, Vijaykumar, A., Bardelli, A.P., Rothberg,
A., Hilboll, A., Kloeckner, A., Scopatz, A., Lee, A., Rokem, A., Woods, C.N., Fulton, C.,
Masson, C., Häggström, C., Fitzgerald, C., Nicholson, D.A., Hagen, D.R., Pasechnik,
D.V., Olivetti, E., Martin, E., Wieser, E., Silva, F., Lenders, F., Wilhelm, F., Young, G.,
Price, G.A., Ingold, G.-L., Allen, G.E., Lee, G.R., Audren, H., Probst, I., Dietrich, J.P.,
Silterra, J., Webber, J.T., Slavič, J., Nothman, J., Buchner, J., Kulick, J., Schönberger,
J.L., De Miranda Cardoso, J.V., Reimer, J., Harrington, J., Rodríguez, J.L.C., Nunez-

Iglesias, J., Kuczynski, J., Tritz, K., Thoma, M., Newville, M., Kümmerer, M.,
Bolingbroke, M., Tartre, M., Pak, M., Smith, N.J., Nowaczyk, N., Shebanov, N., Pavlyk,
O., Brodtkorb, P.A., Lee, P., McGibbon, R.T., Feldbauer, R., Lewis, S., Tygier, S.,
Sievert, S., Vigna, S., Peterson, S., More, S., Pudlik, T., Oshima, T., Pingel, T.J.,
Robitaille, T.P., Spura, T., Jones, T.R., Cera, T., Leslie, T., Zito, T., Krauss, T.,
Upadhyay, U., Halchenko, Y.O., Vázquez-Baeza, Y., 2020. SciPy 1.0: fundamental
algorithms for scientific computing in Python. Nat. Methods 17, 261–272.
https://doi.org/10.1038/s41592-019-0686-2
Wang, P., Robert, L., Pelletier, J., Dang, W.L., Taddei, F., Wright, A., Jun, S., 2010. Robust
Growth of Escherichia coli. Curr. Biol. 20, 1099–1103.
https://doi.org/10.1016/j.cub.2010.04.045
Wang, X., Liu, X., Possoz, C., Sherratt, D.J., 2006. The two *Escherichia coli* chromosome arms
locate to separate cell halves. Genes Dev. 20, 1727–1731.
https://doi.org/10.1101/gad.388406
Waskom, M., 2021. seaborn: statistical data visualization. J. Open Source Softw. 6, 3021.
https://doi.org/10.21105/joss.03021
Weitao, T., Nordstrom, K., Dasgupta, S., 1999. Mutual suppression of mukB and seqA
phenotypes might arise from their opposing influences on the Escherichia coli nucleoid
structure. Mol. Microbiol. 34, 157–168. https://doi.org/10.1046/j.1365-
2958.1999.01589.x
White, S., Quinn, J., Enzor, J., Staats, J., Mosier, S.M., Almarode, J., Denny, T.N., Weinhold,
K.J., Ferrari, G., Chan, C., 2021. FlowKit: A Python Toolkit for Integrated Manual and
Automated Cytometry Analysis Workflows. Front. Immunol. 12, 768541.
https://doi.org/10.3389/fimmu.2021.768541
Wientjes, F.B., Nanninga, N., 1989. Rate and topography of peptidoglycan synthesis during cell
division in Escherichia coli: concept of a leading edge. J. Bacteriol. 171, 3412–3419.
https://doi.org/10.1128/jb.171.6.3412-3419.1989
Wiktor, J., Gynnå, A.H., Leroy, P., Larsson, J., Coceano, G., Testa, I., Elf, J., 2021. RecA finds
homologous DNA by reduced dimensionality search. Nature 597, 426–429.
https://doi.org/10.1038/s41586-021-03877-6
Woldringh, C.L., 2002. The role of co-transcriptional translation and protein translocation
(transertion) in bacterial chromosome segregation. Mol. Microbiol. 45, 17–29.
https://doi.org/10.1046/j.1365-2958.2002.02993.x
Woldringh, C.L., Huls, P., Pas, E., Brakenhoff, G.J., Nanninga, N., 1987. Topography of
Peptidoglycan Synthesis during Elongation and Polar Cap Formation in a Cell Division
Mutant of Escherichia coli MC4100. Microbiology 133, 575–586.
https://doi.org/10.1099/00221287-133-3-575
Wu, F., Japaridze, A., Zheng, X., Wiktor, J., Kerssemakers, J.W.J., Dekker, C., 2019a. Direct
imaging of the circular chromosome in a live bacterium. Nat. Commun. 10, 2194.
https://doi.org/10.1038/s41467-019-10221-0
Wu, F., Swain, P., Kuijpers, L., Zheng, X., Felter, K., Guurink, M., Solari, J., Jun, S., Shimizu,
T.S., Chaudhuri, D., Mulder, B., Dekker, C., 2019b. Cell Boundary Confinement Sets the
Size and Position of the E. coli Chromosome. Curr. Biol. 29, 2131-2144.e4.
https://doi.org/10.1016/j.cub.2019.05.015

Wu, L.J., Lee, S., Park, S., Eland, L.E., Wipat, A., Holden, S., Errington, J., 2020. Geometric
principles underlying the proliferation of a model cell system. Nat. Commun. 11, 4149.
https://doi.org/10.1038/s41467-020-17988-7
Xiang, Y., Surovtsev, I.V., Chang, Y., Govers, S.K., Parry, B.R., Liu, J., Jacobs-Wagner, C.,
2021. Interconnecting solvent quality, transcription, and chromosome folding in
Escherichia coli. Cell 184, 3626-3642.e14. https://doi.org/10.1016/j.cell.2021.05.037
Yamaichi, Y., Fogel, M.A., McLeod, S.M., Hui, M.P., Waldor, M.K., 2007. Distinct
Centromere-Like *parS* Sites on the Two Chromosomes of *Vibrio* spp. J. Bacteriol. 189,
5314–5324. https://doi.org/10.1128/JB.00416-07
Yang, D., Männik, Jaana, Retterer, S.T., Männik, Jaan, 2020. The effects of polydisperse
crowders on the compaction of the *Escherichia coli* nucleoid. Mol. Microbiol. 113, 1022–
1037. https://doi.org/10.1111/mmi.14467
Yao, S., Helinski, D.R., Toukdarian, A., 2007. Localization of the Naturally Occurring Plasmid
ColE1 at the Cell Pole. J. Bacteriol. 189, 1946–1953. https://doi.org/10.1128/JB.01451-
06
Youngren, B., Nielsen, H.J., Jun, S., Austin, S., 2014. The multifork *Escherichia coli*
chromosome is a self-duplicating and self-segregating thermodynamic ring polymer.
Genes Dev. 28, 71–84. https://doi.org/10.1101/gad.231050.113
Yu, W., Herbert, S., Graumann, P.L., Götz, F., 2010. Contribution of SMC (Structural
Maintenance of Chromosomes) and SpoIIIE to Chromosome Segregation in
Staphylococci. J. Bacteriol. 192, 4067–4073. https://doi.org/10.1128/JB.00010-10
Zhang, C., Shao, P.G., Van Kan, J.A., Van Der Maarel, J.R.C., 2009. Macromolecular crowding
induced elongation and compaction of single DNA molecules confined in a nanochannel.
Proc. Natl. Acad. Sci. 106, 16651–16656. https://doi.org/10.1073/pnas.0904741106
Zhou, Z., Siddiquee, M.M.R., Tajbakhsh, N., Liang, J., 2020. UNet++: Redesigning Skip
Connections to Exploit Multiscale Features in Image Segmentation. IEEE Trans. Med.
Imaging 39, 1856–1867. https://doi.org/10.1109/TMI.2019.2959609
Zimmerman, S.B., Trach, S.O., 1991. Estimation of macromolecule concentrations and excluded
volume effects for the cytoplasm of Escherichia coli. J. Mol. Biol. 222, 599–620.
https://doi.org/10.1016/0022-2836(91)90499-V

2115
2116

FIGURE LEGENDS

Figure 1: Correlations between polysome and nucleoid dynamics at the single-cell level. CJW7323

cells were grown in M9gluCAAT in a microfluidic device. **A.** Schematic illustrating observable nucleoid

segregation events. **B.** Fluorescence images of RplA-GFP and HupA-mCherry for a representative cell

(CJW7323) from birth to division. **C.** Ensemble kymographs of the average RplA-GFP and HupA-mCherry

fluorescence during the cell division cycle (>300,000 segmented cell instances from 4122 complete cell

division cycles). The average relative timing of cell constriction initiation was estimated as shown in Figure

1 – figure supplement 2. **D.** Two-dimensional projections of the average RplA-GFP and HupA-mCherry

fluorescence signals in predivisional cells (4564 cells with two nucleoid objects, from 1907 cell division

cycles, 95-100% into the cell division cycle) and their intensity profiles. White arrows indicate RplA-GFP

enrichments at the quarter cell positions, while the black arrow indicates the site of cell constriction. **E.** Plot

showing the dynamics of RplA-GFP accumulation and HupA-mCherry depletion at mid-nucleoid (median ±

IQR) during the nucleoid cycle (see Figure 1 – Figure supplement 2D). Data from 3240 nucleoid segregation

cycles are shown (40 nucleoid cycle bins, 2512-4823 segmented nucleoids per bin). **F.** Correlation

(Spearman ρ = -0.52, p-value < 10^{-10}) between the rate of RplA-GFP accumulation in the middle of the

nucleoid ($\frac{\Delta RplA_{mid-nuc}\ conc.}{\Delta T}$) and the rate of HupA-mCherry depletion in the same region ($\frac{\Delta HupA_{mid-nuc}\ conc.}{\Delta T}$)

between the initiation of nucleoid splitting and just before the end of nucleoid splitting (3214 complete

nucleoid cycles). The colormap indicates the Gaussian kernel density estimation (KDE). Solid black line

indicates the linear-regression fit to the data. **G.** Percentage of cells that continue to accumulate polysomes

in the middle of the cell ($\frac{\Delta RplA_{mid-cell}\ conc.}{\Delta T}$) during four relative time bins (1335 to 1957 cell division cycles

per bin) covering the period from the end of nucleoid splitting until cell division. **H.** Correlation (Spearman

ρ = 0.47, p-value < 10^{-10}) between the rate of RplA-GFP accumulation at mid-cell ($\frac{\Delta RplA_{mid-cell}\ conc.}{\Delta T}$) and the

rate of distance increase between the sister nucleoids ($\frac{\Delta Distance_{nuc}}{\Delta T}$) during the first quartile (0-25%) of the

period between the end of nucleoid splitting and cell division (1376 cell division cycles). The black markers

correspond to nine bins (mean ± SEM, 75 to 177 cell division cycles per bin) within the 5^{th}-95^{th} percentiles

of *x*-axis range. Also shown is the distribution of the cell elongation rates ($\frac{\Delta Length_{cell}}{\Delta T}$) during the same time

interval, with the mean and SD shown by the solid and dashed lines, respectively. **I.** Plot showing the

coefficients of a linear mixed-effects model (see eq. 3 in Methods and Figure 1 – figure supplement 3B) for

four interval bins between the completion of nucleoid splitting and cell division. The coefficients quantify the

relative contribution of polysome accumulation at mid-cell ($\frac{\Delta RplA_{mid-cell}\ conc.}{\Delta T}$) and cell elongation ($\frac{\Delta Length_{cell}}{\Delta T}$)

to the rate of sister nucleoid migration ($\frac{\Delta Distance_{nuc}}{\Delta T}$). All coefficients are significant (Prob(<|Z|) < 10^{-9}), except

for the one marked with an asterisk that is marginally significant (Prob(<|Z|) = 0.02).

2149

Figure 2: Correlation of the extent and relative timing of polysome accumulation with nucleoid

segregation at the population level. A. RplA-mEos2 and DAPI (scaled by the whole cell average)

demographs constructed from snapshots of DAPI-stained CJW6768 cells (815 to 2771 cells per

demograph) expressing RplA-mEos2 and growing in different nutrient conditions (see Table S1 for

abbreviations). The demographs were arranged from smallest (M9mann, top) to biggest average cell area

(M9malaCAAT, bottom). Additional demographs for different ribosomal reporters and nutrient conditions

are shown in Figure 2 – figure supplement 1. **B.** Correlation between average polysome accumulation and

average nucleoid depletion at mid-cell for all tested strains (Spearman ρ_{all} = -0.85, p-value < 10^{-10}) with

different ribosomal and nucleoid reporters, and within each strain (-0.75 < Spearman ρ_{strain} < -0.95, p-values

< 10^{-3}) across nutrient conditions. A linear regression was fitted to all the data.

2160

2161 **Figure 3: Correlations between polysome and nucleoid asymmetries. A.** Distributions of RplA-GFP

2162 concentration in the new (grey) and the old (black) pole regions of newborn cells (0-2.5% of the cell division

cycle, n = 912 cell division cycles). The histograms were smoothed using Gaussian kernel density estimations. **B.** Correlation (Spearman ρ = -0.41, p-value < 10^{-10}) between the polar polysome asymmetry and the position of the nucleoid centroid around the cell center of cells at the beginning of the division cycle (0-10%, n = 1179 cell division cycles). The contour plot consists of 9 levels with a lower data density threshold of 25%. The polar polysome profile for cells with correlations indicated by the numbers 1 and 2 are schematically illustrated in the next panel. A linear regression (solid grey line) was fitted to the data. **C.** Schematic illustrating the effects of the relative polysome abundance between the poles on the position of the nucleoid. **D.** Average 2D projections of the RplA-GFP and HupA-mCherry concentration (conc.) at different cell division cycle intervals (~9440 to 47240 cell images per cell division cycle interval from 4122 cell division cycles). The dotted line indicates the boundary between two cell division cycles. **E.** Density plot comparing the distribution of the HupA-mCherry maximum concentration toward the new pole (gray) to that toward the old pole (black) in newborn cells (0-2.5% of the division cycle, n = 912 cell division cycles). The histograms were smoothed using Gaussian kernel density estimations. **F.** Correlation (Spearman ρ = 0.52, p-value<10^{-10}) between the nucleoid density asymmetry and the relative availability of polysome-free space between the two cell halves early in the cell division cycle (0-10% of the division cycle, n = 2150 cell division cycles). The contour plot consists of 9 levels with a lower data density threshold of 25%. Values above 1 on the *x*-axis indicate more polysome-free space toward the new pole, and values below 1 correspond to cells with more polysome-free space toward the old pole. On the *y*-axis, values above 1 indicate higher DNA density toward the new pole and values below 1 indicate higher DNA density toward the old pole. **G.** Average 2D projections of newborn cells (0-10% into the cell division cycle) from the lower-left quartile in panel C (region 2, n = 223 cell division cycles) and the upper right quartile in panel C (region 1, n = 557 cell division cycles) and their 1D intensity profiles.

Figure 4: Simulation results of the reaction-diffusion model for different growth rates or nucleoid diffusion rates. A. Simulation of a non-growing virtual cell, initialized with homogeneous polysome concentration and a nucleoid spread between the two poles (*t* = 0 s). It reaches steady state with the nucleoid compacted at mid-cell at *t* = 998 s. **B.** 1D simulation of polysome (blue) and nucleoid (red) dynamics during slow growth (growth rate = 0.25 h^{-1}, D_n = 0.001 μm^2/s, cell length at birth = 2.2 μm) at different relative cell division cycle timepoints. The simulation was initialized from the equilibrium polysome and nucleoid distribution (at 0%). **C.** Schematic summarizing how polysomes accumulate in the middle of the elongating nucleoid, causing nucleoid splitting. **D.** Correlation between the relative timing of nucleoid splitting and the growth rate as captured by our reaction-diffusion model (D_n = 0.001 μm^2/sec) across six growth rate bins. The cell and nucleoid lengths for each growth rate bin matched previously published data (Govers et al., 2024). **E.** Deviation between the steady state after infinite relaxation time (dashed curves) and the polysome or nucleoid profiles in newborn cells after one simulation round (solid curves) for increasing nucleoid diffusion constants. The simulations were performed for a growth rate of 0.57 h^{-1}, which is comparable to the average growth rate in our microfluidic experiments (Figure 2 – figure supplement 2A).

Figure 5: Comparison of the nonequilibrium polysome accumulation with freely diffusing particles.

A. Phase contrast and fluorescence (fluor.) images of two representative single cells (CJW7651). The red

circles indicate the position of the mCherry-µNS particle in each cell. **B.** Two-dimensional average cell

projections of the DAPI concentration (conc.) and the RplA-msfGFP concentration, and 2D histogram of

the mCherry-µNS particle density for four cell length bins of CJW7651 cells (~2580 cells per bin) grown in

M9gluCAAT and spotted on an agarose pad containing the same medium. Since the cell pole identity

cannot be inferred from snapshot images, pole assignment was random. **C.** Average 1D profiles of the

scaled DAPI and RplA-msfGFP concentrations and the mCherry-µNS probability density.

2209

Figure 6: Effects of rifampicin treatment and polysome depletion on nucleoid segregation and

compaction. A. Plot showing the average instantaneous growth rate (mean ± SD shown by the solid black

curve and grey shaded region, respectively) of a cell population (n = 2629 cell division cycles) undergoing

two rounds of rifampicin treatment in a microfluidic device supplemented with M9gluCAAT. The distribution

of the average cell cycle growth rate of unperturbed populations is also shown on the right (n = 4122 cell

division cycles from a different microfluidics experiment). The solid horizontal line indicates the average

growth rate. **B.** Plot showing the average distance between nucleoid peaks for a population of cells

(squares, 114 cell division cycles) that were born (-112 to -102 min) and divided before the addition of

rifampicin, and for a population of cells (circles, 112 cell division cycles) that were born just before (-22 to -

12 min) and divided after the addition of rifampicin. A third-degree polynomial was fitted to the data from

the unperturbed population (solid blue curve) and juxtaposed (dashed blue curve) with the data from the

interrupted population. **C.** Average 1D profile and 2D projections of the scaled (divided by the whole cell

average concentration) RplA-GFP and HupA-mCherry signals for cells before and after rifampicin addition

(n = 112 cell division cycles). The red dashed horizontal lines in the 1D intensity profiles and the white

crosses in the 2D profiles mark the nucleoid peaks. **D.** Plot showing the RplA-GFP accumulation relative to

the HupA-mCherry depletion at mid-cell from 0 to 24 min after birth (colormap) for cells that completed their

division cycle before the addition of rifampicin (n = 114 cell division cycles) and for cells that were subjected

to rifampicin 12 min (n = 112 cell division cycles) or 3 min (n = 99 cell division cycles) after birth. **E.** Average

1D and 2D scaled RplA-GFP and HupA-mCherry intensity profiles for newborn cells (0 to 10 min after birth)

before (left, n = 726 cell division cycles), just after (middle, n = 367 cell division cycles), and much after

(right, n = 235 cell division cycles) rifampicin addition.

2231

Figure 7: Effects of ectopic polysome accumulation on nucleoid dynamics. A. Schematic

summarizing the experiment. **B.** Representative phase contrast and mTagBFP2 fluorescence images at

different times after induction with IPTG (100 µM) are shown, next to a plot showing the mTagBFP2

fluorescence of the entire population (mean ± SD, n = 3624 cell trajectories) over time. **C.** Plot showing how

instantaneous growth rate (mean ±SD, n = 3624 mTagBFP2 induction trajectories) decreases following

induction of mTagBFP2 synthesis. **D-F.** Representative kymographs and images of the normalized (divided
by the whole cell average) RplA-msfGFP and HupA-mCherry fluorescence signals in cells (CJW7798) born
during mTagBFP2 over-expression. **F.** Phase contrast images are shown to illustrate the formation of
inclusion bodies (see also Figure 6 – figure supplement 1). Additional cell examples are shown in Movie
S5.

2242

Figure 8: Effects of cell width increase on polysome and nucleoid dynamics. A. Comparison of the
cell width increase during cell growth between CJW7323 cells treated with cephalexin (mean ± SD, 360 cell
growth trajectories, 418 to 1511 segmented cells per bin) and cells treated with both cephalexin (50 µg/mL)
and A22 (4 µg/mL) (grey, mean ± SD, 309 cell growth trajectories, 51 to 1684 segmented cells per bin).
The same cell area bins are compared between the two populations. **B.** Phase contrast and fluorescence
images of a representative cephalexin-treated cell expressing RplA-GFP and HupA-mCherry. **C.** Same as
B but for a short cell growing in the presence of A22 and cephalexin. **D.** Same as C but for a longer cell.
The white arrowheads indicate the polysome bridges that connect polysome accumulations between two
DNA-free regions. Additional examples are shown in Movie S6 **E.** Representative fluorescence images of
RplA-GFP and HupA-mCherry in a cell treated with A22 and cephalexin. The dotted lines indicate the
representative cross-like polysome accumulation, which forms during the fusion of the polysome
accumulation towards the center (see also Movie S6). **F.** Comparison of the segmented polysome
accumulations and nucleoid objects between A22+cephalexin (150 sampled segmented cells from 47
growth trajectories) and cephalexin-treated (150 sampled segmented cells from 100 growth trajectories)
cells. The polysome and nucleoid areas per cell were normalized by the population-average statistic from
cephalexin-treated cells. All differences between the two populations are statistically significant (Mann-
Whitney p-value $< 10^{-10}$).

2260

SUPPLEMENTARY FIGURE LEGENDS
Figure 1 – figure supplement 1: Reproducibility analysis of the dynamic ribosome and nucleoid
distributions between microfluidic experiments. While phase-contrast images were acquired every
minute in all microfluidics experiments, two different intervals (1 min and 3 min) were used for fluorescence
image acquisition. **A.** Kymographs showing the RplA-GFP and HupA-mCherry concentration in CJW7323
cells growing in microfluidic channels in M9gluCAAT. Cell and nucleoid contours are shown in each channel
using a different color for each cell lineage (from dark purple to bright orange). **B-E.** Two biological replicate
experiments were performed for this strain and nutrient condition using different intervals of fluorescence
image acquisition (1 or 3 min). **B.** Plots showing that the instantaneous and average division cycle growth
rates were nearly identical between the two experiments. **C.** Plots showing the distributions of the indicated
fluorescence signals for the 1-min acquisition interval (black) compared to the 3-min frame rate interval
(orange) before and after correction. The difference in the excitation power between the two experiments
was 50% (240 %ms vs. 360 %ms), which was also reflected in the ratio of their average fluorescence values

(1.4 for RplA-GFP and 1.6 for HupA-mCherry). The fluorescence values from the 1-min interval experiment
were corrected by multiplying by these ratios, resulting in a near-perfect overlap between fluorescence
distributions after correction. **D.** Plots showing that the scaled (z-score) RplA-GFP and HupA-mCherry
intensity profiles from birth to division were almost identical between the two experiments. The intensity
profiles are shown for 10 cell division cycle intervals (1489 and 2633 cell division cycles for the 1-min and
3-min interval experiments, respectively). The z-score was calculated for each segmented cell by
subtracting the whole cell average fluorescence and then dividing the difference by the standard deviation.
Each intensity profile corresponds to the average z-score of all segmented cell instances within the
corresponding cell cycle interval. **E.** Plot showing that the average RplA-GFP concentration from birth to
division (averages ± SD across 20 cell division cycle bins) remained constant and was virtually identical
between the two experiments.

2285

2286 **Figure 1 – figure supplement 2: Determination of the relative timing of cell constriction.** The average
2287 cell width at mid-cell was used to estimate the relative timing of cell constriction from birth (0%) to division
2288 (100%). Considering the variability in the constriction onset across single cells, the average timing
2289 corresponds to the elbow of the curve, or the point with the maximum distance from the linear segment that
2290 connects the ends of the cell width curve, also known as the chord.

2291

Figure 1 – figure supplement 3: Intracellular distributions of RplA-GFP in rifampicin-treated cells
following rifampicin removal. A. Representative image of RplA-GFP fluorescence in cells (CJW7323)
treated with rifampicin (100 μg/mL). The cells were treated for 45 min, washed, and spotted on an
M9gluCAAT agarose pad without antibiotic. The presented snapshot corresponds to the first timepoint
immediately after cell spotting showing diffuse distribution of RplA-GFP signal. **B.** Time-lapse fluorescence
(fluor.) images showing the emergence of RplA-GFP signal accumulation at mid-cell (single arrow) or
quarter cell positions (double arrows) during recovery from rifampicin treatment (same experiment as in
panel A).

2300

Figure 1 – figure supplement 4: Tracking nucleoid segregation cycles. Since the timing of nucleoid
segregation varied between cell division cycles, the nucleoid segregation cycles were tracked
independently of the cell division cycles for each cell lineage to measure the relative timing of RplA-GFP
accumulation and HupA-mCherry depletion in the middle of the nucleoid (as shown in Figure 1E). **A.** Plots
showing the frequency of cells with one, two, three, or four detected nucleoid objects across 10 cell division
cycle bins from birth (top) to division (bottom). Data from 4122 cell division cycles are shown. **B.** The polarity
of the cells (+/-) and the relative position of the nucleoid mask (toward the new or old pole) was used to
track the nucleoid segregation cycle. This strategy was used to identify four groups of nucleoid segregation
cycles (group -2, -1, 1 and 2). Group -2 includes nucleoids that were "born" toward the old cell pole in a
mother cell with negative (-) polarity and were inherited at the center of a daughter cell with a negative

polarity. Group -1 includes nucleoids that were "born" toward the new cell pole in a mother cell with negative

(-) polarity and were inherited at the center of a daughter cell with a positive (+) polarity. Group 1 includes

nucleoids that were "born" toward the new cell pole in a mother cell with positive (+) polarity and were

inherited at the center of a daughter cell with negative (-) polarity. Group 2 includes nucleoids that were

"born" toward the old cell pole in a mother cell with positive (+) polarity and were inherited at the center of

a daughter cell with positive (+) polarity. **C.** Plots showing the distributions of nucleoid positions around the

cell center for 10 nucleoid segregation cycle bins, from the time a nucleoid was "born" (top) until it split

(bottom). These plots show the inheritance of the nucleoids from the quarter cell positions of the mother

cells to the middle of their daughters for the four groups of nucleoid segregation cycles. The solid gray line

represents the average density of the nucleoid positions for all four groups. Data from 2286 complete

nucleoid cycles are shown. **D.** Schematic that explains the definition of a nucleoid cycle. The nucleoid cycle

ranges from the end of a nucleoid splitting event, until the next splitting of the sister nucleoids. It usually

extends beyond cell division, into the next cell division cycle.

2324

Figure 1 – figure supplement 5: Correlations used to calculate the relative contribution of polysome

accumulation and cell elongation to nucleoid migration. A. Correlation (Spearman ρ = 0.47, p-value <

10^{-10}) between the rate of RplA-GFP accumulation at mid-cell ($\frac{\Delta RplA_{mid-cell\ conc.}}{\Delta T}$) and the rate of distance

increase between the sister nucleoids minus the rate of cell elongation ($\frac{\Delta Distance_{nuc}}{\Delta T} - \frac{\Delta Length_{cell}}{\Delta T}$) across

cells. The colormap corresponds to a Gaussian kernel density estimation. Binned data are also shown

(mean ± SEM, 75 to 177 cell division cycles per bin, 9 bins in total) within the 5th-95th percentiles of the *x*-

axis range. **B.** Scaled correlations (z-scores) between $\frac{\Delta RplA_{mid-cell\ conc.}}{\Delta T}$ (blue) or $\frac{\Delta Length_{cell}}{\Delta T}$ (grey) and

$\frac{\Delta Distance_{nuc}}{\Delta T}$ (y-axis) during four relative time bins (1335 to 1957 cell division cycles per bin) covering the

period from the end of nucleoid splitting until cell division. Shown are the Spearman correlations (ρ), all with

a p-value below 10^{-6}, except for the one marked with an asterisk (p-value = 0.02).

2335

Figure 1 – figure supplement 6: Examination of the relative timing of the initiation of nucleoid

constriction and the accumulation of polysomes at mid-nucleoid. A. RplA-GFP and HupA-mCherry

concentration (fluorescence arbitrary units) images of four representative single cells (CJW7323) growing

in M9glyT and their fluorescence intensity profiles along the cell length. **B.** Demographs of the scaled RplA-

GFP and HupA-mCherry concentration from 13554 *E. coli* cells grown in M9glyT. **C.** Average 1D intensity

profiles of the scaled (divided by the whole cell average) RplA-GFP and HupA-mCherry concentration for

12 bins of cells lengths (>1100 cells per intensity profile). **D.** Plot showing the scaled RplA-GFP and HupA-

mCherry concentration in the middle of the cells for increasing cell length (~370 cells per bin, 25 bins). Cells

longer than 3 μm were excluded to avoid the effects of cell constriction. The arrow and vertical dashed line

indicate the minimum cell length bin with an apparent HupA-mCherry depletion at mid-cell, marking the

initiation of nucleoid splitting. The error bars indicate mean ± standard error of the mean (SEM).

2347

Figure 2 – figure supplement 1: Demographs of scaled ribosome and nucleoid fluorescence for
strains with different ribosome markers and under various nutrient conditions. Demographs are
ordered according to increasing average cell area, which scales with growth rate (Schaechter et al., 1958),
from left to right and top to bottom. These demographs were constructed from snapshots of the following
strains: **A.** CJW6768 (RplA-mEos2 ribosomal marker, 747 to 2446 cells per condition), **B.** CJW6769 (RpsB-
mEos2 ribosomal marker, 690 to 3169 cells per condition), **C.** CJW7020 (RplA-msfGFP ribosomal marker,
657 to 2432 cells per condition) and **D.** CJW7021 (RpsB-msfGFP ribosomal marker, 788 to 1950 cells per
condition), using the cell length as a proxy for the cell division cycle, with the signal intensity profile sorted
from the shortest newborn cells to the longest predivisional cells. Note that the demographs for the
CJW6768 strain (panel A) in the nutrient conditions M9mann, M9mala, M9malt, M9mannCAAT,
M9glyCAAT and M9malaCAAT are not shown here, as they are presented in Figure 2A. The nutrient
abbreviations and compositions are explained in Table S1.

2360

Figure 2 – figure supplement 2: Correlation of the extent and relative timing of polysome
accumulation with nucleoid segregation at the single-cell level. A. Plot showing the variability in growth
rate (GR) across cell division cycles for cells growing in microfluidics in M9gluCAAT (4114 cell division
cycles, 9 growth rate bins with 93 to 860 cell division cycles each). **B.** Ensemble kymographs of the RplA-
GFP and HupA-mCherry concentration normalized by the average fluorescence for the slowest,
intermediate and fastest growing population bins shown in panel A. **C.** Plot showing the correlation between
the average RplA-GFP accumulation and HupA-mCherry depletion at mid-cell across the growth rate bins
(mean ± SD, 93 to 860 cell division cycles per bin) shown in panel A.

2369

Figure 2 – figure supplement 3: Calculation of RplA-msfGFP concentration after cell curvature
correction. A. Average 2D RplA-msfGFP projections for short (2.5 to 3 µm – left), unconstricted cells and
long (3.5 and 5.5 µm - right) constricted cells. The average RplA-msfGFP signal from 1000 sampled cells
(CJW7651) is shown for each population. **B.** The 2D cell areas were divided in cylindrical sectors with a
height (h) of a single pixel. The number of cylindrical sectors corresponds to the average cell length in pixels
(40 and 60 sectors for the unconstricted and constricted cells, respectively). The radius (r) of each
cylindrical segment corresponds to half the cell diameter for that specific cell length position. The cylindrical
segments at the poles or constriction site have a smaller radius due to the curvature of the cell boundaries.
The division of the cell area into cylindrical segments allowed us to calculate the volume for each segment
($V_{cylinder} = \pi\, r^2\, h$). The sum of the fluorescence per segment was corrected for the 3D curvature of the cell
boundaries by dividing by the volume of the cylinder, either considering the entire cell width or different
depths of view narrower than the maximal cell width. **C.** One-dimensional profiles of the cylinder diameter
are also indicative of the cell curvature at the poles and constriction site. **D.** Comparisons of the 1D average
RplA-msfGFP projection between the poles before (dashed lines) or after correcting for the cell curvature

(solid lines) for different depths of view (colormap). The uncorrected signal (dashed lines) corresponds to

the average fluorescence along each cylindrical segment (mean of pixel intensities). The 1D profiles were

normalized by dividing by the maximal RplA-msfGFP concentration at mid-cell. This correction shows that

the pronounced decrease of the RplA-msfGFP fluorescence at the poles is the result of the curvature of the

cell boundaries. The observed over-correction at the cell poles for larger depths of view (> 8 pixels or px)

is likely due to the over-estimation of the cell segmentation mask that was determined based on the phase

contrast snapshot images of cells on agarose pads.

2391

Figure 3 – figure supplement 1: Quantification of the polysome and nucleoid asymmetries using

fitted Gaussian functions. A. Example images of RplA-GFP and HupA-mCherry fluorescence signals in

a single newborn cell instance. **B.** Plots showing the Gaussian-function fitting on RplA-GFP and HupA-

mCherry intensity profiles from the single cell snapshot shown in panel A. Three Gaussian functions were

fitted to the RplA-GFP fluorescence in newborn cells (eq. 11 in Methods), capturing the accumulation of

polysomes at mid-cell and the poles. Two Gaussian functions were fitted to the HupA-mCherry fluorescence

(eq. 12 in methods) capturing the two lobes of the segregating sister nucleoids. The parameters of the fitted

Gaussian functions provided information about the position (μ: mean) and the concentration (A: amplitude)

of each fluorescence statistic, as well as the cell length range occupied by it (σ: standard deviation). The

Gaussian area (eq. 13 in Methods) corresponds to the abundance of each macromolecule. The RplA-GFP

Gaussians were fitted above the cellular background, which presumably corresponds to the uniform

fluorescence of free ribosomes or ribosomal subunits. For the HupA-mCherry Gaussian fittings, the cellular

background corresponds to the DNA-free cell regions. The Gaussian parameters were also used to

describe the polysome asymmetries between the poles (eq. 14 in Methods) as well as the position of the

nucleoid around the cell center (eq. 15 in Methods). These statistics were used in Figure 3B. **C.** Gaussian

functions were also fitted to the ensemble average RplA-GFP and HupA-mCherry fluorescence (right)

calculated from all the cell segmentation instances of a single cell division cycle, 0-10% from birth to division

(left). **D.** The Gaussian fitting to the ensemble RplA-GFP and HupA-mCherry intensity profiles

(concentration in arbitrary fluorescence units) early in the cell division cycle for the 2D projection shown in

panel C (right).

2412

Figure 3 – figure supplement 2: Correlations between the polysome distribution statistics and the

nucleoid compaction asymmetries. A. Plots showing the nucleoid and polysome density axial asymmetry

determined using the parameters of the fitted Gaussians on the HupA-mCherry and RplA-GFP fluorescence

(eq. 16 and eq. 18 in Methods). **B.** Plots showing the scaled correlations between the polysome Gaussian

parameters (μ, A, and σ for each polysome accumulation) and the nucleoid density asymmetry (eq. 16 in

Method; 2103 cell division cycles). Each marker corresponds to the data for a single cell division cycle (0-

10% into the cell division cycle) as in Figure 3 – figure supplement 1C-D. A Gaussian kernel density

estimation (KDE) was used to illustrate the density of the scatter plots (see Methods). **C.** Bar graph showing

the coefficients of a linear mixed-effects model (see eq. 17 in Methods) used to identify the polysome
statistics that contribute to the nucleoid density asymmetry. The highly contributing polysome statistics were
combined into a compound polysome statistic (eq. 18 in Methods) that describes the available DNA space
between polysome accumulations. This compound statistic correlates with the nucleoid density asymmetry
(eq. 16 in Methods) as shown in Figure 3F. The stars indicate the polysome statistics that most significantly
correlate with the nucleoid density asymmetry (absolute coefficient above 0.3).

2427

Figure 4 – figure supplement 1: Effects of chloramphenicol treatment on nucleoids in non-dividing
cells. A. Representative example of an elongated cephalexin-treated cell that does not fuse its nucleoids
upon chloramphenicol treatment. **B.** Representative example of an elongated cephalexin-treated cell that
fuses two of its nucleoids upon chloramphenicol treatment. **C.** Plot showing the distributions of the minimum
distance between adjacent nucleoids in cells where nucleoid fusion happens (n = 67), compared to cells
that do not display nucleoid fusion (n = 134). The two subpopulations were classified as described in the
Methods section.

2435

2436 **Figure 4 – figure supplement 2: Determination of simulated cell growth parameters based on**
2437 **experimental measurements.** Polysome and nucleoid dynamics were simulated for 10 different growth
2438 rates. The predivisional polysome and nucleoid profiles of simulated trajectories initialized from the steady
2439 state (Figure 4A) were used as initial conditions (D_n = 10^{-3} $\mu m^2/s$). The cell length at birth increased with
2440 growth rate according to published measurements (Govers et al., 2024).

2441

Figure 4 – figure supplement 3: Comparison of cell length at nucleoid splitting between simulations
and experimental data. Across different growth rates, the cell lengths at nucleoids splitting are compared
between our model simulations and the experimentally determined population-averaged statistics (Govers
et al., 2024). The range of the y-axis spans 2 μm (between 3 and 5 μm) to match the ~2 μm range of the
cell areas at birth across nutrient conditions, which spans between 2 and 4 μm in Figure 4 – figure
supplement 2.

2448

2449 **Figure 4 – figure supplement 4: Simulation of an extended model with three different polysome**
2450 **species.** Kymographs of simulated data of an extended version of the theoretical model that considers
2451 three polysome species with different diffusion coefficients $D_p = 0.018,\ 0.023, or\ 0.028\ \mu m^2/s$, representing
2452 polysomes with a decreasing number of ribosomes.

2453

2454

Figure 6 – figure supplement 1: Phenotypic effects of transcription inhibition. Plots showing cellular
parameters (instantaneous growth rate, cell area, nucleoid area, and the ratio between nucleoid area and
cell area, or NC ratio) affected by the rifampicin treatment. The instantaneous growth rate was estimated

using the log-transformed cell area, using a rolling window of 10 min. The black line corresponds to the
averages from 1859 cell division cycles and the gray-shaded area indicates the range of one standard
deviation around the mean. The two cycles of rifampicin treatment (as shown in Figure 5A) were overlaid
for this plot, using the time of rifampicin addition as t = 0 min.

2462

2463 **Figure 7 – figure supplement 1: DNA content during protein overexpression from plasmids.** Plots
2464 showing the scaling between the DNA content (DRAQ5 fluorescence area) and the cell size (side scatter
2465 area) quantified by flow cytometry, comparing uninduced and induced (100μM IPTG for 200 min) conditions
2466 for CJW7798 cells. Data from three biological data are shown.

2467

Figure 7 – figure supplement 2: Simulations of ectopic polysome formations away from the
nucleoid. Polysome formation was introduced at fixed nucleoid-free regions in the theoretical model to
mimic the ectopic polysome formation due to plasmid-driven expression in experiments (as in Figure 7).
The model recapitulates the two main phenotypes observed in experiments. The simulations were initialized
from the steady-state polysome and nucleoid distribution, similar to the steady state in Figure 4A. **A.**
Kymograph of simulated data for which polysomes are ectopically produced near the cell pole, at a rate
equivalent to 46% of total polysome production at cell birth. **B.** Kymograph of simulated data for which
polysomes are produced between the sister nucleoids, at a rate equivalent to 47% of total polysome
production at cell birth.

2477

2478 **Figure 7 – figure supplement 3: Phenotypic effects of prolonged protein overexpression from**
2479 **plasmids.** Representative phase contrast and fluorescence images of cells (CJW7798) after prolonged
2480 induction (100 μM IPTG, for 9 h and 15 min) of mTagBFP2 expression from aT7 promoter on a multi-copy
2481 plasmid.

2482

2483 **Figure 8 – figure supplement 1: Phenotypic of effects of cell growth under A22 and cephalexin**
2484 **treatment.** Representative phase contrast and fluorescence images of cells (CJW7323) treated with A22
2485 (4 μg/mL) alone or with both A22 (4 μg/mL) and cephalexin (50 μg/mL) for 130 min.

2486

2487

Movie legends
Movie S1: Example time-lapse sequence showing ribosome and nucleoid dynamics during cell
growth under steady state condition in a microfluidic channel. Shown are corresponding inverted
phase contrast signal, RplA-GFP signal, HupA-mCherry signal, cell contours (based on phase contrast
signals) and nucleoid contours (based on nucleoid signal segmentation) of *E. coli* cells (CJW7323) grown
in a microfluidic channel supplemented with M9gluCAAT. The white circles indicate the centroid of the cell
segmentation masks and the extending lines indicate the tracked cell traces.

2495

2496 **Movie S2: Video showing the average subcellular distribution of ribosome and nucleoid signals**
2497 **from birth to division.** Shown are 2D average RplA-GFP (top) and HupA-mCherry (bottom) projections
2498 from birth to division, with corresponding average intensity profiles (right). The cell projections are oriented
2499 from the old pole on the left to the new pole on the right. The average cell contour is also drawn. Ensemble
2500 data from 4122 division cycles of CJW7323 cells growing in M9gluCAAT are shown.

2501

2502 **Movie S3: Video showing simulated 1D profiles of polysome and nucleoid concentration during**
2503 **slow and fast cell growth.** Simulations during slow (left) and fast (right) growth are shown. The simulations
2504 were initialized from the equilibrium configuration, with a compact symmetric nucleoid ($D_n = 10^{-3}$ μm^2/sec)
2505 at the cell center.

2506

2507 **Movie S4: Video showing the effects of rifampicin addition on cell growth, ribosome signal**
2508 **heterogeneity, nucleoid segregation, and nucleoid compaction.** Examples of five microfluidic channels
2509 showing the corresponding inverted phase contrast, RplA-GFP and HupA-mCherry signals (from left to
2510 right) of cells (CJW7323) growing within microfluidic channels. Rifampicin was added at 120 and 720 min.
2511 Each antibiotic treatment lasted 120 min.

2512

Movie S5: Video showing the effects of ectopic polysome formation on nucleoid dynamics over
time. The RplA-GFP and HupA-mCherry fluorescence normalized (norm.) by the average cell fluorescence
are shown, together with their corresponding phase contrast image, for multiple cells (CJW7798) in
succession following induction of mTagBFP2 expression from a T7 promoter on a multi-copy plasmid. The
scale bar indicates 1 µm, and the time since cell birth is shown in minutes. The cell contours indicate the
boundaries of the cell masks obtained by cell segmentation of the corresponding phase contrast images.

2519

Movie S6 Video showing how the loss of cell width confinement due to cephalexin and A22
treatment affects ribosome and nucleoid distributions over time. Fluorescence images of RplA-GFP
and HupA-mCherry fluorescence (fluor.) normalized by the average cellular fluorescence images, together
with their corresponding phase contrast images, are shown for cells (CJW7323) following treatment with
A22 (4 µg/mL) and cephalexin (50 µm/mL). The scale bar indicates 1 µm. The time since drug addition is
shown in minutes.

2532 **Supplementary tables**

2533 **Table S1. Growth medium abbreviation and composition.**

Abbreviation	Composition
M9acet	1x M9 salts, 0.2% acetate (sodium salt)
M9acetCAAT	1x M9 salts, 0.2% acetate, 0.1% casamino acids, 1µg/mL thiamine
M9fuc	1x M9 salts, 0.2% fucose
M9fucCAAT	1x M9 salts, 0.2% fucose, 0.1% casamino acids, 1µg/mL thiamine
M9fum	1x M9 salts, 0.2% fumarate (disodium salt)
M9fumCAAT	1x M9 salts, 0.2% fumarate, 0.1% casamino acids, 1µg/mL thiamine
M9GlcNAc	1x M9 salts, 0.2% N-acetylglucosamine
M9GlcNAcCAAT	1x M9 salts, 0.2% N-acetylglucosamine, 0.1% casamino acids, 1µg/mL thiamine
M9glu	1x M9 salts, 0.2% glucose
M9gluCAAT	1x M9 salts, 0.2% glucose, 0.1% casamino acids, 1µg/mL thiamine
M9gly	1x M9 salts, 0.2% glycerol
M9glyCAAT	1x M9 salts, 0.2% glycerol, 0.1% casamino acids, 1µg/mL thiamine
M9Lala	1x M9 salts, 0.2% L-alanine
M9LalaCAAT	1x M9 salts, 0.2% L-alanine, 0.1% casamino acids, 1µg/mL thiamine
M9Lara	1x M9 salts, 0.2% L-arabinose
M9LaraCAAT	1x M9 salts, 0.2% L-arabinose, 0.1% casamino acids, 1µg/mL thiamine
M9mala	1x M9 salts, 0.2% malate (sodium salt)
M9malaCAAT	1x M9 salts, 0.2% malate, 0.1% casamino acids, 1µg/mL thiamine
M9malt	1x M9 salts, 0.2% maltose
M9maltCAAT	1x M9 salts, 0.2% maltose, 0.1% casamino acids, 1µg/mL thiamine
M9mann	1x M9 salts, 0.2% mannose
M9mannCAAT	1x M9 salts, 0.2% mannose, 0,1% casamino acids, 1µg/mL thiamine
M9pyr	1x M9 salts, 0.2% pyruvate (sodium salt)
M9pyrCAAT	1x M9 salts, 0.2% pyruvate, 0,1% casamino acids, 1µg/mL thiamine
M9sorb	1x M9 salts, 0.2% sorbitol
M9sorbCAAT	1x M9 salts, 0.2% sorbitol, 0,1% casamino acids, 1µg/mL thiamine
M9succ	1x M9 salts, 0.2% succinate (disodium salt)
M9succCAAT	1x M9 salts, 0.2% succinate, 0,1% casamino acids, 1µg/mL thiamine
M9treh	1x M9 salts, 0.2% trehalose
M9trehCAAT	1x M9 salts, 0.2% trehalose, 0,1% casamino acids, 1µg/mL thiamine
M9xyl	1x M9 salts, 0.2% xylose
M9xylCAAT	1x M9 salts, 0.2% xylose, 0,1% casamino acids, 1µg/mL thiamine

2534

2535
2536 **Table S2: *Escherichia coli* strains used in this study.**

Strain	Genotype	Source
MG1655	*E. coli* MG1655 (F-lambda- *ilvG- rfb-50 rph-1*)	(Guyer et al., 1981; Jensen, 1993)
MG1655 (DE3)	*E. coli* MG1655 (DE3)	Kind gift from Dr K. Prather (Massachusetts Institute of Technology), (Tseng et al., 2010b)
CJW5158	*E. coli* BW25113 *hupA::hupA-mcherry-frt-kanR-frt*	(Gray et al., 2019)
CJW6723	*E. coli* MG1655 Δ*lacZYA::P_lac-egfp-µNS hupA:hupA-mcherry*	(Gray et al., 2019)
CJW6768	*E. coli* MG1655 *rplA::rplA-meos2*	(Sanamrad et al., 2014)
CJW6769	*E. coli* MG1655 *rspB::rpsB-meos2*	(Sanamrad et al., 2014)
CJW7019	*E. coli* MG1655 *rplA::rplA-msfgfp-frt-kanR-frt*	(Gray et al., 2019)
CJW7020	*E. coli* MG1655 *rplA::rplA-msfgfp*	(Gray et al., 2019)
CJW7021	*E. coli* MG1655 *rpsB::rpsB-msfgfp*	(Gray et al., 2019)
CJW7144	*E. coli* MG1655 Δ*lacZYA::P_lac-mcherry-µNS-frt-kanR-frt*	This study
CJW7145	*E. coli* MG1655 Δ*lacZYA::P_lac-mcherry-µNS-frt-kanR-frt rplA::rplA-msfgfp*	This study
CJW7323	*E. coli* MG1655 *rplA::rplA-msfgfp hupA::hupA-mcherry*	(Xiang et al., 2021)
CJW7466	*E. coli* MG1655 (DE3) *hupA:hupA-mcherry*	This study
CJW7651	*E. coli* MG1655 Δ*lacZYA::P_lac-mcherry-µNS rplA::rplA-msfgfp*	This study
CJW7766	*E. coli* MG1655 (DE3) *hupA:hupA-mcherry rplA::rplA-msfgfp-frt-kanR-frt*	This study
CJW7798	*E. coli* MG1655 (DE3) *hupA:hupA-mcherry rplA::rplA-msfgfp-frt-kanR-frt | pET28:mTagBFP2-CmR*	This study

2537
2538

Table S3: Plasmids used in this study.

Plasmid name	Relevant genetic elements	Source
pKD13	*frt-kanR-frt-R6Kori-ampR*	(Datsenko and Wanner, 2000)
pKD46	*araC-bet-exo-A101(Ts)ori-ampR*	(Datsenko and Wanner, 2000)
pCP20	*cmR-A101(Ts)ori-ampR-flp-λ-repressor(Ts)*	(Datsenko and Wanner, 2000)
pER12	*pBAD322A-gfp-µNS*	Kind gift from Dr A. Janakiraman (City College of New York), (Broering et al., 2005)
pER12-mcherry	*pBAD322A-mcherry-µNS*	This study
pAPG1	*attB-pBAD322A-mcherry-µNS-attB-kanR-ColE1ori*	This study
pSB3C5-proA-B0032-E0051	*PproA-lacZα-GFP-p15Aori-cmR*	(Davis et al., 2011)
pET28:GFP	*PT7-lacO-GFP-kanR-pMB1ori-lacI*	(Shis and Bennett, 2013)
pBAD:TagBFP2	*araC-ParaBAD-mTagBFP2-ampR-pBAD322ori*	(Subach et al., 2011)
pET28:mTagBFP2	*PT7-lacO-mTagBFP2-kanR-pMB1ori-lacI*	This study
pET28:mTagBFP2-CmR	*PT7-lacO-mTagBFP2-cmR-pMB1ori-lacI*	This study

Table S4: DNA oligonucleotides used in this study.

Oligo name	Sequence	Source
ER12-MCR-fwd	5' CTCCATACCCGTTTTTTTGGGCTAGCAGGAGGAATTCATGGTGAGCAAGGGCGAGGAG 3'	This study
ER12-MCR-rev2	5' CACTGGAGGAGCCTGCTTTTTTGTACAAACTTGTTGACTTGTACAGCTCGTCCATG 3'	This study
µNSmCherry fwd	5' CACAGGTTGCTCCGGGCTATGAAATAGAAAAATGAATCCGTTGAAGCCTGATCGATGCATAATGTGCC 3'	This study
µNSmCherry rev	5' AGCTCCAGCCTACACAGAGTTTGTAGAAACGCAAAAAG 3'	This study
FRT_KanR fwd	5' GTTTCTACAAACTCTGTGTAGGCTGGAGCTGC 3'	This study
FRT_KanR Rev	5' TTAAAGGTATTAAAAACAACTTTTTTGTCTTTTTACCTTCCCGTTTCGCTCCTGTCAAACATGAGAATTAATTCC 3'	This study
ColE1 fwd	5' GAGCGAAACGGGAAGGTAAAAAGACAAAAAGTTGTTTTTAATACCTTTAACGTTCCACTGAGCGTC 3'	This study
ColE1 rev	5' CAGGCTTCAACGGATTCATTTTTTCTATTTCATAGCCCGGAGCAACCTGTGGGTTATCCACAGAATCAGG 3'	This study
pAPG1 seq1	5' GATCAAGCAGAGGCTGAAG 3'	This study
pAPG1 seq2	5' CCAGGCATCAAATTAAGC 3'	This study
pAPG1 seq3	5' TCTACGTGTTCCGCTTCC 3'	This study
pAPG1 seq4	5' TTGAAGCCTGATCGATGC 3'	This study
pAPG1 seq5	5' AAGATTAGCGGATCCTACC 3'	This study
lacZYA_redµNS fwd	5' TATGTTGTGTGGAATTGTGAGCGGATAACAATTTCACACAGGAAACAGCTATGGTGAGCAAGGGCGA 3'	This study
lacZYA_redµNS rev	5' CAATTTTTATAATTTAAACTGACGATTCAACTTTATAATCTTTGAAATAAGGATCCGTCGACCTGCAG 3'	This study
LacI fwd	5' GGCCGATTCATTAATGCAGCTGGC 3'	This study
CynX rev	5' GGCCTGATAAGCGCAGCGTATC 3'	This study
mCherry rev	5' GGTGCTTCACGTAGGCCTTGG 3'	This study
KanR fwd	5' CGGAGAACCTGCGTGCAATCC 3'	This study
mTagBFP2 fwd	5' CGGAGCTCGAATTCGGATCCTTAATTAAGCTTGTGCCCCAGTTTG 3'	This study
mTagBFP2 fwd	5' CTTTAAGAAGGAGATATACCATGAGCGAGCTGATTAAGGAGAAC 3'	This study
pET28_one fwd	5' TCCTTAATCAGCTCGCTCATGGTATATCTCCTTCTTAAAGTTAAAC 3'	This study
pET28_one rev	5' GGATAACCGTATTACCGCCTTTGAGTGAGCTGATACCG 3'	This study
pET28_two fwd	5' AGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCC 3'	This study
pET28_two rev	5' TGGGGCACAAGCTTAATTAAGGATCCGAATTCGAGCTCC 3'	This study
pET28mTagBFP2 fwd	5' CTTAGTGACTCGAATTCGCGCGCCAATCCGGATATAGTTCC 3'	This study
pET28mTagBFP2 rev	5' ACTTTCTGGCTGGATGATGGACGTGAGTTTTCGTTCCACTG 3'	This study
cmR fwd	5' AGTGGAACGAAAACTCACGTCCATCATCCAGCCAGAAAGTG 3'	This study
cmR rev	5' GAACTATATCCGGATTGGCGCGCGAATTCGAGTCACTAAGG 3'	This study

2545

Table S5: Chemicals used in this study.

Chemical	Source	Catalog number
Agarose	AmericanBio	Cat#AB00972-00500
4',6-diamidine-2'-phenylindole (DAPI), dihydrochloride fluorescent dye	Thermo Fisher Scientific	Cat#D1306
Isopropyl β-D-1-thiogalactopyranoside (IPTG)	Sigma Aldrich	Cat#I5502
Rifampicin	Sigma Aldrich	Cat#R3501
Cephalexin	Sigma Aldrich	Cat#C4895
A22	Sigma Aldrich	Cat#SML0471
Chloramphenicol	Sigma Aldrich	Cat#C0378
4',6-Diamidine-2-phenylindole dihydrochloride (DAPI)	Thermo Fisher Scientific	Cat#D1306
eBioscience™ DRAQ5™	Thermo Fisher Scientific	Cat#65-0880-92

2546

2547

2548

Table S6: Software used in this study.

Program / library / script	Package	Source
Oufti	www.oufti.org	(Paintdakhi et al., 2016)
MATLAB	www.mathworks.com	Mathworks
Python	www.python.org	Python Software Foundation
Numpy	www.numpy.org	(Harris et al., 2020)
Scipy	www.scipy.org	(Virtanen et al., 2020)
Pytorch	www.pytorch.org	(Paszke et al., 2019)
Scikit-image	www.scikit-image.org	(Van Der Walt et al., 2014)
Scikit-learn	www.scikit-learn.org/stable/	(Pedregosa et al., 2012)
Statsmodels	www.statsmodels.org/stable/index.html	(Seabold and Perktold, 2010)
Shapely	www.pypi.org/project/shapely/	(Gillies, Sean et al., 2023)
Matplotlib	www.matplotlib.org	(Hunter, 2007)
Seaborn	www.seaborn.pydata.org	(Waskom, 2021)
Pandas	www.pandas.pydata.org	(McKinney, 2010)
Python	Omnipose neural network	(Cutler et al., 2022)
MATLAB	SuperSegger	(Stylianidou et al., 2016)
Python	Unet neural network	(Wiktor et al., 2021; Zhou et al., 2020)
snapshots_analysis_UNET_ghv.py	Snapshot image analysis (from UNET masks) – custom Python class	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025)
snapshots_analysis_OUFTI_GrayGovers_ghv.py	Snapshot image analysis (from Oufti masks) – custom Python class	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025)
snapshots_analysis_functions.py	Extraction of fluorescence and morphology statistics from cell snapshots – custom Python functions	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025)
microfluidics_segmentation_ghv.py	Cell segmentation and tracking from time-lapse images in microfluidics – custom Python class	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025)
microfluidics_analysis_functions_ghv.py	Extraction of fluorescence and morphology statistics from time-lapse images	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025)

	in microfluidics – custom Python functions	
Omnipose_to_python_ghv.py	Extraction of segmentation masks and tracked lineages from Omnipose/SuperSegger to Python for further analysis	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025)
Otsu_phase_segmentation_ghv.py	Cell segmentation and tracking using Otsu thresholding (Otsu, 1979)	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025)
LoG_adaptive_image_filter.py	Image filter that applies relative and local thresholding to segment fluorescence objects and particles	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025)
Microfluidics_segmentation_and_tracking_example.pdf	Example for cell segmentation and tracking in microfluidics	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025)
Analysis_of_cell_morphology_and_fluorescence_in_microfluidics.pdf	Example for the extraction of fluorescence and morphology statistics from time-lapse images in microfluidics	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025)
Analysis_of_cell_morphology_and_fluorescence_in_agarose_pads.pdf	Example for the extraction of fluorescence and morphology statistics from cell snapshots	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025)
Example_of_using_Otsu_based_segmentation_ghv.pdf	Example for the segmentation of antibiotic-treated cells using the Otsu threshold	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2025)
tune_diffusion.ipynb	Simulations of the minimal reaction-diffusion model that describes nucleoid segregation in *E. coli*.	This study (https://github.com/qiweiyuu/polysome)
Time_lapse_on_agarose_pad	Python repository that includes the *fluorescence_analysis* class and its associated functions, used to track cell trajectories and analyze their fluorescence statistics in time-lapse images on agarose pads.	This study (https://github.com/alexSysBio/Time_lapse_on_agarose_pad)
flowio_to_pandas	Python repository that includes the *flow_cytometry_class* class and its associated functions, used to parse FCS files into Pandas. This class can also be used for polygon and histogram gating.	This study (https://github.com/alexSysBio/flowio_to_pandas)
nd2_to_array.py	Function used to	This study

	parse .nd2 files into numpy arrays.	(https://github.com/alexSysBio/Adding_ND2_images_to_python)
Bivariate_medial_axis_estimation.py	Functions used to draw the medial axis in segmented cell masks from snapshots or time-lapse images on agarose pads.	This study (https://github.com/alexSysBio/Cell_medial_axis_definitions)
Unever_background_correction.py	Functions used to subtract the background from agarose-pad images.	This study (https://github.com/alexSysBio/Image_background_subtraction)

2549

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 1 - figure supplement 1

Figure 1 - Figure supplement 2

Figure 1 - figure supplement 3

Figure 1 - figure supplement 4

Figure 1 - figure supplement 5

Figure 1 - figure supplement 6

Figure 2 - figure supplement 1

Figure 2 - figure supplement 2

Figure 2 - figure supplement 3

A **Single newborn cell**

old   *new*

1.5 µm

200 300 400 500

RplA-GFP conc. (a.u.)

100 200 300

HupA-mCherry conc. (a.u.)

C **0-10% into the cell division cycle**

Single cell fluorescence   **Average cell projection**

Average cell length: 2.9µm

1.5µm

RplA-GFP conc. (a.u.)
x10² 6 5 4 3 2

RplA-GFP conc. (a.u.)
x10² 6 5 4 3 2

HupA-mCherry conc. (a.u.)
x10² 3 2 1

new

old

old   *new*

HupA-mCherry conc. (a.u.)
x10² 3 2 1

B *Single cell intensity profile*

RplA-GFP concentration (a.u.)
400 350 300 250

$Poly_{old}$ (μ,σ,A)
$Poly_{mid}$ (μ,σ,A)
$Poly_{new}$ (μ,σ,A)

A
σ
µ

old pole   *new pole*

HupA-mCherry concentration (a.u.)
250 200 150 100 50

Nuc_{old} (μ,σ,A)
Nuc_{new} (μ,σ,A)

-0.5   0.5
Relative cell length

D *0-10% into the cell dvision cycle*

RplA-GFP concentration (a.u.)
520 480 440 400

$Poly_{old}$ (μ,σ,A)
$Poly_{mid}$ (μ,σ,A)
$Poly_{new}$ (μ,σ,A)

old pole   *new pole*

HupA-mCherry concentration (a.u.)
200 100

Nuc_{old} (μ,σ,A)
Nuc_{new} (μ,σ,A)

-0.5   0.0   0.5
Relative cell length

Figure 3 - figure supplement 1

Figure 3 - figure supplement 2

Figure 4 - Figure supplement 1

Figure 4 - figure supplement 2

Figure 4 - figure supplement 3

Figure 4 - figure supplement 4

Figure 6 - figure supplement 1

Figure 7 - figure supplement 1

Figure 7 - figure supplement 2

mTagBFP2 over-expression from a plasmid

Figure 7 - figure supplement 3

Phase contrast (a.u.)   **RplA-GFP (a.u.)**   **HupA-mCherry (a.u.)**

Figure 8 - figure supplement 1