DNA/polysome phase separation and cell width confinement couple nucleoid segregation to cell growth in *Escherichia coli*

Alexandros Papagiannakis[1,2,3], Qiwei Yu[5], Sander K. Govers[1,&], Wei-Hsiang Lin[1,2,#], Ned S. Wingreen[5,6], and Christine Jacobs-Wagner[1,2,3,4]

[1]Howard Hughes Medical Institute, Stanford University, Stanford, CA 94305, USA

[2]Sarafan Chemistry, Engineering, and Medicine for Human Health Institute, Stanford University, Stanford, CA 94305, USA

[3]Department of Biology, Stanford University, Stanford, CA 94305, USA

[4]Department of Microbiology and Immunology, School of Medicine, Stanford, CA 94305, USA

[5]Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton NJ 08544, USA

[6]Department of Molecular Biology, Princeton University, Princeton NJ 08544, USA

[&]Current address: Department of Biology, KU Leuven, Leuven, Belgium

[#]Current address: Institute of Molecular Biology, Academia Sinica, Taipei, Taiwan

*Corresponding author: email: jacobs-wagner@stanford.edu

ABSTRACT

Chromosome segregation is essential for cellular proliferation. Unlike eukaryotes, bacteria lack cytoskeleton-based machinery to segregate their chromosomal DNA (nucleoid). The bacterial ParABS system segregates the duplicated chromosomal regions near the origin of replication. However, this function does not explain how bacterial cells partition the rest (bulk) of the chromosomal material. Furthermore, some bacteria, including *Escherichia coli*, lack a ParABS system. Yet, *E. coli* faithfully segregates nucleoids across various growth rates. Here, we provide theoretical and experimental evidence that polysome production during chromosomal gene expression helps compact, split, segregate, and position nucleoids in *E. coli* through phase separation, inherently coupling these processes to biomass growth across nutritional conditions. Halting polysome formation immediately stops sister nucleoid migration while ensuing polysome depletion gradually reverses nucleoid segregation. Redirecting gene expression away from the chromosome and toward plasmids arrests nucleoid segregation and causes ectopic polysome accumulations that drive aberrant nucleoid dynamics. Cell width perturbations show that radial confinement of polysomes and nucleoids spatially controls their phase separation to ensure that nucleoids split along the cell width and segregate along the cell length. Our findings suggest a built-in mechanism for coupling chromosome segregation to cell growth and highlight the importance of cell width regulation in nucleoid segregation.

INTRODUCTION

All cells must segregate their replicated chromosomes to propagate genetic information to daughter cells at division. Remarkably, this essential process is far less understood in bacteria than eukaryotes. Eukaryotic cells use the mitotic spindle, a sophisticated cytoskeleton-based machine. No equivalent structure has been identified in bacteria, which fold their chromosomal material into a membrane-less organelle called the nucleoid. Many bacterial species use a ParABS system to segregate a specific DNA region proximal to the chromosomal origin of replication (*oriC*) (Figge et al., 2003; Ireton et al., 1994; Jalal and Le, 2020; Lim et al., 2014; Livny et al., 2007; Mohl and Gober, 1997). However, this initial step does not explain how the rest of the replicated megabase-long chromosomes partition. Furthermore, molecular dynamics simulations suggest that the segregation of the duplicated *oriC* regions is dispensable for the segregation of the entire chromosome (Harju et al., 2024). Consistent with this idea, the chromosomally encoded *parA* and/or *parB* gene are often not essential for cell viability (Bartosik et al., 2009; Charaka and Misra, 2012; Donczew et al., 2016; Donovan et al., 2010; Du et al., 2016; Ireton et al., 1994; Jakimowicz et al., 2007a, 2007b; Jecz et al., 2015; Kadoya et al., 2011; Kawalek et al., 2020; Kim et al., 2000; Lagage et al., 2016; Lee and Grossman, 2006; Lewis et al., 2002; Li, 2019; Li et al., 2015; Minnen et al., 2011; Santi and McKinney, 2015; Takacs et al., 2022; Yamaichi et al., 2007; Yu et al., 2010). In fact, some bacteria do not even encode a ParABS system for *oriC* partitioning (Livny et al., 2007), with *Escherichia coli* being a prime example. Yet, *E. coli* segregates its duplicated nucleoids faithfully. *E. coli* mutants in which the *mukB* gene is deleted exhibit moderate nucleoid segregation defects linked to the organization and condensation of the nucleoid (Danilova et al., 2007; Mäkelä et al., 2021; Sawitzke and Austin, 2000). However, even in these mutants, most cells successfully segregate and partition their nucleoids between the daughter cells. Furthermore, topoisomerase I mutations or *seqA* deletion suppress the chromosome segregation defects associated with the *mukB* deletion (Danilova et al., 2007; Sawitzke and Austin, 2000; Weitao et al., 1999), suggesting that at least one other mechanism is at play.

Nucleoid segregation must somehow be coordinated with cell growth to accommodate the various growth rates that *E. coli* can display depending on the environmental conditions. In the presence of a poor-quality carbon source, *E coli* initiates nucleoid segregation late in the cell division cycle. As the quality of the carbon source improves and the growth rate increases, nucleoid segregation occurs at an increasingly earlier stage of the cell division cycle until it initiates in the preceding cycle to accommodate faster division times (Bates and Kleckner, 2005; Govers et al., 2024; Tiruvadi-Krishnan et al., 2022). Over the years, multiple mechanisms have been proposed to explain nucleoid segregation in *E. coli*. In 1963, Jacob et al. hypothesized that sister chromosomes migrate from mid-cell to quarter-cell positions through membrane attachment of the sister chromosomes and zonal growth of the cell envelope between the attachment points (Jacob et al., 1963). This early model was largely abandoned following the discovery that *E. coli* cells elongate via a dispersed mode of cell wall growth along the cell length up to the final stage of the division cycle (Cooper and Hsieh, 1988; Navarro et al., 2022; Wientjes and Nanninga, 1989; Woldringh et al., 1987).

Moreover, the segregation of chromosomal loci is faster than the rate of cell elongation (Kuwada et al.,
2013). DNA replication has also been proposed to facilitate chromosome partitioning though a DNA
polymerase-dependent extrusion mechanism (Lemon and Grossman, 2000). Simulations of coarse-grained
polymer models have also suggested that the conformational entropy of confined circular nucleoid chains
could provide an unmixing force that separates two mixed DNA polymers (Jun and Mulder, 2006; Jun and
Wright, 2010; Pande et al., 2023). While these proposed mechanisms likely contribute to the nucleoid
splitting into two objects, they cannot explain how separated sister nucleoids move away from each other.
Additionally, the DNA-demixing models consider the DNA polymers in an empty cylinder, ignoring any
potential effects from crowders. Furthermore, recent modeling work has argued that entropic forces
generated from excluded volume interactions between partially replicated chromosomes inhibit (rather than
promote) their segregation (Harju et al., 2023). This argues for the existence of one or more additional
segregation mechanisms, which could involve mechanical stress between overlapping sister nucleoids
(Fisher et al., 2013), DNA loop extrusion (Harju et al., 2023), or excluded volume effects between the DNA
meshwork and cytoplasmic crowders (Miangolarra et al., 2021; Wu et al., 2019b). While it is likely that
multiple factors contribute to nucleoid segregation and positioning inside the *E. coli* cell, the dominant
mechanism underlying the diffusional bias must be tightly coupled to cellular metabolism to explain how the
timing of nucleoid segregation changes with growth rate. How this is achieved remains unknown.

In contrast to eukaryotes, bacteria do not insulate their DNA in membrane-bound compartments. As a
result, the chromosomal meshwork can freely interact with the macromolecules present in the cytoplasm,
which exhibits a high level of polydispersity and crowding (McGuffee and Elcock, 2010; Zimmerman and
Trach, 1991). Experiments and modeling have suggested that macromolecular crowders exert compaction
forces on the compressible nucleoids through steric (i.e., excluded-volume) interactions (Pelletier et al.,
2012; Wu et al., 2019b; Yang et al., 2020; Zhang et al., 2009). In this context, polysomes, which consist of
mRNAs loaded with translating ribosomes, constitute a sizeable and abundant cytoplasmic crowder
(McGuffee and Elcock, 2010). The proteome fraction allocated to ribosomes increases with growth rate
across nutrient conditions (Chure and Cremer, 2023; Dai et al., 2017; Dourado and Lercher, 2020; Hu et
al., 2020; Molenaar et al., 2009; Scott et al., 2014; Si et al., 2017), reaching up to 20% of the cell mass
during fast growth (Chure and Cremer, 2023). The proportion of ribosomes engaged in translation also
increases with improving nutrient quality, attaining 70-80% under fast growth (Dai et al., 2017;
Forchhammer and Lindahl, 1971; Mohapatra and Weisshaar, 2018; Sanamrad et al., 2014; Varricchio and
Monier, 1971). These polysomes form structures (Brandt et al., 2009) of comparable or larger size than the
average ~50 nm mesh size of the nucleoid (Xiang et al., 2021). Consistent with a steric clash between
polysomes and the nucleoid meshwork, fluorescently-labeled ribosomes accumulate in DNA-free regions
such as the cell poles and between segregated nucleoids (Azam et al., 2000; Bakshi et al., 2015, 2012;
Chai et al., 2014; Gray et al., 2019; Lewis et al., 2000; Mohapatra and Weisshaar, 2018; Robinow and
Kellenberger, 1994; Sanamrad et al., 2014; Xiang et al., 2021).

Interestingly, theoretical work suggests that excluded-volume effects alone may be sufficient for the DNA to spontaneously phase separate from polysomes and compact into its observed nucleoid form (Mondal et al., 2011). Furthermore, a recent nonequilibrium statistical physics model proposes that nucleoid positioning and segregation could potentially be explained by considering the steric interaction (i.e., repulsion) between polysomes and DNA and the nonequilibrium effects associated with mRNA synthesis and degradation (Miangolarra et al., 2021). In this theoretical model, polysome accumulation in the middle of the nucleoid emerges due to polysomes born in the middle of the nucleoid taking longer to diffuse out of the nucleoid than those born closer to the nucleoid edges. Polysome accumulation beyond a certain concentration threshold drives phase separation between the DNA and polysomes at the mid-nucleoid location, resulting in nucleoid splitting. However, we lack experimental validation of this model. Furthermore, the coupling between nucleoid segregation and growth rate has not been addressed either experimentally or theoretically.

Here, we quantitatively characterize the temporal and spatial dynamics of nucleoids and polysomes in *E. coli* under different conditions and perturbations. In this work, we refer to nucleoid segregation as a series of events observable by microscopy (Figure 1A): (i) the initiation of nucleoid splitting, marked by the depletion of a DNA marker near the mid position of the nucleoid, (ii) the end of nucleoid splitting, which results in the generation of two separable nucleoid objects, and (iii) the migration of the sister nucleoids away from each other, marked by the increasing distance between the centroid of the nucleoids. This sequence of events can initiate in the preceding cell division cycle under nutrient-rich (fast-growth) conditions. Our experimental findings, combined with modeling, provide evidence that out-of-equilibrium and asymmetric polysome rearrangements in the cell result in both DNA compaction and the choreography of the nucleoid segregation cycle during growth across cells and nutrient conditions. The dual involvement of polysomes in protein synthesis and nucleoid dynamics ensures self-regulating coordination between cell growth and chromosome segregation across a wide range of growth rates, even under conditions where *E. coli* divides faster than it can replicate its chromosome. We also show that cell width plays a critical role in the directional patterning of nucleoid segregation by spatially regulating the phase separation between polysomes and chromosomes.

RESULTS

The rate of polysome accumulation correlates with the rate of nucleoid segregation across cells.

While the negative correlation between ribosomes and DNA has been well established from snapshot images (Azam et al., 2000; Bakshi et al., 2015, 2012; Chai et al., 2014; Gray et al., 2019; Lewis et al., 2000; Mohapatra and Weisshaar, 2018; Robinow and Kellenberger, 1994; Sanamrad et al., 2014; Xiang et al., 2021), the relative dynamics of these two macromolecules have not been quantified. To examine these dynamics with high temporal resolution (every 1 or 3 min, Figure 1 – figure supplement 1A-E), we ran time-

lapse microfluidic experiments using an *E. coli* strain carrying the 50S ribosomal protein RplA fused to GFP
and the nucleoid-associated protein HupA tagged with mCherry (Xiang et al., 2021). Cells were grown at
30°C in M9 buffer supplemented with glucose, casamino acids and thiamine (M9gluCAAT). While the
ribosome signal displayed the expected polar and inter-nucleoid enrichments, the accrual of ribosome
signal in the middle of the cells was particularly strong compared to the polar regions, as shown by
representative individual cells (Figure 1B and Movie S1) and average kymographs (Figure 1C, n = 4122
cell division cycles). Cell constriction contributed to the apparent depletion of ribosomal signal from the mid-
cell region at the end of the cell division cycle (Figure 1B-C and Movie S1). In predivisional cells, the
ribosomal signal accumulated in the middle of the segregated nucleoids near the ¼ and ¾ cell positions,
as shown in average one-dimensional (1D) and two-dimensional (2D) fluorescence profiles (n = 1907
predivisional cells, Figure 1D). As expected (Bakshi et al., 2012; Sanamrad et al., 2014), the RplA-GFP
signal became homogeneous within cells following their treatment with rifampicin (Figure 1 – figure
supplement 1F), a transcription inhibitor that causes polysome depletion over time (Blundell and Wild, 1971;
Campbell et al., 2001; Hartmann et al., 1967). This indicates that all ribosomal enrichments consist of
polysomes and therefore will be referred to as polysome accumulations hereafter.
163
Under our relatively nutrient-rich growth conditions (M9gluCAAT), nucleoid splitting occurred early in the
division cycle and even sometimes initiated in the segregated nucleoids from the preceding division cycle
(Movie S1 and Figure 1B). Given the variability in nucleoid segregation timing across cells (Figure 1 – figure
supplement 2A), we developed a computational method to track nucleoid dynamics independently of the
cell division cycle (see Methods and Figure 1 – figure supplement 2B-C). Specifically, we focused on the
nucleoid cycle—defined as the period between the ends of two nucleoid splitting events (Figure 1 – figure
supplement 2D)—instead of the division cycle. By tracking the accumulation of RplA-GFP and the depletion
of HupA-mCherry in the middle of nucleoids, we found that polysome accumulation and nucleoid splitting
correlated in time (Figure 1E). Furthermore, the rate of polysome accumulation at mid-nucleoid ($\Delta RplA_{mid\text{-}nuc}/\Delta T$)
correlated with the rate of DNA depletion in the same region ($\Delta HupA_{mid\text{-}cell}/\Delta T$) at the single-cell level
(Figure 1F). This indicates that cells that accumulated polysomes faster also split their nucleoids faster.
Importantly, the fitted linear regression had an intercept close to zero for both axes (Figure 1F), indicating
that when the rate of polysome accumulation approached zero, so did the rate of nucleoid splitting.
177
To examine what happens when sister nucleoids move away from each other, we divided the time between
the end of nucleoid splitting and cell division into four bins. We found that right after nucleoid splitting (0-
25% bin), most (~70%) cells continued to accumulate polysomes at mid-cell, i.e., between the sister
nucleoids (Figure 1G). For these cells, the rate of polysome accumulation ($\Delta RplA_{mid\text{-}cell}/\Delta T$) positively
correlated with the rate of nucleoid migration ($\Delta Distance_{nuc}/\Delta T$) (Figure 1H). Thus, the faster that cells
accumulated polysomes at mid-cell, the faster the sister nucleoids migrated apart (and vice versa).
184

Cell elongation may also contribute to sister nucleoid migration near the end of the division cycle.

We noted that there was a substantial percentage (~30%) of cells with decreasing RplA-GFP signal at mid-cell right after nucleoid splitting ($\Delta RplA_{mid-cell}/\Delta T \leq 0$, Figure 1H). Interestingly, in these cells, nucleoid migration did not stop; instead, its average rate was similar to the average rate of cell elongation ($\Delta Length_{cell}/\Delta T$) (Figure 1H). In fact, when the cell elongation rate was subtracted from the nucleoid migration rate for each single cell (Figure 1- figure supplement 3A), the positive correlation between polysome accumulation and nucleoid migration rates remained, but now the average rate of nucleoid migration was near zero in cells with no polysome accumulation ($\Delta RplA_{mid-cell}/\Delta T \leq 0$). This finding may suggest a mixed contribution between polysome accumulation and cell elongation to nucleoid migration. This was interesting considering that polysomes became less enriched at mid-cell but more enriched in the middle of sister nucleoids in predivisional cells (Figure 1B-D), with the average ribosome concentration remaining constant during the cell division cycle (Figure 1 – figure supplement 1E). The spatial change in polysome enrichments was accompanied with a steady decline in the percentage of cells that continued to accumulate polysomes at mid-cell between the end of nucleoid splitting and cell division (Figure 1G). Concurrent with this decline, the migration rate of sister nucleoids became less correlated with the rate of polysome accumulation and more correlated with the rate of cell elongation (Figure 1 – figure supplement 3B). To examine the relative correlation of polysome accumulation and cell elongation with nucleoid migration over time, we used a linear mixed-effects model to analyze each relative nucleoid migration interval (see Methods). The coefficients of the fitted mixed linear regressions suggest the following hypothesis: Early during nucleoid migration (0-25% between the completion of nucleoid splitting and cell division), polysome accumulation contributes most to the measured variance in the displacement of the sister nucleoids (Figure 1I). This contribution progressively decreases over time while that of cell elongation increases (Figure 1I). Such a switch in relative contribution would be consistent with the spatiotemporal dynamics of polysome accumulation and cell wall synthesis, as described in the Discussion.

Polysome accumulation at mid-cell correlates with the relative timing of nucleoid segregation across nutrient conditions and growth rates.

If polysome production plays a role in nucleoid segregation, it predicts that the timing and amount of polysome accumulation at mid-cell will correlate with the timing of nucleoid segregation across nutrient conditions. To test this prediction, we analyzed images of fluorescently labeled ribosomes and nucleoids in cells grown under 30 different carbon source conditions (Table S1) that vary the doubling times (~40 min to ~4 h) and average cell areas (~1.9 to ~3 μm^2) of *E. coli*. This dataset included both previously published (Gray et al., 2019) and new microscopy snapshots from our laboratory. Demographs generated from these images revealed that the polysome accumulation at mid-cell was reproducible across all conditions and strains, irrespective of the ribosomal subunit protein (RplA or RpsB) or the fluorescent tag (msfGFP, mEos2 or GFP) used to mark ribosomes (Figure 2A and Figure 2 – figure supplement 1). In these experiments,

nucleoids were typically imaged using DAPI (rather than a fluorescent fusion to HupA), indicating that polysome accumulation at mid-nucleoid was independent of the DNA labeling method. Importantly, and consistent with our prediction, the richer the growth conditions (i.e., the larger the average cell area), the earlier the polysome accumulation and the nucleoid splitting occurred in the division cycle based on relative cell lengths (Figure 2A and Figure 2 – figure supplement 1). In addition, the polysome accumulation at mid-cell was more pronounced in nutrient-rich media (e.g., M9malaCAAT) compared to nutrient-poor ones (e.g., M9mann) where the nucleoid segregated later in the division cycle (Figure 2A and Figure 2 – figure supplement 1).

To quantify these phenotypes across strains and nutrient conditions, we extracted and correlated population-level polysome and nucleoid statistics. We found a strong correlation (Spearman correlation ρ = -0.85) between the amplitude of the average ribosomal signal accumulation and the average nucleoid signal depletion at mid-cell (Figure 2B). Since the average cell area (colormap in Figure 2B) correlated with the growth rate of the population (Schaechter et al., 1958), this plot also confirmed that faster growing populations displayed stronger polysome accumulation and greater DNA depletion at mid-cell on average (Figure 2B).

We wondered whether the correlation between polysome and nucleoid dynamics could also be seen across cells growing in the exact same nutritional condition but at different growth rates due to cell-to-cell variability. To examine this possibility, we divided our timelapse microfluidics data into growth rate bins (Figure 2C) and compared their temporal polysome and nucleoid patterns. We found that, compared to the slowest growing subpopulation (bin 1 with a mean growth rate = 0.28 h^{-1}), the fastest growing subpopulation (bin 13 with = 0.74 h^{-1}) displayed a more pronounced and earlier accumulation of polysomes that was associated with earlier nucleoid segregation (Figure 2D). The subpopulation with an intermediate growth rate (bin 7 with = 0.51 h^{-1}) also displayed an intermediate pattern of polysome accumulation and nucleoid segregation. In fact, when we calculated the average polysome accumulation and nucleoid depletion at mid-cell for each growth rate bin, we found that the two statistics correlated (Figure 2E). Faster growing cells accumulated more polysomes at mid-cell and split their nucleoids earlier during the cell division cycle compared to slower growing cells.

Both the average proteome fraction dedicated to ribosomes and the average fraction of ribosomes engaged in translation are known to correlate with growth rate across nutrient conditions (Chure and Cremer, 2023; Dai et al., 2017; Dourado and Lercher, 2020; Hu et al., 2020; Molenaar et al., 2009; Scott et al., 2014; Si et al., 2017). Thus, a role for polysome production in nucleoid segregation may provide a mechanistic link between growth rate and the relative timing of nucleoid splitting both across cells in a given nutrient condition as well as across populations in different nutrient conditions.

Spatial polysome asymmetry correlates with nucleoid positioning.
A surprising result was the apparent higher polysome accumulation at mid-cell relative to the cell pole
regions (Figures 1B-C and 2A), which was not artificially created by the smaller cytoplasmic volumes at the
cell poles or constriction sites due to membrane curvature (Figure 2 – figure supplement 2). This uneven
distribution of polysomes suggested limited diffusion-driven equilibration of polysome concentration
between the DNA-free regions. At division, such a disequilibrium could lead to a higher concentration of
polysomes at the new cell pole relative to the old pole in daughter cells through inheritance. To examine
this possibility, we turned back to the microfluidic experiments in which we traced cell lineages from mother
to daughter cells (Figure 1 – figure supplement 1A), determined the pole identity (new vs. old) of each
tracked cell (Figure 1 – figure supplement 2B), and compared the polysome accumulations between the
new and old poles (see Methods and Figure 3 – figure supplement 1A-D). Old mother cells (located at the
end of the microfluidic channels) and their daughters were excluded from our analysis to avoid cell aging
effects (Chao et al., 2023; Coquel et al., 2013; Koleva and Hellweger, 2015; Łapińska et al., 2019; Lindner
et al., 2008; Proenca et al., 2019). Consistent with the absence of rapid equilibration of polysome
concentration between DNA-free regions through diffusion, we found that newborn cells had more
polysomes at the new pole compared to the old one on average (Figure 3A).
275
Across cells, the polysome distribution asymmetry between cell poles negatively correlated with the position
of the nucleoid in newborn daughter cells (Figure 3B, cells always oriented with the new pole to the right).
The *x* and *y* intercepts of a fitted linear regression to the data were not zero, indicating that a nucleoid
positioned precisely at the cell center did not equate with an even distribution of polysomes between poles.
Instead, newborn cells with centrally located nucleoids tended to have more polysomes at the new pole
compared to the old pole (example #1 in Figure 3B-C) whereas cells born with symmetric polysome
enrichments between poles tended to display an off-center nucleoid closer to the new pole (example #2 in
Figure 3B-C).
284
Spatial polysome asymmetry correlates with asymmetric nucleoid compaction in newborn cells.
Construction of average 2D cell projections of the RplA-GFP signal in cells sorted based on their relative
timing to cell division confirmed the polysome asymmetry between poles in newborn cells (Figure 3D and
Movie S2). Strikingly, the corresponding 2D projections of the HupA-mCherry signal revealed another
spatial asymmetry, this time, in the average DNA mass distribution along the nucleoid length (axial
asymmetry). The average HupA-mCherry signal concentration was higher toward the new pole right after
birth or, correspondingly, toward the middle of the cell (future new pole) in the period prior to division (Figure
3D and Movie S2). Quantification of this axial asymmetry in newborn cells revealed that the HupA-mCherry
concentration is ~20% higher toward the new pole on average (Figure 3E), suggesting that the DNA density
is uneven along the nucleoid length. This nucleoid mass asymmetry emerged late in the nucleoid migration
cycle, typically before cell division such that it was inherited by newborn cells (Figure 3D).

To examine if features of polysome accumulations (e.g., position, amplitude, or fraction of cell length
covered) correlate with the asymmetric nucleoid density and its variability among cells, we used a linear
mixed-effects model (see Methods). We found that in newborn cells, the positions of polysome enrichments
in the old-pole and mid-cell regions significantly correlated with the HupA-mCherry density asymmetry
(Figure 3 – figure supplement 1E-G). Guided by this finding, we hypothesized that the positions of the
accumulating polysomes along the cell length determine the space available for the chromosome to occupy,
and thereby dictate the DNA density distribution along the nucleoid. To examine this hypothesis, we
combined the correlated polysome accumulation characteristics into a compound statistic that describes
the relative polysome-free space between two cell halves (see Methods). Compared to other polysome
statistics, the relative availability of cell space depleted of polysomes (RplA-GFP signal) correlated most
strongly with the asymmetric distribution of DNA (HupA-mCherry signal) in newborn cells (Figure 3F). In
other words, the DNA concentration was higher in cell regions with more space available between polysome
enrichments. In most (~60%) cells, the distance between polysome enrichments, and thus the DNA
concentration along the nucleoid, was greater between mid-cell and the new pole (i.e., *x* and *y* values >1,
example #1 in Figure 3F-G). The opposite pattern was true for a small fraction (~14%) of cells, where the
larger distance between polysome enrichments was located between mid-cell and the old pole (*x* and *y*
values < 1, example #2 in Figure 3F-G). The correlated variability in polysome and nucleoid patterning
across cells suggests that the size of the polysome-depleted spaces helps determine where the
chromosomal DNA is most concentrated along the cell length. This patterning is likely reinforced through
the displacement of the polysomes away from the DNA dense region.

A minimal reaction-diffusion model generates experimentally observed cellular asymmetries and
growth rate-dependent nucleoid segregation.
Our single-cell studies provided experimental support that phase separation between polysomes and DNA
contributes to nucleoid segregation. However, it remained unclear whether the same mechanism could also
explain the growth rate-dependent trends and cellular asymmetries that we observed (Figures 2 and 3).
Therefore, we built a minimal reaction-diffusion model (see Methods) to describe the dynamics of
polysomes and DNA during the cell cycle based on previous findings (Miangolarra et al., 2021). The model
takes into account two ingredients important for nucleoid segregation: steric effects (repulsion) between
DNA and polysomes (described by Cahn-Hilliard theory) and the nonequilibrium processes of polysome
synthesis and degradation (described by linear reaction kinetics). The model is based on realistic
parameters of polysome diffusion, production, and degradation (see Methods). We assume polysome
production (i.e., transcription) to be uniform within the nucleoid and polysome degradation (i.e., mRNA
decay) to be uniform across the entire cell (from pole to pole). These assumptions consider the most trivial
cases (see Discussion for other cases). We also assume that the cell grows exponentially and that the
nucleoid expands in size proportionally to the cell during growth, which has been experimentally verified

(Campos et al., 2014; Govers et al., 2024). Since *E. coli* grows along its long axis and polysomes do not

readily go around the nucleoid to equilibrate (Figure 3A-D and Figure 2 – figure supplement 2), we reduced

the problem to one dimension, the cell length. Importantly, the system is driven out of equilibrium by the

continuous production and degradation of polysomes. It operates at a nonequilibrium steady state even at

fixed cell length.

Initialized from the steady state of a single compacted nucleoid in the middle of the cell with symmetric

polysome distributions at the poles, our model shows that polysomes accumulate in the middle of the

nucleoid during growth (Figure 4A and Movie S3, left panel). This is followed by division of the nucleoid into

two entities, which then move apart from each other as polysomes accumulated between them. The model

provides a minimal mechanism for nucleoid segregation: At any given point, polysomes that form in the

middle of the nucleoid have a lower probability of escaping the nucleoid through diffusion compared to

polysomes born at the edge of the nucleoid (Figure 4B). Thus, the polysome concentration rises

monotonically towards the center of the nucleoid, its peak increasing with nucleoid length (quadratically in

the quasi-steady-state limit). Nucleoid elongation during cell growth promotes polysome accumulation in

the middle of the nucleoid. Phase separation driven by the steric repulsion between DNA and polysomes

results in the nucleoid splits in two (Figure 4A and Movie S3, left panel).

We wondered if this simple reaction-diffusion mechanism could also explain the correlation between growth

rate and the relative timing of nucleoid segregation. Therefore, we performed simulations for different

growth rates, matching the cell and nucleoid length at birth with population-level measurements (Figure 4

– figure supplement 1A). To initialize each simulation in a realistic fashion, we used the last time point (i.e.,

half of the predivisional cell) of the previous simulation as initial conditions, capturing the new/old pole

identity as well as any cellular asymmetries inherited between generations. The relative timing of nucleoid

splitting was measured as nucleoid depletion at mid-cell beyond a threshold of 0.1 arbitrary units (Figure

4A and Figure 4 – figure supplement 1B). We found that our model successfully captures the negative trend

between the growth rate and the relative timing of nucleoid splitting. Simulated cells that grew faster also

split their nucleoids earlier from birth to division (Figure 4C and Movie S3), in agreement with population-

level data (Figure 2 – figure supplement 1).

To examine a potential origin for the asymmetries in polysome distribution and nucleoid compaction that

we observed (Figure 3), we examined the effect of the nucleoid diffusion coefficient D_n, which is a model

parameter describing how fast the nucleoid relaxes towards its equilibrium configuration. Large D_n

represents the quasi-steady-state limit where the nucleoid relaxation time scale is much shorter than the

cell doubling time, and the nucleoid always assumes its (symmetric) equilibrium state. Conversely, small

D_n (i.e., slower relaxation time) can lead to asymmetric concentration profiles that will be inherited by the

daughter cells. Consistent with this expectation, we found that during relatively fast growth (~40 min

doubling time), a lower nucleoid diffusion coefficient results in a larger deviation from the equilibrium
concentration profiles for the nucleoid and polysomes (dotted curves vs. solid curves, Figure 4D). In fact,
at diffusion coefficients below 0.001 μm^2/s, the model (Figure 4D) reproduced the experimentally observed
asymmetries, including the nucleoid position offset towards the new pole at birth, the higher polysome
concentration at the new pole compared to the old one, and the asymmetric nucleoid compaction (Figure
3). Our minimal model thus suggests that the material properties of the nucleoid (e.g., stiffness) may
contribute to the observed nucleoid and polysome asymmetries in *E. coli* (see Discussion).

Polysomes accumulate at mid-nucleoid in DNA regions inaccessible to freely diffusing particles of
similar sizes.
In the model, the early polysome accumulation in the middle of nucleoid is caused by the nonequilibrium
processes of polysomes being born within the nucleoid while being degraded uniformly across the cell (due
to mRNA turnover). It predicts that the early mid-nucleoid enrichment of polysome signal observed in our
experiments is the product of such nonequilibrium processes rather than of polysomes simply diffusing into
undetected DNA-free space. If this is correct, freely diffusing objects of similar sizes to polysomes should
accumulate at mid-cell after polysomes accumulate there, i.e., after free DNA-space has been generated
through nucleoid splitting. To test this expectation, we compared the average distribution of RplA-msfGFP
with that of freely diffusing mCherry-labeled μNS particles from snapshot images of DAPI-stained cells
(Figure 4E). These μNS particles consist of a fragment of a mammalian reovirus protein (Broering et al.,
2005, 2002) that self-assembles into a particle, typically one per cell, when produced orthogonally in *E. coli*
(Parry et al., 2014). They have sizes between 50 and 200 nm (Parry et al., 2014; Xiang et al., 2021) similar
to polysomes (Brandt et al., 2009; Slayter et al., 1968) and are therefore largely excluded by the nucleoid
mesh (Xiang et al., 2021). After sorting cells by length into four bins, the positions of mCherry-labeled μNS
particles from approximately 2580 cells per bin were superimposed using the relative cellular coordinates
to construct particle density maps. Cells were randomly oriented in this analysis (meaning that asymmetries
between poles cannot be observed), as the pole identity cannot be assigned from snapshot images.

We found that in the shortest (i.e., newborn) cells, RplA-msfGFP-labeled polysomes had already
accumulated within the nucleoid (Figure 4F, bin 1). In contrast, mCherry-μNS particles were restricted to
the cell poles and were not able to access the mid-cell region until after nucleoid splitting was clearly visible
(Figure 4F, bins 2-4). This was also shown in the corresponding 1D average concentration and particle
density profiles (Figure 4G). These observations support the notion that the early mid-nucleoid
accumulation of RplA-msfGFP is caused by nonequilibrium effects associated with polysome synthesis and
degradation rather than polysome diffusion into DNA-free space.

Arrest of polysome production immediately stops nucleoid segregation while polysome depletion gradually reverses it.

Our correlative analyses and model (Figures 1-4) support the hypothesis that the interactions and ensuing phase separation between polysomes and nucleoid promote nucleoid segregation and macromolecular asymmetries along the cell length. To probe causality, we first aimed to disrupt the proposed mechanism using rifampicin. Rifampicin treatment is known to homogenize ribosome distribution and expand the nucleoid over time through polysome depletion (Bakshi et al., 2014; Dworsky and Schaechter, 1973; Koch and Gross, 1979; Pettijohn and Hecht, 1974; Xiang et al., 2021). However, our hypothesis predicts a faster effect on nucleoid segregation. Blocking transcription should instantly reduce the rate of polysome production to zero, causing an immediate arrest of nucleoid segregation. Then, gradual depletion of the existing polysomes due to mRNA decay should cause, on a slower time scale, a dissipation of the phase separation.

To test these predictions, we subjected cells growing in M9gluCAAT in microfluidic channels to two rounds of rifampicin treatment (Figure 5A and Movie S4). Rifampicin resulted in growth rate inhibition (Figure 5A) and changes in the nucleoid area and nucleoid-to-cell area (NC) ratio (Figure 5 – figure supplement 1), as previously described (Bakshi et al., 2014; Dworsky and Schaechter, 1973; Koch and Gross, 1979; Pettijohn and Hecht, 1974; Xiang et al., 2021). In cells that completed their division cycle before antibiotic addition (squares and fitted blue curve in Figure 5B), the distance between the intensity peak of each sister nucleoid increased monotonically between birth and division, displaying the dynamics of normal, unperturbed nucleoid segregation. Cells born 22 min to 12 min before the treatment (circles in Figure 5B) experienced the same nucleoid segregation dynamics up to the time of rifampicin addition. Exposure to rifampicin lead to the near-immediate arrest of nucleoid segregation (Figure 5B), consistent with our prediction.

Interestingly, the distance between the sister nucleoids remained the same for close to 30 min into rifampicin treatment, after which it started to decrease (Figure 5B). Average 1D and 2D cell projections of the scaled (divided by the whole cell average) RplA-GFP and HupA-mCherry concentration suggest that the time delay between the arrest in nucleoid segregation and its reversal is likely due to the compressible nature of the nucleoid, which has been demonstrated in vitro (Pelletier et al., 2012). As the polysomes in the middle of the cells started to visibly deplete (> 6 min after rifampicin addition), the DNA signal expanded to fill the emerging available space without affecting the distance between the peaks of the sister nucleoids (white crosses, Figure 5C). This is consistent with the removal of a compaction force exerted by the accumulating polysomes on the soft nucleoid. About 30 min after rifampicin addition and further polysome depletion, the peak signals of the sister nucleoids (white crosses, Figure 5C) started migrating closer to each other. Eventually, after 1 h of rifampicin treatment, when the RplA-GFP fluorescence was homogeneous and there were no polysome accumulations detected, the two nucleoid objects fused into one (Figure 5C), consistent with the dissipation of phase separation.

These results support the notion that in untreated cells, polysome accumulation effectively exerts a force

on the compressible nucleoid, which translates into its observed compaction and translocation (hence,

segregation). Gradual polysome depletion through rifampicin treatment progressively decreased this effect,

reversing the process. This reversal became obvious when we plotted the correlation between the relative

polysome accumulation and nucleoid depletion at mid-cell for two cell lineages that experienced rifampicin

at different times after birth (Figure 5D). Irrespective of their birth time (12 or 3 min before the addition of

rifampicin), the negative correlation between the two variables was reversed ~9 min after the addition of

the antibiotic, following the same path as for the untreated cells but in the opposite direction (Figure 5D).

Polysome depletion during rifampicin treatment also resulted in correlated loss of asymmetric nucleoid

compaction in newborn cells (Figure 5E). Altogether, these results support the notion that the asymmetric

accumulation of polysomes results in an anisotropic force that asymmetrically compacts and segregates

nucleoids.

Ectopic polysome production redirects nucleoid dynamics.

Our second approach to test causality experimentally was to redirect polysome formation away from the

chromosome by over-expressing a useless protein (mTagBFP2) for the cell from a T7 promoter on a multi-

copy pET28 plasmid (Figure 6A). The resulting CJW7798 strain also carried the ribosome (RplA-msfGFP)

and DNA (HupA-mCherry) markers. We reasoned that high expression of BFP2 from the plasmid may slow

polysome production within the nucleoid and create polysome accumulations at ectopic cellular locations

through recruitment of ribosomes to plasmid transcripts. This, in turn, should affect nucleoid dynamics if

our proposed mechanism is correct.

Plasmid expression of mTagBFP2 was induced by addition of 100 µM IPTG and expression of a

chromosomally encoded T7 RNA polymerase (Figure 6A-B). The gradual increase of mTagBFP2

fluorescence in the cells was associated with a concomitant decrease in cell growth rate (Figure 6C),

consistent with reduced gene expression from the chromosome. These cells displayed various patterns of

polysome accumulations (Movie S5), presumably due to stochastic clustering of plasmids in DNA-free

regions as previously reported for other multi-copy plasmids devoid of DNA partitioning genes (Hsu and

Chang, 2019; Reyes-Lamothe et al., 2014; Yao et al., 2007). Importantly, the ectopic accumulations of

polysomes had a drastic effect on nucleoid dynamics. In some cells, polysomes accumulated at one pole

instead of the middle of the nucleoid, preventing the nucleoid from splitting (Figure 6D). Expansion of the

polysome accumulation at a pole effectively pushed the nucleoid toward the opposite pole of the cell. In

other cells, polysome accumulation occurred between sister nucleoids, but did not relocate to the

segregated nucleoids at the ¼ and ¾ cell positions. Rather, polysome accumulation persisted and

expanded between the sister nucleoids, effectively further pushing them apart (Figure 6E). Often, cells

filamented and showcased correlated polysome and nucleoid dynamics that changed in time (Figure 6F),

resulting in transient events of nucleoid fusion, splitting, or changes in migration direction depending on

where polysomes accumulated. Movie S5 shows additional examples of such dependency.

All cells displayed diffuse mTagBFP2 fluorescence (Figure 6B). We also observed the formation of inclusion

bodies (bright phase contrast) that typically remained at a pole or sometimes moved along the edge of a

growing polysome accumulation (Figure 6F). After a long of period of IPTG induction (> 8 h), polysome

accumulation eventually decreased, leading to nucleoid decompaction (Figure 6 – figure supplement 1).

Altogether, these results support the notion that ectopic polysome accumulation drives nucleoid dynamics.

Radial confinement prevents nucleoid splitting along the cell width and fusion of polysome

accumulations from distinct DNA-free regions.

In the model (Figure 4), phase separation between polysomes and nucleoid occurs when the cell and

nucleoid elongate. As *E. coli* cells grow by increasing their length and not their width, we reasoned that

radial confinement forces nucleoids to grow along the cell length, thereby promoting nucleoid segregation

along this cellular dimension. Keeping nucleoids close to the membrane across the cell width may act as a

diffusion barrier for polysomes from distinct DNA-free regions, preventing polysome accumulations to fuse

into larger phase condensates.

To test these ideas, we first treated cultures with A22 to inhibit cell width control through the inactivation of

MreB (Bean et al., 2009; Iwai et al., 2002). This resulted in cells with a polysome phase at the cell center

surrounded by a nucleoid phase around the cell periphery (Figure 7 – figure supplement 1, top). To further

increase the cell width (> 2.5-fold), we exposed cells to the cell division inhibitor cephalexin in addition to

A22 (Figure 7A and Figure 7 – figure supplement 1, bottom). As a control, we showed that cephalexin

treatment alone resulted in filamentous cells (of constant cell width) with multiple nucleoids separated by

polysome accumulations (Figure 7B), consistent with previous reports (Chai et al., 2014; Gray et al., 2019;

Thappeta et al., 2024). In cells treated with both drugs, we observed two types of subcellular

rearrangements. In smaller cells, which started with a single nucleoid, drug treatment resulted in a single

large polysome accumulation at the cell center, displacing the nucleoid into a toroidal shape at the cell

periphery (Figure 7C and Movie S6). In longer cells with two segregated nucleoids, the nucleoids expanded

and often aberrantly segregated along the cell width concomitant with polysome accumulation at the site of

nucleoid splitting (white arrowheads, Figure 7D). Consequently, a polysome "bridge" was formed between

the polysome accumulations flanking the nucleoid. These polysome bridges resulted in a characteristic

cross-like polysome pattern, marking the two axes (longitudinal and radial) of nucleoid segregation (Figure

7E). The loss of radial confinement led to coalescence of polysome accumulations toward the cell center

and fusion of nucleoids around the cell periphery (Figure 7C-E and Figure 7 – figure supplement 1). As a

result, the nucleoids and polysome accumulations decreased in number while increasing in size in

cephalexin/A22-treated cells compared to cephalexin-treated cells with the same cell area distribution but

normal cell width (Figure 7F). These results suggest a critical role for cell width regulation in limiting the
diffusion of polysomes around the nucleoid and thereby spatially controlling the phase separation between
polysomes and nucleoids to promote longitudinal nucleoid segregation.
520

DISCUSSION
The flow of genetic information intrinsically couples nucleoid segregation to cell growth.
Polysomes, a term in which we include monosomes for simplicity, are made of mRNAs in complex with
ribosomes. This study provides experimental and theoretical evidence (Figures 1-7) that polysome
production within the nucleoid—an inherent product of chromosomal gene expression—contributes to
nucleoid segregation and positioning in *E. coli* cells. This may also be true in other bacteria, as reduced or
abrogated transcription via gene deletion or antibiotic treatment causes chromosome segregation defects
in *Streptococcus pneumoniae* (Kjos and Veening, 2014) and *Bacillus subtilis* (Dworkin and Losick, 2002).
529

Due to their anabolic footprint (Belliveau et al., 2021; Chure and Cremer, 2023; Dai et al., 2017; Dourado
and Lercher, 2020; Hu et al., 2020; Molenaar et al., 2009; Scott et al., 2014; Si et al., 2017), polysomes
integrate the rate of nucleoid segregation with that of gene expression and cell growth. The concentration
of polysomes and their rate of accumulation in the cell are a direct reflection of transcriptional and
translational activities in the cell (Balakrishnan et al., 2022). The higher the concentration of polysomes, the
faster is the growth rate. Thus, in our proposed model, an increase in polysome concentration not only
leads to more protein synthesis and faster cell growth, but also results in faster nucleoid segregation. The
reverse is true for a decrease in polysome concentration, inherently coupling these processes without the
help of a dedicated regulatory system. Such coupling was seen across isogenic cells with variable growth
rates in the same nutrient condition (Figures 1F, 1H and 2C-E) as well as across nutrient conditions that
led to a wide range of growth rates (Figure 2A-B and Figure 2 – figure supplement 1).
541

Directional nucleoid splitting requires phase separation and radial confinement.
A key aspect of our proposed mechanism for nucleoid segregation is that polysomes form within nucleoids
due to chromosomal gene expression, which, together with polysome turnover due to mRNA decay, creates
an out-of-equilibrium system (Figure 4) (Miangolarra et al., 2021). Redirecting polysome formation to
plasmid gene expression led to ectopic polysome accumulations that dramatically affected nucleoid
dynamics (Figure 6 and Movie S5).
548

In normal cells, the effective force that segregates nucleoids appears to be linked to the propensity of the
chromosomal meshwork and polysomes to phase separate. Mutual exclusion is, at least in part, caused by
the steric repulsion between the two macromolecules. While ribosomes or ribosomal subunits freely diffuse
across the cell unobstructed by the presence of the nucleoid (Bakshi et al., 2012; Sanamrad et al., 2014),
the larger polysomes are impeded by the chromosomal mesh based on size considerations alone (Xiang

et al., 2021). Modeling studies have suggested that such steric hinderance between large crowders
(polysomes) and a polymeric meshwork (chromosome) can result in phase separation and polymer
compaction (Bakshi et al., 2014; Castellana et al., 2016; Miangolarra et al., 2021; Mondal et al., 2011; Wu
et al., 2019b). It is also possible that phase separation between nucleoids and polysomes involves non-
steric interactions such as electrostatic repulsion between the negatively charged DNA and RNA (mRNA
and rRNA), as previously hypothesized (Joyeux, 2015). Such steric and non-steric interactions may
contribute to the effective poor solvent quality of the polysome-rich cytoplasm for the chromosome (Xiang
et al., 2021). Future research will be necessary to elucidate the precise nature of the interaction between
chromosomes and mRNAs, whether individually or in complex with ribosomes.
563
We found that the width of the cell controls the dynamics of phase separation between polysomes and
nucleoids by radially confining the nucleoid and directing its growth and segregation along a single cellular
dimension, cell length (Figure 7). Radial confinement effectively prevents the diffusion and fusion of
polysome accumulations from distinct DNA-free regions, which leads to alternating enrichments of
polysomes and DNA along the length of elongating cells (Figure 7B). This radial confinement explains why
differences in RplA-GFP signal between DNA-free regions (Figure 3A-B and Figure 2 – figure supplement
2) do not equilibrate rapidly, validating our use of a 1D model. Diffusion restriction around nucleoids is
consistent with our previous report that polysomes diffuse much faster over short distances (within DNA-
free domains) than long distances (across DNA-free domains) (Gray et al., 2019). Loss of radial
confinement results in apparent fusion of polysome accumulations near the cell center and displacement
of nucleoids toward the cell periphery (Figure 7C-E). This phase separation between the polysomes and
the nucleoid likely contributes to the torus topology of nucleoids previously observed in A22-treated *E. coli*
(Wu et al., 2019a). Our findings highlight the importance of cell width regulation and suggests that nucleoid
segregation may have imposed an evolutionary constraint on cell morphology.
578
Nucleoid segregation likely involves multiple factors.
Our models show that the most trivial case of uniform production of polysomes (i.e., uniform mRNA
synthesis) within the nucleoid is sufficient to cause an enrichment of polysomes at mid-nucleoid (Figure 4)
(Miangolarra et al., 2021). Inside cells, this polysome enrichment at mid-nucleoid may be enhanced by a
bias in mRNA synthesis across the nucleoid. For instance, the chromosomal region close to the origin of
replication has been shown to be more highly expressed per gene copy than other regions on the
chromosome (Scholz et al., 2019) and this region is located in the middle of the nucleoid prior to DNA
replication (Bates and Kleckner, 2005; Cass et al., 2016; Fisher et al., 2013; Kuwada et al., 2013, 2013;
Mäkelä et al., 2021; Sadhir and Murray, 2023; Wang et al., 2006). In addition, this highly expressed
chromosomal region is the first one to replicate, which should lead to further local increase in mRNA
expression due to a doubling in gene dosage (Pountain et al., 2022). However, it is important to reiterate

that such localized mRNA synthesis is not expected to be a prerequisite for polysome-based nucleoid

segregation since it is not included in our reaction-diffusion model (Figure 4).

Other factors are likely involved in nucleoid segregation. In fact, our data revives the abandoned sixty-year-

old hypothesis by Jacob et al. (1963) that cell growth and surface expansion separate the sister nucleoids,

but with two notable differences. First, the contribution of cell growth to nucleoid splitting would be partial

(Figure 1I and Figure 1 – figure supplement 3). Second, cell growth would contribute predominantly near

the end of the division cycle (Figure 1I). This late timing makes sense for two reasons. First, it corresponds

to the time when polysomes stop accumulating between the separated sister nucleoids and polysome

accumulations emerge at the middle of these nucleoids (i.e., at the ¼ and ¾ cell positions) to start the next

round of segregation (Figure 1D and Movie S2). Second, this is also when *E. coli* switches its cell wall

growth pattern from dispersed along the cell body to zonal and divisome-dependent at mid-cell (Cooper

and Hsieh, 1988; Gray et al., 2015; Navarro et al., 2022; Wientjes and Nanninga, 1989; Woldringh et al.,

1987). Indeed, zonal cell growth between the sister nucleoids was a key assumption of the original 1963

model (Jacob et al., 1963). Altogether, we propose a working model in which gene expression, which results

in polysome production within the nucleoid, promotes nucleoid splitting and accelerates the migration of the

resulting sister nucleoids apart, with cell elongation facilitating this latter function near the end of the cell

division cycle. DNA replication and processes that contribute to chromosome condensation and

organization (such as DNA loop extrusion, supercoiling, and preferential loading of DNA remodeling

complexes) are also likely to be important for robust chromosome segregation across all cells (Danilova et

al., 2007; Harju et al., 2023; Hofmann et al., 2019; Holmes and Cozzarelli, 2000; Lemon and Grossman,

2000; Mäkelä et al., 2021; Minnen et al., 2011; Sawitzke and Austin, 2000; Weitao et al., 1999).

We note that in our time-lapse experiments, the accumulation of polysome signal appeared to slightly

precede the depletion of DNA signal that marked the initiation of nucleoid splitting (Figure 1E). Polysome

enrichment in the middle of unconstricted nucleoids was also occasionally observed in snapshot images of

cells growing on glycerol, a slow growth condition that results in a single nucleoid segregation event late

during the cell division cycle (Figure 1 – figure supplement 4A-D). This is not seen in our model in which

polysome accumulation and nucleoid splitting occur at the same time (Figure 4A and Movie S3). This small

discrepancy may reflect a limitation of our experimental or modeling approach. For example, it is possible

that the point spread function of our fluorescent DNA marker slightly delays the moment at which we can

detect signal depletion at mid-nucleoid and thereby the initiation of nucleoid splitting. Alternatively, the small

difference in timing may be associated with a model simplification. In our model, the nucleoid is effectively

a solution of DNA fragments. In reality, the nucleoid consists of a circular polymer, crosslinked by nucleoid-

associated proteins. These DNA crosslinks may cause a small resistance that marginally delays the

initiation of nucleoid splitting relative to the polysome enrichment at mid-nucleoid.

E. coli is an asymmetric organism.
The spatial macromolecular asymmetries uncovered in our study contrast with the common perception of
E. coli as a symmetric organism. Under our experimental conditions, the distribution of polysomes at the
new pole in newborn cells was, on average, higher than at the old pole through inheritance of the large mid-
cell accumulation of polysomes from their mother cells (Figure 3A and D). We also observed an asymmetric
distribution in DNA density within nucleoids, which correlated with the availability of polysome-free space
along the cell length and width (Figure 3E-G). This asymmetry emerged before cell division (Figure 3D and
Movie S2). Simulations of our reaction-diffusion model suggest that slower diffusing/relaxing nucleoids are
more likely to reproduce these nucleoid position and compaction asymmetries during the finite course of
the cell division cycle (Figure 4D). A reduction in the apparent nucleoid diffusivity has been linked to
nucleoid-associated proteins, which bridge and thus stiffen the DNA polymer (Subramanian and Murray,
2023). The physiological significance for these dynamic cellular asymmetries is not clear at this time, though
it is conceivable that a difference in DNA compaction within the nucleoid may affect gene expression.
Regardless, our study illustrates how spatial and temporal asymmetries in the cytoplasm can emerge from
the interactions between two of the most important macromolecules.
642
643   **METHODS**
644
645   **Strains and constructs**
646   Strains and plasmids used for this study are listed in Tables S2 and S3, respectively, while the sequences
647   of the oligonucleotides used to make constructs can be found in Table S4.
648
To measure the concentration and spatial heterogeneity of ribosomes inside *E. coli*, we used strains in
which the RplA 50S ribosome subunit protein (strains CJW7323, CJW6768, CJW7020 and CJW7651) or
the RpsB 30S ribosome subunit protein (strains CJW6769 and CJW7021) are fused with mEos2 (strains
CJW6768 and CJW6769), msfGFP (strains CJW7020, CJW7021, CJW7651, and CJW7798) or GFP
(CJW7323) (Gray et al., 2019; Xiang et al., 2021). Nucleoid characteristics were measured using strains in
which HupA, a nucleoid-associated protein, is fused with mCherry (strains CJW7323, CJW6723 and
CJW7798) (Xiang et al., 2021). Alternatively, DAPI was used to stain the DNA (strains CJW6768,
CJW6769, CJW7020, CJW7021, and CJW7651).
657
The mCherry-µNS (CJW7651) particles were chromosomally expressed from the native Lac promoter after
induction with 150 µM IPTG (for 3 h for agarose pad experiments, or continuously for microfluidics
experiments). Strain CJW7651 was constructed as follows. The *gfp* coding sequence in the pER12
(pBAD322A-gfp-µNS) plasmid (kind gift from Dr. A. Janakiraman, City College of New York) was swapped
with the mCherry-coding sequence using megaprimer whole plasmid (MEGAWHOP) cloning (Bryksin and
Matsumura, 2010) and the primer pair ER12-MCR-fwd/ER12-MCR-rev2 to generate plasmid pER12-
mCherry. The mCherry-µNS coding sequence and the *rrnB* transcriptional terminator were amplified from

the pER12-mCherry plasmid (primer pair µNSmCherry fwd/rev) and assembled with the *frt* site flanked with

a kanamycin resistance cassette (amplified from pKD13 (Datsenko and Wanner, 2000) using the primer

pair FRT_KanR fwd/rev) and the ColE1 origin of replication (PCR amplified from the pBAD22A plasmid

(Guzman et al., 1995) using the primer pair ColE1 fwd/rev) using Gibson DNA assembly (Gibson et al.,

2009) to form the pAPG1 plasmid. The pAPG1 plasmid also included two 50 bp sites homologous to the

attB region of the *E. coli* chromosome, introduced as overhangs in the primers used for Gibson DNA

assembly. The *attB* homologous regions allowed for the integration of the arabinose inducible mCherry-

µNS expression cassette into the respective site, though this was not used in this study. The pAPG1

plasmid was verified by sequencing using the pAPG1 seq1-5 primers. The mCherry-µNS coding sequence

was then PCR amplified from the pAPG1 plasmid using primer pair lacZYA_redµNS fwd/rev, which includes

50-bp overhangs homologous to the region upstream and downstream of the *lacZYA* operon, and

integrated downstream of the *lac* promoter in the MG1655 strain (Guyer et al., 1981; Jensen, 1993) using

lambda red recombination and the pKD46 plasmid (Datsenko and Wanner, 2000) for the construction of

the CJW7144 strain. Correct insertion of the mCherry-µNS coding sequence to substitute the *lacZYA*

operon coding sequences was confirmed by colony PCR using primer pairs LacI_fwd/CynX_rev and

LacI_fwd/mCherry_rev. Transduction with a P1 phage lysate of the CJW7144 strain was used to transfer

the mCherry-µNS expression cassette into the CJW7020 strain using kanamycin as a selection marker. As

a result, the CJW7145 strain was constructed, which was verified using colony PCR and the primer pairs

LacI_fwd/CynX_rev and LacI_fwd/mCherry_rev, KanR_fwd/CynX_rev. The CJW7145 strain was then

transformed using the pCP20 plasmid (Datsenko and Wanner, 2000), which encodes the FLP recombinase

(Cherepanov and Wackernagel, 1995) to remove the kanamycin resistance cassette, yielding the CJW7651

strain.

The CJW7798 strain was derived from the MG1655 (DE3) strain (Tseng et al., 2010a), which carries the

T7 RNA polymerase-encoding gene under *lacUV5* control. Transduction with a P1 phage lysate of the

CJW5158 strain (Gray et al., 2019) was used to transfer the gene encoding the HupA-mCherry fusion into

the MG1655 (DE3) strain using kanamycin as a selection marker. The pCP20 plasmid (Datsenko and

Wanner, 2000), which encodes the FLP recombinase (Cherepanov and Wackernagel, 1995), was used to

remove the kanamycin resistance cassette, yielding the CJW7466 strain. Transduction with a P1 phage

lysate of the CJW7019 strain, a kanamycin-resistant intermediate of strain CJW7020 (Gray et al., 2019),

was used to transfer the gene encoding the RplA-msfGFP fusion into the CJW7466 strain using kanamycin

as a selection marker. The resulting strain (CJW7766) was grown in the presence of kanamycin since the

cells tend to lose msfGFP fluorescence in the absence of the antibiotic. The sequence of the CJW7766

strain was confirmed by whole genome sequencing.

The pET28:mTagBFP2 plasmid variant was derived from the pET28:GFP plasmid (Shis and Bennett,

2013), which was a gift from Mathew Bennett (Addgene plasmid # 60733 ; http://n2t.net/addgene:60733;

RRID:Addgene_60733). First, the GFP coding sequence was substituted with the mTagBFP2 coding
sequence from the pBAD-mTagBFP2 plasmid (Subach et al., 2011), a gift from Vladislav Verkhusha
(plasmid # 34632; http://n2t.net/addgene:34632; RRID:Addgene_34632). Specifically, the mTagBFP2
coding sequence was amplified using the primer pair mTagBFP2 fwd/rev and assembled (Gibson DNA
assembly) with the pET28 backbone, which was amplified in two pieces using the pET28_one fwd/rev and
the pET28_two fwd/rev primer pairs to derive the pET28:mTagBFP2 plasmid. Then, the kanamycin
resistance cassette that was originally present in the pET28 backbone was substituted with the
chloramphenicol resistance cassette from the pSB3C5-proA-B0032-E0051 plasmid (Davis et al., 2011)
which was a gift from Joseph Davis and Robert Sauer (Addgene plasmid # 107244;
http://n2t.net/addgene:107244; RRID:Addgene_107244). Specifically, the backbone of the
pET28:mTagBFP2 plasmid excluding the kanamycin resistance cassette was amplified using the
pET28mTagBFP2 fwd/rev primers and assembled (Gibson DNA assembly) with the coding sequence of
the chloramphenicol resistance cassette that was amplified using the cmR fwd/rev primer pair. As a result,
the pET28:mTagBFP2-CmR plasmid was created. The proper assembly of the pET28:mTagBFP2 and
pET28:mTagBFP2-CmR plasmids were confirmed by sequencing.
717
718 The pET28:mTagBFP2-CmR plasmid was introduced into CJW7766 by electroporation to create the
719 CJW7798 strain, which was used to redirect the ribosomes away from the nucleoid onto the plasmid-
720 expressed mTagBFP2 mRNAs after inducing the expression of T7 RNA polymerase with IPTG.
721
Growth conditions
Strains CJW6769, CJW7020 and CJW7021 used in Figure 2 and Figure 2 – figure supplement 1 were
grown as previously described (Gray et al., 2019) using a basic M9 medium formulation without trace
elements and supplemented with 0.2% (w/v) carbon source and when specified with 0.1% (w/v) casamino
acids (CAA) and 1 µg/mL thiamine (T). The strain CJW6768 (used in Figure 2 and and Figure 2 – figure
supplement 1) was grown using the same medium formulation. The abbreviations of the medium growth
conditions presented in Figure 2A and Figure 2 – figure supplement 1 are defined in Table S1. For the rest
of our experiments on agarose pads or in a microfluidic device, strains CJW7323, CJW7651 and 7798 were
grown in M9 salts (final concentrations: 33.7 mM Na_2HPO_4, 22 mM KH_2PO_4, 8.55 mM NaCl, 9.35 mM
NH_4Cl, 1 mM $MgSO_4$, 0.3 mM $CaCl_2$) supplemented with trace elements (Fe, Zn, Cu, Co, B, Mn), 0.1%
(w/v) thiamine, and, when specified, 0.4% (w/v) casamino acids. The pH of the 10x M9 salts was adjusted
to 7.2 with NaOH. The trace elements were added at a final concentration of 13.4 mM ethylene-diamine-
tetra-acetic-acid, 3.1 mM of $FeCl_3$-$6H_2O$, 0.62 mM of $ZnCl_2$, 76 µM of $CuCl_2$-$2H_2O$, 42 µM of $CoCl_2$-$2H_2O$,
162 µM of HBO_3-$2H_2O$, 8.1 µM of $MnCl_2$-$4H_2O$.
736
To achieve steady-state exponential growth prior to imaging, a stationary phase liquid culture in the
appropriate growth medium was diluted at least 10,000 times and grown to an optical density at 600 nm
(OD_{600}) between 0.1 and 0.3. These cells were either loaded in a mother-machine-type microfluidics device

(Lin and Jacobs-Wagner, 2022; Wang et al., 2010) where they were grown under the constant flow of
medium (~0.5 µL/sec), or spotted on 1% agarose pads prepared with the same growth medium. In the
microfluidic device, cells were grown for at least 3 h prior to image acquisition. All precultures and
experiments were performed at 30ºC.

The mCherry-µNS particles (CJW7651 strain) were imaged on agarose pads after inducing exponentially
growing cells with 150 µM IPTG for 3 h (Figure 4E-G). Before spotting on the 1% agarose pad, the induced
cells were stained with DAPI (1 µg/mL) for 5 min. In both experiments, the cells were growing in
M9gluCAAT.

To redirect ribosomes to plasmid-expressed, T7 promoter-driven mTagBFP2 mRNAs, exponentially
growing CJW7798 cells grown in M9glyCAAT to an OD ~ 0.2 were spotted on a 1% agarose pad containing
M9glyCAAT and 100 µM IPTG, the latter to induce the expression of the T7 RNA polymerase.

Rifampicin treatment
The antibiotic rifampicin (see Table S5) was used to block transcription, deplete mRNAs, and release the
mRNA-bound ribosomes (polysomes). Rifampicin treatment was performed either in batch culture (Figure
1 – figure supplement 1F) or in microfluidics (Figure 5 and Movie S4).

For the microfluidic experiment, two rounds of rifampicin treatment were performed by switching between
M9gluCAAT containing antibiotic (100µg/mL) and antibiotic-free medium at 30ºC. For the medium switches,
solenoid valves were used in a custom-built pressurized perfusion system, achieving fast (within 1 min)
changes in the cellular environment. The first switch to rifampicin occurred 2 h after normal growth in
antibiotic-free medium (Figure 5A and Movie S4). Rifampicin treatment lasted 2 h until the system switched
back to antibiotic-free medium, where cells were left to recover for 8 h. Twelve hours into the experiment,
the system switched again to rifampicin-containing medium.

A22 and cephalexin treatment
Cephalexin (50 µg/mL) and A22 (4 µg/mL) (see Table S5) were added to exponentially growing CJW7323
cell cultures (OD_{600} ~ 0.1) in M9gluCAAT just prior to spotting on a 1% agarose pad, which was made with
the same growth medium and contained the same antibiotic concentrations. The radial expansion of the
cells and the polysome and nucleoid signals were tracked over time in a time-lapse experiment. The
selected concentration of A22 has previously been shown to not affect cell growth (Takacs et al., 2010). All
precultures and time-lapse imaging experiments were performed at 30ºC.

Ectopic polysome accumulation experiment

Exponentially growing CJW7798 cells in M9glyCAAT supplemented with kanamycin (50 μg/mL) and

chloramphenicol (35 μg/mL) were washed one time with M9glyCAAT minimal medium lacking antibiotics

but supplemented with 100 μM IPTG (see Table S5). The washed cells were spotted on a 1% agarose pad

prepared with M9glyCAAT also supplemented with IPTG (100 μM) for time-lapse imaging. All pre-cultures

and time-lapse imaging experiments were performed at 30ºC. Since chloramphenicol affects protein

synthesis and thus polysome formation, the agarose pads lacked chloramphenicol, resulting in a fraction

that loss the plasmids based on the absence of blue fluorescence. Therefore, only cells that expressed blue

fluorescence (and thus carried the pET28:mTagBFP2-CmR plasmid) were analyzed.

Microscopy

Strains CJW6769, CJW7020, and CJW7021 used in Figure 2A-B and Figure 2 – figure supplement 1, were

imaged on agarose pads using the same microscopy set-up and optical configurations as previously

described (Gray et al., 2019). Snapshots of the CJW6768 strain (used in Figure 2A-B and Figure 2 – figure

supplement 1) were taken using a Nikon Ti-E microscope equipped with a 100x Plan Apo 1.45NA Ph3 oil

objective, a Hamamatsu Orca-Flash4.0 V2 CMOS camera (16-bit, Slow Scan sensor mode) and a

Lumencor Spectra X LED (Light Emitting Diode) engine. The 395/25nm LED was used to excite DAPI and

the 470/24nm LED was used to excite GFP. DAPI fluorescence was acquired using an ET350/50x

(excitation filter), RT400lp (dichroic mirror), ET460/50m (emission filter) filter cube from Chroma. The

ET470/40x (excitation filter), T495lpxr (dichroic mirror), ET525/50m (emission filter) configuration from

Chroma was used for GFP fluorescence acquisition. The microscope was controlled using the NIS

Elements software by Nikon and the ND acquisition module.

For the time-lapse observation of cells (strain CJW7323) treated with A22 and/or cephalexin, images were

taken using a Nikon Ti-E microscope, equipped with a 100x Plan Apo 1.45NA Ph3 oil objective, a

Hamamatsu Orca-Flash4.0 V2 CMOS camera (16-bit, Slow Scan sensor mode), and a Sola solid state

white light source (Lumencor), in a temperature-controlled enclosure (Okolabs). The AT470/40x excitation

filter, combined with a T495LPXR beam-splitter and an ET525/50m emission filter, was used for GFP. For

mCherry visualization, the ET560/40x excitation filter, combined with a T585lp beam-splitter and a

ET630/75m emission filter, was used. A neutral density filter (32x) was applied in the excitation path to

reduce phototoxicity and photobleaching. Images were taken every 2.5 min in the brightfield channel (phase

contrast) and every 5 min in the fluorescence channels (RplA-GFP and HupA-mCherry).

For the rest of the brightfield and epi-fluorescence wide-field microscopy experiments (strains CJW7323,

CJW7651, and CJW7798), snapshots or time-lapse images were taken using a Nikon Ti2-E inverted

microscope, equipped with a 100x Plan Apo 1.45NA Ph3 oil objective, a Photometrics Prime BSI back-

illuminated sCMOS camera (2048x2048 pixels sensor with a pixel size of 6.5 μm), and a Lumencor Spectra

III LED (Light Emitting Diode) engine, in a temperature-controlled enclosure (Okolabs). The HDR 16-bit
sensor mode was used to acquire images. The microscope was controlled using the NIS Elements software
by Nikon, and the JOBS or the ND acquisition module were used to acquire snapshots or time-lapse
images. The Perfect Focus System (by Nikon) was used to maintain focus in microfluidics experiments.
The auto-focus function in NIS Elements was used to locate the optimal z-position in agarose pad
experiments (snapshots or time-lapse). For the DAPI, BFP, GFP, or mCherry channels, a polychroic mirror
(FF-409/493/596-Di02 by Shemrock) combined with a triple-pass emitter (FF-1-432/523/702-25 by
Shemrock) was used. For DAPI, BFP, and GFP imaging, additional emission filters were applied in the
optical path (FF01-432-36 by Shemrock, FF01-432-36 by Shemrock and ET525/50M by Chroma,
respectively). DAPI, BFP, GFP, and mCherry were excited using a 390/22nm, 390/22nm, 475/28nm and
575/25nm LED, respectively.
825
For the microfluidic experiments (strain CJW7323), the excitation light intensity was reduced to 20% for the
1-min interval and 30% for the 3-min interval imaging of the RplA-GFP and HupA-mCherry. An exposure
time of 120-ms was used for both markers. In all time-lapse experiments, a neutral density filter (absorptive
ND filter, OD:1.3 / 5% transmission, NE13B by Thorlabs) was also applied to reduce the LED power and
minimize phototoxicity and photobleaching. The LED powers (factored by the neutral density filters) were
calibrated using a microscope slide power meter with an 18x18 mm sensor size (S170C by Thorlabs). The
light intensity was measured for each excitation wavelength (475/28nm and 575/25nm) at the end of the
objective (without immersion oil) for different LED intensities (% of maximum intensity). From the generated
calibration curves, the RplA-GFP excitation light power was estimated to be 524 µW and 629 µW for the 1-
min and 3-min fluorescence interval imaging, respectively. HupA-mCherry was excited with 109 µW light
power for the 1-min and 191 µW for the 3-min fluorescence interval imaging.
837
838 The type N immersion oil (by Nikon) was used in all experiments except for the time-lapse observation of
839 A22+cephalexin-treated cells where the type F immersion oil (by Nikon) was used. All imaging (time-lapse
840 and snapshots) was done at 30ºC.
841
Analysis software and code availability
Cropping and alignment of the microfluidic channels were performed with MATLAB (www.mathworks.com)
using a previously published pipeline from our lab (Lin and Jacobs-Wagner, 2022). Supervised classification
and curation of the Oufti cell meshes was also implemented in MATLAB (Campos et al., 2018; Govers et
al., 2024; Gray et al., 2019). For the T7 experiments, segmentation and tracking of the CJW7798 cells was
performed using the Omnipose deep neural network architecture (Cutler et al., 2022), using a previously
trained model (Thappeta et al., 2024) and the SuperSegger MATLAB-based package (Stylianidou et al.,
2016). The remaining analyses were performed using Python 3.9 (www.python.org), that included the
numpy (Harris et al., 2020), *scipy* (Virtanen et al., 2020), *pandas* (McKinney, 2010), *scikit-mage* (Van Der

Walt et al., 2014), *scikit-learn* (Pedregosa et al., 2012)*, shapely* (Gillies, Sean et al., 2023), *statsmodels*

(Seabold and Perktold, 2010) and *pytorch* (Paszke et al., 2019) libraries. The *matplotlib* (Hunter, 2007) and

seaborn (Waskom, 2021) libraries were used for plotting. The analysis pipeline and functions (summarized

in Table S6) are available in the Jacobs-Wagner lab GitHub repository:

www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024.

Cell segmentation and tracking

For the CJW6768, CJW6769, CJW7020, and CJW7021 strains used in Figure 2 and Figure 2 – figure

supplement 1, Oufti (Paintdakhi et al., 2016) was used to draw cell meshes on the phase contrast snapshots

and a MATLAB-based support vector machine model was used to remove the badly segmented cells as

previously described (Campos et al., 2018; Govers et al., 2024; Gray et al., 2019).

The Omnipose deep neural network architecture (Cutler et al., 2022) with a previously trained model

(Thappeta et al., 2024) was used to segment the CJW7798 cells during T7 RNA polymerase induction on

an agarose pad. The segmented cells were then tracked using SuperSegger (Stylianidou et al., 2016). A

custom class (*omnipose_to_python_ghv.py*) was developed to transfer the SuperSegger segmentation and

tracking data into Python for post-processing.

Due to the unusual and variable morphology of the A22/cephalexin-treated cells, a custom Python class

was developed for their segmentation and tracking. The *Otsu_phase_segmentation_ghv.py* class

segments the cells by applying an Otsu-threshold (Otsu, 1979) on the inverted phase images, followed by

binary dilation (*scikit-image.morphology.binary_dilation* Python function) hole filling

(*scipy.ndimage.binary_fill_holes* Python function), and binary closing (*scikit-*

image.morphology.binary_closing Python function). The segmentation masks were tracked between

subsequent timepoints, using cell distance as well as cell area constraints, and linked into cell growth

trajectories. Cell morphology criteria such as the maximum pixel distance from the medial axis and the

medial axis sinuosity were used to remove bad segmentations. The remaining segmentation masks were

manually curated.

For the other experiments on agarose pads, a neural network (Wiktor et al., 2021; Zhou et al., 2020) with a

previously designed and trained U-net architecture (Mäkelä et al., 2024) was used to segment single cells

based on phase contrast snapshots. The generated segmentation masks were further processed by

watershed separation, filling the holes within masks and removing unusually small masks. Finally, a

graphical interface was used to manually remove the badly curated cells (less than 5% of the segmented

cell population).

In contrast to the Oufti software (Paintdakhi et al., 2016) that generated sub-pixel meshes around the cell
boundaries, the other applied segmentation algorithms returned pixel-based cell masks. To deal with this
discrepancy, the sub-pixel cell meshes from Oufti were converted into pixel-based cell masks by collecting
the pixels within the circumscribed single cell area in Python (*oufti_snapshots_GrayGovers* class, included
in the *snapshots_analysis_OUFTI_GrayGovers* Python script). After this conversion, the same Phyton-
based functions were applied for the analysis of the cell fluorescence and morphology statistics regardless
of the segmentation method used.

To segment cells growing in the microfluidic device, a custom library of image analysis functions
(*mother_machine_segmentation* class, included in the *microfluidics_segmentation_ghv* Python script) was
developed in Python. This algorithm was applied on the cropped, aligned, background-subtracted, and
inverted phase contrast images that were produced using a previously published pipeline (Lin and Jacobs-
Wagner, 2022) in MATLAB. Cell segmentation was performed in three steps. First, all the cells within the
microfluidics channel, which were brighter than the microfluidic channel background in the inverted and
background-corrected phase-contrast channel, were segmented using an Otsu intensity threshold (Otsu,
1979). This crude thresholding step, which separated the cell (brighter: 1) from the background (darker: 0)
pixels, yielded at least one masked label for the entire row of stacked cells in each microfluidic channel. A
watershed segmentation was then applied to define the boundaries between individual cells and split the
Otsu-based binary mask(s). The watershed algorithm was guided by the number of cells and their relative
positions in the microfluidic channel, which were determined using an adaptive filter combined with the local
decrease of the inverted phase intensity between the poles of adjacent cells. Finally, a graphical interface
was used to curate the cell masks by manually splitting or merging cell labels.

After segmentation, the curated single-cell masks were tracked over time and linked into trajectories from
birth to division using the centroid distance and the relative area difference between cells at consecutive
time-points (*mother_machine_tracking* class, included in the *microfluidics_segmentation_ghv* Python
script). Examples of cell segmentation and tracking in microfluidics are shown in Movie S1 and Figure 1 –
figure supplement 1A. Regardless of the image acquisition time intervals for the fluorescence channels,
phase-contrast images were acquired every minute, which allowed for accurate cell tracking.

Fluorescence background correction was different between the microfluidic and agarose-pad experiments.
In microfluidics, the average background was measured within two 8-pixel (0.528 µm) wide areas, one on
the left side and one the right side of the channel, at least 10 pixels (0.66 µm) away from the channel
boundaries from top to bottom. The estimated background was then subtracted along the channel length.
The background correction was integrated in the *get_fluorescence_image* function, located in the
*microfluidics_analysis_functions_*ghv Python script.

In agarose-pad experiments, an Otsu threshold (Otsu, 1979) was used on the inverted phase-contrast
image to segment all cells. A binary dilation was then performed on these crude cell masks before
estimating the local average fluorescence of the unmasked pixels (pixels outside the dilated cell masks).
The locally-averaged background was then used to fill in the cell areas and reconstruct the background of
the entire field of view in the absence of cells. The smoothed (Gaussian smoothing) reconstructed
background fluorescence was subtracted from each fluorescence image. This background correction
pipeline (*back_sub* function) is integrated in both the *unet_snapshots* class and the
oufti_snapshots_GoversGray class. For the time-lapse imaging of A22-treated CJW7323 cells (Figure 7)
and CJW7798 cells during induction of T7 RNA polymerase expression (Figure 6), the fluorescence
background was not subtracted since we did not need to quantify the raw pixel values. The normalized
polysome and nucleoid fluorescence normalized by the average whole cell fluorescence is shown instead,
marking the relative macromolecular rearrangements.

mCherry-µNS particle localization and tracking
For the localization of the diffraction-limited mCherry-µNS particles (Figure 4E-G), a set of custom functions
were developed in Python and implemented in the *particle_positions_snapshots* class, included in the
snapshots_analysis_UNET_ghv Python script.

The fluorescent particles were segmented in the background corrected mCherry-µNS fluorescence images
using a Laplace of Gaussian (LoG) and an adaptive filter combined. The particle masks that had an area
larger than a specified threshold (80 pixels for the microfluidics and 90 pixels for the agarose-pad
experiments), which usually included two particles from the same or adjacent cells, were further processed
by applying a relative fluorescence threshold (90^{th} percentile of masked pixel intensity) to find the local
fluorescence maxima and separate the two objects. Additional minimum area and aspect ratio constraints
were applied. To accurately estimate the particle position with sub-pixel resolution, a 2D Gaussian function
(eq. 1) with rotation (eq. 2) was fitted (least square method: *scipy.optimize.leastsq*) to an area of 7x7 pixels
centered at the centroid of the particle mask:

$$g(x,y) = A \exp\left(-\frac{1}{2}\left(\left(\frac{x_0 - x_{rot}}{\sigma_x}\right)^2 + \left(\frac{y_0 - y_{rot}}{\sigma_y}\right)^2\right)\right) \quad \text{[eq. 1]}$$

952
953   and
954   $$x_{rot} = x\cos(\theta) - y\sin(\theta)$$
955   $$y_{rot} = x\sin(\theta) + y\cos(\theta) \qquad \text{[eq. 2]}$$
956
where *x* and *y* are the coordinates in the 2D imaging plane, *A* is the amplitude of the fitted Gaussian, σ is
the standard deviation in each dimension, and θ is the rotation angle in radians, which was used to rotate
the Gaussian distribution (x_{rot}, y_{rot}) around the particle center or the Gaussian mean (x_0, y_0).

Two-dimensional cell mapping and alignment

Ribosome and nucleoid fluorescence statistics, as well as particle positions within cells, were mapped in

relative 2D cellular coordinates from pole to pole and across the cell width. Such mapping allowed us to

project the fluorescence or position statistics in 1 or 2D for specific cell length ranges and cell division cycle

intervals. For this analysis, it was necessary to draw the medial axis for each of the segmented cells, which

was obtained differently in microfluidic and agarose-pad images.

In the microfluidic experiments, where all the cells were stacked in a vertically oriented channel, the medial

axis was drawn by fitting a second-degree polynomial to the most distant coordinates from the cell

boundaries, excluding the cell caps at the poles where the medial axis was linearly extrapolated. The length

of the cells, which was parallel to the length of the microfluidic channel, was used as the independent

variable for the fitting. This medial axis estimation is implemented via the *all_medial_axis* function in the

microfluidics_analysis_functions_ghv Python script.

However, in the agarose-pad experiments, cells were randomly oriented in the imaging plane. Thus, it was

impossible to use one of the coordinates (x or y) as the independent variable as this would bias our medial

axis estimation toward the same coordinate for cells that were not diagonally oriented. Instead, we

developed an algorithm that scanned through the middle of the cells with a fixed sub-pixel step and

directionality constraints to identify nodes at the most distant locations from the cell boundaries. These

ordered (from pole to pole) and numbered nodes were used to fit the x and y coordinates of the medial axis

separately, using the number of the node as an independent variable and its position coordinates as

dependent variables. The degree of the fitted polynomials (*d*) scaled linearly with the total number of nodes

(*N*) based on the relation $d = 0.1N - 5$. Similar to the microfluidic experiments, the medial axis was linearly

extrapolated at the cell caps. This medial axis estimation is implemented using the *get_medial_axis* function

in both the *unet_snapshots* and *oufti_snapshots_GoversGray* classes located in the

snapshots_analysis_UNET_ghv and *snapshots_analysis_OUFTI_GrayGovers_ghv* Python scripts

respectively. The medial axis estimation function is also provided as a separate function in the

Bivariate_medial_axis_estimation.py Python script.

The medial axis of each cell, with a resolution of 0.1 pixel, was used to map the cell pixels and particle

positions along the cell length and width. The projection of each pixel or particle position on the medial axis,

defined as the most proximal node on the central line, was used to determine its cell length coordinate. The

distance between the pixel or particle position and its medial axis projection was used to determine its

absolute cell width coordinate. The sign of the cross product $\overrightarrow{PL} \times \overrightarrow{PC}$ (plus or minus), where *L* was the

location of the particle, *P* was its projection on the medial axis, and *C* was the center of mass of the cell

mask, was used to determine the position of particles or cell pixels on the sagittal plane.

Cell polarity and ages

In time-lapse experiments, the old and new poles of the cells were determined as follows. The pole of the

daughter cell that was closer to the mid-cell position of its predivisional mother cell, where cell constriction

occurs, was defined as the new pole. The medial axis coordinates were adjusted based on this polarity. As

a result, the relative cell length extended from -1 to 1, with -1 corresponding to the old pole, 1 to the new

pole, and zero to the cell center (see Figure 3 – figure supplement 1B, D and E). The implementation of

this method is included in the *get_polarity* function in the *microfluidics_analysis_functions_ghv* Python

script.

1007

Cell lineages and ages relative to the oldest mother cells (first lineage) at the closed end of the microfluidic

channel were determined using the orientation of the cell poles. The daughter cells that had the same

polarity as the oldest mother cells were assigned the same age and lineage as their mothers. The daughter

cells with the opposite polarity, which inherited the new pole of their mother cells, were considered younger

and belonged to the second lineage. The daughter cells of mother cells belonging to the second lineage

were classified as lineages three and four depending on their polarity. Cells from the third lineage had

opposite polarity and those from the fourth lineage had the same polarity as their second lineage mother

cells. Finally, mother cells from the third lineage, which inherited the new pole of their second-lineage

mother cells, divided to yield the fifth and sixth cell lineages. The fifth lineage had the same polarity as its

third-lineage mother and its sister lineage (sixth) had the opposite polarity and inherited the new pole from

its third-lineage mother cell. The implementation of this method is included in the *get_ages* function in the

*microfluidics_analysis_functions_*ghv Python script. For the analysis of the data presented in Figure 3, only

the third, fourth, fifth and sixth lineages were considered to avoid age-related effects at the cell poles (Chao

et al., 2023; Coquel et al., 2013; Koleva and Hellweger, 2015; Łapińska et al., 2019; Lindner et al., 2008;

Proenca et al., 2019).

1023

1024 **Construction of intensity profiles, 2D cell projections, demographs, and kymographs**

1025 With the ribosome and nucleoid fluorescence pixels as well as the µNS particle positions inside the cells

1026 having been mapped, the fluorescence and particle statistics from pole to pole (medial axis projection) and

1027 across the cell width (distance from the medial axis) were plotted for different cell division cycle intervals or

1028 cell length ranges.

1029

An intensity profile represents the change in a fluorescence statistic along the cell from one pole to the

other. Such a fluorescence statistic can be the average fluorescence intensity (as in Figures 1D, 3G, 4G,

and Figure 3- figure supplement 1B), which corresponds to the concentration of the reported protein per

cell area (sum of pixels divided by the number of pixels). In other analysis (as in Figure 5C and Figure 1 –

figure supplement 4C), the protein concentration along the cell length was divided by the whole cell

average. This scaled statistic is expressed as a percentage change relative to average concentration (mean

%). A value above 100 indicates increase above the average, whereas a value below 100 corresponds to
a decrease below the average concentration. This scaled statistic is insensitive to the RplA and HupA
concentration variability between cells, or to the maturation of the GFP or mCherry fluorophores. As a result,
the scaled concentrations showcase rearrangements of the tagged proteins and do not imply changes in
protein synthesis relative to cell growth.

1041

Intensity profiles were plotted for single cells (Figure 3 – figure supplement 1B and Figure 1 – figure
supplement 4A) or populations (as in Figures 1D, 3G, 4G, 5C, 5E). In the second instance, the intensity
profiles represent the average 1D projection of the fluorescence statistic for a specific cell division cycle
interval, cell length range, growth rate range, polysome and nucleoid asymmetry group, or time range (e.g.,
during antibiotic treatment). To generate a single-cell intensity profile, the medial axis length and its
projected pixels were binned, and the average value of the selected fluorescence statistic was calculated
per bin. If the number of bins was higher than the length of the medial axis in pixels, the missing values
were filled in by linear interpolation. To avoid artefacts from the cell boundaries, the intensity profile was
estimated for a box with a width of six pixels (~0.4 µm) centered on the medial axis (i.e., three pixels on
each side of the medial axis). The intensity profile was occasionally smoothed by a moving average of a
specified cell-length window, without excluding the bins at the edges which were not smoothed.
Alternatively, a univariate spline (*scipy.interpolate.UnivariateSpline*) was fitted to the cell-length-binned data
(as in Figure 4G). The univariate splines were fitted to the binned fluorescence intensity along the cell
length. The number of cell length bins (N_{bin}) linearly scaled with the cell length in µm (l_{cell}) using the first-
order relationship: $N_{bin} = \frac{100}{4.3} l_{cell}$ (only the whole part of the decimal was considered). Similar to the
intensity profile where a fluorescence statistic was plotted along the cell length, the density profiles
(histograms of the particle positions along the cell) for the mCherry-µNS (Figure 4G) particles along the cell
were also plotted. Note that the intensity profiles shown in Figures 1D, 3G, 4G, 5C, 5E, Figure 1 – figure
supplement 1D, and Figure 2 – figure supplement 2D are averages of many segmented cells or cell division
cycles. In these averaged profiles, the variability in the timing of nucleoid segregation across cells
attenuates the appearance of polysome accumulation. Furthermore, the point spread function of fluorescent
ribosome markers causes an underestimation of the magnitude of the polysome accumulations in epi-
fluorescence micrographs (Bakshi et al., 2012).

1065

The intensity profiles were also used to construct demographs from agarose-pad experiments (Figure 2A,
Figure 2 – figure supplement 1, and Figure 1 – figure supplement 4B) where cells were sorted by cell length
from the shorter newborn to the longer predivisional cells. Intensity profiles were also used to construct
ensemble kymographs from microfluidic experiments where the cells were sorted by their relative position
in the cell division cycle (Figures 1C and 2D) or single-cell kymographs from agarose pad time-lapse
measurements (Figure 6D-F). Specifically, the average intensity profile was estimated for each cell length
or cell division cycle bin, and the average intensity profiles were stacked from birth to division (left to right)

oriented according to the cell polarity (top to bottom). In the ensemble kymographs, the fluorescence

intensities were projected along the relative cell length (cell length %), whereas in the single-cell

kymographs the length of the projection linearly scales with the cell length. A Gaussian smoothing

(*skimage.filters.gaussian*) was applied to smooth the demographs and ensemble kymographs.

1077

In addition to the intensity or density profiles for a fluorescence statistic or the density of the µNS particles

along the medial axis, 2D projections of the ribosome or nucleoid signal distribution (Figures 1D, 3D, 3G,

4F, 5C, 5E, and Figure 2 – figure supplement 2A) as well as the µNS particle positions (Figure 4F) were

also constructed. These 2D intensity or density maps were constructed by binning the cell pixels or particle

positions not only by cell length, but also by cell width. The average fluorescence statistic or the particle

density was then shown per bin, including data from many single cells within a specified cell division cycle

or cell length range. A Gaussian smoothing (*skimage.filters.gaussian*) was applied to smooth the 2D

projections.

1086

Extraction of population-level statistics of polysomes and nucleoids

To quantify the correlation between polysome accumulation and nucleoid segregation at the population

level (Figure 2A-B and Figure 2 – figure supplement 1), the average polysome accumulation at mid-cell

was extracted from cell snapshot images. This statistic was obtained by averaging the scaled (divided by

the whole cell average concentration) RplA-GFP and HupA-mCherry intensity profiles of all the cells in the

population. Therefore, this statistic describes the average "behavior" of the population under a specific

growth condition. We reasoned that if, in a specific nutrient condition, nucleoid segregation happens earlier

during the division cycle, then a higher fraction of the population will have segregated nucleoids. If so, the

scaled HupA-mCherry concentration should exhibit a stronger depletion at mid-cell on average. This

depletion was measured as the relative concentration difference between the HupA-mCherry peaks, which

corresponded to the two lobes of the constricting nucleoid, and the trough at mid-cell, which corresponds

to the point of nucleoid splitting. Similarly, the average polysome accumulation was measured as the RplA-

GFP relative concentration difference between the mid-cell accumulation (peak) and the polysome depleted

regions near the quarter cell positions, or at the centers of the segregating sister nucleoids (troughs). The

same statistics were used to quantify the relative polysome accumulation at mid-cell and the relative

nucleoid depletion in the same region between cell division cycles with different growth rates in the same

nutrient condition (Figure 2C-E).

1104

Nucleoid segregation cycle tracking

In this work, the nucleoid segregation cycle was defined as the period from the end of a nucleoid splitting

event (i.e., when a segregating nucleoid is segmented by local thresholding (custom Python function:

LoG_adaptive_image_filter.py) as a separate object from its sister nucleoid) until the end of the splitting of

the same nucleoid object (Figure 1 – figure supplement 2D), usually after cell division but sometimes in the

same cell division cycle (Figure 1 – figure supplement 2A).

1111

To track the nucleoid objects through the nucleoid segregation cycle from the mother cells to their

daughters, we used the cell polarity, i.e., based on pole identity (Figure 1 – figure supplement 2B). The

nucleoids of daughter cells that had opposite polarity to their mother cells were inherited from the new pole

of the mother cells (groups 1 and -1 in Figure 1 – figure supplement 2B-C). The nucleoids in the daughter

cells that had the same polarity as their mother cells were inherited from the old pole of the mother cells

(groups 2 and -2 in Figure 1 – figure supplement 2B-C). During a complete nucleoid segregation cycle, the

nucleoid was tracked from the quarter position of the mother cell to the center of its daughter cells (Figure

1 – figure supplement 2C). This method is implemented in the *track_nucleoids* function, included in the

microfluidics_analysis_functions_ghv Python script.

1121

1122   To quantify the polysome accumulation and nucleoid splitting during the nucleoid segregation cycle (Figure

1123   1E-F), we measured the RplA-GFP and HupA-mCherry concentration within the cell length region of 2.5

1124   pixels adjacent to the center of the tracked nucleoid object (5 pixels or 0.33 µm in total).

1125

Quantification and segmentation of polysomes and nucleoids segmentation in cells treated with

cephalexin or cephalexin + A22

The polysome accumulations and nucleoids objects were also segmented in A22 and cephalexin-treated

cells using an adaptive filter (custom Python function: *LoG_adaptive_image_filter.py*) on the masked cell

fluorescence. The number of the detected polysome accumulations or nucleoid objects corresponds to the

number of the segmented labels in the RplA-GFP or HupA-mCherry channel respectively. Similarly, the

average polysome accumulation or nucleoid area corresponds to the average area of all segmented labels

per channel. For a fair comparison between cells treated with A22 + cephalexin and cells treated with

cephalexin alone, cells were randomly sampled without substitution from 12 cell area bins between 10 and

13 µm^2 (bin width = 0.25 µm^2) such that the compared populations in Figure 7F had the same cell area

distributions (same number of cells per cell area bin).

1137

Linear mixed-effects models to determine the relative contribution of polysome accumulation and

cell elongation to the migration of the sister nucleoids

To estimate the relative contribution of the polysome accumulation at mid-cell and that of cell elongation to

the migration of the separated sister nucleoids (first seen in Figure 1H), linear mixed-effects regressions

were fitted to the scaled single cell data shown in Figure 1 – figure supplement 3B (*statsmodel* package in

Python, *statsmodels.formula.api.mixedlm* function). One model was fitted for each relative time interval

between the end of nucleoid splitting and cell division, describing the rate of distance increase between the

separated sister nucleoids (σ, response variable) as a function of the polysome concentration increase rate

(ρ), the rate of cell elongation (λ), their interaction (ρλ), and noise (ε) (*statsmodels.formula.api.ols*) as

follows:

1148
1149
$$\sigma = \beta_0 + \beta_1 \rho + \beta_2 \lambda + \beta_3 (\rho\lambda) + \varepsilon \qquad \text{[eq. 3]}$$
1150

The single cell data were scaled by subtracting the population mean and then dividing by the standard

deviation prior to model fitting for each of the four relative time intervals (in Figure 1 – figure supplement

3B). The coefficients of the linear regression model (shown in Figure 1I) provide a measure of the relative

effect of the independent variables on the dependent variable. The cell elongation (λ) had a statistically

significant effect to the response variable (σ) Prob(>|Z|) < 10^{-9}) across all relative time intervals. The

polysome accumulation coefficients (ρ) had a statistically significant effect to the response variable (σ)

(Prob(>|Z|) < 10^{-14}) for the first three relative time intervals (0-25%, 25-50% and 50-75%) and a marginally

significant one Prob(>|Z|) = 0.02) for the last quartile (75-100%). The interaction term (ρλ) had a marginally

significant effect (Prob(>|Z|) = 0.002) for the first quartile (0-25%) yet with a very small coefficient of 0.06

and did not present any significance (Prob(>|Z|) > 0.05) for the remaining time intervals (25-50%, 50-75%

and 75-100%). Hence, it is not shown in Figure 1I. The most significant effect on the migration of the sister

nucleoids (σ) was presented by the rate of polysome accumulation at mid-cell (ρ), 0-25% from the end of

nucleoid splitting to cell division, with a coefficient of 0.46 and Prob(>|Z|) < 10^{-84}.

1164
1165
Calculation of rates

The rates of the polysome accumulation ($\frac{\Delta RplA_{mid-nuc} \ conc.}{\Delta T}$) or nucleoid depletion ($\frac{\Delta HupA_{mid-nuc} \ conc.}{\Delta T}$) in the

middle of the nucleoid (2.5 pixels adjacent the nucleoid center) (Figure 1F), correspond to the slope of a

linear regression (Python fitting function: *numpy.polyfit*) fitted to the change of the mid-nucleoid RplA-GFP

or HupA-mCherry concentration over time, 40-90% into the nucleoid segregation cycle (as in Figure 1E).

This nucleoid cycle interval corresponds to the average time from the initiation of nucleoid splitting to just

before its completion. Choosing the appropriate range to correlate the two statistics was important since

polysomes appeared to accumulate in the middle of the nucleoid before the onset of nucleoid splitting

(Figures 1E and Figure 1 – figure supplement 4D).

1175

The rate of RplA-GFP concentration increase at mid-cell ($\frac{\Delta RplA_{mid-cell} \ conc.}{\Delta T}$) was calculated (Figure 1H) within

a region covering 10% of the total cell length at the cell center. The same results were obtained when the

rates were calculated within a cell region covering 2.5 pixels adjacent to the cell center. The rate of nucleoid

migration ($\frac{\Delta Distance_{nuc}}{\Delta T}$) corresponds to the slope of a linear regression that describes the natural log-

transformed distance increase between the separated sister nucleoids over time (Figure 1H). Similarly, the

rate of cell elongation ($\frac{\Delta Length_{cell}}{\Delta T}$) was calculated considering the natural log-transformed cell length

increase over time (Figure 1H). The rate of RplA-GFP concentration increase at mid-cell ($\frac{\Delta RplA_{mid-cell} \ conc.}{\Delta T}$),

the rate of nucleoid migration ($\frac{\Delta Distance_{nuc}}{\Delta T}$), and the rate of cell elongation ($\frac{\Delta Length_{cell}}{\Delta T}$) were calculated within

four relative time intervals from the end of nucleoid splitting to cell division (Figure 1G and Figure 1 – figure

supplement 3B). Only single nucleoid migration intervals with more than four timepoints were considered

to ensure a reliable fitting. The small fraction (~8%) of cells that were born with two separately detected

nucleoid objects, which indicates that the end of nucleoid splitting occurred in the previous cell division

cycle, were removed from the analysis. The cell division cycles with a negative cell elongation rate (<1%)

were also excluded from the analysis.

1190

Reaction-diffusion model

Our theoretical model is an extension of previous work (Miangolarra et al., 2021), which showed that steric

effects between the DNA and the transcription-translation machinery (including polysomes) are sufficient

to drive nucleoid segregation. To simply describe these interactions, we employed the Flory-Huggins theory

for regular solutions (Flory, 1942; Huggins, 1941) and modeled the non-dimensionalized free-energy

density f as a function of the local volume fractions of the nucleoid (n) and polysomes (p):

1197

1198 $$f(n,p) = \frac{p}{v_p}\ln p + \frac{n}{v_n}\ln n + (1-p-n)\ln(1-p-n) + \chi_p p(1-p-n) + \chi_n n(1-p-n) + \chi_{np} np -$$

1199 $$\kappa_p \nabla p \nabla(1-p-n) - \kappa_n \nabla n \nabla(1-p-n) - \kappa_{np} \nabla n \nabla p . \qquad \text{[eq. 4]}$$

1200

1201 Here, $(1-n-p)$ is the volume fraction of the cytoplasm not occupied by nucleoid or polysomes, whereas

1202 v_n and v_p are proportional to the molecular volumes of the elementary translational degrees of freedom

1203 (i.e., of the nucleoid element and single polysome, respectively). χ_p, χ_n and χ_{np} are the Flory-Huggins

1204 interaction parameters, and κ_p, κ_n and κ_{np} are the interfacial tensions. We take $\kappa_{ij} = \lambda^2 \chi_{ij}$, with λ being a

1205 characteristic interface width.

1206

1207 The time evolution of the volume fractions is described by a combination of Cahn-Hilliard theory (Cahn and

1208 Hilliard, 1958) and the reactions kinetics of the polysomes. We assume that polysomes are produced inside

1209 the nucleoid at a constant rate k_1 and degraded uniformly along the cell at a rate k_{-1}:

1210

1211 $$\frac{\partial p}{\partial t} = \nabla(M_p \nabla \mu_p) + k_1 n - k_{-1} p , \quad \text{[eq. 5]}$$

1212 $$\frac{\partial n}{\partial t} = \nabla(M_n \nabla \mu_n), \qquad \text{[eq. 6]}$$

1213

1214 where $\mu_p = \frac{\delta F}{\delta p}$ and $\mu_n = \frac{\delta F}{\delta n}$ are the local chemical potentials of polysomes and nucleoid elements

1215 respectively, with $F = \int f[n(x), p(x)]dx$ representing the total free energy. $M_p = v_p D_p p$ and $M_n = v_n D_n n$

1216 are the mobility coefficients of the polysomes and the nucleoid, respectively, which depend on the diffusion

1217 coefficients of the polysomes (D_p) and the nucleoid (D_n). This recovers the Fick law of diffusion in the
1218 noninteracting limit.

1219

Since *E. coli* grows by elongation while maintaining its cylindrical shape and cell width, we reduce the
problem to one dimension, with x representing the position along the long axis of the cell. We assume that
the cell length grows exponentially such that the length at time t is $L(t) = L(0)e^{\gamma t}$, with γ being the growth
rate and $L(0)$ the initial cell length at birth. Exponential growth dilutes the existing polysomes with rate γ.
On the other hand, we assume that the nucleoid is not diluted since DNA replication occurs continuously,
and the nucleoid length is known to grow in proportion to cell length (Gray et al., 2019). Letting $\tilde{x} = x/L(t)$
be the relative position along the cell's long axis, the time evolution is given by:

1227

$$\frac{\partial p}{\partial t} = \partial_{\tilde{x}}\big(\widetilde{M}_p \partial_{\tilde{x}} \mu_p\big) + k_1 n - (k_{-1} + \gamma)p, \qquad \text{[eq. 7]}$$

1228

$$\frac{\partial n}{\partial t} = \partial_{\tilde{x}}\big(\widetilde{M}_n \partial_{\tilde{x}} \mu_n\big), \qquad\qquad\qquad \text{[eq. 8]}$$

1229

1230

1231 where $\partial_{\tilde{x}} = \frac{\partial}{\partial \tilde{x}} = L(t)\frac{\partial}{\partial x}$, and $\widetilde{M}_{n,p} = M_{n,p}L(t)^{-2}$. The chemical potentials are:

1232

$$\mu_p = v_p^{-1}\ln p - \ln(1 - p - n) + \chi_p(1 - 2p - n) + \big(\chi_{np} - \chi_n\big)n$$

1233

$$-\chi_p \tilde{\lambda}^2 \partial_{\tilde{x}}^2 (2p + n) + \big(\chi_{np} - \chi_n\big)\tilde{\lambda}^2 \partial_{\tilde{x}}^2 n, \qquad \text{[eq. 9]}$$

1234

1235

$$\mu_n = v_n^{-1}\ln n - \ln(1 - p - n) + \chi_n(1 - 2n - p) + \big(\chi_{np} - \chi_p\big)p$$

1236

$$-\chi_n \tilde{\lambda}^2 \partial_{\tilde{x}}^2 (2n + p) + \big(\chi_{np} - \chi_p\big)\tilde{\lambda}^2 \partial_{\tilde{x}}^2 p, \qquad \text{[eq. 10]}$$

1237

1238

1239 where $\tilde{\lambda} = \lambda/L(t)$.

1240

We take the non-dimensionalized Flory-Huggins interaction parameters to be $\chi_p = 0.2$, $\chi_n = 0.4$ and $\chi_{np} =$
1.2. The χ_{np} parameter captures the steric repulsion between the nucleoid and the polysomes (Miangolarra
et al., 2021). The diffusion coefficient of the polysome was set to $D_p = 0.015 \mu m^2/s$, which lies within the
same order of magnitude as was previously measured by single-molecule tracking (Bakshi et al., 2012).
Different diffusion coefficients were tested for the nucleoid ($D_n = 0.0003, 0.0005, 0.001$ or $0.005 \,\mu m^2/s$) to
capture different levels of nucleoid stiffness and hence response time to the local changes in polysome
concentration. The polysome degradation rate was set to $k_{-1} = 0.003 s^{-1}$, which corresponds to a half-life
of around 5 min (Bernstein et al., 2004). We take the polysome production rate to be $k_1 = k_{1,0}\left(1 + \frac{\gamma}{k_{-1}}\right)$,
with $k_{1,0} = 0.002 s^{-1}$, to match the observation that the nucleoid length is proportional to cell length (Gray
et al., 2019). The initial cell length $L(0)$ matched the cell length at birth, which has been previously shown
to increase exponentially with the growth rate (Govers et al., 2024). Hence, each of the simulated growth

rates $(\gamma = 0.25, 0.4, 0.55, 0.7, 0.85$ or $1\,\text{h}^{-1})$ was matched to a cell length at birth (Figure 4 – figure

supplement 1A) by fitting a linear regression $(L(0) = l_0 e^{\gamma/\gamma_0}$, with fitted parameters $l_0 = 1.61\mu\text{m}$ and $\gamma_0 =$

$1.84\text{h}^{-1})$ to the experimental data (Govers et al., 2024). Other parameters are chosen as follows: $\lambda =$

$0.03\mu\text{m}$, $v_n = 10$, $v_p = 5$.

1256

The dynamic equations (eqs. 7-8) were solved numerically in a fixed 1D domain $\tilde{x} \in (0,1)$, with no-flux

boundary conditions. Space is discretized into $N = 128$ grid points, and time is discretized to steps of Δt.

Time stepping was implemented with an implicit-explicit scheme as previously demonstrated (Mao et al.,

2020, 2019). The simulations were initialized from the (symmetric) steady-state solution of the system at

the initial cell length $L(0)$ (Figure 4A, dashed curves in Figure 4D, and Movie S3), or using the (asymmetric)

macromolecular distribution of the predivisional cell of the former simulation as the initial condition (Figure

4C, solid curves in Figure 4D) which allowed us to capture cell polarity (new versus old pole). The cell

length $L(t)$ was updated and recorded at each time step during the simulation. The nucleoid splitting time

(shown in Figure 4C) was determined by thresholding the relative decrease of the nucleoid concentration

at the cell center (as shown in in Figure 4 – figure supplement 1B). The code for the model is available at

https://github.com/qiweiyuu/polysome.

1268

Polysome and nucleoid analysis based on fitted Gaussian functions

Quantification of polysome and nucleoid asymmetries within cells along the cell division cycle was achieved

by fitting Gaussian functions (Python least squares optimization: *scipy.optimize.curve_fit*) to the RplA-GFP

or the HupA-mCherry intensity profiles of cells growing in M9gluCAAT in the microfluidic device. Three

Gaussian functions were fitted to the polysome signal concentration early in the cell division cycle. Two of

them captured the nucleoid-excluded polysomes accumulating at the poles and the third captured the

polysome accumulation in the middle of nucleoid (Figure 3 – figure supplement 1A-B):

1276

1277 $$Poly(l) = A_{old} \exp\left(-\frac{1}{2}\left(\frac{l_{old}-l}{\sigma_{old}}\right)^2\right) + A_{mid} \exp\left(-\frac{1}{2}\left(\frac{l_{mid}-l}{\sigma_{mid}}\right)^2\right) + A_{new} \exp\left(-\frac{1}{2}\left(\frac{l_{new}-l}{\sigma_{new}}\right)^2\right) \quad \text{[eq. 11]},$$

1278

1279   where A_{old}, A_{new} and A_{mid} are the amplitudes, l_{old}, l_{new} and l_{mid} are the means, and σ_{old}, σ_{new} and σ_{mid} are the

1280   standard deviations of the respective fitted functions along the relative cell length (l).

1281

1282   Two Gaussian functions were fitted to the concentration of constricting nucleoids (Figure 3 – figure

1283   supplement 1A-B):

1284

1285 $$Nuc(l) = A_{old} \exp\left(-\frac{1}{2}\left(\frac{l_{old}-l}{\sigma_{old}}\right)^2\right) + A_{new} \exp\left(-\frac{1}{2}\left(\frac{l_{new}-l}{\sigma_{new}}\right)^2\right) \quad \text{[eq. 12]}$$

1286

All Gaussian functions were fitted over the cell background fluorescence, which corresponds to the baseline

fluorescence within the masked cell boundaries and along the relative cell length coordinates from the old

pole (-1) via the cell center (0) to the new pole (1). The Gaussian functions were fitted either to single

newborn cells (0-2.5% into the cell division cycle) as shown in Figure 3 – figure supplement 1A-B and as

applied in Figures 3A and 3E, or to the less noisy average intensity profile of the segmented cells within the

0-10% range of their cell division cycle, as shown in Figure 3 – figure supplement 1C-D and used in Figures

3B and 3F. The areas of the Gaussian functions fitted to the polysomes (used in Figure 3A-B) were

estimated using the function:

1295
1296
$$Area = A\, \sigma\, \sqrt{2\pi} \quad \text{[eq. 13]},$$
1297
1298 where A is the amplitude and σ *is* the standard deviation.

1299

1300 The Gaussian areas were used to quantify the polysome asymmetries between the poles:

1301
1302
$$Poly_{asym} = log_{10}\left(\frac{Area_{new}}{Area_{old}}\right) \quad \text{[eq. 14]},$$
1303
1304
1305 where $Area_{new}$ and $Area_{old}$ are the areas of the fitted Gaussians to the nucleoid-excluded polysomes

1306 accumulating in the new and the old pole, respectively. The statistic in eq. 14 corresponds to the x-axis in

1307 Figure 3B. It is a positive value when the new pole contains more polysomes than the old one, and a

1308 negative value when the old pole contains more polysomes than the new one.

1309

1310 The nucleoid Gaussian statistics were used to quantify the nucleoid position asymmetry defined as:

1311
1312
1313
$$Nuc_{pos} = \frac{l_{new} + l_{old}}{2}\, \frac{l_{cell}}{2} \quad \text{[eq. 15]},$$
1314
1315 and the nucleoid compaction asymmetry was calculated as:

1316
1317
$$Nuc_{comp} = \frac{A_{new}}{A_{old}} \quad \text{[eq. 16]},$$
1318
where l_{old} and l_{new} are the means of the nucleoid Gaussians in relative cell coordinates (as in eq. 12) and

l_{cell} is the cell length in μm. The nucleoid position was first estimated in relative cell coordinates from the old

pole (-1) via the cell center (0) to the new pole (1), and then multiplied by half the cell length to convert to

absolute cell length units. A_{old} and A_{new} are the amplitudes of the fitted nucleoid Gaussians (as in eq. 12) in

arbitrary fluorescence units. The nucleoid position corresponds to the mid-point between the two Gaussian

means (eq. 15) and is a positive number when the nucleoid is shifted toward the new pole and a negative

number when it is positioned closer to the old pole (y-axis in Figure 3B).

1326

1327 The nucleoid compaction asymmetry was calculated as the relative nucleoid Gaussian amplitude (eq. 16).

1328 It had a positive value greater than 1 when the nucleoid concentration was higher toward the new pole and

1329 had a positive value smaller than 1 if the nucleoid was more concentrated toward the old pole (y-axis in

1330 Figure 3F).

1331

1332 To determine which polysome Gaussian statistics and related asymmetries may contribute to nucleoid
1333 compaction asymmetry (eq. 16), a linear mixed-effects model was used (*statsmodel* package in Python,
1334 *statsmodels.formula.api.mixedlm* function). The polysome Gaussian statistics were used as independent
1335 variables and the nucleoid compaction asymmetry as a dependent variable:

1336

1337

1338
$$Nuc_{comp} = \beta_0 + \beta_1 l_{old} + \beta_2 l_{mid} + \beta_3 l_{new} + \beta_4 \sigma_{old} + \beta_5 \sigma_{mid}$$
1339
$$+\beta_6 \sigma_{new} + \beta_7 A_{old} + \beta_8 A_{mid} + \beta_9 A_{new} + \varepsilon \qquad \text{[eq. 17]},$$

1340

where β_0 is the constant (intercept), β_n *(n=1,2...,9)* are the coefficients for each of the polysome Gaussian
statistics (same as in eq. 4 and plotted in Figure 3 – figure supplement 1E-F), and ε is the error. The entire
dataset was treated as a single group. Before fitting the linear mixed-effects model, all data were normalized
by subtracting the average and dividing the resulting value by their standard deviation (Figure 3 – figure
supplement 1F). The relative cell length positions of the polysomes at the old pole (l_{old}) and in the middle of
the nucleoid (l_{mid}) were found to significantly influence the asymmetric nucleoid compaction (Figure 3 –
figure supplement 1G). We reasoned that the available spaces between the polysome positions (Figure 3
– figure supplement 1E) may influence the compaction of the nucleoid and devised a new statistic that
compares the available polysome-free space toward the new pole to that of toward the old pole:

1350

1351
$$Poly_{space} = \frac{l_{new} - l_{mid}}{l_{mid} - l_{old}} \qquad \text{[eq. 18]},$$

1352
1353 where l_{new}, l_{mid} and l_{old} are the polysome Gaussian means (as in eq. 11), which represent the positions of
1354 the polysomes at the new pole, mid-cell region, and old pole, respectively, expressed as relative cell
1355 coordinates from the old pole (-1) via the cell center (0) to the new pole (1). The fraction above (eq. 18) has
1356 a value greater than 1 when the polysome-free space is greater toward the new pole and a value smaller
1357 than 1 when the polysome-free space is greater toward the old pole (x-axis in Figure 3F).

1358

Linear regressions
Two types of linear regressions were used in this work. In Figure 2B, a first-degree polynomial was fitted
using the ordinary least squares method and the numpy library in Python (*numpy.polyfit*), always using the
x-axis as the independent variable and the *y*-axis as the dependent variable. In Figures 1F and 3B, a
principal component regression was fitted using the scikit-learn library in Python
(*sklearn.decomposition.PCA*). A principal component analysis was first applied on the two-dimensional z-
transformed data to find the linear regressor that explained most of the variance (the first principal
component). This linear regressor was then rescaled to the original 2D plane. The principal component
regression eliminates the dependent variable bias associated with a traditional univariate fit. This bias is
introduced when calculating the prediction error (sum of squares) only along the dependent variable axis
during the least squares optimization. The elimination of this univariate bias is particularly important for the

fitted regressions in Figures 1F and 3B since their parameters were used to estimate the rate of nucleoid

splitting in the absence of polysome accumulation in the middle (y-intercept in Figure 1F), the position of

the nucleoid with equal polysomes at the poles (y-intercept in Figure 3B), or the polar asymmetry required

for a centered nucleoid (x-intercept in Figure 3B).

1374

1375   **Kernel density estimations**

1376   One-dimensional (Figures 3A, E, and Figure 1- figure supplement 2C) or two-dimensional (Figures 1F, 1H,

1377   Figure 1 – figure supplement 3A-B, and Figure 3 – figure supplement 1F) Gaussian kernel density

1378   estimations were fitted using the *scipy.stats.gaussian_kde* method. The bandwidth was determined using

1379   Scott's rule (Scott, 1992).

1380

1381   **Correlation coefficient calculations**

1382   All Spearman correlation coefficients (referred to as Spearman ρ) were estimated using the *scipy.stats*

1383   Python library and the *spearmanr* function. Only those Spearman correlation coefficients with a p-value

1384   below 0.05 are shown. Otherwise, the correlation is annotated as not-significant (NS) (as in Figure 1 –

1385   figure supplement 3B).

1386

1387   **ADDITIONAL RESOURCES**

1388   This study did not generate any additional resources.

1389

ACKNOWLEDGMENTS

We are grateful to Dr. A Janakiraman (City College of New York) for the pER12 (pBAD322A-gfp-μNS)

plasmid, Dr K. Prather (Massachusetts Institute of Technology) for the MG1655 (DE3) strain, and the

members of the Jacobs-Wagner laboratory for fruitful discussions and critical reading of the manuscript.

This work was part of the research program Rubicon Science 2018-1 with project number 019.181EN.018,

which was financed by the Netherlands Organization for Scientific Research (NWO). C.J.W. is an

investigator of Howard Hughes Medical Institute.

1397

AUTHOR CONTRIBUTIONS

Conceptualization, A.P. and C.J.-W.; methodology, A.P.; software, A.P. and Q.Y.; formal analysis, A.P.;

investigation, A.P., Q.Y, and S.K.G.; resources, A.P., Q.Y., and W.-H.L.; data curation, A.P.; writing –

original draft, A.P. and C.J.-W.; writing – review and editing, A.P., C.J.-W , N. W., Q.Y., S.K.G., and W.-

H.L; visualization, A.P. and C.J.-W.; supervision, N.W. and C.J.-W.; project administration, C.J.-W.; funding

acquisition, A.P., N.W. and C.J.-W.

1404

1405   **DECLARATION OF INTERESTS**

1406   The authors declare no competing interests.

REFERENCES

Azam, T.A., Hiraga, S., Ishihama, A., 2000. Two types of localization of the DNA-binding proteins within the Escherichia coli nucleoid: Intracellular localization of E. coli DNA-binding proteins. Genes Cells 5, 613–626. https://doi.org/10.1046/j.1365-2443.2000.00350.x

Bakshi, S., Choi, H., Mondal, J., Weisshaar, J.C., 2014. Time-dependent effects of transcription- and translation-halting drugs on the spatial distributions of the *E scherichia coli* chromosome and ribosomes: Time-dependent drug effects on *E. coli* chromosome spatial distribution. Mol. Microbiol. 94, 871–887. https://doi.org/10.1111/mmi.12805

Bakshi, S., Choi, H., Weisshaar, J.C., 2015. The spatial biology of transcription and translation in rapidly growing Escherichia coli. Front. Microbiol. 6. https://doi.org/10.3389/fmicb.2015.00636

Bakshi, S., Siryaporn, A., Goulian, M., Weisshaar, J.C., 2012. Superresolution imaging of ribosomes and RNA polymerase in live Escherichia coli cells: Superresolution imaging of ribosome and RNAP. Mol. Microbiol. 85, 21–38. https://doi.org/10.1111/j.1365-2958.2012.08081.x

Balakrishnan, R., Mori, M., Segota, I., Zhang, Z., Aebersold, R., Ludwig, C., Hwa, T., 2022. Principles of gene regulation quantitatively connect DNA to RNA and proteins in bacteria. Science 378, eabk2066. https://doi.org/10.1126/science.abk2066

Bartosik, A.A., Mierzejewska, J., Thomas, C.M., Jagura-Burdzy, G., 2009. ParB deficiency in Pseudomonas aeruginosa destabilizes the partner protein ParA and affects a variety of physiological parameters. Microbiology 155, 1080–1092. https://doi.org/10.1099/mic.0.024661-0

Bates, D., Kleckner, N., 2005. Chromosome and Replisome Dynamics in E. coli: Loss of Sister Cohesion Triggers Global Chromosome Movement and Mediates Chromosome Segregation. Cell 121, 899–911. https://doi.org/10.1016/j.cell.2005.04.013

Bean, G.J., Flickinger, S.T., Westler, W.M., McCully, M.E., Sept, D., Weibel, D.B., Amann, K.J., 2009. A22 Disrupts the Bacterial Actin Cytoskeleton by Directly Binding and Inducing a Low-Affinity State in MreB. Biochemistry 48, 4852–4857. https://doi.org/10.1021/bi900014d

Belliveau, N.M., Chure, G., Hueschen, C.L., Garcia, H.G., Kondev, J., Fisher, D.S., Theriot, J.A., Phillips, R., 2021. Fundamental limits on the rate of bacterial growth and their influence on proteomic composition. Cell Syst. 12, 924-944.e2. https://doi.org/10.1016/j.cels.2021.06.002

Bernstein, J.A., Lin, P.-H., Cohen, S.N., Lin-Chao, S., 2004. Global analysis of *Escherichia coli* RNA degradosome function using DNA microarrays. Proc. Natl. Acad. Sci. 101, 2758–2763. https://doi.org/10.1073/pnas.0308747101

Blundell, M.R., Wild, D.G., 1971. Altered ribosomes after inhibition of *Escherichia coli* by rifampicin. Biochem. J. 121, 391–398. https://doi.org/10.1042/bj1210391

Brandt, F., Etchells, S.A., Ortiz, J.O., Elcock, A.H., Hartl, F.U., Baumeister, W., 2009. The Native 3D Organization of Bacterial Polysomes. Cell 136, 261–271. https://doi.org/10.1016/j.cell.2008.11.016

Broering, T.J., Arnold, M.M., Miller, C.L., Hurt, J.A., Joyce, P.L., Nibert, M.L., 2005. Carboxyl-Proximal Regions of Reovirus Nonstructural Protein µNS Necessary and Sufficient for

Forming Factory-Like Inclusions. J. Virol. 79, 6194–6206.
https://doi.org/10.1128/JVI.79.10.6194-6206.2005

Broering, T.J., Parker, J.S.L., Joyce, P.L., Kim, J., Nibert, M.L., 2002. Mammalian Reovirus Nonstructural Protein µNS Forms Large Inclusions and Colocalizes with Reovirus Microtubule-Associated Protein µ2 in Transfected Cells. J. Virol. 76, 8285–8297. https://doi.org/10.1128/JVI.76.16.8285-8297.2002

Bryksin, A.V., Matsumura, I., 2010. Overlap extension PCR cloning: a simple and reliable way to create recombinant plasmids. BioTechniques 48, 463–465. https://doi.org/10.2144/000113418

Cahn, J.W., Hilliard, J.E., 1958. Free Energy of a Nonuniform System. I. Interfacial Free Energy. J. Chem. Phys. 28, 258–267. https://doi.org/10.1063/1.1744102

Campbell, E.A., Korzheva, N., Mustaev, A., Murakami, K., Nair, S., Goldfarb, A., Darst, S.A., 2001. Structural Mechanism for Rifampicin Inhibition of Bacterial RNA Polymerase. Cell 104, 901–912. https://doi.org/10.1016/S0092-8674(01)00286-0

Campos, M., Govers, S.K., Irnov, I., Dobihal, G.S., Cornet, F., Jacobs-Wagner, C., 2018. Genomewide phenotypic analysis of growth, cell morphogenesis, and cell cycle events in *Escherichia coli*. Mol. Syst. Biol. 14, e7573. https://doi.org/10.15252/msb.20177573

Campos, M., Surovtsev, I.V., Kato, S., Paintdakhi, A., Beltran, B., Ebmeier, S.E., Jacobs-Wagner, C., 2014. A constant size extension drives bacterial cell size homeostasis. Cell 159, 1433–1446. https://doi.org/10.1016/j.cell.2014.11.022

Cass, J.A., Kuwada, N.J., Traxler, B., Wiggins, P.A., 2016. Escherichia coli Chromosomal Loci Segregate from Midcell with Universal Dynamics. Biophys. J. 110, 2597–2609. https://doi.org/10.1016/j.bpj.2016.04.046

Castellana, M., Hsin-Jung Li, S., Wingreen, N.S., 2016. Spatial organization of bacterial transcription and translation. Proc. Natl. Acad. Sci. 113, 9286–9291. https://doi.org/10.1073/pnas.1604995113

Chai, Q., Singh, B., Peisker, K., Metzendorf, N., Ge, X., Dasgupta, S., Sanyal, S., 2014. Organization of Ribosomes and Nucleoids in Escherichia coli Cells during Growth and in Quiescence. J. Biol. Chem. 289, 11342–11352. https://doi.org/10.1074/jbc.M114.557348

Chao, L., Chen, C.K., Shi, C., Rang, C.U., 2023. Spatial and temporal distribution of ribosomes in single cells reveals aging differences between old and new daughters of *Escherichia coli* (preprint). Evolutionary Biology. https://doi.org/10.1101/2023.06.21.545935

Charaka, V.K., Misra, H.S., 2012. Functional Characterization of the Role of the Chromosome I Partitioning System in Genome Segregation in Deinococcus radiodurans. J. Bacteriol. 194, 5739–5748. https://doi.org/10.1128/JB.00610-12

Cherepanov, P.P., Wackernagel, W., 1995. Gene disruption in Escherichia coli: TcR and KmR cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance determinant. Gene 158, 9–14. https://doi.org/10.1016/0378-1119(95)00193-A

Chure, G., Cremer, J., 2023. An optimal regulation of fluxes dictates microbial growth in and out of steady state. eLife 12, e84878. https://doi.org/10.7554/eLife.84878

Cooper, S., Hsieh, M.L., 1988. The Rate and Topography of Cell Wall Synthesis during the Division Cycle of Escherichia coli Using N-Acetylglucosamine as a Peptidoglycan Label. Microbiology 134, 1717–1721. https://doi.org/10.1099/00221287-134-6-1717

Coquel, A.-S., Jacob, J.-P., Primet, M., Demarez, A., Dimiccoli, M., Julou, T., Moisan, L., Lindner,
A.B., Berry, H., 2013. Localization of Protein Aggregation in Escherichia coli Is Governed
by Diffusion and Nucleoid Macromolecular Crowding Effect. PLoS Comput. Biol. 9,
e1003038. https://doi.org/10.1371/journal.pcbi.1003038

Cutler, K.J., Stringer, C., Lo, T.W., Rappez, L., Stroustrup, N., Brook Peterson, S., Wiggins, P.A.,
Mougous, J.D., 2022. Omnipose: a high-precision morphology-independent solution for
bacterial cell segmentation. Nat. Methods 19, 1438–1448.
https://doi.org/10.1038/s41592-022-01639-4

Dai, X., Zhu, M., Warren, M., Balakrishnan, R., Patsalo, V., Okano, H., Williamson, J.R., Fredrick,
K., Wang, Y.-P., Hwa, T., 2017. Reduction of translating ribosomes enables Escherichia
coli to maintain elongation rates during slow growth. Nat. Microbiol. 2, 16231.
https://doi.org/10.1038/nmicrobiol.2016.231

Danilova, O., Reyes-Lamothe, R., Pinskaya, M., Sherratt, D., Possoz, C., 2007. MukB colocalizes
with the oriC region and is required for organization of the two Escherichia coli
chromosome arms into separate cell halves. Mol. Microbiol. 65, 1485–1492.
https://doi.org/10.1111/j.1365-2958.2007.05881.x

Datsenko, K.A., Wanner, B.L., 2000. One-step inactivation of chromosomal genes in *Escherichia*
coli K-12 using PCR products. Proc. Natl. Acad. Sci. 97, 6640–6645.
https://doi.org/10.1073/pnas.120163297

Davis, J.H., Rubin, A.J., Sauer, R.T., 2011. Design, construction and characterization of a set of
insulated bacterial promoters. Nucleic Acids Res. 39, 1131–1141.
https://doi.org/10.1093/nar/gkq810

Donczew, M., Mackiewicz, P., Wróbel, A., Flärdh, K., Zakrzewska-Czerwińska, J., Jakimowicz, D.,
2016. ParA and ParB coordinate chromosome segregation with cell elongation and
division during *Streptomyces* sporulation. Open Biol. 6, 150263.
https://doi.org/10.1098/rsob.150263

Donovan, C., Schwaiger, A., Krämer, R., Bramkamp, M., 2010. Subcellular Localization and
Characterization of the ParAB System from *Corynebacterium glutamicum*. J. Bacteriol.
192, 3441–3451. https://doi.org/10.1128/JB.00214-10

Dourado, H., Lercher, M.J., 2020. An analytical theory of balanced cellular growth. Nat.
Commun. 11, 1226. https://doi.org/10.1038/s41467-020-14751-w

Du, W.-L., Dubarry, N., Passot, F.M., Kamgoué, A., Murray, H., Lane, D., Pasta, F., 2016. Orderly
Replication and Segregation of the Four Replicons of Burkholderia cenocepacia J2315.
PLOS Genet. 12, e1006172. https://doi.org/10.1371/journal.pgen.1006172

Dworkin, J., Losick, R., 2002. Does RNA polymerase help drive chromosome segregation in
bacteria? Proc. Natl. Acad. Sci. 99, 14089–14094.
https://doi.org/10.1073/pnas.182539899

Dworsky, P., Schaechter, M., 1973. Effect of Rifampin on the Structure and Membrane
Attachment of the Nucleoid of *Escherichia coli*. J. Bacteriol. 116, 1364–1374.
https://doi.org/10.1128/jb.116.3.1364-1374.1973

Figge, R.M., Easter, J., Gober, J.W., 2003. Productive interaction between the chromosome
partitioning proteins, ParA and ParB, is required for the progression of the cell cycle in
Caulobacter crescentus: Dominant-negative ParB alleles. Mol. Microbiol. 47, 1225–1237.
https://doi.org/10.1046/j.1365-2958.2003.03367.x

Fisher, J.K., Bourniquel, A., Witz, G., Weiner, B., Prentiss, M., Kleckner, N., 2013. Four-
Dimensional Imaging of E. coli Nucleoid Organization and Dynamics in Living Cells. Cell
153, 882–895. https://doi.org/10.1016/j.cell.2013.04.006
Flory, P.J., 1942. Thermodynamics of High Polymer Solutions. J. Chem. Phys. 10, 51–61.
https://doi.org/10.1063/1.1723621
Forchhammer, J., Lindahl, L., 1971. Growth rate of polypeptide chains as a function of the cell
growth rate in a mutant of Escherichia coli 15. J. Mol. Biol. 55, 563–568.
https://doi.org/10.1016/0022-2836(71)90337-8
Gibson, D.G., Young, L., Chuang, R.-Y., Venter, J.C., Hutchison, C.A., Smith, H.O., 2009.
Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat. Methods 6,
343–345. https://doi.org/10.1038/nmeth.1318
Gillies, Sean, van der Wel, Casper, Van den Bossche, Joris, Taves, Mike W., Arnott, Joshua,
Ward, Brendan C., Others, 2023. Shapely. https://doi.org/10.5281/ZENODO.5597138
Govers, S.K., Campos, M., Tyagi, B., Laloux, G., Jacobs-Wagner, C., 2024. Apparent simplicity and
emergent robustness in the control of the Escherichia coli cell cycle. Cell Syst. 15, 19-
36.e5. https://doi.org/10.1016/j.cels.2023.12.001
Gray, A.N., Egan, A.J., Van'T Veer, I.L., Verheul, J., Colavin, A., Koumoutsi, A., Biboy, J., Altelaar,
A.F.M., Damen, M.J., Huang, K.C., Simorre, J.-P., Breukink, E., Den Blaauwen, T., Typas,
A., Gross, C.A., Vollmer, W., 2015. Coordination of peptidoglycan synthesis and outer
membrane constriction during Escherichia coli cell division. eLife 4, e07118.
https://doi.org/10.7554/eLife.07118
Gray, W.T., Govers, S.K., Xiang, Y., Parry, B.R., Campos, M., Kim, S., Jacobs-Wagner, C., 2019.
Nucleoid Size Scaling and Intracellular Organization of Translation across Bacteria. Cell
177, 1632-1648.e20. https://doi.org/10.1016/j.cell.2019.05.017
Guyer, M.S., Reed, R.R., Steitz, J.A., Low, K.B., 1981. Identification of a Sex-factor-affinity Site in
E. coli as. Cold Spring Harb. Symp. Quant. Biol. 45, 135–140.
https://doi.org/10.1101/SQB.1981.045.01.022
Guzman, L.M., Belin, D., Carson, M.J., Beckwith, J., 1995. Tight regulation, modulation, and
high-level expression by vectors containing the arabinose PBAD promoter. J. Bacteriol.
177, 4121–4130. https://doi.org/10.1128/jb.177.14.4121-4130.1995
Harju, J., Van Teeseling, M.C.F., Broedersz, C.P., 2024. Loop-extruders alter bacterial
chromosome topology to direct entropic forces for segregation. Nat. Commun. 15, 4618.
https://doi.org/10.1038/s41467-024-49039-w
Harju, J., Van Teeseling, M.C.F., Broedersz, C.P., 2023. Loop-extruders alter bacterial
chromosome topology to direct entropic forces for segregation (preprint). Biophysics.
https://doi.org/10.1101/2023.06.30.547230
Harris, C.R., Millman, K.J., Van Der Walt, S.J., Gommers, R., Virtanen, P., Cournapeau, D.,
Wieser, E., Taylor, J., Berg, S., Smith, N.J., Kern, R., Picus, M., Hoyer, S., Van Kerkwijk,
M.H., Brett, M., Haldane, A., Del Río, J.F., Wiebe, M., Peterson, P., Gérard-Marchant, P.,
Sheppard, K., Reddy, T., Weckesser, W., Abbasi, H., Gohlke, C., Oliphant, T.E., 2020.
Array programming with NumPy. Nature 585, 357–362.
https://doi.org/10.1038/s41586-020-2649-2

Hartmann, G., Honikel, K.O., Knüsel, F., Nüesch, J., 1967. The specific inhibition of the DNA-
directed RNA synthesis by rifamycin. Biochim. Biophys. Acta BBA - Nucleic Acids Protein
Synth. 145, 843–844. https://doi.org/10.1016/0005-2787(67)90147-5
Hofmann, A., Mäkelä, J., Sherratt, D.J., Heermann, D., Murray, S.M., 2019. Self-organised
segregation of bacterial chromosomal origins. eLife 8, e46564.
https://doi.org/10.7554/eLife.46564
Holmes, V.F., Cozzarelli, N.R., 2000. Closing the ring: Links between SMC proteins and
chromosome partitioning, condensation, and supercoiling. Proc. Natl. Acad. Sci. 97,
1322–1324. https://doi.org/10.1073/pnas.040576797
Hsu, T.-M., Chang, Y.-R., 2019. High-Copy-Number Plasmid Segregation—Single-Molecule
Dynamics in Single Cells. Biophys. J. 116, 772–780.
https://doi.org/10.1016/j.bpj.2019.01.019
Hu, X.-P., Dourado, H., Schubert, P., Lercher, M.J., 2020. The protein translation machinery is
expressed for maximal efficiency in Escherichia coli. Nat. Commun. 11, 5260.
https://doi.org/10.1038/s41467-020-18948-x
Huggins, M.L., 1941. Solutions of Long Chain Compounds. J. Chem. Phys. 9, 440–440.
https://doi.org/10.1063/1.1750930
Hunter, J.D., 2007. Matplotlib: A 2D Graphics Environment. Comput. Sci. Eng. 9, 90–95.
https://doi.org/10.1109/MCSE.2007.55
Ireton, K., Gunther, N.W., Grossman, A.D., 1994. spo0J is required for normal chromosome
segregation as well as the initiation of sporulation in Bacillus subtilis. J. Bacteriol. 176,
5320–5329. https://doi.org/10.1128/jb.176.17.5320-5329.1994
Iwai, N., Nagai, K., Wachi, M., 2002. Novel S -Benzylisothiourea Compound That Induces
Spherical Cells in Escherichia coli Probably by Acting on a Rod-shape-determining
Protein(s) Other Than Penicillin-binding Protein 2. Biosci. Biotechnol. Biochem. 66,
2658–2662. https://doi.org/10.1271/bbb.66.2658
Jacob, F., Brenner, S., Cuzin, F., 1963. On the Regulation of DNA Replication in Bacteria. Cold
Spring Harb. Symp. Quant. Biol. 28, 329–348.
https://doi.org/10.1101/SQB.1963.028.01.048
Jakimowicz, D., Brzostek, A., Rumijowska-Galewicz, A., Żydek, P., Dołzbłasz, A., Smulczyk-
Krawczyszyn, A., Zimniak, T., Wojtasz, Ł., Zawilak-Pawlik, A., Kois, A., Dziadek, J.,
Zakrzewska-Czerwińska, J., 2007a. Characterization of the mycobacterial chromosome
segregation protein ParB and identification of its target in Mycobacterium smegmatis.
Microbiology 153, 4050–4060. https://doi.org/10.1099/mic.0.2007/011619-0
Jakimowicz, D., Żydek, P., Kois, A., Zakrzewska-Czerwińska, J., Chater, K.F., 2007b. Alignment of
multiple chromosomes along helical ParA scaffolding in sporulating Streptomyces
hyphae. Mol. Microbiol. 65, 625–641. https://doi.org/10.1111/j.1365-
2958.2007.05815.x
Jalal, A.S.B., Le, T.B.K., 2020. Bacterial chromosome segregation by the ParABS system. Open
Biol. 10, 200097. https://doi.org/10.1098/rsob.200097
Jecz, P., Bartosik, A.A., Glabski, K., Jagura-Burdzy, G., 2015. A Single parS Sequence from the
Cluster of Four Sites Closest to oriC Is Necessary and Sufficient for Proper Chromosome
Segregation in Pseudomonas aeruginosa. PLOS ONE 10, e0120867.
https://doi.org/10.1371/journal.pone.0120867

Jensen, K.F., 1993. The Escherichia coli K-12 "wild types" W3110 and MG1655 have an rph
frameshift mutation that leads to pyrimidine starvation due to low pyrE expression
levels. J. Bacteriol. 175, 3401–3407. https://doi.org/10.1128/jb.175.11.3401-3407.1993
Jun, S., Mulder, B., 2006. Entropy-driven spatial organization of highly confined polymers:
Lessons for the bacterial chromosome. Proc. Natl. Acad. Sci. 103, 12388–12393.
https://doi.org/10.1073/pnas.0605305103
Jun, S., Wright, A., 2010. Entropy as the driver of chromosome segregation. Nat. Rev. Microbiol.
8, 600–607. https://doi.org/10.1038/nrmicro2391
Kadoya, R., Baek, J.H., Sarker, A., Chattoraj, D.K., 2011. Participation of Chromosome
Segregation Protein ParAI of *Vibrio cholerae* in Chromosome Replication. J. Bacteriol.
193, 1504–1514. https://doi.org/10.1128/JB.01067-10
Kawalek, A., Wawrzyniak, P., Bartosik, A.A., Jagura-Burdzy, G., 2020. Rules and Exceptions: The
Role of Chromosomal ParB in DNA Segregation and Other Cellular Processes.
Microorganisms 8, 105. https://doi.org/10.3390/microorganisms8010105
Kim, H.-J., Calcutt, M.J., Schmidt, F.J., Chater, K.F., 2000. Partitioning of the Linear Chromosome
during Sporulation of *Streptomyces coelicolor* A3(2) Involves an *oriC* -Linked *parAB*
Locus. J. Bacteriol. 182, 1313–1320. https://doi.org/10.1128/JB.182.5.1313-1320.2000
Kjos, M., Veening, J.-W., 2014. Tracking of chromosome dynamics in live *S treptococcus*
pneumoniae reveals that transcription promotes chromosome segregation:
Transcription promotes chromosome segregation. Mol. Microbiol. 91, 1088–1105.
https://doi.org/10.1111/mmi.12517
Koch, A.L., Gross, G.H., 1979. Growth Conditions and Rifampin Susceptibility. Antimicrob.
Agents Chemother. 15, 220–228. https://doi.org/10.1128/AAC.15.2.220
Koleva, K.Z., Hellweger, F.L., 2015. From protein damage to cell aging to population fitness in E.
coli: Insights from a multi-level agent-based model. Ecol. Model. 301, 62–71.
https://doi.org/10.1016/j.ecolmodel.2015.01.024
Kuwada, N.J., Cheveralls, K.C., Traxler, B., Wiggins, P.A., 2013. Mapping the driving forces of
chromosome structure and segregation in Escherichia coli. Nucleic Acids Res. 41, 7370–
7377. https://doi.org/10.1093/nar/gkt468
Lagage, V., Boccard, F., Vallet-Gely, I., 2016. Regional Control of Chromosome Segregation in
Pseudomonas aeruginosa. PLOS Genet. 12, e1006428.
https://doi.org/10.1371/journal.pgen.1006428
Łapińska, U., Glover, G., Capilla-Lasheras, P., Young, A.J., Pagliara, S., 2019. Bacterial ageing in
the absence of external stressors. Philos. Trans. R. Soc. B Biol. Sci. 374, 20180442.
https://doi.org/10.1098/rstb.2018.0442
Lee, P.S., Grossman, A.D., 2006. The chromosome partitioning proteins Soj (ParA) and Spo0J
(ParB) contribute to accurate chromosome partitioning, separation of replicated sister
origins, and regulation of replication initiation in Bacillus subtilis. Mol. Microbiol. 60,
853–869. https://doi.org/10.1111/j.1365-2958.2006.05140.x
Lemon, K.P., Grossman, A.D., 2000. Movement of Replicating DNA through a Stationary
Replisome. Mol. Cell 6, 1321–1330. https://doi.org/10.1016/S1097-2765(00)00130-1
Lewis, P.J., Thaker, S.D., Errington, J., 2000. Compartmentalization of transcription and
translation in Bacillus subtilis. EMBO J. 19, 710–718.
https://doi.org/10.1093/emboj/19.4.710

Lewis, R.A., Bignell, C.R., Zeng, W., Jones, A.C., Thomas, C.M., 2002. Chromosome loss from par
mutants of Pseudomonas putida depends on growth medium and phase of growth.
Microbiology 148, 537–548. https://doi.org/10.1099/00221287-148-2-537
Li, H., 2019. Random Chromosome Partitioning in the Polyploid Bacterium *Thermus*
thermophilus HB27. G3 GenesGenomesGenetics 9, 1249–1261.
https://doi.org/10.1534/g3.119.400086
Li, H., Angelov, A., Pham, V.T.T., Leis, B., Liebl, W., 2015. Characterization of chromosomal and
megaplasmid partitioning loci in Thermus thermophilus HB27. BMC Genomics 16, 317.
https://doi.org/10.1186/s12864-015-1523-3
Lim, H.C., Surovtsev, I.V., Beltran, B.G., Huang, F., Bewersdorf, J., Jacobs-Wagner, C., 2014.
Evidence for a DNA-relay mechanism in ParABS-mediated chromosome segregation.
eLife 3, e02758. https://doi.org/10.7554/eLife.02758
Lin, W.-H., Jacobs-Wagner, C., 2022. Connecting single-cell ATP dynamics to overflow
metabolism, cell growth, and the cell cycle in Escherichia coli. Curr. Biol. 32, 3911-
3924.e4. https://doi.org/10.1016/j.cub.2022.07.035
Lindner, A.B., Madden, R., Demarez, A., Stewart, E.J., Taddei, F., 2008. Asymmetric segregation
of protein aggregates is associated with cellular aging and rejuvenation. Proc. Natl.
Acad. Sci. 105, 3076–3081. https://doi.org/10.1073/pnas.0708931105
Livny, J., Yamaichi, Y., Waldor, M.K., 2007. Distribution of Centromere-Like *parS* Sites in
Bacteria: Insights from Comparative Genomics. J. Bacteriol. 189, 8693–8703.
https://doi.org/10.1128/JB.01239-07
Mäkelä, J., Papagiannakis, A., Lin, W.-H., Lanz, M.C., Glenn, S., Swaffer, M., Marinov, G.K.,
Skotheim, J.M., Jacobs-Wagner, C., 2024. Genome concentration limits cell growth and
modulates proteome composition in *Escherichia coli*.
https://doi.org/10.1101/2024.02.26.582181
Mäkelä, J., Uphoff, S., Sherratt, D.J., 2021. Nonrandom segregation of sister chromosomes by
Escherichia coli MukBEF. Proc. Natl. Acad. Sci. 118, e2022078118.
https://doi.org/10.1073/pnas.2022078118
Mao, S., Chakraverti-Wuerthwein, M.S., Gaudio, H., Košmrlj, A., 2020. Designing the
Morphology of Separated Phases in Multicomponent Liquid Mixtures. Phys. Rev. Lett.
125, 218003. https://doi.org/10.1103/PhysRevLett.125.218003
Mao, S., Kuldinow, D., Haataja, M.P., Košmrlj, A., 2019. Phase behavior and morphology of
multicomponent liquid mixtures. Soft Matter 15, 1297–1311.
https://doi.org/10.1039/C8SM02045K
McGuffee, S.R., Elcock, A.H., 2010. Diffusion, Crowding & Protein Stability in a Dynamic
Molecular Model of the Bacterial Cytoplasm. PLoS Comput. Biol. 6, e1000694.
https://doi.org/10.1371/journal.pcbi.1000694
McKinney, W., 2010. Data Structures for Statistical Computing in Python. Presented at the
Python in Science Conference, Austin, Texas, pp. 56–61.
https://doi.org/10.25080/Majora-92bf1922-00a
Miangolarra, A.M., Li, S.H.-J., Joanny, J.-F., Wingreen, N.S., Castellana, M., 2021. Steric
interactions and out-of-equilibrium processes control the internal organization of
bacteria. Proc. Natl. Acad. Sci. 118, e2106014118.
https://doi.org/10.1073/pnas.2106014118

Minnen, A., Attaiech, L., Thon, M., Gruber, S., Veening, J.-W., 2011. SMC is recruited to oriC by
ParB and promotes chromosome segregation in Streptococcus pneumoniae:
Chromosome segregation in S. pneumoniae. Mol. Microbiol. 81, 676–688.
https://doi.org/10.1111/j.1365-2958.2011.07722.x

Mohapatra, S., Weisshaar, J.C., 2018. Functional mapping of the *E. coli* translational machinery
using single-molecule tracking. Mol. Microbiol. 110, 262–282.
https://doi.org/10.1111/mmi.14103

Mohl, D.A., Gober, J.W., 1997. Cell Cycle–Dependent Polar Localization of Chromosome
Partitioning Proteins in Caulobacter crescentus. Cell 88, 675–684.
https://doi.org/10.1016/S0092-8674(00)81910-8

Molenaar, D., Van Berlo, R., De Ridder, D., Teusink, B., 2009. Shifts in growth strategies reflect
tradeoffs in cellular economics. Mol. Syst. Biol. 5, 323.
https://doi.org/10.1038/msb.2009.82

Mondal, J., Bratton, B.P., Li, Y., Yethiraj, A., Weisshaar, J.C., 2011. Entropy-Based Mechanism of
Ribosome-Nucleoid Segregation in E. coli Cells. Biophys. J. 100, 2605–2613.
https://doi.org/10.1016/j.bpj.2011.04.030

Navarro, P.P., Vettiger, A., Ananda, V.Y., Llopis, P.M., Allolio, C., Bernhardt, T.G., Chao, L.H.,
2022. Cell wall synthesis and remodelling dynamics determine division site architecture
and cell shape in Escherichia coli. Nat. Microbiol. 7, 1621–1634.
https://doi.org/10.1038/s41564-022-01210-z

Otsu, N., 1979. A Threshold Selection Method from Gray-Level Histograms. IEEE Trans. Syst.
Man Cybern. 9, 62–66. https://doi.org/10.1109/TSMC.1979.4310076

Paintdakhi, A., Parry, B., Campos, M., Irnov, I., Elf, J., Surovtsev, I., Jacobs-Wagner, C., 2016.
Oufti: an integrated software package for high-accuracy, high-throughput quantitative
microscopy analysis: Oufti: image analysis software. Mol. Microbiol. 99, 767–777.
https://doi.org/10.1111/mmi.13264

Pande, S., Mitra, D., Chatterji, A., 2023. Entropy mediated organization of E.coli chromosome in
fast growth conditions. https://doi.org/10.48550/ARXIV.2304.02275

Parry, B.R., Surovtsev, I.V., Cabeen, M.T., O'Hern, C.S., Dufresne, E.R., Jacobs-Wagner, C., 2014.
The Bacterial Cytoplasm Has Glass-like Properties and Is Fluidized by Metabolic Activity.
Cell 156, 183–194. https://doi.org/10.1016/j.cell.2013.11.028

Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z., Gimelshein,
N., Antiga, L., Desmaison, A., Köpf, A., Yang, E., DeVito, Z., Raison, M., Tejani, A.,
Chilamkurthy, S., Steiner, B., Fang, L., Bai, J., Chintala, S., 2019. PyTorch: An Imperative
Style, High-Performance Deep Learning Library.
https://doi.org/10.48550/ARXIV.1912.01703

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Müller, A., Nothman, J., Louppe, G., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas,
J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., Duchesnay, É., 2012. Scikit-learn:
Machine Learning in Python. https://doi.org/10.48550/ARXIV.1201.0490

Pelletier, J., Halvorsen, K., Ha, B.-Y., Paparcone, R., Sandler, S.J., Woldringh, C.L., Wong, W.P.,
Jun, S., 2012. Physical manipulation of the *Escherichia coli* chromosome reveals its soft
nature. Proc. Natl. Acad. Sci. 109. https://doi.org/10.1073/pnas.1208689109

Pettijohn, D.E., Hecht, R., 1974. RNA Molecules Bound to the Folded Bacterial Genome Stabilize
DNA Folds and Segregate Domains of Supercoiling. Cold Spring Harb. Symp. Quant. Biol.
38, 31–41. https://doi.org/10.1101/SQB.1974.038.01.006
Pountain, A.W., Jiang, P., Yao, T., Homaee, E., Guan, Y., Podkowik, M., Shopsin, B., Torres, V.J.,
Golding, I., Yanai, I., 2022. Transcription-replication interactions reveal principles of
bacterial genome regulation (preprint). Systems Biology.
https://doi.org/10.1101/2022.10.22.513359
Proenca, A.M., Rang, C.U., Qiu, A., Shi, C., Chao, L., 2019. Cell aging preserves cellular
immortality in the presence of lethal levels of damage. PLOS Biol. 17, e3000266.
https://doi.org/10.1371/journal.pbio.3000266
Reyes-Lamothe, R., Tran, T., Meas, D., Lee, L., Li, A.M., Sherratt, D.J., Tolmasky, M.E., 2014.
High-copy bacterial plasmids diffuse in the nucleoid-free space, replicate stochastically
and are randomly partitioned at cell division. Nucleic Acids Res. 42, 1042–1051.
https://doi.org/10.1093/nar/gkt918
Robinow, C., Kellenberger, E., 1994. The bacterial nucleoid revisited. Microbiol. Rev. 58, 211–
232. https://doi.org/10.1128/mr.58.2.211-232.1994
Sadhir, I., Murray, S.M., 2023. Mid-cell migration of the chromosomal terminus is coupled to
origin segregation in Escherichia coli. Nat. Commun. 14, 7489.
https://doi.org/10.1038/s41467-023-43351-7
Sanamrad, A., Persson, F., Lundius, E.G., Fange, D., Gynnå, A.H., Elf, J., 2014. Single-particle
tracking reveals that free ribosomal subunits are not excluded from the *Escherichia coli*
nucleoid. Proc. Natl. Acad. Sci. 111, 11413–11418.
https://doi.org/10.1073/pnas.1411558111
Santi, I., McKinney, J.D., 2015. Chromosome Organization and Replisome Dynamics in
Mycobacterium smegmatis. mBio 6, e01999-14. https://doi.org/10.1128/mBio.01999-14
Sawitzke, J.A., Austin, S., 2000. Suppression of chromosome segregation defects of *Escherichia*
coli muk mutants by mutations in topoisomerase I. Proc. Natl. Acad. Sci. 97, 1671–1676.
https://doi.org/10.1073/pnas.030528397
Schaechter, M., MaalOe, O., Kjeldgaard, N.O., 1958. Dependency on Medium and Temperature
of Cell Size and Chemical Composition during Balanced Growth of Salmonella
typhimurium. J. Gen. Microbiol. 19, 592–606. https://doi.org/10.1099/00221287-19-3-
592
Scholz, S.A., Diao, R., Wolfe, M.B., Fivenson, E.M., Lin, X.N., Freddolino, P.L., 2019. High-
Resolution Mapping of the Escherichia coli Chromosome Reveals Positions of High and
Low Transcription. Cell Syst. 8, 212-225.e9. https://doi.org/10.1016/j.cels.2019.02.004
Scott, D.W., 1992. Multivariate Density Estimation: Theory, Practice, and Visualization, 1st ed,
Wiley Series in Probability and Statistics. Wiley.
https://doi.org/10.1002/9780470316849
Scott, M., Klumpp, S., Mateescu, E.M., Hwa, T., 2014. Emergence of robust growth laws from
optimal regulation of ribosome synthesis. Mol. Syst. Biol. 10, 747.
https://doi.org/10.15252/msb.20145379
Seabold, S., Perktold, J., 2010. Statsmodels: Econometric and Statistical Modeling with Python.
Presented at the Python in Science Conference, Austin, Texas, pp. 92–96.
https://doi.org/10.25080/Majora-92bf1922-011

Shis, D.L., Bennett, M.R., 2013. Library of synthetic transcriptional AND gates built with split T7
RNA polymerase mutants. Proc. Natl. Acad. Sci. 110, 5028–5033.
https://doi.org/10.1073/pnas.1220157110
Si, F., Li, D., Cox, S.E., Sauls, J.T., Azizi, O., Sou, C., Schwartz, A.B., Erickstad, M.J., Jun, Y., Li, X.,
Jun, S., 2017. Invariance of Initiation Mass and Predictability of Cell Size in Escherichia
coli. Curr. Biol. 27, 1278–1287. https://doi.org/10.1016/j.cub.2017.03.022
Slayter, H., Kiho, Y., Hall, C.E., Rich, A., 1968. AN ELECTRON MICROSCOPIC STUDY OF LARGE
BACTERIAL POLYRIBOSOMES. J. Cell Biol. 37, 583–590.
https://doi.org/10.1083/jcb.37.2.583
Stylianidou, S., Brennan, C., Nissen, S.B., Kuwada, N.J., Wiggins, P.A., 2016. *SuperSegger* : robust
image segmentation, analysis and lineage tracking of bacterial cells. Mol. Microbiol. 102,
690–700. https://doi.org/10.1111/mmi.13486
Subach, O.M., Cranfill, P.J., Davidson, M.W., Verkhusha, V.V., 2011. An Enhanced Monomeric
Blue Fluorescent Protein with the High Chemical Stability of the Chromophore. PLoS
ONE 6, e28674. https://doi.org/10.1371/journal.pone.0028674
Subramanian, S., Murray, S.M., 2023. Subdiffusive movement of chromosomal loci in bacteria
explained by DNA bridging. Phys. Rev. Res. 5, 023034.
https://doi.org/10.1103/PhysRevResearch.5.023034
Takacs, C.N., Wachter, J., Xiang, Y., Ren, Z., Karaboja, X., Scott, M., Stoner, M.R., Irnov, I.,
Jannetty, N., Rosa, P.A., Wang, X., Jacobs-Wagner, C., 2022. Polyploidy, regular
patterning of genome copies, and unusual control of DNA partitioning in the Lyme
disease spirochete. Nat. Commun. 13, 7173. https://doi.org/10.1038/s41467-022-
34876-4
Thappeta, Y., Cañas-Duarte, S.J., Kallem, T., Fragasso, A., Xiang, Y., Gray, W., Lee, C., Cegelski, L.,
Jacobs-Wagner, C., 2024. Glycogen phase separation drives macromolecular
rearrangement and asymmetric division in *E. coli*.
https://doi.org/10.1101/2024.04.19.590186
Tiruvadi-Krishnan, S., Männik, Jaana, Kar, P., Lin, J., Amir, A., Männik, Jaan, 2022. Coupling
between DNA replication, segregation, and the onset of constriction in Escherichia coli.
Cell Rep. 38, 110539. https://doi.org/10.1016/j.celrep.2022.110539
Tseng, H.-C., Harwell, C.L., Martin, C.H., Prather, K.L., 2010a. Biosynthesis of chiral 3-
hydroxyvalerate from single propionate-unrelated carbon sources in metabolically
engineered E. coli. Microb. Cell Factories 9, 96. https://doi.org/10.1186/1475-2859-9-96
Tseng, H.-C., Harwell, C.L., Martin, C.H., Prather, K.L., 2010b. Biosynthesis of chiral 3-
hydroxyvalerate from single propionate-unrelated carbon sources in metabolically
engineered E. coli. Microb. Cell Factories 9, 96. https://doi.org/10.1186/1475-2859-9-96
Van Der Walt, S., Schönberger, J.L., Nunez-Iglesias, J., Boulogne, F., Warner, J.D., Yager, N.,
Gouillart, E., Yu, T., 2014. scikit-image: image processing in Python. PeerJ 2, e453.
https://doi.org/10.7717/peerj.453
Varricchio, F., Monier, R., 1971. Ribosome Patterns in *Escherichia coli* Growing at Various Rates.
J. Bacteriol. 108, 105–110. https://doi.org/10.1128/jb.108.1.105-110.1971
Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski,
E., Peterson, P., Weckesser, W., Bright, J., Van Der Walt, S.J., Brett, M., Wilson, J.,
Millman, K.J., Mayorov, N., Nelson, A.R.J., Jones, E., Kern, R., Larson, E., Carey, C.J.,

Polat, İ., Feng, Y., Moore, E.W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R.,
Henriksen, I., Quintero, E.A., Harris, C.R., Archibald, A.M., Ribeiro, A.H., Pedregosa, F.,
Van Mulbregt, P., SciPy 1.0 Contributors, Vijaykumar, A., Bardelli, A.P., Rothberg, A.,
Hilboll, A., Kloeckner, A., Scopatz, A., Lee, A., Rokem, A., Woods, C.N., Fulton, C.,
Masson, C., Häggström, C., Fitzgerald, C., Nicholson, D.A., Hagen, D.R., Pasechnik, D.V.,
Olivetti, E., Martin, E., Wieser, E., Silva, F., Lenders, F., Wilhelm, F., Young, G., Price, G.A.,
Ingold, G.-L., Allen, G.E., Lee, G.R., Audren, H., Probst, I., Dietrich, J.P., Silterra, J.,
Webber, J.T., Slavič, J., Nothman, J., Buchner, J., Kulick, J., Schönberger, J.L., De Miranda
Cardoso, J.V., Reimer, J., Harrington, J., Rodríguez, J.L.C., Nunez-Iglesias, J., Kuczynski, J.,
Tritz, K., Thoma, M., Newville, M., Kümmerer, M., Bolingbroke, M., Tartre, M., Pak, M.,
Smith, N.J., Nowaczyk, N., Shebanov, N., Pavlyk, O., Brodtkorb, P.A., Lee, P., McGibbon,
R.T., Feldbauer, R., Lewis, S., Tygier, S., Sievert, S., Vigna, S., Peterson, S., More, S.,
Pudlik, T., Oshima, T., Pingel, T.J., Robitaille, T.P., Spura, T., Jones, T.R., Cera, T., Leslie,
T., Zito, T., Krauss, T., Upadhyay, U., Halchenko, Y.O., Vázquez-Baeza, Y., 2020. SciPy 1.0:
fundamental algorithms for scientific computing in Python. Nat. Methods 17, 261–272.
https://doi.org/10.1038/s41592-019-0686-2

Wang, P., Robert, L., Pelletier, J., Dang, W.L., Taddei, F., Wright, A., Jun, S., 2010. Robust Growth
of Escherichia coli. Curr. Biol. 20, 1099–1103. https://doi.org/10.1016/j.cub.2010.04.045

Wang, X., Liu, X., Possoz, C., Sherratt, D.J., 2006. The two *Escherichia coli* chromosome arms
locate to separate cell halves. Genes Dev. 20, 1727–1731.
https://doi.org/10.1101/gad.388406

Waskom, M., 2021. seaborn: statistical data visualization. J. Open Source Softw. 6, 3021.
https://doi.org/10.21105/joss.03021

Weitao, T., Nordstrom, K., Dasgupta, S., 1999. Mutual suppression of mukB and seqA
phenotypes might arise from their opposing influences on the Escherichia coli nucleoid
structure. Mol. Microbiol. 34, 157–168. https://doi.org/10.1046/j.1365-
2958.1999.01589.x

Wientjes, F.B., Nanninga, N., 1989. Rate and topography of peptidoglycan synthesis during cell
division in Escherichia coli: concept of a leading edge. J. Bacteriol. 171, 3412–3419.
https://doi.org/10.1128/jb.171.6.3412-3419.1989

Wiktor, J., Gynnå, A.H., Leroy, P., Larsson, J., Coceano, G., Testa, I., Elf, J., 2021. RecA finds
homologous DNA by reduced dimensionality search. Nature 597, 426–429.
https://doi.org/10.1038/s41586-021-03877-6

Woldringh, C.L., Huls, P., Pas, E., Brakenhoff, G.J., Nanninga, N., 1987. Topography of
Peptidoglycan Synthesis during Elongation and Polar Cap Formation in a Cell Division
Mutant of Escherichia coli MC4100. Microbiology 133, 575–586.
https://doi.org/10.1099/00221287-133-3-575

Wu, F., Japaridze, A., Zheng, X., Wiktor, J., Kerssemakers, J.W.J., Dekker, C., 2019a. Direct
imaging of the circular chromosome in a live bacterium. Nat. Commun. 10, 2194.
https://doi.org/10.1038/s41467-019-10221-0

Wu, F., Swain, P., Kuijpers, L., Zheng, X., Felter, K., Guurink, M., Solari, J., Jun, S., Shimizu, T.S.,
Chaudhuri, D., Mulder, B., Dekker, C., 2019b. Cell Boundary Confinement Sets the Size
and Position of the E. coli Chromosome. Curr. Biol. 29, 2131-2144.e4.
https://doi.org/10.1016/j.cub.2019.05.015

Xiang, Y., Surovtsev, I.V., Chang, Y., Govers, S.K., Parry, B.R., Liu, J., Jacobs-Wagner, C., 2021.
Interconnecting solvent quality, transcription, and chromosome folding in Escherichia
coli. Cell 184, 3626-3642.e14. https://doi.org/10.1016/j.cell.2021.05.037
Yamaichi, Y., Fogel, M.A., McLeod, S.M., Hui, M.P., Waldor, M.K., 2007. Distinct Centromere-
Like *parS* Sites on the Two Chromosomes of *Vibrio* spp. J. Bacteriol. 189, 5314–5324.
https://doi.org/10.1128/JB.00416-07
Yang, D., Männik, Jaana, Retterer, S.T., Männik, Jaan, 2020. The effects of polydisperse
crowders on the compaction of the *Escherichia coli* nucleoid. Mol. Microbiol. 113, 1022–
1037. https://doi.org/10.1111/mmi.14467
Yao, S., Helinski, D.R., Toukdarian, A., 2007. Localization of the Naturally Occurring Plasmid
ColE1 at the Cell Pole. J. Bacteriol. 189, 1946–1953. https://doi.org/10.1128/JB.01451-
06
Yu, W., Herbert, S., Graumann, P.L., Götz, F., 2010. Contribution of SMC (Structural
Maintenance of Chromosomes) and SpoIIIE to Chromosome Segregation in
Staphylococci. J. Bacteriol. 192, 4067–4073. https://doi.org/10.1128/JB.00010-10
Zhang, C., Shao, P.G., Van Kan, J.A., Van Der Maarel, J.R.C., 2009. Macromolecular crowding
induced elongation and compaction of single DNA molecules confined in a nanochannel.
Proc. Natl. Acad. Sci. 106, 16651–16656. https://doi.org/10.1073/pnas.0904741106
Zhou, Z., Siddiquee, M.M.R., Tajbakhsh, N., Liang, J., 2020. UNet++: Redesigning Skip
Connections to Exploit Multiscale Features in Image Segmentation. IEEE Trans. Med.
Imaging 39, 1856–1867. https://doi.org/10.1109/TMI.2019.2959609
Zimmerman, S.B., Trach, S.O., 1991. Estimation of macromolecule concentrations and excluded
volume effects for the cytoplasm of Escherichia coli. J. Mol. Biol. 222, 599–620.
https://doi.org/10.1016/0022-2836(91)90499-V

1911
1912

FIGURE LEGENDS

Figure 1: Correlations between polysome and nucleoid dynamics at the single-cell level. CJW7323

cells were grown in in M9gluCAAT in a microfluidic device. **A.** Schematic illustrating observable nucleoid

segregation events. **B.** Fluorescence images of RplA-GFP and HupA-mCherry for a representative cell

(CJW7323) from birth to division. **C.** Ensemble kymographs of the average RplA-GFP and HupA-mCherry

fluorescence during the cell division cycle (>300,000 segmented cell instances from 4122 complete cell

division cycles). **D.** Two-dimensional projections of the average RplA-GFP and HupA-mCherry

fluorescence signals in predivisional cells (4564 cells with two nucleoid objects, from 1907 cell division

cycles, 95-100% into the cell division cycle) and their intensity profiles. White arrows indicate RplA-GFP

enrichments at the quarter cell positions, while the black arrow indicates the site of cell constriction. **E.** Plot

showing the dynamics of RplA-GFP accumulation and HupA-mCherry depletion at mid-nucleoid (median ±

IQR) during the nucleoid cycle (see Figure 1 – Figure supplement 2D). Data from 3240 nucleoid segregation

cycles are shown (40 nucleoid cycle bins, 2512-4823 segmented nucleoids per bin). **F.** Correlation

(Spearman ρ = -0.52, p-value < 10^{-10}) between the rate of RplA-GFP accumulation in the middle of the

nucleoid ($\frac{\Delta RplA_{mid-nuc} \; conc.}{\Delta T}$) and the rate of HupA-mCherry depletion in the same region ($\frac{\Delta HupA_{mid-nuc} \; conc.}{\Delta T}$)

between the initiation of nucleoid splitting and just before the end of nucleoid splitting (3214 complete

nucleoid cycles). The colormap indicates the Gaussian kernel density estimation (KDE). Solid black line

indicates the linear-regression fit to the data. **G.** Percentage of cells that continue to accumulate polysomes

in the middle of the cell ($\frac{\Delta RplA_{mid-cell\ conc.}}{\Delta T}$) during four relative time bins (1335 to 1957 cell division cycles

per bin) covering the period from the end of nucleoid splitting until cell division. **H.** Correlation (Spearman

$\rho = 0.47$, p-value $< 10^{-10}$) between the rate of RplA-GFP accumulation at mid-cell ($\frac{\Delta RplA_{mid-cell\ conc.}}{\Delta T}$) and the

rate of distance increase between the sister nucleoids ($\frac{\Delta Distance_{nuc}}{\Delta T}$) during the first quartile (0-25%) of the

period between the end of nucleoid splitting and cell division (1376 cell division cycles). The black markers

correspond to nine bins (mean $\pm$ SEM, 75 to 177 cell division cycles per bin) within the 5^{th}-95^{th} percentiles

of *x*-axis range. Also shown is the distribution of the cell elongation rates ($\frac{\Delta Length_{cell}}{\Delta T}$) during the same time

interval, with the mean and SD shown by the solid and dashed lines, respectively. **I.** Plot showing the

coefficients of a linear mixed-effects model (see eq. 3 in Methods and Figure 1 – figure supplement 3B) for

four interval bins between the completion of nucleoid splitting and cell division. The coefficients quantify the

relative contribution of polysome accumulation at mid-cell ($\frac{\Delta RplA_{mid-cell\ conc.}}{\Delta T}$) and cell elongation ($\frac{\Delta Length_{cell}}{\Delta T}$)

to the rate of sister nucleoid migration ($\frac{\Delta Distance_{nuc}}{\Delta T}$). All coefficients are significant (Prob($<|Z|$) $< 10^{-9}$), except

for the one marked with an asterisk that is marginally significant (Prob($<|Z|$) = 0.02).

1944

Figure 2: Correlation of the extent and relative timing of polysome accumulation with nucleoid

segregation at the population and single-cell levels. A. RplA-mEos2 and DAPI (scaled by the whole

cell average) demographs constructed from snapshots of DAPI-stained CJW6768 cells (815 to 2771 cells

per demograph) expressing RplA-mEos2 and growing in different nutrient conditions (see Table S1 for

abbreviations). The demographs were arranged from smallest (M9mann, top) to biggest average cell area

(M9malaCAAT, bottom). Additional demographs for different ribosomal reporters and nutrient conditions

are shown in Figure 2 – figure supplement 1. **B.** Correlation between average polysome accumulation and

average nucleoid depletion at mid-cell for all tested strains (Spearman ρ_{all} = -0.85, p-value $< 10^{-10}$) with

different ribosomal and nucleoid reporters, and within each strain (-0.75 $<$ Spearman ρ_{strain} $<$ -0.95, p-values

$< 10^{-3}$) across nutrient conditions. A linear regression was fitted to all the data. **D.** Plot showing the variability

in growth rate (GR) across cell division cycles for cells growing in microfluidics in M9gluCAAT (4114 cell

division cycles, 9 growth rate bins with 93 to 860 cell division cycles each). **E.** Ensemble kymographs of the

RplA-GFP and HupA-mCherry concentration normalized by the average fluorescence for the slowest,

intermediate and fastest growing population bins shown in panel D. **F.** Plot showing the correlation between

the average RplA-GFP accumulation and HupA-mCherry depletion at mid-cell across the growth rate bins

(mean $\pm$ SD, 93 to 860 cell division cycles per bin) shown in panel D.

1961

Figure 3: Correlations between polysome and nucleoid asymmetries. A. Distributions of RplA-GFP
concentration in the new (grey) and the old (black) pole regions of newborn cells (0-2.5% of the cell division
cycle, n = 912 cell division cycles). The histograms were smoothed using Gaussian kernel density
estimations. **B.** Correlation (Spearman ρ = -0.41, p-value < 10^{-10}) between the polar polysome asymmetry
and the position of the nucleoid centroid around the cell center of cells at the beginning of the division cycle
(0-10%, n = 1179 cell division cycles). The contour plot consists of 9 levels with a lower data density
threshold of 25%. The polar polysome profile for cells with correlations indicated by the numbers 1 and 2
are schematically illustrated in the next panel. A linear regression (solid grey line) was fitted to the data. **C.**
Schematic illustrating the effects of the relative polysome abundance between the poles on the position of
the nucleoid. **D.** Average 2D projections of the RplA-GFP and HupA-mCherry concentration (conc.) at
different cell division cycle intervals (~9440 to 47240 cell images per cell division cycle interval from 4122
cell division cycles). The dotted line indicates the boundary between two cell division cycles. **E.** Density plot
comparing the distribution of the HupA-mCherry maximum concentration toward the new pole (gray) to that
toward the old pole (black) in newborn cells (0-2.5% of the division cycle, n = 912 cell division cycles). The
histograms were smoothed using Gaussian kernel density estimations. **F.** Correlation (Spearman ρ = 0.52,
p-value<10^{-10}) between the nucleoid density asymmetry and the relative availability of polysome-free space
between the two cell halves early in the cell division cycle (0-10% of the division cycle, n = 2150 cell division
cycles). The contour plot consists of 9 levels with a lower data density threshold of 25%. Values above 1
on the *x*-axis indicate more polysome-free space toward the new pole, and values below 1 correspond to
cells with more polysome-free space toward the old pole. On the *y*-axis, values above 1 indicate higher
DNA density toward the new pole and values below 1 indicate higher DNA density toward the old pole. **G.**
Average 2D projections of newborn cells (0-10% into the cell division cycle) from the lower-left quartile in
panel C (region 2, n = 223 cell division cycles) and the upper right quartile in panel C (region 1, n = 557 cell
division cycles) and their 1D intensity profiles.
1986
Figure 4: Simulation results of the reaction-diffusion model for different growth rates or nucleoid
diffusion rates, and comparison of the nonequilibrium polysome accumulation with freely diffusing
particles. A. 1D simulation of polysome (blue) and nucleoid (red) dynamics during slow growth (growth
rate = 0.25 h^{-1}, D_n = 0.001 μm^2/s, cell length at birth = 1.84 μm) at different relative cell division cycle
timepoints. The simulation was initialized from the equilibrium polysome and nucleoid distribution (at 0%).
B. Schematic summarizing how polysomes accumulate in the middle of the elongating nucleoid, causing
nucleoid splitting. **C.** Correlation between the relative timing of nucleoid splitting and the growth rate as
captured by our reaction-diffusion model (D_n = 0.001 μm^2/sec) across six growth rate bins. The cell and
nucleoid lengths for each growth rate bin matched previously published data shown in Figure 4 – figure
supplement 1A. **D.** Deviation between the steady state after infinite relaxation time (dashed curves) and the
polysome or nucleoid profiles in newborn cells after one simulation round (solid curves) for increasing
nucleoid diffusion constants during fast growth (1 h^{-1}). **E.** Phase contrast and fluorescence (fluor.) images

of two representative single cells (CJW7651). The red circles indicate the position of the mCherry-µNS
particle in each cell. **F.** Two-dimensional average cell projections of the DAPI concentration (conc.) and the
RplA-msfGFP concentration, and 2D histogram of the mCherry-µNS particle density for four cell length bins
of CJW7651 cells (~2580 cells per bin) grown in M9gluCAAT and spotted on an agarose pad containing
the same medium. Since the cell pole identity cannot be inferred from snapshot images, pole assignment
was random. **G.** Average 1D profiles of the scaled DAPI and RplA-msfGFP concentrations and the
mCherry-µNS probability density.

2006

Figure 5: Effects of rifampicin treatment and polysome depletion on nucleoid segregation and
compaction. A. Plot showing the average instantaneous growth rate (mean ± SD shown by the solid black
curve and grey shaded region, respectively) of a cell population (n = 2629 cell division cycles) undergoing
two rounds of rifampicin treatment in a microfluidic device supplemented with M9gluCAAT. The distribution
of the average cell cycle growth rate of unperturbed populations is also shown on the right (n = 4122 cell
division cycles from a different microfluidics experiment). The solid horizontal line indicates the average
growth rate. **B.** Plot showing the average distance between nucleoid peaks for a population of cells
(squares, 114 cell division cycles) that were born (-112 to -102 min) and divided before the addition of
rifampicin, and for a population of cells (circles, 112 cell division cycles) that were born just before (-22 to -
12 min) and divided after the addition of rifampicin. A third-degree polynomial was fitted to the data from
the unperturbed population (solid blue curve) and juxtaposed (dashed blue curve) with the data from the
interrupted population. **C.** Average 1D profile and 2D projections of the scaled (divided by the whole cell
average concentration) RplA-GFP and HupA-mCherry signals for cells before and after rifampicin addition
(n = 112 cell division cycles). The red dashed horizontal lines in the 1D intensity profiles and the white
crosses in the 2D profiles mark the nucleoid peaks. **D.** Plot showing the RplA-GFP accumulation relative to
the HupA-mCherry depletion at mid-cell from 0 to 24 min after birth (colormap) for cells that completed their
division cycle before the addition of rifampicin (n = 114 cell division cycles) and for cells that were subjected
to rifampicin 12 min (n = 112 cell division cycles) or 3 min (n = 99 cell division cycles) after birth. **E.** Average
1D and 2D scaled RplA-GFP and HupA-mCherry intensity profiles for newborn cells (0 to 10 min after birth)
before (left, n = 726 cell division cycles), just after (middle, n = 367 cell division cycles), and much after
(right, n = 235 cell division cycles) rifampicin addition.

2028

2029   **Figure 6: Effects of ectopic polysome accumulation on nucleoid dynamics. A.** Schematic
2030   summarizing the experiment. **B.** Representative phase contrast and mTagBFP2 fluorescence images at
2031   different times after induction with IPTG (100 µM) are shown, next to a plot showing the mTagBFP2
2032   fluorescence of the entire population (mean ± SD, n = 3624 cell trajectories) over time. **C.** Plot showing how
2033   instantaneous growth rate (mean ±SD, n = 3624 mTagBFP2 induction trajectories) decreases following
2034   induction of mTagBFP2 synthesis. **D-F.** Representative kymographs and images of the normalized (divided
2035   by the whole cell average) RplA-msfGFP and HupA-mCherry fluorescence signals in cells (CJW7749) born

2036   during mTagBFP2 over-expression. **F.** Phase contrast images are shown to illustrate the formation of

2037   inclusion bodies (see also Figure 6 – figure supplement 1). Additional cell examples are shown in Movie

2038   S5.

2039

Figure 7: Effects of cell width increase on polysome and nucleoid dynamics. A. Comparison of the

cell width increase during cell growth between CJW7323 cells treated with cephalexin (mean ± SD, 360 cell

growth trajectories, 418 to 1511 segmented cells per bin) and cells treated with both cephalexin (50 µg/mL)

and A22 (4 µg/mL) (grey, mean ± SD, 309 cell growth trajectories, 51 to 1684 segmented cells per bin).

The same cell area bins are compared between the two populations. **B.** Phase contrast and fluorescence

images of a representative cephalexin-treated cell expressing RplA-GFP and HupA-mCherry. **C.** Same as

B but for a short cell growing in the presence of A22 and cephalexin. **D.** Same as C but for a longer cell.

The white arrowheads indicate the polysome bridges that connect polysome accumulations between two

DNA-free regions. Additional examples are shown in Movie S6 **E.** Representative fluorescence images of

RplA-GFP and HupA-mCherry in a cell treated with A22 and cephalexin. The dotted lines indicate the

representative cross-like polysome accumulation, which forms during the fusion of the polysome

accumulation towards the center (see also Movie S6). **F.** Comparison of the segmented polysome

accumulations and nucleoid objects between A22+cephalexin (150 sampled segmented cells from 47

growth trajectories) and cephalexin-treated (150 sampled segmented cells from 100 growth trajectories)

cells. The polysome and nucleoid areas per cell were normalized by the population-average statistic from

cephalexin-treated cells. All differences between the two populations are statistically significant (Mann-

Whitney p-value $< 10^{-10}$).

2057

2058

Figure 1 – figure supplement 1: Reproducibility analysis of the dynamic ribosome and nucleoid

distributions between microfluidic experiments. While phase-contrast images were acquired every

minute in all microfluidics experiments, two different intervals (1 min and 3 min) were used for fluorescence

image acquisition. **A.** Kymographs showing the RplA-GFP and HupA-mCherry concentration in CJW7323

cells growing in microfluidic channels in M9gluCAAT. Cell and nucleoid contours are shown in each channel

using a different color for each cell lineage (from dark purple to bright orange). **B-E.** Two biological replicate

experiments were performed for this strain and nutrient condition using different intervals of fluorescence

image acquisition (1 or 3 min). **B.** Plots showing that the instantaneous and average division cycle growth

rates were nearly identical between the two experiments. **C.** Plots showing the distributions of the indicated

fluorescence signals for the 1-min acquisition interval (black) compared to the 3-min frame rate interval

(orange) before and after correction. The difference in the excitation power between the two experiments

was 50% (240 %ms vs. 360 %ms), which was also reflected in the ratio of their average fluorescence values

(1.4 for RplA-GFP and 1.6 for HupA-mCherry). The fluorescence values from the 1-min interval experiment

were corrected by multiplying by these ratios, resulting in a near-perfect overlap between fluorescence

distributions after correction. **D.** Plots showing that the scaled (z-score) RplA-GFP and HupA-mCherry

intensity profiles from birth to division were almost identical between the two experiments. The intensity

profiles are shown for 10 cell division cycle intervals (1489 and 2633 cell division cycles for the 1-min and

3-min interval experiments, respectively). The z-score was calculated for each segmented cell by

subtracting the whole cell average fluorescence and then dividing the difference by the standard deviation.

Each intensity profile corresponds to the average z-score of all segmented cell instances within the

corresponding cell cycle interval. **E.** Plot showing that the average RplA-GFP concentration from birth to

division (averages ± SD across 20 cell division cycle bins) remained constant and was virtually identical

between the two experiments. **F.** Representative image of RplA-GFP fluorescence in cells (CJW7323)

treated with rifampicin (100 µg/mL). The cells were treated for 45 min, washed, and spotted on an

M9gluCAAT agarose pad without antibiotic. The presented snapshot corresponds to the first time point

immediately after cell spotting showing diffuse distribution of RplA-GFP signal.

2085

Figure 1 – figure supplement 2: Tracking nucleoid segregation cycles. Since the timing of nucleoid

segregation varied between cell division cycles, the nucleoid segregation cycles were tracked

independently of the cell division cycles for each cell lineage to measure the relative timing of RplA-GFP

accumulation and HupA-mCherry depletion in the middle of the nucleoid (as shown in Figure 1E). **A.** Plots

showing the frequency of cells with one, two, three, or four detected nucleoid objects across 10 cell division

cycle bins from birth (top) to division (bottom). Data from 4122 cell division cycles are shown. **B.** The polarity

of the cells (+/-) and the relative position of the nucleoid mask (toward the new or old pole) was used to

track the nucleoid segregation cycle. This strategy was used to identify four groups of nucleoid segregation

cycles (group -2, -1, 1 and 2). Group -2 includes nucleoids that were "born" toward the old cell pole in a

mother cell with negative (-) polarity and were inherited at the center of a daughter cell with a negative

polarity. Group -1 includes nucleoids that were "born" toward the new cell pole in a mother cell with negative

(-) polarity and were inherited at the center of a daughter cell with a positive (+) polarity. Group 1 includes

nucleoids that were "born" toward the new cell pole in a mother cell with positive (+) polarity and were

inherited at the center of a daughter cell with negative (-) polarity. Group 2 includes nucleoids that were

"born" toward the old cell pole in a mother cell with positive (+) polarity and were inherited at the center of

a daughter cell with positive (+) polarity. **C.** Plots showing the distributions of nucleoid positions around the

cell center for 10 nucleoid segregation cycle bins, from the time a nucleoid was "born" (top) until it split

(bottom). These plots show the inheritance of the nucleoids from the quarter cell positions of the mother

cells to the middle of their daughters for the four groups of nucleoid segregation cycles. The solid gray line

represents the average density of the nucleoid positions for all four groups. Data from 2286 complete

nucleoid cycles are shown. **D.** Schematic that explains the definition of a nucleoid cycle. The nucleoid cycle

ranges from the end of a nucleoid splitting event, until the next splitting of the sister nucleoids. It usually

extends beyond cell division, into the next cell division cycle.

2109

Figure 1 – figure supplement 3: Correlations used to calculate the relative contribution of polysome
accumulation and cell elongation to nucleoid migration. A. Correlation (Spearman ρ = 0.47, p-value <
10^{-10}) between the rate of RplA-GFP accumulation at mid-cell ($\frac{\Delta RplA_{mid-cell\ conc.}}{\Delta T}$) and the rate of distance
increase between the sister nucleoids minus the rate of cell elongation ($\frac{\Delta Distance_{nuc}}{\Delta T} - \frac{\Delta Length_{cell}}{\Delta T}$) across
cells. The colormap corresponds to a Gaussian kernel density estimation. Binned data are also shown
(mean ± SEM, 75 to 177 cell division cycles per bin, 9 bins in total) within the 5^{th}-95^{th} percentiles of the *x*-
axis range. **B.** Scaled correlations (z-scores) between $\frac{\Delta RplA_{mid-cell\ conc.}}{\Delta T}$ (blue) or $\frac{\Delta Length_{cell}}{\Delta T}$ (grey) and
$\frac{\Delta Distance_{nuc}}{\Delta T}$ (y-axis) during four relative time bins (1335 to 1957 cell division cycles per bin) covering the
period from the end of nucleoid splitting until cell division. Shown are the Spearman correlations (ρ), all with
a p-value below 10^{-6}, except for the one marked with an asterisk (p-value = 0.02).

2120

Figure 1 – figure supplement 4: Examination of the relative timing of the initiation of nucleoid
constriction and the accumulation of polysomes at mid-nucleoid. A. RplA-GFP and HupA-mCherry
concentration (fluorescence arbitrary units) images of four representative single cells (CJW7323) growing
in M9glyT and their fluorescence intensity profiles along the cell length. **B.** Demographs of the scaled RplA-
GFP and HupA-mCherry concentration from 13554 *E. coli* cells grown in M9glyT. **C.** Average 1D intensity
profiles of the scaled (divided by the whole cell average) RplA-GFP and HupA-mCherry concentration for
12 bins of cells lengths (>1100 cells per intensity profile). **D.** Plot showing the scaled RplA-GFP and HupA-
mCherry concentration in the middle of the cells for increasing cell length (~370 cells per bin, 25 bins). Cells
longer than 3 µm were excluded to avoid the effects of cell constriction. The arrow and vertical dashed line
indicate the minimum cell length bin with an apparent HupA-mCherry depletion at mid-cell, marking the
initiation of nucleoid splitting. The error bars indicate mean ± standard error of the mean (SEM).

2132

Figure 2 – figure supplement 1: Demographs of scaled ribosome and nucleoid fluorescence for
strains with different ribosome markers and under various nutrient conditions. Demographs are
ordered according to increasing average cell area, which scales with growth rate (Schaechter et al., 1958),
from left to right and top to bottom. These demographs were constructed from snapshots of the following
strains: **A.** CJW6768 (RplA-mEos2 ribosomal marker, 747 to 2446 cells per condition), **B.** CJW6769 (RpsB-
mEos2 ribosomal marker, 690 to 3169 cells per condition), **C.** CJW7020 (RplA-msfGFP ribosomal marker,
657 to 2432 cells per condition) and **D.** CJW7021 (RpsB-msfGFP ribosomal marker, 788 to 1950 cells per
condition), using the cell length as a proxy for the cell division cycle, with the signal intensity profile sorted
from the shortest newborn cells to the longest predivisional cells. Note that the demographs for the
CJW6768 strain (panel A) in the nutrient conditions M9mann, M9mala, M9malt, M9mannCAAT,
M9glyCAAT and M9malaCAAT are not shown here, as they are presented in Figure 2A. The nutrient
abbreviations and compositions are explained in Table S1.

2145

2146

Figure 2 – figure supplement 2: Calculation of RplA-msfGFP concentration after cell curvature
correction. A. Average 2D RplA-msfGFP projections for short (2.5 to 3 μm – left), unconstricted cells and
long (3.5 and 5.5 μm - right) constricted cells. The average RplA-msfGFP signal from 1000 sampled cells
(CJW7651) is shown for each population. **B.** The 2D cell areas were divided in cylindrical sectors with a
height (h) of a single pixel. The number of cylindrical sectors corresponds to the average cell length in pixels
(40 and 60 sectors for the unconstricted and constricted cells, respectively). The radius (r) of each
cylindrical segment corresponds to half the cell diameter for that specific cell length position. The cylindrical
segments at the poles or constriction site have a smaller radius due to the curvature of the cell boundaries.
The division of the cell area into cylindrical segments allowed us to calculate the volume for each segment
($V_{cylinder} = \pi\, r^2\, h$). The sum of the fluorescence per segment was corrected for the 3D curvature of the cell
boundaries by dividing by the volume of the cylinder, either considering the entire cell width or different
depths of view narrower than the maximal cell width. **C.** 1D profiles of the cylinder diameter are also
indicative of the cell curvature at the poles and constriction site. **D.** Comparisons of the 1D average RplA-
msfGFP projection between the poles before (dashed lines) or after correcting for the cell curvature (solid
lines) for different depths of view (colormap). The uncorrected signal (dashed lines) corresponds to the
average fluorescence along each cylindrical segment (mean of pixel intensities). The 1D profiles were
normalized by dividing by the maximal RplA-msfGFP concentration at mid-cell. This correction shows that
the pronounced decrease of the RplA-msfGFP fluorescence at the poles is the result of the curvature of the
cell boundaries. The observed over-correction at the cell poles for larger depths of view (> 8 pixels or px)
is likely due to the over-estimation of the cell segmentation mask that was determined based on the phase
contrast snapshot images of cells on agarose pads.

2168

2169

Figure 3 – figure supplement 1: Analysis of the polysome and nucleoid asymmetries using fitted
Gaussian functions. A. Example images of RplA-GFP and HupA-mCherry fluorescence signals in a single
newborn cell instance. **B.** Plots showing the Gaussian-function fitting on RplA-GFP and HupA-mCherry
intensity profiles from the single cell snapshot shown in panel A. Three Gaussian functions were fitted to
the RplA-GFP fluorescence in newborn cells (eq. 11 in Methods), capturing the accumulation of polysomes
at mid-cell and the poles. Two Gaussian functions were fitted to the HupA-mCherry fluorescence (eq. 12 in
methods) capturing the two lobes of the segregating sister nucleoids. The parameters of the fitted Gaussian
functions provided information about the position (μ: mean) and the concentration (A: amplitude) of each
fluorescence statistic, as well as the cell length range occupied by it (σ: standard deviation). The Gaussian
area (eq. 13 in Methods) corresponds to the abundance of each macromolecule. The RplA-GFP Gaussians
were fitted above the cellular background, which presumably corresponds to the uniform fluorescence of
free ribosomes or ribosomal subunits. For the HupA-mCherry Gaussian fittings, the cellular background
corresponds to the DNA-free cell regions. The Gaussian parameters were also used to describe the

polysome asymmetries between the poles (eq. 14 in Methods) as well as the position of the nucleoid around

the cell center (eq. 15 in Methods). These statistics were used in Figure 3B. **C.** Gaussian functions were

also fitted to the ensemble average RplA-GFP and HupA-mCherry fluorescence (right) calculated from all

the cell segmentation instances of a single cell division cycle, 0-10% from birth to division (left). **D.** The

Gaussian fitting to the ensemble RplA-GFP and HupA-mCherry intensity profiles (concentration in arbitrary

fluorescence units) early in the cell division cycle for the 2D projection shown in panel C (right). **E.** Plots

showing the nucleoid and polysome density axial asymmetry determined using the parameters of the fitted

Gaussians on the HupA-mCherry and RplA-GFP fluorescence (eq. 16 and eq. 18 in Methods). **F.** Plots

showing the scaled correlations between the polysome Gaussian parameters (μ, A, and σ for each

polysome accumulation) and the nucleoid density asymmetry (eq. 16 in Method; 2103 cell division cycles).

Each marker corresponds to the data for a single cell division cycle (0-10% into the cell division cycle) as

in panel C (right). A Gaussian kernel density estimation (KDE) was used to illustrate the density of the

scatter plots (see Methods). **G.** Bar graph showing the coefficients of a linear mixed-effects model (see eq.

17 in Methods) used to identify the polysome statistics that contribute to the nucleoid density asymmetry.

The highly contributing polysome statistics were combined to a compound polysome statistic (eq. 18 in

Methods) that describes the available DNA space between polysome accumulations. This compound

statistic correlates with the nucleoid density asymmetry (eq. 16 in Methods) as shown in Figure 3F. The

stars indicate the polysome statistics that most significantly correlate with the nucleoid density asymmetry

(absolute coefficient above 0.3).

2202

Figure 4 – figure supplement 1: Statistics of polysome accumulation and nucleoid depletion during

simulated cell growth. Polysome and nucleoid dynamics were simulated for six different growth rates.

The predivisional polysome and nucleoid profiles of simulated trajectories initialized from the steady state

(infinite relaxation time) were used as initial conditions ($D_n = 10^{-3}$ μm^2/s). **A.** Plot showing cell length at birth

increasing with growth rate according to published measurements (Govers et al., 2024). **B.** Plot showing

the dynamics of polysome accumulation (top) and nucleoid depletion (bottom) at mid-cell from birth to

division for six simulated growth rates (colormap). A nucleoid depletion threshold of 0.1 arbitrary units (a.u.,

dashed horizontal line) was used to determine the relative time of nucleoid splitting that is presented in

Figure 4C.

2212

Figure 5 – figure supplement 1: Phenotypic effects of transcription inhibition. Plots showing cellular

parameters (instantaneous growth rate, cell area, nucleoid area, and the ratio between nucleoid area and

cell area, or NC ratio) affected by the rifampicin treatment. The instantaneous growth rate was estimated

using the log-transformed cell area, using a rolling window of 10 min. The black line corresponds to the

averages from 1859 cell division cycles and the gray-shaded area indicates the range of one standard

deviation around the mean. The two cycles of rifampicin treatment (as shown in Figure 5A) were overlaid

for this plot, using the time of rifampicin addition as t = 0 min.

2220

2221 **Figure 6 – figure supplement 1: Phenotypic effects of prolonged protein over-expression from**
2222 **plasmids.** Representative phase contrast and fluorescence images of cells (CJW7798) after prolonged
2223 induction (100 µM IPTG, for 9 h and 15 min) of mTagBFP2 expression from aT7 promoter on a multi-copy
2224 plasmid.

2225

2226 **Figure 7 – figure supplement 1: Phenotypic of effects of cell growth under A22 and cephalexin**
2227 **treatment.** Representative phase contrast and fluorescence images of cells (CJW7323) treated with A22
2228 (4 µg/mL) alone or with both A22 (4 µg/mL) and cephalexin (50 µg/mL) for 130 min.

2229

2230

Movie legends
Movie S1: Example time-lapse sequence showing ribosome and nucleoid dynamics during cell
growth under steady state condition in a microfluidic channel. Shown are corresponding inverted
phase contrast signal, RplA-GFP signal, HupA-mCherry signal, cell contours (based on phase contrast
signals) and nucleoid contours (based on nucleoid signal segmentation) of *E. coli* cells (CJW7323) grown
in a microfluidic channel supplemented with M9gluCAAT. The white circles indicate the centroid of the cell
segmentation masks and the extending lines indicate the tracked cell traces.

2238

2239 **Movie S2: Video showing the average subcellular distribution of ribosome and nucleoid signals**
2240 **from birth to division.** Shown are 2D average RplA-GFP (top) and HupA-mCherry (bottom) projections
2241 from birth to division, with corresponding average intensity profiles (right). The cell projections are oriented
2242 from the old pole on the left to the new pole on the right. The average cell contour is also drawn. Ensemble
2243 data from 4122 division cycles of CJW7323 cells growing in M9gluCAAT are shown.

2244

2245 **Movie S3: Video showing simulated 1D profiles of polysome and nucleoid concentration during**
2246 **slow and fast cell growth.** Simulations during slow (left) and fast (right) growth are shown. The simulations
2247 were initialized from the equilibrium configuration, with a compact symmetric nucleoid ($D_n = 10^{-3}$ µm^2/sec)
2248 at the cell center.

2249

2250 **Movie S4: Video showing the effects of rifampicin addition on cell growth, ribosome signal**
2251 **heterogeneity, nucleoid segregation, and nucleoid compaction.** Examples of five microfluidic channels
2252 showing the corresponding inverted phase contrast, RplA-GFP and HupA-mCherry signals (from left to
2253 right) of cells (CJW7323) growing within microfluidic channels. Rifampicin was added at 120 and 720 min.
2254 Each antibiotic treatment lasted 120 min.

2255

Movie S5: Video showing the effects of ectopic polysome formation on nucleoid dynamics over

time. The RplA-GFP and HupA-mCherry fluorescence normalized (norm.) by the average cell fluorescence

are shown, together with their corresponding phase contrast image, for multiple cells (CJW7798) in

succession following induction of mTagBFP2 expression from a T7 promoter on a multi-copy plasmid. The

scale bar indicates 1 µm, and the time since cell birth is shown in minutes. The cell contours indicate the

boundaries of the cell masks obtained by cell segmentation of the corresponding phase contrast images.

2262

Movie S6 Video showing how the loss of cell width confinement due to cephalexin and A22

treatment affects ribosome and nucleoid distributions over time. Fluorescence images of RplA-GFP

and HupA-mCherry fluorescence (fluor.) normalized by the average cellular fluorescence images, together

with their corresponding phase contrast images, are shown for cells (CJW7323) following treatment with

A22 (4 µg/mL) and cephalexin (50 µm/mL). The scale bar indicates 1 µm. The time since drug addition is

shown in minutes.

2269

2270

2271 **Supplementary tables**

2272 **Table S1. Growth medium abbreviation and composition.**

Abbreviation	Composition
M9acet	1x M9 salts, 0.2% acetate (sodium salt)
M9acetCAAT	1x M9 salts, 0.2% acetate, 0.1% casamino acids, 1µg/mL thiamine
M9fuc	1x M9 salts, 0.2% fucose
M9fucCAAT	1x M9 salts, 0.2% fucose, 0.1% casamino acids, 1µg/mL thiamine
M9fum	1x M9 salts, 0.2% fumarate (disodium salt)
M9fumCAAT	1x M9 salts, 0.2% fumarate, 0.1% casamino acids, 1µg/mL thiamine
M9GlcNac	1x M9 salts, 0.2% N-acetylglucosamine
M9GlcNacCAAT	1x M9 salts, 0.2% N-acetylglucosamine, 0.1% casamino acids, 1µg/mL thiamine
M9glu	1x M9 salts, 0.2% glucose
M9gluCAAT	1x M9 salts, 0.2% glucose, 0.1% casamino acids, 1µg/mL thiamine
M9gly	1x M9 salts, 0.2% glycerol
M9glyCAAT	1x M9 salts, 0.2% glycerol, 0.1% casamino acids, 1µg/mL thiamine
M9Lala	1x M9 salts, 0.2% L-alanine
M9LalaCAAT	1x M9 salts, 0.2% L-alanine, 0.1% casamino acids, 1µg/mL thiamine
M9Lara	1x M9 salts, 0.2% L-arabinose
M9LaraCAAT	1x M9 salts, 0.2% L-arabinose, 0.1% casamino acids, 1µg/mL thiamine
M9mala	1x M9 salts, 0.2% malate (sodium salt)
M9malaCAAT	1x M9 salts, 0.2% malate, 0.1% casamino acids, 1µg/mL thiamine
M9malt	1x M9 salts, 0.2% maltose
M9maltCAAT	1x M9 salts, 0.2% maltose, 0.1% casamino acids, 1µg/mL thiamine
M9mann	1x M9 salts, 0.2% mannose
M9mannCAAT	1x M9 salts, 0.2% mannose, 0,1% casamino acids, 1µg/mL thiamine
M9pyr	1x M9 salts, 0.2% pyruvate (sodium salt)
M9pyrCAAT	1x M9 salts, 0.2% pyruvate, 0,1% casamino acids, 1µg/mL thiamine
M9sorb	1x M9 salts, 0.2% sorbitol
M9sorbCAAT	1x M9 salts, 0.2% sorbitol, 0,1% casamino acids, 1µg/mL thiamine
M9succ	1x M9 salts, 0.2% succinate (disodium salt)
M9succCAAT	1x M9 salts, 0.2% succinate, 0,1% casamino acids, 1µg/mL thiamine
M9treh	1x M9 salts, 0.2% trehalose
M9trehCAAT	1x M9 salts, 0.2% trehalose, 0,1% casamino acids, 1µg/mL thiamine
M9xyl	1x M9 salts, 0.2% xylose
M9xylCAAT	1x M9 salts, 0.2% xylose, 0,1% casamino acids, 1µg/mL thiamine

2273

2274

2275 **Table S2: *Escherichia coli* strains used in this study.**

Strain	Genotype	Source
MG1655	*E. coli* MG1655 (F-lambda- *ilvG- rfb-50 rph-1*)	(Guyer et al., 1981; Jensen, 1993)
MG1655 (DE3)	*E. coli* MG1655 (DE3)	Kind gift from Dr K. Prather (Massachusetts Institute of Technology), (Tseng et al., 2010b)
CJW5158	*E. coli* BW25113 *hupA::hupA-mcherry-frt-kanR-frt*	(Gray et al., 2019)
CJW6723	*E. coli* MG1655 Δ*lacZYA*::P*lac*-*egfp-µNS* hupA:*hupA-mcherry*	(Gray et al., 2019)
CJW6768	*E. coli* MG1655 *rplA::rplA-meos2*	(Sanamrad et al., 2014)
CJW6769	*E. coli* MG1655 *rspB::rpsB-meos2*	(Sanamrad et al., 2014)
CJW7019	*E. coli* MG1655 *rplA::rplA-msfgfp-frt-kanR-frt*	(Gray et al., 2019)
CJW7020	*E. coli* MG1655 *rplA::rplA-msfgfp*	(Gray et al., 2019)
CJW7021	*E. coli* MG1655 *rpsB::rpsB-msfgfp*	(Gray et al., 2019)
CJW7144	*E. coli* MG1655 Δ*lacZYA*::P*lac*-*mcherry-µNS-frt-kanR-frt*	This study
CJW7145	*E. coli* MG1655 Δ*lacZYA*::P*lac*-*mcherry-µNS-frt-kanR-frt rplA::rplA-msfgfp*	This study
CJW7323	*E. coli* MG1655 *rplA::rplA-msfgfp hupA::hupA-mcherry*	(Xiang et al., 2021)
CJW7466	*E. coli* MG1655 (DE3) hupA:*hupA-mcherry*	This study
CJW7651	*E. coli* MG1655 Δ*lacZYA*::P*lac*-*mcherry-µNS rplA::rplA-msfgfp*	This study
CJW7766	*E. coli* MG1655 (DE3) hupA:*hupA-mcherry rplA::rplA-msfgfp-frt-kanR-frt*	This study
CJW7798	*E. coli* MG1655 (DE3) hupA:*hupA-mcherry rplA::rplA-msfgfp-frt-kanR-frt	pET28:mTagBFP2-CmR*	This study

2276
2277 **Table S3: Plasmids used in this study.**

Plasmid name	Relevant genetic elements	Source
pKD13	*frt-kanR-frt-R6Kori-ampR*	(Datsenko and Wanner, 2000)
pKD46	*araC-bet-exo-A101(Ts)ori-ampR*	(Datsenko and Wanner, 2000)
pCP20	*cmR-A101(Ts)ori-ampR-flp-λ-repressor(Ts)*	(Datsenko and Wanner, 2000)
pER12	*pBAD322A-gfp-µNS*	Kind gift from Dr A. Janakiraman (City College of New York), (Broering et al., 2005)
pER12-mcherry	*pBAD322A-mcherry-µNS*	This study
pAPG1	*attB-pBAD322A-mcherry-µNS-attB-kanR-ColE1ori*	This study
pSB3C5-proA-B0032-E0051	*PproA-lacZα-GFP-p15Aori-cmR*	(Davis et al., 2011)
pET28:GFP	*PT7-lacO-GFP-kanR-pMB1ori-lacI*	(Shis and Bennett, 2013)
pBAD:TagBFP2	*araC-ParaBAD-mTagBFP2-ampR-pBAD322ori*	(Subach et al., 2011)
pET28:mTagBFP2	*PT7-lacO-mTagBFP2-kanR-pMB1ori-lacI*	This study
pET28:mTagBFP2-CmR	*PT7-lacO-mTagBFP2-cmR-pMB1ori-lacI*	This study

2278
2279
2280 **Table S4: DNA oligonucleotides used in this study.**

Oligo name	Sequence	Source
ER12-MCR-fwd	5' CTCCATACCCGTTTTTTTGGGCTAGCAGGAGGAATTCATGGTGAGCAAGGGCGAGGAG 3'	This study
ER12-MCR-rev2	5' CACTGGAGGAGCCTGCTTTTTTGTACAAACTTGTTGACTTGTACAGCTCGTCCATG 3'	This study
µNSmCherry fwd	5' CACAGGTTGCTCCGGGCTATGAAATAGAAAAATGAATCCGTTGAAGCCTGATCGATGCATAATGTGCC 3'	This study
µNSmCherry rev	5' AGCTCCAGCCTACACAGAGTTTGTAGAAACGCAAAAAG 3'	This study
FRT_KanR fwd	5' GTTTCTACAAACTCTGTGTAGGCTGGAGCTGC 3'	This study
FRT_KanR Rev	5' TTAAAGGTATTAAAAACAACTTTTTGTCTTTTTACCTTCCCGTTTCGCTCCTGTCAAACATGAGAATTAATTCC 3'	This study
ColE1 fwd	5' GAGCGAAACGGGAAGGTAAAAAGACAAAAAGTTGTTTTTAATACCTTTAACGTTCCACTGAGCGTC 3'	This study
ColE1 rev	5' CAGGCTTCAACGGATTCATTTTTCTATTTCATAGCCCGGAGCAACCTGTGGGTTATCCACAGAATCAGG 3'	This study
pAPG1 seq1	5' GATCAAGCAGAGGCTGAAG 3'	This study

pAPG1 seq2	5' CCAGGGCATCAAATTAAGC 3'	This study
pAPG1 seq3	5' TCTACGTGTTCCGCTTCC 3'	This study
pAPG1 seq4	5' TTGAAGCCTGATCGATGC 3'	This study
pAPG1 seq5	5' AAGATTAGCGGATCCTACC 3'	This study
lacZYA_redµNS fwd	5' TATGTTGTGTGGAATTGTGAGCGGATAACAATTTCACACAGGAAACAGCTATGGTGAGCAAGGGCGA 3'	This study
lacZYA_redµNS rev	5' CAATTTTTATAATTTAAACTGACGATTCAACTTTATAATCTTTGAAATAAGGATCCGTCGACCTGCAG 3'	This study
LacI fwd	5' GGCCGATTCATTAATGCAGCTGGC 3'	This study
CynX rev	5' GGCCTGATAAGCGCAGCGTATC 3'	This study
mCherry rev	5' GGTGCTTCACGTAGGCCTTGG 3'	This study
KanR fwd	5' CGGAGAACCTGCGTGCAATCC 3'	This study
mTagBFP2 fwd	5' CGGAGCTCGAATTCGGATCCTTAATTAAGCTTGTGCCCCAGTTTG 3'	This study
mTagBFP2 fwd	5' CTTTAAGAAGGAGATATACCATGAGCGAGCTGATTAAGGAGAAC 3'	This study
pET28_one fwd	5' TCCTTAATCAGCTCGCTCATGGTATATCTCCTTCTTAAAGTTAAAC 3'	This study
pET28_one rev	5' GGATAACCGTATTACCGCCTTTGAGTGAGCTGATACCG 3'	This study
pET28_two fwd	5' AGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCC 3'	This study
pET28_two rev	5' TGGGGCACAAGCTTAATTAAGGATCCGAATTCGAGCTCC 3'	This study
pET28mTagBFP2 fwd	5' CTTAGTGACTCGAATTCGCGCGCCAATCCGGATATAGTTCC 3'	This study
pET28mTagBFP2 rev	5' ACTTTCTGGCTGGATGATGGACGTGAGTTTTCGTTCCACTG 3'	This study
cmR fwd	5' AGTGGAACGAAAACTCACGTCCATCATCCAGCCAGAAAGTG 3'	This study
cmR rev	5' GAACTATATCCGGATTGGCGCGCGAATTCGAGTCACTAAGG 3'	This study

2281

2282

2283 **Table S5: Chemicals used in this study.**

Chemical	Source	Catalog number
Agarose	AmericanBio	Cat#AB00972-00500
4',6-diamidine-2'-phenylindole (DAPI), dihydrochloride fluorescent dye	Thermo Fisher Scientific	Cat#D1306
Isopropyl β-D-1-thiogalactopyranoside (IPTG)	Sigma Aldrich	Cat#I5502
Rifampicin	Sigma Aldrich	Cat#R3501
Cephalexin	Sigma Aldrich	Cat#C4895
A22	Sigma Aldrich	Cat#SML0471
4',6-Diamidine-2-phenylindole dihydrochloride (DAPI)	Thermo Fisher Scientific	Cat#D1306

2284

2285

2286 **Table S6: Software used in this study.**

Program / library / script	Package	Source
Oufti	www.oufti.org	(Paintdakhi et al., 2016)
MATLAB	www.mathworks.com	Mathworks
Python	www.python.org	Python Software Foundation
Numpy	www.numpy.org	(Harris et al., 2020)
Scipy	www.scipy.org	(Virtanen et al., 2020)
Pytorch	www.pytorch.org	(Paszke et al., 2019)
Scikit-image	www.scikit-image.org	(Van Der Walt et al., 2014)
Scikit-learn	www.scikit-learn.org/stable/	(Pedregosa et al., 2012)
Statsmodels	www.statsmodels.org/stable/index.html	(Seabold and Perktold, 2010)
Shapely	www.pypi.org/project/shapely/	(Gillies, Sean et al., 2023)
Matplotlib	www.matplotlib.org	(Hunter, 2007)
Seaborn	www.seaborn.pydata.org	(Waskom, 2021)
Pandas	www.pandas.pydata.org	(McKinney, 2010)
Python	Omnipose neural network	(Cutler et al., 2022)
MATLAB	SuperSegger	(Stylianidou et al., 2016)
Python	Unet neural network	(Wiktor et al., 2021; Zhou et al., 2020)

snapshots_analysis_UNET_ghv.py	Snapshot image analysis (from UNET masks) – custom Python class	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
snapshots_analysis_OUFTI_GrayGovers_ghv.py	Snapshot image analysis (from Oufti masks) – custom Python class	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
snapshots_analysis_functions.py	Extraction of fluorescence and morphology statistics from cell snapshots – custom Python functions	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
microfluidics_segmentation_ghv.py	Cell segmentation and tracking from time-lapse images in microfluidics – custom Python class	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
microfluidics_analysis_functions_ghv.py	Extraction of fluorescence and morphology statistics from time-lapse images in microfluidics – custom Python functions	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
Omnipose_to_python_ghv.py	Extraction of segmentation masks and tracked lineages from Omnipose/SuperSegger to Python for further analysis	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
Otsu_phase_segmentation_ghv.py	Cell segmentation and tracking using Otsu thresholding (Otsu, 1979)	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
Biviriate_medial_axis_estimation.py	Function that draws the medial axis in randomly oriented cells on agarose pads	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
LoG_adaptive_image_filter.py	Image filter that applies relative and local thresholding to segment fluorescence objects and particles	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
Microfluidics_segmentation_and_tracking_example.pdf	Example for cell segmentation and tracking in microfluidics	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
Analysis_of_cell_morphology_and_fluorescence_in_microfluidics.pdf	Example for the extraction of fluorescence and morphology statistics from time-lapse images in microfluidics	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
Analysis_of_cell_morphology_and_fluorescence_in_agarose_pads.pdf	Example for the extraction of fluorescence and morphology statistics from cell snapshots	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
Example_of_using_Otsu_based_segmentation_ghv.pdf	Example for the segmentation of antibiotic-treated cells using the Otsu threshold	This study (www.github.com/JacobsWagnerLab/published/tree/master/Papagiannakis_2024)
tune_diffusion.ipynb	Simulations of the minimal reaction-diffusion model that describes nucleoid segregation in *E. coli*.	This study (https://github.com/qiweiyuu/polysome)

2287

2288

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 1 - figure supplement 1

Figure 1 - figure supplement 2

Figure 1 - figure supplement 3

Figure 1 - figure supplement 4

Figure 2 – figure supplement 1

Figure 2 - figure supplement 2

Figure 3 - figure supplement 1

Figure 4 - figure supplement 1

Figure 5 - figure supplement 1

mTagBFP2 over-expression from a plasmid

Figure 6 - figure supplement 1

A22 treatment

2 µm

2 µm

A22 + Cephalexin treatment

2 µm

Phase contrast (a.u.)   RplA-GFP (a.u.)   HupA-mCherry (a.u.)

Signal percentile (%)   99.5   0.5

Figure 7 - figure supplement 1