Thiol Stress Fuels Pyrazinamide Action Against *Mycobacterium tuberculosis*

Lev Ostrer[1]†, Taylor A. Crooks[1]†, Michael D. Howe[1], Sang Vo[1,2], Ziyi Jia[1], Pooja Hegde[2], Courtney C. Aldrich[2], Anthony D. Baughn[1]*

[1]Department of Microbiology and Immunology, University of Minnesota Medical School, Minneapolis, Minnesota USA
[2]Department of Medicinal Chemistry, College of Pharmacy, University of Minnesota Medical School, Minneapolis, Minnesota USA

†These authors contributed equally to this work.
*Correspondence can be made to Anthony D. Baughn.

Email: abaughn@umn.edu

Author Contributions: Lev Ostrer, Taylor A. Crooks and Anthony D. Baughn conceived of the study. Lev Ostrer and Taylor A. Crooks designed and performed microbiology-related experiments, data analysis and interpretation. Sang Vo synthesized and performed quality control for the study compound. Michael D. Howe and Taylor A. Crooks developed Tn-seq and RNA-seq pipelines and conducted computational analyses. Pooja Hegde performed mass spectrometry data acquisition and analysis. Ziyi Jia constructed and validated targeted mutations in mycobacterial strains and conducted growth inhibition studies with these strains. Courtney C. Aldrich provided critical resources and expertise for synthesis and analysis of the study compound and its products. Lev Ostrer and Taylor A. Crooks prepared all display images. All authors discussed the results and contributed to writing and editing of the final manuscript, with major contributions by Anthony D. Baughn, Lev Ostrer and Taylor A. Crooks.

Competing Interest Statement: The authors have no competing interests to disclose.

Classification: Biological Sciences, Microbiology.

Keywords: Tuberculosis, Pyrazinamide, Thiol Stress, Coenzyme A, Drug Discovery

This PDF file includes:

> Main Text
> Figures 1 to 6
> Supplemental Figure S1
> Supplemental Table S1

Abstract

Pyrazinamide (PZA) is a cornerstone of first-line antitubercular drug therapy and is unique in its ability to kill nongrowing populations of *Mycobacterium tuberculosis* through disruption of coenzyme A synthesis. Unlike other drugs, PZA action is conditional and requires potentiation by host-relevant environmental stressors, such as low pH and nutrient limitation. Despite its pivotal role in tuberculosis therapy, the mechanistic basis for PZA potentiation remains unknown and the durability of this crucial drug is challenged by the emergent spread of drug resistance. To advance our understanding of PZA action and facilitate discovery efforts, we characterized the activity of a more potent PZA analog, morphazinamide (MZA). Here, we demonstrate that like PZA, MZA acts in part through impairment of coenzyme A synthesis. Unexpectedly, we find that, in contrast to PZA, MZA does not require potentiation due to aldehyde-mediated disruption of thiol metabolism and maintains bactericidal activity against PZA-resistant strains. Our findings reveal a novel dual action mechanism of MZA that synergistically disrupts coenzyme A synthesis resulting in a faster rate of killing and a higher barrier to resistance relative to PZA. Together, these observations resolve the mechanistic basis for potentiation of a key first-line antitubercular drug and provide new insights for discovery of improved therapeutic approaches for tuberculosis.

Significance Statement

Pyrazinamide is the only antitubercular drug of its kind, capable of targeting persistent *Mycobacterium tuberculosis* through disruption of the coenzyme A biosynthetic pathway. A peculiar feature of this drug is that its activity is conditional, requiring low pH for its action. Despite decades of investigation, the precise basis for this conditional susceptibility has remained elusive which has been a barrier to discovery of more effective next-generation analogs. Using approaches in chemical biology, functional genomics and bacterial physiology we demonstrate that activation of thiol stress is the basis for pyrazinamide potentiation. These findings resolve a long-standing question regarding the mechanistic basis for conditional PZA susceptibility of *M. tuberculosis* and reveal novel avenues for antimicrobial drug discovery efforts.

Main Text

Introduction

For several decades, pyrazinamide (PZA) has been an integral part of the first-line standard short course therapy for tuberculosis (TB)[1]. A unique feature of PZA is its ability to kill non-replicating populations of the causative agent *Mycobacterium tuberculosis*[2]. Use of PZA has played a pivotal role in reducing TB relapse rates and shortening the duration of treatment from 9 to 6 months[1]. In spite of this success, wide use of this treatment regimen has resulted in the emergent spread of PZA resistance. Current rates of PZA resistance in *M. tuberculosis* clinical isolates range from 16 to 42% depending upon patient cohort[3]. The primary mechanism of clinical PZA resistance occurs through spontaneous loss-of-function mutations in the *M. tuberculosis pncA* gene which encodes an amidase that is required for activation of PZA to pyrazinoic acid (POA)[4,5]. Identification of potent next-generation analogs of PZA that circumvent resistance will be critical to maintain use of this important therapeutic tool.

Despite its crucial role in TB therapy the antitubercular mechanism of PZA action is not fully defined. Recent evidence demonstrates that PZA acts through disruption of coenzyme A (CoA) biosynthesis, in part, through destabilization of L-aspartate decarboxylase (PanD) by POA[6,7]. Intriguingly, while CoA plays an indispensable role in *M. tuberculosis* metabolism, PZA susceptibility is conditional and requires activation of the cell envelope stress response through exposure of bacilli to low pH or a variety of other stressors[8]. How disruption of CoA biosynthesis connects with conditional susceptibility of *M. tuberculosis* to PZA has not been resolved[9]. Yet, it has recently been shown that exposure of *M. tuberculosis* to low pH results in thiol stress[10] which likely interferes with CoA synthesis through limitation of cysteine, an essential pathway substrate.

Of note, consistent with an essential role for environmentally driven potentiation of PZA action *in*
vivo, PZA shows no detectable antitubercular activity in mice that lack T cells and are thereby
impaired for macrophage activation which is required for full phagosomal acidification[11].
Identification of means to bypass the need for host-mediated potentiation of PZA action would
advance antitubercular therapy through promoting drug potency in the context of compromised
immunity which is causally associated with progression of TB[9].
To address the unmet needs outlined above, we set out to characterize the mechanism of
action of an intriguing PZA analog, morphazinamide (MZA, Figure 1A). Early clinical studies
demonstrated efficacy of MZA against *M. tuberculosis* in humans with equivalent potency and
safety profiles relative to PZA[12,13]. Further, *in vitro* assays indicated that the antitubercular activities
of MZA and PZA were comparable, with similar levels of growth inhibition in broth culture[14].
However, it was noted that MZA and other aminomethylene analogs did not require low pH to show
inhibitory activity and retained activity against PZA resistant isolates, suggesting that they operate
by a distinct mechism[14]. In this study we applied approaches in chemical biology, functional
genomics and bacterial physiology to characterize the mechanism of action of MZA. Similar to its
sustained activity against *pncA* null strains of *M. tuberculosis*, we find that MZA also retains activity
against *M. tuberculosis* strains with newly described PZA resistance mechanisms[8]. By using
transcriptional profiling and transposon sequencing (Tn-seq) we find many parallels between
responses of *M. tuberculosis* to PZA, POA and MZA that highlight the impact of these drugs on
CoA metabolism. Importantly, we uncover an association between thiol metabolism and
susceptibility to these drugs and find that MZA susceptibility in particular is strongly influenced by
the mycothiol reductase MscR due to release of aldehyde. Together, our findings reveal a central
role for thiol stress as a principal driver of PZA action which resolves the basis for conditional
susceptibility and illuminates new opportunities to improve potency and raise the resistance barrier
to PZA analogs.

104
105 **Results**
106
Impact of MZA on PZA susceptible and resistant *M. tuberculosis*. To assess the antitubercular
activity of MZA, we performed minimum inhibitory concentration assays (MICs) using a panel of
strains with differing mechanisms of PZA resistance and under various conditions that influence
PZA susceptibility. We began by comparing activity between PZA and MZA using standard growth
conditions at circumneutral pH with both H37Ra and H37Rv backgrounds. Consistent with the
known conditional susceptibility of *M. tuberculosis* to PZA, cultures grew unimpaired at
concentrations as high as 4.5 mM in medium at circumneutral pH (Figure 1B; Table S1). In striking
contrast, under these same conditions we observed >90% inhibition of growth in the presence of
280 µM MZA (Figure 1B; Table S1). Next, we assessed whether MZA activity could be enhanced
by conditions that potentiate PZA and diminished under conditions that antagonize PZA. To
evaluate potentiation, MICs were determined under acidic (pH 5.8) conditions (Figure 1C,D). To
determine if MZA activity could be antagonized similarly to PZA, the effect of the CoA intermediate
pantothenate on drug action was evaluated. Consistent with previous work, PZA was fully inhibitory
at 500 µM under acidic conditions (Figure 1C,D; Table S1), while the addition of 230 µM
pantothenate fully antagonized PZA action[15] (Figure 1E; Table S1). Acidic conditions were able to
enhance MZA activity by two-fold, lowering its MIC to 140 µM (Figure 1C,D; Table S1), whereas
addition of pantothenate caused a slightly antagonistic effect on MZA activity at both acidic and
circumneutral pH (Figure 1E; Table S1). These observations indicate that MZA is mildly impacted
by conditions that strongly influence PZA action.
We next sought to determine whether MZA retained activity against several previously
described PZA resistant isolates. These strains included mutants deleted for *pncA* (required for
activation of PZA to POA) and *sigE* (required for potentiation of PZA susceptibility at low pH), as
well as a *clpC1* promoter mutant and a strain with truncation of the 3' end of *panD*[8], both of which
are thought to interfere with POA action through stabilization of PanD[7]. Despite PZA showing
diminished activity against this panel of strains, MZA was highly active with MICs ranging from 140
to 280 µM (Figure 1F,G; Table S1). It is worth noting that the ∆*pncA* strain showed a two-fold

increase in its MZA MIC relative to the parental strain (Figure 1G; Table S1) indicating a minor contribution of PncA to MZA action.

One of the key features of PZA is its unique ability to elicit a bactericidal effect on non-growing *M. tuberculosis*[16]. Thus, we sought to determine whether MZA also has bactericidal activity against actively growing and metabolically quiescent drug tolerant bacteria by performing kill curves under nutrient rich and nutrient limiting conditions. When mid-exponential cultures under either acidic or circumneutral pH growth conditions were treated with 0.9 mM MZA, greater than 99.999% killing was observed after 4 days (Figure 1H,I). After 14 days of incubation following a single treatment, no surviving CFU were observed. We next tested the bactericidal activity of PZA and MZA under nutrient limiting conditions by exposing starved bacteria to either 0.9 mM PZA or MZA. After 56 days of stasis, the majority of untreated bacilli survived, whereas there was more than 99.9% loss in viability of bacilli exposed to 0.9 mM PZA (Figure 1J). In contrast to PZA, MZA bactericidal activity was transient, with 80% killing in the first 2 days followed by a plateau (Figure 1J). To determine whether the plateau in killing was due to drug inactivation or due to enrichment of a drug tolerant population, starved cultures were washed and treated with fresh MZA every 3 days. As before, we observed 80% killing after 3 days for both H37Rv (Figure 1K) and H37Rv Δ*pncA* (Figure 1L). Following the second treatment over 99.5% of bacteria were no longer viable for both strains (Figure 1K,L). The third treatment resulted in sterilization of bacteria with no detectable CFU for either strain (Figure 1K,L). These data indicate that MZA has a potent PncA-independent bactericidal activity against both actively growing and metabolically quiescent *M. tuberculosis*.

To assess the antibacterial spectrum of MZA action, we evaluated its inhibitory activity against a panel of human-associated ESKAPE pathogens[17]. All ESKAPE pathogens that were tested showed MICs greater than 10-fold higher than those for *M. tuberculosis* (Figure 1M,N), suggesting that like PZA, MZA activity is highly selective for mycobacterial species.

Transcriptional responses of *M. tuberculosis* to PZA, POA and MZA exposure. To gain a better understanding of the impact of PZA, POA and MZA on *M. tuberculosis* physiology and to determine similarities and differences in global responses of bacilli to these drugs, we performed genome-wide transcriptional profiling using RNA-seq. Since PZA is only active at low pH, all treatments were performed using *M. tuberculosis* grown in acidic (pH 5.8) media. Cultures of exponentially growing H37Rv were exposed to 200 µM PZA, POA, MZA or vehicle only (DMSO) for 24 hours. The resulting transcriptional profiles revealed strikingly similar results (Figure 2A-C; Figure S1A). Specifically, we observed a significant overexpression of early genes in the CoA biosynthesis pathway, including *panB, panG*, *panC, panD and coaX* (Figure 2A-C). This observation is consistent with the current understanding that POA interferes with CoA biosynthesis through disruption of PanD activity[18,19]. Unexpectedly, expression of most other genes of the CoA pathway were not greatly altered by exposure to PZA, POA and MZA with the exception of the final gene of the pathway *coaE* which was down-regulated by 1.5 to 2-fold relative to the no drug control. Due to the central role of CoA in lipid metabolism, we also observed differential expression of numerous genes involved in fatty acid synthesis (FAS) and utilization. The majority of genes associated with the FAS-I and FAS-II systems that are critical for long-chain and mycolic acid biosynthesis, respectively, were overexpressed under all three treatment conditions (Figure 2A-C,F). Interestingly, we also observed down regulation of the large gene cluster for synthesis and export of phthiocerol dimycocerosate (PDIM) virulence lipids, as well as differential expression of several other less well characterized genes potentially involved in polyketide and fatty acid synthesis (Figure 2F,G). Beyond lipid synthesis and metabolism, there were also indications of altered respiratory activity and oxidative stress with induction of genes for fumarate reductase, the microaerophilic-type cytochrome *bd* oxidase and catalase peroxidase *katG* (Figure 2F).

These similarities in responses of *M. tuberculosis* to MZA, PZA and POA exposure suggests mechanistic conservation in their antimicrobial action. Our observation that deletion of *pncA* conferred a slight increase in the MIC for MZA led us to speculate that POA may be released from MZA in a PncA-dependent manner. To understand the role of PncA in response of *M. tuberculosis* to MZA, we performed RNA-seq using the H37Rv Δ*pncA* strain exposed to PZA or MZA (Figure 2D,E). As anticipated, the transcriptional profile of H37Rv Δ*pncA* showed no

significant differences between the PZA-treated and untreated control (Figure 2D; Figure S1B). In
contrast, the transcriptional profile of H37Rv Δ*pncA* exposed to MZA was distinct from the untreated
control, suggesting that MZA exerts additional stress onto bacilli beyond PZA release (Figure 2E;
Figure S1B). Despite this distinction, there were only 12 transcripts that met a threshold cutoff for
significant differential abundance (2-fold change, adj *p*-value ≤0.05), none of which were
immediately informative regarding the PncA-independent action of MZA.

Release of PZA and POA from MZA. Based on the distinctions in antitubercular activity of PZA
and MZA, and on the profoundly different transcriptional responses of H37Rv and H37Rv Δ*pncA*
to MZA exposure, we hypothesized that MZA likely acts through two interacting mechanisms, one
that is PncA-dependent and another that is PncA-independent. To parse these mechanisms, we
assessed MZA metabolism in strains with and without PncA activity. We began by characterizing
the rate of PZA release and conversion to POA. Exponentially growing H37Ra (PncA-proficient)
cultures were amended with 250 µM MZA and abundance of MZA, PZA and POA was measured
in cell extracts and media over an 8-hour time course via LC-MS. In the first 4 hours, cell-associated
MZA rapidly accumulated then steadily declined, while PZA levels rose rapidly from 17 µM to 138
µM (Figure 3A). With continued incubation, levels of MZA and PZA declined to 34 µM and 121 µM,
respectively. Meanwhile, limited POA accumulation was observed in cell lysates during the time
course. The concentration of MZA present in the supernatant steadily declined from 250 to 100 µM
over the eight-hour exposure, during which we also observed a steady increase of PZA from 4 to
67 µM and POA from 1 to 100 µM, respectively (Figure 3B).
To determine whether PncA is essential for formation of POA from MZA, a similar
experiment was performed with overnight exposure using the vaccine strain *M. bovis* BCG which
lacks pyrazinamidase activity due to a loss-of-function mutation in *pncA*. Following overnight
exposure, cell-associated MZA was near the limit of detection for both BCG and the control strain
H37Ra. Both strains showed accumulation of high levels of PZA, whereas only H37Ra showed an
abundant level of POA (Figure 3C). Together, these data demonstrate a rapid conversion of MZA
to PZA, followed by PncA-dependent conversion of PZA to POA.

MZA promotes thiol stress. To further characterize the PncA-independent mechanism of MZA
action, we selected for spontaneously resistant mutants of H37Rv, H37Rv Δ*pncA* and BCG.
Selections were performed using solid medium containing a geometric series of concentrations
from 0.28 to 0.9 mM MZA. No resistant isolates were recovered from the H37Rv background at any
of the tested concentrations. In contrast, H37Rv Δ*pncA* and BCG strains yielded mutants resistant
to 0.41 mM MZA at a frequency of $10^{-5.9}$, and to 0.61 mM MZA at a frequency of 10^{-7}. No resistant
isolates were observed for either H37Rv Δ*pncA* or BCG at 0.9 mM MZA.
After confirming resistance phenotypes, six independently selected isolates of H37Rv
Δ*pncA* and four of BCG were analyzed using whole genome sequencing. All ten isolates had single
nucleotide mutations within the promoter region of *mscR* (Figure 3D) encoding mycothiol reductase
which is critical for formaldehyde tolerance of mycobacteria[20]. To confirm the protective nature of
mutations in the *mscR* promoter region against aldehyde stress, we exposed wild-type and MZA
resistant mutants to methenamine, a bactericidal antibiotic that acts through the spontaneous
release of formaldehyde and is commonly used to treat urinary tract infections[21]. While the MIC of
methenamine for the wild-type BCG strain was 0.9 mM, the MZA resistant strains showed
resistance to 7.1 mM methenamine (Figure 3E,F).
To confirm whether the identified promoter mutations conferred MZA resistance through
increased MscR expression, we constructed GFP reporter fusions to the *mscR* promoters from
wild-type and C-15T resistant variant in the integrative reporter plasmid pYUB1227. Reporter
constructs were integrated at the L5 attachment site in BCG and confirmed strains were grown to
exponential phase. After fluorescence was normalized to the OD_{600} a 1.7 fold increase in
fluorescence was observed for the *mscR* promoter variant relative to the wild-type *mscR* promoter
(Figure 3G) confirming that resistance was due to increased expression of mycothiol reductase. To
validate the role of *mscR* in intrinsic resistance of *M. tuberculosis* to MZA we constructed a H37Rv
Δ*mscR* strain and assessed MZA susceptibility. This strain showed an eight-fold enhanced level of
susceptibility relative to the parental control (Figure 3H).

Given the role of MscR in MZA susceptibility and resistance, we reasoned that cysteine
supplementation would antagonize MZA action through alleviating thiol stress. To test this
hypothesis, BCG wild-type and *mscR* promoter variants were exposed to inhibitory concentrations
of 0.9 mM methenamine or MZA in the presence of varying concentrations of cystine as a source
of cysteine (Figure 3I,J). For wild-type BCG, 4.1 mM cysteine supplementation was highly
antagonistic for both MZA and methenamine. For the *mscR* promoter mutant, cysteine
supplementation was highly antagonistic for MZA at an eight-fold lower concentration than was
required for the wild-type strain. Together, these data demonstrate a central role for thiol stress in
MZA action and establish increased MscR expression as a mechanism for MZA and methenamine
resistance.

Genome-wide fitness profiling and chemical biology reveal the unique self-potentiating
mechanism of MZA. To better understand pathways that modulate resistance and susceptibility
of *M. tuberculosis* to PZA, POA and MZA, we conducted fitness profiling by using Tn-seq. A
saturated library of ~70,000 transposon mutagenized H37Ra was prepared using the *himar1*
mariner transposon[22]. Colonies were harvested and homogenized to form a pooled library and
aliquots were precultured in the absence of drug. Exponentially growing libraries were then
subcultured and grown for 5 generations in the absence of drug or in the presence of a
concentration of PZA (233 µM), POA (105 µM) or MZA (112 µM) that resulted in a two-fold
extension of the generation time. These concentrations and number of generations were chosen
to allow for the highest resolution of gain- and loss-of-fitness conferred by transposon insertion.
Following growth, cells were harvested, genomic DNA was extracted and deep-sequencing of
transposon junctions was performed to determine relative abundance of each insertion[8]. Intra-
population fitness was determined by comparing abundance of individual insertion sites before and
after enrichment using a previously established methodology as described in Opijnen et al. 2009[23].
To determine the change in fitness of specific Tn-insertion mutants, the relative fitness of each
insertion site was evaluated by calculating the change in fitness with drug compared to the absence
of drug. Insertions with relative fitness scores > 1 were considered gain-of-fitness mutants, while
relative fitness scores < 1 were considered loss-of-fitness mutants. Consistent with our previous
findings on PZA and POA susceptibility[8], insertions in genes involved in phosphate utilization and
transport (*pstA1*, *pstC2*, *pstS3*, *phoT*), cell wall stress response regulator (*sigE*), and acyl-CoA
synthetase (*fadD31*) demonstrated substantially greater fitness in the presence of all 3 drugs
(Figure 4A-C). We also observed a consistent gain-of-fitness for mutants with insertions in the
PDIM locus which correlates with our RNA-seq data and previous indications that PDIM mutants
are moderately resistant to PZA[19]. In addition, *pncA* insertions were found to have a gain-of-fitness
in PZA and MZA treated samples, but not POA treated samples, supporting our observation that
PncA plays a supporting role in MZA activity (Figure 4A-C). Further, insertions in the *mscR* locus
harbored substantial fitness defects in MZA treated samples relative to PZA and POA treated
samples (Figure 4C). This finding is consistent with our observation that MZA susceptibility is
modulated by MscR activity in mycobacteria. Interestingly, strains with insertions in *sseA*, recently
shown to be critical for maintaining thiol homeostasis in *M. tuberculosis*[24], showed a substantial
loss-of-fitness in the presence of PZA, POA and MZA, consistent with a direct connection between
thiol stress and activity of these drugs. Collectively these data indicate MZA, PZA and POA show
conserved interactions within *M. tuberculosis* metabolism, yet, that MZA action is distinct due to its
unique ability to exacerbate thiol stress.

Since MZA action shares many features with PZA but does not require exposure to low pH
for this aspect of its activity, we next evaluated whether aldehyde can potentiate PZA action *in lieu*
of exposure to low pH. To do so, we conducted a DiaMOND assay[25] testing for synergy between
methenamine and PZA at circumneutral pH (Figure 4E). While the individual MIC of PZA at neutral
pH surpassed 8 mM and MIC of methenamine was 0.44 mM, the combination of the two produced
a Fractional Inhibitory Concentration Index (based on the IC_{90} of individual drugs) (FICI) below 0.5
indicative of synergy[25], indicating that aldehyde stress sufficiently potentiates PZA activity. To
further characterize the synergistic dual action of MZA we assessed whether the common point for
synergy between aldehyde stress and PZA action was the depletion of CoA pools[26]. To test this
hypothesis, we measured intracellular CoA abundance using H37Ra treated with formaldehyde,

PZA, MZA, or a combination of PZA and formaldehyde. Reduction of CoA in PZA treated cells was
reproducible and consistent with previous reports[27] (~24%), but did not meet statistical significance
in our assay. CoA reduction in cells treated with MZA (21%), formaldehyde (34%) and the
combination of formaldehyde and PZA (30%) was highly robust and significant (Figure 4F).
Together, these data demonstrate that the synergistic dual action of MZA is driven through
collateral impairment of CoA metabolism.

MZA shows superior potency relative to PZA against intracellular *M. tuberculosis*. Given the
critical role of cell-mediated responses in PZA efficacy and the growing problem of drug resistant
M. tuberculosis, we assessed the ability of MZA to kill intracellular *M. tuberculosis* in resting or IFN-
γ activated macrophages. RAW 264.7 macrophages were exposed to *M. tuberculosis* H37Rv or
H37Rv Δ*pncA* at a multiplicity of infection of 1:1 and extracellular bacteria were removed by
washing adhered macrophages. For assessing the impact of macrophage activation on MZA
efficacy, 5 ng/ml IFN-γ was added one day prior to *M. tuberculosis* infection and replenished every
other day. MZA and PZA were used at a concentration of 0.9 mM and compared to treatment with
no drug (DMSO). At the concentration tested, PZA showed limited activity against H37Rv and no
activity against H37Rv Δ*pncA* in resting and activated macrophages (Figure 5A-D). In contrast,
MZA caused a greater than 3 $\log_{10}$ decrease in H37Rv viability by day four of treatment in both
resting and activated macrophages, and after a week, bacterial loads fell below the detection
threshold (Figure 5A,B). Comparable findings were also observed when macrophages were
infected with H37Rv Δ*pncA* (Figure 5C,D). Taken together these data demonstrate that the robust
bactericidal activity of MZA is both independent of macrophage potentiation and highly effective
against intracellular PZA resistant isolates.

316
317   **Discussion**
318
In this study we employed a combination of approaches in functional genomics, bacterial
physiology and chemical biology to characterize the synergistic dual-action *M. tuberculosis*-specific
mechanism of the PZA analog MZA. Our analyses revealed a striking overlap in a central
component of the mechanisms of action of PZA and MZA while also resolving the underlying basis
for the peculiar difference in their antitubercular activity. These findings led us to identify a self-
potentiating mode of action specific to MZA that is both unique from and more potently bactericidal
than its PZA counterpart. The unique features of MZA, which include rapid killing, high target
specificity, low rates of resistance and independence from environmentally-driven potentiation, can
be attributed to the drug-specific release of aldehyde resulting in disruption of thiol homeostasis.
The rate of aldehyde release, as determined by mass spectrometry, takes place over the course of
hours, which allows for elimination of bacilli from infected macrophages within a matter of days.
The superior feature of MZA is orchestrated by the synergistic actions of its two components,
aldehyde release followed by PZA collateral susceptibility (Figure 6). The rapid engagement of
aldehyde elicits bactericidal activity against both actively growing and dormant cells alike while
simultaneously sensitizing any survivors to the activity of PZA. This potentiation event ensues via
depletion of biologically active thiols, resulting in bacterial growth arrest and increased demand for
CoA with consequent disruption of fatty acid biosynthesis, a pathway known to be perturbed by
PZA[15,28]. The mechanism proposed here is further confirmed by synergistic activity of methenamine
with PZA and the antagonistic activity of cysteine against MZA.
The robust activity of MZA against non-growing *M. tuberculosis* suggests that entry into the
cytoplasm is likely a passive event. Hence, MZA is effective at killing *M. tuberculosis* independent
of metabolic activity or growth state of the cells. Passive entry of MZA into the cytoplasm also
supports a low likelihood for emergence of drug resistant strains via mutations that alter drug
transport. While low-level resistance to aldehyde release can be achieved via activation mutations
in the *mscR* promoter, achieving full resistance to MZA will likely be a challenge due to the
promiscuous nature of aldehyde-mediated damage. Moreover, we find that PZA resistant strains
of *M. tuberculosis* and *M. bovis* BCG with *mscR* promoter-up mutations still show measurable
susceptibility to MZA, suggesting either that the rate of internal aldehyde release is greater than

the MscR turnover rate or that other aldehydes that cannot be effectively neutralized by MscR are
released by MZA. In either case, the bacilli will be faced with a greater adaptive burden to develop
full resistance to MZA (Figure 6).
Given the pivotal role of PZA in TB treatment regimens and the limited activity of this drug
in models of compromised immunity[11], it will be critical to resolve the importance of aldehyde-driven
thiol stress in its antitubercular efficacy. In the present study, aldehyde release from MZA was able
to reduce bacterial loads in resting macrophages to levels below the limit of detection in under a
week. Based on loss of PZA efficacy in the athymic mouse model[11], it has been speculated that
antitubercular activity of PZA may be diminished in the context of T cell deficiency or dysfunction
due to insufficient macrophage activation, resulting in reduced treatment efficacy and higher rates
of drug resistance[9]. Circumnavigating the need for host-driven potentiation of PZA may prove to be
a valuable approach in the treatment of *M. tuberculosis* in the context of compromised immunity.

Interestingly, it has recently been recognized that production of host-derived aldehydes are
modulated via IFN-γ activation and play a role as antimicrobial effectors against intracellular
pathogens, such as *M. tuberculosis* and *Francisella tularensis* [29,30]. Whether host-derived
aldehydes participate in the bactericidal activity of PZA *in vivo* has yet to be determined, but, may
represent an opportunity for host-directed therapy for bolstering the action of this critical
antitubercular agent. Further, microbe-derived aldehyde production may also represent an
opportunity that can be explored for novel therapeutic discovery. For example, selective disruption
of cytokinin metabolism has been shown to result in accumulation of toxic aldehyde species in *M.
tuberculosis* and further sensitize the bacilli to other host stressors[31]. Further, as noted by Darwin
and Stanley[29], mycobacterial glycolysis may represent an exploitable target for driving toxic
aldehyde accumulation in *M. tuberculosis*[32]. Along these lines, it is curious to note that strains
bearing loss-of-function mutations in *glpK* (encoding glycerol kinase) show a selective advantage
both in humans and in mice treated with antitubercular agents[33,34]. Since disruption of glycerol
phosphate synthesis would reduce the level of endogenous metabolic aldehydes through limitation
of glyceraldehyde phosphate and dihydroxyacetone phosphate synthesis, glycerol kinase likely
represents a key mediator in endogenous mycobacterial aldehyde production[29]. In strong support
of this concept, Dick and colleagues previously determined that mutations in *glpK* are associated
with PZA resistance *in vitro*[19]. Our findings, coupled with the emerging role of host- and microbe-
derived aldehydes in compromising the fitness of *M. tuberculosis*, advocate for further evaluation
of the role of aldehyde-mediated potentiation of antimicrobial agents such as PZA for improved
therapeutic approaches.

380
381 **Materials and Methods**
382
Standard growth conditions. Middlebrook 7H9 broth (Difco) supplemented with 10% Middlebrook
OADC (Difco), 0.02% (V/V) glycerol and 0.05% (V/V) tyloxapol were used for all experiments
involving liquid *M. tuberculosis* and *M. bovis* cultures. Depending on experimental requirements,
the media were adjusted to pH 7.2 or pH 5.8. Cultures were incubated shaking at 100 RPM 37ºC.
All solid media used in experiments involving *M. tuberculosis* and *M. bovis* were based on
Middlebrook 7H10 (Difco) supplemented with 10% Middlebrook OADC, 0.02% (V/V) glycerol. For
M. smegmatis the same media were used with the exclusion of OADC and replacing glycerol with
0.2% (W/V) dextrose. ESKAPE pathogens and *E. coli* (for plasmid amplification) were propagated
in Luria-Bertani broth (LB, Difco) under standard growth conditions (37°C shaking at 250 RPM).

Strain construction. The knockout strains (H37Rv Δ*pncA* and H37Rv Δ*mscR*) and were
constructed using the ORBIT system[35]. H37Rv was electroporated with pKM461. After selection on
50 µg/mL kanamycin, H37Rv pKM461 was induced with anhydrotetracycline and electroporated
with pKM464 and the position specific *pncA* oligonucleotide 5'-
TACCTCGGCGCCACGGCGGCGGACCCGGCCCGCGCCCGGTGGCTCCT
GCACTTCGGCATGGTGGGCCGCAGGTTTGTACCGTACACCACTGAGACCGCGGTGGTTGA
CCAGACAAACCCCTCGACTCGCTTCCGACAGCACCTCGAAGACCGCTTCGGGTGCGTGAG

CACGCTGGGCGGTTCGCAGTG-3' or *mscR* 5'-GCGCGCATGGTCAGCGACGCTACACCGT
AGGTTGGACACCATGAGTCAGACGGTGCGCGGTGTGATCGCAGGTTTGTCTGGTCAACCAC
CGCGGTCTCAGTGGTGTACGGTACAAACCAAGGTATTGCGTTCGGTGGTGATGTTGTGATG
GCCGCCATCGAGCGCGTCATCACCCACGGCACCTTCGA-3'. Transformed bacteria were then
plated on supplemented 7H10 medium containing 50 µg/mL hygromycin B and 10% sucrose to
select for recombinants and counter select against pKM461. Individual colonies were then re-
streaked, confirmed via PCR, and full genome sequenced to validate desired modification and no
additional mutations.
Fluorescent reporter *mscR* strains were constructed using a modified integrative reporter
plasmid pYUB1227. Restriction enzymes *Pvu*I and *Mfe*I (NEB) were used to prepare the backbone
and the prompter inserts, followed by a T4 ligation. Primers used to validate promoter changes
were mscRF 5'-TTGCAACGCATCCCTGATCT-3' and mscRR 5'-AGGGCAGATTGTGTGGACAG-
3'. Additional validation was performed via Nanopore sequencing. Plasmid amplification was
performed in *E. coli* DH5α. Plasmids were purified using a QIAprep Spin Miniprep Kit (cat.# 27104).
BCG Pasteur 1173P2 was then transformed using previously prepared plasmids via standard
electroporation protocol. Successful transformants were selected for on standard media amended
with 50 µg/mL hygromycin B and validated via genomic DNA extraction and PCR using validation
primers described above.

Determination of minimum inhibitory concentrations. Bacteria were grown in 30 mL PETG
square media bottles (Nalgene) containing 5 mL of 7H9 media under standard growth conditions
(37°C shaking at 100 RPM) to OD_{600} of 0.15-0.25. Bacteria were then diluted to OD_{600} of 0.01 and
allowed to incubate for 2 weeks in presence of the compound(s) being tested. Concentrations of
compounds were determined as a function of a geometric series (0.05, 0.1, 0.2 ,0.4, 0.81, 1.62,
3.25 and 6.5 mM) for PZA, (0.06, 0.11, 0.22, 0.45 and 0.9 mM) for MZA and (0.06, 0.11, 0.22, 0.44,
0.88, 1.75 and 3.5 mM) for methenamine. MIC values were then calculated by plotting inhibition
curve and calculating based on the appropriate slope intercept.

Antagonism assays. Bacteria were prepared in the same way as for MIC assays and were then
inoculated into 7H9 media containing either 0.9mM methenamine, 0.9mM MZA or DMSO (vehicle
control) and antagonist (pantothenate or cystine) at indicated concentrations. Pantothenate was
used at a single concentration of 230 µM, cystine concentrations ranged from 15 µg/mL to 500
µg/mL in 2-fold increments. The media pH was adjusted to 7.2 and cells were incubated in 96-well
plated for 8 days at 37°C with no shaking. Measurements were performed using a BioTek Synergy
H1 plate reader (Agilent). All experiments were performed in triplicate.

Bacterial kill curves. To assess the bactericidal activity of MZA and methenamine, against
exponentially growing cells, bacteria were grown in 30 mL bottles containing 5 mL of 7H9 media
under standard growth conditions (37°C shaking at 100 RPM) to OD_{600} of 0.15-0.25. Bacteria were
then diluted to OD_{600} of 0.03 (approximately $5X10^6$ bacilli) and plated on 7H10 media to determine
the starting population of bacteria. After addition of (0, 0.28, 0.41, 0.61 or 0.9mM of MZA) and (0.06,
0.11, 0.22, 0.44, 0.88, 1.75 and 3.5 mM) µg/mL Methenamine, CFU counts were determined at 0,
2-, 4-, 7- and 14-day time intervals.
To assess bactericidal activity of PZA and MZA against nutrient limited *M. tuberculosis*,
bacteria were grown in 30 mL bottles containing 5 mL of 7H9 media under standard growth
conditions (37° C shaking at 100 RPM) to OD_{600} of 0.15-0.25. Bacteria were then washed three
times with 1xPBS saline to remove any residual media and resuspended in 1xPBS saline amended
with 0.05% tyloxapol. Prior to addition of the drugs CFU/mL were determined by serial dilution
plating on 7H10 media, 900 µM PZA, MZA or DMSO were then added to the buffer and CFU
measurements were taken at predetermined intervals 0, 2-, 3-, 4-, 7- and every 7 days afterwards
until 62-day timepoint was reached. For MBC measurements under starvation conditions with
multiple treatments, bacteria were prepared in the same way as for a single treatment. However,
at every timepoint culture was washed three times to remove residual antibiotics and split into two
cultures. One of the two cultures were treated with DMSO while the other was amended with fresh
MZA for a final concentration of 900 µM.

RNA-seq. For bulk RNA-seq library preparation and sequencing, exponentially growing H37Rv and H37Rv Δ*pncA* were grown to the OD$_{600}$ of 0.15 and inoculated with 200 µM MZA, PZA or equivalent volume of DMSO (as a vehicle control). Cells were then incubated under standard condition for 24 hours prior to harvesting. Following incubation, TRIzol RNA purification was performed. Cells were chilled, pelleted and resuspended in a 500 µL of TriReagent (Invitrogen) with 1% polyacryl carrier (Molecular Research Center) and lysed using 250 µL of 0.1 mm zirconia beads (BioSpec). Samples were then centrifuged and supernatant was transferred to a clean 2 mL microcentrifuge tube and combined with 50 µL of 5-bromo-3-chloro-propane. Following a 10-minute incubation samples were centrifuged aqueous phase was removed and transferred to a clean 1.5 mL Eppendorf tube where samples were treated with 250 µL of isopropanol. Following a 10-minute incubation samples were centrifuged again, isopropanol was removed and RNA coating the tube was washed with 300 uL of 75% EtOH. After the final wash, RNA was dried and eventually resuspended in 50 uL of DEPC-treated water (Invitrogen). A DNA-free (™) DNAse treatment (Invitrogen) was used to remove residual DNA from the samples prior to sequencing. Samples were treated with Ribo-Zero Plus kit and Libraries were prepared using an Illumina Stranded Total RNA Prep Ligation. Sequencing was performed at the University of Minnesota Genomic Center using a 50 paired-end NovaSeq S-prime platform.

For bulk RNA-seq analysis, the pipeline used for preprocessing raw fastq files can be found at https://github.com/MDHowe4/RNAseq-Pipeline. Quality control of RNA sequencing read quality was assessed with FastQC. Read length thresholding and t-overhang trimming of forward and reverse reads was completed with Cutadapt with minimum read length cutoff of 30 bp. Reads were mapped to the M. tuberculosis H37Rv reference genome (NC_000962.3) using the STAR aligner without spliced alignment detection (--alignIntronMax 1)[36]. Total reads per gene were counted using featureCounts[37]. Genes with <10 reads mapped occurrences across wild-type and mutant experiments were not included in further analysis. Differentially expressed genes (DEGS) were found using the negative binomial generalized linear model of DESeq2[38]. Genes were considered differentially expressed if they displayed a log$_2$ fold-change ≥ 1.5 or ≤ -1.5 and adjusted *P*-value ≤ 1x10^{-6}. Volcano plots were generated using the EnhancedVolcano package.

Mass spectrometry. For mass spectrometry analyte preparation, for each sample, a total of 40 mL of bacteria were grown under standard growth conditions at pH 5.8. Upon reaching OD$_{600}$ of 0.4, bacteria were inoculated with either 250 µM MZA or PZA. After three days of treatment the samples were pelleted. Cell extract samples were prepared for mass spectrometry analysis by resuspending pellets in 1 mL of extraction buffer (40% methanol, 40% acetonitrile and 20% water) and bead beaten using 250 µL of 0.1 mm zirconia beads (BioSpec). Samples were then centrifuged to remove the solids. Liquid fractions were then filtered using 22 µm filter. A portion of the filtrate was then used to determine protein content using a Pierce™ BCA Protein Assay Kit (Thermo Scientific). The remaining samples were further purified using a 3 kDa column (PALL Life Sciences). Supernatant sample preparation was performed in the identical way as cell extract, except for the omission of the first bead beating step. Bacterial cell extracts were diluted 10-fold (10 µL in 100 µL) in DI water. 100 µL of diluted solution was treated with 100 µL methanol: water solution containing *para*-aminosalicylic acid (PAS, internal standard), mixed well by vortexing, centrifuged (2 min, 15000 × g), and supernatant (150 µL) was analyzed by LC-MS/MS.

For LC-MS/MS analysis, reverse-phase LC was performed on a Zorbax Eclipse XDB- C8 column (150 mm, 4.6 mm, 5 µm; Agilent) using Shimadzu UFLC XR instrument. The elution gradient was carried out with binary solvent system consisting of 0.1% formic acid in H$_2$O (solvent A) and 0.1% formic acid in MeCN (solvent B). A linear gradient profile with the following proportions (v/v) of solvent B was applied (t (min), %B): (0, 5), (1, 5), (4, 95), (6, 95), (7, 5), (8, 5) with 2 min for re-equilibration to provide a total run time of 10 min. The flow rate was 0.5 mL/min and the column oven was maintained at 40°C. The injection volume was 10 µL. The retention times for MZA, PZA, POA and PAS (internal standard) were 3.5, 2.6, 3.8 and 6.8 min, respectively. MS/MS analysis were carried out using triple quadrupole/linear ion trap instrument (AB SCIEX QTRAP 5500). MZA, PZA and POA peak areas were calculated using MultiQuant, version 2.0.2. MZA, PZA and POA areas were normalized to p-amino salicylate areas and the MZA, PZA and POA concentration in samples were determined using the standard curves of MZA, PZA and POA. For the preparation

of MZA, PZA and POA standard curve, average area of standard 0 was subtracted from MZA, PZA
and POA of all higher standards prior to area normalization. Sample concentrations were back
calculated to account for dilutions. Each sample was performed in triplicate.

Transposon mutagenesis and transposon sequencing. *M. tuberculosis* H37Ra was
mutagenized with the mariner *himar1* transposon using the phAE180 temperature-sensitive
mycobacteriophage[39]. Approximately 70,000 independent mutants were selected on 7H10 media
amended with 50 μg/mL kanamycin. Transposon mutants were pooled, homogenized, then
aliquoted into 25% (V/V) glycerol stocks and stored at -80°C. Growth curves to establish 50%
inhibition of growth rate using a geometric series of drug concentrations were carried out using 7H9
medium at a pH of 6.0. Cultures were seeded at OD_{600} of 0.01 in 80 mL of medium using transposon
mutants pre-cultured in a shaking incubator for 48 hours at 37°C. Enrichment was carried out in
the presence or absence of drugs for 5 generations as determined by OD_{600}. After 5 generations,
cultures were flash frozen in liquid nitrogen for genomic DNA (gDNA) extraction. gDNA extraction
was carried out using a previously established protocol[40]. Library prep and sequencing was
performed at the University of Minnesota Genomics Center with read mapping performed using a
previously established protocol[8]. DNA fragmentation and Illumina P7
(CAAGCAGAAGACGGCATACGAGAT) ligation was performed. Amplification of transposon site
junction was carried out using a P7 (CAAGCAGAAGACGGCATACGAGAT) and mariner-specific
Mariner_1R_TnSeq_noMm   primer
(TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCGGGGACTTATCAGCCAACC).
Amplification of himar1-enriched samples was performed using a P5 indexing primer
(AATGATACGGCGACCACCGAGATCTACAC[i5]TCGTCGGCAGCGTC; [i5] barcode sequence)
and a P7 primer HotStarTaq master mix kit (Qiagen) after a 1:50 dilution. Sequencing was
performed on an Illumina NovaSeq. The 5' ends of reads were trimmed to remove adapter and
transposon sequence using Cutadapt to leave the TA-gDNA junction intact[41]. Trimmed reads with
less than 18 bp were discarded and reads were mapped to the M. tuberculosis H37Ra genome
(GenBank Accession: CP000611) with 1 bp permitted mismatch using Bowtie 2[42] (PMID:
22388286). Mapped reads were printed in SAM format and counted per TA dinucleotide sight using
SAMreader_Ta script[43]. Scripts used for read processing may be found on GitHub
(https://github.com/MDHowe4/Himar1-TnSeq-Pipeline).
The relative abundance of insertions at TA sites was calculated for each sample. TA sites
lacking mapped insertions at the start of enrichment were omitted. TA sites without mapped
insertions at the end of enrichment that were present at the start of the enrichment were set to the
limit of detection without contributing to the total number of insertions. Relative fitness of site-
specific mutants were calculated with a previously characterized function using the population
expansion factor, as well as the relative number of insertions at the beginning and end of
enrichment[23].

Isolation of spontaneous mutants. To isolate spontaneously MZA resistant mutants, H37Rv
Δ*pncA* and BCG cells were grown up to mid-log phase for selection. 1 mL of culture diluted to OD_{600}
of 0.3 was plated onto standard 7H10 agar supplemented with 409 μM MZA for H37Rv Δ*pncA*, and
607 μM for BCG. Plates were then incubated at 37°C in stationary incubator for 14-21 days.
Resistant mutants were then picked and streaked again for isogenic backgrounds, then grown
again for 14-21 days at 37°C in a stationary incubator. Isogenic colonies were picked, then grown
in standard 7H9 broth for 14 days followed by genomic DNA extraction using a the previously
described DNA extraction methodology. Illumina sequencing was performed at SeqCenter. Read
alignment and variant calling to the H37Rv reference genome (NC_000962.3) and BCG reference
genome (BCG Pasteur 1173P2) was performed using Breseq version 0.36.0.

Coenzyme A detection assays. Bacteria were prepared for extraction in the same way as
described in mass-spectrometry methods. 80 mL per treatment sample of H37Ra was grown to an
OD_{600} of 0.13 in 7H9 media in the shaking incubator at 100 RPM and 37°C. All of the samples were
then treated with 200 μM of either PZA, MZA, formaldehyde or formaldehyde + PZA. After an eight-
hour exposure cells were harvested and resuspended in 400 μL of ice cold 1xPBS and lysed using

zirconia beads. Samples were then centrifuged and filtered using 0.22-µm filter. 50 µl aliquots were
then taken to perform a BCA assay (Pierce™ BCA Protein Assay Kit, ThermoScientific). The
remaining extracts were then filtered one more time using a using a 3 kDa column (PALL Life
Sciences). Twice filtered lysate was then used to perform fluorometric assay using the Free
Coenzyme A Assay kit (Sigma-Aldrich). The readout was then normalized to the protein
abundance. All assays were performed in triplicate.

Macrophage Infections. RAW 264.7 macrophages were prepared by seeding density of 1.0-2.0 x
10^5 cells per well in DMEM/F-12 medium containing 10% FBS without antibiotics (DMEM/F-12
complete) in 12 well (1 mL/well) plates. In order to allow for cells to adhere, macrophages were
incubated overnight in a humidified 5% CO_2 chamber. The following day cells were rinsed with
Hank's buffer three times and replenished with fresh DMEM/F-12 media. A 5 ng/mL of IFN-γ was
added to the well containing macrophages meant to be activated. Following a 14-16h incubation,
macrophages were washed again in Hank's buffer and replenished with fresh, drug and interferon
free DMEM media. Upon transfer of macrophages into BSL-3 space, media was removed, cells
were washed with Hank's buffer three times and fresh DMEM media containing *M. tuberculosis*
H37Rv. An MOI of 1:1 was used for the initial infection. To prepare bacteria, H37Rv was grown
under standard growth conditions. Upon reaching OD_{600} of 0.25, culture was washed three times,
resuspended, and then diluted in DMEM media to the 1.0-2.0 x 10^5 CFU/mL. Following a 2h
infection, infected macrophages were washed three times with Hank's buffer and resuspended in
complete DMEM media amended with 900 µM MZA, PZA or DMSO (as a vehicle control) as well
as 5ng/ml of IFN-γ for activated macrophages. Media was changed daily for the duration of the
experiment with re-induction via IFN-γ taking place on every other day. During every media change,
cells were washed three times with Hank's buffer and adherence and appearance of macrophages
was monitored via a stereoscope. On plating days, cells were washed as described above, and
macrophages were lysed via TritonX-100. The lysate was then serially diluted in 0.05% Tyloxapol
1xPBS pH 7.2 buffer using 10-fold dilution in a 96well plate format and plated on 7H10 media.
Plates were incubated for two weeks and colonies counted for CFU determination. All experiments
were performed in triplicate.

MZA synthesis and NMR analysis. Formaldehyde 37% in water (89 µL, 1.9 mmol) was prepared
with PZA (200 mg, 1.6 mmol) and morpholine (1.1 mL, 12.9 mmol) in 100% ethanol solvent with
one drop of concentrated HCl. The reaction was refluxed for 6 hours and neutralized with one drop
of 1 N NaOH. The crude mixture was evaporated and separated via silica gel flash chromatography
with methanol/dichloromethane solvent system. The pure compound (>95%) was a white solid. The
calculated yield was 63.7%. Observed (M+H) $^+$ of 223.0 m/z with mass spectrometry analysis.
^{1}H NMR (400 MHz, CDCl$_3$-d, ppm): δ = 9.35 (s, 1), 8.71 (d, 1), 8.48 (d,1), 8.12 (s, 1), 4.29
(d, 2), 3.65 (t, 4), and 2.58 (t, 4). ^{13}C NMR (100 MHz, DMSO-d$_6$ ppm): δ = 164.2, 148.0, 145.0,
144.1, 143.7, 66.4, 60.8, and 50.5. Purity: >99%.

Quantification and statistical analyses. Where applicable, 95% confidence intervals were used
to determine statistical significance. Numbers of biological replicates and statistical tests that were
employed are described in respective figure legends.

Materials availability. Materials will be made available upon reasonable request and may require
completion of a Material Transfer Agreement and payment for related expenses.

Data and code availability. All data are available in the main manuscript or supplemental
information. Primary sequencing data from RNA-seq, Tn-seq, and whole-genome sequencing
studies are publicly available through the National Center for Biotechnology Information via SRA
link https://www.ncbi.nlm.nih.gov/sra/PRJNA1104292.
Code for bulk RNA-seq analysis pipeline used for preprocessing raw fastq files can be
found on GitHub at https://github.com/MDHowe4/RNAseq-Pipeline. Code used for read processing
can be found on GitHub https://github.com/MDHowe4/Himar1-TnSeq-Pipeline.
Paste your materials and methods section here.
607

Acknowledgments

We would like to acknowledge the following individuals and institutions for their invaluable contributions to this work. We express our gratitude to the NIH for their generous grant support Grant Numbers: R01 AI123146, T32 GM008244 and T32 HL007741. We are also grateful to the University of Minnesota Genomics Center and University of Minnesota Biosafety Level 3 Program Core Facilities, and SeqCenter for their technical support and expertise. We also would like to acknowledge Biorender which was used for the preparation of the figures. Lastly, we would like to thank Dr. Elise Lamont for her input and support for macrophage work and Dr. Subhankar Panda for determination of MZA purity.

References

1.   Lamont, E.A., Dillon, N.A., and Baughn, A.D. (2020). The Bewildering Antitubercular Action of Pyrazinamide. Microbiol. Mol. Biol. Rev. MMBR *84*, e00070-19. https://doi.org/10.1128/MMBR.00070-19.

2.   Heifets, L., and Lindholm-Levy, P. (1992). Pyrazinamide sterilizing activity in vitro against semidormant Mycobacterium tuberculosis bacterial populations. Am. Rev. Respir. Dis. *145*, 1223–1225. https://doi.org/10.1164/ajrccm/145.5.1223.

3.   Whitfield, M.G., Soeters, H.M., Warren, R.M., York, T., Sampson, S.L., Streicher, E.M., van Helden, P.D., and van Rie, A. (2015). A Global Perspective on Pyrazinamide Resistance: Systematic Review and Meta-Analysis. PloS One *10*, e0133869. https://doi.org/10.1371/journal.pone.0133869.

4.   Zheng, X., Ning, Z., Drobniewski, F., Yang, J., Li, Q., Zhang, Z., and Hu, Y. (2017). pncA mutations are associated with slower sputum conversion during standard treatment of multidrug-resistant tuberculosis. Int. J. Antimicrob. Agents *49*, 183–188. https://doi.org/10.1016/j.ijantimicag.2016.10.012.

5.   Scorpio, A., Lindholm-Levy, P., Heifets, L., Gilman, R., Siddiqi, S., Cynamon, M., and Zhang, Y. (1997). Characterization of pncA mutations in pyrazinamide-resistant Mycobacterium tuberculosis. Antimicrob. Agents Chemother. *41*, 540–543. https://doi.org/10.1128/AAC.41.3.540.

6.   Sun, Q., Li, X., Perez, L.M., Shi, W., Zhang, Y., and Sacchettini, J.C. (2020). The molecular basis of pyrazinamide activity on Mycobacterium tuberculosis PanD. Nat. Commun. *11*, 339. https://doi.org/10.1038/s41467-019-14238-3.

7.   Gopal, P., Sarathy, J.P., Yee, M., Ragunathan, P., Shin, J., Bhushan, S., Zhu, J., Akopian, T., Kandror, O., Lim, T.K., et al. (2020). Pyrazinamide triggers degradation of its target aspartate decarboxylase. Nat. Commun. *11*, 1661. https://doi.org/10.1038/s41467-020-15516-1.

8.   Thiede, J.M., Dillon, N.A., Howe, M.D., Aflakpui, R., Modlin, S.J., Hoffner, S.E., Valafar, F., Minato, Y., and Baughn, A.D. (2021). Pyrazinamide Susceptibility Is Driven by Activation of the SigE-Dependent Cell Envelope Stress Response in Mycobacterium tuberculosis. mBio *13*, e0043921. https://doi.org/10.1128/mbio.00439-21.

9.   Lamont, E.A., and Baughn, A.D. (2019). Impact of the host environment on the antitubercular action of pyrazinamide. EBioMedicine *49*, 374–380. https://doi.org/10.1016/j.ebiom.2019.10.014.

10.   Coulson, G.B., Johnson, B.K., Zheng, H., Colvin, C.J., Fillinger, R.J., Haiderer, E.R., Hammer, N.D., and Abramovitch, R.B. (2017). Targeting Mycobacterium tuberculosis Sensitivity

to Thiol Stress at Acidic pH Kills the Bacterium and Potentiates Antibiotics. Cell Chem. Biol. *24*, 993-1004.e4. https://doi.org/10.1016/j.chembiol.2017.06.018.

11.   Almeida, D.V., Tyagi, S., Li, S.-Y., Wallengren, K., Pym, A.S., Ammerman, N.C., Bishai, W.R., and Grosset, J.H. (2014). Revisiting Anti-tuberculosis Activity of Pyrazinamide in Mice. Mycobact. Dis. Tuberc. Lepr. *4*, 145. https://doi.org/10.4172/2161-1068.1000145.

12.   Cerchiai, E., and Belluomini, R. (1963). [CLINICAL AND BACTERIOLOGICAL FINDINGS ON THE USE OF MORPHAZINAMIDE IN THE TREATMENT OF CHRONIC TUBERCULOSIS]. G. Ital. Chemioter. *10*, 74–78.

13.   Zítková, L., Stastná, J., Tousek, J., and Viklický, J. (1983). Toxicity of morphazinamide compared with pyrazinamide. Czech. Med. *6*, 140–151.

14.   Chung, W.J., Kornilov, A., Brodsky, B.H., Higgins, M., Sanchez, T., Heifets, L.B., Cynamon, M.H., and Welch, J. (2008). Inhibition of M. tuberculosis in vitro in monocytes and in mice by aminomethylene pyrazinamide analogs. Tuberc. Edinb. Scotl. *88*, 410–419. https://doi.org/10.1016/j.tube.2008.06.001.

15.   Dillon, N.A., Peterson, N.D., Rosen, B.C., and Baughn, A.D. (2014). Pantothenate and pantetheine antagonize the antitubercular activity of pyrazinamide. Antimicrob. Agents Chemother. *58*, 7258–7263. https://doi.org/10.1128/AAC.04028-14.

16.   Hu, Y., Coates, A.R., and Mitchison, D.A. (2006). Sterilising action of pyrazinamide in models of dormant and rifampicin-tolerant Mycobacterium tuberculosis. Int. J. Tuberc. Lung Dis. Off. J. Int. Union Tuberc. Lung Dis. *10*, 317–322.

17.   De Oliveira, D.M.P., Forde, B.M., Kidd, T.J., Harris, P.N.A., Schembri, M.A., Beatson, S.A., Paterson, D.L., and Walker, M.J. (2020). Antimicrobial Resistance in ESKAPE Pathogens. Clin. Microbiol. Rev. *33*, e00181-19. https://doi.org/10.1128/CMR.00181-19.

18.   Zhang, S., Chen, J., Shi, W., Liu, W., Zhang, W., and Zhang, Y. (2013). Mutations in panD encoding aspartate decarboxylase are associated with pyrazinamide resistance in Mycobacterium tuberculosis. Emerg. Microbes Infect. *2*, e34. https://doi.org/10.1038/emi.2013.38.

19.   Gopal, P., Yee, M., Sarathy, J., Low, J.L., Sarathy, J.P., Kaya, F., Dartois, V., Gengenbacher, M., and Dick, T. (2016). Pyrazinamide Resistance Is Caused by Two Distinct Mechanisms: Prevention of Coenzyme A Depletion and Loss of Virulence Factor Synthesis. ACS Infect. Dis. *2*, 616–626. https://doi.org/10.1021/acsinfecdis.6b00070.

20.   Vargas, D., Hageman, S., Gulati, M., Nobile, C.J., and Rawat, M. (2016). S-nitrosomycothiol reductase and mycothiol are required for survival under aldehyde stress and biofilm formation in Mycobacterium smegmatis. IUBMB Life *68*, 621–628. https://doi.org/10.1002/iub.1524.

21.   Strom, J.G., and Jun, H.W. (1993). Effect of urine pH and ascorbic acid on the rate of conversion of methenamine to formaldehyde. Biopharm. Drug Dispos. *14*, 61–69. https://doi.org/10.1002/bdd.2510140106.

22.   Rubin, E.J., Akerley, B.J., Novik, V.N., Lampe, D.J., Husson, R.N., and Mekalanos, J.J. (1999). In vivo transposition of mariner-based elements in enteric bacteria and mycobacteria. Proc. Natl. Acad. Sci. U. S. A. *96*, 1645–1650. https://doi.org/10.1073/pnas.96.4.1645.

692 23. van Opijnen, T., Bodi, K.L., and Camilli, A. (2009). Tn-seq: high-throughput parallel
693 sequencing for fitness and genetic interaction studies in microorganisms. Nat. Methods *6*, 767–
694 772. https://doi.org/10.1038/nmeth.1377.

695 24. Nambi, S., Long, J.E., Mishra, B.B., Baker, R., Murphy, K.C., Olive, A.J., Nguyen, H.P.,
696 Shaffer, S.A., and Sassetti, C.M. (2015). The Oxidative Stress Network of Mycobacterium
697 tuberculosis Reveals Coordination between Radical Detoxification Systems. Cell Host Microbe
698 *17*, 829–837. https://doi.org/10.1016/j.chom.2015.05.008.

699 25. Van, N., Degefu, Y.N., and Aldridge, B.B. (2021). Efficient Measurement of Drug
700 Interactions with DiaMOND (Diagonal Measurement of N-Way Drug Interactions). Methods Mol.
701 Biol. Clifton NJ *2314*, 703–713. https://doi.org/10.1007/978-1-0716-1460-0_30.

702 26. Newton, G.L., and Fahey, R.C. (2002). Mycothiol biochemistry. Arch. Microbiol. *178*,
703 388–394. https://doi.org/10.1007/s00203-002-0469-4.

704 27. Gopal, P., Nartey, W., Ragunathan, P., Sarathy, J., Kaya, F., Yee, M., Setzer, C.,
705 Manimekalai, M.S.S., Dartois, V., Grüber, G., et al. (2017). Pyrazinoic Acid Inhibits Mycobacterial
706 Coenzyme A Biosynthesis by Binding to Aspartate Decarboxylase PanD. ACS Infect. Dis. *3*, 807–
707 819. https://doi.org/10.1021/acsinfecdis.7b00079.

708 28. Zimhony, O., Cox, J.S., Welch, J.T., Vilchèze, C., and Jacobs, W.R. (2000).
709 Pyrazinamide inhibits the eukaryotic-like fatty acid synthetase I (FASI) of Mycobacterium
710 tuberculosis. Nat. Med. *6*, 1043–1047. https://doi.org/10.1038/79558.

711 29. Darwin, K.H., and Stanley, S.A. (2022). The aldehyde hypothesis: metabolic
712 intermediates as antimicrobial effectors. Open Biol. *12*, 220010.
713 https://doi.org/10.1098/rsob.220010.

714 30. Berry, S.B., Espich, S., Thuong, N.T.T., Chang, X., Dorajoo, R., Khor, C.-C., Heng, C.-K.,
715 Yuan, J.-M., Fox, D., Anaya-Sanchez, A., et al. (2023). Disruption of Aldehyde Dehydrogenase 2
716 protects against bacterial infection. Preprint at Cold Spring Harbor Laboratory,
717 https://doi.org/10.1101/2023.08.24.554661 https://doi.org/10.1101/2023.08.24.554661.

718 31. Samanovic, M.I., Tu, S., Novák, O., Iyer, L.M., McAllister, F.E., Aravind, L., Gygi, S.P.,
719 Hubbard, S.R., Strnad, M., and Darwin, K.H. (2015). Proteasomal Control of Cytokinin Synthesis
720 Protects Mycobacterium tuberculosis against Nitric Oxide. Mol. Cell *57*, 984–994.
721 https://doi.org/10.1016/j.molcel.2015.01.024.

722 32. Pethe, K., Sequeira, P.C., Agarwalla, S., Rhee, K., Kuhen, K., Phong, W.Y., Patel, V.,
723 Beer, D., Walker, J.R., Duraiswamy, J., et al. (2010). A chemical genetic screen in
724 Mycobacterium tuberculosis identifies carbon-source-dependent growth inhibitors devoid of in
725 vivo efficacy. Nat. Commun. *1*. https://doi.org/10.1038/ncomms1060.

726 33. Safi, H., Gopal, P., Lingaraju, S., Ma, S., Levine, C., Dartois, V., Yee, M., Li, L., Blanc, L.,
727 Ho Liang, H.-P., et al. (2019). Phase variation in  *Mycobacterium tuberculosis glpK*
728 produces transiently heritable drug tolerance. Proc. Natl. Acad. Sci. *116*, 19665–19674.
729 https://doi.org/10.1073/pnas.1907631116.

730 34. Bellerose, M.M., Baek, S.-H., Huang, C.-C., Moss, C.E., Koh, E.-I., Proulx, M.K., Smith,
731 C.M., Baker, R.E., Lee, J.S., Eum, S., et al. (2019). Common Variants in the Glycerol Kinase
732 Gene Reduce Tuberculosis Drug Efficacy. mBio *10*. https://doi.org/10.1128/mbio.00663-19.

35.   Murphy, K.C., Nelson, S.J., Nambi, S., Papavinasasundaram, K., Baer, C.E., and
Sassetti, C.M. (2018). ORBIT: a New Paradigm for Genetic Engineering of Mycobacterial
Chromosomes. mBio *9*, e01467-18. https://doi.org/10.1128/mBio.01467-18.

36.   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinforma.
Oxf. Engl. *29*, 15–21. https://doi.org/10.1093/bioinformatics/bts635.

37.   Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. Bioinforma. Oxf. Engl. *30*, 923–930.
https://doi.org/10.1093/bioinformatics/btt656.

38.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.
https://doi.org/10.1186/s13059-014-0550-8.

39.   Kriakov, J., Lee, S. hee, and Jacobs, W.R. (2003). Identification of a regulated alkaline
phosphatase, a cell surface-associated lipoprotein, in Mycobacterium smegmatis. J. Bacteriol.
185, 4983–4991. https://doi.org/10.1128/JB.185.16.4983-4991.2003.

40.   Larsen, M.H., Biermann, K., Tandberg, S., Hsu, T., and Jacobs, W.R. (2007). Genetic
Manipulation of Mycobacterium tuberculosis. Curr. Protoc. Microbiol. *Chapter 10*, Unit 10A.2.
https://doi.org/10.1002/9780471729259.mc10a02s6.

41.   Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet.journal *17*, 10. https://doi.org/10.14806/ej.17.1.200.

42.   Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2.
Nat. Methods *9*, 357–359. https://doi.org/10.1038/nmeth.1923.

43.   Pritchard, J.R., Chao, M.C., Abel, S., Davis, B.M., Baranowski, C., Zhang, Y.J., Rubin,
E.J., and Waldor, M.K. (2014). ARTIST: high-resolution genome-wide assessment of fitness
using transposon-insertion sequencing. PLoS Genet. *10*, e1004782.
https://doi.org/10.1371/journal.pgen.1004782.

759
760
761

Figure 1. Impact of MZA on PZA susceptible and resistant *M. tuberculosis*. A, Structure of
MZA and activation of PZA to POA by PncA_Mtb. Dose response curves comparing MZA and PZA
susceptibility for *M. tuberculosis* strains **B,** H37Ra at pH 7.2 and **C,** H37Ra at pH5.8 and **D,** H37Rv
at pH 5.8. **E,** Assessment of pantothenate antagonism of MZA action on H37Ra at pH 7.2.
Determination of MZA susceptibility of PZA resistant strains **F,** H37Rv Δ*sigE*, H37Rv P_*clpC1*::*himar1*,
H37Rv *panD*::*himar1* and **G,** H37Rv Δ*pncA* compared to parental H37Rv. Dose dependent kill
curves of H37Rv exposed to MZA at **H,** pH of 7.2 and **I,** pH 5.8 over a 14-day period. **J,** Kill curves
performed with H37Rv under starvation conditions following a one-time treatment with PZA, MZA
or DMSO over a 60-day period at pH of 7.2. Kill curves performed with **K,** H37Rv and **L,** H37Rv
Δ*pncA* (PZA resistant) incubated at neutral pH under starvation conditions with MZA replenished
every 3 days**.** Evaluation of MZA activity against **M,** Gram-positive and **N,** Gram-negative ESKAPE
pathogens. All assays were performed in biological triplicate with mean displayed and error-bars
indicative of standard deviation.

777
778
779

Figure 2. Transcriptional profiling of *M. tuberculosis* exposed to MZA, PZA and POA. Volcano plots showing significantly differentially expressed genes in presence of **A,** PZA, **B,** POA and **C,** MZA. Cells were treated with 200 µM MZA, POA or PZA for 24h prior to RNA purification and sequencing. The MZA, POA and PZA transcriptional profiles share many key features, including upregulation in coenzyme A and lipid metabolism pathways. Transcriptional changes associated with exposure of a H37Rv Δ*pncA* to **D,** PZA and **E,** MZA. Cells were treated with 200 µM MZA or PZA for 24h prior to RNA purification and sequencing. **F,** Venn diagram showing upregulated genes in MZA, POA and PZA treated cultures, with the corresponding GO term analysis of common genes to the right. **G,** Venn diagram showing downregulated genes in MZA, POA and PZA treated cultures, with the corresponding GO term analysis of common genes to the right. RNA-seq was performed in biological triplicate.

793
794
795
796

Figure 3. Metabolism of MZA results in release of PZA and drives thiol stress. Time course
monitoring **A,** intracellular and **B,** extracellular abundance of MZA to PZA and POA from H37Ra
exposed to 250 µM MZA. **C,** Intracellular abundance of MZA, PZA and POA following a 24-hr
treatment of H37Ra (PZA susceptible) and *M. bovis* BCG (PZA resistant) with MZA. **D,** *mscR*
promoter region highlighting MZA resistance mutations in H37Rv Δ*pncA* (blue) and BCG (red)
backgrounds, respectively. Inhibition curve of an **E,** wild-type BCG and a **F,** *mscR* promoter mutant
exposed to MZA and methenamine (MET). **G,** Comparison of GFP expression between wild-type
mscR promoter and *mscR* promoter variant GFP reporter constructs in BCG. All samples were
normalized to the OD_{600}, significance was determined using a 2-tailed Student's t-test using *n*=6.
H, Inhibition curve comparing susceptibility of H37Rv and H37Rv Δ*mscR* to MZA. **I,** BCG wild-type
and **J,** BCG *mscR* promoter mutant exposed to 0.9 mM MZA or MET in the absence or presence
of cystine. All assays were performed in biological triplicate unless indicated otherwise with mean
displayed and error-bars indicative of standard deviation.

812
813
814

Figure 4. Genome-wide fitness profiling of *M. tuberculosis* transposon insertion mutants and chemical biology reveal the unique self-potentiating mechanism of MZA. Saturated library of Tn-insertion mutants treated with **A,** 233 µM PZA **B,** 105 µM POA and **C,** MZA 112 µM MZA at pH 6 to achieve a 50% reduction in growth rate. Each point corresponds to a specific TA site harboring a transposon insertion within the genome. The plot shows every transposon insertion identified. Highly depleted or enriched TA sites are highlighted with genes of interest noted. **D,** Venn diagram showing genes enriched for (left) and depleted (right) shared by all three treatments. **E,** DiaMOND assay showing exposure of *M. tuberculosis* H37Ra to a geometric series of PZA and methenamine (MET) alone and in combination. **F,** Abundance of CoA from *M. tuberculosis* H37Ra treated with DMSO (No Drug), MZA, PZA, formaldehyde (Form) and a combination of PZA and formaldehyde (PZA+form). Values were normalized to total protein. Significance was determined using a one-tailed Dunnett's test with Bonferroni correction. All assays were performed in biological triplicate with mean displayed and error-bars indicative of standard deviation.

832
833
834

Figure 5. MZA efficacy against PZA susceptible and PZA resistant *M. tuberculosis* strains in resting and activated macrophages. A, CFU comparison between RAW 264.7 macrophages infected with H37Rv and treated with either MZA, PZA or DMSO. **B,** H37Rv infected macrophages were activated using IFN-γ. **C,** Same experimental design as in A but using the PZA resistant H37Rv Δ*pncA* strain to test efficacy of MZA against PZA resistant strains. **D,** RAW 264.7 macrophages activated with IFN-γ and infected with PZA resistant H37Rv Δ*pncA*. Cells were treated with 0.9 mM MZA or PZA with daily media exchange. All assays were performed in biological triplicate with mean displayed and error-bars indicative of standard deviation. Significance was determined via Kruskal-Wallis test combined with post hoc Dunn test with *p*<0.01 (∗∗), *p*<0.001 (∗∗∗).

Figure 6. Thiol stress mediated potentiation of PZA action. Exposure to low pH drives thiol stress in *M. tuberculosis* resulting in potentiation of PZA activity. MZA releases aldehyde and PZA in *M. tuberculosis* regardless of pH. PZA is converted to POA which interferes with CoA synthesis. Aldehyde release drives thiol stress resulting in synergistic disruption of CoA metabolism with PZA resulting in rapid bacterial cell death.

858
859

861
862 **Figure S1. PCA for RNA-seq treatment groups**. **A**, PCA plots from data from wildtype *M.*
863 *tuberculosis* H37Rv cells treated with DMSO, PZA, POA or MZA. **B**, PCA plots from the *M.*
864 *tuberculosis* H37Rv Δ*pncA* cells treated with DMSO, PZA or MZA. Each point represents an
865 individual biological replicate.
866

867
868
869
870

Table S1. PZA and MZA MIC values under potentiating and antagonizing conditions.

Strain	Condition	MIC in 7H9 media	
		PZA (mM)	MZA (mM)
H37Ra	pH 5.8	0.5	0.14
H37Ra	pH 5.8, pantothenate	>4.5	0.28
H37Ra	pH 7.2	>4.5	0.28
H37Ra	pH 7.2, pantothenate	>4.5	0.56
H37Rv	pH 5.8	0.5	0.14
H37Rv	pH 7.2	>4.5	0.28
H37Rv Δ*pncA*	pH 5.8	>4.5	0.28
H37Rv Δ*sigE*	pH 7.2	>4.5	0.14
H37Rv P*clpC1*::*himar1*	pH 7.2	>4.5	0.14

Experiments were performed using *M. tuberculosis* strains H37Ra and H37Rv and their derivatives. For potentiation of PZA action, pH of 5.8 was used. For antagonism of PZA action, pantothenate was used at a concentration of 0.23 mM. Strains H37Rv Δ*pncA*, Δ*sigE* and P*clpC1*::*himar1* were previously described[8] for their differing levels of PZA resistance. All assays were performed in biological triplicate with mean displayed.