Gut Bacteria Metabolize Natural and Synthetic Steroid Hormones via the Reductive OsrABC Pathway

Christian Jacoby,[1,2*] Kaylie Scorza,[1,2*] Lia Ecker,[1,2] Mary McMillin,[1] Ramanujam Ramaswamy,[1] Anitha Sundararajan,[1] Ashley M. Sidebottom,[1] Huaiying Lin,[1] Keith Dufault-Thompson,[3] Brantley Hall,[4] Xiaofang Jiang,[3] and Samuel H. Light[1,2#]

[1]Duchossois Family Institute, University of Chicago, Chicago, IL, USA
[2]Department of Microbiology, University of Chicago, Chicago, IL, USA
[3]National Library of Medicine, National Institutes of Health, Bethesda, MD, USA
[4]Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, College Park, MD, USA

*These authors contributed equally
#Address correspondence to samlight@uchicago.edu

ABSTRACT

Steroid hormone metabolism by the gut microbiome has multiple implications for mammalian physiology, but the underlying mechanisms and broader significance of this activity remains largely unknown. Here, we isolate a novel human gut bacterium, *Clostridium steroidoreducens*[T] strain HCS.1, that reduces cortisol, progesterone, testosterone, and related steroid hormones to 3β,5β-tetrahydrosteroid products. Through transcriptomics and heterologous enzyme profiling, we identify and biochemically characterize the *C. steroidoreducens* OsrABC reductive steroid hormone pathway. OsrA is a 3-oxo-Δ^1-steroid hormone reductase that selectively targets the Δ^1-bond present in synthetic steroid hormones, including the anti-inflammatory corticosteroids prednisolone and dexamethasone. OsrB is a promiscuous 3-oxo-Δ^4-steroid hormone reductase that converts steroid hormones to 5β-dihydrosteroid intermediates. OsrC is a 3-oxo-5β-steroid hormone oxidoreductase that reduces 5β-intermediates to 3β,5β-tetrahydro products. We find that *osrA* and *osrB* homologs predict steroid hormone reductase activity in diverse gut bacteria and are enriched in Crohn's disease fecal metagenomes. These studies thus identify the basis of reductive steroid hormone metabolism in the gut and establish a link between inflammatory disease and microbial enzymes that deplete anti-inflammatory corticosteroids.

ACKNOWLEDGEMENTS

Research reported in this publication was supported by funding from the National Institutes of Health (NIGMS R35GM146969 and NIDDK P30DK042086, via the University of Chicago Center for Interdisciplinary Study of Inflammatory Intestinal Disorders) and the Searle Scholars Program (to S.H.L), as well as the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Projektnummer 542537779 (to C.J.).

INTRODUCTION

Steroid hormones encompass a broad class of biologically active molecules that play crucial roles in diverse physiological processes. Corticosteroids, like cortisol, are involved in regulating inflammation, immune response, and metabolism,[1] while sex steroids, including estrogens, androgens, and progestins, regulate reproductive functions and secondary sexual characteristics.[2] Due to their wide-ranging effects, both natural and synthetic steroid hormones are commonly used in medical therapies to manage conditions such as autoimmune diseases, hormone deficiencies, and cancers.

The gut microbiome mediates diverse phenotypes through the modification of host- and diet-derived metabolites, including various steroids. Research on microbiome steroid metabolism has primarily focused on bile acids, products of which are important for multiple host phenotypes.[3-5] However, while less studied, steroid hormones have also been identified as an important class of substrates in the gut. These molecules interact with gut microbes after entering the gastrointestinal tract via therapeutic oral or rectal administration or through excretion in bile.[6] Bile is likely a particularly important source of intestinal steroid hormones, as 9-22% of endogenous cortisol,[7,8] 10-15% of testosterone,[9] 20-30% of corticosterone,[10] and up to 30% of progesterone[11] are eliminated from the body through this route.

Gut microbes generate multiple products from steroid hormones, including cortisol and corticosterone derivatives that serve as fecal biomarkers for stress in animal research studies.[12,13] In addition to passing into feces, intestinal microbial steroid hormone products can reenter the bloodstream through enterohepatic circulation. In humans, this is evidenced by the rectal administration of cortisol leading to an increase in specific circulating cortisol derivatives in a microbiome-dependent manner.[14,15] Other studies provide evidence that microbial products of progesterone metabolism similarly enter systemic circulation.[6,16]

The impact of microbial steroid hormone metabolism has been linked to several aspects of mammalian biology. Microbial inactivation of orally administered steroids, including side-chain cleavage of synthetic corticosteroids, reduces the bioavailability of these drugs.[17,18] Microbial pathways that dehydroxylate corticosterone and convert steroid precursors to androgens generate metabolites that contribute to hypertension in animal models[19,20] and promote castration-resistant prostate cancer,[21,22] respectively. Bacterial metabolisms that alter the concentration of steroid hormones with distinct biological activities thus have diverse consequences for mammalian biology.

Previous studies have reported that some gut bacteria reduce the Δ^4-bond in steroid hormones, generating 5β-steroid derivatives.[23] This activity decreases the anti-inflammatory and androgenic properties of glucocorticoids and androgens, respectively, and converts progestins into a neuroactive form.[24] While it stands to reason that Δ^4-steroid hormone reduction may have implications for host biology, the molecular basis and the broader significance of activities in the gut microbiome remains unknown.

Here, we describe the isolation and characterization of a novel steroid hormone-metabolizing gut bacterium, *Clostridium steroidoreducens* HCS.1. By employing a multidisciplinary approach, integrating genomics, transcriptomics, and metabolomics, we characterize the *osrABC* reductive steroid hormone pathway. These findings provide new insights into the diversity of steroid hormone metabolism in the gut microbiome and its potential impact on host health.

RESULTS

Clostridium steroidoreducens is a novel steroid hormone-enriched gut bacterium

To select for members of the gut microbiome with metabolic capabilities that provide a selective advantage in the presence of steroid hormones, we passaged a human fecal sample in a nutritionally limited base medium supplemented with individual corticosteroids (cortisol or corticosterone) or sex steroids (progesterone or testosterone) (**Figure 1A**). We isolated strains

from the final enrichment passages and cultivated them on steroid hormone-infused solid
media. We identified one strain, HCS.1, that cleared insoluble cortisol or progesterone from the
solid media and accumulated a white precipitate indicative of a potential reaction product at
colony centers on progesterone-infused media (**Figure 1B and 1C**). Consistent with HCS.1
possessing a selective advantage in the presence of steroid hormones, 16S rRNA amplicon
sequencing of the final enrichment passages revealed an amplicon sequence variant matching
the HCS.1 16S rRNA sequence was enriched from below the limit of detection in the fecal
inoculum to 0.7-23.6% of the microbial community following cortisol, corticosterone,
progesterone, or testosterone enrichment (**Figure 1D**).
To facilitate further strain characterization, we sequenced and assembled HCS.1 DNA
into 3 circularized contigs, comprising a genome and two plasmids, that contain 3,816,008 base
pairs with 28.4% G + C content, 3595 protein-coding genes, and 111 RNA genes (**Figure 1E**).
Phylogenetic analyses revealed that HCS.1 was closely related to *Clostridium chrysemydis* but
represents a novel species of the genus *Clostridium,* based on an average nucleotide identity
(ANI) of 92.09% to the closest reference genome and accepted taxonomic assignment criteria
(**Extended Data Figure 1**).[25] In recognition of steroid hormone reductase activities detailed
below, we assigned HCS.1 the species name *Clostridium steroidoreducens.*
109
***C. steroidoreducens* possesses broad steroid hormone reductase activity**
To determine whether the *C. steroidoreducens* HCS.1 steroid clearance phenotype was
indicative of metabolic activity, we employed an LC-MS-based assay to track the fate of cortisol
or progesterone in *C. steroidoreducens* HCS.1 culture. We observed that both steroid hormones
were fully depleted from the media, coinciding with the emergence of a minor and major
product. By comparing to compound reference standards, we confirmed that major and minor
products corresponded to 5β-dihydro- and 3β,5β-tetrahydro-steroid derivatives, respectively
(**Figure 2A, Extended Data Figure 2**).
Tracking *C. steroidoreducens* HCS.1 cortisol metabolism over time, we observed that
5β-dihydrocortisol transiently accumulated, peaking at 60 minutes before decreasing to less
than 2% of the total corticosteroid present by 120 minutes (**Figure 2B**). By contrast, 3β,5β-
tetrahydrocortisol steadily accumulated following the introduction of cortisol, exceeding 98% of
the corticosteroid present by 120 minutes (**Figure 2B**). These results suggest that, in contrast to
previously characterized bacterial steroid dehydroxylation[26] and side chain-cleaving[27] activities,
C. steroidoreducens HCS.1 exclusively reduces cortisol, converting it to 3β,5β-tetrahydrocortisol
via a 5β-dihydrosteroid intermediate (**Figure 2C**).
To address the specificity of *C. steroidoreducens* HCS.1 steroid utilization, we next
tested the strain's activity on a panel of steroids with variable functional groups at multiple
positions on the sterol core (**Figure 2D**). We found that *C. steroidoreducens* tolerated
substitutions at C1, C11, C17 positions, exhibiting activity on distinct corticosteroids
(corticosterone, cortisone, prednisolone) and sex steroids (progesterone, testosterone) (**Figure
2E**). In contrast to these polar steroids, the hydrophobic cholesterol-derivative cholestenone
was a poor substrate (**Figure 2E**). These results establish *C. steroidoreducens* as a steroid
hormone-reducing gut bacterium with broad substrate specificity.
134
Fe-S flavoenzyme family OsrB is a 3-oxo-Δ^4-steroid hormone reductase
We next sought to identify the mechanism of steroid hormone reduction by *C.
steroidoreducens.* As bacterial reductases are often induced in the presence of their
substrate,[26,28] we employed a transcriptomics-based approach to identify candidate steroid
hormone reductases in *C. steroidoreducens.* We performed RNA-seq analysis on *C.
steroidoreducens* cells cultivated in the presence or absence of cortisol and identified 30 genes
that were induced >2-fold when cortisol was present (**Supplementary Table 1**). Two of the most
highly induced genes, which we renamed *osrA* and *osrB* (oxosteroid reductase A and B),

encoded proteins annotated as *fadH*-like 2,4-dienoyl-CoA reductases (**Figure 3A,**
Supplementary Table 1).
E. coli 2,4-dienoyl-CoA reductase is the best characterized member of the "Fe-S
flavoenzyme family" of oxidoreductases that contain a conserved N-terminal substrate-binding
domain (PF00724) and a C-terminal NAD(P)H cofactor-binding domain (PF07992) (**Figure
3B**).[29] Divergent members of the Fe-S flavoenzyme family are widespread in gut bacteria and
possess distinct substrate specificities for host- and diet-derived metabolites.[30] Consistent with
OsrA and OsrB representing novel Fe-S flavoenzyme subtypes with distinct substrates, we
observed that these enzymes exhibited remote sequence homology to previously characterized
Fe-S flavoenzymes, including *Clostridium scindens* Fe-S flavoenzymes, BaiCD and BaiH, which
reduce bile acid intermediates structurally related to steroid hormones (**Figure 3C**).[29,31]
To test steroid reductase activity of OsrA and OsrB, we heterologously produced the
enzymes in anaerobically cultured *E. coli* cells. We found cells expressing *osrB*, but not *osrA*,
reduced cortisol to 5β-dihydrocortisol (**Figure 3D**). Studies with anaerobically purified OsrB
revealed that NADH and NADPH were poor electron donors for OsrB. Using the artificial
electron donor methyl viologen, we observed that OsrB similarly reduced a variety of steroid
hormones substrates (**Figure 3E**). These results thus establish OsrB as a promiscuous 3-oxo-
Δ4-steroid hormone reductase that uses a presently unidentified electron donor.
161
Short chain dehydrogenase OsrC is a 3-oxo-5β-steroid hormone oxidoreductase
We next sought to identify the *C. steroidoreducens* enzyme responsible for reduction
of the 3-oxo group on the 5β-dihydrosteroid intermediate generated by OsrB. Microbial bile
acid oxidoreductases with specificity for 3α-, 3β-, 7α-, 7β-, 12α- and 12β-hydroxyl groups
have been previously identified.[32-34] As these characterized steroid oxidoreductases are
members of the short chain dehydrogenase (SDR) enzyme superfamily, we reasoned the *C.
steroidoreducens* enzyme was likely related to this family. An analysis of the *C.
steroidoreducens* genome identified 12 genes with SDR domains. However, none were
induced by cortisol or exhibited high sequence similarity to previously characterized bile acid
oxidoreductases.
As these analyses failed to identify obvious candidates, we next performed an
unbiased screen of SDR-containing *C. steroidoreducens* proteins for 3-oxo-5β-steroid
hormone reductase activity. We confirmed soluble expression of all 12 SDRs in *E. coli* and
tested the activity of overexpressing *E. coli* strains on 5β-dihydrocortisol (**Figure 4A**,
Extended Data Figure 3). We identified two SDRs (BLEONJ_2554 and BLEONJ_1088) that
produced 3β,5β-tetrahydrocortisol and two others (BLEONJ_2478 and BLEONJ_1414) that
yielded 3α,5β-tetrahydrocortisol (**Figure 4A**).
Studies with anaerobically purified BLEONJ_2554 and BLEONJ _1088 revealed
divergent substrate specificities. BLEONJ_2554 showed a pronounced preference for 5α-
steroids and exhibited weak activity that did not follow classical Michaelis-Menten kinetics with
5β-steroid substrates (**Figure 4B, Extended Data Figure 4**). Conversely, gene BLEONJ _1088
displayed a preference for 5β-steroid hormones and accommodated multiple functional groups
at the C9 or C17 positions (**Figure 4B, Extended Data Figure 5**). We further found that
BLEONJ_1088 exhibited a preference for steroid hormones relative to the comparable bile acid
derivative lithocholic acid. We thus conclude that BLEONJ_1088 is a 3-oxo-Δ⁴-steroid hormone
reductase and, on this basis, renamed it OsrC.
188
**Fe-S flavoenzyme family OsrA is a 3-oxo-Δ¹-steroid hormone reductase active on
synthetic corticosteroids**
Synthetic corticosteroids possess potent anti-inflammatory properties and are used to
treat a range of pathologies, including inflammatory bowel disease.[35] The synthetic
corticosteroids drugs dexamethasone, prednisone, prednisolone, and methylprednisolone

contain a Δ^1-bond that is absent in natural corticosteroids and which significantly extends their
half-life (**Figure 5A**).[36] As our initial screen of steroids identified prednisolone as a substrate for
C. steroidoreducens (**Figure 1**), we sought to address the molecular basis of synthetic
corticosteroid metabolism. We first tested *C. steroidoreducens* activity on additional synthetic
corticosteroids dexamethasone, prednisone, and methylprednisolone and found that all were
reduced to 3β,5β-tetrahydrocortisol derivatives, indicating that the bacterium possesses both Δ^1-
and Δ^4-steroid hormone reductase activities (**Figure 5B**).
Considering the similarity of the Δ^1-reduction to the OsrB catalyzed Δ^4-reduction, we next
tested the activity of OsrA and OsrB and found that the two enzymes generated distinct cortisol
and testosterone isomers from prednisolone and the synthetic androgen boldenone,
respectively (**Figure 5C, Extended Data Figure 6**). Based on comparison to reference
standards, we establish that OsrA and OsrB products reflected Δ^1- and Δ^4-steroid hormone
reductase activities, respectively (**Extended Data Figure 6**). These results demonstrate that
OsrA functions as a Δ^1-steroid hormone reductase that acts in conjunction with OsrB and OsrC
to reduce synthetic steroid hormones to 3β,5β-reduced products (**Figure 5E**).

Steroid hormone reductase activities are common in gut bacteria and correlate with the
distribution of *osrA* and *osrB* homologs
We next sought to address the breadth of steroid hormone reductase activity in the gut
microbiome. We performed BLASTp searches of OsrA, OsrB, and OsrC in the Unified Human
Gastrointestinal Genome catalog of representative genomes and metagenome-assembled
genomes, which includes 4,644 prokaryotic species that colonize the human gastrointestinal
tract.[37] These searches identified homologs with high sequence homology to OsrA, OsrB, and
OsrC in 2, 59, and 90 genomes, respectively (**Supplementary Table 2**). Genomes encoding
osrABC homologs included gram-positive bacterial species from multiple taxa, primarily from the
Erysipelotrichaceae and Lachnospiraceae families.
To determine the association of *osrABC* homologs with observed *C. steroidoreducens*
phenotypes, we selected 117 gut bacteria strains for experimental characterization. We tested
these strains on solid media on steroid clearance/precipitate accumulation and assayed a select
subset for cortisol and prednisolone activity. From these studies we identified 29 strains from 14
species with a steroid clearance/precipitate accumulation phenotype and 6 species isolates with
steroid hormone reductase activity (**Figure 6A-6C and Supplementary Table 3**).
Comparing strain genotypes to observed phenotypes revealed several patterns. First,
presence of an *osrB* homolog in a genome strongly predicted steroid clearance/precipitate
accumulation and steroid hormone Δ^4-reductase activity (**Figure 6B and Supplementary Table**
3). Second, the absence of *osrA* homologs tracked with a consistent lack of steroid hormone Δ^1-
reductase activity (**Figure 6C and Supplementary Table 3**). Third, while the presence of an
osrC homolog tracked with production of 3β,5β-tetrahydrocortisol, absence of an *osrC* homolog
was not predictive of fate of the C3 functional group. Indeed, *osrC*-negative strains varied in
their major cortisol product, generating either 5β-dihydrocortisol, 3α,5β-tetrahydrocortisol, or
3β,5β-tetrahydrocortisol (**Figure 6B and Supplementary Table 3**). These results provide
evidence that *osrA* and *osrB* specifically confer Δ^1- and Δ^4-steroid hormone reductase activities,
respectively, while *osrC* likely represents one of multiple evolutionarily distinct 3-oxo-5β-steroid
hormone oxidoreductases.

***osrB* is prevalent in human fecal metagenomes and associated with active Crohn's**
disease
Having established the relevance of *osrABC* homologs for steroid reductase activity in
gut bacteria, we next sought to determine the prevalence of the pathway in the human gut. We
focused our analysis on *osrA* and *osrB*, since these homologs reliably predicted steroid
hormone reductase activities of assayed strains. We recruited reads to *osrA* and *osrB* homologs

in a collection of 1,491 previously published healthy human fecal metagenomes. We detected at
least one read mapping to *osrA* and *osrB* homologs in 2.2% and >98.9% of samples,
respectively (**Supplementary Table 4**). Within most metagenomes multiple *osrB* homologs
recruited many reads. By contrast, the majority *osrA* reads recruited to *Clostridium tertium osrA*
homologs, often with only one or two reads per metagenome (**Supplementary Table 4**). These
analyses demonstrate that *osrB* homologs are common in the gut but that *osrA* homologs are
confined to bacteria that colonize the gut at a low relative abundance.
Considering that glucocorticoids possess potent anti-inflammatory activities and natural
and synthetic variants, including cortisol and prednisolone, are rectally and orally administered
for the treatment of inflammatory bowel disease, we reasoned that OsrA and OsrB activity could
be clinically relevant in this patient population. We analyzed 314 metagenomes from the Lewis
et al.[38] study of active Crohn's disease patients, including a subset treated with corticosteroids
(**Supplementary Table 5**). We observed *osrB* homologs were elevated in Crohn's disease
patient relative to a healthy control population (**Figure 6D**). Further scrutiny revealed that the
increased abundance *osrB* homologs from *Ruminococcus_B gnavus* and *Clostridium_AQ*
innocuum, two taxa previously associated with Crohn's disease inflammation,[39,40] was the
primary driver of this association (**Figure 6E**).
osrA homologs similarly exhibited elevated abundance in Crohn's disease
metagenomes, but their low prevalence coupled with the relatively small sample size of this
dataset complicated statistical analysis of the significance of this relationship (**Figure 6D**). To
address this issue, we expanded our dataset to include 1537 metagenomes from multiple
separate studies that included Crohn's disease and control populations. Analysis of this larger
dataset confirmed that *osrA* homologs were significantly elevated in Crohn's disease patient
metagenomes and revealed that the *osrA* from *Clostridium tertium* was the primary driver of this
association (**Extended Data Figure 7A and 7B, Supplementary Table 5**).
To determine whether identified associations extended an independent dataset, we
evaluated 569 Crohn's disease patient metagenomes collected as part of Integrative Human
Microbiome Project (**Supplementary Table 6**). We observed a similar association between
elevated abundance of *osrA* and *osrB* homologs and microbiome dysbiosis scores used as a
proxy for active Crohn's disease in this study (**Extended Data Figure 8A-8C, Supplementary**
Table 6).[41] Underscoring the relevance of these observations to active Crohn's disease,
metagenomes from this study with microbiome dysbiosis score consistent with inactive Crohn's
disease exhibited intermediate *osrB* homolog levels between dysbiotic Crohn's disease and
control non-IBD populations (**Extended Data Figure 8A**, **Supplementary Table 6**).
We further investigated the relationship between microbial steroid reductases and
corticosteroid treatment in Crohn's disease patients, as reported in the Lewis et al.[38] study. Our
analysis revealed that corticosteroid therapy was associated with an increase in *osrA* homolog
prevalence, as these homologs were detected in 9.6% of corticosteroid-treated patients
compared to only 2.4% in untreated patients. This suggests a potential selective pressure
favoring bacteria that metabolize synthetic corticosteroids in patients receiving these therapies.
Interestingly, within the metagenomes where *osrA* homologs were present, their abundance did
not significantly differ between treated and untreated patients, suggesting that the effect of
corticosteroid treatment may relate to increased colonization of bacteria with *osrA* homologs
(**Extended Data Figure 9)**.
289
DISCUSSION
In this study, we characterize *Clostridium steroidoreducens* HCS.1, a previously
uncharacterized gut bacterium that encodes a novel reductive pathway, OsrABC, for the
metabolism of steroid hormones. Our work expands on previous studies of microbial steroid
metabolism by demonstrating the widespread prevalence and activity of OsrB and OsrC in the
gut microbiota, which catalyze the reduction of natural steroid hormones into their 3β5β-

tetrahydro derivatives. Notably, the OsrB homologs are prevalent in Crohn's disease-associated
bacterial communities, implicating these enzymes in both health and disease contexts.
One of the most compelling aspects of this work is the link between microbial steroid
metabolism and chronic inflammatory conditions, particularly Crohn's disease. Our data indicate
that *osrB* homologs are enriched in pro-inflammatory taxa such as *Clostridim_AQ innocuum* and
Ruminococcus_B gnavus, both of which have previously been associated with Crohn's
disease.[39,40] Pro-inflammatory gut microbes often exhibit a competitive advantage in
inflammatory conditions and induce inflammation to generate conditions favorable for their
growth.[41] This suggests that these microbes may leverage steroid hormone metabolism to gain
a competitive advantage in the inflamed gut environment. Specifically, OsrB-mediated depletion
of endogenous anti-inflammatory glucocorticoids may represent an adaptive strategy employed
by *Clostridim_AQ innocuum* and *Ruminococcus_B gnavus* to perpetuate inflammation and
support their growth.
In a clinical context, the OsrABC reductase pathway may also have significant
implications for glucocorticoid therapies, which are commonly administered to manage
inflammatory bowel disease. Our findings suggest that the OsrABC reductase pathway could
modulate the effective dose of administered glucocorticoids by degrading these anti-
inflammatory compounds. This underscores the importance of further investigations into the
microbial impact on drug bioavailability in relation to both the efficacy and dosing of steroid
therapies in patients.
Beyond the clinical considerations, our study highlights the broader significance of gut
microbial steroid metabolism in human health. Notably, a concurrently published manuscript
independently identifies the 3-oxo-Δ^4-steroid hormone reductase activity of OsrB homologs,
along with the characterization of additional novel gut bacterial enzymes that metabolize
progestins.[42] Together, these findings represent a crucial step forward in delineating the broader
landscape of microbial steroid hormone metabolism and its potential clinical implications.
322
323
MATERIALS AND METHODS
<u>Steroid hormone enrichment culture</u>
For each enrichment sample, 15 mM steroid hormone (cortisol, corticosterone,
progesterone, or testosterone) suspensions were prepared in 1 mL basal growth medium (Difco
M9 minimal salts, 20 mM acetate, 20 mM formate, tryptone 0.01% w/v, Bacto yeast extract
0.01% w/v, trace vitamins and minerals, $MgSO_4$ 4.09% w/v; pH 6.5). Homogenized fecal
samples were pelleted and washed 3x in phosphate buffer saline (PBS), then resuspended in 1
mL saline. 20 µL cell suspension was added to each enrichment culture condition and incubated
for 72 hours. After 72 hours, 20 µL of each culture was used to inoculate fresh media
supplemented with its respective compound. Cultures were passaged a total of 4 times. After
the final passage, a portion of each condition was preserved in 20% glycerol and frozen at -80
°C. The remaining culture was pelleted and processed for 16S rRNA sequencing.
336
<u>Isolation of HCS.1</u>
Preserved stocks of enrichment culture samples were plated onto fresh brain heart
infusion (BHI) agar and incubated for 4 days at 37 °C under anaerobic conditions (5% H_2, 10%
CO_2, 85% N_2). Distinct colonies were passaged to confirm purity and identified by 16S rRNA V4-
V5 variable region sequencing. Purified isolates of HCS.1 were stored at -80 °C in a 20%
glycerol suspension. Frozen glycerol stocks were deposited in the DFI Symbiotic Bacterial
Strain Bank Repository (https://dfi.cri.uchicago.edu/biobank/).
344
<u>Steroid clearance assay</u>

To prepare steroid clearance assays, progesterone and cortisol amounts for a final
concentration of 12 mM or 16 mM, respectively, were sterilized by suspension in 70% v/v
ethanol, followed evaporation at room temperature for 2 hours. Dried steroid powders were
sifted into autoclaved BHI agar and stirred rapidly, shortly before pouring into plates. Solid
plates were stored under anaerobic conditions at 25 °C for at least 24 hours prior to use.
To test HCS.1 steroid clearance, solid BHI plates were incubated at 37 °C for 2 days.
After 2 days, single colonies were picked and suspended in 200 µL PBS. Aliquots of the cell
suspension were spread onto solid steroid plates and incubated anaerobically for 5 days at 37
°C. To test steroid clearance of other bacterial strains, solid BHI plates were incubated at 37 °C
for 4 days. After 4 days, single colonies from each isolate were picked and suspended in 200 µL
PBS. 2 µL aliquots were spotted onto solid progesterone plates, dried, and incubated
anaerobically for 3 days at 37 °C.
358
16S rRNA sequencing and analysis
Cells from final steroid enrichment passages were collected by centrifugation and their
genomic DNA extracted using the QIAamp PowerFecal Pro DNA kit (Qiagen). Briefly, samples
were suspended in a bead tube (Qiagen) along with lysis buffer and loaded on a bead mill
homogenizer (Fisherbrand). Samples were then centrifuged, and the supernatant was
resuspended in a reagent that effectively removed inhibitors. DNA was then purified routinely
using a spin column filter membrane and quantified using Qubit. The 16S rRNA variable V4-V5
region was amplified using universal bacterial primers, 564F and 926R. Amplicons were purified
using magnetic beads, then quantified and pooled at equimolar concentrations. The Qiagen
QIAseq one-step amplicon library kit was used to ligate Illumina sequencing-compatible
adaptors onto amplicons. Reads were sequenced on an Illumina MiSeq platform to generate 2 x
250 base pair reads, with 5,000-10,000 reads per sample. Amplified 16S rRNA amplicons were
processed through the dada1 pipeline in R. Forward reads were trimmed at 210 bp and reverse
reads were trimmed at 150 bp, to remove low-quality nucleotides. Chimeras were detected and
removed using default parameters. Amplicon sequence variants between 300 and 360 bp in
length were taxonomically assigned to the genus level using the RDP Classifier (v2.13) with a
minimum bootstrap confidence score of 80.
376
377 Sample preparation for whole genome sequencing
378 To prepare HCS.1 for whole genome sequencing, 10 mL BHI broth was inoculated with
379 cells from a single bacterial colony and incubated anaerobically at 37 °C for 48 hours. The
380 culture was centrifuged at 4000 x g for 10 minutes. The resulting pellet was resuspended,
381 washed in phosphate buffer saline (PBS), and re-centrifuged.
382
Whole genome sequencing library preparation: Illumina short reads
Samples for Illumina short sequencing were extracted using the QIAamp PowerFecal
Pro DNA kit (Qiagen), as described in the preceding subsection. Libraries were prepared using
200 ng of genomic DNA using the QIAseq FX DNA library kit (Qiagen). Briefly, DNA was
fragmented enzymatically into shorter fragments and desired insert size was achieved by
adjusting fragmentation conditions. Fragmented DNA was end repaired and 'A's' were added to
the 3'ends to stage inserts for ligation. During ligation step, Illumina compatible Unique Dual
Index (UDI) adapters were added to the inserts and prepared library was PCR amplified.
Amplified libraries were cleaned up, and QC was performed using Tapestation 4200 (Agilent
Technologies). Libraries were sequenced on an Illumina NextSeq 1000/2000 to generate
2x150bp reads.
394
395 Whole genome sequencing library preparation: Oxford Nanopore long reads

396   Samples for Nanopore and Illumina hybrid assemblies were extracted using the high
397 molecular weight NEB Monarch Genomic DNA Purification Kit. DNA was QC'ed using genomic
398 Tapestation 4200. Nanopore libraries were prepared using the Rapid Sequencing Kit (SQK-
399 RAD114) and sequenced on MinION R10.4.1 flow cells. Nanopore reads were base-called
400 using ONT Guppy basecalling software version 6.5.7+ca6d6af, minimap2 version 2.24-r1122,
401 and was demultiplexed using ONT Guppy barcoding software version 6.5.7+ca6d6af using local
402 HPC GPU. N50 of the nanopore long read is 7077 base pairs, the average read length is 4529.4
403 base pairs, while the average read quality is 15.6, which is typical of Nanopore reads. Hybrid
404 assembly was performed with both nanopore and Illumina short reads using Unicylcer
405 v0.5.0.[43,44]
406
407 <u>Taxonomic classification of *Clostridium steroidoreducens* sp. nov. Strain HCS.1^T^</u>
408   The classification of strain HCS.1 as a novel species, *Clostridium steroidoreducens* sp.
409 nov. was performed using GTDB-Tk (version 2.3.2)[45] on the KBase platform. Genome quality
410 assessment, phylogenetic placement, and taxonomic classification were performed according to
411 GTDB guidelines. The HCS.1 genome was uploaded to the KBase website and analyzed using
412 the GTDB-Tk classify workflow, which assigns genomes to the closest known taxa based on
413 conserved marker genes. Strain HCS.1 was classified within the genus *Clostridium*, but did not
414 match any known species in the Genome Taxonomy Database (GTDB, version r207).
415 Phylogenetic placement within the GTDB bacterial tree confirmed that the strain represented a
416 distinct lineage, supporting its designation as a new species.
417
418 <u>Transcriptomic analysis of HCS.1</u>
419   To prepare HCS.1 samples for transcriptomic analysis, six 50 mL Lysogeny broth
420 cultures were inoculated with bacteria cells andshaken anaerobically at 37 °C for 48 hours. After
421 48 hours, cortisol powder was added to 3 cultures, to a final concentration of 8 mM. All 6
422 cultures were incubated for an additional 4 hours, then pelleted at 4000 x g for 10 minutes. The
423 resulting pellets were flash-frozen in a dry ice/ethanol bath and stored at -80 °C until ready for
424 subsequent processing. Cell pellets were thawed and total RNA from biological replicates
425 extracted using the Maxwell RSC instrument (Promega). Extracted RNA was quantified using
426 Qubit, and integrity was measured using TapeStation (Agilent Technologies). Libraries from ribo-
427 depleted samples were constructed using the NEB's Ultra Directional RNA library prep kit for
428 Illumina. First, up to 500 ng total RNA was subjected to ribosomal RNA depletion (for bacteria)
429 using NEBNext rRNA depletion kit. Ribosomal -RNA depleted samples were fragmented based
430 on RNA integrity number (RIN). Post cDNA synthesis, Illumina compatible adapters were ligated
431 onto the inserts and final libraries were QC'ed using TapeStation (Agilent technologies).
432 Libraries were normalized using library size and final library concentration (as determined by
433 Qubit). Library concentration (ng/ul) was converted to nM to calculate dsDNA library
434 concentration. Equimolar libraries were then pooled together at identical volumes to ensure
435 even read distribution across all samples. Normalized libraries were then sequenced on
436 Illumina's NextSeq 1000/2000 at 2x100bp read length.
437   High-quality reads were mapped to the circularized hybrid assembled genome of HCS.1
438 (NCBI: CP170704), using Bowtie2 (v.2.4.5), and sorted with Samtools (v1.6). Read counts were
439 generated using featureCounts (v2.0.1) with Bakta annotations.[46] Gene expression was
440 quantified as the total number of reads uniquely aligning to the reference genome, binned by
441 annotated gene coordinates. Differential gene expression and quality control analyses were
442 performed using DESeq2 in R with Benjamini–Hochberg false discovery rate adjustment applied
443 for multiple testing corrections.[47]
444
445 <u>Bacterial culture steroid reductase assay</u>

A complete list of strains used in this study is provided in **Supplementary Table 2**.
Strains were incubated under anaerobic conditions (85% N_2, 10% CO_2, 5% H_2) at 37 °C in an
anaerobic chamber (Coy Laboratory). Liquid brain-heart infusion (BHI) broth supplemented with
100 µM steroids from 10 mM stocks in methanol was used for growth. Cultures were grown
anaerobically with a 1% (v/v) inoculum from a pre-culture and supplemented with steroids after
4 hours of incubation during the exponential growth phase. Bacterial cultures were extracted by
the addition of 9 volumes of methanol supplemented with 0.5 µM methylprednisolone as an
internal standard for LC-MS analysis.
454
LC-MS-Q-TOF analysis of steroids
Extracted samples were vortexed and centrifuged twice at 21,000 × g for 15 minutes,
with the supernatant transferred to new tubes after each centrifugation step. The methanol
fraction was filtered through 0.2 µm nylon membrane filters prior to LC-MS analysis. Samples
were analyzed using an Agilent 6540 UHD Q-TOF mass spectrometer coupled to an Agilent
1200 Infinity LC system. Separation was performed on a XBridge C18 column (2.1x100mm, 3.5
µm particle size) using 0.1% aqueous formic acid and acetonitrile with 0.1% formic acid as
mobile phases. The separation gradient ranged from 20% to 100% acetonitrile over 4 minutes at
50 °C with a flow rate of 0.5 mL/min. Mass spectra were acquired in negative ion mode for
glucocorticoids ($[M+FA-H]^-$) or positive ion mode for all other steroids ($[M+H]^+$), with an ion spray
voltage of 3500 V and a nozzle voltage of 2000 V. The source temperature was set to 300 °C,
and the gas flow rate was 8 L/min. Data were processed and visualized using MassHunter
software version 10.
468
Molecular biology
Gene transformations were performed by Gibson assembly using 2x NEBuilder® HiFi
DNA Assembly Master Mix (New England Biolabs, NEB, E2621X). Primers were designed using
SnapGene (see **Supplementary Table 7**), incorporating 20 bp flanking regions complementary
to a linearized expression vector (pMCSG53) and the gene of interest from the HCS.1 genome.
PCR, cloning, and transformation were performed according to the protocols provided on the
NEB website. The Gibson assembly reaction was incubated at 50 °C for 1 hour and then
transformed into *E. coli* XL1-Blue competent cells according to the manufacturer's protocol.
Transformed cells were plated on Luria-Bertani (LB) agar plates containing 100 µg/mL
carbenicillin, and successful transformations were confirmed using sequencing primers specific
for the backbone vector (see **Supplementary Table 7**). Positive colonies were validated and
sequenced by the University of Chicago Genomics Facility. The final constructs were then
transformed into chemically competent *E. coli* Rosetta™ (DE3) competent cells (Novagen)
according to NEB protocols. Transformed cells were plated on LB agar plates supplemented
with 100 µg/mL carbenicillin.
484
Protein production in *E. coli*
Protein production in *E. coli* Rosetta cells was performed under aerobic conditions for all
short-chain dehydrogenases (SDRs) and under anaerobic conditions for OsrA and OsrB.
Cultures were grown in either 2x YT medium (20 g/L tryptone, 10 g/L yeast extract, and 5 g/L
NaCl) or TB medium (12 g/L tryptone, 24 g/L yeast extract, 4 mL/L glycerol, 9. 4 g/L K_2HPO_4,
and 2.2 g/L KH_2PO_4) supplemented with 0.5% (w/v) glucose and 1 mM ferric ammonium citrate,
respectively.
Induction of protein expression was initiated during the exponential phase when optical
densities (OD_{600}) reached 0.4-0.6 by the addition of 1 mM isopropyl β-D-1-
thiogalactopyranoside (IPTG). Cultures were incubated for 3-5 hours at 37 °C with shaking at
200 rpm. Cells were harvested by centrifugation at 4500 × g for 20 minutes. Cell pellets were

then frozen at -80 °C for storage prior to subsequent experimental use. Protein production was
confirmed by SDS-PAGE analysis.
498
Purification of heterologous produced proteins
Cell Lysis and Protein Purification. Frozen cell pellets were supplemented with 0.1 mg/mL
DNase and lysed either aerobically (for SDRs) or anaerobically (for OsrA and OsrB) using a
Thermo Spectronic French pressure cell at 1,100 PSI. The crude cell extract was centrifuged at
75,600 x g for 30 minutes, followed by filtration through a 0.2 µm nylon membrane (Fisher
Scientific) before being applied to a purification system.
505
Aerobic purification of SDRs. The filtered extract was applied to an ÄKTA pure system (Cytiva)
using a 1 mL Strep-Tactin®XT 4Flow® column (Iba Lifesciences). The column was equilibrated
with 10 volumes of equilibration buffer (100 mM Tris/HCl, pH 8.0, 150 mM NaCl) at 1 mL/min
and 4°C. Proteins were loaded via a 5 mL loop, followed by washing of non-specifically bound
proteins. Elution was performed with 5 mL elution buffer (100 mM Tris/HCl, pH 8.0, 150 mM
NaCl, and 50 mM biotin). Eluted proteins were collected in 1 mL fractions and were
concentrated using Pierce™ Protein concentrators (10 kDa), desalted, and either used directly
or transferred to storage buffer (50 mM Tris/HCl, pH 7.5, 10% (w/v) glycerol, and 50 mM NaCl)
using PD-10 desalting columns (Cytiva) before storage at -80 °C.
515
Anaerobic purification of OsrB. Anaerobic purification was performed in an anaerobic chamber.
A Strep-Tactin®XT 4Flow® gravity column (Iba Lifesciences) was used with a WET FRED
system (Iba Lifesciences) to maintain a constant flow rate of ~1 mL/min, adjusted using a lab
jack stand (LABALPHA). The column was equilibrated with anaerobic equilibration buffer (100
mM Tris/HCl, pH 8.0, 150 mM NaCl) and elution was performed with anaerobic elution buffer
(100 mM Tris/HCl, pH 8.0, 150 mM NaCl, and 50 mM biotin). Eluted proteins were collected in 1
mL tubes, concentrated and desalted using Pierce™ Protein Concentrators PES, 10K MWCO,
0.5 mL, at 10,000 × g in a microcentrifuge. Proteins were either used directly for enzymatic
assays or transferred to anaerobic storage buffer (50 mM Tris/HCl, pH 7.5, 10% (w/v) glycerol,
and 50 mM NaCl) and frozen at -80 °C.
526
Whole-cell assays of heterologous enzymes
The activity of heterologously expressed proteins was assessed under either aerobic
conditions (for SDRs) or anaerobic conditions (for OsrA and OsrB). *E. coli* Rosetta cells were
grown in media as described above, supplemented with 100 µM steroids (prepared from 10 mM
stock solutions in methanol) using a 1% (v/v) inoculum. Protein production was induced with 1
mM IPTG at an OD of 0.4-0.7 and cultures were incubated overnight at 37 °C without shaking.
Reactions were quenched by the addition of 9 volumes of methanol containing 0.5 µM
methylprednisolone as internal standard (IS). LC-MS samples were prepared as described
previously.
536
Enzymatic assays with purified SDRs
The kinetic properties of purified SDRs were determined using reaction mixtures in 96-
well plates with a total volume of 100 µL. The reaction mixture contained 25 mM Tris/HCl (pH
7.0), 1 mM NADPH, 10 µM to 1 mM steroids (diluted in 20 mM hydroxypropyl-β-cyclodextrin),
and 0.01 to 0.5 mg/mL protein, depending on the enzyme activity. Enzyme activity was
monitored by measuring the reduction of NADPH at 340 nm using a plate reader (BioTek,
Cytation 5) at 37°C. A NADPH standard curve was analyzed under identical conditions with an
extinction coefficient of 1398 M^{-1} for quantitation.
545
Enzymatic assays with purified OsrA and OsrB

The substrate preferences of OsrA and OsrB were analyzed under anaerobic conditions
using a 100 µL reaction mixture containing 50 mM Tris/HCl (pH 7.0), 200 µM methyl viologen,
0.5 mM steroids (from 10x stock solutions in methanol), and 50 µg/mL protein. Enzyme activity
was monitored by measuring electron donor reduction at 605 nm using a plate reader (BioTek,
Epoch 2) at 37°C. Quantification was performed using an electron donor standard curve
generated under the same conditions with an extinction coefficient of 1689 M^{-1} for quantitation.
553

<u>Phylogenetic tree construction</u>
Genome metadata were retrieved from a local UHGG database and used to map
genome IDs to species names. A comparative analysis was performed using BLASTp
(version 2.15.0+) to search for homologs of the target sequence against the UHGP-100
database, limiting results to the top 20000 hits. The BLASTp output was processed to map
genome IDs to species names and format the sequences in FASTA format, removing
duplicates to ensure data quality. Additional sequences were appended to the data set as
needed. Sequence alignment was performed using Clustal Omega (version 1.2.2)[48] with
output formatted as FASTA. Header sanitization was performed to remove special characters,
and duplicate sequences were filtered out using custom Python scripts to maintain alignment
integrity. Phylogenetic analysis was performed using IQ-TREE (version 2.3.6)[49] with
automatic model selection to determine the best-fitting substitution model based on the data.
The reliability of the phylogenetic trees was assessed using 1,000 ultrafast bootstrap
replicates to assess branch support. The final phylogenetic trees were visualized and
interpreted using the Interactive Tree of Life (iTOL)[50] to explore the evolutionary relationships
among the identified protein sequences.
570

<u>Metagenomics</u>
To identify relevant sequences of OsrA or OsrB, a BLAST search was first conducted
against the UHGP-100 database, applying a 49% similarity threshold based on experimental
evidence indicating that this threshold effectively identifies relevant homologs while minimizing
false positives. For OsrB, the identified protein sequences were used to construct a
phylogenetic tree using Clustal Omega for alignment and IQ-TREE with the "mtest" model
selection and 1,000 bootstrap replicates as described above. Based on the initial analysis, 25
sequences that were not phylogenetically related to OsrB were manually excluded. This
curation step ensured that only sequences relevant to the target enzymes were retained for
downstream analysis.
After phylogenetic filtering, genome information was traced back using the UHGP protein
IDs. The corresponding genomes were downloaded from the Unified Human Gastrointestinal
Genome (UHGG) database using FTP links provided in the metadata file. The genomes were
then used for further analysis, where each protein sequence was screened against the
respective genome using tblastn with a stringent e-value threshold of 1e^{-200} to ensure high
specificity. The best nucleotide sequence was selected for each protein based on coverage and
bit score and subsequently compiled into a combined FASTA file.
Metagenomic samples were downloaded from the Sequence Read Archive (SRA). Reads were
quality trimmed to remove adapter sequences using TrimGalore with default settings,[51] and
potential human contamination was removed by mapping the reads to the human reference
genome (T2T-CHM13v2.0) using Bowtie2 (version 2.5.3) and removing the mapped reads with
Samtools (version 1.61.1).[52,53] Samples were then mapped to the gene reference datasets for
osrA and *osrB* using Bowtie2 (version 2.5.3), and copies per million (CPM values) were
calculated for each gene in each sample. Samples with total read counts below 1,000,000 were
excluded.
Metagenomic data were filtered to ensure quality for statistical analyses. Zero values
were replaced with 1e^{-6} for statistical assessment. A 99th percentile filter was applied to CPM

values for each gene (*osrA* and *osrB*) to remove extreme outliers. Normality was assessed
using the Shapiro-Wilk test; if both groups were normal ($p > 0.05$), a two-sided Welch's t-test
was used to determine if there was any significant difference between the groups, regardless of
direction. If normality was not met, a two-sided Mann-Whitney U test was applied. This
conservative approach ensured that differences were detected without assuming the direction of
the effect, providing flexibility in hypothesis testing. Analyses were performed using Python with
Scipy, Pandas, and Seaborn libraries.

605
606
607 **Figure 1.** ***Clostridium steroidoreducens* is a novel steroid-enriched species.** (A) Schematic overview
608 of steroid enrichment and strain isolation experiments. (B) HCS.1 strain colonies on cortisol-infused
609 media. (C) HCS.1 strain colonies on progesterone-infused media. Scale bar, 200 μm. (D) Microbial
610 community profile of the final steroid hormone enrichment passage based on16S rRNA amplicon
611 sequencing. (E) Circular representation of the HCS.1 genome.

612
613
Figure 2. *C. steroidoreducens* HCS.1 possesses promiscuous 3-oxo-Δ⁴-beta steroid hormone
reductase activity. (A) Products formed from *C. steroidoreducens* HCS.1 incubation with cortisol. (B)
Time-course analysis of cortisol metabolism by *C. steroidoreducens* HCS.1. (C) Proposed pathway for
cortisol reduction by *C. steroidoreducens* HCS.1. DHC and THC stand for dihydrocortisol and
tetrahydrocortisol, respectively. (D) Steroid substrates tested for *C. steroidoreducens* HCS.1. (E)
Measured *C. steroidoreducens* HCS.1 steroid substrate consumption.

Figure 3. OsrB is a 3-oxo-Δ⁴-steroid hormone reductase. (A) Gene expression of *C. steroidoreducens* HCS.1 in the presence versus absence of cortisol. Gray and red dashed lines indicate genes with statistical significance and >2-fold induction in response to cortisol, respectively. (B) Crystal structure of Fe-S flavoenzyme 2,4-dienoyl-CoA reductase (FadH) bound to ligands (PDB code: 1PS9). (C) Percent sequence identity of OsrA and OsrB to Fe-S flavoenzymes FadH and bile acid reductases BaiH and BaiCD. (D) Conversion of cortisol to 5β-dihydrocortisol (DHC) by *E. coli* expressing *osrA* or *osrB* versus an empty vector control. (E) Rate of reduction of indicated steroid hormones by purified OsrB.

628
629
Figure 4. OsrC is a 3-oxo-5β-steroid hormone oxidoreductase. (A) Phylogenetic analysis of *C.*
steroidoreducens HCS.1 SDR domain-containing protein sequences. Product formed from 5β-
dihydrocortisol by *E. coli* strains overexpressing SDR domain-containing proteins are shown with their
respective gene identifiers. THC stand for tetrahydrocortisol. (B) Kinetic parameters of reduction of
indicated steroid hormones by purified SDR domain-containing proteins. Abbreviations stand for
dihydrotestosterone (DHT), dihydroprogesterone (DHP), dihydrocortisol (DHC), and dehydrolithocholic
acid (DLCA).

637
638
Figure 5. OsrA is a 3-oxo-Δ¹-reductase essential for complete reduction of synthetic
corticosteroids. (A) Structure of synthetic corticosteroids used in assays. (B) Products formed from
synthetic corticosteroids following incubation with *C. steroidoreducens* HCS.1 cells. (C) Percent
prednisolone conversion to cortisol following incubation of *E. coli* cells with *osrA*- and *osrB*-expressing
plasmids or an empty vector control. (D) *C. steroidoreducens* HCS.1 steroid reduction pathway identified
in this study.

645
Figure 6. Steroid reductase activities are widespread in gut bacteria and elevated in active Crohn's
disease. (A) Representative images of gut bacteria grown on progesterone-infused media, showing
steroid clearance/precipitate accumulation-positive (blue background) and -negative (gray background)
colonies. Scale bar, 200 μm. (B) Corticosteroids produced by gut bacterial isolates after incubation with
cortisol. The protein with the highest sequence identity to OsrB encoded by each genome was used to
generate the tree. (C) Corticosteroids produced by gut bacterial isolates after incubation with
prednisolone. Identity refers to the sequence identity of the protein with the highest sequence identity to
OsrA encoded by each genome. (D) reads mapping to *osrA* and *osrB* homologs in metagenomes from
healthy and Crohn's disease (CD) patients. Only metagenomes with at least one read mapping to a gene
are included in the analysis. NQ refers to the not quantified statistical difference, due to the low number of
healthy metagenomes with reads mapping to *osrA*. CPM refers to copies per million. (E) Difference in
osrB homolog abundance for taxa showing the greatest changes in relative abundance between healthy
and CD metagenomes.

REFERENCES

1. Nussey, S. & Whitehead, S. The adrenal gland. in *Endocrinology: An Integrated Approach* (BIOS Scientific Publishers, 2001).
2. Nussey, S. & Whitehead, S. The gonad. in *Endocrinology: An Integrated Approach* (BIOS Scientific Publishers, 2001).
3. Buffie, C. G. et al. Precision microbiome reconstitution restores bile acid mediated resistance to Clostridium difficile. *Nature* 517, 205–208 (2015).
4. Paik, D. et al. Human gut bacteria produce TH17-modulating bile acid metabolites. *Nature* 603, 907–912 (2022).
5. Campbell, C. et al. Bacterial metabolism of bile acids promotes generation of peripheral regulatory T cells. *Nature* 581, 475–479 (2020).
6. Adlercreutz, H., Martin, F., Järvenpää, P. & Fotsis, T. Steroid absorption and enterohepatic recycling. *Contraception* 20, 201–223 (1979).
7. Decker, H. A. et al. Metabolism of 4-C 14-cortisol in man: body distribution and rates of conjugation. *J. Clin. Endocrinol. Metab.* 16, 1137–1150 (1956).
8. Peterson, R. E., Wyngaarden, J. B., Guerra, S. L., Brodie, B. B. & Bunim, J. J. THE PHYSIOLOGICAL DISPOSITION AND METABOLIC FATE OF HYDROCORTISONE IN MAN. *J. Clin. Invest.* 34, 1779–1794 (1955).
9. Sandberg, A. A. & Slaunwhite, W. R. Metabolism of 4-C14-testosterone in human subjects. I. Distribution in bile, blood, feces and urine. *J. Clin. Invest.* 35, 1331–1339 (1956).
10. Adlercreutz, H. & Martin, F. Biliary excretion and intestinal metabolism of progesterone and estrogens in man. *J. Steroid Biochem.* 13, 231–244 (1980).
11. Migeon, C. J., Paul, A. C., Samuels, L. T. & Sandberg, A. A. Metabolism of 4-C14-corticosterone in man. *J. Clin. Endocrinol. Metab.* 16, 1291–1298 (1956).
12. Möstl, E. & Palme, R. Hormones as indicators of stress. *Domest. Anim. Endocrinol.* 23, 67–74 (2002).
13. Palme, R., Rettenbacher, S., Touma, C., El-Bahr, S. M. & Möstl, E. Stress hormones in mammals and birds: comparative aspects regarding metabolism, excretion, and noninvasive measurement in fecal samples. *Ann. N. Y. Acad. Sci.* 1040, 162–171 (2005).
14. Nabarro, J. D., Moxham, A., Walker, G. & Slater, J. D. Rectal hydrocortisone. *Br. Med. J.* 2, 272–274 (1957).
15. Wade, A. P., Slater, J. D., Kellie, A. E. & Holliday, M. E. Urinary excretion of 17-ketosteroids following rectal infusion of cortisol. *J. Clin. Endocrinol. Metab.* 19, 444–453 (1959).
16. Martin, F., Peltonen, J., Laatikainen, T., Pulkkinen, M. & Adlercreutz, H. Excretion of progesterone metabolites and estriol in faeces from pregnant women during ampicillin administration. *J. Steroid Biochem.* 6, 1339–1346 (1975).
17. Ly, L. K. et al. Bacterial steroid-17,20-desmolase is a taxonomically rare enzymatic pathway that converts prednisone to 1,4-androstanediene-3,11,17-trione, a metabolite that causes proliferation of prostate cancer cells. *J. Steroid. Biochem. Mol. Biol.* 199, 105567 (2020).
18. Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R. & Goodman, A. L. Mapping human microbiome drug metabolism by gut bacteria and their genes. *Nature* 570, 462–467 (2019).
19. Latif, S. A., Sheff, M. F., Ribeiro, C. E. & Morris, D. J. Selective inhibition of sheep kidney 11β-hydroxysteroid dehydrogenase isoform 2 activity by 5α-reduced (but not 5β) derivatives of adrenocorticosteroids. *Steroids* 62, 230–237 (1997).
20. Honour, J. W., Borriello, S. P., Ganten, U. & Honour, P. Antibiotics attenuate experimental hypertension in rats. *J. Endocrinol.* 105, 347–350 (1985).
21. N, P. *et al.* Commensal bacteria promote endocrine resistance in prostate cancer through androgen biosynthesis. *Science* **374**, (2021).

22. Terrisse, S., Zitvogel, L. & Kroemer, G. Effects of the intestinal microbiota on prostate cancer
treatment by androgen deprivation therapy. *Microb. Cell Graz Austria* 9, 202–206 (2022).
23. Stokes, N. A. & Hylemon, P. B. Characterization of delta 4-3-ketosteroid-5 beta-reductase and 3
beta-hydroxysteroid dehydrogenase in cell extracts of Clostridium innocuum. *Biochim. Biophys.*
Acta **836**, 255–261 (1985).
24. Penning, T. M. & Covey, D. F. 5β-Dihydrosteroids: Formation and Properties. *Int. J. Mol. Sci.* **25**,
8857 (2024).
25. Parks, D. H. et al. A standardized bacterial taxonomy based on genome phylogeny substantially
revises the tree of life. *Nat. Biotechnol.* 36, 996–1004 (2018).
26. McCurry, M. D. et al. Gut bacteria convert glucocorticoids into progestins in the presence of
hydrogen gas. *Cell* 187, 2952-2968.e13 (2024).
27. Devendran, S., Mythen, S. M. & Ridlon, J. M. The desA and desB genes from Clostridium
scindens ATCC 35704 encode steroid-17,20-desmolase. *J. Lipid Res.* **59**, 1005–1014 (2018).
28. Maini Rekdal, V. et al. A widely distributed metalloenzyme class enables gut microbial metabolism
of host- and diet-derived catechols. *eLife* 9, e50845 (2020).
29. Hubbard, P. A., Liang, X., Schulz, H. & Kim, J.-J. P. The crystal structure and reaction mechanism
of Escherichia coli 2,4-dienoyl-CoA reductase. *J. Biol. Chem.* 278, 37553–37560 (2003).
30. Little, A. S. et al. Dietary- and host-derived metabolites are used by diverse gut bacteria for
anaerobic respiration. *Nat. Microbiol.* 9, 55–69 (2024).
31. Funabashi, M. et al. A metabolic pathway for bile acid dehydroxylation by the gut microbiome.
Nature 582, 566–570 (2020).
32. Kang, D.-J., Ridlon, J. M., Moore, D. R., Barnes, S. & Hylemon, P. B. Clostridium scindens baiCD
and baiH genes encode stereo-specific 7alpha/7beta-hydroxy-3-oxo-delta4-cholenoic acid
oxidoreductases. *Biochim. Biophys. Acta* 1781, 16–25 (2008).
32. Edenharder, R. & Schneider, J. 12 beta-dehydrogenation of bile acids by Clostridium
paraputrificum, C. tertium, and C. difficile and epimerization at carbon-12 of deoxycholic acid by
cocultivation with 12 alpha-dehydrogenating Eubacterium lentum. *Appl. Environ. Microbiol.* **49**,
964–968 (1985).
33. Doden, H. L. & Ridlon, J. M. Microbial Hydroxysteroid Dehydrogenases: From Alpha to Omega.
Microorganisms 9, 469 (2021).
34. Doden, H. *et al.* Metabolism of Oxo-Bile Acids and Characterization of Recombinant 12α-
Hydroxysteroid Dehydrogenases from Bile Acid 7α-Dehydroxylating Human Gut Bacteria. *Appl.*
Environ. Microbiol. **84**, e00235-18 (2018).
35. Ardizzone, S. & Bianchi Porro, G. Comparative tolerability of therapies for ulcerative colitis. *Drug*
Saf. 25, 561–582 (2002).
36. Yang, R. & Yu, Y. Glucocorticoids are double-edged sword in the treatment of COVID-19 and
cancers. *Int. J. Biol. Sci.* 17, 1530–1537 (2021).
37. Almeida, A. et al. A unified catalog of 204,938 reference genomes from the human gut
microbiome. *Nat. Biotechnol.* 39, 105–114 (2021).
38. Lewis, J. D. et al. Inflammation, Antibiotics, and Diet as Environmental Stressors of the Gut
Microbiome in Pediatric Crohn's Disease. *Cell Host Microbe* 18, 489–500 (2015).
39. Ha, C. W. Y. et al. Translocation of Viable Gut Microbiota to Mesenteric Adipose Drives Formation
of Creeping Fat in Humans. *Cell* 183, 666-683.e17 (2020).
40. Hall, A. B. et al. A novel Ruminococus gnavus clade enriched in inflammatory bowel disease
patients. *Genome Med.* 9, 103 (2017).
41. Lloyd-Price, J. *et al.* Multi-omics of the gut microbial ecosystem in inflammatory bowel diseases.
Nature **569**, 655–662 (2019).
42. Arp, G. *et al.* Gut Bacteria Encode Reductases that Biotransform Steroid Hormones. *bioRxiv*
2024.10.04.616736 (2024) doi:10.1101/2024.10.04.616736.

43. Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial genome
assemblies from short and long sequencing reads. *PLoS Comput. Biol.* 13, e1005595 (2017).
44. Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Completing bacterial genome assemblies with
multiplex MinION sequencing. *Microb. Genomics* 3, e000132 (2017).
45. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. *Bioinformatics* 36, 1925–1927 (2019).
46. O, S. *et al.* Bakta: rapid and standardized annotation of bacterial genomes via alignment-free
sequence identification. *Microb. Genomics* **7**, (2021).
47. Mi, L., W, H. & S, A. Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. *Genome Biol.* **15**, (2014).
48. Sievers, F. et al. Fast, scalable generation of high-quality protein multiple sequence alignments
using Clustal Omega. *Mol. Syst. Biol.* 7, 539 (2011).
49. Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* 32, 268–274
(2015).
50. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display
and annotation. *Bioinformatics* 23, 127–128 (2007).
51. Krueger, F. et. al. FelixKrueger/TrimGalore: v0.6.10. *Zenodo* doi: 10.5281/zenodo.7598955
(2023).
52. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–
359 (2012).
53. H, L. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinforma. Oxf. Engl.* **25**,
(2009).

Extended Data Figure 1. Phylogenetic analysis supporting assignment of HCS.1 within the
***Clostridium* genus.** Tree generated from alignment of conserved marker genes.

785
786
787 **Extended Data Figure 2. Product of *C. steroidoreducens* HCS.1 incubation with progesterone.**
788 Major progesterone product formed by *C. steroidoreducens* HCS.1. Comparison with reference standards
789 confirms 3β-,5β-tetrahydroprogesterone (THP) production.

790
791
Extended Data Figure 3. 12% SDS-PAGE of _E. coli_ lysates expressing SDR domain-containing
proteins. SDS-PAGE analysis showing the heterologous production of _C. steroidoreducens_ SDR domain-
containing proteins in _E. coli_, pre- and post-isopropyl β-D-1-thiogalactopyranoside (IPTG) induction. Red
boxes highlight expressed protein.

Extended Data Figure 4. Reaction rates of enriched BLEONJ_2554 on indicated substrates.
Reaction rates for the 3'-reduction of 5-reduced steroids by anaerobically purified short-chain
dehydrogenase BLEONJ_2554.

800
Extended Data Figure 5. Reaction rates of enriched OsrC on indicated substrates. Reaction rates
for the 3'-reduction of 5-reduced steroids by anaerobically purified short-chain dehydrogenase OsrC.

803
Extended Data Figure 6. OsrA and OsrB products from Δ^1- and Δ^4-steroid hormone substrates. (A)
Products following prednisolone incubation with purified OsrA or OsrB. Comparison to a cortisol reference
standard confirm that OsrA generates the Δ^1-reduced product and show that OsrB produces a distinct
cortisol isomer. (B) Products following boldenone incubation with purified OsrA or OsrB. Comparison to
reference standards confirm that OsrA and OsrB generate Δ^1- and Δ^4-reduced products, respectively.

Extended Data Figure 7. *osrA* levels in expanded dataset of healthy and Crohn's disease
metagenomes. (A) Reads mapping to *osrA* homologs in expanded dataset of Crohn's disease (CD)
patient metagenomes relative to healthy controls. (B) Difference in *osrA* homolog levels in CD relative to
healthy metagenomes.

817
Extended Data Figure 8. Association between *osrA* and *osrB* and Crohn's disease in Integrative
Human Microbiome Project metagenomes. (A) Reads mapping to *osrA* and *osrB* homologs in Crohn's
disease (CD) metagenomes relative to non-IBD controls. CPM stands for copies per million. (B)
Difference in *osrB* homolog levels from taxa with the most significant changes in relative abundance
between CD dysbiotic and non-IBD metagenomes. (C) Difference in *osrA* homolog levels in CD dysbiotic
relative to non-IBD metagenomes.

824
825
826 **Extended Data Figure 9. Association between *osrA* and *osrB* and steroid usage in the Lewis et al**
827 **study.** Reads mapping to *osrA* and *osrB* homologs in Crohn's disease (CD) patient metagenomes
828 grouped based on patient corticosteroid treatment.