rbfox1 loss in zebrafish drives HPI axis hyperactivation via *bdnf/trkb2* pathway and allostatic overload during later life

Adele Leggieri[1], Judit García-González[2], Saeedeh Hosseinian[1], Peter Ashdown[1], Sofia Anagianni[1], Xian Wang[1], William Havelange[1], Noèlia Fernàndez-Castillo[3,4,5,6], Bru Cormand[3,4,5,6] and Caroline H. Brennan[1]*

Centre for Brain and Behaviour, School of Biological and Behavioural Sciences, Queen Mary University of London, Mile End Rd, London, E1 4NS, United Kingdom

Department of Genetics and Genomic Sciences, Icahn School of Medicine, Mount Sinai, New York City, NY 10029, USA

Departament de Genètica, Microbiologia i Estadística, Facultat de Biologia, Universitat de Barcelona, Barcelona, Catalunya, 08028, Spain

Centro de Investigación Biomédica en Red de Enfermedades raras (CIBERER), Spain

Institut de Biomedicina de la Universitat de Barcelona, Barcelona, Catalunya, 08028, Spain

Institut de recerca Sant Joan de Déu, Espluges de Llobregat, Catalunya, 08950, Spain

* Correspondence: Caroline H. Brennan, c.h.brennan@qmul.ac.uk

ABSTRACT

Mutations in the *RBFOX1* gene are associated with psychiatric disorders but how RBFOX1 influences
psychiatric disorder vulnerability remains unclear. Recent studies showed that RBFOX proteins
mediate the alternative splicing of PAC1, a critical HPA axis activator. Further, RBFOX1 dysfunction is
linked to dysregulation of BDNF/TRKB, a pathway promoting neuroplasticity, neuronal survival, and
stress resilience. Hence, RBFOX1 dysfunction may increase psychiatric disorder vulnerability via HPA
axis dysregulation, leading to disrupted development and allostatic overload. To test this hypothesis,
we generated a zebrafish *rbfox1* loss-of-function (LoF) line and examined behavioural and molecular
effects during development. We found that *rbfox1* LoF mutants exhibited hyperactivity, impulsivity
and heightened arousal, alongside alterations in proliferation – traits associated with
neurodevelopmental and stress-related disorders. In adults, loss of *rbfox1* function led to decreased
fertility and survival, consistent with allostatic overload. At the molecular level, at larval stages *rbfox1*
LoF disrupted the expression of *bdnf*, *trkb2*, *pac1a-hop* and HPI axis genes. Pharmacological
intervention targeting TRKB restored HPI axis and *bdnf/trkb2* gene expression. In adults, dysregulation
of *bdnf/trkb2* and HPI axis genes was only seen following acute stress exposure. Our findings revealed
a fundamental role for RBFOX1 in integrating stress responses through its regulation of BDNF/TRKB
and neuroendocrine signalling. By uncovering how RBFOX1 modulates stress adaptation from
development to adulthood, our study provides novel insights into the molecular basis of stress
resilience and psychiatric disease vulnerability, suggesting *RBFOX1* loss predisposes to psychiatric
diseases through HPA axis hyperactivation during development, impairing adaptation and heightening
vulnerability to allostatic overload in later life.

INTRODUCTION

Psychiatric disorders are influenced by multiple genes, with a substantial portion of their heritability
linked to common genetic variations[1]. Identifying gene variants associated with multiple psychiatric
disorders is crucial for understanding their shared genetic underpinnings and enhancing therapeutic
strategies.

RNA binding fox-1 homologue 1 (RBFOX1), also known as FOX1 or ataxin 2-binding protein 1 (A2BP1),
is a splicing factor highly conserved among vertebrates and expressed in the heart, brain and skeletal
muscle, where it contributes to normal development and function[2,3]. Alterations in *RBFOX1* expression
or function have been associated with susceptibility to psychiatric disorders, particularly those
associated with changes in stress-related behaviour[4–6]. In a cross-disorder genome-wide association
study (GWAS), *RBFOX1* emerged as the second most pleiotropic *locus*, showing association with seven
out of the eight disorders studied: schizophrenia, bipolar disorder, depression, attention-
deficit/hyperactivity disorder (ADHD), autism spectrum disorder (ASD), obsessive-compulsive disorder
and Tourette syndrome[1]. In mice and zebrafish, *Rbfox1/rbfox1* knockout caused hyperactivity,
increased anxiety-like behaviour and altered social behaviour[2,6]. However, the mechanisms by which
RBFOX1 genetic variants contribute to psychiatric disease are poorly understood.

In both mice and zebrafish, it has been suggested that Rbfox1 regulates the Pituitary Adenylate
Cyclase-Activating Polypeptide Type 1 Receptor (Pac1) alternative splicing, an important mediator of
corticotropin releasing hormone (CRH) synthesis in the hypothalamus[4,7]. CRH is the first hormone to
be secreted in response to stress by the hypothalamic-pituitary-adrenal (HPA) axis (hypothalamic-
pituitary-interrenal (HPI) axis in fish)[8]. In mammals there are several PAC1 isoforms with the
predominant isoforms in the brain being PAC1-hop (long isoform) and PAC1-short[9]. In mice and

zebrafish, following acute stress exposure, both *Pac1-hop/pac1-hop* and *Pac1-short/pac1-short*
expression increased, while at the late recovery phase, only *Pac1-hop* was still up-regulated[4].

Another regulator of the stress response and a resilience factor against chronic stress-induced
psychopathology is the brain derived neurotrophic factor (BDNF)/Tropomyosin receptor kinase B
(TRKB) pathway[10–12]. BDNF is a neurotrophin possessing a pivotal role in the modulation of
neurotransmission and synaptic plasticity and dysregulation of the BDNF/TRKB pathway has been
associated with several neuropsychiatric diseases, including anxiety/stress disorders[13,14]. In human
neural stem cells, *RBFOX1* knockdown increased *BDNF* expression levels[15], while another study in mice
identified *TrkB* as a target of RBFOX1 within the hippocampus[16]. Further, PAC1 activation can elevate
BDNF levels and potentiate TRKB activity, enhancing BDNF/TRKB neuroprotective and plasticity-
promoting effects, especially in the context of stress response and neuropsychiatric health[17].

Given that both PAC1 and BDNF/TRKB influence how the brain adapts to and manages stress, RBFOX1
variants may increase susceptibility to psychiatric disorders through dysregulation of the stress
response, leading to adaptive plasticity and disrupted development in the short-term and allostatic
overload in the long-term. Allostasis refers to the collective processes by which the body actively
maintains stability (or homeostasis) in response to environmental stressors[12]. When allostasis primary
mediators (e.g., HPA axis hormones, catecholamines, cytokines) are overactivated or fail to return to
normal, it leads to an allostatic state[18,19]. The cumulative results of an allostatic state are referred to
as allostatic load and, when maladaptive, allostatic overload[18]. Excess glucocorticoid exposure during
early life and early life stress have been shown to cause prolonged activation of the allostatic systems,
ultimately leading to allostatic overload[19–22], while neurotrophic factors such as BDNF play a key role
in regulating adaptive plasticity and mechanisms counteracting damage caused by allostatic
overload[12]. Here, to explore the possibility that *RBFOX1* loss of function (LoF) leads to increased
vulnerability to psychiatric disease through HPA axis hyperactivation and allostasis-induced
adaptation during development, we generated a zebrafish line carrying a predicted *rbfox1* LoF
mutation using CRISPR-Cas9 gene editing and assessed behavioural and molecular changes at different
developmental stages. We hypothesized that RBFOX1 regulates HPA axis activity through an effect on
BDNF/TRKB signalling leading to disrupted brain development.

RESULTS

Generation of a loss of function *rbfox1* line

CRISPR-Cas9 genome editing generated a 19 base pair deletion (NM_001005596.1, nucleotides 120-
138, TCCCATCGGCCCAGTTCGC) that introduced a premature termination codon (PTC) at position 58
in the *rbfox1* amino acid sequence. Details regarding the generation of the line, nucleotide and amino
acid sequences can be found in our previous study[2]. Wild type animals are denoted throughout as
rbfox1$^{+/+}$, heterozygous mutants as *rbfox1$^{+/19del}$* and homozygous mutants as *rbfox1$^{19del/19del}$*.

To evaluate whether the PTC elicited mRNA non-sense mediated decay (NMD) and consequent
reduction of *rbfox1* mRNA in mutant fish, we examined *rbfox1* expression by quantitative Real-Time
polymerase chain reaction (qPCR) and by *in situ* hybridisation (ISH). As *RBFOX1* itself is alternatively
spliced to generate nuclear and cytoplasmic isoforms[7], we designed primers targeting all *rbfox1*
zebrafish isoforms available on NCBI (see Supplementary Table1). qPCR showed that *rbfox1* transcript
levels were significantly lower in mutant larvae compared to *rbfox1$^{+/+}$* siblings (p$_{rbfox1}$$^{+/+}$ $_{vs\ rbfox1}$$^{+/19del}$ <
0.05, p$_{rbfox1}$$^{+/+}$ $_{vs\ rbfox1}$$^{19del/19del}$ < 0.01), and ISH showed that *rbfox1* was not detectable in *rbfox1$^{19del/19del}$*
fish, at either larval or adult stages (Supplementary Figure 1A-C).

These results confirm degradation of defective *rbfox1* mRNA in mutant fish.

***rbfox1* mutant fish show hyperactivity, impulsivity and hyperarousal behaviour**

As mutations in *RBFOX1* locus have been linked to several psychiatric diseases in humans, we assessed
zebrafish larvae and adult fish for phenotypic traits associated with such disorders and examined
rbfox1 mRNA distribution in *rbfox1$^{+/+}$* larvae and adults (Tübingen strain).

In humans, *RBFOX1* copy number variants (CNVs) and LoF mutations are risk factors for ADHD[6]. Two
major ADHD traits are hyperactivity and increased impulsivity. We therefore assessed hyperactive and
impulsive behaviour of *rbfox1$^{+/+}$*, *rbfox1$^{+/19del}$* and *rbfox1$^{19del/19del}$* 5 days post fertilization (dpf) larvae
and adults. When we measured larval locomotion, we observed a gene dosage effect on distance
travelled, whereby *rbfox1* mutant larvae travelled greater distances than wild type siblings ($p_{rbfox1^{+/+} vs}$
$_{rbfox1^{+/19del}}$ < 0.05, $p_{rbfox1^{+/+} vs rbfox1^{19del/19del}}$ < 0.0001, $p_{rbfox1^{+/19del} VS rbfox1^{19del/19del}}$ < 0.05) (Figure 1A), and a
significant increase in the swimming speed of *rbfox1$^{19del/19del}$* larvae ($p_{rbfox1^{+/+} vs rbfox1^{19del/19del}}$ < 0.0001, p
$_{rbfox1^{+/19del} VS rbfox1^{19del/19del}}$ < 0.05) (Figure 1B). This is in line with hyperactivity observed in adult
rbfox1$^{19del/19del}$ in our previous study[2]. We also measured larval burst swimming, a parameter
previously used as a measure to predict impulsive behaviour in zebrafish larvae[23]. We found a
significant increase in the number of peaks (acceleration events when the fish travelled > 5 mm in <
12 sec) in *rbfox1$^{19del/19del}$* larvae ($p_{rbfox1^{+/+} vs rbfox1^{19del/19del}}$ < 0.0001, $p_{rbfox1^{+/19del} VS rbfox1^{19del/19del}}$ < 0.0001)
(Figure 1C). Impulsive behaviour was then assessed in adult (7 months old) fish using the 5-choice
serial reaction time task (5-CSRTT)[24]. The 5-CSRTT consists of five stages (Supplementary Table2) each
run for at least a week until fish are promoted to the next stage. We found that 79% *rbfox1$^{+/+}$,* 72%
rbfox1$^{+/19del}$ and 62% *rbfox1$^{19del/19del}$* learned the task within 9 weeks. We found no significant
differences in the correct responses (stages 2-5, Figure 1D-G). In stage 5, we found a significant
difference in the number of premature responses (increased impulsivity) such that *rbfox1$^{19del/19del}$* fish
were more impulsive compared to *rbfox1$^{+/+}$* siblings ($p < 0.05$) (Figure 1H).

RBFOX1 CNVs have also been identified in individuals with schizophrenia[6,15]. Deficits in habituation to
acoustic startle reflex are seen in both schizophrenic patients and animal models of schizophrenia[25].
The acoustic startle assay has been widely employed to measure habituation in animals, including
zebrafish[26]. The magnitude of the startle response and the extent of habituation can also serve as
indicators of hyperarousal linked to heightened stress responses[27]. We therefore tested 5dpf zebrafish
larvae (*rbfox1$^{+/+}$*, *rbfox1$^{+/19del}$* and *rbfox1$^{19del/19del}$*) in the habituation to acoustic startle response assay.
Consistent with our hyperactivity assay, during the baseline (first 10 min), the genotype had a
significant main effect on distance travelled [Effect of genotype: $\chi2(2) = 8.3397$, $p < 0.05$] whereby
rbfox1$^{19del/19del}$ larvae travelled greater distances than *rbfox1$^{+/+}$* siblings ($p < 0.05$). During the startle
stimuli, we observed a main effect of genotype [Effect of genotype: $\chi2(2) =33.536$, $p < 0.0001$] and
stimulus number [Effect of stimulus number: $\chi2(9) = 487.968$, $p < 0.0001$] on distance travelled, and
a significant two-way interaction between genotype and stimulus number [Effect of genotype by
stimuli: $\chi2(18) = 31.511$, $p < 0.05$], whereby *rbfox1$^{19del/19del}$* larvae travelled greater distances than
rbfox1$^{+/+}$ ($p < 0.001$) and *rbfox1$^{+/19del}$* ($p < 0.01$) (Figure 1I). When assessed for the rate of habituation,
larvae showed a habituation response to repeated acoustic startle consistent with previous reports[28]:
100% of *rbfox1$^{+/+}$*animals responded to the first acoustic stimulus, but only 22% responded to the last.
When we examined the rate of habituation over time, we observed a significant genotype effect
[Effect of genotype: $\chi2(2) =7.2676$, $p < 0.05$] and a significant two-way interaction between genotype
and stimulus number [Effect of genotype by stimulus number: $\chi2(18) =132.8476$, $p < 0.001$] whereby

$rbfox1^{19del/19del}$ showed reduced rate of habituation and a greater proportion of responders compared
to $rbfox1^{+/+}$ siblings ($p_{rbfox1^{+/+} \text{ vs } rbfox1^{19del/19del}}$ < 0.05) (Figure 1J).

SNPs in *RBFOX1* locus have also been associated with anxiety[6,29]. Hence, we assessed anxiety-like
behavior in 5dpf zebrafish larvae using the forced light-dark transition (FLDT) assay and in adult
zebrafish using the novel tank diving (NTD) assay. In the FLDT assay, zebrafish are exposed to sudden
transitions in illumination with effects on locomotion and amplitude of response on transition from
dark to light being used as a measure of anxiety-like behaviour: the increased locomotion/startle
response upon the light-dark transition is attributed to increased stress/anxiety level[30,31]. Consistent
with our acoustic startle assay, we found that during the baseline (first 5 min) (Figure 1K) genotype
had a significant main effect on distance travelled [Effect of genotype: $\chi2(2)$ = 8.8262, $p < 0.05$]
whereby $rbfox1^{19del}$ larvae travelled greater distances than $rbfox1^{+/+}$ siblings ($p_{rbfox1^{+/+} \text{ vs } rbfox1^{+/19del}}$ <
0.05, $p_{rbfox1^{+/+} \text{ vs } rbfox1^{19del/19del}}$ < 0.05). During the 1min light flash (min 5-6 of the assay) we found a
significant main effect of genotype on distance travelled [Effect of genotype: $\chi2(2)$ = 12.8139, $p <$
0.01] with $rbfox1^{19del/19del}$ larvae travelling greater distances than $rbfox1^{+/+}$ siblings ($p < 0.01$) (Figure
1K). On transition from dark to light (from sec 240 to sec 360) we found a significant main effect of
time [Effect of time: $\chi2(118)$ = 705.091, $p < 0.001$] and genotype [Effect of genotype: $\chi2(2)$ = 10.289,
$p < 0.01$], and a significant two-way interaction between time and genotype [Effect of time by
genotype: $\chi2(236)$ = 188.008, $p < 0.001$] on the amplitude of response, whereby $rbfox1^{19del/19del}$ larvae
startled more than $rbfox1^{+/+}$ ($p < 0.01$) and $rbfox1^{+/19del}$ ($p < 0.01$) siblings (Figure 1L). As there was a
significant difference in basal locomotion, we also examined the amplitude of response on transition
from dark to light normalizing the data against baseline, as reported previously[32]. We observed similar
results as in absence of normalization: significant main effect of time [Effect of time: $\chi2(118)$ =
1705.943, $p < 0.001$] and genotype [Effect of genotype: $\chi2(2)$ = 12.756, $p < 0.01$], and a significant
two-way interaction between time and genotype [Effect of time by genotype: $\chi2(236)$ = 336.523, $p <$
0.001] whereby $rbfox1^{19del/19del}$ larvae startled more than $rbfox1^{+/+}$ siblings ($p_{rbfox1^{+/+} \text{ vs } rbfox1^{19del/19del}}$ <
0.01).
183
Anxiety-like behaviour in adult animals was assessed using the NTD assay. When introduced to a novel
tank, zebrafish will first dive to the bottom of the tank, to seek protection, and then gradually increase
their swimming over time[33]. We found no significant differences between *rbfox1* genotypes over the
entire duration of the NTD assay ($p > 0.05$). However, during the first minute of the assay, we found a
significant two-way interaction between genotype and the time spent at the bottom of the tank [Effect
of genotype by proportion at the bottom tank: $\chi2(10)$ =22.3333, $p < 0.05$], such that $rbfox1^{19del/19del}$
fish spent less time in the bottom than $rbfox1^{+/+}$ fish ($p < 0.001$) (Supplementary Figure 3A). In line
with previous findings[2], we observed no significant differences in distance travelled ($p > 0.05$)
(Supplementary Figure 3B), nor in the number of the transitions to the top area of the tank between
rbfox1 genotypes ($p > 0.05$) (Supplementary Figure 3C).
194
Thus, our data showed that loss of *rbfox1* resulted in behavioural changes that resemble human
psychiatric disorder traits and suggest that adaptation occurs as the animal develop. Consistent with
our behavioural results and as seen previously[2,3], when we assessed *rbfox1* mRNA distribution in
$rbfox1^{+/+}$ larvae and adults (Tübingen strain), we found that *rbfox1* was expressed in regions of the
brain involved in the response to stress, in social and emotional behaviour, and in reward and learning
(Supplementary Figure 2A-B) in agreement with data in rodents[34] and humans[35].

201
***rbfox1* LoF disrupts zebrafish larvae HPI axis and *bdnf/trkb2* pathway**

As larvae showed altered stress response and RBFOX1 has been linked to regulation of HPA axis and
BDNF/TRKB gene expression[4,15,16], we examined the expression of components of the HPI axis and of
bdnf/trkb2. Since we did not observe significant differences in behaviour between wild type and
heterozygous animals, we employed wild type and homozygous animals only.

We performed qPCR experiments in 5dpf zebrafish larvae to assess changes in the expression levels
of the HPI axis markers corticotropin releasing hormone (*crhb*), mineralocorticoid receptor (*mr*), and
glucocorticoid receptor (*gr*), and of *bdnf* and *trkb2*. In teleosts, the duplication of the genome gave
rise to two *CRH* genes, *crha* and *crhb,* and two *TRKB* genes, *trkb1* and *trkb2*[36,37]. However, only *crhb*
and *trkb2* are regarded as the orthologues of the mammalian *CRH* and *TRKB* respectively[37,38]. Then,
unlike other teleosts, zebrafish possess a single copy of the *MR* (*mr*) and *GR* (*gr*) genes, and a single
copy of the *BDNF* gene (*bdnf*)[37,39]. In mammals, several *TRKB* splicing isoforms are present, but the
most abundant ones are the full-length (TRKB.FL/TK+) and the truncated (TRKB.T1/TK-), this latter
lacking the catalytic tyrosine kinase (TK) domain[40]. As in zebrafish the presence of both *trkb2* full-
length and truncated forms has been demonstrated[41], here we used a pair of *trkb2 TK+/TK-* common
primers, and another pair targeting only *TK+* to distinguish effects on the expression of the two
isoforms.

In *rbfox1*[19del/19del] larvae, in we observed a significant up-regulation of *bdnf* ($p < 0.05$) and *TK+* ($p <$
0.05), and a significant down-regulation of *TK-* ($p < 0.05$) expression levels (Figure 2A). Regarding HPI
axis, in *rbfox1*[19del/19del] larvae we found a significant up-regulation of *crhb* ($p < 0.0001$) and a significant
up-regulation of *mr* ($p < 0.01$) (Figure 2C, D). We observed no significant changes in *gr* expression
levels ($p > 0.05$) (Figure 2C, D).

To assess *rbfox1* LoF effects on zebrafish larvae during early developmental stages, we performed
crhb, *bdnf* and *trkb2* qPCR experiments also in 3dpf larvae. Similarly to 5dpf larvae, in 3dpf
rbfox1[19del/19del] larvae we observed a significant up-regulation of *crhb* ($p < 0.01$), *bdnf* ($p < 0.05$) and
TK+ ($p < 0.05$), and a significant down-regulation of *TK-* ($p < 0.05$) (Figure 2B).

These findings showed that *rbfox1* LoF led to alterations in stress-related gene expression and
hyperactivation of the HPI axis, in line with our behavioural data and implicating *RBFOX1* as a critical
regulator of stress response mechanisms.

Treatment with TRKB agonist/antagonist restores HPI axis gene expression in *rbfox1* LoF larvae

To assess whether changes in HPI axis gene expression seen in *rbfox1* LoF larvae were mediated by
bdnf/trkb2 signalling, we examined the expression of HPI axis components (*crhb*, *mr*, *gr*) in 5dpf
zebrafish larvae following chronic exposure (from 5 h to 5dpf) to the TRKB agonist 7,8-DHF or
antagonist ANA-12.

In *rbfox1*[+/+] larvae, neither TRKB agonist 7,8-DHF or antagonist ANA-12 influenced HPI axis gene
expression ($p > 0.05$) (Figure 2C, D). In *rbfox1*[19del/19del] larvae, chronic exposure to either TRKB agonist
or antagonist significantly reduced *crhb* ($p_{rbfox1^{19del/19del} \text{ vs } rbfox1^{19del/19del} + ANA-12} < 0.0001$, $p_{rbfox1^{19del/19del} \text{ vs}}$
$_{rbfox1^{19del/19del} + 7,8-DHF} < 0.0001$) and *mr* ($p_{rbfox1^{19del/19del} \text{ vs } rbfox1^{19del/19del} + ANA-12} < 0.0001$, $p_{rbfox1^{19del/19del} \text{ vs}}$
$_{rbfox1^{19del/19del} + 7,8-DHF} < 0.05$) expression levels relative to untreated mutant larvae, restoring HPI axis
gene expression to levels comparable with wild type controls ($p > 0.05$) (Figure 2C, D). Following
identification and removal of outliers (1 outlier in the *rbfox1*[+/+] + ANA-12 group, 1 outlier in the
rbfox1[+/+] + 7,8-DHF group and 1 outlier in the *rbfox1*[19del/19del] + 7,8-DHF group), we observed a significant
down-regulation of *mr* expression levels in *rbfox1*[19del/19del] + ANA-12 versus *rbfox1*[+/+] un-exposed
controls ($p < 0.01$) (Figure 2C) and a significant up-regulation of *mr* expression levels in *rbfox1*[19del/19del]+
7,8-DHF versus both exposed ($p < 0.01$) and un-exposed ($p < 0.05$) controls (Figure 2D).

As the hypothalamus is the primary player in the stress response and Rbfox1 had only previously been
shown to affect TRKB expression within the hippocampus[16], we used ISH to determine whether *trkb2*
mRNA expression was reduced within the hypothalamic area. In agreement with previous findings[42],
in *rbfox1*[+/+] we found that *trkb2* is widely expressed in the brain of 5dpf larvae, whereas in
rbfox1[19del/19del] larvae, consistently with our qPCR experiments, we found a significant overall reduction
of *trkb2* mRNA in the whole brain (Figure 2E-H), including in the hypothalamus ($p < 0.01$) (Figure 2F, I,
K, Supplementary Figure 5).

Our findings showed that the dysregulation in HPI axis gene expression caused by *rbfox1* LoF was
prevented by TRKB agonists and antagonists suggesting the increased CRH expression is mediated by
an action of *RBFOX1* LoF on BDNF signalling.

***rbfox1* LoF alters *pac1a* expression levels of zebrafish larvae**

As RBFOX proteins have been shown to be able to regulate *Pac1* alternative splicing to include the *hop*
cassette[7] and PAC1 has been shown to regulate *Bdnf* transcription and potentiate TRKB activity[17], we
examined expression levels of *pac1a,* the zebrafish homologue of the mammalian *PAC1*, in *rbfox1*[+/+]
and *rbfox1*[19del/19del] 5dpf larvae by qPCR.

In mammals there are several PAC1 isoforms and their role in the regulation of stress is poorly
understood[9]. In the brain, the predominant isoforms are *PAC1-hop* (long isoform) and *PAC1-short*[9].
Zebrafish possess two *pac1* genes, *pac1a* and *pac1b*, but only *pac1a* contains the *hop* cassette[9]*. PAC1-
short* enhances *CRH* transcription, while *PAC1-hop* reduces *CRH* synthesis during late stress recovery
phase[4]. We measured expression levels of both *pac1a-short* and *-hop* isoforms and we observed no
differences in *pac1a-short* expression, but we found a significant up-regulation of *pac1a-hop* in
rbfox1[19del/19del] mutant larvae ($p < 0.05$) (Figure 2L).

This finding strengthens previous data[7] suggesting that RBFOX1 is not the main regulator, or at least
not the sole regulator of, *PAC1* alternative splicing and suggests that changes in *crhb* expression seen
here are the result of the altered *bdnf/trkb2* expression.

***rbfox1* mutants undergo adaptive mechanisms and allostatic overload during development**

The HPA axis possesses a vital role in the maintenance of allostasis, the process by which the body
achieves stability in response to stress or environmental challenges. Dysregulation of the HPA axis
often leads to disrupted allostasis during later life, also termed as allostatic load (i.e., the physiological
consequence resulting from the cumulative "wear and tear" of the body in response to chronic
stress)[22]. As we observed dysregulation of the HPI axis gene expression in *rbfox1* mutant larvae and
altered behavioural responses in *rbfox1* mutant larvae but not in adults, we assessed expression of
HPI axis components in adult zebrafish, in the presence and absence of acute stress (NTD), to explore
the possibility of adaptation that may contribute to differences in allostatic load.

We first measured *crhb*, *mr*, *gr*, *bdnf* and *trkb2* mRNA expression levels. In physiological resting
conditions, we observed no differences in HPI nor *bdnf/trkb2* expression levels ($p > 0.05$) (Figure 3A,
B). However, after stress exposure, regarding the HPI axis, in *rbfox1*[19del/19del] adults we observed the
same dysregulation seen in *rbfox1* LoF larvae: we found significant up-regulation of *crhb* ($p < 0.001$)
and *mr* ($p < 0.0001$) and no changes in *gr* expression levels ($p > 0.05$) (Figure 3A), and significant up-
regulation of *bdnf* ($p < 0.01$) and *TK+* ($p < 0.05$), and significant down-regulation of *TK-* ($p < 0.05$) (Figure
3B, C).

Excess glucocorticoid (GC) exposure during early developmental stages or early life stress (ELS) can
lead to vulnerability to allostatic overload in the long-term[19,22]. One of the effects caused by GC- or
ELS-induced allostatic overload is reduced cell proliferation during adulthood[19]. This effect has been

suggested to be developmentally dynamic, since it is often preceded by increased cell proliferation
during early stages[19]. Further, BDNF signalling has been shown to regulate neural stem cell
proliferation through TK-, suggesting that *rbfox1* LoF fish showing altered levels of *TK-* expression, may
show altered proliferation across the life course[43]. We therefore assessed the rate of proliferation at
both larval and adult stages using qPCR for proliferating cell nuclear antigen (*pcna*). We observed a
significant up-regulation of *pcna* expression levels in 5dpf *rbfox1*$^{19del/19del}$ zebrafish larvae ($p < 0.05$)
and a significant down-regulation of *pcna* expression levels in the brains of adult *rbfox1*$^{19del/19del}$ fish (p
< 0.01) when compared to *rbfox1*$^{+/+}$ siblings (Figure 3D).

In adult zebrafish, other effects caused by GC-induced allostatic overload include reduced fertility and
survival rates[19]. Therefore, we assessed fertility and survival of *rbfox1*$^{19del/19del}$ zebrafish and *rbfox1*$^{+/+}$
siblings. We found that both fertility ($p < 0.05$) and survival ($p < 0.001$) rates of *rbfox1*$^{19del/19del}$ fish were
significantly reduced compared to *rbfox1*$^{+/+}$ siblings (Figure 3E, F).

Consistent with our data showing hyperactivation of the stress response system, these findings
suggest that, during development, *rbfox1* mutants undergo adaptive changes like those caused by GC-
exposure, leading to altered development and lasting differences in response to environmental
stressors.

DISCUSSION

In this study we generated a CRISPR-Cas9 LoF *rbfox1* zebrafish line (*rbfox1*^{19del}) to investigate the
mechanisms by which *RBFOX1* LoF increases susceptibility to psychiatric disorders.
311
Allostatic load has been linked with several cognitive disorders including depression, schizophrenia,
anxiety and PTSD[44]. HPA axis overactivity is a key factor in the onset of the allostatic load[22]. RBFOX1
has been shown to influence expression of stress related genes such as PAC1 and BDNF/TRKB.
Therefore, alterations in RBFOX1 function may increase psychiatric disorders vulnerability through
alterations in stress response systems, such as the HPA axis, pre-disposing to allostatic overload
vulnerability in later life (Figure 4).
318
Similarly to *Rbfox1* deficient mice[6], we found that *rbfox1*$^{19del/19del}$ mutants were hyperactive and
impulsive. These findings are consistent with studies implicating common and rare *RBFOX1* genetic
variants as risk factors for psychiatric disorders like ASD, ADHD, schizophrenia and anxiety/stress
disorders[5,6,45]. At larval stages *rbfox1*$^{19del/19del}$ were hyperactive and had increased burst swimming,
indicative of increased impulsivity. In the response and habituation to acoustic startle assay,
rbfox1$^{19del/19del}$ larvae startled more and had a higher response rate over time compared to *rbfox1*$^{+/+}$
siblings. In a forced light-dark assay, although baseline hyperactivity of *rbfox1*$^{19del/19del}$ larvae makes
interpretation difficult, *rbfox1*$^{19del/19del}$ larvae showed an increased startle response upon dark to light
transition, consistent with heightened anxiety. These results suggest that *rbfox1* LoF leads to
heightened arousal, aligning with studies linking *RBFOX1* to mood and anxiety disorders, including
PTSD[46]. Interestingly, *rbfox1*$^{19del/19del}$ adults showed no significant difference in the novel tank diving
assay, an adult measure of stress-reactive behaviour, suggesting behavioural adaptation. In the 5-
CSRTT we found that *rbfox1*$^{19del/19del}$ were more impulsive than *rbfox1*$^{+/+}$ siblings, consistent with
RBFOX1 variants being associated with impulsivity and related disorders such as ADHD and ASD[6].
333
In *rbfox1* LoF larvae we observed dysregulation of the HPI axis, analogous to the HPA axis in mammals.
The HPA axis is crucial for managing stress, with CRH triggering cortisol secretion, affecting
metabolism, immunity, and behaviour[47,48]. In zebrafish larvae under resting physiological conditions,
we found increased *crhb* and *mr*, with no changes in *gr* levels. In adults, although resting levels were
unchanged, *rbfox1* LoF showed an exaggerated molecular response to stress. Elevated *crhb* levels as
seen in *rbfox1*$^{19del/19del}$ larvae suggest a chronic anxiety state. Increased CRH and MR levels are linked

to mood disorders like depression and PSTD[49–52]. These findings support a role for *RBFOX1* in *CRH*
regulation and suggest that *RBFOX1* LoF may contribute to mood disorders via HPA axis
hyperactivation and disrupted stress resilience.

BDNF and TRKB are also key regulators of the stress response[53]. BDNF levels often increase during
stress to promote neuronal survival and plasticity, buffering negative effects of stress on the brain via
TRKB[10,11,13,53–55]. Dysfunction of BDNF/TRKB signalling is linked to several stress-related disorders[13].
The increased *bdnf* expression we observed in *rbfox1^{19del/19del}* larvae suggests a compensatory
response to cope with *crhb* increase and maintain homeostasis. At 3 and 5dpf, in *rbfox1^{+/+}* larvae we
detected both *trkb2* truncated (*TK-*) and full-length (*TK+*) forms, and in *rbfox1^{19del/19del}* larvae we
observed a significant down-regulation of *TK-* and a significant up-regulation of *TK+*, which is predicted
to lead to an increased activation of TRKB signalling, consistent with the increased *crhb* expression as
seen here[56]. Studies in rodents showed that *Rbfox1* up-regulation led to increased TrkB.T1 (*TK-*)
expression in the hippocampus and no changes were observed following *Rbfox1* deletion[16]. In
contrast, we observed down-regulation of *trkb2 TK-* expression in *rbfox1^{19del/19del}* larvae in resting
physiological conditions. The difference between these studies may be due to i) differences in the
tissue examined (zebrafish larvae whole head versus rodent hippocampus), ii) the developmental
stage of the animals (larval versus adult stage), iii) the genotype of the animals studied (*rbfox1^{19del/19del}*
in our study versus *Rbfox1^{+/-}* in the study in rodents), or iv) presence of compensatory mechanisms (in
the hippocampal mouse model *Rbfox1* knockdown resulted in up-regulation of *Rbfox2*). As for this
latter hypothesis, as in previous studies in zebrafish[2], we did not see any changes in *rbfox2* expression
upon *rbfox1* LoF (Supplementary Figure4) suggesting that this compensatory mechanism does not
occur in larval fish.

Consistent with our hypothesis that *RBFOX1* LoF leads to hyperactivation of the HPA axis through the
BDNF/TRKB pathway, chronic exposure to TRKB agonist (7,8-DHF) or antagonist (ANA-12) restored HPI
axis gene expression to that seen in *rbfox1^{+/+}* larvae. ANA-12 is a selective non-competitive antagonist
of TRKB exerting central TRKB blockade and producing rapid and long-lasting anxiolytic and
antidepressant effects[57]. Mice treated with ANA-12 showed reduced anxiety-like behaviour[57] and in
stressed rats, ANA-12 blocked *Crh* increase in the hypothalamus and amygdala[58]. 7,8-DHF is a potent
selective agonist of TRKB used in the treatment of several disorders, including depression and
schizophrenia, and has been shown to enhance memory consolidation and emotional learning in
healthy rodents[59]. In a similar fashion to the effects of ANA-12 seen in rats (blocking *Crh* increase)[57],
after chronic treatment with ANA-12, we observed that the expression of HPI axis genes in
rbfox1^{19del/19del} larvae was restored to *rbfox1^{+/+}* levels. Contrary to expectation, a similar result was
seen in the presence of chronic agonist exposure. However, chronic exposure to agonists may lead to
desensitization and inhibition of TRKB signalling[60]. Interestingly, *rbfox1^{19del/19del}* larvae were
differentially sensitive to the effect of TRKB drugs on *mr* expression, most likely due to the unbalanced
TK+/TK- levels. These findings warrant further investigations. It is of note that neither TRKB agonist
nor antagonist had a significant effect on gene expression in *rbfox1^{+/+}* larvae. One possible explanation
is that, as the larvae were not exposed to any stressors, any effect on Trkb2 signalling had limited
effect on HPI axis activity.

In *rbfox1^{19del/19del}* larvae, we also observed up-regulation of *pac1a-hop* with no changes in *pac1a-short*
levels. Previous studies in zebrafish have shown that Pac1-short gain of function caused persistent *crh*
increase, while overexpression of Pac1-hop prevented stress-induced *crh* transcription activation[4].
These previous studies also suggest that RBFOX1 regulates the alternative splicing of *PAC1* promoting
the formation of *PAC1-hop*[4]. However, the increase in *pac1-hop* seen in our study argues against this
suggestion and suggests an adaptive response to the increased expression of *crh* either through
rbfox2[7] (whose expression was not altered by *rbfox1* LoF (Supplementary Figure4)) or through an
unknown mechanism.

In contrast to larvae, differences in HPI axis and *bdnf/trkb2* gene expression were not seen in adult
animals under resting conditions. However, despite no difference in stress-reactive behaviour as
determined by the novel tank diving assay, *rbfox1[19del/19del]* adults showed exaggerated HPI activation
on challenge with a stressor, aligning with the previous suggestion that Rbfox1 is required for the
termination of the acute endocrine stress response[4] and that chronic stress during development leads
to adult HPI axis adaptation, often at a cost in later life (e.g., reduced fertility and survival)[19,38]. This
latter was evidenced by reduced fertility and survival rates in *rbfox1[19del/19del]* adults. Another long-term
effect of excess GC-exposure/early life stress is reduced neural stem cell proliferation in adult animals,
often accompanied by increased proliferation at early stages[19]. Our findings that *rbfox1[19del/19del]*
showed increased *pcna* levels at larval staged but reduced *pcna* levels during adulthood, corroborate
this latter hypothesis of a dynamic developmental effect of early life stress. Consistent with the *crhb*
increase observed in stressed *rbfox1[19del/19del]* adults, and as seen in larvae, after acute stress,
rbfox1[19del/19del] adults also showed unbalanced *trkb2* expression, with up-regulation of *TK+* and down-
regulation of *TK-* .

In conclusion, given the conservation between fish HPI axis and the mammalian HPA axis, our data
unveils a pivotal new finding in *CRH* regulation, revealing an interplay between RBFOX1 and
BDNF/TRKB in the context of chronic stress and stress resilience and suggests that RBFOX1 plays a
crucial role in adaptive stress mechanisms. In response to stressful challenges, RBFOX1 functions as a
"switch" regulating the balance between short/long isoforms of TRKB receptors: during the stress
recovery phase, RBFOX1 switches the balance towards *TRKB.T1/TK-* isoform, decreasing *CRH* levels
and turning off the HPA axis (Figure 4). Furthermore, as loss of *rbfox1* in zebrafish led to behavioral
phenotypes resembling those associated with human psychiatric diseases (hyperactivity, impulsivity,
reduced habituation and heightened arousal), our findings also suggest that *RBFOX1* contributes to
the liability of psychiatric disorders through HPA axis hyperactivation and BDNF/TRKB dysregulation,
which leads to disrupted development and vulnerability to allostatic overload in later life. The suite of
behavioural effects seen in *rbfox1* LoF animals coupled with exaggerated HPI axis response on
exposure to a stressor are reminiscent of reported traits in passive versus active copers[62] suggesting
that RBFOX1 activity may also play a key role in individual differences in coping styles. Although we
show a key role for RBFOX1 in regulation of the HPA axis consistent with this being a primary
mechanism by which *RBFOX1* variants pre-disposes to psychiatric disorders, RBFOX1 regulates the
splicing of a wide range of additional genes[15] which may also play a part of disease vulnerability.

METHODS

Animal maintenance

All fish were maintained in a recirculating system (Tecniplast, UK) with a 14h:10h light/dark cycle and
a constant temperature of 28°C. Fish were fed with ZM-400 fry food (Zebrafish Management Ltd.) in
the morning and brine shrimps in the afternoon. Breeding was set up in the evening, in sloping
breeding tanks (Tecniplast) provided with dividers for timed mating. The following morning, dividers
were removed to allow spawning. Eggs were collected in Petri dishes (max 50 eggs/dish). Infertile eggs
were removed, and fertile ones were incubated at 28°C. Petri dishes were checked daily to ensure
consistent developmental stage across groups. If reared, larvae were moved to the recirculating
system at 5 days post fertilization (dpf) and fed as stated above.

All procedures were carried out under license in accordance with the Animals (Scientific Procedures)
Act, 1986 and under guidance from the Local Animal Welfare and Ethical Review Board at Queen Mary
University of London.

Generation of *rbfox1* loss of function zebrafish line

The zebrafish *rbfox1* loss of function (LoF) mutant line was generated as described previously using Tübingen strain as background [63]. CRISPR RNA (crRNA) (Merck) was designed to target *rbfox1* exon 2 (**CCC**AGTTCGCTCCCCCTCAGAAC, PAM sequence in bold, MwoI recognition site underlined). A 3 µL injection mix containing 1 µL (F_C 83 ng/µL) crRNA, 1 µL (F_C 83 ng/µL) tracrRNA (Merck, #TRACRRNA05N), 1 µL (F_C 1.67 µM) and 1 µL Cas9 protein (New England Biolabs, #M0646) was freshly prepared on the morning of the injection procedure. Then, 1 nL of the injection mix was injected into one-cell stage zebrafish embryos (~ 100-150 embryos). Injection efficacy was assessed at 24 hours post fertilization (hpf) by polymerase chain reaction (PCR) from genomic DNA (*rbfox1*_Forward, 5'-TAATCAAGACGCCCCAGCAC–3'; *rbfox1*_Reverse, 5'- GTACTCAGCAGGAATGCCGT- 3') followed by MwoI (New England Biolabs, #R0573S) restriction enzyme digestion. Successful injections will introduce indel mutations disrupting the recognition site of the restriction enzyme, preventing this latter from cutting the PCR amplicon. Once reached sexual maturity (at ~ 3 months of age) injected fish (F_0) were outcrossed with wild type to generate F_1 embryos. The progeny may carry different mutations due to the mosaic nature of the F_0 parents. F_1 fish were therefore screened for mutations leading to premature termination codon (PTC) via cloning into pGEM-T Easy vector (Promega, #A1360), followed by transformation, colony PCR, DNA purification (Monarch® Plasmid Miniprep Kit, New England Biolabs, #T1010) and sequencing (Source BioScience PLC). Quantitative real time PCR (qPCR) was used to confirm reduction of *rbfox1* mRNA expression.

Fish breeding and genotyping

For adult experiments, mixed sexes were used. When possible, animals were associated with an identification number and genotype was assigned after data analysis. For larval experiments, fish were generated by heterozygous in-cross and genotyped prior to data analysis. Genomic DNA was extracted from fins and using the HotSHOT method. Briefly, samples were incubated at 95 °C in 50 mM NaOH for 30 min, followed by 1 min at 10 °C. Reaction was stopped using 1M Tris HCl (1/10 of the initial NaOH volume), pH 8.00. Genotyping primers were the same as the ones used to identify founder carriers.

Drug Treatment

Zebrafish embryos of each genotype (*rbfox1^{+/+}* and *rbfox1^{19del/19del}*) were treated from 5hpf to 5dpf with 20 µM TRKB antagonist ANA-12 (abcam, Cat.ab146196), or 20 µM TRKB agonist 7,8-dihydroxyflavone (7,8-DHF) (abcam #ab120996), or dimethylsulfoxide (DMSO, vehicle 0.01%) (Merck #34869). Treatment was performed in Petri dishes (max 30 embryos x dish) and drugs or vehicle were dissolved in fish water. Solutions were replaced daily.

Quantitative Real-Time PCR

Quantitative Real-time PCR (qPCR) of target RNA was performed on 5dpf *rbfox1* wild type (*rbfox1^{+/+}*) and *rbfox1* homozygous (*rbfox1^{19del/19del}*) larvae, using Luna® Universal One-Step RT-qPCR Kit (New England Biolabs #M3005) and a Bio-Rad 96-well qPCR machine (CFX96 Touch Real-Time PCR Detection System). Total RNA was isolated using TRIzol reagent (Thermo Fisher Scientific) following manufacturer's instructions. Briefly, after homogenization, RNA was isolated by precipitation, rinsed in ethanol and resuspended in RNase free water. Total RNA was then quantified using BioDrop

(Biochrom Ltd.), and up to 1 µg was reverse transcribed to cDNA using the ProtoScript II First Strand
cDNA Synthesis Kit (New England Biolabs, #E6560) following manufacturer's instructions. The resulting
cDNA yield and quality were evaluated using BioDrop (Biochrom Ltd.). As for zebrafish *trkb2 TK+* and
TK- isoforms quantification, Ct values resulting from the amplification of the *TK+* specific product were
subtracted from the Ct values resulting from the amplification of the *TK+/TK-* common product. All
reactions included 5 biological replicates and 3 technical replicates. For experiments in larvae, each
biological replicate consisted of 15 larval heads (eyes and jaw removed). For experiments in adults,
each biological replicate consisted of a single brain. Actin – β 2 (*actb2*) and ribosomal protein L13a
(*rpl13a*) were employed as reference genes. For assessing *rbfox1* mRNA expression, *rbfox1* primers
were designed upstream of the CRISPR deletion. Accession numbers, primer sequences and
amplification efficiencies for all the reference and target genes can be found in Supplementary Table
1.

***In situ* hybridization**

In situ hybridization (ISH) was carried out on whole mount zebrafish larvae and on larval (sagittal) and
adult brain (transverse) sections as described previously[2]. For *rbfox1* ISH, the original plasmid used to
generate the riboprobe was provided by Dr William Norton (University of Leicester). The plasmid used
to produce the *trkb2* riboprobe was generated in our laboratory (Forward primer: 5'-
GTTCGTGGAATGGCTTGCTG-3', Reverse primer: 5'-TCTGGCCCACGATGTTTTCA-3') using the pGEM®-T
Easy Vector System (Promega, #A1360) and *in-house* generated *E. coli* DH5α competent cells.
Riboprobes to identify *rbfox1* (NM_001005596) and *trkb2* (NM 01197161.2) mRNA were synthetized
by *in vitro* transcription (IVT) using MAXIscript™ SP6/T7 kit (Invitrogen by Thermo Fisher Scientific,
#AM1322), following manufacturer's instructions and using a DIG RNA Labeling Mix, 10× conc (Roche,
#11277073910) containing digoxigenin labeled uracil.

In situ hybridization on whole mount zebrafish larvae

In situ hybridization was carried out on 28hpf, 2- 3- 4- and 5dpf *rbfox1*$^{+/+}$ and *rbfox1*$^{19del/19del}$ larvae. To
prevent skin pigmentation, embryos were incubated in 0.2 mM 1-phenyl 2-thiourea (PTU) (Sigma,
#S527335) from 24hpf. When they reached the desired age, larvae were fixed in 4% paraformaldehyde
(PFA) (Merck, #158127) overnight (ON) at 4°C. The following day, larvae were rinsed in 1x phosphate
buffered saline (PBS) (Thermo Fisher Scientific, #18912014) supplemented with Tween 20 (Sigma,
#P1379) (0.05% v/v), dehydrated in ascending methanol series (25%, 50%, 70%, 80%, 90%, 100%
methanol, 5 min each) and stored in 100% methanol at -20 °C. To perform ISH experiments, larvae
were rehydrated in descending methanol series (100%, 90%, 80%, 70%, 50%, 25% methanol), 5 min
each, and washed in 1xPBS, 5 min. Larvae were permeabilized using proteinase K (PK) (ITW Reagents,
#A3830) (stock 20 µg/mL in 1xPBS) as follows: 28hpf larvae were permeabilized in PK 1:2000 in 1xPBS
for 20 min at room temperature (RT), older stages were permeabilized in PK 1:1000 at 37°C for at least
30 min. Then, larvae were post fixed in 4% PFA for 20 min and washed in 1xPBS at RT, 5 x 5 min.
Prehybridization was carried out in hybridization solution (HB) containing 50% formamide, 5% saline
sodium citrate buffer (SSC), 50 µg/mL heparin, 0.05 mg/mL yeast RNA, 0.1% Tween 20, and 0.92%
citric acid at 68 °C for 2h. Thereafter, larvae were incubated in HB containing *rbfox1* riboprobe (500
pg/µL), ON at 68°C. Post hybridization washes were performed at 68°C with a gradient of 2xSSC and
formamide (50%, 25% and 0% formamide), 10 min each, and then twice with 0.02xSSC, 30 min each.
Subsequently, larvae were blocked in blocking solution (BS) containing 10% normal sheep serum
(Gibco, #16070096) and 2 µg/µL bovine serum albumin, for 1h at RT. After blocking step, larvae were
incubated in anti-digoxigenin Fab fragments conjugated with alkaline phosphatase (Roche,

#11093274910), 1:2000 in BS, 1h at RT and then ON at 4°C. The following day, larvae were washed in
1xPBS, 6 x 15 min each, and then in alkaline phosphatase (100 mM NaCl, 100 mM Tris HCl, 50 mM
MgCl2, 0.1% Tween 20) (NTMT) buffer, 3 x 5 min each. The chromogenic reaction was carried out by
incubating the larvae in BCIP/NBT solution (Merck, #203790) in NTMT buffer, at RT in the dark, and
were observed every 20 min until the signal detection. After reaching the desired staining, larvae were
washed in 1xPBS at RT, post fixed in 4% PFA for 2h, cleared and stored in 80% glycerol at 4°C. For
sagittal cryosections, larvae were embedded in 1.5% low gelling temperature agarose (Scientific
Laboratories Supplies, #A9414) supplemented with 5% Sucrose in 1xPBS. Sections (5 μm) were
collected on adhesive microscope slides Superfrost® Plus Gold (Epredia).

In situ hybridization on sections

ISH was conducted on paraffin embedded *rbfox1*$^{+/+}$ and *rbfox1*$^{19del/19del}$ adult brains (transverse
sections) and 5dpf *rbfox1*$^{+/+}$ and *rbfox1*$^{19del/19del}$ zebrafish larvae (sagittal sections). Fish were culled by
an overdose of tricaine prior to head removal. Dissected adult brains and larvae were fixed in 4% PFA
in 1xPBS, ON at 4°C. Tissues were then rinsed in 1xPBS and dehydrated in ascending ethanol series (15
min in each of 30%, 50%, 70%, 80%, 90%, 100% ethanol) and embedded in paraffin. Transverse (adult
brains) or sagittal (5 days old larvae) sections of 12 μm (adult brains) or 7 μm (5 days old larvae)
thickness were cut using a microtome (Leica). To perform ISH, slides were de-waxed in xylene (twice,
10 min each), rehydrated in descending ethanol series (2 x 5 min in absolute ethanol, then 90%, 80%,
70%, 50% and 25% ethanol, 5 min each), and rinsed in 1xPBS for 5 min. Then, sections were
permeabilized using PK (0.05 μg/μL) for 8 min at RT, washed with 2 mg/mL glycine twice (5 min each),
post fixed in 4% PFA for 20 min and washed in 1xPBS at RT. Prehybridization was carried out in HB, for
1h at 68°C. Thereafter, sections were incubated in HB containing *rbfox1* riboprobe (500 pg/μL), ON at
68°C. Post hybridization washes were performed at 68°C twice for 20 min in 1xSSC, twice for 20 min
in 0.2xSSC, and several washes were performed in 1xPBS, 5 min each at RT. Then, sections were
blocked in BS for 30 min at RT and incubated in a 1:2000 dilution of anti-digoxigenin Fab fragments
conjugated with alkaline phosphatase in BS, ON at 4°C. The following day, sections were washed in
1xPBS, 5 x 10 min each. The chromogenic reaction was carried out by incubating the slides in BCIP/
NBT solution in NTMT buffer, at RT in the dark, and were observed every 20 min until the signal
detection. When the desired staining was obtained, sections were washed in 1xPBS at RT, dehydrated
in ascending ethanol series (25%, 50%, 70%, 80%, 90%, 100% ethanol, 5 min each), cleared in xylene
(twice, 5 min each) and mounted with dibutyl phthalate polystyrene xylene mounting medium (Sigma,
#06522).

Image acquisition and processing

Pictures of whole mount ISH on zebrafish larvae were acquired by Leica MZ75 microscope. For ISH on
sections, pictures were acquired using a Leica DMRA2 upright epifluorescent microscope with colour
QIClick camera (Leica) and processed with Velocity 6.3.1 software (Quorum Technologies Inc).
Quantification of the ISH staining signal intensity was performed as described previously[64], using Fiji
software[65]. Adult anatomical structures were identified according to the Neuroanatomy of the
Zebrafish Brain by Wullimann [66].

Behavioral Assays

For larvae, all behavioural experiments were conducted on the progeny of a *rbfox1*$^{+/19del}$ in-cross.
Larvae were genotyped prior to data analysis. For adults, were the genotype of the animals was
known, fish were pseudorandomised across testing systems with all trials having an approx. equal

number of each genotype. The adult fish were weight- and age-matched, with approximately equal
numbers of both sexes.

Larval behavioural experiments

Patterns of locomotor activity of 5dpf *rbfox1+/+*, *rbfox1+/19del* and *rbfox119del/19del* mutant zebrafish larvae
were investigated as described previously[23,28,67]. Tests were conducted between 9 a.m. and 4 p.m. At
5 dpf, larvae were placed in individual wells of a 24-well plate. To reduce stress due to manipulation,
larvae were acclimatised for at least 1 h before testing. Then, plates were placed into the DanioVision
observation chamber (Noldus). Locomotion parameters such as distance travelled and swimming
velocity were recorded using EthoVision XT software (Noldus). Data were exported in 1 min and 1 sec
time bins and analysed with R programming language[68]. For the hyperactivity assay, larval basal
locomotion was tracked for 15 min in dark conditions. Larval swimming burst was assessed as
described previously[23] and peaks were considered as the acceleration events when larvae travelled >
5 mm in < 12 sec. The forced light-dark transition assay was performed as described previously[69,70]
with modifications: after an initial 5 min period of dark (baseline), larvae were exposed to one
light/dark cycle of 1 min light (Noldus white light device) followed by 5 min dark. The response and
habituation to acoustic startle stimuli was performed as described previously[28]: after 10 min of
baseline (no stimuli, dark conditions), larvae were subjected to 10 sound/vibration stimuli (Noldus
tapping device) over 20 sec (2 sec intervals between each stimulus).

5-Choice Serial Reaction Time Task

As deficit in impulse control and learning are associated with many psychiatric diseases including
ADHD and bipolar disorder, we measured impulsive action using a zebrafish version of the 5-Choice
Serial Reaction Time Task (5-CSRTT)[24]. Adult *rbfox1+/+*, *rbfox1+/19del* and *rbfox119del/19del* zebrafish, 7
months old, mixed sexes, were singly housed for a week prior to experiment and remained singly
housed for the whole duration of the assay (9 weeks). Fish were tested using the Zantiks AD units.
Each unit was provided with a small tank with five apertures and a food hopper insert. The five
apertures created five different entry points for the fish, acting like the five nose poke holes of the
rodent version of the assay. The food hopper was placed at the opposite side of the five apertures and
formed an area for the fish to enter and collect food reward. Below the testing tank there was an
integrated screen, used to display white light (stimulus) into the five apertures. Responses were
detected when a fish entered these apertures and recorded with an integrated camera placed at the
top of the tank. The experiment consisted of five training stages: i) habituation, ii) initiator training,
iii) stimulus light training, iv) 5-CSRTT/no delay, v) 5-CSRTT/variable-delay. Details of each stage are
provided in Supplementary Table 2.

Novel tank diving

Novel tank diving is a behavioural test to assess anxiety-like behaviour in adult fish. Response to novel
tank was assessed in *rbfox1+/+*, *rbfox1+/19del* and *rbfox119del/19del* 9 months old zebrafish, mixed-sexes, as
described previously[67]. Fish were singly housed for a week prior to performing the experiment and
acclimatized for at least 1 h in the behavioral room on the testing day. Behavioural assays were
conducted between 9 a.m. and 2 p.m. During the test, fish were individually placed into a 1.5 L tank
and their behaviour was tracked and recorded using EthoVision system (Noldus). Data were exported
in 1 min time bin and analysed as previously described[67]. Experimental groups were randomised
during testing. We analyzed three behaviours in response to the novel tank: i) time spent in the bottom

of the tank, ii) total distance traveled, and iii) number of transitions to the top–bottom area of the
tank.

Statistical analysis

For qPCR, relative mRNA expressions were calculated using the Pfaffl method[71]. Outliers were
identified and removed using Dixon's test (α = 0.05)[72]. Differences in gene expression were assessed
using a one-way ANOVA followed by Tukey's post-hoc test using GraphPad Software (Prism). For
behavioural analysis, all data were analysed with R programming language[68]. Scripts used for analysis
are available on GitHub (https://github.com/AdeleLeg). For models where distance moved, distance
to zone, velocity, or top halves visits were the response variable, we fitted data to mixed linear models
using the "*lme4*" package, and where proportion of responders or proportion of time spent in the
bottom third were our response variable, we fitted data to beta regression models using the "*betareg*"
package. In all instances, for all experiments, we used genotype as fixed effect, and fish ID and day of
experiment as random effects. In the response and habituation to acoustic startle, we used also the
stimulus number as fixed effect. As in García-González et al.[67], we reported significant fixed effects as
Type II Wald χ^2 from models using the package "*car*," post hoc Tukey's tests were also conducted as
necessary with the package "*emmeans*". For the 5-CSRTT, overall correct responses (learning) and
anticipatory responses (impulsivity) were assessed using the formulas in Supplementary Table 2.
Statistical analysis of mean intensity in *in situ* hybridization images of zebrafish larval sections was
performed using ImageJ. Mean intensity values were quantified across specified regions of interest,
indicated in Supplementary Figure 5, ensuring that comparable areas of similar dimensions were used
for each sample to calculate differences in expression. The resulting values were compared across
groups using a paired t-test to determine significant differences using GraphPad Software (Prism). For
all experiments, sample size needed to achieve adequate statistical power for detecting a significant
effect was determined based on data from previous research or pilot studies. Accepted α level and
power were respectively ≤ 0.05 and ≥ 0.80.

Fertilisation and survival analysis

Fertilisation and survival rates were measured as described previously[19]. For fertilization rate we
measured the percentage of fertilized eggs when pairing *rbfox1[19del/19del]* and compared it with
fertilization rate of *rbfox1[+/+]* siblings. Data were collected upon 3 different mating trials, each trial
comprising 3-5 trios (1 male and 2 females). Trials were performed on different weeks, using different
trios for each test. Eggs were collected in different Petri dishes, properly labelled to distinguish
between trios. Fertilisation rate was determined by averaging across Petri dishes from the same trio
and calculated at 6-8hpf. For survival analysis, we measured the percentage of surviving animals
(across 3 tanks for each genotype for each batch, to avoid tank effect) from when larvae were added
to the nursery (at 5dpf) until 2 months of age, when fish were transferred into adult aquarium.
656

CONTRIBUTIONS

AL conceived the project, designed and conducted the experiments, analysed the data, and wrote the
manuscript. JGG designed and performed the CRISPR/Cas9 experiment generating the line. SH, PA, SA,
XW and WH contributed to the experiments. NFC and BC edited the manuscript. CHB directed the
study, designed the experiments, edited the manuscript, and secured funding. All authors contributed
to the article and approved the submitted version.

ACKNOWLEDGMENTS

CHB, AL, SH, PA and WH were supported by the NIH grant U01 DA044400-03. JGG was supported by
a 2022 NARSAD Young Investigator Grant (Number #30749) by the Brain & Behavior Research
Foundation. NFC and BC were supported by the grants PID2021-127776OB-I100, 202218 31 and 2021-
SGR-01093 from the Ministry of Science, Innovation and Universities (Spain), Fundació La Marató de
TV3, and AGAUR-Generalitat de Catalunya, respectively, and by ICREA Academia 2021. Authors would
like to acknowledge SciDraw (https://scidraw.io/) for images used to create the graphical abstract.

CONFLICT OF INTEREST
The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.
673

REFERENCES

1. Cross-Disorder Group of the Psychiatric Genomics Consortium. Electronic address:

plee0@mgh.harvard.edu & Cross-Disorder Group of the Psychiatric Genomics Consortium.

Genomic Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric

Disorders. *Cell* **179**, 1469-1482.e11 (2019).

2. Antón-Galindo, E. *et al.* Pleiotropic contribution of rbfox1 to psychiatric and

neurodevelopmental phenotypes in two zebrafish models. *Transl. Psychiatry* **14**, 1–11 (2024).

3. Thisse, B. & Thisse, C. ZFIN Publication: Thisse et al., 2004. *ZFIN* http://zfin.org/ZDB-PUB-

040907-1.

4. Amir-Zilberstein, L. *et al.* Homeodomain Protein Otp and Activity-Dependent Splicing Modulate

Neuronal Adaptation to Stress. *Neuron* **73**, 279–291 (2012).

5. Abi-Dargham, A. *et al.* Candidate biomarkers in psychiatric disorders: state of the field. *World*

Psychiatry **22**, 236–262 (2023).

6. O'Leary, A. *et al.* Behavioural and functional evidence revealing the role of RBFOX1 variation in

multiple psychiatric disorders and traits. *Mol. Psychiatry* **27**, 4464–4473 (2022).

7. Lee, J.-A., Tang, Z.-Z. & Black, D. L. An inducible change in Fox-1/A2BP1 splicing modulates the

alternative splicing of downstream neuronal target exons. *Genes Dev.* **23**, 2284–2293 (2009).

8.   Shaughnessy, C. A., Myhre, V. D., Hall, D. J., McCormick, S. D. & Dores, R. M. Hypothalamus-
pituitary-interrenal (HPI) axis signaling in Atlantic sturgeon (Acipenser oxyrinchus) and sterlet
(Acipenser ruthenus). *Gen. Comp. Endocrinol.* **339**, 114290 (2023).

9.   Biran, J. *et al.* Splice-specific deficiency of the PTSD-associated gene PAC1 leads to a paradoxical
age-dependent stress behavior. *Sci. Rep.* **10**, 9559 (2020).

10. Linz, R. *et al.* Acute psychosocial stress increases serum BDNF levels: an antagonistic relation to
cortisol but no group differences after mental training. *Neuropsychopharmacology* **44**, 1797–
1804 (2019).

11. Jeanneteau, F. D. *et al.* BDNF and glucocorticoids regulate corticotrophin-releasing hormone
(CRH) homeostasis in the hypothalamus. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 1305–1310 (2012).

12. Karatsoreos, I. N. & McEwen, B. S. Psychobiological allostasis: resistance, resilience and
vulnerability. *Trends Cogn. Sci.* **15**, 576–584 (2011).

13. Miao, Z., Wang, Y. & Sun, Z. The Relationships Between Stress, Mental Disorders, and Epigenetic
Regulation of BDNF. *Int. J. Mol. Sci.* **21**, 1375 (2020).

14. Suliman, S., Hemmings, S. & Seedat, S. Brain-Derived Neurotrophic Factor (BDNF) protein levels
in anxiety disorders: systematic review and meta-regression analysis. *Front. Integr. Neurosci.* **7**,
55 (2013).

15. Fogel, B. L. *et al.* RBFOX1 regulates both splicing and transcriptional networks in human
neuronal development. *Hum. Mol. Genet.* **21**, 4171–4186 (2012).

16. Tomassoni-Ardori, F. *et al.* Rbfox1 up-regulation impairs BDNF-dependent hippocampal LTP by
dysregulating TrkB isoform expression levels. *eLife* **8**, e49673 (2019).

17. Solés-Tarrés, I., Cabezas-Llobet, N., Vaudry, D. & Xifró, X. Protective Effects of Pituitary
Adenylate Cyclase-Activating Polypeptide and Vasoactive Intestinal Peptide Against Cognitive
Decline in Neurodegenerative Diseases. *Front. Cell. Neurosci.* **14**, (2020).

18. Mcewen, B. S. Protection and Damage from Acute and Chronic Stress: Allostasis and Allostatic

Overload and Relevance to the Pathophysiology of Psychiatric Disorders. *Ann. N. Y. Acad. Sci.*

1032, 1–7 (2004).

19. Eachus, H. *et al.* Elevated glucocorticoid alters the developmental dynamics of hypothalamic

neurogenesis in zebrafish. *Commun. Biol.* **7**, 1–14 (2024).

20. Danese, A. & McEwen, B. S. Adverse childhood experiences, allostasis, allostatic load, and age-

related disease. *Physiol. Behav.* **106**, 29–39 (2012).

21. Bornstein, S. R. *et al.* Stress-inducible-stem cells: a new view on endocrine, metabolic and

mental disease? *Mol. Psychiatry* **24**, 2–9 (2019).

22. Kinlein, S. A., Wilson, C. D. & Karatsoreos, I. N. Dysregulated Hypothalamic–Pituitary–Adrenal

Axis Function Contributes to Altered Endocrine and Neurobehavioral Responses to Acute Stress.

Front. Psychiatry **6**, (2015).

23. Lange, M. *et al.* The ADHD-susceptibility gene lphn3.1 modulates dopaminergic neuron

formation and locomotor activity during zebrafish development. *Mol. Psychiatry* **17**, 946–954

(2012).

24. Parker, M. O. *et al.* Development and automation of a test of impulse control in zebrafish. *Front.*

Syst. Neurosci. **7**, (2013).

25. Halberstadt, A. L. & Geyer, M. A. Habituation and sensitization of acoustic startle: opposite

influences of dopamine D1 and D2-family receptors. *Neurobiol. Learn. Mem.* **92**, 243–248

(2009).

26. Burgess, H. A. & Granato, M. Sensorimotor gating in larval zebrafish. *J. Neurosci. Off. J. Soc.*

Neurosci. **27**, 4984–4994 (2007).

27. Cohen, H. *et al.* Blunted HPA axis response to stress influences susceptibility to posttraumatic

stress response in rats. *Biol. Psychiatry* **59**, 1208–1218 (2006).

28. Leggieri, A. *et al.* Ankk1 Loss of Function Disrupts Dopaminergic Pathways in Zebrafish. *Front.*

Neurosci. **16**, (2022).

29. Davies, M. N. *et al.* Generalised Anxiety Disorder – A Twin Study of Genetic Architecture,

Genome-Wide Association and Differential Gene Expression. *PLOS ONE* **10**, e0134865 (2015).

30. Basnet, R. M., Zizioli, D., Taweedet, S., Finazzi, D. & Memo, M. Zebrafish Larvae as a Behavioral

Model in Neuropharmacology. *Biomedicines* **7**, E23 (2019).

31. Lee, H. B. *et al.* Novel zebrafish behavioral assay to identify modifiers of the rapid, nongenomic

stress response. *Genes Brain Behav.* **18**, e12549 (2019).

32. Hillman, C., Kearn, J. & Parker, M. O. A unified approach to investigating 4 dpf zebrafish larval

behaviour through a standardised light/dark assay. *Prog. Neuropsychopharmacol. Biol.*

Psychiatry **134**, 111084 (2024).

33. Baiamonte, M., Parker, M. O., Vinson, G. P. & Brennan, C. H. Sustained Effects of Developmental

Exposure to Ethanol on Zebrafish Anxiety-Like Behaviour. *PLoS ONE* **11**, e0148425 (2016).

34. Gene Detail :: Allen Brain Atlas: Mouse Brain. https://mouse.brain-map.org/gene/show/92602.

35. Brain tissue expression of RBFOX1 - Summary - The Human Protein Atlas.

https://www.proteinatlas.org/ENSG00000078328-RBFOX1/brain.

36. Grone, B. P. & Maruska, K. P. Divergent evolution of two corticotropin-releasing hormone (CRH)

genes in teleost fishes. *Front. Neurosci.* **9**, 365 (2015).

37. Sahu, M. P. *et al.* Neurotrophin receptor Ntrk2b function in the maintenance of dopamine and

serotonin neurons in zebrafish. *Sci. Rep.* **9**, 2036 (2019).

38. Eachus, H., Ryu, S., Placzek, M. & Wood, J. Zebrafish as a model to investigate the CRH axis and

interactions with DISC1. *Curr. Opin. Endocr. Metab. Res.* **26**, 100383 (2022).

39. Faught, E. & Vijayan, M. M. The mineralocorticoid receptor is essential for stress axis regulation

in zebrafish larvae. *Sci. Rep.* **8**, 18081 (2018).

40. Tessarollo, L. & Yanpallewar, S. TrkB Truncated Isoform Receptors as Transducers and

Determinants of BDNF Functions. *Front. Neurosci.* **16**, (2022).

41. Martin, S. C., Marazzi, G., Sandell, J. H. & Heinrich, G. Five Trk Receptors in the Zebrafish. *Dev.*

Biol. **169**, 745–758 (1995).

42. Thisse, C. & Thisse, B. High-resolution in situ hybridization to whole-mount zebrafish embryos.

Nat. Protoc. **3**, 59–69 (2008).

43. Islam, O., Loo, T. X. & Heese, K. Brain-Derived Neurotrophic Factor (BDNF) has Proliferative

Effects on Neural Stem Cells through the Truncated TRK-B Receptor, MAP Kinase, AKT, and

STAT-3 Signaling Pathways. *Curr. Neurovasc. Res.* **6**, 42–53.

44. Finlay, S., Rudd, D., McDermott, B. & Sarnyai, Z. Allostatic load and systemic comorbidities in

psychiatric disorders. *Psychoneuroendocrinology* **140**, 105726 (2022).

45. Elia, J. *et al.* Rare structural variants found in attention-deficit hyperactivity disorder are

preferentially associated with neurodevelopmental genes. *Mol. Psychiatry* **15**, 637–646 (2010).

46. van der Kolk, B. Posttraumatic stress disorder and the nature of trauma. *Dialogues Clin.*

Neurosci. **2**, 7–22 (2000).

47. Karin, O. *et al.* A new model for the HPA axis explains dysregulation of stress hormones on the

timescale of weeks. *Mol. Syst. Biol.* **16**, (2020).

48. Herman, J. P. *et al.* Regulation of the hypothalamic-pituitary-adrenocortical stress response.

Compr. Physiol. **6**, 603–621 (2016).

49. Holsboer, F. The Corticosteroid Receptor Hypothesis of Depression. *Neuropsychopharmacology*

23, 477–501 (2000).

50. Arborelius, L., Owens, M. J., Plotsky, P. M. & Nemeroff, C. B. The role of corticotropin-releasing

factor in depression and anxiety disorders. *J. Endocrinol.* **160**, 1–12 (1999).

51. Yehuda, R. Status of Glucocorticoid Alterations in Post-traumatic Stress Disorder. *Ann. N. Y.*

Acad. Sci. **1179**, 56–69 (2009).

52. Sheng, J. A. *et al.* The Hypothalamic-Pituitary-Adrenal Axis: Development, Programming Actions

of Hormones, and Maternal-Fetal Interactions. *Front. Behav. Neurosci.* **14**, (2021).

53. Notaras, M. & van den Buuse, M. Neurobiology of BDNF in fear memory, sensitivity to stress,

and stress-related disorders. *Mol. Psychiatry* **25**, 2251–2274 (2020).

54. Berton, O. *et al.* Essential Role of BDNF in the Mesolimbic Dopamine Pathway in Social Defeat

Stress. *Science* **311**, 864–868 (2006).

55. Eckert, A. *et al.* The link between sleep, stress and BDNF. *Eur. Psychiatry* **41**, S282 (2017).

56. Fenner, B. M. Truncated TrkB: Beyond a dominant negative receptor. *Cytokine Growth Factor*

Rev. **23**, 15–24 (2012).

57. Cazorla, M. *et al.* Identification of a low–molecular weight TrkB antagonist with anxiolytic and

antidepressant activity in mice. *J. Clin. Invest.* **121**, 1846–1857 (2011).

58. Azogu, I. & Plamondon, H. Blockade of TrkB receptors in the nucleus accumbens prior to

heterotypic stress alters corticotropin-releasing hormone (CRH), vesicular glutamate transporter

2 (vGluT2) and glucocorticoid receptor (GR) within the mesolimbic pathway. *Horm. Behav.* **90**,

98–112 (2017).

59. Bollen, E. *et al.* 7,8-Dihydroxyflavone improves memory consolidation processes in rats and

mice. *Behav. Brain Res.* **257**, 8–12 (2013).

60. Haapasalo, A., Koponen, E., Hoppe, E., Wong, G. & Castrén, E. Truncated trkB.T1 Is Dominant

Negative Inhibitor of trkB.TK+-Mediated Cell Survival. *Biochem. Biophys. Res. Commun.* **280**,

1352–1358 (2001).

61. Maussion, G. *et al.* Regulation of a truncated form of tropomyosin-related kinase B (TrkB) by

Hsa-miR-185* in frontal cortex of suicide completers. *PloS One* **7**, e39301 (2012).

62. Koolhaas, J. M., de Boer, S. F., Coppens, C. M. & Buwalda, B. Neuroendocrinology of coping

styles: Towards understanding the biology of individual variation. *Front. Neuroendocrinol.* **31**,

307–321 (2010).

63. Keatinge, M. *et al.* CRISPR gRNA phenotypic screening in zebrafish reveals pro-regenerative

genes in spinal cord injury. *PLOS Genet.* **17**, e1009515 (2021).

64. Dobrzycki, T., Krecsmarik, M. & Monteiro, R. Genotyping and Quantification of In Situ

Hybridization Staining in Zebrafish. *JoVE J. Vis. Exp.* e59956 (2020) doi:10.3791/59956.

65. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**,

676–682 (2012).

66. Wullimann, M. F., Rupp, B. & Reichert, H. The brain of the zebrafish Danio rerio: a

neuroanatomical atlas. in *Neuroanatomy of the Zebrafish Brain: A Topological Atlas* (eds.

Wullimann, M. F., Rupp, B. & Reichert, H.) 19–87 (Birkhäuser, Basel, 1996). doi:10.1007/978-3-

0348-8979-7_5.

67. García-González, J., de Quadros, B., Havelange, W., Brock, A. J. & Brennan, C. H. Behavioral

Effects of Developmental Exposure to JWH-018 in Wild-Type and Disrupted in Schizophrenia 1

(disc1) Mutant Zebrafish. *Biomolecules* **11**, 319 (2021).

68. R Development Core Team. R: A language and environment for statistical computing.

https://stat.ethz.ch/pipermail/r-help/2008-May/161481.html (2008).

69. Gutiérrez, H. C. *et al.* Screening for drugs to reduce zebrafish aggression identifies caffeine and

sildenafil. *Eur. Neuropsychopharmacol.* **30**, 17–29 (2020).

70. Parker, M. O., Brock, A. J., Sudwarts, A. & Brennan, C. H. Atomoxetine reduces anticipatory

responding in a 5-choice serial reaction time task for adult zebrafish. *Psychopharmacology*

(Berl.) **231**, 2671–2679 (2014).

71. Pfaffl, M. W. A new mathematical model for relative quantification in real-time RT–PCR. *Nucleic*

Acids Res. **29**, e45 (2001).

72. Dean, R. B. & Dixon, W. J. Simplified Statistics for Small Numbers of Observations. *Anal. Chem.*

23, 636–638 (1951).

Figure 1. *rbfox1* **mutant fish show hyperactivity, impulsivity and hyperarousal behaviour. A-C)** Locomotion assay in 5 days post fertilization (dpf) zebrafish larvae (*rbfox1^{+/+}*, *rbfox1^{+/19del}*, *rbfox1^{19del/19del}*): **A)** distance travelled during 15 min; **B)** average speed; **C)** number of peaks/swimming burst (acceleration events when the fish travelled > 5 mm in < 12 s); N (A-C) = *rbfox1^{+/+}* 24; *rbfox1^{+/19del}* 24; *rbfox1^{19del/19del}* 24. **D-H)** 5-choice serial reaction time task in adult (7 months old) zebrafish (*rbfox1^{+/+}*, *rbfox1^{+/19del}*, *rbfox1^{19del/19del}*): **D)** number of initiated trials during stage 2 (weeks 2-4); all

rbfox1$^{+/+}$ (19), all *rbfox1*$^{+/19del}$ (18) and all *rbfox1*$^{19del/19del}$ (40) moved to the next stage; **E-G)** average

correct responses for stages 3-5 (weeks 3-9); **E)** at the end of stage 3 (weeks 3-7), all *rbfox1*$^{+/+}$ (19), 17

rbfox1$^{+/19del}$ and 39 *rbfox1*$^{19del/19del}$ moved to the next stage; **F)** at the end of stage 4 (weeks 4-8), all

rbfox1$^{+/+}$ (19), all *rbfox1*$^{+/19del}$ (17) and 33 *rbfox1*$^{19del/19del}$ moved to the next stage; **F)** at the end of stage

5 (weeks 5-9), 15 *rbfox1*$^{+/+}$, 13 *rbfox1*$^{+/19del}$ and all *rbfox1*$^{19del/19del}$ (33) showed ≥ 50% correct responses;

H) premature responses during stage 5 (weeks 5-9). Each dot, square or triangle in D-H) represents a

single adult zebrafish *rbfox1*$^{+/+}$, *rbfox1*$^{+/19del}$ and *rbfox1*$^{19del/19del}$ respectively; N = *rbfox1*$^{+/+}$ 19;

rbfox1$^{+/19del}$ 18; *rbfox1*$^{19del/19del}$ 40. **I-J)** Response and habituation to acoustic startle assay in 5dpf

zebrafish larvae (*rbfox1*$^{+/+}$, *rbfox1*$^{+/19del}$, *rbfox1*$^{19del/19del}$): **I)** mean distance travelled during the assay; **J)**

rate of habituation over time/stimuli; N = *rbfox1*$^{+/+}$ 40; *rbfox1*$^{+/19del}$ 39; *rbfox1*$^{19del/19del}$ 35. **K-L)** Forced

light-dark transition assay in 5dpf zebrafish larvae (*rbfox1*$^{+/+}$, *rbfox1*$^{+/19del}$, *rbfox1*$^{19del/19del}$): **K)** distance

travelled during the 10 min assay; **I)** 1 sec time bin resolution plots of the dark-light transition; N =

rbfox1$^{+/+}$ 43; *rbfox1*$^{+/19del}$ 34; *rbfox1*$^{19del/19del}$ 37. All larvae employed in behavioural experiments were

progeny of *rbfox1*$^{+/19del}$ in-cross and were genotyped after experiments and prior to data analysis. In

all graphs: bars represent standard error of the mean (SEM); * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$;

**** $p < 0.0001$.

Figure 2. *rbfox1* **LoF disrupts zebrafish larvae HPI axis,** *bdnf/trkb2* **pathway and** *pac1a* **expression**.
TRKB modulation restore HPI gene expression. A) Expression levels of *bdnf* and *trkb2* full-length (*TK+*)
and truncated (*TK-*) in 5 days post fertilisation (dpf) larvae and of **B)** *crhb*, *bdnf*, *TK+* and *TK-* in 3dpf
larvae (*rbfox1*+/+ and *rbfox1*19del/19del). **C-D)** Expression levels of HPI axis genes *crhb*, *mr* and *gr* in 5dpf
larvae (*rbfox1*+/+ and *rbfox1*19del/19del) in presence or absence of the TRKB selective **C)** antagonist ANA-
12 and **D)** agonist 7,8-DHF. **E-G)** Whole mount *in situ* hybridisation (ISH) for *trkb2* anti-sense riboprobe
in **E)** *rbfox1*+/+ and **F)** *rbfox1*19del/19del and **G)** *trkb2* sense riboprobe in *rbfox1*+/+ 5dpf larvae (lateral view
in the main boxes and dorsal view in the smaller boxes on the top right corner). **H, I)** Sagittal
cryosection of *trkb2* ISH in **H)** *rbfox1*+/+ and **I)** *rbfox1*19del/19del 5 dpf larvae. Black boxes in **H-a)** and **I-a)**
represent the region of the brain showed in higher magnification panels in **H-a')** and **I-a')**. Scale bars:

200 μm in E), F) and G); 100 μm in F-a) and I-a); 50 μm in F-a) and I-a). **J)** Schematic depiction (sagittal)
of zebrafish larval brain indicating position of levels illustrated by ISH on sagittal cryosections. **K)** *trkb2*
ISH intensity mean in the hypothalamus of 5dpf zebrafish larvae, *rbfox1$^{+/+}$* versus *rbfox1$^{19del/19del}$*. N = 4
larvae x genotype. **L)** Expression levels of *pac1-hop* and *pac1-short* in *rbfox1$^{+/+}$* and *rbfox1$^{19del/19del}$* 5 dpf
larvae. For qPCR experiments, reference genes were actin – β 2 (*actb2*) and ribosomal protein L13a
(*rpl13a*). Each green dot/pink triangle in A-D, K-L) represents a pool of 15 larval heads (eyes and jaw
removed), while yellow ones represent larvae exposed to TRKB drugs. Where indicated, we used Log$_{10}$
transformation to normalize the data facilitating a clearer visualization of trends within the dataset.
All larvae employed were progeny of *rbfox1$^{+/19del}$* in-cross and were genotyped after experiments and
prior to data analysis. In all graphs: bars represent standard error of the mean (SEM); * p < 0.05; ** p
< 0.01; **** p < 0.0001.

Figure   3. *rbfox1* mutants undergo adaptive mechanisms and allostatic overload during
development. Expression levels of **A)** HPI axis genes corticotropin releasing hormone b (*crhb*), mineral
corticoid receptor (*mr*) and glucocorticoid receptor (*gr*) and of **B)** *bdnf* and *trkb2* truncated/full-length
(*TK-/TK+*) (common primers) in adult zebrafish brain (*rbfox1$^{+/+}$* and *rbfox1$^{19del/19del}$*) in normal resting
conditions and after exposure to a stressor (novel tank diving). **C)** Expression levels of the *trkb2 TK+*
and *TK-* in adult zebrafish brain (*rbfox1$^{+/+}$* and *rbfox1$^{19del/19del}$*) after exposure to a stressor. **D)**
Expression levels of proliferating cell nuclear antigen (*pcna*) in 5 days post fertilisation zebrafish larvae
and adult zebrafish brain (*rbfox1$^{+/+}$* and *rbfox1$^{19del/19del}$*) in normal resting conditions. **E)** Fertility rate
and **F)** survival rate of *rbfox1$^{+/+}$* and *rbfox1$^{19del/19del}$*. Each green dot/pink triangle in A-B) represents a
single adult *rbfox1$^{+/+}$* or *rbfox1$^{19del/19del}$* brain under resting physiological conditions respectively, while

yellow dots/triangles represent $rbfox1^{+/+}$ or $rbfox1^{19del/19del}$ brain after stress exposure respectively. In
B) each green dot/pink triangle represents single adult $rbfox1^{+/+}$ or $rbfox1^{19del/19del}$ brain after stress
exposure respectively. In D) for larvae each green dot/pink triangle represents a pool of $rbfox1^{+/+}$ or
$rbfox1^{19del/19del}$ 15 larval heads (eyes and jaw removed) respectively; for adults each green dot/pink
triangle represents a single adult $rbfox1^{+/+}$ or $rbfox1^{19del/19del}$ brain respectively. In E) each dot/triangle
represents average fertility of 3-5 trios (1 male and 2 females) assessed over 2-3 petri dish (50 embryos
per dish) per trio. Each trio belonged to a different tank (for each genotype for each batch) to avoid
tank effect). In F) each dot/triangle represents the percentage of survival of a single fish stock
comprising 50 larvae. For qPCR experiments, reference genes were actin – β 2 (*actb2*) and ribosomal
protein L13a (*rpl13a*). Where indicated, we used Log_{10} transformation to normalize the data
facilitating a clearer visualization of trends within the dataset. All larvae employed were progeny of
$rbfox1^{+/19del}$ in-cross and were genotyped after experiments and prior to data analysis. In all graphs:
bars represent standard error of the mean (SEM); * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p <$
0.0001.

A)

B)

912
913
Figure 4. RBFOX1 role in stress resilience and chronic stress recovery: proposed mechanism of
action. A) In response to stressful challenges, BDNF expression levels increase, stimulating *CRH*
transcription via TRKB.FL/TK+ (yellow pathway). At the same time, stress triggers also *PACAP*
transcription (red pathway) and OTP-mediated transcription of *RBFOX1* and *CRH* (blue pathway).
PACAP stimulates *CRH* transcription via PAC1-short and inhibits *CRH* transcription via PAC1-hop.
PACAP binding to PAC1-short also increases *BDNF* transcription (red dashed line) and enhances
TRKB.FL/TK+ activity (red dashed line). RBFOX1 functions as a "switch" maintaining the balance
between full-length/truncated TRKB isoforms (promoting TRKB mRNA stability and/or alternative
splicing): during the stress recovery phase, *RBFOX1* switches the balances in support of *TRKB.T1/TK-*

isoform, decreasing *CRH* levels and turning off the HPA axis. **B)** *RBFOX1* deletion leads to an increase
in *BDNF* expression levels and an unbalance between *TRKB.FL/TK+* (increased) and *TRKB.T1/TK-*
(decreased) isoforms. Such dysregulation ultimately leads to an increase in *CRH* transcription and
hyperactivation of the HPA axis. Our data suggest that RBFOX1-mediated regulation of *TRKB.T1/TK-*
mRNA stability and/or alternative splicing is necessary to promote neuroplasticity and stress
resilience.

Supplementary Figure 1. Generation of a *rbfox1* loss of function line. A) *rbfox1* mRNA expression
in *rbfox1+/+*, *rbfox1+/19del* and *rbfox1 19del/19del* 5 days post fertilisation (dpf) larvae. *In situ* hybridisation
on **B)** whole mount zebrafish larvae at 3dpf, a) *rbfox1+/+* and b) *rbfox1 19del/19del* and **C)** adult transverse
sections of *rbfox1+/+* (a) fore- and (c) mid- brain and *rbfox1 19del/19del* (b) fore- and (d) mid-brain. In A)
each dot/square/triangle represents a pool of *rbfox1+/+* or *rbfox1 19del/19del* 15 larval heads (eyes and jaw
removed) respectively. For qPCR experiments, reference genes were actin – β 2 (*actb2*) and ribosomal
protein L13a (*rpl13a*). All larvae employed were progeny of *rbfox1+/19del* in-cross and were genotyped
after experiments and prior to data analysis. In A): bars represent standard error of the mean (SEM);
* p < 0.05; ** p < 0.01. Scale bars: 200 μm in B); 100 μm in C). Abbreviations: D, dorsal area of dorsal
telencephalon; Hv, ventral hypothalamus; TeO, optic tectum; Tl, longitudinal tori; Vd, ventral area of
dorsal telencephalon; Vv, ventral area of ventral telencephalon.

Supplementary Figure   2.   *rbfox1* is expressed in regions of the brain associated with stress response, social and emotional behaviour, reward and learning. A) *rbfox1* whole mount *in situ* hybridization (ISH) on zebrafish larvae at 28 hours post fertilization and 3-4-5 days post fertilization. **B)** *rbfox1* ISH on adult zebrafish (a - c) forebrain and (d-e) mid-/hind-brain transverse sections. Black boxes represent the region of the brain showed in higher magnification panels. Schematic depictions of the lateral view of the zebrafish brain indicate position of levels illustrated by ISH. Scale bars: 200 µm in A), 100 µm in B) a-d); 50 µm in B) a', c, d', d'', e. Abbreviations: A, anterior tuberal nucleus; ATN, anterior thalamic nucleus; CZ, central zone of the TeO; D, dorsal area of dorsal telencephalon; Dm, dorsal area of medial telencephalon; ECL, external cellular layer; GL, glomerular layer; HaV, ventral area of the habenula; Hv, ventral hypothalamus; ICL, internal cellular layer; PGZ, periventricular gray zone; TeO, optic tectum; TL, longitudinal tori; TPp, periventricular nucleus of the posterior tuberculum; Val, valvula cerebelli; Vd, ventral area of dorsal telencephalon; Vl, central area of lateral telencephalon; VM, ventromedial nucleus; Vv, ventral area of ventral telencephalon.

Supplementary Figure 3. Novel tank diving assay in adult *rbfox1*⁺/⁺, *rbfox1*⁺/¹⁹ᵈᵉˡ and *rbfox1*¹⁹ᵈᵉˡ/¹⁹ᵈᵉˡ fish. A) Time spent at the bottom of the tank. **B)** Distance travelled during the assay and **C)** number of transitions between the top and bottom of the tank. N = *rbfox1*⁺/⁺, 50; *rbfox1*⁺/¹⁹ᵈᵉˡ 54; *rbfox1*¹⁹ᵈᵉˡ/¹⁹ᵈᵉˡ 54. Bars represent standar error of the mean (SEM).

Supplementary Figure 4. A) *rbfox2* expression levels in 5 days old larvae, *rbfox1*⁺/⁺ and *rbfox1*¹⁹ᵈᵉˡ/¹⁹ᵈᵉˡ. Actin – β 2 (*actb2*) and ribosomal protein L13a (*rpl13a*) were used as reference genes. Dots/triangles represent pool of 15 larval heads (eyes and jaw removed). Bars represent standard error of the mean (SEM). N = *rbfox1*⁺/⁺ 5; *rbfox1*¹⁹ᵈᵉˡ/¹⁹ᵈᵉˡ 5.

Supplementary Figure 5. Details of *in situ* hybridisation experiments for *trkb2* anti-sense riboprobe on 5 days post fertilisation zebrafish larvae, sagittal sections. **A)** Overview of larval sagittal sections, *rbfox1*$^{+/+}$ (four panels on the left) and *rbfox1*$^{19del/19del}$ (four panels on the right). Each panel corresponds to a different animal. N = 4 per genotype. Sections have been placed on the same slide and were stained under the same conditions. **B)** *In situ* hybridisation for *trkb2* sense riboprobe on 5 days post fertilisation zebrafish larva, sagittal section. **C)** Example of areas selected (areas enclosed by the black dotted lines) to calculate *trkb2* mean intensity in the hypothalamus of zebrafish *rbfox1*$^{+/+}$ (left panel) and *rbfox1*$^{19del/19del}$ (right panel) larval sections.

Supplementary Table 1. List of primers used for qPCR experiments.

Gene (purpose)	Gene acronym	Accession number	Sequence (5' to 3')	Product size (bp)
actin, β 2 (housekeeping)	*actb2*	NM_181601.5	Forward: CGAGCTGTCTTCCCATCCA	86
			Reverse: TCACCAACGTAGCTGTCTTTCTG	
ribosomal protein L13a (housekeeping)	*rpl13a*	NM_212784.1	Forward: TCTGGAGGACTGTAAGAGGTATGC	148
			Reverse: AGACGCACAATCTTGAGAGCAG	
RNA binding fox-1 homolog 1	*rbfox1**	NM_001005596.1	Forward: AACAGCATACAGGCGGTCTC	110
			Reverse: GGCTGCGTTTTGATTTCTGT	
corticotropin releasing hormone	*crhb*	NM_001007379.1	Forward: CGAGACATCCCAGTATCCAAAAAG	60
			Reverse: TCCAACAGACGCTGCGTTAA	
mineralcorticoid receptor	*mr*	NM_001100403	Forward: CTTCCAGGTTTCCGCAGTCTAC	75
			Reverse: GGAGGAGAGACACATCCAGGAT	
glucocorticoid receptor	*gr*	NM_001020711.3	Forward: ACTCCATGCACGACTTGGTG	90
			Reverse: GCATTTCGGGAAACTCCACG	
brain derived neurotrophic factor	*bdnf*	XM_009303234.3	Forward: AGCAGACTGTCTTTTGTTGTGTT	123
			Reverse: CTTAAGCTCACACAGGTGCC	
neurotrophic tyrosine kinase, receptor, type 2b (common primers)	*trkb2.TK+/TK-*	NM_001197161.2	Forward: CGCATACCTGACGTGGATCA	130
			Reverse: AAGCAACCCCAGCTATTCCC	
neurotrophic tyrosine kinase, receptor, type 2b (tyrosine kinase domain primers)	*trkb2.TK+*	NM_001197161.2	Forward: TGAAAGTGGGCGAGCAGATT	173
			Reverse: CATCTGGACCATGAGCCCTG	
adenylate cyclase activating	*pac1a-hop*	NM_001142926.1	Forward: CAGAAATGCTTCAGTGAGCCC	82

polypeptide 1a (pituitary) receptor type I (adcyap1r1a) transcript variants 1 and 2		NM_001013444 .2	Reverse: CCAGACGTAGCGTGATGGTA	
adenylate cyclase activating polypeptide 1a (pituitary) receptor type I (adcyap1r1a), transcript variant 3	*pac1a-short*	NM_001142925 .1	Forward: GATTCTGGCTGCTGGGACAT	183
			Reverse: GCGCGCCAGACGTAGATAAA	
RNA binding fox-1 homolog 2	*rbfox2*	NM_214746.2	Forward: CCGTATGCTAACGGGGACAG	106
			Reverse: ACCCTGGAACTGCGTAGAAC	

992 * These primers amplify all *Danio rerio rbfox1* isoforms available on NCBI: XM_068216713.1,
993 XM_068216714.1, XM_017354041.3, XM_068216717.1, XM_068216718.1, XM_017354043.3,
994 XM_017354044.3, XM_068216722.1, XM_017354045.3, XM_068216724.1, XM_068216725.1,
995 XM_068216726.1, XM_068216728.1, XM_017354049.3   XM_068216731.1.
996
997
998 **Supplementary Table 2.** 5-Choice Serial Reaction Time Task stages (5-CSRTT) and corresponding
999 descriptions. Source: https://zantiks.com
1000

Stage	Name of the stage	Description
Stage 1	Habituation	All lights are illuminated (initiator light, five choice lights and food light) for 20 sec and the food is released. All fish are moved to the next stage of training after 1 week.
Stage 2	Initiator training	Initiator light is on for 20 sec. If the fish swims to the illuminated initiation area, an initiated trial is recorded and the feeder releases food with the food light on for 20 sec. This is followed by 10 sec ITI. The initiator training is repeated for 30 trials. Fish move to the next stage when the average of initiated trials to total trials over five days is > 50%.
Stage 3	Stimulus light training	The initiation light is on for 20 sec. If the fish swims to the illuminated initiation area, an initiated trial is recorded and all five choice lights are illuminated for 20 sec. If the fish swims to any illuminated choice area whilst the light is on, a correct response is recorded and the feeder releases food with the food light on for 20 sec. This is followed by 10 sec ITI. The stimulus light training is repeated for 30 trials. Fish move to the next stage when

		both the average of initiated trials to total trials and correct responses to initiated trials are > 50%.
Stage 4	5-CSRTT/no-delay	The initiation light is on for 20 sec. If the fish swims to the initiation area, an initiated trial is recorded and one of five choice lights is illuminated for 20 sec. If the fish swims to the illuminated choice area, a correct response is recorded and the feeder releases food with the food light on for 20 sec. This is followed by 10 sec ITI. The 5-CSRTT/no-delay stage is repeated for 30 trials. Fish move to the next stage when the average of both initiated trials to total trials and correct responses to initiated trials are > 50%.
Stage 5	5-CSRTT/variable-delay	The initiation light is on for 20 sec. If the fish swims to the initiation area, an initiated trial is recorded and after a variable pre-stimulus interval of 0-8 sec, one of five choice lights is illuminated for 20 sec. This is followed by 10 sec ITI. If during the pre-stimulus interval, the fish swims to any unilluminated choice area, a premature response is recorded. The average of premature responses to initiated trials over four days is reported as a measure of impulsive action.

1001