rbfox1 LoF mutants show disrupted *bdnf/trkb2* and *crhb/nr3c2* expression and increased
cortisol levels during development coupled with signs of allostatic overload in adulthood

Adele Leggieri[1], Judit García-González[2], Saeedeh Hosseinian[1], Peter Ashdown[1], Sofia Anagianni[1], Xian
Wang[1], William Havelange[1], Noèlia Fernàndez-Castillo[3,4,5,6], Bru Cormand[3,4,5,6] and Caroline H.
Brennan[1]*

1 Centre for Brain and Behaviour, School of Biological and Behavioural Sciences, Queen Mary
University of London, Mile End Rd, London, E1 4NS, United Kingdom

2 Department of Genetics and Genomic Sciences, Icahn School of Medicine, Mount Sinai, New York
City, NY 10029, USA

3 Departament de Genètica, Microbiologia i Estadística, Facultat de Biologia, Universitat de
Barcelona, Barcelona, Catalunya, 08028, Spain

4 Centro de Investigación Biomédica en Red de Enfermedades raras (CIBERER), Spain

5 Institut de Biomedicina de la Universitat de Barcelona, Barcelona, Catalunya, 08028, Spain

6 Institut de recerca Sant Joan de Déu, Espluges de Llobregat, Catalunya, 08950, Spain

* Correspondence: Caroline H. Brennan, c.h.brennan@qmul.ac.uk

ABSTRACT

Mutations in the RBFOX1 gene are associated with psychiatric disorders but how RBFOX1 influences
psychiatric disorder vulnerability remains unclear. Recent studies showed that RBFOX proteins
mediate the alternative splicing of PAC1, a critical HPA axis activator. Further, RBFOX1 dysfunction is
linked to dysregulation of BDNF/TRKB, a pathway promoting neuroplasticity, neuronal survival, and
stress resilience. Hence, RBFOX1 dysfunction may increase psychiatric disorder vulnerability via HPA
axis dysregulation, leading to disrupted development and allostatic overload. To test this hypothesis,
we generated a zebrafish *rbfox1* loss of function (LoF) line and examined behavioural and molecular
effects during development. We found that *rbfox1* LoF mutants exhibited hyperactivity, impulsivity
and heightened arousal, alongside alterations in proliferation – traits associated with
neurodevelopmental and stress-related disorders. In adults, loss of *rbfox1* function led to decreased
fertility and survival, consistent with allostatic overload. At the molecular level, at larval stages
rbfox1 mutants showed increased cortisol levels and disrupted expression of key stress-related
genes (*bdnf, trkb2, pac1a-hop, crhb, nr3c2*). Pharmacological intervention targeting TRKB restored
crhb and *nr3c2* gene expression and hyperactive and hyperarousal behaviours. In adults,
dysregulation of *crhb*, *nr3c2* and *bdnf/trkb2* genes was only seen following acute stress exposure.
Our findings reveal a fundamental role for RBFOX1 in integrating stress responses through its
regulation of BDNF/TRKB and neuroendocrine signalling.

INTRODUCTION

Psychiatric disorders are influenced by multiple genes, with a substantial portion of their heritability
linked to common genetic variations[1]. Identifying gene variants associated with multiple psychiatric
disorders is crucial for understanding their shared genetic underpinnings and enhancing therapeutic
strategies.

RNA binding fox-1 homologue 1 (RBFOX1), also known as FOX1 or ataxin 2-binding protein 1
(A2BP1), is a splicing factor highly conserved among vertebrates and expressed in the heart, brain
and skeletal muscle, where it contributes to normal development and function[2,3]. Alterations in
RBFOX1 expression or function have been associated with susceptibility to psychiatric disorders,
particularly those associated with changes in stress-related behaviour[4–6]. In a cross-disorder
genome-wide association study (GWAS), *RBFOX1* emerged as the second most pleiotropic *locus*,
showing association with seven out of the eight disorders studied: schizophrenia, bipolar disorder,
depression, attention-deficit/hyperactivity disorder (ADHD), autism spectrum disorder (ASD),
obsessive-compulsive disorder and Tourette syndrome[1]. In mice and zebrafish, *Rbfox1/rbfox1*
knockout caused hyperactivity, increased anxiety-like behaviour and altered social behaviour[2,6].
However, the mechanisms by which *RBFOX1* genetic variants contribute to psychiatric disease are
poorly understood.

In both mice and zebrafish, it has been suggested that Rbfox1 regulates the Pituitary Adenylate
Cyclase-Activating Polypeptide Type 1 Receptor (Pac1) alternative splicing, an important mediator of
corticotropin releasing hormone (CRH) synthesis in the hypothalamus[4,7]. CRH is the first hormone to
be secreted in response to stress by the hypothalamic-pituitary-adrenal (HPA) axis (hypothalamic-
pituitary-interrenal (HPI) axis in fish)[8]. In mammals there are several PAC1 isoforms with the
predominant isoforms in the brain being PAC1-hop (long isoform) and PAC1-short[9]. In mice and
zebrafish, following acute stress exposure, both *Pac1-hop/pac1-hop* and *Pac1-short/pac1-short*
expression increased, while at the late recovery phase, only *Pac1-hop* was still up-regulated[4].

Another regulator of the stress response and a resilience factor against chronic stress-induced
psychopathology is the brain derived neurotrophic factor (BDNF)/Tropomyosin receptor kinase B
(TRKB) pathway[10–12]. BDNF is a neurotrophin possessing a pivotal role in the modulation of
neurotransmission and synaptic plasticity and dysregulation of the BDNF/TRKB pathway has been
associated with several neuropsychiatric diseases, including anxiety/stress disorders[13,14]. In human
neural stem cells, *RBFOX1* knockdown increased *BDNF* expression levels[15], while another study in
mice identified *TrkB* as a target of RBFOX1 within the hippocampus[16]. Further, PAC1-short activation
can elevate *BDNF* levels and potentiate TRKB activity, enhancing BDNF/TRKB neuroprotective and
plasticity-promoting effects, especially in the context of stress response and neuropsychiatric
health[17].

Given that both PAC1 and BDNF/TRKB influence how the brain adapts to and manages stress,
RBFOX1 variants may increase susceptibility to psychiatric disorders through dysregulation of the
stress response, leading to adaptive plasticity and disrupted development in the short-term and
allostatic overload in the long-term. Allostasis refers to the collective processes by which the body
actively maintains stability (or homeostasis) in response to environmental stressors[12]. When
allostasis primary mediators (e.g., HPA axis hormones, catecholamines, cytokines) are overactivated
or fail to return to normal, it leads to an allostatic state[18,19]. The cumulative results of an allostatic
state are referred to as allostatic load and, when maladaptive, allostatic overload[18]. Excess
glucocorticoid exposure during early life and early life stress have been shown to cause prolonged
activation of the allostatic systems, ultimately leading to allostatic overload[19–22], while neurotrophic
factors such as BDNF play a key role in regulating adaptive plasticity and mechanisms counteracting
damage caused by allostatic overload[12]. Here, to explore the possibility that *RBFOX1* loss of function
(LoF) leads to increased vulnerability to psychiatric disease through HPA axis hyperactivation and
allostasis-induced adaptation during development, we generated a zebrafish line carrying a
predicted *rbfox1* LoF mutation using CRISPR-Cas9 gene editing and assessed behavioural and
molecular changes at different developmental stages. We hypothesized that RBFOX1 regulates HPA
axis activity through an effect on BDNF/TRKB signalling leading to disrupted brain development.

RESULTS

Generation of a loss of function *rbfox1* line

CRISPR-Cas9 genome editing generated a 19 base pair deletion (NM_001005596.1, nucleotides 120-
138, TCCCATCGGCCCAGTTCGC) that introduced a premature termination codon (PTC) at position 58
in the *rbfox1* amino acid sequence. This line is recorded on ZFIN as qm4, ZDB-ALT-240222-6. Details
regarding the generation of the line, nucleotide and amino acid sequences can be found in our
previous study[2]. Wild type animals are denoted throughout as *rbfox1$^{+/+}$*, heterozygous mutants as
rbfox1$^{+/19del}$ and homozygous mutants as *rbfox1$^{19del/19del}$*.

To evaluate whether the PTC elicited mRNA non-sense mediated decay (NMD) and consequent
reduction of *rbfox1* mRNA in mutant fish, we examined *rbfox1* expression by quantitative Real-Time
polymerase chain reaction (qPCR) and by *in situ* hybridisation (ISH). As *RBFOX1* itself is alternatively
spliced to generate nuclear and cytoplasmic isoforms[7], we designed primers targeting all *rbfox1*
zebrafish isoforms available on NCBI (see Supplementary Table1). qPCR showed that *rbfox1*
transcript levels were significantly lower in mutant larvae compared to *rbfox1$^{+/+}$* siblings ($p_{rbfox1}{}^{+/+}$ vs
$_{rbfox1}{}^{+/19del}$ < 0.05, $p_{rbfox1}{}^{+/+}$ vs $_{rbfox1}{}^{19del/19del}$ < 0.01), and ISH showed that *rbfox1* was not detectable in
rbfox1$^{19del/19del}$ fish, at either larval or adult stages (Supplementary Figure 1A-C).

These results confirm degradation of defective *rbfox1* mRNA in mutant fish.

rbfox1 mutant fish show hyperactivity, impulsivity and hyperarousal behaviour

As mutations in _RBFOX1_ locus have been linked to several psychiatric diseases in humans, we assessed zebrafish larvae and adult fish for phenotypic traits associated with such disorders and examined _rbfox1_ mRNA distribution in _rbfox1$^{+/+}$_ larvae and adults (Tübingen strain).

In humans, _RBFOX1_ copy number variants (CNVs) and LoF mutations are risk factors for ADHD[6]. Two major ADHD traits are hyperactivity and increased impulsivity. We therefore assessed hyperactive and impulsive behaviour of _rbfox1$^{+/+}$_, _rbfox1$^{+/19del}$_ and _rbfox1$^{19del/19del}$_ 5 days post fertilization (dpf) larvae and adults. When we measured larval locomotion, we observed a gene dosage effect on distance travelled, whereby _rbfox1_ mutant larvae travelled greater distances than wild type siblings ($p_{rbfox1^{+/+} \text{ vs } rbfox1^{+/19del}} < 0.05$, $p_{rbfox1^{+/+} \text{ vs } rbfox1^{19del/19del}} < 0.0001$, $p_{rbfox1^{+/19del} \text{ VS } rbfox1^{19del/19del}} < 0.05$) (Supplementary Figure 2A), and a significant increase in the swimming speed of _rbfox1$^{19del/19del}$_ larvae ($p_{rbfox1^{+/+} \text{ vs } rbfox1^{19del/19del}} < 0.0001$, $p_{rbfox1^{+/19del} \text{ VS } rbfox1^{19del/19del}} < 0.05$) (Supplementary Figure 2B). This is in line with hyperactivity observed in adult _rbfox1$^{19del/19del}$_, in our previous study and in other _rbfox1_ model[2,23]. We also measured larval burst swimming, a parameter previously used as a measure to predict impulsive behaviour in zebrafish larvae[24]. We found a significant increase in the number of peaks (acceleration events when the fish travelled > 5 mm in < 12 sec) in _rbfox1$^{19del/19del}$_ larvae ($p_{rbfox1^{+/+} \text{ vs } rbfox1^{19del/19del}} < 0.0001$, $p_{rbfox1^{+/19del} \text{ VS } rbfox1^{19del/19del}} < 0.0001$) (Figure 1A). Impulsive behaviour was then assessed in adult (7 months old) fish using the 5-choice serial reaction time task (5-CSRTT)[25]. The 5-CSRTT assay measures sustained attention and impulsive action by requiring an animal to detect a brief visual cue (white light stimulus) presented randomly across one of five apertures and "nosepoke" that aperture to signal recognition (Supplementary Figure 3). If correct, a "food" signal light comes on (at the opposite end of the test chamber) and the animal collects food reward (Supplementary Figure 3). Crucially, a pause occurs prior to the onset of the stimulus light (called a pre-stimulus interval [PSI]), during which a 'premature' response can be interpreted as 'impulsivity'- impulsive action[25]. The assay consists of five stages (Supplementary Table2) each run for at least a week until fish are promoted to the next stage. We found that 79% _rbfox1$^{+/+}$_, 72% _rbfox1$^{+/19del}$_ and 62% _rbfox1$^{19del/19del}$_ learned the task within 9 weeks. We found no significant differences in the correct responses (stages 2-5, Figure 1B-E). In stage 5, we found a significant difference in the number of premature responses (increased impulsivity) such that _rbfox1$^{19del/19del}$_ fish were more impulsive compared to _rbfox1$^{+/+}$_ siblings ($p < 0.05$) (Figure 1F).

RBFOX1 CNVs have also been identified in individuals with schizophrenia[6,15]. Deficits in habituation to acoustic startle reflex are seen in both schizophrenic patients and animal models of schizophrenia[26]. The acoustic startle assay has been widely employed to measure habituation in animals, including zebrafish[27]. The magnitude of the startle response and the extent of habituation can also serve as indicators of hyperarousal linked to heightened stress responses[28]. We therefore tested 5dpf zebrafish larvae (_rbfox1$^{+/+}$_, _rbfox1$^{+/19del}$_ and _rbfox1$^{19del/19del}$_) in the habituation to acoustic startle response assay. Consistent with our hyperactivity assay, during the baseline (first 10 min), the genotype had a significant main effect on distance travelled [Effect of genotype: $\chi2(2) = 8.3397$, $p < 0.05$] whereby _rbfox1$^{19del/19del}$_ larvae travelled greater distances than _rbfox1$^{+/+}$_ siblings ($p < 0.05$). During the startle stimuli, we observed a main effect of genotype [Effect of genotype: $\chi2(2) = 33.536$, $p < 0.0001$] and stimulus number [Effect of stimulus number: $\chi2(9) = 487.968$, $p < 0.0001$] on distance travelled, and a significant two-way interaction between genotype and stimulus number [Effect of genotype by stimuli: $\chi2(18) = 31.511$, $p < 0.05$], whereby _rbfox1$^{19del/19del}$_ larvae travelled greater distances than _rbfox1$^{+/+}$_ ($p < 0.001$) and _rbfox1$^{+/19del}$_ ($p < 0.01$) (Supplementary Figure 2C). When assessed for the rate of habituation, wild type larvae showed a habituation response to repeated acoustic startle consistent with previous reports[29]: 100% of _rbfox1$^{+/+}$_animals responded to the first acoustic stimulus, but only 22% responded to the last. When we examined the rate of

habituation over time, in line with previous findings[30], we observed a significant genotype effect
[Effect of genotype: $\chi2(2)$ =7.2676, $p < 0.05$] and a significant two-way interaction between
genotype and stimulus number [Effect of genotype by stimulus number: $\chi2(18)$ =132.8476, $p <$
0.001] whereby $rbfox1^{19del/19del}$ showed reduced rate of habituation and a greater proportion of
responders compared to $rbfox1^{+/+}$ siblings ($p_{rbfox1^{+/+} vs\ rbfox1^{19del/19del}} < 0.05$) (Supplementary Figure
2D).

SNPs in *RBFOX1* locus have also been associated with anxiety[6,31]. Hence, we assessed anxiety-like
behavior in 5dpf zebrafish larvae using the forced light-dark transition (FLDT) assay and in adult
zebrafish using the novel tank diving (NTD) assay. In the FLDT assay, zebrafish are exposed to sudden
transitions in illumination with effects on locomotion and amplitude of response on transition from
dark to light being used as a measure of anxiety-like behaviour: the increased locomotion/startle
response upon the light-dark transition is a measure of anxiety-like behaviour[29,32,33]. Consistent with
our acoustic startle assay, we found that during the baseline (first 5 min) (Figure 1G) genotype had a
significant main effect on distance travelled [Effect of genotype: $\chi2(2)$ = 14.4878, $p < 0.001$]
whereby $rbfox1^{19del}$ larvae travelled greater distances than $rbfox1^{+/+}$ siblings ($p_{rbfox1^{+/+} vs\ rbfox1^{+/19del}} <$
0.05, $p_{rbfox1^{+/+} vs\ rbfox1^{19del/19del}} < 0.01$). During the 1min light flash (min 5-6 of the assay) we found a
significant main effect of genotype on distance travelled [Effect of genotype: $\chi2(2)$ = 8.8518, $p <$
0.05] with $rbfox1^{19del}$ larvae travelling greater distances than $rbfox1^{+/+}$ siblings ($p_{rbfox1^{+/+} vs\ rbfox1^{+/19del}} <$
0.05, $p_{rbfox1^{+/+} vs\ rbfox1^{19del/19del}} < 0.05$) (Figure 1H). On transition from dark to light (analysed at 1 sec
time bin resolution from sec 240 to sec 360 as the startle response during the transition is visible
only at this resolution) we found a significant main effect of time [Effect of time: $\chi2(118)$ =
1051.3482, $p < 0.001$] and genotype [Effect of genotype: $\chi2(2)$ = 8.9155, $p < 0.05$], and a significant
two-way interaction between time and genotype [Effect of time by genotype: $\chi2(236)$ = 514.8308, p
< 0.001] on the amplitude of response, whereby $rbfox1^{19del/19del}$ larvae startled more than $rbfox1^{+/+}$ (p
< 0.05) and $rbfox1^{+/19del}$ ($p < 0.05$) siblings (Figure 1H). As there was a significant difference in basal
locomotion, we also examined the amplitude of response on transition from dark to light
normalizing the data against baseline, as reported previously[34]. We observed similar results as in
absence of normalisation: significant main effect of time [Effect of time: $\chi2(118)$ = 1249.918, $p <$
0.001] and genotype [Effect of genotype: $\chi2(2)$ = 37.716, $p < 0.001$], and a significant two-way
interaction between time and genotype [Effect of time by genotype: $\chi2(236)$ = 4532.966, $p < 0.001$]
whereby $rbfox1^{19del/19del}$ larvae startled more than $rbfox1^{+/+}$ ($p < 0.0001$) and $rbfox1^{+/19del}$ ($p < 0.001$)
siblings.
193
Anxiety-like behaviour in adult animals was assessed using the NTD assay. When introduced to a
novel tank, zebrafish will first dive to the bottom of the tank, to seek protection, and then gradually
increase their swimming over time[35]. Over the entire duration of the NTD assay, we found no
significant differences between $rbfox1$ genotypes($p > 0.05$). However, during the first minute of the
assay, we found a significant two-way interaction between genotype and the time spent at the
bottom of the tank [Effect of genotype by proportion at the bottom tank: $\chi2(10)$ =22.3333, $p <$
0.05], such that $rbfox1^{19del/19del}$ fish spent less time in the bottom than $rbfox1^{+/+}$ fish ($p < 0.001$)
(Supplementary Figure 2E). In line with previous findings[2], we observed no significant differences in
distance travelled ($p > 0.05$) (Supplementary Figure 3F), nor in the number of the transitions to the
top area of the tank between $rbfox1$ genotypes ($p > 0.05$) (Supplementary Figure 3E).
204
205   Thus, our data showed that loss of $rbfox1$ resulted in behavioural changes in zebrafish that are
206   relevant to core domains often altered in human psychiatric disorders (such as hyperactivity,
207   impulsivity, reduced habituation and heightened arousal) and suggest that adaptation occurs as the
208   animal develops.

rbfox1 is expressed in the developing neural and cardiac tissues and in adult brain regions involved in stress, social and emotional behaviours, reward and learning

210
211
212

Consistent with our behavioural results and as seen previously[2,3], when we assessed *rbfox1* mRNA
distribution in *rbfox1⁺/⁺* larvae and adults (Tübingen strain), we found that *rbfox1* was expressed in
regions of the brain involved in the response to stress, in social and emotional behaviour, and in
reward and learning (Figure 1A-B) in agreement with data in rodents[36] and humans[37].

In wild type larvae *rbfox1* is expressed in the spinal cord and hindbrain lateral neurons (Figure 1A, 28
hpf), whereas at later developmental stages, *rbfox1* expression was detected in the mid- and
hindbrain (Figure 1A, 2-5 dpf) and in the heart (Figure 1A, 5dpf) as described previously[38].

In adult fish, *rbfox1* is expressed along the entire whole rostro-caudal brain axis. In the forebrain,
rbfox1 was detected in the glomerular (GL), internal (ICL), external (ECL) cellular layers of the
olfactory bulbs (Figure 1B a, a'). More caudally, *rbfox1* expression was detected in the medial zone of
the dorsal telencephalic area (Dm), and in the dorsal (Vd), lateral (Vl) and ventral (Vv) nuclei of the
ventral telencephalic area (Figure 1B b). In the diencephalon, *rbfox1* expression was detected in the
ventral habenular nucleus (HaV), in the anterior (A) and ventromedial (VM) thalamic nuclei (Figure
1B c). In the midbrain, *rbfox1* was detected in the posterior part of the parvocellular preoptic
nucleus (PPp, Figure 1B d), in the ventral zone of the periventricular hypothalamus (Hv, Figure 1B d,
e'), in the periventricular gray zone (PGZ,) and in the central zone (CZ) of the optic tectum (Figure 1B
e'), in the torus longitudinalis (TL) (Figure 1B e, e', g), in the periventricular nucleus of posterior
tuberculum (TPp), in the anterior tuberal nucleus (ATN, Figure 2B e, e''), in the posterior tuberal
nucleus of the hypothalamus (PTN, Figure 1B f) and in the paraventricular organ (PVO, Figure 1B f).
In the hindbrain, *rbfox1* was observed in the lateral division of the valvula cerebelli (Val) (Figure 1B
g).

rbfox1 mutant larvae show elevated cortisol levels and altered expression of crhb, nr3c2, bdnf and trkb2

235
236

As RBFOX1 has been linked to regulation of HPA axis and *BDNF/TRKB* gene expression[4,15,16], we
examined the expression of components of the HPI axis and of *bdnf/trkb2*. In parallel, we measured
whole-body cortisol levels, in resting physiological conditions, at both larval and adult stages to
assess physiological outcomes. Since we did not observe significant differences in behaviour
between wild type and heterozygous animals, we employed wild type and homozygous animals only.

We performed qPCR experiments in 5dpf zebrafish larvae to assess changes in the expression levels
of the HPI axis markers corticotropin releasing hormone (*crhb*), mineralocorticoid receptor (*nr3c2*),
glucocorticoid receptor (*nr3c1*), corticotropin releasing hormone receptor 1 (*crhr1*), corticotropin
releasing hormone receptor 2 (*crhr2*), mineralocorticoid receptor 1 (*mc1r*), proopiomelanocortin a
(*pomca*), Krüppel-like factor 9 (*klf9*), as well as of *bdnf* and *trkb2*. In teleosts, the duplication of the
genome gave rise to two *CRH* genes, *crha* and *crhb*, and two *TRKB* genes, *trkb1* and *trkb2*[39,40]. Among
these, *crhb* and *trkb2* are most commonly referred to as the functional orthologues of mammalian
CRH and *TRKB* respectively, whereas *crha* and *trkb1* function appear more divergent and remain less
well understood [39,40]. Zebrafish also possess two *pomc* paralogues (*pomca* and *pomcb*) but only
pomca is functionally relevant for the activation of the HPI axis[41], and six melanocortin receptors
(*mc1r, mc2r, mc3r, mc4r, mc5ra, mc5rb*). Among these, we selected *mc1r* as representative of
melanocortin signalling. Then, unlike other teleosts, zebrafish possess a single copy of the *MR*

(*nr3c2*), *GR* (*nr3c1*) and *KLF9* (*klf9*) genes, and a single copy of the *BDNF* gene (*bdnf*)[40,42]. In
mammals, several *TRKB* splicing isoforms are present, but the most abundant ones are the full-
length (TRKB.FL/TK+) and the truncated (TRKB.T1/TK-), this latter lacking the catalytic tyrosine kinase
(TK) domain[43]. As in zebrafish the presence of both *trkb2* full-length and truncated forms has been
demonstrated[44], here we used a pair of *trkb2 TK+/TK-* common primers, and another pair targeting
only *TK+* to distinguish effects on the expression of the two isoforms.

In *rbfox1*[19del/19del] larvae, inwe observed a significant up-regulation of *bdnf* (p < 0.05) and *TK+* (p <
0.05), and a significant down-regulation of *TK-* (p < 0.05) expression levels (Figure 3A). Regarding HPI
axis, in *rbfox1*[19del/19del] larvae we found a significant up-regulation of *crhb* (p < 0.0001) and a
significant up-regulation of *nr3c2* (p < 0.01) (Figure 2B). We observed no significant changes in the
expression levels of the other genes examined (p > 0.05) (Figure 3B).

To assess *rbfox1* LoF effects on zebrafish larvae during early developmental stages, we performed
crhb, *bdnf* and *trkb2* qPCR experiments also in 3dpf larvae. Similarly to 5dpf larvae, in 3dpf
rbfox1[19del/19del] larvae we observed a significant up-regulation of *crhb* (p < 0.01), *bdnf* (p < 0.05) and
TK+ (p < 0.05), and a significant down-regulation of *TK-* (p < 0.05) (Supplementary Figure 4A).

We measured whole-body cortisol levels under resting physiological conditions in both larvae and
adult zebrafish to evaluate the physiological impact of *rbfox1* loss and observed changes in *crhb*
expression. We found a significant increase in cortisol levels in *rbfox1*[19del/19del] larvae (p < 0.05)
(Figure 3C). In contrast, we observed no significant differences in adult fish (p > 0.05) (Figure
3C).These findings showed that *rbfox1* LoF led to alterations in stress-related gene expression and
cortisol levels in larvae, in line with our behavioural data and implicating *RBFOX1* as a critical
regulator of stress response mechanisms.

Pharmacological treatment targeting TrkB signalling restores *crhb* and *nr3c2* gene expression and
disrupted hyperactivity, startle response and habituation in *rbfox1* LoF larvae

To assess whether molecular and behavioural changes seen in *rbfox1* LoF larvae were mediated by
bdnf/trkb2 signalling, we repeated qPCRs and behavioural experiments in 5dpf zebrafish larvae
following chronic exposure (from 5 h to 5dpf) to the TRKB agonist 7,8-DHF or antagonist ANA-12. As
in our line *rbfox1* expression was knocked down throughout development, we initiated TRKB drug
treatment from gastrulation.

In *rbfox1*[+/+] larvae, neither TRKB agonist 7,8-DHF or antagonist ANA-12 had a significant effect on
crhb and *nr3c2* gene expression (p > 0.05) (Figure 3D, E). In *rbfox1*[19del/19del] larvae, chronic exposure
to either TRKB agonist or antagonist significantly reduced *crhb* and *nr3c2* expression levels relative
to untreated mutant larvae (p < 0.001), restoring gene expression to levels comparable with wild
type controls (p > 0.05) (Figure 3D, E). Following identification and removal of outliers (1 outlier in
the *rbfox1*[+/+] + ANA-12 group, 1 outlier in the *rbfox1*[+/+] + 7,8-DHF group and 1 outlier in the
rbfox1[19del/19del]+ 7,8-DHF group), we observed a significant down-regulation of *nr3c2* expression
levels in *rbfox1*[19del/19del] + ANA-12 versus *rbfox1*[+/+] un-exposed controls (p < 0.01) (Figure 2D) and a
significant up-regulation of *nr3c2* expression levels in *rbfox1*[19del/19del]+ 7,8-DHF versus both exposed
(p < 0.01) and un-exposed (p < 0.05) controls (Figure 3D, EE). Interestingly, while having the same
effect on *crhb* and *nr3c3* expression, ANA-12 and 7,8-DHF had different outcomes on *bdnf*
expression. Chronic treatment with ANA-12 reduced *bdnf* expression in *rbfox1*[+/+] (p < 0.05) but had
no effect on *rbfox1*[19del/19del]. On the other hand, chronic treatment with 7,8-DHF increased bdnf levels
in both *rbfox1*[+/+] and *rbfox1*[19del/19del] larvae (Supplementary Figure C).

Then, we examined the effects of ANA-12 and 7,8-DHF on behaviour to assess which behavioural
alterations observed in *rbfox1* mutant were influenced by bdnf/trkb2 signalling. In presence of 7,8-

DHF we found a significant two-way interaction between genotype and drug on distance travelled
[Effect of genotype by drug: χ2(2) = 47.3782, p < 0.05] and on swimming velocity [Effect of genotype
by drug: χ2(2) = 31.3951, p < 0.05] such that treatment with 7,8-DHF restored hyperactive
behaviour observed in *rbfox1^19del/19del* larvae (p < 0.01) (Figure 3F, G). Further, following chronic
exposure to 7,8-DHF there was no significant difference between *rbfox1^19del/19del* and *rbfox1^+/+* larvae
in response to startle with white light (Figure 3H) or in the proportion of responders to the acoustic
startle (p > 0.05) (Figure 3 I). ANA-12 had no significant effects on *rbfox1^19del/19del* behavioural
alterations.

As the hypothalamus is the primary player in the stress response and Rbfox1 had only previously
been shown to affect TRKB expression within the hippocampus[16], we used ISH to determine whether
trkb2 mRNA expression was reduced within the hypothalamic area. In agreement with previous
findings[45], in *rbfox1^+/+* we found that *trkb2* is widely expressed in the brain of 5dpf larvae, whereas in
rbfox1^19del/19del larvae, consistently with our qPCR experiments, we found a significant overall
reduction of *trkb2* mRNA in the whole brain (Figure 3J-L), including in the hypothalamus (p < 0.01)
(Figure 3M, N, P, Supplementary Figure 5).Our findings showed that the dysregulation in *crhb* and
nr3c2 expression caused by *rbfox1* LoF was prevented by TRKB agonists and antagonists. Further, we
showed that hyperactivity, startle hypersensitivity and heightened proportion of responders to the
acoustic startles observed in *rbfox1* mutants are at least partially mediated by disrupted bdnf/trkb2
signalling.

rbfox1 LoF alters *pac1a* expression levels of zebrafish larvae

As RBFOX proteins have been shown to be able to regulate *Pac1* alternative splicing to include the
hop cassette[7] and PAC1 has been shown to regulate *Bdnf* transcription and potentiate TRKB
activity[17], we examined expression levels of *pac1a,* the zebrafish homologue of the mammalian
PAC1, in *rbfox1^+/+* and *rbfox1^19del/19del* 5dpf larvae by qPCR.
323
In mammals there are several PAC1 isoforms and their role in the regulation of stress is poorly
understood[9]. In the brain, the predominant isoforms are *PAC1-hop* (long isoform) and *PAC1-short*[9].
Zebrafish possess two *pac1* genes, *pac1a* and *pac1b*, but only *pac1a* contains the *hop* cassette[9].
PAC1-short enhances *CRH* transcription, while *PAC1-hop* reduces *CRH* synthesis during late stress
recovery phase[4]. We measured expression levels of both *pac1a-short* and *-hop* isoforms and we
observed no differences in *pac1a-short* expression, but we found a significant up-regulation of
pac1a-hop in *rbfox1^19del/19del* mutant larvae (p < 0.05) (Figure 3Q).

This finding shows that elevated pac1a-hop/Pac1a-hop alone is not sufficient to counteract *crhb*
increase and strengthens previous data[7] suggesting that RBFOX1 is not the main regulator (or at
least not the sole regulator of) *PAC1* alternative splicing..

rbfox1 mutants undergo adaptive mechanisms and allostatic overload during development

The HPA axis possesses a vital role in the maintenance of allostasis, the process by which the body
achieves stability in response to stress or environmental challenges. Dysregulation of the HPA axis
often leads to disrupted allostasis during later life, also termed as allostatic load (i.e., the
physiological consequence resulting from the cumulative "wear and tear" of the body in response to
chronic stress)[22]. As we observed dysregulation of the HPI axis gene expression in *rbfox1* mutant
larvae and altered behavioural responses in *rbfox1* mutant larvae but not in adults, we assessed
expression of HPI axis components in adult zebrafish, in the presence and absence of acute stress
(NTD), to explore the possibility of adaptation that may contribute to differences in allostatic load.

We first measured *crhb, nr3c2, nr3c1, bdnf* and *trkb2* mRNA expression levels. In physiological
resting conditions, we observed no differences in HPI nor *bdnf/trkb2* expression levels (p > 0.05)

(Figure 4A, B). However, after stress exposure, regarding the HPI axis, in *rbfox1[19del/19del]* adults we
observed the same dysregulation seen in *rbfox1* LoF larvae: we found significant up-regulation of
crhb ($p < 0.001$) and *mr* ($p < 0.0001$) and no changes in *nr3c1* expression levels ($p > 0.05$) (Figure 4A),
and significant up-regulation of *bdnf* ($p < 0.01$) and *TK+* ($p < 0.05$), and significant down-regulation of
TK- ($p < 0.05$) (Figure 4B, C).

Excess glucocorticoid (GC) exposure during early developmental stages or early life stress (ELS) can
lead to vulnerability to allostatic overload in the long-term[19,22]. One of the effects caused by GC- or
ELS-induced allostatic overload is reduced cell proliferation during adulthood[19]. This effect has been
suggested to be developmentally dynamic, since it is often preceded by increased cell proliferation
during early stages[19]. Further, BDNF signalling has been shown to regulate neural stem cell
proliferation through TK-, suggesting that *rbfox1* LoF fish showing altered levels of *TK-* expression,
may show altered proliferation across the life course[46]. We therefore assessed the rate of
proliferation at both larval and adult stages using qPCR for proliferating cell nuclear antigen (*pcna*).
We observed a significant up-regulation of *pcna* expression levels in 5dpf *rbfox1[19del/19del]* zebrafish
larvae ($p < 0.05$) and a significant down-regulation of *pcna* expression levels in the brains of adult
rbfox1[19del/19del] fish ($p < 0.01$) when compared to *rbfox1[+/+]* siblings (Figure 4D).

In adult zebrafish, other effects caused by GC-induced allostatic overload include reduced fertility
and survival rates[19]. Therefore, we assessed fertility and survival of *rbfox1[19del/19del]* zebrafish and
rbfox1[+/+] siblings. We found that both fertility ($p < 0.05$) and survival ($p < 0.001$) rates of
rbfox1[19del/19del] fish were significantly reduced compared to *rbfox1[+/+]* siblings (Figure 4E, F).

Consistent with our findings (disrupted HPI axis gene expression and increased cortisol in *rbfox1*
mutant larvae under resting physiological conditions and in *rbfox1* adult mutant only following stress
exposure), these results suggest that *rbfox1* mutants may engage compensatory mechanisms during
development similar to those triggered by early-life GC exposure. While these mechanisms may
transiently restore homeostasis, they likely lead to enduring alterations in stress responsivity and
neurodevelopmental trajectories.

DISCUSSION

In this study we generated a CRISPR-Cas9 LoF *rbfox1* zebrafish line (*rbfox1[19del]*) to investigate the
mechanisms by which *RBFOX1* LoF increases susceptibility to psychiatric disorders.
374
Allostatic load has been linked with several cognitive disorders including depression, schizophrenia,
anxiety and PTSD[47]. HPA axis overactivity is a key factor in the onset of the allostatic load[22]. RBFOX1
has been shown to influence expression of stress-related genes such as PAC1 and BDNF/TRKB.
Therefore, alterations in RBFOX1 function may increase psychiatric disorder vulnerability through
alterations in stress response systems, such as the HPA axis, pre-disposing to allostatic overload
vulnerability in later life (Figure 5).
381
Similarly to *Rbfox1* deficient mice[6] and as seen in previous studies employing *rbfox1* mutant
zebrafish[2,23], we found that *rbfox1[19del/19del]* mutants were hyperactive and impulsive. These findings
are consistent with studies implicating common and rare *RBFOX1* genetic variants as risk factors for
psychiatric disorders like ASD, ADHD, schizophrenia and anxiety/stress disorders[5,6,48]. At larval stages
rbfox1[19del/19del] were hyperactive and had increased burst swimming, indicative of increased
impulsivity. In the response and habituation to acoustic startle assay, *rbfox1[19del/19del]* larvae startled
more and had a higher response rate over time compared to *rbfox1[+/+]* siblings, consistent with
previous findings in *rbfox1* mutant zebrafish[30]. In a forced light-dark assay, although baseline
hyperactivity of *rbfox1[19del/19del]* larvae makes interpretation difficult, *rbfox1[19del/19del]* larvae showed an
increased startle response upon dark to light transition, consistent with heightened anxiety. These

results suggest that *rbfox1* LoF leads to heightened arousal, aligning with studies linking *RBFOX1* to mood and anxiety disorders, including PTSD[49]. Interestingly, *rbfox1*[19del/19del] adults showed reduced bottom dwelling only during the first minute of the novel tank diving assay (an adult measure of anxiety-like behaviour) while there were no significant differences during the remaining 4 minutes of the assay This latter result suggests behavioural adaptation possibly coupled with increased impulsivity or impaired threat assessment (decreased bottom dwelling during the first minute of assay. This aligns with the broader *rbfox1*[19del/19del] behavioural profile (hyperactivity, heightened startle and poor habituation at larval stages) indicative of dysregulated arousal rather than reduced anxiety. Similarly, in the 5-CSRTT we found that *rbfox1*[19del/19del] were more impulsive than *rbfox1*[+/+] siblings, consistent with *RBFOX1* variants being associated with impulsivity and related disorders such as ADHD and ASD[6].

In *rbfox1* LoF larvae, under resting physiological conditions, we observed increased cortisol levels and dysregulation of *crhb* and *nr3c2*, both key components of fish HPI axis, which is analogous to the HPA axis in mammals. The HPA axis is crucial for managing stress, with CRH triggering cortisol secretion, affecting metabolism, immunity, and behaviour[50,51]. In zebrafish larvae under resting physiological conditions, we found increased cortisol, *crhb* and *nr3c2*, with no changes in the expression levels of the other HPI axis genes assessed (*nr3c1, crhr1, crhr2, mc1r, pomca, kf19*). In adults, although resting levels were unchanged, *rbfox1* LoF showed an exaggerated molecular response to stress. Notably, both CRH and MR have important roles in the brain beyond their canonical endocrine functions via the HPA axis. CRH regulates anxiety and arousal acting as a neuromodulator in limbic regions such as the amygdala and hippocampus[52]. Similarly, hippocampal MRs play a key role in early stress responses by supporting cognitive appraisal of novel situations and promoting behavioural flexibility, thus shaping the initial reaction to stress[53]. Elevated cortisol and *crhb* levels as seen in *rbfox1*[19del/19del] larvae under resting physiological conditions, suggest a chronic anxiety state. Increased cortisol, CRH and MR levels are linked to mood disorders like depression and PTSD[54–57]. These findings support a role for *RBFOX1* in *CRH* regulation and suggest that *RBFOX1* LoF may contribute to mood disorders and stress resilience via HPA axis dysregulation and/or non-canonical brain-specific effects of CRH and MR.

BDNF and TRKB are also key regulators of the stress response[58]. BDNF levels often increase during stress to promote neuronal survival and plasticity, buffering negative effects of stress on the brain via TRKB[10,11,13,58–60]. Dysfunction of BDNF/TRKB signalling is linked to several stress-related disorders[13]. The increased *bdnf* expression we observed in *rbfox1*[19del/19del] larvae may suggest a compensatory response to cope with *crhb* increase and maintain homeostasis. Given BDNF's established role in modulating neuronal plasticity and stress adaptation, its upregulation may serve to stabilise or buffer hypothalamic circuits disrupted by excessive *crhb* transcription[58]. This response could represent an intrinsic attempt to mitigate possible overactivation of the HPI axis or the reduced activity within the trkb2 signalling (a suggestion supported by finding that incubation in the TRKB agonist 7,8-DHF rescues the behaviour as well as the expression of *crhb*) and maintain functional equilibrium in the face of impaired regulatory input caused by *rbfox1* loss. Alternatively, the observed increase in *bdnf* expression could result from either dysregulation of BDNF repressors or an imbalance in TRKB isoform expression caused by RBFOX1 loss, where upregulation of the full-length isoform may enhance BDNF expression through a positive feedback mechanism[40,61]. At 3 and 5dpf, in *rbfox1*[+/+] larvae we detected both *trkb2* truncated (*TK-*) and full-length (*TK+*) forms, and in *rbfox1*[19del/19del] larvae we observed a significant down-regulation of *TK-* and a significant up-regulation of *TK+*, which is predicted to lead to an increased activation of TRKB signalling, consistent with the increased *crhb* expression as seen here[62]. Studies in rodents showed that *Rbfox1* up-regulation led to increased TrkB.T1 (TK-) expression in the hippocampus and no changes were observed following *Rbfox1* deletion[16]. In contrast, we observed down-regulation of *trkb2 TK-* expression in *rbfox1*[19del/19del] larvae in resting physiological conditions. The difference between these

studies may be due to i) differences in the tissue examined (zebrafish larvae whole head versus
rodent hippocampus), ii) the developmental stage of the animals (larval versus adult stage), iii) the
genotype of the animals studied ($rbfox1^{19del/19del}$ in our study versus $Rbfox1^{+/-}$ in the study in rodents),
or iv) presence of compensatory mechanisms (in the hippocampal mouse model $Rbfox1$ knockdown
resulted in up-regulation of $Rbfox2$). As for this latter hypothesis, as in previous studies in zebrafish[2],
we did not see any changes in $rbfox2$ expression upon $rbfox1$ LoF (Supplementary Figure4)
suggesting that this compensatory mechanism does not occur in larval fish.
450
Consistent with the possibility that $RBFOX1$ LoF may contribute to dysregulation of the HPA axis
through a direct or indirect effect on BDNF/TRKB pathway, chronic exposure to TRKB agonist (7,8-
DHF) or antagonist (ANA-12) restored HPI axis gene expression to that seen in $rbfox1^{+/+}$ larvae. ANA-
12 is a selective non-competitive antagonist of TRKB exerting central TRKB blockade and producing
rapid and long-lasting anxiolytic and antidepressant effects[63]. Mice treated with ANA-12 showed
reduced anxiety-like behaviour[63] and in stressed rats, ANA-12 blocked Crh increase in the
hypothalamus and amygdala[64]. 7,8-DHF is a potent selective agonist of TRKB used in the treatment
of several disorders, including depression and schizophrenia, and has been shown to enhance
memory consolidation and emotional learning in healthy rodents[65]. In a similar fashion to the effects
of ANA-12 seen in rats (blocking Crh increase)[57], after chronic treatment with ANA-12, we observed
that the expression of HPI axis genes in $rbfox1^{19del/19del}$ larvae was restored to $rbfox1^{+/+}$ levels. [66]
Interestingly, ANA-12, despite restoring $crhb$ and $nr3c2$ levels to wild type, did not rescue
$rbfox1^{19del/19del}$ larvae behaviour, indicating that blockade of TRKB can normalise some aspects of HPI
axis output but is insufficient to correct the behavioural phenotype (perhaps because it further
disrupts necessary TRKB-mediated signalling in the brain). In contrast, 7,8-DHF, normalises $crhb$ and
$nr3c2$ expression and rescues some behavioural aspects altered in $rbfox1^{19del/19del}$ larvae (i.e.,
hyperactivity, startle response to white light stimulus and proportion of fish responding to acoustic
stimuli), suggesting that enhancing TRKB signalling through the full-length receptor may restore
functional signalling balance and downstream pathways (e.g., synaptic plasticity, habituation
mechanisms). Finally, the fact that 7,8-DHF increases $bdnf$ expression in $rbfox1^{19del/19del}$ larvae while
ANA-12 has no effect suggests a potential positive feedback loop whereby TRKB activation enhances
BDNF expression, helping restore adaptive neuroendocrine and behavioral responses. Future studies
employing acute or temporally restricted treatments (e.g., 3–5 dpf), or employing different TRKB
agonist/antagonist, will help delineate the specificity of TRKB signalling effects. Interestingly,
$rbfox1^{19del/19del}$ larvae were differentially sensitive to the effect of TRKB drugs on $nr3c2$ expression,
most likely due to the unbalanced $TK+/TK-$ levels. These findings warrant further investigations. It is
of note that neither TRKB agonist nor antagonist had a significant effect on $crhb$ nor $n3cr2$
expression in $rbfox1^{+/+}$ larvae. One possible explanation is that, as the larvae were not exposed to
any stressors, any effect on Trkb2 signalling had limited effect on HPI axis activity.
480
In $rbfox1^{19del/19del}$ larvae, we also observed up-regulation of $pac1a-hop$ with no changes in $pac1a-$
$short$ levels. Previous studies in zebrafish have shown that Pac1-short gain of function caused
persistent crh increase, while overexpression of Pac1-hop prevented stress-induced crh transcription
activation[4]. These previous studies also suggest that RBFOX1 regulates the alternative splicing of
$PAC1$ promoting the formation of $PAC1-hop$[4]. However, the increase in $pac1a-hop$ seen in our study
argues against this suggestion and suggests an adaptive response to the increased expression of crh
either through $rbfox2$[7] (whose expression was not altered by $rbfox1$ LoF (Supplementary Figure4)) or
through an unknown mechanism. Elevated $pac1a-hop$/Pac1a-hop alone might be not sufficient to
counteract $crhb$ upregulation, given the concurrent increase in $bdnf$ levels and the shift in $trkb2$
isoform expression (upregulation of the excitatory long form and downregulation of the inhibitory
short form). This interpretation is further supported by the decreased in $crhb$ levels observed
following chronic TRKB modulation. Another option is that the upregulation of $pac1a-hop$ mRNA
may not reflect increased functional Pac1a-hop receptor activity, particularly in the context of

disrupted splicing regulation following *rbfox1* loss, suggesting that RBFOX1 might be involved in
PAC1 mRNA stability rather than alternative splicing.
496
In contrast to larvae, differences in HPI axis and *bdnf/trkb2* gene expression and cortisol levels were
not seen in adult animals under resting conditions. However, *rbfox1*$^{19del/19del}$ adults showed increased
crhb and *nr3c2* on challenge with a stressor, aligning with the previous suggestion that Rbfox1 is
required for the termination of the acute endocrine stress response[4] and that chronic stress during
development leads to adult HPI axis adaptation, often at a cost in later life (e.g., altered stress
responsivity, reduced fertility and survival, decreased *pcna* expression)[19,67]. This interpretation is
consistent with previous work by Eachus et al., which demonstrated that early-life exposure to
excess GC can lead to long-term changes in stress-regulatory circuits and adult behavioural
dysfunction[19]. Supporting this, *rbfox1* LoF adults show heightened *crhb*, *nr3c2* and *bdnf* expression,
and imbalance in *trkb2* isoform expression, specifically in response to a stressor (the novel tank
diving assay). Together, these findings suggest that while baseline homeostasis may be restored
through developmental adaptation, stress responsivity remains dysregulated, pointing to an
allostatic shift toward a passive coping style for *rbfox1* mutants[68,69]. Passive (cautious, risk-averse,
high CRH) and active (bold, exploratory, low CRH) coping styles refer to behavioural strategies that
animals employ to respond to stress and that are underpinned by distinct neuroendocrine pathways,
providing a valuable framework for interpreting stress susceptibility, resilience and the behavioral
consequences of genetic mutations[68,69]. The shift toward a passive coping style of *rbfox1* mutants
support a role for RBFOX1 in shaping stress responsivity and behavioural strategies. In *rbfox1*$^{19del/19del}$
adults we observed also reduced fertility and reduced survival rates, two long-term effects often
associated with excess GC-exposure/early life stress[19]. Another long-term effect of excess GC-
exposure/early life stress is reduced neural stem cell proliferation in adult animals, often
accompanied by increased proliferation at early stages[19]. Our findings that *rbfox1*$^{19del/19del}$ showed
increased *pcna* levels at larval staged but reduced *pcna* levels during adulthood, corroborate this
latter hypothesis of a dynamic developmental effect of early life stress. Consistent with the *crhb*
increase observed in stressed *rbfox1*$^{19del/19del}$ adults, and as seen in larvae, after acute stress,
rbfox1$^{19del/19del}$ adults also showed unbalanced *trkb2* expression, with up-regulation of *TK+* and down-
regulation of *TK-*.
524
In conclusion, given the conservation between fish HPI axis and the mammalian HPA axis, our data
unveils a pivotal new finding in *CRH* regulation, revealing an interplay between RBFOX1 and
BDNF/TRKB in the context of chronic stress and stress resilience and suggests that RBFOX1 plays a
crucial role in adaptive stress mechanisms. In response to stressful challenges, RBFOX1 functions as a
"switch" regulating the balance between short/long isoforms of TRKB receptors: during the stress
recovery phase, RBFOX1 switches the balance towards *TRKB.T1/TK-* isoform, decreasing *CRH* levels
and turning off the HPA axis (Figure 5). In *rbfox1*$^{19del/19del}$, this regulatory switch appears disrupted
(increased cortisol, *crhb*, *nr3c2*, *bdnf*, and full-length *trkb* expression alongside reduced truncated
trkb) suggesting a failure to terminate the stress response. Behaviourally, mutants exhibited
hyperactivity, impulsivity, heightened arousal, and reduced habituation, phenotypes that parallel
core symptoms seen in several human psychiatric disorders. Pharmacological experiments support a
functional consequence of this dysregulation. Although both ANA-12 (TRKB antagonist) and 7,8-DHF
(TRKB agonist) normalised *crhb* and *nr3c2* levels, only 7,8-DHF rescued behavioural abnormalities.
These findings suggest that restoring balanced TRKB signalling (by selectively enhancing the activity
of residual inhibitory TRKB.T1 receptors without compromising the beneficial effects mediated by
excitatory TRKB.FL receptors) is critical for functional recovery. This supports the idea that an
imbalance in TRKB isoform expression, rather than overall receptor activity, may underlie the
observed behavioural pathology. The differential effects of these compounds also point to a possible

role for positive feedback in BDNF expression via TRKB-FL signalling, which may be essential for
behavioral homeostasis. Together these results suggest that *RBFOX1* contributes to the liability of
psychiatric disorders through CRH, MR and BDNF/TRKB dysregulation, which leads to disrupted
development and vulnerability to allostatic overload in later life. The suite of behavioural effects
seen in *rbfox1* LoF animals coupled with exaggerated response on exposure to a stressor are
reminiscent of reported traits in passive versus active coping styles[68] suggesting that RBFOX1 activity
may also play a key role in individual differences in coping strategies. Although we show a key role
for RBFOX1 in regulation of the HPA axis, consistent with this being a primary mechanism by which
RBFOX1 variants pre-disposes to psychiatric disorders, RBFOX1 regulates the splicing of a wide range
of additional genes[15] which may also play a part of disease vulnerability.
553

554 **METHODS**
555

Animal maintenance

All fish were maintained in a recirculating system (Tecniplast, UK) with a 14h:10h light/dark cycle
and a constant temperature of 28°C. Fish were fed with ZM-400 fry food (Zebrafish Management
Ltd.) in the morning and brine shrimps in the afternoon. Breeding was set up in the evening, in
sloping breeding tanks (Tecniplast) provided with dividers for timed mating. The following morning,
dividers were removed to allow spawning. Eggs were collected in Petri dishes (max 50 eggs/dish).
Infertile eggs were removed, and fertile ones were incubated at 28°C. Petri dishes were checked
daily to ensure consistent developmental stage across groups. If reared, larvae were moved to the
recirculating system at 5 days post fertilization (dpf) and fed as stated above.

All procedures were carried out under license in accordance with the Animals (Scientific Procedures)
Act, 1986 and under guidance from the Local Animal Welfare and Ethical Review Board at Queen
Mary University of London.

Although our previous studies examined two distinct *rbfox1* mutant lines (*rbfox1*^{del19} vs
rbfox1^{sa15940})[2], we chose to focus on the *rbfox1*^{del19} line for the current study due to its more
pronounced phenotype.

Generation of *rbfox1* loss of function zebrafish line

The zebrafish *rbfox1* loss of function (LoF) mutant line was generated as described previously using
Tübingen strain as background [71]. CRISPR RNA (crRNA) (Merck) was designed to target *rbfox1* exon 2
(**CCC**<u>AGTTCGC</u>TCCCCCTCAGAAC, PAM sequence in bold, Mwol recognition site underlined). A 3 μL
injection mix containing 1 μL (F_c 83 ng/μL) crRNA, 1 μL (F_c 83 ng/μL) tracrRNA (Merck,
#TRACRRNA05N), 1 μL (F_c 1.67 μM) and 1 μL Cas9 protein (New England Biolabs, #M0646) was
freshly prepared on the morning of the injection procedure. Then, 1 nL of the injection mix was
injected into one-cell stage zebrafish embryos (~ 100-150 embryos). Injection efficacy was assessed
at 24 hours post fertilization (hpf) by polymerase chain reaction (PCR) from genomic DNA
(*rbfox1*_Forward, 5ꞌ-TAATCAAGACGCCCCAGCAC–3ꞌ; *rbfox1*_Reverse, 5ꞌ-
GTACTCAGCAGGAATGCCGT- 3ꞌ) followed by Mwol (New England Biolabs, #R0573S) restriction
enzyme digestion. Successful injections will introduce indel mutations disrupting the recognition site
of the restriction enzyme, preventing this latter from cutting the PCR amplicon. Once reached sexual
maturity (at ~ 3 months of age) injected fish (F_0) were outcrossed with wild type to generate F_1
embryos. The progeny may carry different mutations due to the mosaic nature of the F_0 parents. F_1
fish were therefore screened for mutations leading to premature termination codon (PTC) via
cloning into pGEM-T Easy vector (Promega, #A1360), followed by transformation, colony PCR, DNA

purification (Monarch® Plasmid Miniprep Kit, New England Biolabs, #T1010) and sequencing (Source
BioScience PLC). Quantitative real time PCR (qPCR) was used to confirm reduction of *rbfox1* mRNA
expression.
591
592 **Fish breeding and genotyping**
593
For adult experiments, mixed sexes were used. When possible, animals were associated with an
identification number and genotype was assigned after data analysis. For larval experiments, fish
were generated by heterozygous in-cross and genotyped prior to data analysis. Genomic DNA was
extracted from fins and using the HotSHOT method. Briefly, samples were incubated at 95 °C in 50
mM NaOH for 30 min, followed by 1 min at 10 °C. Reaction was stopped using 1M Tris HCl (1/10 of
the initial NaOH volume), pH 8.00. Genotyping primers were the same as the ones used to identify
founder carriers.
601
602 **Drug Treatment**
603
604 Zebrafish embryos of each genotype (*rbfox1*$^{+/+}$ and *rbfox1*$^{19del/19del}$) were treated from 5hpf to 5dpf
605 with 20 μM TRKB antagonist ANA-12 (abcam, Cat.ab146196), or 20 μM TRKB agonist 7,8-
606 dihydroxyflavone (7,8-DHF) (abcam #ab120996), or dimethylsulfoxide (DMSO, vehicle 0.01%) (Merck
607 #34869). Treatment was performed in Petri dishes (max 30 embryos x dish) and drugs or vehicle
608 were dissolved in fish water. Solutions were replaced daily.
609
Quantitative Real-Time PCR
Quantitative Real-time PCR (qPCR) of target RNA was performed on 5dpf *rbfox1* wild type (*rbfox1*$^{+/+}$)
and *rbfox1* homozygous (*rbfox1*$^{19del/19del}$) larvae, using Luna® Universal One-Step RT-qPCR Kit (New
England Biolabs #M3005) and a Bio-Rad 96-well qPCR machine (CFX96 Touch Real-Time PCR
Detection System). Total RNA was isolated using TRIzol reagent (Thermo Fisher Scientific) following
manufacturer's instructions. Briefly, after homogenization, RNA was isolated by precipitation, rinsed
in ethanol and resuspended in RNase free water. Total RNA was then quantified using BioDrop
(Biochrom Ltd.), and up to 1 μg was reverse transcribed to cDNA using the ProtoScript II First Strand
cDNA Synthesis Kit (New England Biolabs, #E6560) following manufacturer's instructions. The
resulting cDNA yield and quality were evaluated using BioDrop (Biochrom Ltd.). As for zebrafish
trkb2 TK+ and *TK-* isoforms quantification, Ct values resulting from the amplification of the *TK+*
specific product were subtracted from the Ct values resulting from the amplification of the *TK+/TK-*
common product. All reactions included 5 biological replicates and 3 technical replicates. For
experiments in larvae, each biological replicate consisted of 15 larval heads (eyes and jaw removed).
For experiments in adults, each biological replicate consisted of a single brain. Actin – β 2 (*actb2*) and
ribosomal protein L13a (*rpl13a*) were employed as reference genes. For assessing *rbfox1* mRNA
expression, *rbfox1* primers were designed upstream of the CRISPR deletion. Accession numbers,
primer sequences and amplification efficiencies for all the reference and target genes can be found
in Supplementary Table 1.
629
In situ **hybridization**
In situ hybridization (ISH) was carried out on whole mount zebrafish larvae and on larval (sagittal)
and adult brain (transverse) sections as described previously[2]. For *rbfox1* ISH, the original plasmid
used to generate the riboprobe was provided by Dr William Norton (University of Leicester). The
plasmid used to produce the *trkb2* riboprobe was generated in our laboratory (Forward primer: 5'-

GTTCGTGGAATGGCTTGCTG-3', Reverse primer: 5'-TCTGGCCCACGATGTTTTCA-3') using the pGEM®-T
Easy Vector System (Promega, #A1360) and *in-house* generated *E. coli* DH5α competent cells.
Riboprobes to identify *rbfox1* (NM_001005596) and *trkb2* (NM 01197161.2) mRNA were synthetized
by *in vitro* transcription (IVT) using MAXIscript™ SP6/T7 kit (Invitrogen by Thermo Fisher Scientific,
#AM1322), following manufacturer's instructions and using a DIG RNA Labeling Mix, 10× conc
(Roche, #11277073910) containing digoxigenin labeled uracil.

In situ hybridization on whole mount zebrafish larvae

In situ hybridization was carried out on 28hpf, 2- 3- 4- and 5dpf *rbfox1*$^{+/+}$ and *rbfox1*$^{19del/19del}$ larvae.
To prevent skin pigmentation, embryos were incubated in 0.2 mM 1-phenyl 2-thiourea (PTU) (Sigma,
#S527335) from 24hpf. When they reached the desired age, larvae were fixed in 4%
paraformaldehyde (PFA) (Merck, #158127) overnight (ON) at 4°C. The following day, larvae were
rinsed in 1x phosphate buffered saline (PBS) (Thermo Fisher Scientific, #18912014) supplemented
with Tween 20 (Sigma, #P1379) (0.05% v/v), dehydrated in ascending methanol series (25%, 50%,
70%, 80%, 90%, 100% methanol, 5 min each) and stored in 100% methanol at -20 °C. To perform ISH
experiments, larvae were rehydrated in descending methanol series (100%, 90%, 80%, 70%, 50%,
25% methanol), 5 min each, and washed in 1xPBS, 5 min. Larvae were permeabilized using
proteinase K (PK) (ITW Reagents, #A3830) (stock 20 µg/mL in 1xPBS) as follows: 28hpf larvae were
permeabilized in PK 1:2000 in 1xPBS for 20 min at room temperature (RT), older stages were
permeabilized in PK 1:1000 at 37°C for at least 30 min. Then, larvae were post fixed in 4% PFA for 20
min and washed in 1xPBS at RT, 5 x 5 min. Prehybridization was carried out in hybridization solution
(HB) containing 50% formamide, 5% saline sodium citrate buffer (SSC), 50 µg/mL heparin, 0.05
mg/mL yeast RNA, 0.1% Tween 20, and 0.92% citric acid at 68 °C for 2h. Thereafter, larvae were
incubated in HB containing *rbfox1* riboprobe (500 pg/µL), ON at 68°C. Post hybridization washes
were performed at 68°C with a gradient of 2xSSC and formamide (50%, 25% and 0% formamide), 10
min each, and then twice with 0.02xSSC, 30 min each. Subsequently, larvae were blocked in blocking
solution (BS) containing 10% normal sheep serum (Gibco, #16070096) and 2 µg/µL bovine serum
albumin, for 1h at RT. After blocking step, larvae were incubated in anti-digoxigenin Fab fragments
conjugated with alkaline phosphatase (Roche, #11093274910), 1:2000 in BS, 1h at RT and then ON
at 4°C. The following day, larvae were washed in 1xPBS, 6 x 15 min each, and then in alkaline
phosphatase (100 mM NaCl, 100 mM Tris HCl, 50 mM MgCl2, 0.1% Tween 20) (NTMT) buffer, 3 x 5
min each. The chromogenic reaction was carried out by incubating the larvae in BCIP/NBT solution
(Merck, #203790) in NTMT buffer, at RT in the dark, and were observed every 20 min until the signal
detection. After reaching the desired staining, larvae were washed in 1xPBS at RT, post fixed in 4%
PFA for 2h, cleared and stored in 80% glycerol at 4°C. For sagittal cryosections, larvae were
embedded in 1.5% low gelling temperature agarose (Scientific Laboratories Supplies, #A9414)
supplemented with 5% Sucrose in 1xPBS. Sections (5 µm) were collected on adhesive microscope
slides Superfrost® Plus Gold (Epredia).

In situ hybridization on sections

ISH was conducted on paraffin embedded *rbfox1*$^{+/+}$ and *rbfox1*$^{19del/19del}$ adult brains (transverse
sections) and 5dpf *rbfox1*$^{+/+}$ and *rbfox1*$^{19del/19del}$ zebrafish larvae (sagittal sections). Fish were culled by
an overdose of tricaine prior to head removal. Dissected adult brains and larvae were fixed in 4%
PFA in 1xPBS, ON at 4°C. Tissues were then rinsed in 1xPBS and dehydrated in ascending ethanol
series (15 min in each of 30%, 50%, 70%, 80%, 90%, 100% ethanol) and embedded in paraffin.
Transverse (adult brains) or sagittal (5 days old larvae) sections of 12 µm (adult brains) or 7 µm (5

days old larvae) thickness were cut using a microtome (Leica). To perform ISH, slides were de-waxed
in xylene (twice, 10 min each), rehydrated in descending ethanol series (2 x 5 min in absolute
ethanol, then 90%, 80%, 70%, 50% and 25% ethanol, 5 min each), and rinsed in 1xPBS for 5 min.
Then, sections were permeabilized using PK (0.05 µg/µL) for 8 min at RT, washed with 2 mg/mL
glycine twice (5 min each), post fixed in 4% PFA for 20 min and washed in 1xPBS at RT.
Prehybridization was carried out in HB, for 1h at 68°C. Thereafter, sections were incubated in HB
containing *rbfox1* riboprobe (500 pg/µL), ON at 68°C. Post hybridization washes were performed at
68°C twice for 20 min in 1xSSC, twice for 20 min in 0.2xSSC, and several washes were performed in
1xPBS, 5 min each at RT. Then, sections were blocked in BS for 30 min at RT and incubated in a
1:2000 dilution of anti-digoxigenin Fab fragments conjugated with alkaline phosphatase in BS, ON at
4°C. The following day, sections were washed in 1xPBS, 5 x 10 min each. The chromogenic reaction
was carried out by incubating the slides in BCIP/ NBT solution in NTMT buffer, at RT in the dark, and
were observed every 20 min until the signal detection. When the desired staining was obtained,
sections were washed in 1xPBS at RT, dehydrated in ascending ethanol series (25%, 50%, 70%, 80%,
90%, 100% ethanol, 5 min each), cleared in xylene (twice, 5 min each) and mounted with dibutyl
phthalate polystyrene xylene mounting medium (Sigma, #06522).

Image acquisition and processing

Pictures of whole mount ISH on zebrafish larvae were acquired by Leica MZ75 microscope. For ISH
on sections, pictures were acquired using a Leica DMRA2 upright epifluorescent microscope with
colour QIClick camera (Leica) and processed with Velocity 6.3.1 software (Quorum Technologies Inc).
Quantification of the ISH staining signal intensity was performed as described previously[72], using Fiji
software[73]. Adult anatomical structures were identified according to the Neuroanatomy of the
Zebrafish Brain by Wullimann [74].

Behavioral Assays

For larvae, all behavioural experiments were conducted on the progeny of a *rbfox1*[+/19del] in-cross.
Larvae were genotyped prior to data analysis. For adults, where the genotype of the animals was
known, fish were pseudorandomised across testing systems with all trials having an approx. equal
number of each genotype. The adult fish were weight- and age-matched, with approximately equal
numbers of both sexes. Sex was not treated as a biological variable in the statistical analyses.

Larval behavioural experiments

Patterns of locomotor activity of 5dpf *rbfox1*[+/+], *rbfox1*[+/19del] and *rbfox1*[19del/19del] mutant zebrafish
larvae were investigated as described previously[24,29,33]. Tests were conducted between 9 a.m. and 4
p.m. At 5 dpf, larvae were placed in individual wells of a 24-well plate. To reduce stress due to
manipulation, larvae were acclimatised for at least 1 h before testing. Then, plates were placed into
the DanioVision observation chamber (Noldus). Locomotion parameters such as distance travelled
and swimming velocity were recorded using EthoVision XT software (Noldus). Data were exported in
1 min and 1 sec time bins and analysed with R programming language[75]. For the hyperactivity assay,
larval basal locomotion was tracked for 15 min in dark conditions. Larval swimming burst was
assessed as described previously[24] and peaks were considered as the acceleration events when
larvae travelled > 5 mm in < 12 sec. The forced light-dark transition assay was performed as
described previously[76,77] with modifications: after an initial 5 min period of dark (baseline), larvae
were exposed to one light/dark cycle of 1 min light (Noldus white light device) followed by 5 min

dark. The response and habituation to acoustic startle stimuli was performed as described
previously[29]: after 10 min of baseline (no stimuli, dark conditions), larvae were subjected to 10
sound/vibration stimuli (Noldus tapping device) over 20 sec (2 sec intervals between each stimulus).

5-Choice Serial Reaction Time Task

We measured impulsive action using a zebrafish version of the 5-Choice Serial Reaction Time Task (5-
CSRTT)[25]. Adult *rbfox1*[+/+], *rbfox1*[+/19del] and *rbfox1*[19del/19del] zebrafish, 7 months old, mixed sexes, were
singly housed for a week prior to experiment and remained singly housed for the whole duration of
the assay (9 weeks). Fish were tested using the Zantiks AD units. Each unit was provided with a small
tank with five apertures and a food hopper insert. The five apertures created five different entry
points for the fish, acting like the five nose poke holes of the rodent version of the assay. The food
hopper was placed at the opposite side of the five apertures and formed an area for the fish to enter
and collect food reward. Below the testing tank there was an integrated screen, used to display
white light (stimulus) into the five apertures. Responses were detected when a fish entered these
apertures and recorded with an integrated camera placed at the top of the tank. The experiment
consisted of five training stages: i) habituation, ii) initiator training, iii) stimulus light training, iv) 5-
CSRTT/no delay, v) 5-CSRTT/variable-delay. Details of each stage are provided in Supplementary
Table 2.

Novel tank diving

Novel tank diving is a behavioural test to assess anxiety-like behaviour in adult fish. Response to
novel tank was assessed in *rbfox1*[+/+], *rbfox1*[+/19del] and *rbfox1*[19del/19del] 9 months old zebrafish, mixed-
sexes, as described previously[33]. Fish were singly housed for a week prior to performing the
experiment and acclimatized for at least 1 h in the behavioral room on the testing day. Behavioural
assays were conducted between 9 a.m. and 2 p.m. During the test, fish were individually placed into
a 1.5 L tank and their behaviour was tracked and recorded using EthoVision system (Noldus). Data
were exported in 1 min time bin and analysed as previously described[33]. Experimental groups were
randomised during testing. We analyzed three behaviours in response to the novel tank: i) time
spent in the bottom of the tank, ii) total distance traveled, and iii) number of transitions to the top–
bottom area of the tank.

Whole-body cortisol extraction

Larvae

Larval cortisol was extracted from pools of 12 larvae x sample using a modified protocol based on
Baiamonte et al. (2016)[35]. Briefly, larvae were snap-frozen upon collection and stored at -20 °C until
extraction was performed. On the day of extraction, frozen samples were thawed on ice in and
homogenised in 400 µl of ice-cold 1×PBS for 30 seconds using a Bioruptor (MP Biomedicals). Cortisol
was extracted by adding 500 µl of ethyl acetate (Sigma) to each homogenate. Samples were
vortexed for 30 seconds, centrifuged at 5000 rpm for 10 minutes and then snap frozen at −80 °C.
The organic phase was carefully decanted into glass vials (VWR). The extraction was repeated twice
and the combined organic phases were pooled. The pooled ethyl acetate extracts were evaporated
at 60 °C using a rotary evaporator (IKA). The dried cortisol was resuspended in 200 µl of 1xPBS,
vortexed for 30 seconds and stored at −20 °C until cortisol quantification using a human salivary
cortisol ELISA kit (Salimetrics).

Adults

Adult cortisol was extracted from 4 adult zebrafish whole-body. Fish were weighed and
homogenised in 2× volumes (w/v) of 1xPBS using a Bioruptor (MP Biomedicals). Cortisol was
extracted using the same ethyl acetate procedure as for larvae. However, after evaporation, to
prevent any interference with the assay, residual lipids were eliminated by partitioning the dried
extracts between 500µl 1xPBS and 500µl hexane (Sigma). The upper organic layer was discarded and
the aqueous phase was retained and stored at −20°C until cortisol quantification using a human
salivary cortisol ELISA kit (Salimetrics).

Cortisol measurement

Samples were thawed on ice and 50µl of each was assayed using the Salimetrics human salivary
cortisol ELISA kit, following the manufacturer's protocol. Cortisol values for adults were normalised
to body weight (ng/g), while larval and juvenile values were normalised to total tissue protein
(ng/mg). This assay was validated for use with zebrafish whole-body extracts previously[35,78].
Standards, high and low controls and zebrafish samples (25µL) were added to the plate provided
with the Salimetric kit, along with Assay Diluent (25µL) for the zero standard. The Enzyme
Conjugate was diluted 1:1600 in Assay Diluent and 200µL was dispensed into each well. After
mixing for 5minutes at 500rpm, the plate was incubated at RT for 1h and washed four times with
1× wash buffer (300µL x well x wash). Next, 200µL of TMB Substrate Solution was added, followed
by a 30-minute dark incubation with first 5min mixing. The reaction was stopped with 50µL of Stop
Solution, mixed until wells turned yellow, and absorbance was read at 450nm within 10minutes
using an OMEGA plate reader (BMG Labtech).

Statistical analysis

For qPCR, relative mRNA expressions were calculated using the Pfaffl method[79]. Outliers were
identified and removed using Dixon's test ($\alpha = 0.05$)[80]. Differences in gene expression were assessed
using a one-way ANOVA followed by Tukey's post-hoc test using GraphPad Software (Prism). For
behavioural analysis, all data were analysed with R programming language[75]. Scripts used for
analysis are available on GitHub (https://github.com/AdeleLeg). For models where distance moved,
distance to zone, velocity, or top halves visits were the response variable, we fitted data to mixed
linear models using the "*lme4*" package, and where proportion of responders or proportion of time
spent in the bottom third were our response variable, we fitted data to beta regression models using
the "*betareg*" package. In all instances, for all experiments, we used genotype as fixed effect, and
fish ID and day of experiment as random effects. In the response and habituation to acoustic startle,
we used also the stimulus number as fixed effect. As in García-González et al.[33], we reported
significant fixed effects as Type II Wald χ^2 from models using the package "*car*," post hoc Tukey's
tests were also conducted as necessary with the package "*emmeans*". For the 5-CSRTT, overall
correct responses (learning) and anticipatory responses (impulsivity) were assessed using the
formulas in Supplementary Table 2. Statistical analysis of mean intensity in *in situ* hybridization
images of zebrafish larval sections was performed using ImageJ. Mean intensity values were
quantified across specified regions of interest, indicated in Supplementary Figure 5, ensuring that
comparable areas of similar dimensions were used for each sample to calculate differences in
expression. The resulting values were compared across groups using a paired t-test to determine
significant differences using GraphPad Software (Prism). For all experiments, sample size needed to
achieve adequate statistical power for detecting a significant effect was determined based on data
from previous research or pilot studies. Accepted α level and power were respectively ≤ 0.05 and ≥
0.80.

Fertilisation and survival analysis

Fertilisation and survival rates were measured as described previously[19]. For fertilization rate we
measured the percentage of fertilized eggs when pairing $rbfox1^{19del/19del}$ and compared it with
fertilization rate of $rbfox1^{+/+}$ siblings. Data were collected upon 3 different mating trials, each trial
comprising 3-5 trios (1 male and 2 females). Trials were performed on different weeks, using
different trios for each test. Eggs were collected in different Petri dishes, properly labelled to
distinguish between trios. Fertilisation rate was determined by averaging across Petri dishes from
the same trio and calculated at 6-8hpf. For survival analysis, we measured the percentage of
surviving animals (across 3 tanks for each genotype for each batch, to avoid tank effect) from when
larvae were added to the nursery (at 5dpf) until 2 months of age, when fish were transferred into
adult aquarium.
824

CONTRIBUTIONS

AL conceived the project, designed and conducted the experiments, analysed the data, and wrote
the manuscript. JGG designed and performed the CRISPR/Cas9 experiment generating the line. SH,
PA, SA, XW and WH contributed to the experiments. NFC and BC edited the manuscript. CHB
directed the study, designed the experiments, edited the manuscript, and secured funding. All
authors contributed to the article and approved the submitted version.

ACKNOWLEDGMENTS

CHB, AL, SH, PA and WH were supported by the NIH grant U01 DA044400-03. JGG was supported by
a 2022 NARSAD Young Investigator Grant (Number #30749) by the Brain & Behavior Research
Foundation. NFC and BC were supported by the grants PID2021-127776OB-I100, 202218 31 and
2021-SGR-01093 from the Ministry of Science, Innovation and Universities (Spain), Fundació La
Marató de TV3, and AGAUR-Generalitat de Catalunya, respectively, and by ICREA Academia 2021.
Authors would like to acknowledge SciDraw (https://scidraw.io/) for images used to create the
graphical abstract. We thank Prof. Marina Resmini for kindly providing the equipment used to dry
cortisol extracts.

CONFLICT OF INTEREST
The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.
843

REFERENCES

1.   Cross-Disorder Group of the Psychiatric Genomics Consortium. Electronic address:

plee0@mgh.harvard.edu & Cross-Disorder Group of the Psychiatric Genomics Consortium.

Genomic Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric

Disorders. *Cell* **179**, 1469-1482.e11 (2019).

2.   Antón-Galindo, E. *et al.* Pleiotropic contribution of rbfox1 to psychiatric and

neurodevelopmental phenotypes in two zebrafish models. *Transl. Psychiatry* **14**, 1–11 (2024).

3.   Thisse, B. & Thisse, C. ZFIN Publication: Thisse et al., 2004. *ZFIN* http://zfin.org/ZDB-PUB-

040907-1.

4.   Amir-Zilberstein, L. *et al.* Homeodomain Protein Otp and Activity-Dependent Splicing Modulate

Neuronal Adaptation to Stress. *Neuron* **73**, 279–291 (2012).

5.   Abi-Dargham, A. *et al.* Candidate biomarkers in psychiatric disorders: state of the field. *World*

Psychiatry **22**, 236–262 (2023).

6.   O'Leary, A. *et al.* Behavioural and functional evidence revealing the role of RBFOX1 variation in

multiple psychiatric disorders and traits. *Mol. Psychiatry* **27**, 4464–4473 (2022).

7.   Lee, J.-A., Tang, Z.-Z. & Black, D. L. An inducible change in Fox-1/A2BP1 splicing modulates the

alternative splicing of downstream neuronal target exons. *Genes Dev.* **23**, 2284–2293 (2009).

8.   Shaughnessy, C. A., Myhre, V. D., Hall, D. J., McCormick, S. D. & Dores, R. M. Hypothalamus-

pituitary-interrenal (HPI) axis signaling in Atlantic sturgeon (Acipenser oxyrinchus) and sterlet

(Acipenser ruthenus). *Gen. Comp. Endocrinol.* **339**, 114290 (2023).

9.   Biran, J. *et al.* Splice-specific deficiency of the PTSD-associated gene PAC1 leads to a paradoxical

age-dependent stress behavior. *Sci. Rep.* **10**, 9559 (2020).

10. Linz, R. *et al.* Acute psychosocial stress increases serum BDNF levels: an antagonistic relation to

cortisol but no group differences after mental training. *Neuropsychopharmacology* **44**, 1797–

1804 (2019).

11. Jeanneteau, F. D. *et al.* BDNF and glucocorticoids regulate corticotrophin-releasing hormone

(CRH) homeostasis in the hypothalamus. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 1305–1310 (2012).

12. Karatsoreos, I. N. & McEwen, B. S. Psychobiological allostasis: resistance, resilience and

vulnerability. *Trends Cogn. Sci.* **15**, 576–584 (2011).

13. Miao, Z., Wang, Y. & Sun, Z. The Relationships Between Stress, Mental Disorders, and Epigenetic

Regulation of BDNF. *Int. J. Mol. Sci.* **21**, 1375 (2020).

14. Suliman, S., Hemmings, S. & Seedat, S. Brain-Derived Neurotrophic Factor (BDNF) protein levels

in anxiety disorders: systematic review and meta-regression analysis. *Front. Integr. Neurosci.* **7**,

55 (2013).

15. Fogel, B. L. *et al.* RBFOX1 regulates both splicing and transcriptional networks in human

neuronal development. *Hum. Mol. Genet.* **21**, 4171–4186 (2012).

16. Tomassoni-Ardori, F. *et al.* Rbfox1 up-regulation impairs BDNF-dependent hippocampal LTP by

dysregulating TrkB isoform expression levels. *eLife* **8**, e49673 (2019).

17. Solés-Tarrés, I., Cabezas-Llobet, N., Vaudry, D. & Xifró, X. Protective Effects of Pituitary

Adenylate Cyclase-Activating Polypeptide and Vasoactive Intestinal Peptide Against Cognitive

Decline in Neurodegenerative Diseases. *Front. Cell. Neurosci.* **14**, (2020).

18. Mcewen, B. S. Protection and Damage from Acute and Chronic Stress: Allostasis and Allostatic

Overload and Relevance to the Pathophysiology of Psychiatric Disorders. *Ann. N. Y. Acad. Sci.*

1032, 1–7 (2004).

19. Eachus, H. *et al.* Elevated glucocorticoid alters the developmental dynamics of hypothalamic

neurogenesis in zebrafish. *Commun. Biol.* **7**, 1–14 (2024).

20. Danese, A. & McEwen, B. S. Adverse childhood experiences, allostasis, allostatic load, and age-

related disease. *Physiol. Behav.* **106**, 29–39 (2012).

21. Bornstein, S. R. *et al.* Stress-inducible-stem cells: a new view on endocrine, metabolic and

mental disease? *Mol. Psychiatry* **24**, 2–9 (2019).

22. Kinlein, S. A., Wilson, C. D. & Karatsoreos, I. N. Dysregulated Hypothalamic–Pituitary–Adrenal

Axis Function Contributes to Altered Endocrine and Neurobehavioral Responses to Acute Stress.

Front. Psychiatry **6**, (2015).

23. Adel, M. R. *et al.* Decreased Brain Serotonin in rbfox1 Mutant Zebrafish and Partial Reversion of

Behavioural Alterations by the SSRI Fluoxetine. *Pharmaceuticals* **17**, 254 (2024).

24. Lange, M. *et al.* The ADHD-susceptibility gene lphn3.1 modulates dopaminergic neuron

formation and locomotor activity during zebrafish development. *Mol. Psychiatry* **17**, 946–954

(2012).

25. Parker, M. O. *et al.* Development and automation of a test of impulse control in zebrafish. *Front.*

Syst. Neurosci. **7**, (2013).

26. Halberstadt, A. L. & Geyer, M. A. Habituation and sensitization of acoustic startle: opposite

influences of dopamine D1 and D2-family receptors. *Neurobiol. Learn. Mem.* **92**, 243–248

(2009).

27. Burgess, H. A. & Granato, M. Sensorimotor gating in larval zebrafish. *J. Neurosci. Off. J. Soc.*

Neurosci. **27**, 4984–4994 (2007).

28. Cohen, H. *et al.* Blunted HPA axis response to stress influences susceptibility to posttraumatic

stress response in rats. *Biol. Psychiatry* **59**, 1208–1218 (2006).

29. Leggieri, A. *et al.* Ankk1 Loss of Function Disrupts Dopaminergic Pathways in Zebrafish. *Front.*

Neurosci. **16**, (2022).

30. Zhong, Y. *et al.* RBFOX1 and Working Memory: From Genome to Transcriptome Revealed

Posttranscriptional Mechanism Separate From Attention-Deficit/Hyperactivity Disorder. *Biol.*

Psychiatry Glob. Open Sci. **3**, 1042–1052 (2023).

31. Davies, M. N. *et al.* Generalised Anxiety Disorder – A Twin Study of Genetic Architecture,

Genome-Wide Association and Differential Gene Expression. *PLOS ONE* **10**, e0134865 (2015).

32. Basnet, R. M., Zizioli, D., Taweedet, S., Finazzi, D. & Memo, M. Zebrafish Larvae as a Behavioral

Model in Neuropharmacology. *Biomedicines* **7**, E23 (2019).

33. García-González, J., de Quadros, B., Havelange, W., Brock, A. J. & Brennan, C. H. Behavioral

Effects of Developmental Exposure to JWH-018 in Wild-Type and Disrupted in Schizophrenia 1

(disc1) Mutant Zebrafish. *Biomolecules* **11**, 319 (2021).

34.   Hillman, C., Kearn, J. & Parker, M. O. A unified approach to investigating 4 dpf zebrafish larval

behaviour through a standardised light/dark assay. *Prog. Neuropsychopharmacol. Biol.*

Psychiatry **134**, 111084 (2024).

35.   Baiamonte, M., Parker, M. O., Vinson, G. P. & Brennan, C. H. Sustained Effects of Developmental

Exposure to Ethanol on Zebrafish Anxiety-Like Behaviour. *PLoS ONE* **11**, e0148425 (2016).

36.   Gene Detail⬛:: Allen Brain Atlas: Mouse Brain. https://mouse.brain-map.org/gene/show/92602.

37.   Brain tissue expression of RBFOX1 - Summary - The Human Protein Atlas.

https://www.proteinatlas.org/ENSG00000078328-RBFOX1/brain.

38.   Thisse, B., Thisse, C. Fast Release Clones: A High Throughput Expression Analysis. (2004)

doi:ZDB-PUB-040907-1.

39.   Grone, B. P. & Maruska, K. P. Divergent evolution of two corticotropin-releasing hormone (CRH)

genes in teleost fishes. *Front. Neurosci.* **9**, 365 (2015).

40.   Sahu, M. P. *et al.* Neurotrophin receptor Ntrk2b function in the maintenance of dopamine and

serotonin neurons in zebrafish. *Sci. Rep.* **9**, 2036 (2019).

41.   Shi, C. *et al.* Hyperandrogenism in POMCa-deficient zebrafish enhances somatic growth without

increasing adiposity. *J. Mol. Cell Biol.* **12**, 291–304 (2020).

42.   Faught, E. & Vijayan, M. M. The mineralocorticoid receptor is essential for stress axis regulation

in zebrafish larvae. *Sci. Rep.* **8**, 18081 (2018).

43.   Tessarollo, L. & Yanpallewar, S. TrkB Truncated Isoform Receptors as Transducers and

Determinants of BDNF Functions. *Front. Neurosci.* **16**, (2022).

44.   Martin, S. C., Marazzi, G., Sandell, J. H. & Heinrich, G. Five Trk Receptors in the Zebrafish. *Dev.*

Biol. **169**, 745–758 (1995).

45.   Thisse, C. & Thisse, B. High-resolution in situ hybridization to whole-mount zebrafish embryos.

Nat. Protoc. **3**, 59–69 (2008).

46. Islam, O., Loo, T. X. & Heese, K. Brain-Derived Neurotrophic Factor (BDNF) has Proliferative

Effects on Neural Stem Cells through the Truncated TRK-B Receptor, MAP Kinase, AKT, and

STAT-3 Signaling Pathways. *Curr. Neurovasc. Res.* **6**, 42–53.

47. Finlay, S., Rudd, D., McDermott, B. & Sarnyai, Z. Allostatic load and systemic comorbidities in

psychiatric disorders. *Psychoneuroendocrinology* **140**, 105726 (2022).

48. Elia, J. *et al.* Rare structural variants found in attention-deficit hyperactivity disorder are

preferentially associated with neurodevelopmental genes. *Mol. Psychiatry* **15**, 637–646 (2010).

49. van der Kolk, B. Posttraumatic stress disorder and the nature of trauma. *Dialogues Clin.*

Neurosci. **2**, 7–22 (2000).

50. Karin, O. *et al.* A new model for the HPA axis explains dysregulation of stress hormones on the

timescale of weeks. *Mol. Syst. Biol.* **16**, (2020).

51. Herman, J. P. *et al.* Regulation of the hypothalamic-pituitary-adrenocortical stress response.

Compr. Physiol. **6**, 603–621 (2016).

52. Bale, T. L. & Vale, W. W. CRF and CRF Receptors: Role in Stress Responsivity and Other

Behaviors. *Annu. Rev. Pharmacol. Toxicol.* **44**, 525–557 (2004).

53. Joëls, M., Karst, H., DeRijk, R. & de Kloet, E. R. The coming out of the brain mineralocorticoid

receptor. *Trends Neurosci.* **31**, 1–7 (2008).

54. Holsboer, F. The Corticosteroid Receptor Hypothesis of Depression. *Neuropsychopharmacology*

23, 477–501 (2000).

55. Arborelius, L., Owens, M. J., Plotsky, P. M. & Nemeroff, C. B. The role of corticotropin-releasing

factor in depression and anxiety disorders. *J. Endocrinol.* **160**, 1–12 (1999).

56. Yehuda, R. Status of Glucocorticoid Alterations in Post-traumatic Stress Disorder. *Ann. N. Y.*

Acad. Sci. **1179**, 56–69 (2009).

57. Sheng, J. A. *et al.* The Hypothalamic-Pituitary-Adrenal Axis: Development, Programming Actions

of Hormones, and Maternal-Fetal Interactions. *Front. Behav. Neurosci.* **14**, (2021).

58. Notaras, M. & van den Buuse, M. Neurobiology of BDNF in fear memory, sensitivity to stress,

and stress-related disorders. *Mol. Psychiatry* **25**, 2251–2274 (2020).

59. Berton, O. *et al.* Essential Role of BDNF in the Mesolimbic Dopamine Pathway in Social Defeat

Stress. *Science* **311**, 864–868 (2006).

60. Eckert, A. *et al.* The link between sleep, stress and BDNF. *Eur. Psychiatry* **41**, S282 (2017).

61. Nakagomi, A. *et al.* Role of the central nervous system and adipose tissue BDNF/TrkB axes in

metabolic regulation. *Npj Aging Mech. Dis.* **1**, 1–11 (2015).

62. Fenner, B. M. Truncated TrkB: Beyond a dominant negative receptor. *Cytokine Growth Factor*

Rev. **23**, 15–24 (2012).

63. Cazorla, M. *et al.* Identification of a low–molecular weight TrkB antagonist with anxiolytic and

antidepressant activity in mice. *J. Clin. Invest.* **121**, 1846–1857 (2011).

64. Azogu, I. & Plamondon, H. Blockade of TrkB receptors in the nucleus accumbens prior to

heterotypic stress alters corticotropin-releasing hormone (CRH), vesicular glutamate transporter

2 (vGluT2) and glucocorticoid receptor (GR) within the mesolimbic pathway. *Horm. Behav.* **90**,

98–112 (2017).

65. Bollen, E. *et al.* 7,8-Dihydroxyflavone improves memory consolidation processes in rats and

mice. *Behav. Brain Res.* **257**, 8–12 (2013).

66. Haapasalo, A., Koponen, E., Hoppe, E., Wong, G. & Castrén, E. Truncated trkB.T1 Is Dominant

Negative Inhibitor of trkB.TK+-Mediated Cell Survival. *Biochem. Biophys. Res. Commun.* **280**,

1352–1358 (2001).

67. Eachus, H., Ryu, S., Placzek, M. & Wood, J. Zebrafish as a model to investigate the CRH axis and

interactions with DISC1. *Curr. Opin. Endocr. Metab. Res.* **26**, 100383 (2022).

68. Koolhaas, J. M., de Boer, S. F., Coppens, C. M. & Buwalda, B. Neuroendocrinology of coping

styles: Towards understanding the biology of individual variation. *Front. Neuroendocrinol.* **31**,

307–321 (2010).

69. Koolhaas, J. M. *et al.* Coping styles in animals: current status in behavior and stress-physiology.

Neurosci. Biobehav. Rev. **23**, 925–935 (1999).

70. Maussion, G. *et al.* Regulation of a truncated form of tropomyosin-related kinase B (TrkB) by

Hsa-miR-185* in frontal cortex of suicide completers. *PloS One* **7**, e39301 (2012).

71. Keatinge, M. *et al.* CRISPR gRNA phenotypic screening in zebrafish reveals pro-regenerative

genes in spinal cord injury. *PLOS Genet.* **17**, e1009515 (2021).

72. Dobrzycki, T., Krecsmarik, M. & Monteiro, R. Genotyping and Quantification of In Situ

Hybridization Staining in Zebrafish. *JoVE J. Vis. Exp.* e59956 (2020) doi:10.3791/59956.

73. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**,

676–682 (2012).

74. Wullimann, M. F., Rupp, B. & Reichert, H. The brain of the zebrafish Danio rerio: a

neuroanatomical atlas. in *Neuroanatomy of the Zebrafish Brain: A Topological Atlas* (eds.

Wullimann, M. F., Rupp, B. & Reichert, H.) 19–87 (Birkhäuser, Basel, 1996). doi:10.1007/978-3-

0348-8979-7_5.

75. R Development Core Team. R: A language and environment for statistical computing.

https://stat.ethz.ch/pipermail/r-help/2008-May/161481.html (2008).

76. Gutiérrez, H. C. *et al.* Screening for drugs to reduce zebrafish aggression identifies caffeine and

sildenafil. *Eur. Neuropsychopharmacol.* **30**, 17–29 (2020).

77. Parker, M. O., Brock, A. J., Sudwarts, A. & Brennan, C. H. Atomoxetine reduces anticipatory

responding in a 5-choice serial reaction time task for adult zebrafish. *Psychopharmacology*

(Berl.) **231**, 2671–2679 (2014).

78. Alderman, S. L. & Bernier, N. J. Ontogeny of the corticotropin-releasing factor system in

zebrafish. *Gen. Comp. Endocrinol.* **164**, 61–69 (2009).

79. Pfaffl, M. W. A new mathematical model for relative quantification in real-time RT–PCR. *Nucleic*

Acids Res. **29**, e45 (2001).

80. Dean, R. B. & Dixon, W. J. Simplified Statistics for Small Numbers of Observations. *Anal. Chem.*

23, 636–638 (1951).

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

Figure 1. *rbfox1* mutant fish show hyperactivity, impulsivity and hyperarousal behaviour. A
number of peaks/swimming burst (acceleration events when the fish travelled > 5 mm in < 12 s); N =
rbfox1$^{+/+}$ 24; *rbfox1*$^{+/19del}$ 24; *rbfox1*$^{19del/19del}$ 24. **B-F)** 5-choice serial reaction time task in adult (7
months old) zebrafish (*rbfox1*$^{+/+}$, *rbfox1*$^{+/19del}$, *rbfox1*$^{19del/19del}$): **B)** number of initiated trials during
stage 2 (weeks 2-4); all *rbfox1*$^{+/+}$ (19), all *rbfox1*$^{+/19del}$ (18) and all *rbfox1*$^{19del/19del}$ (40) moved to the
next stage; **C-E)** average correct responses for stages 3-5 (weeks 3-9); **C)** at the end of stage 3 (weeks
3-7), all *rbfox1*$^{+/+}$ (19), 17 *rbfox1*$^{+/19del}$ and 39 *rbfox1*$^{19del/19del}$ moved to the next stage; **D)** at the end of
stage 4 (weeks 4-8), all *rbfox1*$^{+/+}$ (19), all *rbfox1*$^{+/19del}$ (17) and 33 *rbfox1*$^{19del/19del}$ moved to the next

stage; **E)** at the end of stage 5 (weeks 5-9), 15 *rbfox1*^{+/+}, 13 *rbfox1*^{+/19del} and all *rbfox1*^{19del/19del} (33) showed ≥ 50% correct responses; **F)** premature responses during stage 5 (weeks 5-9). Each dot, square or triangle in D-H) represents a single adult zebrafish *rbfox1*^{+/+}, *rbfox1*^{+/19del} and *rbfox1*^{19del/19del} respectively; N = *rbfox1*^{+/+} 19; *rbfox1*^{+/19del} 18; *rbfox1*^{19del/19del} 40. **G-H)** Forced light-dark transition assay in 5dpf zebrafish larvae (*rbfox1*^{+/+}, *rbfox1*^{+/19del}, *rbfox1*^{19del/19del}): **G)** distance travelled during the 10 min assay; **H)** 1 sec time bin resolution plots of the dark-light transition; N = *rbfox1*^{+/+} 43; *rbfox1*^{+/19del} 34; *rbfox1*^{19del/19del} 37. All larvae employed in behavioural experiments were progeny of *rbfox1*^{+/19del} in-cross and were genotyped after experiments and prior to data analysis. In all graphs: bars represent standard error of the mean (SEM); * p < 0.05; ** p < 0.01; *** p < 0.001; **** p < 0.0001.

1054

Figure 2. *rbfox1* is expressed in regions of the brain associated with stress response, social and emotional behaviour, reward and learning. A) *rbfox1* whole mount *in situ* hybridization (ISH) on zebrafish larvae at 28 hours post fertilization and 3-4-5 days post fertilization. **B)** *rbfox1* ISH on adult zebrafish (a - d) forebrain and (e-g) mid-/hind-brain transverse sections. Black boxes represent the region of the brain showed in higher magnification panels. Schematic depictions of the lateral view of the zebrafish brain indicate position of levels illustrated by ISH. Scale bars: 200 μm in A), 100 μm in B) a-c, e, f); 50 μm in B) a', d, e', e'', g. Abbreviations: A, anterior tuberal nucleus; ATN, anterior thalamic nucleus; CZ, central zone of the TeO; D, dorsal area of dorsal telencephalon; DIL, diffuse nucleus of the inferior lobe of the hypothalamus; Dm, dorsal area of medial telencephalon; ECL, external cellular layer; GL, glomerular layer; HaV, ventral area of the habenula; Hv, ventral hypothalamus; ICL, internal cellular layer; PGZ, periventricular gray zone; PPp, parvocellular preoptic nucleus, posterior part; PTN, posterior tuberal nucleus of the hypothalamus; PVO, paraventricular organ; TeO, optic tectum; TL, longitudinal tori; TPp, periventricular nucleus of the posterior tuberculum; Val, valvula cerebelli; Vd, ventral area of dorsal telencephalon; Vl, central area of lateral telencephalon; VM, ventromedial nucleus; Vv, ventral area of ventral telencephalon.

1072
1073 **Figure 3.** *rbfox1* **LoF disrupts zebrafish larvae HPI axis,** *bdnf/trkb2* **pathway and** *pac1a*

expression. TRKB modulation restore HPI gene expression and behaviour. A) Expression levels of *bdnf* and *trkb2* full-length (*TK+*) *and* truncated (*TK-*) in 5 days post fertilisation (dpf) larvae (*rbfox1$^{+/+}$* and *rbfox1$^{19del/19del}$*). *trkb2* expression is shown in a stacked bar format, where the directly measured *TK+* (long form) is shown as the lower portion and the inferred *TK-* (short form) is shown on top. **B)** Expression levels of HPI axis genes *crhb, nr3c2, nr3c1, crhr1, crhr2, mc1r, pomca* and *kf19* in 5dpf larvae (*rbfox1$^{+/+}$* and *rbfox1$^{19del/19del}$*). **C)** Whole-body cortisol measurement of 5dpf larvae (left panel) and adult fish (right panel) in resting physiological conditions. Cortisol values are normalised per whole-body homogenate (g) (*rbfox1$^{+/+}$* and *rbfox1$^{19del/19del}$*). **D-E)** Expression levels of *crhb* and *nr3c2* in presence or absence of the TRKB selective **D)** antagonist ANA-12 and **E)** agonist 7,8-DHF in 5dpf larvae (*rbfox1$^{+/+}$* and *rbfox1$^{19del/19del}$*). **F-I)** Behavioural assays in presence and absence of the TRKB agonist 7,8-DHF in *rbfox1* larvae (*rbfox1$^{+/+}$* and *rbfox1$^{19del/19del}$*): **F)** mean distance travelled, **G)** average speed, **H)** startle response to white light stimulus and **I)** Proportion of responders over time in the response and habituation to acoustic startle assay. **J-N)** Whole mount *in situ* hybridisation (ISH) for *trkb2* anti-sense riboprobe in **J)** *rbfox1$^{+/+}$* and **K)** *rbfox1$^{19del/19del}$* and **L)** *trkb2* sense riboprobe in *rbfox1$^{+/+}$* 5dpf larvae (lateral view in the main boxes and dorsal view in the smaller boxes on the top right corner). **M, N)** Sagittal cryosection of *trkb2* ISH in **M)** *rbfox1$^{+/+}$* and **N)** *rbfox1$^{19del/19del}$* 5 dpf larvae. Black boxes in **M-a)** and **N-a)** represent the region of the brain showed in higher magnification panels in **M-a')** and **N-a')**. Scale bars: 200 µm in J), K) and L); 100 µm in M-a) and N-a); 50 µm in M-a') and N-a'). **O)** Schematic depiction (sagittal) of zebrafish larval brain indicating position of levels illustrated by ISH on sagittal cryosections. **P)** *trkb2* ISH intensity mean in the hypothalamus of 5dpf zebrafish larvae, *rbfox1$^{+/+}$* versus *rbfox1$^{19del/19del}$*. N = 4 larvae x genotype. **Q)** Expression levels of *pac1-hop* and *pac1-short* in *rbfox1$^{+/+}$* and *rbfox1$^{19del/19del}$* 5 dpf larvae. For qPCR experiments, reference genes were actin − β 2 (*actb2*) and ribosomal protein L13a (*rpl13a*). Each green dot/pink triangle in A-D, K-L) represents a pool of 15 larval heads (eyes and jaw removed), while yellow ones represent larvae exposed to TRKB drugs. Where indicated, we used Log$_{10}$ transformation to normalize the data facilitating a clearer visualisation of trends within the dataset. All larvae employed were progeny of *rbfox1$^{+/19del}$* in-cross and were genotyped after experiments and prior to data analysis. For cortisol measurement (G), each dot/triangle represent a pool of 12 larvae in the left panel (larvae) and a single whole zebrafish in the right panel (adult fish). In all graphs: bars represent standard error of the mean (SEM); * p < 0.05; ** p < 0.01; **** p < 0.0001.

1105

Figure 4. *rbfox1* mutants undergo adaptive mechanisms and allostatic overload during development. Expression levels of **A)** HPI axis genes corticotropin releasing hormone b (*crhb*), mineral corticoid receptor (*mr*) and glucocorticoid receptor (*gr*) and of **B)** *bdnf* and *trkb2* truncated/full-length (*TK-/TK+*) (common primers) in adult zebrafish brain (*rbfox1+/+* and *rbfox1^19del/19del*) in normal resting conditions and after exposure to a stressor (novel tank diving). **C)** Expression levels of the *trkb2 TK+* and *TK-* in adult zebrafish brain (*rbfox1+/+* and *rbfox1^19del/19del*) after exposure to a stressor.). *trkb2* expression is shown in a stacked bar format, where the directly measured *TK+* (long form) is shown as the lower portion and the inferred *TK-* (short form) is shown on top. **D)** Expression levels of proliferating cell nuclear antigen (*pcna*) in 5 days post fertilisation zebrafish larvae and adult zebrafish brain (*rbfox1+/+* and *rbfox1^19del/19del*) in normal resting conditions. **E)** Fertility rate and **F)** survival rate of *rbfox1+/+* and *rbfox1^19del/19del*. Each green dot/pink triangle in A-B) represents a single adult *rbfox1+/+* or *rbfox1^19del/19del* brain under resting physiological conditions respectively, while yellow dots/triangles represent *rbfox1+/+* or *rbfox1^19del/19del* brain after stress exposure respectively. In B) each green dot/pink triangle represents single adult *rbfox1+/+* or *rbfox1^19del/19del* brain after stress exposure respectively. In D) for larvae each green dot/pink triangle represents a pool of *rbfox1+/+* or *rbfox1^19del/19del* 15 larval heads (eyes and jaw removed) respectively; for adults each green dot/pink triangle represents a single adult *rbfox1+/+* or *rbfox1^19del/19del* brain respectively. In E) each dot/triangle represents average fertility of 3-5 trios (1 male and 2 females) assessed over 2-3 petri dish (50 embryos per dish) per trio. Each trio belonged to a different tank (for each genotype for each batch) to avoid tank effect). In F) each dot/triangle represents the percentage of survival of a single fish stock comprising 50 larvae. For qPCR experiments, reference genes were actin − β 2 (*actb2*) and ribosomal protein L13a (*rpl13a*). Where indicated, we used Log_{10} transformation to normalize the data facilitating a clearer visualization of trends within the dataset. All larvae employed were progeny of *rbfox1+/19del* in-cross and were genotyped after experiments and prior to data analysis. In all graphs: bars represent standard error of the mean (SEM); * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$.

1132

1133

1134
1135

Figure 5. RBFOX1 role in stress resilience and chronic stress recovery: proposed mechanism
of action. A) In response to stressful challenges (yellow flash), BDNF expression levels increase,
stimulating *CRH* transcription via TRKB.FL/TK+ (yellow pathway). At the same time, stress triggers
also *PACAP* transcription (red pathway) and OTP-mediated transcription of *RBFOX1* and *CRH* (blue
pathway). PACAP stimulates *CRH* transcription via PAC1-short and inhibits *CRH* transcription via
PAC1-hop. PACAP binding to PAC1-short also increases *BDNF* transcription (red line) and enhances
TRKB.FL/TK+ activity (red line). RBFOX1 functions as a "switch" maintaining the balance between
full-length/truncated TRKB isoforms (promoting TRKB mRNA stability and/or alternative splicing):
during the stress recovery phase, *RBFOX1* switches the balances in support of *TRKB.T1/TK-* isoform,
decreasing *CRH* levels and turning off the HPA axis. **B)** *RBFOX1* deletion leads to an increase in *BDNF*
expression levels (yellow arrow and white box) and an unbalance between *TRKB.FL/TK+* (increased)
and *TRKB.T1/TK-* (decreased) isoforms. Such dysregulation ultimately leads to an increase in *CRH*
transcription (and possible hyperactivation of the HPA axis), and increase in PAC1-hop levels as
possible compensatory mechanism. Our data suggest that RBFOX1-mediated regulation of
TRKB.T1/TK- mRNA stability and/or alternative splicing is necessary to promote neuroplasticity and
stress resilience.
1152
1153
1154
1155

1156

Supplementary Figure 1. Generation of a *rbfox1* loss of function line. A) *rbfox1* mRNA expression in *rbfox1*$^{+/+}$, *rbfox1*$^{+/19del}$ and *rbfox1*$^{19del/19del}$ 5 days post fertilisation (dpf) larvae. *In situ* hybridisation on **B)** whole mount zebrafish larvae at 3dpf, a) *rbfox1*$^{+/+}$ and b) *rbfox1*$^{19del/19del}$ and **C)** adult transverse sections of *rbfox1*$^{+/+}$ (a) fore- and (c) mid- brain and *rbfox1*$^{19del/19del}$ (b) fore- and (d) mid-brain. In A) each dot/square/triangle represents a pool of *rbfox1*$^{+/+}$ or *rbfox1*$^{19del/19del}$ 15 larval heads (eyes and jaw removed) respectively. For qPCR experiments, reference genes were actin − β 2 (*actb2*) and ribosomal protein L13a (*rpl13a*). All larvae employed were progeny of *rbfox1*$^{+/19del}$ in-cross and were genotyped after experiments and prior to data analysis. In A): bars represent standard error of the mean (SEM); * $p < 0.05$; ** $p < 0.01$. Scale bars: 200 µm in B); 100 µm in C). Abbreviations: D, dorsal area of dorsal telencephalon; Hv, ventral hypothalamus; TeO, optic tectum; Tl, longitudinal tori; Vd, ventral area of dorsal telencephalon; Vv, ventral area of ventral telencephalon. N = 4 x genotype.

1168

1169

Supplementary Figure 2. *rbfox1* mutant fish show hyperactivity, impulsivity and hyperarousal behaviour. A-B) Locomotion assay in 5 days post fertilization (dpf) zebrafish larvae (*rbfox1*$^{+/+}$, *rbfox1*$^{+/19del}$, *rbfox1*$^{19del/19del}$): **A)** distance travelled during 15 min; **B)** average speed; N (A-B) = *rbfox1*$^{+/+}$ 24; *rbfox1*$^{+/19del}$ 24; *rbfox1*$^{19del/19del}$ 24. 5-9). **C-D)** Response and habituation to acoustic startle assay in 5dpf zebrafish larvae (*rbfox1*$^{+/+}$, *rbfox1*$^{+/19del}$, *rbfox1*$^{19del/19del}$): **C)** mean distance travelled during the assay; **D)** rate of habituation over time/stimuli; N = *rbfox1*$^{+/+}$ 40; *rbfox1*$^{+/19del}$ 39; *rbfox1*$^{19del/19del}$ 35. E-G) Novel tank diving assay in *rbfox1*$^{+/+}$, *rbfox1*$^{+/19del}$ and *rbfox1*$^{19del/19del}$. **E)** Time spent at the bottom of the tank. **F)** Distance travelled during the assay and **G)** number of transitions

1178 between the top and bottom of the tank. N = *rbfox1$^{+/+}$*, 50; *rbfox1$^{+/19del}$* 54; *rbfox1$^{19del/19del}$* 54. Bars
1179 represent standar error of the mean (SEM). All larvae employed in behavioural experiments were
1180 progeny of *rbfox1$^{+/19del}$* in-cross and were genotyped after experiments and prior to data analysis. In
1181 all graphs: bars represent standard error of the mean (SEM); * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$;
1182 **** $p < 0.0001$.

1183

1184

Supplementary Figure   3. 5-choice serial reaction time task. A) Zantiks behavioural testing
apparatus used in the laboratory (left panel) and a schematic representation of a single Zantiks
chamber (right panel). **B)** Schematic overview of the pipeline in a single 5-choice trial, including
initiation, stimulus presentation and response outcome. **C-E)** Illustrative diagrams of representative
trial outcomes as depicted in the pipeline in panel B), including C) correct response, D) incorrect
response and E) omission.

1191

1192

1193

Supplementary Figure 4. A) *crhb* and *bdnf* (left) and trkb2 short and long isoform (right) expression
levels in 3 days old larvae, *rbfox1*$^{+/+}$ and *rbfox1*$^{19del/19del}$. **B)** *rbfox2* expression levels in 5 days old
larvae, *rbfox1*$^{+/+}$ and *rbfox1*$^{19del/19del}$. **C)** *bdnf* expression levels in 5 days old larvae, *rbfox1*$^{+/+}$ and
rbfox1$^{19del/19del}$, following chronic exposure to either ANA-12 (left) or 7,8-DHF (right). Actin – β 2
(*actb2*) and ribosomal protein L13a (*rpl13a*) were used as reference genes. Dots/triangles represent
pool of 15 larval heads (eyes and jaw removed). Bars represent standard error of the mean (SEM). N
= *rbfox1*$^{+/+}$ 5; *rbfox1*$^{19del/19del}$ 5.

1203

1204

Supplementary Figure 5. Details of *in situ* hybridisation experiments for *trkb2* anti-sense riboprobe on 5 days post fertilisation zebrafish larvae, sagittal sections. **A)** Schematic representation indicating the level of the anatomical section (red line). **B)** Representative sagittal section of zebrafish larva stained for *rbfox1* (anti-sense riboprobe), with anatomical regions labelled in black according to ZFIN atlas of zebrafish larvae (https://zfin.org/zf_info/anatomy.html). **C-D)** Overview of zebrafish larvae

hypothalamus (sagittal sections), *rbfox1*$^{+/+}$ (four panels on the left) and *rbfox1*$^{19del/19del}$ (four panels
on the right). Each panel corresponds to a different animal. N = 4 per genotype. Sections have been
placed on the same slide and were stained under the same conditions. **E)** *In situ* hybridisation for
trkb2 sense riboprobe on 5 days post fertilisation zebrafish larva, sagittal section. **F)** Example of
areas selected (areas enclosed by the black dotted lines) to calculate *trkb2* mean intensity in the
hypothalamus of a) zebrafish *rbfox1*$^{+/+}$ (left panel) and a') *rbfox1*$^{19del/19del}$ (right panel) larval sections.
G) Comparison of *trkb2* and crhb expression in the larval zebrafish brain. The left panel displays
crhb-expressing cells in the larval brain from the publicly available MapZebrain atlas
(https://mapzebrain.org/atlas/2d). The right panel shows whole-mount *in situ* hybridisation for *trkb2*
expression in 5 days old zebrafish larva. Scale bars: F) 50 μm; G) 100 μm.

1230 **Supplementary Table 1.** List of primers used for qPCR experiments.
1231

Gene (purpose)	Gene acronym	Accession number	Sequence (5' to 3')	Product size (bp)
actin, β 2 (housekeeping)	*actb2*	NM_181601.5	Forward: CGAGCTGTCTTCCCATCCA	86
			Reverse: TCACCAACGTAGCTGTCTTTCTG	
ribosomal protein L13a (housekeeping)	*rpl13a*	NM_212784.1	Forward: TCTGGAGGACTGTAAGAGGTATGC	148
			Reverse: AGACGCACAATCTTGAGAGCAG	
RNA binding fox-1 homolog 1	*rbfox1**	NM_001005596.1	Forward: AACAGCATACAGGCGGTCTC	110
			Reverse: GGCTGCGTTTTGATTTCTGT	
corticotropin releasing hormone	*crhb*	NM_001007379.1	Forward: CGAGACATCCCAGTATCCAAAAAG	60
			Reverse: TCCAACAGACGCTGCGTTAA	
mineralcorticoid receptor	*nr3c2*	NM_001100403	Forward: CTTCCAGGTTTCCGCAGTCTAC	75
			Reverse: GGAGGAGAGACACATCCAGGAAT	
glucocorticoid	*nr3c1*	NM_001020711	Forward:	90

receptor		.3	ACTCCATGCACGACTTGGTG	
			Reverse: GCATTTCGGGAAACTCCACG	
brain derived neurotrophic factor	*bdnf*	XM_009303234.3	Forward: AGCAGACTGTCTTTTGTTGTGTT	123
			Reverse: CTTAAGCTCACACAGGTGCC	
neurotrophic tyrosine kinase, receptor, type 2b (common primers)	*trkb2.TK+/TK-*	NM_001197161.2	Forward: CGCATACCTGACGTGGATCA	130
			Reverse: AAGCAACCCCAGCTATTCCC	
neurotrophic tyrosine kinase, receptor, type 2b (tyrosine kinase domain primers)	*trkb2.TK+*	NM_001197161.2	Forward: TGAAAGTGGGCGAGCAGATT	173
			Reverse: CATCTGGACCATGAGCCCTG	
adenylate cyclase activating polypeptide 1a (pituitary) receptor type I (adcyap1r1a) transcript variants 1 and 2	*pac1a-hop*	NM_001142926.1 NM_001013444.2	Forward: CAGAAATGCTTCAGTGAGCCC	82
			Reverse: CCAGACGTAGCGTGATGGTA	
adenylate cyclase activating polypeptide 1a (pituitary) receptor type I (adcyap1r1a), transcript variant 3	*pac1a-short*	NM_001142925.1	Forward: GATTCTGGCTGCTGGGACAT	183
			Reverse: GCGCGCCAGACGTAGATAAA	
RNA binding fox-1 homolog 2	*rbfox2*	NM_214746.2	Forward: CCGTATGCTAACGGGGACAG	106
			Reverse: ACCCTGGAACTGCGTAGAAC	
corticotropin releasing hormone receptor 1	*crhr1*	XM_691254.7	Forward: TCGGTGTTGAGTCTCATGCC	186
			Reverse: CGCGTGCGGGTAAAATGTAG	
corticotropin releasing hormone receptor 2	*crhr2*	NM_001113644.1	Forward: CTGACGAGCCACCTGTTTCT	207
			Reverse: TTACCCGTCCACCGATCAAC	
melanocortin	*mc1r*	NM_180970.1	Forward: TGCTGGTCACCGCAAAAATG	115

			Reverse: ATATAACGGTCCGCCGCAAT	
propiomelanocortin a	*pomca*	NM_181438.3	Forward: CTGCCATAGATGATGCCGGT	140
			Reverse: TGCGTTTGCTCGTCTCTGAT	
Krüppel-like factor 9	*klf9*	NM_001128729.1	Forward: GACAAACTAACCGGCGAGGA	153
			Reverse: GTTCTCCCGATTCTCCGCAT	

1232   * These primers amplify all *Danio rerio rbfox1* isoforms available on NCBI: XM_068216713.1,
1233   XM_068216714.1, XM_017354041.3, XM_068216717.1, XM_068216718.1, XM_017354043.3,
1234   XM_017354044.3, XM_068216722.1, XM_017354045.3, XM_068216724.1, XM_068216725.1,
1235   XM_068216726.1, XM_068216728.1, XM_017354049.3   XM_068216731.1.
1236
1237

1238   **Supplementary Table 2.** 5-Choice Serial Reaction Time Task stages (5-CSRTT) and corresponding
1239   descriptions. Source: https://zantiks.com
1240

Stage	Name of the stage	Description
Stage 1	Habituation	All lights are illuminated (initiator light, five choice lights and food light) for 20 sec and the food is released. All fish are moved to the next stage of training after 1 week.
Stage 2	Initiator training	Initiator light is on for 20 sec. If the fish swims to the illuminated initiation area, an initiated trial is recorded and the feeder releases food with the food light on for 20 sec. This is followed by 10 sec ITI. The initiator training is repeated for 30 trials. Fish move to the next stage when the average of initiated trials to total trials over five days is > 50%.
Stage 3	Stimulus light training	The initiation light is on for 20 sec. If the fish swims to the illuminated initiation area, an initiated trial is recorded and all five choice lights are illuminated for 20 sec. If the fish swims to any illuminated choice area whilst the light is on, a correct response is recorded and the feeder releases food with the food light on for 20 sec. This is followed by 10 sec ITI. The stimulus light training is repeated for 30 trials. Fish move to the next stage when both the average of initiated trials to total trials and correct responses to initiated trials are > 50%.
Stage 4	5-CSRTT/no-delay	The initiation light is on for 20 sec. If the fish swims to the initiation area, an initiated trial is recorded and one of five choice lights is illuminated for 20 sec. If the fish swims to the illuminated choice area, a correct response is recorded and the feeder releases food with the food light on for 20 sec. This is followed by 10 sec ITI. The 5-CSRTT/no-delay stage is repeated for 30 trials. Fish move

		to the next stage when the average of both initiated trials to total trials and correct responses to initiated trials are > 50%.
Stage 5	5-CSRTT/variable-delay	The initiation light is on for 20 sec. If the fish swims to the initiation area, an initiated trial is recorded and after a variable pre-stimulus interval of 0-8 sec, one of five choice lights is illuminated for 20 sec. This is followed by 10 sec ITI. If during the pre-stimulus interval, the fish swims to any unilluminated choice area, a premature response is recorded. The average of premature responses to initiated trials over four days is reported as a measure of impulsive action.

1241