Effects of SSRIs on the spatial transcriptome of dorsal raphe serotonin neurons

Henningson C.[1], Mlost, J.[1], Pollak Dorocic I.[1]

[1]Science for Life Laboratory, Department of Biochemistry and Biophysics, Stockholm University, Sweden

Correspondence: iskra.pollak@scilifelab.se

Abstract

The serotonin system is the main therapeutic target for selective serotonin reuptake inhibitors (SSRIs) in treating depression, yet the mechanism of action of SSRIs remains incompletely understood. To investigate the molecular and transcriptional effects of SSRI administration on serotonin neurons, we performed spatial transcriptomics, a spatially resolved RNA-sequencing method in intact brain tissue. Mouse brain sections containing the dorsal raphe nucleus and adjacent midbrain structures were analyzed, revealing six distinct serotonergic subpopulations with unique molecular signatures and spatial distributions. Both acute and chronic fluoxetine treatment induced a large number of changes in gene expression in the dorsal raphe nucleus. Notably, *Htr1a* expression increased following acute treatment but decreased after chronic administration, supporting previous findings on serotonin transporter blockade effects on 5HT1a autoreceptors. Gene enrichment and network analysis identified key pathways modulated by SSRI administration, including Ras, MAPK and cAMP signalling pathways as well as pathways involved in axonal guidance. Additionally, we observed treatment-dependent opposing transcriptional changes in neuropeptides, particularly thyrotropin-releasing hormone (*Trh*) and Prodynorphin (*Pdyn*), with distinct spatial localization within the dorsal raphe nucleus. Collectively, our transcriptomic and in situ hybridization analyses reveal spatial and cell-type-specific heterogeneity in SSRI action within the dorsal raphe nucleus, providing new insights into the molecular basis of SSRI treatment effects.

Introduction

Depression is a leading contributor to the global disability burden, with high lifetime prevalence, elevated risk of suicide, and a critically increasing incidence[1]. The serotonin system in the brain is the primary pharmacological target of the most common antidepressant treatment, the Selective Serotonin Reuptake Inhibitors (SSRIs). These drugs are routinely prescribed as a first-line treatment for depression and a range of comorbid conditions including anxiety and bipolar disorder, and are thus one of the most prescribed pharmaceutical drugs in the world. A large number of patients, in some countries as many as 10-15% of the total population[2], currently rely on antidepressants to alleviate their suffering, with SSRIs being the most common form of treatment[3]. Despite this extensive clinical use, the molecular mechanisms underlying SSRI action remain incompletely understood. Furthermore, SSRIs exhibit variable treatment outcomes, delayed onset of therapeutic effects, and numerous side effects[4]. Understanding the mechanism of action represents a critical challenge in modern psychopharmacology and is essential for developing more effective therapeutic strategies.

Serotonin (5-hydroxytryptamine) is one of the major neuromodulators in the central nervous system, and plays a role in both normal function and disease. Serotonin interacts with at least 14 different receptor subtypes and therefore influences a wide range of physiological processes, from homeostatic maintenance to complex roles in mood and motivation[5]. In the mammalian brain, serotonergic neurons are exclusively located within the brainstem raphe nuclei, with the dorsal raphe nucleus (DR) containing the majority of forebrain-projecting serotonin neurons. These neurons are embedded within an extensive neural network, receiving inputs from multiple forebrain regions[6-8] and projecting widely throughout the brain to modulate diverse behaviors[9]. One limitation of our current knowledge of the serotonin system, as well as its pharmacological targeting, is that the system has been considered unitary, *ie.* comprising all neurons that synthesize serotonin. However, recent single-cell RNA sequencing studies have described the heterogeneity of the serotonin system and elucidated a number of molecular serotonergic subpopulations within the DR[10-12]. The DR cytoarchitecture is arranged in a broad-level topography, with neuron types spatially arranged in the dorsal, ventral and later DR sub-compartments and linked to distinct forebrain-circuits. This cellular complexity is further enhanced by the presence of other neuronal types, including dopaminergic, glutamatergic, and GABAergic neurons, as well as various neuropeptide-expressing cells[13,14]. Understanding the spatial organization and molecular signatures of these distinct neuronal populations is crucial for elucidating their specific functional roles and responses to pharmacological interventions.

SSRIs exert their action by blocking the serotonin transporter (SERT), leading to rapid increases in extracellular serotonin concentration within hours[15]. However, clinical efficacy is only achieved after several weeks of chronic administration[16]. This delay in therapeutic effect is hypothesized to be related to 5-HT_{1A} receptor desensitization. 5-HT_{1A} are Gi-coupled receptors expressed both post- and pre-synaptically, with pre-synaptic autoreceptors providing crucial feedback inhibition of serotonin neurons. The initial SSRI-induced elevation of synaptic serotonin activates these autoreceptors, resulting in decreased serotonin neuron firing and release[17,18]. Chronic SSRI treatment leads to 5-HT_{1A} autoreceptor desensitization and subsequent loss of feedback inhibition, ultimately increasing serotonin neuron activity[19-21]. While this model has been influential in explaining the delayed therapeutic onset of SSRIs, alternative mechanisms have also been proposed, including enhanced hippocampal neurogenesis and increased synaptic plasticity[23].

Antidepressants induce widespread transcriptional changes across multiple brain regions, including the prefrontal cortex, hippocampus, and amygdala[24-26]. However, most studies have focused on limited candidate genes in specific regions. Recent unbiased profiling using bulk RNA sequencing revealed that chronic fluoxetine treatment produces region-specific transcriptional effects across the brain, with the most pronounced changes occurring in the

109 DR[27]. However, the specific effects on serotonin neurons, their spatial subtypes, and other cell
110 types in the DR are unknown.
111
Here, we employed spatial transcriptomics and in-situ hybridization to analyze the
molecular architecture of the mouse dorsal raphe nucleus under baseline conditions and
following acute or chronic fluoxetine administration. We first established a high-resolution
spatial map of the DR transcriptional landscape, identifying molecular markers that define
distinct serotonergic subpopulations. We then characterized the transcriptional responses to
SSRI treatment across these neuronal populations, revealing divergent effects of acute versus
chronic administration. Our findings provide insight into the spatial and cell type-specific
heterogeneity of SSRI action within the DR, with implications for understanding the complex
mechanisms underlying antidepressant action. The data is accessible through an interactive
app available at https://st-ssri.serve.scilifelab.se/app/st-ssri.
122
123 **Results**
124
125 *Spatial transcriptomic profiling of midbrain and DR*
126
Using unbiased and untargeted spatial transcriptomics, we profiled gene expression
in coronal brain sections containing the DR and neighboring regions, in adult mice (**Fig. 1A**).
We detected, on average, 3892 genes and 10,239 reads per spot (3892 ± 1133 unique genes,
10,239 ± 4978 reads). The complete dataset after quality control contained information on the
expression of 30,413 unique genes across 8369 spots. Shared Nearest Neighbor (SNN)
clustering of gene expression across the whole brain section revealed 13 distinct clusters (**Fig.**
1B). We did not observe batch nor sex mediated effects on clustering (**Fig. S1**). Aligning the
molecular clusters with spatial location in brain tissue largely recapitulated anatomical brain
regions, but also provided a higher-resolution delineation within the same brain region (**Fig.**
1C). For example, the periaqueductal gray (PAG) can be subdivided into the dorsal, lateral,
ventrolateral and aqueductal PAG subregions with distinct molecular gene expression profiles.
Spatial sequencing also clearly reveals the DR anatomical cluster, as well as a cluster flanking
the DR bilaterally with distinct molecular signatures (**Fig. 1D**).
140
The DR cluster is clearly identified with the expression of canonical serotonin markers,
relating to serotonin synthesis, reuptake, release and regulation. These include: *Tph2*
(Tryptophan hydroxylase 2), *Slc6a4* (serotonin transporter, SERT), *Slc18a2* (Vesicular
monoamine transporter 2, VMAT2), *Ddc* (dopa decarboxylase), and *Gchfr* (GTP
cyclohydrolase I feedback regulatory protein) that is involved in post-translational regulation
of the serotonin synthesis pathway[28] (**Fig. 1E**). Another molecular cluster (lateral DR-GABA),
flanks the DR bilaterally and is marked by *Gad2* expression. Overall, the spatial sequencing
approach revealed molecular markers for midbrain anatomical structures, and reliably
identified the DR based on serotonin-associated gene expression. The subsequent analysis
focuses on the DR cluster, in order to define spatial subpopulations of serotonin neurons, and
determine the effects of acute and chronic fluoxetine treatment.
152
153 *Spatial subpopulations within DR serotonin system*
154
The DR exhibits complex molecular heterogeneity across diverse cell populations.
While single-cell RNA sequencing (scRNA-seq) studies have identified distinct serotonergic
subpopulations[10–12], this approach cannot preserve spatial information. Previous studies
attempted to map the molecular subtypes to anatomical locations using visualization of limited
candidate genes identified through scRNA-seq. However, no fully unbiased analysis of linking
complete transcriptomic profiles directly to anatomical location in tissue has been performed
in the DR.

Spatial transcriptomics analysis revealed 6 distinct molecular subclusters within the
DR, each showing specific anatomical distributions along the rostro-caudal axis (**Fig. 2A,B**).
These clusters demonstrated unique spatial organization: one midline, one ventral, one
caudal, two lateral clusters, and one cluster with reduced serotonergic marker expression (**Fig.**
2C). All serotonergic clusters show strong molecular identity based on distinct marker genes,
while the non-serotonergic cluster is defined by weak expression of canonical serotonin
markers, likely containing less serotonin neurons compared to other DR spots. We identified
cluster-specific marker genes, with several showing DR-restricted expression patterns:
midline (*Slc17a8, Pdyn*), caudal (*Met*), and lateral cluster 1 (*Wfdc12, Trh*) (**Fig. 2D**). Other
markers, including ventral cluster-associated *Apod* and lateral cluster 2-associated *Avp* and
Slc5a7, showed broader expression patterns extending beyond the DR. Notably, lateral
cluster 2 exhibited a distinct molecular profile, likely influenced by its proximity to cholinergic
neurons of the rostral lateral dorsal tegmental nucleus, as supported by *Slc5a7* expression.
175
To validate our spatial transcriptomics findings at single-cell resolution, we performed
multiplexed in-situ hybridization (ISH) for key marker genes in the DR (**Fig. 2E, F**). The midline
cluster is defined by the expression of *Slc17a8*, the gene coding for Vesicular Glutamate
Transporter 3 (VGLUT3). This neural subpopulation has been extensively characterized, and
has been shown to contain serotonin neurons that co-express VGLUT3 and are able to co-
release glutamate in several forebrain regions[29–31]. We found that approximately half of
Slc17a8 neurons in the DR express *Slc6a4*, the gene coding for SERT (59.9% +/- 8.3% of
Slc17a8 cells express *Slc6a4*, mean ± SD, n = 1979 *Slc17a8* neurons, n = 6 mice) consistent
with previous characterizations of this co-releasing population[13]. The GABAergic marker gene
Gad2, which codes for Glutamate Decarboxylase 2, showed widespread expression
throughout the DR and adjacent regions. A minority of DR *Gad2* cells co-expressed *Slc6a4*
(11.1% +/- 5.1% of *Gad2* cells express *Slc6a4*, n = 3718 *Gad2* neurons, n = 5 mice). Analysis
of the serotonin autoreceptor *Htr1a* revealed extensive but not exclusive co-expression with
Slc6a4 (71.0% ± 10.2% of *Htr1a* cells express *Slc6a4*, n = 3114 *Htr1a* neurons from 6 mice),
confirming its predominant but not exclusive association with serotonergic neurons.
191
Our spatial transcriptomics analysis identified additional molecular markers with
distinct spatial distributions in the DR. *Pdyn*, encoding the neuropeptide prodynorphin, showed
strong midline expression and substantial co-localization with serotonergic neurons (76.9%
+/- 6.5% of *Pdyn* cells express *Slc6a4*, n = 1235 *Pdyn* neurons, n = 6 mice). *Trh* (thyrotropin-
releasing hormone) exhibited a specific expression pattern in the caudal and ventral lateral
wings of the DR, with moderate serotonergic overlap (37.5% +/- 16.8% of *Trh* cells express
Slc6a4, n = 483 *Trh* neurons, n = 5 mice). *Bdnf* (brain derived neurotrophic factor), linked to
the regulation of stress and development of mood disorders[32], showed predominantly lateral
expression with limited serotonergic co-expression (11.5% +/- 6.9% of *Bdnf* cells express
Slc6a4, n = 2709 *Bdnf* neurons, n = 5 mice). *Vip* (vasoactive intestinal peptide) was
concentrated in the dorsal DR along the aqueduct, with minimal serotonergic overlap (3.7%
+/- 3.0% of *Vip* cells express *Slc6a4*, n = 356 *Vip* neurons, n =6 mice). We integrated these
findings into a comprehensive molecular map showing the spatial distribution and serotonergic
co-expression patterns of each marker along the rostro-caudal axis (**Fig. S2**). These results
reveal distinct molecular signatures that define spatially segregated neuronal populations
within the DR.
208
Through integration of previously published single-cell RNA sequencing data from the
DR[11], we mapped five previously identified serotonergic subtypes (5-HT-I through V) onto our
spatial transcriptomics dataset using CytoSPACE[33] (**Fig. S3A**). CytoSPACE is a
computational approach to integrate scRNA-seq and spatial transcriptomics data by assigning
single cells to spatial locations based on gene expression similarity to determine cell type
composition of each spot. Analysis revealed substantial overlap between certain subclusters
and subpopulations, particularly evident between 5-HT-V and the Caudal subcluster (**Fig.**
S3A, B) Analysis of top spatial cluster gene expression patterns demonstrated distinct

molecular signatures across serotonergic subpopulations (**Fig. S3C**). The 5-HT-V subtype and
Caudal spatial cluster exhibited concordant expression profiles, characterized by expression
of *Met*, *Calcr*, and several other genes. The 5-HT-IV subtype aligned with the Midline spatial
cluster, sharing *Slc17a8* expression while showing minimal expression of Lateral 1 marker
genes. Similarly, 5-HT-I subtype and the Lateral 1 spatial cluster demonstrated shared
expression of multiple markers, including *Prph*, *Cdh13*, and *Ctsl*. In contrast, subpopulations
5-HT-II and III displayed mixed expression patterns, incorporating markers characteristic of
both midline and lateral populations (**Fig. S3C, D**). Quantitative spatial distribution analysis
revealed preferential localization patterns, with 5-HT-II predominantly present in Lateral 1
(56% ± 13.3% of mapped 5-HT-II cells, mean ± SD, n = 37 of 66 neurons) and 5-HT-III showing
enrichment in Midline regions (71% ± 4.5% of mapped 5-HT-III cells, mean ± SD, n = 17 of 24
neurons). These findings demonstrate that while serotonergic subtypes exhibit clear spatial
preferences, each DR spatial cluster represents a distinct microenvironment characterized by
specific combinations of 5-HT subtypes (**Fig. S5E**).
231
232 *Effects of acute and chronic SSRI treatment on DR serotonin system*
233
SSRIs directly target serotonergic neurons by blocking SERT, yet their effects on gene
expression within these neurons are not known. Previous transcriptional studies have focused
primarily on downstream brain regions innervated by serotonergic projections, leaving a
critical gap in our understanding of SSRI action at its primary site of effect. To address this,
we employed spatial transcriptomics to analyze gene expression changes in the DR following
fluoxetine administration. Given the well-documented temporal disparity between SSRI
administration and therapeutic efficacy, we examined both acute and chronic treatment
conditions to capture the dynamic transcriptional landscape underlying this therapeutic delay.
242
Three treatment protocols using gavage administration were administered to healthy,
naïve mice: acute (single dose fluoxetine, 10mg/kg), chronic (22 days daily administration,
10mg/kg), and control (single dose of saline) (n=3/treatment group) (**Fig. 3A**). We sought to
administer a dose and treatment schedule comparable to clinical use in treatment of
depression[34,35]. Our analysis identified a range of differentially expressed genes (DEGs)
between the control, acute and chronic treatment protocols in both the whole midbrain tissue
clusters (**Fig. S4A)** and the DR clusters (**Fig. S4B,C**). Focusing specifically within the DR,
acute and chronic SSRI administration resulted in DEGs enriched in several important
signalling pathways and biological processes, including: Ras, MAPK, cAMP and axon
guidance pathways (**Fig. 3B**). Chronic treatment led to widespread downregulation of axon
guidance genes, most notably *Plxnc1*, which encodes a member of the plexin family and
regulates axon guidance, cell motility and migration[36] (**Fig. 3C**). Conversely, chronic treatment
upregulated *Rasgrp1*, an activator of the Erk/MAP kinase cascade. The serotonin
autoreceptor gene *Htr1a* showed biphasic regulation, with upregulation following acute
treatment and subsequent downregulation after chronic administration, consistent with
established SSRI-induced autoreceptor desensitization[19-21]. Similarly, the immediate early
transcription factor *Fos* used as a proxy for neural activity, is downregulated over chronic
treatment (**Fig. 3C**). These findings reveal broad temporal dynamics in gene expression
across multiple functional pathways during SSRI treatment (**Fig. 3D, Fig. S4C**).
262
We investigated how many of the DEGs are uniquely modulated in each treatment
condition by analyzing the degree of overlap between the DEGs in control, acute and chronic
fluoxetine administration condition (**Fig. 3E, F**). From a total of 232 DEGs identified, 192
showed downregulation and 153 showed upregulation across all different treatment
conditions. Comparing the treatments, the number of downregulated DEGs in the acute vs
chronic treatment is highest, while each treatment comparison shows a large number of
unique upregulated DEGs. Interestingly, very few up- and down-regulated DEGs are shared
between all three treatment conditions.
271

In order to find key factors among DEGs, we performed network analysis of protein-protein interactions and selected the top three up- or down-regulated genes (**Fig. 4**). Following acute SSRI treatment, *Bdnf* was the only hub gene detected but among the top three upregulated genes were *Bdnf, Pdyn* and *Rgs5*, whereas the top three downregulated genes were *Myl9, Nptx2* and *Adgrl4* (**Fig. 4A**). Following chronic SSRI treatment, we found 7 hub genes: *Nras, Fos, Calb1, Rras, Fcer1g, Pdyn* and *Csf1r* (**Fig. 4B**). Among the top three upregulated genes were *Med1, Fcer1g, Csf1r*, whereas top three downregulated genes were *Pdyn, Nptx2* and *Fxyd5* (**Fig. 4B**). When comparing acute vs chronic SSRI treatment, we found 10 hub genes: *Fos, Met, Kit, Hcrt, Rbfox3, Trh, Ntrk1, Vip, Pdyn, Nkx2-2* (**Fig. 4C**). Among the top three upregulated genes were *Socs4, Bmp1* and *Bhlhe22*, whereas top three downregulated genes were *Vip, Tnnt1* and *Nkx6-1* (**Fig. 4C**). Interestingly, the two neuropeptide genes, *Pdyn* and *Trh*, showed opposite dynamic regulation over the time course of SSRI treatment. *Pdyn* was upregulated after acute and downregulated after chronic treatment, compared to the baseline. By contrast, *Trh* was initially downregulated, and then upregulated after the course of treatment (**Fig. 3D**).

Analysis of neuronal activity patterns revealed distinct temporal responses to SSRI treatment across DR subregions (**Fig. S3D**). Using NEUROeSTIMator[37], a bioinformatic approach that generates normalized activity scores based on the expression of 22 immediate early genes, we observed region-specific alterations in neuronal activity. Chronic SSRI administration resulted in significantly reduced overall activity in the DR compared to control conditions. This reduction in activity was specifically localized to the midline subcluster, while the lateral subcluster maintained baseline activity levels. Notably, this activity differential was only observed when comparing acute versus chronic treatment conditions. Analysis of the aggregate dataset showed no significant treatment effect on overall neuronal activity.

In summary, temporal analysis of SSRI administration revealed extensive transcriptional remodeling within the DR, with the most pronounced differences emerging between acute and chronic treatment conditions. These changes encompassed individual genes, molecular pathways, and region-specific activity patterns across DR subdomains. The transition from acute to chronic fluoxetine treatment was characterized by the largest number of differentially expressed genes, affecting both up- and down-regulated transcriptional programs and associated signaling networks.

Dynamic gene expression changes of DR neuropeptides in acute and chronic SSRI treatment

Transcriptomic analysis revealed opposing regulation of *Pdyn* and *Trh* during SSRI treatment. To confirm these findings, we performed ISH on DR tissue after either acute or chronic fluoxetine treatment as well as with vehicle control. We found that acute administration increased both the number of *Pdyn*-expressing cells and total *Pdyn* mRNA spots in the midline DR (**Fig. 5A**). This acute upregulation was partially reversed following chronic treatment. Acute treatment also increased the number of neurons co-expressing *Pdyn* and *Slc6a4*. ISH analysis of *Trh* expression also corroborated our transcriptomic findings: chronic treatment increased both *Trh*-expressing cells and total *Trh* mRNA spots in the lateral DR (**Fig. 5B**). We observed no changes in *Trh* expression following acute treatment, nor in the number of neurons co-expressing *Trh* and *Slc6a4*. These findings demonstrate distinct regulatory mechanisms for *Pdyn* and *Trh*: SSRI-induced changes in *Pdyn* expression occur in serotonergic neurons, while *Trh* regulation appears independent of its co-expression with serotonin.

The gene *Slc17a8* (VGLUT3) has been speculated to play a role in the treatment effect of SSRIs[38]. We sought to clarify whether the expression levels of *Slc17a8* alone, or in cells co-expressing *Slc6a4*, are affected by SSRI treatment and found no evidence of either (**Fig. 5C**). Though identified as DEGs in the transcriptomic analysis, ISH revealed no significant

treatment effects upon *Gad2*, *Htr1a*, *Bdnf*, or *Vip* expression in the DR (**Fig. S5**). Additionally,
we investigated potential sex-specific effects of acute and chronic SSRI administration. The
treatment-dependent regulation of *Pdyn* and *Trh* expression was consistent across male and
female brains, while other markers showed no treatment effects in either sex. Overall, ISH
analysis revealed no statistically significant sex-specific differences in the expression of *Pdyn*,
Trh, *Slc17a8*, *Gad2*, *Vip*, *Bdnf*, and *Htr1a* across control, acute, or chronic treatment
conditions (**Fig. S6**).
333
334 **Discussion**
335
Using unbiased spatial transcriptomics, we generated a comprehensive molecular
map of the DR, identifying six distinct subpopulations with unique spatial distributions. Our
findings extend previous single-cell RNA-sequencing studies[10–12] by providing crucial spatial
context to DR molecular heterogeneity. We identified several region-specific markers: *Slc17a8*
and *Pdyn* in the midline, *Met* in the caudal region, and *Wfdc12* and *Trh* in the lateral DR. These
markers may be valuable candidates for targeted genetic manipulation of specific DR
subpopulations using transgenic or viral approaches. Additionally, our analysis of
neuropeptide expression revealed distinct spatial patterns, with *Pdyn*, *Trh*, and *Vip* showing
region-specific distributions and varying degrees of co-expression with serotonergic markers.
345
Further, we investigated how DR gene expression is modified by SSRI treatment.
Comparison of acute and chronic treatment conditions revealed extensive transcriptional
changes. Gene enrichment and network analysis identified key modulated pathways among
the 232 differentially expressed genes in the DR, with 10-12 genes annotated to each of the
Ras, MAPK, and cAMP signaling cascades, as well as axonal guidance pathways. SSRI-
associated deficits in axonal guidance have been documented, particularly during
development[39], and chronic fluoxetine treatment reduces both the density and morphology of
serotonergic projections to the PFC and hippocampus[40,41]. Our findings show that chronic
treatment decreased expression of several axonal guidance genes, including *Plxnc1*. Network
analysis identified multiple RAS pathway regulators as hub genes. Changes in these hub
genes and other highly regulated genes including *Bdnf*, *Csf1r*, *Nras*, *Rras*, *Fos*, *Met*, *Kit*, *Myl9*,
and *Ntrk1* align with the neurotrophic hypothesis of antidepressant action[42].
358
Our analysis revealed significant changes in neuropeptide expression, potentially
underlying direct behavioral effects of SSRIs. Of particular interest was the regulation of *Pdyn*,
which encodes prodynorphin, an opioid peptide that activates kappa opioid receptors (KOR).
While KOR activation produces analgesia, it also induces dysphoria and depressive-like
behaviors, distinct from other opioid receptor subtypes[43]. *Pdyn* expression occurs in multiple
limbic regions, including the amygdala, hippocampus, nucleus accumbens, and
hypothalamus, and has recently been identified in a subpopulation of DR serotonin neurons[10-
12.] Although KOR signaling has been implicated in stress-induced depressive symptoms
through its interaction with the dopaminergic system, its relationship with serotonergic function
in mood regulation remains poorly understood. Studies using the forced swim test have shown
that KOR disruption reduces depression-like behavior[44], supporting KOR signaling's pro-
depressive role. Our findings revealed biphasic regulation of *Pdyn*, with increased expression
following acute SSRI treatment and subsequent decrease after chronic administration. This
initial upregulation may contribute to the documented adverse effects during early SSRI
treatment, including increased anxiety and suicide risk[45,46].
374
The neuropeptide-coding gene *Trh* showed opposing temporal regulation compared to
Pdyn, with downregulation after acute treatment and upregulation following chronic SSRI
administration. While thyrotropin-releasing hormone (TRH) is primarily known for its role in the
hypothalamic–pituitary–thyroid axis, it is also expressed in limbic regions, including the
hippocampus, amygdala, and lateral DR. TRH signals through two receptors (TRHR1 and
TRHR2) expressed throughout these regions, suggesting broader functions beyond endocrine

regulation. Evidence supports TRH involvement in cognitive and emotional processing[47], with both TRHR1 and TRHR2 knockout mice exhibiting depression- and anxiety-like behaviors[48,49]. Clinically, depression is associated with blunted TRH response and hypothyroidism[47], though the distinct roles of region-specific TRH systems remain unclear. Our observation that chronic SSRI treatment upregulates *Trh* expression in the DR suggests a potential mechanism contributing to therapeutic efficacy. Notably, neither *Trh* nor *Pdyn* regulation showed sex-specific effects during SSRI treatment.

Our findings provide molecular support for the $5HT_{1a}$ desensitization hypothesis, with *Htr1a* showing upregulation following acute treatment and downregulation after chronic administration. We observed reduced *Fos* expression after chronic SSRI treatment, suggesting decreased neuronal activity. Chronic treatment particularly suppressed activity in the midline DR cluster. These transcriptional changes present an interesting contrast with previous electrophysiological and immunohistochemical studies, which showed acute SSRI treatment initially decreases serotonergic neuron activity before returning to baseline during chronic treatment, with variable region-dependent effects[17,50-52]. This apparent discrepancy might be explained in two ways. First, our spatial transcriptomics data represents mini-bulk measurements, potentially capturing SSRI-induced changes in non-serotonergic neurons within the analyzed regions. Alternatively, distinct serotonergic subtypes may respond differently to SSRI treatment, with some populations showing decreased activity while others maintain or increase their firing rates. Future cell type-specific studies will be crucial to resolve these possibilities and elucidate the temporal dynamics of SSRI effects across distinct neuronal populations.

Our analysis reveals spatial molecular heterogeneity within the DR and provides new insights into region-specific SSRI actions. While our findings support established theories of SSRI mechanisms, including $5HT_{1a}$ desensitization and neurotrophic effects, we also uncovered a potential novel neuropeptide-based mechanism involving dynamic regulation of *Pdyn* and *Trh*. The opposing temporal patterns of these neuropeptides may contribute to both the initial adverse effects and subsequent therapeutic benefits of SSRI treatment. The candidate genes, pathways and cell types uncovered pave the way for deeper mechanistic studies to unravel the underpinnings of depression, and reveal novel targets for more specific pharmacological targeting.

Acknowledgements

This study was supported by SciLifeLab and Wenner Gren Foundations grants to I.P.D, and Polish National Agency for Academic Exchange (The Bekker Programme - BPN/BEK/2021/1/00152) and SciLifeLab RED Postdoctoral Fellowship to J.M. The authors acknowledge support from the National Genomics Infrastructure in Stockholm funded by Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish Research Council, and SNIC/Uppsala Multidisciplinary Center for Advanced Computational Science for assistance with massively parallel sequencing and access to the UPPMAX computational infrastructure. This work was supported by the National Bioinformatics Infrastructure Sweden (NBIS) at SciLifeLab. We thank A.R. Costa, and all the members of the Pollak Dorocic lab for thoughtful discussion, and C. Broberger for generous use of technical equipment.

Author contributions

C.H. performed the experiments, analysed and visualised the data. J.M. contributed to spatial transcriptomic analysis and produced the network analysis. I.P.D. conceived and supervised the study. All authors wrote and edited the manuscript.

Data availability
Data has been deposited at 10.5281/zenodo.14859692 and is also accessible through an interactive app available at https://st-ssri.serve.scilifelab.se/app/st-ssri. Any additional information and code details are available from the lead contacts upon request.

Competing interests
The authors declare no competing interests.

Figure 1. Spatial transcriptomic profiling reveals distinct molecular domains in the mouse midbrain and DR

A. Experimental strategy: Coronal sections of mouse midbrain including the DR were cut and multiple sections were mounted on Visium slide capture areas, followed by RNA sequencing and tissue alignment.

B. UMAP plot based on the transcriptional signature of each spot, covering the whole midbrain section, revealed 13 molecular clusters. Acronyms in cluster names: PAG, periaqueductal gray; DR, dorsal raphe; LDTg, laterodorsal tegmental nucleus; LC, locus coeruleus.

C. Representative midbrain tissue section with overlaid sequencing spots, color represents molecular cluster in (B).

D. Visualization of four anatomically distinct spatio-molecular clusters, including DR (top left), lateral DR (top right), lateral PAG (LPAG, bottom left) and dorsal PAG (bottom right).

E. Dot plot for each molecular cluster with expression level of individual genes. Diameter of the dot corresponds to the percentage of spots in the cluster expressing the gene, color indicates average expression of the gene.

Figure 2. Spatio-molecular mapping of DR reveals specific spatial markers

A. UMAP plot based on the transcriptional signature of each DR spot, cluster names refer to the spatial location or molecular identity.

B. Molecular clusters from A overlaid with tissue section, showing spatial localization of each cluster in DR and rosto-causal expression patterns.

C. Dot plot showing expression of 10 cluster marker genes per cluster. Diameter of the dot corresponds to the percentage of spots in the cluster expressing the gene, color indicates average expression of the gene.

D. Spatial heatmap of gene expression in DR for selected marker genes. Example of DR-specific markers (eg. midline-*Slc17a8, Pdyn*) and non-DR-specific marker genes (ventral-*Apod*).

E. In-situ hybridization in DR (area within broken white border) with 8 probes*: Slc6a4, Gad2, Slc17a8, Htr1a, Pdyn, Trh, Bdnf, Vip,* each displaying a distinct single-cell distribution pattern within DR. Scale bar = 100µm.

F. Quantification of the co-expression of each marker gene from (E) with serotonin marker *Slc6a4* (SERT).

Figure 3. Acute and chronic SSRI treatment elicits distinct gene expression changes in the DR

A. Schematic representation of the pharmacological experiment.

B. Gene enrichment analysis in the KEGG database of DEGs in the DR. Number in the bar represents the number DEGs observed across all treatment conditions.

C. Expression profiles of single DEGs in the DR, categorized by the involvement in specific KEGG pathways.

D. List of DEGs in the DR categorized by function from the Mouse Genome Informatics database and their expression levels across control, acute and chronic treatment conditions.

E. Upset plot of upregulated genes in the DR, between treatment conditions.

F. Upset plot of downregulated genes in the DR, between treatment conditions.

Figure 4. Network analysis of DR DEGs identifies key regulatory hubs across SSRI treatment
Protein-protein interaction network based on DEGs in DR in STRING database for changes in each comparison condition: control vs. acute (A), control vs. chronic (B) and acute vs. chronic (C). Hub genes are marked in a diamond node (◆). Red color denotes upregulation, blue denotes downregulation in comparison to control (A-B) or acute (C) group.

Figure 5. Dynamic gene expression changes of *Pdyn*, *Trh* and *Slc17a8* in DR following acute and chronic SSRI treatment

A. Representative images of in-situ hybridization of *Pdyn* and *Slc6a4* co-expression and *Pdyn* in the DR. Box plots comparing treatment groups: sum of *Pdyn*+ cells per section (one-way ANOVA, F(2,15) = 6.64, p = 0.009**. Pairwise t-test with Holms correction; Acute vs Control, p=0.0072**), sum of *Pdyn* spots (transcripts) per section (one-way ANOVA, F(2,15) = 5.735, p = 0.014*. Pairwise t-test with Holms correction; Acute vs Control, p=0.0125*), sum of cells co-expressing *Pdyn* and *Slc6a4* per section (one-way ANOVA, F(2,15) = 5.004, p = 0.022*. Pairwise t-test with Holms correction; Acute vs Control, p=0.0255*), and sum of *Pdyn* spots in *Slc6a4* cells. Control n = 6, acute n=6, chronic n=6 mice per group.

B. Representative images of in-situ hybridization of *Trh* and *Slc6a4* co-expression and *Trh* in the DR. Box plots comparing treatment groups of sum of *Trh* cells per section (one-way ANOVA, F(2,13) = 10.29, p = 0.002**. Pairwise t-test with Holms correction; Acute vs Chronic, p = 0.00132**; Chronic vs Control, p=0.00418**, control n = 5, acute n=6, chronic n=5 mice per group), sum of *Trh* spots (transcripts) per slice (one-way ANOVA, F(2,11) = 5.79, p = 0.019*. Pairwise t-test with Holms correction; Acute vs Chronic, p = 0.039*; Chronic vs Control, p=0.039*, control n = 4, acute n=5, chronic n=5 mice per group), sum of cells co-expressing *Trh* and *Slc6a4* per section (control n = 5, acute n=6, chronic n=4 mice per group) and sum of *Trh* spots in *Slc6a4* cells (control n = 4, acute n=5, chronic n=4 mice per group).

C. Representative images of in-situ hybridization of *Slc17a8* and *Slc6a4* co-expression and *Slc17a8* in the DR. Box plots comparing treatment groups of sum of *Slc17a8* cells per section, sum of *Slc17a8* spots (transcripts) per slice, sum of cells co-expressing *Slc17a8* and *Slc6a4* per and sum of *Slc17a8* spots in *Slc6a4* cells. Control n = 6, acute n=6, chronic n=6 mice per group.

Dots represent individual animals, p-values shown follow p < 0.05 = *, p < 0.01 = **, p < 0.001 = ***. Scale bar = 100µm. Boxplots show the data distribution in each group.

Figure S1. UMAP plot of midbrain transcriptomic data show no effect of sample, sex, and treatment in clustering

Lack of the technical- or batch- driven effects on clustering results in UMAP plot (**A**) was visualized by overlaying sample ID from different animals (**B**), their sex (**C**) or treatment group (**D**).

Figure S2. Regional expression and co-localization with serotonin for multiple markers in the DR along the rostro-caudal axis

Spatio-molecular map for each marker gene (*Pdyn, Trh, Slc17a8, Htr1a, Vip, Bdnf, Gad2*) in the DR, along the rostro-caudal axis (-4.3mm to -4.6mm). Co-expression of each specific marker with *Slc6a4* (SERT) is marked by a black circle (each circle correspond to a single cell), Slc6a4 positive cells lacking marker co-expression are marked in red, and the specific marker cells lacking Slc6a4 expression are shown in their respective color.

Figure S3. Mapping and deconvolution of serotonin subtypes reveals distinct spatial domains in DR

A. Integration of five molecular subtypes of DR serotonin neurons from published scRNA-seq dataset (5-HT-I through V from Huang, et al. 2019) mapped to spatial location across the rostro-caudal axis, using CytoSPACE computational deconvolution.

B. Spatial transcriptomic clusters overlaid with tissue section, showing spatial localization of each cluster in DR and rostro-caudal expression patterns.

C. Expression of top genes from spatial clusters (Caudal, Lateral 1 and Midline) mapped to five serotonin subtypes (5-HT-I through V). Diameter of the dot corresponds to the percentage of spots in the cluster expressing the gene, colour indicates average expression of the gene.

D. Expression of top genes from spatial clusters (Caudal, Lateral 1 and Midline). Diameter of the dot corresponds to the percentage of spots in the cluster expressing the gene, colour indicates average expression of the gene.

E. The distribution of serotonergic subtypes (5-HT-I through V) was quantified across three spatial domains (Caudal, Lateral 1, and Midline), expressed as the percentage of each subtype's total mapped cells within these clusters.

Figure S4. SSRI-induced changes in gene expression and activity in midbrain and DR

A. Heatmap visualization of the number of DEGs in clusters from whole midbrain section, across treatment conditions. The clusters correspond to UMAP in Fig. 1B.

B. Results of gene enrichment analysis in Gene Ontology, Biological Processes of DEGs in the DR across all treatment conditions.

C. Volcano plots representing top up- and down-regulated genes in the DR across treatment conditions.

D. Box plot comparing DR cluster neural activity between treatment groups, activity per transcriptomic spot generated by NEUROeSTIMator (one-way ANOVA followed by Tukey multiple comparisons of means, $F(2,443) = 4.101$, $p = 0.0172^*$; Control vs Chronic, $p = 0.0229^*$), DR Midline vs Lateral 1 cluster (one-way ANOVA followed by Tukey multiple comparisons of means, $F(2,219) = 6.022$, $p = 0.0028^{**}$; Chronic vs Acute, $p = 0.0466^*$), and the whole dataset (no difference). Control n = 3, acute n=3, chronic n=3 mice per group. Dots represent individual spots in the ST dataset, p-values $p < 0.05 = ^*$, $p < 0.01 = ^{**}$, $p < 0.001 = ^{***}$.

Figure S5. Gene expression of Bdnf, Gad2, Htr1a and Vip in the DR across SSRI treatment

A. Representative images of in-situ hybridization of *Bdnf* and *Slc6a4* co-expression and *Bdnf* in the DR. Box plots comparing treatment groups: sum of *Bdnf*+ cells per section, sum of *Bdnf* spots (transcripts) per section, sum of cells co-expressing *Bdnf* and *Slc6a4* per section, and sum of *Bdnf* spots in *Slc6a4* cells.
B. Representative images of in-situ hybridization of *Gad2* and *Slc6a4* co-expression and *Gad2* in the DR. Box plots comparing treatment groups: sum of *Gad2*+ cells per section, sum of *Gad2* spots (transcripts) per section, sum of cells co-expressing *Gad2* & *Slc6a4* per section and sum of *Gad2* spots in *Slc6a4*+ cells.
C. Representative images of in-situ hybridization of *Htr1a* and *Slc6a4* co-expression and *Htr1a* in the DR. Box plots comparing treatment groups of sum of *Htr1a*+ cells per section, sum of *Htr1a* spots per slice, sum of cells co-expressing *Htr1a* & *Slc6a4* per and sum of *Htr1a* spots in *Slc6a4* + cells.
D. Representative images of in-situ hybridization of *Vip* and *Slc6a4* co-expression and *Vip* in the DR. Box plots comparing treatment groups of sum of *Vip*+ cells per section and sum of cells co-expressing *Vip* and *Slc6a4* per slice.
Scale bar = 100μm, control n = 5-6, acute n=5-6, chronic n=6 mice per group.

Figure S6. Lack of sex-specific modulation of gene expression in DR by SSRI treatment
No sex effects are observed due to acute nor chronic SSRI treatment in the expression of *Pdyn*, *Trh*, *Slc17a8*, *Gad2*, *Bdnf*, *Htr1a* and *Vip*.
A. Box plots comparing treatment groups divided by sex: sum of *Pdyn*+ cells per section, sum of cells co-expressing *Pdyn* and *Slc6a4* per section.
B. Box plots comparing treatment groups divided by sex: sum of *Trh*+ cells per section, sum of cells co-expressing *Trh* and *Slc6a4* per section. Control n = 5, acute n=6, chronic n=5/n=4 mice per group.
C. Box plots comparing treatment groups divided by sex: sum of *Slc17a8*+ cells per section, sum of cells co-expressing *Slc17a8* and *Slc6a4* per section.
D. Box plots comparing treatment groups divided by sex: sum of *Gad2*+ cells per section, sum of cells co-expressing *Gad2* and *Slc6a4* per section. Control n = 5, acute n=6, chronic n=6 mice per group.
E. Box plots comparing treatment groups divided by sex: sum of *Bdnf*+ cells per section, sum of cells co-expressing *Bdnf* and *Slc6a4* per section.
F. Box plots comparing treatment groups divided by sex: sum of *Htr1a*+ cells per section, sum of cells co-expressing *Htr1a* and *Slc6a4* per section.
G. Box plots comparing treatment groups divided by sex: sum of *Vip*+ cells per section, sum of cells co-expressing *Vip* and *Slc6a4* per section.
Two-way ANOVA using Sidak multiple comparisons. Unless otherwise specified control n = 6, acute n=6, chronic n=6 mice per group.

Methods

Animals

All experiments were carried out in adult C57BL/6J (Charles River) male and female mice. Mice were maintained under standard housing conditions with a 12-hour light cycle and with ad libitum access to food and water. All animal experiments had received approval from the local ethical board, Stockholms Norra Djurförsöksetiska Nämnd, and were performed in accordance with the European Communities Council Directive 2010/63. For spatial transcriptomics experiments, 9 mice (5 male, 4 female) were used, for *in-situ* hybridization 17 mice (9 females, 8 males) were used.

Fluoxetine administration

Fluoxetine Hydrochloride (Sigma-Aldrich, F132) was dissolved in saline and stored at -20C° until use. For administration the fluoxetine hydrochloride was diluted to 2 mg/ml and administered to the mice by gavage in 10 mg/kg dose. The dose was calculated to be within the range of human equivalent doses prescribed for depression. The standard dose of fluoxetine for treatment of depression in humans is 20mg/day, with the highest being 80 mg/day[1]. Assuming an adult human weighs 60kg,the range of doses are 0.333-1.333 mg/kg. converting the 10 mg/kg mouse dose into the human equivalent is 0.81 mg/kg[2].

The gavage procedure was performed with soft flexible feeding tubes to minimize the discomfort and potential damage to the animal. Mice in the acute treatment group received a single 10mg/kg dose of fluoxetine and brain tissue was harvested after 1,5h. Fluoxetine has been shown to produce behavioural effects in C57BL/6J mice after 30 min[3]. Mice in the chronic treatment group received 10 mg/kg/day for 22 consecutive days. The brain was harvested 5 h after the final dose to avoid any acute dose effect from the final dose. The control group received a single dose of saline instead at the equivalent volume to weight followed by brain harvest after 1.5 h.

Tissue preparation

Isoflourane was used to render the animals unresponsive before cervical dislocation. The animals were decapitated, the brain extracted and immediately placed in ice cold saline for a maximum of 15 sec. For spatial transcriptomics experiments, the midbrain was dissected on an ice cold metal block of approximately 5x5x2 mm. The brains utilized for RNAscope experiments remained whole. The tissue was flash frozen using powdered dry ice and was stored airtight at -80C° prior to sectioning. All tools and surfaces were cleaned with 70% ethanol and RNAase away (Thermo Scientific, 10666421) prior to and during the harvesting.

For spatial transcriptomics, the dissected midbrains were trimmed, labelled and embedded together in O.C.T. (VWR, 361603E). The blocks were sectioned on a cryostat in 10 μm coronal sections that were placed on the Visium slide capture areas. 3-4 trimmed brain sections were placed in single capture areas. For RNAscope experiments, whole brains were embedded in O.C.T., cut in 20 μm coronal sections on the cryostat and placed on SuperFrost+ slides and allowed to dry in the cryostat cold chamber. The slides were stored airtight at -80C° until either RNAscope or Visium experiment start.

Fluorescent in-situ hybridization

In situ hybridization to detect expression of genes of interest was performed using Hiplex RNAscope technology following the manufacturer's protocol for fresh frozen tissue (Advanced Cell Diagnostics, ACD, Hayward, CA). In brief, 20μm sections were fixed in 4% PFA (Sigma-Aldrich, 252549) solution (60 min, RT), dehydrated for 5 min each in 50%, 75%, 99% and 99% laboratory grade ethanol (Fisher Chemical E/0600DF/F17). The slides were stored overnight in 99% ethanol at -20C°. The next day the slides dried at RT and a hydrophobic barrier was drawn around each tissue section. Protease IV was added to permeabilize the

tissue sections (30 min, RT). After washing, each section was covered in probe dilution of 8 probes, or with positive and negative control probes, and incubated at 40C° for 2h in a humidity controlled environment. With wash steps between each incubation three amplification steps were performed, each 30 min 40C°. To reduce autofluorescence the slices were incubated with 3% FFPE reagent mix 30 min RT prior to 'HiPlex Fluoro T1–T4 v2', 15 min 40C°. After washing, DAPI staining was applied followed by clear mounting media and the first round of imaging.

The next day the sections were washed, and fresh cleaving solution was applied for 15 min RT two consecutive times, with washes in between to ensure removal of signal from previous imaging round. The slides were washed and incubated with 3% FFPE reagent mix 30 min RT, followed by 'HiPlex Fluoro T5–T8 v2 ' for 15 min 40C°. After washing, clear mounting media was added, followed by the second imaging round. The following probes were used: Htr1a-T1, Vip-T2, Bdnf-T3, Pdyn-T4, Trh-T5, Slc6a4-T6, Slc17a8-T7, Gad2-T8.

Microscopy

Imaging was performed on a Leica DM6 B stand up microscope with 40x oil immersion lens using the Leica developed LasX software. Intensities and exposure were setup on a separate tissue to minimize tissue bleaching. Each section was scanned in brightfield and the image ROI was selected and saved. Several focus points were added for each ROI to ensure uniform focus throughout the stitched image. Each slice was imaged in 5 channels (DAPI, 488, 550, 650, 750) with a z-stack of 8x8 um depth over two separate imaging rounds. The LasX Mosaic Merge stitching tool was used to merge the individual images together, using the blend 'smooth' setting.

Image analysis

All images were imported to Fiji (ImageJ 2.14.0), using Bio-Formats importer to create a single max intensity projection image for all four channels. The max intensity image from imaging round 1 & 2 of the same slice was uploaded into a Qupath (0.5.0) project[4]. Using the Fiji/Qupath plugin, BigWarp in Fiji was utilized to create coordinate maps to overlap the two rounds of imaging based on the DAPI staining in the same slice.

Cell segmentation was performed in Qupath with Stardist scripts using their dsb2018_heavy_augment.pb model, identifying nuclei based in DAPI staining with 5.0 cell expansion and 0.5 threshold[5]. A region of interest was selected that included all Slc6a4+ cells, and corresponded with the Dorsal Raphe Nucleus location. Within this ROI Qupath's own 'Object Classification' was utilized to assign cell identity based on marker expression. Object classifiers were trained either on the image itself or on other images in the data set. Individual RNA puncta were quantified using Qupaths subcellular detection feature. The detection threshold was set individually for the image based on a combination of the autofluorescence of the image and the fluorescence strength seen in the negative control for the specific channel.

Objects and annotations and their measurements were transferred to the Round 2 image using the Warpy plugin "transfer annotations and measurements" .

Two rounds of exports were merged based on the shared cell object ID into a single data frame saved per animal and slice. All slices were combined into a larger data frame from which summary tables were produced to assess sum or cells, sum of spots, both per marker and co-expressing subpopulations. We tested our data for normal distribution and initially performed ANOVA to compare our three treatment groups, followed by pairwise t-test adjusted with the Holm-correction, if the ANOVA was statistically significant ($p < 0.05$).

Spatial transcriptomics

The RNA integrity of the frozen tissue sections was measured with a bioanalyzer (Agilent) and was above 8 RIN for all samples. The 10x Genomics Visium protocol was performed by National Genomics Infrastructure at SciLifeLab, Sweden. Briefly, the sections were fixed with methanol at -20C° before performing H&E staining and bright field imaging. The tissue was then permeabilized enzymatically and the RNA was captured on the barcoded probes attached to the glass slide by the poly-A tail. The remaining tissue was washed away, and libraries were generated for each capture area and sequenced. Manual alignment of the H&E stained histology images to the expression spot grid was performed using the 10x Genomics Loupe Browser software (v. 5.1.0). The raw sequencing data files (FASTQ files) for the sequenced library samples were processed using the 10x Genomics Space Ranger software (v. 1.3.0) using the mouse genome reference transcriptome version GRCh38 2020-A (July 7, 2020) provided by 10x Genomics.

Analysis of Visium sequencing data

For each capture area the respective histological image was used for data alignment. The data was processed in R 4.2.2 with Seurat (V4.3.0).

Spots were filtered based on cutoff values on several QC parameters: spots containing less than 1000 genes, spots containing less than 1000 nFeatures, spots containing more than 24 % mitochondrial genes. Genes from a number of categories were removed from further analysis, which included haemoglobin genes, mitochondrial genes, ribosomal genes and X/Y chromosome genes. For each capture area, data was normalized using SCTransform with 'glmGamPoi' method, v2 vst.flavor and regressing out % mitochondrial genes and % ribosomal genes. Only a set of variable genes was returned for further analysis. In order to minimize batch effects, data across the capture areas was integrated using Reciprocal PCA based on top 3000 features and top 30 principal components. To find integration anchors we used capture area containing slices from the control-treated group as a reference and K = 10.

Clustering was performed using the original Louvain algorithm and resolution = 0.9. To visualize data points in two-dimensional space, we have performed Uniform Manifold Approximation and Projection (UMAP) on the first 20 principal components using 50 closest neighbours and spread = 2. Three clusters were excluded based on criteria that it must be present on a minimum two slices per condition. After initial clustering, we selected a cluster with elevated Tph2 expression as a DR cluster and subjected it to subclustering by using the same algorithms but changed the number of dimensions to top 8 and resolution to 0.8. We have carefully selected above mentioned algorithms and parameters by comparing clustering stability, purity and silhouette width. To find markers and perform differential gene expression analysis, we used FindMarkers() function on "data" slot from integrated dataset and compared medians by Wilcoxon Rank Sum test. We have set "logfc" threshold to 0.5 and minimum expression in at least 20% of cluster's spots. Differential gene expression was analysed comparing a single cluster with the rest of the dataset to find cluster marker genes and within the same cluster between treatment conditions to find treatment affected genes. For multiple comparison correction, we used False Discovery Rate and only changes with p-value < 0.05 were considered significant.

Network analysis: To identify key genes in DR affected by SSRI treatment, we created a Protein–Protein Interaction (PPI) Network with the STRING database using the list of all DEGs detected in DR, with significance cutoff value of FDR < 0.05 and minimum expression in at least 20% of the cluster spots. Network was then visualized and analyzed in Cytoscape (3.10.0). Based on the degree distribution in all three networks, the degree threshold value for hub genes was set to 6. The PPI network incorporates protein interactions known from experiments, text mining, databases, co-expression, co-occurrence, neighbourhood and gene fusion.

Gene enrichment analysis: In order to investigate the potential functional effects of the SSRI treatment in the DR we performed gene enrichment analysis through The Database for Annotation, Visualization and Integrated Discovery (DAVID)[6,7] bioinformatics. We uploaded all DEGs detected in the DR, with significance cutoff value of FDR < 0.05 and minimum expression in at least 20% of the cluster spot, as a list in the DAVID analysis wizard and selected annotation categories KEGG pathways and GO terms Biological processes_5.

Activity analysis with NEUROeSTIMator: Using the gene-level counts per Visium spot as input for the NEUROeSTIMator[8] v0.1. One-way Anova with following Tukey's test were performed for statistical analysis.

Deconvolution with CytoSPACE, mapping single cell seq data to Visium: Huang et al, 2019 single cell sequencing data was used for cell type deconvolution performed with CytoSPACE[9] v1.0.5, default settings. Estimated number of cells was set to 9 based on VistoSeg[10]. All available cell types were mapped, but only serotonin subtypes were isolated for visualization and further quantification.

Reproducibility and sample size

Sample size based on other previous research with no calculations done prior to the experiments. All experiments contained between 3-6 biological replicates per group. Data was excluded based on raw data quality, if dissection of the slices did not contain the same clusters in all treatment groups, or if the cluster contained too few data points for further analysis.

References for methods:

1. Altamura, A. C., Moro, A. R. & Percudani, M. Clinical Pharmacokinetics of Fluoxetine. *Clin. Pharmacokinet.* **26**, 201–214 (1994).

2. Nair, A. B. & Jacob, S. A simple practice guide for dose conversion between animals and human. *J Basic Clin Pharm* **7**, 27–31 (2016).

3. ZI, J. *et al.* Mouse strain differences in SSRI sensitivity correlate with serotonin transporter binding and function. *Scientific reports* **7**, (2017).

4. Bankhead, P. *et al.* QuPath: Open source software for digital pathology image analysis. *Sci Rep* **7**, 16878 (2017).

5. Stevens, M. *et al.* StarDist Image Segmentation Improves Circulating Tumor Cell Detection. *Cancers (Basel)* **14**, 2916 (2022).

6. Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and functional annotation of gene lists (2021 update). *Nucleic Acids Res* **50**, W216–W221 (2022).

7. Huang, D. W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44–57 (2009).

8. Bahl, E. et al. Using deep learning to quantify neuronal activation from single-cell and spatial transcriptomic data. Nat Commun **15**, 779 (2024).

9. Vahid, M. R. et al. High-resolution alignment of single-cell and spatial transcriptomes with CytoSPACE. Nat Biotechnol **41**, 1543–1548 (2023).

10. Tippani, M. et al. VistoSeg: Processing utilities for high-resolution images for spatially resolved transcriptomics data. Biological Imaging **3**, e23 (2023).

References

1.   Moreno-Agostino, D. *et al.* Global trends in the prevalence and incidence of depression:a systematic review and meta-analysis. *Journal of Affective Disorders* **281**, 235–243 (2021).

2.   OECD. *Pharmaceutical Market*. https://www.oecd-ilibrary.org/content/data/data-00545-en (2016).

3.   Luo, Y., Kataoka, Y., Ostinelli, E. G., Cipriani, A. & Furukawa, T. A. National Prescription Patterns of Antidepressants in the Treatment of Adults With Major Depression in the US Between 1996 and 2015: A Population Representative Survey Based Analysis. *Frontiers in Psychiatry* **11**, (2020).

4.   Ferguson, J. M. SSRI Antidepressant Medications: Adverse Effects and Tolerability. (2001).

5.   Cools, R., Roberts, A. C. & Robbins, T. W. Serotoninergic regulation of emotional and behavioural control processes. *Trends Cogn Sci* **12**, 31–40 (2008).

6.   Pollak Dorocic, I. *et al.* A Whole-Brain Atlas of Inputs to Serotonergic Neurons of the Dorsal and Median Raphe Nuclei. *Neuron* **83**, 663–678 (2014).

7.   Weissbourd, B. *et al.* Presynaptic Partners of Dorsal Raphe Serotonergic and GABAergic Neurons. *Neuron* **83**, 645–662 (2014).

8.   Ogawa, S. K., Cohen, J. Y., Hwang, D., Uchida, N. & Watabe-Uchida, M. Organization of monosynaptic inputs to the serotonin and dopamine neuromodulatory systems. *Cell Rep* **8**, 1105–1118 (2014).

9.   Muzerelle, A., Scotto-Lomassese, S., Bernard, J. F., Soiza-Reilly, M. & Gaspar, P. Conditional anterograde tracing reveals distinct targeting of individual serotonin cell groups (B5–B9) to the forebrain and brainstem. *Brain Struct Funct* **221**, 535–561 (2016).

10.   Ren, J. *et al.* Single-cell transcriptomes and whole-brain projections of serotonin neurons in the mouse dorsal and median raphe nuclei. *eLife* **8**, e49424 (2019).

11.   Huang, K. W. *et al.* Molecular and anatomical organization of the dorsal raphe nucleus. *eLife* **8**, e46464 (2019).

12.   Okaty, B. W. *et al.* A single-cell transcriptomic and anatomic atlas of mouse dorsal raphe Pet1 neurons. *eLife* **9**, e55523 (2020).

13.   Cardozo Pinto, D. F. *et al.* Characterization of transgenic mouse models targeting neuromodulatory systems reveals organizational principles of the dorsal raphe. *Nat Commun* **10**, 4633 (2019).

14.   Fu, W. *et al.* Chemical neuroanatomy of the dorsal raphe nucleus and adjacent structures of the mouse brain. *Journal of Comparative Neurology* **518**, 3464–3494 (2010).

15.   Hervás, I. & Artigas, F. Effect of fluoxetine on extracellular 5-hydroxytryptamine in rat brain. Role of 5-HT autoreceptors. *Eur J Pharmacol* **358**, 9–18 (1998).

16.   Stahl, S. M. Mechanism of action of serotonin selective reuptake inhibitors: Serotonin receptors and pathways mediate therapeutic effects and side effects. *Journal of Affective Disorders* **51**, 215–235 (1998).

17.   Czachura, J. F. & Rasmussen, K. Effects of acute and chronic administration of fluoxetine on the activity of serotonergic neurons in the dorsal raphe nucleus of the rat. *Naunyn-Schmied Arch Pharmacol* **362**, 266–275 (2000).

18.   Kreiss, D. S. & Lucki, I. Effects of acute and repeated administration of antidepressant drugs on extracellular levels of 5-hydroxytryptamine measured in vivo. *J Pharmacol Exp Ther* **274**, 866–876 (1995).

19.   Hensler, J. Differential regulation of 5-HT1A receptor-G protein interactions in brain following chronic antidepressant administration. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* **26**, (2002).

20.   Artigas, F., Romero, L., de Montigny, C. & Blier, P. Acceleration of the effect of selected antidepressant drugs in major depression by 5-HT1A antagonists. *Trends in Neurosciences* **19**, 378–383 (1996).

21.   de Montigny, C. & Blier, P. Effects of antidepressant treatments on 5-HT neurotransmission: electrophysiological and clinical studies. *Adv Biochem Psychopharmacol* **39**, 223–239 (1984).

22.   Commons, K. G. & Linnros, S. E. Delayed Antidepressant Efficacy and the Desensitization Hypothesis. *ACS Chem Neurosci* **10**, 3048–3052 (2019).

23.   Harmer, C. J., Duman, R. S. & Cowen, P. J. How do antidepressants work? New perspectives for refining future treatment approaches. *The Lancet Psychiatry* **4**, 409–418 (2017).

24.   Carrillo-Roa, T. *et al.* Common genes associated with antidepressant response in mouse and man identify key role of glucocorticoid receptor sensitivity. *PLOS Biology* **15**, e2002690 (2017).

25.   Conti, B. *et al.* Region-specific transcriptional changes following the three antidepressant treatments electro convulsive therapy, sleep deprivation and fluoxetine. *Mol Psychiatry* **12**, 167–189 (2007).

26.   Surget, A. *et al.* Corticolimbic Transcriptome Changes are State-Dependent and Region-Specific in a Rodent Model of Depression and of Antidepressant Reversal. *Neuropsychopharmacol* **34**, 1363–1380 (2009).

27.   Rayan, N. A. *et al.* Integrative multi-omics landscape of fluoxetine action across 27 brain regions reveals global increase in energy metabolism and region-specific chromatin remodelling. *Mol Psychiatry* 1–16 (2022) doi:10.1038/s41380-022-01725-1.

28.   Wyler, S. C., Donovan, L. J., Yeager, M. & Deneris, E. Pet-1 Controls Tetrahydrobiopterin Pathway and Slc22a3 Transporter Genes in Serotonin Neurons. *ACS Chem. Neurosci.* **6**, 1198–1205 (2015).

29.   Gras, C. *et al.* A Third Vesicular Glutamate Transporter Expressed by Cholinergic and Serotoninergic Neurons. *J. Neurosci.* **22**, 5442–5451 (2002).

30.   Liu, Z. *et al.* Dorsal Raphe Neurons Signal Reward through 5-HT and Glutamate. *Neuron* **81**, 1360–1374 (2014).

31.   Sengupta, A., Bocchio, M., Bannerman, D. M., Sharp, T. & Capogna, M. Control of Amygdala Circuits by 5-HT Neurons via 5-HT and Glutamate Cotransmission. *J. Neurosci.* **37**, 1785–1796 (2017).

32.   Cavaleri, D. *et al.* The role of BDNF in major depressive disorder, related clinical features, and antidepressant treatment: Insight from meta-analyses. *Neuroscience & Biobehavioral Reviews* **149**, 105159 (2023).

33. Vahid, M. R. et al. High-resolution alignment of single-cell and spatial transcriptomes with CytoSPACE. Nat Biotechnol **41**, 1543–1548 (2023).

34.   Altamura, A. C., Moro, A. R. & Percudani, M. Clinical Pharmacokinetics of Fluoxetine. *Clin. Pharmacokinet.* **26**, 201–214 (1994).

35.   Nair, A. B. & Jacob, S. A simple practice guide for dose conversion between animals and human. *J Basic Clin Pharm* **7**, 27–31 (2016).

36.   Yuasa-Kawada, J., Kinoshita-Kawada, M., Tsuboi, Y. & Wu, J. Y. Neuronal guidance genes in health and diseases. *Protein & Cell* **14**, 238–261 (2023).

37. Bahl, E. et al. Using deep learning to quantify neuronal activation from single-cell and spatial transcriptomic data. Nat Commun **15**, 779 (2024).

38.   Fischer, A. G., Jocham, G. & Ullsperger, M. Dual serotonergic signals: a key to understanding paradoxical effects? *Trends in Cognitive Sciences* **19**, 21–26 (2015).

39.   Homberg, J. R., Schubert, D. & Gaspar, P. New perspectives on the neurodevelopmental effects of SSRIs. *Trends in Pharmacological Sciences* **31**, 60–65 (2010).

40.   Nazzi, S. *et al.* Reversible Morphological Remodeling of Prefrontal and Hippocampal Serotonergic Fibers by Fluoxetine. *ACS Chem. Neurosci.* **15**, 1702–1711 (2024).

41.   Nazzi, S., Maddaloni, G., Pratelli, M. & Pasqualetti, M. Fluoxetine Induces Morphological Rearrangements of Serotonergic Fibers in the Hippocampus. *ACS Chem. Neurosci.* **10**, 3218–3224 (2019).

42.   Casarotto, P. C. *et al.* Antidepressant drugs act by directly binding to TRKB neurotrophin receptors. *Cell* **184**, 1299-1313.e19 (2021).

43.   Knoll, A. T. & Carlezon, W. A. Dynorphin, stress, and depression. *Brain Research* **1314**, 56–73 (2010).

44.   McLaughlin, J. P., Marton-Popovici, M. & Chavkin, C. κ Opioid Receptor Antagonism and Prodynorphin Gene Disruption Block Stress-Induced Behavioral Responses. *J. Neurosci.* **23**, 5674–5683 (2003).

45.   Ramos, R. T., Gentil, V. & Gorenstein, C. Clomipramine and initial worsening in panic disorder: beyond the 'jitteriness syndrome'. *J Psychopharmacol* **7**, 265–269 (1993).

46.   Grillon, C., Levenson, J. & Pine, D. S. A Single Dose of the Selective Serotonin Reuptake Inhibitor Citalopram Exacerbates Anxiety in Humans: A Fear-Potentiated Startle Study. *Neuropsychopharmacol* **32**, 225–231 (2007).

47.   Joffe, R. T. & Marriott, M. Thyroid Hormone Levels and Recurrence of Major Depression. *AJP* **157**, 1689–1691 (2000).

48.   Zeng, H. *et al.* Thyrotropin-Releasing Hormone Receptor 1-Deficient Mice Display Increased Depression and Anxiety-Like Behavior. *Molecular Endocrinology* **21**, 2795–2804 (2007).

49.   Sun, Y. *et al.* TRH-Receptor-Type-2-Deficient Mice are Euthyroid and Exhibit Increased Depression and Reduced Anxiety Phenotypes. *Neuropsychopharmacol* **34**, 1601–1608 (2009).

50.   Payet, J. M. *et al.* Exposure to Acute and Chronic Fluoxetine has Differential Effects on Sociability and Activity of Serotonergic Neurons in the Dorsal Raphe Nucleus of Juvenile Male BALB/c Mice. *Neuroscience* **386**, 1–15 (2018).

51.   Payet, J. M. *et al.* The Role of Dorsal Raphe Nucleus Serotonergic Systems in Emotional Learning and Memory in Male BALB/c Mice. *Neuroscience* **534**, 1–15 (2023).

52.   Payet, J. M. *et al.* Involvement of dorsal raphe nucleus serotonergic systems in social approach-avoidance behaviour and in the response to fluoxetine treatment in peri-adolescent female BALB/c mice. *Behavioural Brain Research* **408**, 113268 (2021).