Shaping Kale Morphology and Physiology Using Precision LED Light Recipes

Sabine Scandola[1,2,*], Lauren E. Grubb[1,*], Brigo Castillo[1], Lexyn Iliscupidez[1], Curtis Kennedy[1,3], Nicholas Boyce[1], Mohana Talasila[1], and R. Glen Uhrig[1, 4,#]

[1] Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada

[2] Lethbridge Research and Development Centre, Agriculture and Agri-Food Canada, Lethbridge, Canada

[3] Department of Computer Science, University of Alberta, Edmonton, Alberta, Canada

[4] Department of Biochemistry, University of Alberta, Edmonton, Alberta, Canada

[*] These authors contributed equally (S.S. and L.E.G.)

[#] Senior and corresponding author:

Dr. R. Glen Uhrig
Department of Biological Sciences,
University of Alberta,
Edmonton, Alberta, Canada, T6G 2E9
Email: ruhrig@ualberta.ca
Phone: +1 780 492 3088
Fax : +1 780 492-9234ruhrig@ualberta.ca

Running Title: Light condition effects on kale growth and development

Key Words: Kale, Anthocyanins, Light emitting diodes (LEDs), Horticulture, PhotosynQ, Quantitative Proteomics, Metabolomics

Abstract

Light emitting diodes (LEDs) present an innovative opportunity for manipulating plant growth and development. Light serves as a fundamental energy source and an environmental cue, providing an ability to control plant traits through the precise manipulation of light intensity and quality. Here, we assess effects of light intensity and spectral composition on the growth and physiology of the important model plant: Kale (*Brassica oleracea*), known for its phenotypic plasticity and nutritional composition, kale is a crop well-suited for indoor cultivation. Here, we employ a combination of advanced phenotyping, computer vision, gas chromatography-mass spectrometry (GC-MS) metabolomics, and liquid chromatography-mass spectrometry (LC-MS)-based quantitative proteomics to characterize the molecular changes that underpin light-dictated differences in the growth and metabolism of two different kale cultivars under different light intensity and spectral composition scenarios. We identify time-of-day and cultivar-specific light intensity and spectral composition-induced changes related to growth, shade avoidance, photosynthesis and several nutritional components including amino acids, glucosinolates and carotenoids. Our results offer a key resource to the plant community and demonstrate the translational potential of light manipulation in tailoring kale growth and nutritional content for enhanced crop productivity and/or health benefits, while simultaneously offering a more cost-effective solution for contemporary agricultural challenges.

1 Introduction

Kale (*Brassica oleracea*), also known as leaf cabbage, belongs to the large *Brassicaceae* family that includes cauliflower, broccoli and collard greens. Established as a superfood, this leafy green is rich in fiber and essential vitamins, as well as antioxidants such as flavonoids, each of which have been shown to positively affect cardiovascular and gastrointestinal health[1]. To date, the health benefits of kale have been heavily studied, with researchers continuing to identify new anti-cancer (e.g. sinigrin, spirostanol)[2] and neuroprotective (e.g. sulforaphane)[3] compounds, which neutralize reactive oxygen species. Further, kale demonstrates excellent temperature resilience resulting from its waxy coating and high production of polyamines[4], rendering it a valuable agricultural crop in the context of climate change[5].

Kale is also well-suited for indoor growing facilities. As a leafy green, it is a favored choice for horticultural growers as it has a quick harvest cycle with minimal space and resource requirements[6], appealing to both growers as well as consumers looking for fresh, locally-sourced produce[7]. Indoor / vertical farms have strongly co-evolved with new technologies as the promise of providing fresh and locally grown food largely relies on reducing production costs through technological advancements[8]. In this context, LED technology has revolutionized indoor farming, providing a wide range of energy-efficient and versatile lighting options. For example, unlike traditional light sources (e.g. incandescent bulbs, fluorescent tubes, metal halide or high pressure sodium bulbs), LED lighting systems offer an ability to fine-tune the spectral composition and intensity of the applied light[9]. Light is critical for photosynthesis, which drives plant metabolism[10], while plant photoreceptors, such as phytochromes and cryptochromes, respond to light quality, impacting processes such as plant growth (e.g. biomass) and development (e.g. flowering)[11]. However, despite our understanding of kale at a phenotypic level, we lack knowledge of how light impacts kale at the molecular level, limiting our ability to enhance genetics through targeted breeding or biotechnology approaches.

To date, elementary changes in spectral composition have been shown to affect kale phenotypic and metabolic profiles[12]. Blue light enhances growth and nutritional qualities[13], while red light affects kale pre- and post-harvest by altering nutrient and chlorophyll content[14]. Similar to red light, far-red light enhances photosynthetic rates through the Emerson effect[15]. Beyond specific light types, light wavelength ratios also play an essential role in how phytochromes

perceive and transduce light signals[16], offering opportunities to build precise light recipes that
maximize traits of interest.

To better exploit currently available kale genetics, we endeavored to precisely resolve the
impacts of light intensity and spectral composition on kale growth, development and nutritional
content by contrasting the diel molecular changes in two related kale cultivars: Dwarf curled
scotch (K3) and Red Scarlet (K9). While there are many kale cultivars, each with unique
morphologies, growth parameters and nutritional characteristics[17], our previous study examining
nine commercially available kale cultivars found that K3 and K9, both *Brassica oleracea var.*
sabellica, showed the most diversity in their phenotypes and molecular characteristics[17],
allowing us here to capture an in-depth understanding of how light properties impact kale growth
and development in order to establish a compendium of molecular changes that can be utilized
by the broader community. We specifically took a diel experimental approach to analyzing the
effects of light intensity and spectral composition by harvesting tissues at end-of-day (ED, ZT11
(zeitgeber)) and end-of-night (EN, ZT23) time-points as both the proteome and metabolome are
known to undergo substantial diel changes[17]. Understanding how external factors such as time-
of-day intersect with differences in light intensity and quality to impact kale physiology is
essential for determining optimal horticultural growing conditions and harvesting times to extract
maximize value, as well as inform future efforts in breeding more productive and nutritious kale
varieties.

2 Materials and Methods

2.1 Plant material and cultivation

Dwarf Curled Scotch (K3) and Red Scarlet (K9) kale seeds (*B. oleracea var. sabellica*) were
purchased from West Coast Seeds (https://www.westcoastseeds.com/). Seeds were surface
sterilized with incubations in 70% (v/v) ethanol and 30% (v/v) bleach solutions at room
temperature beginning with a 4-minute incubation in 3 mL of 70% (v/v) ethanol. The seeds were
then rinsed with 3 mL of sterile milli-Q water and 3 mL of 30% (v/v) bleach was added onto each
tube for a 7-minute incubation with periodical shaking every 2 minutes. Finally, the bleach
solution was decanted, and the seeds were rinsed three times with 3 mL of sterile Milli-Q water.
Following sterilization, seeds from each cultivar were plated onto 0.5 x Murashige and Skoog
(MS) media plates containing 1% (w/v) sucrose. The seeds were then stratified at 4 °C in total
darkness for 4 days. Following stratification, the kale seedlings were entrained to their

respective light treatment for 7 days. To prepare the soil for seedling transplant, ~25 kg of soil
(Sun Gro Sunshine Mix #1) was mixed with 8 L of H_2O and then transferred onto 4" square
polypropylene pots. Transplantation occurred once most plants had reached the cotyledon
stage at 11 days post-imbibition (dpi).

2.2 Light Conditions

The programmable LED (Perihelion Model) lights used for this experiment were provided by
G2V Optics Inc. and were installed inside ventilated growth chambers lined with reflective mylar.
The kale plants were grown under a 12-hour light and 12-hour dark regiment. Three different
light intensities were tested: 75, 125, 175 PPFD ($\mu mol.m^{-2}.s^{-1}$) (**Supp Figure 1A**). Spectral light
conditions were assigned in spectral ratios of R/B and R/FR of low R/B or R/FR, balanced light
and high R/B or R/FR (**Supp Figure 1B**).

2.3 Measurement of Kale Photosynthetic Performance via MultisynQ

Photosynthetic performance was measured following the Photosynthesis RIDES 2.0 protocol
using a MultispeQ v1.0 (PhotosynQ LLC, USA)[18]. At mid-day (zeitgeber; ZT6) 34 dpi, the
photosynthetic performance of the kale plants was measured by clamping the second true leaf
between the MultisynQ sensors ensuring full coverage. The leaf remained in the chamber until
the data collection process was completed (~10 s). A total of 10 randomly selected plants were
analyzed for each cultivar and light condition.

2.4 Phenotyping measurements

Plants were grown in separate chambers under different light intensity and spectral conditions
(**Supp Figure 1**). Each chamber was equipped with single-board Raspberry Pi 3 B+
computers and ArduCam Noir Cameras (OV5647 1080p) with a motorized IR-CUT filter and two
infrared LEDs. Pictures of plants were taken every 5 minutes. Plant surface area analysis
across light conditions was performed by averaging the area of each plant measured during a 1-
hour period (between ZT6 and ZT7) at 22 dpi. Plant surface areas were extracted using Python
scripts from the open-source software package PlantCV (https://plantcv.readthedocs.io/),
following the methodology previously described[19].

2.5 Harvesting and Sample Processing

At 35 dpi, kale leaves were harvested at two time points: end-of-night (ZT23) and end-of-day
(ZT11). This was done to capture the expansive metabolic changes occurring during the
transitions between night-to-day and day-to-night, respectively. Leaves from three kale plants of

the same cultivar were pooled into a labelled 50 mL conical tube and promptly frozen in liquid

nitrogen to preserve the proteins and metabolites present. In total, four tubes were collected

from each cultivar per chamber during harvest. Two metal beads were then introduced into each

conical tube and the samples were ground by subjecting them to 3x – 30 s/1200 rpm using a

Geno/Grinder[®] - Automated Tissue Homogenizer and Cell Lyser. 100 mg (±1 mg) of each finely-

ground sample was then portioned into each of two separate 2 mL Eppendorf Safe-Lock tubes

for further processing.

2.6 Anthocyanin Extraction and Quantification:

Anthocyanin analysis was performed as previously described by Neff and Chory (1998), with the

following modifications. Specifically, 500 µL of 100% (v/v) methanol-1% (v/v) hydrochloric acid

was added to each tube containing the finely ground samples and incubated at 4 °C for 24

hours in darkness. From there, 200 µL of H_2O and 500 µL of chloroform were added and the

tubes were centrifuged at 18,000 x g for 5 minutes at room temperature. 400 µL of the

supernatant was then transferred over to new tubes and adjusted to a total volume of 800 µL

with 60% (v/v) methanol-1% (v/v) hydrochloric acid. A total of 200 µL from each volume-

corrected sample was transferred into a well in a 96-well plate and absorbance at 530 nm and

657 nm was measured with a Tecan Spark® plate reader using 60% (v/v) methanol -1% (v/v)

hydrochloric acid as the background. The relative anthocyanin content was then derived

mathematically using: $Antho = (Abs_{530} - Abs_{657}) \times 1000 \times powder\ weight\ (mg)^{-1}$.

2.7 Metabolite analysis via Gas Chromatography Mass Spectrometry (GCMS)

2.7.1 Metabolite extraction: A total of 700 µL of ice-cold 100% (v/v) methanol was added to the

tubes containing 100 mg of finely ground leaf sample and promptly vortexed for ~20 seconds.

Using a thermomixer, these were then incubated for 15 minutes at 70 °C with shaking at 1200

rpm. From there, the tubes were centrifuged at 18,000 rpm for 15 minutes and the supernatants

were transferred into new tubes. The pellet was then re-extracted using 700 µL of 50% (v/v)

methanol containing Ribitol at 25 µL per sample at 0.4 mg/mL in water and vortexed for ~20

seconds until re-suspended. These tubes were then centrifuged at 14,000 rpm for 10 minutes

and 600 µL of the supernatant was combined with the previous supernatant. Finally, 100 µL was

transferred into new tubes and dried using a vacuum centrifuge at 1 torr at room temperature for

1.5 - 2 hours.

2.7.2 Gas chromatography mass spectrometry analysis:

The dried samples were derivatized by adding 100 µL of methoxamine hydrochloride into each
tube and then incubated at 30 °C at 850 rpm for 1 hour using a Thermomixer F2.0 (Eppendorf).
Subsequently, 50 µL of N,O-bis(trimethylsilyl)trifluoroacetamide (BSTFA) was added to each
tube and incubated at 50 °C at 850 rpm for 3 hours using a Thermomixer F2.0 (Eppendorf). A
total of 150 µL of each sample was then transferred to vials in preparation for injection in the
GC-MS. Samples were analyzed via a 7890A gas chromatograph (Agilent) paired to a 5975C
quadrupole mass detector (Aglient). The initial GC oven temperature was set at 70 °C and then
increased to 325 °C at 7 °C per minute two minutes post-injection where the temperature was
then maintained at 325 °C for 3.6 minutes. The injection and ion source temperatures were set
to 300 °C and 230 °C, respectively with a solvent delay of 5 minutes. The helium flow rate was
also adjusted to a rate of 1 mL/minute. Measurements were performed with electron impact
ionization (70 eV) in full scan mode (m/z 33-600). Identification of metabolites was performed
based on their mass spectral and retention time index, and how closely they matched to those
acquired from the National Institute of Standards and Technology library and the Golm
Metabolome Database.

2.8 *Protein extraction and LC-MS analysis*

2.8.1 Protein extraction and LC-MS data acquisition

Kale leaf tissue harvested at ZT23 and ZT11 was flash-frozen in liquid N_2. Tissue was ground in
liquid N_2 using a mortar and pestle and aliquoted into 400 mg samples (n=4 replicates).
Samples were extracted with 50 mM HEPES-KOH pH 8.0, 50 mM NaCl, and 4% (w/v) SDS at a
ratio of 1:2 (w/v). Next, samples were vortexed followed by incubation for 15 min at 95 °C using
a Thermomixer F2.0 (Eppendorf) shaking at 1,100 rpm. Samples were subsequently incubated
at room temperature for an additional 15 min of shaking followed by clarification at 20,000 x g
for 5 min at room temperature, where the supernatant was then retained in fresh Eppendorf
microtubes. Protein concentration was estimated by bicinchoninic (BCA) assay (23225; Thermo
Scientific), followed by sample reduction using 10 mM dithiothreitol (DTT) for 5 min at 95 °C.
Following cooling to room temperature, samples were alkylated using 30 mM iodoacetamide
(IA) in dark for 30 min at room temperature without shaking. The alkylation was stopped by
addition of 10 mM DTT, followed by a brief vortex and 10 min room temperature incubation
without shaking. Samples were digested overnight with 1:100 sequencing grade trypsin (V5113;
Promega), followed by quantification of the generated peptide pools using a Nanodrop (Thermo
Scientific). Subsequently, samples were acidified with formic acid to a final concentration of 5%
(v/v) followed by drying by vacuum centrifugation. Peptide desalting was then performed using

ZipTip C18 pipette tips (ZTC18S960; Millipore), with peptides subsequently dried and dissolved
in 3% (v/v) ACN / 0.1% (v/v) FA prior to MS analysis. A Fusion Lumos Tribrid Orbitrap mass
spectrometer (Thermo Scientific) was used to analyse the digested peptides in data
independent acquisition (DIA) mode using the BoxCarDIA method as previously described[20]. 1
µg of dissolved peptide was injected using an Easy-nLC 1200 system (LC140;
ThermoScientific) and a 50 cm Easy-Spray PepMap C18 column (ES903; ThermoScientific).
Liquid chromatography and BoxCarDIA acquisition was performed as previously described[20].

2.8.2 Proteomic Data Analysis

All acquired BoxCarDIA data was analyzed using a library-free approach in Spectronaut v14
(Biognosys AG) under default settings. The *B. oleracea* var. *oleracea* proteome
(Uniprot:https://www.uniprot.org/ containing 58,545 proteins) was used for data searching.
Default search parameters were used for proteome quantification including: a protein, peptide
and PSM FDR of 1%, trypsin digestion with 1 missed cleavage, fixed modification including
carbamidomethylation of cysteine residues and variable modification including methionine
oxidation. Data was Log2 transformed and median normalized with significantly changing
differentially abundant proteins determined and corrected for multiple comparisons (Bonferroni-
corrected p-value <0.05; q-value).

2.9 *Bioinformatics*

Gene Ontology (GO) enrichment analysis was performed using the Database for Annotation,
Visualization and Integrated Discovery (DAVID; v 6.8; https://david.ncifcrf.gov/home.jsp).A)
using a significance threshold of p-value < 0.01. Conversion of *B. oleracea* gene identifiers to
Arabidopsis was performed using Uniprot (https://www.uniprot.org/) followed by Ensembl
Biomart (https://plants.ensembl.org/biomart) for STRING association network analysis and
metabolic pathway analysis. STRING association network analysis was performed using
Cytoscape v3.10.1 (https://cytoscape.org) in combination with the String DB plugin stringApp
and an edge threshold of 0.95. Visualization of GO dotplot was performed using R version 4.3.1
and *ggplot2* package. Metabolic pathway enrichment was performed using the Plant Metabolic
Network (PMN; https://plantcyc.org) with enrichment determined using a Fisher's exact test (p-
value < 0.01). Final figures were assembled using Affinity Designer (v.2.4.1).

3 Results and Discussion
3.1 Light intensity and spectral composition influence kale morphology

We first investigated the effects of light intensity and spectral composition on the morphology of

kale. For our light intensity experiments, plants were grown under three different light intensities:

75, 125 and 175 PPFD (**Supp Figure 1A**). We found a relatively linear relationship between

light intensity with plant fresh weight and surface area in both cultivars at 35 dpi (**Figure 1A-B**).

Consistent with our observations, previous studies have reported a linear relationship between

light intensity and fresh weight in pea microgreens[21], as well as kale, cabbage, arugula and

mustard[22]. Importantly, these studies used PPFD up to 600 $\mu mol \cdot m^{-2} \cdot s^{-1}$, with increases in fresh

weight even at the upper PPFD values, suggesting our highest light intensity of 175 PPFD does

not invoke a high light-induced stress response. As light intensities used in vertical farming are

typically lower, around 200 PPFD[23–25], our study is highly relevant for informing practices of kale

grown in indoor farms, in addition to examining the feasibility of growing kale on a smaller, more

economical scale for home growers or with lower energy usage / footprint. We next examined

the effect of light intensity on diel leaf anthocyanin content, which provides plant coloration to

attract pollinators, but also acts as an antioxidant with benefits to human health[26]. Although

anthocyanins could not be detected in K3, the K9 cultivar, which is known to produce

anthocyanins, typically showing red colouration, did not show a significant difference in

anthocyanin concentration with increasing light intensity (**Figure 1D**). Anthocyanins are typically

produced in response to stressful conditions[27], thus it is possible that our use of a lower light

intensity than that experienced outdoors, combined with the stage of growth, resulted in lower

anthocyanin content.

Our spectral composition experiments consisted of modulating the ratio of red to blue

(R/B) and red to far-red (R/FR) light, while maintaining an otherwise balanced spectra and a

light intensity of 125 PPFD (**Supp Figure 1B**). Previous studies have shown a high R/B ratio

promotes plant growth[28], whereas a low R/B ratio promotes accumulation of beneficial

anthocyanins[29]. In our experiments, K3 exhibited an increase in fresh weight with increasing R/B

with no corresponding change in surface area, whereas K9 showed no change in fresh weight,

but an increase in plant surface area with increasing R/B and R/FR (**Figure 2A-B**), indicating

that red light may be particularly important for K9 growth. Supporting this possibility are

observed increases in dry mass and seedling area specifically in red kale varieties under

elevated red light, whereas blue light had a stronger effect on green kale dry mass[30], suggesting

spectral composition impacts on growth are cultivar specific. This emphasizes the need to

generate cultivar-specific molecular response data, such as that generated here, to uncover

unique breeding opportunities and inform growers choosing genetics for indoor farming. We

additionally observed changes in K9 anthocyanin content with modulated spectral composition

(**Figure 2C**). Consistent with previous reports, an increase in R/B induced a decrease in
anthocyanin content, whereas an increase in R/FR stimulated anthocyanin production at both
timepoints (**Figure 2C**). Therefore, our results show that changing light intensity and spectral
composition impacts kale morphological characteristics and anthocyanin levels.
301
3.2 Modulating light intensity and spectral composition impacts kale photosynthetic performance
We next measured light intensity- and spectra- induced changes in a suite of photosynthetic
parameters alongside relative chlorophyll levels using a MultispeQ device[18] (detailed in **Table 1;**
Supp Figure 2). In particular, ΦPSII, which represents how much light energy is absorbed by
photosystem II[18], and chlorophyll content (SPAD), were both impacted by changes in light
intensity and spectral composition. Increases in light intensity (**Figure 1C**) and R/FR (**Figure**
2C) resulted in increasing SPAD in both cultivars, whereas increasing R/B resulted in a SPAD
decrease (**Figure 2C**). Similarly, ΦPSII increased under increasing light intensity, but in the K9
cultivar only (**Figure 1C**), while an increase in ΦPSII was found in both cultivars with increasing
R/FR (**Figure 2C**). Conversely, an increasing R/B ratio resulted in a decrease in ΦPSII (**Figure**
2C). Our finding of increased ΦPSII with increasing light intensity aligns with previous
observations in lettuce where the highest ΦPSII was found at 200 PPFD across a range of 100
to 800 PPFD, which is comparable to our 175 PPFD treatment[31]. Furthermore, blue light is
found to be more important than red light in modulating photosystem I and II activity and
photosynthetic electron transport capacity in cucumber[32], consistent with our increasing R/B
ratio showing lower SPAD and ΦPSII. Overall, our results indicate that kale photosynthetic
parameters and chlorophyll content can be modulated by changes in light intensity and spectral
composition.
320
3.3 Analysis of total proteome changes with altered light intensity and spectral composition
To contextualize the observed phenotypic changes, we used quantitative proteomics to assess
the protein-level changes in both K3 and K9 over the varying light intensities and spectral
compositions. This analysis quantified a total of 5,493 proteins across our intensity experiments,
and 6,527 proteins across spectra experiments (**Supp Table 1**). We queried this data for
significantly changing proteins by comparing each varied light intensity or spectra to the lowest
treatment condition. Thus,125 PPFD and 175 PPFD were compared to 75 PPFD (**Figure 3A;**
Supp Table 1) and balanced and high spectral compositions compared to low (**Figure 3B-C**
Supp Table 1).

Using these significantly changing proteins, we undertook gene ontology (GO)
enrichment analysis to resolve biological processes underpinning plant response to the
changing light intensity and spectral composition (**Figure 3D**). Here, we find an enrichment in
biological processes related to cell wall and cell cycle, response to oxidative stress and
hormone signaling, along with extensive changes in multiple aspects of metabolism including:
sterol biosynthesis, photosynthesis and pigment metabolism, carbohydrate metabolism,
glycolysis and glucosinolate production. We next refined our GO analysis results by performing
association network analysis using STRING-DB (https://string-db.org/; edge score > 0.7) to
define hubs of associated proteins exhibiting cultivar or time-of-day responses. To do this, we
used the significantly changing proteins induced by a change in light intensity (**Figure 4**), R/B
(**Figure 5A**) and R/FR (**Figure 5B**). As the closely related kale relative, *Arabidopsis thaliana*
(Arabidopsis), has more extensive bioinformatic resources, we first converted the *B. oleracea*
gene identifiers for all quantified proteins in the study to Arabidopsis gene identifiers (AGI) using
a   combination   of   UniProt   (https://www.uniprot.org/)   and   Ensembl   Biomart
(https://plants.ensembl.org/biomart). This identified unique AGIs for 90% of the significantly
changing proteins in our intensity experiments (2395/2662), along with 88.9% (2222/2499) and
89.4% (2520/2819) of proteins in our R/B and R/FR spectra experiments; respectively. Along
with validating enriched biological processes, the association network also indicated changes in
amino acid metabolism, phenylpropanoid metabolism, proteolysis and RNA splicing.

Interestingly, several of the enriched biological processes could be attributed to the plant
shade avoidance response, or relief thereof, including the changes in hormone signaling such
as brassinosteroid (BR) and auxin signaling, cell wall organization / biogenesis and
photosynthesis. Plant shade avoidance response is associated with conditions that mimic
shading, including low light intensity, low blue light and low R/FR ratios[33] and has been linked to
increased BR[34–36] and auxin signaling[37]; both of particular importance for shade-induced
hypocotyl elongation. Within our dataset, we find a down-regulation of BR signaling in K3 with
increasing light intensity at ZT11, and an up-regulation of auxin-activated signaling pathway with
increasing R/B at ZT11. Both observations are consistent with a relief of shade avoidance
responses with our increasing intensity experiments, and an induction upon growth under
decreased blue light.

Further, BR shade avoidance response is related to changes in cell wall structure,
particularly through the action of xyloglucan endotransglucosylase/hydrolases (XTHs) and
expansins (EXPAs), with cell wall modification promoting elongation of the hypocotyl to drive
growth toward light[38]. Our data find a multitude of cell wall proteins changing in abundance in

response to both light intensity and spectral composition, potentially driving shade relief-like responses in both kale cultivars. We specifically note a down-regulation of cell wall organization proteins, primarily XTHs, EXPAs, and pectinesterases (PMEs) with increasing light intensity and R/FR ratio, combined with an up-regulation of XTHs with increasing R/B, which is consistent with changes in the plant perception of shade and hypocotyl elongation. Interestingly, more members of the XTH family were down-regulated in K9 with increasing light intensity and K3 with increasing R/FR, suggesting K9 may be more sensitive to changes in intensity and K3 more sensitive to changes in red light wavelengths, a finding that agrees with our growth and morphology observations.

Beyond its impact on shade avoidance, light intensity and quality critically impact plant photosynthesis and light harvesting, as well as pigment metabolism (e.g. chlorophyll and carotenoids). Intriguingly, our proteomics data reveal a concurrent down-regulation of light harvesting proteins under higher light intensity conditions at ZT23 in K3, with LHCA1 (A0A0D3CTW2; A0A0D3DT59), LHCA2 (A0A0D3DUU9; A0A0D3BWE5), LHCA3 (A0A0D3E520), LHCB2.4 (A0A0D3DBD3), LHCB4.2 (A0A0D3CLT0) and LHCB5 (A0A0D3BQY2) all down-regulated. Alternatively, we only find LHCA2 and LHCB5 were down-regulated in the K9 cultivar, indicating a stronger response in K3. This may result from sufficient, or even excessive light energy under the 175 PPFD light intensity, such that light harvesting potential has been reached, a phenomenon which is well-established in response to light/shade levels[39–41]. Correspondingly, under a higher R/FR ratio we also find the down-regulation of light harvesting proteins, consistent with relief from conditions that promote shade avoidance responses. By contrast, under a balanced ratio we see an up-regulation of light harvesting and photosynthesis-related proteins, particularly in K9 at ZT11, suggesting an optimal ratio of R/B light is best for light harvesting and photosynthetic responses in kale. This is consistent with excessive amounts of red light resulting in an imbalance in PSI and PSII, and thus blue light being required for optimal photosynthetic efficiency[42]. Overall, our data suggest light-dependent differences exist between the two cultivars, with K3 light harvesting proteins showing a stronger response to light intensity changes, and K9 more responsive to changes in spectral composition.

3.4 Analysis of metabolite changes with modified light intensity and spectral composition

With a multitude of metabolic pathways enriched in our GO and STRING-DB analyses, we next used the Arabidopsis orthologs for the significantly changing proteins with increasing intensity, R/B and R/FR to perform metabolic pathway analysis using the Plant Metabolic Network (PMN;

https://plantcyc.org; **Supp Tables 2-4**). Here, enriched metabolic pathways further corroborated
our GO and STRING-DB association network analyses, resolving pathways for photosynthesis
and respiration, amino acid metabolism, pigment metabolism, glycolysis, carbohydrate
metabolism, oxidative stress responses, glucosinolate production and sterol biosynthesis. We
next sought to determine how these changes in metabolic pathway protein abundance translate
into bonified plant metabolic outcomes, using GC-MS to quantify changes in the metabolite
content of each cultivar under each light conditions (**Figure 6; Supp Figure 3; Supp Table 5**),
R/B ratio (**Figure 7; Supp Figure 4-6**). Amino acids (16), fatty acids (3), organic acids (11),
polyamines (2), sterols (2), sugars (5), and sugar alcohols (2) were detected and quantified.

Amino acids - Our proteome analysis indicated changes in amino acid metabolism for both
cultivars in response to changing light intensity and spectral composition (**Supp Figure 6**). Our
metabolite analysis found extensive changes in the glutamate-derived family of amino acids,
which includes glutamic acid, glutamine, GABA, proline and arginine. These amino acids
typically account for the highest amino acid proportions in plants, with glutamic acid and proline
present at the highest quantity in kale[43]. Glutamate metabolism also represents a critical link
between nitrogen and carbon metabolism through the TCA cycle, with several products of
glutamate catabolism, including proline and GABA, associated with plant stress response[44,45].
In both cultivars, we observed a general increase in glutamic acid and proline at both
time-points with increasing light intensity. GABA increased in ZT11 in both cultivars at ZT11, but
at ZT23 there was a decrease in GABA in K3 and an increase in K9. Increased glutamic acid
abundance with increasing light intensity coincided with our proteome-based metabolic pathway
analysis, where we identified down-regulation of proteins involved in glutamate catabolism. This
included GLUTAMATE DEHYDROGENASE1 (GDH1; A0A0D3EEJ0) and GDH2
(A0A0D3AI31), which convert glutamate to glutamine, DELTA-1-PYRROLINE-5-
CARBOXYLATE SYNTHASE B (P5CSB; A0A0D3BXP3), involved in proline biosynthesis, N-
ACETYLGLUTAMATE SYNTHASE (NAGS; A0A0D3A172) and N-ACETYLGLUTAMATE
KINASE (NAGK; A0A0D3BX89), which are steps toward ornithine biosynthesis, and
CARBAMOYL-PHOSPHATE SYNTHASE ALPHA (CARA; A0A0D3CZA9), which drive arginine
biosynthesis. We found this trend in both K3 and K9 and at both time-points, suggesting
increasing the light intensity in our experiments may relieve stress responses associated with
low light. The increase in proline concentration under higher light intensities, however, may
suggest sufficient levels of proline with downregulation of P5CSB to prevent further
accumulation.

We find similar results in our increased R/B ratio proteome data with proteins associated
with proline biosynthesis and glutamate metabolism primarily down-regulated at ZT23 in both
K3 and K9. This manifested as an increase in glutamic acid in both cultivars at both time-points.
Proline levels were not responsive to R/B increase, while glutamine peaked under balanced
light condition at ZT11 and decreased at ZT23. Alternatively, increasing R/FR spectral
composition induced the most changes in metabolite content of glutamate family amino acids.
Interestingly, metabolite data indicated peak glutamic acid, glutamine and GABA levels in both
K3 and K9 under balanced light conditions. This may indicate that balanced light conditions are
associated with the lowest stress and most balanced C/N metabolism during the day.

The aspartate-derived amino acids are also of great importance for human nutrition as
they cannot be synthesized by the human body and must be obtained through diet[46]. This
includes the essential amino acids lysine, threonine, methionine and isoleucine. Of particular
interest is lysine, as it is generally of very low abundance in plants[47]. Within our proteome
metabolic pathway analysis, lysine biosynthesis is enriched among down-regulated proteins as
light intensity increases, particularly in K3 at ZT11, with a down-regulation of ASPARTATE
KINASE (AK2) and of DIHYDRODIPICOLINATE SYNTHASE1 (DHDPS1); both involved in
lysine biosynthesis. However, within our metabolite data, there is a significantly higher amount
of lysine present in K3 under 125 PPFD than either 75 or 175 PPFD, suggesting more complex
regulation mechanisms. Regulatory complexity has previously been observed in attempts to
engineer greater lysine production, where lysine has been shown to allosterically inhibit the
activity of DHDPS1[48], as well as impact the abundance of other aspartate-derived amino acids,
such as threonine[49]. In contrast to increasing light intensity, increasing R/B and R/FR ratios
showed enriched lysine biosynthesis metabolic pathway among up-regulated proteins,
suggesting a greater proportion of red light may promote lysine biosynthesis. Our metabolite
data suggested that lysine levels could be modulated by changing light intensity and spectra,
with a peak at 125 PPFD for both cultivars at ZT23, and a peak under balanced light in K3 at
ZT23. Thus, our results suggest the difficulty in engineering lysine levels in kale may be
possible by carefully modulating a combination of light intensity and spectral composition.

Finally, we noted cultivar-specific differences in glycine content with increasing intensity.
We observed decreased glycine abundance with increasing light intensity in K3, but increased
glycine in K9. Previously, we observed differential accumulation of glycine forming the major
difference between sub-groups containing K3 and K9[17]. Although we did not observe much
change in serine content, both glycine and serine are key to the functioning of primary

metabolism and photorespiration[50], and thus this difference may point to cultivar specific
mediation of primary metabolism response to differing light intensity.

Carbohydrates – Our proteomic analysis revealed changes in carbohydrate metabolic proteins
across each light intensity and spectral composition manipulation treatment. We also find that
this manifested in the differential abundance of several sugars in our metabolite analysis,
including fructose, glucose, mannose-6-phosphate (M6P) and trehalose. In our proteomic data,
we observe a general down-regulation of proteins involved in glycolysis and starch degradation,
as well as trehalose biosynthetic proteins with increasing light intensity. As trehalose has been
associated with shade avoidance[51] and regulation of the circadian clock[52], this is consistent with
reduced stress under low light intensities associated with shade as we increase the light
intensity from 75 to 175 PPFD. We also find that increasing the R/B ratio down-regulates
proteins associated with glycolysis in both cultivars, as well as mannose metabolism in the K9
cultivar only. This was consistent with the lowest M6P levels found in K9 under balanced light
conditions. Finally, increasing R/FR was associated with the down-regulation of glycolytic
proteins in both cultivars, as well as myo-inositol biosynthetic proteins in K9. Conversely, myo-
inositol biosynthesis was up-regulated in K3 with increasing R/FR. In K3, this correlates with
corresponding metabolite levels, as the myo-inositol level was significantly higher in K3 at ZT11
under high R/FR. With myo-inositol being involved in numerous biological processes, including
auxin biosynthesis, membrane biogenesis, light and abiotic stress responses[53], its modulation
likely impacts several of the morphological responses we observe.

Other nutritional compounds - In addition to amino acids and carbohydrates, our metabolite and
proteome analyses identified light intensity- and spectra-induced changes in several other
compounds beneficial for human nutrition, with some cultivar and time-of-day specific
differences. Specifically, we note an increase in ascorbic acid (vitamin C) with increasing light
intensity, particularly in K3 at ZT11. Vitamin C has antioxidant properties[54], and this finding
corresponds with the finding of superoxide metabolic processes being enriched amongst up-
regulated proteins in K3 at ZT11. In addition, the amount of vitamin C in K3 in our R/FR
experiments appeared to decrease under the balanced light condition, but increase with an
increase in R/FR, indicative of the balanced light condition having the lowest oxidative stress.
Conversely, in K9, ascorbic acid increased with increasing R/FR, suggesting different responses
to oxidative stress among cultivars, and potentially resulting in increased anthocyanin within K9.
It is possible that K3 deals with oxidative stress by increasing ascorbic acid and K9 by

increasing anthocyanin content. The K3 ascorbic acid content is consistent with previous

findings of red and far-red light both inducing increases in ascorbic acid content in green pepper

fruit[55].

In addition to ascorbic acid, anthocyanins are a natural source of antioxidants, beneficial

for human health[56,57], with the generation of anthocyanins typically driven by light[58]. In our study,

we were only able to detect anthocyanins within the K9 cultivar, with no significant difference

under the changing light intensity conditions applied, but a significant decrease with increasing

R/B, and a significant increase with increasing R/FR. Within our metabolite analysis, we observe

changes associated with anthocyanin and phenylpropanoid metabolism: including chlorogenic

acid and quininic acid with changing R/B, and sinapinic acid, chlorogenic acid, quininic acid and

3-coumaroyl-D-quinic acid changing in R/FR.

Additionally, within our metabolic pathway analysis, we observed significant pathway

enrichment associated with phenylpropanoid and flavonoid metabolism with increasing R/B and

R/FR composition. Correspondingly, we also see a significant up-regulation of flavonoid

biosynthetic pathways, which is more pronounced in the K9 cultivar and consistent with our

ability to detect anthocyanins in this cultivar. Intriguingly, we find an up-regulation of

phenylpropanoid biosynthesis pathways in K9 with increasing R/B at ZT23, which contrasts our

observed anthocyanin levels. It is possible that at ZT23 the phenylpropanoid pathway is

diverting the flux toward other phenylpropanoid products. For example, we see the coumarin

biosynthesis pathway enriched in our metabolic pathway analysis, coupled with observed up-

regulation of DEFECTIVE IN CUTICULAR RIDGES (DCR; A0A0D3DD85), ENHACED

PSEUDOMONAS SUSCEPTIBILITY1 (EPS1; A0A0D3D884) and COUMARIN SYNTHASE

(COSY; A0A0D3CEC8) in K9 at ZT23, indicating possible increased coumarin biosynthesis[59–61].

Thus, our results suggest the ideal conditions for kale anthocyanin production under lower

intensities involves the use of a high R/FR spectral composition, with higher R/B ratios having

negative impacts by possibly diverting the phenylpropanoid pathway toward other products. This

has substantial application potential for low energy indoor farming.

Although not measured in our metabolomics analysis, our proteomic analysis identified

changes in abundance of proteins involved in production of glucosinolates and carotenoids.

Individual glucosinolates are typically detected using HPLC, not GC-MS[62,63]. Playing a role in

kale flavour and plant defense, glucosinolates can also act as anti-oncogenic compounds for

enhanced human health[64]. Our proteomic analyses indicated changes in glucosinolate

metabolism with increasing light intensity and R/B ratio. In our intensity experiments, the K9

cultivar showed a down-regulation of proteins associated with glucosinolate catabolism as light

intensity increased. Our metabolic pathway analysis also indicated increased biosynthesis of glucosinolates in K3, thus the cultivars might achieve the same elevated levels of glucosinolates at higher light intensities through differing mechanisms. In our bioinformatics assessment of R/B responses, we find glucosinolate catabolism to be decreased and glucosinolate biosynthesis increased in K3. This has previously been observed with levels of the glucosinolates sinigrin, glucobrassicin, and 4-methoxy glucobrassicin in *B. napus* sprouts exposed to increasing red light levels[65], while another study found differential cultivar responses with changing light intensities in Chinese cabbage[66].

In addition to flavonoids and glucosinolates, carotenoids are also antioxidant compounds beneficial to human health[67]. Within our proteomic data, we find K3 has lower carotenoid production with increasing light intensity, combined with decreasing protein abundance of CAROTENE CIS-TRANS-ISOMERASE (CRTISO; A0A0D3DYP6), 15-CIS-PHYTOENE DESATURASE (PDS; A0A0D3BJ64) and PHYTOENE SYNTHASE1 (PSY1; A0A0D3AJG7). In our spectral experiments, it appears that increasing red light in both the R/B and R/FR experiments drives increased carotenoid biosynthesis in the K9 cultivar, along with increasing protein abundance of CRTISO (A0A0D3DYP6), LYCOPENE EPSILON CYCLASE2 (LUT2; A0A0D3B320), PSY1 and PDS in R/B experiments and PDS, CRTISO and CYP97B3 under R/FR conditions. These results suggest that K3 might be more sensitive to light intensity changes and K9 to levels of red light. We thought it interesting to find lower carotenoid biosynthesis with increasing light intensity in K3 as carotenoids tend to increase with higher light intensity[68], however there is evidence in pepper of lower carotenoids with increasing light intensity that is suggested to be due to a higher photo-oxidation rate than that of carotenoid synthesis[69,70].

Conclusions

We present an integrated, systems-level analysis of two phylogenetically related kale cultivars under increasing light intensity, R/B and R/FR spectral ratios with time-of-day precision. Our dataset reveals that light intensity and spectral composition can modulate myriad processes within kale, eliciting changes in growth, and photosynthetic parameters, as well as cell wall and hormones related to shade avoidance (**Figure 8**). Further, we identify changes in nutritional content such as amino acids, carbohydrates, glucosinolates, and anthocyanins with subtle differences between the K3 and K9 cultivar (**Figure 8**). Collectively, these results provide wide-ranging knowledge to indoor farmers of what genetics offer precise nutritional benefits when

566 grown under certain conditions. This offers substantial value to indoor farming operations
567 looking to grow diverse leafy greens of enhanced nutritional value.
568
569

REFERENCES

1. Satheesh, N. & Workneh Fanta, S. Kale: Review on nutritional composition, bio-active compounds,

anti-nutritional factors, health beneficial properties and value-added products. *Cogent Food Agric.*

6, 1811048 (2020).

2. Rachwał, K. *et al.* Red Kale (Brassica oleracea L. ssp. acephala L. var. sabellica) Induces Apoptosis in

Human Colorectal Cancer Cells In Vitro. *Molecules* **28**, 6938 (2023).

3. Bowen-Forbes, C. *et al.* Broccoli, Kale, and Radish Sprouts: Key Phytochemical Constituents and

DPPH Free Radical Scavenging Activity. *Molecules* **28**, 4266 (2023).

4. Cao, D. *et al.* Spermidine enhances chilling tolerance of kale seeds by modulating ROS and

phytohormone metabolism. *PLOS ONE* **18**, e0289563 (2023).

5. Bauer, N. *et al.* Mechanisms of Kale (Brassica oleracea var. acephala) Tolerance to Individual and

Combined Stresses of Drought and Elevated Temperature. *Int. J. Mol. Sci.* **23**, 11494 (2022).

6. Fussy, A. & Papenbrock, J. An Overview of Soil and Soilless Cultivation Techniques—Chances,

Challenges and the Neglected Question of Sustainability. *Plants* **11**, 1153 (2022).

7. https://www.alliedmarketresearch.com, A. M. R. Kale Microgreen Market Size, Share, Price, Trends

|Research Report 2030. *Allied Market Research* https://www.alliedmarketresearch.com/kale-

microgreen-market-A16137.

8. Hati, A. J. & Singh, R. R. Smart Indoor Farms: Leveraging Technological Advancements to Power a

Sustainable Agricultural Revolution. *AgriEngineering* **3**, 728–767 (2021).

9. Nájera, C., Gallegos-Cedillo, V. M., Ros, M. & Pascual, J. A. LED Lighting in Vertical Farming Systems

Enhances Bioactive Compounds and Productivity of Vegetables Crops. *Biol. Life Sci. Forum* **16**, 24

(2022).

10. McCree, K. J. The action spectrum, absorptance and quantum yield of photosynthesis in crop plants.

Agric. Meteorol. **9**, 191–216 (1971).

11. Krahmer, J., Ganpudi, A., Abbas, A., Romanowski, A. & Halliday, K. J. Phytochrome, Carbon Sensing,

Metabolism, and Plant Growth Plasticity. *Plant Physiol.* **176**, 1039–1048 (2018).

12. Chen, J. *et al.* Effect of Photoperiod on Chinese Kale (Brassica alboglabra) Sprouts Under White or

Combined Red and Blue Light. *Front. Plant Sci.* **11**, (2021).

13. Kong, Y., Schiestel, K. & Zheng, Y. Pure blue light effects on growth and morphology are slightly

changed by adding low-level UVA or far-red light: A comparison with red light in four microgreen

species. *Environ. Exp. Bot.* **157**, 58–68 (2019).

14. Deng, M. *et al.* Influence of pre-harvest red light irradiation on main phytochemicals and antioxidant

activity of Chinese kale sprouts. *Food Chem.* **222**, 1–5 (2017).

15. Zhen, S. & van Iersel, M. W. Far-red light is needed for efficient photochemistry and photosynthesis.

J. Plant Physiol. **209**, 115–122 (2017).

16. Furuya, M. & Schäfer, E. Photoperception and signalling of induction reactions by different

phytochromes. *Trends Plant Sci.* **1**, 301–307 (1996).

17. Scandola, S., Mehta, D., Castillo, B., Boyce, N. & Uhrig, R. G. Systems-level proteomics and

metabolomics reveals the diel molecular landscape of diverse kale cultivars. *Front. Plant Sci.* **14**,

(2023).

18. Kuhlgert, S. *et al.* MultispeQ Beta: a tool for large-scale plant phenotyping connected to the open

PhotosynQ network. *R. Soc. Open Sci.* **3**, 160592 (2016).

19. Gehan, M. A. *et al.* PlantCV v2: Image analysis software for high-throughput plant phenotyping.

PeerJ **5**, e4088 (2017).

20. Mehta, D., Scandola, S. & Uhrig, R. G. BoxCar and Library-Free Data-Independent Acquisition

Substantially Improve the Depth, Range, and Completeness of Label-Free Quantitative Proteomics.

Anal. Chem. **94**, 793–802 (2022).

21. Balázs, L. *et al.* Quantifying the Effect of Light Intensity Uniformity on the Crop Yield by Pea

Microgreens Growth Experiments. *Horticulturae* **9**, 1187 (2023).

22. Jones-Baumgardt, C., Llewellyn, D., Ying, Q. & Zheng, Y. Intensity of Sole-source Light-emitting

Diodes Affects Growth, Yield, and Quality of Brassicaceae Microgreens. (2019)

doi:10.21273/HORTSCI13788-18.

23. Dou, H. *et al.* Supplementary Far-Red Light for Photosynthetic Active Radiation Differentially

Influences the Photochemical Efficiency and Biomass Accumulation in Greenhouse-Grown Lettuce.

Plants **13**, 2169 (2024).

24. Gudžinskaitė, I., Laužikė, K., Pukalskas, A. & Samuolienė, G. The Effect of Light Intensity during

Cultivation and Postharvest Storage on Mustard and Kale Microgreen Quality. *Antioxidants* **13**, 1075

(2024).

25. Boucher, L., Nguyen, T.-T.-A., Brégard, A., Pepin, S. & Dorais, M. Optimizing Light Use Efficiency and

Quality of Indoor Organically Grown Leafy Greens by Using Different Lighting Strategies. *Agronomy*

13, 2582 (2023).

26. Zhang, Y., Butelli, E. & Martin, C. Engineering anthocyanin biosynthesis in plants. *Curr. Opin. Plant*

Biol. **19**, 81–90 (2014).

27. Kaur, S. *et al.* Protective and defensive role of anthocyanins under plant abiotic and biotic stresses:

An emerging application in sustainable agriculture. *J. Biotechnol.* **361**, 12–29 (2023).

28. Dou, H., Niu, G. & Gu, M. Photosynthesis, Morphology, Yield, and Phytochemical Accumulation in

Basil Plants Influenced by Substituting Green Light for Partial Red and/or Blue Light. (2019)

doi:10.21273/HORTSCI14282-19.

29. Anum, H., Cheng, R. & Tong, Y. Improving plant growth, anthocyanin production and oxidative

status of red lettuce (*Lactuca sativa* cv. Lolla Rossa) by optimizing red to blue light ratio with a

constant green light fraction in a plant factory. *Sci. Hortic.* **338**, 113832 (2024).

30. Frąszczak, B. *et al.* Morphological and Photosynthetic Parameters of Green and Red Kale

Microgreens Cultivated under Different Light Spectra. *Plants* **12**, 3800 (2023).

31. Fu, W., Li, P. & Wu, Y. Effects of different light intensities on chlorophyll fluorescence characteristics

and yield in lettuce. *Sci. Hortic.* **135**, 45–51 (2012).

32. Miao, Y., Wang, X., Gao, L., Chen, Q. & Qu, M. Blue light is more essential than red light for

maintaining the activities of photosystem II and I and photosynthetic electron transport capacity in

cucumber leaves. *J. Integr. Agric.* **15**, 87–100 (2016).

33. Pierik, R. & de Wit, M. Shade avoidance: phytochrome signalling and other aboveground neighbour

detection cues. *J. Exp. Bot.* **65**, 2815–2824 (2014).

34. Lv, M. & Li, J. Molecular Mechanisms of Brassinosteroid-Mediated Responses to Changing

Environments in Arabidopsis. *Int. J. Mol. Sci.* **21**, 2737 (2020).

35. Keuskamp, D. H. *et al.* Blue-light-mediated shade avoidance requires combined auxin and

brassinosteroid action in Arabidopsis seedlings. *Plant J.* **67**, 208–217 (2011).

36. Klahre, U. *et al.* The Arabidopsis DIMINUTO/DWARF1 Gene Encodes a Protein Involved in Steroid

Synthesis. *Plant Cell* **10**, 1677–1690 (1998).

37. Ma, L. & Li, G. Auxin-Dependent Cell Elongation During the Shade Avoidance Response. *Front. Plant*

Sci. **10**, (2019).

38. Sasidharan, R. *et al.* Light Quality-Mediated Petiole Elongation in Arabidopsis during Shade

Avoidance Involves Cell Wall Modification by Xyloglucan Endotransglucosylase/Hydrolases. *Plant*

Physiol. **154**, 978–990 (2010).

39. Teramoto, H., Nakamori, A., Minagawa, J. & Ono, T. Light-Intensity-Dependent Expression of Lhc

Gene Family Encoding Light-Harvesting Chlorophyll-a/b Proteins of Photosystem II in

Chlamydomonas reinhardtii. *Plant Physiol.* **130**, 325–333 (2002).

40. Anderson, J. M., Chow, W. S. & Goodchild, D. J. Thylakoid Membrane Organisation in Sun/Shade

Acclimation. *Funct. Plant Biol.* **15**, 11–26 (1988).

41. Anderson, J. M., Chow, W. S. & Park, Y.-I. The grand design of photosynthesis: Acclimation of the

photosynthetic apparatus to environmental cues. *Photosynth. Res.* **46**, 129–139 (1995).

42. Jin, D. *et al.* Effect of Red and Blue Light on Cucumber Seedlings Grown in a Plant Factory.

Horticulturae **9**, 124 (2023).

43. Lisiewska, Z., Kmiecik, W. & Korus, A. The amino acid composition of kale (*Brassica oleracea* L. var.

acephala), fresh and after culinary and technological processing. *Food Chem.* **108**, 642–648 (2008).

44. Okumoto, S., Funck, D., Trovato, M. & Forlani, G. Editorial: Amino Acids of the Glutamate Family:

Functions beyond Primary Metabolism. *Front. Plant Sci.* **7**, (2016).

45. Kavi Kishor, P. B. & Sreenivasulu, N. Is proline accumulation per se correlated with stress tolerance

or is proline homeostasis a more critical issue? *Plant Cell Environ.* **37**, 300–311 (2014).

46. Galili, G. & Amir, R. Fortifying plants with the essential amino acids lysine and methionine to

improve nutritional quality. *Plant Biotechnol. J.* **11**, 211–222 (2013).

47. Yang, Q., Zhao, D. & Liu, Q. Connections Between Amino Acid Metabolisms in Plants: Lysine as an

Example. *Front. Plant Sci.* **11**, (2020).

48. Hall, C. J. *et al.* Differential lysine-mediated allosteric regulation of plant dihydrodipicolinate

synthase isoforms. *FEBS J.* **288**, 4973–4986 (2021).

49. Van Bochaute, P., Novoa, A., Ballet, S., Rognes, S. E. & Angenon, G. Regulatory mechanisms after

short- and long-term perturbed lysine biosynthesis in the aspartate pathway: the need for isogenes

in Arabidopsis thaliana. *Physiol. Plant.* **149**, 449–460 (2013).

50. Rosa-Téllez, S. *et al.* The serine–glycine–one-carbon metabolic network orchestrates changes in

nitrogen and sulfur metabolism and shapes plant development. *Plant Cell* **36**, 404–426 (2024).

51. Hwang, G. *et al.* Trehalose-6-phosphate signaling regulates thermoresponsive hypocotyl growth in

Arabidopsis thaliana. *EMBO Rep.* **20**, e47828 (2019).

52. Avidan, O. *et al.* Direct and indirect responses of the Arabidopsis transcriptome to an induced

increase in trehalose 6-phosphate. *Plant Physiol.* **196**, 409–431 (2024).

53. Sharma, N., Chaudhary, C. & Khurana, P. Role of myo-inositol during skotomorphogenesis in

Arabidopsis. *Sci. Rep.* **10**, 17329 (2020).

54. Gęgotek, A. & Skrzydlewska, E. Antioxidative and Anti-Inflammatory Activity of Ascorbic Acid.

Antioxidants **11**, 1993 (2022).

55. Pashkovskiy, P. *et al.* Post-Harvest Red- and Far-Red-Light Irradiation and Low Temperature Induce

the Accumulation of Carotenoids, Capsaicinoids, and Ascorbic Acid in Capsicum annuum L. Green

Pepper Fruit. *Foods* **12**, 1715 (2023).

56. Tena, N., Martín, J. & Asuero, A. G. State of the Art of Anthocyanins: Antioxidant Activity, Sources,

Bioavailability, and Therapeutic Effect in Human Health. *Antioxidants* **9**, 451 (2020).

57. Pojer, E., Mattivi, F., Johnson, D. & Stockley, C. S. The Case for Anthocyanin Consumption to

Promote Human Health: A Review. *Compr. Rev. Food Sci. Food Saf.* **12**, 483–508 (2013).

58. Liu, C., Yao, X., Li, G., Huang, L. & Xie, Z. Transcriptomic profiling of purple broccoli reveals light-

induced anthocyanin biosynthetic signaling and structural genes. *PeerJ* **8**, e8870 (2020).

59. Kim, C. Y. *et al.* Emergence of a proton exchange-based isomerization and lactonization mechanism

in the plant coumarin synthase COSY. *Nat. Commun.* **14**, 597 (2023).

60. Vanholme, R. *et al.* COSY catalyses trans–cis isomerization and lactonization in the biosynthesis of

coumarins. *Nat. Plants* **5**, 1066–1075 (2019).

61. Rani, S. H., Krishna, T. H. A., Saha, S., Negi, A. S. & Rajasekharan, R. Defective in Cuticular Ridges

(*DCR*) of *Arabidopsis thaliana*, a Gene Associated with Surface Cutin Formation, Encodes a Soluble

Diacylglycerol Acyltransferase*. *J. Biol. Chem.* **285**, 38337–38347 (2010).

62. Grosser, K. & van Dam, N. M. A Straightforward Method for Glucosinolate Extraction and Analysis

with High-pressure Liquid Chromatography (HPLC). *J. Vis. Exp. JoVE* 55425 (2017)

doi:10.3791/55425.

63. Theunis, M. *et al.* Optimization and validation of analytical RP-HPLC methods for the quantification

of glucosinolates and isothiocyanates in *Nasturtium officinale* R. Br and *Brassica oleracea*. *LWT* **165**,

113668 (2022).

64. Ltd, wflpublisher. Comparative study on functional components, antioxidant activity and color

parameters of selected colored leafy vegetables as affected by photoperiods. *wflpublisher.com*

https://www.wflpublisher.com/Abstract/2605.

65. Park, C. H. *et al.* Effects of Light-Emitting Diodes on the Accumulation of Glucosinolates and Phenolic

Compounds in Sprouting Canola (Brassica napus L.). *Foods* **8**, 76 (2019).

66. Zhou, B. *et al.* Effects of light intensity on the biosynthesis of glucosinolate in Chinese cabbage

plantlets. *Sci. Hortic.* **316**, 112036 (2023).

67. Eggersdorfer, M. & Wyss, A. Carotenoids in human nutrition and health. *Arch. Biochem. Biophys.*

652, 18–26 (2018).

68. Flores, M., Hernández-Adasme, C., Guevara, M. J. & Escalona, V. H. Effect of different light

intensities on agronomic characteristics and antioxidant compounds of Brassicaceae microgreens in

a vertical farm system. *Front. Sustain. Food Syst.* **8**, (2024).

69. Simkin, A. J., Zhu, C., Kuntz, M. & Sandmann, G. Light-dark regulation of carotenoid biosynthesis in

pepper (*Capsicum annuum*) leaves. *J. Plant Physiol.* **160**, 439–443 (2003).

70. Pizarro, L. & Stange, C. Light-dependent regulation of carotenoid biosynthesis in plants. *Cienc. E*

Investig. Agrar. **36**, 143–162 (2009).

Figure 1. Effect of light intensities on kale growth and development.
A. Pictures of K3 and K9 plants at 35 dpi grown under different light intensities (75, 125, and 175 PPFD).
B. K3 and K9 weight at 35 dpi (n=6), leaf area at 22 dpi (n=15).
C. Relative chlorophyll content (SPAD) and photosynthetic parameters (Phi2) measured with MultispeQ at 35 dpi (n=10).
D. K9 anthocyanin content at 35 dpi (n=4), at different light intensities (75, 125, and 175 PPFD).
Letters shows significant differences using a one-way ANOVA and Tukey's post-hoc test (adjusted p-value <0.05).

Figure 2. Effect of light spectra on kale growth and development.
A. Pictures of K3 and K9 plants at 35 dpi grown under different light spectra (low, balanced, high R/B and R/FR).
B. K3 and K9 weight at 35 dpi (n=6), leaf area at 22 dpi (n=15), relative chlorophyll content (SPAD) and photosynthetic parameters (Phi2) measured with MultispeQ at 35 dpi (n=10) at different light spectra (low, balanced, high R/B and R/FR).
C. K9 anthocyanin content at 35 dpi (n=4), at different light spectra (low, balanced, high R/B and R/FR).
Letters shows significant differences using a one-way ANOVA and Tukey's post-hoc test (adjusted p-value <0.05).

Figure 3. Analysis of significantly changing proteins with changing light intensity and spectral composition.

A-C. Number of proteins with a significant change in abundance with increase in light intensity from 75 PPFD (A), from low R/B (B) and from low R/FR (C).

D. Dot plot representation of enriched GO terms for biological process of significantly changing proteins (q-value < 0.05; Log2FC > 0.58 or < -0.58) that are up- or down-regulated with increasing light intensity (top panel), increasing R/B ratio (middle panel), or increasing R/FR ratio (bottom panel) at ZT23 (left) or ZT11 (right). Dot size indicates the gene ratio (number of proteins in the changing in the condition/total quantified proteins in the category in the study). Dot colour represents the Log10 (p-value). Only GO terms with p-value <0.01 and a total count < 500 are included.

Figure 4. Association network analysis of significantly changing proteins with increasing light intensity
An association network analysis was performed using STRING-DB to depict significantly changing proteins in K3 and K9 kale under increasing light intensity at ZT23 and ZT11. Edge thickness indicates strength of connection between the nodes. Minimum edge threshold was set to 0.95. Protein nodes are labelled either by primary gene name or Arabidopsis gene identifier (AGI). Outer circle surrounding each node represents the standardized relative Log2FC of the indicated significantly changing protein with an increase from 75 PPFD to either 125 PPFD or 175 PPFD in K3 or K9 as indicated by the legend. The scale of blue to yellow indicates the relative decrease or increase in abundance, respectively from 75 PPFD to either 125 PPFD or 175 PPFD. Node groupings are indicated by a grey circle representing proteins involved in the same biological process.

Figure 5. Association network analysis of significantly changing proteins with changing light quality
An association network analysis was performed using STRING-DB to depict significantly changing proteins in K3 and K9 kale under increasing R/B spectral ratio (A) or R/FR ratio (B) at ZT23 and ZT11. Edge thickness indicates the strength of connection between the nodes. Minimum edge threshold was set to 0.95. Protein nodes are labelled either by primary gene name or Arabidopsis gene identifier (AGI). Outer circle surrounding each node represents the standardized relative Log2FC of the indicated significantly changing protein with an increase from low R/B (A) or low R/FR (B) to balanced (R = B) or high R/B (R > B) (A) or balanced (R = FR) or high R/FR (R > FR) (B) in K3 or K9 at ZT23 or ZT11 as indicated by the legend. The scale of blue to yellow indicates the relative decrease or increase in abundance, respectively. Node groupings are indicated by a grey circle representing proteins involved in the same biological process.

Figure 6: GC-MS analysis of kale cultivars K3 and K9 under different light intensities (75 PPFD, 125 PPFD and 175 PPFD) and at different time of the day (ZT11 and ZT23).
Heatmap of relative cultivars metabolite changes at ZT11 and ZT23 for 125 PPFD/75 PPFD and for 175 PPFD/ 75PPFD. Scale represents log2FC, n=4

Figure 7: GC-MS analysis of kale cultivars K3 and K9 under different spectral compositions
A) Heatmap of relative cultivars metabolite changes at ZT11 and ZT23 for changing R/B ratio Balanced/Low and High/Low. Scale represents log2FC, n=4
B) Heatmap of relative cultivars metabolite changes at ZT11 and ZT23 for changing R/FR ratio Balanced/Low and High/Low. Scale represents log2FC, n=4

Main proteomic and metabolic pathways influenced by light in Kale cultivars

Figure 8. Summary of proteomic and metabolic changes with varying light intensity and spectra
Summary of main proteomic and metabolic pathways influenced by light in kale cultivars K3 and K9 at ZT11 and ZT23 with increasing PPFD, R/B and R/FR.

Table. Effect of light intensity and spectra on photosynthetic parameters measured with the MultispeQ (PhotosynQ Inc.). Relative chlorophyll content (SPAD) and photosynthetic parameters (Phi2, PhiNO, PhiNPQ, LEF, gH^+, ECSt, vH^+) were measured with the MultispeQ at 35 DPI. n=10.

		K3		K9	
	Intensity	**Mean**	**SEM**	**Mean**	**SEM**
SPAD	75 PPFD	32.97	3.73	30.24	2.61
	125 PPFD	37.77	6.14	35.22	4.47
	175 PPFD	41.90	3.95	39.93	2.90
Phi2	75 PPFD	0.56	0.02	0.58	0.02
	125 PPFD	0.61	0.02	0.60	0.02
	175 PPFD	0.59	0.02	0.60	0.01
PhiNO	75 PPFD	0.32	0.02	0.30	0.02
	125 PPFD	0.31	0.03	0.31	0.02
	175 PPFD	0.31	0.01	0.31	0.02
PhiNPQ	75 PPFD	0.109	0.015	0.111	0.008
	125 PPFD	0.095	0.022	0.096	0.009
	175 PPFD	0.093	0.019	0.096	0.014
LEF	75 PPFD	23.91	2.15	20.37	2.18
	125 PPFD	39.06	4.02	36.72	5.72
	175 PPFD	53.40	9.03	49.33	7.33
gH+	75 PPFD	119.61	30.59	115.37	19.10
	125 PPFD	175.33	50.61	110.91	35.23
	175 PPFD	153.41	56.18	117.67	51.88
ECSt	75 PPFD	0.008	0.002	0.009	0.002
	125 PPFD	0.006	0.002	0.008	0.003
	175 PPFD	0.007	0.002	0.008	0.006
vH+	75 PPFD	0.058	0.013	0.051	0.017
	125 PPFD	0.071	0.018	0.059	0.026
	175 PPFD	0.068	0.023	0.045	0.025

		K3		K9	
	R/B	**Mean**	**SEM**	**Mean**	**SEM**
SPAD	Low	44.70	2.84	47.06	3.23
	Balanced	43.80	2.29	45.10	2.09
	High	44.14	2.11	44.35	2.47
Phi2	Low	0.66	0.01	0.65	0.01
	Balanced	0.58	0.02	0.58	0.01
	High	0.57	0.01	0.57	0.01
PhiNO	Low	0.26	0.01	0.26	0.01
	Balanced	0.31	0.02	0.33	0.01
	High	0.33	0.02	0.33	0.02
PhiNPQ	Low	0.086	0.020	0.093	0.009
	Balanced	0.089	0.013	0.101	0.006
	High	0.112	0.019	0.104	0.017
LEF	Low	28.12	1.80	26.60	2.78
	Balanced	36.03	4.55	34.02	3.94
	High	48.07	6.07	44.81	6.75
gH+	Low	147.92	42.54	178.52	50.97
	Balanced	170.01	13.77	144.49	42.57
	High	162.13	42.77	151.98	47.40
ECSt	Low	0.007	0.002	0.006	0.002
	Balanced	0.006	0.001	0.007	0.002
	High	0.006	0.002	0.007	0.002
vH+	Low	0.056	0.017	0.062	0.022
	Balanced	0.083	0.016	0.069	0.025
	High	0.081	0.012	0.082	0.032

		K3		K9	
	R/FR	**Mean**	**SEM**	**Mean**	**SEM**
SPAD	Low	41.98	1.59	44.08	2.14
	Balanced	44.57	1.66	46.30	2.70
	High	49.09	2.11	48.98	2.71
Phi2	Low	0.54	0.01	0.54	0.01
	Balanced	0.62	0.01	0.60	0.02
	High	0.66	0.01	0.65	0.01
PhiNO	Low	0.35	0.02	0.35	0.01
	Balanced	0.30	0.02	0.30	0.02
	High	0.24	0.01	0.25	0.00
PhiNPQ	Low	0.110	0.016	0.099	0.014
	Balanced	0.086	0.020	0.096	0.010
	High	0.097	0.013	0.107	0.014
LEF	Low	36.91	1.13	31.47	9.58
	Balanced	42.16	4.99	39.07	2.77
	High	39.98	1.17	37.25	3.36
gH+	Low	152.08	27.80	174.73	56.02
	Balanced	173.91	60.05	145.10	52.68
	High	150.88	107.37	137.71	48.16
ECSt	Low	0.007	0.001	0.006	0.002
	Balanced	0.006	0.001	0.007	0.002
	High	0.007	0.003	0.007	0.002
vH+	Low	0.068	0.017	0.066	0.047
	Balanced	0.076	0.023	0.075	0.018
	High	0.071	0.015	0.072	0.030