Shaping Kale Morphology and Physiology Using Different LED Light Recipes

Sabine Scandola[1,2,*], Lauren E. Grubb[1,*], Brigo Castillo[1], Lexyn Iliscupidez[1], Curtis Kennedy[1,3], Nicholas Boyce[1], and R. Glen Uhrig[1, 4,#]

[1] Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada

[2] Lethbridge Research and Development Centre, Agriculture and Agri-Food Canada, Lethbridge, Canada

[3] Department of Computer Science, University of Alberta, Edmonton, Alberta, Canada

[4] Department of Biochemistry, University of Alberta, Edmonton, Alberta, Canada

[*] These authors contributed equally (S.S. and L.G.)

[#] Senior and corresponding author:

Dr. R. Glen Uhrig
Department of Biological Sciences,
University of Alberta,
Edmonton, Alberta, Canada, T6G 2E9
Email: ruhrig@ualberta.ca
Phone: +1 780 492 3088
Fax : +1 780 492-9234ruhrig@ualberta.ca

Running Title: Light condition effects on kale growth and development

Key Words: Kale, Anthocyanins, Light emitting diodes (LEDs), Horticulture, PhotosynQ, Quantitative Proteomics, Metabolomics

Abstract

In the evolving landscape of horticultural science, light emitting diodes (LEDs) present an innovative opportunity for manipulating plant growth and development. Light serves as a fundamental energy source and an environmental cue for plant life, providing us an ability to control essential plant traits through the precise manipulation of light intensity and quality. In this study, we assess the effects of light intensity and spectral composition on the growth and physiology of a horticulturally significant model plant: Kale (*Brassica oleracea*). Selected for its phenotypic plasticity and nutritional composition, kale is a crop well-suited for indoor cultivation using LEDs. Here, we employ a combination of advanced phenotyping, computer vision, gas chromatography-mass spectrometry (GC-MS) metabolomics, and liquid chromatography-mass spectrometry (LC-MS)-based quantitative proteomics to characterize the molecular changes that underpin light-dictated differences in the growth and metabolism of two different kale cultivars under different light intensity and spectral composition scenarios. Our results not only offer a key resource to the plant community, but also demonstrate the translational potential of light manipulation in tailoring kale growth and nutritional content for enhanced crop productivity and/or nutritional content, while simultaneously offering a more cost-effective solution for contemporary agricultural challenges.

1 Introduction

Kale (*Brassica oleracea*), also known as leaf cabbage, belongs to the large *Brassicaceae* family that includes cauliflower, broccoli and collard greens. Established as a superfood, this leafy green is rich in fiber and essential vitamins, as well as antioxidants such as flavonoids, each of which have been shown to positively affect cardiovascular and gastrointestinal health[1–3]. To date, the health benefits of kale have been heavily studied, with researchers continuing to identify new anti-cancer (e.g. sinigrin, spirostanol)[4] and neuroprotective (e.g. sulforaphane)[5] compounds, which neutralize reactive oxygen species. Further, kale demonstrates excellent temperature resilience as a result of its waxy coating and high production of polyamines[6], rendering it a valuable agricultural crop in the context of climate change[7].

Kale is also suited for indoor / vertical farming. As a leafy green, it is a favored choice for horticultural growers as it has a quick harvest cycle with minimal space and resource requirements[8], appealing to both growers as well as consumers looking for fresh, locally-sourced produce[9]. Indoor / vertical farms have strongly co-evolved with new technologies as the promise of providing fresh and locally grown food largely relies on reducing production costs through technological advancements[10,11]. In this context, LED technology has revolutionized indoor farming, providing a wide range of energy-efficient and versatile lighting options. For example, unlike traditional light sources (e.g. incandescent bulbs, fluorescent tubes, metal halide or high pressure sodium bulbs), LED lighting systems offer an ability to fine-tune the spectral composition and intensity of the applied light[12,13]. Light is critical for photosynthesis, which drives plant metabolism and growth[14], while plant photoreceptors, such as phytochromes and cryptochromes, respond to light quality, impacting processes such as plant growth (e.g. biomass) and development (e.g. flowering)[15]. However, despite our understanding of kale at a phenotypic level, we lack knowledge of how light impacts kale at the molecular level, limiting our ability to enhance genetics through targeted breeding or biotechnology approaches.

To date, elementary changes in spectral composition have been shown to affect kale phenotypic and metabolic profiles[16]. Blue light enhances growth and nutritional qualities[17], while red light affects kale pre- and post-harvest by altering metabolites and chlorophyll content[18]. Similar to red light, far-red light enhances photosynthetic rates through the Emerson effect[19]. Beyond specific light types, light wavelength ratios also play an essential role in how phytochromes perceive and transduce light signals[20], offering opportunities to build precise light recipes that maximize traits of interest.

To better exploit currently available kale genetics, we endeavored to precisely resolve the
impacts of light intensity and spectral composition on kale growth, development and nutritional
content by contrasting the diel molecular changes in two related kale cultivars: Dwarf curled
scotch (K3) and Red Scarlet (K9) (**Figure 1**). While there are many kale cultivars, each with
unique morphologies, growth parameters and nutritional characteristics[21], our previous study
examining nine commercially available kale cultivars found that K3 and K9, both *Brassica*
oleracea var. sabellica, showed the most diversity in their phenotypes and molecular
characteristics[21], allowing us here to capture an in-depth understanding of how light properties
impact kale growth and development in order to establish a compendium of molecular changes
that can be utilized by the broader community. We specifically took a diel experimental
approach to analyzing the effects of light intensity and spectral composition by harvesting
tissues at end-of-day (ED, ZT11 (zeitgeber)) and end-of-night (EN, ZT23) time-points as both
the proteome and metabolome are known to undergo substantial diel changes[21]. Understanding
how external factors such as time-of-day intersect with differences in light intensity and quality
to impact kale physiology, is essential for determining optimal horticultural growing conditions
and harvesting times in order to maximize value, as well as inform future efforts in breeding
more productive and nutritious kale varieties.

2 Materials and Methods

2.1 Plant material and cultivation

Dwarf Curled Scotch (K3) and Red Scarlet (K9) kale seeds (*B. oleracea var. sabellica*) were
purchased from West Coast Seeds (https://www.westcoastseeds.com/). Seeds were surface
sterilized with incubations in 70% (v/v) ethanol and 30% (v/v) bleach solutions at room
temperature beginning with a 4-minute incubation in 3 mL of 70% (v/v) ethanol. The seeds were
then rinsed with 3 mL of sterile milli-Q water and 3 mL of 30% (v/v) bleach was added onto each
tube for a 7-minute incubation with periodical shaking every 2 minutes. Finally, the bleach
solution was removed, and the seeds were rinsed three times with 3 mL of sterile Milli-Q water.
Following sterilization, the seeds from each cultivar were plated onto 0.5 x Murashige and
Skoog (MS) media plates containing 1% (w/v) sucrose. The seeds were then stratified at 4 °C in
total darkness for 96 hours. Following stratification, the kale seedlings were entrained to their
respective light treatment for 7 days. To prepare the soil for seedling transplant, ~25 kg of soil
(Sun Gro Sunshine Mix #1) was mixed with 8 L of H_2O and then transferred onto 4" square

polypropylene pots. Transplantation occurred once most plants had reached the cotyledon
stage at 11 days post-imbibition (dpi).

2.2 Light Conditions

The programmable LED (Perihelion Model) lights used for this experiment were provided by
G2V Optics Inc. and were installed inside ventilated growth chambers lined with reflective mylar.
The kale plants were grown under a 12-hour light and 12-hour dark regiment. Three different
light intensities were tested: 75, 125, 175 PPFD ($\mu mol.m^{-2}.s^{-1}$) (**Supp Figure 1A**). Spectral light
conditions were assigned in spectral ratios of R/B and R/FR of low R/B or R/FR, balanced R/B
or R/FR and high R/B or R/FR (**Supp Figure 1B**).

2.3 Measurement of Kale Photosynthetic Performance via MultisynQ

Photosynthetic performance was measured following the Photosynthesis RIDES 2.0 protocol
using a MultispeQ v1.0 (PhotosynQ LLC, USA)[22]. At mid-day (zeitgeber; ZT6) 34 dpi, the
photosynthetic performance of the kale plants was measured by clamping the second true leaf
between the MultisynQ sensors ensuring full coverage. The leaf remained in the chamber until
the data collection process was completed (~10 s). A total of 10 randomly selected plants were
analyzed for each cultivar and light condition.

2.4 Phenotyping measurements

Plants were grown in separate chambers under different light intensity and spectral conditions
(**Supp Figure 1**). Each chamber was equipped with single-board Raspberry Pi 3 B+
computers and ArduCam Noir Cameras (OV5647 1080p) with a motorized IR-CUT filter and two
infrared LEDs. Pictures of plants were taken every 5 minutes. Plant surface area analysis
across light conditions was performed by averaging the area of each plant measured during a 1-
hour period (between ZT6 and ZT7) at 22 dpi. Plant surface areas were extracted using Python
scripts from the open-source software package PlantCV (https://plantcv.readthedocs.io/),
following the methodology described by Gehan *et al.*, 2017[23].

2.5 Harvesting and Sample Processing

At 35 dpi, kale leaves were harvested at two time points: end-of-night (ZT23) and end-of-day
(ZT11). This was done to capture the expansive metabolic changes occurring during the
transitions between night-to-day and day-to-night, respectively. Leaves from three kale plants of
the same cultivar were pooled into a labelled 50 mL conical tube and promptly frozen in liquid
nitrogen to preserve the proteins and metabolites present. In total, four tubes were collected

165 from each cultivar per chamber during harvest. Two metal beads were then introduced into each
166 conical tube and the samples were ground by subjecting them to 3x – 30 s/1200 rpm using a
167 Geno/Grinder® - Automated Tissue Homogenizer and Cell Lyser. 100 mg (±1 mg) of each finely-
168 ground sample was then portioned into each of two separate 2 mL Eppendorf Safe-Lock tubes
169 for further processing.

170 *2.6 Anthocyanin Extraction and Quantification:*

171 Anthocyanin analysis was performed as previously described by Neff and Chory (1998), with the
172 following modifications. Specifically, 500 µL of 100% (v/v) methanol-1% (v/v) hydrochloric acid
173 was added to each tube containing the finely ground samples and incubated at 4 °C for 24
174 hours in darkness. From there, 200 µL of H_2O and 500 µL of chloroform were added and the
175 tubes were centrifuged at 18,000 x g for 5 minutes at room temperature. 400 µL of the
176 supernatant was then transferred over to new tubes and adjusted to a total volume of 800 µL
177 with 60% (v/v) methanol-1% (v/v) hydrochloric acid. A total of 200 µL from each volume-
178 corrected sample was transferred into a well in a 96-well plate and absorbance at 530 nm and
179 657 nm was measured with a Tecan Spark® plate reader using 60% (v/v) methanol -1% (v/v)
180 hydrochloric acid as the background. The relative anthocyanin content was then derived
181 mathematically using: $Antho = (Abs_{530} - Abs_{657}) \times 1000 \times powder\ weight\ (mg)^{-1}$.

182 *2.7 Metabolite analysis via Gas Chromatography Mass Spectrometry (GCMS)*

183 *2.7.1 Metabolite extraction:* A total of 700 µL of ice-cold 100% (v/v) methanol was added to the
184 tubes containing 100 mg of finely ground leaf sample and promptly vortexed for ~20 seconds.
185 Using a thermomixer, these were then incubated for 15 minutes at 70 °C with shaking at 1200
186 rpm. From there, the tubes were centrifuged at 18,000 rpm for 15 minutes and the supernatants
187 were transferred into new tubes. The pellet was then re-extracted using 700 µL of 50% (v/v)
188 methanol containing Ribitol at 25 µL per sample at 0.4 mg/mL in water and vortexed for ~20
189 seconds until re-suspended. These tubes were then centrifuged at 14,000 rpm for 10 minutes
190 and 600 µL of the supernatant was combined with the previous supernatant. Finally, 100 µL was
191 transferred into new tubes and dried using a vacuum centrifuge at 1 torr at room temperature for
192 1.5 - 2 hours.

193 *2.7.2 Gas chromatography mass spectrometry analysis:*

194 The dried samples were derivatized by adding 100 µL of methoxamine hydrochloride into each
195 tube and then incubated at 30 °C at 850 rpm for 1 hour using a Thermomixer F2.0 (Eppendorf).

Subsequently, 50 µL of N,O-bis(trimethylsilyl)trifluoroacetamide (BSTFA) was added to each

tube and incubated at 50 °C at 850 rpm for 3 hours using a Thermomixer F2.0 (Eppendorf). A

total of 150 µL of each sample was then transferred to vials in preparation for injection in the

GC-MS. Samples were analyzed via a 7890A gas chromatograph (Agilent) paired to a 5975C

quadrupole mass detector (Aglient). The initial GC oven temperature was set at 70 °C and then

increased to 325 °C at 7 °C per minute two minutes post-injection where the temperature was

then maintained at 325 °C for 3.6 minutes. The injection and ion source temperatures were set

to 300 °C and 230 °C, respectively with a solvent delay of 5 minutes. The helium flow rate was

also adjusted to a rate of 1 mL/minute. Measurements were performed with electron impact

ionization (70 eV) in full scan mode (m/z 33-600). Identification of metabolites was performed

based on their mass spectral and retention time index, and how closely they matched to those

acquired from the National Institute of Standards and Technology library and the Golm

Metabolome Database.

2.8 *Protein extraction and LC-MS analysis*

2.8.1 Protein extraction and LC-MS data acquisition

Kale leaf tissue harvested at ZT23 and ZT11 was flash-frozen in liquid N_2. Tissue was ground in

liquid N_2 using a mortar and pestle and aliquoted into 400 mg samples (n=4 replicates).

Samples were extracted with 50 mM HEPES-KOH pH 8.0, 50 mM NaCl, and 4% (w/v) SDS at a

ratio of 1:2 (w/v). Next, samples were vortexed followed by incubation for 15 min at 95 °C using

a Thermomixer F2.0 (Eppendorf) shaking at 1,100 rpm. Samples were subsequently incubated

at room temperature for an additional 15 min of shaking followed by clarification at 20,000 x g

for 5 min at room temperature, where the supernatant was then retained in fresh Eppendorf

microtubes. Protein concentration was estimated by bicinchoninic (BCA) assay (23225; Thermo

Scientific), followed by sample reduction using 10 mM dithiothreitol (DTT) for 5 min at 95 °C.

Following cooling to room temperature, samples were alkylated using 30 mM iodoacetamide

(IA) in dark for 30 min at room temperature without shaking. The alkylation was stopped by

addition of 10 mM DTT, followed by a brief vortex and 10 min room temperature incubation

without shaking. Samples were digested overnight with 1:100 sequencing grade trypsin (V5113;

Promega), followed by quantification of the generated peptide pools using a Nanodrop (Thermo

Scientific). Subsequently, samples were acidified with formic acid to a final concentration of 5%

(v/v) followed by drying by vacuum centrifugation. Peptide desalting was then performed using

ZipTip C18 pipette tips (ZTC18S960; Millipore), with peptides subsequently dried and dissolved

in 3% (v/v) ACN / 0.1% (v/v) FA prior to MS analysis. A Fusion Lumos Tribrid Orbitrap mass

spectrometer (Thermo Scientific) was used to analyse the digested peptides in data

independent acquisition (DIA) mode using the BoxCarDIA method as previously described[24]. 1

µg of dissolved peptide was injected using an Easy-nLC 1200 system (LC140;

ThermoScientific) and a 50 cm Easy-Spray PepMap C18 column (ES903; ThermoScientific).

Liquid chromatography and BoxCarDIA acquisition was performed as previously described[24].

2.8.2 Proteomic Data Analysis

All acquired BoxCarDIA data was analyzed using a library-free approach in Spectronaut v14

(Biognosys AG) under default settings. The *B. oleracea* var. *oleracea* proteome

(Uniprot:https://www.uniprot.org/ containing 58,545 proteins) was used for data searching.

Default search parameters were used for proteome quantification including: a protein, peptide

and PSM FDR of 1%, trypsin digestion with 1 missed cleavage, fixed modification including

carbamidomethylation of cysteine residues and variable modification including methionine

oxidation. Data was Log2 transformed and median normalized with significantly changing

differentially abundant proteins determined and corrected for multiple comparisons (Bonferroni-

corrected p-value <0.05; q-value).

2.9 *Bioinformatics*

Gene Ontology (GO) enrichment analysis was performed using the Database for Annotation,

Visualization and Integrated Discovery (DAVID; v 6.8; https://david.ncifcrf.gov/home.jsp).A)

using a significance threshold of p-value < 0.01. Conversion of *B. oleracea* gene identifiers to

Arabidopsis was performed using Uniprot (https://www.uniprot.org/) followed by Ensembl

Biomart (https://plants.ensembl.org/biomart) for STRING association network analysis and

metabolic pathway analysis. STRING association network analysis was performed using

Cytoscape v3.10.1 (https://cytoscape.org) in combination with the String DB plugin stringApp

and an edge threshold of 0.95. Visualization of GO dotplot was performed using R version 4.3.1

and *ggplot2* package. Metabolic pathway enrichment was performed using the Plant Metabolic

Network (PMN; https://plantcyc.org) with enrichment determined using a Fisher's exact test (p-

value < 0.01). Final figures were assembled using Affinity Designer (v.2.4.1).

3 Results

3.1 Kale morphology under modified light intensity and spectral composition

To test the effects of light intensity and spectral composition, kale seedlings were grown under
three different light intensities: 75 PPFD, 125 PPFD and 175 PPFD (**Supp Figure 1A**), while
light spectra experimentation consisted of changes in the ratio of red to blue (R/B) (**Supp**
Figure 1B) and red to far-red (R/FR) light (**Supp Figure 1B)**. We found a significant correlation
between plant size and increasing light intensity (**Figure 1A-B**), with both K3 and K9 exhibiting
an increase in size as a function of the light intensity increasing from 75 PPFD to 125 PPFD
(Figure 1B). We also found that fresh weight changes ranged from 2.5 to 7.5 g (a 3-fold
increase) 35 dpi (**Figure 1B**). Consistent with this, there was also an increase in plant area that
correlated with increasing light intensity for both cultivars (**Figure 1B**). Next, we examined the
effect of light intensity on diel leaf anthocyanin content, as anthocyanins provide coloration to
plants to attract pollinators, as well as acting as antioxidants, which can be beneficial for human
health[25,26]. As expected, anthocyanin was not detected in K3, however, there was not a
significant difference in anthocyanin content with increasing light intensity in K9, which is known
to produce anthocyanins (**Figure 1B**). It is possible that lower intensity than would otherwise be
experienced in outdoor growing conditions, the stage of growth and/or a general low stress
growth environment resulted in lower anthocyanin production than would be expected.

Next, we tested the effects of light spectral changes (**Figure 1C-D**), while maintaining light
intensity under constant 125 PPFD. Here, we found that light differentially modifies the
morphology of K3 plants (**Figure 1C**). R/B spectral variation affected K3 fresh weight (**Figure**
1D), with no corresponding change in surface area (**Figure 1D**). Conversely, K9 did not exhibit
spectral-dependent weight differences (**Figure 1D**), but did have increasing plant surface area
with changes in R/B and R/FR ratios (**Figure 1D**), indicating that red light is particularly
important for K9 growth. Despite anthocyanin content below detection level in K3, we do
observe significant diel spectral composition differences in K9 (**Figure 1D**). As R/B ratio
increases, we find anthocyanins decrease, while we alternatively find that red light (high R/FR
ratio) stimulates anthocyanin production over FR (low R/FR ratio). Therefore, changing light
intensity and spectral composition impacts morphological characteristics and anthocyanin levels
in kale.

3.2 Kale photosynthetic performance with changing light intensity and quality

Next, using a MultispeQ device, we measured a suite of photosynthetic parameters alongside
relative chlorophyll levels. These included fluorescence-based parameters such as ΦPSII,
ΦNPQ, ΦNO, and Linear Electron Flow (LEF). ΦPSII, or realized steady-state efficiency, which
measures how much of the light energy is absorbed by PSII (photosystem II). In other words,

ΦPSII represents the number of electrons transported via PSII per quantum absorbed by PSII[27].
ΦNPQ (non-photochemical quenching) represents the proportion of absorbed light energy that
is dissipated as heat, a process that helps protect the plant from excess energy that would
otherwise damage PSII[28], while ΦNO (non-regulatory energy dissipation) represents energy that
is neither used for photochemistry nor dissipated via regulatory quenching[28]. The other
parameters measured were the proton conductivity (gH^+), the electrochromic gradient across
the thylakoid membrane (ESCt) and the proton flux (vH^+) which are parameters linked to the
ATP synthase efficiency[22].

For the light intensity experiments (**Figure 2A**), an increase in PPFD influenced the chlorophyll
content (SPAD) in both K3 and K9. LEF also increased as a function of the light intensity for
both K3 and K9 (**Supp Figure 2A**). For K3, we observe a slight increase in proton conductivity
(gH^+) and decrease of the electrochromic shift decay across the thylakoid membrane (ESCt)
(**Supp Figure 2A**), indicating an increase in conductance[29]. LEF (Linear electron flow) and gH+
are linked to the flow of electrons from PSI to PSII and the flow of protons through the ATP
synthase[22], which supports the observed increase in plant fresh weight and area as the plant
produces more ATP required for plant growth. For the K9 cultivar, ΦPSII seemed to reach a
plateau under 125 PPFD (**Figure 2A**), with ATP synthase parameters demonstrating limited
light intensity-induced changes.

However, when light spectral composition was modified (**Figure 2B**), both K3 and K9 exhibited
a decrease in SPAD and ΦPSII with increasing R/B. Conversely, SPAD and ΦPSII increased as
R/FR increased for both cultivars. No change in the ATP synthase related parameters were
observed for K3 or K9, except LEF, which increased with increasing R/B (**Supp Figure 2B**).
Thus, kale photosynthetic parameters and chlorophyll content can be modulated by changes in
light intensity and spectral composition.

3.3 Assessing changes in kale metabolic content with varied light intensity and spectral
composition

To uncover a potential molecular rationale for the observable phenotypic changes, we next
analyzed the metabolite content of each cultivar by GC-MS when grown under the different light
intensities (**Figure 3**), R/B ratios (**Figure 4**) and R/FR ratios (**Figure 5**). Amino acids (15), fatty
acids (3), organic acids (11), polyamines (2), sterols (2), sugars (5), and sugar alcohols (2) were
detected and quantified. Under differing light intensities, we observed cultivar and light intensity
specific differences for Asn, Asp, Glu, Lys, Pro, Ser, Gly, and Ile. Asp, Glu and Pro that

correlated with an increase in light intensity in both cultivars and was more noticeable at ZT23 compared to ZT11. Another noticeable trend was a peak in Asn, Gly, Ile, Lys and Ser at both time-points under 125 PPFD. Conversely, change in R/B ratio (**Figure 4**) resulted in several amino acids having changes at ZT11. In particular, Gly which increased with increasing red light at ZT11 in both cultivars. Other changes seemed cultivar and time of day specific, such as Asn, which increased in K3 at ZT11 under high R/B, or Asp, which increased in K9 at ZT11 under high R/B. Alternatively, our main observations with modified R/FR included an increase in Ala, Asn and Lys levels in K3 and K9 at both time-points under balanced R/FR. We also find GABA and Gln quantities to be highest at ZT11 under balanced R/FR in both cultivars.

In the organic acids and polyamines, chlorogenic acid (related to the phenylpropanoid pathway, which includes lignin, flavonoids, coumarins, and lignans), glyceric acid, succinic acid, 3-O-coumaroyl-quinic acid and quininic acid were all positively affected by increasing light intensity (**Figure 3**). Additionally, citric acid and malic acid showed a large increase in both K3 and K9 under 175 PPFD, with malic acid higher at ZT11 and citric acid higher at ZT23 (**Figure 3**). With citric acid linked to plant growth and photosynthesis by alleviating plant stress[30], this may indicate that 175 PPFD represents a light intensity that initiates stress in kale. For our spectra experiments, increases quininic acid at ZT23 and citric acid, succinic acid, malic acid, and glyceric acid at both ZT11 and ZT23 were found with increasing R/B in both K3 and K9 (**Figure 4**), however, glyceric acid only increased at ZT23 in K9. We also found Spermidine, a polyamine, to be lower in K9 at ZT23 under balanced R/B (**Figure 4**). As well, Chlorogenic acid, which is important for anthocyanin production, was higher in K9, compared to K3, likely aligning with its known purpling phenotype (**Figure 5**). Lastly, with increasing R/FR, we find that K9 exhibits a greater overall number of organic acid / sterol compounds changing in abundance compared to K3 (**Figure 5**), except for quininic acid and 3-O-coumaroyl-D-quinic acid (**Supp Figure 3-5**).

With links to plant stress response, auxin perception and cell wall biosynthesis[31], the sugar alcohol Myo-inositol increases as a function of light intensity in both cultivars at ZT11 (**Figure 3**). We find Myo-inositol in K3 to be more affected by high PPFD than K9. However, Myo-inositol was higher in K3 at ZT23 under low R/B relative to K9 and higher at ZT11 in all the conditions relative to K9 (**Figure 4**). Glycerol decreases at ZT23 when R/B increased (**Figure 4**), which contrasts most changes we observe when R/B increased. Myo-inositol was the only sugar alcohol with a significant change in K3 over K9, which manifested under high R/FR at ZT11 (**Figure 5**).

Among sugar metabolites, fructose and glucose exhibited similar trends, showing a slight
increase with light intensity (**Figure3**). Mannose-6-phosphate (M6P) increases with intensity in
both cultivars and seems greatly affected in K3 with intensity under 175 PPFD. M6P is
associated with protein sorting and transport[32], stress, and the cell wall[33], all of which may
contribute to adaptation to higher light intensity[34]. The sugars fructose and glucose increased as
R/B increased in K9, while K3 showed no statistically significant changes (**Figure 4**).
Interestingly, M6P was higher in K3 at ZT23 under balanced R/B, while it was lower as R/B
increased at ZT11 (**Figure 4**). Sucrose showed very little change at either time-point. Finally,
trehalose in K9 seemed to increase with the R/B, while K3 only exhibited an increase at ZT11
under high R/B. Amongst the sugars, fructose and glucose was maximal in K3 at ZT23 under
balanced R/FR, while in K9, it was largely unchanged. At ZT11, K9 was increased in fructose
and glucose under low and balanced R/FR compared to high R/FR. Finally, both M6P and
sucrose were higher in K9 under balanced R/FR at ZT23 (**Figure 5**). Collectively, our results
indicate that changing light intensity and quality has dramatic impacts on overall plant
metabolism in kale.

3.4 Analysis of kale light intensity- and spectra- responsive proteome

We sought to further contextualize the observed phenotypic and metabolite changes by using
quantitative proteomics to examine the protein-level changes in K3 and K9 over the varying light
intensities and spectral compositions. We were able to quantify a total of 5,493 proteins across
our intensity experiments, and 6,527 proteins across spectra experiments (**Supp Table 3**). We
next identified significantly changing proteins using a fold-change analysis comparing each
varied light intensity or spectra to a control, considering 75 PPFD as control for light intensity
compared to 125 PPFD and 175 PPFD (**Figure 6A; Supp Table 3**) and low R/B (**Figure 6B;**
Supp Table 3) and low R/FR (**Figure 6C; Supp Table 3**) ratios compared to balanced and high
R/B and R/FR ratios, respectively. For our intensity experiments, K3 exhibited the most
significantly changing proteins from 75 PPFD to 125 PPFD at ZT11 and from 75 PPFD to 175
PPFD at ZT23. For spectral experiments, K9 appeared more responsive to increasing R/B and
R/FR ratios, possessing more significantly changing proteins than K3. Specifically, this involved
low to high R/B and R/FR ratios at ZT11, and increases under both balanced and high R/B and
R/FR ratios at ZT23.

We next performed gene ontology (GO) enrichment analysis using the significantly changing
proteome to resolve what biological processes are responsive to changes in either light intensity
or spectral composition (**Figure 6D**). This analysis revealed biological processes related to the

cell wall, photosynthesis / pigment metabolism, oxidative stress, carbohydrate metabolism, and

lipid metabolism across both intensity and spectra experiments. With increasing intensity, we

observed an enrichment of more biological processes when considering down-regulated versus

up-regulated proteins at both ZT23 and ZT11. Alternatively, we increased R/B ratio, we

observed more enriched biological processes among up-regulated proteins at both time-points.

Finally, with increasing R/FR ratio, we observed more enriched biological processes among up-

regulated proteins at ZT23, and among down-regulated proteins at ZT11.

Plant growth is associated with changes in cell wall composition and structure[35]. In

accordance with the changes in growth parameters detailed above, we observed changes in the

abundance of proteins involved in cell wall processes. In our intensity experiments, we primarily

observed the down-regulation of proteins involved in cell wall related biological processes with

increasing light intensity at both ZT23 and ZT11. This included the GO terms 'Plant-Type

Secondary Cell Wall Biogenesis' (GO:0009834), 'Cell Wall Organization' (GO:0071555)

'Xyloglucan Metabolic Process' (GO:0010411) and 'Cell Wall Biogenesis' (GO:0042546).

Xyloglucan is an abundant component of primary cell walls[36]. Among the down-regulated

proteins in K3 we find enrichment of 'Pectin Catabolic Processes' (GO:0045490) at ZT11 under

125 PPFD, with pectin representing another major component of plant primary cell walls[37]. With

increasing R/B, we also identified an enrichment of the GO term 'Xyloglucan Metabolic Process'

among up-regulated proteins in K3 at ZT11. In our R/FR spectral experiments, increasing R/FR

down-regulated proteins involved in 'Xyloglucan Metabolic Process', 'Cell Wall Organization',

and Cell Wall Biogenesis' in K3 at ZT11 under high R/FR. Conversely, a more complex

response was observed at ZT23 in K3, with 'Xyloglucan Metabolic Process and 'Cell Wall

Biogenesis' enriched among up-regulated proteins under balanced R/FR, and down-regulated

proteins enriched for 'Plant-Type Secondary Cell Wall' and 'Cell Wall Modification'

(GO:0042545) at the highest R/FR. Taken together, these results indicate that light conditions

substantially impact cell wall related processes in kale.

While the MultiSynQ measurements shed light on alterations in photosynthesis and chlorophyll

content with changing light intensity, our GO analysis revealed enrichment of terms associated

with photosynthesis and pigment metabolism. In our intensity experiments, down-regulated

proteins were enriched for 'Photosynthesis, Light Harvesting in Photosystem I' (GO:0009768)

under the highest PPFD in K3 at ZT23. In our R/B spectra experiments, a balanced R/B ratio in

K9 seemed to be optimal for photosynthetic performance, with up-regulated proteins enriched

for the GO terms 'Photosynthetic Electron Transport in Photosystem I' (GO:0009773) and

'Photosynthesis, Light Harvesting in Photosystem I' at ZT11. Additionally, we observed an enrichment of 'Carotenoid Metabolic Process' (GO:0016117) among down-regulated proteins at ZT11 in K3 under balanced R/B, and among up-regulated proteins under high R/B in K9 at ZT23. Finally, up-regulated proteins at ZT11 in K9 under high R/FR were enriched for the GO term 'Photosynthesis' (GO:0015979). These results suggest that kale photosynthetic processes are impacted by both light intensity and spectra composition, with increasing light intensity generally resulting in a down-regulation of light harvesting proteins and increasing red light ratios in higher photosynthetic processes.

In accordance with our measured changes in the abundance of proteins related to photosynthetic processes, we also observe protein-level changes associated with general metabolic processes, such as carbohydrate metabolism and lipid metabolism. In our intensity experiments, we observed a down-regulation of proteins enriched for 'Lipid Catabolic Process' (GO:0016042) and 'Glycolytic Process' (GO:0006096) in K3 at ZT11. With increasing R/B light, we observed an increase in proteins related to lipid biosynthesis in K3. This included enriched GO terms 'Wax Biosynthetic Process' (GO:0010025), 'Lipid Transport' (GO:0006869), and 'Lipid Biosynthetic Process' (GO:0008610) at ZT11, and of 'Lipid Oxidation' (GO:0034440) at ZT23. 'Glycolytic Process' was enriched among down-regulated proteins at ZT11 and ZT23 in K9 under high R/B, but among up-regulated proteins under balanced R/B in K3 at ZT23. Finally, at ZT11 down-regulated proteins were enriched for 'Carbohydrate Metabolic Process' (GO:0005975) in K3 under balanced R/B, and at ZT23 in both K3 and K9 there was an enrichment for the GO term 'Carbohydrate Metabolic Process' under balanced R/B ratio for up-regulated proteins. In our increasing R/FR experiments, at ZT11 down-regulated K3 proteins were enriched for 'Fatty Acid Biosynthetic Process' (GO:0006633), and at ZT23, down-regulated K9 proteins were enriched for 'Fatty Acid Metabolic Process' (GO:0006631). Additionally, 'Lipid Oxidation' was enriched among up-regulated K3 proteins under balanced R/FR. Glycolysis and TCA cycle proteins appeared to be down-regulated in K3 at both ZT11 and ZT23 under balanced R/FR, with enrichment of 'Succinyl-CoA Metabolic Process' (GO:0006104) and 'Glycolytic Process' at ZT11 and 'Mitochondrial Respiratory Chain Complex I Assembly' (GO:0032981), along with 'Glycolytic Process' at ZT23 in K9. Finally, 'Carbohydrate Metabolism' was enriched amongst down-regulated proteins in K9 at ZT11 under balanced R/FR, while 'Glucan Catabolic Process' (GO:0009251) was enriched amongst up-regulated under high R/FR in K9 at ZT11. Thus, light intensity and spectral composition play a large role in protein levels associated with metabolic processes in kale.

High light intensity can be associated with oxidative stress, and we observed changes in
proteins associated with oxidative stress responses in our intensity and spectra experiments.
For instance, in our intensity experiments, both kale cultivars had up-regulated proteins at ZT11
showing enrichment for 'Superoxide Metabolic Process' (GO:0006801) under 175 PPFD.
Interestingly, under 175 PPFD, the K3 cultivar showed an enrichment of the terms 'Toxin
Catabolic Process' (GO:0009407), 'Response to Oxidative Stress' (GO:0006979), and
'Glutathione Metabolic Process' (GO:0006749) amongst down-regulated proteins at ZT23,
indicating some diel regulation and cultivar-specific oxidative responses. We additionally
observed potential cultivar-specific oxidative stress response in our R/B spectral experiments at
ZT11, with K3 down-regulated proteins enriched for 'Glutathione Metabolic Process' under high
R/B and K9 up-regulated proteins enriched for 'Hydrogen Peroxide Catabolic Process'
(GO:0042744) under high R/B. At ZT23, only K3 showed altered oxidative stress response, with
down-regulated proteins enriched for 'Superoxide Metabolic Process' under high R/B and up-
regulated proteins enriched for 'Toxin Catabolic Process' and 'Glutathione Metabolic Process'
under high R/B. In our R/FR spectral experiments, K3 up-regulated proteins were enriched for
the terms 'Toxin Catabolic Process' and 'Glutathione Metabolic Process' under high R/FR at
ZT11, while K9 down-regulated proteins were enriched for 'Response to Oxidative Stress' under
balanced R/FR. At ZT23, K3 down-regulated proteins were similarly enriched for 'Response to
Oxygen-Containing Compound' (GO:19011700) and 'Response to Hydrogen Peroxide'
(GO:0042542), while up-regulated proteins were enriched for 'Glutathione Metabolic Process',
under balanced R/FR. Overall, these results indicate diel regulated, cultivar-specific responses
to oxidative stress with altered red light intensity and quality.

To further contextualize our GO analysis results, we performed an association network analysis
using STRING-DB (https://string-db.org/) for each of the Intensity, R/B and R/FR spectra
experimental conditions in order to further contextualize proteome-level responses (**Figures 7-**
8). As *Arabidopsis thaliana* (Arabidopsis) has more bioinformatic resources, we first converted
the *B. oleracea* gene identifiers for all quantified proteins in the study to Arabidopsis gene
identifiers (AGI) using UniProt (https://www.uniprot.org/) and Ensembl Biomart
(https://plants.ensembl.org/biomart). We identified unique AGIs for 90% of the significantly
changing proteins in our intensity experiments (2395/2662), along with 88.9% (2222/2499) and
89.4% (2520/2819) in our R/B and R/FR spectra experiments; respectively. Using a highly
stringent STRING-DB edge score of 0.95, this analysis corroborated our phenotyping
experiments and GO analysis. For photosynthesis and pigment metabolism, we observed a
down-regulation of protein with increasing intensity and R/FR, but an up-regulation with

increasing R/B ratio. Regarding oxidative stress responses, we observed complex changes in detoxification, glutathione metabolism and ascorbate metabolism, with proteins both up- and down-regulated across both intensity and spectra experiments. We additionally observe hubs of changing proteins related to carbohydrate metabolism, fatty acid biosynthesis and degradation, glycolysis and TCA cycle that align with our GO enrichment findings. In addition to our GO enrichment analysis, the association network analysis uniquely revealed large hubs of differentially expressed proteins related to spliceosome, ubiquitin / proteasome and phenylpropanoid metabolism.

Finally, using the Arabidopsis gene identifiers for the significantly changing proteins with increasing intensity, R/FR and R/B, we performed metabolic pathway enrichment using the Plant Metabolic Network (PMN; https://plantcyc.org; **Supp Tables 4-6**). The enriched pathways corroborated our findings with metabolite analysis, GO analysis and association networks, with enriched pathways being resolved for amino acid metabolism, lipid metabolism, carbohydrate metabolism, oxidative stress response, glucosinolate production, general metabolic processes, growth-associated hormones, photosynthesis and pigment metabolism (chlorophyll and carotenoids).

4 Discussion

Light intensity and spectral composition are major determinants of plant growth, metabolite content and yield. Given the health benefits of kale and its increased usage in vertical farm settings, it is important to understand how exposure to different light intensities and R/B or R/FR spectral compositions impact growth and nutritional quality. Several studies have identified changes in growth, yield and metabolite content in *Brassica* microgreens under different light intensities[38,39], and under different spectral compositions[16–18], however, there is a notable lack of knowledge about their effects at the molecular level, which is critical for future breeding endeavors to yeild gains. As a result, our combined use of plant phenotyping, physiology, metabolomics and quantitative proteomics offers a comprehensive analysis of the effects of varying light intensity and quality on two closely-related kale cultivars. Overall, our results point to changes in shade avoidance response, with shade presenting as lower light intensity, low blue light and low R/FR ratios[40], and manifesting as changes in growth and development, photosynthesis, phenylpropanoid metabolism, cell wall processes, amino acid production, carbohydrate metabolism, lipid and fatty acid metabolism and organic acid production. We further contrast the phenotypic, physiological and molecular responses of two kale cultivars with time-of-day precision, offering new levels of insight for precision vertical farming.

4.1 *Photoreceptors*

Changes in light quality are detected by plant photoreceptors, which invoke signal transduction
pathways that impact plant growth and development. These photoreceptors include
phytochromes (PhyA-E), which sense R/FR light, cryptochromes (CRY1-3) and phototropins
(PHOT1/2), which sense blue light, and UVR8, which senses UV-B[41]. We quantified several of
these in our experiments, including PhyB (A0A0D3BBQ1), CRY1 (A0A0D3E8M6), PHOT1
(A0A0D3BGL7; A0A0D3BGL8), PHOT2 (A0A0D3ECB2), and UVR8 (A0A0D3E235). However,
only a few of them showed significant changes in abundance across a few conditions. These
photoreceptors typically undergo regulation by phosphorylation[42–47]; thus, it is likely post-
translational modifications, rather than simply protein abundance that govern their responses to
changes in light intensity and spectral composition.

4.2 Proteome changes with changing light intensity and quality are associated with changes in
shade avoidance responses

4.2.1 Growth, development and hormone signaling

Previous studies have reported a linear relationship between light intensity and fresh weight in
pea microgreens[48], as well as kale, cabbage, arugula and mustard[49], similar to our
observations. Importantly, several of these studies used PPFD up to 600 µmol·m^{-2}·s^{-1} with
increases in fresh weight at the upper PPFD values, suggesting our highest light intensity of 175
PPFD is not inducing a high light stress response. As light intensities used in vertical farming
are typically lower, around 200 PPFD[50], our study is highly relevant to informing practices for
kale grown in vertical farms. With our spectra experiments, we only observed an increase in K3
fresh weight with increasing R/B ratio, and an increase in K9 surface area with both increasing
R/B and R/FR ratios. In alignment with this, a previous report found an increase in dry mass and
seedling area specifically with red kale varieties, whereas blue light had a stronger effect on
green kale dry mass[51], suggesting growth parameters under different light qualities are cultivar-
dependent. This emphasizes the need to generate cultivar-specific molecular response data,
such as that generated here, to uncover unique breeding opportunities.

Our proteomic data revealed that under the conditions tested, hormone pathways
associated with plant growth and development, including brassinosteroids (BR) and auxin, play
a key role. For example, we observed down-regulation of K3 proteins involved in BR signaling
with increasing light intensity. This is consistent with previous reports of Arabidopsis BR
biosynthetic genes *CONSTITUTIVE PHOTOMORPHIC DWARF* (*CPD)*, and *DWF4* being dark

induced, but light repressed[52–55], as well as the transcription of BR signaling proteins *bri1 EMS*
SUPPRESSOR1 (BES1) and *BRASSINOZOLE RESISTANT1 (BZR1)* elevation at night and
repression during the day[54]. Additionally, BR signaling appears to play a role in shade
avoidance response[56,57]. In our study, we observe the down-regulation of kale DIMUNUTO
(DIM)/DWF1 protein (A0A0D3CI99) in K9 with increasing light intensity, which has been
reported to be involved in shade-induced hypocotyl elongation in Arabidopsis[58]. We additionally
observe down-regulation of BZR1 (A0A0D3CWJ3) in K9 at ZT11 and up-regulation at ZT23 with
increasing light intensity, suggesting differential regulation of its protein stability and abundance.
The observed down-regulation within our dataset with increasing light intensity indicates relief
from perceived shade conditions. In addition to BR signaling proteins changing with light
intensity, we observed an up-regulation of proteins involved in auxin signaling with an increasing
R/B ratio in K3. Previous work has identified red light as inducing auxin flux from leaves to roots,
possibly through increased abundance of *PINOID1* (*PIN)1, PIN3,* and *PIN4*[59], thus our observed
up-regulation of PIN3 (A0A0D3CZR1) in K3 under high R/B may indicate a similar response.
Additionally, up-regulation of auxin signaling is consistent with increased shade avoidance
response, where auxin signaling has been shown to play a role in the hypocotyl elongation[60].
Overall, our results indicate that modulation of light intensity and spectral quality induce
changes in kale growth and development, possibly through changes in key hormone signaling
pathways.

4.2.2 Cell wall

Our observation of down-regulated BR biosynthesis and signaling proteins under increasing
light intensity suggests a down-regulation of shade avoidance responses. Related to BR shade
avoidance response is changes in cell wall structure, particularly by the action of xyloglucan
endotransglucosylase/hydrolases (XTHs) and expansins[61,62], with cell wall modification
promoting elongation of the hypocotyl to drive growth toward light. Within our dataset, we
observe a multitude of cell wall proteins changing in abundance in response to light intensity
and spectral composition, potentially relieving shade-like conditions in both kale cultivars.
Specifically, we note a down-regulation of cell wall organization proteins, primarily XTHs,
expansins (EXPAs), and pectinesterases (PMEs) with increasing light intensity and R/FR ratio,
but an up-regulation of XTHs with increasing R/B, consistent with changes in the plant
perception of shade. Interestingly, more members of the XTH family were down-regulated in K9
with increasing light intensity and K3 with increasing R/FR, suggesting K9 may be more

sensitive to changes in intensity and K3 more sensitive to changes in red light wavelengths.
This is also consistent with our growth observations.

4.2.3 Photosynthesis

Light intensity and quality also critically impact plant photosynthesis and light harvesting, as well
as pigment metabolism (e.g. chlorophyll and carotenoids). Our LEF measure, used as an
approximation for photosynthetic rate[22], indicates a higher photosynthetic rate with increasing
intensity and R/B ratio in both K3 and K9, and with increasing R/FR in K9. Intriguingly, our
proteomics data implies a concurrent down-regulation of light harvesting proteins at higher
intensities at ZT23 in K3, with LHCA1 (A0A0D3CTW2; A0A0D3DT59), LHCA2 (A0A0D3DUU9;
A0A0D3BWE5), LHCA3 (A0A0D3E520), LHCB2.4 (A0A0D3DBD3), LHCB4.2 (A0A0D3CLT0)
and LHCB5 (A0A0D3BQY2) all down-regulated. In the K9 cultivar, we only find LHCA2 and
LHCB5 were down-regulated, indicating a stronger response in K3. This is likely due to
sufficient, or even excessive light energy under the 175 PPFD light intensity, such that light
harvesting potential has been reached, a phenomenon which is well-established in response to
light/shade levels[63–65]. Correspondingly, under a higher R/FR ratio we find the down-regulation
of light harvesting proteins, consistent with the relief of shade avoidance conditions. By contrast,
under a balanced R/B ratio we see an up-regulation of light harvesting and photosynthesis-
related proteins, particularly in K9, suggesting an optimal ratio of R/B light is best for light
harvesting and photosynthetic responses. Red light can act to promote photosynthesis and
plant growth, but excessive amounts of red light can cause an imbalance in PSI and PSII, and
thus blue light is needed[66]. Overall, our data suggest light-dependent differences exist between
the two cultivars, with K3 light harvesting proteins showing a stronger response to light intensity
changes, and K9 more to changes in spectral composition.

4.3 Alteration of light intensity and spectral composition changes kale nutritional profile

Kale consumption is associated with myriad health benefits as a result of anthocyanins and
flavonoids, carotenoids, amino acids, glucosinolates, vitamins, sugars and phenolic
compounds[18,67,68]). Several studies have assessed the impact of light intensity and quality on
nutritional composition in a range of *Brassica* microgreens[38,69], however, none provide data
linking phenotypes with metabolic and proteomic profiles with time-of-day precision.

4.3.1 Anthocyanins / phenylpropanoids

Anthocyanins are a natural antioxidant source, beneficial for human health as reactive oxygen
species scavangers[70,71] with the generation of anthocyanins typically driven by light[72]. In our
study, we were only able to detect anthocyanins within the K9 cultivar, with no significant
difference with changing light intensity, but a significant decrease with increasing R/B, and
increase with increasing R/FR. Within our metabolite analysis, we observed changes associated
with anthocyanin and phenylpropanoid metabolism: including chlorogenic acid and quininic acid
with changing R/B, and sinapinic acid, chlorogenic acid, quininic acid and 3-coumaroyl-D-quinic
acid in R/FR. Additionally, within our metabolic pathway analysis, we observed significant
pathway enrichment associated with phenylpropanoid and flavonoid metabolism with increasing
R/B and R/FR composition. As expected with increasing R/FR, we see a significant up-
regulation of flavonoid biosynthetic pathways, which is more pronounced in the K9 cultivar and
consistent with our ability to detect anthocyanins in this cultivar. Intriguingly, we observe an up-
regulation of phenylpropanoid biosynthesis pathways in K9 with increasing R/B at ZT23, which
contrasts with our observed anthocyanin levels. It is possible that at ZT23 the phenylpropanoid
pathway is diverting the flux toward other phenylpropanoid products. For example, we see the
coumarin biosynthesis pathway enriched in our metabolic pathway analysis, coupled with
observed up-regulation of DEFECTIVE IN CUTICULAR RIDGES (DCR; A0A0D3DD85),
ENHACED PSEUDOMONAS SUSCEPTIBILITY1 (EPS1; A0A0D3D884) and COUMARIN
SYNTHASE (COSY; A0A0D3CEC8) in K9 at ZT23, indicating possible increased coumarin
biosynthesis[73–75]. Thus, our results suggest the ideal conditions for kale anthocyanin production
under lower intensities involves the use of a high R/FR spectral composition, with higher R/B
ratios having negative impacts by possibly diverting the phenylpropanoid pathway toward other
products.

4.3.2 Glucosinolates

Another component of kale consumption beneficial for human health is that of glucosinolates,
which are amino acid-derived with sulphur and thioglucosidic moieties[76,77]. Playing a role in kale
flavour and plant defense, glucosinolates can also act as anticancer compounds for human
health[78]. Our proteomic analyses indicated changes in glucosinolate metabolism with increasing
light intensity and R/B ratio. In our intensity experiments, the K9 cultivar showed a down-
regulation of proteins associated with glucosinolate catabolism as light intensity increased. Our
metabolic pathway analysis indicated increased biosynthesis of glucosinolates in K3, thus the
cultivars might achieve the same elevated levels of glucosinolates at higher light intensities
through differing mechanisms. In our assessment of R/B responses, we observe a decrease in

glucosinolate catabolism by GO analysis and an increase in glucosinolate biosynthesis by

metabolic pathway analysis in K3. This has previously been observed with levels of the

glucosinolates sinigrin, glucobrassicin, and 4-methoxy glucobrassicin in *B. napus* sprouts

exposed to increasing red light levels[79], while another study found differential cultivar responses

with changing light intensities in Chinese cabbage[80].

4.3.3 Carotenoids

In addition to flavonoids and glucosinolates, carotenoids are also antioxidants compounds

beneficial to human health[81]. Within our proteomic data, we find K3 has lower carotenoid

production with increasing light intensity, combined with decreasing protein abundance of

CAROTENE CIS-TRANS-ISOMERASE (CRTISO; A0A0D3DYP6), 15-CIS-PHYTOENE

DESATURASE (PDS; A0A0D3BJ64) and PHYTOENE SYNTHASE1 (PSY1; A0A0D3AJG7). In

our spectral experiments, it appears that increasing red light in both the R/B and R/FR

experiments drives increased carotenoid biosynthesis in the K9 cultivar, along with increasing

protein abundance of CRTISO (A0A0D3DYP6), LYCOPENE EPSILON CYCLASE2 (LUT2;

A0A0D3B320), PSY1 and PDS in R/B experiments and PDS, CRTISO and CYP97B3 under

R/FR conditions. These results suggest that K3 might be more sensitive to light intensity

changes and K9 to levels of red light. We thought it interesting to find lower carotenoid

biosynthesis with increasing light intensity in K3 as carotenoids tend to increase with higher light

intensity[82], however there is evidence in pepper of lower carotenoids with increasing light

intensity that is suggested to be due to a higher photo-oxidation rate than that of carotenoid

synthesis[83,84].

4.3.4 Amino acids

Our metabolite profiling identified several changes in the amino acid content of K3 and K9 kale

in response to changing light intensity and quality. We observe extensive changes in our

metabolic and proteomic analyses associated with the glutamate family of amino acids,

including glutamine, glutamate, GABA, proline, arginine. These amino acids represent the

highest amino acid quantities in plants, with glutamic acid and proline present at the highest

quantity in kale[85]. Glutamate metabolism also represents a critical link between nitrogen and

carbon metabolism through the TCA cycle, with several glutamate catabolism products,

including proline and GABA, associated with plant stress responses[86,87]. With increasing light

intensity, we observe a down-regulation of proteins involved in glutamate catabolism, including

GLUTAMATE DEHYDROGENASE1 (GDH1; A0A0D3EEJ0) and GDH2 (A0A0D3AI31), which

converts glutamate to glutamine, DELTA-1-PYRROLINE-5-CARBOXYLATE SYNTHASE B

(P5CSB; A0A0D3BXP3), involved in proline biosynthesis, N-ACETYLGLUTAMATE SYNTHASE

(NAGS; A0A0D3A172) and N-ACETYLGLUTAMATE KINASE (NAGK; AT3G57560), which are

steps toward ornithine biosynthesis, and CARBAMOYL-PHOSPHATE SYNTHASE ALPHA

(CARA; A0A0D3CZA9), toward arginine biosynthesis. We found this trend in both K3 and K9

and at both ZT11 and ZT23, suggesting increasing the light intensity in our experiments may

relieve stress responses associated with low light. This also coincides with increased glutamic

acid accumulation at higher light intensities (**Figure 3**). However, we also observe an increase

in proline in our metabolomics data under higher light intensities, with a significant increase in

K3 at ZT11, which may suggest sufficient levels of proline and the down-regulation of P5CSB to

prevent further accumulation. Similar findings are present in our proteome data from increasing

R/B ratio, with proteins associated with proline biosynthesis and glutamate metabolism primarily

down-regulated at ZT23 in both K3 and K9. However, although we did not find a significant

difference in the levels of glutamic acid or proline in either cultivar. Thus, increasing amounts of

red light in the presence of blue light may be associated with less stressful conditions, however

not to the extent of our light intensity manipulation. The situation was slightly more complex with

increasing R/FR light. Our finding of P5CSA up-regulation suggests increased proline

biosynthesis, while increase in ARGINASE 1 (ARGAH1; A0A0D3B8E0) and CARA, suggest

enhanced flux through the glutamate pathway in K3, whereas glutamate metabolic proteins are

down-regulated in K9. The metabolite data displayed a peak at balanced R/FR in K3 for the

amino acids glutamate, glutamine and GABA, but a less clear trend in K9, suggesting

differential amino acid regulation between the cultivars with changing R/FR light ratios.

The aspartate-derived amino acids are also of great importance for human nutrition as they

cannot be synthesized by the human body and must be obtained through diet[88]. This includes

the essential amino acids lysine, threonine, methionine and isoleucine. Of particular interest is

lysine, as it is generally of very low abundance in plants[89]. We observe a general down-

regulation of proteins associated with lysine biosynthesis as light intensity increases, particularly

in K3 at ZT11, with a down-regulation of ASPARTATE KINASE (AK2) and of

DIHYDRODIPICOLINATE SYNTHASE1 (DHDPS1), both involved in lysine biosynthesis.

However, within our metabolite data, there is a significantly higher amount of lysine present in

K3 under 125 PPFD than either 75 or 175 PPFD, suggesting more complex regulation of lysine

biosynthesis. This has previously been observed in attempts to engineer greater lysine

production, where lysine has been shown to allosterically inhibit the activity of DHDPS1[90], as

well as impacting other aspartate-derived amino acids, such as threonine[91]. In contrast to

increasing light intensity, increasing R/B ratio induced an up-regulation of proteins involved in lysine biosynthesis, which only somewhat mirrored an increase in lysine metabolite abundance at ZT11. Finally, proteomic data suggested an increase in R/FR ratio leads to an up-regulation of lysine biosynthesis in K9, which along with the increasing R/B trend suggests greater proportion of red light may promote lysine synthesis, whereas balanced R/FR seems to have the highest measurable lysine levels.

We previously identified diel regulation of branched-chain amino acids (BCAA) in kale[21], of which we observed changes in the levels of isoleucine with intensity (highest under 125 PPFD), which mirrored down-regulation of branched chain amino acids catabolic enzymes in our proteomic data. Additionally, valine trending higher under balanced R/FR, mirrored the down-regulation of valine catabolic enzymes in our proteomic data under balanced R/FR. Finally, we observe altered serine levels correlating with changing light intensity and spectral composition. Given serine is a key substrate for other essential amino acids, such as methionine and cysteine[92], this is likely of interest for human nutrition.

4.3.5 Carbohydrates

Aligning with our observed changes in amino acids and corresponding metabolic enzymes, we also observe changes in carbohydrate metabolism, manifesting in the differential abundance of sugars and of enzymes involved in carbohydrate metabolism. In our proteomic data, we observe a general down-regulation of proteins involved in glycolysis and starch degradation, as well as trehalose biosynthetic proteins. As trehalose has been associated with shade avoidance[93] and regulation of the circadian clock[94], this is consistent with reduced stress under low light intensities associated with shade as we increase the light intensity from 75 PPFD to 125 and 175 PPFD. Specifically, we find that increasing the R/B ratio down-regulates proteins associated with glycolysis in both cultivars, and mannose metabolism in K9. This was consistent with the lowest M6P levels found in K9 under balanced R/B. Finally, increasing R/FR was associated with the down-regulation of glycolytic proteins in both cultivars, as well as myo-inositol biosynthetic proteins in K9. Conversely, myo-inositol biosynthesis was up-regulated in K3 with increasing R/FR. In K3, this showed moderate correlation with corresponding metabolite levels, as myo-inositol level was significantly higher in K3 at ZT11 under high R/FR. As myo-inositol is involved in many biological processes, including auxin biosynthesis, membrane biogenesis, light and abiotic stress responses[95], its modulation may impact several of the morphological responses we observe.

4.4 Conclusions

In summary, we present an integrated systems-level analysis of two closely-related kale cultivars under increasing light intensity, R/B and R/FR spectral ratios with time-of-day precision. Our data reveals that light intensity and spectral composition can modulate myriad processes within kale, including changes in photosynthesis, cell wall, nutritional components such as amino acids, carbohydrates, phenylpropanoid metabolism among others. Overall, this rich dataset serves not only as a fundamentally interesting dataset for further characterization of kale, it also offers to inform future opportunities for enhanced kale cultivation in vertical farms for desired morphological and nutritional traits.

ACKNOWLEDGEMENTS

The authors thank the Natural Sciences and Engineering Research Council of Canada (NSERC), Alberta Innovates (AI) and Canada Foundation for Innovation (CFI) for funding. The authors thank G2V Optics Inc. for providing their programmable LED lighting system as well as Jack Moore of the Alberta Proteomics and Mass Spectrometry Facility for assistance with mass spectrometer operation and maintenance.

AUTHOR CONTRIBUTIONS

R.G.U. and S.S. conceptualized the study. S.S., B.C., L.I., C.K., N.B. performed research and data acquisition. S.S., L.E.G. and R.G.U. analyzed the data and wrote the paper with input from all authors.

DATA AVAILABILITY

All raw data files have been uploaded to ProteomeXchange (http://www.proteomexchange.org/) via the PRoteomics IDEntification Database (PRIDE; https://www.ebi.ac.uk/pride/) with the dataset identifier PXD055752.

CONFLICTS OF INTEREST

None to declare

REFERENCES

1. Satheesh, N. & Workneh Fanta, S. Kale: Review on nutritional composition, bio-active
compounds, anti-nutritional factors, health beneficial properties and value-added products.
Cogent Food Agric. **6**, 1811048 (2020).
2. Kim, S. Y., Yoon, S., Kwon, S. M., Park, K. S. & Lee-kim, Y. C. Kale juice improves coronary
artery disease risk factors in hypercholesterolemic men. *Biomed. Environ. Sci.* **21**, 91–97
(2008).
3. Raychaudhuri, S., Fan, S., Kraus, O., Shahinozzaman, M. & Obanda, D. N. Kale
supplementation during high fat feeding improves metabolic health in a mouse model of
obesity and insulin resistance. *PLOS ONE* **16**, e0256348 (2021).
4. Rachwał, K. *et al.* Red kale (*Brassica oleracea* L. *ssp. acephala* L. var. *sabellica*) induces
apoptosis in human colorectal cancer cells *in vitro. Molecules* **28**, 6938 (2023).
5. Bowen-Forbes, C. *et al.* Broccoli, kale, and radish sprouts: key phytochemical constituents
and DPPH free radical scavenging activity. *Molecules* **28**, 4266 (2023).
6. Cao, D. *et al.* Spermidine enhances chilling tolerance of kale seeds by modulating ROS and
phytohormone metabolism. *PLOS ONE* **18**, e0289563 (2023).
7. Bauer, N. *et al.* Mechanisms of kale (*Brassica oleracea var. acephala*) tolerance to
individual and combined stresses of drought and elevated temperature. *Int. J. Mol. Sci.* **23**,
11494 (2022).
8. Fussy, A. & Papenbrock, J. An overview of soil and soilless cultivation techniques—
chances, challenges and the neglected question of sustainability. *Plants* **11**, 1153 (2022).
9. https://www.alliedmarketresearch.com, A. M. R. Kale Microgreen Market Size, Share, Price,
Trends |Research Report 2030. *Allied Market Research*
https://www.alliedmarketresearch.com/kale-microgreen-market-A16137.
10. Avgoustaki, D. D. & Xydis, G. Chapter One - How energy innovation in indoor vertical
farming can improve food security, sustainability, and food safety? in *Advances in Food*
Security and Sustainability (ed. Cohen, M. J.) vol. 5 1–51 (Elsevier, 2020).
11. Hati, A. J. & Singh, R. R. Smart indoor farms: Leveraging technological advancements to
power a sustainable agricultural revolution. *AgriEngineering* **3**, 728–767 (2021).
12. Nájera, C., Gallegos-Cedillo, V. M., Ros, M. & Pascual, J. A. LED lighting in vertical farming
systems enhances bioactive compounds and productivity of vegetables crops. *Biol. Life Sci.*
Forum **16**, 24 (2022).
13. Ma, Y., Xu, A. & Cheng, Z.-M. (Max). Effects of light emitting diode lights on plant growth,
development and traits a meta-analysis. *Hortic. Plant J.* **7**, 552–564 (2021).
14. McCree, K. J. The action spectrum, absorptance and quantum yield of photosynthesis in
crop plants. *Agric. Meteorol.* **9**, 191–216 (1971).
15. Krahmer, J., Ganpudi, A., Abbas, A., Romanowski, A. & Halliday, K. J. Phytochrome, carbon
sensing, metabolism, and plant growth plasticity. *Plant Physiol.* **176**, 1039–1048 (2018).
16. Chen, J. *et al.* Effect of photoperiod on chinese kale (*Brassica alboglabra*) sprouts under
white or combined red and blue light. *Front. Plant Sci.* **11**, (2021).
17. Kong, Y., Schiestel, K. & Zheng, Y. Pure blue light effects on growth and morphology are
slightly changed by adding low-level UVA or far-red light: A comparison with red light in four
microgreen species. *Environ. Exp. Bot.* **157**, 58–68 (2019).
18. Deng, M. *et al.* Influence of pre-harvest red light irradiation on main phytochemicals and
antioxidant activity of Chinese kale sprouts. *Food Chem.* **222**, 1–5 (2017).
19. Zhen, S. & van Iersel, M. W. Far-red light is needed for efficient photochemistry and
photosynthesis. *J. Plant Physiol.* **209**, 115–122 (2017).

20. Furuya, M. & Schäfer, E. Photoperception and signalling of induction reactions by different
phytochromes. *Trends Plant Sci.* **1**, 301–307 (1996).
21. Scandola, S., Mehta, D., Castillo, B., Boyce, N. & Uhrig, R. G. Systems-level proteomics
and metabolomics reveals the diel molecular landscape of diverse kale cultivars. *Front.*
Plant Sci. **14**, (2023).
22. Kuhlgert, S. *et al.* MultispeQ Beta: a tool for large-scale plant phenotyping connected to the
open PhotosynQ network. *R. Soc. Open Sci.* **3**, 160592 (2016).
23. Gehan, M. A. *et al.* PlantCV v2: Image analysis software for high-throughput plant
phenotyping. *PeerJ* **5**, e4088 (2017).
24. Mehta, D., Scandola, S. & Uhrig, R. G. BoxCar and library-free data-independent acquisition
substantially improve the depth, range, and completeness of label-free quantitative
proteomics. *Anal. Chem.* **94**, 793–802 (2022).
25. He, J. & Giusti, M. M. Anthocyanins: Natural colorants with health-promoting properties.
Annu. Rev. Food Sci. Technol. **1**, 163–187 (2010).
26. Zhang, Y., Butelli, E. & Martin, C. Engineering anthocyanin biosynthesis in plants. *Curr.*
Opin. Plant Biol. **19**, 81–90 (2014).
27. Genty, B., Briantais, J.-M. & Baker, N. R. The relationship between the quantum yield of
photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochim.*
Biophys. Acta BBA - Gen. Subj. **990**, 87–92 (1989).
28. Baker, N. R., Harbinson, J. & Kramer, D. M. Determining the limitations and regulation of
photosynthetic energy transduction in leaves. *Plant Cell Environ.* **30**, 1107–1125 (2007).
29. Asher, A. H. & Theg, S. M. Electrochromic shift supports the membrane destabilization
model of Tat-mediated transport and shows ion leakage during Sec transport. *Proc. Natl.*
Acad. Sci. **118**, e2018122118 (2021).
30. Mallhi, Z. I. *et al.* Citric acid enhances plant growth, photosynthesis, and phytoextraction of
lead by alleviating the oxidative stress in castor beans. *Plants* **8**, 525 (2019).
31. Eckardt, N. A. Myo-Inositol biosynthesis genes in Arabidopsis: Differential patterns of gene
expression and role in cell death. *Plant Cell* **22**, 537 (2010).
32. Castonguay, A. C., Olson, L. J. & Dahms, N. M. Mannose 6-phosphate receptor homology
(MRH) domain-containing lectins in the secretory pathway. *Biochim. Biophys. Acta BBA -*
Gen. Subj. **1810**, 815–826 (2011).
33. Verbančič, J., Lunn, J. E., Stitt, M. & Persson, S. Carbon supply and the regulation of cell
wall synthesis. *Mol. Plant* **11**, 75–94 (2018).
34. Díaz, M., De Haro, V., Muñoz, R. & Quiles, M. J. Chlororespiration is involved in the
adaptation of Brassica plants to heat and high light intensity. *Plant Cell Environ.* **30**, 1578–
1585 (2007).
35. Cosgrove, D. J. Growth of the plant cell wall. *Nat. Rev. Mol. Cell Biol.* **6**, 850–861 (2005).
36. Wilson, L. F. L. *et al.* The biosynthesis, degradation, and function of cell wall β-xylosylated
xyloglucan mirrors that of arabinoxyloglucan. *New Phytol.* **240**, 2353–2371 (2023).
37. Tong, L. *et al.* Powerful cell wall biomass degradation enzymatic system from saprotrophic
Aspergillus fumigatus. *Cell Surf. Amst. Neth.* **11**, 100126 (2024).
38. Kamal, K. Y. *et al.* Evaluation of growth and nutritional value of Brassica microgreens grown
under red, blue and green LEDs combinations. *Physiol. Plant.* **169**, 625–638 (2020).
39. He, R. *et al.* The combination of selenium and LED light quality affects growth and
nutritional properties of broccoli sprouts. *Molecules* **25**, 4788 (2020).
40. Pierik, R. & de Wit, M. Shade avoidance: phytochrome signalling and other aboveground
neighbour detection cues. *J. Exp. Bot.* **65**, 2815–2824 (2014).

41. Paik, I. & Huq, E. Plant photoreceptors: Multi-functional sensory proteins and their signaling
networks. *Semin. Cell Dev. Biol.* **92**, 114–121 (2019).
42. Liu, W. *et al.* Phosphorylation of Arabidopsis UVR8 photoreceptor modulates protein
interactions and responses to UV-B radiation. *Nat. Commun.* **15**, 1221 (2024).
43. Liu, X. *et al.* FERONIA coordinates plant growth and salt tolerance via the phosphorylation
of phyB. *Nat. Plants* **9**, 645–660 (2023).
44. Viczián, A. *et al.* Differential phosphorylation of the N-terminal extension regulates
phytochrome B signaling. *New Phytol.* **225**, 1635–1650 (2020).
45. Gao, L. *et al.* Blue light-induced phosphorylation of Arabidopsis cryptochrome 1 is essential
for its photosensitivity. *J. Integr. Plant Biol.* **64**, 1724–1738 (2022).
46. Schumacher, P. *et al.* A phosphorylation switch turns a positive regulator of phototropism
into an inhibitor of the process. *Nat. Commun.* **9**, 2403 (2018).
47. Inoue, S. *et al.* Blue light-induced autophosphorylation of phototropin is a primary step for
signaling. *Proc. Natl. Acad. Sci.* **105**, 5626–5631 (2008).
48. Balázs, L. *et al.* Quantifying the effect of light intensity uniformity on the crop yield by pea
microgreens growth experiments. *Horticulturae* **9**, 1187 (2023).
49. Jones-Baumgardt, C., Llewellyn, D., Ying, Q. & Zheng, Y. Intensity of sole-source light-
emitting diodes affects growth, yield, and quality of Brassicaceae microgreens. (2019)
doi:10.21273/HORTSCI13788-18.
50. Dou, H. *et al.* Supplementary far-red light for photosynthetic active radiation differentially
influences the photochemical efficiency and biomass accumulation in greenhouse-grown
lettuce. *Plants* **13**, 2169 (2024).
51. Frąszczak, B. *et al.* Morphological and photosynthetic parameters of green and red kale
microgreens cultivated under different light spectra. *Plants* **12**, 3800 (2023).
52. Bancos, S. *et al.* Diurnal regulation of the brassinosteroid-biosynthetic CPD gene in
Arabidopsis. *Plant Physiol.* **141**, 299–309 (2006).
53. Pan, Y. *et al.* Cytochrome P450 monooxygenases as reporters for circadian-regulated
pathways. *Plant Physiol.* **150**, 858–878 (2009).
54. Martínez, C. *et al.* PIF4-induced BR synthesis is critical to diurnal and thermomorphogenic
growth. *EMBO J.* **37**, e99552 (2018).
55. Asahina, M. *et al.* Blue light-promoted rice leaf bending and unrolling are due to up-
regulated brassinosteroid biosynthesis genes accompanied by accumulation of
castasterone. *Phytochemistry* **104**, 21–29 (2014).
56. Lv, M. & Li, J. Molecular mechanisms of brassinosteroid-mediated responses to changing
environments in Arabidopsis. *Int. J. Mol. Sci.* **21**, 2737 (2020).
57. Keuskamp, D. H. *et al.* Blue-light-mediated shade avoidance requires combined auxin and
brassinosteroid action in Arabidopsis seedlings. *Plant J.* **67**, 208–217 (2011).
58. Klahre, U. *et al.* The Arabidopsis DIMINUTO/DWARF1 gene encodes a protein involved in
steroid synthesis. *Plant Cell* **10**, 1677–1690 (1998).
59. Yun, F., Liu, H., Deng, Y., Hou, X. & Liao, W. The role of light-regulated auxin signaling in
root development. *Int. J. Mol. Sci.* **24**, 5253 (2023).
60. Ma, L. & Li, G. Auxin-dependent cell elongation during the shade avoidance response.
Front. Plant Sci. **10**, (2019).
61. Sasidharan, R. *et al.* Light quality-mediated petiole elongation in Arabidopsis during shade
avoidance involves cell wall modification by xyloglucan endotransglucosylase/hydrolases.
Plant Physiol. **154**, 978–990 (2010).

62. Sasidharan, R., Chinnappa, C. C., Voesenek, L. A. C. J. & Pierik, R. The regulation of cell
wall extensibility during shade avoidance: a study using two contrasting ecotypes of
Stellaria longipes. Plant Physiol. **148**, 1557–1569 (2008).
63. Teramoto, H., Nakamori, A., Minagawa, J. & Ono, T. Light-intensity-dependent expression
of Lhc gene family encoding light-harvesting chlorophyll-a/b proteins of photosystem II in
Chlamydomonas reinhardtii. Plant Physiol. **130**, 325–333 (2002).
64. Anderson, J. M., Chow, W. S. & Goodchild, D. J. Thylakoid membrane organisation in
sun/shade acclimation. *Funct. Plant Biol.* **15**, 11–26 (1988).
65. Anderson, J. M., Chow, W. S. & Park, Y.-I. The grand design of photosynthesis: Acclimation
of the photosynthetic apparatus to environmental cues. *Photosynth. Res.* **46**, 129–139
(1995).
66. Jin, D. *et al.* Effect of red and blue light on cucumber seedlings grown in a plant factory.
Horticulturae **9**, 124 (2023).
67. Podsędek, A. Natural antioxidants and antioxidant capacity of Brassica vegetables: A
review. *LWT - Food Sci. Technol.* **40**, 1–11 (2007).
68. Sun, B., Yan, H., Liu, N., Wei, J. & Wang, Q. Effect of 1-MCP treatment on postharvest
quality characters, antioxidants and glucosinolates of Chinese kale. *Food Chem.* **131**, 519–
526 (2012).
69. Liu, K. *et al.* Light intensity and photoperiod affect growth and nutritional quality of Brassica
microgreens. *Molecules* **27**, 883 (2022).
70. Tena, N., Martín, J. & Asuero, A. G. State of the art of anthocyanins: antioxidant activity,
sources, bioavailability, and therapeutic effect in human health. *Antioxidants* **9**, 451 (2020).
71. Pojer, E., Mattivi, F., Johnson, D. & Stockley, C. S. The case for anthocyanin consumption
to promote human health: A review. *Compr. Rev. Food Sci. Food Saf.* **12**, 483–508 (2013).
72. Liu, C., Yao, X., Li, G., Huang, L. & Xie, Z. Transcriptomic profiling of purple broccoli reveals
light-induced anthocyanin biosynthetic signaling and structural genes. *PeerJ* **8**, e8870
(2020).
73. Kim, C. Y. *et al.* Emergence of a proton exchange-based isomerization and lactonization
mechanism in the plant coumarin synthase COSY. *Nat. Commun.* **14**, 597 (2023).
74. Vanholme, R. *et al.* COSY catalyses trans–cis isomerization and lactonization in the
biosynthesis of coumarins. *Nat. Plants* **5**, 1066–1075 (2019).
75. Rani, S. H., Krishna, T. H. A., Saha, S., Negi, A. S. & Rajasekharan, R. Defective in
Cuticular Ridges (*DCR*) of *Arabidopsis thaliana*, a gene associated with surface cutin
formation, encodes a soluble diacylglycerol acyltransferase*. *J. Biol. Chem.* **285**, 38337–
38347 (2010).
76. Agerbirk, N. & Olsen, C. E. Glucosinolate structures in evolution. *Phytochemistry* **77**, 16–45
(2012).
77. Biegańska-Marecik, R., Radziejewska-Kubzdela, E. & Marecik, R. Characterization of
phenolics, glucosinolates and antioxidant activity of beverages based on apple juice with
addition of frozen and freeze-dried curly kale leaves (*Brassica oleracea* L. var. *acephala* L.).
Food Chem. **230**, 271–280 (2017).
78. Ltd, wflpublisher. Comparative study on functional components, antioxidant activity and
color parameters of selected colored leafy vegetables as affected by photoperiods.
wflpublisher.com https://www.wflpublisher.com/Abstract/2605.
79. Park, C. H. *et al.* Effects of light-emitting diodes on the accumulation of glucosinolates and
phenolic compounds in sprouting canola (*Brassica napus L.*). *Foods* **8**, 76 (2019).

80. Zhou, B. *et al.* Effects of light intensity on the biosynthesis of glucosinolate in Chinese cabbage plantlets. *Sci. Hortic.* **316**, 112036 (2023).

81. Eggersdorfer, M. & Wyss, A. Carotenoids in human nutrition and health. *Arch. Biochem. Biophys.* **652**, 18–26 (2018).

82. Flores, M., Hernández-Adasme, C., Guevara, M. J. & Escalona, V. H. Effect of different light intensities on agronomic characteristics and antioxidant compounds of Brassicaceae microgreens in a vertical farm system. *Front. Sustain. Food Syst.* **8**, (2024).

83. Simkin, A. J., Zhu, C., Kuntz, M. & Sandmann, G. Light-dark regulation of carotenoid biosynthesis in pepper (*Capsicum annuum*) leaves. *J. Plant Physiol.* **160**, 439–443 (2003).

84. Pizarro, L. & Stange, C. Light-dependent regulation of carotenoid biosynthesis in plants. *Cienc. E Investig. Agrar.* **36**, 143–162 (2009).

85. Lisiewska, Z., Kmiecik, W. & Korus, A. The amino acid composition of kale (*Brassica oleracea* L. var. *acephala*), fresh and after culinary and technological processing. *Food Chem.* **108**, 642–648 (2008).

86. Okumoto, S., Funck, D., Trovato, M. & Forlani, G. Editorial: Amino acids of the glutamate family: functions beyond primary metabolism. *Front. Plant Sci.* **7**, (2016).

87. Kavi Kishor, P. B. & Sreenivasulu, N. Is proline accumulation per se correlated with stress tolerance or is proline homeostasis a more critical issue? *Plant Cell Environ.* **37**, 300–311 (2014).

88. Galili, G. & Amir, R. Fortifying plants with the essential amino acids lysine and methionine to improve nutritional quality. *Plant Biotechnol. J.* **11**, 211–222 (2013).

89. Yang, Q., Zhao, D. & Liu, Q. Connections between amino acid metabolisms in plants: lysine as an example. *Front. Plant Sci.* **11**, (2020).

90. Hall, C. J. *et al.* Differential lysine-mediated allosteric regulation of plant dihydrodipicolinate synthase isoforms. *FEBS J.* **288**, 4973–4986 (2021).

91. Van Bochaute, P., Novoa, A., Ballet, S., Rognes, S. E. & Angenon, G. Regulatory mechanisms after short- and long-term perturbed lysine biosynthesis in the aspartate pathway: the need for isogenes in Arabidopsis thaliana. *Physiol. Plant.* **149**, 449–460 (2013).

92. Stein, O. & Granot, D. An overview of sucrose synthases in plants. *Front. Plant Sci.* **10**, (2019).

93. Hwang, G. *et al.* Trehalose-6-phosphate signaling regulates thermoresponsive hypocotyl growth in Arabidopsis thaliana. *EMBO Rep.* **20**, e47828 (2019).

94. Avidan, O. *et al.* Direct and indirect responses of the Arabidopsis transcriptome to an induced increase in trehalose 6-phosphate. *Plant Physiol.* **196**, 409–431 (2024).

95. Sharma, N., Chaudhary, C. & Khurana, P. Role of myo-inositol during skotomorphogenesis in Arabidopsis. *Sci. Rep.* **10**, 17329 (2020).

96. Godoy Herz, M. A. *et al.* Light regulates plant alternative splicing through the control of transcriptional elongation. *Mol. Cell* **73**, 1066-1074.e3 (2019).

FIGURES

Figure 1. Effect of light intensities and spectra on kale growth and development.

A. Pictures of K3 and K9 plants at 35 dpi grown under different light intensities (75, 125, and 175 PPFD).

B. K3 and K9 weight at 35 dpi (n=6), leaf area at 22 dpi (n=15) and K9 anthocyanin content at
35 dpi (n=4), at different light intensities (75, 125, and 175 PPFD).
C. Pictures of K3 and K9 plants at different light spectra (low, balanced, high R/B and R/FR)
grown for 35 dpi.
D. K3 and K9 weight at 35 dpi (n=6), leaf area at 22 dpi (n=28) and K9 anthocyanin content at
35 dpi (n=4), at different light spectra (low, balanced, high R/B and R/FR).
Letters shows significant differences using a one-way ANOVA and Tukey's post-hoc test
(adjusted p-value <0.05).
1019

Figure 2. Effect of Light Intensity and spectra on Photosynthetic Parameters Measured
with MultispeQ.
A. Light intensity effect on relative chlorophyll content (SPAD) and photosynthetic parameters
(Phi2, PhiNO, PhiNPQ) were measured with MultispeQ at 35 dpi (n=10).
B. Light spectra effect on relative chlorophyll content (SPAD) and photosynthetic parameters
(Phi2, PhiNO, PhiNPQ) were measured with MultispeQ at 35 dpi (n=10).
Letters show significant differences using a one-way ANOVA and Tukey's post-hoc test
(adjusted p-value < 0.05).
1028

1029   **Figure 3. Effect of light intensity treatment on the metabolite composition of Kale**
1030   **cultivars.**
1031   GC-MS analysis of primary metabolites in kale cultivars K3 and K9 under different light intensity
1032   75, 125 and 175 PPFD and at ZT11 / ZT23 (n=4). Different letters indicate significant
1033   differences using a one-way ANOVA and Tukey's post-hoc test (adjusted p-value < 0.05).
1034

1035   **Figure 4. Effect of light spectra treatment on the metabolite composition of Kale**
1036   **cultivars.**
1037   GC-MS analysis of primary metabolites in kale cultivars K3 and K9 under different light spectra
1038   (low, balanced and high R/B) and at ZT11 / ZT23 (n=4). Different letters indicate significant
1039   differences using a one-way ANOVA and Tukey's post-hoc test (adjusted p-value < 0.05).
1040

1041   **Figure 5. Effect of light spectra treatment on the metabolite composition of Kale**
1042   **cultivars.**
1043   GC-MS analysis of primary metabolites in kale cultivars K3 and K9 under different light spectra
1044   (low, balanced and high R/FR) and at ZT11 / ZT23 (n=4). Different letters indicate significant
1045   differences using a one-way ANOVA and Tukey's post-hoc test (adjusted p-value < 0.05).
1046

Figure 6. Analysis of significantly changing proteins with changing light intensity and
spectral composition.
A-C. Number of proteins with a significant change in abundance with increase in light intensity
from 75 PPFD (A), from low R/B (B) and from low R/FR (C).
D. Dot plot representation of enriched GO terms for biological process of significantly changing
proteins (q-value < 0.05; Log2FC > 0.58 or < -0.58) that are up- or down-regulated with
increasing light intensity (top panel), increasing R/B ratio (middle panel), or increasing R/FR
ratio (bottom panel) at ZT23 (left) or ZT11 (right). Dot size indicates the gene ratio (number of
proteins in the changing in the condition/total quantified proteins in the category in the study).
Dot colour represents the Log10 (p-value). Only GO terms with p-value <0.01 and a total count
< 500 are included.
1058

Figure 7. Association network analysis of significantly changing proteins with increasing light intensity
An association network analysis was performed using STRING-DB to depict significantly changing proteins in K3 and K9 kale under increasing light intensity at ZT23 and ZT11. Edge thickness indicates strength of connection between the nodes. Minimum edge threshold was set to 0.95. Protein nodes are labelled either by primary gene name or Arabidopsis gene identifier (AGI). Outer circle surrounding each node represents the standardized relative Log2FC of the indicated significantly changing protein with an increase from 75 PPFD to either 125 PPFD or 175 PPFD in K3 or K9 as indicated by the legend. The scale of blue to yellow indicates the relative decrease or increase in abundance, respectively from 75 PPFD to either 125 PPFD or 175 PPFD. Node groupings are indicated by a grey circle representing proteins involved in the same biological process.

Figure 8. Association network analysis of significantly changing proteins with changing light quality
An association network analysis was performed using STRING-DB to depict significantly changing proteins in K3 and K9 kale under increasing R/B spectral ratio (A) or R/FR ratio (B) at ZT23 and ZT11. Edge thickness indicates the strength of connection between the nodes. Minimum edge threshold was set to 0.95. Protein nodes are labelled either by primary gene name or Arabidopsis gene identifier (AGI). Outer circle surrounding each node represents the standardized relative Log2FC of the indicated significantly changing protein with an increase from low R/B (A) or low R/FR (B) to balanced R/B (R = B) or high R/B (R > B) (A) or balanced R/FR (R = FR) or high R/FR (R > FR) (B) in K3 or K9 at ZT23 or ZT11 as indicated by the legend. The scale of blue to yellow indicates the relative decrease or increase in abundance, respectively. Node groupings are indicated by a grey circle representing proteins involved in the same biological process.

SUPPLEMENTAL DATA

Table S1. Gas chromatography metabolite data for Intensity experiments

Table S2. Gas chromatography metabolite data for Spectra experiments

Table S3. Liquid chromatography proteomic data

Table S4. Pathway enrichment analysis using the Plant Metabolic Network (PMN;

https://plantcyc.org) for Intensity experiments

Table S5. Pathway enrichment analysis using the Plant Metabolic Network (PMN;

https://plantcyc.org) for R/B Spectra experiments

Table S6. Pathway enrichment analysis using the Plant Metabolic Network (PMN;

https://plantcyc.org) for R/FR Spectra experiments

Supplemental Figure 1. Light spectra of the different light treatments
A. Light spectra of different light intensity (75, 125 and 175 PPFD).
B. Light spectra of the different light ratios (low R/B, balanced and high R/FR). Light intensity is 125 PPFD for all spectra treatment.

Supplemental Figure 2: Effect of Light Intensity and spectra on Photosynthetic
Parameters Measured with MultispeQ.
A. Light intensity effect on photosynthetic parameters (LEF, gH+, ECSt, vH+) were measured
with MultispeQ at 35 dpi (n=10).
B. Light spectra effect on photosynthetic parameters (LEF, gH+, ECSt, vH+) were measured
with MultispeQ at 35 dpi (n=10).
Letters show significant differences using a one-way ANOVA and Tukey's post-hoc test
(adjusted p-value < 0.05).
1110

Supplemental Figure 3: GC-MS analysis of kale cultivars K3 and K9 under different light
intensities (75 PPFD, 125 PPFD and 175 PPFD) and at different time of the day (ZT11 and
ZT23).
A. Heatmap of relative cultivars metabolite changes at ZT11 and ZT23.
B. Eucledian distance clustered heatmap of relative cultivars metabolite changes.
C. Heatmap of relative diel metabolite changes in K3 and K9.
D.Eucledian distance clustered heatmap of relative diel metabolite changes in K3 and K9.
Heatmaps were generated using the R package *Pheatmaps*. Scale represents Log2 fold-
change (FC); n=4.
1120

Supplemental Figure 4: GC-MS analysis of kale cultivars K3 and K9 under different light
spectra (low, balanced, and high R/B ratios) and at different time of the day (ZT11 and
ZT23).
A. Heatmap of relative cultivars metabolite changes at ZT11 and ZT23.
B. Eucledian distance clustered heatmap of relative cultivars metabolite changes.
C. Heatmap of relative diel metabolite changes in K3 and K9.
D. Eucledian distance clustered heatmap of relative diel metabolite changes in K3 and K9.
Heatmaps were generated using the R package *Pheatmaps*. Scale represents Log2 fold-
change (FC); n=4.
1130

Supplemental Figure 5: GC-MS analysis of kale cultivars K3 and K9 under different light
spectra (low, balanced, and high R/FR ratios) and at different time of the day (ZT11 and
ZT23).
A. Heatmap of relative cultivars metabolite changes at ZT11 and ZT23.
B. Eucledian distance clustered heatmap of relative cultivars metabolite changes.
C. Heatmap of relative diel metabolite changes in K3 and K9.
D. Eucledian distance clustered heatmap of relative diel metabolite changes in K3 and K9.
Heatmaps were generated using the R package *Pheatmaps*. Scale represents Log2 fold-
change (FC); n=4.

Figure 1. Effect of light intensities and spectra on kale growth and development.
A. Pictures of K3 and K9 plants at 35 dpi grown under different light intensities (75, 125, and 175 PPFD).
B. K3 and K9 weight at 35 dpi (n=6), leaf area at 22 dpi (n=15) and K9 anthocyanin content at 35 dpi (n=4), at different light intensities (75, 125, and 175 PPFD).
C. Pictures of K3 and K9 plants at different light spectra (low, balanced, high R/B and R/FR) grown for 35 dpi.
D. K3 and K9 weight at 35 dpi (n=6), leaf area at 22 dpi (n=28) and K9 anthocyanin content at 35 dpi (n=4), at different light spectra (low, balanced, high R/B and R/FR).
Letters shows significant differences using a one-way ANOVA and Tukey's post-hoc test (adjusted p-value <0.05).

Figure 2. Effect of Light Intensity and spectra on Photosynthetic Parameters Measured with MultispeQ.

A.Light intensity effect on relative chlorophyll content (SPAD) and photosynthetic parameters (Phi2, PhiNO, PhiNPQ) were measured with MultispeQ at 35 dpi (n=10).

B.Light spectra effect on relative chlorophyll content (SPAD) and photosynthetic parameters (Phi2, PhiNO, PhiNPQ) were measured with MultispeQ at 35 dpi (n=10).

Letters show significant differences using a one-way ANOVA and Tukey's post-hoc test (adjusted p-value < 0.05).

Figure 3. Effect of light intensity treatment on the metabolite composition of Kale cultivars.

GC-MS analysis of primary metabolites in kale cultivars K3 and K9 under different light intensity 75, 125 and 175 PPFD and at ZT11 / ZT23 (n=4). Different letters indicate significant differences using a one-way ANOVA and Tukey's post-hoc test (adjusted p-value < 0.05).

Figure 4. Effect of light spectra treatment on the metabolite composition of Kale cultivars. GC-MS analysis of primary metabolites in kale cultivars K3 and K9 under different light spectra (low, balanced and high R/B) and at ZT11 / ZT23 (n=4). Different letters indicate significant differences using a one-way ANOVA and Tukey's post-hoc test (adjusted p-value < 0.05).

Figure 5. Effect of light spectra treatment on the metabolite composition of Kale cultivars. GC-MS analysis of primary metabolites in kale cultivars K3 and K9 under different light spectra (low, balanced and high R/FR) and at ZT11 / ZT23 (n=4). Different letters indicate significant differences using a one-way ANOVA and Tukey's post-hoc test (adjusted p-value < 0.05).

Figure 6. Analysis of significantly changing proteins with changing light intensity and spectral composition.

A-C. Number of proteins with a significant change in abundance with increase in light intensity from 75 PPFD (A), from low R/B (B) and from low R/FR (C).

D. Dot plot representation of enriched GO terms for biological process of significantly changing proteins (q-value < 0.05; Log2FC > 0.58 or < -0.58) that are up- or down-regulated with increasing light intensity (top panel), increasing R/B ratio (middle panel), or increasing R/FR ratio (bottom panel) at ZT23 (left) or ZT11 (right). Dot size indicates the gene ratio (number of proteins in the changing in the condition/total quantified proteins in the category in the study). Dot colour represents the Log10 (p-value). Only GO terms with p-value <0.01 and a total count < 500 are included.

Figure 7. Association network analysis of significantly changing proteins with increasing light intensity.
An association network analysis was performed using STRING-DB to depict significantly changing proteins in K3 and K9 kale under increasing light intensity at ZT23 and ZT11. Edge thickness indicates strength of connection between the nodes. Minimum edge threshold was set to 0.95. Protein nodes are labelled either by primary gene name or Arabidopsis gene identifier (AGI). Outer circle surrounding each node represents the standardized relative Log2FC of the indicated significantly changing protein with an increase from 75 PPFD to either 125 PPFD or 175 PPFD in K3 or K9 as indicated by the legend. The scale of blue to yellow indicates the relative decrease or increase in abundance, respectively from 75 PPFD to either 125 PPFD or 175 PPFD. Node groupings are indicated by a grey circle representing proteins involved in the same biological process.

Figure 8. Association network analysis of significantly changing proteins with changing light quality.
An association network analysis was performed using STRING-DB to depict significantly changing proteins in K3 and K9 kale under increasing R/B spectral ratio (A) or R/FR ratio (B) at ZT23 and ZT11. Edge thickness indicates the strength of connection between the nodes. Minimum edge threshold was set to 0.95. Protein nodes are labelled either by primary gene name or Arabidopsis gene identifier (AGI). Outer circle surrounding each node represents the standardized relative Log2FC of the indicated significantly changing protein with an increase from low R/B (A) or low R/FR (B) to balanced R/B (R = B) or high R/B (R > B) (A) or balanced R/FR (R = FR) or high R/FR (R > FR) (B) in K3 or K9 at ZT23 or ZT11 as indicated by the legend. The scale of blue to yellow indicates the relative decrease or increase in abundance, respectively. Node groupings are indicated by a grey circle representing proteins involved in the same biological process.

Supplemental Figure 1. Light spectra of the different light treatments.

A. Light spectra of different light intensity (75, 125 and 175 PPFD).
B. Light spectra of the different light ratios (low R/B, balanced and high R/FR). Light intensity is 125 PPFD for all spectra treatment.

Supplemental Figure 2: Effect of Light Intensity and spectra on Photosynthetic Parameters Measured with MultispeQ.

A. Light intensity effect on photosynthetic parameters (LEF, gH+, ECSt, vH+) were measured with MultispeQ at 35 dpi (n=10).

B. Light spectra effect on photosynthetic parameters (LEF, gH+, ECSt, vH+) were measured with MultispeQ at 35 dpi (n=10).

Letters show significant differences using a one-way ANOVA and Tukey's post-hoc test (adjusted p-value < 0.05).

A.

B.

C.

D.

Supplemental Figure 3: GC-MS analysis of kale cultivars K3 and K9 under different light intensities (75 PPFD, 125 PPFD and 175 PPFD) and at different time of the day (ZT11 and ZT23).

A. Heatmap of relative cultivars metabolite changes at ZT11 and ZT23.
B. Eucledian distance clustered heatmap of relative cultivars metabolite changes.
C. Heatmap of relative diel metabolite changes in K3 and K9.
D. Eucledian distance clustered heatmap of relative diel metabolite changes in K3 and K9. Heatmaps were generated using the R package *Pheatmaps*. Scale represents Log2 fold-change (FC); n=4.

Supplemental Figure 4: GC-MS analysis of kale cultivars K3 and K9 under different light spectra (low, balanced, and high R/B ratios) and at different time of the day (ZT11 and ZT23).

A. Heatmap of relative cultivars metabolite changes at ZT11 and ZT23.
B. Eucledian distance clustered heatmap of relative cultivars metabolite changes.
C. Heatmap of relative diel metabolite changes in K3 and K9.
D. Eucledian distance clustered heatmap of relative diel metabolite changes in K3 and K9. Heatmaps were generated using the R package *Pheatmaps*. Scale represents Log2 fold-change (FC); n=4.

Supplemental Figure 5: GC-MS analysis of kale cultivars K3 and K9 under different light spectra (low, balanced, and high R/FR ratios) and at different time of the day (ZT11 and ZT23).

A. Heatmap of relative cultivars metabolite changes at ZT11 and ZT23.
B. Eucledian distance clustered heatmap of relative cultivars metabolite changes.
C. Heatmap of relative diel metabolite changes in K3 and K9.
D. Eucledian distance clustered heatmap of relative diel metabolite changes in K3 and K9. Heatmaps were generated using the R package *Pheatmaps*. Scale represents Log2 fold-change (FC); n=4.