Cardiolipin deficiency disrupts electron transport chain to drive steatohepatitis

Marisa J. Brothwell,[1,2] Guoshen Cao,[1,3] J. Alan Maschek,[1,2,4] Annelise M. Poss,[1,2] Alek D.

Peterlin,[1,2] Liping Wang,[1,2] Talia B. Baker,[5,6] Justin L. Shahtout,[1,7] Piyarat Siripoksup,[1,7] Quentinn

J. Pearce,[4] Jordan M. Johnson,[1,2] Fabian M. Finger,[8,9] Alexandre Prola,[10] Sarah A. Pellizzari,[3,11]

Gillian L. Hale,[5,11] Allison M. Manuel,[4] Shinya Watanabe,[1,2] Edwin R. Miranda,[1,2,12] Kajsa E.

Affolter,[5,10] Trevor S. Tippetts,[1,2] Linda S. Nikolova,[13] Ran Hee Choi,[1,2,12] Stephen T. Decker,[1,2,12]

Mallikarjun Patil,[1,2,12] J. Leon Catrow,[4] William L. Holland,[1,2,3,12] Sara M. Nowinski,[14] Daniel S.

Lark,[15,16] Kelsey H. Fisher-Wellman,[17] Patrice N. Mimche,[18] Kimberley J. Evason,[5,11] James E.

Cox,[1,3,4] Scott A. Summers,[1,2,3,5,12] Zach Gerhart-Hines,[8,9] Katsuhiko Funai[1,2,3,5,7,12]*

[1]Diabetes & Metabolism Research Center; University of Utah; Salt Lake City, UT; USA.

[2]Department of Nutrition and Integrative Physiology; University of Utah; Salt Lake City, UT;

USA.

[3]Department of Biochemistry; University of Utah; Salt Lake City, UT; USA.

[4]Metabolomics Core Research Facility; University of Utah; Salt Lake City, UT; USA.

[5]Huntsman Cancer Institute; University of Utah, Salt Lake City, UT; USA.

[6]Division of Transplantation and Advanced Hepatobiliary Surgery, Department of Surgery;

University of Utah; Salt Lake City, UT; USA.

[7]Department of Physical Therapy and Athletic Training; University of Utah; Salt Lake City, UT;

USA.

[8]Novo Nordisk Foundation Center for Basic Metabolic Research; University of Copenhagen;

Copenhagen; DK.

[9]Center for Adipocyte Signaling (ADIPOSIGN); University of Southern Denmark; Odense; DK.

[10]Laboratory of Fundamental and Applied Bioenergetics; University of Grenoble Alpes, Inserm

U1055; Grenoble; FR.

[11]Department of Pathology; University of Utah; Salt Lake City, UT; USA.

[12]Molecular Medicine Program; University of Utah; Salt Lake City, UT; USA.

[13]Electron Microscopy Core Facility; University of Utah; Salt Lake City, UT; USA.

[14]Department of Metabolism and Nutritional Programming; Van Andel Institute; Grand Rapids,

MI; USA.

[15]College of Health and Human Sciences; Colorado State University; Fort Collins, CO; USA.

[16]Columbine Health Systems Center for Healthy Aging; Colorado State University; Fort Collins,

CO; USA

[17]Department of Cancer Biology, Wake Forest University School of Medicine; Atrium Health

Wake Forest Baptist Comprehensive Cancer Center; Winston-Salem, NC; USA.

[18]Departments of Dermatology and Medicine; Division of Gastroenterology and Hepatology,

Indiana University School of Medicine; Indianapolis, IN; USA.

*Lead Contact

Correspondence:

Katsuhiko Funai, Ph.D.

kfunai@health.utah.edu

Abstract

Metabolic dysfunction-associated steatotic liver disease (MASLD) is a progressive disorder marked by lipid accumulation, leading to metabolic dysfunction-associated steatohepatitis (MASH). A key feature of the transition to MASH involves oxidative stress resulting from defects in mitochondrial oxidative phosphorylation (OXPHOS). Here, we show that pathological alterations in the lipid composition of the inner mitochondrial membrane (IMM) directly instigate electron transfer inefficiency to promote oxidative stress. Specifically, mitochondrial cardiolipin (CL) was downregulated with MASLD/MASH in humans and in mice. Hepatocyte-specific CL synthase knockout (CLS-LKO) led to spontaneous and robust MASH with extensive steatotic and fibrotic phenotype. Loss of CL paradoxically increased mitochondrial respiratory capacity but also reduced the formation of $I+III_2+IV$ respiratory supercomplex, promoted electron leak primarily at sites III_{QO} and II_F of the electron transport chain, and disrupted the propensity of coenzyme Q (CoQ) to become reduced. Thus, low mitochondrial CL disrupts electron transport chain to promote oxidative stress and contributes to pathogenesis of MASH.

Main

Metabolic dysfunction-associated steatotic liver disease (MASLD) is a growing global health

concern with an increasing prevalence that parallels the rise in obesity.[1] In the United States,

annual medical costs related to MASLD exceed $103 billion.[2] A large portion of patients with

MASLD only exhibit steatosis, a silent and relatively benign early stage characterized by lipid

accumulation in hepatocytes without hepatocellular inflammation.[3] Steatosis can then progress

to metabolic dysfunction-associated steatohepatitis (MASH), determined by hepatocyte injury

and tissue fibrosis.[4] MASH is the last stage of MASLD that may be reversible, making

intervention at this stage particularly important.[3,5] Although extensive clinical and basic research

have been conducted in this field, the underlying mechanisms by which fatty liver transitions to

MASH remain poorly understood.[6-8]

A defect in mitochondrial function is considered one of the hallmarks of MASLD progression in

both mice and humans.[9-12] MASLD is initially associated with an increase in mitochondrial

respiratory capacity, followed by a subsequent impairment in oxidative phosphorylation

(OXPHOS), and increased production of mitochondrial reactive oxygen species (ROS).[11,13]

Mitochondrial ROS is thought to be caused by an inefficient electron transport chain (ETC) that

increases the propensity for electron leak. However, the mechanisms by which mitochondrial

electron leak promotes MASLD are unknown.

Cardiolipin (CL) is a phospholipid with four acyl chains conjugated to two phosphatidylglycerol

moieties linked by another glycerol molecule.[14] CL resides almost exclusively in the inner

mitochondrial membrane (IMM), constituting approximately 15–20% of the mitochondrial

phospholipids.[15] CL is synthesized by the condensation of phosphatidylglycerol (PG) and

cytidine diphosphate-diacylglycerol (CDP-DAG) at the IMM via the enzyme cardiolipin synthase

(CLS).[16,17] Structural studies indicate that CL is essential for the activities of OXPHOS

enzymes.[18-22] In non-hepatocytes, decreased CL leads to compromised oxidative capacity,[23,24]

impaired membrane potential,[25] and altered cristae morphology.[26] In particular, low CL is

associated with increased H_2O_2 production.[27,28]

In this manuscript, we set out to examine the changes in liver mitochondrial lipidome induced by

MASH. In both mice and in humans, mitochondrial CL was downregulated in livers with

MASLD/MASH compared to healthy controls. We then performed a targeted deletion of CLS in

hepatocytes and studied its effects on liver, mitochondrial bioenergetics, and potential

mechanisms that drive these changes.

Results

Mitochondrial cardiolipin levels are decreased in MASLD/MASH

Previous research from our lab in non-hepatocytes indicated that mitochondrial phospholipid

composition affects OXPHOS electron transfer efficiency to alter electron leak.[15,29,30] MASLD

has been shown to alter the total cellular lipidome in liver.[31] However, MASLD may also

influence mitochondrial content in the hepatocytes, making it difficult to discern whether these

are changes in the lipid composition of mitochondrial membranes and/or changes in cellular

mitochondrial density. Thus, we performed liquid chromatography-tandem mass spectrometry

(LC-MS/MS) lipidomics specifically on mitochondria isolated from human liver samples with or

without MASH. Samples from patients undergoing liver transplant or resection due to end-stage

MASH (Figure 1A, and Figure 1– table supplement 1) were compared to samples from patients

without MASH, undergoing resection for benign liver tumors or metastases (Figure 1B).

Strikingly, mitochondrial CL levels were reduced, primarily due to a decrease in tetralinoleoyl CL

(Figure 1C), commonly thought as the mature and functional form of CL. Additionally, levels of

mitochondrial PG, an essential precursor to CL production, were also reduced, suggesting a

disruption in CL biosynthesis (Figure 1D-E).

We next performed lipidomic analyses on four distinct mouse models of MASLD/MASH (Figure

1F-Q). These included: 1) mice given a Western high-fat diet (HFD, Envigo TD.88137) or

standard chow diet for 16 weeks (Figure 1F), 2) ob/ob mice or their wildtype littermates at 20

weeks of age (Figure 1G), 3) mice given the Gubra Amylin NASH diet for 30 weeks (GAN,

Research Diets D09100310) or standard chow (Figure 1H), 4) mice injected with carbon

tetrachloride (CCl_4) or vehicle (corn oil) for 6 weeks (Figure 1I). Importantly, none of these

interventions appear to alter the protein abundances of OXPHOS subunits or citrate synthase

(Figure 1J-M), suggesting that these interventions did not alter mitochondrial density in

hepatocytes. Nevertheless, we performed all mitochondrial lipidomic analyses by quantifying

lipids per mg of mitochondrial proteins. Each intervention appeared to alter different subsets of

mitochondrial lipid classes (Figure 1N-Q, and Figure 1—figure supplement 1 and Figure 1—

figure supplement 2), as seen with our previous studies in skeletal muscle and brown adipose

tissues.[29,30] We take these observations to mean that most physiological interventions induce

multiple systemic and local responses that are not mechanistically directly related to the

phenotype of interest (e.g., cold exposure or exercise can increase food intake, obesity could

affect locomotion and insulation, etc.). Although several phospholipid classes were altered

among the four models, strikingly, mitochondrial CL was reduced in all four MASLD/MASH

models (Figure 1N-R). Furthermore, PG was significantly increased in all MASLD/MASH models

(Figure 1N-R). These changes coincided with decreased transcript levels for CLS (Figure 1S-V).

These observations further suggest that an insult in CL synthesis may be a key factor to

disrupting mitochondrial function in MASLD/MASH.

Hepatocyte-specific deletion of cardiolipin synthase promotes MASH

CL is thought to be exclusively synthesized in the IMM where CLS is localized. To study the role

of CL in hepatocytes, we generated mice with hepatocyte-specific knockout of CLS (CLS-LKO

for *CLS liver knockout*, driven by albumin-Cre) (Figure 2A,B), which successfully decreased

mitochondrial CL levels (Figure 2C, and Figure 2—figure supplement 1). Consistent with our

previous studies in non-hepatocytes, CLS deletion does not completely reduce CL levels to

zero, suggesting that CL generated in other tissues may be imported. Our results showed that

decreased levels of CL did not significantly impact body weight or composition (Figure 2D,E) but

resulted in significantly less liver mass (Figure 2F).

We sought to further characterize livers from control and CLS-LKO mice. Histological analyses

revealed that CLS deletion was sufficient to promote steatosis (Figure 2G) and fibrosis (Figure

2H) in standard chow-fed and high-fat fed conditions (Figure 2—figure supplement 2A,B). To

more comprehensively describe the effects of loss of hepatic CLS on gene expression, we

performed RNA sequencing on these livers. CLS deletion increased the expression of 713

genes and decreased 1026 genes (Figure 2—figure supplement 2C). Pathway analyses

revealed that many of the signature changes that occur with MASLD/MASH also occurred with

CLS deletion (Figure 2I, and Figure 2—figure supplement 2D). This MASLD/MASH phenotype

in our CLS knockout model was further confirmed with an elevation of the liver enzymes AST

and ALT (Figure 2J,K) as well as increased mRNA levels of inflammatory markers (Figure 2L).

We then proceeded to confirm these data by further phenotyping liver tissues from control and

CLS-LKO mice.

In steatohepatitis, immune cell populations in the liver become altered to activate pathological

immune response.[32] Flow cytometry on livers from control and CLS-LKO mice indicated that the

loss of CL promotes a robust classic immune response found in MASH (Figure 2M, and Figure

2—figure supplement 2E). cDC2 cells are a broad subset of dendritic cells with specific surface

markers (e.g., CD11b, CD172a) that allow them to be distinguished from other dendritic cell

populations.[33] This broad population of dendritic cells was not different between control and

CLS-LKO mice (Figure 2N). Notably, there was a marked reduction in the Kupffer cell

population (Figure 2O) - traditionally involved in maintaining liver homeostasis whose

dysfunction can lead to dysregulated immune response.[34] This reduction appears to be

counterbalanced by a concomitant increase in Ly6C[hi] population, which are known to typically

go on to become inflammatory monocytes (Figure 2M,P). The replacement of Kupffer cells with

other inflammatory cell populations suggests a shift towards a more pro-inflammatory

environment, which may exacerbate liver injury and promote fibrosis. Nonetheless, the MHC-II

cell population and neutrophils were not increased (Figure 2Q,R) with neutrophils actually

decreased (Figure 2R). The cDC1 cell population was not different, which is traditionally

elevated in response to cytotoxic T cells and might not be directly related to liver fibrosis.[35]

Together, these findings suggest that even on a chow diet, CLS-deficient livers exhibit

inflammatory cell infiltration, a hallmark often associated with early signs of MASH.

CLS deletion promotes fatty liver but increases mitochondrial respiratory capacity

Hepatocyte lipid accumulation may suggest defects in substrate handling, which is often

manifested in systemic substrate handling. Indeed, CLS deletion modestly reduced glucose or

pyruvate handling, even in chow-fed conditions (Figure 3A-D). Lipid accumulation in

hepatocytes can occur due to an increase in lipogenesis, a decrease in VLDL secretion, or a

decrease in β-oxidation. However, mRNA levels for lipogenesis genes trended lower (not

higher), and mostly unchanged for VLDL secretion or β-oxidation (Figure 3E, and Figure 3—

figure supplement 1A). Circulating triglycerides were not lower in CLS-LKO mice compared to

control mice (Figure 3—figure supplement 1B).

MASLD is known to be associated with reduced mitochondrial oxidative capacity, and such an

effect may also occur with CL deficiency to induce lipid accumulation. Indeed, mRNA levels of

several genes in the ETC were downregulated with CLS deletion, particularly those associated

with structural components of the ETC complexes and the electron carrier CoQ (Figure 3F).

Given that CL is located in the IMM where it binds to enzymes involved in OXPHOS,[36-39] we

reasoned that the loss of CL could reduce mitochondrial oxidative capacity to promote steatosis.

Consistent with subcellular localization of CL, CLS deletion resulted in mitochondria with

disorganized membrane structures and poorly developed cristae (Figure 3G). However,

mitochondrial density quantified with western blots for respiratory complex subunits and citrate

synthase (Figure 3H), as well as mtDNA/nucDNA (Figure 3I), showed no differences in livers

from control and CLS-LKO mice. We thus speculated that CL lowers respiratory capacity not by

reducing the total number of mitochondria or OXPHOS respirasomes, but by reducing the

activity of respiratory enzymes. To our surprise, CLS deletion increased, rather than decreased,

mitochondrial respiration (JO_2), as measured by high-resolution Oroboros respirometry (Figure

3J), using both with Krebs cycle substrates (Figure 3K) as well as fatty acyl substrates (Figure

3L), to a similar degree. Importantly, these changes occurred in the absence of differences in

OXPHOS subunit abundance per unit of mitochondria (Figure 3M), ruling out the possibility that

changes in the abundance of respiratory enzymes are contributing to the change in respiration.

A caveat to these findings is that CLS deletion promotes reduction in respiratory capacity after

HFD-feeding (Figure 3—figure supplement 1C,D) However, CLS-LKO mice are steatotic in

standard chow-fed condition, indicating that reduced mitochondrial fatty acid oxidation cannot

be the cause of steatosis at baseline. The initial increase in respiration followed by its

subsequent decrease is reminiscent of what is thought to occur with liver's mitochondrial

respiration over the course of MASLD progression.[40]

High-resolution respirometry experiments were performed in isolated mitochondria from

hepatocytes by providing exogenous supraphysiological concentrations of substrates. While

these assays provide robust measurements of respiratory capacity (the potential of

mitochondria), they do not necessarily reflect their endogenous activity. To address this point,

we performed stable isotope tracing experiments using uniformly labeled ^{13}C-palmitate (Figure

4A)[41] in murine hepa1-6 cells without (scrambled shRNA or shSC) or with CLS knockdown

(shRNA targeting CLS or shCLS). Surprisingly, but consistent with the JO_2 data, CLS deletion

increased, not decreased, the incorporation of palmitate into TCA intermediates (Figure 4B-D).

We also performed a similar tracing experiment using uniformly labeled ^{13}C-glucose (Figure

4E-J, and Figure 4—figure supplement 1A-E) and observed increased labeling towards

pyruvate (Figure 4E,F), reduced labeling towards lactate and alanine (Figure 4G, and Figure

4—figure supplement 1D), and normal labeling towards TCA intermediates except for reduced

labeling towards succinate (Figure 4H-J, and Figure 4—figure supplement 1B,C,E). Overall,

despite the altered substrate incorporation, a decrease in TCA flux does not appear to account

for the steatotic phenotype observed with CLS deletion.

Low hepatic CL paradoxically increases mitochondrial respiratory capacity and promotes

electron leak

Oxidative stress is thought to play a critical role in the transition from MASLD to MASH, wherein

sustained metabolic insult leads to hepatocellular injury and collagen deposition resulting in

fibrosis.[7] CLS deletion promotes liver fibrosis in standard chow-fed condition (Figure 5A and

Figure 2H) and in HFD-fed condition (Figure 2—figure supplement 2) that coincided with

increased mRNA levels for fibrosis (Figure 5B and 2I). Tissue fibrosis is often triggered by

apoptosis, and CLS deletion appeared to activate the caspase pathway (Figure 5C,D). How

does deletion of CLS, a mitochondrial enzyme that produces lipids for IMM, activate apoptosis?

Cytochrome c is an electron carrier that resides in IMM, which shuttles electrons between

complexes III and IV.[42] Under normal physiological conditions, cytochrome c is anchored to the

IMM by its binding to cardiolipin.[39] During the initiation of intrinsic apoptosis, CL can undergo

oxidation and redistribution from the IMM to the outer membrane space (OMM). CL oxidation

weakens its binding affinity for cytochrome c, releasing it from the IMM and into the OMM where

it signals apoptosis.[13] However, neither mitochondrial nor cytosolic cytochrome c abundance

appeared to be influenced by CLS deletion (Figure 5E,F, and Figure 5—figure supplement

1A,B).

Mitochondrial ROS has been implicated in apoptosis and fibrosis with MASLD.[43-45] Using high-

resolution fluorometry in combination with high-resolution respirometry, we quantified electron

leak from liver mitochondria with the assumption that almost all electrons that leak react with

molecular O_2 to produce O_2^-. Using recombinant superoxide dismutase, we ensure that all O_2^-

produced is converted into H_2O_2, which was quantified with the AmplexRed fluorophore.[46] There

were striking increases in mitochondrial electron leak in CLS-LKO mice compared to control

mice on both standard chow (Figure 5G) and high-fat diet (Figure 5—figure supplement 1C). It

is noteworthy that endogenous antioxidant pathways were insufficient to completely suppress

oxidative stress induced by CLS deletion (H_2O_2 emission shown in the 1^{st} and 2^{nd} bars in Figure

5G, and Figure 5—figure supplement 1C). We also confirmed that JH_2O_2/JO_2 was elevated with

CLS knockdown in mitochondria from murine hepa1-6 cell line (Figure 5—figure supplement

1D) suggesting that low CL induces oxidative stress in a cell-autonomous manner.

While unknown, CLS may possess an enzymatic activity independent of CL synthesis that may

contribute to electron leak. To more conclusively show that the loss of mitochondrial CL

contributes to oxidative stress, we supplied exogenous CL to isolated mitochondria by fusing

them with small unilamellar vesicles (SUVs) (Figure 5H).[47] Isolated mitochondria from control

and CLS-LKO mice were fused with SUVs containing either CL or phosphatidylcholine (PC)

(Figure 5I, and Figure 5—figure supplement 1E). Remarkably, reintroducing CL to mitochondria

from CLS-LKO mice reduced H_2O_2 production back to baseline, whereas PC had no effect.

Thus, loss of CL drives the increased mitochondrial leak observed with CLS deletion.

Loss of CL attenuates coenzyme Q interface with complex III

How does the loss of CL promote electron leak? We first addressed CL's influence on the

formations of respiratory supercomplexes. Respiratory supercomplexes exist in several

combinations of multimers of Complex I, III, IV, and V and are thought to form either transiently

or stably to improve electron transfer efficiency.[48,49] CL may play an essential role in the stability

of ETC supercomplexes.[50,51] Using blue native polyacrylamide gel electrophoresis, we

investigated supercomplex assembly in isolated hepatic mitochondria from control and CLS-

LKO mice (Figure 6). When probing indiscriminately for all supercomplexes, there were many

bands whose intensity were influenced by CLS deletion (Figure 6A). In particular, $I+III_2+IV$

supercomplex appeared to be substantially and consistently reduced in samples from CLS-LKO

mice compared to control. To gain a more granular and definitive picture on the influence of

CLS deletion on individual supercomplexes, we quantified supercomplexes using subunit-

specific western blotting (Figure 6B-M). Strikingly, CLS deletion substantially reduced the

abundance of $I+III_2+IV$ supercomplex when probing for UQCRSF1/complex III (Figure 6H,I),

while increasing the abundances for CIII dimer (Figure 6I) and CIV monomer (Figure 6J,K), as

well as trends for increase in CI monomer (Figure 6B-E). These findings suggest that loss of CL

reduces $I+III_2+IV$ supercomplex assembly to attenuate electron flux through these complexes,

promoting electron leak.

To better understand how low CL promotes inefficient electron transfer, we quantified electron

leak at four specific sites of the ETC: 1) quinone-binding site in complex I (I_Q), 2) flavin-

containing site in complex I (I_F), 3) succinate-dehydrogenase-associated site in complex II (II_F),

and 4) the ubiquinol oxidation site in complex (III_{Qo}) Figure 7A). Electron leak at each of these

sites can be quantified separately using substrates and inhibitors that restrict electron flow

specific to these sites (Figure 7B-E). Among complex I, III, and IV (the components of $I+III_2+IV$

supercomplex), CLS deletion robustly increased electron leak at sites II_F and III_{Qo} (Figure 7D,E).

In addition, electron leak either increased or trended to increase sites on complex I (Figure

7B,C). These findings raise a possibility that low CL may globally suppress the efficiency of the

electron transport chain to promote electron leak.

To further examine the efficiency of the electron transport chain, we studied the redox state of

CoQ, a carrier that feeds electrons from complex I and II into complex III. It is noteworthy that

CoQ, like CL, is a lipid molecule (Figure 7F), and we thought it was conceivable that CL

somehow interacts with CoQ to influence its electron transfer efficiency. Using redox mass

spectrometry, we measured CoQ levels in whole liver tissues from control and CLS-LKO mice

and found no difference in whole liver tissue CoQ levels (Figure 7—figure supplement 1A-I).

However, since CoQ may also be found outside of mitochondria, we performed CoQ redox

mass spectrometry in isolated mitochondria fractions of livers from control or CLS-LKO mice.

Indeed, oxidized CoQ levels were increased (Figure 7G, and Figure 7—figure supplement

1J,L,N) in CLS-LKO mice compared to their controls. In contrast, reduced forms of CoQ were

not lower in CLS-LKO mice compared to control mice (Figure 7H, and Figure 7—figure

supplement 1K,M,O). Our findings that the normal level of reduced CoQ yielded greater

oxidized CoQ, combined with increased electron flux (i.e., increase JO_2, Figure 3J-L) can be

interpreted to mean that there is a defect at the interface of CoQ-complex III that disrupts the

efficient transfer of electrons promoting electron leak at site III_{QO}. Conversely, it may also be

interpreted that there are defects at complex I and II where oxidized CoQ are unable to become

reduced due to lack of their ability to efficiently charge CoQ, also promoting electron leak.

To extrapolate these findings into humans, we went back to liver samples from patients

undergoing liver transplant or resection due to end-stage MASH (Figure 1A) to quantify the

abundance of mitochondrial CoQ. MASH was associated with a striking ~40% reduction in the

abundance of mitochondrial CoQ (Figure 7I) (tissue samples were not large enough to perform

CoQ redox mass spectrometry on mitochondria). A Pearson correlation analysis showed a

highly robust correlation between the abundances of mitochondrial CL and CoQ (Figure 7J, $R^2 =$

0.64), indicating that the variability in the abundance of CL explains 64% of the variability of the

abundance of CoQ. Based on our findings that CL is reduced with MASLD/MASH and that loss

of CL influences CoQ electron transfer efficiency, we interpret these findings to mean that loss

of CL destabilizes CoQ in humans.

Discussion

In hepatocytes, disruptions of mitochondrial bioenergetics lead to and exacerbate metabolic-

associated steatohepatitis.[52] CL, a key phospholipid in the inner mitochondrial membrane, plays

a critical role in mitochondrial energy metabolism.[23] In this manuscript, we examined the role of

CL in the pathogenesis of MASLD. In mice and in humans, MASLD/MASH coincided with a

reduction in mitochondrial CL levels. Hepatocyte-specific deletion of CLS was sufficient to

spontaneously induce MASH pathology, including steatosis and fibrosis, along with shift in

immune cell populations towards a more pro-inflammatory profile, all of which occurred in mice

given a standard chow diet. Paradoxically, high-resolution respirometry and stable isotope

experiments showed that CLS deletion promotes, instead of attenuates, mitochondrial oxidative

capacity in a fashion reminiscent of temporal changes that occur with mitochondrial

bioenergetics in human MASH.[40] Our principal finding on the role of hepatocyte CL in

bioenergetics is that its loss robustly increases electron leak. This was likely due to: 1) direct

interaction of CL with complex III, 2) lower abundance of I+III$_2$+IV respiratory supercomplex, 3)

global electron leak from the electron transport chain, and 4) loss of CL potentially interfering

with the ability of CoQ to transfer electrons. In humans, mitochondrial CL and CoQ were co-

downregulated in MASH patients compared to healthy controls, with a strong correlation ($R^2 =$

0.64) between CL and CoQ. Together, these results implicate CL as a key regulator of MASH

progression, particularly through its effect on the electron transport chain to promote oxidative

stress.

How might CL regulate CoQ? CoQ is the main electron transporter between complex I/II and III.

CLS deletion in hepatocytes appeared to disrupt CoQ's ability to cycle between its oxidized and

reduced forms. There are several ways in which low CL might directly or indirectly influence

CoQ's redox state. The primary suspect is CL interacting with complex III, as eight or nine CL

molecules are tightly bound to complex III[38] and are thought to be essential to its function.[53]

While CL has been found to bind to other respiratory complexes, our data suggest that loss of

CL might disproportionately influence complex III. This is also supported by our findings that the

loss of CL reduced the formation of complex III-dependent supercomplex, without influencing

other supercomplexes. Reduced capacity for complex III to efficiently accept electrons from

CoQ might explain the increased electron leak at site III_{Q0} and increased level of oxidized CoQ.

Meanwhile, loss of CL also increased electron leak at complex I and II. CL has also been

implicated to bind to these complexes.[20,21,36] Thus, complex III dysfunction is perhaps unlikely to

entirely explain electron leaks at sites I_F, I_Q, and II_F.[54] Conversely, complex I and II are unlikely

to be the only primary sites of defect as such defects probably will not promote electron leak at

site III_{QO}. Moreover, CLS deletion did not have an effect on the abundance of reduced CoQ

despite there being greater abundance of oxidized CoQ, suggesting that complex I/II might also

have reduced ability to transfer electrons onto CoQ. Another potential mechanism by which CL

influences CoQ redox state is by CL directly interacting with CoQ. As they are both lipid

molecules in the IMM, low CL may reduce the lateral diffusability of CoQ between respiratory

complexes. Low CL might also indirectly influence CoQ by contributing to changes in membrane

properties, distribution of ETC in the cristae, and the cristae architecture.[55] Finally, increased

electron leak, regardless of their origin, could have a feed-forward effect by which oxidative

stress disrupts redox homeostasis in other components of ETC.

MASH is a progressive liver disease characterized by lipid accumulation, inflammation, and

fibrosis in the liver.[56] The progression to MASH involves a complex interplay of metabolic stress,

mitochondrial defects, and immune responses that collectively promote hepatocellular injury.[57]

Our findings suggest that a low mitochondrial CL level directly induces key pathological features

of MASH, including steatosis, fibrosis, and immune cell infiltration, even in the absence of

dietary or environmental stressors, such as high-fat diet. When mice were fed a standard chow

diet, CLS-LKO mice were more prone to lipid droplet accumulation than control mice. This

phenotype was exacerbated when the mice were challenged with a high-fat diet. We primarily

interrogated the mitochondrial bioenergetics of standard chow-fed control or CLS-LKO mice. A

lower respiration rate would partially explain the lipid droplet accumulation, but to our surprise,

CLS deletion increased JO_2 regardless of substrates. Similarly, experiments using uniformly

labeled ^{13}C-palmitate or ^{13}C-glucose showed that CLS deletion promoted an overall increase in

the flux toward TCA intermediates, particularly for palmitate. CLS deletion did not appear to

increase de novo lipogenesis or reduce VLDL secretion. Thus, it is unclear what mechanisms

contribute to steatosis induced by CLS deletion.

Liver fibrosis is characterized by the accumulation of excess extracellular matrix components,

including type I collagen, which disrupts liver microcirculation and leads to injury.[55] Livers from

CLS-LKO mice exhibited more fibrosis compared to control mice, even on a standard chow diet,

which was worsened when fed a high-fat diet. Indeed, transcriptomic analyses revealed that

CLS deletion activates pathways for fibrosis and degeneration, with many of the collagen

isoforms upregulated. In the MASH liver, collagen deposition is accompanied by inflammatory

cell infiltrate promoting an overall inflammatory environment. Flow cytometry experiments

further confirmed that CLS deletion led to an increase in Ly6C[hi] cell population, suggesting that

dying resident Kupffer cells are being replaced by Ly6C[hi] monocytes in the livers of CLS-LKO

mice.[33]

Early in the MASLD disease progression, mitochondria adapt to the increased energy demands

by increasing their respiratory capacity. In the later stages of disease progression to MASH,

mitochondrial respiration diminishes.[58] This pattern was reminiscent of our observations in the

CLS-LKO mice. Livers from CLS-LKO mice fed a standard chow diet exhibited greater

respiratory capacity compared to that of control mice. Conversely, livers from CLS-LKO mice on

a high-fat diet exhibited lower respiratory capacity compared to that of control mice. We interpret

these findings to suggest that liver mitochondria in chow-fed CLS-LKO mice are more

representative of early stage of MASLD, while high-fat-diet fed CLS-LKO mice resemble later

stages of MASLD.

In non-hepatocytes, low CL levels have been linked to electron leak in the context of a

deficiency of the tafazzin gene, a CL transacylase, whose mutation promotes Barth

syndrome.[15,26,59,60] Paradoxically, we previously showed that CLS deletion in brown adipocytes

does not increase electron leak.[30] It is important to note that CLS deletion in our current or

previous study does not completely eliminate CL (likely due to an extracellular source). We do

not believe that CL is dispensable for efficient electron transfer in adipocytes. Rather, due to

unclear mechanisms, different cell types likely exhibit varying tolerances to low CL influencing

their bioenergetics, with brown adipocytes appearing more tolerant than hepatocytes.

Regardless, electron leak was elevated with CLS deletion in both standard chow and high-fat

diet-fed conditions. These observations mirror what has been shown in MASLD progression.[61]

The effect of CLS deletion on electron leak was due to reduced CL levels, as the reintroduction

of cardiolipin via SUVs completely rescued the electron leak.

CL is reported to be essential for the formation and stability of supercomplexes.[14,50,51,62] CL has

a distinctive conical shape with four fatty acyl chains, which allows it to create a highly curved

membrane environment in the IMM, promoting close packing of protein complexes that likely

facilitates supercomplex formation.[50] CL also directly interacts with various subunits of the ETC

complexes through electrostatic interactions, which help stabilize the supercomplexes by

anchoring them together in a specific spatial orientation to optimize electron flow.[48] Liver-

specific deletion of CLS only resulted in a lower abundance of the $I+III_2+IV$ supercomplex. We

interpret these findings to mean that $I+III_2+IV$ supercomplex is particularly sensitive to the

reduced CL level in hepatocytes.

In our study, we observed a striking correlation between CL levels and CoQ in human liver

samples from healthy/MASH patients (R_2 of 0.64). In contrast, low mitochondrial CL induced by

CLS deletion coincided with a greater mitochondrial CoQ content in CLS-LKO mice compared to

controls. These data likely suggest that acute and robust reduction in CLS or CL level might

trigger a compensatory CoQ production in mice. Conversely, because the samples from

healthy/MASH patients were from those who had cirrhosis, MASH samples likely came from

subjects that had suffered from years of MASLD pathology. In those samples, where reduction

in CL was quantitatively modest compared to what was induced with CLS knockout, CoQ might

have gradually decreased coincidental to the decrease in CL. Regardless of these differences in

mice and humans, it is clear that there is a relationship between CL and CoQ that is worth

further exploration.

In conclusion, our findings identify a critical role for CL in regulating ETC efficiency to promote

oxidative stress. In both humans and mice, MASLD is associated with a decrease in hepatic

mitochondrial CL, suggesting that low CL may be the cause of the obligatory increase in

oxidative stress known to occur with MASLD progression. We further link CL deficiency to

increased electron leak from the electron transport chain, and as the primary defect that

promotes oxidative stress. We believe that these bioenergetic changes underlie the

pathogenesis of MASLD, as CL deletion was sufficient to cause steatosis, fibrosis, and

inflammation, phenocopying many changes that occur with MASLD/MASH progression. Further

research will be needed to fully uncover how CL regulates CoQ, and to test whether rescuing

the CL/CoQ axis might be effective in treating patients with MASLD/MASH.

Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
GRIM19	Abcam	ab110240
SDHA	Abcam	ab14715
UQCRFS1	Abcam	ab14746
MTCO1	Abcam	ab14705
ATP5a	Abcam	Ab14748
NDUFA9	Abcam	Ab14713
Total OxPhos Antibody cocktail	Abcam	MS604-300
Citrate Synthetase	Abcam	Ab96600
Cytochrome c	Cell Signaling	11940S
Caspase-3	Cell Signaling	9661S
Caspase-7	Cell Signaling	9491S
CD16/32 (Fc Block, clone 93)	Biolegend	101308
CD45 (FITC, clone S18009F)	Biolegend	157214
CD11b (BVC421, clone M1/70)	Biolegend	101202
F4/80 (APC, clone BM8)	Biolegend	123102
TIM4 (PerCP/Cy5.5, clone RMT4-54)	Biolegend	130020
Ly6C (PE, clone HK1.4)	Biolegend	128008
MHCII (BV605, clone M5/114.15.2)	Biolegend	107639
CD11c (BV785, clone N418)	Biolegend	117302
Ly6G (PE/Cy7, clone 1A8)	Biolegend	164504
OxPhos Blue Native WB Antibody Cocktail	ThermoFisher	45-7999
Bacterial and virus strains		
Second-generation lentiviral-mediated knockdown system		
NEB Stable Competent E. Coli	NEB	C3040H
Biological samples		
Chemicals, peptides, and recombinant proteins		
Amplex Red Reagent	ThermoFisher Scientific	A12222
Auranofin	Sigma Aldrich	A6733
Carmustine (BCNU)	Sigma Aldrich	C0400
SPLASH Mix	Avanti Polar Lipids	330707
Cardiolipin Mix I	Avanti Polar Lipids	LM6003
Bovine Serum Albumin	Sigma Aldrich	A7030

Protease Inhibitor Cocktail	Thermo Scientific	78446
Tamoxifen	Sigma Aldrich	T5648
Sunflower Oil	Sigma Aldrich	S5007
TRIzol	Thermo Scientific	15596018
Mini-PROTEAN TGX Gels	BioRad	4561086
ECL	PerkinElmer	104001EA
Malate	Sigma Aldrich	M7397
Pyruvate	Sigma Aldrich	P2256
GDP	Sigma Aldrich	G7127
CL 316,243 hydrate	Sigma Aldrich	C5976
ADP	Sigma Aldrich	A5285
ATP	Sigma Aldrich	A9187
Glutamate	Sigma Aldrich	G5889
Succinate	Sigma Aldrich	S3674
Carnitine	Sigma Aldrich	8.40092
Palmitoyl-CoA	Sigma Aldrich	P9716
Palmitoyl-L-carnitine	Sigma Aldrich	P1645
SYBR Green	Thermo Scientific	A25776
4% Paraformaldehyde	Thermo	J19943-K2
Opti-MEM	Gibco	31985
DMEM	Gibco	1195-092
FBS	Gibco	10082-147
Penicillin-streptomycin	Gibco	15140122
CL for SUVs	Avanti Polar Lipids	840012
PC for SUVs	Avanti Polar Lipids	850375C
PAGE 3-12% Bis-Tris Gels	Thermo Scientific	BN1001BOX
NativePage 20x Cathode Buffer	Invitrogen	BN2002
NativePage 5% G-250 Sample additive	Invitrogen	BN2004
Nativepage 4x Sample Buffer	Invitrogen	BN2003
CoQ Standard	Cambridge Isotope Labs	CIL DLM-10279
Palmitic acid (U-$^{13}C_{16}$)	Cambridge Isotope Labs	CLM-409-0.1
D-Glucose ((U-$^{13}C_6$)	Cambridge Isotope Labs	CLM-1396-5
Critical commercial assays		
Pierce BCA Protein Assay Kit	Thermo Scientific	23227
iScript cDNA Synthesis Kit	BioRad	1708891
SDH Detection Assay Kit	Abcam	ab228560
Direct-zol RNA Miniprep Plus Kit	Zymo	R2070
DNeasy Blood and Tissue Kit	Qiagen	69504

Deposited data		
Experimental models: Cell lines		
HEK293T cells	ATCC	CTRL-3216
Hepa 1-6 murine hepatoma cells	ATCC	CRL-1830
Experimental models: Organisms/strains		
Mouse: CLS conditional knockout (CLS-cKO)	Sustarsic et al. 2018	N/A
Mouse: CLS-LKO	This paper	N/A
Mouse: Alb-Cre	Jackson Laboratory	003574
Oligonucleotides		
RT qPCR Primer ATGL F (CCACTCACATCTACGGAGCC)	www.IDTDNA.com	
RT qPCR Primer ATGL R (TAATGTTGGCACCTGCTTCA)	www.IDTDNA.com	
RT qPCR Primer DGAT1 F (GACGGCTACTGGGATCTGA)	www.IDTDNA.com	
RT qPCR Primer DGAT1 R (TCACAACACACCAATTCAGG)	www.IDTDNA.com	
RT qPCR Primer FAS F (GGATAGCTGTGTAGTGTAACCAT)	www.IDTDNA.com	
RT qPCR Primer FAS R (GGTCATCGTGATAACCACACA)	www.IDTDNA.com	
RT qPCR Primer SCD1 F (GCTCTACACCTGCCTCTTCG)	www.IDTDNA.com	
RT qPCR Primer SCD1 R (CAGCCGAGCCTTGTAAGTTC)	www.IDTDNA.com	
RT qPCR Primer CLS F (TGACCTATGCAGATCTTATTCCA)	Johnson et al. 2019	
RT qPCR Primer CLS R (TGGCAGAGTTCGGTATCTGA)	Johnson et al. 2019	
RT qPCR Primer TNFa F (CCACCACGCTCTTCTGTCTAC)	www.IDTDNA.com	
RT qPCR Primer TNFa R (AGGGTCTGGGCCATAGAACT)	www.IDTDNA.com	
RT qPCR Primer Taz F (CCCTCCATGTGAAGTGGCCATTCC)	Johnson et al. 2019	
RT qPCR Primer Taz R (TGGTGGTTGGAGACGGTGATAAGG)	Johnson et al. 2019	
mtDNA F: (TTAAGACACCTTGCCTAGCCACAC)	Mouse Primer Depot NCI/NIH	
mtDNA R: (CGGTGGCTGGCACGAAATT)	Mouse Primer Depot NCI/NIH	

nucDNA F: (ATGACGATATCGCTGCGCTG)	Mouse Primer Depot NCI/NIH	
nucDNA R: (TCACTTACCTGGTGCCTAGGGC)	Mouse Primer Depot NCI/NIH	
Recombinant DNA		
Sc	Addgene	1864
Crls1	Sigma Aldrich	TRCN0000123937
psPAX2	Addgene	12260
pMD2.G	Addgene	12259
Software and algorithms		
GraphPad Prism 9.0	GraphPad	N/A
Other		

Experimental model and subject details

Human participants

De-identified liver samples were acquired from the University of Utah Biorepository and

Molecular Pathology Shared Resource from patients undergoing liver transplantation or

resection due to end-stage liver disease and/or liver tumor(s). Informed consent was obtained

from patients. All patients were classified to their respective diagnosis by a pathologist at the

time of initial collection. The diagnosis for individual tissue samples was confirmed by a

pathologist based on histology review of formalin-fixed, paraffin-embedded sections takes from

the same location as the tissue analyzed by targeted lipid mass spectrometry.

Mice

All mice (male and female) used in this study were bred onto C57BL/6J background. CLS-LKO

mice were generated by crossing the CLS conditional knockout (CLScKO[+/+]) generously

donated by Dr. Zachary Gerhart-Hines (University of Copenhagen)[24] with mice heterozygous for

Albumin promotor Cre (Alb-Cre$^{+/-}$) to produce liver-specific deletion of the CLS gene

(CLScKO$^{+/+}$, Alb-Cre$^{+/-}$) or control (CLScKO$^{+/+}$, no Cre) mice. CLScKO$^{+/+}$ mice harbor loxP sites

that flank exon 4 of the CLS gene. For high-fat diet studies, 8 wk CLS-LKO and their respective

controls began high-fat diet (HFD, 42% fat, Envigo TD.88137) feeding for 8 wks. Mice were

fasted 4 hours and given an intraperitoneal injection of 80 mg/kg ketamine and 10 mg/kg

xylazine prior to terminal experiments and tissue collection. All animal experiments were

performed with the approval of the Institutional Animal Care and Use Committees at the

University of Utah.

Cell lines

Hepa 1-6 murine hepatoma cells were grown in high-glucose DMEM (4.5 g/L glucose, with L-

glutamine; Gibco 11965-092) supplemented with 10% FBS (heat-inactivated, certified, US

origin; Gibco 10082-147), and 0.1% penicillin-streptomycin (10,000 U/mL; Gibco 15140122). For

lentivirus-mediated knockdown of CLS, CLS expression was decreased using the pLKO.1

lentiviral-RNAi system. Plasmids encoding shRNA for mouse *Crls1* (shCLS: TRCN0000123937)

was obtained from MilliporeSigma. Packaging vector psPAX2 (ID 12260), envelope vector

pMD2.G (ID 12259), and scrambled shRNA plasmid (SC: ID 1864) were obtained from

Addgene. HEK293T cells in 10 cm dishes were transfected using 50 µL 0.1% polyethylenimine,

200 µL, 0.15 M sodium chloride, and 500 µL Opti-MEM (with HEPES, 2.4 g/L sodium

bicarbonate, and l-glutamine; Gibco 31985) with 2.66 µg of psPAX2, 0.75 µg of pMD2.G, and 3

µg of either scrambled or *Crls1* shRNA plasmid. Cells were selected with puromycin throughout

differentiation to ensure that only cells infected with shRNA vectors were viable.

Method details

Body composition

To assess body composition, mice were analyzed using a Bruker Minispec NMR (Bruker,

Germany) 1 week prior to terminal experiments. Body weights were measured and recorded

immediately prior to terminal experiments.

RNA quantification

For quantitative polymerase chain reaction (qPCR) experiments, mouse tissues were

homogenized in TRIzol reagent (Thermo Fisher Scientific) and RNA was isolated using

standard techniques. The iScript cDNA Synthesis Kit was used to reverse transcribe total RNA,

and qPCR was performed with SYBR Green reagents (Thermo Fisher Scientific). Pre-validated

primer sequences were obtained from mouse primer depot

(https://mouseprimerdepot.nci.nih.gov/). All mRNA levels were normalized to RPL32. For RNA

sequencing, liver RNA was isolated with the Direct-zol RNA Miniprep Plus kit (Zymo Cat#:

R2070). RNA library construction and sequencing were performed by the High-Throughput

Genomics Core at the Huntsman Cancer Institute, University of Utah. RNA libraries were

constructed using the NEBNext Ultra II Directional RNA Library Prep with rRNA Depletion Kit

(human, mouse rat). Sequencing was performed using the NovaSeq S4 Reagent Kit v1.5

150x150 bp Sequencing with >25 million reads per sample using adapter read 1:

AGATCGGAAGAGCACACGTCTGAACTCCAGTCA and adapter read 2:

AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT. Pathway analyses were performed by the

Bioinformatics Core at the Huntsman Cancer Institute, University of Utah using the Reactome

Pathway Database. For differentially expressed genes, only transcripts with Padj < 0.05 and

baseMean > 100 are included.

DNA isolation and quantitative PCR

Genomic DNA for assessments of mitochondrial DNA (mtDNA) was isolated using a

commercially available kit according to the manufacturer's instructions (69504, Qiagen).

Genomic DNA was added to a mixture of SYBR Green (Thermo Fisher Scientific) and primers.

Sample mixtures were pipetted onto a 3840well plate and analyzed with QuantStudio 12K Flex

(Life Technologies). The following primers were used: mtDNA fwd, TTAAGA-CAC-CTT-GCC-

TAG-CCACAC; mtDNA rev, CGG-TGG-CTG-GCA-CGA-AAT-T; nucDNA fwd, ATGACG-ATA-

TCG-CTG-CGC-TG; nucDNA rev, TCA-CTT-ACC-TGGTGCCTA-GGG-C.

Western blot analysis

For whole liver lysate, frozen liver was homogenized in a glass homogenization tube using a

mechanical pestle grinder with homogenization buffer (50 mM Tris pH 7.6, 5 mM EDTA, 150

mM NaCl, 0.1% SDS, 0.1% sodium deoxycholate, 1% triton X-100, and protease inhibitor

cocktail). After homogenization, samples were centrifuged for 15 min at 12,000 × g. Protein

concentration of supernatant was then determined using a BCA protein Assay Kit (Thermo

Scientific). Equal protein was mixed with Laemmeli sample buffer and boiled for 5 mins at 95°C

for all antibodies except for OXPHOS cocktail antibody (at room temp for 5 mins), and loaded

onto 4–15% gradient gels (Bio-Rad). Transfer of proteins occurred on a nitrocellulose

membrane and then blocked for 1 hr. at room temperature with 5% bovine serum albumin in

Tris-buffered saline with 0.1% Tween 20 (TBST). The membranes were then incubated with

primary antibody (see Extended Data Table 2), washed in TBST, incubated in appropriate

secondary antibodies, and washed in TBST. Membranes were imaged utilizing Western

Lightning Plus-ECL (PerkinElmer) and a FluorChem E Imager (Protein Simple). For isolated

mitochondria, identical procedures were taken with equal protein of mitochondrial preps.

Single cell preparation of liver tissue for flow cytometry

After mice were euthanized using isoflurane, blood was collected by cardiac puncture, the

abdomen was exposed and the liver collected, rinsed with PBS and weighed. Liver was

subsequently transferred in approximately 3ml of serum-free RPMI-1640 containing

Collagenase D (10mg/ml; Sigma) and DNase (1mg/ml; Sigma) and incubated in a rocking

platform for 45 min at 37°C. The liver extract was mashed through a 70µm filter and re-

suspended in RPMI-1640 containing 10% FBS and centrifuged at 1600 rpm for 5 min. The

supernatant was discarded and the pellet re-suspended in approximately 4 ml of 70% Percoll,

then transferred to 15 ml conical tube, carefully overlayed with 4 ml of 30% Percoll and

centrifuged 1600 rpm for 25 min with the brake turned off. The non-parenchymal cell

suspension from the Percoll interface was removed and mixed with 10 mL of RPMI-1640

containing 10% FBS and the cells were centrifuged at 1600 rpm for 5 min. Red blood cells

(RBC) were removed from the pelleted single cell suspensions of livers non-parenchymal cells

by incubation in an ammonium chloride -based 1x RBC lysis buffer (Thermofisher, eBioscience).

The cells were again pelleted and mixed with FACS buffer (2% BSA, 2mM EDTA in PBS), then

stained with Zombie-NIR viability dye (BioLegend) per manufacturer's instructions to

discriminate live vs dead cells. To prevent non-specific Fc binding, the cells were incubated with

Fc Block (anti-mouse CD16/32, clone 93, Biolegend) for 15 min followed by the indicated

antibodies cocktail for 60 min in the dark on ice: CD45 (FITC, clone S18009F, Biolegend),

CD11b (BVC421, clone M1/70, Biolegend), F4/80 (APC, clone BM8, Biolegend), TIM4

(PerCP/Cy5.5, clone RMT4-54, Biolegend), Ly6C (PE, clone HK1.4, Biolegend), MHCII (BV605,

clone M5/114.15.2, Biolegend), CD11c (BV785, clone N418, Biolegend) and Ly6G (PE/Cy7,

clone 1A8, Biolegend). After surface staining, cells were fixed with a paraformaldehyde-based

fixation buffer (BioLegend). Flow cytometric acquisition was performed on a BD Fortessa X20

flow cytometer (BD Biosciences) and data analyzed using FlowJo software (Version 10.8.1;

Tree Star Inc).

Glucose tolerance test

Intraperitoneal glucose tolerance tests were performed by injection of 1 mg glucose per gram

body mass at least 6 days prior to sacrifice. Mice were fasted for 4 hours prior to glucose

injection. Blood glucose was measured 30 minutes before glucose injection and at 0, 15, 30, 60,

90, and 120 minutes after injection via tail bleed with a handheld glucometer (Bayer Contour

7151H).

Pyruvate tolerance test

Pyruvate tolerance tests were performed by injection of 2 mg pyruvate per gram of body mass

in PBS adjusted to pH 7.3-7.5 at least 6 days prior to sacrifice. Blood glucose was measured 30

minutes before pyruvate injection and at 0, 15, 30, 45, 60, 75, 90, 105, and 120 minutes after

injection via tail bleed with a handheld glucometer (Bayer Contour 7151H).

Electron microscopy

To examine mitochondrial ultrastructure and microstructures, freshly dissected liver tissues from

CLS-LKO and their controls were sectioned into ≈2 mm pieces and processed by the Electron

Microscopy Core at University of Utah. To maintain the ultrastructure of the tissue via

irreversible cross-link formation, each section was submerged in fixative solution (1%

glutaraldehyde, 2.5% paraformaldehyde, 100 mM cacodylate buffer pH 7.4, 6 mM CaCl2, 4.8%

sucrose) and stored at 4°C for 48 hours. Samples then underwent 3 × 10-minute washes in 100

mM cacodylate buffer (pH 7.4) prior to secondary fixation (2% osmium tetroxide) for 1 hour at

room temperature. Osmium tetroxide as a secondary fixative has the advantage of preserving

membrane lipids which are not preserved using aldehyde alone. After secondary fixation,

samples were subsequently rinsed for 5 minutes in cacodylate buffer and distilled H_2O, followed

by prestaining with saturated uranyl acetate for 1 hour at room temperature. After prestaining,

each sample was dehydrated with a graded ethanol series (2 × 15 minutes each: 30%, 50%,

70%, 95%; then 3 × 20 minutes each: 100%) and acetone (3 × 15 minutes) and were infiltrated

with EPON epoxy resin (5 hours 30%, overnight 70%, 3 × 2-hour 40 minute 100%, 100% fresh

for embed). Samples were then polymerized for 48 hours at 60°C. Ultracut was performed using

Leica UC 6 ultratome with sections at 70 nm thickness and mounted on 200 mesh copper grids.

The grids with the sections were stained for 20 minutes with saturated uranyl acetate and

subsequently stained for 10 minutes with lead citrate. Sections were examined using a JEOL

1200EX transmission electron microscope with a Soft Imaging Systems MegaView III CCD

camera.

Histochemistry

A fresh liver tissue was taken from each mouse and immediately submerged in 4%

paraformaldehyde for 12 hours and then 70% ethanol for 48 hours. Tissues were sectioned at

10-µm thickness, embedded in paraffin, and stained for Masson's Trichrome to assess fibrosis

or hematoxylin and eosin (H&E) to determine fat droplet accumulation. Samples were imaged

on Axio Scan Z.1 (Zeiss).

Native PAGE

Isolated mitochondria (100 µg) suspended in MIM were pelleted at 12,000 x g for 15 min and

subsequently solubilized in 20 µL sample buffer (5 µL of 4x Native Page Sample Buffer, 8 µL

10% digitonin, 7 µL ddH$_2$O per sample) for 20 min on ice and then centrifuged at 20,000 x g for

30 mins at 4°C. 15 µL of the supernatant (75 µg) was collected and placed into a new tube and

mixed with 2 µL of G-250 sample buffer additive. Dark blue cathode buffer (50 mLs 20X Native

Page running buffer, 50 mLs 20x cathode additive, 900 mLs ddH$_2$O) was carefully added to the

front of gel box (Invitrogen Mini Gel Tank A25977) and anode buffer (50 mLs 20x Native Page

running buffer to 950 mL ddH$_2$O) was carefully added to the back of the gel box making sure to

not mix. The samples were then loaded onto a native PAGE 3-12% Bis-Tris Gel (BN1001BOX,

Thermo Fisher Scientific), and electrophoresis was performed at 150 V for 1 hour on ice. The

dark blue cathode buffer was carefully replaced with light blue cathode buffer (50 mLs 20X

Native Page running buffer, 5 mL 20X cathode additive to 945 mLs ddH$_2$O) and run at 30 V

overnight at 4°C. Gels were subsequently transferred to PVDF at 100 V, fixed with 8% acetic

acid for 5 min, washed with methanol, and blotted with the following primary antibodies Anti-

GRIM19 (mouse monoclonal; ab110240), Anti-SDHA (mouse monoclonal; ab14715), Anti-

UQCRFS1 (mouse monoclonal; ab14746), Anti-MTCO1 (mouse monoclonal; ab14705), Anti-

ATP5a (mouse monoclonal; ab14748), Anti-NDUFA9 (mouse monoclonal; ab14713) in 5% non-

fat milk in TBST. Secondary anti-mouse HRP antibody listed in the key resources table and

Western Lightning Plus-ECL (PerkinElmer NEL105001) was used to visualize bands.

Isolation of mitochondrial and cytosolic fractions

Liver tissues were minced in ice-cold mitochondrial isolation medium (MIM) buffer [300 mM

sucrose, 10 mM Hepes, 1 mM EGTA, and bovine serum albumin (BSA; 1 mg/ml) (pH 7.4)] and

gently homogenized with a Teflon pestle. To remove excess fat in the samples, an initial high-

speed spin was performed on all samples: homogenates were centrifuged at 12,000 x g for 10

mins at 4°C, fat emulsion layers were removed and discarded, and resulting pellets were

resuspended in MIM + BSA. Samples were then centrifuged at 800 x g for 10 min at 4°C. The

supernatants were then transferred to fresh tubes and centrifuged again at 1,300 x g for 10 min

at 4°C. To achieve the mitochondrial fraction (pellet), the supernatants were again transferred to

new tubes and centrifuged at 12,000 x g for 10 min at 4°C. The resulting crude mitochondrial

pellets were washed three times with 0.15 M KCl to remove catalase, and then spun a final time

in MIM. The final mitochondrial pellets were resuspended in MIM buffer for experimental use.

For cytosolic fraction, the supernatant from 12,000 x g spin was ultracentrifuged at 180,000 x g

for 16 hours in a fixed angle rotor (FIBERLite, F50L-24x1.5). The top half of the supernatant

was collected as analyzed as the cytosolic fraction for western blotting analyses.

Mitochondrial respiration measurements

Mitochondrial O_2 utilization was measured using Oroboros O2K Oxygraphs. Isolated

mitochondria (50 µg for TCA substrate respiration and 100 µg for fatty acid respiration) were

added to the oxygraph chambers containing assay buffer Z (MES potassium salt 105 mM, KCl

30 mM, KH_2PO_4 10 mM, $MgCl_2$ 5 mM, BSA 1 mg/ml). Respiration was measured in response to

the following substrates: 0.5 mM malate, 5 mM pyruvate, 2.5 mM ADP, 10 mM succinate, 1.5

µM FCCP, 0.02 mM palmitoyl-carnitine, 0.5 mM malate.

Mitochondrial JH_2O_2

Mitochondrial H_2O_2 production was determined in isolated mitochondria from liver tissue using

the Horiba Fluoromax-4/The Amplex UltraRed (10 µM)/horseradish peroxidase (3 U/ml)

detection system (excitation/emission, 565:600, HORIBA Jobin Yvon Fluorolog) at 37°C.

Mitochondrial protein was placed into a glass cuvette with buffer Z supplemented with 10 mM

Amplex UltraRed (Invitrogen), 20 U/mL CuZn SOD). Since liver tissue is capable of producing

resorufin from amplex red (AR), without the involvement of horseradish peroxidase (HRP) or

H_2O_2, phenylmethylsulfonyl fluoride (PMSF) was included to the experimental medium due to its

ability to inhibit HRP-independent conversion of AR to resorufin.[63] PMSF was added to the

cuvette immediately prior to measurements and at a concentration that does not interfere with

biological measurements (100 µM).[63] A 5-min background rate was obtained before adding 10

mM succinate to the cuvette to induce H_2O_2 production. After 4 min, 100 µM 1,3-bis(2-

chloroethyl)-1-nitrosourea (BCNU) was added to the cuvette with 1 µM auranofin to inhibit

glutathione reductase and thioredoxin reductase, respectively. After an additional 4 min, the

assay was stopped, and the appearance of the fluorescent product was measured.

Site-specific electron leak was measured by systematically stimulating each site while inhibiting

the other three.[54] Site I_F was investigated in the presence of 4 mM malate, 2.5 mM ATP, 5 mM

aspartate, and 4 µM rotenone; site I_Q was measured as a 4 µM rotenone-sensitive rate in the

presence of 5 mM succinate; site III_{QO} was measured as a 2 µM myxothiazol-sensitive rate in

the presence of 5 mM succinate, 5 mM malonate, 4 µM rotenone, and 2 µM antimycin A; and

site II_F was measured as the 1 mM malonate-sensitive rate in the presence of 0.2 mM succinate

and 2 µM myxothiazol. As previously mentioned, electron leak is quantified using Amplex Red in

the presence of excess superoxide dismutase, such that both superoxide and hydrogen

peroxide production are accounted for by a change in fluorescence intensity (JH_2O_2) using high-

resolution fluorometry (Horiba Fluoromax4®).

Lipid mass spectrometry for reduced and oxidized CoQ

Liver tissue was homogenized in ice cold isolation buffer (250 mM sucrose, 5 mM Tris-HCl, 1

mM EGTA, 0.1% fatty acid free BSA, pH 7.4, 4°C) using a tissuelyser. Mitochondria were then

isolated via differential centrifugation (800 x g for 10 min, 1300 x g for 10 min, 12,000 x g for 10

min at 4°C), flushing each step under a stream nitrogen to prevent oxidation. Protein content

was determined by bicinchoninic acid assay using the Pierce BCA protein assay with bovine

serum albumin as a standard. To extract CoQ from mitochondria, incubations of 100 µg

mitochondrial protein in 250 µL ice-cold acidified methanol, 250 µL hexane, and 1146 pmol per

sample of CoQ standard (Cambridge Isotope Laboratories, CIL DLM-10279) were vortexed. The

CoQ-containing hexane layer was separated by centrifugation (10 min, 17,000 x g, 4°C) and

then dried down under a stream of nitrogen. Dried samples were then resuspended in methanol

containing 2 mM ammonium formate and transferred to 1.5 mL glass mass spectrometry vials.

Liquid chromatography-mass spectrometry (LC-MS/MS) was then performed on the

reconstituted lipids using an Agilent 6530 UPLC-QTOF mass spectrometer.

U-[13]C-glucose and U-[13]C-palmitate labeling in hepa1-6 cell line

Stable isotope tracing experiments were conducted using U-[13]C-glucose or U-[13]C-palmitate to

assess metabolic fluxes in hepa1-6 cells with or without CLS knockdown. Cells were incubated

in either U-^{13}C-glucose or U-^{13}C-palmitate for 4 hours. For palmitate incubations, U-^{13}C-

palmitate was conjugated to fatty acid-free bovine serum albumin (BSA) and supplemented with

1 mM carnitine prior to being added to the culture medium. Following the 4-hour incubation

period, cells were washed once with 1 mL of ice-cold Dulbecco's Phosphate Buffered Saline

(dPBS) and then scraped in 0.7 mL ice-cold dPBS. The cells were pelleted by centrifugation at

500 x g for 5 minutes at 4°C, and the supernatant was carefully removed. Cell pellets were

resuspended in 215 µL of ice-cold dPBS and sonicated on ice using a 1/8" probe sonicator at

30% amplitude for 6 seconds. Debris was lightly pelleted by centrifugation at 500 x g for 2

minutes at 4°C. From the resulting supernatant, 25 µL was transferred to a separate tube for

protein quantification using the bicinchoninic acid (BCA) assay. Samples incubated with U-^{13}C-

glucose were submitted for gas chromatography-mass spectrometry (GC-MS) and liquid

chromatography-mass spectrometry (LC-MS) analysis, while those incubated with U-^{13}C-

palmitate were submitted for GC-MS analysis only.

GC-MS

All GC-MS analysis was performed with an Agilent 5977b GC-MS MSD-HES and an Agilent

7693A automatic liquid sampler. Dried samples were suspended in 40 µL of a 40 mg/mL O-

methoxylamine hydrochloride (MOX) (MP Bio #155405) in dry pyridine (EMD Millipore

#PX2012-7) and incubated for one hour at 37 °C in a sand bath. 25 µL of this solution was

added to auto sampler vials. 60 µL of N-methyl-N-trimethylsilyltrifluoracetamide (MSTFA with

1% TMCS, Thermo #TS48913) was added automatically via the auto sampler and incubated for

30 minutes at 37 °C. After incubation, samples were vortexed and 1 µL of the prepared sample

was injected into the gas chromatograph inlet in the split mode with the inlet temperature held at

250 °C. A 10:1 split ratio was used for analysis for most metabolites. Any metabolites that

saturated the instrument at the 10:1 split was analyzed at a 100:1 split ratio. The gas

chromatograph had an initial temperature of 60 °C for one minute followed by a 10 °C/min ramp

to 325 °C and a hold time of 10 minutes. A 30-meter Agilent Zorbax DB-5MS with 10 m

Duraguard capillary column was employed for chromatographic separation. Helium was used as

the carrier gas at a rate of 1 mL/min.

Data were collected using MassHunter software (Agilent). Metabolites were identified and their

peak area was recorded using MassHunter Quant. This data was transferred to an Excel spread

sheet (Microsoft, Redmond WA). Metabolite identity was established using a combination of an

in-house metabolite library developed using pure purchased standards, the NIST library and the

Fiehn library. There are a few reasons a specific metabolite may not be observable through GC-

MS.

LC-MS

A pooled quality control (QC) sample for each sample type was prepared by combining 10 µL of

each sample. Samples were randomized prior to analysis. A SCIEX 7600 Zeno-ToF (AB SCIEX

LLC, Framingham, MA, USA) coupled to an Agilent 1290 Infinity II HPLC system in positive-

ionization modes was used for analysis. Separation was achieved using a Waters BEH zhilic 2.1

x 100 mm column (Waters Corporation, Milford, MA, USA) with Phenomenex Krudkatcher Ultra

(Phenomenex, Torrence, CA, USA). Buffers consisted of 99% ACN with 5% ddH2O (buffer B)

and 5% 25 mM ammonium carbonate in ddH2O (buffer A). An initial concentration of 99% buffer

B was decreased to 85% over 2 min, then further decreased to 75%, and 60% over 3-minute

intervals. Next, buffer B was decreased to 40% and held for 1 min. Then Buffer B was

decreased to 1% and held for 1 min. Eluents were returned to initial conditions over 0.1 min,

and the system was allowed to equilibrate for 5 min between runs. Mass spectrometry analysis

was performed by high-resolution multiple reaction monitoring (MRM HR).

Data was collected using SCIEX analyst. Chromatogram integration was performed using

SCIEX MultiQuant. This data was transferred to an Excel spread sheet (Microsoft, Redmond

WA). Statistical analysis was performed using Microsoft Excel Data Analysis add-in. The

metabolite lists were curated to the metabolites of interest for the researcher as applicable.

Metabolite identity was established using a combination of an in-house metabolite library

developed using pure purchased standards and the human metabolite data base (hmdb.ca).

Isotope correction was performed using in-house software.

Mitochondrial phospholipids enrichment

Isolated mitochondria (500 µg) from 2-month-old mice were incubated in fusion buffer [220 mM

mannitol, 70 mM sucrose, 2 mM Hepes, 10 mM KH2PO4, 5 mM MgCl2, 1 mM EGTA, 10 mM

glutamate, 2 mM malate, 10 mM pyruvate, and 2.5 mM ADP (pH 6.5)] for 20 min at 30°C under

constant stirring agitation in the presence of 15 nmol of small unilamellar vesicles (SUVs) (CL

from Avanti Polar Lipids, 840012; PC from Avanti Polar Lipids, 850375C). After fusion,

mitochondria were layered on a sucrose gradient (0.6 M) and centrifuged 10 min at 10,000g at

4°C to remove SUV. Pellet was then washed in mitochondrial buffer [250 mM sucrose, 3 mM

EGTA, and 10 mM tris-HCl, (pH 7.4)].

Serum AST and ALT

Mice were sacrificed by CO_2 inhalation and blood samples collected via cardiac puncture into

heparin-coated tubes and centrifuged for collection of plasma within 1 hour of blood collection

and frozen at -80°C until analysis. Plasma samples from mice were processed in a single batch

for determination of serum alanine aminotransferase (ALT) and aspartate aminotransferase

(AST) levels using a DC Element chemistry analyser (HESKA).

Quantification and statistical analyses

The level of significance was set at $p < 0.05$. Student's T tests were used to determine the

significance between experimental groups and two-way ANOVA analysis followed by a within-

row pairwise comparison post hoc test where appropriate. For Student's T tests, Gaussian

distribution was assumed as well as assuming both populations had the same standard

deviation. The sample size (n) for each experiment is shown in the figure legends and

corresponds to the sample derived from the individual mice or for cell culture experiments on an

individual batch of cells. Unless otherwise stated, statistical analyses were performed using

GraphPad Prism software.

Materials availability

Plasmids utilized by this study are available from Sigma Aldrich. Mouse lines generated by this

study may be available at personal request from the lead contact. No new reagents were

created or used by this study.

Data and code availability

All data are available in the main text or the Extended Data. Code for RNA sequencing can be

retrieved upon request.

 Figure Legends

Figure 1. Hepatic mitochondrial phospholipidome in models of MASLD/MASH

(a) Schematic for mitochondrial phospholipidomic analyses in human liver samples. (b)

Representative H&E staining of healthy or MASH livers. Arrows indicate fibrotic liver tissue. (c,

d) Hepatic mitochondrial CL and PG levels in healthy controls or individuals with MASH (n=10

and 16 per group). (e) Pathway for CL biosynthesis. (f–i) Representative H&E and Masson's

Trichrome staining of livers from mice under various MAFLD conditions: chow vs. Western HFD

(16 wks), wildtype vs. ob/ob (20 wks), chow vs. GAN diet (30 wks), and vehicle vs. carbon

tetrachloride (6 wks). (j–m) Western blot of OXPHOS subunits and citrate synthase in liver

tissues from the same MAFLD models. (n–q) Heatmaps of hepatic mitochondrial

phospholipidomes from the same MAFLD models. (r) Venn diagram comparing hepatic

mitochondrial phospholipidomes across all four MASLD models. (s–v) CLS mRNA levels in

livers from the same MAFLD models (n=4–6 per group). Statistical significance was determined

by 2-way ANOVA with within-row pairwise comparison (c, d, n–q) and unpaired, two-sided

Student's t-test (c, d total lipid panels, s–v). All measurements were taken from distinct samples.

Figure 2. Hepatocyte-specific deletion of CLS induces MASLD/MASH

(a) Schematic for hepatocyte-specific deletion of CLS in mice. (b, c) CLS mRNA and

mitochondrial CL levels in livers from control or CLS-LKO mice (n=5–7 per group). (d–f) Body

mass, body composition, and liver mass of control or CLS-LKO mice (n=6–13 per group). (g, h)

Representative H&E and Masson's Trichrome staining of livers from control or CLS-LKO mice.

(i) RNAseq-derived heatmap of select genes associated with MASH, liver regeneration, and

HCC (n=5–7 per group). (j, k) Serum AST and ALT levels (n=6–7 per group). (l) mRNA levels of

TNFα, TGFβ, IL-12, and MCP1 in livers from control or CLS-LKO mice (n=5–7 per group). (m)

Representative flow cytometry gating for liver cell populations in control or CLS-LKO mice. (n–s)

Quantification of cDC2, F4/80+, Ly6C^{hi} inflammatory monocytes, MHC-II+, neutrophils, and

cDC1 cell populations in livers (n=5–7 per group). All results are from mice fed standard chow.

Statistical significance was determined by 2-way ANOVA with within-row pairwise comparison

(c, l) and unpaired, two-sided Student's t-test (b, d–f, i [adjusted for FDR], and j–s). Data

represent mean ± SEM (b–f, j–l, n–s). All measurements were taken from distinct samples.

Figure 3. CLS deletion increases mitochondrial respiratory capacity

(a, b) Glucose tolerance test (IPGTT) and area under the curve (AUC) for control or CLS-LKO

mice fed standard chow (n=6–7 per group). (c, d) Pyruvate tolerance test (PTT) and AUC (n=6–

8 per group). (e, f) RNAseq-derived heatmaps of genes associated with lipogenesis, VLDL, β-

oxidation, and ETS complex structure/function (n=5–6 per group). (g) Representative electron

microscopy images of liver mitochondria from control or CLS-LKO mice. (h) Western blot of

whole liver lysate using OXPHOS cocktail and citrate synthase for control or CLS-LKO mice. (i)

Mitochondrial-to-nuclear DNA ratio in liver tissue from control or CLS-LKO mice (n=8 per group).

(j) Representative tracing from high-resolution respirometry during maximal respiration using

TCA cycle intermediates. (k, l) JO_2 consumption in isolated liver mitochondria from control or

CLS-LKO mice in response to malate, pyruvate, ADP, succinate, FCCP (k), or palmitoyl-

carnitine, malate, and ADP (l) (n=6 per group). (m) Western blot of isolated mitochondria from

livers of control or CLS-LKO mice using OXPHOS cocktail. All results are from mice fed

standard chow. Statistical significance was determined by 2-way ANOVA with within-row

pairwise comparison (a, c, k, l) and unpaired, two-sided Student's t-test (b, d, e, f [adjusted for

FDR], and i). Data represent mean ± SEM (a–d, k, l). Data in box-and-whiskers plot (i) represent

median with min-to-max. All measurements were taken from distinct samples.

Figure 4. Stable isotope tracing with [U-^{13}C] palmitate and [U-^{13}C] glucose in hepa1-6 cells

with or without CLS deletion

(a) Schematic of stable isotope tracing with [U-^{13}C] palmitate or [U-^{13}C] glucose, showing key

intermediates in β-oxidation and the TCA cycle. (b–d) Levels of labeled succinate, malate, and

fumarate from palmitate tracing in hepa1-6 cells without (shSC) or with CLS deletion (shCLS)

(n=6 per group). (e–j) Levels of labeled pyruvate, acetyl-CoA, lactate, succinate, fumarate, and

citrate from glucose tracing in hepa1-6 cells without or with CLS deletion (n=6 per group).

Statistical significance was determined by unpaired, two-sided Student's t-test. Data represent

mean ± SEM. All measurements were taken from distinct samples.

Figure 5. CL deficiency promotes mitochondrial electron leak

(a) Representative electron microscopy images of liver fibrosis in control or CLS-LKO mice (red

arrows). (b) mRNA levels of fibrotic markers (Col1a1 and Desmin) in liver tissues (n=5–7 per

group). (c–f) Levels of cleaved caspase-3, cleaved caspase-7, mitochondrial cytochrome c, and

cytosolic cytochrome c in liver tissues (n=4–7 per group). (g) H_2O_2 emission in isolated liver

mitochondria stimulated with succinate, or succinate with auranofin and BCNU (n=3–4 per

group). (h) Schematic of small unilamellar vesicles (SUVs) containing cardiolipin (CL) or

phosphatidylcholine (PC) for mitochondrial enrichment. (i) H_2O_2 production in liver mitochondria

enriched with CL or PC SUVs (n=4 per group). All results are from mice fed standard chow.

Statistical significance was determined by 2-way ANOVA with within-row pairwise comparison

(b, g, i) and unpaired, two-sided Student's t-test (c–f). Data represent mean ± SEM. All

measurements were taken from distinct samples.

Figure 6. Influence of CL deficiency on respiratory supercomplex formation

(a–j) Representative western blots of respiratory supercomplexes in isolated liver mitochondria

from control or CLS-LKO mice, detected using antibodies for supercomplexes, monomers,

dimers, or oligomers, GRIM19/complex I (b), NDUFA9/complex I (d), SDHA2/complex II (f),

UQCRFS1/complex III (h), MTCO1/complex IV (j), and ATP5A/complex V (l). Complex II is not

thought to form respiratory supercomplexes. (c, e, g, i, k, m) Quantification of blots (n=4 per

group). All results are from mice fed standard chow. Statistical significance was determined by

2-way ANOVA with within-row pairwise comparison (first panels in c, e, i, k) and unpaired, two-

sided Student's t-test (second panels in c, e, g, i, k, and both panels in m). Data represent mean

± SEM. All measurements were taken from distinct samples.

Figure 7. CL deficiency disrupts coenzyme Q homeostasis in ~~mice and humans~~**humans**

and mice

(a) Schematic of site-specific electron leak. (b–e) Electron leak at sites I_Q, I_F, II_F, and III_{Q0} in liver

mitochondria from control or CLS-LKO mice (n=7 per group). (f) Oxidized CoQ (ubiquinone) can

be reduced to $CoQH_2$ (ubiquinol). (g, h) Oxidized and reduced CoQ levels in isolated liver

mitochondria from control or CLS-LKO mice (n=7 per group). (i) CoQ levels in liver mitochondria

from healthy human controls or patients with advanced steatohepatitis (n=10 and 16 per group).

(j) Pearson correlation of CL and CoQ levels in human liver samples ($R^2 = 0.64$). Data in (b–e,

g, h) are from mice fed standard chow. Statistical significance was determined by unpaired, two-

sided Student's t-test (b–e, panel of total levels in i) and 2-way ANOVA with within-row pairwise

comparison (g, h, i). Data represent mean ± SEM (b–e, g, h, i). All measurements were taken

from distinct samples.

Author Contributions

MJB and KF contributed to conceptualization, experimental design, investigation, formal

analysis, writing the original draft, and reviewing and editing the manuscript. GC, ADP, LW, and

RHC performed investigation and formal analysis. JAM, QJP, and JMJ were involved in

investigation, methodology, and formal analysis. AMP, TBB, JLS, SW, ERM, AMM, and STD

contributed to investigation. PS, FMF, and ZGH contributed to experimental design, with PS and

FMF also contributing to investigation, and ZGH contributing to resources and methodology. AP,

SAP, GLH, and PNM were involved in investigation and methodology, with SAP also

contributing to formal analysis. KEA and KJE contributed to investigation and resources. TST

and MP contributed to conceptualization, with TST also contributing to investigation. LSN, SMN,

DSL, and KHF-W contributed to methodology, with LSN also contributing to investigation. JLC

contributed to methodology and formal analysis. WLH and SAS contributed to experimental

design and resources. JEC contributed to methodology. All authors read and approved the

paper.

Corresponding Author

Correspondance to Katsuhiko Funai

E-mail: kfunai@health.utah.edu

Ethics Declarations

Competing interests: the authors declare no competing interests.

References

1. Younossi, Z.M.*, et al.* Global epidemiology of nonalcoholic fatty liver disease—Meta-analytic assessment of prevalence, incidence, and outcomes. *Hepatology* **64**, 73-84 (2015).
2. Miao, L.T., Giovanni; Byrne, Christopher D; Cao, Ying-Ying; Zheng, Ming-Hua. Current status and future trends of the global burden of MASLD. *Trends in Endocrinology & Metabolism* **29**, 1043-2760 (2024).
3. Saito, K.*, et al.* Characterization of hepatic lipid profiles in a mouse model with nonalcoholic seatohepatitis and subsequent fibrosis. *Scientific Reports* **5**(2015).
4. John, B.S.S. Nonalcoholic Steatohepatitis (NASH). in *StatPearls* (StatPearls Publishing, Treasure Island, 2023).
5. Vigano, L., Lleo, A. & Aghemo, A. Non-alcoholic fatty liver disease, non-alcoholic steatohepatitis, metabolic syndrome and hepatocellular carcinoma—a composite scenario. *Hepatobiliary Surgery and Nutrition* **7**, 130-133 (2018).
6. Fan, J.-G. & Cao, H.-X. Role of diet and nutritional management in non-alcoholic fatty liver disease. *Gastroenterology and Hepatology* **28**, 87-87 (2013).
7. Gordon, D.L.*, et al.* Biomarkers of NAFLD progression: a lipidomics approach to an epidemic. *Journal of Lipid Research* **56**, 722-736 (2015).
8. Fu, S.*, et al.* Aberrant lipid metabolism disrupts calcium homeostasis causing liver endoplasmic reticulum stress in obesity. *Nature* **473**, 528-531 (2011).
9. Simoes, I.C.M., Fontes, A., Pinton, P., Zischka, H. & Wieckowski, M.R. Mitochondria in non-alcoholic fatty liver disease. *The International Journal of Biochemistry & Cell Biology* **95**, 93-99 (2018).
10. Leveille, M. & Estall, J.L. Mitochondrial Dysfunction in the Transition from NASH to HCC. *Metabolites* **9**, 233 (2019).
11. Einer, C.*, et al.* Mitochondrial adaptation in steatotic mice. *Mitochondrion* **59**, 483-495 (2018).
12. Sunny, N.E., Bril, F. & Cusi, K. Mitochondrial Adaptation in Nonalcoholic Fatty Liver Disease: Novel Mechanisms and Treatment Strategies. *Trends in Endocrinology & Metabolism* **28**(2018).
13. Noureddin, M. & Sanyal, A.J. Pathogenesis of NASH: the Impact of Multiple Pathways. *Current Hepatology Reports* **17**, 350-360 (2018).
14. Pfeiffer, K.*, et al.* Cardiolipin stabilizes respiratory chain supercomplexes. *Metabolism and Bioenergetics* **278**, 52873-52880 (2003).
15. Johnson, J.M.*, et al.* Targeted overexpression of catalase to mitochondria does not prevent cardioskeletal myopathy in Barth syndrome. *Journal of Molecular and Cellular Cardiology* **121**, 94-102 (2018).
16. Houtkooper, R.H. & Vaz, F.M. Cardiolipin, the heart of mitochondrial metabolism. *Cell. Mol. Life Sci.* **65**, 2493-2506 (2008).
17. Schlame ab, M. & Hostetler, K.Y. Cardiolipin synthase from mammalian mitochondria. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* **1348**, 207-213 (1997).
18. Paradies, G., Paradies, V., Ruggiero, F.M. & Petrosillo, G. Role of Cardiolipin in Mitochondrial Function and Dynamics in Health and Disease: Molecular and Pharmacological Aspects. *Cells* **8**, 728 (2019).
19. Funai, K., Summers, S.A. & Rutter, J. Reign in the membrane: How common lipids govern mitochondrial function. *Current Opinion in Cell Biology* **63**, 162-173 (2020).
20. Jussupow, A., Luca, A.D. & Kaila, V.R. How cardiolipin modulates the dynamics of respiratory complex I. *Science Advances* **5**(2019).

21. Schwall, C.T., Greenwood, V.L. & Alder, N.N. The stability and activity of respiratory
Complex II is cardiolipin-dependent. *Biochimica et Biophysica Acta (BBA) - Molecular*
and Cell Biology of Lipids **1817**, 1588-1596 (2012).
22. Bazan, S*., et al.* Cardiolipin-dependent Reconstitution of Respiratory Supercomplexes
from Purified Saccharomyces cerevisiae Complexes III and IV*. **288**, 401-411 (2013).
23. Dudek, J*., et al.* Cardiolipin deficiency affects respiratory chain function and organization
in an induced pluripotent stem cell model of Barth syndrome. *Stem Cell Research* **11**,
806-819 (2013).
24. Sustarsic, E.G*., et al.* Cardiolipin Synthesis in Brown and Beige Fat Mitochondria Is
Essential for Systemic Energy Homeostasis. *Cell Metab* **28**, 159-174 (2018).
25. Jiang, F*., et al.* Absence of cardiolipin in the crd1 null mutant results in decreased
mitochondrial membrane potential and reduced mitochondrial function. *Journal of*
Biological Chemistry **275**, 22387-22394 (2000).
26. Xu, Y., Sutachan, J.J., Plesken, H., Kelley, R.I. & Schlame, M. Characterization of
lymphoblast mitochondria from patients with Barth syndrome. *Laboratory Investigation; a*
journal of technical methods and pathology **85**, 823-830 (2005).
27. Hsu, P*., et al.* Cardiolipin remodeling by TAZ/tafazzin is selectively required for the
initiation of mitophagy. *Autophagy* **11**, 643-652 (2015).
28. Le, C.H*., et al.* Tafazzin deficiency impairs CoA-dependent oxidative metabolism in
cardiac mitochondria. *Journal of Biological Chemistry* **295**, 12485-12497 (2020).
29. Heden, T.D*., et al.* Mitochondrial PE potentiates respiratory enzymes to amplify skeletal
muscle aerobic capacity. *Science Advances* **5**, 8352 (2019).
30. Johnson, J.M.P., Alek D; Balderas, Enrique; Sustarsic, Elahu G; Maschek, J Alan;
Brothwell, Marisa J; Alejandro, Jara-Ramos; Panic, Vanja; Morgan, Jeffrey T; Villanueva,
Claudio J; Sanchez, Alejandro; Rutter, Jared; Lodhi, Irfan J; Cox, James E; Fisher-
Wellman, Kelsey H; Chaudhuri, Dipayan; Gerhart-Hines, Zachary; Funai, Katsuhiko.
Mitochondrial phosphatidylethanolamine modulates UCP1 to promote brown adipose
thermogenesis. *Science Advances* **9**, 7864 (2023).
31. Kartsoli, S., Kostara, C.E., Tsimihodimos, V., Bbairaktari, E.T. & Christodoulou, D.K.
Lipidomics in non-alcoholic fatty liver disease. *World Journal of Hepatology* **12**, 436-450
(2020).
32. Ehud Zigmond, C.V. Two Roads Diverge in the Sick Liver, Monocytes Travel Both. *Cell*
Press **10**, 1016 (2020).
33. Zahao, X.-A.C., Guang-Mei; Liu, Yong; Chen, Yu-Xin; Wu, Hong-Yan; Chen, Jin; Xiong,
Ya-Li; Tian, Chen; Wang, Gui-Yang; Jia, Bei; Xia, Juan; Wang, Jian; Yan, Xiao-Min;
Zhang, Zhao-Ping; Huang, Rui; Wu, Chao. Inhibitory effect of silymarin on CCl4-induced
liver fibrosis by reducing Ly6Chi monocytes infiltration. *International Journal of Clinical &*
Experimental Pathology **10**, 11941-11951 (2017).
34. Daemen, S.G., Anastasiia; Kalugotla, Gowri; He, Li; Chan, Mandy M.; Beals, Joseph W.;
Liss, Kim H.; Klein, Samuel; Feldstein, Ariel E.; Finck, Brian N.; Artyomov, Maxim N.;
Schilling, Joel D. Dynamic Shifts in the Composition of Resident and Recruited
Macrophages Influence Tissue Remodeling in NASH. *Cell Reports* **34**, 108626 (2021).
35. Liyanage, S.E.G., Peter J.; Ribeiro, Joana; Cristante, Enrico; Sampson, Robert D.;
Luhmann, Ulrich F.O.; Ali, Robin R.; Bainbridge, James W. Flow cytometric analysis of
inflammatory and resident myeloid populations in mouse ocular inflammatory models.
Experimental Eye Research **151**, 160-170 (2016).
36. Fry, M. & Green, D.E. Cardiolipin requirement for electron transfer in complex I and III of
the mitochondrial respiratory chain. *Journal of Biological Chemistry* **256**, 1874-1880
(1981).

37. Paradies, G., Petrosillo, G., Pistolese, M. & Ruggiero, F.M. Reactive oxygen species
affect mitochondrial electron transport complex I activity through oxidative cardiolipin
damage. *Gene* **286**, 135-141 (2002).
38. Lange, C., Nett, J.H., Trumpower, B.L. & Hunte, C. Specific roles of protein–
phospholipid interactions in the yeast cytochrome bc1 complex structure. *The EMBO*
Journal **20**, 6591-6600 (2001).
39. Robinson, N.C. Functional binding of cardiolipin to cytochrome c oxidase. *Journal of*
Bioenergetics and Biomembranes **25**, 153-163 (1993).
40. Koliaki, C.*, et al.* Adaptation of hepatic mitochondrial function in humans with non-
alcoholic fatty liver is lost in steatohepatitis. *Cell Metab* **21**, 739-746 (2015).
41. Siripoksup, P.*, et al.* Sedentary behavior in mice induces metabolic inflexibility by
suppressing skeletal muscle pyruvate metabolism. *J Clin Invest* **134**(2024).
42. Drose, S. & Brandt, U. The Mechanism of Mitochondrial Superoxide Production by the
Cytochrome bc1 Complex. *Journal of Biological Chemistry* **283**, 21649-21654 (2008).
43. Marra, F. & Svegliati-Baroni, G. Lipotoxity and the gut-liver axis in NASH pathogenesis.
Hepatology **68**, 280-295 (2018).
44. Buzzetti, E., Pinzani, M. & Tsochatzis, E.A. The multiple-hit pathogenesis of non-
alcoholic fatty liver disease (NAFLD). *Metabolism* **65**, 1038-1048 (2016).
45. Turkseven, S.*, et al.* Mitochondria-targeted antioxidant mitoquinone attenuates liver
inflammation and fibrosis in cirrhotic rats. *American Physiological Society* **318**, 298-304
(2020).
46. Karakuzu, O.C., Melissa R.; Liu, Yi; Garsin, Danielle A. Amplex Red Assay for
Measuring Hydrogen Peroxide Production from Caenorhabditis elegans. *Bio-protocol* **9**,
3409 (2019).
47. Prola, A.B., Jordan; Vandestienne, Aymeline; Piquereau, Jerome; Denis, Raphael GP;
Guyot, Stephane; Chauvin, Hadrien; Mourier, Arnaud; Maurer, Marie; Henry, Celine;
Khadhraoui, Nahed; Gallerne, Cindy; Molinie, Thibaut; Courtin, Guillaume; Gressette,
Melanie; Solgadi, Audrey; Dumont, Florent; Castel, Julien; Ternacle, Julien; Demarquoy,
Jean; Malgoyre, Alexandra; Koulmann, Nathalie; Derumeaux, Genevieve; Giraud, Marie-
France; Joubert, Frederic; Veksler, Vladimir; Luquet, Serve; Relaix, Frederic; Tiret,
Laurent; Pilot-Storck, Fanny. Cardiolipin content controls mitochondrial coupling and
energetic efficiency in muscle. *Science Advances* **7**(2021).
48. Mileykovskaya, E. & Dowhan, W. Cardiolipin-Dependent Formation of Mitochondrial
Respiratory Supercomplexes. *Chemistry and Physics of Lipids* **179**, 42-48 (2014).
49. Pennington, E.R., Funai, K., Brown, D.A. & Shaikh, S.R. The role of cardiolipin
concentration and acyl chain composition on mitochondrial inner membrane molecular
organization and function. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell*
Biology of Lipids **1864**, 1039-1052 (2019).
50. Arnarez, C., Marrink, S.J. & Periole, X. Molecular mechanism of cardiolipin-mediated
assembly of respiratory chain supercomplexes. *Chemical Science* (2016).
51. Bottinger, L.*, et al.* Phosphatidylethanolamine and Cardiolipin Differentially Affect the
Stability of Mitochondrial Respiratory Chain Supercomplexes. *Journal of Molecular*
Biology **5**, 677-686 (2012).
52. Dornas, W. & Schuppan, D. Mitochondrial oxidative injury: a key player in nonalcoholic
fatty liver disease. *Gastrointestinal and Liver Physiology* **319**, G400-G411 (2020).
53. Gomez, B. & Robinson, N.C. Quantitative Determination of Cardiolipin in Mitochondrial
Electron Transferring Complexes by Silicic Acid High-Performance Liquid
Chromatography. *Analytical Biochemistry* **267**, 212-216 (1999).
54. Brand, M.D. Mitochondrial generation of superoxide and hydrogen peroxide as the
source of mitochondrial redox signaling. *Free Radical Biology and Medicine* **100**, 14-31
(2016).

55. Decker, S.T.F., Katsuhiko. Mitochondrial membrane lipids in the regulation of
bioenergetic flux. *Cell Metab* **36**, 1963-1978 (2024).
56. Bessone, F., Razori, M.V. & Roma, M.G. Molecular pathways of nonalcoholic fatty liver
disease development and progression. *Cell. Mol. Life Sci.* **76**, 99-128 (2019).
57. Ines C. M. Simoes, R.A., Jose Teixeira, Agnieszka Karkucinska-Wieckowska, Adriana
Carvalho, Susana P. Pereira, Rui F. Simoes, Sylwia Szymanska, Michal Dabrowski,
Justyna Janikiewicz, Agnieszka Dobrzyn, Paulo J. Oliveira, Yaiza Potes, Mariusz R.
Wieckowski. The Alterations of Mitochondrial Function during NAFLD Progression-- An
Independent Effect of Mitochondrial ROS Production. *International Journal of Molecular
Sciences* **22**, 6848 (2021).
58. Simoes, I.C., Fontes, A., Pinton, P., Zischka, H. & Wieckowski, M.R. Mitochondria in
non-alcoholic fatty liver disease. *The International Journal of Biochemistry & Cell Biology*
95, 93-99 (2017).
59. Goncalves, R.L.S., Schlame, M., Bartelt, A.L., Brand, M.D. & Hotamisligil, G.S.
Cardiolipin deficiency in Barth syndrome is not associated with increased
superoxide/H2O2 production in heart and skeletal muscle mitochondria. *FEBS Letters*
595, 415-432 (2021).
60. Xu, Y.*, et al.* Loss of protein association causes cardiolipin degradation in Barth
syndrome. *Nature Chemical Biology* **12**, 641-647 (2016).
61. Simoes, I.C.A., Ricardo; Teixeira, Jose; Karkucinska-Wieckowska, Agnieszka; Carvalho,
Adriana; Pereira, Susana P.; Simoes, Rui F.; Szymanska Sylwia; Dabrowski, Michal;
Janikiewicz, Justyna; Dobrzyn, Agnieszka; Oliveira, Paulo J.; Potes, Yaiza; Wieckowski,
Mariusz R. . The Alterations of Mitochondrial Function during NAFLD Progression-- An
Independent Effect of Mitochondrial ROS Production. *International Journal of Molecular
Sciences* **22**, 6848 (2021).
62. Bazan, S.*, et al.* Cardiolipin-dependent Reconstitution of Respiratory Supercomplexes
from Purified Saccharomyces cerevisiae Complexes III and IV*. *Journal of Biological
Chemistry* **288**, 401-411 (2013).
63. Miwa, S.*, et al.* Carboxylesterase converts Amplex red to resorufin: Implications for
mitochondrial H2O2 release assays. *Free Radical Biology and Medicine* **90**(2016).

1094

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7

Figure 1– figure supplement 1

Figure 1– figure supplement 2

Figure 2– figure supplement 1

Figure 2– figure supplement 2

High-fat Diet

A H&E — Ctrl / CLS-LKO — 50 μm / 50 μm

B Masson's Trichome — Ctrl / CLS-LKO — 50 μm / 50 μm

C Volcano plot: -Log10 (p Adjusted Value) vs Log2 Fold Change; Vmn2r3, Tiam2, Knstrn, Dlgap1, Nupr1, Tenm4, Gpnmb; 1026 / 713

D Normalized Enrichment Score

- Glyoxylate metabolism and glycine degradation
- Complex I biogenesis
- Branched chain amino acid catabolism
- Respiratory electron transport ATP synthesis
- Citric acid TCA cycle and respiratory electron transport
- Respiratory electron transport
- Metabolism of amino acids and derivatives
- Golgi to ER retrograde transport
- Keratinization
- RMTs methylate histone arginines
- TP53 transcription of cell cycle genes
- Lymphoid and non-lymphoid immunoregulatory interactions
- Signaling by PDGF
- Mitotic prometaphase
- Megakaryocyte differentiation and platelet function
- NCAM signaling for neurite out-growth
- Megakaryocyte development and platelet production
- PRC2 methylates histones and DNA
- Rho GTPase effectors
- COPI-dependent golgi to ER retrograde traffic
- Chromosome maintenance
- O-glycosylation of TSR domain containing proteins
- Non-integrin membrane ECM interactions
- Collagen chain trimerization
- Mitotic spindle checkpoint
- Molecules associated with elastic fibres
- Deposition of new CENPA containing nucleosomes
- Condensation of prophase chromosomes
- ECM proteoglycans
- Amplification of signal from the kinetochores
- Elastic fibre formation
- Assembly of collagen fibrils
- Kinesins
- Resolution of sister chromatid cohesion
- Rho GTPases activate formins
- Integrin cell surface interactions
- Collagen biosynthesis and modifying enzymes
- Collagen degradation
- Collagen formation
- Extracellular matrix organization
- Degradation of the extracellular matrix

E Flow cytometry gating: SSC-A vs FSC-A; FSC-H vs FSC-A; Comp-APC-Cy7-A :: Live_Dea vs FSC-A (Live Cells 91.8); Comp-FITC-A :: CD45 vs FSC-A (CD45); Comp-BV421-A :: CD11b vs Comp-APC-A :: F4 80 (CD11b); Comp-BV786-A :: CD11c vs Comp-BV605-A :: MHC-II; Comp-PE-A :: Ly6C vs Comp-PE-Cy7-A :: Ly6G (Ly6Chi, Ly6Clow, PMN); Comp-PE-A :: Ly6C vs Comp-BV605-A :: MHC-II (Inflammatory monocytes); Comp-PE-A :: Ly6C vs Comp-BV605-A :: MHC-II; Comp-BV786-A :: CD11c vs Comp-APC-A :: F4 80 (cDC2, F4/80)

Figure 3– figure supplement 1

Figure 4– figure supplement 1

A Glycine M+2   **B** Aspartate M+2   **C** 3-Phosphoglyceric acid M+3   **D** Alanine M+3   **E** Malate M+2

Figure 5– figure supplement 1

Figure 7– figure supplement 1

Figure 1– table supplement 1

Patient demographic information

	Healthy	MASH
Age at time of collection	50.3 ± 9.6 yrs	62.2 ± 7.1 yrs
Sex	Male: 1 Female: 10	Male: 9 Female: 8
Alcohol use?	N/A	Yes: 0 No: 17
Race	White: 9 African American: 1 Asian: 1	White: 10 Unknown: 7

Supplemental Figure Legends

Figure 1 – figure supplement 1. Mitochondrial phospholipidome from Figure 1N&O. (A-H)

Abundance of mitochondrial lipids (CL, PC, PE, PI, PS, LPC, PG, LPE) in livers of mice fed

standard chow or HFD for 16 weeks (n=5 per group). (I-P) Abundance of mitochondrial lipids

(CL, PC, PE, PI, PS, LPC, PG, LPE) in livers of wildtype or leptin-deficient mice, 30 weeks old

(n=6 per group). Statistical significance was determined by 2-way ANOVA with within-row

pairwise comparison. Data represent mean ± SEM. All measurements were taken from distinct

samples.

Figure 1 – figure supplement 2. Mitochondrial phospholipidome from Figure 1P&Q. (A-H)

Abundance of mitochondrial lipids (CL, PC, PE, PI, PS, LPC, PG, LPE) in livers of mice fed

standard chow or Gubra-Amylin MASH diet for 30 weeks (n=6 per group). (I-P) Abundance of

mitochondrial lipids (CL, PC, PE, PI, PS, LPC, PG, LPE) in livers of mice injected with corn oil or

carbon tetrachloride for 10 weeks (n=5–7 per group). Statistical significance was determined by

2-way ANOVA with within-row pairwise comparison. Data represent mean ± SEM. All

measurements were taken from distinct samples.

Figure 2– figure supplement 1. Mitochondrial phospholipidome from standard chow or

high-fat diet fed control and CLS-LKO livers. (A-G) Abundance of mitochondrial lipids (PC,

PE, PI, PS, LPC, PG, LPE) in livers of control or CLS-LKO mice fed standard chow, 8 weeks old

(n=5–6 per group). (H-O) Abundance of mitochondrial lipids (CL, PC, PE, PI, PS, LPC, PG,

LPE) in livers of control or CLS-LKO mice fed a high-fat diet for 8 weeks (n=11–12 per group).

Statistical significance was determined by 2-way ANOVA with within-row pairwise comparison.

Data represent mean ± SEM. All measurements were taken from distinct samples.

Figure 2– figure supplement 2. Additional histological and transcriptomic data from

control and CLS-LKO mice. (A) Representative H&E staining of livers from control or CLS-

LKO mice fed a high-fat diet (HFD) for 8 weeks. (B) Representative Masson's Trichrome

staining of livers from control or CLS-LKO mice fed a HFD for 8 weeks. (C) Volcano plot of

differentially expressed genes in livers from control or CLS-LKO mice fed standard chow (n=5–7

per group). (D) Pathway analysis of transcriptomic data in livers from control or CLS-LKO mice

fed standard chow (n=5–7 per group). (E) Gating strategy used for flow cytometry experiments

in control and CLS-LKO mice. Pathway analysis was performed using the Reactome Pathway

Database. All measurements were taken from distinct samples.

Figure 3– figure supplement 1. Additional metabolic and mitochondrial phenotyping data

with CLS deletion. (A) mRNA levels of lipogenic genes in livers from control or CLS-LKO mice

fed standard chow (n=6–7 per group). (B) Serum triglycerides in control or CLS-LKO mice fed

standard chow (n=6–7 per group). (C, D) JO_2 consumption in isolated liver mitochondria from

control or CLS-LKO mice fed a high-fat diet (HFD) for 8 weeks in response to substrates: (C)

malate, pyruvate, ADP, succinate, and FCCP (n=11–12 per group); (D) palmitoyl-carnitine,

malate, and ADP (n=7 per group). Statistical significance was determined by 2-way ANOVA

with within-row pairwise comparison (A, C, D) and unpaired Student's t-test (B). Data represent

mean ± SEM. All measurements were taken from distinct samples.

Figure 4– figure supplement 1. Additional fluxomic phenotyping data with CLS deletion.

(A-D) Levels of labeled metabolites (glycine, aspartate, 3-phosphoglyceric acid, alanine, and

malate) from glucose tracing in Hepa1-6 cells without (shSC) or with CLS deletion (shSLC) (n=6

per group). Statistical significance was determined by unpaired Student's t-test. Data represent

mean ± SEM. All measurements were taken from distinct samples.

Figure 5– figure supplement 1. Additional mitochondrial phenotyping data with CLS

deletion. (A, B) Levels of mitochondrial and cytosolic cytochrome c in livers of control or CLS-

LKO mice fed a high-fat diet (HFD) for 8 weeks (n=6 per group). (C, D) JH_2O_2 emission and

production in isolated liver mitochondria from (C) control or CLS-LKO mice fed a HFD for 8

weeks, stimulated with succinate or succinate plus auranofin and BCNU (n=9–8 per group), and

(D) Hepa1-6 cells without (shSC) or with CLS deletion (shCLS), stimulated with succinate or

succinate plus auranofin and BCNU (n=3 per group). (E) Electron leak of liver mitochondria from

control mice fused with SUVs (n=4 per group). Statistical significance was determined by 2-way

ANOVA with within-row pairwise comparison (C, D, E) and unpaired Student's t-test (A, B). All

measurements were taken from distinct samples.

Figure 7– figure supplement 1. Additional data on coenzyme Q (A-C) Total CoQ_8, CoQ_9,

and CoQ_{10} levels in whole liver tissue from control and CLS-LKO mice (n=7 per group). (D-I)

Oxidized and reduced CoQ_8, CoQ_9, and CoQ_{10} levels in whole liver tissue from control and CLS-

LKO mice (n=7 per group). (J-O) Oxidized and reduced CoQ_8, CoQ_9, and CoQ_{10} levels in

isolated mitochondria from control and CLS-LKO mice (n=7 per group). Statistical significance

was determined by unpaired Student's t-test. Data represent mean ± SEM. All measurements

were taken from distinct samples.