A sensory approach to turbidity: How sources and levels shape aquatic light environments and fish visual ecology

Anna Garcia[1]*, Adelaide Sibeaux[1], Theresa Burt de Perera[1], Cait Newport[1]

[1]Department of Biology, University of Oxford, Oxford, United Kingdom

*Corresponding author: anna.garcia.connect@gmail.com

Abstract

Turbidity is a ubiquitous source of sensory pollution that is likely to impact the appearance of the visual stimuli that animals rely on for survival and reproduction. Understanding how different turbidity sources impact the appearance of the ambient light environment is the foundational first step towards predicting whether and how animals will cope with the global increases in the severity and frequency of high turbidity events caused by anthropogenic disturbance. Here, we measured how four common turbidity sources (algae, bentonite, calcium carbonate, and kaolin), and variable turbidity levels, changed the appearance of the ambient light environment. We measured total number of photons (luminance), hue, chroma, and image contrast, and we evaluated the effect of each turbidity source and level on settling rate, pH, and KH. Both turbidity source and turbidity level impacted the appearance of the ambient light environment. With increasing turbidity level, calcium carbonate and kaolin increased luminance while algae decreased luminance, bentonite caused the greatest change in hue, and algae caused the greatest change in chroma. This demonstrates that the impacts of different turbidity sources on the ambient light environment are not uniform, giving a potential explanation for the discrepancies between studies on the effects of turbidity on fish behaviour. Consideration of the effect of specific turbidity sources on ambient light is crucial for the design of experiments that seek to investigate how changes in turbidity impact the perception of important visual information, which underpins the survival and reproductive success of aquatic organisms around the world.

29   **Keywords**: sensory pollution, vision, teleost, algae, suspended sediment, behaviour

Summary Statement:

31   Different turbidity sources and levels uniquely affect aquatic light environments and fish visual
32   perception, emphasizing the need for these factors when interpreting fish behaviour and the
33   ecological consequences of turbidity.
34

Introduction

Turbidity refers to organic or inorganic particles in water which can alter both water clarity and
quality. While the impacts of turbidity on aquatic ecosystems are likely to be wide ranging, its most
visible impact is on the light environment. Particles in the water scatter and absorb light and can
significantly alter the appearance of visual features, potentially making visual signals more difficult
to detect or identify. Turbidity levels therefore play an important role in shaping the visual
environment of aquatic species. Turbidity occurs naturally to some degree in all aquatic
ecosystems and fluctuates over both short and long-time scales, suggesting that visually reliant
animals likely have some mechanisms to cope with the associated changing visual conditions.
However, natural turbidity conditions are being increasingly altered through human activities
(Burke, Reytar et al. 2011, Seers and Shears 2015, Cartwright, Fearns et al. 2021), changing
both the concentration of turbidity as well as the source. For example, more frequent flooding in
Australia is causing plumes of turbidity that are visible from space to drain onto the Great Barrier
Reef (Malthus, Lehmann et al. 2019), and sewage and chemical discharges in British rivers and
coasts are turning normally clear waters to an opaque brown (Bojko, Lipp et al. 2020, Love,
Crooks et al. 2020, Albini, Lester et al. 2023). For example, more frequent flooding in Australia is
causing plumes of turbidity that are visible from space to drain onto the Great Barrier Reef
(Malthus, Lehmann et al. 2019) and sewage and chemical discharges in British rivers and coasts
are turning normally clear waters to an opaque brown (Bojko, Lipp et al. 2020, Love, Crooks et
al. 2020, Albini, Lester et al. 2023). It is not just turbidity levels that are changing; in some
ecosystems, the sources of turbidity have also shifted, leading to the presences of multiple
sources in some ecosystems. For example, in addition to the inorganic turbidity associated with
coastal run-off, the Great Barrier Reef is likely to experience an increase in organic turbidity as
bleaching events can lead to eventual algae blooms (Fukunaga, Burns et al. 2022). In the UK,

nutrient runoff from intensive agricultural practices can accelerate the export of carbon and other
organic nutrients resulting in algal blooms (Glendell and Brazier 2014). There is some evidence
that turbidity can negatively impact fish. For example, studies have shown that turbidity can alter
species richness of coral reef fish (Moustaka, Langlois et al. 2018) and disrupt fish community
structures in shallow lakes (Hayes, Rutledge et al. 1992). Additionally, turbidity has also been
linked to increased gill epithelium thickness in clownfish larvae (Amphiprion percula) (Hess,
Wenger et al. 2015), reduced foraging success and growth rate in smallmouth bass (Micropterus
dolomieu) (Sweka and Hartman 2011), decreased survival of shed larvae (Alosa sapidissima)
(Auld and Schubel 1978), and altered metabolism in rainbow trout (Berli, Gilbert et al. 2014).
Furthermore, turbidity can contribute to eutrophication and anoxia (Smith 2003, Budria and
Candolin 2015, Cumming and Herbert 2016), impairment of metabolic reduced physiological
condition (Budria and Candolin 2015, Hasenbein, Fangue et al. 2016), and negatively affect
spawning success (Burkhead and Jelks 2001, Sutherland and Meyer 2007). While it seems likely
that the effects of turbidity could be wide-ranging, its impact on visually-guided fish behaviour has
received less attention, leaving it unclear whether turbidity presents a serious threat to fish
populations.
75
Turbidity has been shown to significantly alter some visually-guided behaviours in fish, including
foraging (Lunt and Smee 2015) and searching (Newport, Padget et al. 2021), collective behaviour
(Chamberlain and Ioannou 2019), predator-prey interactions (Ortega, Figueiredo et al. 2020), and
swimming rates (Hildebrandt and Parsons 2016). For guppies (*Poecilia reticulata*), increased
turbidity causes individuals to be less active, more solitary (Borner, Krause et al. 2015), and more
vulnerable to predation (Kimbell and Morrell 2015). But some species acclimate to changing visual
conditions; the same guppy species raised in turbid water, display a shift in spectral sensitivity
and an increase in activity (Ehlman, Sandkam et al. 2015). African cichlids (*Pseudocrenilabrus*
multicolour) reared in turbid conditions exhibited an ontogenetic shift in the relative size of their
eyes and brains, with young fish raised in turbid water developing relatively larger eyes, and older
fish developing relatively larger brains compared to those raised in clear water (Tiarks, Gray et
al. 2024). Studies exploring the impacts of turbidity on the expression of sexually selected traits
have revealed complex effects on both sexes, making it difficult to predict the outcomes of such
interactions (Järvenpää and Lindström 2004, Järvenpää and Lindström 2011, Sundin, Rosenqvist
et al. 2016, Sundin, Aronsen et al. 2017, Järvenpää, Pauli et al. 2019). For example, turbidity
interferes with the visual inspection of conspicuous red nuptial coloration by female three-spined
sticklebacks (*Gasterosteus aculeatus*), altering both female mate choice preference, and male

courtship behaviour (Wong, Candolin et al. 2007, Tuomainen and Candolin 2011, Candolin and
Jensen 2021). However, there is also evidence that turbidity can have no impact on sexual
selection behaviour, as shown studies on yellow perch (*Perca flavescens*), wreakfish (*Cynoscion*
regalis), white perch (*Morone americana*), and tessellated darter (*Etheostoma olmstedi*), (Grecay
and Targett 1996, Shoji, North et al. 2005, Pangle, Malinich et al. 2012, Kellogg and Leipzig-Scott
2017). In sum, the effects of turbidity on fish are neither unidirectional nor easily predicted, and
the evidence suggests that not all fish species or behaviours are equally affected by turbidity
(Rodrigues, Ortega et al. 2023).

The disparity in the types and magnitude of effects observed between studies may also arise from
methodological differences, particularly the lack of consideration for how specific turbidity
conditions alter the light environment and, consequently the transmission of visual information
that experimenters aim to manipulate. Differences in turbidity particle size, shape, composition,
and concentration influence the scattering of light. Small particles whose radii are less than 10%
of the wavelength of light obey Rayleigh scattering. In this case, the scattering of light is strongly
wavelength and intensity-dependent, and shorter wavelengths (e.g. violet) are scattered more
than longer wavelengths (e.g. red) (Johnsen 2012). Larger particles that are comparable to or
larger, than the wavelength of light, follow Mie scattering which is relatively wavelength-
independent (Bohren and Huffman 2008). Both forms of scattering are dependent on the total
light intensity (i.e. number of photons available) whereby smaller particles scatter more light than
larger particles. Moreover, light absorption is highly dependent on the properties of the turbidity
source. For example, photopigments like chlorophylls in algal blooms selectively absorb light in
the 440-475 nm and 630-675 nm ranges (Bricaud, Claustre et al. 2004). On the other hand,
mineral and sediment-derived turbidity sources absorb in the shorter 400-550 nm wavelength
range (Babin, Morel et al. 2003). Despite the source-dependent effects of turbidity on the
transmission of important visual information, previous research seldom considers how different
types of turbidity may influence experimental outcomes (but see (Radke and Gaupisch 2005, Li,
Zhang et al. 2013, Cano-Rocabayera, Vargas-Amengual et al. 2020, Figueiredo, Granzotti et al.
2020)).

To understand how different sources and levels of turbidity have been addressed in the literature,
we conducted a review of existing studies. We identified 200 papers that explore the impact of
turbidity on a range of fish behaviours including anti-predator, foraging efficiency, movement,
sexual selection, social/collective behaviour (Fig. 1). In this review, we also included papers

assessing the impact of different turbidity sources on fish physiological conditions and biodiversity

as these parameters are directly related to fish behavioural ecology. See supplementary materials

S1 for details of the literature search process. We found that the turbidity sources used in

experiments can be broadly categorised into eight groups (Fig. 1) and that natural sediment,

bentonite clay, and algae/phytoplankton were the most used turbidity sources. We also found that

the use of a particular turbidity source is not field- or species-specific. For example, the studies

examining the impacts of turbidity on foraging efficiency, used all eight of our turbidity groups.

Similarly, different studies involving the same species, Trinidadian guppies (*Poecilia reticulata),*

used different turbidity source and concentrations. While Borner, Krause et al. (2015) explored

the effects of turbidity on social dynamics using Kaolin, a white clay with a fine particle size (>1

µm), at concentrations of up to 1000 Nephelometric Turbidity Units (NTU), Kimbell and Morrell

(2015) tested behavioural responses to predation using Bentonite, a typically reddish clay with

even smaller particles (>1 µm), at levels of up to 200 NTU. Interestingly, the two groups cite very

different reasons for their choice of turbidity source. While Borner, Krause et al. (2015) made their

selection based on practical considerations (kaolin was thought to stay suspended for longer),

Kimbell and Morrell (2015) chose a source that was ecologically relevant. While both reasons are

valid, surprisingly neither of author considered how each turbidity source would alter the

appearance or detectability of conspecifics or predators, which was at the heart of the behaviours

being tested in their studies. This is a common weakness of many studies on this topic including

our own (Newport, Padget et al. 2021).

Figure 1 - The frequency of use of different turbidity sources in 200 scientific papers exploring the impact
of turbidity on fish behaviour, physiological condition, and biodiversity. Papers are grouped by turbidity
source. "Clay" and "natural sediment" are often sourced directly from the study site, with their specific
composition unknown. More specific turbidity sources, such as bentonite, were grouped separately. Within
each turbidity source, papers are categorized by research focus. Studies using multiple turbidity sources or
response categories were counted separately for each.

The aim of this paper is to describe how different sources and levels of turbidity impact the
ambient light environment and conspicuousness of visual features, as well as some practical
considerations associated with using turbidity in laboratory behavioural experiments. We hope
this will inspire other researchers to consider turbidity sources carefully in their experimental
design and empower them to select the appropriate turbidity sources based on their experimental
investigation and design.

The ability of an animal to perceive visual features depends not only on its visual system, but also
on four additional optical factors that determine the light available to its eyes: 1) the light
environment, 2) the amount of light reflected by the stimulus, 3) the light transmitted between the
feature and the eye , 4) and the degree of veiling light (light scattered from particles directly into
the observer's eye) (Endler 1990). These four factors determine the amount of light that reaches
the eye and marks the beginning of the visual processing pathway. This study focuses on how
turbidity affects the first factor – the light environment – aiming to provide an in-depth
understanding of how turbidity influences the appearance of visual features within the
environment. Absolute irradiance is commonly used to describe the light environment. It is a
measure of the number of photons per wavelength within a light environment and is visualized as
a spectral curve. Changes to the shape of this curve indicate changes in the light environment.
For example, shifts in the position of the curve's peaks indicate changes in colour composition,
while flattening peaks shows reduced light availability and lower spectral saturation. The shape
of the curve can be described using three characteristics: luminance, hue, and chroma (Endler
1990). Luminance is calculated from the area under the spectral curve, hue is determined by the
wavelength range at which the number of photons peaks, and chroma represents the difference
between the lowest and highest peaks across all wavelengths (see Endler 1990 for mathematical
formulas). These characteristics describe how the available light will influence the appearance of
visual features within the light environment and are independent any species' specific visual
system. The specific perception and interpretation of visual signals will depend on the visual

system of individual species, but the raw optical input dictates what is possible and serves as a

basis for visual modelling. Therefore, this study provides a critical foundation for predicting how

turbidity might alter the appearance of visual stimuli by examining the effects of different turbidity

levels and sources on the light environment.

Methods:

We tested how four sources of turbidity commonly used in behavioural experiments – algae,

bentonite, kaolin, calcium carbonate – alters the light environment. We specifically measured

changes in absolute irradiance, and image contrast. Changes to luminance, hue, and chroma

were calculated from absolute irradiance measurements. We also measured turbidity settling rate,

as well as changes in pH and water hardness (calcium carbonate concentration, KH) over time to

assess the usability of each turbidity source in behavioural experiments with fish. The latter two

were measurements help determine whether dissolving particles significantly alter water

chemistry, potentially exceeding the healthy limits for fish. We tested turbidity levels between 0

and 100 Nephelometric Turbidity Units (NTU) which reflect those commonly found in nature

(Cyrus and Blaber 1992, Miranda 2011, Li, Zhang et al. 2013, Macdonald, Ridd et al. 2013,

Ehlman, Sandkam et al. 2015) and used in experiments (Ajemian, Sohel et al. 2015, Borner,

Krause et al. 2015, Kimbell and Morrell 2015, Newport, Padget et al. 2021).

General Apparatus:

Experiments were conducted in a temperature-controlled laboratory (21.2 ± 0.587 °C; mean ±

standard deviation) with artificial fluorescent lighting that followed a 12-hour light/dark cycle

(Clean A-MP LED6000-840 M600Q LDO, total power 46.3 W LED, Zumtobel). A glass tank with

dimensions 84 cm x 30 cm x 57.5 cm (l x h x w) (Fig. 2) was filled to a height of 28 cm (volume =

135 litres). RO water was used to provide a standardised, replicable mineral composition of the

water. Opaque white PVC sheets were secured outside the tank against the walls and under the

base to standardize the source of incoming light. A tripod was secured inside the tank to the

bottom to hold either 1) an optic fibre cable at 15 cm from the bottom of the tank allowing to

measure total photons and spectral composition of the light in the water, or 2) a GoPro (HERO8

Black, wide lens, photo, standard output, zoom x1. GoPro, Inc.) for image contrast

measurements. The GoPro was attached to the tripod facing the image standard, with its lens at

a height of 15 cm.

Directly across from the tripod, a laminated (Office Depot laminating pouches, matt, 75 μm) image
standard printed on paper (Canon, Matte Photo Paper), was attached to the back wall with Blu
Tack (Bostick Smart Adhesives) (see Supplementary Fig. 1, S2 for an illustration of the
apparatus). We included an image standard to ensure that colour was present in the tank, allowing
us to measure changes in colour caused by turbidity. The image standard was designed with
colours selected based on the cone sensitivity maxima of a coral reef fish species *(Rhinecanthus*
aculeatus) and goldfish, ensuring the colours are visible to two species that inhabit very different
visual environments. The orange (Hex colour: #ff6b00) and blue (#0035ff) squares match the
goldfish cone sensitivity peaks at 623 nm and 447 nm, respectively (Palacios, Varela et al. 1998).
The green (#56ff00) and purple (#7900e6) squares correspond to the *R. aculeatus* cone
sensitivity peaks at 528 nm and 413 nm, respectively (Cheney, Newport et al. 2013). The width
of the black and white stripes was chosen to be discriminable from a viewing distance of 1 m for
both species (Neumeyer 2003, Cheney, Hudson et al. 2022).

Turbidity preparation:

Turbidity was measured in Nephelometric Turbidity Units (NTU) using a turbidimeter (2100Q
Portable Turbidimeter, Hach). For all turbidity measurements, three 15 ml water samples were
collected and averaged. Prior to measurement in the turbidimeter, samples were inverted several
times to ensure particles remained suspended. Inorganic turbidity sources were purchased in dry
powder form (bentonite clay: Fullers Earth, Trustleaf, Cambridgeshire UK; calcium carbonate:
Intra Laboratories, Plymouth, UK; kaolin clay: Trustleaf, Cambridgeshire UK). Each turbidity
source was mixed by hand with small additions of RO water and sieved through a hand net. This
step was required to prevent particles from clumping and causing uneven particle distribution. A
stock solution of known concentration was created and used to gradually increase turbidity within
the experimental tank to the desired concentration of 100 NTU. The amount of inorganic turbidity
(bentonite: 5.04×10^{-2} grams per litre (g/L); calcium carbonate: 2.22×10^{-2} g/L; kaolin: 1.19×10^{-2} g/L).
^{2} g/L).

Live freshwater algae, *Scenedesmus quadricauda* (Blades Biological Ltd) was purchased and
grown over a 12-week period to produce 135L of water at 100 NTU. *S. quadricauda* was chosen
because of its fast growth rate and preferred temperature range which matched that of our lab.

To start, 240 mL of culture starter was added to the experimental tank which was half filled
(approximately 70L), along with 80 mL of Alga-Gro growth medium (Blades Biological Ltd). A
glass lid was placed loosely over the tank to reduce contamination, and a grow light (9 hour on/
15 hour off) was attached over the tank (40 cm above tank). The mixture was gently stirred by
hand every other day to increase aeration and reduce algal clumping. Over the four weeks of
growth, the maximum turbidity level reached was 9 NTU so the tank was moved outside to a
greenhouse on February 6th, 2024. A commercial aquatic plant fertiliser (AquaDesign PLANT+
1000ML Aquarium Plant "all in one" Complete Liquid Fertiliser) was added and stirred weekly,
and the algae was left to grow for a further 8 weeks until it had reached an average turbidity level
measurement of 50 NTU. Algal turbidity levels plateaued at 12 weeks, so algae turbidity
measurements were reduced from 0-100 to 0-52 NTU for this turbidity source.

Light measurements:

259 The side-welling absolute irradiance was measured with an Ocean Insights Flame-S-UV-VIS
260 spectrometer with a 400 μm fibre and a cosine corrector (CC-3 Spectralon optical diffuser, Ocean
261 Insight). The fibre optic cable was held in place with a 3D printed mount and measurements were
262 taken from the centre of the tank, 25 cm from the image standard, 15 cm above the bottom of the
263 tank, with the cosine corrector pointing towards the image standard.

Experimental procedure:

The following procedure was repeated for each turbidity source. The experimental tank was first
filled with RO water and the tripod secured to the bottom of the tank 50 cm from the image
standard (Supplementary Fig. S2). A wavemaker (TMC Reef Flow 2.0 16000 DC Wavemaker;
intensity setting: 2, mode: 1) was used throughout tests with inorganic turbidity sources to ensure
particles remained suspended. The wavemaker was removed for tests with algae to prevent
damage to the algal cells. Prior to adding any turbidity, baseline measurements and absolute
irradiance were taken in clear water. Approximately 100 mL of turbidity solution was then added
to the experimental tank to achieve a turbidity level between 0-5 NTU. The two measurements
(NTU and absolute irradiance) were repeated. This process continued, increasing the turbidity in
increments of 10 NTU, up to a maximum of 100 NTU. Once the final measurement had been
taken, the tank was emptied, cleaned, and refilled with RO water to test the next turbidity source.
276

A similar procedure was used to measure image contrast, but the camera was mounted on the
tripod instead of the fibre optic cable. The camera was positioned 15 cm from the image standard,
at an optimal distance determined by pilot experiments to best capture image contrast at different
turbidity levels. As previously described, baseline measurements were taken (i.e. turbidity level,
contrast photo) in clear water and repeated as turbidity was gradually increased. Once a turbidity
of 100 NTU was reached, measurements were taken and then repeated every hour over a total
of six hours. The final turbidity level measurement of a trial day determined the starting turbidity
level of the next trial day. The tank was emptied, cleaned, and refilled and this protocol repeated
the next trial day at the next starting turbidity level. This process was repeated until the turbidity
level dropped to 10 NTU by the final hour of an experimental day. The number of trial days
required to reach this criterion varied for each turbidity source (algae: n = 1 trial day, bentonite: n
= 5 trial days or 35 measurements; calcium carbonate: n = 2 trial days or 14 measurements; kaolin
clay: n = 5 trial days or 35 measurements).
290
291   Finally, the procedure was repeated to measure how turbidity level, pH and water hardness
292   changed over a six-hour period. The experimental tank was initially set to a turbidity of 100 NTU
293   measured. The following four measurements were taken every hour: 1) turbidity, 2) pH (Chauvin
294   Arnoux waterproof C.A 10001 pH metre), 3), temperature, and 4) KH (API KH test, API Fish Care).

Analysis:

Data Treatment:

Absolute irradiance: The number of photons at each wavelength (photons cm^{-2} s^{-1} $nm^{-1)}$ were
extracted from raw spectrometer measurements using OceanView software (Ocean Insight;
Version 2.0.8). Absolute irradiance was plotted as a spectral curve for each experimental
condition and allowed the shape of the curves to be compared. Spectral curves were also
standardized so that the area under each curve equals 1, effectively normalizing the overall light
levels. Four turbidity level categories were chosen to compare spectral curve shape
characteristics: 1) 10 NTU (±3 NTU), 2) 33 NTU (±2 NTU), 3) 52 NTU (±4.5 NTU), and 4) 94 NTU
(±2 NTU). The final category did not include spectral measurements for algae.
305
Absolute irradiance spectrum values were used to calculate luminance, hue and chroma (see
Endler 1990). Luminance refers to the total number of photons and is the achromatic property of
the light environment. Hue and chroma are determined by the relative number of photons in

different wavebands of the light spectrum and are the chromatic properties of the light
environment. Hue is determined by the wavelength region with the steepest slope in the spectra,
as well as whether the slope is positive or negative. The hue of the ambient light environment is
the category of colour, calculated using the segment classification method (Endler, 1990). This
method is used to assess changes to the chromatic properties of hue and chroma independently
of a specific visual system while still preserving the fundamental processes of 'opponency' and
'lateral inhibition' common to many invertebrate and invertebrate species. This produces a two-
dimensional colour space with the opponent pairs of long to medium-short wavelengths on the
vertical axis, and short to medium-long wavelengths on the horizontal axis. Hue is calculated as
the clockwise angle from the vertical axis about the origin, where the origin represents full
spectrum white light. This value indicates the waveband in the spectra with the greatest number
of photons. Chroma is the chromatic saturation of the light environment and is determined by the
magnitude of the difference in the number of photons between different parts of the spectrum and
the maximum slope of the spectra. A spectrum with more photons in one waveband compared
to its opponent pair waveband, will have a higher chroma than a spectrum with equal/similar
numbers of photons within opponent pairs.
325

Image Contrast: The image contrast of the photos of the image standard was calculated using
Michelson contrast (Michelson 1927), which measures the contrast gradient of a sinusoidal
pattern on a scale of 0 to 1. Michelson contrast gives the relation between the spread and the
sum of two luminance values and is commonly used in signal processing theory to determine the
quality of a signal relative to its noise level (e.g. scattered light) (Cronin, Johnsen et al. 2017,
Hunter, Godde et al. 2018). This method involves calculating the luminance difference between
two patches of an image and dividing it by the total luminance of both patches. Luminance values
were determined using a custom Matlab script, which averaged pixel intensity values from
specified coordinates for all black and white stripes.

Statistical Analysis:

We tested if turbidity source and level had a statistically significant effect on luminance, hue,
chroma, image contrast, pH, and KH. We also tested whether turbidity source had a significant
effect on turbidity level over time, which could indicate whether some types of turbidity remain
suspended more than others. In total, we conducted seven different statistical tests (see Table 1
for model parameters). All statistical analyses were conducted using R Studio (v.4.3.2) using the
lme4 (linear mixed models) package (Bates 2015). To identify models that best fit our data, we

tested various model parameters and interactions and compared the resulting models using

Akaike Information Criterion (AIC). We selected the model with the lowest AIC value and then

validated it using the DHARMa package, which tests for residual normality, over-dispersion, and

outliers (Hartig and Lohse 2022). Some models did not meet the assumption of residual normality

(see table 1), but each model was selected as the best possible fit model amongst those tested

(Hartig and Lohse 2022) (see Supplementary S3 for model selection). Results were plotted using

the ggplot2 package (Wickham 2016).

Table 1: Summary of linear model parameters. For well-fitting models, residual normality is

expected ('Yes'), no overdispersion ('No'), and no residual outliers ('No').

Model number	Model parameters			Model validation		
	Response Variable	Fixed effects	Random intercept	Residual normality	Over-dispersion	Residual outliers
1	Luminance	Turbidity source*concentration	Turbidity source	No	No	No
2	Hue	Turbidity source*concentration	Turbidity source	No	No	No
3	Chroma	Turbidity source*concentration	Turbidity source	No	No	No
4	Image contrast	Turbidity source + concentration^2	Turbidity source + camera location	Yes	No	No
5	pH	Turbidity source * time + turbidity source	Turbidity source + Initial turbidity level	Yes	No	No
6	KH	Turbidity source + time	Turbidity source + Initial turbidity level	No	No	No
7	Turbidity level	Turbidity source * time + turbidity source	Turbidity source + Initial turbidity level	Yes	No	No

Results:

Effect of turbidity on absolute irradiance:

Changes in the shape of a spectral curve can provide insights into the light environment. Fig. 2A-
D shows that turbidity source and level can alter the total irradiance (i.e. total number of photons).
Increasing turbidity level generally increases the number of photons for bentonite, calcium
carbonate, and kaolin, whereas algae have the opposite effect. Changes to the shape of the curve
can provide information on the number of photons at different wavelengths and indicate an overall
change in the colour of the light environment. These changes in shape can be seen in Fig 2A-D
but are more obvious in Fig 2E-H where total irradiance has been standardized. In the presence
of algae, there are fewer photons in the short-wavelength region (~400-525 nm) and more
photons in the medium-to-long wavelengths (~525-650 nm). Bentonite similarly reduced the
number of photons at shorter wavelengths while increasing them at longer wavelengths (~575-
700 nm), with a slight shift toward longer wavelengths.
365
366   Turbidity level has an effect for bentonite, as the shape of the curve between ~575-700 nm, differs
367   from 10 NTU to 94 NTU. Algae also shows an effect with turbidity, as the curve shape between
368   ~400-525 nm and ~525-650 nm differs between 10 NTU and 52 NTU. In contrast, calcium
369   carbonate and kaolin show minimal change from the clear water condition, with only slight
370   changes at the highest turbidity levels.
371

Effect of turbidity on luminance, hue, and chroma:

We found a significant, non-linear relationship between luminance and turbidity level
(Supplementary S4, Table 2). There was a significant interaction between all turbidity sources
and turbidity level on luminance: increasing turbidity level caused an increase in luminance for
kaolin (8.02×10^{18} photons ± SE = 8.64×10^{17}, p = <0.001), followed by calcium carbonate ($7.15 \times$
10^{18} photons ± SE = 8.00×10^{17}, p = <0.001). Bentonite resulted in a smaller but still significant
increase (3.77×10^{18} photons ± SE = 7.97×10^{17}, p = <0.001), while algae led to a significant
decrease in luminance (-1.99×10^{18} photons ± SE = 8.13×10^{17}, p = 1.47×10^{-2}) (F**ig. 2I**).
380

There was a significant interaction between all turbidity sources and turbidity level on hue:
increasing turbidity level caused the greatest shift in hue for bentonite ($6.31 \times 10^{-3\circ}$ ± SE = $3.35 \times$

10^{-4}, p= <0.001), followed by algae ($5.40 \times 10^{-3\circ} \pm SE = 8.53 \times 10^{-4}$, p= <0.001), then calcium

carbonate ($1.96 \times 10^{-3}{}^{\circ} \pm SE = 3.99 \times 10^{-4}$, p= <0.001), then kaolin ($1.51 \times 10^{-3\circ} \pm SE = 4.11 \times 10^{-4}$

^{4}, p= <0.001) (**Fig. 2J**, Supplementary S4, Table 3).

Furthermore, there was a significant interaction between some turbidity sources and their level

on chroma: increasing turbidity level caused an increase in chroma for algae ($3.32 \times 10^{-3} \pm SE =$

2.68×10^{-4}, p= <0.001) and bentonite ($3.71 \times 10^{-4} \pm SE = 1.05 \times 10^{-4}$, p = 1.39×10^{-3}). Neither

calcium carbonate ($-1.34 \times 10^{-4} \pm SE = 1.25 \times 10^{-4}$, p = 0.29) nor kaolin ($-1.68 \times 10^{-4} \pm SE = 1.29$

$\times 10^{-4}$, p = 0.20) turbidity level had a significant impact on chroma (**Fig. 2K**, Supplementary S4,

Table 4).

We found a significant, relationship between image contrast (Michelson contrast) and turbidity

level ($6.86 \times 10^{-3} \pm SE = 7.84 \times 10^{-4}$, p <0.001) (**Fig. 2L,** Supplementary S4, Table 5). When

considering individual turbidity sources, algae had the greatest effect on image contrast ($-0.69 \pm$

$SE = 6.06 \times 10^{-2}$, p <0.001), followed by kaolin ($-0.59 \pm SE = 5.97 \times 10^{-2}$, p <0.001), then bentonite

($-0.46 \pm SE = 5.88 \times 10^{-2}$, p <0.001), and calcium carbonate ($-0.37 \pm SE = 5.96 \times 10^{-2}$, p <0.001).

Figure 2: (A-**H) Effect of turbidity on absolute irradiance:** A-D show raw light spectra and E-H show

standardised spectra with the area under the curve set to one. (**I-L) Effect of turbidity on luminance, hue,**

chroma, and image contrast. Individual points indicate observed values. In panel L, lines represent the

best-fitting model, which shows a significantly non-linear decrease in image contrast with increasing

turbidity. In panels I-K, algae were only measured up to 50 NTU as we could not achieve a turbidity level

above this under our growing conditions. In panel L, contrast measurements were taken from 0-120 NTU,

but only 0-50 NTU are displayed as contrast values > 50 NTU approached zero. (**M-O) Turbidity settling**

rate, and the effects of turbidity on pH and water hardness (KH) over time. For visualisation purposes,

settling rate, pH, and KH data points are presented as the average percentage change in each value

between time points, then re-transformed to fit the original scale.

Turbidity settling rate, pH and water hardness over time:

The settling rate of each turbidity was described by repeatedly measuring suspended turbidity

over time. All turbidity led to some degree of settling, but there was a significant effect of turbidity

source on the settling rate (**Fig. 2M**, Supplementary S4, Table 6). Calcium carbonate had the

most rapid settling rate (-0.24 NTU $\pm$ SE = 2.73 x 10^{-2}, p <0.001), followed by algae (-0.15 NTU $\pm$

SE = 3.79 x 10^{-2}, p <0.001), kaolin (-8.21 NTU x 10^{-2} $\pm$ SE = 1.73 x 10^{-2}, p <0.001), and bentonite

(-6.47 x 10^{-2} NTU $\pm$ SE = 1.73 x 10^{-2}, p <0.001).

Some turbidity sources had a significant effect on pH, as well as a significant interaction between

turbidity source and time. Compared to pure RO water (pH=6.73 $\pm$ SE = 0.22), both algae and

calcium carbonate significantly increased pH (1.91 $\pm$ SE = 0.37, p < 0.001 and 1.33 $\pm$ SE = 0.32,

p < 0.001, respectively) but bentonite and kaolin had non-significant effects (p > 0.05) (**Fig. 2N**,

Supplementary S4, Table 7). Over time, calcium carbonate significantly increased pH (0.19 $\pm$SE

= 4.60 x 10^{-2}, p < 0.001), while kaolin decreased pH (0.12 $\pm$SE = 2.91 x 10^{-2}, p < 0.001). The

observed trend was that changes in pH generally occurred within the first hour, except in the case

of algae (**Fig. 2N**).

There was a significant impact of some turbidity sources on the water KH. KH increased for

calcium carbonate (1.50 $\pm$SE = 0.19, p < 0.001) and algae (1.00 $\pm$SE = 0.23, p < 0.001), but not

bentonite or kaolin (p>0.05) **(Fig. 2O,** Supplementary S4, Table 8)**.**

Discussion:

As land use changes and global temperatures rise, turbidity conditions in both freshwater and

marine habitats are expected to shift, impacting not only turbidity levels but also their sources. As

a result, fish will increasingly encounter turbidity levels and sources beyond what they have

evolved to cope with. Our results demonstrate that turbidity's effects on the light environment are

significant and dependent on both turbidity level and source. Algae reduce luminance, shift hue

towards medium-long wavelengths, and increase chroma, creating a darker, greener, and more

saturated light environment. Bentonite slightly increases luminance, shifts hue towards medium-

long and long wavelengths, and slightly increases chroma, resulting in a brighter, redder light
environment. Calcium carbonate and kaolin increase luminance but have minimal effects on hue
and chroma, producing a much brighter environment without altering chromatic properties. These
findings have implications for assessing the risks of changing turbidity to natural environments.
For example, Engström-Öst and Candolin (2007) found that increasing algae-based turbidity can
alter courtship behaviour in three-spined sticklebacks, and males in turbid water had to work
harder for female attention. As part of courtship, males of this species develop a bright red belly
during the breeding season, which serves as an important fitness signal (Seehausen, Terai et al.
2008). Our results here predict that the red patches would be less conspicuous in algal turbidity
but remain visible in bentonite-induced turbidity. Thus, for this species, nutrient runoff leading to
algal blooms may impair courtship behaviours, while clay soil runoff is less likely to have such an
impact. This study highlights the need to consider turbidity's impact on the visual environment
when trying to understand and predict its effects on aquatic species and their behaviour.
455
Although turbidity is generally associated with reduced light levels, our experiments show that the
total light available was actually higher in turbid water for the inorganic turbidity sources
(Bowmaker 1995) tested. Both irradiance (Fig 2A-D) and luminance (Fig 2I) were greater in turbid
water than in clear water. This increase in brightness is due to light scattering, which is influenced
by factors such as particle composition, size, concentration (Utne-Palm 2002, Johnsen 2012), as
well as other physical water properties including temperature, salinity, and pressure (Zhang and
Hu 2021, Tomperi, Isokangas et al. 2022). In contrast, algal turbidity reduces light availability, as
photosynthetic cells absorb rather than scatter light. The achromatic component of the light
environment (i.e. luminance) guides many fundamental visual tasks such motion detection, depth
perception, and pattern and texture discrimination (Siebeck, Wallis et al. 2014, van den Berg,
Hollenkamp et al. 2020). A shift in luminance could prevent the detection of critical visual cues.
Furthermore, a significant decrease in luminance may cause a shift from cone-based to rod-based
vision (Wolfe and Ali 2015), which would substantially impact visual abilities. As rods generally
have lower acuity than cones (Bowmaker 1995), significantly reduced luminance could impair a
species' ability to detect smaller visual cues or fine features. This is likely to happen with some
turbidity sources, such as charcoal, that lead to a rapid drop in luminance (e.g. by 1190% at 13.3
NTU, unpublished data).
473
Light consists of both achromatic (luminance) and chromatic (hue, chroma) components. Spectra
with a decrease in the area under the curve will have a lower luminance, making the environment

appear darker. The luminance of a visual feature is proportional to how closely its reflectance
spectrum matches the ambient light spectrum. For this reason, a blue fish in blue water will appear
brighter (higher luminance) and more visible than a red fish. Changes to the chromatic component
of the light environment shift the relative availability of different wavelengths, altering how visual
stimuli reflecting those wavelengths appear (Endler 1990). Spectra with differing wavelengths of
greatest slope are perceived as different hues (Endler 1990), and spectra with a greater difference
between the maximum and minimum peaks of the curve will have a greater chroma, resulting in
more saturated colours. Additionally, the perceived hue of a visual feature can change based on
the spectral properties of the turbidity source, with these effects being wavelength-specific.
Turbidity sources like algae, which absorb longer wavelengths (e.g. red, orange), reduce the
visibility of red and orange features, while green and blue features will remain conspicuous. In
contrast, bentonite turbidity scatters shorter wavelengths (e.g. blue, green), limiting the visibility
of blue and green features but not red, orange or brown ones. Both algae and bentonite also
significantly increase the chroma of the light environment, so that the light environment is more
saturated in specific regions of the spectrum. The saturated region is dependent on the turbidity
source. Regardless of a species' visual system, spectra with a greater magnitude of change
between different parts of the spectrum will typically be perceived as having greater chroma, and
spectra with differing wavelengths of greatest slope will be perceived as different hues (Endler
1990). Finally, we observed a significant decrease in image contrast with the addition of all
turbidity sources and increasing turbidity levels. Image contrast measurements incorporate
estimates from all four optical factors (i.e. the ambient light environment, the spectral reflectance
of the visual stimuli, the transmission of light through the medium, and the amount of veiling light)
in the light pathway to provide an approximation of how turbidity affects the visibility of a stimulus
and therefore can serve as a practical tool for assessing the impacts of turbidity without complex
visual modelling. In our setup, the greatest changes in image contrast, and consequently stimulus
visibility, occurred between 0-15 NTU, with effects stabilizing at 20 NTU. The combined chromatic
changes to the light environment are likely to significantly affect visually-guided behaviours in fish,
particularly those relying on colour signals such as sexual selection, aposematism, spatial
cognition, camouflage, and predator-prey interactions (Endler 1993).

While we suggest that experimental turbidity sources should be selected for their potential effects
on the visual environment, not all sources will be suitable for all species. When turbidity sources
were added to RO water, we observed a significant drop in turbidity level over time that was
independent of the turbidity source. Calcium carbonate had the greatest turbidity level change

over time, followed by algae, kaolin, then bentonite. The difference in turbidity level change over time between turbidity sources found here is useful to consider when designing turbidity experiments. Scientists wishing to take advantage of quickly changing turbidity levels, for example where changing turbidity levels between trials is logistically challenging, might choose to use a turbidity source that settles quickly. Turbidity levels that change more slowly on the other hand, might be more useful to experiments with longer trial durations. Both pH and KH were significantly impacted in a turbidity source-dependent manner. Algae, calcium carbonate and kaolin lead to significant increases or decreases (for kaolin only) in the water pH. While the change in pH was immediate with algae, it was observed as soon as the second measurement was taken (one hour after the turbidity source was added). This indicates that these calcium carbonate and kaolin need a short time in the water to modify its chemistry. Calcium carbonate and algae significantly increased the water KH. The results of this study are not intended to allow researchers to predict the precise impact of each turbidity source on pH and KH in their experimental design, as small changes in their design such as water mineral content and temperature can impact pH and KH (Irwin, Stoner et al. 2013). However, it demonstrates that pH and KH must be taken into account when designing turbidity experiments and monitored during experiments as they may change with turbidity source and time and could crucially impact the welfare of the study species and the output of behavioural experiments (Kleinhappel, Burman et al. 2019, Cleal, Gibbon et al. 2020). This paper is intended to serve as a guide to key considerations and potential effect sizes, acknowledging that each environment, natural or laboratory, has unique characteristics.

This study explores how turbidity influences the light environment, and demonstrates that luminance, hue, chroma, and image contrast are significantly affected by turbidity, with variation depending on turbidity source and level. Our findings indicate that different sources and levels of turbidity cannot be assumed to uniformly affect the appearance of visual stimuli or the aquatic environment. We also examined particle setting rate by turbidity source, along with their effects on pH and KH. Understanding turbidity's impact on animals requires considering how it alters visual information and may help us identify the mechanisms species have evolved to cope with changing visual conditions. Moreover, this study highlights the need for conservation ecology studies to consider turbidity levels, sources, and ecological contexts when interpreting the potential consequences of turbidity for wild fish populations.

Acknowledgements:

We thank animal technicians and field station support staff: Helen Sanders, Christine Soper, Allex Turner, Phil Smith, John Hogg, and Angela Bishop.

Competing interests:

 The authors declare no conflict of interest.

Funding:

This research was support by a Leverhulme Trust Early Career Fellowship (CN), the University of Oxfords Hannah Elizabeth Jenkinson Research Fund (AG), and University of Oxford Hester Cordelia Parsons Fund (AG), and a Human Frontier Science Program grant RGP0016/2019 (AS and TB).

Data and resource availability:

We are creating a data repository link at the time of submission linked to the manuscript. In the meantime, all data and code are available on demand.

References:

Ajemian, M. J., S. Sohel and J. Mattila (2015). "Effects of turbidity and habitat complexity on antipredator behavior of three-spined sticklebacks (*Gasterosteus aculeatus*)." Environmental Biology of Fishes **98**(1): 45-55.

Albini, D., L. Lester, P. Sanders, J. Hughes and M. C. Jackson (2023). "The combined effects of treated sewage discharge and land use on rivers." Global Change Biology **29**(22): 6415-6422.

Auld, A. H. and J. R. Schubel (1978). "Effects of suspended sediment on fish eggs and larvae - Laboratory assessment." Estuarine and Coastal Marine Science **6**(2): 153-164.

Babin, M., A. Morel, V. Fournier-Sicre, F. Fell and D. Stramski (2003). "Light scattering properties
of marine particles in coastal and open ocean waters as related to the particle mass
concentration." Limnology and Oceanography **48**(2): 843-859.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). "Fitting linear mixed-effects models Using
lme4." Journal of Statistical Software **67**(1).

Berli, B. I., M. J. H. Gilbert, A. L. Ralph, K. B. Tierney and P. Burkhardt-Holm (2014). "Acute
exposure to a common suspended sediment affects the swimming performance and physiology
of juvenile salmonids." Comparative Biochemistry and Physiology Part A: Molecular & Integrative
Physiology **176**: 1-10.

Bohren, C. F. and D. R. Huffman (2008). Absorption and scattering of light by small particles,
John Wiley & Sons.

Bojko, J., E. K. Lipp, A. T. Ford and D. C. Behringer (2020). Pollution can drive marine diseases.
Marine Disease Ecology. Oxford, Oxford University Press.

Borner, K. K., S. Krause, T. Mehner, S. Uusi-Heikkilä, I. W. Ramnarine and J. Krause (2015).
"Turbidity affects social dynamics in Trinidadian guppies." Behavioral Ecology and Sociobiology
69(4): 645-651.

Bowmaker, J. K. (1995). "The visual pigments of fish." Progress in retinal and eye research **15**(1):
1-31.

Bricaud, A., H. Claustre, J. Ras and K. Oubelkheir (2004). "Natural variability of phytoplanktonic
absorption in oceanic waters: Influence of the size structure of algal populations." Journal of
Geophysical Research: Oceans **109**(C11).

Budria, A. and U. Candolin (2015). "Human-induced eutrophication maintains high parasite
prevalence in breeding threespine stickleback populations." Parasitology **142**(5): 719-727.

Burke, L., K. Reytar, M. Spalding and A. Perry (2011). Reefs at Risk Revisited.

Burkhead, N. M. and H. L. Jelks (2001). "Effects of suspended sediment on the reproductive
success of the tricolor shiner, a crevice-spawning minnow." Transactions of the American
Fisheries Society **130**(5): 959-968.

Candolin, U. and I. Jensen (2021). "Phenotypic plasticity in courtship exposed to selection in a
human-disturbed environment." Evolutionary Applications **14**(10): 2392-2401.

Cano-Rocabayera, O., S. Vargas-Amengual, C. Aranda, A. de Sostoa and A. Maceda-Veiga
(2020). "Mosquito larvae consumption in turbid waters: the role of the type of turbidity and the
larval stage in native and invasive fish." Hydrobiologia **847**(5): 1371-1381.

Cartwright, P. J., P. R. C. S. Fearns, P. Branson, M. V. W. Cuttler, M. O'Leary, N. K. Browne and
R. J. Lowe (2021) "Identifying metocean drivers of turbidity using 18 ears of MODIS satellite data:
Implications for marine ecosystems under climate change." Remote Sensing **13** DOI:
10.3390/rs13183616.

Chamberlain, A. C. and C. C. Ioannou (2019). "Turbidity increases risk perception but constrains
collective behaviour during foraging by fish shoals." Animal Behaviour **156**: 129-138.

Cheney, K. L., J. Hudson, F. de Busserolles, M. Luehrmann, A. Shaughnessy, C. van den Berg,
N. F. Green, N. J. Marshall and F. Cortesi (2022). "Seeing Picasso: an investigation into the visual
system of the triggerfish *Rhinecanthus aculeatus*." Journal of Experimental Biology **225**(7).

Cheney, K. L., C. Newport, E. C. McClure and N. J. Marshall (2013). "Colour vision and response
bias in a coral reef fish." Journal of Experimental Biology **216**(Pt 15): 2967-2973.

Cleal, M., A. Gibbon, B. D. Fontana and M. O. Parker (2020). "The importance of pH: How
aquarium water is affecting behavioural responses to drug exposure in larval zebrafish."
Pharmacology Biochemistry and Behavior **199**: 173066.

Cronin, T. W., S. Johnsen, N. J. Marshall and E. Warrant (2017). Visual ecology. Princeton,
Princeton University Press.

Cumming, H. and N. A. Herbert (2016). "Gill structural change in response to turbidity has no
effect on the oxygen uptake of a juvenile sparid fish." Conservation Physiology **4**: 12.

Cyrus, D. P. and S. J. M. Blaber (1992). "Turbidity and salinity in a tropical northern Australian
estuary and their influence on fish distribution." Estuarine, Coastal and Shelf Science **35**(6): 545-
563.

Ehlman, S. M., B. A. Sandkam, F. Breden and A. Sih (2015). "Developmental plasticity in vision
and behavior may help guppies overcome increased turbidity." Journal of Comparative
Physiology A **201**(12): 1125-1135.

Endler, J. A. (1990). "On the measurement and classification of colour in studies of animal colour
patterns." Biological Journal of the Linnean Society **41**(4): 315-352.

Endler, J. A. (1993). "The color of light in forests and its implications." Ecological Monographs
63(1): 1-27.

Figueiredo, B. R. S., R. V. Granzotti, L. F. Fiori, C. L. Nieman and E. Benedito (2020). "Cascading
effects of predation risk under high organic and inorganic turbidity: impacts on individuals and
shoals of a mesopredator fish." Aquatic Ecology **54**(3): 855-868.

Fukunaga, A., J. H. R. Burns, K. H. Pascoe and R. K. Kosaki (2022). "A remote coral reef shows
macroalgal succession following a mass bleaching event." Ecological Indicators **142**: 109175.

Glendell, M. and R. E. Brazier (2014). "Accelerated export of sediment and carbon from a
landscape under intensive agriculture." Science of The Total Environment **476-477**: 643-656.

Grecay, P. A. and T. E. Targett (1996). "Effects of turbidity, light level and prey concentration on
feeding of juvenile weakfish *Cynoscion regalis*." Marine Ecology Progress Series **131**: 11-16.

Hartig, F. and L. Lohse (2022). "DHARMa: Residual diagnostics for hierarchical (multi-level /
mixed) regression models."

Hasenbein, M., N. A. Fangue, J. Geist, L. M. Komoroske, J. Truong, R. McPherson and R. E.
Connon (2016). "Assessments at multiple levels of biological organization allow for an integrative
determination of physiological tolerances to turbidity in an endangered fish species."
Conservation Physiology **4**: 16.

Hayes, J. W., M. J. Rutledge, B. L. Chisnall and F. J. Ward (1992). "Effects of elevated turbidity
on shallow lake fish communities." Environmental Biology of Fishes **35**(2): 149-168.

Hess, S., A. S. Wenger, T. D. Ainsworth and J. L. Rummer (2015). "Exposure of clownfish larvae
to suspended sediment levels found on the Great Barrier Reef: Impacts on gill structure and
microbiome." Scientific Reports **5**: 8.

Hildebrandt, E. K. and G. R. Parsons (2016). "Effect of turbidity on the swimming performance of

the golden shiner, *Notemigonus crysoleucas*." Copeia **104**(3): 752-755.

Hunter, M., B. Godde and B. Olk (2018). "Effects of absolute luminance and luminance contrast

on visual search in low mesopic environments." Attention, Perception, & Psychophysics **80**(5):

1265-1277.

Irwin, M. D., J. B. Stoner and A. M. Cobaugh (2013). Zookeeping : an introduction to the science

and technology. Chicago ;, The University of Chicago Press.

Järvenpää, M. and K. Lindström (2004). "Water turbidity by algal blooms causes mating system

breakdown in a shallow-water fish, the sand goby *Pomatoschistus minutus*." Proceedings of the

Royal Society of London. Series B: Biological Sciences **271**(1555): 2361-2365.

Järvenpää, M. and K. Lindström (2011). "Algal blooms decrease care but increase egg survival

in a fish with paternal care." Behavioral Ecology and Sociobiology **65**(11): 2023-2028.

Järvenpää, M., B. D. Pauli and K. Lindström (2019). "Water turbidity constrains male mating

success in a marine fish." Behavioral Ecology and Sociobiology **73**(10): 7.

Johnsen, S. (2012). The optics of life: A biologist's guide to light in nature, Princeton University

Press.

Kellogg, K. A. and P. Leipzig-Scott (2017). "The influence of turbidity on prey consumption in the

tessellated darter." Transactions of the American Fisheries Society **146**(3): 508-511.

Kimbell, H. S. and L. J. Morrell (2015). "Turbidity influences individual and group level responses

to predation in guppies, *Poecilia reticulata*." Animal Behaviour **103**: 179-185.

Kleinhappel, T. K., O. H. P. Burman, E. A. John, A. Wilkinson and T. W. Pike (2019). "The impact

of water pH on association preferences in fish." Ethology **125**(4): 195-202.

Li, W., T. L. Zhang, C. W. Zhang, Z. J. Li, J. S. Liu and B. J. Hicks (2013). "Effects of turbidity and

light intensity on foraging success of juvenile mandarin fish *Siniperca chuatsi* (Basilewsky)."

Environmental Biology of Fishes **96**(8): 995-1002.

Love, A. C., N. Crooks and A. T. Ford (2020). "The effects of wastewater effluent on multiple

behaviours in the amphipod, *Gammarus pulex*." Environmental Pollution **267**: 115386.

Lunt, J. and D. L. Smee (2015). "Turbidity interferes with foraging success of visual but not
chemosensory predators." Peerj **3**: 12.

Macdonald, R. K., P. V. Ridd, J. C. Whinney, P. Larcombe and D. T. Neil (2013). "Towards
environmental management of water turbidity within open coastal waters of the Great Barrier
Reef." Marine Pollution Bulletin **74**(1): 82-94.

Malthus, T. J., E. Lehmann, X. Ho, E. Botha and J. Anstee (2019) "Implementation of a satellite
based inland water algal bloom alerting system using analysis ready data." Remote Sensing **11**
DOI: 10.3390/rs11242954.

Michelson, A. A. (1927). Studies in optics. Illinois, University of Chicago Press.

Miranda, L. E. (2011). "Depth as an organizer of fish assemblages in floodplain lakes." Aquatic
Sciences **73**(2): 211-221.

Moustaka, M., T. J. Langlois, D. McLean, T. Bond, R. Fisher, P. Fearns, P. Dorji and R. D. Evans
(2018). "The effects of suspended sediment on coral reef fish assemblages and feeding guilds of
north-west Australia." Coral Reefs **37**(3): 659-673.

Neumeyer, C. (2003). Color vision in fishes and Its neural basis. Sensory Processing in Aquatic
Environments. S. P. Collin and N. J. Marshall. New York, NY, Springer New York**:** 223-235.

Newport, C., O. Padget and T. B. de Perera (2021). "High turbidity levels alter coral reef fish
movement in a foraging task." Scientific Reports **11**(1): 5976.

Ortega, J. C. G., B. R. S. Figueiredo, W. J. da Graça, A. A. Agostinho and L. M. Bini (2020).
"Negative effect of turbidity on prey capture for both visual and non-visual aquatic predators."
Journal of Animal Ecology **89**(11): 2427-2439.

Palacios, A. G., F. J. Varela, R. Srivastava and T. H. Goldsmith (1998). "Spectral sensitivity of
cones in the goldfish, *Carassius auratus*." Vision Research **38**(14): 2135-2146.

Pangle, K. L., T. D. Malinich, D. B. Bunnell, D. R. DeVries and S. A. Ludsin (2012). "Context-
dependent planktivory: interacting effects of turbidity and predation risk on adaptive foraging."
Ecosphere **3**(12): 18.

700 Radke, R. and A. Gaupisch (2005). "Effects of phytoplankton-induced turbidity on predation

701 success of piscivorous Eurasian perch (*Perca fluviatilis*): possible implications for fish community

702 structure in lakes." Naturwissenschaften **92**(2): 91-94.

703 Seehausen, O., Y. Terai, I. S. Magalhaes, K. L. Carleton, H. D. J. Mrosso, R. Miyagi, I. van der

704 Sluijs, M. V. Schneider, M. E. Maan, H. Tachida, H. Imai and N. Okada (2008). "Speciation

705 through sensory drive in cichlid fish." Nature **455**(7213): 620-626.

706 Seers, B. M. and N. T. Shears (2015). "Spatio-temporal patterns in coastal turbidity – Long-term

707 trends and drivers of variation across an estuarine-open coast gradient." Estuarine, Coastal and

708 Shelf Science **154**: 137-151.

709 Shoji, J., E. W. North and E. D. Houde (2005). "The feeding ecology of *Morone americana* larvae

710 in the Chesapeake Bay estuarine turbidity maximum: the influence of physical conditions and prey

711 concentrations." Journal of Fish Biology **66**(5): 1328-1341.

712 Siebeck, U. E., G. M. Wallis, L. Litherland, O. Ganeshina and M. Vorobyev (2014). "Spectral and

713 spatial selectivity of luminance vision in reef fish." Frontiers in Neural Circuits **8**.

714 Smith, V. H. (2003). "Eutrophication of freshwater and coastal marine ecosystems a global

715 problem." Environmental Science and Pollution Research **10**(2): 126-139.

716 Sundin, J., T. Aronsen, G. Rosenqvist and A. Berglund (2017). "Sex in murky waters: algal-

717 induced turbidity increases sexual selection in pipefish." Behavioral Ecology and Sociobiology

718 **71**(5).

719 Sundin, J., G. Rosenqvist, S. Myhren and A. Berglund (2016). "Algal turbidity hampers ornament

720 perception, but not expression, in a sex-role-reversed pipefish." Ethology **122**(3): 215-225.

721 Sutherland, A. B. and J. L. Meyer (2007). "Effects of increased suspended sediment on growth

722 rate and gill condition of two southern Appalachian minnows." Environmental Biology of Fishes

723 **80**(4): 389-403.

724 Sweka, J. and K. Hartman (2011). "Effects of turbidity on prey consumption and growth in Brook

725 trout and implications for bioenergetics modeling." Canadian Journal of Fisheries and Aquatic

726 Sciences **58**: 386-393.

Tiarks, J. H., S. M. Gray and L. J. Chapman (2024). "Turbidity drives plasticity in the eyes and
brains of an African cichlid." Journal of Experimental Biology **227**(7): jeb246708.

Tomperi, J., A. Isokangas, T. Tuuttila and M. Paavola (2022). "Functionality of turbidity
measurement under changing water quality and environmental conditions." Environmental
Technology **43**(7): 1093-1101.

Tuomainen, U. and U. Candolin (2011). "Behavioural responses to human-induced environmental
change." Biological Reviews **86**(3): 640-657.

Utne-Palm, A. C. (2002). "Visual feeding of fish in a turbid environment: Physical and behavioural
aspects." Marine and Freshwater Behaviour and Physiology **35**(1-2): 111-128.

van den Berg, C. P., M. Hollenkamp, L. J. Mitchell, E. J. Watson, N. F. Green, N. J. Marshall and
K. L. Cheney (2020). "More than noise: context-dependent luminance contrast discrimination in a
coral reef fish (*Rhinecanthus aculeatus*)." Journal of Experimental Biology **223**(21): jeb232090.

Wickham, H. (2016). ggplot2: Elegant graphics for data analysis. New York, Springer-Verlag.

Wolfe, U. and N. Ali (2015). "Dark adaptation and purkinje shift: a laboratory exercise in perceptual
neuroscience." Journal of Undergraduate Neuroscience Education **13**(2): A59-63.

Wong, B. B. M., U. Candolin and K. Lindström (2007). "Environmental deterioration compromises
socially enforced signals of male quality in three-spined sticklebacks." American Naturalist **170**(2):
184-189.

Zhang, X. and L. Hu (2021). "Light scattering by pure water and seawater: Recent development."
Journal of Remote Sensing.
747
748

Figure legends:

Figure 1 - The frequency of use of different turbidity sources in 200 scientific papers exploring the impact of turbidity on fish behaviour, physiological condition, and biodiversity. Papers are grouped by turbidity source. "Clay" and "natural sediment" are often sourced directly from the study site, with their specific composition unknown. More specific turbidity sources, such as bentonite, were grouped separately. Within each turbidity source, papers are categorized by research focus. Studies using multiple turbidity sources or response categories were counted separately for each.

Figure 2: (A-H) Effect of turbidity on absolute irradiance: A-D show raw light spectra and E-H show standardised spectra with the area under the curve set to one. (I-L) Effect of turbidity on luminance, hue, chroma, and image contrast. Individual points indicate observed values. In panel L, lines represent the best-fitting model, which shows a significantly non-linear decrease in image contrast with increasing turbidity. In panels I-K, algae were only measured up to 50 NTU as we could not achieve a turbidity level above this under our growing conditions. In panel L, contrast measurements were taken from 0-120 NTU, but only 0-50 NTU are displayed as contrast values > 50 NTU approached zero. (M-O) Turbidity settling rate, and the effects of turbidity on pH and water hardness (KH) over time. For visualisation purposes, settling rate, pH, and KH data points are presented as the average percentage change in each value between time points, then re-transformed to fit the original scale.