TENM4 is an essential transduction component for touch

Mohammed A. Khallaf[1,2], Angela Tzu-Lun Huang[1,3], Letizia Dalmasso[1,3], Sampurna Chakrabarti[1],

Rowena Groeneveld[1], Yinth Andrea Bernal Sierra[1], Jonathan Alexis Garcia-Contreras[1], Emilie

Nello[1], Anja Schütz[4], Valérie Bégay[1], Alessandro Santuz[5], Sarah Hedtrich[6,7,8], Wei Zhong[1],

Christina Schiel[9], Steven J. Middleton[1], Oliver Popp[6,10], Niccolò Zampieri[5], Philipp Mertins[6,10],

Severine Kunz[9] and Gary R. Lewin[1,11,12*]

Affiliations:

[1]Max Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), Molecular Physiology of Somatic Sensation, Robert-Rössle Str. 10, 13125 Berlin-Buch, Germany

[2]Department of Zoology and Entomology, Faculty of Science, Assiut University, Assiut, 71516 Egypt

[3]These authors contributed equally to this work

[4]Max Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), Protein Production and Characterization, Robert-Rössle Str. 10, 13125 Berlin-Buch, Germany

[5]Max Delbrück Center for Molecular Medicine, Laboratory of Development and Function of Neural Circuits, Robert-Rössle-Str. 10, 13125 Berlin, Germany

[6]Berlin Institute of Health at Charité - Universitätsmedizin Berlin, Germany Charité – Universitätsmedizin Berlin, Germany

[7]School of Biomedical Engineering, The University of British Columbia, Vancouver, BC, Canada

[8]Max-Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), 13125

[9]Max Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), Electron microscopy Technology Platform, Robert-Rössle Str. 10, 13125 Berlin-Buch, Germany

[10]Max Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), Proteomics, Robert-Rössle Str. 10, 13125 Berlin-Buch, Germany

[11]Charité-Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany

[12]German Center for Mental Health (DZPG), partner site Berlin

*Correspondence glewin@mdc-berlin.de

SUMMARY

Gentle touch is conveyed to the brain by fast-conducting sensory fibers. Mechanosensitive ion channels at the terminals of these neurons are thought to be gated by extracellular tethers that transmit force from the surrounding matrix to the channel complex, but the molecular identity of such tethers has remained unknown. Here, we identify Teneurin-4 (TENM4) as an essential extracellular linker protein for mechanotransduction in mechanoreceptors. Sensory neuron–specific deletion of *Tenm4* in mice caused profound touch insensitivity, while acute and reversible proteolytic disassembly of TENM4 at sensory endings confirmed its direct role in force transduction. Ultrastructural analyses revealed TENM4 localization to filamentous structures at the neurite–laminin interface, defining it as a structural component of the mechanosensory tether. These findings identify TENM4 as a core element of fast somatic sensation and provide molecular insight into how extracellular forces are coupled to ion channel activation.

INTRODUCTION

All organisms, from bacteria to humans, react within a fraction of a second to mechanical stimuli.[1,2] In all branches of life, a molecularly diverse range of mechanosensitive ion channels are rapidly gated by force to initiate signaling.[2] The fastest and most sensitive mechanosensitive channels detect nanometer scale movements that enable hearing and touch perception.[3–5] Interestingly, mechanosensitive channels in fruit flies, worms and vertebrate hair cells require intracellular or extracellular tether proteins that relay force to the channel.[5–11] Vertebrate touch sensation also requires mechanosensitive ion channels, of which PIEZO2 and ELKIN1 share roles in mechanosensation.[4,12,13] The skin is supplied with a rich variety of molecularly and physiologically distinct mechanoreceptors,[14] the cell bodies of which reside in the dorsal root ganglia (DRG). Despite their functional diversity, all mechanoreceptors transduce mechanical stimuli into electrical signals at specialized neuroglial endings in the skin.[15,16] There are several lines of evidence that extracellular proteins at these neuroglial endings are directly involved in transferring force to mechanically gated channels. First, mechanically activated currents in sensory neurons can be acutely ablated by proteases, including blisterase, an enzyme that specifically cleaves furin consensus sites.[17,18] Mechanosensitive currents are restored 24 hours after protease treatment in cultured sensory neurons, suggesting that the protease target is synthesized by sensory neurons.[17] In the skin, protease treatment of the receptive fields of individual mechanoreceptors rapidly abolishes mechanosensitivity indicating an effect at the neuroglial ending.[17] Protease treatment was also associated with the loss of tether-like structures synthesized by sensory neurons.[17] These data, although suggestive, did not identify the molecular nature of the extracellular protein(s) that are necessary for transduction. Here, we identify TENM4 as structural component in mechanotransduction complexes the integrity of which is necessary for transduction in essentially all cutaneous mechanoreceptors.

RESULTS

Screening for tether candidates

Blisterase cleaves a furin consensus site RX(K/R)R, which was predicted to occur in at least 11,302 mouse proteins; but only a small minority of these possess putative cleavage sites located in protease accessible extracellular domains. Candidates with predicted extracellular domains were selected using the Database for Annotation, Visualization, and Integrated Discovery (DAVID).[19] Using gene paint (www.genepaint.org),[20] we selected 34 of these genes for further analysis based on the presence of in situ hybridization signals in the developing and adult DRG. Sympathetic neurons located in the superior cervical ganglion (SCG) do not possess mechanically activated currents (MA currents).[17,21] We thus reasoned that our candidate(s) should be expressed in the DRG, but not in the SCG. Using qPCR, we could not detect the mRNA for six of the 34 genes in the SCG, but all showed robust mRNA expression in the DRG (Table S1). The six genes coded for the following proteins: Teneurin-4 (TENM4), FRAS1 Related Extracellular Matrix Protein 2 (FREM2), Proprotein Convertase Subtilisin/Kexin Type 5 (PCSK5), Scribble Homolog (SCRIB), Alpha-Tectorin (TECTA), and Slit Guidance Ligand 1 (SLIT1). We used antibodies directed against the six candidates and obtained signals in cultured DRG neurons for each candidate, but not in SCG cultures, validating the results of the qPCR experiment (Figure S1A). We chose to study TENM4 and FREM2 first, partly because these proteins had the largest extracellular domains of the candidates (mouse proteins 2771 aa and 3160 aa, respectively), consistent with the large tether structures we had previously observed.[17,22] We generated two mouse strains with floxed *Frem2* and *Tenm4* alleles (Figures S2A and S2B) as both *Frem2* and *Tenm4* appear to be essential genes during embryonic development.[23,24]

Sensory neuron-specific conditional knockouts were created using an adeno-associated virus that is neurotropic for DRG neurons carrying a Cre/GFP cassette (AAV-PHP.S-Cre-eGFP).[25,26] We

examined *Frem2^{fl/fl}* mice 4-6 weeks after virus application, a time point chosen to allow sufficient

Cre-mediated excision.[26] Paw withdrawal thresholds to von Frey hair stimulation, responses to

cotton swab and brush stimuli indicated no loss of touch sensitivity in these mice (Figure S1B).

We thus turned our attention to the potential role of TENM4 in sensory mechanotransduction.

Mechanoreceptors express the TENM4 protein

TENM4 is a member of the Teneurin family, consisting of four proteins in vertebrates, which are

evolutionarily conserved type II membrane proteins known to be essential during neural

development.[27,28] Teneurins are dimeric proteins and there is some evidence that hetero-dimers

between two different Teneurin proteins may be formed.[29] We identified a commercially available

TENM4 antibody directed against the intracellular N-terminus, which is divergent between the

four mouse Teneurins (Anti-TENM4-N).[30] But we also generated an antibody against a large

recombinant extracellular fragment, which included the extracellular compact YD shell region of

the protein (Anti-TENM4-C). We first looked for TENM4 at the neuroglial endings of a variety

of mechanoreceptor types in the skin where touch stimuli are transduced (Figure 1A),[15,16] and

found strong immunolabeling of sensory endings that were also positive for neurofilament-200

(NF200[+]), a marker of all myelinated sensory fibers (Figure 1B). Sensory end-organs innervated

by rapidly-adapting mechanoreceptors (RAMs) like Meissner corpuscles and hair follicles were

all strongly positive for TENM4 (Figure 1B). Antibodies against S100β were used to label terminal

Schwann cells that make up the Meissner corpuscle and are closely associated with lanceolate

endings around hair follicles,[15,31] however TENM4 labelling was not seen in these cells (Figure

1B). We found that slowly-adapting mechanoreceptors (SAMs) positive for NF200[+] that innervate

CK20[+] Merkel cells were also co-labelled with TENM4 (Figure 1B). Using the same strategy as

for *Frem2*, we generated mice with a sensory neuron specific deletion of *Tenm4* (Figures 2A and

S2B) and saw essentially no TENM4 labeled sensory endings associated with RAM or SAM

innervated end-organs (Figure 1B), demonstrating the specificity of the antibody (Anti-TENM4-N). We also detected TENM4+ axons in wild type (WT) skin that terminated as free nerve endings (Figure S1C). Thus, TENM4 appeared to be a good marker of myelinated sensory endings including low threshold mechanoreceptors (LTMRs).

Figure 1

118
Figure 1. TENM4 is present at sites of transduction in mechanoreceptors. (**A**) Schematic of mechanoreceptor and nociceptor endings in hairy and glabrous skin. (**B**) TENM4+ and NF200+ sensory endings associated with rapidly-adapting mechanoreceptors (RAMs) like Meissner's corpuscles, lanceolate

endings around hair follicles and slowly adapting mechanoreceptors (SAMs) innervating CK20+ Merkel
cells. S100+ terminal Schwann cells were not TENM4+. TENM4 immunostaining was largely absent from
RAM and SAM endings in *Tenm4^CKO* mice. Scale bars 20μm. (**C**) TENM4+ staining was absent in cultured
sensory neurons from *Tenm4^CKO* mice. Scale bar 10μm. (**D**) Representative DRG sections showing double
labelling of *Tenm4* smFISH probe with NF200, TRKC, *Kcnq4-mRNA*, CGRP, TH or IB4, Scale bar 100μm.
Bottom shows quantification Tenm4+ sensory neurons co-labeled with different sensory markers. For each
co-labelling 2 male and 3 female mice were used and more than 700 neurons counted.
129

Our validated antibody revealed the presence of TENM4 in large diameter DRG neurons in culture

as well as a specific localization to the mechanoreceptor endings associated with specific end-

organs in the skin (Figures 1B,C and S1E). We next analyzed *Tenm4* expression by single-

molecule fluorescence in situ hybridization (smFISH) and assessed its co-localization with

markers of nociceptors and mechanoreceptors in DRG sections (Figure 1D). Consistent with our

results from the skin, all NF200$^+$ cells were also *Tenm4$^+$*. However, many nociceptive sensory

neurons positive for calcitonin-gene related peptide (CGRP) were also co-labeled with *Tenm4*

(Figure 1D). TRKC and *Kcnq4* expression are highly specific to RAMs,[32,33] and all these cells

were also positive for *Tenm4* (Figure 1D). Small sensory neurons positive for tyrosine hydroxylase

(TH), a marker of low threshold tactile C-fibers[34], were also largely positive for *Tenm4*. However,

we saw very little, or no overlap with nociceptor markers such as isolectin-B4 (IB4) (Figure 1D).

As expected, the TENM4 signal was largely absent in cultured sensory neurons and sections from

Tenm4^CKO mice (Figures 1C and S1E). Together, these data indicate that TENM4 is selectively

present in virtually all mechanoreceptors including nociceptors with myelinated axons, suggesting

its involvement in touch and rapid pain sensation.

***Tenm4* deletion abolishes mechanosensitivity and causes sensory deficits**

146 We next tested whether TENM4 is necessary for the mechanosensory function of identified mouse

147 mechanoreceptors using *Tenm4^CKO* mice (Figure 2A). Four weeks after retro-orbital injection of

148 AAV-PHP.S-Cre-EGFP virus in *Tenm4^fl/fl* mice we cultured sensory neurons from the DRG and

149 found that ~80% showed green fluorescence. Thus, the vast majority of the sensory neurons had

probably undergone Cre-mediated excision at the *Tenm4* locus (*Tenm4^{CKO}* mice). From the same mice we used an ex vivo saphenous skin-nerve preparation to trace the trajectory of single units by means of an electrical stimulus until the point of exit from the nerve branch. Normally, single identified Aβ- and Aδ-fibers always have a mechanosensitive receptive field in the skin near to the nerve exit point, which we confirmed here in control mice (*Tenm4^{fl/fl}*) (Figure 2B).[4,12,35,36] However, blinded recordings made from *Tenm4^{CKO}* mice revealed that >75% of the Aβ-fibers and 60% of Aδ-fibers had no detectable mechanosensitive receptive field (Figure 2B). Recordings from C-fiber nociceptors revealed no significant differences in the presence of mechanosensitive receptive fields between control and *Tenm4^{CKO}* mice (Figure 2B). The deletion of the *Tenm4* gene in the majority of DRG neurons did not lead to axon loss as determined with transmission electron microscopy (Figure S3A) nor to any changes in axonal conduction velocity (Figures S3B and S3C).

Figure 2. TENM4 is required for mechanoreceptor function and touch sensation. (**A**) Left, schematic for generation of *Tenm4^{CKO}* mice via retro-orbital injection of AAV-PHP.S-Cre-EGFP of *Tenm4^{fl/fl}* mice. Top right, representative image of DRG sensory neurons from *Tenm4^{CKO}* mice 4 weeks after being transduced with AAV-PHP.S-Cre-EGFP. Scale bar is 50 µm. Bottom right, quantification of the percentage of GFP positive cultures DRG neurons after viral transduction. Dots represendata from 3 mice. (**B**) Electrical search technique (schematic) used to identify saphenous fibers with and without a mechanosensitive receptive field. *Tenm4^{CKO}* mice exhibited a substantial increase in Aβ-fibers and Aδ-fibers without a mechanosensitive receptive field, C-fibers remained unaffected. Significance calculated with Fisher's exact test, data from 5 *Tenm4^{fl/fl}* (control) and 10 *Tenm4^{CKO}* mice. (**C**) Behavioral tests on touch evoked behaviors. *Tenm4^{CKO}* mice exhibited reduced responses to a cotton swab and a dynamic brush. Additionally, *Tenm4^{CKO}* mice exhibited three-fold higher withdrawal thresholds and significant deficits in mean force required for

a 50% withdrawal using von Frey hairs, compared to controls. Data were collected from 15 *Tenm4*[fl/fl] and

15 *Tenm4*[CKO] mice. Significance calculated with Student's T-Test.

Expression of the virally expressed GFP marker in sensory neurons of *Tenm4*[CKO] mice indicated

that around 80% of the cells were transduced 4 weeks post-injection (Figure 2A). However, the

presence of Cre protein will not always lead to complete gene excision 4 weeks after transduction.

We therefore asked whether the remaining mechanosensitive fibers may have impaired function

due to incomplete ablation of the TENM4 protein in sensory neurons. We thus examined the

stimulus response characteristics of the remaining non-silenced sensory fibers identified using a

mechanical search stimulus in both glabrous and hairy skin.[16,37] We used a 20Hz vibrating stimulus

with continuously increasing amplitude to stimulate RAMs in hairy and glabrous skin. We found

that all RAMs displayed significantly reduced firing rates in *Tenm4*[CKO] mice compared to control

mice (Figure S3D). Velocity sensitivity was also measured using ramp stimuli of different

velocities and spike rates to these stimuli were also significantly impaired in RAMs from

Tenm4[CKO] mice (Figure S3E). Low threshold mechanoreceptors with Aδ-fiber conduction

velocities, so called D-hair receptors, showed significantly higher vibration thresholds in

Tenm4[CKO] mice compared to controls (Figure S3D). Merkel cells are innervated by slowly adapting

mechanoreceptors (SAMs) and are found in hairy and glabrous skin. Stimulus response functions

calculated with a series of 10s long ramp and hold stimuli showed that SAMs fired significantly

less and had higher mechanical thresholds for activation in *Tenm4*[CKO] mice compared to controls

(Figures S3E and S3F). We also recorded from Aδ mechanonociceptors (AMs) which detect

sustained high threshold mechanical stimuli and contribute to fast mechanical pain sensation.[38] In

Tenm4[CKO] AMs displayed reduced sustained firing to high forces and stimulus response functions

were reduced, but this was not significantly different compared to controls (Figures S3F and S3G).

Taken together, our data show that normal levels of TENM4 are necessary for the mechanosensory

function of almost all cutaneous sensory receptors, with the exception of unmyelinated C-fibers

(Figure 2B). We predicted that the dramatic functional loss of sensory neuron mechanosensitivity

in *Tenm4^{CKO}* should lead to profound behavioral deficits, especially to touch. Indeed, *Tenm4^{CKO}*

mice displayed substantially reduced responses to a cotton swab and dynamic brush stimuli (Figure

2C). Additionally, paw withdrawal thresholds to a series of von Frey hairs were substantially

elevated in *Tenm4^{CKO}* mice (Figures 2C and S4A). Sensory deficits varied in degree between

individuals probably due to differences efficiency of gene excision, but in many cases, animals

completely failed to respond to mechanical stimuli indicating a profound loss of sensation (Figures

2C and S4A).

We confirmed these results using a complementary strategy in which we generated mice in which

the *Tenm4$^{fl/fl}$* alleles were crossed with mice carrying an *Advillin-CreERT2* allele[4] allowing the

inducible deletion of *Tenm4* only in sensory neurons following tamoxifen treatment. Around 4

weeks after tamoxifen injections *Avil^{CreERT2}:Tenm4$^{fl/fl}$* mice were tested for paw withdrawal using

a series of von Frey hairs of increasing force. As for virus injected mice, these treated animals

displayed profoundly elevated paw withdrawal thresholds (Figure S4C). Using a radiant heat

stimulus, we measured thermal thresholds in both *Avil^{CreERT2}:Tenm4$^{fl/fl}$* and virus injected

Tenm4^{CKO} mice and found no significant deficits (Figures S4B and S4C). Loss of touch sensation

may also lead to altered coordination and indeed we noted that *Tenm4^{CKO}* mice showed hindlimb

clasping that was rarely seen in control mice (Figure S4D). We found that essentially all

Parvalbumin positive DRG soma were also stained positive for TENM4 (Figure S4E),

Parvalbumin is a marker for proprioceptors, although it also probably found in some cutaneous

sensory neurons.[39,40] We generated conditional *Tenm4* mutant in which the gene was deleted in

proprioceptors by crossing our *Tenm4$^{fl/fl}$* mice with a *Parvalbumin-Cre* line (*Pv-Cre:Tenm4$^{fl/fl}$*

mice).[41] We found that *Pv-Cre:Tenm4$^{fl/fl}$* mice showed mild impairments in their ability to remain

on a rotating wheel (rotarod) test compared to controls (*Tenm4$^{fl/fl}$* mice) (Figure S4D). These

deficits could be due to alterations in proprioceptors, but may also reflect the reduced input from

cutaneous afferents that also contribute to normal proprioception.[39,40,42] This prompted us to ask

whether we could detect TENM4 protein at the endings of proprioceptors that innervate the muscle

spindle. We visualized proprioceptor endings in muscle spindles from the mouse gastrocnemius

muscle using *Pv^{cre};Rx3^{Flpo};Ai65D* mice[43] (see methods). We failed to detect TENM4 signal in

identified proprioceptor endings (Fig. S4F). It thus appears likely that motor coordination deficits

seen following conditional *Tenm4* deletion were due to a loss in cutaneous sensory feedback in

these animals. These data also suggest that not all sensory neurons that express TENM4 can

transport the protein to the sensory end-organ.

TENM4 cleavage abolishes touch sensitivity

Cre-mediated excision using AAV vectors leads to the removal of *Tenm4* over several weeks.

TENM4 has well described roles in intercellular signaling,[30,44] the alteration of which could have

indirect effects on transduction. Therefore, we wished to directly test the idea that TENM4 forms

part of a complex that facilitates force transfer from the extracellular matrix (ECM) to transduction

channels. Ideally, specific cleavage of proteins linking the channel to ECM should abolish

transduction.[17] We thus developed a new technology that enabled us to specifically ablate TENM4

function within minutes. We named this method PreScission Protein Disassembly, and it is based

on the fact that the plant virus derived PreScission protease cleaves a specific consensus sequence

(LEVLFQ↓GP),[45] that rarely occurs in the extracellular exposed residues of mouse proteins (Table

S2). This method is conceptually similar to a method recently used in nematodes to genetically

target proteolytic cleavage of transduction molecules.[10] Using homologous recombination in

mouse ES cells we inserted the PreScission cleavage sequence into *Tenm4*. We replaced 9 codons

leading to the replacement of 7 amino acids which expose a PreScission sequence on the surface

of the extracellular YD shell domain of TENM4.[44] This manipulation was predicted to make

TENM4 newly susceptible to cleavage by the PreScission enzyme (Figures 3A and S2C). Mice

carrying the PreScission site in the *Tenm4* gene were bred, but out of 569 newborns from

heterozygote crosses we obtained no homozygote animals. The failure to obtain homozygotes

indicates that altering both *Tenm4* alleles likely causes embryonic lethality, consistent with

previous studies.[30,46] We thus decided to work with heterozygotes which we termed *Tenm4*$^{KI/+}$

mice. Considering the observed homozygote embryonic lethality, it was important to examine

whether *Tenm4*$^{KI/+}$ mice show deficits in baseline sensory function. *Tenm4*$^{KI/+}$ mice exhibited

uncleaved full-length TENM4 (Figure 3B), and showed no deficits in behavioral responses to

mechanical and thermal stimuli (Figures S5A-C). We also made recordings from the saphenous

nerve of *Tenm4*$^{KI/+}$ mice and, using the electrical search method, demonstrated that Aβ-fibers, Aδ-

fibers and C-fibers exhibited normal mechanosensitivity, with unaltered stimulus response

properties and normal axonal conduction velocities (Figures 3C and S6A). Thus, inclusion of

mutant TENM4 protein into the mechanotransduction complex does not impact on baseline

sensory function. Since TENM4 forms obligate homodimers,[29,44] a stochastic mixing of mutated

and wild type proteins would ensure that the vast majority of dimers would have at least one

PreScission cleavable site. Indeed, we directly confirmed that the PreScission enzyme cleaves

TENM4 in vivo. Western blots from glabrous skin revealed a 300 kDa band corresponding to

uncleaved TENM4 in WT skin treated with PreScission protease. However, in skin samples taken

from *Tenm4*$^{KI/+}$ mice treated with PreScission, we detected not only the full-length protein but also

a ~180 kDa fragment, consistent with the predicted size of the cleavage product (Figure 3D). Thus

endogenous dimeric complexes at sensory endings consist of both WT and PreScission cleavable

protein. We next applied the PreScission protease directly to the receptive fields of single identified

mechanoreceptors to ask whether acute TENM4 ablation can rapidly abolish mechanosensitivity

as we previously saw using the Blisterase enzyme.[17] Once a mechanoreceptor was identified and

classified,[37] the receptive field was isolated with a metal ring and repeatedly stimulated every

minute for up to 45 mins. We used a vibration stimulus (20Hz gradually increasing to an amplitude

of 100 mN) for all LTMRs (RAMs, SAMs and D-hair receptors) or a 150mN amplitude ramp and

hold stimulus for mechanonociceptors (AMs or C-fiber nociceptors). After baseline recordings for

three minutes the receptive field was exposed to the PreScission enzyme (2 units/µL) in

physiological ringer solution (Figure 3E). Protease treated LTMRs from WT mice showed

remarkably robust spiking for the entire 45 mins with barely any diminution of mechanosensitivity

(Figures 3E and 3F). In contrast, all LTMRs (RAMs, SAMs and D-hair receptors) recorded from

$Tenm4^{KI/+}$ mice showed progressively decreasing spike responses to the constant mechanical

stimuli starting just minutes after protease application, until no response was evoked between 30

and 45 minutes (Figure 3F). Complete loss of mechanosensitivity was also seen in RAMs

innervating the Meissner's corpuscles in glabrous skin (Figure 3F). For each mechanoreceptor

examined we used an electrical stimulus at the end of the experiment and could show in every case

that the electrically evoked spike was always still present, indicating a specific loss of

mechanosensitivity. The rapid loss of LTMR mechanosensitivity could in theory be due to a toxic

effect of the PreScission cleavage product on the mechanoreceptors. This idea was directly tested

by adding the fluid containing the cleavage product from PreScission treated $Tenm4^{KI/+}$ skin to the

isolated receptive fields of WT neurons (Figure S7A). The buffer containing the cleavage product

had no effect on the mechanosensitivity of WT RAMs (Figure S7A).

Figure 3

Figure 3. Rapid loss of mechanoreceptor function following TENM4 cleavage. (**A**) Generation of *Tenm4KI/+* mice: insertion of a 7 residue PreScission consensus sequence into genomic locus encoding the TENM4 extracellular domain. (**B**) western blot of endogenous TENM4 from glabrous skin revealed uncleaved full-length TENM4 in WT and *Tenm4KI/+* mice (**C**) Electrical search technique revealed that most

Aβ, Aδ, and C-fibers retained mechanosensitivity in *Tenm4^{KI/+}* mice as in WT mice. Significance calculated

with Fisher's exact test. Data collected from 4 WT (control) and 5 *Tenm4^{KI/+}* mice. (**D**) Left, experimental

design. Right, western blot of endogenous TENM4 from glabrous skin after PreScission protease treatment

revealed uncleaved full-length TENM4 in WT mice, a cleaved product of the predicted size was detected

in *Tenm4^{KI/+}* mice. Loading control β-actin. (**E**) Representative example of a single unit (RAM) responding

to repeated 20Hz vibration before and after application of PreScission to the receptive field in a WT (left)

or *Tenm4^{KI/+}* (right) mouse. Note the progressive decrease in stimulus induced firing following PreScission

treatment only in *Tenm4^{KI/+}* (right) mouse. (**F**) Mean mechanoreceptor responses plotted as a percentage of

the starting value for all myelinated Aβ and Aδ types, before and after PreScission. Note mechanoreceptors

in WT all retained mechanosensitivity (grey), but all mechanoreceptor types in *Tenm4^{KI/+}* mice,

progressively lost responsiveness during the 45 mins of observation (blue). (**G**) Experimental design. (**H**)

Electrical search 1 hour after PreScission protease injections. Substantial and significant decrease in the

proportion of Aβ and Aδ-fibers displaying mechanosensitivity in PreScission treated *Tenm4^{KI/+}*, but not in

WT mice. The proportion of mechanosensitive C-fibers remained unchanged in both groups. Significance

calculated with Fisher's exact test. Data collected from 4 WT and 4 *Tenm4^{KI/+}* mice. (**I**) Behavioral assays

used to assess the effects of TENM4 cleavage 1 hr after PreScission injection to the plantar hind paw.

Mechanical withdrawal threshold measured with von Frey hairs were unchanged in WT mice treated with

saline or PreScission. Additionally, *Tenm4^{KI/+}* mice injected with PreScission showed decreased responses

to both a cotton swab and a dynamic brush, no changes were observed in treated WT mice. Data collected

from 13 WT and 12 *Tenm4^{KI/+}* mice. Two-way repeated measures ANOVA followed by Sidak's post-test.

ns indicates $p > 0.05$; * $p < 0.05$; ** $p < 0.01$; ***; $p < 0.001$, means ± s.e.m.

Notably, we observed a slower, but robust loss of the ability of AMs in *Tenm4^{KI/+}* to transduce

force, reaching its maximum 45 mins after PreScission application (Figure 3F). However, in

contrast to LTMRs, most AMs retained residual responses to intense mechanical stimuli at the end

of the 45-minute observation period (5/6 tested). C-fiber nociceptors in WT and *Tenm4^{KI/+}* mice

retained functionality after exposure to PreScission protease (Figure S7B), indicating a

fundamentally distinct transduction mechanism in AM nociceptors compared to C-fiber

nociceptors.

We next performed behavioral and electrophysiological assays on mice subjected to subcutaneous

injections of the PreScission enzyme into either the glabrous skin or hairy skin innervated by the

saphenous nerve (Figure 3G). One hour after subcutaneous injections of the PreScission enzyme

we observed a substantial and significant decrease in the proportion of Aβ- and Aδ-fibers with

mechanosensitive receptive fields in *Tenm4^{KI/+}* mice compared to WT mice in the saphenous nerve

(Figure 3H). Not all mechanoreceptors lost their mechanosensitivity, likely because the subcutaneously injected enzyme did not reach the entire innervation territory of the saphenous nerve. However, no loss of C-fiber mechanosensitivity was observed in either genotype treated with the PreScission protease (Figures 3H and S7B). We next asked whether touch evoked behaviors which are routinely measured using stimuli to the glabrous skin are impacted by the application of the PreScission protease. In WT mice PreScission injections into the glabrous skin had no significant impact on paw withdrawal thresholds to von Frey stimuli, cotton swab or brush responses. However, *Tenm4*$^{KI/+}$ mice exhibited a substantial and significant increase in paw withdrawal thresholds as well as much reduced responses to cotton swab and brush stimuli (Figures 3I and S7C). Noxious heat withdrawal latencies were not significantly impacted after PreScission injection in *Tenm4*$^{KI/+}$ mice compared to saline injected controls (Figure S7D).

TENM4 is essential for touch sensation

Cleavage of the TENM4 protein clearly leads to an acute loss of touch sensation in *Tenm4*$^{KI/+}$ mice, but in theory the TENM4 protein should be re-synthesized to reinstate function. We thus investigated the temporal dynamics of mechanosensitivity and touch sensation recovery in *Tenm4*$^{KI/+}$ mice following cutaneous PreScission injections (Figure 4A). We found that a substantial proportion of Aβ and Aδ-fibers were still mechanically silent 24h following PreScission injection in *Tenm4*$^{KI/+}$ mice. However, at 72h almost all Aβ and Aδ-fibers had mechanosensitive receptive fields, as was observed in PreScission injected WT control mice at 24 and 72h (Figure 4B). C-fiber nociceptors identified with an electrical search retained normal mechanosensitivity at all time points and conditions (Figure 4B). To assess how this recovery manifests behaviorally, we monitored paw withdrawal thresholds over time following PreScission injection. Consistent with the electrophysiological data, we found that PreScission-injected *Tenm4*$^{KI/+}$ mice displayed elevated paw withdrawal thresholds at 12, 24, and 48h which had then

355 returned to normal levels at 72h (Figure 4C). Control WT mice injected with the PreScission

356 protease showed no changes in paw withdrawal thresholds over the same period of time (Figure

357 4C). Together, these results reveal that in sensory endings the TENM4 protein is likely both

358 necessary and sufficient for touch transduction.

359
Figure 4. TENM4 is at sensory endings is necessary and sufficient for touch sensation. (**A**)
Experimental design. (**B**) *Tenm4$^{KI/+}$* mice exhibit a significant decrease in the proportion of Aβ and Aδ
fibers displaying mechanosensitivity only after 24h, but not at 72h, following PreScission injection, while
C-fibers remain unaffected at both time points. All WT fibers showed no change. Fisher's exact test. Data
were collected from 4 mice per group. (**C**) WT mice showed constant mean withdrawal thresholds to von
Frey filaments before and up to 72h following PreScission treatment. In contrast, PreScission-injected
Tenm4$^{KI/+}$ mice displayed significantly elevated paw withdrawal thresholds at 12, 24, and 48 hours, which
returned to normal by 72h. Statistical significance calculated with Friedman test followed by Dunn's
multiple comparisons test. Data collected from 10 mice per group.

TENM4 is necessary and sufficient for MA-currents.

To connect behavioral recovery with underlying transduction mechanisms, we assessed

mechanically activated currents in identified sensory neurons. Cultured large-diameter sensory

neurons possess sensitive and large amplitude indentation-induced currents that are also sensitive

to protease treatment.[12,17,47] We made whole cell patch clamp recordings and measured MA

currents in isolated large sensory neurons (diameter >30μm) from the two mouse models

(*Tenm4^{CKO}* and *Tenm4$^{KI/+}$*) using cell indentation (Figures 5A and 5B). As expected almost all

large neurons from control mice positive for GFP (*Tenm4^{fl/fl}* injected with AAV-PHP.S-eGFP

virus, lacking Cre) displayed robust MA currents (Figure 5A)[12,47]. In contrast, over half of the large

diameter GFP-positive neurons recorded from *Tenm4^{CKO}* mice lacked any MA current (Figure 5A).

However, the incidence and characteristics of MA currents recorded from small diameter sensory

neurons, which give rise to unmyelinated C-fibers, were comparable between neurons from

Tenm4^{CKO} mice and controls (Figure S8A). Similarly, a substantial proportion of PreScission-

treated large-diameter DRG neurons in *Tenm4^{KI/+}* showed no MA current which was significantly

different from WT treated neurons (Figure 5B). Importantly, the loss of MA-currents was not

accompanied by detectable changes in electrical excitability as reflected by unchanged resting

membrane potentials or action potential thresholds in sensory neurons isolated from both mouse

models (Figures S8B and S8C).

Figure 5

Figure 5. TENM4 is necessary and sufficient for MA-currents. (**A**) Poking-induced currents in GFP
positive sensory neurons infected with the AAV-PHP.S-Cre-EGFP. Scale bar 20μm. *Tenm4^{CKO}* mice had a
deficit in MA-currents in large neurons compared to control *Tenm4^{fl/fl}* mice. Significance calculated with
Chi-squared test. Data collected from 4 *Tenm4^{fl/fl}* and 5 *Tenm4^{CKO}* mice. (**B**) Recordings from large diameter
sensory in culture (scale bar = 20μm) showed a substantial decrease in the proportion of cells with poking-

induced currents, specifically in cells from *Tenm4$^{KI/+}$* treated with the PreScission protease. Significance
calculated with Chi-squared test. (**C**) Cartoon of pillar stimulation. Example of transfected N2a$^{Piezo1-/-}$ cells
expressing *Tenm4-ALFA* (red) and *pRK8-eGFP* (black) on pillar array. Scale bar 5 μm. (**D**) Representative
pillar stimulation with current traces evoked from single pilus movements in N2a$^{Piezo1-/-}$ cells expressing
Tenm4-ALFA (red) compared to control *pRK8-eGFP* transfected cells (black). Deflection-current amplitude
plot showing large increase in the amplitude and sensitivity of MA-currents in cells transfected with
TENM4 (red) compared to control (black). Statistical significance calculated with Kruskal-Wallis test.
Right shows that MA-currents were evoked at much higher frequencies, 20.7± 4.2 % in *Tenm4-ALFA*
expressing cells, compared to controls, 7.9 ± 3.4%. Statistical significance calculated with Mann-Whitney
test.
404

Ion channels like PIEZO2 and ELKIN1[48,49] are efficiently gated by substrate deflection and we

asked whether TENM4 can modulate this gating.[3,12,49] We generated full length *Tenm4* expression

constructs with a C-terminal ALFA-tag that can be recognized by fluorescently labeled nanobodies

(Figure S9A).[50] We used Neuro2a-PIEZO1-Knockout (N2a$^{Piezo1-/-}$) cells that have minimal, but

detectable deflection induced currents,[51,52] and transfected these cells with the *Tenm4-Alfa* or

control *pRK8-eGFP* constructs. Transfected cells were plated on pillar arrays and then live labelled

with anti-ALFA tag nanobodies which bind to the extracellular C-terminus of the expressed

TENM4 protein. We made targeted whole cell patch clamp recordings from cells that expressed

TENM4 on the membrane and measured the sensitivity and amplitude of currents evoked by single

pillar displacements (Figure 5C). As observed previously, control GFP transfected cells exhibited

occasional small amplitude currents to pillar deflection[51,52]. Strikingly, we observed large

amplitude mechanically-gated currents in cells with plasma membrane-localized TENM4, and a

quantitative analysis revealed a large increase in the incidence, amplitude and frequency of pilli-

evoked currents in these cells compared to GFP controls (Figure 5D).

This data suggests that N2a$^{Piezo1-/-}$ cells normally express MA channels that interact with TENM4

to facilitate gating of channels via the extracellular matrix. We therefore carried out a quantitative

proteomic analysis of N2a cells to reveal expressed ion channels potentially targeted by TENM4.

We compared the proteome of N2a cells with that of N2a$^{Piezo1-/-}$ cells and could quantify the levels

of more than 8000 proteins in both samples (Figures S9B and S9C). Interestingly, a large

proportion of the proteome showed significant changes in the absence of PIEZO1, e.g. >16% of

the proteome showed significantly higher abundances. Included in the significantly up-regulated

protein set were TENM4 itself as well as the MA ion channel ELKIN1.[12,49] We did not detect any

peptides from PIEZO2, suggesting that unlike ELKIN1 this channel is not expressed in N2A cells.

Interestingly, many of the highest up-regulated proteins were themselves components of the

extracellular matrix.

TENM4 associates with mechanically activated ion channels

Given the TENM4 dependence of MA currents, we examined whether TENM4 physically

associates with candidate mechanosensitive channels like PIEZO2 or ELKIN1. Due to the

structural similarity among Teneurins especially in the C-terminal domain, we designed a

truncated construct covering the entire N-terminus and a portion of the extracellular region

including sequences involved in dimerization to assess binding between Teneurin family members.

Utilizing a tripartite-GFP-based protein complementation assay in HEK293T cells, we confirmed

that the Teneurins can interact with each other (Figures S10A and S10B). We also measured a

robust complementation signal (GFP fluorescence) between ELKIN1 and TENM4, and the

strength of the interaction signal was notably much higher than that observed for ELKIN1 with

other Teneurins (Figures S10C and S10D). We next asked whether TENM4 is associated with MA

channels on the plasma membrane using Airyscan super-resolution microscopy. In N2A cells we

co-expressed our ALFA-tagged *Tenm4* construct with either mScarlet-tagged Elkin1 or Piezo2

constructs. Nanobody-labeled TENM4 exhibited a punctate distribution on the plasma membrane

as did mScarlet-tagged ELKIN1 or PIEZO2 (Figures 6A and S10E). Quantification indicated that

approximately 75% of TENM4 was colocalized with ELKIN1, while around 30% of TENM4

localized together with PIEZO2 channels (Figures 6A and S10E). We next used HEK-293 cells

transfected either with *Elkin1* or *Piezo2* expression plasmids and ALFA-tagged *Tenm4*. We used

ALFA nanobody conjugated resins (ALFA SelectorPE^{TM}) to pull down the expressed TENM4

protein complexes which were then eluted using high concentrations of the purified ALFA peptide.

Using Western blotting, we could detect co-elution of ELKIN1 specifically in cases where

TENM4-ALFA protein had been expressed, but not PIEZO2 (Figures 6B and S10F). Together,

these results demonstrate specific and direct interactions between TENM4 and MA channels like

ELKIN1. Our data does not exclude a potential interaction between PIEZO2 and TENM4 in

sensory neurons as it is possible that molecular factors necessary for this interaction are not present

in HEK293 or N2A cells.

These data suggested that the clear increase in MA currents in N2a$^{Piezo1-/-}$ cells overexpressing

TENM4 could be due to increased interaction with endogenous ELKIN1 channels. We tested this

idea by measuring deflection induced currents in N2a$^{Piezo1-/-}$ cells treated with siRNAs targeting

the *Elkin1* transcript. Knockdown of *Elkin1* transcripts in N2a$^{Piezo1-/-}$ cells over expressing TENM4

did not prevent the dramatic increase in MA currents in these cells compared to controls (Figure

S10F,G). This negative result may mean that TENM4 increases mechanosensitive currents in N2A

cells via interaction with an as yet unknown ion channel. However, it is also possible that siRNA

targeting in this case was not sufficiently efficient to remove all the ELKIN1 protein in these cells.

Figure 6

464
Figure 6. TENM4 interacts with ELKIN1. (**A**) Representative super-resolution images of N2a cells co-
transfected with a *Tenm4-ALFA* expression plasmid with mScarlet-ELKIN1 (purple in middle), arrows
indicate co-localization signals. Right higher magnification of co-localized protein region (from white dash
boxes). Scale bar is 10μm for cells, 1μm magnified region. Quantification was carried out and a co-
localization correlation, ranging from -1 to 1, signifies the strength of overlap. Values near 1 indicate strong
co-localization, near -1 indicate separation. TENM4-ELKIN1 displays strong co-localization. Right,
TENM4 shows 75% overlap with ELKIN1 on the cell surface. (**B**) Co-elution of TENM4-ALFA and
ELKIN1 using an ALFA selector resins. Western blot of ALFA pull-down elution from HEK293T cell
transfected with TENM4-ALFA and ELKIN1 (right panel) or ALFA vector plasmid and ELKIN1 (left
panel). * indicates $p < 0.05$; ** indicates $p < 0.01$; *** indicates $p < 0.001$ means ± s.e.m.
475

TENM4 is a part of a mechanosensory tether

These molecular interactions prompted us to test a structural model in which TENM4 links

mechanically activated (MA) channels to the extracellular matrix within discrete membrane

domains. We have already shown that large extracellular tethers made by mouse sensory neurons

in culture appear to be necessary for the expression of mechanosensitive currents.[17] Furthermore,

specific cleavage of TENM4 in mouse sensory neurons is sufficient to abolish mechanosensitive

currents (Figure 5B). We therefore next asked if we could localize the protein to matrix-neurite

structures in cultured mouse sensory neurons. We performed immunogold labeling with our

antibody targeting the extracellular domain of TENM4 followed by transmission electron

microscopy (TEM). Remarkably, immunogold labelled TENM4 was localized at the neurite-

laminin interface, often associated with filamentous extracellular tether-like structures (Figures

7A, 7B, S11A, and S11B; arrowheads). Over 90% of the gold particles within 200 nm of the neurite

membrane were localized to the neurite laminin interface, with only a small fraction found

elsewhere, indicating strong enrichment of TENM4 at sites of extracellular contact (Figure 7C;

423 at interface vs. 45 outside). Analysis of 215 neurites revealed that while the majority of

neurites harbored a single gold-labeled tether, more than 30% of neurites displayed multiple

labelled tether structures in close proximity to each other on the same neurite (Figure 7B,D). This

latter finding suggests that TENM4-positive tethers may often be enriched in discrete domains

along the neurite (Figure 7D). Of the 325 total gold-labelled tethers analyzed, most (~70%) were

labeled with a single gold particle, however a notable fraction of labelling (~30%) consisted of

multiple gold particles on a single structure. Additionally, we calculated a density of ~2 gold-

labeled tethers/µm² of neurite laminin interface (Figures S11A and S11B), consistent with our

earlier findings that contrast enhanced tether structures have a density of ~ 2-4/μm^2.[17] These

findings provide compelling ultrastructural evidence that TENM4 is a key component of tether-

like structures at the neurite–laminin interface (Figure 7F).

Figure 7. TENM4 is localized at sensory tethers. (**A**) Immunofluorescence of mouse sensory neurons stained with the custom antibody recognizing extracellular epitopes. TENM4 displays a punctate membrane pattern. Scale bar is 20μm. (**B**) Representative TEM images of two neurites showing variable Tenm4 labeling. The left neurite contains two tethers, one with a single gold particle and one with four, while the right neurite shows three tethers with one, two, and three gold particles, respectively (arrowheads) Scale bar 100 nm (**C**) Gold particles within 200 nm of the neurite membrane are highly enriched (>90%) at the neurite-laminin interface. (**D**) ~70% of neurites have a single gold-labeled tether; ~30% have multiple. (**E**) ~70% of tethers carry one gold particle; ~30% have multiple. (**F**) Cartoon illustrating how TENM4 transmits extracellular mechanical forces to ELKIN1 channels to drive mechanosensitive currents. Loss of TENM4 or its proteolytic cleavage with a protease (Green Pacman) disrupts this mechanical coupling and reduces mechanosensitive currents.

DISCUSSION

Almost all multicellular organisms respond rapidly to the tiniest of mechanical cues. Classic

genetic screens in model organisms like *Drosophila melanogaster* and *Caenorhabditis elegans*

have shown that not just mechanically gated ion channels are required for mechanoreceptor

function, but also a number of extracellular matrix proteins[6,9,53,54]. Body touch receptors in *C.*

elegans are probably the best characterized mechanoreceptors at the molecular level and here at

least a dozen proteins are required for normal mechanotransduction including three ECM proteins

MEC-1, MEC-5 and MEC-9[9,53–55]. Recently, compelling evidence was provided that the Kunitz

domain protein MEC-1 may link the mechanosensitive ion channel MEC-4 to specialized basal

lamina at sites of transduction.[9] Mammalian cutaneous touch receptors are diverse in terms of end-

organ morphology[14], but also in terms of the mechanosensitive ion channels required for their

function. Thus in mice, genetic ablation of genes encoding the mechanically activated channels

PIEZO2 or ELKIN1 or their modulator STOML3 leads to silencing of mechanoreceptor function,

but in each case large sub-populations of mechanoreceptors retain normal function.[4,12,35,36] In flies

and worms genetic rescue of touch insensitivity has been routinely shown to provide evidence that

the ion channel and ECM components are directly involved in transduction.[6,53,54,56–58] Genetic

rescue experiments in mouse models are more difficult, but in some cases reintroduction of

mechanotransduction genes into isolated sensory neurons was shown to reconstitute

mechanosensitive currents.[12,35] In the present study we used an unbiased approach to identify a

novel component of mammalian touch namely the type II membrane protein TENM4. Using

sensory neuron specific *Tenm4* gene deletion in adult mice we demonstrate that presence of this

protein is essential for transduction in virtually all mechanoreceptors that innervate both hairy and

glabrous skin (Figure 2). In addition, genetic ablation of *Tenm4* also led to the functional silencing

of fast conducting mechanonociceptors, but had no discernible effect on unmyelinated C-fiber

afferents (Figure 2). These phenotypes were consistent with the relatively restricted expression of

TENM4 in sensory neurons with myelinated axons (Figure 1). There were small diameter neurons

positive for TENM4 e.g. small diameter tyrosine hydroxylase positive neurons (Figure 1D).

However, our data on Parvalbumin positive proprioceptors suggested that not all sensory neurons

are capable of transporting TENM4 to sensory terminals where transduction takes place.

TENM4 has very well characterized signaling roles within the developing CNS [59] and so it was

important to establish that the profound sensory phenotypes we observed were not due to a failure

of inter-cellular signaling at neuroglial endings. For this reason, we established a novel technology

termed PreScission Protein Disassembly which relies on the use of a highly specific protease to

acutely manipulate proteins in vivo with near perfect specificity. We show that acute proteolytic

cleavage of TENM4 molecules at the site of transduction is sufficient to completely abolish

mechanoreceptor function within minutes (Figure 3). After PreScission cleavage of TENM4, it

took 72 hours for newly synthesized TENM4 to restore both mechanoreceptor function and normal

touch evoked behavior. The rescue of mechanoreceptor function with endogenously expressed un-

cleaved TENM4 demonstrated that the presence of intact TENM4 at the sensory ending is both

necessary and sufficient for touch receptor function (Figure 4).

We hypothesized that there are protein tethers that couple mechanically gated ion channels to the

ECM to facilitate fast mechanotransduction. Using cultured sensory neurons as a model we had

previously shown that large tether like proteins that connect the sensory neuron membrane with

the ECM appeared to be necessary for fast mechanotransduction.[17,22] These tether-like structures

were ablated by enzymes that cleave furin consensus sequences and this formed the basis of our

unbiased screen which led to the identification of TENM4. Using PreScission Protein Disassembly

we now show that fast mechanosensitive currents in isolated sensory neurons depend on intact

TENM4 (Figure 5). This data strongly suggests that protein tethers synthesized by sensory neurons

are at least in part made up of TENM4 molecules. Indeed, using immunogold TEM we show a

dense localization of TENM4 at the neurite–laminin interface, where it clusters along filamentous

tethers. These findings identify TENM4 as a core structural element of a putative mechanosensory

tether linking the ECM to mechanosensitive channels. Interestingly, the density of TENM4

positive tethers was very similar to our previous estimates of tether density.[17,22] Gold labelling of

tethers was often found very close to the neurite membrane, consistent with the known compact

structure of the extracellular teneurin domain, recognized by our antibody.[44,60] However, we also

frequently observed multiple gold labels along filaments often at distances longer than the

dimensions of a single extracellular domain (~30 nm) (Figure 7). This observation suggests that

some of the labelled TENM4 may undergo regulated cleavage and then be incorporated into

tethers[59]. It is intriguing in this context to note that a recent study on fly Teneurin-M suggested

that this protein can form oligomeric zippers via protein interfaces.[61] Cultured sensory neurons

have served as a very useful model for mechanotransduction at the sensory ending. However, it is

increasingly clear that specialized sensory Schwann cells at a variety of neuroglial endings actively

participate in the transduction of mechanical stimuli.[15,62,63] This means that examining sensory

cells alone may only reveal part of the mechanistic basis of transduction.[15,64] Here we show that

TENM4 is synthesized by sensory neurons and is present at their endings in a variety of end-organ

structures including Meissner's corpuscles, Merkel cell complexes and lanceolate endings around

hair follicles (Figure 1). Indeed, acute TENM4 cleavage abolishes mechanotransduction at all

these sensory endings in the skin (Figure 3). It was striking that TENM4 integrity is required for

transduction at such a wide variety of sensory endings suggesting that this protein is truly central

to virtually all touch transduction.

One implication of our findings is that there is likely a connection between TENM4 molecules and

mechanically gated ion channels. Consistently, we show that in a heterologous expression system

TENM4 strongly associates with the mechanically activated ion channel ELKIN1 (Figure 6). In

contrast, we were not able to co-purify PIEZO2 protein with TENM4 in the same system, but this

does not necessarily mean there is no interaction in vivo. Indeed, it will be important to determine

what other proteins TENM4 interacts with in sensory neurons given it's central role in touch

transduction. We also made the striking observation that introduction of TENM4 into N2a[*Piezo1-/-*]

cells was sufficient to induce mechanically activated currents, however, we were not able to

unequivocally determine whether this effect was exerted via ELKIN1 or by other ion channels

expressed in these cells.

Teneurins play essential roles in nervous system development, in particular during neuronal

migration, oligodendrocyte differentiation, and establishment of neural circuits.[44,65] Our data

provide an unexpected and surprising link between one of these proteins and fast

mechanotransduction, as we show that the integrity of TENM4 is absolutely necessary for almost

all somatic sensation. Indeed, the mechanisms by which Teneurins coordinate neurodevelopmental

events may well include interactions with mechanosensitive ion channels the roles of which in

development have only just started to be investigated.[66] In conclusion, here we have identified a

new essential component of touch transduction with a novel methodology that allowed us to

definitively establish an essential role for intact TENM4 in rapid touch transduction. The central

role of TENM4 for mechanotransduction in virtually all myelinated mechanoreceptors makes this

molecule an ideal starting point to identifying further touch transduction components.

References

1. Cox, C.D., Bavi, N., and Martinac, B. (2018). Bacterial Mechanosensors. Annual Review of
Physiology *80*, 71–93. https://doi.org/10.1146/annurev-physiol-021317-121351.

2. Jin, P., Jan, L.Y., and Jan, Y.N. (2020). Mechanosensitive Ion Channels: Structural Features
Relevant to Mechanotransduction Mechanisms. Annu Rev Neurosci *43*, 207–229.
https://doi.org/10.1146/annurev-neuro-070918-050509.

3. Poole, K., Herget, R., Lapatsina, L., Ngo, H.D., and Lewin, G.R. (2014). Tuning Piezo ion
channels to detect molecular-scale movements relevant for fine touch. Nature
communications *5*, 3520. https://doi.org/10.1038/ncomms4520.

4. Ranade, S.S., Woo, S.H., Dubin, A.E., Moshourab, R.A., Wetzel, C., Petrus, M., Mathur, J.,
Bégay, V., Coste, B., Mainquist, J., et al. (2014). Piezo2 is the major transducer of mechanical
forces for touch sensation in mice. Nature *516*, 121-U330.
https://doi.org/10.1038/nature13980.

5. Qiu, X., and Müller, U. (2022). Sensing sound: Cellular specializations and molecular force sensors. Neuron *110*, 3667–3687. https://doi.org/10.1016/j.neuron.2022.09.018.

6. Chung, Y.D., Zhu, J., Han, Y., and Kernan, M.J. (2001). nompA encodes a PNS-specific, ZP domain protein required to connect mechanosensory dendrites to sensory structures. Neuron *29*, 415–428.

7. Goodman, M.B., and Sengupta, P. (2019). How Caenorhabditis elegans Senses Mechanical Stress, Temperature, and Other Physical Stimuli. Genetics *212*, 25–51. https://doi.org/10.1534/genetics.118.300241.

8. Göpfert, M.C., Humphris, A.D.L., Albert, J.T., Robert, D., and Hendrich, O. (2005). Power gain exhibited by motile mechanosensory neurons in Drosophila ears. Proc Natl Acad Sci U S A *102*, 325–330. https://doi.org/10.1073/pnas.0405741102.

9. Das, A., Franco, J.A., Mulcahy, B., Wang, L., Chapman, D., Jaisinghani, C., Pruitt, B.L., Zhen, M., and Goodman, M.B. (2023). Conserved basal lamina proteins, laminin and nidogen, are repurposed to organize mechanosensory complexes responsible for touch sensation. Preprint at bioRxiv, https://doi.org/10.1101/2022.02.11.479800 https://doi.org/10.1101/2022.02.11.479800.

10. Sanfeliu-Cerdán, N., Català-Castro, F., Mateos, B., Garcia-Cabau, C., Ribera, M., Ruider, I., Porta-de-la-Riva, M., Canals-Calderón, A., Wieser, S., Salvatella, X., et al. (2023). A MEC-2/stomatin condensate liquid-to-solid phase transition controls neuronal mechanotransduction during touch sensing. Nat Cell Biol *25*, 1590–1599. https://doi.org/10.1038/s41556-023-01247-0.

11. Zhang, W., Cheng, L.E., Kittelmann, M., Li, J., Petkovic, M., Cheng, T., Jin, P., Guo, Z., Göpfert, M.C., Jan, L.Y., et al. (2015). Ankyrin Repeats Convey Force to Gate the NOMPC Mechanotransduction Channel. Cell *162*, 1391–1403. https://doi.org/10.1016/j.cell.2015.08.024.

12. Chakrabarti, S., Klich, J.D., Khallaf, M.A., Hulme, A.J., Sánchez-Carranza, O., Baran, Z.M., Rossi, A., Huang, A.T.-L., Pohl, T., Fleischer, R., et al. (2024). Touch sensation requires the mechanically gated ion channel ELKIN1. Science *383*, 992–998. https://doi.org/10.1126/science.adl0495.

13. Sánchez-Carranza, O., Chakrabarti, S., Kühnemund, J., Schwaller, F., Bégay, V., García-Contreras, J.A., Wang, L., and Lewin, G.R. (2024). Piezo2 voltage-block regulates mechanical pain sensitivity. Brain *147*, 3487–3500. https://doi.org/10.1093/brain/awae227.

14. Abraira, V.E., and Ginty, D.D. (2013). The sensory neurons of touch. Neuron *79*, 618–639. https://doi.org/10.1016/j.neuron.2013.07.051.

15. Ojeda-Alonso, J., Calvo-Enrique, L., Paricio-Montesinos, R., Kumar, R., Zhang, M.-D., Poulet, J.F.A., Ernfors, P., and Lewin, G.R. (2024). Sensory Schwann cells set perceptual thresholds

for touch and selectively regulate mechanical nociception. Nat Commun *15*, 898.
https://doi.org/10.1038/s41467-024-44845-8.

16. Schwaller, F., Begay, V., Garcia-Garcia, G., Taberner, F.J., Moshourab, R., McDonald, B.,
Docter, T., Kuhnemund, J., Ojeda-Alonso, J., Paricio-Montesinos, R., et al. (2021). USH2A is a
Meissner's corpuscle protein necessary for normal vibration sensing in mice and humans.
Nat Neurosci *24*, 74–81. https://doi.org/10.1038/s41593-020-00751-y.

17. Hu, J., Chiang, L.Y., Koch, M., and Lewin, G.R. (2010). Evidence for a protein tether involved
in somatic touch. Embo J *29*, 855–867. https://doi.org/10.1038/emboj.2009.398.

18. Poole, C.B., Jin, J., and McReynolds, L.A. (2003). Cloning and biochemical characterization of
blisterase, a subtilisin-like convertase from the filarial parasite, Onchocerca volvulus. J Biol
Chem *278*, 36183–36190. https://doi.org/10.1074/jbc.M302601200.

19. Sherman, B.T., Hao, M., Qiu, J., Jiao, X., Baseler, M.W., Lane, H.C., Imamichi, T., and Chang, W.
(2022). DAVID: a web server for functional enrichment analysis and functional annotation of
gene lists (2021 update). Nucleic Acids Res *50*, W216–W221.
https://doi.org/10.1093/nar/gkac194.

20. Visel, A., Thaller, C., and Eichele, G. (2004). GenePaint.org: an atlas of gene expression
patterns in the mouse embryo. Nucleic Acids Res *32*, D552-6.
https://doi.org/10.1093/nar/gkh029.

21. McCarter, G.C., Reichling, D.B., and Levine, J.D. (1999). Mechanical transduction by rat
dorsal root ganglion neurons in vitro. Neurosci Lett *273*, 179–182.
https://doi.org/10.1016/s0304-3940(99)00665-5.

22. Chiang, L.-Y., Poole, K., Oliveira, B.E., Duarte, N., Sierra, Y.A.B., Bruckner-Tuderman, L., Koch,
M., Hu, J., and Lewin, G.R. (2011). Laminin-332 coordinates mechanotransduction and
growth cone bifurcation in sensory neurons. Nat. Neurosci. *14*, 993–1000.
https://doi.org/10.1038/nn.2873.

23. Pavlakis, E., Chiotaki, R., and Chalepakis, G. (2011). The role of Fras1/Frem proteins in the
structure and function of basement membrane. Int J Biochem Cell Biol *43*, 487–495.
https://doi.org/10.1016/j.biocel.2010.12.016.

24. Nakamura, H., Cook, R.N., and Justice, M.J. (2013). Mouse Tenm4 is required for mesoderm
induction. BMC Dev Biol *13*, 9. https://doi.org/10.1186/1471-213X-13-9.

25. Chakrabarti, S., Pattison, L.A., Doleschall, B., Rickman, R.H., Blake, H., Callejo, G.,
Heppenstall, P.A., and Smith, E.S.J. (2020). Intraarticular Adeno-Associated Virus Serotype
AAV-PHP.S-Mediated Chemogenetic Targeting of Knee-Innervating Dorsal Root Ganglion
Neurons Alleviates Inflammatory Pain in Mice. Arthritis Rheumatol *72*, 1749–1758.
https://doi.org/10.1002/art.41314.

26. Chan, K.Y., Jang, M.J., Yoo, B.B., Greenbaum, A., Ravi, N., Wu, W.-L., Sánchez-Guardado, L.,
Lois, C., Mazmanian, S.K., Deverman, B.E., et al. (2017). Engineered AAVs for efficient
noninvasive gene delivery to the central and peripheral nervous systems. Nat Neurosci *20*,
1172–1179. https://doi.org/10.1038/nn.4593.

27. Mosca, T.J. (2015). On the Teneurin track: a new synaptic organization molecule emerges.
Front Cell Neurosci *9*, 204. https://doi.org/10.3389/fncel.2015.00204.

28. Mosca, T.J., Hong, W., Dani, V.S., Favaloro, V., and Luo, L. (2012). Trans-synaptic Teneurin
signalling in neuromuscular synapse organization and target choice. Nature *484*, 237–241.
https://doi.org/10.1038/nature10923.

29. Feng, K., Zhou, X.H., Oohashi, T., Morgelin, M., Lustig, A., Hirakawa, S., Ninomiya, Y., Engel,
J., Rauch, U., and Fassler, R. (2002). All four members of the Ten-m/Odz family of
transmembrane proteins form dimers. J Biol Chem *277*, 26128–26135.
https://doi.org/10.1074/jbc.M203722200.

30. Tucker, R.P., and Chiquet-Ehrismann, R. (2006). Teneurins: a conserved family of
transmembrane proteins involved in intercellular signaling during development. Dev Biol
290, 237–245. https://doi.org/10.1016/j.ydbio.2005.11.038.

31. Li, L., and Ginty, D.D. (2014). The structure and organization of lanceolate mechanosensory
complexes at mouse hair follicles. Elife *3*, e01901.

32. Heidenreich, M., Lechner, S.G., Vardanyan, V., Wetzel, C., Cremers, C.W., De Leenheer, E.M.,
Aránguez, G., Moreno-Pelayo, M.Á., Jentsch, T.J., and Lewin, G.R. (2012). KCNQ4 K(+)
channels tune mechanoreceptors for normal touch sensation in mouse and man. Nat.
Neurosci. *15*, 138–145. https://doi.org/10.1038/nn.2985.

33. Marmigere, F., and Ernfors, P. (2007). Specification and connectivity of neuronal subtypes in
the sensory lineage. Nat Rev Neurosci *8*, 114–127.

34. Li, L., Rutlin, M., Abraira, V.E., Cassidy, C., Kus, L., Gong, S., Jankowski, M.P., Luo, W., Heintz,
N., Koerber, H.R., et al. (2011). The Functional Organization of Cutaneous Low-Threshold
Mechanosensory Neurons. Cell *147*, 1615–1627. https://doi.org/10.1016/j.cell.2011.11.027.

35. Wetzel, C., Hu, J., Riethmacher, D., Benckendorff, A., Harder, L., Eilers, A., Moshourab, R.,
Kozlenkov, A., Labuz, D., Caspani, O., et al. (2007). A stomatin-domain protein essential for
touch sensation in the mouse. Nature *445*, 206–209. https://doi.org/10.1038/nature05394.

36. Murthy, S.E., Loud, M.C., Daou, I., Marshall, K.L., Schwaller, F., Kühnemund, J., Francisco,
A.G., Keenan, W.T., Dubin, A.E., Lewin, G.R., et al. (2018). The mechanosensitive ion channel
Piezo2 mediates sensitivity to mechanical pain in mice. Science Translational Medicine *10*,
eaat9897. https://doi.org/10.1126/scitranslmed.aat9897.

37. Walcher, J., Ojeda-Alonso, J., Haseleu, J., Oosthuizen, M.K., Rowe, A.H., Bennett, N.C., and
Lewin, G.R. (2018). Specialized mechanoreceptor systems in rodent glabrous skin. J Physiol
596, 4995–5016. https://doi.org/10.1113/JP276608.

38. Arcourt, A., Gorham, L., Dhandapani, R., Prato, V., Taberner, F.J., Wende, H., Gangadharan,
V., Birchmeier, C., Heppenstall, P.A., and Lechner, S.G. (2017). Touch Receptor-Derived
Sensory Information Alleviates Acute Pain Signaling and Fine-Tunes Nociceptive Reflex
Coordination. Neuron *93*, 179–193. https://doi.org/10.1016/j.neuron.2016.11.027.

39. Dietrich, S., Company, C., Song, K., Lowenstein, E.D., Riedel, L., Birchmeier, C., Gargiulo, G.,
and Zampieri, N. (2022). Molecular identity of proprioceptor subtypes innervating different
muscle groups in mice. Nat Commun *13*, 6867. https://doi.org/10.1038/s41467-022-34589-
8.

40. Santuz, A., and Zampieri, N. (2024). Making sense of proprioception. Trends in Genetics *40*,
20–23. https://doi.org/10.1016/j.tig.2023.10.006.

41. Hippenmeyer, S., Vrieseling, E., Sigrist, M., Portmann, T., Laengle, C., Ladle, D.R., and Arber,
S. (2005). A Developmental Switch in the Response of DRG Neurons to ETS Transcription
Factor Signaling. PLoS Biol *3*, e159. https://doi.org/10.1371/journal.pbio.0030159.

42. de Nooij, J.C., Doobar, S., and Jessell, T.M. (2013). Etv1 Inactivation Reveals Proprioceptor
Subclasses that Reflect the Level of NT3 Expression in Muscle Targets. Neuron *77*, 1055–
1068. https://doi.org/10.1016/j.neuron.2013.01.015.

43. Oliver, K.M., Florez-Paz, D.M., Badea, T.C., Mentis, G.Z., Menon, V., and de Nooij, J.C. (2021).
Molecular correlates of muscle spindle and Golgi tendon organ afferents. Nat Commun *12*,
1451. https://doi.org/10.1038/s41467-021-21880-3.

44. Del Toro, D., Carrasquero-Ordaz, M.A., Chu, A., Ruff, T., Shahin, M., Jackson, V.A., Chavent,
M., Berbeira-Santana, M., Seyit-Bremer, G., Brignani, S., et al. (2020). Structural Basis of
Teneurin-Latrophilin Interaction in Repulsive Guidance of Migrating Neurons. Cell *180*, 323-
339.e19. https://doi.org/10.1016/j.cell.2019.12.014.

45. Kumar, P., and Aradhyam, G.K. (2014). Easy and efficient protocol for purification of
recombinant peptides. Protein Expr Purif *95*, 129–135.
https://doi.org/10.1016/j.pep.2013.12.004.

46. Lossie, A.C., Nakamura, H., Thomas, S.E., and Justice, M.J. (2005). Mutation of l7Rn3 shows
that Odz4 is required for mouse gastrulation. Genetics *169*, 285–299.
https://doi.org/10.1534/genetics.104.034967.

47. Hu, J., and Lewin, G.R. (2006). Mechanosensitive currents in the neurites of cultured mouse
sensory neurones. J Physiol *577*, 815–828. https://doi.org/10.1113/jphysiol.2006.117648.

48. Kang, H., Han, A.-R., Zhang, A., Jeong, H., Koh, W., Lee, J.M., Lee, H., Jo, H.Y., Maria-Solano,
M.A., Bhalla, M., et al. (2024). GolpHCat (TMEM87A), a unique voltage-dependent cation

channel in Golgi apparatus, contributes to Golgi-pH maintenance and hippocampus-dependent memory. Nat Commun *15*, 5830. https://doi.org/10.1038/s41467-024-49297-8.

49. Patkunarajah, A., Stear, J.H., Moroni, M., Schroeter, L., Blaszkiewicz, J., Tearle, J.L., Cox, C.D., Furst, C., Sanchez-Carranza, O., Ocana Fernandez, M.D.A., et al. (2020). TMEM87a/Elkin1, a component of a novel mechanoelectrical transduction pathway, modulates melanoma adhesion and migration. Elife *9*. https://doi.org/10.7554/eLife.53308.

50. Götzke, H., Kilisch, M., Martínez-Carranza, M., Sograte-Idrissi, S., Rajavel, A., Schlichthaerle, T., Engels, N., Jungmann, R., Stenmark, P., Opazo, F., et al. (2019). The ALFA-tag is a highly versatile tool for nanobody-based bioscience applications. Nat Commun *10*, 4403. https://doi.org/10.1038/s41467-019-12301-7.

51. Ojeda-Alonso, J., Bégay, V., Garcia-Contreras, J.A., Campos-Pérez, A.F., Purfürst, B., and Lewin, G.R. (2022). Lack of evidence for participation of TMEM150C in sensory mechanotransduction. J Gen Physiol *154*, e202213098. https://doi.org/10.1085/jgp.202213098.

52. Moroni, M., Servin-Vences, M.R., Fleischer, R., Sánchez-Carranza, O., and Lewin, G.R. (2018). Voltage gating of mechanosensitive PIEZO channels. Nature Communications *9*, 1096. https://doi.org/10.1038/s41467-018-03502-7.

53. Emtage, L., Gu, G.Q., Hartwieg, E., and Chalfie, M. (2004). Extracellular proteins organize the mechanosensory channel complex in touch receptor neurons. Neuron *44*, 795–807. https://doi.org/10.1016/j.neuron.2004.11.010.

54. Du, H., Gu, G., William, C.M., and Chalfie, M. (1996). Extracellular proteins needed for C. elegans mechanosensation. Neuron *16*, 183–194.

55. Chalfie, M., and Sulston, J. (1981). Developmental genetics of the mechanosensory neurons of Caenorhabditis elegans. Dev. Biol. *82*, 358–370.

56. Hehlert, P., Effertz, T., Gu, R.-X., Nadrowski, B., Geurten, B.R.H., Beutner, D., de Groot, B.L., and Göpfert, M.C. (2025). NOMPC ion channel hinge forms a gating spring that initiates mechanosensation. Nat Neurosci *28*, 259–267. https://doi.org/10.1038/s41593-024-01849-3.

57. Walker, R.G., Willingham, A.T., and Zuker, C.S. (2000). A Drosophila mechanosensory transduction channel. Science (New York, N.Y.) *287*, 2229–2234.

58. O'Hagan, R., Chalfie, M., and Goodman, M.B. (2005). The MEC-4 DEG/ENaC channel of Caenorhabditis elegans touch receptor neurons transduces mechanical signals. Nat. Neurosci *8*, 43–50. https://doi.org/10.1038/nn1362.

59. Tucker, R.P., and Chiquet-Ehrismann, R. (2006). Teneurins: a conserved family of transmembrane proteins involved in intercellular signaling during development. Dev Biol *290*, 237–245. https://doi.org/10.1016/j.ydbio.2005.11.038.

60. Meijer, D.H., Frias, C.P., Beugelink, J.W., Deurloo, Y.N., and Janssen, B.J.C. (2022). Teneurin4
dimer structures reveal a calcium-stabilized compact conformation supporting homomeric
trans-interactions. EMBO J *41*, e107505. https://doi.org/10.15252/embj.2020107505.

61. Li, J., Bandekar, S.J., and Arac, D. (2023). The structure of fly Teneurin-m reveals an
asymmetric self-assembly that allows expansion into zippers. Embo Rep *24*, e56728.
https://doi.org/10.15252/embr.202256728.

62. Abdo, H., Calvo-Enrique, L., Lopez, J.M., Song, J., Zhang, M.-D., Usoskin, D., El Manira, A.,
Adameyko, I., Hjerling-Leffler, J., and Ernfors, P. (2019). Specialized cutaneous Schwann cells
initiate pain sensation. Science (New York, N.Y.) *365*, 695–699.
https://doi.org/10.1126/science.aax6452.

63. Nikolaev, Y.A., Feketa, V.V., Anderson, E.O., Schneider, E.R., Gracheva, E.O., and Bagriantsev,
S.N. (2020). Lamellar cells in Pacinian and Meissner corpuscles are touch sensors. Sci Adv *6*.
https://doi.org/10.1126/sciadv.abe6393.

64. Sánchez-Carranza, O., Bégay, V., Chakrabarti, S., Pampols-Perez, M., Wang, L., García-
Contreras, J.A., Hammes, A., and Lewin, G.R. (2025). Mechanically gated currents in mouse
sensory neurons lacking PIEZO2. Biophys J, S0006-3495(25)00734-9.
https://doi.org/10.1016/j.bpj.2025.10.042.

65. Hor, H., Francescatto, L., Bartesaghi, L., Ortega-Cubero, S., Kousi, M., Lorenzo-Betancor, O.,
Jimenez-Jimenez, F.J., Gironell, A., Clarimon, J., Drechsel, O., et al. (2015). Missense
mutations in TENM4, a regulator of axon guidance and central myelination, cause essential
tremor. Hum Mol Genet *24*, 5677–5686. https://doi.org/10.1093/hmg/ddv281.

66. Pillai, E.K., and Franze, K. (2024). Mechanics in the nervous system: From development to
disease. Neuron *112*, 342–361. https://doi.org/10.1016/j.neuron.2023.10.005.

820
821

ACKNOWLEDGEMENTS

We thank Maria Braunschweig, Franziska Bartelt, Kathleen Barda and Madlen Driesner for excellent technical assistance. Franziska Binder, Franziska Westphal, and Florian Keim for animal care taking. Séverine Kunz and Christina Schiel for electron microscopy and Protein Production & Characterization Technology Platform (MDC) team for excellent technical assistance. We thank members of Lewin lab for constructive discussions. We thank Stefan Lechner and Clément Verkest the mouse Piezo2-mScarlet construct. **Funding:** This research was funded by an ERC grant to GRL (Sensational Tethers 789128) and from the Deutsche Forschungsgemeinschaft SFB958. A.T-L. H was a recipient for MOST postdoctoral research abroad program (111-2917-I-564-011). S.C. was a recipient of an Alexander von Humboldt research fellowship. **Author contributions:** Conceptualization: M.A.K., A.T-L. H and G.R.L. Mouse model selection, design and validation: G.R.L, V.B. A.S. N.Z. and M.A.K; Patch clamp electrophysiology: S.C (poking) and J.A.G-C. (Pillar array); Antibody validation: MAK, L.D., R.G. A.S. and A.T-L.H; Tripartite GFP design and implementation; candidate screening and molecular biology: YASB, L.D and A.T-L. H. Skin nerve electrophysiology: M.A.K, R.G, W.Z, S.J.M. and G.R.L; Super resolution microscopy and biochemistry: A.T-L.H, M.A.K and S.H.; immunogold labelling-TEM: L.D., S.K. C.S., and E.N.; Behavioral assessments M.A.K. with R.G. Protein Production: A.S. Proteomics J.A.G-C, S.C, O,P and P.M. Writing: M.A.K. and G.R.L with input from all authors; Supervision and funding: G.R.L.

Competing interests: The authors declare that they have no competing interests.

SUPPLEMENTAL INFORMATION: