# Rapid speciation in the holopelagic ctenophore *Mnemiopsis* following glacial recession

Authors: Remi N. Ketchum[1,2]*, Edward G. Smith[3], Leandra M. Toledo[1], Whitney B. Leach[4], Natalia Padillo-Anthemides[1], Andreas D. Baxevanis[5], Adam M. Reitzel[6], Joseph F. Ryan[1]*

[1] Whitney Laboratory for Marine Bioscience, University of Florida, St. Augustine, Florida, USA
[2] Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA
[3] School of Life Sciences, The University of Warwick, Coventry, UK
[4] Department of Biological Sciences, Lehigh University, Bethlehem, Pennsylvania, USA
[5] Division of Intramural Research, National Human Genome Research Institute, National Institutes of Health, Bethesda, Maryland, USA
[6] Department of Biological Sciences, University of North Carolina at Charlotte, Charlotte, North Carolina, USA

## Abstract

Understanding how populations diverge is one of the oldest and most compelling questions in evolutionary biology. An in depth understanding of how this process operates in planktonic marine animals, where barriers for gene flow are seemingly absent, is critical to understanding the past, present, and future of ocean life. *Mnemiopsis* plays an important ecological role in its native habitat along the Atlantic coast of the Americas and is highly destructive in its non-native habitats in European waters. Although historical literature described three species of *Mnemiopsis*, the lack of stable morphological characters has led to the collapse of this group into a single species, *Mnemiopsis leidyi*. We generate high-quality reference genomes and use a whole-genome sequencing approach to reveal that there are two species of *Mnemiopsis* along its native range and show that historical divergence between the two species coincides with historical glacial melting. We define a hybridization zone between species and highlight that environmental sensing genes likely contribute to the invasive success of *Mnemiopsis*. Overall, this study provides insights into the fundamental question of how holopelagic species arise without clear barriers to gene flow and sheds light on the genomic mechanisms important for invasion success in a highly invasive species.

## Introduction

Speciation and species delineation is of fundamental importance to evolutionary biology, yet the vast majority of speciation literature is based on terrestrial species with small population sizes, clear physical barriers to gene flow, and external fertilization (1–3). Additionally, what we know about speciation in the ocean is largely dominated by active swimmers with well-developed dispersal abilities (i.e., fish) and coastal epibenthic fauna that are relatively sedentary in their adult stages (4). To better understand mechanisms driving speciation in the oceans, it is critical to expand this set of organisms to include animals that employ a wide range of life history strategies and inhabit other distinct marine ecosystems. Holopelagic invertebrate species spend their entire life in the water column and represent a common suite of life history traits in the ocean with high fecundities (5), large population sizes (6), and high dispersal potential.

*Mnemiopsis* (A. Agassiz, 1865) is a holopelagic genus within the phylum Ctenophora with a native
range extending continuously along more than 25,000 kilometers of the Atlantic coast of the
Americas from Kittery, Maine to the Valdes Peninsula in Argentina (7–9). These comb jellies are
considered one of the 100 most invasive species in the world (10) and their introductions into
European waters have triggered community trophic cascades and collapse of fisheries (11). The
extensive geographic distribution and invasive abilities of *Mnemiopsis* are driven in part by its
broad environmental tolerance (12), flexible planktivorous diet (13), extensive regeneration
abilities (14), high fertility levels (15), and a capacity for self-fertilization (16). Risk management
for invasive species requires genetic information to establish management units and accurate
taxonomic classification is a fundamental component of effective control measures.
52
Despite its rich 100-year history (17) as an evo devo model and the availability of extensive
genomics resources (18–21), our understanding of species boundaries in *Mnemiopsis* is subject to
debate (22–26). Within the World Register of Marine Species (WoRMS), *Mnemiopsis* is currently
divided into two accepted species, *Mnemiopsis leidyi* (A. Agassiz, 1865; type location
Massachusetts) and *Mnemiopsis gardeni* (L. Agassiz, 1860; type location South Carolina). A third
species, *Mnemiopsis mccradyi* (Mayer, 1900; type location South Carolina) is recognized as a
synonym for *Mnemiopsis leidyi* (27). However, there is much confusion as to the ranges and
differences in morphology between these two named species and most modern literature considers
all ctenophores along the Atlantic and Gulf coasts of the United States as a single species:
*Mnemiopsis leidyi* (28). Although there have been previous studies that suggest the possibility of
multiple *Mnemiopsis* species (22–26), the molecular data from these studies are based on a limited
set of markers and/or limited population sampling and have been insufficient to resolve this
question. However, the rapid expansion of next-generation sequencing (NGS) technologies has
revolutionized the study of genomic variation within and between species and presents us with a
unique opportunity to test for multiple species, revisit the naming system, and explore the
evolutionary changes that contribute to invasion success of holopelagic species.
69
We conducted whole-genome short-read sequencing of 118 individuals from 13 populations within
*Mnemiopsis's* native range along the United States Atlantic coast and generated two near
chromosome-level genome assemblies of one individual from Florida and one individual from
Massachusetts. We used a suite of genomic approaches to understand holopelagic population
dynamics, demographic histories, changes in genomic architecture, and evolutionary adaptations.
Taken together, these findings highlight barriers to gene flow in holopelagic species, mechanisms
underpinning invasion success, and the importance of high spatial resolution sampling in
identifying units for invasive species management.
78
**Results and Discussion**
**Whole-genome sequencing of *Mnemiopsis* populations**

We sequenced the genomes of 118 *Mnemiopsis* individuals from 13 populations along the US
Atlantic coast (Figure 1A, Supplementary Table 1). We generated a total of 730 Gb of Illumina
reads and an average genome depth of 18.89x. From this, we generated three datasets that were
appropriate for different types of analyses: the All-Sites dataset with 81,758,954 single nucleotide
polymorphisms (SNPs), the Population-specific dataset with 3,856,382 SNPs, and an LD-Filtered
dataset with 89,750 SNPs.

**Figure 1. A)** Sampling map for *Mnemiopsis* collections. Each site is colored according to the MDS plot
color scheme with the site abbreviations in parentheses. Stars indicate populations that were sampled for
PacBio genome sequencing and assembly. **B)** Phylogenetic tree based on nuclear SNPs, estimated by
ngsDist and FastMe using 1,302 SNPs and rooted with *Bolinopsis infundibulum.* The size of the circle on
the branch represents the bootstrap support. The northern and southern lineages both receive 100%
bootstrap support. The colors of the sample names indicate which lineage the sample belongs to (blue;
northern, red; southern). **C)** Mitochondrial genome tree generated using an alignment of the whole
mitogenome and IQ-TREE v2.3.2. The colors of the sample names indicate which lineage the sample
belongs to.

**Population genomic evidence of two *Mnemiopsis* species**
We initially performed phylogenetic analyses using nuclear and mitochondrial markers from our
*Mnemiopsis* samples collected along the Atlantic Coast. Our nuclear phylogenetic tree (Figure 1B)
revealed the presence of two distinct clades that we will herein refer to as the northern lineage
(Roanoke Island, NC to Woods Hole, MA) and the southern lineage (Pivers Island, NC to Panacea,
FL). For the most part, these two distinct clades are also observed in our mitochondrial
phylogenetic tree (Figure 1C). However, we identified six individuals from populations near the
geographic interface of these two clades that exhibit mitonuclear discordance. Specifically, we
observed one individual collected at Gloucester Point, Virginia (VAV10) and four individuals

collected in Roanoke Island, North Carolina (NCX3–5,8) with northern nuclear and southern
mitochondrial genotypes (Figure 1B–C). Likewise, we observed one individual collected from
Wrightsville Beach, North Carolina (NCW15) with southern nuclear and northern mitochondrial
genotypes. This level of mitochondrial discordance could be consistent with geographically
localized hybridization and adaptive introgression, secondary contact, or demographic disparities
such as sex-biased dispersal (29, 30).
112
To further investigate these patterns, we performed admixture, multidimensional scaling (MDS),
and population genetic analyses. We evaluated the presence of 2–7 populations (by adjusting the
K parameter) within our dataset using ngsADMIX (31). Setting K=2 separates individuals into
northern and southern lineages with the Roanoke Island, NC (NCX) and Gloucester Point, VA
(VAV) samples showing signs of admixture (Figure 2A). When K=3, the Roanoke Island samples
form their own subpopulation and at higher K values, more substructure within the two lineages
becomes evident. MDS plots (Figure 2B) support our ngsADMIX results with the first principal
component mirroring the patterns observed in K=2 and accounting for 39.4% of the variance in
the data. Our observations of distinct lineages can also be found at the population level, with
pairwise $F_{ST}$ comparisons revealing that the average $F_{ST}$ between the northern and southern lineage
was 0.278 when the putative hybrid population (Roanoke Island) was included in the northern
lineage and 0.315 when it was removed (Table 1). Due to these cumulative results, we also tested
for evidence of hybridization in our data using $f_3$ statistics (32). We compared Roanoke Island, NC
(NCX) to the northern and southern lineage and notably, this resulted in a negative $f3$ statistic and
an absolute Z score >= 3 (threshold used for significance) highlighting that Roanoke Island, NC is
a hybrid population (Supplementary Table 2). A recent study (26) detected evidence of
hybridization in two populations in Virginia: Gloucester Point and Wachapreague. Because we
also sampled from Gloucester Point, we tested for evidence of hybridization in this population by
comparing it to the northern and southern lineages (Supplementary Table 3) and did not detect
evidence for hybridization. This can likely be explained by the increased statistical power of using
4.8 million SNPs versus the 96 SNPs used for testing in Pujolar et al. 2023 or could be related to
temporal shifts in the hybrid zone.

**Figure 2. A)** Admixture plots for the 118 *Mnemiopsis* samples collected for K=2-4 where each bar
represents a single individual, and the colors represent each of the K components. **B)** MDS plot for PC1
and PC2 with the percent of total variance explained by each component shown above the plot in
parentheses. Samples are colored according to their sampling location with northern lineage samples
clustering on the left in blues and purples and southern lineage samples clustering on the right in reds. **C)**
Stairway plot results show different demographic histories of the two lineages with generation time of one
year and a mutation rate of 6.85 x $10^{-8}$ per base pair per year.

Taken together, our analyses clearly reveal two divergent genetic lineages in US Atlantic
*Mnemiopsis*. These results are congruent with evidence of genetic differentiation between
Massachusetts and Florida in a previous study (22). However, unlike previous studies, our high-
resolution spatial sampling has revealed a clear genetic breakpoint at Roanoke Island, NC. This
breakpoint is located where the Gulf Stream leaves the coast and moves offshore into the open
Atlantic Ocean. Contrary to expectations for a holopelagic species, we do not find evidence of

gene flow between lineages, except at the hybrid zone. Importantly, this hybrid zone is particularly
narrow, with admixture observed at Roanoke Island but absent from Piver's Island, NC located
~200 km away, highlighting that despite their holopelagic nature, clear barriers exist preventing
panmixia in *Mnemiopsis*.

**Table 1.** $F_{ST}$ values across all populations of *Mnemiopsis* collected based on the LD-Filtered SNP dataset.
These locations included: Woods Hole (MAW), Little Egg Harbor Township (NJR), Berlin (MDA),
Gloucester Point (VAV), Roanoke Island (NCX), Pivers Islands (NCB), Wrightsville Beach (NCW),
Charleston (SCS), Savannah (GAS), Savannah offshore (GAB), Flagler Beach (FLM), Fort Pierce (FLF),
and Panacea (FLP).

|     | MAW | NJR | MDA | VAV | NCX | NCB | NCW | SCC | GAS | GAB | FLM | FLF | FLP |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| MAW | X | X | X | X | X | X | X | X | X | X | X | X | X |
| NJR | 0.039 | X | X | X | X | X | X | X | X | X | X | X | X |
| MDA | 0.056 | -0.003 | X | X | X | X | X | X | X | X | X | X | X |
| VAV | 0.215 | 0.097 | 0.075 | X | X | X | X | X | X | X | X | X | X |
| NCX | 0.327 | 0.230 | 0.210 | 0.113 | X | X | X | X | X | X | X | X | X |
| NCB | 0.400 | 0.328 | 0.314 | 0.262 | 0.230 | X | X | X | X | X | X | X | X |
| NCW | 0.402 | 0.330 | 0.317 | 0.266 | 0.235 | 0.001 | X | X | X | X | X | X | X |
| SCC | 0.402 | 0.331 | 0.318 | 0.266 | 0.236 | 0.0003 | 0.0002 | X | X | X | X | X | X |
| GAS | 0.407 | 0.335 | 0.322 | 0.271 | 0.240 | 0.001 | 0.001 | 0.001 | X | X | X | X | X |
| GAB | 0.402 | 0.330 | 0.317 | 0.265 | 0.235 | -0.0003 | 0.001 | 0.00004 | −0.00002 | X | X | X | X |
| FLM | 0.391 | 0.318 | 0.305 | 0.253 | 0.223 | 0.029 | 0.028 | 0.027 | 0.032 | 0.027 | X | X | X |
| FLF | 0.404 | 0.331 | 0.318 | 0.265 | 0.235 | 0.061 | 0.058 | 0.087 | 0.064 | 0.057 | 0.03 | X | X |
| FLP | 0.422 | 0.349 | 0.336 | 0.283 | 0.254 | 0.089 | 0.086 | 0.087 | 0.092 | 0.086 | 0.059 | 0.039 | X |

## Demographic history of *Mnemiopsis*

Stairway plot (33) analysis shows that the demographic histories of *Mnemiopsis* could be traced
back from very recent history (<100 generations/years ago) to approximately 100,000 years ago
(Figure 2C). The demographic trajectories of the two lineages were remarkably different with the
southern lineage experiencing constant population expansion, especially at about 10,000 years ago.
The northern lineage has experienced two population bottlenecks, occurring roughly 10,000 years
ago and 2,000-3,000 years ago. More recently, the northern lineage is experiencing a population
expansion. Interestingly, the split in the effective population size curves between the two lineages
occurred ~10,000 years ago which coincides roughly with the retreat of the Laurentide Glacier (by
~8,000 years ago this glacier was confined mostly to Canada) and the flood that was caused when
Lake Agassiz burst and flooded several parts of North America, including the northern Atlantic
Ocean (34). This influx of cold, freshwater may have reduced the current in the ocean conveyor
belt and resulted in less warm seawater traveling poleward and subsequently causing temperatures
to drop in the Northern Hemisphere (34, 35). This population size split may have been a result of
a reduction in the Gulf Stream current during this time which could have resulted in *Mnemiopsis*
to move northward and begin a population expansion as temperatures slowly rose. Finally, the
population size decline around 2,000 – 3,000 years ago in the northern lineage may have been
caused by the Little Ice Age which occurred ca. 1300-1850 CE (36, 37).

Our results differ from the  pairwise sequentially Markovian coalescent (PSMC) results reported
in Jaspers et al (22). Although they also reported population expansions about 50,000 generations

ago and a population bottleneck about 1,000-2,000 generations ago, their northern lineage did not
show evidence of oscillations in population size. In contrast, ours showed a steady increase in
population size beginning at that time. Their southern lineage showed a steady increase in
population size from 500,000-50,000 generations ago and then a decrease in population size from
then onwards whereas ours has shown a steady increase from 100,000 generations ago to present.
The difference in findings can likely be explained by the increased geospatial resolution with our
study drawing conclusions from whole-genome level sampling of 5 northern populations and 8
southern populations versus one northern and southern population in the previous study. In
addition, while the PSMC method (38) performs well for inferring ancient histories, the Stairway
Plot method performs better for inference of more recent population histories and we see the most
discordance in our results when looking at more recent population histories (33, 39).
194

**Structural genomic evidence of speciation**
To investigate genomic differences between the two species, we produced two high-quality, near
chromosome-level genome assemblies of one Massachusetts *Mnemiopsis* sample and one Florida
sample using PacBio Hifi sequencing technology. When assembled, the Florida genome was 206
Mb and half of the genome was contained in contigs 3.6 Mb or larger (N50). There were 174
contigs total and the 100 largest contigs made up 98% of the assembly. The Massachusetts genome
was 215 Mb and half of the genome was contained in contigs 2 Mb or larger. There were 207
contigs total and the 100 largest contigs made up 89% of the assembly. We identified 27,570 and
26,046 protein-coding genes for Florida and Massachusetts, respectively and were able to
functionally annotate 27,476 genes for the Florida genome. Benchmarking Universal Single-Copy
Orthologs (40) analysis revealed that our assemblies exhibit a similar degree of completeness to
other ctenophore genomes (Table 2).
207

Chromosome-scale linkages of orthologous genes tend to be conserved even over very large time
scales (41–45). As such, chromosomal rearrangements are major evolutionary events that can
either cause speciation events and/or provide evidence of long periods of isolation between
lineages. We conducted macrosynteny analyses to check for evidence of chromosomal
rearrangements between the Florida and Massachusetts near-chromosomal genomes. In addition,
we compared both genomes to the recently published chromosome-level genome assembly of
*Bolinopsis microptera* (44), a lineage that is closely related to *Mnemiopsis* (46). We identified at
least one chromosomal rearrangement between the Florida and Massachusetts genomes (Figure
3A-C). The Massachusetts scaffold L006 maps to two Florida scaffolds, G002, which maps
completely to Chromosome 1 of *B. microptera*, and G004, which maps completely to chromosome
8 of *B. microptera*. The fact that this linkage group is present in both Florida *Mnemiopsis* and *B.*
*microptera*, suggests that the genomic rearrangement occurred in the Massachusetts lineage. There
are potentially other rearrangements involving the Massachusetts scaffolds L012 and L018, but a
chromosome-scale assembly would be needed to confirm or refute those possibilities.
222

223  **Table 2**. Comparison of genome assembly benchmarks between the *Mnemiopsis leidyi* published genome
224  (18) and the two new genome assemblies.

| | *Mnemiopsis leidyi* 2013 | *Mnemiopsis leidyi* 2024 | *Mnemiopsis gardeni* 2024 |
|---|---|---|---|
| Assembly size | 155,865,547 | 215,818,400 | 206,611,897 |
| No. of scaffolds | NA | 5,100 | NA |
| No. of contigs | NA | 207 | 174 |
| Max length (bases) | | 6,852,836 | 9,685,951 |
| N50 | 187,314 (scaff N50) | 2,034,471 (contig N50) | 3,675,794 (contig N50) |
| BUSCO Complete | 204 (80%) | 211 (83%) | 214 (84%) |
| BUSCO Complete + Partial | 233 (91%) | 231 (91 %) | 235 (91 %) |
| BUSCO missing | 22 (8.6%) | 24 (9%) | 20 (8%) |
| BUSCO duplicated | 1.47 | 1.9 | 3.74 |

**Figure 3.** Macrosynteny on a near-chromosome scale between *Mnemiopsis leidyi* (top), *M.* gardeni (middle), and *Bolinopsis microptera* (bottom). **(A)** Chord diagram of macrosynteny between *M. leidyi* (MA), *M. gardeni* (FL), and *B. macropteri*. The coordinates of 12,134 orthologs (based on reciprocal best BLAST hits) are plotted. **(B)** Plot of macrosynteny between *M. leidyi* and *M. gardeni*. **(C)** Oxford dot plot between *M. leidyi* (MA) and *M. gardeni* (FL). Each dot represents an ortholog.

**The combined evidence supports two species along the US Atlantic coast**

Both the population genomic and structural genomic evidence are consistent with there being one species of *Mnemiopsis* (*Mnemiopsis leidyi*) that inhabits the coastal waters stretching from the Outer Banks region of North Carolina to the northern limits of the native range (i.e., at least as north as Woods Hole, MA). Likewise, there is a second species that stretches from at least the Gulf of Mexico to Piver's Island, NC.

The position of this break corresponds with the location where the swift-flowing Gulf Stream diverges from the North Carolina coast as it collides with the Labrador current (47). This is a known dispersal barrier, and it is thought that these ocean flows are the source of range limits for many species (48). This seems to serve as the likely biogeographic barrier separating *Mnemiopsis leidyi* (north) from *Mnemiopsis gardeni* (south). During the last glacial maximum, the Gulf Stream followed its present-day course, but its flow was relaxed by at least 10% (49). As the Laurentide glacier receded and led to a habitat hospitable to *Mnemiopsis*, this reduction of flow may have been sufficient to allow a small founder population to overcome this barrier and these individuals may have initiated the present day *Mnemiopsis leidyi* lineage. A small founding population would be consistent with drastic genomic changes like chromosomal rearrangements being tolerated and spreading to fixation.

**Genomic signatures of selection in genes mediating environmental sensing**

Due to the substantial divergence between these two species, their geographical distribution that spans a significant environmental gradient, and their success as an invasive species, we investigated signatures of selection and local adaptation between the two lineages using the cross-population composite-likelihood ratio (XP-CLR) test (50). To highlight functional categories of these genes, we conducted gene ontology (GO) analysis (51) of the 2,093 genes that overlap with the genomic windows ranked in the top 10% of XP-CLR values. These genes were significantly enriched (P<=0.05) for GO molecular function categories (Table 3). Interestingly, many of these categories include categories associated with sensory processing, neural activity, and/or diet, for example G protein-coupled receptor signaling pathways (P=0.0004), extracellularly glutamate-gated ion channel activity (P=0.012), Golgi vesicle transport (P=0.0078), neurotransmitter transport (P=0.0321), and chitin metabolic processes (P=0.0043).

The membrane category, which was significantly enriched (P=0.0122), included several key sensory related and neural genes found in regions with high XP-CLR values such as six TRP channels, four glutamate receptors, the single *Mnemiopsis* voltage-gated calcium channel, one of the two *Mnemiopsis* voltage-gated sodium channels, nine voltage-gated potassium channels, two potassium 2-pore domain channels, two calcium channels, a calcium-activated potassium channel, a voltage-gated chloride channel, and two ASIC channels. The chitin metabolic processes category includes two chitinase genes, which are critical for digesting prey. The membrane category also includes syntaxin-6, which is in the genomic region with the highest fixation index of the XP-CLR

analysis (Figure 4). Within the XP-CLR outlier regions, we also observe *O*-GLcNAcase (Figure
4) which is a post-translational modifier of serine and threonine residues on a wide range of
proteins (52). This protein has been associated with stress responses ranging from heat shock to
oxidative stress. As both syntaxin-6 and *O*-GLcNAcase have been implicated in modulating
autophagic flux in response to nutrient status in *Caenorhabditis elegans* (53, 54)*,* we hypothesize
that overwintering in northern populations may be a driver of these selective sweeps as organisms
are likely nutrient deficient during this time. In addition, our outlier dataset also contains many
other genes that are relevant for nutrient sensing and metabolism such as mechanistic target of
rapamycin (mTOR), acetyl coenzyme A synthase, malate dehydrogenase, and cytochrome c
oxidase subunit 2. Combined, these results suggest that genes involved in sensing a changing
environment and adapting to changes in diet were under high positive selection in the northern
lineage as it invaded new environments. This finding provides important candidate genes for future
functional verification and sheds light on the importance of nutrient sensing and resilience to
nutrient deficiency which may make *Mnemiopsis* such a prolific invader.
286
287 **Table 3.** Gene Ontology (GO) term analysis reveals enrichment in relevant biological processes.

| GO TERM | P-Value | Description |
|---|---|---|
| GO:0005515 | 0 | protein binding |
| GO:0003723 | 0.0002 | RNA binding |
| GO:0004553 | 0.0002 | hydrolase activity |
| GO:0007186 | 0.0004 | G protein-coupled receptor signaling pathway |
| GO:0005215 | 0.001 | transporter activity |
| GO:0005524 | 0.0014 | ATP binding |
| GO:0006468 | 0.0014 | protein phosphorylation |
| GO:0005488 | 0.0014 | binding |
| GO:0004930 | 0.0016 | G protein-coupled receptor activity |
| GO:0004672 | 0.0016 | protein kinase activity |
| GO:0007018 | 0.0039 | microtubule-based movement |
| GO:0006030 | 0.0043 | chitin metabolic process |
| GO:0003755 | 0.0044 | peptidyl-prolyl cis-trans isomerase activity |
| GO:0000413 | 0.0044 | protein peptidyl-prolyl isomerization |
| GO:0007165 | 0.0046 | signal transduction |
| GO:0006366 | 0.0064 | transcription by RNA polymerase II |
| GO:0030117 | 0.0074 | membrane coat |
| GO:0048193 | 0.0078 | Golgi vesicle transport |
| GO:0003676 | 0.0082 | nucleic acid binding |
| GO:0045454 | 0.0083 | cell redox homeostasis |
| GO:0008081 | 0.0098 | phosphoric diester hydrolase activity |

| | | |
|---|---|---|
| GO:0005234 | 0.012 | extracellularly glutamate-gated ion channel activity |
| GO:0016020 | 0.0122 | membrane |
| GO:0055085 | 0.0128 | transmembrane transport |
| GO:0005868 | 0.0143 | cytoplasmic dynein complex |
| GO:0008061 | 0.0147 | chitin binding |
| GO:0009055 | 0.0156 | electron transfer activity |
| GO:0006452 | 0.0157 | translational frameshifting |
| GO:0045905 | 0.0157 | positive regulation of translational termination |
| GO:0045901 | 0.0157 | positive regulation of translational elongation |
| GO:0043022 | 0.0159 | ribosome binding |
| GO:0016462 | 0.0164 | pyrophosphatase activity |
| GO:0016021 | 0.0174 | |
| GO:0005216 | 0.0176 | monoatomic ion channel activity |
| GO:0005096 | 0.019 | GTPase activator activity |
| GO:0010181 | 0.0192 | FMN binding |
| GO:0005978 | 0.0196 | glycogen biosynthetic process |
| GO:0005622 | 0.0198 | intracellular anatomical structure |
| GO:0006811 | 0.0223 | monoatomic ion transport |
| GO:0015035 | 0.0231 | protein-disulfide reductase activity |
| GO:0004970 | 0.0235 | glutamate-gated receptor activity |
| GO:0008017 | 0.0243 | microtubule binding |
| GO:0005576 | 0.0256 | extracellular region |
| GO:0003777 | 0.026 | microtubule motor activity |
| GO:0004252 | 0.0268 | serine-type endopeptidase activity |
| GO:0004177 | 0.0282 | aminopeptidase activity |
| GO:0051056 | 0.0289 | regulation of small GTPase-mediated signal transduction |
| GO:0046933 | 0.0292 | proton-transporting ATP synthase activity |
| GO:0006820 | 0.0311 | monoatomic anion transport |
| GO:0006836 | 0.0321 | neurotransmitter transport |
| GO:0005328 | 0.0326 | neurotransmitter:sodium symporter activity |
| GO:0006810 | 0.033 | transport |
| GO:0016651 | 0.033 | oxidoreductase activity |
| GO:0019001 | 0.0395 | guanyl nucleotide binding |
| GO:0031683 | 0.0395 | G-protein beta/gamma-subunit complex binding |
| GO:0005452 | 0.0398 | solute:inorganic anion antiporter activity |
| GO:0004674 | 0.0412 | protein serine/threonine kinase activity |

| | | |
|---|---|---|
| GO:0006270 | 0.0423 | DNA replication initiation |
| GO:0008603 | 0.0426 | cAMP-dependent protein kinase regulator activity |
| GO:0005952 | 0.0426 | cAMP-dependent protein kinase complex |
| GO:0006383 | 0.0426 | transcription by RNA polymerase III |
| GO:0001932 | 0.0426 | regulation of protein phosphorylation |
| GO:0051287 | 0.0429 | NAD binding |
| GO:0005815 | 0.0442 | microtubule organizing center |
| GO:0046872 | 0.0444 | metal ion binding |
| GO:0005975 | 0.0449 | carbohydrate metabolic process |
| GO:0043130 | 0.0452 | ubiquitin binding |
| GO:0046856 | 0.0455 | phosphatidylinositol dephosphorylation |
| GO:0051920 | 0.0461 | peroxiredoxin activity |

**Figure 4**. The two genes: syntaxin-6 (top) and O-GlcNAcase (bottom) are under selection. The XP-CLR, $F_{ST}$, $D_{XY,}$ and nucleotide diversity values were calculated with nonoverlapping 10kb windows and step sizes. The location of the two genes is shown with a dotted gray line.

## **Conclusion**

Our knowledge of evolution is heavily biased across the tree of life and expansions in taxon sampling often leads to a much richer understanding of how evolutionary forces shape the genomic variation between populations and species. Our findings in the holopelagic ctenophore, *Mnemiopsis* show that despite having life history traits that should promote panmixia (i.e., high fecundity, large population sizes, and high dispersal potential), we instead find evidence of a strong genetic breakpoint that is likely due to barriers created by strong currents off the North Carolina

330  coast. We provide evidence that implicates the retreat of the Laurentide glacier and the subsequent
331  flood of Lake Agassiz in the lineage splitting of the northern and southern species. Finally, we
332  show evidence for selection on environmental sensing genes and genes implicated in surviving
333  stressful conditions, all of which likely contribute to the invasive success of *Mnemiopsis*.
334
**<u>Materials and Methods</u>**
**Sampling**
*Populations*. A total of 118 *Mnemiopsis* samples were collected from thirteen locations along the
Atlantic and Gulf coast of the United States. These locations included: Woods Hole (MA), Little
Egg Harbor Township (NJ), Berlin (MD), Gloucester Point (VA), Roanoke Island (NC), Pivers
Islands (NC), Wrightsville Beach (NC), Charleston (SC), Savannah (GA), Savannah offshore
(GA), Flagler Beach (FL), Fort Pierce (FL), and Panacea (FL). Samples were collected between
August 2021 and November 2022 from surface waters (<10m) using either a cteno-dipper or a tow
net. Samples immediately preserved in ethanol and placed in a -20°C freezer until further
processing.
345
346  *Genome*. One *Mnemiopsis* individual was collected from two locations for genome assembly:
347  Woods Hole (MA) and Flagler Beach (FL). These samples were collected in August 2023 from
348  surface waters (<2m) using a cteno-dipper. Samples were preserved in ethanol and placed in a -
349  20°C freezer until further processing.
350
**DNA Extraction and Sequencing**
*Populations*. High molecular weight DNA was extracted according to the CTAB/Chloroform
protocol detailed in Ehrlich et al. 2017 (55). All DNA extractions were visualized on a 1% agarose
gel for quality control. One hundred and nineteen individually indexed, paired-end libraries with
an approximate insert size of 550 bp were constructed using the NEBNext Ultra II FS DNA Library
Preparation kit. Whole-genome sequencing was performed at the University of Florida's ICBR
Bioinformatics Core Facility on a NovaSeq6000.
358
*Genome*. High molecular weight DNA was extracted using a modified SDS-based lysis described
in Moritz et al (55). DNA extractions were visualized on a 1% agarose gel for quality control and
quantified using a Qubit dsDNA High Sensitivity Assay Kit on a Qubit 2.0 Fluorometer. Genome
sequencing was performed on a Pacific Biosciences Sequel IIe platform, in the Circular Consensus
sequencing (CCS) mode. A total of 17.7 - 20.2 Gb of Hifi reads were generated using CCS for the
two individuals. PacBio library preparation and sequencing were performed at NIH Intramural
Sequence Center (NISC) in Rockville, MD.
366
**De novo Genome Assembly**
Hifiasm v0.19.8 (56) was used to assemble the two genomes. For the Flagler Beach, FL individual,
the `--hom-cov` flag was set to 90. To reduce the amount of duplication, PurgeHaplotigs v1.1.2

(57) was run with the 'contigcov' option for low coverage, low point between the peaks, and high
coverage cutoffs set to 10, 70, and 175 respectively. The percent cutoff for identifying a contig as
a haplotig was set to 50% and this assembly was subsequently run through PurgeHaplotigs 'clip'
option that identifies and trims overlapping contig ends. For the Woods Hole, MA individual, the
`--hom-cov` flag was set to 101, the 'contigcov' options were set to 25, 70, and 155, and the `-j`
and `-s` flags were both set to 55. The percent cutoff `-a` flag was set to 60% and the final assembly
was also clipped to remove overlapping contig ends.
377
**Gene Prediction and Annotation**
Genomic repetitive elements were identified with RepeatModeler v1.0.11 (58) to generate a
*Mnemiopsis*-specific repeat element library. Repetitive regions were soft masked prior to gene
prediction and annotation using RepeatMasker v4.0.8. We aligned previously published bulk
RNA-Seq reads from tissues (PRJEB28334) to the above genome assembly using Bowtie2 (59).
Aligned RNA-seq data and gene models for *M. leidyi* (ML2.2.aa) were input into Braker v2.1.5
(60). Assembled gene/protein models were functionally annotated using BLASTP v2.9.0+ with
three protein databases: UniProt Knowledgebase Swiss-Prot protein models v2021-03, RefSeq
invertebrate protein models, and the *M. leidyi* protein models (18). Finally, BUSCO v5 (40) was
used to measure the completeness of the genome assembly and protein using the something
database.
389
**MacrosyntR**
We analyzed conserved chromosome-scale synteny (macrosynteny) between these new Florida
and Massachusetts *Mnemiopsis* assemblies as well as the published *Bolinopsis microptera* genome
assembly. To identify conserved genes, we performed diamond BLAST (61) to compute the
reciprocal best BLAST hits between the 2013 ML2.2 *Mnemiopsis* gene models and the gene
models of each of the new *Mnemiopsis* assemblies and the published *B. microptera* gene models.
These reciprocal best BLAST hits were used to identify significantly conserved macrosyntenic
blocks in MacrosyntR (version 0.3.3; (62)).
398
**Data processing and filtering of 118 short-read genomes**
Raw sequence data was trimmed using Trimmomatic v0.39 (63) with a minimum phred score of
33. Once the data was filtered and adaptors were removed, the paired sequences were aligned to
the Flagler Beach, FL genome using BWA-mem v0.7.12 (38). Sequence alignments were filtered
using Samtools v1.9 (64) for a minimum quality score of 30, Sambamba v0.6.6 (65) was used to
filter duplicate reads with default parameters, and bamutil v1.0.15 (66) was used to clip
overlapping read pairs.
406
We generated three datasets for our different analyses. For each dataset, we used the Flagler Beach,
FL genome as our reference and ancestral genome. *LD-Filtered Dataset.* The first was a highly
filtered dataset that was produced by using ANGSD v0.921 (67) to call SNPs and estimate

genotype likelihoods using the samtools model (`-GL 1`). We filtered the SNP data for 1) positions
that were present in at least 94 out of 118 samples, 2) a maximum coverage of 2360x per site across
all individuals to avoid calling sites in highly repetitive regions, 3) a minimum of 354x coverage
per site across all individuals, 4) a minimum base quality score <20, 5) a p-value cut off of 10-
6 for calling polymorphic loci and 5) retained only SNPs with a minor global allele frequency
>5%. The ANGSD output was then subsampled for one SNP out of every 50 and pairwise linkage
disequilibrium (LD) was estimated with ngsLD v1.1.1 (68) on this subsampled dataset assuming
a maximum distance of 10Kb between SNPs. Linked sites were pruned with the script
'prune_graph.pl' with a maximum distance of 10 Kb between SNPs and a minimum weight of 0.5
(69). The final LD-filtered dataset was used to generate phylogenetic trees, ngsadmix and MDS
plots, and calculate $F_{ST}$ measures. *Population-Specific Filtered Dataset.* The second dataset was
filtered using the same ANGSD commands as the first dataset (excluding the LD filtering steps)
but then underwent a second round of population- specific filtering. Specifically, sampling
locations from Woods Hole, MA to Roanoke Island, NC were included in the 'Northern lineage'
group and sampling locations from Piver's, NC to Panacea, FL were included in the 'Southern
lineage' group. These two groups were filtered separately and for the northern lineage, we filtered
the dataset for 1) positions that were present in at least 36 out of 45 individuals, 2) a maximum
coverage of 900x coverage per site across all individuals, 3) a minimum of 36x coverage per site
across all individuals. For the southern lineage we filtered the dataset for 1) positions that were
present in at least 58 out of 73 individuals, 2) a maximum coverage of 1,460x coverage per site
across all individuals, 3) a minimum of 175x coverage per site across all individuals. We restricted
the secondary lineage-specific filtering step to the subset of positions that were retained in the first
round of filtering using the -sites option in ANGSD. Filtering in this manner guarantees that sites
with an allele that is fixed in one lineage will be retained. The final population-specific filtered
dataset was used to generate stairway plots, sliding window $F_{st}$ values, XP-CLR values, and to
calculate $f_3$ statistics. *All-Sites Filtered Dataset.* The third dataset was filtered with the same
method as the population-specific dataset, but we kept all positions including non-polymorphic
sites. This dataset was used to calculate nucleotide diversity and $D_{XY}$ values.
438
**Population Structure Analysis**
We used the ngsDist option in ngsTools (70) on the LD-Filtered dataset to compute genetic
distances between individuals and perform multiple dimensional scaling (MDS). We then looked
for signatures of admixture using ngsAdmix (31) with runs ranging from K =2 to K = 7. Weighted
pairwise $F_{STS}$ were estimated between the individuals from the North (Woods Hole, MA to
Roanoke Island, NC) and the individuals from the South (Pivers Island, NC to Panacea, FL) using
ANGSD with the `-doSaf 1` flag. We then used realSFS to estimate the 2D site frequency
spectrum for each population and calculated the average pairwise weighted $F_{ST}$ using 'realSFS
fst.' Weighted pairwise $F_{ST}$ estimates were also calculated between each sampling location using
ANGSD with the same parameters.
449

NgsDist was used to compute genetic distances from genotype probabilities with *Bolinopsis*
*infundibulum* as the outgroup (SRX14647678). Raw *Bolinopsis* sequences were filtered according
to the same protocol as was applied to *Mnemiopsis* and then we ran ANGSD with *Bolinopsis*
included, a minor allele frequency filter of 0.05 and a SNP pval filter of 1e-6. This set of filters
resulted in 1,302 SNPs that were used to generate the phylogenetic tree based on the nuclear
genome. NgsDist was run with 100 bootstrap replicates and a boot block size of 20. Finally,
FastME v2.1.6.1 (71) was used to visualize the pairwise genetic distances and RAxML-NG v0.9.0
(72) was used to place bootstrap supports onto the tree. Final figures were generated using the
interactive tree of life tool v6.9.1 (73).
459

Novoplasty v4.3.1 (74) was used to extract the mitochondrial genome from each trimmed sample
using the published *Mnemiopsis leidyi* COI sequence (KF435121.1(75)). When this was
unsuccessful, we instead used the published *Bolinopsis microptera* (SRX10473759 (46)) whole-
genome sequences and ran Novoplasty with the published *B. microptera* COI seed (MW735734.1
(76)) to extract the complete mitochondrial genome. The complete *B. microptera* mitochondrial
genome was then used as a seed for the remaining samples that initially failed Novoplasty. Using
this approach, we were able to successfully assemble 112 out of 118 *Mnemiopsis* mitochondrial
genomes. Next, we used MARS (77) to ensure that each mitochondrial genome began at the same
location using the branch and bound method and the block length set to 100. We used MAFFT
v7.480 (78) to generate an alignment and IQ-TREE v2.3.2 (79) to generate a phylogenetic tree.
470

**Demographic Analysis**
StairwayPlot v2 (33) was used to reconstruct effective population sizes based on the folded SFSs
generated from the All-Sites Filtered dataset. We used a mutation rate of $6.85 \times 10^{-8}$ per base pair
per year (22) and a generation time of 1 year (80). A one-year generation time was chosen based
on Charlesworth et al 1994 (81) who asserted that generation time can be estimated as the mean
age at reproduction for a female. The hermaphroditic *Mnemiopsis* can reproduce within the first
month of its lifespan (82) its lifespan is approximately 2 years (80) and there is no information
indicating that they ever lose the ability to reproduce. As such a mean age of reproduction was set
to 1 year.
480

*$F_3$ Statistics*
We used $f_3$ statistics, included as part of the ADMIXTOOLS package v.7.0.2 (32), as a direct test
for detecting hybridization. The $f_3$ statistic estimates whether allele frequency differences between
each population combination deviates more than expected because of incomplete lineage sorting,
which implicates admixture. If the $f_3$ statistic is negative, this suggests that admixture has likely
occurred. For this analysis, all populations which belonged to the northern mitochondrial lineage
were combined into one population and all the populations which belonged to the southern
mitochondrial lineage were combined into the second population. These two lineages were tested
against Roanoke Island, NC and Gloucester Point, VA.

**Genomic Signatures of Selection**

To detect regions of the genome under natural selection between the two lineages, we used XP-CLR (50) which uses the AFS to calculate local deviations between populations. We used a sliding window and step size of 10kb with the maxSNPs flag set to 200. The genomic regions in the top 10% of the XP-CLR values were considered to represent selective sweeps and were associated with overlapping genes using Bedtools v.2.29.0 (83). We conducted a Monte Carlo analysis to identify overrepresented gene ontology terms as was conducted in Babonis et al. 2018 (84).

To further investigate genomic patterns of regions under selection, we generated sliding window scans for $F_{ST}$, $D_{XY}$, and nucleotide diversity values. To generate the sliding window scans, we used the population-specific filtered dataset with the Roanoke Island, NC population removed. We then used realSFS $F_{ST}$ index in ANGSD to compute per-site $F_{ST}$ indexes and realSFS $F_{ST}$ stats2 to perform a window-sliding analysis with the window and step size set to 10 kb to ensure nonoverlapping windows. The All-Sites Filtered dataset, with the Roanoke Island population removed, was used to generate nucleotide diversity and $D_{XY}$ values. For nucleotide diversity calculations, we used realSFS saf2theta in ANGSD to generate folded SFS's and then generated these values along a sliding window with thetaStat do_stat and the window and step size set to 10 kb. For $D_{XY}$ calculations, we used the CalcDxy.R code by Joshua Penalba then averaged the resulting values over the same 10kb windows as the $F_{ST}$ and nucleotide calculations.

**Acknowledgements**

This work was funded with a Postdoctoral Research Fellowship in Biology (PRFB) from the National Science Foundation (award number: 2109712). We would like to thank Dr. Ann Tarrant, Roland Hagan, Robert Aguilar, Dr. Deborah Steinberg, Tor Mowatt-Larssen, Eli at Outer Banks Kayaking Adventures, Dr. Joshua Osterberg, Dr. Jacob Warner, Dr. Daniel Sasson, Dr. Marc Frischer, Dr. Adam Greer, Dr. Holly Sweat, and Dr. Yuriy Bobkov for help collecting *Mnemiopsis* samples along the US Atlantic coast. We would like to acknowledge the ICBR for their excellent sequencing and service (RRID:SCR_019152). We would like to thank Dr. James Thomas at the NIH Intramural Sequencing Center (NISC) for his help with the PacBio sequencing. We would also like to thank Dr. Gustav Paulay and Dr. Claudia Mills for their help navigating the complex taxonomy of *Mnemiopsis*. Finally, we want to thank Dr. Mark Ravinet for his help brainstorming ideas pertaining to the hybridization of these two lineages.

**Citations**

1.  L. Q. Choo, *et al.*, Genome-wide phylogeography reveals cryptic speciation in the circumglobal planktonic calcifier Limacina bulimoides. *Molecular Ecology* **32**, 3200–3219 (2023).

2.    C. Breusing, *et al.*, Allopatric and Sympatric Drivers of Speciation in Alviniconcha
Hydrothermal Vent Snails. *Mol Biol Evol* **37**, 3469–3484 (2020).

3.    D. Schluter, D. J. Howard, S. H. Berlocher, *Endless forms: species and speciation* (Oxford
Univ. Press, 1998).

4.    M. P. Miglietta, A. Faucci, F. Santini, Speciation in the Sea: Overview of the Symposium
and Discussion of Future Directions. *Integrative and Comparative Biology* **51**, 449–455
(2011).

5.    C. Jaspers, J. H. Costello, S. P. Colin, Carbon content of Mnemiopsis leidyi eggs and
specific egg production rates in northern Europe. *Journal of Plankton Research* **37**, 11–15
(2015).

6.    T. Shiganova, Changes in appendicularian Oikopleura dioica abundance caused by invasion
of alien ctenophores in the Black Sea. *Journal of the Marine Biological Association of the*
*United Kingdom* **85**, 477–494 (2005).

7.    H. W. Mianzan, P. Martos, J. H. Costello, R. A. Guerrero, Avoidance of hydrodynamically
mixed environments by Mnemiopsis leidyi (Ctenophora: Lobata) in open-sea populations
from Patagonia, Argentina. *Jellyfish Blooms: New Problems and Solutions* 113–124 (2010).

8.    G. R. Harbison, S. P. Volovik, THE CTENOPHORE, MNEMIOPSIS LEIDYI, IN THE
BLACK SEA: A HOLOPLANKTONIC ORGANISM TRANSPORTED in *Nonindigenous*
*Estuarine & Marine Organisms (NEMO): Proceedings of the Conference and Workshop,*
*Seattle, Washington, April 1993*, (US Department of Commerce, National Oceanic and
Atmospheric Administration, 1994), p. 25.

9.    H. W. Mianzan, P. Martos, J. H. Costello, R. A. Guerrero, Avoidance of hydrodynamically
mixed environments by Mnemiopsis leidyi (Ctenophora: Lobata) in open-sea populations
from Patagonia, Argentina. *Jellyfish Blooms: New Problems and Solutions* 113–124 (2010).

10.   S. Lowe, M. Browne, S. Boudjelas, M. De Poorter, *100 of the world's worst invasive alien*
*species: a selection from the global invasive species database* (Invasive Species Specialist
Group Auckland, 2000).

11.   L. Doyle, Mnemiopsis leidyi. *AN INTRODUCTION TO MARINE INVASIVE SPECIES* **52**
(1984).

12.   T. A. Shiganova, Adaptive strategies of Mnemiopsis leidyi A. Agassiz 1865 in different
environments of the Eurasian seas. *Marine Pollution Bulletin* **161**, 111737 (2020).

13. C. Jaspers, L. F. Møller, T. Kiørboe, Reproduction rates under variable food conditions and
starvation in Mnemiopsis leidyi: significance for the invasion success of a ctenophore.
*Journal of Plankton Research* **37**, 1011–1018 (2015).

14. A. Edgar, D. G. Mitchell, M. Q. Martindale, Whole-body regeneration in the lobate
ctenophore Mnemiopsis leidyi. *Genes* **12**, 867 (2021).

15. D. A. Sasson, J. F. Ryan, The sex lives of ctenophores: the influence of light, body size, and
self-fertilization on the reproductive output of the sea walnut, Mnemiopsis leidyi. *PeerJ* **4**,
e1846 (2016).

16. D. A. Sasson, A. A. Jacquez, J. F. Ryan, The ctenophore Mnemiopsis leidyi regulates egg
production via conspecific communication. *BMC Ecology* **18**, 1–10 (2018).

17. C. M. Child, The gradient in susceptibility to cyanides in the meridional conducting path of
the ctenophore Mnemiopsis. *American Journal of Physiology-Legacy Content* **43**, 87–112
(1917).

18. J. F. Ryan, *et al.*, The genome of the ctenophore Mnemiopsis leidyi and its implications for
cell type evolution. *Science* **342**, 1242592 (2013).

19. R. T. Moreland, *et al.*, A customized Web portal for the genome of the ctenophore
Mnemiopsis leidyi. *BMC Genomics* **15**, 1–13 (2014).

20. R. T. Moreland, A.-D. Nguyen, J. F. Ryan, A. D. Baxevanis, The Mnemiopsis Genome
Project Portal: integrating new gene expression resources and improving data visualization.
*Database* **2020**, baaa029 (2020).

21. R. T. Moreland, S. Zhang, S. N. Barreira, J. F. Ryan, A. D. Baxevanis, An AI-generated
proteome-scale dataset of predicted protein structures for the ctenophore Mnemiopsis
leidyi. *Proteomics* 2300397 (2024).

22. C. Jaspers, *et al.*, Invasion genomics uncover contrasting scenarios of genetic diversity in a
widespread marine invader. *Proceedings of the National Academy of Sciences* **118**,
e2116211118 (2021).

23. K. M. Bayha, *et al.*, Worldwide phylogeography of the invasive ctenophore Mnemiopsis
leidyi (Ctenophora) based on nuclear and mitochondrial DNA data. *Biological Invasions*
**17**, 827–850 (2015).

24. C. Verwimp, *et al.*, Population genomic structure of the gelatinous zooplankton species
Mnemiopsis leidyi in its nonindigenous range in the North Sea. *Ecology and Evolution* **10**,
11–25 (2020).

25.   T. B. Reusch, S. Bolte, M. Sparwel, A. G. Moss, J. Javidpour, Microsatellites reveal origin
and genetic diversity of Eurasian invasions by one of the world's most notorious marine
invader, Mnemiopsis leidyi (Ctenophora). *Molecular ecology* **19**, 2690–2699 (2010).

26.   J. M. Pujolar, D. Breitburg, J. Lee, M. B. Decker, C. Jaspers, Hybridization and adaptive
introgression in a marine invasive species in native habitats. *Iscience* **26** (2023).

27.   S. Ahyong, *et al.*, World Register of Marine Species (WoRMS). WoRMS Editorial Board.
Deposited 14 September 2024.

28.   L. N. Seravin, The systematic revision of the genus Mnemiopsis (Ctenophora, Lobata).
*Zoologičeskij žurnal* **73**, 9–18 (1994).

29.   M. Domènech, L. C. Crespo, A. Enguídanos, M. A. Arnedo, Mitochondrial discordance in
closely related Theridion spiders (Araneae, Theridiidae), with description of a new species
of the T. melanurum group. *Zoosystematics and Evolution* **96**, 159–173 (2020).

30.   D. W. Foltz, *et al.*, Global population divergence of the sea star Hippasteria phrygiana
corresponds to the onset of the last glacial period of the Pleistocene. *Marine Biology* **160**,
1285–1296 (2013).

31.   L. Skotte, T. S. Korneliussen, A. Albrechtsen, Estimating individual admixture proportions
from next generation sequencing data. *Genetics* **195**, 693–702 (2013).

32.   N. Patterson, *et al.*, Ancient Admixture in Human History. *Genetics* **192**, 1065–1093
(2012).

33.   X. Liu, Y.-X. Fu, Stairway Plot 2: demographic history inference with folded SNP
frequency spectra. *Genome Biology* **21**, 280 (2020).

34.   H. (Kikki) F. Kleiven, *et al.*, Reduced North Atlantic deep water coeval with the glacial
Lake Agassiz freshwater outburst. *science* **319**, 60–64 (2008).

35.   J. T. Teller, D. W. Leverington, J. D. Mann, Freshwater outbursts to the oceans from glacial
Lake Agassiz and their role in climate change during the last deglaciation. *Quaternary*
*Science Reviews* **21**, 879–887 (2002).

36.   L. Martínez-García, *et al.*, Historical demographic processes dominate genetic variation in
ancient Atlantic cod mitogenomes. *Frontiers in Ecology and Evolution* **9**, 671281 (2021).

37.   L. M. Atmore, *et al.*, Population dynamics of Baltic herring since the Viking Age revealed
by ancient DNA and genomics. *Proceedings of the National Academy of Sciences* **119**,
e2208703119 (2022).

38.　H. Li, R. Durbin, Inference of human population history from individual whole-genome
sequences. *Nature* **475**, 493–496 (2011).

39.　X. Liu, Y.-X. Fu, Exploring population size changes using SNP frequency spectra. *Nature*
*genetics* **47**, 555–559 (2015).

40.　M. Manni, M. R. Berkeley, M. Seppey, E. M. Zdobnov, BUSCO: assessing genomic data
quality and beyond. *Current Protocols* **1**, e323 (2021).

41.　N. H. Putnam, *et al.*, Sea anemone genome reveals ancestral eumetazoan gene repertoire
and genomic organization. *science* **317**, 86–94 (2007).

42.　N. H. Putnam, *et al.*, The amphioxus genome and the evolution of the chordate karyotype.
*Nature* **453**, 1064–1071 (2008).

43.　O. Simakov, *et al.*, Insights into bilaterian evolution from three spiralian genomes. *Nature*
**493**, 526–531 (2013).

44.　D. T. Schultz, *et al.*, Ancient gene linkages support ctenophores as sister to other animals.
*Nature* **618**, 110–117 (2023).

45.　T. D. Lewin, I. J.-Y. Liao, Y.-J. Luo, Annelid comparative genomics and the evolution of
massive lineage-specific genome rearrangement in bilaterians. *Molecular Biology and*
*Evolution* **41**, msae172 (2024).

46.　S. B. Johnson, *et al.*, Speciation of pelagic zooplankton: Invisible boundaries can drive
isolation of oceanic ctenophores. *Frontiers in Genetics* **13**, 970314 (2022).

47.　L. D. Talley, M. S. McCartney, Distribution and circulation of Labrador Sea water. *Journal*
*of Physical Oceanography* **12**, 1189–1205 (1982).

48.　B. Gaylord, S. D. Gaines, Temperature or transport? Range limits in marine species
mediated solely by flow. *The American Naturalist* **155**, 769–789 (2000).

49.　K. Matsumoto, J. Lynch-Stieglitz, Persistence of Gulf Stream separation during the Last
Glacial Period: Implications for current separation theories. *Journal of Geophysical*
*Research: Oceans* **108** (2003).

50.　H. Chen, N. Patterson, D. Reich, Population differentiation as a test for selective sweeps.
*Genome research* **20**, 393–402 (2010).

51.　M. Ashburner, *et al.*, Gene ontology: tool for the unification of biology. *Nature Genetics*
**25**, 25–29 (2000).

52.  C.-R. Torres, G. W. Hart, Topography and polypeptide distribution of terminal N-
acetylglucosamine residues on the surfaces of intact lymphocytes. Evidence for O-linked
GlcNAc. *Journal of Biological Chemistry* **259**, 3308–3317 (1984).

53.  B. Guo, *et al.*, O-GlcNAc-modification of SNAP-29 regulates autophagosome maturation.
*Nature Cell Biology* **16**, 1215–1226 (2014).

54.  N. Mizushima, Sugar modification inhibits autophagosome–lysosome fusion. *Nature Cell
Biology* **16**, 1132–1133 (2014).

55.  M. Ehrlich, "Invasion Genomics: population structure and diversity patterns in the invasive
ctenophore Mnemiopsis leidyi based on whole-genome re-sequencing," Christian-
Albrechts-Universität Kiel. (2017).

56.  H. Cheng, G. T. Concepcion, X. Feng, H. Zhang, H. Li, Haplotype-resolved de novo
assembly using phased assembly graphs with hifiasm. *Nature methods* **18**, 170–175 (2021).

57.  M. J. Roach, S. A. Schmidt, A. R. Borneman, Purge Haplotigs: allelic contig reassignment
for third-gen diploid genome assemblies. *BMC Bioinformatics* **19**, 1–10 (2018).

58.  J. M. Flynn, *et al.*, RepeatModeler2 for automated genomic discovery of transposable
element families. *Proceedings of the National Academy of Sciences* **117**, 9451–9457
(2020).

59.  B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nature methods*
**9**, 357–359 (2012).

60.  T. Brůna, K. J. Hoff, A. Lomsadze, M. Stanke, M. Borodovsky, BRAKER2: automatic
eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a
protein database. *NAR genomics and bioinformatics* **3**, lqaa108 (2021).

61.  B. Buchfink, K. Reuter, H.-G. Drost, Sensitive protein alignments at tree-of-life scale using
DIAMOND. *Nature methods* **18**, 366–368 (2021).

62.  S. El Hilali, R. R. Copley, macrosyntR: Drawing automatically ordered Oxford Grids from
standard genomic files in R. *bioRxiv* 2023.01. 26.525673 (2023).

63.  A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114–2120 (2014).

64.  H. Li, *et al.*, The sequence alignment/map format and SAMtools. *bioinformatics* **25**, 2078–
2079 (2009).

65.   A. Tarasov, A. J. Vilella, E. Cuppen, I. J. Nijman, P. Prins, Sambamba: fast processing of
NGS alignment formats. *Bioinformatics* **31**, 2032–2034 (2015).

66.   G. Jun, M. K. Wing, G. R. Abecasis, H. M. Kang, An efficient and scalable analysis
framework for variant extraction and refinement from population-scale DNA sequence
data. *Genome research* **25**, 918–925 (2015).

67.   T. S. Korneliussen, A. Albrechtsen, R. Nielsen, ANGSD: analysis of next generation
sequencing data. *BMC bioinformatics* **15**, 1–13 (2014).

68.   E. A. Fox, A. E. Wright, M. Fumagalli, F. G. Vieira, ngsLD: evaluating linkage
disequilibrium using genotype likelihoods. *Bioinformatics* **35**, 3855–3856 (2019).

69.   R. N. Lou, A. Jacobs, A. P. Wilder, N. O. Therkildsen, A beginner's guide to low-coverage
whole genome sequencing for population genomics. *Molecular Ecology* **30**, 5966–5993
(2021).

70.   M. Fumagalli, F. G. Vieira, T. Linderoth, R. Nielsen, ngsTools: methods for population
genetics analyses from next-generation sequencing data. *Bioinformatics* **30**, 1486–1487
(2014).

71.   V. Lefort, R. Desper, O. Gascuel, FastME 2.0: a comprehensive, accurate, and fast
distance-based phylogeny inference program. *Molecular biology and evolution* **32**, 2798–
2800 (2015).

72.   A. M. Kozlov, D. Darriba, T. Flouri, B. Morel, A. Stamatakis, RAxML-NG: a fast, scalable
and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35**,
4453–4455 (2019).

73.   I. Letunic, P. Bork, Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic
tree display and annotation tool. *Nucleic Acids Research* gkae268 (2024).

74.   N. Dierckxsens, P. Mardulyn, G. Smits, NOVOPlasty: de novo assembly of organelle
genomes from whole genome data. *Nucleic acids research* **45**, e18–e18 (2017).

75.   S. Ghabooli, *et al.*, Invasion pathway of the ctenophore Mnemiopsis leidyi in the
Mediterranean Sea. *PloS one* **8**, e81067 (2013).

76.   L. M. Christianson, S. B. Johnson, D. T. Schultz, S. H. Haddock, Hidden diversity of
Ctenophora revealed by new mitochondrial COI primers and sequences. *Molecular Ecology*
*Resources* **22**, 283–294 (2022).

77.   L. A. Ayad, S. P. Pissis, MARS: improving multiple circular sequence alignment using
refined sequences. *BMC genomics* **18**, 1–10 (2017).

78.   K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular biology and evolution* **30**, 772–780
(2013).

79.   B. Q. Minh, *et al.*, IQ-TREE 2: new models and efficient methods for phylogenetic
inference in the genomic era. *Molecular biology and evolution* **37**, 1530–1534 (2020).

80.   J. S. Presnell, M. Bubel, T. Knowles, W. Patry, W. E. Browne, Multigenerational laboratory
culture of pelagic ctenophores and CRISPR–Cas9 genome editing in the lobate Mnemiopsis
leidyi. *Nature Protocols* **17**, 1868–1900 (2022).

81.   B. Charlesworth, *Evolution in age-structured populations* (Cambridge University Press
Cambridge, 1994).

82.   A. Edgar, J. M. Ponciano, M. Q. Martindale, Ctenophores are direct developers that
reproduce continuously beginning very early after hatching. *Proceedings of the National*
*Academy of Sciences* **119**, e2122052119 (2022).

83.   A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842 (2010).

84.   L. S. Babonis, *et al.*, Integrating embryonic development and evolutionary history to
characterize tentacle-specific cell types in a ctenophore. *Molecular Biology and Evolution*
**35**, 2940–2956 (2018).