Phylogenetic Measures of the Core Microbiome

Sharon Anne Bewick[1]*, Benjamin Thomas Camper[1]

[1]Department of Biological Sciences, Clemson University, Clemson, South Carolina 29631, USA

*sbewick@clemson.edu

Abstract

Background: A fundamental concept in microbial ecology is the 'core microbiome.' Typically, core microbiomes are defined as the microbial taxa, genes, or functions shared by a threshold number of microbiome samples from a particular type of habitat (e.g., a particular type of host or a particular type of environment/ecosystem). In defining the core microbiome, the goal is to capture the portion of the microbial community that is conserved across samples from the focal habitat. Recently, there has been growing interest in developing methods to better characterize core microbiomes. As a result, numerous occurrence- and abundance-based measures have been defined. However, none have included phylogenetically aware metrics for analyzing core microbiomes.

Results: In this paper, we develop the concept of the 'core community phylogeny' – a phylogeny where branches are selected based on their presence in multiple samples from a single type of habitat. We then use the core community phylogeny to define phylogenetic metrics for the diversity of core microbiomes from a single type of habitat, for turnover between core microbiomes from two different types of habitat, and for shared diversity across core microbiomes from two or more different types of habitat. As compared to non-phylogenetic metrics, our phylogenetic metrics show greater consistency across taxonomic rank and less

sensitivity to strain variation across microbiome samples. Thus, our metrics address key

challenges in the interpretation of core microbiomes.

Conclusions: We provide a phylogenetic framework for characterizing and comparing core

microbiomes. Importantly, the methods that we propose allows integration of microbiome

properties across taxonomic rank. Ultimately, this will provide both a more consistent picture of

the core microbiome, as well as novel biological insight into the conserved components of

microbial communities.

Introduction

Core microbiomes have become increasingly important as a dimension for studying and

characterizing sets of microbial communities.[1,2] Broadly speaking, a core microbiome is the

group of microbial taxa, genes, or functions shared by a large number of microbiome samples

from a specific type of habitat. Often, the habitat is defined based on a host species[3–5] (e.g., the

core microbiome of the human gut[6]), or a specific type of ecosystem (e.g., the core microbiome

of a particular soil type[7]). However, much broader[8] (e.g., the core microbiome of ruminant guts[9])

and much narrower (e.g., the core microbiome of a particular host population[10] or host

genotype[11]) habitat classes can be used as well. Because core microbial taxa are commonly, if

not always, associated with a specific habitat, they are thought to be important for identifying

microbial persistence requirements within the habitat, predicting ecosystem function[12–15] or, in

the case of host-associated microbiomes, understanding symbiotic relationships essential to host

performance.[16–19] By contrast, it is generally assumed that microbial taxa occurring only

sporadically in a given habitat are more likely to be transients or other community members that

do not contribute substantially to host or ecosystem function.

While the concept of a core microbiome is both simple and appealing, how to best define the core microbiome of any given habitat remains unclear.[1,2] In general, three strategies have been proposed[1] based on 1) the frequency of occurrence (i.e., incidence) of microbial taxa across samples, 2) the abundance of microbial taxa in individual samples or across a set of samples, or 3) a combination of occurrence and abundance criteria. Within these three strategies, however, there are a wide range of parameters that can influence interpretation. One obvious example is the threshold occurrence rate or abundance that a microbial taxon must achieve to be considered part of the core. In some cases, microbial taxa are included in the core if they are found in at least two samples from a given habitat. In other cases, however, much higher core thresholds have been used, such as requiring that a microbial taxon be present in half or even all of the samples from a particular habitat. Some studies even present analyses across a range of core thresholds so that results can be interpreted in context.[20,21]

Beyond the abundance or occurrence threshold, another decision that can influence identification of core microbial taxa is the taxonomic rank at which the core microbiome is defined. Studies exist spanning a wide range of taxonomic ranks from phyla to amplicon sequence variants (ASVs).[1] Not surprisingly, examining the core microbiome at different taxonomic scales often results in different conclusions. Higher taxonomic ranks like phyla and order tend to be more conserved across samples, resulting in a larger proportion of microbial taxa being identified as part of the core. However, neither ecology nor function are widely conserved at these higher taxonomic ranks, making the core microbiome less useful for informing host or ecosystem ecology. By contrast, at very low taxonomic ranks, it is unclear whether variation between

microbial taxa has significant ecological or functional consequences. ASVs, for example, differ

by one base pair in the portion of the marker gene used for sequencing.[22] However, in microbial

taxa with multiple copies of the 16S rRNA marker gene, individual gene copies can vary

intragenomically by >1%.[23,24] Thus, ASVs may be much more diverse than the populations or

functions that they are being used to represent.[25] This can impact the core microbiome in one of

two ways. First, if multiple closely related ASVs are present in the threshold number of samples,

they may all be counted individually as part of the core, artificially inflating core diversity.

Second, if individual samples each contain unique but closely related ASVs, none of the ASVs

may occur in sufficient samples to be included in the core, artificially deflating core diversity.

Perhaps because of the challenges associated with defining core microbiomes at very high and

very low taxonomic ranks, most analyses of core microbiomes have used intermediate taxonomic

ranks, for example 97% operational taxonomic units (OTUs) or genera.[1]

The importance of taxonomic scale, and the apparent variation in study outcomes as a function of

taxonomic rank are not unique to analysis of core microbiomes.[1] Indeed, these same challenges

occur in more general types of ecological analyses of both microbial and macrobial systems, for

example, calculation of α- and β-diversity in and between ecological communities. This has led

to the development of different phylogenetically aware metrics including Faith's phylogenetic

diversity[26] (PD), Rao's quadratic entropy[27] and phylogenetic entropy[28] for α-diversity, and

UniFrac distances[29] for β-diversity. Phylogenetically aware metrics are advantageous because

they integrate information across taxonomic ranks based on the branch lengths in a phylogenetic

tree. This removes the need to select an arbitrary taxonomic rank for any particular ecological

analysis. Further, because some phylogenetic properties can serve as indicators of both

functional diversity[30] and evolutionary potential[31] (i.e., capacity to evolve in response to a

perturbation, for example environmental change[32–34]), phylogenetically aware metrics are

thought to provide more informative comparisons of properties like diversity than do analyses at

a single taxonomic scale.

Despite the many benefits of phylogenetically aware ecological analyses, phylogenetically aware

metrics for characterizing the core microbiome are currently lacking. In this paper, we develop a

framework for analyzing core microbiomes using phylogenetic methods. To do this, we first

develop the concept of a 'core community phylogeny' - a phylogeny where branches are defined

based on being present in multiple samples from a single type of host or environment. We then

use this phylogeny to examine common measures of core communities, including Venn

diagrams, as well as metrics of within-habitat and between-habitat diversity.

Core Community Phylogenies

All phylogenetically aware metrics of ecological communities, for example Faith's phylogenetic

diversity (PD) and UniFrac distance, are based on an underlying phylogenetic tree comprising

taxa present in the community. Logically, the analogous phylogenetic tree for core microbiomes

should comprise all microbial taxa that are conserved across microbiomes from a particular type

of habitat. However, for core microbiomes, there are two possible ways of defining such a tree: a

'tip-based' approach, and a 'branch-based' approach.

Tip-based phylogeny: One way to build a core community phylogeny is to first identify core

microbial taxa based on their occurrence across a threshold number of microbiome samples from

a particular type of habitat. Once these microbial taxa have been identified, a phylogenetic tree

can be reconstructed based on the sequences (or other traits) of the conserved taxa. Notably,

conserved microbial taxa can be identified at any pre-selected taxonomic rank (e.g., phyla,

genera, 97% OTUs, ASVs, etc.) provided that a phylogenetic tree can be built connecting the

various microbial lineages.

Branch-based phylogeny: An alternative way to build a core community phylogeny is to first

build a phylogeny of all microbial taxa present in a set of microbiome samples from a particular

type of habitat. This phylogenetic tree can then be examined, branch-by-branch, to determine

which branches are present on/in a threshold number of microbiome samples. Branches shared

by a threshold number of samples can then be retained, while branches not shared by the

threshold number of samples can be discarded. Note that, at least for a rooted tree, this results in

discarding shallow branches connected to deeper branches that have also been discarded, such

that the remaining phylogeny is fully connected (but see below for issues that arise with unrooted

trees). Like the tip-based phylogeny, the branch-based phylogeny can have leaf nodes at any

taxonomic rank. However, because the branch-based phylogeny integrates information on shared

microbial lineages across all depths of the phylogeny both for defining the core and calculating

core metrics, there is less reason to use high taxonomic levels (see 'Examples' below). Thus, we

recommend ASVs as the default choice.

Importantly, the tip-based phylogeny and the branch-based phylogeny for a given system will not

always be identical. Whereas the branch-based phylogeny will always have all of the branches

present in the tip-based phylogeny, the tip-based phylogeny will often be missing some of the

branches present in the branch-based phylogeny (see Figure 1). More specifically, the tip-based

phylogeny will be missing branches whenever a set of microbiome samples share closely related,

but not identical, taxa. In this case, none of the shallow branches will be considered core for

either the tip-based or the branch-based phylogeny. However, the deeper branches from which

all of the related tips are derived may be present on the threshold number of samples and thus

may be retained in the branch-based phylogeny. By contrast, these branches will be discarded in

the tip-based phylogeny, since definition of core is solely based on shared leaf nodes. This

fundamental difference is what makes the branch-based approach less sensitive to variations in

the precise taxonomic scale considered, at least when making relative comparisons between two

or more core communities (see 'Examples' below).

Figure 1 shows the construction of both a tip-based phylogeny (A) and a branch-based

phylogeny (B) from the same set of microbiome samples. In Figure 1, the non-core branches of

the microbial phylogeny are shown in black, while the core branches are shown in red. Notice

that there are differences between the tip-based phylogeny and the branch-based phylogeny (see

starred branches). In particular, the branch-based phylogeny includes additional internal branches

that are missing from the tip-based phylogeny. This occurs because some clades are represented

in the threshold number of samples, and thus are part of the branch-based core microbiome, even

though the individual taxa within the clades are not present in the threshold number of samples.

Figure 1. Examples of a tip-based phylogeny (A) and a branch-based phylogeny (B) for the same microbial phylogeny. Branches of the microbial phylogeny that are not part of the core community phylogeny are shown in black, while branches that are part of the core community phylogeny are shown in red. For this example dataset, there are 6 samples (red, orange, yellow, green, blue, and purple squares) and the core threshold is set at 0.5. Thu,s being part of the core requires being present in 3 samples. Colored boxes at the end of each terminal branch indicate the samples that a particular microbial taxon is present on. Stars are used to mark branches that are included in the branch-based phylogeny but not in the tip-based phylogeny.

Another consideration in defining the core community phylogeny is whether to assume that the

root node is the most recent common ancestor (MRCA) of all taxa in the sample. The root node

should only be used as the MRCA when a rooted tree is provided. Otherwise, the root node is

arbitrary, and there is no reason to believe that the root node is any more basal than the other

nodes in the tree. Fortunately, microbial phylogenies generated using pipelines like Qiime2[35]

usually have the option for rooting trees (although this is typically done through midpoint

rooting, which involves multiple assumptions). Similar to UniFrac, we strongly encourage using

a rooted tree and forcing the root node to be the MRCA. This is particularly true for branch-

based core community phylogenies. The reason is as follows: using the branch-based approach,

branches are counted based on their inclusion in individual phylogenies from each sample. When

a root node is defined, these individual sample phylogenies necessarily include the root. In

absence of a root, however, individual sample phylogenies only span the taxa present in each

sample (core or otherwise). Consequently, basal branches may not be counted if entire clades are

differentially distributed across subsets of samples. When this happens, distinct clades are not

necessarily connected in the phylogeny that emerges. To remedy this, we connect the final core

community phylogeny based on the MRCAs for all of the distinct clades. This, however, does

not guarantee that branches will not be lost in the process. For tip-based phylogenies, the choice

to use an unrooted tree is less problematic because the branches of the core community

phylogeny are selected based on the occurrence rate of microbial taxa (tip nodes), not on the

occurrence rate of the branches themselves. Even for tip-based trees, however, we recommend

using a rooted tree and assuming that the root node is the MRCA.

Phylogenetic Metrics for Core Microbiomes

Once a core community phylogeny has been constructed, it can be used to perform phylogenetic

analyses of microbiome samples from different types of habitats. Here we consider several

common analysis methods, including Venn diagrams, a core microbiome equivalent of Faith's

phylogenetic diversity (α-diversity), and a core microbiome equivalent of UniFrac distance (β-

diversity).

Venn Diagrams: Venn diagrams are one of the most common methods for visualizing core

microbiomes across habitats. Non-phylogenetic Venn diagrams show counts of the core

microbial taxa shared by different combinations of habitat types. This can be extended to a

phylogenetic framework by replacing counts of shared microbial taxa with shared phylogenetic

branch lengths. Briefly, the core community phylogenies from different habitat types can be

compared, branch-by-branch, to determine which branches appear in which habitat

combinations. The total lengths of these shared branch segments can then be calculated and

reported either as an absolute value or as percentages of the total branch length spanning the core

community phylogenies of all habitats in the analysis.

***Core Faith's Phylogenetic Diversity* (PD):** The phylogenetic equivalent to richness (a count of

microbial taxa present in a community), Faith's PD is the sum of the branch lengths of all

branches spanning the taxa present in a particular microbiome sample or ecological community.

This definition can easily be extended to core microbiomes by summing the branch lengths of all

branches in the core community phylogeny. Doing so gives a measure of how much

phylogenetic diversity is conserved across a threshold number of microbiome samples from a

particular type of habitat.

***Core UniFrac Distances*:** The phylogenetic equivalent to Jaccard's index[36] (the fraction of

unique microbial taxa between two communities), the UniFrac distance[29,37] between two

communities is the sum of the branch lengths that are unique to each microbiome sample or

ecological community divided by the sum of the branch lengths of all branches spanning taxa

present in both microbiome samples or ecological communities. Again, this definition can be

extended to core microbiomes by dividing the branch lengths unique to each core community

phylogeny by the total branch lengths of the two core community phylogenies combined. Doing

so gives a measure of how phylogenetically distinct the core microbiomes are from two different

types of habitats.

 Examples

 Additional file 1 outlines the calculation of Venn diagrams, core Faith's PD, and core UniFrac

 distances using a set of highly simplified core community phylogenies. Specifically, we show

 differences between tip- and branch-based approaches, and outline calculations with both rooted-

 and unrooted trees. In the remainder of the main text, we apply our methods to more realistic

 systems using rooted trees. More specifically, we consider two datasets - one comprising the skin

 microbiota from four lizard populations and the other comprising the gut microbiota from the

 same four lizard populations (see Materials and Methods). In doing so, we show how our

 phylogenetic metrics provide additional and/or alternate information compared to existing

 metrics. We further demonstrate how our approach resolves some of the outstanding issues that

 have been raised regarding choice of taxonomic rank for defining the core microbiome.

 Figure 2 shows standard Venn diagrams based on ASV counts (A, D) along with our new

 phylogenetic Venn diagrams for tip-based (B, E) and branch-based (C, F) trees. As compared to

 our phylogenetic Venn diagrams, non-phylogenetic Venn diagrams (A, D) show the lowest

 percentage of system diversity in the compartment representing intersection of all four lizard

 populations (21.4% for skin and 0% for gut) and the highest percentage of system diversity in the

 four compartments unique to single lizard populations (42.5% for skin and 65% for gut). This is

 not surprising. Using a non-phylogenetic approach, microbial taxa from different host

 populations must be identical to be counted in intersection compartments. This biases against

 finding shared microbial diversity, particularly when using ASVs, because microbial lineages

 with small differences in 16S rRNA gene sequences (e.g., one base pair substitutions) are

deemed unique. By contrast, with a phylogenetic approach, even if different host populations

Figure 2. Venn diagrams for lizard skin (A-C) and gut (D-F) microbiota showing the percentage of the total ASV counts shared by the core microbiota from each lizard population (A, D) and the phylogenetic branch length shared by the core microbiota from each lizard population using either tip-based (B, E) or branch-based (C, F) phylogenies. In all Venn diagrams, lizard populations are as follows: *A. arizonae* (yellow), *A. neomexicanus* from SBluG (green), *A. neomexicanus* from SNW (blue) and *A. marmoratus* (red).

have slightly different ASVs, this only contributes a small amount of diversity to the unique

compartments of the Venn diagram. Meanwhile, the majority of the diversity is allotted to

intersection compartments because the ASVs from the different host populations share most of

their phylogenetic history. Comparing branch-based and tip-based phylogenetic approaches,

branch-based phylogenies show a higher percentage of system diversity in the compartment

representing intersection across all four populations (57.64% for skin and 21.45% for gut) and a

lower percentage of system diversity in the four compartments unique to single lizard

populations (18.24% for skin and 31.22% for gut). This reflects the inclusion of additional, deep

internal branches in branch-based phylogenies that are dropped from tip-based phylogenies (see

Figure 1). These basal branches are more likely to be conserved across different host populations

(i.e., phyla are more likely to be shared than ASVs), inflating diversity in shared compartments

of the Venn diagram.

Figure 3 (A, C) shows the tip-based and branch-based core Faith's PD, along with core richness

(count of microbial taxa) as a function of taxonomic rank for lizard skin and gut microbiota. For

the sake of comparison across metrics and taxonomic rank, we have plotted the core Faith's PD

of three lizard populations (*A. marmoratus* at site SNW as red, *A. neomexicanus* at SNW as blue,

and *A. neomexicanus* at SBluG as green) relative to the fourth lizard population (*A. arizonae* at

SBluG; see Materials and Methods). Figure 3 (B, D) shows the range of values obtained for

relative branch-based and tip-based core Faith's PD and core richness across all taxonomic ranks

considered. Importantly, for both skin and gut microbiota, phylogenetic measures, and the

branch-based core Faith's PD in particular, show much less variation across taxonomic ranks.

For example, whereas richness indicates that the core gut microbiota of the *A. neomexicanus*

population at SNW (blue) is anywhere from 2.28 (microbial order) to 9.25 (microbial species)

times as diverse as the core gut microbiota of the *A. arizonae* population at SBluG, branch-based

core Faith's PD suggests a much narrower range from 1.73 (microbial species) to 1.80 (microbial

family). Not all differences in metric stability across taxonomic rank are as dramatic. Still, the

branch-based core Faith's PD typically shows the least variation across taxonomic ranks, the tip-

based core Faith's PD the second least variation, and richness the most variation.

Figure 3. Comparison of alpha diversity across taxonomic ranks (A, C) for the branch-based
Core Faith's PD (solid, squares), the tip-based Core Faith's PD (dotted, triangles), and richness
(dashed, circles). In both panels, we show the diversity of the focal population relative to the
diversity of the *A. arizonae* population. Focal populations are *A. marmoratus* (red), *A.
neomexicanus* at SNW (blue) and *A. neomexicanus* at SBluG (green). Bar graphs (B, D) showing
minimum and maximum values of alpha diversity across taxonomic rank for *A. marmoratus*
(red), *A. neomexicanus* at SNW (blue), and *A. neomexicanus* at SBluG (green) relative to *A.
arizonae*. Diversity metrics are indicated as follows: 'BranchPD' is the branch-based phylogeny
approach, 'TipPD' is the tip-based phylogeny approach, and 'Richness' is a count of taxa for
each taxonomic rank.

Figure 4. Comparison of beta diversity across taxonomic ranks (A, C) for the branch-based Core UniFrac distance (solid, squares), the tip-based Core UniFrac distance (dotted, triangles), and Jaccard distance (dashed, circles). In both panels, we show turnover of the focal population with respect to the *A. arizonae* population. Focal populations are *A. marmoratus* (red), *A. neomexicanus* at SNW (blue), and *A. neomexicanus* at SBluG (green). Bar graphs (B, D) showing minimum and maximum values of beta diversity (turnover or distance) across taxonomic rank between *A. arizonae* and *A. marmoratus* (red), *A. neomexicanus* at SNW (blue), and *A. neomexicanus* at SBluG (green) with respect to *A. arizonae*. Diversity metrics are as follows: where 'BranchUniFrac' is the branch-based phylogeny approach, 'TipUniFrac' is the tip-based phylogeny approach, and 'Jaccard' is a count of taxa for each taxonomic rank.

Figure 4 (A, C) shows the tip-based and branch-based UniFrac distances, along with the Jaccard

distance (turnover) between the core skin and gut microbiota for *A. arizonae* at SBluG and the

core skin and gut microbiota for all three other whiptail populations (see 'Materials and

Methods') as a function of the taxonomic rank. Figure 4 (B, D) shows the range of values

obtained for tip-based and branch-based UniFrac distances as well as Jaccard distances across all

taxonomic ranks considered. Broadly speaking, the non-phylogenetic approach (i.e., Jaccard

dissimilarity) gives the largest dissimilarities/distances, while the branch-based phylogenetic

approach gives the smallest dissimilarities. With respect to variation in dissimilarity across

taxonomic rank, we again see that the branch-based phylogenetic distance measure gives the

least variation, while the tip-based distance gives intermediate variation and the Jaccard distance

gives the most variation.

The holobiont R package

To facilitate the use of our new phylogenetic measures of core microbiomes, we developed a

new R package, holobiont, that includes functions for identifying the core community

phylogeny in any microbiome, drawing phylogenetic Venn diagrams, calculating the core Faith's

PD for a set of communities, and calculating the core UniFrac distance between two sets of

communities. Briefly, the coretree function takes a phyloseq object[38] and a value for the core

threshold and identifies the core community phylogeny, coloring the branches of the core

community phylogeny red, while leaving the remaining branches of the community tree black.

The corePhyloVenn function takes a phyloseq object, a vector defining the habitat class for

each microbiome sample, and a core threshold and draws a phylogenetic Venn diagram of either

branch lengths or percentages of branch lengths shared by the core community phylogenies of

different combinations of habitat classes. The `coreVennTree` function takes the same input as

the `corePhyloVenn` function and colors the branches of the phylogenetic tree based on the

combination of habitat class core community phylogenies that they belong to. The `coreFaithsPD`

function takes a phyloseq object and a core threshold and calculates the core Faith's PD. Finally,

the `coreUniFrac` function takes a phyloseq object and a vector specifying which samples belong

to which habitat class and calculates the UniFrac distance between the two core community

phylogenies. Additional details outlining the methods and use of our R package can be found at

https://bewicklab.weebly.com/databases--packages.html.

Discussion

In this paper, we propose a framework for performing phylogenetically aware analyses of core

microbiomes. More specifically, we develop the concept of a core community phylogeny - a

phylogeny based on the microbial lineages that are core to a specific habitat class. We then

propose two different methods for building this tree - a tip-based approach and a branch-based

approach. For the tip-based approach, core membership is decided based on occurrence of the

leaf (tip) nodes. This is akin to the standard, non-phylogenetic method for defining a core

microbiome. However, once members of the core microbiome have been identified, a

phylogenetic tree is constructed. This allows for calculation of phylogenetically aware metrics

that leverage the relatedness of the leaf nodes to determine overall properties (e.g., diversity) of

the core microbiome. For the branch-based approach, core membership is decided independently

for each branch of the phylogenetic tree. Therefore, in the branch-based approach, phylogeny

informs not only the calculation of metrics but also, the identification of the core microbiome

itself. The downside of the branch-based approach, however, is that the 'core microbiome' often

includes some internal or ancestral nodes without their associated leaf nodes. Thus, metrics

calculated using the branch-based approach are not necessarily based on a set of extant taxa.

Rather, they are based on a set of complete and partial microbial lineages. Still, because it uses

phylogenetic information to both define the core microbiome and to calculate core metrics, we

suggest using the branch-based approach except when there is a strong desire to identify specific

microbial taxa within the core (e.g., to select for culturing experiments).

As compared to existing metrics, the primary advantage of our phylogenetic metrics is that they

integrate core microbiome properties (i.e., richness, turnover, shared taxa) across taxonomic

scales. Thus, our phylogenetic metrics are more likely to provide biologically relevant

information. This is especially true when non-phylogenetic metrics are calculated at taxonomic

ranks that fail to capture meaningful biological variation.[39] If, for example, the chosen taxonomic

rank is too low, substantial biological variation may be identified that has no functional or

ecological consequences. Alternatively, if the chosen taxonomic rank is too high, large amounts

of meaningful biological variation may be pooled into the same taxonomic unit. Depending on

the system, the type of variation considered and the metric calculated, it can be difficult to *a*

priori determine (or even *a posteriori* assess) which taxonomic ranks are more or less likely to

be biologically informative. Further, in certain instances multiple taxonomic ranks exhibit

substantial meaningful biological variation. In this case, a non-phylogenetic metric at any single

taxonomic rank - even a well-chosen taxonomic rank - cannot fully capture the underlying

signal. Phylogenetic metrics circumvent these challenges by summarizing the full spectrum of

biological variation across all taxonomic ranks, with variation at lower taxonomic ranks

weighted less than variation at higher taxonomic ranks.

Another benefit of our branch-based approach (but not our tip-based approach) is that it

overcomes challenges associated with defining the core microbiome in the face of high levels of

strain variation. This is common in certain microbiome systems, including many host-associated

microbiomes. The pervasive human skin isolate, *Staphylococcus epidermidis*, for example,

evolves on individual humans over time,[40,41] resulting in strain level variation across the human

population. Another common human skin isolate, *Cutibacterium acnes*, exhibits even more

dramatic strain-level variation, showing differences in sequences across pores from a single

human individual.[42] Indeed, in-depth profiling of the human microbiome has shown that there

are numerous personalized microbial taxa that are found on only one or several individuals.[41,43]

Depending on the taxonomic rank at which the core microbiome is defined and the degree to

which any single microbial strain is shared among individual hosts, natural strain-level variation

can cause broadly shared microbial clades to be obscured from the non-phylogenetic core

microbiome. By contrast, if we apply our branch-based approach to these same scenarios, only

the terminal branches of the tree are omitted. Meanwhile, the more basal branches of the clades

are retained because they are shared across all strains.

Because our phylogenetic metrics summarize microbiome measures across taxonomic rank and

because the definition of the core microbiome - at least when using the branch-based approach -

is less sensitive to strain-level variation, our phylogenetic metrics tend to be more consistent than

comparable non-phylogenetic metrics when calculated at different taxonomic ranks. This is true

both in a relative (i.e., comparison between microbiome samples) and an absolute sense. To see

why phylogenetic metrics are more consistent, consider a habitat with hundreds of closely related

core microbial strains within a single phylum. Using a non-phylogenetic approach this system

will have a very high α-diversity at the ASV-level, but a very low α-diversity at the phylum-

level. By contrast, using our phylogenetic approach, α-diversity will be low regardless of

whether the leaf nodes are defined at the ASV- or phylum-level. This is because, even when the

leaf nodes are ASVs all of the different ASV strains contribute very short branch lengths.

Consequently, they do not substantially inflate α-diversity beyond the diversity of the single

phylum. The greater consistency of our phylogenetic metrics across taxonomic ranks means that

it is less important to choose the correct taxonomic rank, and that it is rarely disadvantageous to

use ASVs. This addresses one of the challenges that has been raised about interpreting core

microbiomes when outcomes are variable across taxonomic rank.[1]

The primary goal of this paper is to introduce a framework for phylogenetic analysis of core

microbiomes. For this reason, we have focused on the concepts required to define a core

community phylogeny and have outlined the application of a core community phylogeny to three

types of microbiome analyses - Venn diagrams, α-diversity (Faith's PD), and β-diversity

(UniFrac). Our broad approach, however, is not limited to these particular types of analyses.

Indeed, additional analyses could be developed within the same framework. A straightforward

example would be to use UniFrac distances between core microbiomes to project into ordination

space (e.g., Principal Coordinate Analysis or Nonmetric Multidimensional Scaling). This would

provide phylogenetically informed visualization of the relatedness of different core microbiomes.

A more challenging extension would be to develop a method for applying our phylogenetic

framework to core microbiomes based on abundance thresholds[44] or even to develop metrics

where phylogenetic branches are weighted based on average abundances or occurrence rates in

the samples across which the core microbiome is defined.

While initially developed for microbiome analysis, beyond microbiomes, our community

phylogenetics approach and associated R package may ultimately have wide application to

macrobial systems as well. Currently, very few macrobial studies attempt to identify core

communities (but see Murdock et al.[45]). This is likely because, at least historically, macrobial

systems have not been studied with many replicates. Indeed, even when replicate macrobial

communities are characterized, the effort required to identify all taxa in a single community

means that replicate numbers are low. This makes it difficult to identify a core community.

Further, in species poor macrobial systems where characterization of replicate communities is

possible, the core community concept is not as necessary or useful because the systems are

simple enough to analyze in their entirety. As sequencing techniques improve, however, and

more and more macrobial systems are characterized in a manner similar to microbiomes (e.g.,

through the use of eDNA[46,47]), it seems likely that core communities will emerge as a common

form of analysis for species rich macrobial systems (e.g. tropical rainforests, litter insects) as

well. There is no reason why our proposed framework cannot be applied to macrobial

communities. In fact, our framework can be applied to any system including prokaryotic and

eukaryotic communities, host-associated and free-living communities, and microbial (virus,

bacteria, fungi, etc.) and macrobial communities. It can also be applied to systems characterized

through a wide variety of methods, even including non-sequencing techniques. The only

requirements for our approach are tables of presences/absences of different taxa across different

communities, along with a phylogenetic tree comprising all taxa present across all communities.

Thus, occurrence data can be based on sequencing, but can also be based on any other form of

observation, including transect or plot surveys. Likewise, phylogenies can be constructed using

sequence data, including sequencing of any kind (single marker genes, combinations of marker

genes, whole genomes), but can also be generated based on morphology or other traits

commonly used to build trees (though the validity of any core community analysis will depend

on the accuracy of both the occurrence table and the phylogeny used).

The identification and characterization of core microbiomes has become a hallmark of

microbiome analysis. However, to date, most analyses of core microbiomes have focused on

taxonomy at specified taxonomic ranks. By contrast, phylogenetically aware metrics for

characterizing core microbiomes are largely lacking. Phylogenetically aware metrics have the

benefit of integrating information across taxonomic ranks. This can be particularly helpful when

analysis at different taxonomic ranks yields different outcomes - a common problem in ecology

and evolution[48–52] including the study of core microbiomes.[1] Our phylogenetic framework

overcomes this challenge. Indeed, even when our analysis is performed at different taxonomic

ranks, it typically yields more consistent results than non-phylogenetic methods. With the

development of our new framework and associated R package, we provide a mechanism to bring

phylogenetically aware community analyses to the study of core microbiomes. Ultimately, this

will allow for novel insight into the conserved components of microbial communities both within

and between different types of habitats.

Methods

Data collection

We sampled the microbiota from several whiptail lizard species (genus *Aspidoscelis*, family

Teiidae) at two locations (SBluG and SNW; see Camper et al.[53] for more site details) within

Sevilleta National Wildlife Refuge. Each location represents a site of syntopy between hybrid *A.*

neomexicanus and a progenitor species, *A. marmoratus* (♀) and *A. arizonae* (♂, recently split

from *A. inornatus*[54]). Following taxonomic guidelines by Tucker et al.[55] and Walker et al.,[56] we

refer to *Aspidoscelis* spp. using the masculine epithet. We captured lizards using a rapidly

deployable drift fence trapping array lined with box funnel traps, further outlined in Camper et

al.,[53] and by hand or lizard noose when possible. Upon capture, we sampled the gut (=cloacal)

and skin microbiota of each species following a swabbing protocol in Camper et al.[20] We

sampled each location until we captured at least 15 female lizards of each whiptail species. All

samples were sent to ZymoBIOMICS Targeted Sequencing Service for Microbiota Analysis

(Zymo Research) using 16S rRNA gene sequencing. Additional details are available from

https://www.zymoresearch.com/pages/16s-its-amplicon-sequencing and Camper et al.[20] FASTQ

files for lizard microbiomes, as well as environmental and sequencing controls, are available at

(These will be uploaded to NCBI upon publication).

Data analysis

Using the .biom files supplied by Zymo, we first removed 16S rRNA gene sequences that could

not be identified as Bacteria or Archaea based on the proprietary Zymo database. Additionally,

we removed all samples with a read depth lower than 10,000 reads. Finally, we rarefied all

samples to the read depth of the lowest sample from each particular body site using the

rarefy_even_depth function from the phyloseq package.[38] All analyses were performed using

the R programming language[57] (version 4.2.1) available on our Github page.

For Venn diagrams, we used the `corePhyloVenn` function in our `holobiont` package, selecting

habitat classes based on species x site. We generated Venn diagrams separately for skin and gut

microbiota. In all cases, we reported percentages of the total branch length and used rooted

phylogenies. To compare standard Venn diagrams to our phylogenetic Venn diagrams, we

counted the number of core taxa shared by each species x site combination and used these counts

(richness) to generate a standard core microbiome Venn diagram. All Venn diagrams were

calculated based on ASVs using a core threshold of 0.5 (i.e., a microbial taxon or branch must be

present in 50% of the samples from a particular lizard species x site habitat class).

For Faith's PD, we used the `coreFaithsPD` function in our `holobiont` package. To compare

across taxonomic ranks, we took the original ASV .biom file supplied by Zymo to generate an

ASV phyloseq object.[38] We then used the `tax_glom` function from the `phyloseq` package to

generate OTU tables and phylogenies at taxonomic ranks from species ('Rank7') through to

order ('Rank4'). This allowed us to calculate Faith's PD at all ranks, as well as the related non-

phylogenetic richness. For the latter, we summed up the number of microbial taxa at any given

taxonomic rank that were present in the core microbiome of the focal habitat. All Faith's PD and

richness calculations were performed on rooted trees assuming a core threshold of 0.5.

For Unifrac distances, we used the `coreUniFrac` function in our `holobiont` package.

Comparisons across taxonomic ranks were performed as described for Faith's PD. To compare to

the analogous non-phylogenetic metric, we calculated the Jaccard distance based on taxonomic

co-occurrence patterns between pairs of core microbiomes from different habitats. This

functionality is included as the `coreJaccard` function in our `holobiont` package. Again, all

UniFrac and Jaccard calculations were performed on rooted trees assuming a core threshold of

0.5.

All data and code necessary to recalculate our results are available through GitHub at:

https://github.com/bewicklab/CoreTrees

Declarations

Ethics approval and consent to participate

All research was approved by Clemson University under IACUC protocol numbers #2020-015

and #2021-047. We completed this work under the Sevilleta National Wildlife Refuge Special

Use Permit #SEV_Bewick_Camper_2022_59 and the New Mexico Department of Game and

Fish permit authorization #3772.

Consent for publication

not applicable

Funding

This study was funded by NSF awards #2105604 and #2025541, a Clemson University Support

for Early Exploration and Development (CUSEED) Grant, and the Clemson University Creative

Inquiry (CI) Program.

Availability of data and materials

All sequence data will be deposited in the National Center for Biotechnology Information

(NCBI) database upon acceptance. BIOM and metadata files are included in Additional file 2.

Additionally, BIOM files, metadata files, and all code necessary for the analyses presented in

this manuscript are available at https://github.com/bewicklab/CoreTrees. The holobiont R

package is available at https://bewicklab.weebly.com/databases--packages.html.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

BTC and SAB conceived the idea and wrote the first draft of the manuscript. BTC performed all

related fieldwork. SAB developed the code and data analysis. All authors contributed

substantially to the content of the manuscript.

Acknowledgements

We thank Thomas Dempster, August Spencer, Eva Purcell, Riley Manuel, Drew Kanes and

Zachary Laughlin for their field assistance in New Mexico as well as Lily Margeson, Simon

Dunn, Georgianna Bellinger, Henry Egloff, Kaila Hodges, Camryn Lachica, and Savannah Utz

for their assistance assembling drift fence trapping arrays.

References

1. Neu AT, Allen EE, Roy K. Defining and quantifying the core microbiome: Challenges and prospects. *Proceedings of the National Academy Sciences*. 2021;118(51):1-10. doi:10.1073/pnas.2104429118/-/DCSupplemental

2. Custer GF, Gans M, van Diepen LTA, Dini-Andreote F, Buerkle CA. Comparative Analysis of Core Microbiome Assignments: Implications for Ecological Synthesis. *mSystems*. 2023;8(1):1-14. doi:10.1128/msystems.01066-22

3. Huse SM, Ye Y, Zhou Y, Fodor AA. A core human microbiome as viewed through 16S rRNA sequence clusters. *PLoS One*. 2012;7(6):1-12. doi:10.1371/journal.pone.0034242

4. Zaura E, Keijser BJ, Huse SM, Crielaard W. Defining the healthy "core microbiome" of oral microbial communities. *BMC Microbiol*. 2009;9(259):1-12. doi:10.1186/1471-2180-9-259

5. Muletz Wolz CR, Yarwood SA, Campbell Grant EH, Fleischer RC, Lips KR. Effects of host species and environment on the skin microbiome of Plethodontid salamanders. *Journal of Animal Ecology*. 2018;87(2):341-353. doi:10.1111/1365-2656.12726

6. Turnbaugh PJ, Gordon JI. The core gut microbiome, energy balance and obesity. *Journal of Physiology*. 2009;587(17):4153-4158. doi:10.1113/jphysiol.2009.174136

7. Pershina E V., Ivanova EA, Korvigo IO, et al. Investigation of the core microbiome in main soil types from the East European plain. *Science of the Total Environment*. 2018;631-632:1421-1430. doi:10.1016/j.scitotenv.2018.03.136

8. Salzman S, Whitaker M, Pierce NE. Cycad-feeding insects share a core gut microbiome. *Biological Journal of the Linnean Society*. 2018;123:728-738. https://academic.oup.com/biolinnean/article/123/4/728/4920834

9. Henderson G, Cox F, Ganesh S, et al. Rumen microbial community composition varies with diet and host, but a core microbiome is found across a wide geographical range. *Sci Rep*. 2015;5:1-15. doi:10.1038/srep14567

10. Hamady M, Knight R. Microbial community profiling for human microbiome projects: Tools, techniques, and challenges. *Genome Res*. 2009;19(7):1141-1152. doi:10.1101/gr.085464.108

11. Hernandez-Teran A, Navarro-Diaz M, Benitez M, Lira R, Wegier A, Escalante AE. Host genotype explains rhizospheric microbial community composition: The case of wild cotton metapopulations (Gossypium hirsutum L.) in Mexico. *FEMS Microbiol Ecol*. 2020;96(8):1-11. doi:10.1093/femsec/fiaa109

12. Fan K, Chu H, Eldridge DJ, et al. Soil biodiversity supports the delivery of multiple ecosystem functions in urban greenspaces. *Nat Ecol Evol*. 2023;7:113-126.

13. Sun L, Li G, Zhao J, Zhang T, Liu J, Zhang J. Core microbiota drive multi-functionality of the soil microbiome in the Cinnamomum camphora coppice planting. *BMC Microbiol*. 2024;24(1). doi:10.1186/s12866-023-03170-8

14. Björk JR, O'hara RB, Ribes M, Coma R, Montoya JM. The dynamic core microbiome:
structure, stability and resistance. *bioRxiv*. Published online 2017:1-50.
doi:10.1101/137885
15. De Filippis F, La Storia A, Stellato G, Gatti M, Ercolini D. A selected core microbiome
drives the early stages of three popular Italian cheese manufactures. *PLoS One*.
2014;9(2):1-8. doi:10.1371/journal.pone.0089680
16. Risely A. Applying the core microbiome to understand host–microbe systems. *Journal of
Animal Ecology*. 2020;89(7):1549-1558. doi:10.1111/1365-2656.13229
17. Zaura E, Keijser BJ, Huse SM, Crielaard W. Defining the healthy "core microbiome" of
oral microbial communities. *BMC Microbiol*. 2009;9(259):1-12. doi:10.1186/1471-2180-
9-259
18. Ganz HH, Jospin G, Rojas CA, et al. The Kitty Microbiome Project: Defining the Healthy
Fecal "Core Microbiome" in Pet Domestic Cats. *Vet Sci*. 2022;9(635):1-21.
doi:10.3390/vetsci9110635
19. Holyoak GR, Premathilake HU, Lyman CC, et al. The healthy equine uterus harbors a
distinct core microbiome plus a rich and diverse microbiome that varies with geographical
location. *Sci Rep*. 2022;12(14790):1-14. doi:10.1038/s41598-022-18971-6
20. Camper BT, Kanes AS, Laughlin ZT, Manuel RT, Bewick SA. Transgressive Hybrids as
Hopeful Holobionts. *bioRxiv*. Published online 2024:1-68.
doi:10.1101/2024.08.17.607925
21. Astudillo-García C, Bell JJ, Webster NS, et al. Evaluating the core microbiota in complex
communities: A systematic investigation. *Environ Microbiol*. 2017;19(4):1450-1462.
doi:10.1111/1462-2920.13647
22. Callahan BJ, McMurdie PJ, Holmes SP. Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. *ISME Journal*.
2017;11(12):2639-2643. doi:10.1038/ismej.2017.119
23. Větrovský T, Baldrian P. The Variability of the 16S rRNA Gene in Bacterial Genomes
and Its Consequences for Bacterial Community Analyses. *PLoS One*. 2013;8(2):1-10.
doi:10.1371/journal.pone.0057923
24. Coenye T, Vandamme P. Intragenomic heterogeneity between multiple 16S ribosomal
RNA operons in sequenced bacterial genomes. *FEMS Microbiol Lett*. 2003;228(1):45-49.
doi:10.1016/S0378-1097(03)00717-1
25. Schloss PD. Amplicon Sequence Variants Artificially Split Bacterial Genomes into
Separate Clusters. *mSphere*. 2021;6(4):1-6. doi:10.1128/msphere.00191-21
26. Faith DP. Conservation evaluation and phylogenetic diversity. *Biol Conserv*. 1992;61:1-
10.
27. Rao CR. Diversity and Dissimilarity Coefficients: A Unified Approach. *Theor Popul Biol*.
1982;21:24-43.

28. Allen B, Kon M, Bar-Yam Y. A new phylogenetic diversity measure generalizing the
shannon index and its application to phyllostomid bats. *American Naturalist*.
2009;174(2):236-243. doi:10.1086/600101
29. Lozupone C, Knight R. UniFrac: A new phylogenetic method for comparing microbial
communities. *Appl Environ Microbiol*. 2005;71(12):8228-8235.
doi:10.1128/AEM.71.12.8228-8235.2005
30. Tucker CM, Davies TJ, Cadotte MW, Pearse WD. On the relationship between
phylogenetic diversity and trait diversity. *Ecology*. 2018;99(6):1473-1479.
doi:10.1002/ecy.2349
31. Winter M, Devictor V, Schweiger O. Phylogenetic diversity and nature conservation:
Where are we? *Trends Ecol Evol*. 2013;28(4):199-204. doi:10.1016/j.tree.2012.10.015
32. Mouquet N, Devictor V, Meynard CN, et al. Ecophylogenetics: Advances and
perspectives. *Biological Reviews*. 2012;87(4):769-785. doi:10.1111/j.1469-
185X.2012.00224.x
33. Lankau R, Jørgensen PS, Harris DJ, Sih A. Incorporating evolutionary principles into
environmental management and policy. *Evol Appl*. 2011;4(2):315-325.
doi:10.1111/j.1752-4571.2010.00171.x
34. Lavergne S, Mouquet N, Thuiller W, Ronce O. Biodiversity and climate change:
Integrating evolutionary and ecological responses of species and communities. *Annu Rev
Ecol Evol Syst*. 2010;41:321-350. doi:10.1146/annurev-ecolsys-102209-144628
35. Estaki M, Jiang L, Bokulich NA, et al. QIIME 2 Enables Comprehensive End-to-End
Analysis of Diverse Microbiome Data and Comparative Studies with Publicly Available
Data. *Curr Protoc Bioinformatics*. 2020;70(1):1-46. doi:10.1002/cpbi.100
36. Jaccard P. Distribution de la flore alpine dans le bassin des Dranses et dans quelques
régions voisines. *Bulletin de la Société vaudoise des sciences naturelles*. 1901;37:241-
272.
37. Lozupone C, Lladser ME, Knights D, Stombaugh J, Knight R. UniFrac: An effective
distance metric for microbial community comparison. *ISME Journal*. 2011;5(2):169-172.
doi:10.1038/ismej.2010.133
38. McMurdie PJ, Holmes S. phyloseq: An R package for reproducible interactive analysis
and graphics of microbiome census data. *PLoS One*. 2013;8(4).
39. Dubois A. Phylogeny, taxonomy and nomenclature: The problem of taxonomic categories
and of nomenclatural ranks. *Zootaxa*. 2007;(1519):27-68. doi:10.11646/zootaxa.1519.1.3
40. Zhou W, Spoto M, Hardy R, et al. Host-Specific Evolutionary and Transmission
Dynamics Shape the Functional Diversification of Staphylococcus epidermidis in Human
Skin. *Cell*. 2020;180(3):454-470. doi:10.1016/j.cell.2020.01.006
41. Both A, Huang J, Qi M, et al. Distinct clonal lineages and within-host diversification
shape invasive staphylococcus epidermidis populations. *PLoS Pathog*. 2021;17(2):1-39.
doi:10.1371/JOURNAL.PPAT.1009304

42.   Conwill A, Kuan AC, Damerla R, et al. Anatomy promotes neutral coexistence of strains
680   in the human skin microbiome. *Cell Host Microbe*. 2022;30(2):171-182.
681   doi:10.1016/j.chom.2021.12.007

682   43.   Lloyd-Price J, Mahurkar A, Rahnavard G, et al. Strains, functions and dynamics in the
683   expanded Human Microbiome Project. *Nature*. 2017;550(7674):61-66.
684   doi:10.1038/nature23889

685   44.   Shade A, Stopnisek N. Abundance-occupancy distributions to prioritize plant core
686   microbiome membership. *Curr Opin Microbiol*. 2019;49:50-58.
687   doi:10.1016/j.mib.2019.09.008

688   45.   Murdock SA, Tunnicliffe V, Boschen-Rose RE, Juniper SK. Emergent "core
689   communities" of microbes, meiofauna and macrofauna at hydrothermal vents. *ISME*
690   *Communications*. 2021;1(27):1-13. doi:10.1038/s43705-021-00031-1

691   46.   García-Machado E, Laporte M, Normandeau E, et al. Fish community shifts along a
692   strong fluvial environmental gradient revealed by eDNA metabarcoding. *Environmental*
693   *DNA*. 2022;4(1):117-134. doi:10.1002/edn3.221

694   47.   Gelis ERE, Kamal MM, Subhan B, Bachtiar I, Sani LMI, Madduppa H. Environmental
695   biomonitoring of reef fish community structure with eDNA metabarcoding in the Coral
696   Triangle. *Environ Biol Fishes*. 2021;104(8):887-903. doi:10.1007/s10641-021-01118-3

697   48.   Cavender-Bares J, Kozak KH, Fine PVA, Kembel SW. The merging of community
698   ecology and phylogenetic biology. *Ecol Lett*. 2009;12(7):693-715. doi:10.1111/j.1461-
699   0248.2009.01314.x

700   49.   Graham CH, Storch D, Machac A. Phylogenetic scale in ecology and evolution. *Global*
701   *Ecology and Biogeography*. 2018;27(2):175-187. doi:10.1111/geb.12686

702   50.   Wanntorp HE, Brooks DR, Nilsson T, et al. Phylogenetic approaches in ecology. *Oikos*.
703   1990;57(1):119-132. https://www.jstor.org/stable/3565745

704   51.   Mouquet N, Devictor V, Meynard CN, et al. Ecophylogenetics: Advances and
705   perspectives. *Biological Reviews*. 2012;87(4):769-785. doi:10.1111/j.1469-
706   185X.2012.00224.x

707   52.   Hernández CE, Rodríguez-Serrano E, Avaria-Llautureo J, et al. Using phylogenetic
708   information and the comparative method to evaluate hypotheses in macroecology.
709   *Methods Ecol Evol*. 2013;4(5):401-415. doi:10.1111/2041-210X.12033

710   53.   Camper BT, Laughlin ZT, Kanes AS, Manuel RT. A portable and wind resistant drift
711   fence array for arid environments. *bioRxiv*. Published online 2023:1-25.
712   doi:10.1101/2023.09.28.560034

713   54.   Barley AJ, Reeder TW, Nieto-Montes De Oca A, Cole CJ, Thomson RC. A New Diploid
714   Parthenogenetic Whiptail Lizard from Sonora, Mexico, Is the "Missing Link" in the
715   Evolutionary Transition to Polyploidy. *Am Nat*. 2021;198(2):295-309.
716   doi:10.5061/dryad.2z34tmpm2

717   55.   Tucker DB, Colli GR, Giugliano LG, et al. Methodological congruence in phylogenomic
718   analyses with morphological support for teiid lizards (Sauria: Teiidae). *Mol Phylogenet*
719   *Evol*. 2016;103:75-84.
720   56.   Walker JM, Sullivan BK, Cordes JE. Sanctioned Nomenclature for Triploid
721   Parthenogenetic Colorado Checkered Whiptail and for Other Taxa of the Lizard Genus
722   Aspidoscelis (Family Teiidae). *Herpetol Rev*. 2021;52(3):550-558.
723   57.   R Core Team. R: A language and environment for statistical computing. Published online
724   2021.

Electronic Supplementary Material

core_community_phylogeny_example.docx

Additional file 1: A simplified system demonstrating the construction of core community

phylogenies and calculations of associated metrics.

biom_files.zip

Additional file 2: BIOM files, read counts, taxonomy assignments, microbial phylogenies, and

lizard metadata for gut and skin samples used in this study.

A. ASV Counts (Skin)

B. Tip-based (Skin)

C. Branch-based (Skin)

D. ASV Counts (Gut)

E. Tip-based (Gut)

F. Branch-based (Gut)