Phylogenetic Measures of the Core Microbiome

Sharon Anne Bewick[1]*, Benjamin Thomas Camper[1]

[1]Department of Biological Sciences, Clemson University, Clemson, South Carolina 29631, USA

*sbewick@clemson.edu

Abstract

Background: A useful concept in microbial ecology is the 'core microbiome.' Typically, core microbiomes are defined as the microbial taxa, genes, or functions shared by a threshold number of microbiome samples from a particular type of habitat (e.g., a particular type of host or a particular type of environment/ecosystem). In defining the core microbiome, the goal is to capture the portion of the microbial community that is conserved across samples from the focal habitat. Recently, there has been growing interest in developing methods to better characterize core microbiomes. As a result, numerous occurrence- and abundance-based measures have been defined. However, few have included phylogeny-aware metrics for analyzing core microbiomes. Results: In this paper, we develop the concept of the 'core community phylogeny' – a phylogeny where branches are selected based on their presence in multiple samples from a single type of habitat. We then use the core community phylogeny to define phylogenetic metrics describing the diversity of core microbiomes from a single type of habitat, the turnover between core microbiomes from two different types of habitats, and the shared diversity across core microbiomes from two or more different types of habitats. As compared to non-phylogenetic metrics, our phylogenetic metrics show greater consistency across taxonomic rank and/or

phylogenetic level, as well as less sensitivity to strain variation across microbiome samples.

Thus, our metrics address key challenges in the interpretation of core microbiomes.

Conclusions: We provide a phylogenetic framework for characterizing and comparing core

microbiomes. Importantly, the methods that we propose allow seamless integration of

microbiome properties across taxonomic rank and/or phylogenetic level. Ultimately, this will

provide both a more consistent picture of the core microbiome, as well as novel biological

insight into the conserved components of microbial communities.

Introduction

Core microbiomes have become increasingly important as a dimension for studying and

characterizing sets of microbial communities.[1,2] Broadly speaking, a core microbiome is the

group of microbial taxa, genes, or functions shared by a large number of microbiome samples

from a specific type of habitat. Often, the habitat is defined based on a host species[3–5] (e.g., the

core microbiome of the human gut[6]), or a specific type of ecosystem (e.g., the core microbiome

of a particular soil type[7]). However, much broader[8] (e.g., the core microbiome of ruminant guts[9])

and much narrower (e.g., the core microbiome of a particular host population[10] or host

genotype[11]) habitat classes can be used as well. Because core microbial taxa are commonly, if

not always, associated with a specific habitat, they are thought to be important for identifying

microbial persistence requirements within the habitat, predicting ecosystem function[12–15] or, in

the case of host-associated microbiomes, understanding symbiotic relationships that may be

important to host performance.[16–19] By contrast, it is generally assumed that microbial taxa

occurring only sporadically in a given habitat are more likely to be transients or other community

members that do not contribute substantially to host or ecosystem function.

While the concept of a core microbiome is both simple and appealing, how to best define the

core microbiome of any given habitat remains unclear.[1,2] In general, three different core criteria

have been proposed[1] based on 1) the occupancy (i.e., incidence rate, frequency of occurrence) of

microbial taxa across samples, 2) the abundance of microbial taxa in individual samples or

across a set of samples, or 3) a combination of occupancy and abundance criteria.[20,21] Within

these three strategies, however, there are a wide range of parameters that can influence

interpretation. One obvious example is the threshold occupancy or abundance that a microbial

taxon must achieve to be considered part of the core. In some cases, microbial taxa are included

in the core if they are found in at least two samples from a given habitat.[1] In other cases, much

higher core thresholds have been used, such as requiring that a microbial taxon be present in half

or even all of the samples from a particular habitat. Some studies even present analyses across a

range of core thresholds so that results can be interpreted in context.[22,23]

Beyond the occupancy and abundance thresholds, another decision that can influence

identification of core microbial taxa is the taxonomic rank or phylogenetic level at which the

core microbiome is defined. Studies exist spanning a wide range of taxonomic ranks and

phylogenetic levels from phyla to genera and from 97% operational taxonomic units (OTUs) to

amplicon sequence variants (ASVs).[1] Not surprisingly, examining the core microbiome at

different taxonomic and/or phylogenetic scales often results in different conclusions. Higher

taxonomic ranks like phyla and order tend to be more conserved across samples, resulting in a

larger proportion of microbial taxa being identified as part of the core. However, ecology and

function tend to be less conserved at these higher taxonomic ranks[24,25], making the core

microbiome less useful for informing host or ecosystem ecology. By contrast, at very low

taxonomic ranks, it is unclear whether variation between microbial taxa has significant

ecological or functional consequences. ASVs, for example, may only differ by one base pair in

the portion of the marker gene used for sequencing.[26] However, in microbial taxa with multiple

copies of the 16S rRNA marker gene, individual gene copies can vary intragenomically by

>1%.[27,28] Thus, ASVs may be much more diverse than the populations or functions that they are

being used to represent.[29] This can impact the core microbiome in one of two ways. First, if

multiple closely related ASVs are present in a threshold number of samples, they may all be

counted individually as part of the core, inflating core diversity. Second, if individual samples

each contain unique but closely related ASVs, none of the ASVs may occur in sufficient samples

to be included in the core, reducing core diversity. Perhaps because of the challenges associated

with defining core microbiomes at very high and very low taxonomic and/or phylogenetic scales,

most analyses of core microbiomes have used intermediate scales, for example 97% operational

taxonomic units (OTUs) or genera.[1]

The importance of taxonomic and/or phylogenetic scale, and the apparent variation in study

outcomes as a function of the taxonomic/phylogenetic scale considered are not unique to analysis

of core microbiomes.[1] Indeed, these same challenges occur in more general types of ecological

analyses of both microbial and macrobial systems, for example, calculation of α- and β-diversity

in and between ecological communities. This has led to the development of different phylogeny-

aware metrics including Faith's phylogenetic diversity[30] (PD), Rao's quadratic entropy[31] and

phylogenetic entropy[32] for α-diversity, and UniFrac distances[33] for β-diversity. Phylogeny-aware

metrics are advantageous because they integrate information across taxonomic ranks based on

the branch lengths in a phylogenetic tree. This removes the need to select an arbitrary taxonomic

rank or phylogenetic level for any particular ecological analysis. Further, because some

phylogenetic properties can serve as indicators of both functional diversity[34] and evolutionary

potential[35] (i.e., capacity to evolve in response to a perturbation, for example environmental

change[36–38]), phylogeny-aware metrics often provide more informative comparisons[39] of

properties like diversity than do analyses at any single taxonomic or phylogenetic scale.

Despite the many benefits of phylogeny-aware ecological analyses, few phylogeny-aware

methods for characterizing the core microbiome exist. Several recent papers have begun to

account for phylogeny when characterizing core microbiomes[40–42], for instance by defining core

taxa based on a series of clustering thresholds.[43] Nevertheless, there remains a lack metrics that

smoothly integrate phylogenetic information across taxonomic and phylogenetic scales. In this

paper, we develop a framework for analyzing core microbiomes using phylogenetic methods. To

do this, we first develop the concept of a 'core community phylogeny' - a phylogeny where

branches are defined based on being present in multiple samples from a single type of host or

environment. We then use this phylogeny to examine common measures of core communities,

including Venn diagrams, as well as metrics of within- and between-habitat diversity.

Core Community Phylogenies

All phylogeny-aware metrics of ecological communities, for example Faith's phylogenetic

diversity (PD) and UniFrac distance, are based on an underlying phylogenetic tree comprising

taxa present in the community. Logically, the analogous phylogenetic tree for core microbiomes

should comprise all microbial taxa that are conserved across microbiomes from a particular type

of habitat. However, for core microbiomes, there are two possible ways of defining such a tree: a

'tip-based' approach, and a 'branch-based' approach.

Tip-based phylogeny: One way to build a core community phylogeny is to first identify core

microbial taxa based on their relative abundance and/or occupancy across a threshold number of

microbiome samples from a particular type of habitat. Once these microbial taxa have been

identified, a phylogenetic tree can be reconstructed based on the sequences (or other traits) of the

conserved taxa. Notably, conserved microbial taxa can be identified at any pre-selected

taxonomic rank (e.g., phyla, genera, 97% OTUs, ASVs, etc.) provided that a phylogenetic tree

can be built connecting the various microbial lineages. The tip-based phylogeny is useful in that

it incorporates phylogeny while still identifying extant microbial taxa that comprise the core.

However, because the tip-based phylogeny identifies core taxa without reference to their

phylogenetic relationships, it does not capitalize fully on phylogenetic information.

Consequently, metrics calculated using the tip-based phylogeny are more susceptible to variation

across taxonomic rank and/or phylogenetic level as compared to metrics calculated using the

branch-based approach that we discuss next (see 'Examples' and 'Discussion' for more details).

Branch-based phylogeny: An alternative way to build a core community phylogeny is to first

build a phylogeny of all microbial taxa present in a set of microbiome samples from a particular

type of habitat. This phylogenetic tree can then be examined, branch-by-branch, to determine

which branches satisfy core criteria based occupancy and/or abundance thresholds (i.e., which

branches occur in sufficient numbers of samples or have sufficient abundance associated with

them). Branches that satisfy the core criteria can then be retained, while branches that do not can

be discarded. Note that, at least for a rooted tree, this results in discarding shallow branches

connected to deeper branches that have also been discarded, such that the remaining phylogeny

is fully connected (but see below for issues that arise with unrooted trees). Like the tip-based

phylogeny, the branch-based phylogeny can have leaf nodes at any taxonomic rank. However,

because the branch-based phylogeny integrates information on shared microbial lineages across

all depths of the phylogeny both for defining the core and calculating core metrics, there is less

reason to use high taxonomic levels (see 'Examples' below). Thus, we recommend ASVs as the

default choice.

Importantly, the tip-based phylogeny and the branch-based phylogeny for a given system will not

always be identical. Whereas the branch-based phylogeny will always have all of the branches

present in the tip-based phylogeny, the tip-based phylogeny will often be missing some of the

branches present in the branch-based phylogeny (see Figure 1). More specifically, the tip-based

phylogeny will be missing branches whenever a set of microbiome samples share closely related,

but not identical, taxa. In this case, none of the shallow branches will be considered core for

either the tip-based or the branch-based phylogeny. However, the deeper branches from which

all of the related tips are derived may be present on the threshold number of samples and thus

may be retained in the branch-based phylogeny. By contrast, these branches will be discarded in

the tip-based phylogeny, since definition of core is solely based on shared leaf nodes. This

fundamental difference is what makes the branch-based approach less sensitive to variations in

the precise taxonomic scale considered, at least when making relative comparisons between two

or more core communities (see 'Examples' below).

Figure 1. Examples of a tip-based phylogeny (A) and a branch-based phylogeny (B) for the
same microbial phylogeny. Branches of the microbial phylogeny that are not part of the core
community phylogeny are shown in black, while branches that are part of the core community
phylogeny are shown in red. For this example dataset, there are 6 samples (red, orange, yellow,
green, blue, and purple squares) and we use an occupancy threshold of 0.5 as our core criterion.
Thus being part of the core requires being present in 3 samples. Colored boxes at the end of each
terminal branch indicate the samples that a particular microbial taxon is present on. Stars are
used to mark branches that are included in the branch-based phylogeny but not in the tip-based
phylogeny.

Figure 1 shows the construction of both a tip-based phylogeny (A) and a branch-based

phylogeny (B) from the same set of microbiome samples and based on an occurrence threshold

of 0.5 (i.e., a taxon must occur in at least 50% of samples to be considered part of the core). In

Figure 1, the non-core branches of the microbial phylogeny are shown in black, while the core

branches are shown in red. Notice that there are differences between the tip-based phylogeny and

the branch-based phylogeny (see starred branches). In particular, the branch-based phylogeny

includes additional internal branches that are missing from the tip-based phylogeny. This occurs

because some clades are represented in 50% of samples, and thus are part of the branch-based

core microbiome, even though the individual taxa within the clades are not present in 50% of

samples.

Rooted versus Unrooted Trees: An important consideration in defining the core community

phylogeny is whether to assume that the root node is the most recent common ancestor (MRCA)

of all taxa in the sample. The root node should only be used as the MRCA when a rooted tree is

provided. Otherwise, the root node is arbitrary, and there is no reason to believe that the root

node is any more basal than the other nodes in the tree. Fortunately, microbial phylogenies

generated using pipelines like Qiime2[44] usually have the option for rooting trees (although this is

typically done through midpoint rooting, which involves multiple assumptions). Similar to

UniFrac, we strongly encourage using a rooted tree and forcing the root node to be the MRCA.

This is particularly true for branch-based core community phylogenies. The reason is as follows:

using the branch-based approach, branches are counted based on their inclusion in individual

phylogenies from each sample. When a root node is defined, these individual sample

phylogenies necessarily include the root. In absence of a root, however, individual sample

phylogenies only span the taxa present in each sample (core or otherwise). Consequently, basal

branches may not be counted if entire clades are differentially distributed across subsets of

samples. When this happens, distinct clades are not necessarily connected in the phylogeny that

emerges. To remedy this, we connect the final core community phylogeny based on the MRCAs

for all of the distinct clades. This, however, does not guarantee that branches will not be lost in

the process. For tip-based phylogenies, the choice to use an unrooted tree is less problematic

because the branches of the core community phylogeny are selected based on the occupancy of

microbial taxa (tip nodes), not on the occupancy of the branches themselves. Even for tip-based

trees, however, we recommend using a rooted tree and assuming that the root node is the MRCA.

Criteria for Inclusion: Another important consideration in defining the core community

phylogeny is the criterion used to determine whether a taxon or branch is part of the core. The

simplest option is to use an occupancy threshold. Core taxa could, for example be selected based

on being present in 50%, 90% or 100% of samples from a particular type of habitat.

Alternatively, various relative abundance thresholds could be used, either by themselves or in

combination with an occupancy threshold. Core taxa could, for instance, be selected based on

attaining a mean or minimum relative abundance across all samples or based on reaching a

maximum relative abundance in at least one sample. Whereas occupancy thresholds imply a

degree of conservation across samples, abundance thresholds indicate numerical dominance

(with the caveat that measured dominance may be different from true dominance due to biases in

extraction, PCR, etc.). When used in combination, occupancy and abundance thresholds allow

the core to be selected based on both conservation and numerical dominance.

While occupancy and abundance thresholds are the simplest forms of core criteria, several other

definitions have been developed as well. Recently, for instance, Shade and Stopnisek[20] have

pioneered a method that identifies core taxa based on being both strongly conserved within a

habitat and contributing to variation across that habitat. Fundamentally different from the

threshold approach, this more nuanced method can, in some cases, be more powerful than simple

occupancy and abundance thresholds. Importantly, identification of a core community phylogeny

does not depend on the use of any particular core criterion. Rather, core community phylogenies

can, at least theoretically, be defined based on any criterion that can be applied to individual

taxonomic units. For the tip-based tree, this means applying the chosen criterion when selecting

core taxa. For the branch-based tree, this means applying the chosen criterion on a branch-by-

branch basis.

Phylogenetic Metrics for Core Microbiomes

Once a core community phylogeny has been constructed, it can be used to perform phylogenetic

analyses of microbiome samples from different types of habitats. Here we consider several

common analysis methods, including Venn diagrams, a core microbiome equivalent of Faith's

phylogenetic diversity (α-diversity), and a core microbiome equivalent of UniFrac distance (β-

diversity).

Venn Diagrams: Venn diagrams are one of the most common methods for visualizing core

microbiomes across habitats. Non-phylogenetic Venn diagrams show counts of the core

microbial taxa shared by different combinations of habitat types. This can be extended to a

phylogenetic framework by replacing counts of shared microbial taxa with shared phylogenetic

branch lengths. Briefly, the core community phylogenies from different habitat types can be

compared, branch-by-branch, to determine which branches appear in which habitat

combinations. The total lengths of these shared branch segments can then be calculated and

reported either as an absolute value or as percentages of the total branch length spanning the core

community phylogenies of all habitats in the analysis.

Core Faith's Phylogenetic Diversity **(PD):** The phylogenetic equivalent to richness (a count of

microbial taxa present in a community), Faith's PD is the sum of the branch lengths of all

branches spanning the taxa present in a particular microbiome sample or ecological community.

This definition can easily be extended to core microbiomes by summing the branch lengths of all

branches in the core community phylogeny. Doing so gives a measure of how much

phylogenetic diversity is conserved across microbiome samples from a particular type of habitat.

*Core UniFrac Distances***:** The phylogenetic equivalent to Jaccard's index[45] (the fraction of

unique microbial taxa between two communities), the UniFrac distance[33,46] between two

communities is the sum of the branch lengths that are unique to each microbiome sample or

ecological community divided by the sum of the branch lengths of all branches spanning taxa

present in both microbiome samples or ecological communities. Again, this definition can be

extended to core microbiomes by dividing the branch lengths unique to each core community

phylogeny by the total branch lengths of the two core community phylogenies combined. Doing

so gives a measure of how phylogenetically distinct the core microbiomes are from two different

types of habitats.

Examples

Additional file 1: Simple Example System (see Figures S1.1 and S1.2) outlines the calculation of

Venn diagrams, core Faith's PD, and core UniFrac distances using a set of highly simplified core

community phylogenies. In this additional file, we show differences between tip- and branch-

based approaches, and outline calculations with both rooted and unrooted trees. Throughout the

Figure 2 *Staphylococcus* phylogenies for 10 skin microbiome samples from two individuals - Individual One (A) and Individual Two (B) - using ASVs (left) and OTUs (right). Circles at the end of each terminal branch indicate the number of samples that contained the corresponding ASV or OTU. These circles are colored based on assigned species, with grey being used when multiple species are contained within the same OTU. Black stars indicate taxa that would be part of each individual's core microbiome based on a non-phylogenetic approach using an occupancy threshold of 50%. Red lines indicate branches that are part of each individual's core microbiome based on a branch-based phylogenetic approach using the same occupancy threshold of 50%. Because the terminal branches on these trees are extremely short, trees are visualized using a square root transformation. This transformation was not used in calculating core microbiota.

remainder of the main text, we apply our methods to more realistic systems using rooted trees.

First, we consider the *Staphylococcus* component of the human skin microbiota based on a

dataset from Joglekar et al.[47] For this analysis, we use the branch-based phylogeny combined

with a simple occupancy threshold of 50% (i.e., a taxon/branch must be present in 50% of

samples to be considered part of the core, for alternate core criteria see Additional file 1:

Additional *Staphylococcus* Analyses, Figures S2.1-S2.3). Figure 2 shows *Staphylococcus* taxa

present in skin samples from two different people, first based on amplicon sequence variants

(ASVs) and then based on operational taxonomic units (OTUs, see 'Methods' for more details).

Core taxa that would be identified based on a non-phylogenetic approach are identified with

black stars. Core branches identified based on our phylogenetic approach are colored red. The

Staphylococcus microbiota on these two individuals demonstrate the challenges faced by non-

phylogenetic definitions of the core microbiome: whereas Individual One (Fig. 2A) has a lower

core *Staphylococcus* diversity based on ASVs (3 versus 6), Individual Two (Fig. 2B) has a lower

core *Staphylococcus* diversity based on OTUs (3 versus 4). Two separate phenomena explain this

diversity reversal. First, on Individual One, there are two ASVs from the CCE Clade (see Figure

2A) that do not, independently, reach an occupancy threshold of 50% and thus fail to contribute

to core ASV richness. When these ASVs are pooled into a single OTU, however, the OTU is

present in 50% of samples and thus is counted as part of the core. This leads to an increase in

diversity at the OTU scale and reflects the fact that strain variation across samples from

Individual One obscured the relatively high occupancy of closely related *Staphylococcus* in the

CCE clade. Second, on Individual Two, three closely related *S. capitis/caprae* strains and two

closely related *S. hominis* strains separately contribute to core ASV richness. These ASVs are,

however, pooled and counted as single core taxa at the OTU scale. This leads to a reduction in

OTU diversity and reflects the fact that most of the ASV diversity on Individual Two was a

result of shallow strain variation within the *S. capitis/caprae* and *S. hominis* clades. Unlike non-

phylogenetic measures of core *Staphylococcus* diversity, phylogenetic measures do not vary as

much across phylogenetic levels. Rather, phylogenetic measures suggest that the core microbiota

of the first person is more diverse at both the ASV (0.169 vs. 0.113) and OTU (0.180 vs. 0.105)

scales. The explanation is simple: the relatively fewer core ASVs on Individual One span a wider

phylogenetic breadth than the numerous but closely related ASVs on Individual Two (See

Additional file 1: Additional *Staphylococcus* Analyses for further comparisons across individuals

from the Joglekar et al.[47] study). Rank reversals in diversity estimates as a function of

phylogenetic scale can be problematic because they can make it difficult to interpret differences

between treatment groups. Using a non-phylogenetic approach, for instance, it might be difficult

to assess whether or not core *Staphylococcus* diversity increases resistance to pathogens like

methicillin-resistant *S. aureus* (MRSA).[48,49] This is because the people considered to have high

versus low core *Staphylococcus* diversity changes depending on the phylogenetic scale

considered. By contrast, the stability of our phylogenetic metrics across all phylogenetic scales

makes them more amenable to exploring relationships between core *Staphylococcus* diversity

and disease resistance.

Figure 3 shows a comparison of the core *Staphylococcus* microbiota across skin habitats from the

same Jonglekar et al.[47] study, along with turnover (β-diversity) in core *Staphylococcus*

composition among habitats. Again, differences in the phylogenetic and non-phylogenetic

approaches are apparent. Based on the non-phylogenetic Jaccard distance, for instance, the core

Staphylococcus microbiota of the nares is highly distinct from the core *Staphylococcus*

microbiota of all other skin habitats. This distinction, however, is much less obvious when using

a tip-based UniFrac distance and non-existent when using a branch-based UniFrac distance.

Indeed, the branch-based UniFrac distance suggests that the core *Staphylococcus* microbiota of

Figure 3 (A) Tip-based and (B) branch-based *Staphylococcus* phylogenies showing the shared and unique core community phylogenies, along with Jaccard distances (C), tip-based UniFrac distances (D) and branch-based UniFrac distances (E) among skin habitats. All panels in this figure used a 50% occupancy threshold as the core criterion. Because the terminal branches on these trees are extremely short, trees are visualized using a square root transformation. This transformation was not used in calculating core microbiota.

the nares is actually quite similar to the core *Staphylococcus* microbiota of both moist and

sebaceous sites. Inspection of the phylogenetic trees for the tip- and branch-based approaches

explains these differences. In particular, the nares are missing an *S. hominis* ASV found in the

core *Staphylococcus* microbiota all other skin habitats. The nares are not, however, entirely

lacking *S. hominis* ASVs. Rather, the nares are colonized by a range of other *S. hominis* ASVs,

though none individually reaches the threshold for core inclusion. Nevertheless, because multiple

S. hominis ASVs are present in nares samples, the branch-based approach counts most of the

diversity associated with the *S. hominis* clade towards the core microbiota of both the nares and

the other four habitat classes (sebaceous, moist, dry, feet). It is only the shallow tip branches that

do not contribute towards the core microbiota of the nares. By contrast, for the tip-based and

non-phylogenetic approaches, the missing *S. hominis* ASV drives significant turnover between

the nares and all other sites.

Figure 4. Venn diagrams for lizard gut (A-C) and soil (D-F) microbiota showing the percentage
of the total ASV counts shared by the core microbiota from each lizard species or soil type (A,
D) and the phylogenetic branch length shared by the core microbiota from each lizard species or
soil type using either tip-based (B, E) or branch-based (C, F) phylogenies. In all Venn diagrams,
lizard populations and soil types colored are as follows: *Aspidoscelis arizonae* (yellow), *A.*
neomexicanus (orange), *A. marmoratus* (red), bog soil (brown), permafrost soil (blue) and
seasonally thawed active layer soil (green).

Having demonstrated the benefits of our phylogenetic approach on a relatively simple and low

diversity *Staphylococcus* system, we next turn our attention to two additional datasets – one host-

associated and the other environmental – both of which contain significant diversity and multiple

bacterial phyla. Specifically, we consider the gut microbiota of three different whiptail lizard

(*Aspidoscelis*) species and the soil microbiota of three different soil types (see Materials and

Methods). Figure 4 shows standard Venn diagrams based on ASV counts (A, D) along with our

new phylogenetic Venn diagrams for tip-based (B, E) and branch-based (C, F) trees. For all

Venn diagrams, we define the core based on a 50% occupancy threshold (for alternate core

criteria, see Additional file 1: Additional Venn Diagrams). As compared to our phylogenetic

Venn diagrams, non-phylogenetic Venn diagrams (A, D) show the lowest percentage of system

diversity in the compartments representing the intersections of (A) gut microbiota of all three

lizard species (0%) and (D) microbiota of all three soil types (8.02%). Non-phylogenetic Venn

diagrams also show the highest percentage of system diversity in (A) the three compartments

unique to single lizard species (80.00%) and (D) the three compartments unique to individual

soil types (76.88%). This is not surprising. Using a non-phylogenetic approach, microbial taxa

from different host species/environments must be identical to be counted in intersection

compartments. This biases against finding shared microbial diversity, particularly when using

ASVs, because microbial lineages with small differences in 16S rRNA gene sequences (e.g., one

base pair substitutions) are deemed unique. By contrast, with a phylogenetic approach, even if

different host populations/environments have slightly different ASVs, this only contributes a

small amount of diversity to the unique compartments of the Venn diagram. Meanwhile, the

majority of the diversity is allotted to intersection compartments because the ASVs from the

different host populations/environments share most of their phylogenetic history. Comparing

branch-based and tip-based phylogenetic approaches, branch-based phylogenies show a higher

percentage of system diversity in the compartments representing intersection across (B, C) all

three lizard species (27.10% vs. 19.16%) and (E, F) soil types (29.41% vs. 27.14%). This reflects

the inclusion of additional, deep internal branches in branch-based phylogenies that are dropped

from tip-based phylogenies (see Figure 1). These basal branches are more likely to be conserved

across different host species/environments (i.e., phyla are more likely to be shared than ASVs),

inflating diversity in shared compartments of the Venn diagram (see Additional file 1: Additional

Venn Diagrams Figures S3.1-S3.4 for several additional datasets, including lizard skin and lake

water).

Figure 5 (A-C) shows core richness, and tip- and branch-based core Faith's PD as a function of

taxonomic rank for lizard gut microbiota based on an occupancy threshold (A), an occupancy

and abundance threshold (B) and the Shade and Stopnisek core criterion (C). For the sake of

comparison across metrics and taxonomic rank, we have plotted core diversity of two lizard

species (yellow: *A. arizonae*; red: *A. marmoratus*) relative to the third lizard species (*A.*

neomexicanus; see Materials and Methods). Figure 5 (E-G) shows similar plots for soil

microbiota, this time agglomerating taxa based on cophenetic distance, rather than taxonomy.

Again, for the sake of comparison, we have plotted core diversity of bog (brown) and active

layer (green) soils relative to permafrost soils. Figure 5 (D, H) shows the range of values

obtained for core richness and tip- and branch-based Faith's PD across all taxonomic ranks/tip

agglomerations considered. In most cases, phylogenetic measures of lizard gut and soil

microbiota show less variation across taxonomic rank/tip agglomeration as compared to non-

phylogenetic measures. Using an occupancy threshold of 50%, for instance, the richness of the

core gut microbiota of *A. marmoratus* (red) is anywhere from 0.64 (microbial species) to 2.75

(microbial ASV) times as diverse as the core gut microbiota of *A. neomexicanus*. By contrast, for

the same occupancy threshold, the branch-based core Faith's PD suggests a much narrower range

from 0.52 (microbial ASV) to 0.62 (microbial order). Likewise, using the Shade and Stopnisek

criterion, the richness of the core microbiota of active layer soil is anywhere from 0.92

(agglomeration: $h = 0.15$) to 1.5 (ASV) times as diverse as the core microbiota of permafrost

soil. Meanwhile, for the same Shade and Stopnisek criterion, the branch-based core Faith's PD

exhibit a much narrower range from 1.19 (agglomeration: $h = 0.15$) to 1.22 (ASV). Not all

differences in metric stability across taxonomic rank/tip agglomeration are as dramatic. Still, the

branch-based core Faith's PD typically shows the least variation across taxonomic ranks/tip

agglomeration, while the tip-based core Faith's PD shows the second least variation, and

richness shows the most variation. Nevertheless, there are deviations to this general trend,

particularly for the Shade and Stopnisek criterion (see Additional File 1: Additional α- and β-

diversity Analyses Figures S4.1 and S4.2 for several additional datasets, including lizard skin

and lake water).

429
430

Figure 5. Comparison of alpha diversity across taxonomic ranks (A-C) and degrees of tip agglomeration (E-G, see Materials and Methods) for the branch-based Core Faith's PD (solid, squares), the tip-based Core Faith's PD (dotted, triangles), and richness (dashed, circles) assuming the following core criteria: (A,E) an occupancy threshold of 50% (gold, red, dark brown, bright green), (B,F) an occupancy threshold of 50% and a mean abundance threshold of 0.1% (yellow, dark pink, medium brown, medium green), (C,G) the Shade and Stopnisek method (light yellow, light pink, light brown, light green). In (A-C), we show the core diversity of *Aspidoscelis arizonae* (gold, yellow, pale yellow) and *A. marmoratus* (red, dark pink, light pink) gut microbiota relative to the core diversity of *A. neomexicanus* gut microbiota. In (E-G) we show the core diversity of bog (dark, intermediate and light brown) and active layer (bright, intermediate and light green) soil microbiota relative to the core diversity of permafrost soil microbiota. Bar graphs (D,H) showing minimum and maximum values of alpha diversity across taxonomic rank for *A. arizonae* (gold, yellow, light yellow) and *A. marmoratus* (red, dark pink, light pink) relative to *A. neomexicanus*, and across tip agglomeration for bog soil (dark, intermediate and light brown) and active layer soil (bright, intermediate and light green) relative to permafrost soil. Diversity metrics are indicated as follows: 'BranchPD' is the branch-based phylogeny approach, 'TipPD' is the tip-based phylogeny approach, and 'Richness' is a count of taxa for each taxonomic rank. Core criteria are indicated as follows: 'occ' is the 50% occupancy threshold, 'occ-ab' is the 50% occupancy and 0.1% abundance threshold, and 'shade' is the Shade and Stopnisek method.

454
455
456

Figure 6. Comparison of beta diversity across taxonomic ranks (A-C) and degrees of tip agglomeration (E-G, see Materials and Methods) for the branch-based Core UniFrac distance (solid, squares), the tip-based Core UniFrac distance (dotted, triangles), and Jaccard distance (dashed, circles) assuming the following core criteria: (A,E) an occupancy threshold of 50% (gold, red, dark brown, bright green), (B,F) an occupancy threshold of 50% and a mean abundance threshold of 0.1% (yellow, dark pink, medium brown, medium green), (C,G) the Shade and Stopnisek method (light yellow, light pink, light brown, light green). In panels (A-C), we show turnover of *A. arizonae* (gold, yellow, light yellow) and *A. marmoratus* (red, dark pink, light pink) gut microbiota relative to *A. neomexicanus* gut microbiota. In panels (E-G) we show turnover of bog (dark, intermediate and light brown) and active layer (bright, intermediate and light green) soil microbiota relative to permafrost soil microbiota. Bar graphs (D,H) showing minimum and maximum values of beta diversity (turnover or distance) across taxonomic rank for *A. arizonae* (gold, yellow, light yellow) and *A. marmoratus* (red, dark pink and light pink) relative to *A. neomexicanus* and across tip agglomeration for bog soil (dark, intermediate and light brown) and active layer soil (bright, intermediate and light green) relative to permafrost soil. Diversity metrics are as follows: where 'BranchUniFrac' is the branch-based phylogeny approach, 'TipUniFrac' is the tip-based phylogeny approach, and 'Jaccard' is a count of taxa for each taxonomic rank/tip agglomeration. Core criteria are indicated as follows: 'occ' is the 50% occupancy threshold, 'occ-ab' is the 50% occupancy and 0.1% abundance threshold, and 'shade' is the Shade and Stopnisek method.

Figure 6 (A-C) shows the Jaccard, tip- and branch-based UniFrac distances between core gut

microbiota of *A. neomexicanus* and both *A. arizonae* (yellow) and *A. marmoratus* (red) based on

an occupancy threshold (A), an occupancy and abundance threshold (B) and the Shade and

Stopnisek core criteria (C). Figure 6 (E-G) shows similar plots for soil microbiota, this time

agglomerating taxa based on cophenetic distance, rather than taxonomy. Figure 6 (D, E) shows

the range of values obtained for Jaccard, tip- and branch-based UniFrac distances across all

taxonomic ranks/tip agglomerations considered. Broadly speaking, the branch-based approach

gives the smallest distances, as well as the least variation across taxonomic rank/tip

agglomeration. Meanwhile the tip-based approach gives intermediate distances and intermediate

variation and the non-phylogenetic approach (i.e., Jaccard distance) gives the largest distances

489 and the most variation (see Additional file 1: Additional α- and β-diversity Analyses Figures

490 S4.1 and S4.2 for several additional datasets, including lizard skin and lake water).

The holobiont R package

To facilitate the use of our new phylogenetic measures of core microbiomes, we developed a

new R package, holobiont, that includes functions for identifying the core community

phylogeny in any microbiome, drawing phylogenetic Venn diagrams, calculating the core Faith's

PD for a set of communities, and calculating the core UniFrac distance between two sets of

communities. Briefly, the coreEdges function takes a phyloseq object[50] and the core criterion

and identifies the branches of the core community phylogeny. The coreTree function takes the

same information and colors the branches of the core community phylogeny red, while leaving

the remaining branches of the community tree black. The corePhyloVenn function takes a

phyloseq object, a vector defining the habitat class for each microbiome sample, and the core

criterion and draws a phylogenetic Venn diagram of either branch lengths or percentages of

branch lengths shared by the core community phylogenies of different combinations of habitat

classes. The coreVennTree function takes the same input as the corePhyloVenn function and

colors the branches of the phylogenetic tree based on the combination of habitat class core

community phylogenies that they belong to. The coreFaithsPD function takes a phyloseq object

and the core criterion and calculates the core Faith's PD. Finally, the coreUniFrac function takes

a phyloseq object and a vector specifying which samples belong to which habitat class

(maximum 2) and calculates the UniFrac distance between the core community phylogenies. In

addition to the phylogenetic functions, the holobiont package also provides corresponding non-

phylogenetic functions, including coreTaxa, coreVenn, coreRichness and coreJaccard.

Additional details outlining the methods and use of our R package can be found on the CRAN

website at https://cran.r-project.org/web/packages/holobiont/index.html. A brief discussion of

our modified Shade and Stopnisek algorithm can be found in Additional file 1: Shade and

Stopnisek Algorithm.

Discussion

In this paper, we propose a framework for performing phylogeny-aware analyses of core

microbiomes. More specifically, we develop the concept of a core community phylogeny - a

phylogeny based on the microbial lineages that are core to a specific habitat class. We then

propose two different methods for selecting this tree - a tip-based approach and a branch-based

approach. For the tip-based approach, core membership is decided based on occupancy and/or

abundance of the leaf (tip) nodes. This is akin to the standard, non-phylogenetic method for

defining a core microbiome. However, once members of the core microbiome have been

identified, a phylogenetic tree is constructed. This allows for calculation of phylogeny-aware

metrics that leverage the relatedness of the leaf nodes to determine overall properties (e.g.,

diversity) of the core microbiome. For the branch-based approach, core membership is decided

independently for each branch of the phylogenetic tree. Therefore, in the branch-based approach,

phylogeny informs not only the calculation of metrics but also, the identification of the core

microbiome itself. The downside of the branch-based approach, however, is that the 'core

microbiome' often includes internal nodes while excluding their associated leaf nodes. Thus,

metrics calculated using the branch-based approach are not necessarily based on a set of extant

taxa. Rather, they are based on a set of complete and partial microbial lineages (see, for example,

Figure 2A, where branches leading to the CCE clade are included in the branch-based core

phylogeny, even though all terminal branches from this clade are excluded). Still, because the

branch-based approach uses phylogenetic information to both define the core microbiome and to

calculate core metrics, we suggest using the branch-based approach except when there is a strong

desire to identify specific microbial taxa within the core (e.g., to select for culturing

experiments).

As compared to existing metrics, the primary advantage of our phylogenetic metrics is that they

seamlessly integrate core microbiome properties (i.e., richness, turnover, shared taxa) across

taxonomic scales/phylogenetic levels. Thus, our phylogenetic metrics are more likely to provide

biologically relevant information. This is especially true when non-phylogenetic metrics are

calculated at taxonomic ranks that fail to capture meaningful biological variation.[51] If, for

example, the chosen taxonomic rank is too low, substantial biological variation may be identified

that has no functional or ecological consequences. Alternatively, if the chosen taxonomic rank is

too high, large amounts of meaningful biological variation may be pooled into the same

taxonomic unit. Depending on the system, the type of variation considered and the metric

calculated, it can be difficult to *a priori* determine (or even *a posteriori* assess) which taxonomic

ranks are more or less likely to be biologically informative. Further, in certain instances multiple

taxonomic ranks exhibit substantial meaningful biological variation. In this case, a non-

phylogenetic metric at any single taxonomic rank - even a well-chosen taxonomic rank - cannot

fully capture the underlying signal. Phylogenetic metrics circumvent these challenges by

summarizing the full spectrum of biological variation across all taxonomic ranks, with variation

at lower taxonomic ranks weighted less than variation at higher taxonomic ranks.

Another benefit of our branch-based approach (but not our tip-based approach) is that it

overcomes challenges associated with defining the core microbiome in the face of high levels of

strain variation. This is common in certain microbiome systems, including many host-associated

microbiomes. The pervasive human skin isolate, *Staphylococcus epidermidis*, for example,

evolves on individual humans over time,[52,53] resulting in strain level variation across the human

population. Another common human skin isolate, *Cutibacterium acnes*, exhibits even more

dramatic strain-level variation, showing differences in sequences across pores from a single

human individual.[54] Indeed, in-depth profiling of the human microbiome has shown that there

are numerous personalized microbial taxa that are found on only one or several individuals.[53,55]

Depending on the taxonomic rank at which the core microbiome is defined and the degree to

which any single microbial strain is shared among individual hosts, natural strain-level variation

can cause broadly shared microbial clades to be obscured from the non-phylogenetic core

microbiome. By contrast, if we apply our branch-based approach to these same scenarios, only

the terminal branches of the tree are omitted. Meanwhile, the more basal branches of the clades

are retained because they are shared across all strains. We see this in our simple *Staphylococcus*

example in Figure 2A, where the basal branches of the CCE clade are retained in the branch-

based approach despite none of the extant ASVs from this clade reaching the non-phylogenetic

threshold for core inclusion.

Because our phylogenetic metrics summarize microbiome measures across taxonomic

rank/phylogenetic level and because the definition of the core microbiome - at least when using

the branch-based approach - is less sensitive to strain-level variation, our phylogenetic metrics

tend to be more consistent than comparable non-phylogenetic metrics when calculated at

different taxonomic and/or phylogenetic scales. This is true both in a relative (i.e., comparison

between microbiome samples) and an absolute sense. To see why phylogenetic metrics are more

consistent, consider a habitat with hundreds of closely related core microbial strains within a

single phylum. Using a non-phylogenetic approach, this system will have a very high ASV α-

diversity (i.e., ASV richness), but a very low phylum α-diversity (i.e., phylum richness). By

contrast, using our phylogenetic approach, α-diversity (i.e., Faith's PD) will be low regardless of

whether the leaf nodes are defined at the ASV- or phylum-level. This is because, even when the

leaf nodes are ASVs, all of the different ASV strains contribute very short branch lengths.

Consequently, they do not substantially inflate α-diversity beyond the diversity of the single

phylum. The greater consistency of our phylogenetic metrics across taxonomic ranks means that

it is less important to choose the correct taxonomic rank, and that it is rarely disadvantageous to

use ASVs. This addresses one of the challenges that has been raised about interpreting core

microbiomes when outcomes are variable across taxonomic rank or phylogenetic level.[1]

While our phylogenetic metrics typically yield lower variation across taxonomic rank and/or

phylogenetic level, this is not always the case, particularly when using the Shade and Stopnisek

criterion. Exactly why phylogenetic metrics are more variable when combined with the Shade

and Stopnisek criterion is unclear. This may have to do with the emphasis that the Shade and

Stopnisek method places on microbiome variation rather than microbiome conservation.

Alternatively, it may be related to the stopping criterion used to delineate the Shade and

Stopnisek core. Notably, changing the stopping criterion can impact the relative variation of

phylogenetic and non-phylogenetic measures as a function of taxonomic rank (see Additional

File 1: Shade and Stopnisek Algorithm Figure S5.1). While our phylogenetic metrics may not

provide as much stabilization for the Shade and Stopnisek method, they still appear to perform as

well or better in most cases. In our calculations of core α- and β-diversity for lizard guts and

soils, for instance, our branch-based phylogenetic metric was more stable in 7/8 scenarios

(though the differences were marginal in 2/8 scenarios), and only less stable in 1/8 scenarios (see

Figures 5 and 6). Thus, even for the Shade and Stopnisek method, the branch-based approach

appears to be advantageous.

The primary goal of this paper is to introduce a framework for phylogenetic analysis of core

microbiomes. For this reason, we have focused on the concepts required to define a core

community phylogeny and have outlined the application of a core community phylogeny to three

types of microbiome analyses - Venn diagrams, α-diversity (Faith's PD), and β-diversity

(UniFrac). Our broad approach, however, is not limited to these particular types of analyses.

Indeed, additional analyses could be developed within the same framework. A straightforward

example would be to use UniFrac distances between core microbiomes to project into ordination

space (e.g., Principal Coordinate Analysis or Nonmetric Multidimensional Scaling). This would

provide phylogenetically informed visualization of the relatedness of different core microbiomes.

While initially developed for microbiome analysis, beyond microbiomes, our community

phylogenetics approach and associated R package may ultimately have wide application to

macrobial systems as well. Currently, very few macrobial studies attempt to identify core

communities (but see Murdock et al.[56]). This is likely because, at least historically, macrobial

systems have not been studied with many replicates. Indeed, even when replicate macrobial

communities are characterized, the effort required to identify all taxa in a single community

means that replicate numbers are low. This makes it difficult to identify a core community.

Further, in species poor macrobial systems where characterization of replicate communities is

possible, the core community concept is not as necessary or useful because the systems are

simple enough to analyze in their entirety. As sequencing techniques improve, however, and

more and more macrobial systems are characterized in a manner similar to microbiomes (e.g.,

through the use of eDNA[57,58]), it seems likely that core communities will emerge as a common

form of analysis for species rich macrobial systems (e.g., tropical rainforests or litter insects) as

well. There is no reason why our proposed framework cannot be applied to macrobial

communities. In fact, our framework can be applied to any system including prokaryotic and

eukaryotic communities, host-associated and free-living communities, and microbial (virus,

bacteria, fungi, etc.) and macrobial communities. It can also be applied to systems characterized

through a wide variety of methods, even including non-sequencing techniques. The only

requirements for our approach are tables of presences/absences of different taxa across different

communities, along with a phylogenetic tree comprising all taxa present across all communities.

Thus, occupancy data can be based on sequencing but can also be based on any other form of

observation, including transect or plot surveys. Likewise, phylogenies can be constructed using

sequence data, including sequencing of any kind (single marker genes, combinations of marker

genes, whole genomes), but can also be generated based on morphology or other traits

commonly used to build trees. While any phylogeny can be used for our phylogenetic analysis of

core communities, the validity of the analysis will depend on the accuracy of the phylogeny;

thus, care should be taken to select the most appropriate approach to tree building given the data

available. For 16S rRNA gene sequencing, Qiime2 provides guidance and pipelines for both

alignment-based and fragment-insertion methods of tree reconstruction.[44] Likewise, for shotgun

metagenomics sequencing, the Genome Taxonomy Database tookit (GTDB-Tk)[59,60] provides

different options for both *de novo* tree reconstruction and placement of genomes into a reference

tree.

The identification and characterization of core microbiomes has become a hallmark of

microbiome analysis. However, to date, most analyses of core microbiomes have focused on

taxonomy at specified taxonomic ranks. By contrast, phylogeny-aware metrics for characterizing

core microbiomes are largely lacking. Phylogeny-aware metrics have the benefit of integrating

information across taxonomic ranks. This can be particularly helpful when analysis at different

taxonomic ranks yields different outcomes - a common problem in ecology and evolution[61–65]

including the study of core microbiomes.[1] Our phylogenetic framework overcomes this

challenge. Indeed, even when our analysis is performed at different taxonomic ranks, it typically

yields more consistent results than non-phylogenetic methods. With the development of our new

framework and associated R package, we provide a mechanism to bring phylogeny-aware

community analyses to the study of core microbiomes. Ultimately, this will allow for novel

insight into the conserved components of microbial communities both within and between

different types of habitats.

Methods

Datasets

Skin *Staphylococcus* microbiota: We downloaded the ps_staph_ASV_filter_final.rds file from

https://github.com/skinmicrobiome/Joglekar_Staphylococcus_2023/tree/main/phyloseq_object.

We then loaded this file into R and used it to write out a FASTA file of representative sequences.

From the FASTA file, we used the `align-to-tree-mafft-fasttree` pipeline in Qiime2 to construct

a phylogenetic tree for all *Staphylococcus* ASVs based on their 16S rRNA sequences. The

FASTA file of representative sequences is available on our GitHub page. Next, we rarified

samples to a read depth of 500 using the `rarefy_even_depth` function from the `phyloseq`

package (version 1.50.0).[50] For our analysis of individuals, we randomly selected 10 skin

samples from all 9 individuals with 10+ skin samples available (recognizing that different body

sites were used for different people). We then compared core microbiota across individuals based

on ASVs and OTUs. For the latter, we agglomerated closely related taxa using the `tip_glom`

function from the `phyloseq` package (h = 0.5). For our analysis of skin habitats, we randomly

selected 20 skin samples from all 5 habitats with 20+ skin samples available (sebaceous, moist,

dry, feet, nares). We performed all analyses of the *Staphylococcus* dataset using an unscaled

phylogenetic tree. However, because many of the terminal branches were short, for the sake of

visualization, we plotted the *Staphylococcus* phylogenies using a square-root transformation.

<u>Lizard gut microbiota:</u> We sampled the microbiota from several whiptail lizard species (genus

Aspidoscelis, family *Teiidae*) at two locations (see Camper et al.[66] for more site details) within

Sevilleta National Wildlife Refuge. Each location represents a site of syntopy between hybrid *A.*

neomexicanus and a progenitor species, *A. marmoratus* (♀; a heterogenous grassland-mesquite

shrubland site: 34.397882°N, 106.867637°W, WGS84) and *A. arizonae* (♂, recently split from

A. inornatus[67]; a grassland site with sparse cholla cacti: 34.331178°N, 106.636717°W, WGS84).

Following taxonomic guidelines by Tucker et al.[68] and Walker et al.,[69] we refer to *Aspidoscelis*

spp. using the masculine epithet. We captured lizards using a rapidly deployable drift fence

trapping array lined with box funnel traps, further outlined in Camper et al.,[66] and by hand or

lizard noose when possible. Upon capture, we sampled the gut (=cloacal) and skin microbiota

(for analysis of the skin dataset, see Additional file 1) of each species following a swabbing

protocol in Camper et al.[22] We sampled each location until we captured at least 15 female lizards

of each whiptail species. All samples were sent to ZymoBIOMICS Targeted Sequencing Service

for Microbiota Analysis (Zymo Research) using 16S rRNA gene sequencing. Additional details

are available from https://www.zymoresearch.com/pages/16s-its-amplicon-sequencing and

Camper et al.[22] FASTQ files for lizard microbiomes, as well as environmental and sequencing

controls, are available at (these will be uploaded to NCBI upon publication).

We used FASTA files of representative sequences, supplied to us by ZymioBIOMICS Targeted

Sequencing Service (available on our GitHub page: https://github.com/bewicklab/CoreTrees/), to

create phylogenetic trees, as described above. We then combined these trees with the .biom files,

also supplied to us by the ZymoBIOMICS Targeted Sequencing Service (also available

at: https://github.com/bewicklab/CoreTrees/) for all downstream analysis. Details on the Zymo

pipeline can be found in our previous description of the dataset.[22] Prior to analysis, we removed

16S rRNA gene sequences that could not be identified as Bacteria or Archaea based on the

proprietary Zymo database. Additionally, we removed all samples with a read depth lower than

10,000 reads. Finally, we rarefied all samples to the read depth of the lowest sample. [49]

Soil dataset: We downloaded dataset 1036 from redbiom

(https://qiita.ucsd.edu/study/description/1036).[70] We then used the feature-table.qza file and

the all.seqs.fa files from the 130932 folder. From the FASTA file, we constructed a phylogeny,

again using the same Qiime2 pipeline that was used for the *Staphylococcus* dataset. We then read

719 the resulting feature table and phylogeny into R for analysis. For α-diversity, we rarefied

720 samples to the size of the sample with the lowest read depth. For β-diversity, we rarefied samples

721 to 5000 reads because of the computational time required by the tip_glom function for larger

722 datasets (see below).

724 Lake dataset: We downloaded dataset 1041 from redbiom

725 (https://qiita.ucsd.edu/study/description/1041). We then used the feature-table.qza file and the

726 reference-hit.seqs.fa files from the 131527 folder. From the FASTA file, we constructed a

727 phylogeny, again using the same Qiime2 pipeline that was used for the *Staphylococcus* dataset.

728 We then read the resulting feature table and phylogeny into R for analysis. For both α-and β-

729 diversity, we rarefied samples to the size of the sample with the lowest read depth. Analysis of

730 the lake dataset can be found in Additional file 1.

732 ***Dataset Analysis***

733 Core Trees: We used the coreEdges function from our holobiont package to identify branches

734 of the *Staphylococcus* phylogeny that were core to each individual. We then visualized the trees

735 using a modification of the plot_phylo function from the phyloseq package (included in the

736 code on our Github page). We used the coreVennTree function from our holobiont package to

737 identify and visualize branches of the *Staphylococcus* phylogeny that were shared by the core

738 microbiota of different skin habitats. For all figures in the main text, we used an occupancy

739 threshold of 0.5 as our core criterion.

<u>Venn Diagrams:</u> We used the `coreVenn` and `corePhyloVenn` functions from our `holobiont`

package to create non-phylogenetic and phylogenetic Venn Diagrams respectively. In all cases,

we reported percentages of the total branch length and used rooted phylogenies. All Venn

diagrams in the main text were calculated based on rooted trees for ASVs using an occupancy

threshold of 0.5 as our core criterion. Additional core criteria are included in Additional file 1:

Additional Venn Diagrams.

<u>α-diversity</u>: We used the `coreRichness` and `coreFaithsPD` functions from our `holobiont`

package to calculate non-phylogenetic and phylogenetic α-diversity respectively. To compare

across taxonomic ranks, we took the original ASV .biom file supplied by Zymo or .qza file

downloaded from redbiom to generate an ASV phyloseq object.[50] We then used the `tax_glom`

and `tip_glom` functions from the `phyloseq` package to generate OTU tables and phylogenies for

the lizard and soil/lake datasets respectively. For `tax_glom`, we considered taxonomic ranks from

species ('Rank7') through to order ('Rank4'). For `tip_glom` we considered cophenetic distances

from h = 0.05 to h = 0.2. This allowed us to calculate Faith's PD at different taxonomic

ranks/levels of tip agglomeration, as well as related non-phylogenetic richness. All Faith's PD

and richness calculations were performed on rooted trees using the following core criteria: (1) an

occupancy threshold of 0.5, (2) an occupancy threshold of 0.5 and a mean relative abundance

threshold of 0.1% and (3) a modified Shade and Stopnisek method (see Additional File 1: Shade

and Stopnisek Algorithm).

<u>β-diversity</u>: We used the `coreJaccard` and `coreUniFrac` functions from our `holobiont` package

to calculate non-phylogenetic and phylogenetic β-diversity respectively. Comparisons across

taxonomic ranks/tip agglomeration were performed as described for Faith's PD. Again, all

UniFrac and Jaccard calculations were performed on rooted trees using the following core

criteria: (1) an occupancy threshold of 0.5, (2) an occupancy threshold of 0.5 and a mean relative

abundance threshold of 0.1% and (3) a modified Shade and Stopnisek method.

All analyses were performed using the R programming language[71] (version 4.2.1) available on

our Github page at: https://github.com/bewicklab/CoreTrees

Declarations

Ethics approval and consent to participate

All research was approved by Clemson University under IACUC protocol numbers #2020-015

and #2021-047. We completed this work under the Sevilleta National Wildlife Refuge Special

Use Permit #SEV_Bewick_Camper_2022_59 and the New Mexico Department of Game and

Fish permit authorization #3772.

Consent for publication

not applicable

Funding

This study was funded by NSF awards #2105604 and #2025541, a Clemson University Support

for Early Exploration and Development (CUSEED) Grant, and the Clemson University Creative

Inquiry (CI) Program.

Availability of data and materials

All data and code (BIOM files, metadata files, and all R code) necessary for the analyses

presented in this manuscript are available at https://github.com/bewicklab/CoreTrees unless

otherwise noted in the main text. All data (BIOM and metadata files) are further included as a

zipped file as Additional file 2. The holobiont R package is publicly available through CRAN,

available at https://cran.r-project.org/web/packages/holobiont/index.html. We have uploaded all

raw sequence data (lizard microbiota and controls) to the National Center for Biotechnology

Information's SRA database

(https://dataview.ncbi.nlm.nih.gov/object/PRJNA1188019?reviewer=9hrl0g7nl4fmdqf8icgjglbes

2), and this information will be made publicly available upon acceptance.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

BTC and SAB conceived the idea and wrote the first draft of the manuscript. BTC performed all

related fieldwork. SAB developed the code and data analysis. All authors contributed

substantially to the content of the manuscript.

Acknowledgements

809 We thank Thomas Dempster, August Spencer, Eva Purcell, Riley Manuel, Drew Kanes and

810 Zachary Laughlin for their field assistance in New Mexico as well as Lily Margeson, Simon

811 Dunn, Georgianna Bellinger, Henry Egloff, Kaila Hodges, Camryn Lachica, and Savannah Utz

812 for their assistance assembling drift fence trapping arrays.

References

1. Neu AT, Allen EE, Roy K. Defining and quantifying the core microbiome: Challenges
and prospects. *Proceedings of the National Academy Sciences*. 2021;118(51):1-10.
doi:10.1073/pnas.2104429118/-/DCSupplemental
2. Custer GF, Gans M, van Diepen LTA, Dini-Andreote F, Buerkle CA. Comparative
Analysis of Core Microbiome Assignments: Implications for Ecological Synthesis.
mSystems. 2023;8(1):1-14. doi:10.1128/msystems.01066-22
3. Huse SM, Ye Y, Zhou Y, Fodor AA. A core human microbiome as viewed through 16S
rRNA sequence clusters. *PLoS One*. 2012;7(6):1-12. doi:10.1371/journal.pone.0034242
4. Zaura E, Keijser BJ, Huse SM, Crielaard W. Defining the healthy "core microbiome" of
oral microbial communities. *BMC Microbiol*. 2009;9(259):1-12. doi:10.1186/1471-2180-
9-259
5. Muletz Wolz CR, Yarwood SA, Campbell Grant EH, Fleischer RC, Lips KR. Effects of
host species and environment on the skin microbiome of Plethodontid salamanders.
Journal of Animal Ecology. 2018;87(2):341-353. doi:10.1111/1365-2656.12726
6. Turnbaugh PJ, Gordon JI. The core gut microbiome, energy balance and obesity. *Journal*
of Physiology. 2009;587(17):4153-4158. doi:10.1113/jphysiol.2009.174136
7. Pershina E V., Ivanova EA, Korvigo IO, et al. Investigation of the core microbiome in
main soil types from the East European plain. *Science of the Total Environment*.
2018;631-632:1421-1430. doi:10.1016/j.scitotenv.2018.03.136
8. Salzman S, Whitaker M, Pierce NE. Cycad-feeding insects share a core gut microbiome.
Biological Journal of the Linnean Society. 2018;123:728-738.
https://academic.oup.com/biolinnean/article/123/4/728/4920834
9. Henderson G, Cox F, Ganesh S, et al. Rumen microbial community composition varies
with diet and host, but a core microbiome is found across a wide geographical range. *Sci*
Rep. 2015;5:1-15. doi:10.1038/srep14567

10.   Hamady M, Knight R. Microbial community profiling for human microbiome projects:
843   Tools, techniques, and challenges. *Genome Res*. 2009;19(7):1141-1152.
844   doi:10.1101/gr.085464.108

845   11.   Hernandez-Teran A, Navarro-Diaz M, Benitez M, Lira R, Wegier A, Escalante AE. Host
846   genotype explains rhizospheric microbial community composition: The case of wild
847   cotton metapopulations (Gossypium hirsutum L.) in Mexico. *FEMS Microbiol Ecol*.
848   2020;96(8):1-11. doi:10.1093/femsec/fiaa109

849   12.   Fan K, Chu H, Eldridge DJ, et al. Soil biodiversity supports the delivery of multiple
850   ecosystem functions in urban greenspaces. *Nat Ecol Evol*. 2023;7:113-126.

851   13.   Sun L, Li G, Zhao J, Zhang T, Liu J, Zhang J. Core microbiota drive multi-functionality of
852   the soil microbiome in the Cinnamomum camphora coppice planting. *BMC Microbiol*.
853   2024;24(1). doi:10.1186/s12866-023-03170-8

854   14.   Björk JR, O'hara RB, Ribes M, Coma R, Montoya JM. The dynamic core microbiome:
855   structure, stability and resistance. *bioRxiv*. Published online 2017:1-50.
856   doi:10.1101/137885

857   15.   De Filippis F, La Storia A, Stellato G, Gatti M, Ercolini D. A selected core microbiome
858   drives the early stages of three popular Italian cheese manufactures. *PLoS One*.
859   2014;9(2):1-8. doi:10.1371/journal.pone.0089680

860   16.   Risely A. Applying the core microbiome to understand host–microbe systems. *Journal of
861   Animal Ecology*. 2020;89(7):1549-1558. doi:10.1111/1365-2656.13229

862   17.   Zaura E, Keijser BJ, Huse SM, Crielaard W. Defining the healthy "core microbiome" of
863   oral microbial communities. *BMC Microbiol*. 2009;9(259):1-12. doi:10.1186/1471-2180-
864   9-259

865   18.   Ganz HH, Jospin G, Rojas CA, et al. The Kitty Microbiome Project: Defining the Healthy
866   Fecal "Core Microbiome" in Pet Domestic Cats. *Vet Sci*. 2022;9(635):1-21.
867   doi:10.3390/vetsci9110635

868   19.   Holyoak GR, Premathilake HU, Lyman CC, et al. The healthy equine uterus harbors a
869   distinct core microbiome plus a rich and diverse microbiome that varies with geographical
870   location. *Sci Rep*. 2022;12(14790):1-14. doi:10.1038/s41598-022-18971-6

871   20.   Shade A, Stopnisek N. Abundance-occupancy distributions to prioritize plant core
872   microbiome membership. *Curr Opin Microbiol*. 2019;49:50-58.
873   doi:10.1016/j.mib.2019.09.008

874   21.   Shade A, Handelsman J. Beyond the Venn diagram: The hunt for a core microbiome.
875   *Environ Microbiol*. 2012;14(1):4-12. doi:10.1111/j.1462-2920.2011.02585.x

876   22.   Camper BT, Kanes AS, Laughlin ZT, Manuel RT, Bewick SA. Transgressive Hybrids as
877   Hopeful Holobionts. *bioRxiv*. Published online 2024:1-68.
878   doi:10.1101/2024.08.17.607925

879   23.   Astudillo-García C, Bell JJ, Webster NS, et al. Evaluating the core microbiota in complex
880   communities: A systematic investigation. *Environ Microbiol*. 2017;19(4):1450-1462.
881   doi:10.1111/1462-2920.13647

24.   Martiny JBH, Jones SE, Lennon JT, Martiny AC. Microbiomes in light of traits: A
phylogenetic perspective. *Science (1979)*. 2015;350(6261):1-8.
doi:10.1126/science.aac9323

25.   Martiny AC, Treseder K, Pusch G. Phylogenetic conservatism of functional traits in
microorganisms. *ISME Journal*. 2013;7(4):830-838. doi:10.1038/ismej.2012.160

26.   Callahan BJ, McMurdie PJ, Holmes SP. Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. *ISME Journal*.
2017;11(12):2639-2643. doi:10.1038/ismej.2017.119

27.   Větrovský T, Baldrian P. The Variability of the 16S rRNA Gene in Bacterial Genomes
and Its Consequences for Bacterial Community Analyses. *PLoS One*. 2013;8(2):1-10.
doi:10.1371/journal.pone.0057923

28.   Coenye T, Vandamme P. Intragenomic heterogeneity between multiple 16S ribosomal
RNA operons in sequenced bacterial genomes. *FEMS Microbiol Lett*. 2003;228(1):45-49.
doi:10.1016/S0378-1097(03)00717-1

29.   Schloss PD. Amplicon Sequence Variants Artificially Split Bacterial Genomes into
Separate Clusters. *mSphere*. 2021;6(4):1-6. doi:10.1128/msphere.00191-21

30.   Faith DP. Conservation evaluation and phylogenetic diversity. *Biol Conserv*. 1992;61:1-
10.

31.   Rao CR. Diversity and Dissimilarity Coefficients: A Unified Approach. *Theor Popul Biol*.
1982;21:24-43.

32.   Allen B, Kon M, Bar-Yam Y. A new phylogenetic diversity measure generalizing the
shannon index and its application to phyllostomid bats. *American Naturalist*.
2009;174(2):236-243. doi:10.1086/600101

33.   Lozupone C, Knight R. UniFrac: A new phylogenetic method for comparing microbial
communities. *Appl Environ Microbiol*. 2005;71(12):8228-8235.
doi:10.1128/AEM.71.12.8228-8235.2005

34.   Tucker CM, Davies TJ, Cadotte MW, Pearse WD. On the relationship between
phylogenetic diversity and trait diversity. *Ecology*. 2018;99(6):1473-1479.
doi:10.1002/ecy.2349

35.   Winter M, Devictor V, Schweiger O. Phylogenetic diversity and nature conservation:
Where are we? *Trends Ecol Evol*. 2013;28(4):199-204. doi:10.1016/j.tree.2012.10.015

36.   Mouquet N, Devictor V, Meynard CN, et al. Ecophylogenetics: Advances and
perspectives. *Biological Reviews*. 2012;87(4):769-785. doi:10.1111/j.1469-
185X.2012.00224.x

37.   Lankau R, Jørgensen PS, Harris DJ, Sih A. Incorporating evolutionary principles into
environmental management and policy. *Evol Appl*. 2011;4(2):315-325.
doi:10.1111/j.1752-4571.2010.00171.x

38.   Lavergne S, Mouquet N, Thuiller W, Ronce O. Biodiversity and climate change:
Integrating evolutionary and ecological responses of species and communities. *Annu Rev*
Ecol Evol Syst. 2010;41:321-350. doi:10.1146/annurev-ecolsys-102209-144628

39.   Lozupone C, Hamady M, Knight R. UniFrac - An online tool for comparing microbial
community diversity in a phylogenetic context. *BMC Bioinformatics*. 2006;7.
doi:10.1186/1471-2105-7-371

40.   Aguirre De Cárcer D. A conceptual framework for the phylogenetically constrained
assembly of microbial communities. *Microbiome*. 2019;7(1):1-11. doi:10.1186/s40168-
019-0754-y

41.   Talavera-Marcos S, Parras-Moltó M, Aguirre de Cárcer D. Leveraging phylogenetic
signal to unravel microbiome function and assembly rules. *Comput Struct Biotechnol J*.
2023;21:5165-5173. doi:10.1016/j.csbj.2023.10.039

42.   Parras-Moltó M, Aguirre de Cárcer D. Assessment of phylo-functional coherence along
the bacterial phylogeny and taxonomy. *Sci Rep*. 2021;11(1):1-6. doi:10.1038/s41598-021-
87909-1

43.   Aguirre de Cárcer D. The human gut pan-microbiome presents a compositional core
formed by discrete phylogenetic units. *Sci Rep*. 2018;8(1):1-8. doi:10.1038/s41598-018-
32221-8

44.   Estaki M, Jiang L, Bokulich NA, et al. QIIME 2 Enables Comprehensive End-to-End
Analysis of Diverse Microbiome Data and Comparative Studies with Publicly Available
Data. *Curr Protoc Bioinformatics*. 2020;70(1):1-46. doi:10.1002/cpbi.100

45.   Jaccard P. Distribution de la flore alpine dans le bassin des Dranses et dans quelques
régions voisines. *Bulletin de la Société vaudoise des sciences naturelles*. 1901;37:241-
272.

46.   Lozupone C, Lladser ME, Knights D, Stombaugh J, Knight R. UniFrac: An effective
distance metric for microbial community comparison. *ISME Journal*. 2011;5(2):169-172.
doi:10.1038/ismej.2010.133

47.   Joglekar P, Conlan S, Lee-Lin SQ, et al. Integrated genomic and functional analyses of
human skin–associated Staphylococcus reveal extensive inter- and intra-species diversity.
Proc Natl Acad Sci U S A. 2023;120(47):1-12. doi:10.1073/pnas.2310585120

48.   Yang JJ, Chang TW, Jiang Y, et al. Commensal staphylococcus aureus provokes
immunity to protect against skin infection of methicillin-resistant staphylococcus aureus.
Int J Mol Sci. 2018;19(5):1-14. doi:10.3390/ijms19051290

49.   Parlet CP, Brown MM, Horswill AR. Commensal Staphylococci Influence
Staphylococcus aureus Skin Colonization and Disease. *Trends Microbiol*. 2019;27(6):497-
507. https://www.sciencedirect.com/science/article/pii/S0966842X19300216

50.   McMurdie PJ, Holmes S. phyloseq: An R package for reproducible interactive analysis
and graphics of microbiome census data. *PLoS One*. 2013;8(4).

51.   Dubois A. Phylogeny, taxonomy and nomenclature: The problem of taxonomic categories
and of nomenclatural ranks. *Zootaxa*. 2007;(1519):27-68. doi:10.11646/zootaxa.1519.1.3

52.   Zhou W, Spoto M, Hardy R, et al. Host-Specific Evolutionary and Transmission
Dynamics Shape the Functional Diversification of Staphylococcus epidermidis in Human
Skin. *Cell*. 2020;180(3):454-470. doi:10.1016/j.cell.2020.01.006

53. Both A, Huang J, Qi M, et al. Distinct clonal lineages and within-host diversification
shape invasive staphylococcus epidermidis populations. *PLoS Pathog*. 2021;17(2):1-39.
doi:10.1371/JOURNAL.PPAT.1009304
54. Conwill A, Kuan AC, Damerla R, et al. Anatomy promotes neutral coexistence of strains
in the human skin microbiome. *Cell Host Microbe*. 2022;30(2):171-182.
doi:10.1016/j.chom.2021.12.007
55. Lloyd-Price J, Mahurkar A, Rahnavard G, et al. Strains, functions and dynamics in the
expanded Human Microbiome Project. *Nature*. 2017;550(7674):61-66.
doi:10.1038/nature23889
56. Murdock SA, Tunnicliffe V, Boschen-Rose RE, Juniper SK. Emergent "core
communities" of microbes, meiofauna and macrofauna at hydrothermal vents. *ISME
Communications*. 2021;1(27):1-13. doi:10.1038/s43705-021-00031-1
57. García-Machado E, Laporte M, Normandeau E, et al. Fish community shifts along a
strong fluvial environmental gradient revealed by eDNA metabarcoding. *Environmental
DNA*. 2022;4(1):117-134. doi:10.1002/edn3.221
58. Gelis ERE, Kamal MM, Subhan B, Bachtiar I, Sani LMI, Madduppa H. Environmental
biomonitoring of reef fish community structure with eDNA metabarcoding in the Coral
Triangle. *Environ Biol Fishes*. 2021;104(8):887-903. doi:10.1007/s10641-021-01118-3
59. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk v2: memory friendly
classification with the genome taxonomy database. *Bioinformatics*. 2022;38(23):5315-
5316. doi:10.1093/bioinformatics/btac672
60. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: A toolkit to classify
genomes with the genome taxonomy database. *Bioinformatics*. 2020;36(6):1925-1927.
doi:10.1093/bioinformatics/btz848
61. Cavender-Bares J, Kozak KH, Fine PVA, Kembel SW. The merging of community
ecology and phylogenetic biology. *Ecol Lett*. 2009;12(7):693-715. doi:10.1111/j.1461-
0248.2009.01314.x
62. Graham CH, Storch D, Machac A. Phylogenetic scale in ecology and evolution. *Global
Ecology and Biogeography*. 2018;27(2):175-187. doi:10.1111/geb.12686
63. Wanntorp HE, Brooks DR, Nilsson T, et al. Phylogenetic approaches in ecology. *Oikos*.
1990;57(1):119-132. https://www.jstor.org/stable/3565745
64. Mouquet N, Devictor V, Meynard CN, et al. Ecophylogenetics: Advances and
perspectives. *Biological Reviews*. 2012;87(4):769-785. doi:10.1111/j.1469-
185X.2012.00224.x
65. Hernández CE, Rodríguez-Serrano E, Avaria-Llautureo J, et al. Using phylogenetic
information and the comparative method to evaluate hypotheses in macroecology.
Methods Ecol Evol. 2013;4(5):401-415. doi:10.1111/2041-210X.12033
66. Camper BT, Laughlin ZT, Kanes AS, Manuel RT. A portable and wind resistant drift
fence array for arid environments. *bioRxiv*. Published online 2023:1-25.
doi:10.1101/2023.09.28.560034

67. Barley AJ, Reeder TW, Nieto-Montes De Oca A, Cole CJ, Thomson RC. A New Diploid
Parthenogenetic Whiptail Lizard from Sonora, Mexico, Is the "Missing Link" in the
Evolutionary Transition to Polyploidy. *Am Nat*. 2021;198(2):295-309.
doi:10.5061/dryad.2z34tmpm2
68. Tucker DB, Colli GR, Giugliano LG, et al. Methodological congruence in phylogenomic
analyses with morphological support for teiid lizards (Sauria: Teiidae). *Mol Phylogenet*
Evol. 2016;103:75-84.
69. Walker JM, Sullivan BK, Cordes JE. Sanctioned Nomenclature for Triploid
Parthenogenetic Colorado Checkered Whiptail and for Other Taxa of the Lizard Genus
Aspidoscelis (Family Teiidae). *Herpetol Rev*. 2021;52(3):550-558.
70. Hultman J, Waldrop MP, Mackelprang R, et al. Multi-omics of permafrost, active layer
and thermokarst bog soil microbiomes. *Nature*. 2015;521(7551):208-212.
doi:10.1038/nature14238
71. R Core Team. R: A language and environment for statistical computing. Published online
2021.

1017

1018

1019

1020

1021 **Electronic Supplementary Material**

1022

1023 *core_community_phylogeny_example.docx*

1024 Additional file 1: A simplified system demonstrating the construction of core community

1025 phylogenies and calculations of associated metrics.

1026

1027 *biom_files.zip*

1028 Additional file 2: BIOM files, read counts, taxonomy assignments, microbial phylogenies, and

1029 lizard metadata for gut and skin samples used in this study.