Male is the default sex: functional significance of the sex determination cascade in horned dung beetles

London C. Mitchell,[1] Armin P. Moczek,[1] Erica M. Nadolski[1]

Address: 1. Department of Biology, Indiana University, Bloomington, 915 East 3rd Street, Bloomington IN 47405, USA

Keywords: sexual dimorphism, Transformer, Intersex, Doublesex, *Digitonthophagus*

ORCID:
EMN: 0000-0003-3314-5305
APM: 0000-0002-3478-9949

Email:
London C. Mitchell: lonmitch@iu.edu
Armin P. Moczek: armin@iu.edu
Erica M. Nadolski: emnadols@iu.edu

1 Abstract

Sex-specific trait expression represents a striking dimension of morphological variation
within and across species. The mechanisms instructing sex-specific organ development
have been well studied in a small number of insect model systems, suggesting striking
conservation in some parts of the somatic sex determination pathway while hinting at
possible evolutionary lability in others. However, further resolution of this phenomenon
necessitates additional taxon sampling, particularly in groups in which sexual
dimorphisms have undergone significant elaboration and diversification. Here, we
functionally investigate the somatic sex determination pathway in the gazelle dung
beetle *Digitonthophagus gazella*, an emerging model system in the study of the
development and evolution of sexual dimorphisms. We find that RNA interference
(RNAi) targeting *transformer1 (tra1)* caused chromosomal females to develop
morphological traits largely indistinguishable from those normally only observed in
males, and that $tra1^{RNAi}$ is sufficient to induce splicing of the normally male-specific
isoform of *doublesex* in chromosomal females, while leaving males unaffected. Further,
intersex^{RNAi} was found to phenocopy previously described RNAi phenotypes of
doublesex in female but not male beetles. These findings match predictions derived
from models of the sex determination cascade as developed largely through studies in
Drosophila melanogaster. In contrast, *transformer2*^{RNAi} resulted in larval mortality and
was not sufficient to affect *doublesex* splicing, whereas RNAi targeting *Sex-lethal* and
two putative orthologs of *hermaphrodite* yielded no obvious phenotypic modifications in
either males or females, raising the possibility that the function of a subset of sex
determination genes may be derived in Diptera and thus non-representative of their
roles in other holometabolous orders. Our results help illuminate how the differential
evolutionary lability of the somatic sex determination pathway has contributed to the
extraordinary morphological diversification of sex-specific trait expression found in
nature.

Introduction

Sexual dimorphism, or the presence of trait differences between the sexes, is widespread across eukaryotes and represents one of the most striking dimensions of phenotypic variation both within and across species. Sexual dimorphisms are among the fastest evolving traits and as such constitute the trait class most often used to morphologically distinguish closely related species. Furthermore, because sexual dimorphisms may involve traits greatly elaborated in or even limited to one sex only, sex-specific trait expression has been hypothesized to be an important early stepping-stone in the initiation of novelty in development and evolution. Consequently, secondary sexual traits used as weapons during mate competition and the exaggerated ornaments critical for male displays and female choice have been traditional foci of functional ecology and evolutionary theory (Andersson & Simmons 2006). However, the mechanisms underlying their development have remained generally understudied, except in select model organisms with tools for genetic manipulation. While these investigations have been critical in advancing our understanding of the developmental genetic mechanisms underlying *e.g.* somatic sex determination, they also encounter limitations, especially in insects. Developmental genetic model systems such as the fruit fly *Drosophila melanogaster* and the red flour beetle *Tribolium castaneum* possess only relatively modest degrees of morphological sexual dimorphism generally not reflective of the extraordinary degrees of elaboration and diversification we see in nature, and their ecological or other significance may be difficult to discern in the wild. Furthermore, even within the small sampling of model taxa undertaken to date, a surprising diversity of mechanisms underlying sex-specific development have been documented, suggesting that investigation of additional taxa may be warranted (Hopkins and Kopp 2021). Therefore, here we investigate the molecular genetic basis of morphological sexual dimorphisms in a non-traditional insect model system with striking and highly diversified degrees of morphological sexual dimorphisms.

The mechanisms instructing sex-specific organ formation and/or elaboration have been particularly well studied in the fruit fly *Drosophila melanogaster*. In female flies, the so-called sex determination cascade begins with factors activated by the double X chromosome dosage, which result in the splicing of the active isoform of Sex-lethal (Sxl), which in turn results in the splicing of the active isoform of Transformer1 (Tra1, Salz & Erickson 2010). The resulting Tra1 protein, in combination with the constitutively spliced Transformer2 (Tra2) protein, then regulate the splicing of the female isoform of Doublesex (DsxF; Hoshijima *et al.* 1991). Along with two necessary cofactors, Intersex and Hermaphrodite, the female Doublesex protein then instructs

the development of female phenotypes (Li & Baker 1998, Garret-Engele *et al.* 2002). In male

flies, which possess XY sex chromosomes, Sex-lethal is instead spliced into an inactive form,

which results in the baseline splicing of an inactive form of Transformer1. Subsequently, only the

male Doublesex isoform (DsxM) is spliced, which does not require any co-factors to regulate

development of male phenotypes. In male (XY) flies, expression of *transformer2, intersex, and*

hermaphrodite can be detected, but functional analyses indicate that these genes are not

involved in specifying somatic sex in males (Figure 1A). Outside of flies, members of the

doublesex-mab3-related transcription factor (DMRT) gene family have been implicated across

holometabolous insect orders as a conserved genetic switch with sex-specific isoforms

regulating sexual differentiation and sexually dimorphic development throughout the organism

(silkworm, Ohbayashi *et al.* 2002; honeybee, Cho *et al.* 2007; jewel wasp, Beukeboom & van de

Zande 2010, red flour beetle, Shukla & Palli 2012a; for evolution of this mechanism and

Hemipteran data see Wexler *et al.* 2019).

Despite this deep conservation of DMRT genes acting as a genetic switch via the expression

of alternative, sex-specific isoforms, the pathway upstream of this switch appears to be much

more evolutionarily labile. Across arthropods, significant variation exists both at the

chromosomal level – with an XX-XY system in many flies and beetles, a ZW-ZZ system in

butterflies, and haplodiploidy in hymenopterans – and at the level of the cascade's first

molecular signal, for example via sex-dependent activation of the *Sex-lethal (Sxl)* locus in

Drosophila (Salz & Erickson 2010) or sex-specific expression of a piRNA, *feminizer,* in *Bombyx*

(Kiuchi *et al.* 2014). Yet despite the appreciable diversity in these upstream mechanisms, in all

species studied thus far the pathway then converges on the sex-specific splicing of *dsx*, which is

relayed throughout the body as a signal for sex-specific differentiation and growth (Verhulst and

van de Zande 2015). Likewise, *Transformer1*, the splicing factor directly upstream of *dsx,* has

also been demonstrated to be functionally conserved across all orders studied thus far

(Geuverink & Beukeboom 2014, Verhulst *et al.* 2010); however, the mechanisms by which its

transcription is activated and maintained has been found to differ across groups (Gempe *et al.*

2009, Verhulst *et al.* 2010a). Thus, data to date suggest a remarkable differential evolutionary

lability of the somatic sex determination cascade, characterized by striking conservation in some

parts and rapid evolution in others. However, further characterization of this phenomenon and

its contributions to organismal diversity will require additional taxon sampling, in particular in

groups in which sexual dimorphisms have undergone significant elaboration and diversification.

Here, we investigate the functional significance of an array of members of the somatic sex

determination cascade in the gazelle dung beetle, an emerging model system in the study of the
development and evolution of sexual dimorphisms.

Onthophagine beetles represent a powerful model system to investigate the molecular,
ontogenetic, and evolutionary underpinnings of sex-specific development due to the diversity of
experimentally accessible sexual dimorphisms within and among taxa (Ledon-Rettig *et al.* 2016,
Davidson *et al.* 2023). Onthophagine beetles possess an XX-XY sex chromosome system,
typical of most Coleoptera. Past functional genetic studies have confirmed the role of a single
ortholog of *doublesex* as a sex-specifically spliced transcription factor regulating sex-biased
development in the bull-headed dung beetle *Onthophagus taurus*, its close relative *O.*
sagittarius, and the more distantly related *Digitonthophagus gazella* (Figure 1B, see Kijimoto *et*
al. 2012, Casasa *et al.* 2020, Rohner *et al.* 2021). Additional studies have investigated
downstream targets of doublesex acting as trait-specific effector genes (Ledon-Rettig *et al.*
2017). However, the role and significance - if any - of other members of the sex determination
pathway in the regulation of sexual dimorphisms remain to be investigated. Here, dung beetles
offer a promising opportunity to investigate the means by which information about chromosomal
sex is transmitted to instruct the development of various degrees of sexual dimorphism. In this
study we focused on the role of five cardinal members of the sex determination pathway as
established in *D. melanogaster,* and how they may be regulating the development of sexually
dimorphic traits in the gazelle dung beetle *Digitonthophagus gazella*.

Specifically, we focused our efforts on six morphological traits exhibiting varying degrees of
sexual dimorphism in *D. gazella*: (i) paired posterior head horns present in males but entirely
absent in females (Fig 1B); (ii) paired prothoracic protrusions present in females but absent in
males (Fig 1B); (iii) foretibiae adapted for sex-specific behaviors (females possess a short, wide
foreleg with large tibial teeth used for subterranean tunneling, while the male foreleg is much
thinner, conspicuously elongated, and with short tibial teeth used during copulatory encounters,
Fig 2B); (iv) relative length of the pygidium (the sclerite covering the opening of the genital tract)
and the posterior-most abdominal sclerite (Fig S1); (v) the presence of obvious bilateral arched
cuticular grooves on the inside of the pygidium in females but not males (Fig S2); and (vi) the
sex-specific interior genital components: the male aedeagus and the female vagina and
receptaculum seminis or spermatheca (Fig S3). Together, these traits comprise a spectrum of
'degrees of sexual dimorphism' in the level of morphological differences discernible between
adult males and females. Focusing on these six traits we then sought to identify and functionally

characterize the role of key sex determination pathway genes upstream of *doublesex* (*Sxl, tra1,*
tra2, hermaphrodite, & *intersex).* Specifically, we sought to establish whether cardinal sex
determination pathway members characterized in other insect taxa play conserved or divergent
roles in horned dung beetles by acting as regulators or cofactors of *doublesex* using a
combination of functional genetic analyses and by directly investigating *doublesex* splicing in a
subset of our treatments.

Methods

Beetle husbandry

Digitonthophagus gazella individuals were collected near Barber County, Kansas and reared in
a lab colony as described previously (Moczek & Nagy 2005). Reproductively active adults were
transferred from the colony into a breeding container and allowed to reproduce. Eclosed larvae
were transferred from their natal brood balls into twelve-well plates and provided with organic
cow dung from Marble Hill Farm (Bloomington, IN) as described in Shafiei *et al.* (2001) and kept
in 16:8h light/dark cycle at 28°C pre- and post-injection treatments until eclosion. Eclosed adults
were sacrificed and preserved in 70% ethanol.

BLAST ortholog identification

We generated a custom BLASTP database for the *D. gazella* proteome (Davidson & Moczek
2024) using BLAST (Altschul *et al.* 1990). Query protein sequences of known *D. melanogaster*
sex determination genes were collected from Flybase (version FB2024_02, Öztürk-Çolak *et al.*
2024), and their predicted orthologs from the *Tribolium castaneum* and *O. taurus* genomes were
collected from OrthoDB (Kuznetsov *et al.* 2023). The query sequences were used to search the
custom *D. gazella* proteome database to identify target protein sequences for Sex-lethal,
Transformer1, Transformer2, Intersex, and Hermaphrodite. We established an e-value cutoff of
1×10^{-5} for selecting the best hit from each list of potential targets, and prioritized targets that
appeared as the top hit above the e-value cutoff for all queries.

Double-stranded RNA synthesis for RNA interference

Gene fragment constructs (Table S1) and fragment-specific primers (Table S2) were designed
using the *D. gazella* reference genome and ordered from Integrated DNA Technologies, Inc.
Fragments of each gene suitable as targets for RNA interference were chosen by using BLAST

to query 250bp portions of each gene against a custom *D. gazella* transcriptome database and

selecting those with zero off-target hits. Note that *transformer1* and *transformer2* are not

paralogs despite the naming convention. Synthesis of double-stranded RNA (dsRNA) for gene

knockdown via RNA interference was performed using a protocol optimized for coleopteran

larvae (Philip & Tomoyasu 2011). In brief, for each gene, the DNA template for dsRNA synthesis

was synthesized via PCR to add T7 RNA polymerase binding sequences flanking the gene

fragment. These constructs were purified using a Qiagen QIAquick PCR Purification kit. In vitro

transcription of dsRNA was performed using an Ambion MEGAscript T7 kit, and each target-

specific dsRNA product was purified using an Ambion MEGAclear kit and an ethanol

precipitation step.

Injection of constructs for RNA interference

dsRNA constructs were diluted to 1 ug/uL with injection buffer (Philip & Tomoyasu 2011). Prior to

injection, we sexed each larva to confirm chromosomal sex as described in Moczek and Nijhout

(2002). During the late second to third larval instar stage, chromosomal male larvae exhibit

increasingly prominent genital primordia visible underneath the cuticle in the ventrocaudal

abdomen. These tissues are absent in chromosomal females, allowing for unambiguous sexing

of each larva (Fig S4). A 3µl dose of dsRNA targeting a single gene was injected through the

abdominal cuticle into the hemolymph of sexed second and third-instar larvae using a Hamilton

brand syringe and small 32-gauge removable needle. Control individuals were randomly

selected from each round of developing larvae and injected with pure injection buffer. Previous

work in this system has shown that injection of either pure buffer or nonsense RNA can serve as

suitable controls for dsRNA injection, as neither buffer- nor nonsense RNA-injected adults show

any detectable phenotypic differences compared to wildtype adults (Moczek & Rose 2009,

Kijimoto *et al.* 2012).

Phenotype scoring and photography

We analyzed the phenotypes of RNAi and control individuals post eclosion. We specifically

focused our analysis on the following six sexually dimorphic traits in control and RNAi

individuals. First, we assessed (i) the presence or absence of head horns (normally only seen in

wildtype males), (ii) prothoracic protrusions (normally only seen in wildtype females), and (iii) the

shape and size of foretibiae (drastically elongated and thinner in males compared to females).

In addition, we assessed the morphology of the pygidial flap, a sexually dimorphic projection

attached to the terminal abdominal segment which covers access to the genital track.

Specifically, we evaluated the (iv) exterior of the pygidium and the closure it forms against the

subsequent abdominal sclerite, which is consistently sexually dimorphic across the

Onthophagine clade: males possess an elongated pygidium, resulting in a conspicuous medial

narrowing of the closure between pygidium and the neighboring abdominal sclerite, whereas

females display a consistent spacing between pygidium and neighboring sclerite. Additionally,

we evaluated the presence of (v) bilateral arched cuticular grooves on the interior of the

pygidium, which are present in females and nearly absent in males. Components of the (iv)

internal genitalia were also dissected from representative individuals of both sexes for each

sample group, following the terminology of Roggero *et al.* 2017; specifically, males were

examined for the presence of an aedeagus comprised of a proximal phallobase and distal

parameres ending in conspicuous paired projections, whereas females were examined for the

presence of a vagina and spermatheca. Representative individuals from each sample group

were photographed using a Leica MZ16 microscope with a PLANAPO 2.0x objective and a

PixeLINK PL-D7912CU-T camera; multiple photos of each sample were taken across different

planes of focus and overlaid using Adobe Photoshop.

Doublesex RT-PCR following tra1 & tra2 manipulation

To assess if *Dg-tra1*[RNAi] and *Dg-tra2*[RNAi] are sufficient to affect *doublesex* isoform splicing in *D.*

gazella, additional *Dg-tra1*[RNAi], *Dg-tra2*[RNAi], and control injected larvae were reared for RNA

extraction and RT-PCR targeting the *doublesex* coding region. Larvae were injected as

described above, and upon metamorphosis the head and thoracic region was dissected and

stored in TriZOL at −20°C. RNA extraction was performed using a Zymo Direct-zol RNA

Miniprep kit. Extracted RNA samples were used (i) to perform RT-PCR with primers targeting

the *doublesex* coding region (Table S2) using a ThermoFisher SuperScript TM IV One-Step RT-

PCR kit to assess isoform size, and (ii) sent for Sanger and/or Illumina amplicon sequencing

based on the number of bands per sample, to assess isoform sequence. Illumina amplicon

sequencing was performed after Nextera Small Genome DNA Library preparation on a NextSeq

2000 with a P2 2x100 flow cell. Sequencing data were pre-processed using FastQC (Andrews

2010) and Trimmomatic (Bolger *et al.* 2014), then using Trinity (Haas *et al.* 2013) to perform

genome-guided de-novo transcriptome assembly to assemble transcripts from sequencing

reads. The assembled sequences were then aligned to *O. taurus* and *O. sagittarius* doublesex

isoform sequences from Kijimoto *et al.* (2012) and manually annotated to determine predicted

exon boundaries.

219 Results

We sought to investigate the function of the cardinal sex determination genes *Sex-lethal,*
transformer1, transformer2, intersex, and *hermaphrodite* in the sexually dimorphic gazelle dung
beetle, *D. gazella,* through identification of target homologs, assessment of RNAi phenotypes
resulting from RNA interference targeting each gene, and examination of doublesex splicing
patterns following a subset of RNAi treatments. Below we discuss each of our findings in turn.

BLAST Ortholog Identification

We identified *D. gazella* target homologs for all five genes of interest using BLAST with queries
from *D. melanogaster, T. castaneum,* and *O. taurus.* For the Intersex, Sex-lethal, and
Transformer2 proteins, there was an unambiguous top hit matching to each query (Table S2).
For the Transformer1 protein, the top hits diverged across the queries, so an additional query
from *Trypoxylus dichotomus* (a closer relative than *D. melanogaster* or *T. castaneum*) was used,
and the top hit found to be matching both the *T. dichotomus* and *O. taurus* query was used as
the target gene (Table S2). For Hermaphrodite, which has not been experimentally validated
outside of the *Drosophila* genus, predicted homology of the hits to each query was far lower. We
therefore carefully examined those sequence hits appearing above our e-value cutoff for both *T.*
castaneum and *D. melanogaster* and chose two sequences with the highest conservation as
target genes to proceed with functional genetic characterization (Table S2), as detailed next for
each target gene.

Sex-lethal RNAi

RNA interference targeting *Dg-Sxl* resulted in no change in phenotype in any of the sexually
dimorphic body regions, nor any obvious morphological defects in monomorphic body regions.
Specifically, the head horn, prothorax, foreleg, and genital phenotypes of *Dg-Sxl*[RNAi] injected
individuals matched those of control individuals in both sexes. No other obvious morphological
defects were observed (Fig S5A, Table S3).

Transformer1 RNAi

Dg-tra1 RNA injection resulted in a dramatic degree of masculinization of the female head,
prothorax, and foreleg. Specifically, females injected with the *Dg-tra1*[RNAi] construct developed
prominent head horns similar to those of same-sized males, featured a smooth prothorax

missing bilateral protrusions otherwise typical of female morphology, and developed skinnier,

longer, male-like fore tibiae (Fig 2, Table S3). Additionally, the shape of the pygidium closure –

i.e. the abdominal sclerite covering the entrance to the genital tract – and the internal pygidial

cuticle morphology underwent significant masculinization in *Dg-tra1*^{RNAi} females; specifically, the

closure of the pygidium of RNAi females narrowed significantly in the medial abdomen,

matching control males (Fig S1B), and the internal pygidial cuticle grooves were conspicuously

smaller than those of control females (Fig S2B). Finally, typical female internal genitalia were

absent, but in contrast to the other body regions, masculinization in this body region was not

observed – i.e. no male-like internal genital features were found (Fig S3). In stark contrast, RNAi

targeting *Dg-tra1* in male larvae resulted in no change in the adult phenotypes of the head,

prothorax, foretibiae (Fig 2), closure of the pygidium or the internal pygidial cuticle (Fig S1B,

S2B); similarly, the internal genitalia of *Dg-tra1*^{RNAi} males matched those of control males

without exception (Fig S3).

Transformer2 RNAi

RNAi targeting *Dg-tra2* caused 93% lethality throughout the larval and pupal stages (Table S3).

Surviving adults showed no obvious phenotypic differences compared to control-injected or

wildtype individuals, in either the sexually dimorphic regions of interest or otherwise (head,

prothorax, foretibiae: Fig S5B, pygidium closure: Fig S1E, internal pygidium morphology: Fig

S2E). Note that the single surviving *Dg-tra2*^{RNAi} male was small-bodied, and could not be size

matched to other control and RNAi males; *D. gazella* male head horns and fore tibiae scale with

body size in a hyper-allometric manner, so the relatively small horns and short tibiae of this *Dg-*

tra2^{RNAi} male can most likely be accounted for solely by body size differences and therefore do

not represent an RNAi phenotype.

Intersex RNAi

Female individuals injected with the *Dg-ix*^{RNAi} construct exhibited a pair of small but obvious

head horns, reduced prothoracic protrusions, and moderately longer, thinner forelegs compared

to control-injected females (Fig 3, Table S3). In the posterior abdomen, the pygidium and inner

pygidial grooves displayed a slight phenotypic shift toward a more masculine shape (Fig S1C,

Fig S2C). Male *Dg-ix*^{RNAi} individuals matched controls across all phenotypes: head, prothorax,

foretibiae (Fig 3), pygidium closure (Fig S1C), internal pygidium morphology (Fig S2C), and

internal genitalia (Fig S3).

 Hermaphrodite RNAi

The first potential hermaphrodite ortholog tested, *Dg-jg1708*, caused 100% lethality in the larval

stage in both sexes (Table S3). In contrast, the second potential hermaphrodite ortholog tested,

Dg-jg4744, resulted in only 15% lethality (Table S3). However, the surviving adults displayed no

obvious phenotypic changes in the sexually dimorphic body regions or in other monomorphic

traits in males or females (pygidium closure: Fig S1F, internal pygidium morphology: Fig S2F,

head, prothorax, foretibiae: Fig S5C).

Regulation of doublesex splicing

To further characterize the functions of *Dg-tra1* and *Dg-tra2* in the dung beetle sex

determination pathway, we performed an RT-PCR experiment to assess if *Dg-tra1*^{RNAi} or *Dg-*

tra2^{RNAi} could individually affect *doublesex* isoform splicing. After injection of *Dg-tra1*^{RNAi} and *Dg-*

tra2^{RNAi} constructs, RNAi and control larvae were monitored until metamorphosis, and were then

utilized for RNA extraction and RT-PCR targeting all *doublesex* isoforms. Matching results from

earlier studies on the closely related *O. taurus* (Kijimoto *et al.* 2012), we found that control

males expressed a single, smaller *dsx* isoform, while females express multiple larger isoforms

(Fig 4, Table S4). *Dg-tra1*^{RNAi} females, in contrast, were found to lack the larger female isoforms

and instead, to express the smaller male isoform, whereas *Dg-tra1*^{RNAi} males showed the same

band as control males. In contrast to our results for *tra1*, RNAi targeting *tra2* did not appear to

affect *dsx* splicing in either females or males.

298 Discussion

In this study we sought to investigate the potential conservation of cardinal insect sex

determination factors in regulating sexual differentiation in the sexually dimorphic horned beetle

D. gazella. We performed single-gene RNAi treatments targeting male and female larvae to

determine phenotypic effects in the adults and followed with RT-PCR experiments for a subset

of genes to more directly assess regulatory links between *doublesex* and its potential direct

upstream regulators. Four salient results emerged.

Transformer1 RNAi masculinizes chromosomal females

RNA interference targeting *Dg-tra1* generated females that appeared morphologically nearly

indistinguishable from their chromosomal male counterparts. Head horn lengths of *Dg-tra1*^{RNAi}

females were in a similar size range of control and RNAi males. Similarly, the prominent

prothoracic protrusions normally observed in wildtype or control females were virtually

eliminated in *Dg-tra1*^RNAi^ females, leading to a simpler, rounded prothorax shape matching that

of typical male *D. gazella*. Likewise, foretibiae size and shape of *Dg-tra1*^RNAi^ females also

transformed to the longer, thinner form normally observed only in males. Additionally, the length

between the closure of the pygidium and the adjacent abdominal sclerite decreased

significantly, and the inner pygidial grooves normally present in females decreased substantially

in RNAi females, matching control male phenotypes. These results indicate that *Dg-tra1* is

necessary for the regulation of female development, in line with earlier findings in *D.*

melanogaster as well as more recent work in the red flour beetle *T. castaneum* (Shukla & Palli

2012) and the stag beetle *Dorcus rectus* (Gotoh *et al.* 2024). More generally, our results confirm

that *Dg-tra1* plays a conserved role in the sex determination pathway in horned beetles.

In partial contrast, internal female genitalia did not undergo complete masculinization after

Dg-tra1^RNAi^. While the normal female genital structures (vagina and spermatheca) were absent

from *Dg-tra1*^RNAi^ females as expected, no instances of ectopic male genitalia were observed.

We hypothesize that this may be due to major differences in the timing of growth and

differentiation of genitalia compared to our other focal traits. Specifically, the cells and tissues

that give rise to the adult head including horns, prothorax, foretibiae, and abdominal sclerites do

not initiate sexually dimorphic growth until the end of larval development following epidermal

apolysis from the larval cuticle and entry into the larval-to-pupal molting cycle (Moczek and

Nagy 2005). In contrast, the cells forming male and female internal genitalia begin proliferation

and morphogenetic arrangements as early as during the transition from the second to third

larval instar (S□vácha 1992; Moczek & Nijhout 2000). Thus, the temporal window during which

the signals necessary to alter sex-specific internal genital formation may have been already

closed prior to our mid-third larval instar RNAi treatment, leading to only an incomplete

masculinization of female genital primordia. This early specification of genital primordia relative

to other appendages appears to be widespread (stag beetles: Gotoh *et al.* 2024) and may also

explain the absence of RNAi phenotypes in other studies of sex determination.

Intersex RNAi phenocopies female but not male doublesex RNAi

Our intersex RNAi treatment reduced sex differences in sexually dimorphic regions in females,

but not males. Female *Dg-ix*^RNAi^ individuals exhibited a pair of small ectopic head horns,

reduced but still discernible prothoracic protrusions, and slightly masculinized forelegs (Fig 3).

Additionally, the distance between the closure of the pygidium and the adjacent abdominal
sclerite decreased, and the inner pygidial grooves decreased in size in *Dg-ix*[RNAi] females, all
indicating a morphological state *intermediate* between typical female and male phenotypes.
Collectively, Dg-ix[RNAi] treatment thus phenocopies the effects of *Dg-doublesex*[RNAi] in females,
but not in males (Fig 1B; also see Rohner *et al.* 2021), indicating that Intersex, along with
Transformer1, is necessary for the proper regulation of female but not male somatic sex
determination and sex-biased morphology. These findings also match the results of functional
analyses in the stag beetle *Cyclommatus metallifer* (Gotoh *et al.* 2016) and are concordant with
the logic of the sex determination pathway (Fig1A), which posits that the female Doublesex
protein requires Intersex as a cofactor, whereas the male Doublesex protein can function alone.

Evolutionary lability of the sex determination pathway in holometabolous insects

RNAi targeting the other putative sex determination genes examined in this study – *Dg-Sxl, Dg-*
tra2, Dg-jg1708, and *Dg-jg4474* – yielded no obvious phenotypic modifications in either males
or females, and regardless of whether sexually dimorphic or monomorphic body regions were
examined. These null results have a range of implications.

First, the absence of *Dg-Sxl*[RNAi] phenotypes matches results obtained in other taxa outside of
Dipterans (*Cyclommatus metallifer*, Gotoh *et al.* 2016; *Bombyx mori*, Niimi *et al.* 2006). This
supports the possibility that despite strong conservation of the Sxl coding sequence across
Holometabola, its function in Drosophilid sex determination is likely derived and divergent from
that executed in other Holometabola (Sanchez 2008).

In partial contrast, previous work outside of flies has indicated some divergence of the
functions of the Transformer2 protein. While in the fruit fly it functions primarily alongside Tra1 to
facilitate splicing of female *doublesex*, in other dipterans it also has been shown to function in
splicing *tra1* RNA (*Musca domestica*, Burghardt *et al.* 2005; *Lucilia cuprina*, Concha & Scott
2009; *Ceratitis capitata*, Salvemini *et al.* 2009; *Anastrepha suspensa*, Sarno *et al.* 2010). In the
honeybee *Apis mellifera*, Tra2 has also been shown to splice *doublesex* in addition to the
honeybee ortholog of *transformer1*, *feminizer.* However, in the honeybee Tra2 was additionally
documented to affect embryogenesis and resulted in lower embryonic viability (Nissen *et al.*
2012). Recent investigations of Transformer2 within beetles have yielded mixed results. In *T.*
castaneum (Tenebrionidae), *tra2* RNAi results in improper *doublesex* splicing and additionally
leads to larval lethality, suggesting that in both hymenopterans and coleopterans, Tra2 may be
executing critical functions in juvenile development independent of sex determination alone

(Shukla & Palli 2012). However, in *C. metallifer* (Lucanidae), larvae were able to survive tra2

RNAi administered very late in larval development, yet failed to display any phenotypic effects

(Gotoh *et al.* 2016). In this study we observed high levels of larval and pupal lethality following

Dg-tra2[RNAi] in *D. gazella* (Scarabaeidae), matching the lethality data obtained in the distantly

related *T. castaneum*, but also matching the lack of sex determination effects observed in the

more closely related *C. metallifer*. This raises the possibility that a putative juvenile viability

function of Tra2 may be ancestral in the Coleoptera and divergent in Lucanidae, while the

ancestral function in sex determination may have been lost in the superfamily Scarabaeoidea.

Future work targeting the role of Tra2 across a larger sampling of beetle families could resolve

these complexities, in addition to careful consideration of developmental timing during

treatments in order to target appropriate developmental windows.

Lastly, our investigation into potential homologs of the *Drosophila* sex determination gene

Hermaphrodite failed to yield conclusive results. Hermaphrodite homologs (with either

conserved sequence or function) have yet to be identified outside the *Drosophila* genus,

although the *D. melanogaster* protein is part of the large family of C2H2 zinc-finger transcription

factors. BLAST ortholog identification did not yield a single obvious homolog for *Dmel-her*, so

we chose to proceed with functional analysis of two *D. gazella* zinc-finger transcription factors,

Dg-jg1708 and *Dg-jg4474*. *Dg-jg1708*[RNAi] resulted in 100% mortality during the larval stage,

indicating a crucial role in juvenile development, but likely also indicating an absence of true

homology to *Dmel-her*. In contrast, *Dg-jg1708*[RNAi] individuals survived to adulthood, but

exhibited no obvious phenotypic effects, indicating again an absence of homology to *Dmel-her*.

Currently available data suggest that *hermaphrodite* may indeed be a Drosophilid-specific gene,

and that the female *Doublesex* isoform in horned beetles may only require *Intersex* as a

cofactor, rather than multiple interacting cofactor proteins. Future work tracing the evolution of

Dmel-her across Drosophilids and the entire Dipteran order could elucidate when the necessity

of Hermaphrodite for female sex determination evolved.

Transformer1-Doublesex splicing mechanism is conserved

Based on the work of Kijimoto *et al.* (2012), we predicted that control males would express a

single *dsx* isoform around 900 base pairs in length, while females would express multiple larger

isoforms ranging from 1300-1500 base pairs. Furthermore, based on the RNAi phenotypes

observed, we hypothesized that *tra1* RNAi would eliminate proper isoform splicing in females

but not males.

Our RT-PCR results confirmed that doublesex is spliced in a sex-specific manner in *D.*
gazella, with males expressing a single shorter isoform, and females expressing multiple longer
isoforms (Fig 4). Sequencing results indicated the male isoform to be 900 basepairs in length,
and the longest female isoform is predicted to be 1338 basepairs long. However, our
sequencing approach was not able to resolve the full sequence of the smaller female isoform
that appeared on the gel (Figure 4, Table S4). Concordant with our morphological findings, we
found that *Dg-tra1*[RNAi] females did not express bands typical of wildtype chromosomal females
and instead produced the single smaller isoform characteristic of chromosomal males,
suggesting that the widespread mechanism of female Doublesex splicing via Transformer1 is
conserved in *D. gazella*. As predicted, this effect was restricted to females and *Dg-tra1*[RNAi]
treatment did not affect isoform production in males (Fig 4). In contrast, *Dg-tra2*[RNAi] did not affect
splicing in either males or females, with each sex producing bands identical to their controls, in
contrast to earlier findings in *Tribolium* beetles (Fig4).

These findings further illuminate why and how the reduction of sex differences through the
extreme masculizination of females after *Dg-tra1*[RNAi] differs from the reduction of sex differences
following *doublesex* [RNAi], which instead results in the production of morphologies *intermediate* to
typical male and female phenotypes in both sexes: RNAi targeting *dsx* eliminates active
isoforms in both sexes, thereby eliminating the sex-specific instructions for the regulation of
growth and differentiation relevant for *both* male and female morphologies. In contrast, RNAi
targeting *tra1* eliminates a function necessary only for the regulatory cascade underlying
female-specific development, effectively re-routing the somatic sex differentiation pathway down
the alternative path of masculinization via the male *dsx* isoform.

Taken together, the RNAi phenotypes and the results of the RT-PCR experiment indicate that
transcriptional maleness is a default state in horned beetles, as the male *doublesex* isoform is
produced in the absence of any regulatory inputs from known members of the sex determination
cascade. Earlier work in *Tribolium* has posited the existence of a dominant male "M factor"
present on the Y chromosome in beetles that may act to suppress female sex determination
factors to allow male development, but data confirming the existence of such a factor and how it
may regulate or splice male *doublesex* have not yet materialized (Shukla & Palli 2012). Future
work may elucidate whether male *doublesex* is expressed and spliced in a similar manner to
'housekeeping' genes. Such a mechanism may be explained by the putative evolution of the
sex-specific splicing of *doublesex* in insects: in non-insect arthropods, *doublesex* produces only
a single isoform that primarily regulates masculinization of male tissues through both the

upregulation of masculinizing genes and the repression of feminizing genes (Kato *et al.* 2011).

Current research across arthropods suggests that the derived insect mechanism may have

evolved via a subdivision of ancestral functions (Kopp 2012); thus, it may be possible that the

regulation of the male isoform in insects is still achieved through a more ancestral, non-sex-

specific mechanism.

Conclusion

Here, we investigated the conservation of function of insect sex determination genes in the

sexually dimorphic horned beetle *D. gazella,* through single-gene RNAi experiments and an RT-

PCR experiment to assess evolutionary lability in the insect sex determination cascade. Our

results document that Transformer1 acts as a direct splicing regulator of the female isoforms of

doublesex in horned beetles (Fig 5). Additionally, the gene knockdown experiments confirmed

that Intersex is required for proper female development in horned beetles, and that the role of

Sex-lethal in sex determination is likely derived in Diptera. Finally, this work also suggests that

the role of Transformer2 in sex determination may be particularly evolutionarily labile, as it

appears to have maintained a function ensuring juvenile viability across Coleoptera but lost its

role in sexual differentiation in Scarabaeidae after divergence from Tenebrionidae. (Fig 5). One

standing question that remains regards the factors upstream of Transformer1 in beetles; the

regulatory cascade instructing Tra1 splicing is known in many dipterans and hymenopterans but

remains to be discovered in Coleoptera.

Taken together, this work provides evidence that transcriptional 'maleness' may be the default

state during beetle ontogeny in absence of other regulatory inputs, a finding that matches

results across Holometabola. An exciting open avenue of inquiry concerns the mechanisms that

regulate this default transcription of the male *doublesex* isoform, resulting in default

transcriptional maleness across Holometabola. In general, the lability of both sex determination

cascades on a molecular level and of sexually dimorphic traits on a morphological level

suggests that greater sampling of these phenomena across orders, and of their potential

regulatory links, may be poised to uncover potential important causal connections between the

two. Lastly, this work contributes to a growing number of studies hinting that redeployment of

deeply conserved regulatory pathways in a modular manner may be one mechanism by which

even rapidly evolving traits such as sexual dimorphisms can diversify independently in

development and evolution. Future work in this exciting open area may uncover the

mechanisms involved in re-deployment of the sex determination cascade in new cell types or at

new developmental timepoints.

Acknowledgements

We would like to thank Max Proctor for beetle collecting, Cale Whitworth for sharing his

expertise on the *Drosophila* sex determination pathway, Anna Macagno for her advice on

genitalia dissections, and the Indiana University Center for Genomics and Bioinformatics for

library preparation and sequencing. Earlier drafts of this manuscript benefited from the

comments of P. Davidson, R. Westwick, K. Givens, J. Jones, I. Manley, E. Pieri, and S. Kidd.

This work was supported in part through generous funding from the National Science

Foundation [Grant no. 2243725 and 1901680 to APM] and was performed while EMN was

funded by the National Institutes of Health [T32-HD049336]. Additional support was provided by

the Bloomington High School South Senior Internship Program to LCM.

References

1. Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J. (1990) "Basic local alignment search tool." J. Mol. Biol. 215:403-410.

2. Andersson, M., & Simmons, L. W. (2006). Sexual selection and mate choice. Trends in ecology & evolution, 21(6), 296-302.

3. Andrews S. (2010). FastQC: a quality control tool for high throughput sequence data. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc

4. Arbeitman MN, New FN, Fear JM, Howard TS, Dalton JE, Graze RM. 2016. Sex Differences in Drosophila Somatic Gene Expression: Variation and Regulation by doublesex. G3 Genes|Genomes|Genetics. 6(7):1799–1808. https://doi.org/10.1534/g3.116.027961

5. Bell LR, Horabin JI, Schedl P, Cline TW. 1991. Positive autoregulation of Sex-lethal by alternative splicing maintains the female determined state in Drosophila. Cell. 65(2):229–239. https://doi.org/10.1016/0092-8674(91)90157-T

6. Beukeboom LW, Van De Zande L. 2010. Genetics of sex determination in the haplodiploid wasp Nasonia vitripennis (Hymenoptera: Chalcidoidea). J Genet. 89(3):333–339. https://doi.org/10.1007/s12041-010-0045-7

7. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 30(15):2114–2120. https://doi.org/10.1093/bioinformatics/btu170

8. Bopp D, Saccone G, Beye M. 2013. Sex Determination in Insects: Variations on a Common Theme. Sexual Development. 8(1–3):20–28. https://doi.org/10.1159/000356458

9. Burghardt, G., Hediger, M., Siegenthaler, C., Moser, M., Dübendorfer, A., & Bopp, D. (2005). The transformer2 gene in Musca domestica is required for selecting and maintaining the female pathway of development. Development genes and evolution, 215, 165-176. https://doi.org/10.1007/s00427-004-0464-7

10. Burtis KC, Baker BS. 1989. Drosophila *doublesex* gene controls somatic sexual differentiation by producing alternatively spliced mRNAs encoding related sex-specific polypeptides. Cell. 56(6):997–1010. https://doi.org/10.1016/0092-8674(89)90633-8

11. Casasa S, Zattara EE, Moczek AP. 2020. Nutrition-responsive gene expression and the developmental evolution of insect polyphenism. Nature Ecology & Evolution. 4(7):970–978. https://doi.org/10.1038/s41559-020-1202-x

12. Cho S, Huang ZY, Zhang J. 2007. Sex-Specific Splicing of the Honeybee doublesex Gene Reveals 300 Million Years of Evolution at the Bottom of the Insect Sex-Determination Pathway. Genetics. 177(3):1733–1741. https://doi.org/10.1534/genetics.107.078980

13. Clough E, Jimenez E, Kim Y-A, Whitworth C, Neville MC, Hempel LU, Pavlou HJ, Chen Z-X, Sturgill D, Dale RK, et al. 2014. Sex- and Tissue-Specific Functions of Drosophila Doublesex Transcription Factor Target Genes. Developmental Cell. 31(6):761–773. https://doi.org/10.1016/j.devcel.2014.11.021

14. Concha C, Scott MJ. 2009. Sexual Development in Lucilia cuprina (Diptera, Calliphoridae) Is Controlled by the Transformer Gene. Genetics. 182(3):785–798. https://doi.org/10.1534/genetics.109.100982

15. Davidson PL, Moczek AP. 2024. Genome evolution and divergence in cis-regulatory architecture is associated with condition-responsive development in horned dung beetles. PLOS Genetics. 20(3):e1011165. https://doi.org/10.1371/journal.pgen.1011165

16. Davidson PL, Nadolski EM, Moczek AP. 2023. Gene regulatory networks underlying the development and evolution of plasticity in horned beetles. Current Opinion in Insect Science. 60:101114. https://doi.org/10.1016/j.cois.2023.101114

17. Garrett-Engele CM, Siegal ML, Manoli DS, Williams BC, Li H, Baker BS. 2002. intersex, a gene required for female sexual development in Drosophila, is expressed in both sexes and functions together with doublesex to regulate terminal differentiation. Development. 129(20):4661–4675. https://doi.org/10.1242/dev.129.20.4661

18. Gempe T, Hasselmann M, Schiøtt M, Hause G, Otte M, Beye M. 2009. Sex Determination in Honeybees: Two Separate Mechanisms Induce and Maintain the Female Pathway. PLOS Biology. 7(10):e1000222. https://doi.org/10.1371/journal.pbio.1000222

19. Geuverink E, Beukeboom LW. 2014. Phylogenetic Distribution and Evolutionary Dynamics of the Sex Determination Genes doublesex and transformer in Insects. SXD. 8(1–3):38–49. https://doi.org/10.1159/000357056

20. Gotoh H, Ohtsu I, Umino T, Yamasaki YY, Minakuchi Y, Ito T, Toyoda A, Kitano J. 2024. Induction of male-like mandibles in XX individuals of a stag beetle by gene knockdown of a feminizer gene transformer. Journal of Experimental Zoology Part B: Molecular and Developmental Evolution. https://doi.org/10.1002/jez.b.23274

21. Gotoh H, Zinna RA, Warren I, DeNieu M, Niimi T, Dworkin I, Emlen DJ, Miura T, Lavine LC. 2016. Identification and functional analyses of sex determination genes in the sexually dimorphic stag beetle Cyclommatus metallifer. BMC Genomics. 17(1):250. https://doi.org/10.1186/s12864-016-2522-8

22. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B, Lieber M, et al. 2013. De novo transcript sequence reconstruction from RNA-

seq using the Trinity platform for reference generation and analysis. Nat Protoc. 8(8):1494–1512. https://doi.org/10.1038/nprot.2013.084

23. Hopkins BR, Kopp A. 2021. Evolution of sexual development and sexual dimorphism in insects. Current Opinion in Genetics & Development. 69:129–139. https://doi.org/10.1016/j.gde.2021.02.011

24. Hoshijima K, Inoue K, Higuchi I, Sakamoto H, Shimura Y. 1991. Control of doublesex Alternative Splicing by transformer and transformer-2 in Drosophila. Science. 252(5007):833–836.

25. Ito Y, Niimi T. 2013. The role of doublesex in the evolution of exaggerated horns in the Japanese rhinoceros beetle. EMBO reports. 14(6):561–567. https://doi.org/10.1038/embor.2013.50

26. Kato Y, Kobayashi K, Watanabe H, Iguchi T. 2011. Environmental Sex Determination in the Branchiopod Crustacean Daphnia magna: Deep Conservation of a Doublesex Gene in the Sex-Determining Pathway. PLOS Genetics. 7(3):e1001345. https://doi.org/10.1371/journal.pgen.1001345

27. Kijimoto T, Moczek AP, Andrews J. 2012. Diversification of doublesex function underlies morph-, sex-, and species-specific development of beetle horns. PNAS. 109(50):20526–20531. https://doi.org/10.1073/pnas.1118589109

28. Kiuchi T, Koga H, Kawamoto M, Shoji K, Sakai H, Arai Y, Ishihara G, Kawaoka S, Sugano S, Shimada T, et al. 2014. A single female-specific piRNA is the primary determiner of sex in the silkworm. Nature. 509(7502):633–636. https://doi.org/10.1038/nature13315

29. Kopp A. 2012. Dmrt genes in the development and evolution of sexual dimorphism. Trends in Genetics. 28(4):175–184. https://doi.org/10.1016/j.tig.2012.02.002

30. Kuznetsov, Dmitry, Fredrik Tegenfeldt, Mosè Manni, Mathieu Seppey, Matthew Berkeley, Evgenia V Kriventseva, Evgeny M Zdobnov, OrthoDB v11: annotation of orthologs in the widest sampling of organismal diversity, Nucleic Acids Research, Volume 51, Issue D1, 6 January 2023, Pages D445–D451, https://doi.org/10.1093/nar/gkac998

31. Ledón-Rettig CC, Moczek AP. 2016. The transcriptomic basis of tissue- and nutrition-dependent sexual dimorphism in the beetle Onthophagus taurus. Ecology and Evolution. 6(6):1601–1613. https://doi.org/10.1002/ece3.1933

32. Ledón-Rettig CC, Zattara EE, Moczek AP. 2017. Asymmetric interactions between doublesex and tissue- and sex-specific target genes mediate sexual dimorphism in beetles. Nat Commun. 8(1):14593. https://doi.org/10.1038/ncomms14593

33. Li H, Baker BS. 1998. her, a gene required for sexual differentiation in Drosophila, encodes a zinc finger protein with characteristics of ZFY-like proteins and is expressed independently

of the sex determination hierarchy. Development. 125(2):225–235.
https://doi.org/10.1242/dev.125.2.225

34. Moczek AP, Nagy LM. 2005. Diverse developmental mechanisms contribute to different levels of diversity in horned beetles. Evolution & Development. 7(3):175–185. https://doi.org/10.1111/j.1525-142X.2005.05020.x

35. Moczek AP, Nijhout HF. 2002. A Method for Sexing Final Instar Larvae of the Genus Onthophagus Latreille (Coleoptera: Scarabaeidae). cole. 56(2):279–284. https://doi.org/10.1649/0010-065X(2002)056[0279:AMFSFI]2.0.CO;2

36. Moczek AP, Rose DJ. 2009. Differential recruitment of limb patterning genes during development and diversification of beetle horns. PNAS. 106(22):8992–8997. https://doi.org/10.1073/pnas.0809668106

37. Niimi T, Sahara K, Oshima H, Yasukochi Y, Ikeo K, Traut W. 2006. Molecular cloning and chromosomal localization of the Bombyx Sex-lethal gene. Genome. 49(3):263–268. https://doi.org/10.1139/g05-108

38. Nissen I, Müller M, Beye M. 2012. The Am-tra2 Gene Is an Essential Regulator of Female Splice Regulation at Two Levels of the Sex Determination Hierarchy of the Honeybee. Genetics. 192(3):1015–1026. https://doi.org/10.1534/genetics.112.143925

39. Ohbayashi F, Suzuki MG, Shimada T. 2002. Sex determination in Bombyx mori. Current Science. 83(4):466–471.

40. Öztürk-Çolak et al. (2024). FlyBase: updates to the Drosophila genes and genomes database. Genetics 227(1): iyad211.

41. Philip, Benjamin N., and Yoshinori Tomoyasu. (2011) "Gene knockdown analysis by double-stranded RNA injection." Molecular methods for evolutionary genetics: 471-497.

42. Pultz MA, Baker BS. 1995. The dual role of hermaphrodite in the Drosophila sex determination regulatory hierarchy. Development. 121(1):99–111. https://doi.org/10.1242/dev.121.1.99

43. Roggero A, Barbero E, Palestrini C. 2017. Revised classification and phylogeny of an Afrotropical species group based on molecular and morphological data, with the description of a new genus (Coleoptera: Scarabaeidae: Onthophagini). Org Divers Evol. 17(1):181–198. https://doi.org/10.1007/s13127-016-0297-z

44. Rohner PT, Linz DM, Moczek AP. 2021. Doublesex mediates species-, sex-, environment- and trait-specific exaggeration of size and shape. Proceedings of the Royal Society B: Biological Sciences. 288(1953):20210241. https://doi.org/10.1098/rspb.2021.0241

45. Salvemini M, Robertson M, Aronson B, Atkinson P, Polito C, Saccone G. 2009. Ceratitis capitata transformer-2 gene is required to establish and maintain the autoregulation of Cctra, the master gene for female sex determination. Int J Dev Bio. 53(1): 109-120. https://doi.org/10.1387/ijdb.082681ms

46. Salz H, Erickson JW. 2010. Sex determination in Drosophila: The view from the top. Fly. 4(1):60–70. https://doi.org/10.4161/fly.4.1.11277

47. Sanchez L. 2008. Sex-determining mechanisms in insects. Int J Dev Biol. 52(7):837–856. https://doi.org/10.1387/ijdb.072396ls

48. Shafiei M, Moczek AP, Nijhout HF. 2001. Food availability controls the onset of metamorphosis in the dung beetle Onthophagus taurus (Coleoptera: Scarabaeidae). Physiological Entomology. 26(2):173–180. https://doi.org/10.1046/j.1365-3032.2001.00231.x

49. Shukla JN, Palli SR. 2012a. Doublesex target genes in the red flour beetle, Tribolium castaneum. Sci Rep. 2(1):948. https://doi.org/10.1038/srep00948

50. Shukla JN, Palli SR. 2012b. Sex determination in beetles: Production of all male progeny by Parental RNAi knockdown of transformer. Sci Rep. 2(1):602. https://doi.org/10.1038/srep00602

51. Shukla JN, Palli SR. 2013. Tribolium castaneum Transformer-2 regulates sex determination and development in both males and females. Insect Biochemistry and Molecular Biology. 43(12):1125–1132. https://doi.org/10.1016/j.ibmb.2013.08.010

52. Shukla JN, Palli SR. 2014. Production of all female progeny: evidence for the presence of the male sex determination factor on the Y chromosome. Journal of Experimental Biology. 217(10):1653–1655. https://doi.org/10.1242/jeb.100438

53. S□vácha P. 1992. What are and what are not imaginal discs: Reevaluation of some basic concepts (insecta, holometabola). Developmental Biology. 154(1):101–117. https://doi.org/10.1016/0012-1606(92)90052-I

54. Verhulst EC, Beukeboom LW, van de Zande L. 2010a. Maternal Control of Haplodiploid Sex Determination in the Wasp Nasonia. Science. 328(5978):620–623. https://doi.org/10.1126/science.1185805

55. Verhulst EC, van de Zande L. 2015. Double nexus—Doublesex is the connecting element in sex determination. Briefings in Functional Genomics. 14(6):396–406. https://doi.org/10.1093/bfgp/elv005

56. Verhulst EC, van de Zande L, Beukeboom LW. 2010b. Insect sex determination: it all evolves around transformer. Current Opinion in Genetics & Development. 20(4):376–383. https://doi.org/10.1016/j.gde.2010.05.001

57. Wexler J, Delaney EK, Belles X, Schal C, Wada-Katsumata A, Amicucci MJ, Kopp A. 2019. Hemimetabolous insects elucidate the origin of sexual development via alternative splicing.Wittkopp PJ, Desplan C, Perry M, editors. eLife. 8:e47490. https://doi.org/10.7554/eLife.47490

Figure 1. Schematic of core sex determination cascade in *Drosophila melanogaster* and the conserved role of *doublesex* in *Digitonthophagus gazella*. (A) The female sex determination cascade is depicted on the left side of the diagram: animals with two X chromosomes regulate splicing of the active isoform of Sex-lethal, which splices the active isoform of Transformer1. This protein, in combination with Transformer2, regulates the splicing of the female isoform of Doublesex. Along with two required cofactors, Intersex and Hermaphrodite, the female Doublesex protein regulates development of female phenotypes. The male cascade is depicted on the right side of the diagram: in animals with XY sex chromosomes, Sex-lethal is spliced into an inactive form of the protein, which results in the splicing of only the inactive form of Transformer1. In turn, this results in the transcription and splicing of only the male Doublesex isoform, which alone regulates development of male phenotypes. In XY flies, expression of transformer2, intersex, and hermaphrodite can be detected, but functional analyses indicate that these genes are not involved in the male sex determination cascade. For information on the X-linked signal elements (XSEs) that are the primary links between X chromosome dosage and regulation of Sex-lethal splicing in *Drosophila*, see Salz & Erickson 2010. For information on the downstream targets of Doublesex isoforms, see Clough et al. 2014 (*Drosophila melanogaster*) and Ledón-Rettig et al. 2017 (*Onthophagus taurus*). (B) Wildtype adult male and female *Digitonthophagus gazella* display multiple novel sexually dimorphic traits: (i) male-specific paired, straight posterior head horns (black arrowhead, bottom left), (ii) female-specific paired, rounded prothoracic protrusions (white arrowhead, top left), and (iii) differences in tibiae shape and size with females possessing short, wide forelegs woth large, wide tibial teeth and males displaying much longer, more slender tibiae with small, more rounded tibial teeth (not shown). RNAi targeting all *doublesex* isoforms eliminates these sex differences, generating beetles with intermediate phenotypes by (i) inducing horn formation in females but decreasing horn size in males, (ii) inducing prothoracic protrusions in males but decreasing their size in females, and (iii) decreasing foretibia length in males and increasing it in females (not pictured, see Rohner et al 2021). These *dsx*[RNAi] phenotypes in D. gazella and other studies (Kijimoto et al 2012) establish conservation of its function as a key sex-determination factor in horned beetles. Scale bars = 1mm.

Figure 2. Effects of *transformer1* RNAi on adult *D. gazella.* Representative animals obtained after control injections (black labels) and *tra1* dsRNA injections (red labels) showing (A) head (horns highlighted with black arrowheads) and prothorax (protrusions highlighted with white arrowheads) and (B) fore tibiae. *Dg-tra1*[RNAi] did not affect adult male traits, but in females substantially reduced prothoracic protrusions and induced conspicuous, paired ectopic head horns (panel A upper right; black arrowhead). Likewise, the length and teeth size of female foretibiae transformed to resemble the longer, thinner morphology normally only observed in males (panel B second from top). Scale bars = 1mm.

Figure 3. Effects of *intersex* RNAi on adult *D. gazella*. Representative animals obtained after control injections (black labels) and *ix* dsRNA injections (red labels) showing (A) head (horns highlighted with black arrowheads) and prothorax (protrusions highlighted with white arrowheads) and (B) fore tibiae. *Dg-ix*^RNAi did not affect adult male traits, but in females the treatment moderately reduced prothoracic protrusions, induced small ectopic head horns, and modestly masculinized the foretibiae by transforming their size and shape toward an elongated and thinner morphology, intermediate between control males and females. Scale bars = 1 mm.

Figure 4. Expression and inferred structure of *D. gazella doublesex* isoforms. (A) RT-PCR results from left to right: 1kb ladder, control-injected female bands ~1300-1350 basepairs, control-injected male band of ~900bp, *Dg-tra1*^{RNAi} female band of 900bp, *Dg-tra1*^{RNAi} male band of 900bp, *Dg-tra2*^{RNAi} female bands of ~1300-1350bp, *Dg-tra2*^{RNAi} male band of ~900bp. These results confirm (i) sex-specific splicing pattern of *D. gazella doublesex* in control males and females, and (ii) the role of Transformer1 in simultaneously promoting the splicing of female – while preventing the splicing the male isoform – of *dsx: Dg-tra* ^{RNAi} females produced the male isoform band while female isoform bands are absent. RT-PCR primer pairs correspond to those shown in B. **(B)** Diagram of *doublesex* isoform structures in *D. gazella:* the single shorter male isoform (M) and the longer female isoform (F) sequences obtained from sequenced cDNA are indicated by rectangles, with inferred ORFs shaded either blue (male) or pink (female). Stop codons are indicated by red lines, with putative 3' untranslated regions shown in light pink (after Kijimoto et al 2012). The horizontal grey bars indicate regions used as primers for RT-PCR.

Figure 5. Model of the sex determination cascade in *D. gazella*. This study and past work (Rohner *et al.* 2021) have established partial conservation and partial divergence of the sex determination cascade in horned beetles compared to other holometabolous insects. The female sex determination cascade is depicted on the left side of the diagram, and the male cascade is depicted on the right. Data to date indicate that *Sex-lethal* is conserved in the beetle genome but does not function in sex determination. In female beetles, Transformer1 acts as a direct splicing regulator of female *doublesex* isoforms, and DsxF requires Intersex as a cofactor to regulate proper female development. Transformer2 was found to be necessary for survival through the larval and pupal stages in both males and females but did not affect *doublesex* splicing. Finally, a Hermaphrodite ortholog was not found in horned beetles. In male beetles, the male doublesex isoform can regulate male development without any known cofactors. At present, the regulatory factors upstream of Transformer1 in beetles are unknown.