Combined stem cell and predictive models reveal flavin cofactors as targets in metabolic liver
dysfunction

Julian Weihs[&,1], Fatima Baldo[&,2,3,4], Alessandra Cardinali[&,2,5], Gehad Youssef[&,2,3,4], Katarzyna Ludwik[6], Nils
Haep[7,8], Peter Tang[7], Pavitra Kumar[9], Cornelius Engelmann[8,9], Susanna Quach[1], Mijuna Meindl[1], Martin
Kucklick[10,11], Susanne Engelmann[10,11], Bruno Chillian[12], Michael Rothe[13], David Meierhofer[14], Isabella
Lurje[9,15], Linda Hammerich[9], Prakash Ramachandran[16], Timothy J. Kendall[16], Jonathan A. Fallowfield[16],
Harald Stachelscheid[6], Igor Sauer[7], Frank Tacke[9], Philip Bufler[1], Christian Hudert[1], Namshik Han[2,3,4*], Milad
Rezvani[1,8,17,18, 19*]

1. Charité Universitätsmedizin Berlin, Department of Pediatrics, Division of Gastroenterology,
Nephrology and Metabolic Medicine, Augustenburger Platz 1, 13353 Berlin
2. Milner Therapeutics Institute, University of Cambridge
3. Cambridge Centre for AI in Medicine, University of Cambridge
4. Cambridge Stem Cell Institute, University of Cambridge
5. LifeArc, Lynton House, 7-12 Tavistock Square, London, WC1H 9LT
6. Berlin Institute of Health (BIH) at Charité – Universitätsmedizin Berlin, BIH Core Unit pluripotent
Stem Cells and Organoids (CUSCO), Berlin, Germany
7. Department of Surgery, Campus Virchow-Klinikum, Experimental Surgery, CCM|CVK, Charité –
Universitätsmedizin Berlin, Augustenburger Platz 1, 13353 Berlin, Germany
8. Clinician Scientist Program, Berlin Institute of Health at Charité (BIH), Anna-Louisa-Karsch-Str. 2,
10178 Berlin, Germany
9. Department of Hepatology and Gastroenterology, Charité, Universitätsmedizin Berlin, Campus
Virchow-Klinikum (CVK) and Campus Charité Mitte (CCM), Berlin, Germany
10. Institute for Microbiology, Technische Universität Braunschweig, Braunschweig, Germany
11. Microbial Proteomics, Helmholtz Center for Infection Research, Braunschweig, Germany
12. TRI Thinking Research Instruments GmbH
13. Lipidomix, 13125 Berlin, Germany.
14. Max Planck Institute for Molecular Genetics, Ihnestraße 63-73, 14195 Berlin, Germany
15. Department of General, Visceral and Transplantation Surgery, Heidelberg University Hospital,
Heidelberg, Germany
16. Centre for Inflammation Research, Institute for Regeneration and Repair, University of Edinburgh,
Edinburgh, UK
17. Berlin Institute of Health (BIH), 10178 Berlin, Germany
18. Berlin Institute of Health (BIH) at Charité-Universitätsmedizin Berlin, BIH Center for Regenerative
Therapies (BCRT), Berlin, Germany
19. Division of Gastroenterology, Hepatology and Nutrition, Cincinnati Children's Hospital Medical
Center, Cincinnati, OH 45229, USA

[&]These authors contributed equally
*Corresponding authors, milad.rezvani@charite.de; nh417@cam.ac.uk

Drug discovery for multifactorial diseases like metabolic dysfunction-associated steatotic liver
disease (MASLD) remains challenging due to inadequate models and untargeted drug screenings.
We combined stem-cell-based modeling with computational drug predictions identifying flavin
pathways as therapeutic targets in MASLD. For disease stage-specific discovery, we established a
MASLD testing model, compounding metabolic triggers to intensify mitochondrial crisis. *In vitro*
injuries included adipo- and myokines, immune cell co-culture, and genomic risk factors.
Benchmarking experiments revealed similarities with advanced human MASLD. To query therapeutic
compounds, protein-protein-interaction networks, weighted gene co-expression, and knowledge
graph-based analyses independently predicted flavin adenine dinucleotide (FAD) as an anti-MASLD
factor. Dysregulated flavoproteomes *in vitro* and *in vivo*–in pediatric and adult MASLD patients–
supported our flavin network-focused strategy. We established therapeutic FAD concentrations to
mitigate metabolic injury and fibro-inflammation in human multicellular liver organoids and other
assays. We enhanced therapeutic FAD effects through genetic mitochondrial biogenic augmentation
and identified orally available flavo-active compounds—including Aspirin—restoring mitochondrial
respiration. Our study demonstrates how integrating stem cell-derived disease modeling with
computed drug predictions can expedite therapeutic discovery.

Disease modeling of multifactorial diseases remains challenging, and the target discovery process is
inefficient[1]. For example, cancer cells are suitable for large-scale drug screenings but lack physiological

relevance. In contrast, induced pluripotent stem cells (iPSCs) can generate versatile disease models that
clarify pharmacological targets in difficult-to-culture cell types[2]. However, the limited scalability of iPSC-
based systems poses challenges to effectively screen large compound libraries, which themselves have
restricted read-out dimensions. Computationally enhanced drug discovery aims to overcome these
limitations by prioritizing compounds for experimental validation. Machine learning (ML)-based algorithms
harness clinical and molecular data, to predict therapeutic efficacy of drug compounds, enabling more
focused candidate lists. However, the potential of ML- and iPSC-based systems, along with integrating
clinical cohort data to expedite experimental validation feedback loops, has not been fully realized.
Metabolic dysfunction-associated steatotic liver disease (MASLD) is a prime example of a multifactorial
disease with high unmet need. It affects ~30% of the global population[3] and is projected to increase[4]. In
MASLD, nutrient overload, extrahepatic signaling, and genetic susceptibility (e.g. in *PNPLA3*) cause hepatic
steatosis and lipotoxic cell stress. Hepatocyte injury activates both innate and adaptive immunity[5],
progressively leading to fibrogenesis, potentially developing to cirrhosis and liver failure. Although the
disease sequence is understood[6], the multiple injury mechanisms hamper drug discovery. MASLD models
typically expose hepatocytes to free fatty acids (FFAs), emphasizing early-stage steatosis[7]. While relevant,
this method mainly tests anti-steatotic drugs, overlooking subsequent pathological drivers like metabolic
crisis and mitochondrial dysfunction. Additionally, MASLD is systemic, involving signals from adipose or
muscle tissue, not captured by FFAs. Notably, adipokines like resistin (Res)[8] and myokines, including
myostatin (Myo)[9–11], have been linked to MASLD. However, their precise impact on human hepatocytes
remains unclear. Thus, incorporating various MASLD triggers may reveal relevant pathological features *in*
vitro for drug discovery, while tailoring drug predictions to disease-relevant pathologies enhances the
likelihood of successful therapeutic development.
Here, we integrate multi-modal human stem cell-based modeling with ML-based drug predictions and
extensive benchmarking against clinical cohorts to identify therapeutic targets for a pressing need in
hepatology – MASLD drug development. We establish a drug testing framework using iPSC-derived
hepatocytes (iHeps) exposed to a compounded sequence of MASLD-related triggers. Furthermore, we
utilize three independent computational drug prediction pipelines that analyze clinical data tailored to
MASLD hallmark pathologies, as observed *in vitro*. By integrating these efforts and other pipeline
innovations, including e.g. label-free iHep phenotyping by ML-based morphological imaging analysis, we
identified the therapeutic potential of targeting flavin network biology for MASLD. Our functional validation
confirmed the therapeutic potential of restoring multicellular flavin homeostasis, and we established a
previously unknown functional connection to a well-known pharmaceutical, namely Aspirin.
92
Results:
<u>Establishing a human iPSC-derived MASLD drug testing model capturing the multi-hit disease pathology</u>
We established an integrated *in vitro* and clinically stratified *in vivo* data-based and computed drug
discovery pipeline to identify anti-MASLD compounds **(Fig 1A)**. First, we derived a human iHep-based
MASLD model suitable for drug testing. Conventional models rely on a single injury, like FFA exposure[12,13].
We cumulatively exposed iHeps to FFAs (oleic and palmitic acid (OA/PA)), inter-organ signaling molecules
(Res/Myo), and inflammatory signaling via indirect leukocyte co-culture (peripheral blood mononuclear cells
(PBMCs)) **(Fig. 1B)**. To consider genomic risk modifiers, we incorporated iHeps with and without
homozygous risk mutations for the *PNPLA3*-I148M variant. To contextualize all cellular injuries, we derived
an overall metric based on key MASLD pathologies **(Fig. 1B & Supp. Fig. 1A)**. Each additional injury
compounded effects in both cell lines, albeit certain metabolic tipping points stood out. FFAs affected
steatosis but did not induce major transcriptomic shifts or other functional hallmarks in *PNPLA3*-wildtype
locus-iHeps (*PNPLA3*-I148I) **(Fig. 1B & Supp. Fig. 1B)**. Yet, *PNPLA3*-I148M iHeps displayed higher
susceptibility to lipotoxic stress alone and a higher steatotic propensity **(Supp. Fig. 1A & 2A&B)**. Res/Myo
addition led to the first metabolic crisis, impairing mitochondrial complex gene expression, loss of
mitochondrial membrane potential ($\Delta\Psi m$) and significant down-regulation of metabolic or up-regulation of
cytoskeleton pathways **(Supp. Fig. 1C, Supp. Table 1)**. Notably, without additional triggers, Res and Myo
individually significantly reduced $\Delta\Psi m$ in iHeps and when combined, mitochondrial spare respiratory
capacity **(Supp. Fig. 1D, Fig. 1C)**. Because gut-derived inflammation and immune cell trafficking is well
established in MASLD[14,15], we incorporated endotoxin-primed leukocytes. This marked another key
transition, characterized by the secretion of TNFα and IL-1β, maximum release of the hepatocyte injury
marker aspartate aminotransferase (AST), and maximum metabolic dysregulation, including further
downregulation of metabolic pathways and upregulation of inflammatory pathways **(Fig. 1B, Supp. Fig. 1C,**
Supp. Table 1). Mass Spec analysis of the cellular secretome confirmed elevated MASLD-related markers
such as DPP4[16] or heat shock proteins, indicating cellular stress **(Supp. Fig. 1E, Supp. Table 2)**. PathfindR
analysis revealed donor-specific disease trigger responses: *PNPLA3*-I148M iHeps displayed perturbations
in mRNA processing-related pathways, whereas *PNPLA3*-I148I iHeps displayed mitochondrial-predominant

impairment **(Supp. Fig. 2C)**. Shared pathways included DNA damage and apoptosis **(Supp. Table 3)**.
Importantly, benchmarking the iHep transcriptomes against clinical samples using Gene Set Enrichment
Analysis revealed similar clustering of the Res/Myo and leukocyte incorporating models with the F4 disease
stage **(Fig. 1D)**. Shared pathways included enhanced wound healing and diminished metabolic and
mitochondrial processes. Thus, we established a multifactorial MASLD model capable of mimicking
advanced pathophysiology by combining multiple factors, resulting in defined metabolic tipping points,
such as mitochondrial dysfunction.
127
<u>Computational discovery of FAD as a MASLD-modifying candidate via multimodal network analysis</u>
To identify candidate compounds modifying MASLD progression, we developed a computational drug
repurposing pipeline that integrates three complementary network-based approaches: (1) Protein–protein
interaction (PPI)-based network proximity analysis grounded in early-stage transcriptomics, (2) fibrosis-
resolved weighted gene co-expression network analysis (WGCNA) to capture disease progression signals,
and (3) knowledge graph (KG) inference using PrimeKG to uncover systems-level relationships linking drugs
and hepatic phenotypes **(Fig. 2A-C)**. Each method independently leverages network biology principles, but
their integration, anchored in a shared biological hypothesis around mitochondrial dysfunction, adds both
depth and rigor to candidate prioritization. We tailored this unified strategy to identify high-confidence
therapeutic candidates for testing in our human iPSC-derived MASLD model. For both PPI and WGCNA,
we used transcriptomic data from the Govaere et al. (2020)[17] cohort defining disease-relevant gene
signatures and co-expression modules. For the PPI approach specifically, we refined these signatures using
MitoCarta 3.0[18] focusing on mitochondrial processes. For the KG approach, we used disease phenotypes
as entry points into PrimeKG, a large-scale, multimodal KG. Together, these three layers offer distinct yet
convergent views on MASLD biology, enabling a systems-level prediction pipeline that links clinical data,
network theory, and therapeutic inference. The following subsections describe all pipelines in detail.
PPI network-based drug proximity analysis: Differential expression analysis of liver bulk RNA-seq data
identified 1,927 genes dysregulated in MASL vs. control (**Fig. 3A, Supp. Fig. 3A&B**). To refine for
hepatocyte specificity, we filtered the upregulated genes using four single-cell and single-nucleus liver
RNA-seq datasets from mouse and human, retaining only genes consistently elevated in hepatocytes (**Fig.**
3B). This yielded a robust hepatocyte-specific MASLD signature, with strong classification performance to
effectively separate MASL from controls (**Fig. 3C, Supp. Fig. 3C**). We built a PPI network from this
signature using STRING and identified topological hubs using centrality metrics. We refined them through
permutation testing, yielding 21 high-confidence nodes that were both central and transcriptionally
upregulated (**Fig. 3D**). These genes formed a compact subnetwork that distinguished MASL from control
samples (**Fig. 3E**) and were significantly enriched for the KEGG pathway "Non-alcoholic fatty liver disease"
(FDR < 0.05; **Supp. Fig. 3D**). We performed network-based drug proximity analysis to rank FDA-approved
compounds based on their topological closeness to the 21-node disease module within the human
interactome. This identified 34 compounds with significant proximity, listed in **Supp. Table 4**, including
several that modulate mitochondrial and metabolic regulators disrupted early in MASLD (**Fig. 3F**).
WGCNA-guided mitochondrial module expansion and drug repurposing: WGCNA enables the
identification of transcriptionally coordinated gene modules associated with disease phenotypes. We
applied WGCNA to the Govaere et al. dataset[17] to detect modules correlated with fibrosis severity and
extract key targets for therapeutic inference. WGCNA resolved 19 co-expression modules, of which six
showed significant correlation with fibrosis progression **(Fig. 4A)**. These modules showed enrichment for
biological processes implicated in MASLD. Notably, ME4 was enriched in extracellular matrix genes,
aligning with fibrotic remodelling **(Supp. Fig. 3F)**. We prioritized ME8, enriched for mitochondrial translation
and oxidative phosphorylation (OxPhos), due to its relevance to early MASLD and our MASLD drug testing
model **(Fig. 4B)**.

A

1. *In vitro*–clinical integration for drug prediction

In vitro

In vivo (F1-F4)

Mitochondrial dysfunction

Mitochondria-tailored drug repurposing predictions

FAD

2. Linking Flavo-disruption to clinical outcomes

Pediatric cohort (Berlin)

Adult SteatoSITE cohort (Scotland)

Flavo-processes

Norleucine
Aspirin
3P

FAD

Mitochondrial function ↑

Fibro-inflammation↓
Hepatocyte damage↓

Steatosis↓
Ω-3/Ω-6 PUFAs↑
Label-free rescue

Leukocytes Organoids Hepatocytes

Flavin-manipulation

4. Flavo-active compounds rescue mitochondria

3. Therapeutic *in vitro* flavo-pathway manipulation

B

+OA/PA +Res/Myo +PBMCs

Normalized Injury Score

-50

Pooled injury fit ····

OxPhos expression loss
I II III
IV V

Proteins and function
TNFa IL-1B AST
Steatosis ΔΨm-loss

C

OCR [pmol/min/μg]

7.5

2.5

*

Res   +   +
Myo   +   +

D

GSEA

Shared pathways

F0

GSEA

F1 ←→ F4

Reg. of cell morphogenesis
Reg. of cytoskeleton organization
Response to wounding
Cell Junction assembly
Neg. reg. of organelle organization
Pos. reg. of cellular component biogenesis
Small GTPase mediated signal transduction
Taxis
Est. or maintenance of cell polarity
Striated muscle cell differentiation
Pos. reg. of locomotion
Reg. of cellular component size
Actin filament organization
Cell projection assembly
Protein depolymerization
Cell matrix adhesion
Integrin mediated signaling pathways
Axon development
Cartilage development
Cell substrate adhesion
Mitochondrion organization
Small molecule biosynth. process
Gen. of precursor metabolites and energy
Organic acid biosynth. process
Fatty acid metabolic process
Sulfur compound metabolic process
Ribonucleoprotein complex biogenesis
Ribosome biogenesis
Aerobic respiration
Cellular respiration
Electron transport chain
Energy derivation by ox. of org. compounds
Cytoplasmic translation
Oxidative phosphorylation
Respiratory electron transport chain
ATP synth. coupled electron transport
NADH dehydrogenase complex assembly
Detoxification of copper ion
Dicarboxylic acid metabolic process
Mito. resp. chain complex assembly
Cellular amino acid biosynth. process
Mito. electron transport NADH to UQ
Alpha amino acid catabolic process
Cellular amino acid catabolic process
Alpha amino acid metabolic process
Organic acid catabolic process
Monocarboxylic acid catabolic process
Cellular amino acid metabolic process
Small molecule catabolic process

2
1
0
-1
-2
-3

Group=	F1	F3	F2	F4
n=	48	54	4	54	4	4	14

Fig. 1: Establishing a metabolically-associated steatotic liver disease (MASLD) drug testing model in stem cell-derived hepatocytes (iHeps) A) Schematic of the integrated drug repurposing discovery pipeline. F1-F4 = Fibrosis stage 1-4; PUFAs = polyunsaturated fatty acids **B)** Injury index in iHeps exposed to compounded metabolic injuries (Oleic and palmitic acid = OA/PA; Resistin and myostatin = Res/Myo; peripheral blood mononuclear cells = PBMCs). All parameters were standardized, ranging from Healthy (0%) to the most complex disease condition including all disease triggers (100%). Nuclear-encoded oxidative phosphorylation (OXPHOS) gene expression loss was quantified for each gene encoding individual mitochondrial complex proteins. TNFa, IL1B and AST protein data from supernatant. Steatosis and mitochondrial membrane potential ($\Delta\Psi m$) of live-dye based assayed are displayed. Each parameter was assessed in at least 3 samples. **C)** Mitochondrial spare respiratory capacity of iHeps treated with Res, Myo, or both, assessed for oxygen consumption rate (OCR). Data from Seahorse XF Cell Mito Stress Test shown. Significance was assessed using multiple t-test, corrected according to the Benjamini-Hochberg method. Each data point represents an independent sample **D)** Clinical benchmarking of the *in vitro* MASLD models to clinical disease stages from Goavaere et al (n=206) stratified by fibrosis using Gene Set Enrichment Analysis (GSEA)-based clustering. GSEA was performed for all *in vitro* disease conditions compared to the untreated control. For the clinical dataset, all stages were compared to stage F0. Mitochondrial processes are marked in red.

168
To capture indirect regulators within the interactome, we expanded ME8 by identifying the shortest paths
between ME8 genes using Dijkstra's algorithm. The resulting network comprised 7,545 proteins, with 70
key nodes identified through network centrality analysis **(Fig. 4C-D)**. These key nodes remained enriched
for mitochondrial pathways **(Fig. 4E)**, confirming the importance of the pathways after expansion. We
applied a network-based drug proximity model[19] to these 70 nodes, identifying 132 FDA-approved
compounds with significant proximity ($Z < -1.96$) (**Supp. Table 5**), including a set of mitochondrial-related
compounds prioritized for validation.
KG-based inference using PrimeKG: PrimeKG is a multimodal KG integrating curated biological data from
20 sources. We imported the graph into Neo4j for query-based inference, visualization, and the application
of graph algorithms. The graph schema and initial setup are shown in **Fig. 5A**. To identify repurposing
candidates, we used two complementary queries. First, we calculated shortest paths from "non-alcoholic
steatohepatitis" and "hepatic steatosis" nodes to drug nodes **(Fig. 5B–C)**, returning two unranked drug lists
(500 drugs each) with 97 compounds at their intersection. These included redox cofactors, fibrates, and
dual PPAR agonists. Second, we created a metabolic dysfunction-associated steatohepatitis (MASH)-
specific subgraph containing all nodes and relationships connected to "non-alcoholic steatohepatitis"
(**Supp. Fig. 3G, Fig. 5D**). Graph centrality algorithms applied to this 7,847-node network revealed additional
candidates such as DPP-4 inhibitors. Combining the three query outputs yielded 1,153 unique drugs, with
48 shared across multiple lists **(Supp. Table 6)**. Several compounds, e.g. aspirin, have prior links to MASH,
supporting the graph's biological relevance. We excluded Troglitazone from the list due to known
hepatotoxicity.
Despite differences in data sources and analytical layers, all strategies independently converged on a small
subset of three compounds (**Fig. 2, Supp. Table 7**): FAD, pyridoxal phosphate (PLP), and NADH (**Supp. Fig.
3H**). These redox-cofactors emerged from entirely distinct methodologies, yet their convergence
underscores a shared mechanistic axis involving mitochondrial and metabolic regulation in MASLD. From
these compounds, we prioritized FAD and flavin-related processes for further validation in MASLD, due to
their ill-defined role. This multi-modal strategy provides a robust framework for rational, systems-level drug
repurposing in complex metabolic diseases.
196

Fig. 2: Multimodal computational framework for drug repurposing in MASLD.
A unified pipeline integrating three network-based approaches was developed to identify candidate compounds targeting mitochondrial dysfunction in MASLD. Each method follows four key steps: data integration and signature selection, network construction, network embedding and analysis, and drug prioritization. **A) PPI-based pipeline:** Differentially expressed genes (DEGs) were derived from bulk RNA-seq (Govaere et al.) and filtered using single-cell and single-nucleus liver transcriptomes to retain hepatocyte-enriched genes. A subset overlapping with mitochondrial genes from MitoCarta 3.0 was used to construct a protein–protein interaction (PPI) network. Network hubs were identified by topological analysis, and drug proximity scores were computed to rank candidate compounds (**Supp. Table 4**). **B) WGCNA-based pipeline:** Co-expression modules associated with fibrosis stage were identified using weighted gene co-expression network analysis (WGCNA). Module ME8, enriched for mitochondrial functions, was prioritized. Network expansion identified a 'hidden layer' of intermediate nodes between module genes. Centrality analysis defined 70 key targets, and network-based proximity was used to prioritize compounds (**Supp. Table 5**). **C) Knowledge graph-based pipeline:** PrimeKG, a multimodal biomedical knowledge graph, was queried using phenotype nodes ("hepatic steatosis" and "non-alcoholic steatohepatitis"). Graph traversal, embedding-based scoring, and subgraph analysis identified drugs linked to MASLD-related nodes. Outputs from three complementary KG queries were combined (**Supp. Table 6**). **Bottom row:** A Venn diagram shows the overlap among drugs predicted by each approach. Three compounds, flavin adenine dinucleotide (FAD), pyridoxal phosphate (PLP), and nicotinamide adenine dinucleotide (NADH), were consistently identified across all pipelines (**Supp. Table 7**).

199
200

Fig. 3: Identification of hepatocyte-specific MASLD signatures and drug targets via integrated transcriptomic and network-based analysis. A multi-step pipeline was applied to derive a hepatocyte-specific disease module and prioritize drug candidates based on network proximity.

A) Heatmap showing the expression of 1,927 differentially expressed genes (DEGs) between MASL and control samples from human liver bulk RNA-seq data (GSE135251). Columns represent individual samples grouped by condition (MASL in red, control in turquoise), and rows represent DEGs scaled by row Z-score. Genes are hierarchically clustered. **B)** UMAP projection of a representative liver single-cell RNA-seq dataset showing major cell populations colored by group: MASLD (green), control (blue), and MASL signature-expressing cells (red overlay). MASL-specific DEGs are predominantly expressed in hepatocyte clusters. **C)** Heatmap of hepatocyte-specific upregulated genes retained after cross-validation with four single-cell and single-nucleus RNA-seq datasets (mouse and human). This refined 50-gene MASL signature displays strong separation between MASL and control samples, confirming disease relevance and cell type specificity. **D)** PPI network constructed from the refined MASL signature using STRING. Nodes represent proteins; edges indicate interactions. Red nodes highlight genes identified as network hubs by topological metrics (degree, closeness, betweenness, eigenvector centrality), filtered using permutation-based significance testing (N = 1,000). **E)** Expression heatmap of the 21 hub genes in the bulk RNA-seq dataset. Z-score-normalized rows illustrate consistent upregulation across MASL samples (red) and lower expression in controls (blue), further validating their disease association. **F)** Subnetwork showing the 21 hub genes (orange) and one of the 34 FDA-approved drugs identified via drug–target network proximity analysis. Green node indicates the example drug (FAD), and edges represent known drug–target associations within the human interactome.

Fig. 4: Fibrosis-correlated co-expression network analysis reveals mitochondrial module ME8 as a MASLD-relevant therapeutic target.

Weighted gene co-expression network analysis (WGCNA) was applied to transcriptomic profiles from the Govaere et al. cohort to identify gene modules associated with fibrosis progression in MASLD and nominate mitochondrial drug targets. **A)** Correlation heatmap between module eigengene expression and fibrosis stage. Each row represents a gene module (ME1–ME19), with Pearson correlation coefficients shown in the heatmap. Asterisks indicate statistical significance (*p < 0.05, **p < 0.01, ***p < 0.001). Representative enriched biological processes for significantly correlated modules are indicated with colored labels: ECM organization (blue), mitochondrial gene expression (purple), mRNA transport (orange), proteasomal degradation (green), vesicle transport (yellow), and RNA splicing (pink). **B)** Top 20 enriched Gene Ontology (GO) terms for module ME8, ranked by adjusted p-value. ME8 was enriched for mitochondrial and translational processes, including NADH dehydrogenase complex assembly, oxidative phosphorylation, and ATP metabolic processes. Gene ratio represents the proportion of ME8 genes annotated to each GO term. **C)** Network visualization of ME8 showing both co-expression (grey edges) and physical protein–protein interactions (red edges). Nodes represent ME8 genes, colored by differential expression: red = upregulated, blue = downregulated, white = unchanged (relative to control). The network shows that only a subset of co-expressed genes physically interact, suggesting a layered structure of molecular coordination. **D)** PPI subnetwork constructed from ME8 hub genes after network expansion. Nodes are scaled by degree and colored by betweenness centrality (light to dark purple). This subnetwork includes direct and indirect interactors connecting ME8 genes, used for drug proximity analysis. **E)** GO term enrichment for the 70 key nodes extracted from the ME8 expanded network. Terms include mitochondrial translation, respiratory chain assembly, ATP biosynthesis, and nucleotide metabolic processes. Enrichment remains highly specific to mitochondrial function, indicating that key biological themes are preserved after network expansion.

Fig. 5: Knowledge graph-based inference of MASLD drug candidates using PrimeKG.
Graph-based drug repurposing was performed using PrimeKG, a multimodal knowledge graph integrating curated biomedical relationships across diseases, drugs, phenotypes, and genes. Three query strategies were used to extract compound lists relevant to hepatic steatosis and MASH. **A)** Schema of PrimeKG highlighting node types (e.g., gene, drug, disease, phenotype, pathway) and edge relationships (e.g., "associated with," "target," "interacts with," "phenotype present"). The subset of the graph used for this study is indicated in the blue box. Directed edges define relationships used to construct query paths from disease or phenotype nodes to drug targets. **B)** First traversal strategy starting from the MASH node. Gene nodes directly "associated with" MASH were retrieved, and drug nodes targeting these genes were extracted to create unranked Drug List 1. **C)** Second strategy starting from the "hepatic steatosis" phenotype node. A path was followed via "phenotype present" to reach relevant disease nodes, then further traversed through "associated with" to gene nodes, and finally to their targeting drugs. This yielded unranked Drug List 2, enriched for compounds connected through multiple phenotypic and disease layers. **D)** Third strategy involving subgraph extraction and algorithmic ranking. A MASH-specific subgraph was generated by extracting all nodes and edges connected to the "non-alcoholic steatohepatitis" node. Network centrality metrics (PageRank, betweenness centrality, and closeness centrality) were applied to score and rank drug nodes based on their topological influence within the disease module. This produced ranked Drug List 3, offering a prioritised view of candidate compounds. Stea = Hepatic steatosis; SCAD = Short-Chain Acyl-CoA Dehydrogenase; FAD = Flavin adenine dinucleotide; MASH = metabolic dysfunction-associated steatohepatitis; NQO1 = NAD(P)H:quinone oxidoreductase 1; ACADS = Acyl-CoA Dehydrogenase Short Chain

Flavoproteome perturbations in MASLD
Due to our inferred prediction of FAD´s anti-MASLD activity, we assessed flavin perturbations both *in vitro*
and *in vivo* across independent populations **(Fig. 6A)**. First, metabolite enrichment analysis revealed
significant enrichment *in vitro* for FAD-associated genes among those that were down-regulated **(Fig. 6B)**.
To examine these changes further, we filtered the *in vitro* gene expression data for the human
flavoproteome[20]. Cluster heatmap analysis displayed Res/Myo as a tipping point in flavoproteome
perturbations, especially for the gene expression perturbation of *SARDH*, *PRODH*, *LDHD, DPYD*, and most
of the *FMO* gene family members **(Fig. 6C, Supp. Fig 4A&B)**. *In vivo*, we validated flavoproteome
perturbations using liver tissue proteomics data from our clinically stratified pediatric MASLD cohort in
Berlin, Germany[21]. Using pediatric data reduces the confounding effects of cardiometabolic diseases,
alcohol intake, and concomitant drug treatment present in adults. We observed significant down-regulation
of flavoproteins and FAD-related processes in advanced disease stage, focusing on subcellular processes
(Fig. 6D). All detected proteins of genes affected *in vitro* had, on average, lower abundance in advanced
disease, with DPYD being significantly reduced **(Supp. Fig. 4C & Fig. 6E)**. Next, to connect flavin
processes with clinical outcomes in a large adult cohort, we performed survival analysis using our MASLD
SteatoSITE cohort (n = 508)[22]. As most of the *FMO* gene family was affected *in vitro*, we prioritized these
genes in our analysis. While *FMO1* showed no significant association **(Supp. Fig. 4D)**, increased expression
of *FMO3, FMO4*, and *FMO5* correlated significantly with a higher survival **(Fig. 6F)**. Notably, high *FMO2*
expression correlated with decreased survival **(Supp. Fig. 4D)**, contradicting its recently described
protective role[23]. Survival analysis of the FAD-homeostasis genes *RFT3, RFK,* and *FLAD1* reinforced flavin
perturbances in MASLD pathology **(Supp. Fig. 4E)**. While *RFK* expression correlated significantly with
increased survival, elevated *RFT3* and *FLAD1* expression were associated with adverse outcomes. Thus,
indicating a partial compensatory regulation of synthesizing enzymes that overcomes a co-factor shortage
in advanced disease stages. To evaluate the predictive value of flavoproteome activity, we applied LASSO-
penalized Cox regression, identifying a 10-gene signature that stratified patients by hepatic events (n = 446)
(Fig. 6G, Supp. Table 8). Key contributors included the downregulation of *DPYD* and *RFK*, and the
upregulation of *FMO2*, associated with an increased likelihood of adverse hepatic events. Ultimately, to
enable the experimental interrogation of flavin processes in MASLD via manipulation of flavin homeostasis,
we demonstrate that high-dose FAD supplementation (FAD^H, 50 µM) significantly raises intracellular FAD
levels in our *in vitro* model by 1.3-fold **(Supp. Fig. 4F)**. Thus, we demonstrate the clinical relevance of flavin
processes in MASLD pathology and establish a framework for their functional interrogation using our *in vitro*
MASLD model.

Fig. 6: FAD-dependent processes are perturbed in MASLD-progression. **A)** Scheme of flavoprotein analysis *in vitro* and different *in vivo* populations **B)** Enrichment analysis for the down-regulated differentially expressed genes of the MASLD *in vitro* model incorporating OA/PA, Res/Myo and PBMCs, assessed for the associated metabolites using the human Metabolome database. **C)** Heatmap of normalized gene counts showing clustering of FAD-dependent protein gene expression across different MASLD models in iHeps **D)** Enrichment analysis of shotgun proteomics data from the Berlin pediatric MASLD patients (cohort as in Hudert et al.[21]) Enrichment analysis was done for advanced disease stage (NAS 5-8) in relation to early disease stage (NAS 1-2) **E)** Abundance of DPYD protein in the pediatric MASLD cohort described by Hudert et al.[21] Significance was assessed using multiple t-tests, corrected by the Benjamini-Hochberg method. Each dot represents an individual patient **F)** Survival analysis of FMO-gene expression and human MASLD patients from the SteatoSITE cohort[22]. Significance was assessed using a log-rank test. **G)** Predictive flavoprotein model for hepatic event probability using LASSO-penalized Cox regression. Significance was assessed using a log-rank test.

Flavin modulation therapeutically intercepts metabolic crisis

We assessed the therapeutic potential of flavin manipulation using iHeps and primary hepatocytes **(Fig.**
7A). Titrating FAD in iHeps revealed anti-steatotic effects at 10 µM **(Fig. 7B & Supp. Fig. 5A).** Yet, FAD^H (50
µM) restored mitochondrial function –known to be perturbed in MASH[24,25]– shown by increased coupling
efficiency, decreased mitochondrial uncoupling and enhanced Complex II activity **(Fig. 7C, Supp. Fig. 5B).**
We confirmed the pro-mitochondrial and anti-steatotic effects using primary human hepatocytes **(Supp.**
Fig. 5C-E). FAD^H reduced hepatocyte injury, assessed by the clinical liver injury marker AST, and altered
the fatty acid profile in iHeps **(Supp. Fig. 5F&G)**. Specifically, FAD^H reversed injury-induced changes in
saturated-to-unsaturated and omega-3-to-omega-6 fatty acid ratios, demonstrating beneficial effects on
fatty acid metabolism[26] **(Fig. 7D)**. To pioneer a pipeline that can identify rescue effects without extensive
cellular phenotyping, we assessed FAD^H's effects using label-free artificial neural network-based imaging
analysis of plane brightfield images **(Supp. Fig. 6).** Thereby, FAD^H mitigated injury-induced iHep
morphological perturbations, indicated by tSNE clustering, with multiple mitigated features, including
cytoplasm or droplet inclusion modalities **(Supp. Fig. 6C & Supp. Table 9).** This highlights a rapid and
scalable approach for detecting rescue effects. To dissect therapeutic FAD-mediated effects from
mitochondrial biogenesis, we tested additional overexpression of the mitochondrial biogenesis-inducer
PPARGC1A. *PPARGC1A* is downregulated in MASH patients[27] and specifically in pro-inflammatory *in vitro*
conditions in which its overexpression restores OxPhos gene expression **(Supp. Fig. 7A-C).** RNA-Seq
results revealed that *PPARGC1A*-induction amplified FAD-mediated effects. However, FAD^H alone already
induced significant transcriptomic changes **(Supp. Fig. 7D&E)** in iHeps, underscoring its pharmacological
properties. FAD^H restored pathways suppressed in the disease model and MASH patients[28], like the
complement and coagulation system, while inhibiting disease-induced pathways, including TGF-b signaling
(Fig. 7E, Supp. Fig. 1C). Combinational *PPARGC1A* overexpression further increased metabolic pathway
activity, e.g. TCA cycle **(Supp. Fig. 7F).** Functionally, combining FAD^H with *PPARGC1A* significantly
reduced steatosis compared to FAD^H alone **(Supp. Fig. 7G)**, supporting synergistic effects. To validate
FAD's anti-fibrogenic and anti-TGF-modulating effects, we harnessed fibro-competent multi-cellular liver
organoids from iPSCs containing hepatocytes, macrophages, stellate and endothelial cells[29] **(Supp. Fig.**
8A-C). FAD^H reduced fibrogenic gene signature in organoids injured by the secretome of the inflammatory
iHep MASLD model **(Supp. Fig. 8D)**. Next, we evaluated FAD's immunomodulatory effects in primary
immune cells exposed to the mentioned secretome **(Supp. Fig. 9)**. FAD^H intercepted the activation of innate
and adaptive immune cell compartments, indicated by suppressed surface expression of MASLD-
associated inflammatory activation markers[30] in monocytes and both CD4^+ and CD8^+ T-cells. Notably, this
anti-inflammatory effect persisted in LPS-induced inflammation (**Supp. Fig 9D**). To expand the therapeutic
scope beyond direct FAD supplementation, we predicted pharmacological compounds targeting similar
proteins as FAD (considered flavo-active) of which norleucine, 3P, and aspirin fully restored mitochondrial
respiration in primary hepatocytes **(Fig. 7F)**. These results support the therapeutic potential of modulating
flavin pathways and point to additional clinically tractable compounds.
Thus, we demonstrate functional evidence for therapeutic manipulation of flavin-processes for alleviating
hepatocyte-centric pathologies in MASLD.

A FAD — Flavoactive compounds: 3P, Aspirin, Norleucine → iHeps or primary hepatocytes

B OA/PA + Res/Myo; Healthy, Vehicle, FAD [µM]: 2, 10, 50, 250; Merge LipidSpot Hoechst

C Coupling Efficiency, Proton Leak — Oxygen consumption rate norm. to vehicle; / FAD^H; OA/PA + Res/Myo

D Saturated/Unsaturated; Ω-3/Ω-6; Healthy, / FAD^H; Res/Myo + PBMCs

E Gene Ratio; Up, Down; Gene Count 10, 15, 20; Focal adhesion, MicroRNAs in cancer, ECM-receptor interaction, Protein digestion and absorption, TGF-b signaling pathway, Butanoate metabolism, Tryptophan metabolism, Valine, leucine and isoleucine degradation, PPAR signaling pathway, Thyroid hormone synthesis, Cholesterol metabolism, Peroxisome, Complement and coagulation cascades

F Oxygen consumption rate norm. to vehicle; Healthy, / FAD^H Nor 3P Asp; OA/PA + Res/Myo

Fig. 7: Manipulating FAD-homeostasis alleviates MASLD hallmarks *in vitro*. FAD[H] = high-dose FAD (50 µM) **A)** Schematic of flavin-augmentation experiments in iPSC-derived (iHeps) and primary hepatocytes. **B)** Representative images of fluorescent staining of lipid droplets and nuclei in iHeps that are untreated or exposed to oleic acid and palmitic acid (OA/PA) + resistin and myostatin (Res/Myo) exposed to different FAD concentrations. **C)** Mitochondrial coupling efficiency and proton leak of iHeps (●) and primary mouse hepatocytes (▲) treated exposed to OA/PA + Res/Myo with or without FAD[H] (50 µM) assessed using the Seahorse XF Cell Mito Stress Test. Significance was assessed using multiple t-tests, corrected by the Benjamini-Hochberg method. Each dot represents an independent sample **D)** Quantification of the saturated over non-saturated and omega-3 over omega-6 fatty acid ratios normalized to protein concentration in healthy iHeps or exposed to Res/Myo and peripheral blood mono-nuclear cells (PBMCs) with or without FAD[H] (50µM). Significance was assessed using multiple t-tests, corrected by the Benjamini-Hochberg method. Each dot represent an independent sample. **E)** Top KEGG pathways enriched in iHeps exposed to the 3-Hit disease model treated with FAD[H] (50 µM) relative to the vehicle control (both transduced with an EGFP control). **F)** Testing of mitochondrial spare respiratory capacity for additional flavo-active compounds. Mitochondrial spare respiratory capacity was assessed using the Seahorse XF Cell Mito Stress Test. Norleucine (Nor) was used in a 1 mM, Aspirin (Asp) in 5 µM and 3-Pyridin-4-yl-1H-indazol-5-ylamine dihydrochloride (3P) in 1.5 µM concentration. Significance was assessed using multiple t-tests, corrected by the Benjamini-Hochberg method. Each dot represents an independent sample. / = Vehicle

Discussion:
MASLD is the most common chronic liver disease, yet treatment options are limited[31]. Although several Phase 3 clinical trials were conducted, only Resmetirom received FDA approval. Still, Resmetirom improved fibrosis only in a small number of patients, emphasizing the need for additional treatment strategies. Herein, we integrate stem cell-derived systems with computational drug predictions, uncovering the therapeutic potential of targeting the flavoproteome in MASLD. We established a framework that couples human-relevant *in vitro* disease modeling with network-based and ML-driven hypothesis generation, based on stratified molecular patient data. Specifically, we developed a donor-adjustable drug-testing pipeline capturing hallmark pathologies of MASLD, from inter-organ metabolic triggers to a fibro-inflammatory response. We developed a computational toolbox that predicts biological pathway networks and drug candidates with high accuracy to intercept pathophysiological processes. We increased the chances of successful experimental validation by using a closed-loop iterative feedback system, reducing the need for high-throughput screening. This strategy identified FAD and other compounds active in the flavoprotein network, including Aspirin, which was recently described to improve steatosis in MASLD through an unknown mechanism[32].
Previous MASLD drug repurposing was limited in both models and computational thoroughness, utilizing microarray data from a small patient cohort and evaluating anti-steatotic drugs in HepG2 cells[33]. Contrarily, we harnessed recent advances in ML, not only in drug prioritization but also in detecting rescue effects by label-free imaging based solely on brightfield images. This approach enables straightforward rescue detection and highlights potentially affected morphological structures, as demonstrated for endothelial cells[34]. Further, harmonizing pathology- and cell-type-tailored multi-modal computational predictions, clinical validation and authentic pre-clinical modeling, advanced drug repurposing frameworks from earlier reports[33,35].
This work pioneers the integration of three complementary computational strategies: PPI-based drug proximity analysis, fibrosis-stage-resolved WGCNA, and multimodal KG inference into a unified pipeline. Translating these predictions into relevant *in vitro* models establishes a link between computational findings and experimental validation. Focusing on mitochondrial dysfunction as a central driver of MASLD enabled the biologically meaningful filtering of gene signatures and improved the specificity of compound prioritization. The multimodal KG introduced a systems-level perspective, identifying compounds associated with hepatic phenotypes through complex network relationships. Notably, all three strategies converged on FAD, highlighting the methodology's robustness and supporting a mechanistic role. Yet, limitations may persist in the KG-based component. KGs may offer biased inferences, relying on curated databases that are not routinely updated and are constrained by existing knowledge. Furthermore, integrating non-topological features, like drug chemical structure, could enhance the predictive power. Nonetheless, the KG remains a valuable hypothesis-generation tool when combined with orthogonal computational and experimental approaches, as demonstrated here.
Regarding *in vitro* modeling, we established a roadmap of compounded hepatocellular injury in iHeps. Dissecting the injury response revealed a crucial role for Res and Myo in metabolic and mitochondrial dysfunction in human inter-organ biology, supporting their debated role in humans as reported in mice and carcinoma cell lines[36–40]. Furthermore, leukocyte co-culture led to a second metabolic tipping point,

emphasizing the importance of liver-immune cell crosstalk in MASLD, similarly demonstrated in insulin
resistance modeling[41]. Our modular disease model, combined with multi-cellular organoids and primary
immune cells, and rigorous computational drug prioritization, enabled stage-specific interventions in
complex systems, capturing inter-organ signaling and fibro-inflammatory pathology. Although our Res/Myo
focus as extrahepatic signals further uncovered their role in MASLD, incorporating gut-derived signals or
more defined insulin resistance signaling may further advance our model[42,43]. Furthermore, large-scale
multi-donor analysis[44] can leverage the model´s full potential by enabling donor-specific drug responses for
personalized therapy assessment.
By revealing the importance of flavin processes in MASLD, we demonstrate a promising axis for therapeutic
intervention. Clinically, riboflavin, the precursor of FAD, serves as a treatment for genetic disorders of flavin
metabolism[45] or migraine prevention[46]. Riboflavin deficiency increases MASLD activity in rodents, zebrafish,
and HepG2 cells[47–49], and induces a MASLD phenotype via decreased *PPARA* activity in mice[50]. Stringently,
we show increased PPAR signaling upon FAD supplementation **(Fig. 7E),** indicating a reciprocal interplay.
Despite early reports of flavin reduction in cirrhotic liver disease[51], underlined by our survival analysis, this
is, to our knowledge, the first study to show flavo-protective effects for human MASLD. Previous studies
lacked human relevance, showing decreased hepatic lipid accumulation under riboflavin and selenium co-
supplementation in rats[52], not studying fibro-inflammatory responses. For better clinical translation, we
tested more bioavailable flavo-proteome-modulating drugs, like Aspirin or norleucine, an experimental non-
proteinogenic amino acid commonly used for protein engineering, for which exact molecular targets need
to be identified in future studies. Future translational studies will determine whether manipulating flavin
pathways alone suffices for treating MASLD in a clinical setting or if additional strategies, like mitochondrial
biogenesis stimulus via *PPARGC1A,* are required.
Thus, we showcase how integrating complex stem-cell-derived models with computational drug predictions
enables rapid preclinical discovery and human model-centric functional validation of therapeutic
candidates. Expanding this framework by incorporating patient-specific cells or combinatorial drug
approaches could further advance the dissection of complex diseases and enable more precise therapeutic
interventions.
371
Online Methods:
<u>iPSCs lines</u>
We used the two human iPSC donor lines BIHi001-B major homozygous for PNPLA3 rs738409 variant
(PNPLA3- I148I) and BIHi005-A minor homozygous for the PNPLA3 rs738409 variant (PNPLA3-I148M),
provided by the BIH Core Unit Pluripotent Stem Cells and Organoids. Both cell lines are registered in the
Human Pluripotent Stem Cell Registry (RRID:CVCL_YN70 and RRID:CVCL_IT56). We cultured cells on
vitronectin-coated (Thermo) plates in mTeSR medium (Stem Cell Tech). All experiments were performed at
37°C in a 5% O_2 and 5% CO_2 hypoxic atmosphere.
380
<u>Differentiation into iHeps</u>
We adapted a 28-day protocol[53], optimized by Seifert & Ludwik et al[54] and modified by us for the maturation
of iHeps, as following:
Generation and thawing of ROCKi-Adapted iPSCs:
iPSCs were expanded in mTeSR medium (StemCell Technologies) containing 10 µM ROCKi (Tocris),
detached with 0.5 mM EDTA in PBS for 5 min at room temperature, resuspended in Bambanker freezing
medium (NIPPON), frozen at -80°C using a Mr. Frosty container, and transferred to vapor-phase liquid
nitrogen (-135°C). For thawing, vials were warmed in a 37°C water bath until a small ice crystal remained,
diluted dropwise with 10× volume of mTeSR + ROCKi, centrifuged (3 min, 300 g), and resuspended in fresh
ROCKi-containing mTeSR.
Hepatic Differentiation Protocol:
Upon reaching ~80% confluency, iPSCs were single-cell seeded onto Geltrex-coated 6-, 12-, or 24-well
plates (Day 0) and differentiated as follows:
Days 0–5: DE-induction medium (RPMI1640, 2% B27, 1% PenStrep, 100 ng/ml Activin A, 50 ng/ml Wnt3,
0.5 mM sodium butyrate), followed by 1 day without sodium butyrate.
Days 6–12: KO-medium (KO-DMEM, 20% KO Serum, 1% PenStrep, 1% Glutamax, 1% NEAA, 1% DMSO,
0.1 mM β-mercaptoethanol) for hepatic commitment.
Days 13–19: HBM medium (Lonza) supplemented with 20 ng/ml HGF, 20 ng/ml OSM, and 1% PenStrep.
Days 20–28: Maturation in HCM Bulletkit medium (Lonza, excluding EGF), with supplementation of 5 nM T3
(Tocris), 100 nM sodium selenite (Sigma), and 100 µM CDCA (Sigma)[55,56].
401
402 <u>Primary hepatocytes</u>

Primary hepatocytes were isolated from C57BL/6J mice (12–20 weeks old) using a well-established two-step perfusion technique. Initially, the liver was perfused with a calcium-free buffer to remove blood and non-parenchymal cells, followed by perfusion with collagenase (LiberaseTM, Roche, Switzerland) to dissociate the tissue enzymatically. The resulting cell suspension was purified using a Percoll gradient, yielding hepatocytes with >90% viability as previously described[57]. Purified cells were seeded onto collagen type I-coated culture plates (5 µg/cm²) at densities suitable for experimental needs. Cells were maintained in William's E medium supplemented with 2 mM L-glutamine, 5% PenStrep (penicillin–streptomycin), and hepatocyte growth supplements (CM4000; Thermo Fisher Scientific). This complete medium supported optimal growth and maintenance of hepatocytes *in vitro*.
All animal experiments complied with ethical regulations and were approved by the Committee on Research Animal Care of the Charité Universitätsmedizin Berlin, Berlin, Germany (License number: G0174/20).
The primary human hepatocytes were isolated from a 63-year-old male patient undergoing liver surgery due to colorectal liver metastasis. Cell isolation was done under informed consent and the ethics statement EA1/257/14 according to the protocol described in Horner et al[58]. PHHs were frozen in FCS with 10% DMSO and stored in vapor phase liquid nitrogen below -135°C until further use. For the experiments, we utilized a recently described protocol [59]. Briefly, cells were plated on collagen-coated plates (collagen type 1, rat tail, Gibco), with a Matrigel overlay, and maintained in HCM medium (Lonza).

Primary human peripheral blood mononuclear cells

Blood isolation was done in accordance with ethical standards of the Charité Universitätsmedizin Berlin under the ethical approval EA2_065_21. We obtained healthy human PBMCs using Ficoll-Paque (Sigma). To minimize PBMC donor variability, we derived them from a standard "master" donor who a physician clinically monitored to ensure the absence of clinical illness or medication intake. Specifically, we mixed whole blood with PBS (w/o Ca^{2+} Mg^{2+}) in a 1:1 ratio and overlaid this dilution on the Ficoll in a 3 to 2 ratio, respectively. We then centrifuged the mixture for 30 minutes at 300 g at room temperature, using the lowest levels of brake and acceleration. The leukocyte layer was transferred to a new tube and washed once with PBS. After lysing red blood cells by incubating the cells for 10 min in ACK lysis buffer, we washed twice, counted and froze cells down in a 10% DMSO solution in FCS using Mr. Frosty. The cells were stored in vapor phase liquid nitrogen below -135°C until further use.

Compounded metabolic injury model

We tested different combinations of cumulative metabolic injury hits in iHeps. Generally, we exposed iHeps for 3 days to either FFAs alone (1 Hit), combined with the inter-organ signaling molecules Res and Myo (2 Hit) or in addition with co-cultured PBMCs (3 Hits). For rescue experiments, we also included combinations as indicated.
For FFA injury, we exposed cells to oleic and palmitic acid (OA and PA) in a 5 to 2 ratio on day 28 for 3 days. Oleic and palmitic acid (both Merck) were coupled to the HCM bullet kit-derived BSA by incubation at 37°C for at least 2 h. We used 125 µM as total fatty acid concentration.
To account for inter-organ signaling in our model, we exposed iHeps to recombinant resistin and myostatin (100 ng/ml each, Peprotech) for three days.
For modeling inflammation, we included PBMCs in an indirect trans-well co-culture using Millicell trans-well inserts (Millipore) with a 0.4 µm pore size, 1 day after the addition of the other injuries (Day 29). We thawed the PBMCs 18 h before co-culture and cultured them in RPMI medium containing 10% heat-inactivated fetal calf serum (FCS) and Pen-strep. Within this 18 h pre-incubation, PBMCs incubated with 5 µg/ml lipopolysaccharide (LPS, eBioscience or Stem Cell Technologies) or without. Before adding them to the inner compartments of the trans-wells, we washed of the RPMI medium and re-suspended them in the respective HCM culture medium. PBMC viability after coculture was consistently over 90%, as assessed via Trypan Blue staining. To derive the overall injury metric, we normalized each disease marker for all conditions to untreated iHeps (0%) and iHeps exposed to 3 Hits (100%) and then applyed the geom_smooth function (method='loess', formula= y~x) in RStudio.
Due to our focus on multiple acquired metabolic injuries, compounding their mitochondrial dysfunction, we selected PNPLA3-I148I iHeps for in-depth characterization and mechanistic rescue experiments.

Live dyes

We incubated the cells for 30 min with LipidSpot 488/610 and MitoTracker™ Red CMXRos staining to assess steatosis and mitochondrial membrane potential. We used all dyes according to manufacturer´s instructions. Specifically, Mitotracker was used with a 200 nM end concentration. Eventually, we either performed live-cell imaging or fixed cells in the culture plate using 4% paraformaldehyde, followed by subsequent PBS washes before imaging.

Fluorescence imaging and image analysis

We imaged cells in their respective well plates using the Phenix High-Content Imaging system (Revity). We
captured at least 10 individual images per well, which we analyzed using the Signals Image Artist software
(Version 1.3.29). For measuring steatosis or mitochondrial membrane potential, we manually set upper and
lower thresholds for LipidSpot and Mitotracker staining in each experiment to eliminate background and
artefacts. For Mitotracker staining, we then measured intensity for the selected image region. To assess
steatosis, we measured LipidSpot area and divided it by the respective Mitotracker area or number of
nuclei for each image. Respective signals for steatosis or Mitotracker we then further normalized as
indicated.
472
RNA Isolation
We isolated RNA using a hybrid protocol of trizol RNA isolation combined with column-based purification
(Qiagen or Thermo). We lysed cells in 1 ml of trizol (Thermo) to which we subsequently added 0.2 ml
chloroform. All chloroform-trizol mixtures were then centrifuged for 10 min, using 12,000 g at 4°C. Then, we
transferred the aqueous phase to a new tube and added 70% ethanol in a 1:1 ratio. This mixture we loaded
on RNeasy Micro Kit columns (Qiagen). From this point, we performed every step according to the
manufacturer´s recommendations (including the on-column DNAse treatment). We assessed the
concentration of the isolated RNA using Nanodrop.
481
qPCR
For quantifying gene expression using qPCR, we reverse-transcribed RNA into cDNA using a cDNA
Synthesis kit (High-Capacity cDNA Reverse Transcription Kit (Thermo, Cat. No. 4368813), biotechrabbitTM
or Applied Biosystems™ High-Capacity RNA-to-cDNA™ Kit) according to manufacturer´s instructions. We
then performed RT-qPCR using the PowerUp SYBR Green Master Mix (Applied Biosystems, Cat. No.
A25776) on a QuantStudio 3 Real-Time PCR System (Thermo Fisher Scientific Inc.) according to the
manufacturer's instructions. Primers were synthesized by Eurofins Genomics, with sequences listed in
Table 1. Gene expression changes were calculated using the ddCT method.
490
491 **Table 1: Oligos used in this study**

hNDUFA8_F	ACTGGTCAACAAGTGTGCTTTGG
hNDUFA8_R	CACACTCGTCAAACTTTGCCTGC
hNDUFB1_F	TGAGTTCCCTGTTGCCCTTGGT
hNDUFB1_R	GGGACAAGAACATGAACCCAGTG
hSDHA_F	GAGATGTGGTGTCTCGGTCCAT
hSDHA_R	GCTGTCTCTGAAATGCCAGGCA
hSDHB_F	GCAGTCCATAGAAGAGCGTGAG
hSDHB_R	TGTCTCCGTTCCACCAGTAGCT
hUQCRB_F	TAAGAGGGCACTGGACCTGAAC
hUQCRB_R	CATGATTACTTCTTTGCCCATTCTT
hCox7A2_F	CGGAGAAGCAAAAACTGTTCCAG
hCox7A2_R	CCAACTGTAAGAATCATGGTGGC
hCox5B_F	GGAGATCATGCTGGCTGCAAAG
hCox5B_R	GCAGCCTACTATTCTCTTGTTGG
hATP5A1_F	GCTCCTTACTCTGGCTGTTCCA
hATP5A1_R	GCGGAGCAACAGAGACATCTGA
hATP5G3_F	GCAACAGTAGGAGTGGCTGGTT
hATP5G3_R	GCTTCAGACAAGGCAAATCCCAG
PPARGC1A_F	CCAAAGGATGCGCTCTCGTTCA
PPARGC1A_R	CGGTGTCTGTAGTGGCTTGACT
COL1A1_F	GATTCCCTGGACCTAAAGGTGC
COL1A1_R	AGCCTCTCCATCTTTGCCAGCA
COL4A1_F	TGTTGACGGCTTACCTGGAGAC
COL4A1_R	GGTAGACCAACTCCAGGCTCTC
COL4A2_F	GGATAACAGGCGTGACTGGAGT
COL4A2_R	CTTTGCCACCAGGCAGTCCAAT
TGFB1_F	TACCTGAACCCGTGTTGCTCTC
TGFB1_R	GTTGCTGAGGTATCGCCAGGAA
RPS18_F	GCAGAATCCACGCCAGTACAAG
RPS18_R	GCTTGTTGTCCAGACCATTGGC

<u>Supernatant analysis</u>
We collected supernatant from cells exposed to the indicated disease models with or without rescue conditions after 18-24h, albeit consistent timing for ensured for individual experiments. We stored the samples at -80°C until further use. The molecules TNFa, IL1B and aspartate aminotransferase (AST) were measured by Labor Berlin–Charité Vivantes GmbH, Berlin, Germany, using standard clinical biochemical procedures that can be detailed upon request.

<u>Secretome analysis using Liquide chromatography – tandem mass spectrometry (LC-MS/MS)</u>
For secretome analysis, we analyzed the conditioned supernatant of cells using mass spectrometry. For more sensitive detection of secreted proteins, we added HCM medium without additional growth factors or BSA for 24 h after exposure to the respective disease model. We centrifuged the samples and stored the supernatant at -80°C until the analysis.
For the analysis, 18 µl of each culture supernatant were separated by one-dimensional SDS-polyacrylamide gel electrophoresis according to Beier et al.[60]. Each sample lane was divided into three sub-samples (all protein bands above the bovine serum albumin (BSA) band, the BSA band and all protein bands below the BSA band). In-gel digestion of proteins, extraction of peptides, desalting and preparation of the samples for LC-MS/MS analysis were performed as described by Beier et al [60].
The LC-MS/MS analysis was performed using a nanoAQUITY UPLC system (Waters corporation, Milford, MA, USA) coupled to an LTQ Orbitrap Velos Pro mass spectrometer (Thermo Fisher Scientific Inc., Waltham, Massachusetts; USA). The LC gradient used is described in Lerch et al[61]. A m/z of 350-1750 was used for primary MS scans. Except for the exclusion time, the other MS parameters were taken from Beier et al. (2021). An exclusion time of 12 s was chosen.
MS/MS raw files were analysed using MaxQuant (Max Planck Institute of Biochemistry, Martinsried, Germany, www.maxquant.org, version 2.4.3.0 according to the parameters described previously (Beier et al.[60]) using N-terminal acetylation as an additional variable modification against a database of human proteins (taxonomy id 3A9606_29) and isoforms downloaded from Uniprot on 2023-06-27 and a database of Bos taurus proteins downloaded from Uniprot on 2023-12-20.
Proteins are considered to be reliably identified if they are identified by at least two unique peptides, each of them detected by at least one MS/MS scan in two different samples/replicates. In addition, their total intensity must be greater than 100.000. Protein quantification was performed using MaxQuant (version 2.4.3.0) intensity-based absolute quantification (iBAQ) (Schwanhäusser et al.)[62]. The means of two biological replicates (for iHeps without any trigger only one replicate was measured) were normalised and plotted as a heatmap using the R package tidyHeatmap2 after subtracting the means of the blank samples (culture medium). The data input is provided in **Supplementary Table 2.**
The mass spectrometry proteomic data have been deposited in the ProteomeXchange Consortium via the PRIDE partner repository (Perez-Riverol et al[63]) with the dataset identifier PXD063495. Reviewers will have access to the MS data at project accession: PXD063495 using the token lSRc2SuWKYkF. Alternatively, reviewer can access the dataset by logging in to the PRIDE website using the following account details: Username: reviewer pxd063495@ebi.ac.uk, Password: 7jwiwNNKtUYa

<u>Bulk RNA Sequencing</u>
mRNA sequencing for characterizing the MASLD *in vitro* model was performed at the BIH Genomics Core facility using Illumina TruSeq stranded mRNA library preparation and for sequencing using 50 mln of PE100nt reads per sample on a 25B flowcell of the Illumina NovaSeq X Plus. For exploring transcriptomic changes by FAD treatment (FAD rescue) we used the services from Novogene. The sequencing of the FAD rescue samples was performed using the PE150 sequencing strategy on NovaSeq Xplus sequencer 25B sequencing flow cell (Illumina, San Diego) by Novogene Germany (Munich, Germany).

Features of the MASLD *in vitro* model dataset:
The MASLD *in vitro* model bulk rna-seq dataset consisted of 32 samples from two donor lines *PNPLA3*-I148I and -I148M (16 for each donor line, four of each sample), see Table 2.

Table 2: Sample conditions used for bulk RNA-sequencing and downstream analysis for characterizing the *in vitro* MASLD models.

Name	Condition
HCM	healthy control (untreated cells)
OA PA	cells that were solely treated with an excess of free fatty acids

OA PA ResMyo	cells that in addition to free fatty acid excess also were exposed to 2 MASLD associated signaling molecules from other tissues (adipose and muscle tissue)
OA PA ResMyo PBMCs	cells that experienced both of the above-mentioned conditions + a co-culture of stimulated primary immune cells (immune cells were not included in the RNA sequencing)

548
549 Processing and analysis of MASLD *in vitro* model RNA-seq dataset:
550 We started by carrying out a whole genome decoy-aware indexing and transcript quantification using
551 Salmon v1.10.1 [64]. The Salmon pipeline estimates transcript-level abundance from RNA-seq read data. We
552 then used DESeq2 [65], to test for differentially expressed genes between the following conditions:
    I.   OA PA vs HCM
    II.   OA PA ResMyo vs HCM
    III. OA PA ResMyo PBMCs vs HCM

For derivation of the OXPHOS expression loss in the overall injury index, the counts of all genes of each
oxidative phosphorylation complex were normalized to the total counts and normalized to the untreated
control, to derive the expression changes. A similar strategy was used for deriving the inflammatory gene
expression changes.

Signaling pathway enrichment in MASLD *in vitro* model dataset:
For each set of differentially expressed genes, we tested for enriched KEGG pathways[66].
To identify the common and distinct signalling pathways enriched in the two cell lines, we used the
pathfindR pipeline[67]. pathfindR leverages interaction information from a protein-protein interaction network
(PIN) to help us identify enriched biological pathways from gene expression data.
We mapped the differentially expressed genes with p values less than 0.05 onto BioGrid[68] as the PIN, then
performed an active subnetwork search using a Greedy algorithm[69]. The resulting active subnetworks were
then filtered based on the following scores: (a) quantile threshold greater than 0.8 and (b) number of
significant genes they contain greater than 10. Using the genes in each of the active subnetworks, the
significantly enriched terms (pathways/gene sets) were identified. Gene Ontology "biological processes"[70]
were used as the gene sets for enrichment analysis. Enriched terms with Bonferroni adjusted p values larger
than 0.05 were discarded and the lowest adjusted p value (over all active subnetworks) for each term was
kept. This process of active subnetwork search and enrichment analyses was repeated for 10 iterations,
performed in parallel.

Clinical benchmarking using gene set enrichment analysis:
To functionally characterize the *in vitro* models, we used gene set enrichment analysis (GSEA). GSEA is a
computational method used to identify whether a set of predefined genes shows statistically significant
differences in expression between two biological states (e.g., disease vs. control). Unlike traditional
approaches that focus on individual genes, GSEA evaluates the expression of entire gene sets (groups of
functionally related genes) to see if they are more likely to be upregulated or downregulated in a particular
condition. In the context of this study GSEA helps in functional characterization by providing insights into
the biological processes that are associated with the observed changes in gene expression.
Differentially expressed genes (DEGs) from each of the MASLD *in vitro* model datasets were subjected to
gene set enrichment analysis (GSEA) using the "GSEA" function in R. The "msigdb" package in R was used
to extract predefined gene sets (GO:BP) from the Molecular Signatures Database. Gene Ontology (GO)
biological processes were enriched with a significance threshold of $p < 0.05$. This analysis was also
performed for the DEGs from the Govaere et al. dataset[17], stratified by the fibrosis stages described in their
study. The analyzed gene sets are summarized in Table 3.

Table 3: Datasets used for clinical benchmarking

MASLD *in vitro* model	Govaere et al Fibrosis Stage
OA PA vs HCM	Control
OA PA ResMyo vs HCM	F1

OA PA ResMyo PBMCs vs HCM	F2
	F3
	F4

Features of the FAD rescue dataset:

The FAD rescue bulk rna-seq dataset initially consisted of 12 samples (4x samples per condition) derived

from the *PNPLA3*-I148I line, see Table 4. Due to poor sample quality of one sample from the OA/PA +

Res/Myo + PBMCs + EGFP+ FAD condition which did not pass the sample QC, we had to remove it from

the analysis.

Table 4: Sample conditions used for bulk RNA-sequencing and downstream analysis of the FAD

rescue.

Name	Condition
OA PA ResMyo PBMCs + EGFP	iHeps exposed to the OA/PA, Res/Myo and PBMC disease model which were lentivirally transduced with a vehicle control
OA PA ResMyo PBMCs + EGFP + FAD	iHeps exposed to the OA/PA, Res/Myo and PBMC disease model which were lentivirally transduced with a vehicle control and treated with 50 µM of FAD
OA PA ResMyo PBMCs + *PPARGC1A* + FAD	iHeps exposed to the OA/PA, Res/Myo and PBMC disease model which were lentivirally transduced with *PPARGC1A*-expressing construct and treated with 50 µM of FAD

The processing and data preparation was carried out by Novogene´s bioinformatic services. The generation

of DEGs which were also used as an input for KEGG pathway analysis we carried out using the Novogene

magic web application for the following comparisons:

I.   OA PA ResMyo PBMCs + EGFP vs OA PA ResMyo PBMCs + EGFP + FAD

II.   OA PA ResMyo PBMCs + EGFP + FAD vs OA PA ResMyo PBMCs + *PPARGC1A* + FAD

<u>Shiny Go analysis</u>

For identifying the metabolites that are associated with the down-regulated genes in the MASLD *in vitro*

model we used the Shiny Go Web application[71]. We used the complete list of down-regulated DEGs from

the comparison of PNPLA3-I48I iHeps exposed to OA/PA+Res/Myo+PBMCs to HCM condition as an input.

We analyzed the enrichment for the human metabolome database using an FDR cut-off of 0.05 and

minimum pathway size of 10.

<u>Drug repurposing approaches in a clinical MASLD dataset</u>

Our computational drug repurposing pipeline has three approaches (Table 5): (1) Analysis of single cell and

single nucleus MASLD datasets to identify genes enriched in diseased hepatocytes and early stage MASLD

patients, (2) Weighted gene co-expression network analysis of fibrosis stages in MASH and (3) interrogation

of a biomedical knowledge graph to identify drug compounds associated with hepatic steatosis. We

implemented approaches (1 and 2) using a published clinical MASLD dataset (Govaere et al., 2020). For the

following sections and the results, we utilize the new nomenclature of MASLD. Thus, MASL, refers to NAFL,

MASLD to NAFLD and MASH to NASH in the respective data sets that use the old nomenclature as

identifier.

Table 5: Computational drug repurposing pipelines

Approach	Sequence of analysis
Single cell approach	1. MASLD Bulk RNA sequencing (early-stage vs control) 2. Single cell and single nucleus RNA-sequencing datasets 3. Protein-protein interaction networks 4. Drug-disease proximity analysis

WCGNA approach	1. MASLD Bulk RNA (fibrosis stages)
	2. Protein-protein interaction networks
	3. Drug-disease proximity analysis
Knowledge Graph approach	1. PrimeKG – Breadth-First Search (all shortest paths)
	2. PrimeKG – Drug proximity search

627
628
629 Re-analysis of a bulk RNA-sequencing MASLD dataset:
630 Publicly available bulk RNA sequencing data from frozen liver biopsies were downloaded from the NCBI
631 GEO repository (Home - GEO - NCBI (nih.gov), accession: GSE135251[17]). This dataset comprises 10
632 healthy controls, 51 MASL samples, and 155 MASH patients with varying stages of fibrosis.

633
To analyze this dataset, we utilised the Nextflow workflow system. We first used nf-core/fetchngs v 1.10.0
to fetch the metadata and raw FastQ files from the Gene Expression Omnibus. We then used nf-
core/rnaseq (version 3.12.0) to analyse the RNA sequencing data and produce a gene expression matrix.
Here we utilized STAR to map the raw FastQ reads to the reference genome (GRCh38.p14.) and project the
alignments onto the transcriptome followed by Salmon for quantification. Differential expression analysis
(DESeq2) was performed comparing MASL and healthy samples with an absolute log fold change greater
than 1 and a multiple testing corrected p-value lower than 0.05 was deemed significant. Visualisations
included volcano plots, MA plots, and heatmaps of the top DEGs.

642
Single-cell and single-nucleus RNA-seq MASLD datasets
To refine and validate the hepatocyte specificity of the bulk RNA-seq signature, four single-cell or single-
nucleus RNA-seq datasets were analyzed: GSE210501, GSE166504, GSE212837, and GSE189600. These
datasets include mouse models of MASLD as well as human liver samples at various stages of disease.
Each dataset was processed using the Seurat package (v4.3.0) following standard quality control
procedures, including filtering of cells with high mitochondrial content or low feature counts. Normalization,
identification of highly variable features, PCA, UMAP dimensionality reduction, and clustering were
performed. Hepatocytes were identified based on canonical markers such as human and mouse genes for
Alb, Cyp2e1, and Ttr. For each dataset, the refined MASLD gene signature was tested for differential
expression within hepatocyte populations comparing disease versus control conditions. AUC analyses and
UMAP-based visualizations were used to assess the discriminatory power of the signature, and expression
scores were computed based on average expression of the upregulated targets.

655
Protein-protein interaction (PPI) network construction in MASLD samples:
To identify an MASL signature, we restricted the analysis of the GSE135251 dataset to MASL patients and
healthy control samples. MASH patients were not considered as we wanted to identify the early changes
underpinning steatosis, prior to the onset of fibrosis and inflammation.
To prioritize functionally central regulators within the MASLD transcriptional landscape, we constructed a
protein–protein interaction (PPI) network using STRING v11.0 (combined score > 400). The input consisted
of upregulated hepatocyte-specific genes derived from the bulk analysis. Shortest paths were computed
between all gene pairs, yielding a high-confidence undirected network. Ensembl protein identifiers were
converted to HGNC gene symbols using a curated alias dictionary. Subcellular localization data from the
COMPARTMENTS database were used to annotate membrane and nuclear-associated genes.

666
Weighted gene co-expression network analysis segmented by MASH fibrosis stage:
Gene co-expression network analysis was performed using the WGCNA R package[72,73] (version 1.73). Since
the aim of this work was to construct disease specific co-expression network, the non-MASLD control
samples were not included in this analysis.
First, the co-expression similarity matrix, calculated using a Pearson correlation coefficient for all possible
pairs of genes, was transformed into an adjacency matrix using a soft-thresholding power β such that the
resulting network structure satisfied the scale-free topology criterion. For this study, $\beta = 28$ was chosen.
The adjacency matrix was then converted to a topological overlap matrix (TOM). Co-expression module
identification was accomplished with the function cutreeDynamic from the dynamicTreeCut R package
(version 1.63) using TOM dissimilarity (1-TOM) as the distance measure and a minimum module size cutoff
of 30. Genes that were not assigned to any co-expression module were located in the 'grey' module. The
resulting co-expression modules contained genes that were highly interconnected (co-expressed with each
other).

To evaluate the correlation of co-expression module to fibrosis stage, the 216 MASLD samples were
divided into the following subgroups using the clinical information provided by Govaere et al.[17]: MASLD (51
samples), MASH F0-F1 (34 samples), MASH F2 (53 samples), MASH F3 (54 samples) and MASH F4 (14
samples), where 'F' indicates the degree of fibrosis in each group, while MASLD patients displayed no
fibrosis. We then converted the fibrosis stage metadata into a continuous numerical vector. Specifically,
samples with no fibrosis were assigned a value of 0, samples with stage 1 fibrosis were assigned a value of
1, and so forth up to stage 4. The moduleEigengenes function (WGCNA R package) was used to calculate
module eigengenes.
We then calculated the Pearson correlation coefficients between module eigengene expression and fibrosis
stage, as well as its statistical significance, using functions from the WGCNA package. This approach
allowed us to quantify the strength and direction of the association between gene module expression and
fibrosis severity. Modules were considered significantly associated to fibrosis if p value < 0.05.

<u>Pathway enrichment analysis of WGCNA modules</u>
We performed overrepresentation analysis (ORA) to identify the main biological pathways representative of
each module. First, the ensembl IDs for the genes in each co-expression module were converted to entrez
IDs using the biomaRt R package (version 2.28.0). Enriched Gene Ontology terms, KEGG and Reactome
pathways were then identified using the gProfiler2 R package (version 0.2.2). Enriched pathways were
defined as those with FDR-adjusted p value < 0.05.

<u>WGCNA-guided disease network construction</u>
After having identified disease-relevant co-expression modules via WGCNA, we proceeded to expand such
networks using the Dijkstra algorithm to identify all shortest paths between any two genes in a selected co-
expression module. Each selected co-expression module was analyzed separately. This approach ensured
that the resulting drug predictions could be directly linked to a biological process affected in MASH.
First, the module genes were mapped to the human interactome. The human interactome was assembled
from the following databases: STRING, IntAct, BioGRID, HI-Union, Interactome3D, HINT, APID, BioPlex3.0,
SignaLink3.0. Were necessary, the databases were filtered to: include only interactions between human
proteins, include only interactions validated experimentally and remove self-loops (protein interaction with
itself. For data obtained from the string database, we also removed all interactions with confidence score <
0.4. This resulted in a network of 18748 proteins and 773143 edges. All shortest paths between all pairs of
module genes in the human interactome were found using the all_shortest_paths function from the
NetworkX python package (version 3.0).

<u>Topological analysis of the PPI and WCGNA disease networks</u>
Eigenvector centrality, degree, betweenness centrality and random walk with restart (RWR) were utilized to
identify key proteins in the MASL PPI network and in each WCGNA disease module. To identify significant
key proteins, permutation tests were performed 1000 times for each of the network centrality algorithms.
Each test generated a random network with a preserved degree distribution of the original disease network.
These cumulative results were used to calculate the empirical p-value of the network algorithm. Proteins
with empirical p value < 0.001 (from WGCNA module analysis) and empirical p value < 0.01 (from the single
cell approach analysis) for either centrality measure were considered key nodes. The R and Python igraph
packages were used for topological analysis respectively.

<u>Drug-disease proximity analysis</u>
FDA approved compounds and their interactions were sourced from ChEMBL[74], DrugBank[75], STITCH[76] and
the Barabási group[77]. The network-based distance between drug targets and key disease proteins in the
human interactome was calculated as described by Guney et al[19]. Given K, a set of disease proteins, T a set
of drug targets, and d(K, T), the shortest path length between nodes k ∈ K and t ∈ T in the human
interactome, we define:

$$d_c(K,T) = \frac{1}{||T||} \sum_{t \in T} min_{k \in K} d(K,T) \ (1)$$

The closest distance measure d(K, T) (average shortest path length between any drug target and its closest
disease target, eq. 1) was used here because it showed the best performance in drug-disease pair
prediction in the study by Guney et al[19].
To assess the statistical significance of the distance between a key protein and a drug dc(K, T), we
performed 1000 permutation tests, each time with two randomly selected gene sets. Selection of random
gene sets was performed so that the degree distribution of the random sets was similar to the real sets.
However, due to the scale-free nature of biological networks, there are only a few high-degree nodes in the
human interactome. To avoid repetitive selection of the same, high degree nodes during random selection,

we used a binning approach in which nodes within a certain degree interval were grouped together such
that there were at least 100 nodes in each bin. When selecting the random gene sets, we sampled nodes
from the same bins as the real drug/disease targets. The mean and standard deviation of the reference
distribution was then used to calculate the Z-score of the observed distance. Drugs with Z-score <= -1.96
(for WGCNA-driven drug discovery) and Z-score <= -2 (for PPI-driven drug discovery) were considered
significantly proximal.
745
<u>Network visualization and annotation</u>
Final network visualisations were generated using Cytoscape (v3.9.1), iGraph and NetworkX (v2.6.3) with
Force Directed, Fruchterman–Reingold and circular layouts. In the WGCNA network modules **(Fig. 4C-D)**,
nodes were colored by differential expression and betweenness centrality respectively. Differential
expression was conducted using edgeR (v4.4.2) comparing all MAFLD to control cases. Betweenness
centrality computation was performed in Cytoscape. To illustrate the interaction landscape surrounding
FAD, the top convergent candidate from our pipeline, a subnetwork was constructed including all key
regulators and their 1-hop neighbours. FAD was manually introduced into the network and connected to its
known targets. Nodes were coloured by relevance (the drug in green, direct targets in orange, background
in grey) and annotated using custom scripts. Figures were rendered at high resolution using CairoSVG and
Matplotlib.
757
758   <u>Knowledge graph analysis</u>
759   To enable graph traversal and algorithm calculation, we imported PrimeKG into Neo4j, a graph database
760   management system that stores data as nodes, edges, and their associated attributes (Neo4j Technology
761   Inc., 2015).
762   PrimeKG - Simple Graph Queries of complete graph network
To identify potential drug repurposing candidates, we performed Breadth-First Search (BFS) algorithm to
compute the shortest paths starting from our nodes of interest. In the first approach, we initiated the query
from the disease node "non-alcoholic steatohepatitis" (NASH, referring to the new nomenclature MASH),
identified genes associated with the disease, and then queried for drugs known to interact with these
genes. In a second approach, we began with the phenotype "hepatic steatosis" node and expanded our
search to nearby disease nodes connected to hepatic steatosis, extracting genes associated with these
diseases and identifying drugs that interact with them. In both queries, we excluded relationships that were
labelled "side effect" and "contraindication".
771   PrimeKG – Drug proximity scoring to disease-relevant nodes in disease subgraph
Creation of the Disease Subgraph for Non-Alcoholic Steatohepatitis (NASH):
Starting with the NASH node, we performed a three-hop traversal to expand the subgraph, which ensured a
more comprehensive representation of the network by incorporating interactions from nodes directly or
indirectly connected to NASH. To ensure the quality of the subgraph, we filtered the relationships by
excluding those related to "side effects" and "contraindications", which are not directly relevant to the
therapeutic context of NASH. We will henceforth refer to this subgraph as "nash_subgraph".
Graph Algorithms for Proximity Analysis:
To identify possible repurposing candidate drugs for NASH, we applied three graph centrality algorithms to
"nash_subgraph": PageRank (using a damping factor of 0.85 to ensure convergence and a maximum of
1,000 iterations), Betweenness Centrality, and Closeness Centrality using the Neo4j Graph Data Science
(GDS) library. For each measure, we filtered for Drug nodes, ranked them in descending order of PageRank
score to obtain one list, and retained the top 500 drugs for further analysis.
To obtain a prioritized drug list for downstream integration into the WGCNA and PPI-network based
pipelines, we carried out a union of the drugs from the lists returned from the simple graph query and graph
algorithm proximity calculations.
787
<u>Pediatric MASLD cohort Mass Spec analysis</u>
We validated the perturbations of FAD-dependent processes in pediatric MASLD patients using the cohort
described in Hudert et al[21]. The normalized liver proteomic profiles of the pediatric cohort (see clinical
characterization and methods in Hudert et al[21]) were grouped into two categories according to their NAS
scores 0-2 versus 5-8 for pathway analyses. For plotting individual proteins, the following three categories
were established: NAS 0-2, 3-4, and 5-8.
Pathway analyses: The log2 fold change between the two NAS groups was calculated for each protein
(70% valid values in at least one group, imputation of missing values from normal distribution) (Perseus
v1.6.0.2), which served as input to the String database https://string-db.org/. The annotated keywords
(Uniprot) pathway was chosen to identify significantly regulated pathways between the two NAS groups.
798
<u>Analysis of survival and liver-related events in adult MASLD cohort</u>

We analyzed the SteatoSITE cohort to correlate survival and hepatic events (decompensation) with
flavoproteome gene expression. Kaplan-Meier Plots were generated as described in Schwabe et al and
tested for significance using a log-rank test. Specifically, the survival plot for individual genes uses only
SteatoSITE biopsy cases with successful RNAseq (n = 508). For the predicted flavo-model, we performed
lasso-penalised cox regression including all genes from the flavoproteome for all cases with successful
RNAseq where there was no decompensation coding event prior to the biopsy (n = 446). For the
coefficients from the resulting 10-gene signature, we derived a single metric value for each case by
multiplying the coefficient by the cpm for the named gene and summing them. We then derived a single
cutpoint for that new metric that best divided patients into high and low risk of decompensation using the
surv_cutpoint function in R. Based on that, we then classified them into high and low risk groups to
generate the Kaplan-Meier plot.
811
812 <u>Quantification of intracellular FAD</u>
813 We quantified intracellular FAD levels using the fluorescence-based detection of the FAD assay kit (Abcam,
814 ab204710) according to the manufacturer's instructions. Before harvesting cells for the assay, we washed
815 3x times with PBS to ensure no significant amounts of exogenous FAD remained. We performed the assay
816 read-out using the Spectramax i3X Plate reader.
817
<u>Identification of compounds with similar network topology to FAD</u>
Given the impact of mitochondrial dysfunction in MASLD[78], we refined MASL signatures to a subset of the
MASLD DEG samples associated with mitochondrial structure or function. The list of mitochondrial-
associated genes was sourced from the MitoCarta 3.0 database[18]. We then selected compounds with
shared nodes to FAD. We further expanded the drug search to include 24,000 investigational and
experimental compounds in phase 1-3 clinical trials **(Supp. Table 10)**. From these, we prioritized norleucine
(Nor), aspirin (Asp), 3-Pyridin-4-yl-1H-indazol-5-ylamine dihydrochloride (3P), A-443654 and BCTC for
rescue experiments.
826
<u>Mechanistic rescue experiments</u>
For the rescue experiments, we co-exposed the cells to the indicated concentrations of FAD (Roth), Nor
(Thermo), Asp (Sigma), 3P (Sigma), A-443654 (biomol), BCTC (Sigma), plus the indicated injury condition
using the duration and concentrations described for the iHeps. Thus, the total exposure to each compound
was 72 h. After the co-exposure, we harvested the respective material for the analysis as indicated. BCTC
and A-443654 did not show significant differences, which is why they are omitted in the plot.
833
<u>Measurement of the oxygen consumption rate (OCR)</u>
For measuring the OCR in iHeps, we detached them from one full 6-well plate and seeded them on one full
Geltrex (Thermo) coated Seahorse Plate on day 28 of differentiation. For detachment, we washed iHeps
with PBS and then incubated the cells for at least 10 min with 10x TrypLE (Gibco) until they rounded up. We
added 5-fold volume of PBS with 20% KO-Serum, centrifuged the cells for 3 min at 200 g, and
resuspended them in HCM complete media containing all supplements as described. We first added 50 µl
of the cells for attachment, followed by 50 µl of more medium 5 h later. For testing Res and Myo, we
exposed cells to each factor individually or both combined for 3 days, compared to the vehicle only. For
testing the rescue effect by FAD or other compounds, we co-exposed iHeps and primary mouse
hepatocytes to the compound plus OA/PA and Res/Myo for three days. We measured OCR using the
Seahorse Extracellular Flux (XF)-96 Bioanalyzer from Agilent Technologies, as described previously[79].
Briefly, the XF basal medium from Agilent Technologies was supplemented with 10 mM glucose, 1 mM
pyruvate, and 1 mM glutamine. The electron transport chain complexes inhibitors (oligomycin (2 µM), FCCP
(1 µM), rotenone (1 µM), and antimycin (1 µM)) were titrated and delivered through injection at designated
ports A, B, and C of the Bioanalyzer. The resulting data were normalized to the protein content in each well
using the Pierce™ BCA assay (Thermo Fisher Scientific).
850
<u>Measurement of electron transport chain complex activities via respirometry-based assay</u>
The electron transport chain complex I, II, and IV activities were measured in frozen hepatocytes using
Seahorse XF-96 Extracellular Flux Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA) as described[80].
Primary mouse hepatocytes were treated with the respective treatments for 72 hours. After treatment, the
medium was removed, cells were washed with PBS three times. Then, the cells were harvested and frozen
at -70°C. On the day of assay, cells were homogenized in MAS buffer (70 mM sucrose, 220 mM mannitol, 5
mM KH2PO4, 5 mM MgCl2, 1 mM EGTA, 2 mM HEPES, pH 7.4) using 10 strokes in a glass Dounce
homogenizer. The homogenate was centrifuged at 200 g for 10 minutes at 4°C, and the supernatant was
collected. Protein concentration was measured using the Pierce™ BCA assay (Thermo Fisher Scientific).

For the Seahorse XF-96 assay, 15 µg of homogenate protein was loaded into each well in 20 µl of MAS
buffer. The plate was centrifuged at 2,000 g for 5 minutes at 4°C, after which 130 µl of MAS buffer
containing cytochrome c (10 µg/ml) was added to each well. Substrates were delivered through port A
(NADH, 1 mM; or succinate + rotenone, 5 mM + 2 µM). Inhibitors rotenone and antimycin A (2 µM and 4 µM,
respectively) were delivered through port B, TMPD + ascorbic acid (0.5 mM + 1 mM) through port C, and
azide (50 mM) through port D. The data was normalized to the proteins content. OCR signal after azide
injection was subtracted from NADH (complex I activity), succinate + rotenone (complex II activity), and
TMPD+ascorbic acid (complex IV activity).
868
Lipidomics
We harvested iHeps that we left untreated or exposed to Res/Myo and PBMCs with or without 50 µM of
FAD. We omitted exogenous FFAs to focus on cell-intrinsic changes. After exposure, we harvested the cells
by washing them twice with PBS and subsequently incubating them in 10X TryPLE. After the cells
dissociated from the dish, we inactivated TryPLE using PBS with 20% KO-Serum and centrifuged them for
3 min using 200 g. The Pellets were then stored in -80°C for 24 h until the analysis.
For the analysis cells were hydrolyzed with 100 µL 10 mol/L NaOH within 120 min at 80 °C. The samples
were neutralized with 100 µL acetic acid and filled up to 5 mL with methanol containing internal standards
(Cayman Chemical, Ann Arbor MI).
HPLC measurement was performed using an Agilent 1290 HPLC system with binary pump, autosampler
and column thermostat equipped with a Phenomenex Kinetex-C18 column 2.6 µm, 2.1 x 150 mm column
(Phenomenex, Aschaffenburg, DE) using a solvent system of acetic acid (0.05%) and acetonitrile. All
solvents and buffers in LC-MS-grade were purchased from VWR Germany
The solvent gradient started at 70 % acetonitrile and was increased to 98 % within 10 min and hold until 14
min with a flow rate of 0.4 mL/min and 5 µL injection volume.
The HPLC was coupled with an Agilent 6470 triplequad mass spectrometer with electrospray ionisation
source operated in negative selected ion mode.
886
Label-free brightfield imaging analysis
For the cell morphology analysis by label-free brightfield imaging using the VAIDR technology platform (TRI
Thinking Research Instruments GmbH, Hamburg, Germany), we took a minimum of 5 images per condition
as input.
Image segmentation:
The VAIDR image analysis software was used to develop a segmentation method to detect the relevant
sub-cellular structures of the hepatocytes on the label-free images. To this end, [six] image segments [of
512x512 pixels] were manually annotated using the software's label editor. For the annotation, the following
classes were used: Nucleus, membrane, cytoplasm, lipid, and mitochondria. The software then
automatically trained a segmentation algorithm based on the annotated images. This algorithm was applied
to all [96 full-size] images in the dataset, yielding one segmentation mask per image, which assigns each
pixel to one of the classes mentioned above.
Feature extraction:
Based on these masks and the original images, 185 numerical features were calculated as follows: For each
contiguous region, called "object", belonging to a particular class, the area, roundness, as well as the
image intensity and texture intensity, were calculated. [For example, groups of pixels of the "nucleus" class
represent the detected nuclei. For each such group, the area, roundness, intensity and texturation yield a
rich phenotypic profile of the respective nucleus.] The object-level results were aggregated per well by
calculating the mean and standard deviation of each parameter and each object class over all objects in a
well. The object count for each class was used as an additional feature.
Feature evaluation:
The resulting well-level phenotypic features were evaluated in two ways: (i) visually, by calculating the t-SNE
(1) plot. (ii) For each feature, a T-test was performed to assess whether untreated diseased cells could be
distinguished from FAD-treated diseased cells based on that feature alone. The T-test was perfomed using
the function ttest_ind from the Python package scipy.stats.
912
Lentivirus production and transduction experiments
We generated lentiviral particles by co-transfecting HEK293T cells with the plasmids VSV-G, pCMV8.9
(both provided by the Viral Core Facility of the Charité Universitätsmedizin Berlin) and either EGFP (cloned
into the pLX304 backbone) or *PPARGC1A* (#Addgene #143709). Two days after transfection, supernatant
was harvested, filtered and concentrated 20-fold using the Takara Lenti-X-Concentrator. Virus was
aliquoted and frozen in -80 until further use.
For transduction, iHeps were exposed to a low-level viral titer in their native medium for at least 4 hours,
with the addition of 5 µg/ml polybrene (approximately 0.1 functional MOI) 2 days before exposure to the

disease model. After transduction, the cells were treated with fresh medium and used for further
experiments as indicated.

FAD rescue in fetal liver-like organoids (FLOs)
For generating FLOs, we used the protocol from Quach et al[29] using the *PNPLA3*-I148M line. At day 18, we
added supernatants from healthy iHeps or supernatant from iHeps exposed to OA/PA, Res/Myo and
PBMCs in a to 2 / 5 ratio to the FLO medium, to which either a vehicle control or 50 μM FAD was freshly
added before the culture. Medium was changed every other day. After 120 h, cells were harvested for
analysis.

FAD rescue in primary immune cells
We freshly isolated healthy PBMCs as described above, used for disease modeling, and cultured 50000
cells in 100 μl of RPMI with 10% heat-inactivated FCS and Pen-Strep in a 96-well plate. To these, we either
added 50 μl of supernatant from healthy iHeps or from iHeps exposed to OA/PA, Res/Myo and PBMCs with
or without freshly added FAD (50 μM). We included 5 μg/ml of LPS (Stem Cell Technologies) as a positive
control, with or without 50 μM FAD. After 18 h, we harvested the cells for full-spectrum flow cytometry,
stained for the Panel in Table 6. Briefly, we transferred cells into FACS tubes and washed off the medium
using FACS buffer (PBS with 2 mM EDTA). After centrifugation, we discarded the supernatant, and the
viability dye was added to the residual volume. After 10 minutes, we added the antibody mix that we
prepared in blocking buffer. After 20 min, we washed the samples using FACS buffer. Subsequently, we
fixed the cells in 2% formalin for 10 min, which we washed off using FACS buffer. Then, samples were
acquired on a Cytek® Aurora Cytometer with 3 lasers (Violet, Blue, Red) and 38 detectors with the
Spectroflo® v3.0.3 software. Daily QC was performed according to the manufacturer's instructions. After
unmixing, cells were analyzed with the gating indicated.

Table 6: Flow cytometry-antibodies and dyes used in this study

Channel	Marker	Target	Clone	Manufacturer	Ordering No
BV421	CD80	Myeloid activation	2D10	BioLegend	305222
PacB	CD66b	Neutrophils	G10F5	BioLegend	305112
BV480	CD4	CD4TC	RPA-T4	BD	746541
V500	CD16	Neutrophils/monocytes	3G8	BD	561394
SV538	CD3	TC	UCHT1	BioLegend	300484
BV570	HLA-DR	MHC-II	L243	BioLegend	307638
BV711	CD40L	TC activation	24-31	BioLegend	310838
BV750	CD56	NK cells	5.1H11	BioLegend	362556
BB515	CD8	CD8TC	RPA-T8	BD	564526
Alexa532	CD45	Pan-immune	HI30	eBio	58-0459-42
PE-Cy5	CD19	B cells	HIB19	BioLegend	302210
PerCP-Cy5.5	CD14	Monocytes	M5E2	BioLegend	301824
PE-Fire810	CD86	Myeloid activation	IT2.2	BioLegend	305455
APC	CD40	Myeloid activation	5C3	BioLegend	334310
Al647	PD1	TC activation	NAT105	BioLegend	367420
Al700	CD11c	APC/DC	Bu15	BioLegend	337220
Zombie NIR	Viability			BioLegend	423106
APC Fire 810	CD25	TC activation	M-A251	BioLegend	356150

Multiplexed immunofluorescence (mIF) of organoids
mIF was performed on 5-μm thick formalin-fixed paraffin-embedded organoid sections as previously
described[81]. Briefly, antigen retrieval was performed in heated citrate buffer (pH=6.0). Antibodies were
eluted after each cycle of staining using a 2-mercaptoethanol/SDS (2-ME/SDS) buffer. Primary and
secondary antibodies used in this study are listed in Table 7. Images were acquired on a Zeiss
AxioObserver 7 and processed in FIJI, as previously reported[82].

Table 7: Antibodies used for mIF

Marker	Source	Identifier
Rabbit anti-CD31	abcam	Cat#ab28364
Mouse anti-HepPar1	Dako	Cat#OCHIE5
Goat anti-IBA-1	abcam	Cat#ab5076
Goat anti-PDGFRA	R&D Systems	Cat#AF-307-SP
Donkey anti-Goat IgG Alexa Fluor 488	Sigma-Aldrich	Cat#SAB4600032
Donkey anti-Mouse IgG Alexa Fluor 647	Invitrogen	A32787
Goat anti-Rabbit IgG Alexa Fluor 750	Cell Signaling	Cat#4418S

Statistical testing

We performed data visualization and statistical significance assessment using RStudio or GraphPad Prism (Version 10). We used the statistical tests as indicated, two-sided if not stated otherwise.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

The main data supporting the results in this study are available within the Article and its Supplementary Information. Raw data of the *in vitro* transcriptomics will be provided to the reviewers if requested and uploaded to a public repository upon publication. The SteatoSITE transcriptomic data are deposited in the European Nucleotide Archive (https://www.ebi.ac.uk/ena; study accession number: PRJEB58625). Gene expression data are also freely available for user-friendly interactive browsing online at https://shiny.igc.ed.ac.uk/SteatoSITE_gene_explorer/. Full data access can be requested at https://regeneration-repair.ed.ac.uk/research/research-resources/steatosite. Mass Spec raw data can be found as indicated. Proteomic data from the pediatric cohort was utilized from Hudert et al[21].

References

1. Loewa, A., Feng, J. J. & Hedtrich, S. Human disease models in drug development. *Nat. Rev. Bioeng.* **1**, 545–559 (2023).

2. Cerneckis, J., Cai, H. & Shi, Y. Induced pluripotent stem cells (iPSCs): molecular mechanisms of induction and applications. *Signal Transduct. Target. Ther.* **9**, 112 (2024).

3. Younossi, Z. M. *et al.* The global epidemiology of nonalcoholic fatty liver disease (NAFLD) and nonalcoholic steatohepatitis (NASH): a systematic review. *Hepatology* **77**, 1335–1347 (2023).

4. Le, P. *et al.* Estimated Burden of Metabolic Dysfunction–Associated Steatotic Liver Disease in US Adults, 2020 to 2050. *JAMA Netw. Open* **8**, e2454707 (2025).

5. Koyama, Y. & Brenner, D. A. Liver inflammation and fibrosis. *J. Clin. Invest.* **127**, 55–64 (2017).

6. Buzzetti, E., Pinzani, M. & Tsochatzis, E. A. The multiple-hit pathogenesis of non-alcoholic fatty liver disease (NAFLD). *Metabolism* **65**, 1038–1048 (2016).

7.   Hendriks, D. *et al.* Engineered human hepatocyte organoids enable CRISPR-based target discovery and

drug screening for steatosis. *Nat. Biotechnol.* **41**, 1567–1581 (2023).

8.   Ajmera, V. *et al.* Novel plasma biomarkers associated with liver disease severity in adults with

nonalcoholic fatty liver disease. *Hepatology* **65**, 65–77 (2017).

9.   García, P. S., Cabbabe, A., Kambadur, R., Nicholas, G. & Csete, M. Elevated Myostatin Levels in Patients

with Liver Disease: A Potential Contributor to Skeletal Muscle Wasting. *Anesth. Analg.* **111**, 707–709

(2010).

10.   Delogu, W. *et al.* Myostatin regulates the fibrogenic phenotype of hepatic stellate cells via c-jun N-

terminal kinase activation. *Dig. Liver Dis.* **51**, 1400–1408 (2019).

11.   Wilkes, J. J., Lloyd, D. J. & Gekakis, N. Loss-of-Function Mutation in Myostatin Reduces Tumor Necrosis

Factor α Production and Protects Liver Against Obesity-Induced Insulin Resistance. *Diabetes* **58**, 1133–

1143 (2009).

12.   Jeon, J., Lee, S. H., Kim, D. & Sung, J. H. In vitro hepatic steatosis model based on gut–liver-on-a-chip.

Biotechnol. Prog. **37**, e3121 (2021).

13.   Ricchi, M. *et al.* Differential effect of oleic and palmitic acid on lipid accumulation and apoptosis in

cultured hepatocytes. *J. Gastroenterol. Hepatol.* **24**, 830–840 (2009).

14.   Carpino, G. *et al.* Increased Liver Localization of Lipopolysaccharides in Human and Experimental

NAFLD. *Hepatology* **72**, 470–485 (2020).

15.   An, L. *et al.* The Role of Gut-Derived Lipopolysaccharides and the Intestinal Barrier in Fatty Liver

Diseases. *J. Gastrointest. Surg.* **26**, 671–683 (2022).

16.   Baumeier, C. *et al.* Elevated hepatic DPP4 activity promotes insulin resistance and non-alcoholic fatty

liver disease. *Mol. Metab.* **6**, 1254–1263 (2017).

17.   Govaere, O. *et al.* Transcriptomic profiling across the nonalcoholic fatty liver disease spectrum reveals

gene signatures for steatohepatitis and fibrosis. *Sci. Transl. Med.* **12**, eaba4448 (2020).

18.   Rath, S. *et al.* MitoCarta3.0: an updated mitochondrial proteome now with sub-organelle localization

and pathway annotations. *Nucleic Acids Res.* **49**, D1541–D1547 (2021).

19. Guney, E., Menche, J., Vidal, M. & Barábasi, A.-L. Network-based in silico drug efficacy screening. *Nat.*

Commun. **7**, 10331 (2016).

20. Lienhart, W.-D., Gudipati, V. & Macheroux, P. The human flavoproteome. *Arch. Biochem. Biophys.* **535**,

150–162 (2013).

21. Hudert, C. A. *et al.* Genetic determinants of steatosis and fibrosis progression in paediatric non-

alcoholic fatty liver disease. *Liver Int.* **39**, 540–556 (2019).

22. Kendall, T. J. *et al.* An integrated gene-to-outcome multimodal database for metabolic dysfunction-

associated steatotic liver disease. *Nat. Med.* **29**, 2939–2953 (2023).

23. Ke, C. *et al.* FMO2 ameliorates nonalcoholic fatty liver disease by suppressing ER-to-Golgi transport of

SREBP1. *Hepatology* **81**, 181–197 (2025).

24. Koliaki, C. *et al.* Adaptation of Hepatic Mitochondrial Function in Humans with Non-Alcoholic Fatty Liver

Is Lost in Steatohepatitis. *Cell Metab.* **21**, 739–746 (2015).

25. Pérez-Carreras, M. *et al.* Defective Hepatic Mitochondrial Respiratory Chain in Patients With

Nonalcoholic Steatohepatitis. *Hepatology* **38**, 999–1007 (2003).

26. Simopoulos, A. P. The importance of the ratio of omega-6/omega-3 essential fatty acids. *Biomed.*

Pharmacother. **56**, 365–379 (2002).

27. Arconzo, M. *et al.* Hepatic-specific Pgc-1α ablation drives fibrosis in a MASH model. *Liver Int.* **44**, 2738–

2752 (2024).

28. Subudhi, S. *et al.* Distinct Hepatic Gene-Expression Patterns of NAFLD in Patients With Obesity.

Hepatol. Commun. **6**, 77–89 (2022).

29. Rezvani, M. *et al.* Fetal Liver-like Organoids Recapitulate Blood-Liver Niche Development and

Multipotent Hematopoiesis from Human Pluripotent Stem Cells. Preprint at

https://doi.org/10.1101/2024.10.11.617794 (2024).

30. Zhang, C. & Yang, M. Targeting T Cell Subtypes for NAFLD and NAFLD-Related HCC Treatment: An

Opinion. *Front. Med.* **8**, 789859 (2021).

31. Tacke, F. *et al.* EASL–EASD–EASO Clinical Practice Guidelines on the management of metabolic

dysfunction-associated steatotic liver disease (MASLD). *J. Hepatol.* **81**, 492–542 (2024).

32. Simon, T. G. *et al.* Aspirin for Metabolic Dysfunction–Associated Steatotic Liver Disease Without

Cirrhosis: A Randomized Clinical Trial. *JAMA* **331**, 920 (2024).

33. Zareifi, D. S. *et al.* A network-based computational and experimental framework for repurposing

compounds toward the treatment of non-alcoholic fatty liver disease. *iScience* **25**, 103890 (2022).

34. Kusumoto, D. *et al.* Anti-senescent drug screening by deep learning-based morphology senescence

scoring. *Nat. Commun.* **12**, 257 (2021).

35. Seagle, H. M. *et al.* Genomics-Informed Drug Repurposing Strategy Identifies Novel Therapeutic Targets

for Metabolic Dysfunction-Associated Steatotic Liver Disease. Preprint at

https://doi.org/10.1101/2025.02.18.25321035 (2025).

36. Wen, F. *et al.* Acute Elevated Resistin Exacerbates Mitochondrial Damage and Aggravates Liver

Steatosis Through AMPK/PGC-1α Signaling Pathway in Male NAFLD Mice. *Horm. Metab. Res.* **53**, 132–

144 (2021).

37. Yang, H.-M. *et al.* Resistin impairs mitochondrial homeostasis via cyclase-associated protein 1-mediated

fission, leading to obesity-induced metabolic diseases. *Metabolism* **138**, 155343 (2023).

38. Zhou, L. *et al.* Resistin reduces mitochondria and induces hepatic steatosis in mice by the protein kinase

C/protein kinase G/p65/PPAR gamma coactivator 1 alpha pathway. *Hepatology* **57**, 1384–1393 (2013).

39. Liu, X., Pan, J. P., Bauman, W. A. & Cardozo, C. P. AdipoRon prevents myostatin-induced upregulation of

fatty acid synthesis and downregulation of insulin activity in a mouse hepatocyte line. *Physiol. Rep.* **7**,

(2019).

40. Acquarone, E., Monacelli, F., Borghi, R., Nencioni, A. & Odetti, P. Resistin: A reappraisal. *Mech. Ageing*

Dev. **178**, 46–63 (2019).

41. Groeger, M. *et al.* Modeling and therapeutic targeting of inflammation-induced hepatic insulin

resistance using human iPSC-derived hepatocytes and macrophages. *Nat. Commun.* **14**, 3902 (2023).

42. Yang, J. *et al.* Integrated-gut-liver-on-a-chip platform as an in vitro human model of non-alcoholic fatty

liver disease. *Commun. Biol.* **6**, 310 (2023).

43. Kozyra, M. *et al.* Human hepatic 3D spheroids as a model for steatosis and insulin resistance. *Sci. Rep.*

8, 14297 (2018).

44. Kimura, M. *et al.* En masse organoid phenotyping informs metabolic-associated genetic susceptibility to

NASH. *Cell* **185**, 4216-4232.e16 (2022).

45. Balasubramaniam, S., Christodoulou, J. & Rahman, S. Disorders of riboflavin metabolism. *J. Inherit.*

Metab. Dis. **42**, 608–619 (2019).

46. Chen, Y.-S. *et al.* Effect of Vitamin B2 supplementation on migraine prophylaxis: a systematic review

and meta-analysis. *Nutr. Neurosci.* **25**, 1801–1812 (2022).

47. Park, K. *et al.* Flavin Adenine Dinucleotide Depletion Caused by electron transfer flavoprotein subunit

alpha Haploinsufficiency Leads to Hepatic Steatosis and Injury in Zebrafish. *Hepatol. Commun.* **5**, 976–

991 (2021).

48. Tang, J. *et al.* Severe riboflavin deficiency induces alterations in the hepatic proteome of starter Pekin

ducks. *Br. J. Nutr.* **118**, 641–650 (2017).

49. Wang, Y. *et al.* Effects of riboflavin deficiency and high dietary fat on hepatic lipid accumulation: a

synergetic action in the development of non-alcoholic fatty liver disease. *Nutr. Metab.* **21**, 1 (2024).

50. Masschelin, P. M. *et al.* Vitamin B2 enables regulation of fasting glucose availability. *eLife* **12**, e84077

(2023).

51. Jusko, W. J. & Lewis, G. P. Effect of hepatic disease on liver flavin and protein levels. *Am. J. Clin. Nutr.*

25, 265–268 (1972).

52. Zhang, D. *et al.* Selenium and vitamin B $_2$ cosupplementation alleviates high-fat-diet-induced

nonalcoholic fatty liver disease in rats by regulating hepatic lipid metabolism. *Eur. J. Lipid Sci. Technol.*

126, 2300020 (2024).

53. Kajiwara, M. *et al.* Donor-dependent variations in hepatic differentiation from human-induced

pluripotent stem cells. *Proc. Natl. Acad. Sci.* **109**, 12538–12543 (2012).

54. Seifert*, J. *et al.* Low nanomolar Triiodothyronine Concentrations Drive Further Maturation of hiPSC-

Derived Hepatocyte-Like Cells. Preprint at https://doi.org/10.21203/rs.3.rs-6595009/v1 (2025).

55. Ma, H. *et al.* The nuclear receptor THRB facilitates differentiation of human PSCs into more mature

hepatocytes. *Cell Stem Cell* **29**, 795-809.e11 (2022).

56.   Nell, P. *et al.* Identification of an FXR-modulated liver-intestine hybrid state in iPSC-derived hepatocyte-

like cells. *J. Hepatol.* **77**, 1386–1398 (2022).

57.   Charni-Natan, M. & Goldstein, I. Protocol for Primary Mouse Hepatocyte Isolation. *STAR Protoc.* **1**,

100086 (2020).

58.   Horner, R. *et al.* Impact of Percoll purification on isolation of primary human hepatocytes. *Sci. Rep.* **9**,

6542 (2019).

59.   Kwon, Y. *et al.* Induction of steatosis in primary human hepatocytes recapitulates key

pathophysiological aspects of metabolic dysfunction-associated steatotic liver disease. *J. Hepatol.* **82**,

18–27 (2025).

60.   Beier, N. *et al.* Adaptation of Dinoroseobacter shibae to oxidative stress and the specific role of RirA.

PloS One **16**, e0248865 (2021).

61.   Lerch, M. F. *et al.* A non-coding RNA from the intercellular adhesion (ica) locus of Staphylococcus

epidermidis controls polysaccharide intercellular adhesion (PIA)-mediated biofilm formation. *Mol.*

Microbiol. **111**, 1571–1591 (2019).

62.   Schwanhäusser, B. *et al.* Global quantification of mammalian gene expression control. *Nature* **473**,

337–342 (2011).

63.   Perez-Riverol, Y. *et al.* The PRIDE database at 20 years: 2025 update. *Nucleic Acids Res.* **53**, D543–D553

(2025).

64.   Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-aware

quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).

65.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. *Genome Biol.* **15**, 550 (2014).

66.   Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* **28**, 27–30

(2000).

67.   Ulgen, E., Ozisik, O. & Sezerman, O. U. pathfindR: An R Package for Comprehensive Identification of

Enriched Pathways in Omics Data Through Active Subnetworks. *Front. Genet.* **10**, 858 (2019).

68. Oughtred, R. *et al.* The BioGRID interaction database: 2019 update. *Nucleic Acids Res.* **47**, D529–D541

(2019).

69. Ideker, T., Ozier, O., Schwikowski, B. & Siegel, A. F. Discovering regulatory and signalling circuits in

molecular interaction networks. *Bioinformatics* **18**, S233–S240 (2002).

70. Gene Ontology Consortium. The Gene Ontology (GO) database and informatics resource. *Nucleic Acids*

Res. **32**, 258D – 261 (2004).

71. Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals and plants.

Bioinformatics **36**, 2628–2629 (2020).

72. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network analysis. *BMC*

Bioinformatics **9**, 559 (2008).

73. Zhang, B. & Horvath, S. A general framework for weighted gene co-expression network analysis. *Stat.*

Appl. Genet. Mol. Biol. **4**, Article17 (2005).

74. Mendez, D. *et al.* ChEMBL: towards direct deposition of bioassay data. *Nucleic Acids Res.* **47**, D930–

D940 (2019).

75. Wishart, D. S. *et al.* DrugBank 5.0: a major update to the DrugBank database for 2018. *Nucleic Acids*

Res. **46**, D1074–D1082 (2018).

76. Szklarczyk, D. *et al.* STITCH 5: augmenting protein-chemical interaction networks with tissue and

affinity data. *Nucleic Acids Res.* **44**, D380-384 (2016).

77. Cheng, F., Kovács, I. A. & Barabási, A.-L. Network-based prediction of drug combinations. *Nat. Commun.*

10, 1197 (2019).

78. Myint, M. *et al.* Inflammatory signaling in NASH driven by hepatocyte mitochondrial dysfunctions. *J.*

Transl. Med. **21**, 757 (2023).

79. Kumar, P., Hassan, M., Tacke, F. & Engelmann, C. Delineating the heterogeneity of senescence-induced-

functional alterations in hepatocytes. *Cell. Mol. Life Sci.* **81**, 200 (2024).

80. Juanola, O. *et al.* Intestinal microbiota drives cholestasis-induced specific hepatic gene expression

patterns. *Gut Microbes* **13**, 1911534 (2021).

81. Guillot, A., Kohlhepp, M., Bruneau, A., Heymann, F. & Tacke, F. Deciphering the Immune

Microenvironment on A Single Archival Formalin-Fixed Paraffin-Embedded Tissue Section by An

Immediately Implementable Multiplex Fluorescence Immunostaining Protocol. *Cancers* **12**, 2449

(2020).

82. Guillot, A. *et al.* Mapping the hepatic immune landscape identifies monocytic macrophages as key

drivers of steatohepatitis and cholangiopathy progression. *Hepatology* **78**, 150–166 (2023).

Acknowledgements
We thank the members of the BIH Core Unit pluripotent Stem cells and Organoids, specifically Valeria
Fernandez Vallone, Janine Cernoch and Claudia Schaar for their technical support. We also thank Janine
Altmüller and Tatiana Borodina of the MDC/BIH Genomics Platform, Berlin for technical support.
Additionally, we would like to thank the HPC for Research/Clinic cluster at the Berlin Institute of Health for
providing data storage. The authors would like to acknowledge Novogene UK (Munich, Germany) for
providing bulk RNA sequencing and bioinformatics analysis services.
We are also grateful for the support provided by all the current and former members of the Rezvani Lab,
specifically Natalia Martagón Calderón and Reinhild Dünnebacke. We would also like to thank the following
people for sharing their expertise: Marième Gueye, Amr Al Shebel, Tilman Breiderhoff, Kerstin Sommer,
Sandra Gerstenberg, Verena Klämbt, Marah Alhabahbeh, Adrien Guillot, Hanyang Liu, Tian Lan and Guo
Yin. Further, we want to thank Aras Mattis and lab members from the University of California San Francisco
for discussion. Furthermore, we would like to extend our deepest gratitude to the individuals who have
donated materials to this biomedical research.

Author contributions
M.R. and Na.H. conceived the study. M.R., J.W., Na.H., F.B., G.Y. and A.C. designed and supervised the
study. K.L. and H.S. contributed to protocol establishment. Ni.H., I.S. and P.T. provided and isolated clinical
specimens. P.K. and C.E. contributed SeaHorse analysis. S.Q. and M.M. contributed to *in vitro* FLO and
iHep assays. M.K. and S.E. contributed the Mass spec analysis. B.C. established the label-free AI
brightfield imaging analysis. M.Ro. contributed the Lipidomics analysis. D.M. and C.H. contributed the
characterization and proteomic analysis of the pediatric MASLD co-hort. I.L. and L.H. contributed with the
full spectrum flow cytometry analysis. T.K. and J.F. analyzed the SteatoSITE co-hort. F.T., P.R. and P.B.
participated in discussions and provided intellectual input. J.W., M.R., Na.H., F.B., G.Y. and A.C. wrote the
paper.

Funding
The authors of this scientific article would like to acknowledge the funding provided by the Deutsche
Forschungsgemeinschaft (DFG) Emmy Noether Programme to M. R. (RE 3749/2-1) and the Young
Investigator Award awarded by Charite 3R to J.W. The Berlin-based authors (especially M.R., J.W. and F.T.)
would also like to acknowledge the "CRC/TR 412"/"TRR 412" funded by the DFG. Na.H. is funded by
LifeArc. SteatoSITE was funded by Innovate UK (Precision medicine: impacting through innovative
technology - Ref: TS/R017581/1). P.R. is funded by an MRC Senior Clinical Fellowship (MR/W015919/1).

Competing interests
Na.H., M.R. and J.W. are inventors of the patent application EP24206430. Na.H. is a cofounder of KURE.ai
and CardiaTec Biosciences. J.A.F serves as a consultant or advisory board member for Stellaris, Resolution
Therapeutics, Kynos Therapeutics, Ipsen, River 2 Renal Corp., Stimuliver, and Global Clinical Trial Partners,
has received speaker fees from HistoIndex and Resolution Therapeutics, and research grant funding from
GlaxoSmithKline and Genentech. P.R. has served as a consultant for MSD and Macomics and has received
research support from Genentech, Intercept and Neogenomics.

Code availability
Publicly available packages were used in all analyses, all of which are referenced in the Methods section
along with the specific parameters applied.

1237
Supplementary Information
Supplementary Tables
Supplementary Table 1: KEGG pathways perturbed in PNPLA3-I148I iHeps exposed to different MASLD
models
Supplementary Table 2: Detected human proteins of Mass Spec analysis of conditioned medium from
PNPLA3-I148I iHeps exposed to different MASLD models.
Supplementary Table 3: PathfindR analysis for identifying cell line specific or shared pathways in iHeps
exposed to the 3-Hit MASLD model
Supplementary Table 4: Compound prediction derived from PPIs
Supplementary Table 5: Compound prediction derived from WGCNA
Supplementary Table 6: Compound prediction derived from KG
Supplementary Table 7: Compounds predicted in multiple computational approaches
Supplementary Table 8: Predictive 10-Gene flavo-signature
Supplementary Table 9: Features derived by label-free brightfield imaging different between diseased cells
with or without FAD^H assessed for significance
Supplementary Table 10: Flavo-active compounds
1254
1255 **Supplementary Figures:**

1257

Supplementary Fig. 1: Characterizing the MASLD drug testing model in stem cell-derived hepatocytes (iHeps). A) MASLD Injury index in PNPLA3-I148M iHeps exposed to different MASLD models. All parameters were standardized, ranging from Healthy (0%) to the most complex disease condition including all disease triggers (100%). Inflammatory (Infla.) gene expression includes the genes *TNFA*, *CXCL8*, *CXCL1* and *CCL2*. $\Delta\Psi m$ = mitochondrial membrane potential **B)** Venn diagram displaying the number of differentially expressed genes (DEGs) from bulk RNA sequencing across disease conditions relative to the untreated control. Circle sizes are based on the log2-transformed DEG number. **C)** KEGG Pathway enrichment analysis for the up- and down-regulated genes in *PNPLA3*-I148I iHeps, for the 2-Hit and 3-Hit MASLD model. The 1-Hit model did not have enough DEGs for enrichment analysis and is therefore not shown. **D)** Mitotracker signal of *PNPLA3*-I148M iHeps treated with resistin (Res), myostatin (Myo), or both. Significance was assessed using multiple t-test, corrected according to the Benjamini-Hochberg method. Each dot represents an independent sample **F)** Heatmap of the mean IBAQ values of the identified human proteins. Proteins were identified in the supernatants of untreated iHeps and iHeps exposed to different stimuli (OA/PA = oleic and palmitic acid, Res/Myo, PBMCs = peripheral blood mononuclear cells) using LC-MS/MS. Heatmap shows the mean IBAQ values of n = 2 independent wells (with the exception of untreated iHeps (n=1)) (see **Supp. Table 2**).

Supplementary Fig. 2: Identifying donor specific effects in stem cell-derived hepatocytes (iHeps) with or without genetic MASLD predisposition A) Mitotracker signal in *PNPLA3*-I148I and -I148M iHeps, normalized to the respective untreated control. Significance was assessed using a t-test. Each dot represent an independent sample. **B)** Steatotic area in *PNPLA3*-I148I and -I148M iHeps, normalized to the mitochondrial area. Significance was assessed using a t-test. Each dot represent an independent sample. **C)** PathfindR analysis for enriched Pathways in the 3-Hit disease model specific the *PNPLA3*-I148I or -148M iHeps.

1300
1301

Supplementary Fig. 3: Validation of disease signature, module enrichment, and compound convergence across computational pipelines. Supporting analyses for transcriptomic signature refinement, co-expression module evaluation, and compound overlap across the PPI, WGCNA, and KG pipelines. **A)** UMAP projection of bulk RNA-seq samples (GSE135251) based on global gene expression profiles. Samples cluster by condition, with MASL (red) clearly separated from control (black), indicating robust transcriptomic differences. **B)** Volcano plot showing differential expression analysis between MASL and control samples. Significantly upregulated (red) and downregulated (blue) genes are highlighted (adjusted $p < 0.05$), with log2 fold change on the x-axis and –log10 p-value on the y-axis. **C)** Receiver operating characteristic (ROC) curve demonstrating the classification performance of the refined hepatocyte-specific MASL signature. The area under the curve (AUC) is 0.929, indicating strong discriminative power. **D)** UMAP embedding of bulk RNA-seq samples based on the expression of 21 key network nodes identified from the PPI analysis. The filtered signature maintains separation between MASL (red) and control (black) groups. **E)** KEGG pathway enrichment analysis of the 21 key PPI nodes. The only significantly enriched term was "Non-alcoholic fatty liver disease," supporting the specificity of the derived subnetwork. **F)** Top 20 enriched Gene Ontology (GO) terms for WGCNA module ME4, which was positively correlated with fibrosis severity. Terms include extracellular matrix organization, integrin signaling, and cellular migration. Bars are ranked by gene ratio and colored by adjusted p-value. **G)** Schematic overview of the PrimeKG graph schema showing major node types (e.g., Drug, Disease, Gene/Protein, Exposure, Phenotype, etc.) and example edge types (e.g., "interacts with," "associated with," "phenotype present"). This reflects the ontology structure used in KG-based query design. **H)** Venn diagram summarizing the overlap among predicted compounds from the three computational pipelines. Out of 1,199 unique compounds, three—flavin adenine dinucleotide (FAD), pyridoxal phosphate (PLP), and NADH—were identified by all three methods (KG, WGCNA, and PPI).

1347

Supplementary Fig. 4: Perturbance and manipulation of FAD homeostasis in MASLD A) Heatmap of normalized gene counts showing clustering of FAD-dependent protein gene expression across different MASLD models in an independent differentiation of PNPLA3-I148I iHeps **B)** DEGs the 3-Hit *in vitro* disease model in relation to healthy control filtered for the human flavoproteome. **C)** Abundance of *FMO3*, *FMO5* and *SARDH* proteins in the pediatric MASLD cohort described by Hudert et al. Significance was assessed using multiple t-tests, corrected by the Benjamini-Hochberg method. Each dot represents and independent patient **D)** Survival analysis of FMO-gene expression and human MASLD patients from the SteatoSITE co-hort. **E)** Survival analysis of FAD-homeostasis gene expression and human MASLD patients from the SteatoSITE co-hort. **F)** Concentration of FAD in iHeps exposed to the 3-Hit MASLD model after 3-day treatment with or without 50 μM of FAD. Each dot represents an independent sample.

1348

Supplementary Fig. 5: Characterizing the MASLD alleviating effects of FAD. A) Quantification of the steatotic area normalized to the nucleus count in iHeps exposed to the 2-Hit MASLD model with different concentrations of FAD. Significance was used using a one-sided one sample t-test using the 100% vehicle control marked by the dotted line as a reference. Each dot represents an independent sample **B)** Mitochondrial complex activity (I, II and IV) in mouse hepatocytes exposed to the 2 Hit MASLD model with or without 50 μM FAD[H] treatment. Significance was assessed using multiple t-test, corrected according to the Benjamini-Hochberg method. Each dot represents an independent sample **C)** Scheme, representative images and quantification of mitochondrial membrane potential **(D)** and steatotic area **(E)** for validating FAD effects in primary human hepatocytes exposed to OA/PA, Res/Myo and the pro-inflammatory cytokines IL1B and TNFA. Mitochondrial membrane potential and steatotic area were normalized to the Vehicle control (plus steatotic area was normalized to the nucleus count). Each dot represents an independent sample **F)** Concentration of AST in culture supernatant from iHeps exposed to the 3-Hit disease model, with or without 50 μM of FAD[H]. Cells were transduced with EGFP before disease modeling. Each dot represents an independent sample **G)** Lipidomic heatmap analysis of iHeps either untreated (healthy control) or exposed to the disease trigger Res/Myo and trans-well PBMC co-culture with or without FAD[H] (50 μM). Fatty acid species abundance was normalized by protein concentration and to total amount of measured fatty acid content. Heatmap shows the mean of n = 4-6 independent samples.

1392

Supplementary Fig. 6: Label-free brightfield imaging analysis for detecting rescue effects of FAD.
A) Scheme of label-free brightfield imaging analysis. **B)** tSNE plot of iHeps exposed to Res/Myo and PBMCs with or without FAD^H (50 μM) or being left untreated. **C)** Top 5 morphological parameter different between iHeps exposed to Res/Myo + PBMCs with or without FAD treatment, assessed for significance using t-tests. Each dot represents an independent sample.

Supplementary Fig. 7: Characterizing changes and additional rescuing effects of *PPARGC1A* and its overexpression in MASLD iHeps. A) Gene expression levels of *PPARGC1A* in iHeps exposed to different PBMC co-culture conditions, normalized to untreated cells. Dots represent independent samples. Significance was assessed using one-way anova with Tukey-Post Hoc **B)** Gene expression levels of *PPARGC1A* in iHeps exposed to OA/PA or OA/PA with Res/Myo. Gene expression levels were normalized to the untreated control. Dots represent independent samples. **C)** Expression levels of mitochondrial complex genes in iHeps transduced with *PPARGC1A* or EGFP control assessed via qPCR. Significance was assessed using t-tests. Dots represent independent samples **D)** Schematic of the RNA seq experiment analyzing iHeps transduced with *PPARGC1A* or EGFP control, exposed to the 3-Hit disease model, with or without 50 μM FAD^H. **E)** PCA plot, displaying all tested conditions. **F)** Top KEGG pathways enriched in iHeps exposed to FAD^H plus *PPARGC1A overexpression* relative to FAD^H-treated EGFP control. **G)** Steatotic area in iHeps exposed to the 3-Hit disease model, transduced with EGFP or *PPARGC1A* with different concentrations of FAD. Statistical significance was assessed using a t-test. Dots represent independent samples. Steatotic area was normalized to nuclei count and is relative to the non-transduced vehicle control of each condition.

1463

Supplementary Fig. 8: Characterizing FAD effects in multi-cellular fetal liver organoids (FLOs). A) Scheme of evaluating anti-TGFB effects of FAD^H in FLOs exposed to supernatant from injured iHeps. **B)** Brightfield images of FLOs assessing encapsulation, treated with conditioned medium from healthy iHeps or iHeps undergoing the 3-hit MASLD injury. **C)** Immunofluorescent images of FLOs, showing the abundance of hepatocytes (HepPar1$^+$), stellate cells (PDGFRA$^+$), endothelial cells (CD31$^+$) and macrophages (IBA1$^+$). **D)** Gene expression analysis of fibrosis signature genes in FLOs exposed to conditioned medium from healthy of injured iHeps (3-Hit MASLD model) with or without FAD^H (50 μM). For each gene, each dot represents an independent organoid. Significance was assessed using multiple t-test corrected according to the Benjamini-Hochberg method.

1482

Supplementary Fig. 9: Characterizing anti-inflammatory effects by FAD^H in primary human immune cells A) Full spectrum flow cytometry gating strategy for CD4 and CD8 T-cell characterization in PBMCs isolated from a healthy individual **B)** Representative dot plots of T-cell activation marker in CD4$^+$ and CD8$^+$ T-cells stimulated with iHep supernatant or LPS exposure with or without FAD^H. **C)** Quantification of T-cells double positive for the activation marker PD1 and CD25 after exposure to iHep supernatants with or without FAD^H. Multiple t-test to assess significance, corrected according to the Benjamini-Hochberg method. **D)** Quantification of T-cells double positive for the activation marker PD1 and CD25 after exposure to LPS with or without FAD^H. Multiple t-test to assess significance, corrected according to the Benjamini-Hochberg method. **E)** Full spectrum flow cytometry gating strategy for CD14$^+$ monocyte characterization in PBMCs isolated from a healthy individual. **F)** Representative histogram plots for the activation marker CD40 and CD80 in CD14$^+$ monocytes after exposure to iHep supernatants with or without FAD^H. **G)** Quantification of the parameters shown in F). Multiple t-test to assess significance, corrected according to the Benjamini-Hochberg method.

1489