1

**Noncovarying storage effect: balancing and positive directional selection on mutant alleles**
**that amplify random fitness and demographic fluctuations**

Yuseob Kim

Division of EcoScience, Ewha Womans University, Seoul, Korea 03760

yuseob@ewha.ac.kr

Running head: Eco-evolutionary fluctuation dynamics

Keywords: fluctuation, storage effect, population size, positive feedback, eco-evolutionary

Correspondence:

Yuseob Kim, Science Hall B308, Ewhayeodae-gil 52, Seodaemun-gu, Seoul, Korea 03760

Tel: +82 10 5138 5175

ABSTRACT

Temporally variable environments in natural populations generate fluctuations in not only the

population size but also the fitness effects of mutant alleles. The theory of storage effect, a

species/allelic diversity-promoting mechanism discovered in ecology, predicts that rare mutants

with fluctuating fitness can be positively selected and then maintained in balanced polymorphism

if the population is subdivided into two parts that are respectively exposed to and protected from

fluctuating environment. Recent study found that, under pre-exisiting oscillation in population

size, the mutant is positively selected to fixation if its fitness change correlates with the rate of

population growth, which further amplifies population size oscillation. To further understand

these eco-evolutionary dynamics and elucidate their generality in natural populations, this study

built more realistic models that assume randomly, not cyclically, fluctuating selection and

common demographic features, including heterogeneous ecological patches or an age-structured

population. Mathematical analysis revealed that this novel evolutionary force is generated when

the size of the subpopulation subject to weak selection does not covary with that of the other

subpopulation. Simulations showed that this 'noncovarying storage effect' is robust to various

perturbations in the model. Multi-locus simulations revealed that oscillatory polymorphism at

many loci can be simultaneously maintained and that positive feedback between demography and

selection can accelerate the sequential fixations of fluctuation-amplifying mutations and thus lead

to a drastic amplification of population size fluctuation. These results suggest that the

noncovarying storage effect is a potentially prevalent evolutionary force in nature for maintaining

genetic variation and causing large demographic fluctuations.

ARTICLE SUMMARY

Natural populations experience rapid environmental changes that cause fluctuations in the fitness of genetic variants and population size. This study investigates how evolutionary changes due to fitness differences interact with concurrent demographic changes. The theory of the storage effect, which is a mechanism for promoting genetic diversity, was extended to add random population size fluctuation with heterogeneous environmental patches or in a population divided into different age classes. A novel evolutionary force that either increases genetic variation or drives evolution toward further amplification of population size fluctuation, an unconventional contributor to large demographic fluctuations in nature, was identified and analyzed.

INTRODUCTION

Natural populations experience temporal changes in abiotic and biotic environments. A randomly

variable environment is expected to produce fluctuations in the fitness effects of non-neutral or

phenotype-changing alleles segregating in the population. If the magnitude of fluctuation in the

relative fitness is large, evolutionary changes may occur rapidly at the pace of short-term

ecological/demographic fluctuations. Recent developments in evolutionary biology have revealed

that such rapid evolutionary changes are widespread (Bell 2010; Thompson 2013; Messer *et al.*

2016; Lynch *et al*. 2024). Temporally variable environments should also generate fluctuations in

various aspects of demography, most importantly the number of reproducing individuals (i.e.,

population size). One may then ask whether mutual interaction between demographic and

selective fluctuations creates qualitatively novel eco-evolutionary dynamics that cannot arise if

one of the demographic or population genetic variables is held constant.

Considering that a fluctuation in population size reflects a fluctuation in the reproductive success

of individuals and that long-term reproductive success is maximized if its variance in the offspring

number over time is minimized (Slatkin 1974; Gillespie 1977; Tuljapurkar 1982), one may predict

that a population will evolve to reduce size fluctuation as much as possible (Pfister 1998). For

example, a variant increasing the egg size rather than the egg number at reproductionwhen

plentiful resources are available may be positively selected as the offspring can survive better

during an unfavorable period. Then, when a large fluctuation in population density, such as mast

seeding or insect outbreak, does occur despite the advantage of a small temporal variance in

offspring number, this may be seen as a non-adaptive response of the population to a heavy

fluctuation of the biotic and/or abiotic environment (Myers 1988; Dwyer *et al.* 2004; White 2008;

Bogdziewicz *et al.* 2020). However, recent studies suggested that a species may evolve to undergo

a larger fluctuation in its population size. Liu *et al.* (2019) found that a reproductive strategy

increasing the temporal variance of offspring number can be advantageous in populations

reproducing in overlapping generations under various patterns of environmental fluctuations.

Kim (2023) showed that a mutation amplifying the cyclic fluctuation of population size can be

positively selected if this population is partially protected from the cyclical selection (i.e. fitness

fluctuation occurs only in a subset of the population).

Partial protection from fluctuating selection commonly occurs in many species. In spatially

structured species the amplitude of fluctuation in selective pressure is likely to be heterogeneous

over the geographic range. In plants and animals that have their life cycles divided into several

stages and reproduce in overlapping generations, selective pressure is unlikely to act uniformly over individuals of all stages or ages. For example, if a trait under fluctuating selection is expressed in the insect larval stage, alleles affecting this trait currently carried by adults are not visible to the selection. Similarly, alleles affecting traits that appear after germination remain neutral in the seed bank of a plant species. Such a subset of the population protected from fluctuating selection may be called a "refuge." The presence of a refuge of various kinds was found to generate the advantage of rare alleles, a form of balancing selection commonly referred to as the storage effect (Ellner & Hairston 1994; Reinhold 2000; Gulisija & Kim 2015; Gulisija *et al.* 2016; Bertram & Masel 2019; Yamamichi *et al.* 2023), as the refuge acts to temporarily store the variants that otherwise will be removed from the population during unfavorable periods. The storage effect was first theorized in community ecology, shown to arise not only with partial protection from fluctuation provided by a refuge but with general covariance between environment and competition, and is now one of the most important ecological theories for explaining the stable coexistence of species (Chesson & Warner 1981; Chesson 1994; Barabás *et al.* 2018). Imported into population genetics, it was recently proposed as one of a few mechanisms that may produce the genome-wide occurrence of oscillatory polymorphism, as observed in North American *Drosophila melanogaster* and other species, which appear to be maintained under fluctuating environments (Bergland et al. 2014; Park and Kim 2019; Johnson et al. 2023).

However, Kim (2023) showed that if the oscillation of population size is added to the model of storage effect a variant whose fitness also oscillates in the same directions can be either maintained in balanced polymorphism or positively selected to reach fixation. Mathematical analysis showed that whether balancing or directional selection occurs on this fitness-amplifying allele depends on the relative strengths of fitness versus population size fluctuations. Previous studies on balancing selection by the storage effect in a constant-sized population required the coexisting variants to have identical geometric mean fitness. However, when the population size was allowed to oscillate, it led to negative frequency-dependent selection on mutants with even smaller geometric mean fitness than the wild-type allele. It was also shown that positive directional selection on such variants, which cannot arise in the previous models of the storage effect, occurs if demographic fluctuation is relatively stronger than fitness fluctuation. Furthermore, Kim (2023) found that, under a weak regulation of population density under which the population size can over- or undershoot the carrying capacity, the positive selection leads to the amplification of oscillation in mean absolute fitness, thus population size. In summary, the addition of population size fluctuation, either externally imposed by ecology or due to the

segregation of fitness-amplifying mutants, in the model was found to generate novel eco-evolutionary dynamics not predicted by the conventional model of the storage effect.

The generality of novel eco-evolutionary dynamics discovered in Kim (2023) may however be questioned since the major result was obtained from the simulation of cyclically (seasonally) and symmetrically changing environments under which the population size and fitness oscillations are fully correlated. Although a mathematical analysis found general conditions for determining which evolutionary outcome (the loss, balanced polymorphism, or fixation of the fitness-amplifying allele) is expected, it was based on the model of only two alternating seasons with fixed sizes. Therefore further investigation using more realistic models is needed to obtain better theoretical understanding and to find out whether the novel eco-evolutionary dynamics can arise in a substantially wider range of biological settings.

This study broadens the scope of investigation in the following directions. First, mathematical models assuming random, rather than deterministic and symmetrically seasonal, fluctuations in the environment were built and analyzed. A model with seasonal fluctuation is suitable for cases in which one cycle of environmental changes spans multiple generations. However, for many species, one generation is not shorter than a seasonal cycle. Random fluctuation of carrying capacities and the direction/strength of selection over generations, with a varying degree of correlation between these fluctuations, is probably relevant to a wider range of species including plants that exhibit mast seeding. Second, new models of population structures and inter-deme differences in selection intensity were investigated. Whereas Kim (2023) focused on a population with a seed bank (Turelli *et al.* 2001; Gremer and Venable 2014) in which the mutant is completely protected from selection in the refuge, this study uses a model (the two-patch or TP model) that assumes two demes or ecological patches, both reproducing and undergoing changes in the carrying capacity, under general heterogeneity in selection intensity. This TP model was also analyzed extensively by simulation to test its robustness against perturbations. Another model (the larval-subadult-adult or LSA model) assuming a population with an age structure, in which the life-cycle is divided into the larval (juvenile), subadult, and adult stages, was also explored by simulation. Third, for both TP and LSA models, multi-locus stochastic simulations were performed. Simulations showed that mutant alleles causing wider fluctuation in offspring number can reach fixation at multiple loci in a wide range of conditions. These fixations were found to occur at faster rates with increasing numbers of loci, as fluctuating population size and fluctuating selection reinforced each other in positive feedback.

As will be clarified in the mathematical analysis, a key ingredient for this novel eco-evolutionary

dynamics is the presence of a refuge whose size (or carrying capacity) remains relatively invariant

or does not covary with the size of the rest of the population. One may therefore call this

evolutionary mechanism the "noncovarying storage effect". The result suggests that the

noncovarying storage effect arising under the general condition of heterogeneous fluctuating

selection in structured populations likely contributes to maintaining non-neutral genetic

variation and to increasing demographic variability. Analytical and simulation results for the TP

model will be first presented, followed by simulation results for the LSA model.

Figure 1. Schematic diagrams of population models. **A**. Two-patch (TP) model assumes a population of haploid individuals occupying a field, where the mutant allele $A_2$ has selective advantage $1 + S_t$ in generation $t$, and a refuge, where the advantage is $1 + S^*_t$. Each generation starts with young individuals that either migrated across or stayed in subpopulations after being reproduced in the previous generation. They also undergo mutation and recombination before selection starts. Then, they undergo selection such that the total number of reproducing individual is constrained to the carrying capacity $K_F^{(t)}$ for the field and $K_R^{(t)}$ for the refuge assuming soft selection. In case of hard selection the carrying capcities are effectively $(1 + qS_t)K_F^{(t)}$ and $(1 + qS^*_t)K_F^{(t)}$. **B**. In the larval-subadult-adult (LSA) model, at each time step haploid individuals move from larval to subadult stage with probability $e_L$ and from subadult to adult stage with probability $e_S$. Reproduction, followed immediately by mutation and reproduction, occurs only in the subadult and adult stages. A new larval subpopulation at time $t$ starts with individuals produced at time $t$-1. At the end of time interval $t$, $K_L^{(t)}$ individuals remain in the larval stage before moving to the next stage.

**Two-patch (TP) model**

Consider a haploid population that reproduces in discrete generations. The outline of the life cycle is illustrated in Fig. 1A. The population is divided into two subpopulations or patches that are subject to different levels and/or phases of fluctuating selection (thus two-patch or TP model). For convenience, these subpopulations will be called the field and the refuge following the terminology of Kim (2023). The latter refers to a subpopulation where selection on a mutant allele is weaker. It is assumed that the population is initially fixed for the ancestral $A_1$ allele at a locus. Then, a new mutant allele $A_2$, subject to fluctuating selection arises. The fitness of an individual carrying $A_2$ relative to $A_1$ is given by $1 + S_t$ in the field and $1 + S_t^*$ in the refuge, where $\mathrm{E}[S_t] = \mathrm{E}[S_t^*] = 0$, $\mathrm{Var}[S_t] > 0$, and $\mathrm{Var}[S_t^*] \geq 0$. Notably, while the arithmetic mean of the fitness of $A_2$ over time is 1, its geometric mean is less than 1, in both subpopulations. Natural selection is thus expected to eliminate such an allele from a simple panmictic (unstructured) population.

A fluctuating environment causes fluctuations in the carrying capacities of both the field and refuge, which are given by $K_F^{(t)}$ and $K_R^{(t)}$ respectively for generation $t$. The carrying capacity is defined as the expected number of (reproductive) individuals at the end of each generation. It is assumed that the carrying capacities of both subpopulations depend on common environmental factors that fluctuate randomly over time but that subpopulations may differ in the strengths of the dependence. The sizes of subpopulations, $N_F^{(t)}$ and $N_R^{(t)}$, after the reproduction step in generation $t$, may or many not match the corresponding carrying capacities depending on the strength of population density regulation (see below). Before a new generation starts, each individual in the field migrates to the refuge or remains in the field with probability $m_{FR}$ and 1 - $m_{FR}$. Similarly, each individual in the refuge either migrates or not with probability $m_{RF}$ and 1 - $m_{RF}$. Next, the size of the field (refuge) changes from $\widetilde{N}_F^{(t)}$ ($\widetilde{N}_R^{(t)}$) at the beginning of generation $t$ to $N_F^{(t)}$ ($N_R^{(t)}$) at the end, where $\widetilde{N}_F^{(t)} = (1 - m_{FR})N_F^{(t-1)} + m_{RF}N_R^{(t-1)}$ and $\widetilde{N}_R^{(t)} = m_{FR}N_F^{(t-1)} + (1 - m_{RF})N_R^{(t-1)}$.

Let $n_{1F}$ and $n_{2F}$ be the numbers of individuals in the field carrying $A_1$ and $A_2$ alleles at the beginning (before the reproduction step) of generation $t$ (thus $n_{1F} + n_{2F} = \widetilde{N}_F^{(t)}$). In the reproduction step, two different assumptions can be made regarding the effect of the mean fitness of individuals on the total population size. When the allele with higher or lower fitness increases in frequency, the total population size may not change due to an ecological constraint. In other words, the absolute fitness of the allele is frequency-dependent while its relative fitness remains constant. Conversely

the mutant allele may increase or decrease the absolute reproductive output of the carriers

regardless of its frequency such that the population size can increase over or decrease below the

carrying capacity, which is now interpreted as the expected population size assuming that all

individuals carry the ancestral ($A_1$) allele. For convenience, these two modes of natural selection

will be simply referred to as soft and hard selection, respectively, following the terminology in

Wallace (1975). First, under soft selection, very strong density regulation occurs so that the final

size of the field is constrained to $K_F^{(t)}$. The number of progeny born to each parent is given by

$W K_F^{(t)} / \bar{N}_F^{(t)}$, where the fitness $W$ is 1 for $A_1$ and $1 + S_t$ for $A_2$, and $\bar{N}_F^{(t)} = n_{1F} + (1 + S_t) n_{2F}$.

Second, under hard selection, fitness-changing mutations lead to changes in not only allele

frequencies but also in the population size. To this effect, the number of progeny per parent is

now $W K_F^{(t)} / \tilde{N}_F^{(t)}$. Namely, when $q = n_{2F} / (n_{1F} + n_{2F})$ is the relative frequency of $A_2$ at the

beginning of a generation, the size of the field is expected to reach $(1 + q S_t) K_F^{(t)}$, which becomes

the effective carrying capacity of the field under hard selection. Similarly, reproduction in the

refuge occurs either with soft or hard selection, where subscript $F$ in the above expressions is

replaced by $R$ and $W$ is 1 for $A_1$ and $1 + S_t^*$ for $A_2$. The model so far described changes in the

number of individuals without genetic drift. When a finite population size is assumed, as in

simulation below, each parent in both subpopulations is modeled to produce a Poisson number

of offspring with the mean specified above, which approximates genetic drift in the Wright-Fisher

model.

Finally, how the environment fluctuates over time and how demographic and selective parameters

— $K_F^{(t)}$, $K_R^{(t)}$, $S_t$, and $S_t^*$ — respond need to be specified. The environmental condition is

assumed to fluctuate randomly in each generation, therefore with no temporal autocorrelation.

More specifically, fluctuations occur such that $K_F^{(t)} = K_{F0} e^{U_t}$ and $K_R^{(t)} = K_{R0} e^{V_t}$ where $U_t$ and

$V_t$ are drawn each generation from a joint normal distribution satisfying $E[U_t] = E[V_t] = 0$. The

oscillation of carrying capacities is therefore symmetrical in the log scale, which is generally

predicted in demographic models and observed in nature (Tuljapurkar & Orzack 1980; Myers

1988). It is assumed that $S_t$, $S_t^*$, $U_t$, and $V_t$ may be correlated to each other as they respond to

the fluctuation of common environmental variables. However, the correlation among them

satisfies that $\mathrm{Cov}[S_t, U_t - V_t]$ is greater than $\mathrm{Cov}[S_t^*, U_t - V_t]$, which gives the mathematical

definition of the field and refuge.

**Analytic results: the fate of the $A_2$ allele**

This section aims to predict the fate of the rare $A_2$ allele — whether it will be lost, fixed, or increased to intermediate frequencies and maintained polymorphic in the population — as a function of the demographic and fitness parameters in the TP model. For convenience in the derivation, genetic drift is ignored and a fixed scheme of migration, $m_{FR} = 1 - m_{RF} = r$, that sends a constant fraction $r$ of the population to the refuge regardless of the current location is assumed. (Below the dynamics will be examined in simulation without this restriction in the migration rate.) First, soft selection, under which $N_F^{(t)} = K_F^{(t)}$ and $N_R^{(t)} = K_R^{(t)}$, is assumed. Let $n_1$ and $n_2$ be the total numbers of individuals (counted after reproduction/selection and before migration) carrying $A_1$ and $A_2$ in generation $t - 1$ and $n'_1$ and $n'_2$ be their expected numbers in generation $t$. Then, with $n_1 \gg n_2$,

$$\frac{n'_2}{n_2} \approx (1-r)\frac{K_F^{(t)}}{\widetilde{N}_F^{(t)}}(1 + S_t) + r\frac{K_R^{(t)}}{\widetilde{N}_R^{(t)}}(1 + S_t^*) = \frac{(1+S_t)K_F^{(t)}+(1+S_t^*)K_R^{(t)}}{K_F^{(t-1)}+K_R^{(t-1)}} \tag{1}$$

(Appendix A). This equation takes into account that the change in the absolute number of $A_2$ is determined first by demography, as the size of field changes from $\widetilde{N}_F^{(t)}$ to $K_F^{(t)}$ and that of refuge from $\widetilde{N}_R^{(t)}$ to $K_R^{(t)}$, and also by fitness $1 + S_t$ and $1 + S_t^*$. Whether the copy number of rare allele $A_2$ is expected to increase or decrease is determined by the expectation of $\log\left[\frac{n'_2}{n_2}\right]$ over demographic and fitness fluctuations. It is shown in Appendix A that the condition for the $A_2$ allele being positively selected (i.e. successfully invading the population initially fixed for $A_1$), $E\left[\log\left[\frac{n'_2}{n_2}\right]\right] > 0$, is simplified to

$$\frac{\text{Cov}[S-S^*,U-V]}{\text{Var}[S]+\rho^2\text{Var}[S^*]+2\rho\text{Cov}[S,S^*]} = \frac{\text{Cov}[S-S^*,\log[K_F/K_R]]}{\text{Var}[S]+\rho^2\text{Var}[S^*]+2\rho\text{Cov}[S,S^*]} > \frac{1}{2\rho} \tag{2}$$

where $\rho = K_{R0}/K_{F0}$ (note that $K_F^{(t)} = K_{F0}\mathrm{e}^{U_t}$ and $K_R^{(t)} = K_{R0}\mathrm{e}^{V_t}$) represents the relative size of the refuge and super-/subscripts for time $t$ in variables were omitted.

If it is assumed that selection on $A_2$ in the refuge is weaker than that in the field by a ratio δ, namely $S_t^* = \delta S_t$, the above inequality is simplified to

$$\Phi \equiv \frac{\text{Cov}[S,\log[K_F/K_R]]}{\text{Var}[S]} > \frac{(1+\delta\rho)^2}{2(1-\delta)\rho} \tag{3}$$

Therefore, the rare $A_2$ allele is positively selected if the covariance between its fitness and the log-ratio of the field to refuge size is large relative to the amplitude of the fitness fluctuation. This

requires that the demographic fluctuation in the population be sufficiently large, as already shown

in Kim (2023), and that, to allow the substantial fluctuation of $K_F^{(t)}/K_R^{(t)}$, the carrying capacity of

the refuge should not covary strongly with that of the field. Since this requirement for the refuge

is essential and distinguishes the current from the previous models of storage effect, the

evolutionary force driving this positive selection may be called the noncovarying storage effect

(see Discussion for examples of noncovarying and covarying storages). The conditions (2) and (3)

are easier to meet as δ becomes small well below 1, namely when the fitness fluctuation of $A_2$

becomes either smaller in the refuge compared to the field or negatively correlated to $S_t$.

In a similar manner, the condition for the $A_1$ allele to invade a population initially fixed for $A_2$ under

soft selection becomes

$$\frac{\mathrm{Cov}[S-S^*, U-V]}{(1+2\rho)\mathrm{Var}[S]+(2\rho+\rho^2)\mathrm{Var}[S^*]-2\rho\mathrm{Cov}[S,S^*]} < \frac{1}{2\rho} \tag{4}$$

257   (Appendix A) or

$$\Phi < \frac{1+\delta^2}{1-\delta} + \frac{(1+\delta\rho)^2}{2(1-\delta)\rho} \tag{5}$$

if $S_t^* = \delta S_t$.

In a simpler case with δ = 0, from inequalities (3) and (5), it is predicted that the $A_1$ and $A_2$ alleles

coexist in the population if $1/(2\rho) < \Phi < 1 + 1/(2\rho)$ under soft selection. With $\Phi < 1/(2\rho)$,

the $A_2$ allele is eliminated from the population, as the fitness fluctuation that reduces its geometric

mean in the field is more important than population size fluctuation. With $\Phi > 1 + 1/(2\rho)$, the

$A_2$ allele is predicted to reach fixation in the population.

Using the same approach, the fate of the $A_2$ allele under hard selection can be analyzed. Note that

the carrying capacities of the field and refuge are modified to $(1 + qS_t)K_F^{(t)}$ and $(1 + qS_t^*)K_R^{(t)}$

with hard selection where $q$ is the frequency of $A_2$. The frequency change of $A_2$ when rare is

therefore effectively identical to that in soft selection. Therefore, the condition for $A_2$ invading the

population fixed for $A_1$ is again given by inequalities (2) and (3). However, the condition for the

fixation of $A_2$ is drastically different from that under soft selection. It is shown in Appendix A that

the condition for $\mathrm{E}\left[\log\left[\frac{n_1'}{n_1}\right]\right] > 0$ for rare $A_1$ is exactly the condition for $\mathrm{E}\left[\log\left[\frac{n_2'}{n_2}\right]\right] < 0$ for rare

$A_2$. Therefore, with $S_t^* = \delta S_t$, the condition for the $A_1$ allele to invade the population fixed for $A_2$

is approximately $\Phi < (1 + \delta\rho)^2/(2(1-\delta)\rho)$. This means that the fixation of $A_2$ is ensured if

$\Phi > (1 + \delta\rho)^2/(2(1-\delta)\rho)$; once positively selected after its appearance by mutation, the $A_2$

allele is expected to increase in frequency all the way to fixation. This leads to the amplification of

fluctuation in the size of the field, from $K_F^{(t)}$ to $(1 + S_t)K_F^{(t)}$. This result is consistent with the

finding in Kim (2023) that, under weak density regulation of the population size, an allele with a

wider oscillation in absolute fitness reaches fixation in the presence of refuge.

## One-locus simulation

To validate the above analysis and explore further eco-evolutionary dynamics in the TP model,

the one-locus stochastic simulation was performed. The demographic and fitness fluctuations

were modeled by $U_t = a_U Z_1 + b_U Z_2$, $V_t = a_V Z_1 + b_V Z_3$, $S_t = a_S Z_1 + b_S Z_4$, and $S_t^* = a_{S^*} Z_1 +$

$b_{S^*} Z_5$ where $a_X$ ($b_X$) is a constant for determining the magnitude of change in variable $X$ due to a

common environmental factor (other factors) and the $Z_i$s are standard normal variates that were

drawn independently each generation. Therefore, $Z_1$ represents the state of the environment

affecting both population sizes and fitness.

Simulations were first performed for a scenario in which $A_2$ allele is completely protected from

selection in the refuge ($S_t^* = 0$) and migration follows a simple scheme ($m_{FR} = m_{RF} = 0.5$). As

suggested by the analysis above, the major parameter of eco-evolutionary dynamics in this case

is $\Phi = \text{Cov}[S, U - V]/\text{Var}[S] = (a_S a_U - a_S a_V)/(a_S^2 + b_S^2)$. Simulation results were obtained for

various values of $\Phi$ under soft (Fig. 2A, B) and hard (Fig. 2C, D) selection. Initially the numbers of

$A_1$ and $A_2$ individuals changed deterministically once the demographic and fitness parameters ($U$,

$V$, and $S$) for a given generation were sampled from their joint distribution, effectively assuming

that the sizes of the field and refuge were finite and fluctuating but very large so that genetic drift

can be ignored (Fig. 2A). Various values of $a_S$ and $b_S$ were chosen so that $\Phi$ ranged from 0.3 to 2,

while $\rho$ was set to 1. Starting from 0.5, the frequency of the $A_2$ allele at the 5,000th generation was

recorded for 1000 replicates per parameter set. As predicted, with $\Phi \le 0.5$, the frequency of $A_2$

approached 0, thus confirming negative selection, and with $\Phi \ge 1.5$ the frequency approached 1,

thus confirming positive directional selection. The distribution of the allele frequency was

approximately uniform between 0 and 1 when $\Phi = 1$. Next, genetic drift and bidirectional

mutation between two alleles were added to the above simulations (Fig. 2B). This weakened the

strengths of all modes of selection. With $\Phi = 1$, more simulation runs ended with allele frequencies

close to 0 and 1. However, the proportion of runs ending with intermediate allele frequencies was

still larger than that observed when the alleles were set as neutral (dashed curve in Fig. 2B).

Therefore, an evolutionary force to maintain polymorphism beyond the level under the neutrality,

namely balancing selection, was confirmed.

Figure 2. Distribution of $A_2$ allele frequencies ($p$) at the 5,000th generation in the stochastic simulations of soft (A, B) and hard (C, D) selection in the TP model. The initial frequency of $A_2$ is 0.5 in both field and refuge. Reproduction occurred while $K_F^{(t)}$, $K_F^{(t)}$ and $S_t$ varied stochastically without (A, C) or with (B, D) genetic drift and bidirectional mutation ($\mu = 5 \times 10^{-5}$). Proportions of simulation replicates (out of 1000) with the frequencies in the interval $0 \leq p < 0.1$, $0.1 \leq p < 0.2$, …, $0.8 \leq p < 0.9$, and $0.9 \leq p \leq 1$ were plotted and connected by lines. Fluctuations of population sizes are given by $K_{R0} = K_{F0} = 1000$ ($\rho = 1$), $a_U = 0.3$, $b_U = 0.1$, $a_V = 0.05$, $b_V = 0.1$. Parameters of fluctuation were chosen to yield $\Phi = 0.3, 0.5, 0.77, 1, 1.2, 1.5$, and $2$: $(a_S, b_S) = (0.07, 0.23), (0.1, 0.2), (0.14, 0.16), (0.2, 0.1), (0.2, 0.04), (0.15, 0.05)$, and $(0.1, 0.05)$. In the case of hard selection with $\Phi = 1$ and no genetic drift (panel C), simulation ran only upto 2,500 generations: longer simulations or larger $\Phi$ resulted in all replicates with final frequencies > 0.9. The results connected by dashed lines in B and D represent the cases of neutral evolution ($a_S = b_S = 0$).

310 Similarly, the fate of the $A_2$ allele under hard selection was examined. Stochastic simulation

311 confirmed the approximation above predicting that $A_2$ is positively selected towards fixation if

312 $\Phi > \frac{1}{2\rho} = 0.5$, although genetic drift and mutation weakened the trend (Fig 2C,D). Positive

selection on both rare and common $A_2$ under the same condition means that this allele may not remain polymorphic for long in the population. Indeed, across all values of $\Phi$ there were fewer simulation runs ending with intermediate frequencies of $A_2$ compared to the simulation of neutral alleles subject to genetic drift and bidirectional mutations only (dashed curve in Fig. 2D). Therefore, balancing selection that was observed with soft selection did not arise with hard selection.

Next, the cases in which $A_2$ allele is not completely neutral in the refuge was examined. A simple model setting $S_t^* = \delta S_t$ was used. This time, the $A_2$ allele is initially absent in the population but introduced with recurrent mutation. For a given parameter set simulation was run over at least $10^5$ generations and the mean $\bar{H}$ of expected heterozygosity ($2q(1-q)$, where $q$ is the relative frequency of $A_2$) and the proportion of time when $0 \le q < 0.1$, $P_{01}$, and $0.9 < q \le 1$, $P_{90}$, during simulation were recorded. Simulation with $S_t = S_t^* = 0$ but under the same demographic structure yielded $\bar{H}$ = 0.063 (average over $6 \times 10^5$ generations). Then, as balancing selection is defined to be a force yielding genetic variation above the neutral level, a result was classified as balanced polymorphism if $\bar{H} > 0.075$. If $\bar{H} < 0.05$ and $P_{01} > P_{90}$, it suggests that negative selection against $A_2$ was the dominant evolutionary force. Such results were therefore classified as the loss of $A_2$, since recurrent losses of $A_2$ should be the major events in the simulation (confirmed by visual inspection of allele frequency trajectories; not shown). Conversely, results yielding $\bar{H} <$ 0.05 and $P_{90} > 0.7$, which indicates that positive directional selection on $A_2$ was the dominant force, were classified as the fixation of $A_2$. All others were classified as not distinguishable from the neutral evolution. Simulations largely confirmed the analytic approximations for the conditions for the loss, fixation, or balanced polymorphism of $A_2$ in a population initially fixed for $A_1$ (e.g., $\frac{(1+\delta\rho)^2}{2(1-\delta)\rho} < \Phi < \frac{1+\delta^2}{1-\delta} + \frac{(1+\delta\rho)^2}{2(1-\delta)\rho}$ for the balanced polymorphism under soft selection and $\Phi > \frac{(1+\delta\rho)^2}{2(1-\delta)\rho}$ for the fixation under hard selection; Fig. 3). It is shown that increasing $\delta$ above 0 makes it harder for balancing and positive directional selection on $A_2$ to occur; for a large value of $\delta$, the fluctuation (variance) of $S_t$ needs to be much smaller than its covariance with the fluctuation of $\log[K_F^{(t)}/K_R^{(t)}]$ (eq. 3). Small Var[$S_t$] then means that the relative strength of genetic drift affecting the evolutionary dynamics of $A_2$ is large, explaining the results not distinguisable from the neutrality with large $\delta$ and $\Phi$ in Figure 3. Other results not distinguishable from neutral evolution were obtained when the predicted effects are near the boundary between balancing and directional selection. Another important result is that negative $\delta$, thus selection in the refuge being in the opposite direction, widens the parameter range for balancing selection.

Figure 3. Simulation results for non-zero values of δ. For a given parameter set simulation started with the $A_1$ allele fixed in the population and ran for $10^5$ generations with recurrent bidirectional mutations and soft (left) or hard (right) selection. Four different outcomes regarding the overall behavior of the $A_2$ allele (balanced, lost, fixed, neutral) were marked by different symbols and plotted as functions of δ and Φ. The demographic parameters are identical to those in Figure 1. Five different values of δ (-0.3, -0.1, 0.1, 0.3, and 0.3) and ten different values of Φ were chosen: 0.246, 0.5, 0.8, 1, 1.25, 1.5, 1.786, 2.027, 2.5 and 2.94: $(a_S, b_S)$ = (0.05, 0.22), (0.1, 0.2), (0.2, 0.15), (0.2, 0.1), (0.2, 0), (0.15, 0.05), (0.14, 0), (0.12, 0.02), (0.1, 0) and (0.085, 0). Gray and dashed curves plot $\Phi = \frac{(1+\delta\rho)^2}{2(1-\delta)\rho}$ and $\Phi = \frac{1+\delta^2}{1-\delta} + \frac{(1+\delta\rho)^2}{2(1-\delta)\rho}$ with $\rho = 1$.

Finally, the robustness of balancing and positive directional selection above to deviations from the simple migration scheme, $m_{FR} = 1 - m_{RF} = r$, was examined. In actual populations, migration rates may vary over time and may not be symmetrical between the field and refuge, at least in a short time scale. New simulations therefore sampled migration rates $m_{FR}$ and $m_{RF}$ from uniform distributions in ranges $[m_1, m_2]$ and $[m_3, m_4]$, respectively, each generation (Table S1). With ρ = 1 ($K_{F0} = K_{R0}$), limiting both $m_{FR}$ and $m_{RF}$ close to 0.5 yielded the highest level of balanced polymorphism under soft selection. However, allowing random deviation from 0.5 (e.g. $[m_1, m_2]$ = $[m_3, m_4]$ = [0.3, 0.7]) only slightly lowered the level of polymorphism. Smaller migration rates generally disrupted balancing selection. Interestingly, the effect of $m_{FR}$ and $m_{RF}$ was not symmetric; balancing selection remained effective with $m_{RF}$ close to 0.5 and $m_{FR}$ less than 0.2, however not with the opposite ratio of rates (compare cases with $[[m_1, m_2], [m_3, m_4]]$ = [[0.1, 0.2], [0.4, 0.5]] and [[0.4, 0.5], [0.1, 0.2]] in Table S1). Similar effects of $m_{FR}$ and $m_{RF}$ on the directional increase of $A_2$ allele frequency under hard selection were observed.

360 In summary, one-locus simulations for the TP model not only confirmed the analytic
361 approximations for the conditions under which the noncovarying storage effect leads to either
362 balanced polymorphism or the fixation of the fluctuation-amplifying allele but also showed that
363 the effect is robust to perturbations in the model such as genetic drift and
364 randomness/asymmetry in the migration rate.

**Multi-locus simulation**

To investigate how the eco-evolutionary dynamics generated by the noncovarying storage effect
in the TP model play out when there are many such loci in the genome, new sets of simulations
were performed in which a haploid individual is modeled to have $L$ linked loci, each carrying
either the $A_1$ or $A_2$ allele described above. To avoid exploring too many parameter dimensions
while focusing on the effect of increasing the number of loci, $S_t^* = 0$ and a fixed migration
scheme ($m_{FR} = m_{RF} = 0.5$) are assumed. The simulation tracks the numbers of individuals of
different haplotypes (see Appendix B). Epistasis is not considered; the relative fitness of
haplotype $i$ is given by $W_i = \prod_{l=1}^{L}(1 + S_{t,l})$, where $S_{t,l}$ is 0 if the $l$th locus of the haplotype
carries $A_1$ and $a_S Z_1 + b_S Z_4$ if $A_2$. $Z_1$ is drawn once for all loci while $Z_4$ is drawn separately for
different loci. Individuals randomly pair and undergo recombination by crossing-over such that
the probability of recombination between adjacent loci is $c$ per generation. The simulation starts
with a population fixed for $A_1$ at all loci. Then $A_2$ arises by mutations with probability
μ/locus/generation.

Two major outcomes of the noncovarying storage effect – balancing selection and positive
directional selection – were addressed. First, considering that (1) the negative frequency-
dependent selection on the $A_2$ allele was caused primarily by its fitness correlated to the
fluctuation in the carrying capacity (population size) of the field and that (2) this demographic
fluctuation should be "felt" at all loci in the genome, one may expect that such balancing selection
can occur simultaneously at many loci harboring such variants. To verify if this is true, simulations
were run under soft selection and demographic/selective parameters that promoted
polymorphism in the single-locus model ($\Phi$ = 1 for $\rho$ = 1) with $L$ = 1, 4, or 10. Exemplary
trajectories of allele frequencies are shown in Fig. S1. Each run lasted at least $10^5$ generations over
which $\bar{H}$, the expected heterozygosity ($2p(1-p)$) averaged over time and loci, and the
proportion of time the $A_2$ alleles spent in frequency between 0.1 and 0.5, $P_{15}$, and between 0.5 and
0.9, $P_{59}$, were recorded (Table 1). In all cases with fluctuating fitness ($a_S > 0$), $\bar{H}$ was above the

Table 1. The long-term level of polymorphism in the multi-locus simulation

| demography | selection mode | $L$ | $c$ | $\bar{H}$ | $P_{15}$ | $P_{59}$ |
|---|---|---|---|---|---|---|
| $a_U = 0.2$, $a_V = b_U = b_V = 0.1$ | neutral ($a_S = b_S = 0$) | 1 | - | 0.0639 | 0.069 | 0.074 |
| | soft selection, $\Phi = 1$ ($a_S = 0.1$, $b_S = 0$) | 1 | - | 0.132 | 0.15 | 0.16 |
| | | 4 | $10^{-4}$ | 0.0960 | 0.099 | 0.12 |
| | | 4 | $10^{-3}$ | 0.103 | 0.12 | 0.12 |
| | | 4 | 0.01 | 0.0977 | 0.11 | 0.12 |
| | | 4 | 0.1 | 0.136 | 0.11 | 0.14 |
| | | 10 | $10^{-4}$ | 0.080 | 0.087 | 0.095 |
| | | 10 | $10^{-3}$ | 0.102 | 0.12 | 0.12 |
| | | 10 | 0.01 | 0.109 | 0.15 | 0.10 |
| | | 10 | 0.1 | 0.107 | 0.13 | 0.12 |
| $a_U = 0.3$, $a_V = 0.05$, $b_U = b_V = 0.1$ | neutral ($a_S = b_S = 0$) | 1 | - | 0.0633 | 0.081 | 0.074 |
| | soft selection, $\Phi = 1$ ($a_S = 0.2$, $b_S = 0.1$) | 1 | - | 0.177 | 0.24 | 0.18 |
| | | 4 | $10^{-4}$ | 0.162 | 0.19 | 0.19 |
| | | 4 | $10^{-3}$ | 0.130 | 0.21 | 0.11 |
| | | 4 | 0.01 | 0.168 | 0.22 | 0.18 |
| | | 4 | 0.1 | 0.183 | 0.25 | 0.19 |
| | | 10 | $10^{-4}$ | 0.120 | 0.14 | 0.14 |
| | | 10 | $10^{-3}$ | 0.141 | 0.17 | 0.17 |
| | | 10 | 0.01 | 0.0934 | 0.076 | 0.14 |
| | | 10 | 0.1 | 0.105 | 0.093 | 0.15 |
| $a_U = 0.05$, $a_V = b_U = b_V = 0$ | neutral ($a_S = b_S = 0$) | 1 | - | 0.0660 | 0.085 | 0.063 |
| | soft selection, $\Phi = 1$ ($a_S = 0.05$, $b_S = 0$) | 1 | - | 0.0816 | 0.11 | 0.081 |
| | | 10 | $10^{-4}$ | 0.0728 | 0.085 | 0.079 |
| | | 10 | $10^{-3}$ | 0.0982 | 0.12 | 0.11 |
| | | 10 | 0.01 | 0.0953 | 0.12 | 0.096 |
| | | 10 | 0.1 | 0.0952 | 0.13 | 0.092 |

- Simulations were run for $5 \times 10^5$, $2 \times 10^5$, and $10^5$ generations with $L = 1$, 4, and 10, respectively.
- Other parameters: $K_{R0} = K_{F0} = 1{,}000$ ($\rho = 1$), $\mu = 2 \times 10^{-5}$.

level observed in equivalent simulations with neutral alleles ($S_t = 0$), thus confirming multi-locus balancing selection. For cases of stronger selection (larger $\text{Var}[S_t]$), the level of polymorphism per locus declined as $L$ increased from 1 to 10. The combined effects of the number of loci and the rate of recombination were not easy to interpret as they depended on the parameters of selection (for sets of fitness/demographic parameters that yield the same $\Phi = 1$, $\bar{H}$ responded differently with varying $L$ and $c$). One possible explanation for generally lower variation with 10 loci instead of 1 or 4 is that the simultaneous segregation of alleles at many loci increased the variance of the

400  offspring numbers, thus decreasing the effective population size and weakening the negative
401  frequency-dependent selection that is needed for protected polymorphism. Further exploration
402  of multi-locus polymorphism at a larger scale (in both number of loci and population size) may
403  not be feasible with the current simulation approach based on haplotype frequencies.

Figure 4. Changes of mutant ($A_2$) allele frequencies and subpopulation sizes (A. TP model, blue: $N_F^{(t)}$, green: $N_R^{(t)}$; B. LSA model, blue: larval, green: subadult, gray: adult population) in the multi-locus simulations of hard selection. Allele frequency trajectories of 8 loci were plotted in different colors. Parameters: A (TP model): $L = 8$, $c = 0.02$, $K_{R0} = K_{F0} = 1000$, $\mu = 2\times10^{-5}$, and $\Phi = 1$ ($a_S = 0.15$, $b_S = 0$, $a_U = 0.15$, $a_V = 0$, $b_U = 0$, $b_V = 0.1$). B (LSA model): $L = 8$, $c = 0.02$, $K_{L0} = 5000$, $\mu = 2\times10^{-5}$, and $\Phi' = 1$ ($a_S = 0.1$, $b_S = 0.1$, $a_L = 0.2$, $b_L = 0.1$).

Second, simulation was performed with parameter values under which directional selection for
the $A_2$ allele is expected in the single-locus model ($\Phi > 1 + 1/(2\rho)$ for soft selection and $\Phi >$
$1/(2\rho)$ for hard selection). As expected, substitutions occurred sequentially until $A_2$ reached
fixation at all $L$ loci (Fig. 4A and Fig. S2 for example). Times (in generations) when the frequency

of $A_2$ first exceeded 0.9 at each locus were recorded and sorted into $T_1$ to $T_L$. There is a large

difference in substitution dynamics between soft and hard selection. With soft selection, the mean

waiting time, $\bar{T} = (T_1 + \cdots + T_L)/L$, increased moderately as $L$ increased from 1 to 8 (Fig. 5A).

This may be explained by a reduction in the efficacy of positive selection on $A_2$ at a given locus

when selection at other loci results in genome-wide reduction in the effective population size.

Figure 5. Mean fixation time ($\bar{T}$) with varying number of loci (A; $L$ = 1, 2, 4, and 8) and the coefficient of variation in fixation times with $L$ = 8 in the multi-locus simulation of the TP model. Error bars show two times the standard errors. A total of 100 replicates of simulations were run under soft selection with $\Phi$ = 2 (cyan; $a_S$ = 0.1, $b_S$ = 0, $a_U$ = 0.25, $b_U$ = $a_V$ = $b_V$ = 0.05), hard selection with $\Phi$ = 1 (brown; $a_S$ = 0.1, $b_S$ = 0, $a_U$ = 0.1, $a_V$ = 0, $b_U$ = $b_V$ = 0.1), or hard selection with $\Phi$ = 0.75 (orange; $a_S$ = 0.2, $b_S$ = 0, $a_U$ = 0.25, $a_V$ = 0.1, $b_U$ = $b_V$ = 0.05). Other parameters are identical to those in Figure 4A.

With hard selection, however, $\bar{T}$ decreased as $L$ increased. Furthermore, the coefficient of

variation, the standard deviation of $T_i$'s divided by $\bar{T}$, was significantly smaller with hard selection

than with soft selection (Fig. 5B). As expected for hard selection, the amplitude of the population

size fluctuation progressively increased as the number of loci fixed for $A_2$ increased (Fig. 4A).

These results clearly demonstrate that positive feedback between demographic and selective

fluctuations under hard selection led to a great amplification of population size fluctuation; an

initial fluctuation in $K_F^{(t)}$ positively selected the mutant allele $A_2$ at one locus whose fitness

fluctuates in positive correlation with $K_F^{(t)}$, leading to the fixation of $A_2$ and therefore a larger

fluctuation of the field (given by $(1 + S_t)K_F^{(t)}$), which makes it easier for the $A_2$ allele at another

locus to be positively selected. This process could continue to increase the fluctuation of the field

until its carrying capacity is given by $(1 + S_t)^L K_F^{(t)}$, where $L$ is the number of loci available in the

genome for mutation into $A_2$. There was also a moderate effect of the recombination rate in
accelerating the multi-locus fixation of $A_2$ with hard selection (Table S2). Fixation times generally
increased with decreasing recombination rates for both soft and hard selection, possibly because
tighter linkage increased the occurrence of clonal interference.

**Larval-subadult-adult (LSA) model**

Althoug the refuge in the TP model is expected to play the role of various demographic/genetic
elements that were previously shown to produce the storage effect, such as seed banks,
diapausing stages, or a particular life-stage in a population with overlapping generations, the
noncovarying storage effect found in the TP model may not be replicated in other population
structures as the nature of refuge (i.e. whether it undergoes its own reproduction) and the pattern
of 'migration' in and out of the refuge can be qualitatively different. I therefore investigate an
explict discrete-time model of an age-structured population with overlapping generations. Three
stages – 'larval', 'subadult', and 'adult' – in the life cycle of a haploid species are assumed. Time is
counted in intervals of the length required for one life stage to advance to the next one. At the end
of each interval, the final size of larval population is regulated to its carrying capacity ($K_L^{(t)}$ at
time $t$). Probabilities for an individual to advance from larval to subadult and from subadult to
adult stages are given by $e_L$ and $e_S$, respectively. Therefore, the sizes of the subadult and adult
populations are expected to be $e_L K_L^{(t-1)}$ and $e_L e_S K_L^{(t-2)}$ at time $t$. $K_L^{(t)}$ is assumed to vary
randomly over time with no autocorrelation. Specifically it is modeled to be $K_L^{(t)} =$
$K_{L0} \exp(a_L Z_1 + b_L Z_2)$, where $Z_1$ and $Z_2$ are standard normal variables drawn each time.

The population is made of haploids carrying either the $A_1$ or $A_2$ allele at each of $L$ linked loci. The
(marginal) fitness of $A_2$ relative to $A_1$ at locus $j$ is given by $1 + S_{j,t}$ ($j = 1, ..., L$), where $S_{j,t} =$
$a_S Z_1 + b_S Z_{3,j}$. $Z_{3,j}$ is drawn independently for different loci at each interval $t$. Then, the fitness
of haplotype $i$, $W_i^{(t)}$, is given by the multiplication of allelic fitness across all loci. It is assumed
that this haplotype fitness determine the viability in the larval stage only; the carriers of two
alleles are not different either in the probability of advancing to the next stage or in fertility in the
subadult and adult stages. If there are only larval and subadult stages, they constitute a population
that reproduces in effectively discrete generations. The addition of the adult stage provides a
storage for alleles regardless of their reproductive success in the previous (subadult) stage.
Therefore, the adult stage plays the role of the refuge and is an example of a noncovarying storage

because its size is independent of the concurrent subadult stage.

Multi-locus simulations, allowing not only genetic drift in the reproduction step but also
stochastic advances from one to the next life stage (see Appendix B for method), were performed
with limited sets of parameters, each starting with a population fixed for haplotype 0, which
carries $A_1$ at all loci. The evolutionary dynamics of the $A_2$ alleles in each simulation run over
50,000 time units was summarized by the mean heterozygosity and the proportion of time $A_2$
remained close to relative frequency 0 or 1 (Table S3). Major results produced in the TP model –
multi-locus balancing selection under soft selection or positive directional selection on $A_2$ under
hard selection – were readily obtained with this model. In agreement with Kim (2023) and the TP
model, the relative strength of demographic vs. fitness fluctuation determined the outcome; when
we define $\Phi' = \mathrm{Cov}\left[\log[K_L^{(t)}/K_{L0}], S_{j,t}\right]/\mathrm{Var}[S_{j,t}] = a_L a_S/(a_S^2 + b_S^2)$, which is analogous to $\Phi$ in
the TP model, positive selection on rare $A_2$ occurred when $\Phi'$ was above 0.7. With soft selection
and above this threshold of $\Phi'$, the frequency of $A_2$ underwent oscillations that are indicative of
balancing selection (Fig. S3). Then, as $\Phi'$ continued to increase, balancing selection transitioned
into positive directional selection. With hard selection, positive directional selection occurred
once $\Phi'$ increased above the threshold. Exemplary replicates show that, as the $A_2$ alleles reach
fixation across loci, fluctuations in the sizes of all three life stages become amplified (Fig. 4B, Fig.
S3). Negative (positive) correlation between mean fixation times and the number of loci under
hard (soft) selection and the clustering of fixation times with hard selection were also oberved in
other sets of simulations (Fig. S4), which basically replicated the results from the TP model. This
confirmed that positive feedback between demographic and fitness fluctuations under hard
selection occurs in both models. In conclusion, all major aspects of the noncovarying storage effect
found in the TP model were shown to arise in the LSA model as well.

DISCUSSION

A key assumption in this study is that a tradeoff (antagonistic environmental pleiotropy; Zhang
2023) commonly occurs for the fitness effects of a mutant allele between different time points in
the environmental fluctuation. Namely, a mutant phenotype that increases the reproductive
output relative to others when the environment is generally favorable is likely to decrease it
relative to others during an unfavorable period. Such a mutant allele increases the variance of the
offspring number over time while the mean may remain unchanged. For example, as postulated

by Slatkin (1974), a mutation that makes a female bird lay 10 eggs in one batch instead of

spreading them over time in a randomly and rapidly fluctuating environment will increase the

variance of her fitness while the mean remains the same. Setting the fitness of a mutant to $1 + S_t$

with $E[S_t] = 0$ models such a tradeoff. Classical theory predicts that such a mutant (termed an

arithmetic-mean-preserving or AMF allele in Kim (2023)) is eliminated from a population as its

geometric mean fitness is less than 1.

Kim (2023) and this study however found that the AMF mutant at a low frequency is positively

selected (with both soft and hard selection) if a correlation between demographic and fitness

fluctuations exists in the presence of a refuge whose carrying capacity oscillates in a narrower

range and/or out of phase relative to the field. A heuristic argument for this noncovarying storage

effect on the fitness-amplifying mutant allele may be given. In the TP model, the mean number of

offspring per parent, *i.e.* the absolute fitness, of the rare mutant in the field is given by $W_{2f} =$

$\frac{K_F^{(t)}}{\widetilde{N}_F^{(t)}}(1 + S_t)$ (eq. 1), where $K_F^{(t)}/\widetilde{N}_F^{(t)}$ is the absolute fitness given under demographic

fluctuation alone. With positive correlation between $K_F^{(t)}$ and $S_t$, the fluctuation of $W_{2f}$

becomes greater than that of $1 + S_t$ and, more importantly, its arithmetic mean can now exceed

1. Still, the geometric mean of $W_{2f}$ is not greater than 1 due to its fluctuation over time. However,

the presence of a refuge can reduce the variance of mutant's fitness averaged across the whole

population such that the geometric mean of this average fitness becomes greater than 1. Here, to

generate this dampening effect, the refuge should not undergo the same fluctuation in size with

the field; the absolute fitness in the refuge $W_{2r} = \frac{K_R^{(t)}}{\widetilde{N}_R^{(t)}}(1 + S_t^*)$ in eq. (1) should fluctuate

differently than $W_{2f}$. With parameter values used in of the TP model, $K_R^{(t)}$ was given to covary

incompletely with $K_F^{(t)}$. In the LSA model, the fluctuation of the refuge (the adult stage) is

uncorrelated to the fluctuation of the field (the subadult stage). In Kim (2023), the seed bank

fluctuates in size similar to the subadult and adult stages in the LSA model. In summary, such

refuges, or noncovarying storages, turn the heightened arithmetic mean of the mutant allele

raised by the demographic fluctuation of the field into an elevation of the geometric mean in the

total population.

Not all refuges in the previously proposed mechanisms of the storage effect are noncovarying. For

example, it was suggested that if a gene is expressed in one sex in a dioecious species the other

sex could be a refuge (Reinhold 2000; Kim 2023). However, as equal numbers of male and female

gametes are used in reproduction, the subpopulation sizes of reproductive males and females are

effectively equal while the sex ratio of the adults may fluctuate. Therefore, as the size of the field

relative to the refuge does not fluctuate at all, the noncovarying storage effect should not arise for

the genes of the sex-limited expressions.

One may also model the fluctuating fitness of a mutant by $\exp[S_t]$, making the mutant (termed a

geometric-mean-preserving or GMF allele) quasi-neutral as its geometric mean fitness is identical

to that of the wild-type allele even without demographic fluctuation. Analyzing the model of cyclic

environmental oscillation, Kim (2023) showed that the parameter range of positive selection for

a GMF allele is wider than that for an AMF allele. Similarly, GMF alleles should be subject to the

noncovarying storage effect under broader conditions. However, a GMF allele may be considered

simply a beneficial allele as the arithmetic mean of offspring number is greater than the wild-type

allele. Such a mutation is probably rare and therefore was not considered in this study. One should

however note that GMF alleles might be still important for genetic variation in nature since they

can be maintained in balanced polymorphism in the conventional models of the storage effect, for

example without demographic fluctuations or in the case of sex-limited gene expressions.

Analytic approximation for the one-locus TP model predicts that, under soft selection or strict

density regulation that keeps the population size to the carrying capacity of the field at a given

time regardless of its genetic composition, balancing selection arises if the demographic

fluctuation (a major contributor to $\text{Cov}[S, \log[K_F/K_R]]$) is not so weak or strong compared to

fitness fluctuation ($\text{Var}[S]$ in case of $S^* = 0$). Here, balancing selection is defined as a force that

generates polymorphism at a level greater than expected for neutral alleles. Multi-locus

simulation showed that balanced polymorphism can be simultaneously maintained at many loci,

as expected since the demographic fluctuation (together with the presence of refuge) that turns

the geometric mean fitness of an AMF allele from < 1 to > 1 should apply equally to all loci in the

genome. Previously, Park and Kim (2019) showed that, in the presence of a refuge but without

fluctuation in population size, polymorphism can arise at multiple loci that contribute to the

expression of a phenotype under fluctuating selection. However, the number of loci where

balanced polymorphism emerged increased very slowly as the total number of loci in the model

increased. In contrast, in the current simulation, balanced polymorphism was observed at all loci

available for mutations to arise. However, the model containing only up to 10 loci was examined,

due to the limitation of the simulation method used, and the 10-loci polymorphism at each locus

was generally lower than the level observed in the one-locus simulation. This is probably because

the segregation of alleles at multiple loci elevates variance in the offspring number well above

that of the Poisson distribution, thus increasing genetic drift that weakens the positive selection

on rare alleles. The possibility that this mechanism can maintain the genome-wide polymorphism

over 1,000 loci, currently observed in several natural populations (Bergland, et al. 2014; Johnson,

et al. 2023), should be addressed by a simulation on a large scale in terms of both the number of

loci and the census population size.

This study focused more on the other evolutionary outcome of the noncovarying storage effect,

the fixation of mutations that amplify fitness fluctuations. In the TP model, the condition for the

fixation is less restrictive compared to that for balanced polymorphism, because it requires

relatively weaker selection (smaller variance in relative fitness) for a given demographic

fluctuation to satisfy, assuming $S^* = 0$, $\Phi > 1 + 1/(2\rho)$ with soft selection and $\Phi > 1/(2\rho)$

with hard selection. Similar ranges of conditions (for $\Phi'$) seem to apply for the LSA model (Table

S3). Demographic fluctuation that generates the noncovarying storage effect should apply equally

to all loci in the genome. Therefore, fixations occur at all available loci on which proper mutations

(*i.e.* the $A_2$ allele) arise, as observed in the multi-locus simulation. With soft selection, the pattern

of demographic fluctuation that generates the noncovarying storage effect does not change over

time regardless of any evolutionary change at any locus. However, with hard selection, fluctuation

in the relative size of the field generating the nocovarying storage effect at a given locus intensifies

as more fluctuation-amplifying mutations reach fixation at other loci. Since the carrying capacity

of the field in the TP model is effectively $(1 + S_t)^l K_F^{(t)}$, where $l$ is the number of loci where $A_2$

reached fixation, and wider fluctuation generates stronger positive selection on the rare $A_2$ (larger

$E[\Delta_2]$ in Appendix A), fixation at one locus facilitates the subsequent fixations at other loci. This

positive feedback between demographic fluctuation and positive selection drastically increases

the amplitude of fluctuation in the total population size (Fig. 4). How much the amplitude will

increase should depend on the number of loci at which mutations subject to hard selection arise.

There should be multiple independent ways in which changes in mutants' relative fitness can be

translated into changes in the mean absolute fitness of the entire population. For various life

history traits, one may imagine phenotypes (smaller eggs, smaller adult body, earlier onset of first

reproduction, etc.) that increase the absolute number of surviving offspring during a favorable,

resource-rich period but decrease the number in a bad period while not affecting the amount of

resources that the carriers of the alternative, ancestral alleles acquire. Then, many loci may be

available for hard selection, the combined effect of which may lead to a very large fluctuation in

population size.

The size of a population and its fluctuation should be governed not only by extrinsic factors (the

environmental variables such as temperature and parasite prevalence) but also by intrinsic

factors (species-specific responses to those extrinsic factors). The intrinsic factors such as various
life-history traits are the product of the species' evolutionary history. For example, phenotypic
plasticity responding to environmental cues can reduce the amplitude of the population size
fluctuation, thus acting as a mechanism of population regulation (Caswell 1983; Reed *et al.* 2010),
which according to the classical theory is adaptive as it enhances long-term reproductive success
by increasing the geometric mean fitness (Gillespie 1977; Pfister 1998). If life history traits
evolved in the direction toward suppressing random or cyclic fluctuations in the population size,
fluctuations observed in nature should be interpreted as the effect of variable environments that
the evolved mechanisms of population regulation were unable to contain. However, this study
suggests that evolution can proceed in the opposite direction. A large random fluctuation in
population size can be the product of adaptive evolution driven by positive feedback between
demographic and selective fluctuations. When the degrees of population size fluctuation are
different across species, the eco-evolutionary dynamics proposed in this study might be
particularly important for those exhibiting large fluctuations. Interestingly, annual desert plants
experiencing wider fitness fluctuations after germination have greater proportions of their
population remaining in the seed bank (Gremer & Venable 2014). This observation is in
agreement with the analytic results in this study because a larger seed bank corresponds to a
larger value of $\rho$, which makes it easier for the fixation of the fluctuation-amplifying alleles (i.e.,
lowering the threshold of $\Phi$ for positive selection).

DATA AVAILABILITY

Mathematica notebook files for conducting all simulations reported are available from
https://github.com/YuseobKimLab/RandomFluctuationEcoEvo

ACKNOWLEDGMENT

This study was supported by the National Research Foundation (NRF) grant 2020R1A2C1009261
funded by the Korean government.

CONFLICT OF INTEREST

The author declares no conflict of interest.

REFERENCES

Barabás G, D'Andrea R, Stump SM. (2018). Chesson's coexistence theory. *Ecological monographs*
88:277-303.

Bell, G. (2010). Fluctuating selection: the perpetual renewal of adaptation in variable
environments. *Philos Trans R Soc Lond B Biol Sci*, 365, 87-97.

Bergland AO, Behrman EL, O'Brien KR, Schmidt PS, Petrov DA. (2014). Genomic evidence of rapid
and stable adaptive oscillations over seasonal time scales in Drosophila. *PLoS Genet,* 10,
e1004775.

Bertram, J. & Masel, J. (2019). Different mechanisms drive the maintenance of polymorphism at
loci subject to strong versus weak fluctuating selection. *Evolution*, 73, 883-896.

Bogdziewicz, M., Ascoli, D., Hacket-Pain, A., Koenig, W.D., Pearse, I., Pesendorfer, M. *et al.* (2020).
From theory to experiments for testing the proximate mechanisms of mast seeding: an
agenda for an experimental ecology. *Ecology Letters*, 23, 210-220.

Caswell, H. (1983). Phenotypic plasticity in life-history traits: demographic effects and
evolutionary consequences. *American Zoologist*, 23, 35-46.

Chesson, P.L. (1994). Multispecies competition in variable environments. *Theoretical Population*
*Biology*, 45, 227-276.

Chesson, P.L. & Warner, R.R. (1981). Environmental variability promotes coexistence in lottery
competitive systems. *The American Naturalist*, 117, 923-943.

Dwyer, G., Dushoff, J. & Yee, S.H. (2004). The combined effects of pathogens and predators on
insect outbreaks. *Nature*, 430, 341-345.

Ellner, S. & Hairston, N.G. (1994). Role of overlapping generations in maintaining genetic variation
in a fluctuating environment. *The American Naturalist*, 143, 403-417.

Gillespie, J.H. (1977). Natural selection for variances in offspring numbers: a new evolutionary
principle. *The American Naturalist*, 111, 1010-1014.

Gremer, J.R. & Venable, D.L. (2014). Bet hedging in desert winter annual plants: optimal
germination strategies in a variable environment. *Ecol Lett*, 17, 380-387.

Gulisija, D. & Kim, Y. (2015). Emergence of long-term balanced polymorphism under cyclic
selection of spatially variable magnitude. *Evolution*, 69, 979-992.

Gulisija, D., Kim, Y. & Plotkin, J.B. (2016). Phenotypic plasticity promotes balanced polymorphism
in periodic environments by a genomic storage effect. *Genetics*, 202, 1437-1448.

Johnson OL, Tobler R, Schmidt JM, Huber CD. (2023). Fluctuating selection and the determinants

of genetic variation. *Trends in Genetics,* 39, 491-504

Kim, Y. (2023). Partial protection from fluctuating selection leads to evolution towards wider

population size fluctuation and a novel mechanism of balancing selection. *Proceedings of*

*the Royal Society B*, 290, 20230822.

Liu, M., Rubenstein, D.R., Liu, W.-C. & Shen, S.-F. (2019). A continuum of biological adaptations to

environmental fluctuation. *Proceedings of the Royal Society B*, 286, 20191623.

Lynch, M., Wei, W., Ye, Z. & Pfrender, M. (2024). The genome-wide signature of short-term

temporal selection. *Proceedings of the National Academy of Science USA*,

121:e2307107121.

Messer, P.W., Ellner, S.P. & Hairston, N.G. (2016). Can population genetics adapt to rapid evolution?

*Trends in Genetics*, 32, 408-418.

Myers, J.H. (1988). Can a general hypothesis explain population cycles of forest Lepidoptera? In:

*Advances in Ecological Research*. Elsevier, pp. 179-242.

Park Y, Kim Y. 2019. Partial protection from cyclical selection generates a high level of

polymorphism at multiple non-neutral sites. *Evolution*, 73, 1564-1577.

Pfister, C.A. (1998). Patterns of variance in stage-structured populations: evolutionary predictions

and ecological implications. *Proceedings of the National Academy of Sciences*, 95, 213-218.

Reed, T.E., Waples, R.S., Schindler, D.E., Hard, J.J. & Kinnison, M.T. (2010). Phenotypic plasticity and

population viability: the importance of environmental predictability. *Proceedings of the*

*Royal Society B: Biological Sciences*, 277, 3391-3400.

Reinhold, K. (2000). Maintenance of a genetic polymorphism by fluctuating selection on sex-

limited traits. *Journal of Evolutionary Biology*, 13, 1009-1014.

Slatkin, M. (1974). Hedging one's evolutionary bets. *Nature*, 250, 704-705.

Thompson, J.N. (2013). *Relentless evolution*. University of Chicago Press.

Tuljapurkar, S. & Orzack, S.H. (1980). Population dynamics in variable environments I. Long-run

growth rates and extinction. *Theoretical Population Biology*, 18, 314-342.

Tuljapurkar, S.D. (1982). Population dynamics in variable environments. III. Evolutionary

dynamics of r-selection. *Theoretical population biology*, 21, 141-165.

Turelli, M., Schemske, D.W. & Bierzychudek, P. (2001). Stable two-allele polymorphisms

maintained by fluctuating fitnesses and seed banks: protecting the blues in Linanthus

parryae. *Evolution*, 55, 1283-1298.

Wallace, B. (1975). Hard and soft selection revisited. *Evolution*, 465-473.

White, T. (2008). The role of food, weather and climate in limiting the abundance of animals.

*Biological Reviews*, 83, 227-248.

Yamamichi, M., Letten, A.D. & Schreiber, S.J. (2023). Eco-evolutionary maintenance of diversity in

fluctuating environments. *Ecology Letters*, 26, S152-S167.

Zhang, J. (2023). Patterns and evolutionary consequences of pleiotropy. *Ann Rev Ecol Evol Syst*

54:1-19.

**Appendix A. Derivation of conditions for balancing and directional selection on $A_2$ in the**
**TP model**

Here only deterministic changes in allele frequencies are tracked as a very large population is assumed. At the end of reproduction but before migration into subpopulations at generation $t$ - 1, there are $n_1$ and $n_2$ copies (individuals) of $A_1$ and $A_2$ in the entire population. After migration, the numbers of $A_1$ and $A_2$ copies are $n_{1F} = (1-r)n_1$ and $n_{2F} = (1-r)n_2$ in the field and $n_{1R} = rn_1$ and $n_{2R} = rn_2$ in the refuge. Therefore, the relative frequencies of alleles are identical between the field and refuge before reproduction at the beginning of generation $t$.

Soft selection is first considered. Then, each $A_2$ allele produces $(1 + S_t)K_F^{(t)}/\bar{N}_F^{(t)}$ copies of descendants, where $\bar{N}_F^{(t)} = n_{1F} + (1 + S_t)n_{2F} = (1-r)(n_1 + (1 + S_t)n_2)$. Since $n_1 \gg n_2$ after $A_2$ appears by mutation, $n_1 + (1 + S_t)n_2 \approx n_1$ and $\bar{N}_F^{(t)} \approx \tilde{N}_F^{(t)} = (1-r)(n_1 + n_2) = (1-r)\left(K_F^{(t-1)} + K_R^{(t-1)}\right)$. Similar approximations apply to the refuge. Let $n'_2$ be the expected number of $A_2$ in generation $t$. This leads to eq. (1). Namely,

$$\frac{n'_2}{n_2} = (1-r)\frac{(1+S_t)K_F^{(t)}}{\bar{N}_F^{(t)}} + r\frac{(1+S_t^*)K_R^{(t)}}{\bar{N}_R^{(t)}} \approx (1-r)\frac{(1+S_t)K_F^{(t)}}{\tilde{N}_F^{(t)}} + r\frac{(1+S_t^*)K_R^{(t)}}{\tilde{N}_R^{(t)}} = \frac{(1+S_t)K_F^{(t)}+(1+S_t^*)K_R^{(t)}}{K_F^{(t-1)}+K_R^{(t-1)}}$$

Then, with $K_F^{(t-1)} = K_{F0}e^U$, $K_R^{(t-1)} = K_{R0}e^V$, $K_F^{(t)} = K_{F0}e^{U'}$, and $K_R^{(t)} = K_{R0}e^{V'}$,

$$\Delta_2 \equiv \log\left[\frac{n'_2}{n_2}\right] = \log[e^{U'}(1 + S) + \rho e^{V'}(1 + S^*)] - \log[e^U + \rho e^V],$$

where $S = S_t$, $S^* = S_t^*$, and $\rho = K_{R0}/K_{F0}$. If $E[\Delta_2] > 0$, where the expectation is over demographic and fitness fluctuations, the rare allele $A_2$ is expected to increase its copy number over time. As the expectations of $U$, $V$, $U'$, $V'$, $S$, and $S^*$ are all zero and non-zero correlations exist only among $U'$, $V'$, $S$, and $S^*$ and between $U$ and $V$, from the Taylor expansion of $\Delta_2$ up to the second order, it can be shown that

$$\mathrm{E}[\Delta_2] \approx \frac{1}{2}\alpha_{U'U'}\mathrm{Var}[U'] + \frac{1}{2}\alpha_{V'V'}\mathrm{Var}[V'] + \frac{1}{2}\alpha_{UU}\mathrm{Var}[U] + \frac{1}{2}\alpha_{VV}\mathrm{Var}[V] + \frac{1}{2}\alpha_{SS}\mathrm{Var}[S]$$

$$+ \frac{1}{2}\alpha_{S^*S^*}\mathrm{Var}[S^*] + \alpha_{U'V'}\mathrm{Cov}[U',V'] + \alpha_{UV}\mathrm{Cov}[U,V] + \alpha_{SS^*}\mathrm{Cov}[S,S^*]$$

$$+ \alpha_{SU'}\mathrm{Cov}[S,U'] + \alpha_{SV'}\mathrm{Cov}[S,V'] + \alpha_{S^*U'}\mathrm{Cov}[S^*,U'] + \alpha_{S^*V'}\mathrm{Cov}[S^*,V']$$

where $\alpha_{XY} = \frac{\partial^2 \Delta_2}{\partial X \partial Y}$ is the Taylor coefficient for variables $X$ and $Y$ evaluated at

$(U,V,U',V',S,S^*) = (0,0,0,0,0,0)$. Since $\alpha_{U'U'} = \alpha_{V'V'} = \alpha_{UV} = \alpha_{SU'} = \alpha_{S^*V'} = \frac{\rho}{(1+\rho)^2}$, $\alpha_{UU} =$

$\alpha_{VV} = \alpha_{U'V'} = \alpha_{SV'} = \alpha_{S^*U'} = \alpha_{SS^*} = -\frac{\rho}{(1+\rho)^2}$ , $\alpha_{SS} = -\frac{1}{(1+\rho)^2}$, $\alpha_{S^*S^*} = -\frac{\rho^2}{(1+\rho)^2}$, $\mathrm{Var}[U] =$

$\mathrm{Var}[U'], \mathrm{Var}[V] = \mathrm{Var}[V']$, and $\mathrm{Cov}[U,V] = \mathrm{Cov}[U',V']$, the above equation is simplified to

$$\mathrm{E}[\Delta_2] = \frac{2\rho\mathrm{Cov}[S-S^*,U'-V'] - \mathrm{Var}[S] - \rho^2\mathrm{Var}[S^*] - 2\rho\mathrm{Cov}[S,S^*]}{2(1+\rho)^2}.$$

Therefore, the condition for the $A_2$ allele being positively selected (i.e. invading the population

initially fixed for $A_1$), is $\mathrm{E}[\Delta_2] > 0$ or

$$\frac{\mathrm{Cov}[S-S^*,U'-V']}{\mathrm{Var}[S] + \rho^2\mathrm{Var}[S^*] + 2\rho\mathrm{Cov}[S,S^*]} > \frac{1}{2\rho}$$

assuming $(\mathrm{Var}[S] + \rho^2\mathrm{Var}[S^*])/\rho > -2\mathrm{Cov}[S,S^*]$. If $S_t^* = \delta S_t$, the above condition becomes

$$\Phi = \frac{\mathrm{Cov}[S,U'-V']}{\mathrm{Var}[S]} = \frac{\mathrm{Cov}[S,\log[K_F/K_R]]}{\mathrm{Var}[S]} > \frac{(1+\delta\rho)^2}{2(1-\delta)\rho}.$$

A similar result can be obtained for the condition for the $A_1$ allele to invade a population initially

fixed for $A_2$. The expected change in the number of $A_1$, $n'_1/n_1$, is obtained with a procedure

identical to that leading to eq. (1) except that the relative fitness is flipped. Therefore

$$\frac{n'_1}{n_1} \approx \frac{K_F^{(t)}}{K_F^{(t-1)} + K_R^{(t-1)}}\frac{1}{1+S_t} + \frac{K_R^{(t)}}{K_F^{(t-1)} + K_R^{(t-1)}}\frac{1}{1+S_t^*} = \frac{(1+S_t^*)e^{U'} + (1+S_t)\rho e^{V'}}{(1+S_t)(1+S_t^*)(e^U + \rho e^V)}$$

for $n_1 \ll n_2$, the dynamics of the rare $A_1$ allele depends on the expectation of

$$\Delta_{1,\mathrm{soft}} \equiv \log\left[\frac{n'_1}{n_1}\right] = \log[(1+S_t^*)e^{U'} + (1+S)\rho e^{V'}] - \log[(1+S)(1+S_t^*)(e^U + \rho e^V)].$$

Applying the quadratic approximation used for obtain $\mathrm{E}[\Delta_2]$ above, one obtains

$$\mathrm{E}[\Delta_{1,\mathrm{soft}}] = \frac{(1+2\rho)\mathrm{Var}[S] + (2\rho+\rho^2)\mathrm{Var}[S^*] - 2\rho\mathrm{Cov}[S,S^*] - 2\rho\mathrm{Cov}[S-S^*,U'-V']}{2(1+\rho)^2}$$

The condition for the rare $A_1$ allele to increase in frequency is $\mathrm{E}[\Delta_{1,\mathrm{soft}}] > 0$ or

730 
$$\frac{\mathrm{Cov}[S-S^*,U'-V']}{(1+2\rho)\mathrm{Var}[S]+(2\rho+\rho^2)\mathrm{Var}[S^*]-2\rho\mathrm{Cov}[S,S^*]}<\frac{1}{2\rho}.$$

731 If $S_t^*=\delta S_t$, the above condition becomes

732 
$$\Phi<1-\delta+\frac{(1+\delta\rho)^2}{2(1-\delta)\rho}.$$

With hard selection, the frequency change of $A_2$ when rare is again given by eq. (1). Therefore, the

condition for $A_2$ invading the population fixed for $A_1$ ($\mathrm{E}[\Delta_2]>0$ above) is no different from that

for soft selection. However, because the carrying capacities of the field and refuge fixed for $A_2$ at

generation $t$ is given by $(1+S_t)K_F^{(t)}$ and $(1+S_t^*)K_R^{(t)}$ and $\widetilde{N}_F^{(t)}=(1-r)((1+S_{t-1})K_F^{(t-1)}+$

$(1+S_{t-1}^*)K_R^{(t-1)})$ and $\widetilde{N}_R^{(t)}=r((1+S_{t-1})K_F^{(t-1)}+(1+S_{t-1}^*)K_R^{(t-1)})$ under hard selection, the

opposite condition for $A_1$ invading $A_2$ is given by

739 
$$\frac{n_1'}{n_1}=(1-r)\frac{(1+S_t)K_F^{(t)}}{\widetilde{N}_F^{(t)}}\frac{1}{1+S_t}+r\frac{(1+S_t^*)K_R^{(t)}}{\widetilde{N}_R^{(t)}}\frac{1}{1+S_t^*}=\frac{K_F^{(t)}+K_R^{(t)}}{(1+S_{t-1})K_F^{(t-1)}+(1+S_{t-1}^*)K_R^{(t-1)}}=\frac{e^{U'}+\rho e^{V'}}{(1+S_{t-1})e^U+(1+S_{t-1}^*)\rho e^V}$$

740 Then,

741 
$$\Delta_{1,\mathrm{hard}}\equiv\log\left[\frac{n_1'}{n_1}\right]=\log[e^{U'}+\rho e^{V'}]-\log[(1+S_{t-1})e^U+(1+S_{t-1}^*)\rho e^V].$$

Now, given that $(U',V',S_t,S_t^*)$ and $(U,V,S_{t-1},S_{t-1}^*)$ are independent samples of a joint

distribution, the expectation of $-\Delta_{1,\mathrm{hard}}$ is exactly that of $\Delta_2$ above. Therefore, the condition for

the fixation of $A_2$ ($\mathrm{E}[\Delta_{1,\mathrm{hard}}]<0$) is exactly the condition for positive selection on a rare $A_2$

($\mathrm{E}[\Delta_2]>0$).

**Appendix B. Methods of stochastic simulation**

Stochastic simulations of the single- and multi-locus eco-evolutionary models were implemented

in Mathematica ver. 11.3. In both TP and LSA models, an individual has $L$ loci that are arranged

linearly with probability $c$ for recombination between adjacent loci per generation. The

simulation keeps track of two sets of lists: one containing the sets of haplotypes carried by non-

zero individuals in the field and in the refuge ($\boldsymbol{H}_F$ and $\boldsymbol{H}_R$) and the other containing the

corresponding numbers of individiuals ($\boldsymbol{N}_F$ and $\boldsymbol{N}_R$).

In the TP model, generation $t$ begins after the migration step (re-distribution of individuals

produced in the previous generation into the field and refuge) is completed. Haplotype numbers

are first updated by mutation and recombination. A Poisson variate with mean $NL\mu$, where $N$ is

the current size of a subpopulation at hand, is drawn as the total number of mutation events and

each event is assigned to haplotypes and loci proportional to their current frequencies. Similarly,

for $L > 1$, a Poisson variate with mean $N(L-1)c$ is drawn as the total number of recombination

events in the subpopulation. For each recombination event, two haplotypes are chosen

proportional to their frequencies and one of the $L$-1 intervals is chosen to be the position of

crossing-over between the two haplotypes. Newly created or lost haplotypes during mutation and

recombination steps are tracked by updating ($H_F$, $H_R$) and ($N_F$, $N_R$). Then, the updated lists are

further updated in the step of reproduction according to their fitness. Under soft selection, the

number of individuals with haplotype $i$ born in the next generation is Poisson distributed with

mean $W_i \widetilde{N}_{F,i}^{(t)} K_F^{(t)} / \sum_j W_j \widetilde{N}_{F,j}^{(t)}$ in the field and $\widetilde{N}_{R,i}^{(t)} K_R^{(t)} / \sum_j \widetilde{N}_{R,j}^{(t)}$ in the refuge, where $W_i$ is the

relative fitness of haplotype $i$ and $\widetilde{N}_{F,i}^{(t)}$ ($\widetilde{N}_{R,i}^{(t)}$) is the number of haplotype $i$ individuals in the field

(refuge) after the haplotypes are updated by mutation and recombination. Under hard selection,

the corresponding mean number in the field is $W_i \widetilde{N}_{F,i}^{(t)} K_F^{(t)} / \sum_j \widetilde{N}_{F,j}^{(t)}$. Finally, when there are $N_F$

and $N_R$ individuals in the field and refuge, Poisson variates with the mean $N_F m_{FR}$ ($N_R m_{RF}$) truncated

at the upper bound $N_F$ ($N_R$) is drawn to determine the numbers of migrants from field to refuge

(from refuge to field). Migrants are chosen proportional to the haplotype frequencies and then

merged with the residents of the destination. This completes a single-generation iteration in

updating haplotype lists.

Simulation for the LSA model is similarly performed by tracking the haplotypes. Let $l_i$, $m_i$, and $n_i$

be the number of individuals with haplotype $i$ ($i$ = 1, ..., $2^L$) at the larval, subadult, and adult stages

at time $t$. The following steps are taken to produce the corresponding numbers, $l_i'$, $m_i'$, and $n_i'$, at

time $t$+1. As in the TP model, bidirectional mutations with probability $\mu$ per locus and

recombination with probability $c$ for adjacent loci occur but in subadult and adult stages only.

Recombination must occur among subadults and among adults only. The updated numbers of

individuals by haplotypes in subadults and adults after mutations and recombination are $m_i$*, and

$n_i$*. Next, the numbers of individuals by haplotypes in the next generation are obtained. $l_i'$ is given

by a Poisson number with mean $W_i^{(t)}(m_i^* + n_i^*)K_L^{(t)} / \sum_i(W_i^{(t)}(m_i^* + n_i^*))$ under soft selection

and $W_i^{(t)}(m_i^* + n_i^*)K_L^{(t)} / \sum_i(m_i^* + n_i^*)$ under hard selection. $m_i'$ and $n_i'$ are given by Poisson

numbers with mean $e_L l_i$ and $e_L e_S m_i$*, respectively.

 Table S1. One-locus simulation with variable migration rates in the TP model

| mode | $m_{FR}$ | $m_{RF}$ | $\bar{H}$ | $P_{01}$ | $P_{90}$ |
|---|---|---|---|---|---|
| neutral ($a_S = b_S = 0$, $a_U = 0.3$, $a_V = 0.05$, $b_U = b_V = 0.1$) | [0.495, 0.505] | [0.495, 0.505] | 0.0633 | 0.60 | 0.24 |
| | [0.1, 0.2] | [0.1, 0.2] | 0.0657 | 0.61 | 0.24 |
| soft selection, $\Phi = 1$ ($a_S = 0.2$, $b_S = 0.1$, $a_U = 0.3$, $a_V = 0.05$, $b_U = b_V = 0.1$) | [0.495, 0.505] | [0.495, 0.505] | 0.196 | 0.28 | 0.25 |
| | [0.3, 0.7] | [0.3, 0.7] | 0.177 | 0.45 | 0.14 |
| | [0.3, 0.5] | [0.5, 0.7] | 0.135 | 0.42 | 0.26 |
| | [0.5, 0.7] | [0.3, 0.5] | 0.147 | 0.42 | 0.25 |
| | [0.1, 0.5] | [0.1, 0.5] | 0.142 | 0.59 | 0.08 |
| | [0.2, 0.4] | [0.2, 0.4] | 0.118 | 0.70 | 0.03 |
| | [0.1, 0.2] | [0.1, 0.2] | 0.056 | 0.88 | 0.00 |
| | [0.1. 0.2] | [0.4, 0.5] | 0.186 | 0.34 | 0.22 |
| | [0.4, 0.5] | [0.1, 0.2] | 0.052 | 0.88 | 0.01 |
| Hard selection, $\Phi = 0.8$ ($a_S = 0.2$, $b_S = 0.15$, $a_U = 0.3$, $a_V = 0.05$, $b_U = b_V = 0.1$) | [0.495, 0.505] | [0.495, 0.505] | 0.030 | 0.05 | 0.89 |
| | [0.3, 0.7] | [0.3, 0.7] | 0.020 | 0.01 | 0.95 |
| | [0.2, 0.4] | [0.2, 0.4] | 0.074 | 0.45 | 0.39 |
| | [0.1, 0.2] | [0.1, 0.2] | 0.034 | 0.93 | 0.00 |
| | [0.1. 0.2] | [0.4, 0.5] | 0.038 | 0.02 | 0.90 |
| | [0.4, 0.5] | [0.1, 0.2] | 0.036 | 0.91 | 0.01 |

792 Table S2. Mean time to the fixation ($q > 0.9$) of $A_2$ in the multi-locus simulation of the TP model

| mode | $L$ | $c$ | $\bar{T}$ | c. v.* |
|---|---|---|---|---|
| | 1 | - | 5863 | - |
| soft selection, $\Phi = 2$ | 5 | $10^{-4}$ | 9287 | $0.66 \pm 0.21$ |
| ($a_S = 0.1$, $b_S = 0$, $a_U = 0.25$, | 5 | $10^{-3}$ | 8212 | $0.67 \pm 0.20$ |
| $a_V = 0.05$, $b_U = b_V = 0.05$) | 5 | 0.01 | 8621 | $0.70 \pm 0.26$ |
| | 5 | 0.1 | 8162 | $0.65 \pm 0.26$ |
| | 1 | - | 6242 | - |
| hard selection, $\Phi = 0.75$ | 5 | $10^{-4}$ | 3776 | $0.34 \pm 0.17$ |
| ($a_S = 0.2$, $b_S = 0$, $a_U = 0.25$, | 5 | $10^{-3}$ | 3228 | $0.32 \pm 0.16$ |
| $a_V = 0.1$, $b_U = b_V = 0.05$) | 5 | 0.01 | 3030 | $0.31 \pm 0.20$ |
| | 5 | 0.1 | 3459 | $0.28 \pm 0.19$ |
| | 1 | - | 5177 | - |
| hard selection, $\Phi = 1$ | 5 | $10^{-4}$ | 4648 | $0.35 \pm 0.15$ |
| ($a_S = 0.15$, $b_S = 0$, $a_U = 0.15$, | 5 | $10^{-3}$ | 4018 | $0.35 \pm 0.17$ |
| $b_U = 0.05$, $a_V = 0$, $b_V = 0.1$) | 5 | 0.01 | 3969 | $0.32 \pm 0.20$ |
| | 5 | 0.1 | 3467 | $0.37 \pm 0.21$ |

793 *Coefficient of variation for 5 consecutive times ($T_1$, ..., $T_5$) observed in a single simulation run,
794 averaged over replicates ($\pm$ standard deviation).

795 All results are based on 100 replicates for each parameter set

797      Table S3. Long-term polymorphism in multi-locus simulations of the LSA model

| mode | $a_S$ | $b_S$ | $\Phi'$ | $\bar{H}$ | $P_{01}$ | $P_{90}$ |
|------|-------|-------|---------|-----------|----------|----------|
| neutral | 0 | 0 | - | 0.0757 | 0.42 | 0.40 |
| soft selection | 0.05 | 0.2 | 0.294 | 0.013 | 1 | 0 |
| | 0.1 | 0.2 | 0.5 | 0.017 | 0.96 | 0 |
| | 0.1 | 0.15 | 0.769 | 0.040 | 0.91 | 0.00 |
| | 0.2 | 0.1 | 1 | 0.152 | 0.63 | 0.08 |
| | 0.2 | 0 | 1.25 | 0.116 | 0.31 | 0.41 |
| | 0.15 | 0.05 | 1.5 | 0.054 | 0.17 | 0.71 |
| | 0.1 | 0.05 | 2 | 0.023 | 0.22 | 0.73 |
| hard selection | 0.05 | 0.2 | 0.294 | 0.006 | 0.99 | 0 |
| | 0.1 | 0.2 | 0.5 | 0.013 | 0.97 | 0 |
| | 0.1 | 0.15 | 0.769 | 0.009 | 0.22 | 0.76 |
| | 0.2 | 0.1 | 1 | 0.003 | 0.07 | 0.93 |
| | 0.15 | 0.087 | 1.25 | 0.003 | 0.24 | 0.76 |
| | 0.15 | 0.05 | 1.5 | 0.005 | 0.06 | 0.92 |

- Each simulation ran for 50,000 time units, except for the neutral simulation that ran 500,000
time units.

- Other parameters: $\mu = 2\times10^{-5}$, $K_{L0} = 5000$, $a_L = 0.25$, $b_L = 0.1$, $e_L = 0.7$, $e_S = 0.7$, $L = 4$, $c = 0.03$.

Figure S1

Figure S1 legend: Exemplary trajectories of allele ($A_2$) frequencies in simulations. Allele
frequencies are plotted in different colors for different loci. The number of loci ($L$) and
recombination rate ($c$) are shown on the right side of graphs. Other parameters: $K_{R0} = K_{F0} =$
1000, $\mu = 2\times10^{-5}$, $\Phi = 1$ ($a_S = 0.2$ [0 for neutral], $b_S = 0.1$, $a_U = 0.3$, $b_U = 0.1$, $a_V = 0.05$, $b_V = 0.1$).

Figure S2

Figure S2 (continued)

Figure S2 legend: Examples of simulation runs in which the mutant allele ($A_2$) reaches fixation

sequentially at all 8 loci used in the TP model. The change of population size is shown below

that of allele frequencies, plotted in different colors for different loci. Parameter values are

shown on the right side of graphs. Other parameters: $K_{R0} = K_{F0} = 1000$, $\mu = 2 \times 10^{-5}$.

Figure S3

Figure S3 Legend: Exemplary allele frequency changes in the simulation of the LSA model. Each

simulation started with a population fixed for the $A_1$ allele and ran for 50,000 time units, except

for the case of hard selection with $\Phi' = 1$. The $A_2$ allele appears by recurrent mutations at each of

$L = 4$ loci. Allele frequencies are plotted in different colors for different loci. For hard selection,

the change of population size (blue: larval, green: subadult, gray: adult) is shown below the

allele frequency trajectories. Other parameters: $K_{L0} = 5000$, $a_L = 0.25$, $b_L = 0.1$, $e_L = e_S = 0.7$, $\mu =$

$2 \times 10^{-5}$, $c = 0.03$.

Figure S4

Figure S4 legend: Mean fixation time ($\bar{T}$) with varying number of loci (A; $L$ = 1, 2, and 4) and the
coefficient of variation in fixation times with $L$ = 4 in the multi-locus simulation of the LSA model.
Error bars show two times the standard errors. A total of 100 replicates of simulations were run
under soft selection with $\Phi'$ = 2 (cyan; $a_S$ = 0.1, $b_S$ = 0.05, $a_L$ = 0.25, $b_L$ = 0.1, $\mu$ = 3×10⁻⁵), hard
selection with $\Phi'$ = 1 (brown; $a_S$ = 0.2, $b_S$ = 0.1, $a_L$ = 0.25, $b_L$ = 0.1, $\mu$ = 2×10⁻⁵), or hard selection with
$\Phi'$ = 0.77 (orange; $a_S$ = 0.1, $b_S$ = 0.15, $a_L$ = 0.25, $b_L$ = 0.1, $\mu$ = 3×10⁻⁵). Other parameters are identical
to those in Figure S3.