Biosynthesis of glucosaminyl phosphatidylglycerol in *Pseudomonas aeruginosa*

Fabiha Zaheen Khan[1], Kelli L. Palmer[1]*, and Ziqiang Guan[2]*

[1]Department of Biological Sciences, The University of Texas at Dallas, Richardson, Texas, USA;

[2]Department of Biochemistry, Duke University School of Medicine, Durham, North Carolina, USA.

*Corresponding authors

Email: Kelli Palmer (Kelli.Palmer@utdallas.edu) or Ziqiang Guan (ziqiang.guan@duke.edu)

Abstract

Glucosaminyl phosphatidylglycerol (GlcN-PG) was first identified in bacteria in the 1960s and was recently reported in *Pseudomonas aeruginosa*. Despite the important implications in altering membrane charge (by the modification of anionic PG with cationic glucosamine), the biosynthesis and functions of GlcN-PG have remained uncharacterized. Using bioinformatic and lipidomic analysis, we identified a 3-gene predicted operon, renamed as *gpgSDF*, that is responsible for the biosynthesis and potential transport of GlcN-PG in *P. aeruginosa*: *gpgS* encodes a novel glycotransferase that is responsible for the modification of phosphatidylglycerol (PG) with *N*-acetylglucosamine (GlcNAc) to produce GlcNAc-PG, and *gpgD* encodes a novel deacetylase that removes the acetyl group from GlcNAc-PG to produce GlcN-PG. The third gene, *gpgF*, is predicted to encode a flippase whose activity remains to be experimentally verified. A *P. aeruginosa gpgD* transposon mutant accumulates GlcNAc-PG and lacks GlcN-PG, and as expected, the complementation of *gpgD* restores the production of GlcN-PG. Moreover, the heterologous expression of *gpgSDF* in *Escherichia coli* resulted in production of both GlcNAc-PG and GlcN-PG. The identification of the biosynthetic genes of GlcN-PG paves the way for the investigation of its biological and pathological functions, which has significant implications in our understanding of the unique membrane physiology, pathogenesis and antimicrobial resistance of *P. aeruginosa.*

Importance

The identification of the biosynthetic genes of GlcN-PG paves the way for the investigation of its biological and pathological functions, which has significant implications in our understanding of the unique membrane physiology, pathogenesis and antimicrobial resistance of *P. aeruginosa.*

Introduction

Pseudomonas aeruginosa is a major Gram-negative pathogen widely responsible for pneumonia, surgical infection, bacteremia, and other life-threatening infections in immunocompromised individuals with underlying diseases such as cystic fibrosis (CF) (1, 2) and cancer (3). *P. aeruginosa* possesses an unusual ability to colonize diverse environments and rapidly develop antibiotic resistance. Multi-drug-resistant (MDR) and extreme drug-resistant (XDR) high-risk strains are widespread in healthcare settings, making the treatment of certain *P. aeruginosa* infections extremely challenging (4-6).

An important mechanism for bacteria to cope with antimicrobial stresses and develop antibiotic resistance is by altering the charge of their membrane lipids. A highly conserved and perhaps most studied enzyme responsible for modifying lipids in bacteria is MprF (multiple peptide resistance factor), which modifies lipids via the transfer of amino acids from charged tRNAs to the head groups of anionic phosphatidylglycerol (PG) and cardiolipin (CL) (7, 8), as well as to neutral glycolipids (9, 10). *P. aeruginosa* encodes MprF and produces alanine-modified PG (11). Additionally in *P. aeruginosa*, modifications of the phosphate groups of lipid A (the lipid anchor of lipopolysaccharide) with positively charged moieties such as aminoarabinose and phosphoethanolamine are critical for conferring resistance to cationic antimicrobial peptides and polymyxin (2, 12-16).

The modification of PG with cationic glucosamine, producing zwitterionic glucosaminyl phosphatidylglycerol (GlcN-PG), was recently reported in *P. aeruginosa* (17). Despite the obvious implications in altering membrane charge, and the fact that GlcN-PG was first observed in various bacteria in the 1960s (18-21), the biosynthesis and functions of GlcN-PG have remained uncharacterized.

In this study, by combining bioinformatic, genetic, and lipidomic approaches, we identified the biosynthetic genes required for GlcN-PG synthesis in *P. aeruginosa*. Using the PA14 strain as a model, we show that a 3-gene predicted operon (consisting of PA14_56030, PA14_56040 and PA14_56050) is required for the synthesis and possible transport of GlcN-PG. We have thus renamed this operon as *gpgSDF* (GlcN-PG synthase, deacetylase and flippase). Since *gpgSDF* is conserved in other strains of *P. aeruginosa*, this work has significant implications for understanding the mechanisms of physiology and pathogenesis of *P. aeruginosa*.

Experimental Procedures

Bacterial Strains and Routine Growth Conditions

The bacterial strains used in the study are described in Table 1. For lipidomic analysis, bacterial cells were grown in Luria Bertani (LB) broth (10g/L tryptone, 5g/L yeast extract, 5g/L NaCl) at 37 °C with shaking at 220-250 rpm for 18-20 hours. The optical density (OD) was measured in a disposable cuvette (Thermo Fisher Scientific, Waltham, MA) as absorption at 600 nm by a spectrophotometer (Thermo Scientific Genesys 30).

The *P. aeruginosa* PA14 MAR2xT7 transposon mutants carrying a gentamicin resistance cassette were previously reported (22) and were grown in LB broth containing 15 µg/ml gentamicin. *Escherichia coli* DH5α (Thermo Fisher Scientific) with the plasmid pGEM-T (Promega PR-A3600) was grown in LB broth containing 50 µg/ml ampicillin.

Complementation of the *gpgD* transposon mutant of PA14

The PA14 *gpgD* gene was amplified using primers 1 and 2 (Table S1), which introduced regions complementary to the plasmid pMQ72 (23) (obtained from Dr. Peter Jorth). The pMQ72 plasmid was digested using EcoRI and HindIII. The *gpgD* amplicon and digested pMQ72 were used for Gibson assembly. The reaction was transformed into chemically competent *E. coli* DH5α cells. Gentamicin at 30 µg/mL was used for selection of pMQ72 in *E. coli*. The pMQ72-*gpgD* plasmid was extracted using the Thermo Scientific GeneJET Plasmid Extraction Kit following the manufacturer's protocol, and the *gpgD* sequence was confirmed by Sanger sequencing at the University of Texas at Dallas Genome Center using primer pairs 3 and 4, 5 and 6, 7 and 8, and 9 and 10 (Table S1). After confirmation, the pMQ72-*gpgD* plasmid was amplified using primers 11 and 12 (Table S1) to generate a linear fragment that excluded the gentamicin resistance cassette. The tetracycline resistance cassette from pEX18Tc (24) was amplified using primers 13 and 14, which introduced overhangs corresponding to pMQ72 (Table S1). The pMQ72 fragment and the tetracycline resistance cassette were assembled as described above. Successful insertion of the tetracycline cassette was confirmed using primers 15 and 16 (Table S1). The resulting plasmid, pMQ72-Tet-*gpgD*, was chemically transformed into *E. coli* S17 cells (obtained from Dr. Trusha Parekh), and transformants were selected with tetracycline at 10 µg/mL. The S17 transformants were replica-plated on LB plates with 30 µg/mL gentamicin, where no growth was observed, as expected. The pMQ72-Tet-*gpgD* plasmid was conjugated into PA14 *gpgD*::Tn cells using *E. coli*

S17 as the donor. Transconjugant colonies were selected on Vogel-Bonner minimal medium (VBMM) agar with 100 µg/mL ampicillin and 100 µg/mL tetracycline (25). Transconjugant colonies were confirmed by colony PCR using primer pairs 3 and 10, and 15 and 16 (Table S1). To induce expression of *gpgD* from the *araBAD* promoter of pMQ72, PA14 *gpgD*::Tn(pMQ72-Tet-*gpgD*) were grown to late exponential phase in LB (OD$_{600}$~1.0) with 100 µg/mL tetracycline +/- 1% L-arabinose (26). Cells were pelleted for acidic Bligh-Dyer lipid extractions and normal-phase LC/MS analysis.

Heterologous expression of *gpgSDF* in *E. coli*

The *gpgSDF* predicted operon was amplified by PCR using specific primers (Table S1) and cloned into pGEM-T (27) by TA cloning per the manufacturer's instructions. The newly assembled pWO-GEM-T plasmid was confirmed through PCR reactions with T7 and SP6 and sequencing primer pairs with agarose gel electrophoresis analysis of products. The plasmid was transformed into *E. coli* DH5α and maintained by growth in the presence of 50 µg/ml of ampicillin. Cells were pelleted for acidic Bligh-Dyer lipid extractions and normal-phase LC/MS analysis.

Acidic Bligh-Dyer Lipid Extraction

The acidic Bligh-Dyer lipid extractions were performed as previously described (9, 28). Briefly, bacterial cells were grown in 20 mL of LB broth for ~20 hours; 19 mL of each culture was pelleted at 7000 rpm for 10 min. The cell pellet was washed twice with 1X phosphate buffer saline (PBS, Sigma-Aldrich, St. Louis, MO). Each cell pellet was resuspended in 0.8 mL of 1X PBS and transferred to a 17-mL glass tube with a Teflon-lined cap (Corning Pyrex^{TM}, VWR, Radnor, PA). Afterwards, 1 mL of chloroform and 2 mL of methanol were added to generate a single-phase Bligh-Dyer solution. This solution was incubated for 20 min at room temperature with intermittent mixing. After centrifugation at 1700 rpm for 10 min, the supernatant was transferred to a new glass tube, which was followed by the addition of 100 µl HCl (37%), 0.9 mL 1X PBS and 1 mL chloroform to generate a two-phase Bligh-Dyer solution. After mixing by a vortex, the solution was separated into two phases by centrifugation at 1700 rpm for 5 min at room temperature. The lower phase was recovered and dried under a stream of nitrogen gas before being stored at −80˚C prior to lipidomic analysis.

Normal Phase LC/MS/MS Analysis

Lipidomic analysis by normal phase LC/MS/MS was described previously (9, 28). Briefly, normal phase LC was performed on an Agilent 1200 Quaternary LC system equipped with an Ascentis Silica HPLC column, 5 μm, 25 cm x 2.1 mm (Sigma-Aldrich, St. Louis, MO). Mobile phase A consisted of chloroform/methanol/aqueous ammonium hydroxide (800:195:5, v/v); mobile phase B consisted of chloroform/methanol/water/aqueous ammonium hydroxide (600:340:50:5, v/v); mobile phase C consisted of chloroform/methanol/water/aqueous ammonium hydroxide (450:450:95:5, v/v). The elution program consisted of the following: 100% mobile phase A was held isocratically for 2 min and then linearly increased to 100% mobile phase B over 14 min and held at 100% B for 11 min. The LC gradient was then changed to 100% mobile phase C over 3 min and held at 100% C for 3 min, and finally returned to 100% A over 0.5 min and held at 100% A for 5 min. The LC eluent (with a total flow rate of 300 μl/min) was introduced into the ESI source of a high resolution TripleTOF5600 mass spectrometer (Sciex, Framingham, MA). The instrumental settings for negative ion ESI/MS and MS/MS analysis of lipid species were as follows: IS = -4500 V; CUR = 20 psi; GSI = 20 psi; DP = -55 V; and FP = -150 V. The MS/MS analysis used nitrogen as the collision gas. Analyst TF1.5 software (Sciex, Framingham, MA) was used for data analysis. All analysis was performed by at least duplicate experiments.

Results

Identification of GlcN-PG in *P. aeruginosa* by LC/MS/MS

The lipid extract of *P. aeruginosa* strain PA14 cells was analyzed by normal phase LC/MS in both negative ion and positive ion modes. As shown by the negative total ion chromatogram (TIC) (Fig. 1A), the major detected lipids include alkyl quinolones (AQs), diacylglycerol (DAG), phosphatidylglycerol (PG), phosphatidylethanolamine (PE), cardiolipin (CL) and phosphatidylcholine (PC). These common lipids and their biosynthesis have been extensively studied in *P. aeruginosa* and other bacteria (29-31).

An uncommon lipid with a major [M-H]⁻ molecular ion at *m/z* 908.5 was eluted at ~18.5 min (Fig. 1B). Based on tandem mass spectrometry (MS/MS) (Fig. 1C) and exact mass measurement, it was identified as glucosaminyl-phosphatidylglycerol (GlcN-PG) recently reported in *P. aeruginosa* (17). GlcN-PG was previously found in other bacteria (18, 19, 21, 32).

To our knowledge, there have been no reports on the studies of the biosynthetic genes and functions of GlcN-PG in *P. aeruginosa* or any other bacteria.

Identification of the biosynthetic genes of GlcN-PG

Enzymatically, GlcN-PG is most likely produced from PG via the covalent modification with GlcN. Indeed, this was supported by the observation that during the pellicle growth of *P. aeruginosa*, the increase of GlcN-PG was accompanied by the decrease of PG (17). Previously it was also observed in *Bacillus megaterium* that GlcN-PG was increased, while PG was decreased, when cells were grown in acidic conditions (32).

Given the biochemical mechanisms of other amino sugar modifications (33, 34), we hypothesize that GlcN-PG is likely produced from its immediate precursor, GlcNAc-PG, by *N*-deacetylation, an enzymatic process that is known to be involved in several biosynthetic pathways. For examples, LpxC is the UDP-3-*O*-(*R*-3-hydroxymyristoyl)-*N*-acetylglucosamine deacetylase required for the biosynthesis of lipid A, the lipid anchor of lipopolysaccharide (LPS) in Gram-negative bacteria (35-37). Deacetylation mediated by carbohydrate esterase family 4 proteins (38) plays a vital role in exopolysaccharide processing in different bacterial species, including *P. aeruginosa*.

The potential involvement of deacetylation in the biosynthetic process of GlcN-PG prompted us to carefully examine the PA14 lipidomic data and identify a low level of GlcNAc-PG (Fig. 2A). The [M-H]⁻ ion of GlcNAc-PG is observed at *m/z* 950.59, corresponding to the addition of an acetyl group (42 Da) to GlcN-PG (*m/z* 908.58). The detection of GlcNAc-PG provides an important clue for the existence of a deacetylase that converts GlcNAc-PG to GlcN-PG.

To search for the GlcNAc-PG deacetylase, we queried the PA14 genomic database and found six genes annotated as deacetylases. Among them, four have been well characterized with identified substrates, including LpxC (35, 36). However, the other two putative deacetylases, with locus ID PA14_44830 and PA14_56040 (*gpgD*) (Table 2), have not been experimentally characterized and their substrates have remained unknown.

To determine whether GlcNAc-PG is a substrate of PA14_44830 and/or PA14_56040, we performed lipidomic analysis of their respective transposon mutants from a previously reported mutant collection (22). As shown in Fig. 2B, PA14_44830 mutant contains both GlcN-PG and GlcNAc-PG, with their relative levels similar to the wild-type (WT), suggesting that PA14_44830

is not a GlcNAc-PG deacetylase. By sharp contrast, the PA14_56040 (*gpgD*) mutant lacks GlcN-PG, but contains drastically elevated GlcNAc-PG (Fig. 2C), strongly supporting that GlcNAc-PG and GlcN-PG are the substrate and product of GpgD, respectively. The identification of the accumulated GlcNAc-PG (Fig. 3A) is further confirmed by MS/MS (Fig. 3B). The complete depletion of GlcN-PG in the PA14_56040 (*gpgD*) mutant also indicates that GpgD is the sole deacetylase responsible for the conversion of GlcNAc-PG to GlcN-PG.

PA14_56030 (*gpgS*) is encoded upstream of *gpgD* and is predicted to be a glycosyltransferase (Table 2). Its predicted function and genomic position strongly suggest that GpgS is likely the glycotransferase responsible for transferring the GlcNAc group onto PG to produce GlcNAc-PG. Indeed our lipidomic analysis shows that the PA14_56030 (*gpgS*)::Mar2xT7 mutant lacks both GlcNAc-PG and GlcN-PG (Fig. 2D).

Overall, the lipidomic analysis of the *gpgD* and *gpgS* transposon mutants reveals that GlcN-PG biosynthesis involves two steps: GpgS first transfers the GlcNAc group to PG to form GlcNAc-PG, which is then deacetylated by GpgD to produce GlcN-PG (Fig. 4). Although not experimentally verified, UDP-GlcNAc is likely the GlcNAc donor based on its utilization in other GlcNAc-modifications (39, 40).

We performed complementation of PA14 *gpgD*::Tn mutant, for which a pMQ72 derivative with *gpgD* under the control of an arabinose inducible promoter was constructed (Fig. 5A). Lipidomic analysis showed that arabinose-induced complementation of PA14 *gpgD*::Tn resulted in the production of GlcN-PG (Fig. 5B-C). The complementation of PA14 *gpgS*::Tn mutant remains to be demonstrated. We generated pMQ72-Tet-*gpgS* by cloning the *gpgS* gene in a similar vector system to *gpgD* but have been unable to obtain colonies of the PA14 *gpgS*::Tn mutant bearing pMQ72-Tet-*gpgS* despite trying various approaches, including electroporation, chemical transformation, and conjugation.

PA14_56050 (renamed *gpgF*), predicted to be a flippase (Table 2), is downstream of *gpgS* and *gpgD*. To experimentally demonstrate its flippase function will require the development of an assay that can monitor the membrane translocation of GlcN-PG (41, 42). Preliminary lipidomic analysis of the PA14_56050:Tn mutant shows that it contains both GlcN-PG and GlcNAc-PG (Fig. S1). Compared to WT, the level of GlcN-PG seems significantly lower (by several folds) in the PA14_56050::Tn mutant, whereas GlcNAc-PG is slightly increased. The mechanism of the effects of GpgF on GlcN-PG and GlcNAc-PG production remains to be investigated. The presence of

GlcN-PG in the *gpgF* mutant seems to suggest that GpgD functions cytosolically (Fig. 4), a hypothesis assuming that GpgF is the only flippase for GlcNAc-PG or GlcN-PG. The function of *gpgSDF* in PA14 is analogous to the function of *hexSDF* in *C. difficile*, which is involved in the synthesis of a novel glycolipid, aminohexosyl-hexosyldiradylglycerol (HNHDRG) (34, 43). It is worth noting that the localization of HexSDF was not experimentally determined; both intracellular and extracellular activities of HexD were proposed (34).

P. aeruginosa gpgSDF is sufficient for the synthesis of GlcNAc-PG and GlcN-PG in *E. coli*

E. coli contains PG as one of its most abundant phospholipids but does not have *gpgSDF* and does not produce GlcNAc-PG or GlcN-PG (28, 31, 44). To confirm the biochemical activities of *gpgSDF*, the plasmid pWO-GEM-T with the entire *gpgSDF* predicted operon of PA14 was constructed from pGEM-T vector and was then transformed into *E. coli* DH5α (Fig. 6A). Lipidomic analysis detected both GlcNAc-PG and GlcN-PG (Fig. 5B-D) in *E. coli* with pWO-GEM-T, confirming that *gpgSDF* possesses the expected glycotransferase and deacetylase activities. The identification of GlcNAc-PG and GlcN-PG produced in *E. coli* are supported by exact measurement and MS/MS. It is worth noting that the acyl chain profiles of the GlcNAc-PG and GlcN-PG molecular species heterologously produced in *E. coli* are slightly shorter (about 2 carbon atoms less) than those in *P. aeruginosa.* As expected, the expression of *gpgS* alone in *E. coli* resulted in the production of GlcNAc-PG, but not GlcN-PG (Fig. S2).

GlcN-PG is much less abundant than GlcNAc-PG in the *gpgSDF*-transformed *E. coli* (Fig. 6B-D), compared to *P. aeruginosa*. It is unknown whether this is due to altered codon usage in *P. aeruginosa* relative to *E. coli*, a non-optimal Shine-Dalgarno sequence for *gpgD* in the heterologous host, or other factors. Further optimization of the translation and activity of GpgD will be needed to improve the conversion of GlcNAc-PG to GlcN-PG to facilitate further studies of these lipids in *E. coli*.

Identification of GlcN-PG in other *P. aeruginosa* strains

Our bioinformatic analysis indicates that the *gpgSDF* operon is conserved in all sequenced strains of *P. aeruginosa.* Indeed, our lipidomic analysis confirmed the presence of GlcN-PG in three other strains of *P. aeruginosa* (Fig. 7 and Table 3).

Discussion

Using a combination of lipidomic and bioinformatic approaches, we identified two biosynthetic enzymes that are involved in GlcN-PG synthesis in *P. aeruginosa*. GpgS is a novel glycotransferase that catalyzes the modification of phosphatidylglycerol (PG) with *N*-acetylglucosamine (GlcNAc) to produce GlcNAc-PG, and GpgD is a novel deacetylase that removes the acetyl group of GlcNAc-PG to produce GlcN-PG. The functions of these enzymes were further confirmed by the synthesis of GlcNAc-PG and GlcN-PG from the heterologous expression of *gpgSDF* in *E. coli*.

The membrane distribution of GlcN-PG remains to be determined. *gpgF* is predicted to encode a flippase. Bioinformatic analysis predicts that GpgF has transmembrane spanning helices with inner membrane localization, whereas both GpgS and GpgD lack transmembrane spanning helices and are predicted to be cytoplasmic (Table 2). Thus, the biochemical activities mediated by *gpgSDF* are analogous to MprF which catalyzes Ala-PG synthesis on the cytosolic side of the inner membrane and subsequent translocation of Ala-PG to the periplasmic side of inner membrane (45, 46). In *P. aeruginosa*, it is unknown whether GlcN-PG, if flipped across the inner membrane, would be further transported to the outer membrane via proteins such as the AsmA-like family (47).

The identification of the GlcN-PG biosynthetic genes will enable the study of the biological and pathological functions of GlcN-PG in *P. aeruginosa*. Previously it was observed that GlcN-PG was dramatically increased under acidic conditions (32) for *Bacillus megaterium* and at stationary phase (17) for *P. aeruginosa*, suggesting the production of GlcN-PG is associated with coping with environmental and nutritional stresses. Previous studies have shown the *gpgS* gene (PA4311 in PAO1 strain) to be involved with phenazine production (48), differentially regulated under oxidative stress (49), implicated in exopolysaccharide alginate production (50), and up-regulated in acute and chronic infections (51).

The replacement of anionic PG with zwitterionic GlcN-PG is expected to lower the net negative charge of the bacterial membrane and thus may help confer resistance to cationic antimicrobials. In gram-positive *C. difficile*, GlcN-modification of a glycolipid impacts resistance to cationic antibiotics including daptomycin and bacitracin (34). It is plausible that GlcN-PG plays a similar role in defending against cationic compounds that are toxic to *P. aeruginosa*.

Remarkably, the lipidome of *P. aeruginosa* is more diverse than *E. coli* which has been the model organism for studying lipid composition and metabolism in Gram-negative bacteria. In particular, *P. aeruginosa* contains a multitude of zwitterionic phospholipids, including GlcN-PG, while *E. coli* has only one zwitterionic phospholipid (PE) in addition to two major anionic phospholipids (PG and CL) (28, 31, 44) (Fig. 8). The lipid diversity of *P. aeruginosa* reflects its relatively large genome size (52, 53) and likely contributes to its unusual metabolic flexibility and membrane adaptability. The elucidation of the functions of GlcN-PG and other *P. aeruginosa* zwitterionic phospholipids will be important for fully understanding the molecular mechanisms underlying the unique membrane properties, environmental resilience and adaptability, and high propensity for antibiotic resistance development of *P. aeruginosa*.

Acknowledgements

This work was supported by grants R01AI148366 and R01AI178692 from the National Institutes of Health to K.L.P. and Z.G. and the Cecil H. and Ida Green Chair in Systems Biology Science to K.P. The authors would like to thank the Dillon lab at the University of Texas at Dallas for providing the PA103 and ATCC 27853 strains of *P. aeruginosa*, Dr. Peter Jorth for providing pMQ72, and Dr. Trusha Parekh for providing *E. coli* S17.

Figure 1. Identification of GlcN-PG in *P. aeruginosa* PA14 by LC/MS/MS

The lipid extract of PA14 strain is analyzed by normal phase LC/MS/MS, with GlcN-PG being identified by exact mass measurement and MS/MS. A) Negative total ion chromatogram (TIC) of normal phase LC/MS of the lipid extract of *P. aeruginosa* strain PA14. B) Negative ion ESI mass spectrum (at ~18.5 min) showing the [M-H]$^-$ ions of GlcN-PG. The observed exact mass (908.586) matches with the predicted exact mass (908.587) for the [M-H]$^-$ ion of GlcN-PG (16:0/18:1). C) MS/MS of [M-H]$^-$ ion at *m/z* 908.5 produces fragment ions consistent with GlcN-PG (16:0/18:1) whose chemical structure is shown. The linkage positions of fatty acids and GlcN as depicted in the chemical structure of GlcN-PG (Fig. 1C) have not been determined experimentally and are for illustrative purposes.

Figure 2. PA14_56030 (*gpgS*) and PA14_56040 (*gpgD*) genes are required for the synthesis of GlcNAc-PG and GlcN-PG in *P. aeruginosa*.

LC/MS analysis of the transposon mutants of PA14_56030 (*gpgS*, putative glycotransferase) and PA14_56040 (*gpgD*, putative deacetylase) confirm that they are responsible for the synthesis of GlcNAc-PG and GlcN-PG in *P. aeruginosa,* respectively. A, B) GlcN-PG and a low level of GlcNAc-PG are present in the wild-type PA14 and PA14_44830::Tn mutant. C) GlcN-PG is depleted, and GlcNAc-PG is drastically accumulated in the PA14_56040::Tn mutant. D) Both GlcN-PG and GlcNAc-PG are absent in the PA14_56030::Tn mutant.

Shown are the extracted ion chromatograms (EICs) of NPLC/MS in the negative ion mode.

Figure 3. MS analysis of GlcNAc-PG accumulated in the PA14_56040 (*gpgD*)::Tn mutant.
A) Negative ion mass spectrum of [M-H]⁻ ion at *m/z* 950.6 for GlcNAc-PG (16:0/18:1). B) MS/MS
of [M-H]⁻ ion at *m/z* 950.6 produces fragment ions consistent with GlcNAc-PG (16:0/18:1)

Figure 4. Proposed biosynthesis and translocation of GlcN-PG in *P. aeruginosa*.

A) Diagram of the *gpgSDF* predicted operon and its neighboring genes in *P. aeruginosa* PA14. B) The biosynthesis of GlcN-PG from PG involves two enzymatic steps. First, PG is modified by GlcNAc by a glycosyltransferase (PA14_56030, GpgS), presumably using UDP-GlcNAc as the sugar donor. Second, GlcNAc-PG is de-acetylated by a deacetylase (PA14_56040, GpsD) to produce GlcN-PG. C) Cartoon illustration of the proposed biosynthesis and membrane translocation of GlcN-PG in *P. aeruginosa*. The localization of GpgSDF has not been experimentally determined.

Figure 5. Complementation of the *gpgD*::Tn mutant of PA14 with *gpgD in trans* restores GlcN-PG production.

A) Construction of a pMQ72 derivative with *gpgD* under the control of an arabinose-inducible promoter and containing a tetracycline resistane cassette. B) GlcN-PG is absent without arabinose induction. C) GlcN-PG is produced with arabinose (1%) induction. For better illustration, the GlcN-PG peak is magnified by 5-fold.

Figure 6. The heterologous expression of the PA14 *gpgSDF* genes produces GlcNAc-PG and GlcN-PG in *E. coli*.

A) The plasmid pWO-GEM-T with the *gpgSDF* genes of PA14 was constructed from pGEM-T vector and was then transformed into *E. coli* DH5α. B) Total ion chromatogram of the normal phase LC/MS in the negative ion mode of the lipid extract of *E. coli* expressing the pWO-GMT-T. C) Negative ion mass spectrum of the [M-H]⁻ ions of GlcNAc-PG species (appearing at ~12.5-13.0 min). D) Negative ion ass spectrum of the [M-H]⁻ ions of GlcN-PG species (appearing at ~14.5-15.0 min). The identification of GlcNAc-PG (16:0/16:1) and GlcN-PG (16:0/16:1) is confirmed by MS/MS.

Figure 7. Detection of GlcN-PG in various *P. aeruginosa* laboratory and clinical strains by LC/MS.

Shown are the negative ion mass spectra of the [M-H]⁻ ions of GlcN-PG species in A) PA103, B) ATCC 27853, C) PAO1 (Table 2). The identification of GlcN-PG molecular species is confirmed by exact mass measurement and MS/MS.

Figure 8. Comparison of phospholipid biosynthetic pathways in *P. aeruginosa* and *E. coli*. *P. aeruginosa* possesses more complex phospholipids than *E coli*. Major zwitterionic phospholipids (red) and anionic phospholipids (blue) are in bold fonts. Their biosynthetic genes are green. The synthesis of GlcN-PG from PG in *P. aeruginosa* (the subject of this study) consists of two enzymatic steps catalyzed by GpgS and GpgD.

Abbreviations: G3-P, glycerol-3-phosphate; L-PA, lyso-phosphatidic acid; PA, phosphatidic acid; CDP-DAG, CDP-diacylglycerol; PS, phosphatidylserine; PE, phosphatidylethanolamine; PC, phosphatidylcholine; PGP, phosphatidylglycerol-3-P; PG, phosphatidylglycerol; Ala-PG, alanyl-phosphatidylglycerol; CL, cardiolipin.

Table 1. Bacterial Stains and Plasmid Used in This Study

Species and strains	Description	Source or reference
P. aeruginosa		
PAO1	Wild type (first isolated in 1954 from a wound in Melbourne, Australia)	(54, 55)
PA14	Wild type (first isolated in early 1970s from the blood of a burn patient at Mercy Hospital in Pittsburg, PA, USA)	(56)
PA103 (ATCC 29260)	Wild type	(57)
ATCC 27853_NP446	Wild type (isolated in 1971 from a blood specimen in Peter Bent Brigham Hospital, Boston, USA)	(58)
PA14_56030::Mar2xT7 (*gpgS*::Tn)	PA14 with Mar2xT7 Tn insertion at nucleotide (nt) position 896 of 1116 nt PA14_56030 (*gpgS*) locus	(22)
PA14_56040::Mar2xT7 (*gpgD*::Tn)	PA14 with Mar2xT7 Tn insertion at nt position 226 of 762 nt PA14_56040 (*gpgD*) locus	(22)
PA14_56050::MarxT7 (*gpgF*::Tn)	PA14 with Mar2xT7 Tn insertion at nt position 403 of 1002 nt PA14_56050 (*gpgF*) locus	(22)
PA14_56040::Mar2xT7(pMQ72-Tet-*gpgD*)	PA14_56040::Mar2xT7 with the arabinose-inducible complementation vector pMQ72-Tet-*gpgD*	This work
E. coli		
E. coli DH5α	Cloning host strain	Thermo Fisher Scientific
DH5α (pWO-GEM-T)	pWO-GEM-T with *gpgSDF* predicted operon	This work

Table 2. Bioinformatic Analysis of Function and Localization of GpgS, GpgD and GpgF proteins.

PA14 locus	Predicted protein	Subcellular localization prediction[1]	Conserved domain (CD) description	CD accession[2]	CD e-value[2]
PA14_56030	GpgS	Cytoplasmic	Glycosyltransferase family 4 proteins	cd03814	1.23×10^{-123}
PA14_56040	GpgD	Cytoplasmic	Predicted deacetylase	COG3233	1.15×10^{-95}
PA14_56050	GpgF	Cytoplasmic Membrane	Predicted membrane flippase AglD2/YbhN, UPF0104 family	COG0392	4.10×10^{-24}

[1]Determined by analysis with PSORTb, version 3.0.3 (59)

[2]Determined by analysis with the NCBI Conserved Domains CD-Search tool against the CDD database (60); the hit with the lowest e-value for each protein is shown.

Table 3. Genomic and lipidomic analysis of GlcN-PG in *P. aeruginosa* stains

Strains	*gpgSDF* operon (Accession numbers shown for each)			GlcN-PG
	gpgS	*gpgD*	*gpgF*	
PAO1	PA4311 (NP_253001.1)	PA4312 (NP_253002.1)	PA4313 (NP_253003.1)	✓
PA14	PA14_56030 (YP_792656.1)	PA14_56040 (YP_792657.1)	PA14_56050 (YP_792658.1)	✓
PA103	PA103_5903 (WP_003110035.1)	PA103_5904 (WP_003112786.1)	PA103_5905 (WP_019485776.1)	✓
ATCC 27853	RS24835 (WP_003101181.1)	RS24840 (WP_003101179.1)	RS24845 (WP_016852096.1)	✓

References

1.   Murray, T. S., Egan, M., andKazmierczak, B. I. (2007) Pseudomonas aeruginosa chronic colonization in cystic fibrosis patients Curr Opin Pediatr **19**, 83-88 10.1097/MOP.0b013e3280123a5d
2.   Ernst, R. K., Yi, E. C., Guo, L., Lim, K. B., Burns, J. L., Hackett, M. *et al.* (1999) Specific lipopolysaccharide found in cystic fibrosis airway Pseudomonas aeruginosa Science **286**, 1561-1565 10.1126/science.286.5444.1561
3.   Paprocka, P., Durnas, B., Mankowska, A., Krol, G., Wollny, T., andBucki, R. (2022) Pseudomonas aeruginosa Infections in Cancer Patients Pathogens **11**, 10.3390/pathogens11060679
4.   Silby, M. W., Winstanley, C., Godfrey, S. A., Levy, S. B., andJackson, R. W. (2011) Pseudomonas genomes: diverse and adaptable FEMS Microbiol Rev **35**, 652-680 10.1111/j.1574-6976.2011.00269.x
5.   Kunz Coyne, A. J., El Ghali, A., Holger, D., Rebold, N., andRybak, M. J. (2022) Therapeutic Strategies for Emerging Multidrug-Resistant Pseudomonas aeruginosa Infect Dis Ther **11**, 661-682 10.1007/s40121-022-00591-2
6.   Hawkey, P. M., andLivermore, D. M. (2012) Carbapenem antibiotics for serious infections BMJ **344**, e3236 10.1136/bmj.e3236
7.   Peschel, A., Jack, R. W., Otto, M., Collins, L. V., Staubitz, P., Nicholson, G. *et al.* (2001) Staphylococcus aureus resistance to human defensins and evasion of neutrophil killing via the novel virulence factor MprF is based on modification of membrane lipids with l-lysine J Exp Med **193**, 1067-1076 10.1084/jem.193.9.1067
8.   Ernst, C. M., Kuhn, S., Slavetinsky, C. J., Krismer, B., Heilbronner, S., Gekeler, C. *et al.* (2015) The lipid-modifying multiple peptide resistance factor is an oligomer consisting of distinct interacting synthase and flippase subunits mBio **6**, 10.1128/mBio.02340-14
9.   Joyce, L. R., Manzer, H. S., da, C. M. J., Villarreal, R., Nagao, P. E., Doran, K. S. *et al.* (2022) Identification of a novel cationic glycolipid in *Streptococcus agalactiae* that contributes to brain entry and meningitis PLoS Biol **20**, e3001555 10.1371/journal.pbio.3001555
10.   Christensen, P. M., Martin, J., Uppuluri, A., Joyce, L. R., Wei, Y., Guan, Z. *et al.* (2023) Lipid discovery enabled by sequence statistics and machine learning bioRxiv 10.1101/2023.10.12.562061
11.   Hebecker, S., Arendt, W., Heinemann, I. U., Tiefenau, J. H., Nimtz, M., Rohde, M. *et al.* (2011) Alanyl-phosphatidylglycerol synthase: mechanism of substrate recognition during tRNA-dependent lipid modification in Pseudomonas aeruginosa Mol Microbiol **80**, 935-950 10.1111/j.1365-2958.2011.07621.x
12.   Raetz, C. R., Reynolds, C. M., Trent, M. S., andBishop, R. E. (2007) Lipid A modification systems in gram-negative bacteria Annu Rev Biochem **76**, 295-329 10.1146/annurev.biochem.76.010307.145803
13.   Poirel, L., Jayol, A., andNordmann, P. (2017) Polymyxins: Antibacterial Activity, Susceptibility Testing, and Resistance Mechanisms Encoded by Plasmids or Chromosomes Clin Microbiol Rev **30**, 557-596 10.1128/CMR.00064-16
14.   Yan, A., Guan, Z., andRaetz, C. R. (2007) An undecaprenyl phosphate-aminoarabinose flippase required for polymyxin resistance in Escherichia coli J Biol Chem **282**, 36077-36089 10.1074/jbc.M706172200

15.   Han, M. L., Velkov, T., Zhu, Y., Roberts, K. D., Le Brun, A. P., Chow, S. H. *et al.* (2018) Polymyxin-Induced Lipid A Deacylation in Pseudomonas aeruginosa Perturbs Polymyxin Penetration and Confers High-Level Resistance ACS Chem Biol **13**, 121-130 10.1021/acschembio.7b00836

16.   Zhang, H., Srinivas, S., Xu, Y., Wei, W., andFeng, Y. (2019) Genetic and Biochemical Mechanisms for Bacterial Lipid A Modifiers Associated with Polymyxin Resistance Trends Biochem Sci **44**, 973-988 10.1016/j.tibs.2019.06.002

17.   Abbes, I., Rihouey, C., Hardouin, J., Jouenne, T., De, E., andAlexandre, S. (2018) Identification by mass spectrometry of glucosaminylphosphatidylglycerol, a phosphatidylglycerol derivative, produced by Pseudomonas aeruginosa Rapid Commun Mass Spectrom **32**, 2113-2121 10.1002/rcm.8277

18.   Phizackerley, P. J., MacDougall, J. C., andFrancis, M. J. (1966) Phosphatidylglycerylglucosamine Biochem J **99**, 21C-22C 10.1042/bj0990021c

19.   Bertsch, L. L., Bonsen, P. P., andKornberg, A. (1969) Biochemical studies of bacterial sporulation and germination. XIV. Phospholipids in Bacillus megaterium J Bacteriol **98**, 75-81 10.1128/jb.98.1.75-81.1969

20.   den Kamp, J. O., Bonsen, P. P., andvan Deenen, L. L. (1969) Structural investigations on glucosaminyl phosphatidylglycerol from Bacillus megaterium Biochim Biophys Acta **176**, 298-305 10.1016/0005-2760(69)90187-8

21.   MacDougall, J. C., andPhizackerley, P. J. (1969) Isomers of glucosaminylphospatiylglycerol in Bacillus megaterium Biochem J **114**, 361-367 10.1042/bj1140361

22.   Liberati, N. T., Urbach, J. M., Miyata, S., Lee, D. G., Drenkard, E., Wu, G. *et al.* (2006) An ordered, nonredundant library of Pseudomonas aeruginosa strain PA14 transposon insertion mutants Proc Natl Acad Sci U S A **103**, 2833-2838 10.1073/pnas.0511100103

23.   Shanks, R. M., Caiazza, N. C., Hinsa, S. M., Toutain, C. M., andO'Toole, G. A. (2006) Saccharomyces cerevisiae-based molecular tool kit for manipulation of genes from gram-negative bacteria Appl Environ Microbiol **72**, 5027-5036 10.1128/AEM.00682-06

24.   Hoang, T. T., Karkhoff-Schweizer, R. R., Kutchma, A. J., andSchweizer, H. P. (1998) A broad-host-range Flp-FRT recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked Pseudomonas aeruginosa mutants Gene **212**, 77-86 10.1016/s0378-1119(98)00130-9

25.   Hmelo, L. R., Borlee, B. R., Almblad, H., Love, M. E., Randall, T. E., Tseng, B. S. *et al.* (2015) Precision-engineering the Pseudomonas aeruginosa genome with two-step allelic exchange Nat Protoc **10**, 1820-1841 10.1038/nprot.2015.115

26.   McMackin, E. A. W., Corley, J. M., Karash, S., Marden, J., Wolfgang, M. C., andYahr, T. L. (2021) Cautionary Notes on the Use of Arabinose- and Rhamnose-Inducible Expression Vectors in Pseudomonas aeruginosa J Bacteriol **203**, e0022421 10.1128/JB.00224-21

27.   Zhao, Y., Liu, Z., Yu, S., Wen, S., Hammarstrom, L., andRabbani, H. (2009) Construction of a High Efficiency PCR Products Cloning T Vector Using pGEM-5zf (+) Avicenna J Med Biotechnol **1**, 37-39, https://www.ncbi.nlm.nih.gov/pubmed/23407719

28.   Tan, B. K., Bogdanov, M., Zhao, J., Dowhan, W., Raetz, C. R., andGuan, Z. (2012) Discovery of a cardiolipin synthase utilizing phosphatidylethanolamine and phosphatidylglycerol as substrates Proc Natl Acad Sci U S A **109**, 16504-16509 10.1073/pnas.1212797109

29. Sohlenkamp, C., andGeiger, O. (2016) Bacterial membrane lipids: diversity in structures and pathways FEMS Microbiol Rev **40**, 133-159 10.1093/femsre/fuv008

30. Dowhan, W. (1997) Molecular basis for membrane phospholipid diversity: why are there so many lipids? Annu Rev Biochem **66**, 199-232 10.1146/annurev.biochem.66.1.199

31. Dowhan, W. (2013) A retrospective: use of Escherichia coli as a vehicle to study phospholipid synthesis and function Biochim Biophys Acta **1831**, 471-494 10.1016/j.bbalip.2012.08.007

32. den Kamp, J. O., Houtsmuller, U. M., andvan Deenen, L. L. (1965) On the phospholipids of Bacillus megaterium Biochim Biophys Acta **106**, 438-441 10.1016/0005-2760(65)90059-7

33. Biely, P. (2012) Microbial carbohydrate esterases deacetylating plant polysaccharides Biotechnol Adv **30**, 1575-1588 10.1016/j.biotechadv.2012.04.010

34. Pannullo, A. G., Guan, Z., Goldfine, H., andEllermeier, C. D. (2023) HexSDF Is Required for Synthesis of a Novel Glycolipid That Mediates Daptomycin and Bacitracin Resistance in C. difficile mBio **14**, e0339722 10.1128/mbio.03397-22

35. Jackman, J. E., Raetz, C. R., andFierke, C. A. (1999) UDP-3-O-(R-3-hydroxymyristoyl)-N-acetylglucosamine deacetylase of Escherichia coli is a zinc metalloenzyme Biochemistry **38**, 1902-1911 10.1021/bi982339s

36. Coggins, B. E., Li, X., McClerren, A. L., Hindsgaul, O., Raetz, C. R., andZhou, P. (2003) Structure of the LpxC deacetylase with a bound substrate-analog inhibitor Nat Struct Biol **10**, 645-651 10.1038/nsb948

37. Zhao, J., Cochrane, C. S., Najeeb, J., Gooden, D., Sciandra, C., Fan, P. *et al.* (2023) Preclinical safety and efficacy characterization of an LpxC inhibitor against Gram-negative pathogens Sci Transl Med **15**, eadf5668 10.1126/scitranslmed.adf5668

38. Le Mauff, F., Razvi, E., Reichhardt, C., Sivarajah, P., Parsek, M. R., Howell, P. L. *et al.* (2022) The Pel polysaccharide is predominantly composed of a dimeric repeat of alpha-1,4 linked galactosamine and N-acetylgalactosamine Commun Biol **5**, 502 10.1038/s42003-022-03453-2

39. Hart, G. W., Slawson, C., Ramirez-Correa, G., andLagerlof, O. (2011) Cross talk between O-GlcNAcylation and phosphorylation: roles in signaling, transcription, and chronic disease Annu Rev Biochem **80**, 825-858 10.1146/annurev-biochem-060608-102511

40. Chen, M., Bridges, A., andLiu, J. (2006) Determination of the substrate specificities of N-acetyl-d-glucosaminyltransferase Biochemistry **45**, 12358-12365 10.1021/bi060844g

41. Klein, S., Lorenzo, C., Hoffmann, S., Walther, J. M., Storbeck, S., Piekarski, T. *et al.* (2009) Adaptation of Pseudomonas aeruginosa to various conditions includes tRNA-dependent formation of alanyl-phosphatidylglycerol Mol Microbiol **71**, 551-565 10.1111/j.1365-2958.2008.06562.x

42. Ernst, C. M., andPeschel, A. (2011) Broad-spectrum antimicrobial peptide resistance by MprF-mediated aminoacylation and flipping of phospholipids Mol Microbiol **80**, 290-299 10.1111/j.1365-2958.2011.07576.x

43. Guan, Z., Katzianer, D., Zhu, J., andGoldfine, H. (2014) Clostridium difficile contains plasmalogen species of phospholipids and glycolipids Biochim Biophys Acta **1842**, 1353-1359 10.1016/j.bbalip.2014.06.011

44.   Li, C., Tan, B. K., Zhao, J., andGuan, Z. (2016) In Vivo and in Vitro Synthesis of Phosphatidylglycerol by an Escherichia coli Cardiolipin Synthase J Biol Chem **291**, 25144-25153 10.1074/jbc.M116.762070

45.   Hebecker, S., Krausze, J., Hasenkampf, T., Schneider, J., Groenewold, M., Reichelt, J. *et al.* (2015) Structures of two bacterial resistance factors mediating tRNA-dependent aminoacylation of phosphatidylglycerol with lysine or alanine Proc Natl Acad Sci U S A **112**, 10691-10696 10.1073/pnas.1511167112

46.   Hankins, M. T. K., Parrag, M., Garaeva, A. A., Earp, J. C., Seeger, M. A., Stansfeld, P. J. *et al.* (2025) MprF from Pseudomonas aeruginosa is a promiscuous lipid scramblase with broad substrate specificity Sci Adv **11**, eads9135 10.1126/sciadv.ads9135

47.   Sposato, D., Mercolino, J., Torrini, L., Sperandeo, P., Lucidi, M., Alegiani, R. *et al.* (2024) Redundant essentiality of AsmA-like proteins in Pseudomonas aeruginosa mSphere **9**, e0067723 10.1128/msphere.00677-23

48.   Liang, H., Li, L., Dong, Z., Surette, M. G., andDuan, K. (2008) The YebC family protein PA0964 negatively regulates the Pseudomonas aeruginosa quinolone signal system and pyocyanin production J Bacteriol **190**, 6217-6227 10.1128/JB.00428-08

49.   Salunkhe, P., Smart, C. H., Morgan, J. A., Panagea, S., Walshaw, M. J., Hart, C. A. *et al.* (2005) A cystic fibrosis epidemic strain of Pseudomonas aeruginosa displays enhanced virulence and antimicrobial resistance J Bacteriol **187**, 4908-4920 10.1128/JB.187.14.4908-4920.2005

50.   Wood, L. F., andOhman, D. E. (2012) Identification of genes in the sigma(2)(2) regulon of Pseudomonas aeruginosa required for cell envelope homeostasis in either the planktonic or the sessile mode of growth mBio **3**, 10.1128/mBio.00094-12

51.   Turner, K. H., Everett, J., Trivedi, U., Rumbaugh, K. P., andWhiteley, M. (2014) Requirements for Pseudomonas aeruginosa acute burn and chronic surgical wound infection PLoS Genet **10**, e1004518 10.1371/journal.pgen.1004518

52.   Klockgether, J., Cramer, N., Wiehlmann, L., Davenport, C. F., andTummler, B. (2011) Pseudomonas aeruginosa Genomic Structure and Diversity Front Microbiol **2**, 150 10.3389/fmicb.2011.00150

53.   Stover, C. K., Pham, X. Q., Erwin, A. L., Mizoguchi, S. D., Warrener, P., Hickey, M. J. *et al.* (2000) Complete genome sequence of Pseudomonas aeruginosa PAO1, an opportunistic pathogen Nature **406**, 959-964 10.1038/35023079

54.   Holloway, B. W. (1955) Genetic recombination in Pseudomonas aeruginosa J Gen Microbiol **13**, 572-581 10.1099/00221287-13-3-572

55.   Wilderman, P. J., Vasil, A. I., Martin, W. E., Murphy, R. C., andVasil, M. L. (2002) Pseudomonas aeruginosa synthesizes phosphatidylcholine by use of the phosphatidylcholine synthase pathway J Bacteriol **184**, 4792-4799 10.1128/JB.184.17.4792-4799.2002

56.   Rahme, L. G., Stevens, E. J., Wolfort, S. F., Shao, J., Tompkins, R. G., andAusubel, F. M. (1995) Common virulence factors for bacterial pathogenicity in plants and animals Science **268**, 1899-1902 10.1126/science.7604262

57.   Liu, P. V. (1966) The roles of various fractions of Pseudomonas aeruginosa in its pathogenesis. II. Effects of lecithinase and protease J Infect Dis **116**, 112-116 10.1093/infdis/116.1.112

58.   Cao, H., Lai, Y., Bougouffa, S., Xu, Z., andYan, A. (2017) Comparative genome and transcriptome analysis reveals distinctive surface characteristics and unique physiological

potentials of Pseudomonas aeruginosa ATCC 27853 BMC Genomics **18**, 459 10.1186/s12864-017-3842-z

59. Yu, N. Y., Wagner, J. R., Laird, M. R., Melli, G., Rey, S., Lo, R. *et al.* (2010) PSORTb 3.0: improved protein subcellular localization prediction with refined localization subcategories and predictive capabilities for all prokaryotes Bioinformatics **26**, 1608-1615 10.1093/bioinformatics/btq249

60. Marchler-Bauer, A., andBryant, S. H. (2004) CD-Search: protein domain annotations on the fly Nucleic Acids Res **32**, W327-331 10.1093/nar/gkh454

Supplemental Information

Figure S1.   LC/MS analysis of GlcNAc-PG and GlcN-PG in the PA14_56050 (*gpgF*) mutant of *P. aeruginosa*

PA14_56050 (*gpgF*) is bioinformatically predicted to be a lipid flippase. Lipidomic analysis of the PA14 56050::Tn mutant shows that it contains both GlcN-PG and GlcNAc-PG. Compared to WT, the level of GlcN-PG is significantly lower (roughly by several folds), whereas GlcNAc-PG is slightly increased in the PA14 56050::Tn mutant. The presence of GlcN-PG (at a much higher level than GlcNAc-PG) in the *gpgF* mutant suggests that GpgD functions cytosolically, assuming that GpgF is the only flippase for GlcNAc-PG or GlcN-PG.

Figure S2. Heterologous synthesis of GlcNAc-PG in *E. coli* by expressing PA14_56030 (*gpgS*) from *P. aeruginosa*.

A) *E. coli* DH5a lacks both GlcN-PG and GlcNAc-PG.

B) GlcNAc-PG, but not GlcN-PG, is produced by expressing PA14_56050 (GpgS) in *E. coli* DH5a.

C) Negative ion mass spectrum of [M-H]⁻ ions of GlcNAc-PG species produced by expressing PA14_56050 (GpgS) in *E. coli* DH5a.

Table S1. Primers for *gpgD* complementation in PA14 *gpgD*::Tn mutant

	Primers	Sequence (5'-->3') Plasmid-Gene
1	gpgD_For	CTCATCCGCCAAAACAGCCAATGGCTGAACGCAGCCTGC
2	gpgD_Rev	CCCGTTTTTTTGGGCTAGCGTCATGCCGCGCGGTCCAG
3	Screen_pMQ_MCS_For	GCAAATTCTGTTTTATCAGACCGC
4	pMQ_ScreennRev_1	GGCGGGTTTCCTCGTCCAGGCCGAGTT
5	pMQ_ScreenFor_2	GTCGCCGACGTGCATGTAC
6	pMQ_ScreenRev_2	TCAGCCATAGCGCGGGAACC
7	pMQ_ScreenFor_3	CACGTCGAGCGCCAGCATGC
8	pMQ_ScreenRev_3	CGACGCCAGGCACTGCGC
9	pMQ_ScreenFor_4	GAGCGCCGGCACCCGCCAGGC
10	Screen_pMQ_MCS_Rev	CATAAGATTAGCGGATCCTACCTG
11	Lin_For_pMQ72	CGTTGCTGCTCCATAACATCAAAC
12	Lin_Rev_pMQ72	TAACAATTCGTTCAAGCCGAGATC
13	Tetgene_For	GAACGAATTGTTATCAGGTCGAGGTGG
14	Tetgene_Rev	GGAGCAGCAACGATGAAATCTAACAATG
15	Tet_scr_pMQ_For	CGCGGCGTTGTGACAATTTAC
16	Tet_scr_pMQ_Rev	CCAAGCAGCAAGCGCGTTACG