Original Article

The polyketide pathway in sporopollenin biosynthesis is specific to land plants (Embryophyta)

Damanpreet K. Sraan, Neil W. Ashton and Dae-Yeon Suh*

Department of Chemistry and Biochemistry, University of Regina, Regina, SK S4S 0A2, Canada

Running title: Embryophyte specific sporopollenin biosynthesis

*For correspondence. E-mail suhdaey@uregina.ca

•   **Background and Aims:** Sporopollenin (SP) is a complex biopolymer in the outer wall of

spores and pollen and provides protection from environmental stresses. Its extraordinary chemical

resistance, especially to acetolysis, was widely used to identify SP in biological specimens. This

broad definition of SP led to claims for its widespread occurrence among diverse embryophyte

and non-embryophyte taxa. We previously proposed a biochemical definition that can be used to

distinguish genuine SP from other chemically resistant cell wall materials. The definition was

centred on ASCL (Anther-Specific Chalcone synthase-Like), an embryophyte-specific enzyme of

the polyketide pathway that provides precursors for SP biosynthesis. Herein, we examine the

evolution and distribution of all five enzymes (CYP703A, CYP704B, ACOS, ASCL and TKPR)

of the polyketide pathway and propose a new, more comprehensive definition of SP.

•   **Methods:** We performed BLASTp searches, phylogenetic tree construction, protein

modeling and sequence analysis to determine the presence or absence of ACOS and TKPR in

embryophytes and streptophytic algae.

•   **Key Results:** We found evidence that all five enzymes of the polyketide pathway evolved

from ancestral enzymes of primary metabolism and ACOS, ASCL and TKPR were co-selected

during evolution. The dosage of all five genes has been subjected to strict evolutionary control and,

in some taxa, synteny has provided a selective advantage. All five enzymes are present in

embryophytes but absent in green algae, indicating that the polyketide pathway and therefore SP

is embryophyte-specific.

•   **Conclusions:** The addition of the polyketide pathway in the definition of genuine SP will

allow separation of SP from algaenans and other chemically resistant 'SP-like' algal spore wall

substances. This study further signifies SP as an evolutionary innovation unique to the

embryophyte lineage and encourages research on possible evolutionary relationship between algal

spore wall 'SP-like materials' and embryophyte SP.

Key words: Sporopollenin, Embryophyta, CYP450, acyl-CoA synthetase, anther-specific

chalcone synthase-like, tetraketide α-pyrone reductase, enzyme-specific residue, plant evolution

and terrestrialisation, polyketide pathway, gene copy number, synteny

INTRODUCTION

We previously proposed a new definition of sporopollenin (SP) as follows:

"Sporopollenin is a chemically resistant complex heteropolymer present in the outer walls of

spores and pollen grains and is composed partly of hydroxylated polyketides derived from the

conserved polyketide pathway, which involves ASCL." (Suh and Ashton, 2022).

As a protectant of progeny, SP is extraordinarily resistant to biological and chemical treatments,

and it has been challenging to determine its chemical composition and structure. Hence, prior to

our definition, resistance to acetolysis had been the sole criterion for the identification of SP

(Rowley and Southworth, 1967; Heslop-Harrison, 1969). This imprecise definition resulted in

claims for the existence of SP in algae and various microorganisms and also misguided research

on the evolution of SP and its roles during terrestrialisation (Delwiche *et al*., 1989; Feofilova, 2010;

de Vries and Archibald, 2018).

Recently, genetic and biochemical research has uncovered the biosynthesis pathways for

various SP precursors (Grienenberger and Quilichini, 2021; Xue *et al*., 2023). The polyketide

pathway comprised of five enzymes, CYP703A, CYP704B, ACOS (acyl-CoA synthetase), ASCL

(anther-specific chalcone synthase-like) and TKPR (tetraketide α-pyrone reductase), provides

polyketide SP precursors and is conserved in embryophytes. ASCL belongs to the type III

polyketide synthase family, and previous phylogenetic studies of the family indicated that ASCL

is specific to embryophytes (Jiang *et al*., 2008; Shimizu *et al*., 2017). Indeed, we discovered at

least one *ASCL* in representative embryophytes from all lineages that produce spores or pollen

with sporopolleninous exine. In sharp contrast, we failed to find any *ASCL* in the genomes of all

algal species for which data are available, including Charophytes that were considered to produce

SP-like substances. This finding led us to propose the above-mentioned definition for SP to

distinguish genuine SP from other similarly resistant substances and to use *ASCL* as an identifying marker for the presence of SP in spore and pollen walls. We recognised that there are other enzymes than ASCL in the polyketide pathway and exploring algal genomes for orthologues encoding these enzymes had the potential for augmenting our SP definition (Suh and Ashton, 2022).

In this study, we investigated the evolution of the genes in the polyketide pathway and examined whether CYP703A, CYP704B, ACOS and TKPR are conserved in embryophytes and whether they are absent in green algae.

METHODS AND MATERIALS

Database searches and phylogenetic analysis

BLASTp searches were performed against 35 embryophytes selected from all major lineages from hornworts to eudicots in Phytozome 13 (https://phytozome-next.jgi.doe.gov/), the National Center for Biotechnology Information (https://www.ncbi.nlm.nih.gov/), the *Takakia* project (https://www.takakia.com/blast/blast_cs.html), the *Anthoceros* genomes (https://www.hornworts.uzh.ch/en/hornwort-genomes.html) and the FernBase (https://fernbase.org) using PpACOS6, AtACOS5, PpTKPR1 and AtTKPR1 as queries. Similarly, BLASTp searches were performed against each algal genome in PhycoCosm (https://phycocosm.jgi.doe.gov/phycocosm/home) (accessed on 12 March 2024) using the same query sequences. The cutoff E value was $1.0 \times \mathrm{E}^{-20}$ and truncated and redundant sequences were removed. Representative sequences of the 4CL (4-coumarate:CoA ligase) and CCR (cinnamoyl-CoA reductase) families were retrieved from Phytozome 13. The sequences used for phylogenetic tree construction (Supplementary Data Table S1, S2) were aligned by Clustal Omega

(https://www.ebi.ac.uk/jdispatcher/msa/clustalo), and Maximum Likelihood (ML) trees were built

in MEGA 11 under the WAG+G+I substitution model. Support for the tree was measured using

1,000 bootstrap replicates.

Protein modeling and sequence analysis of ACOS and TKPR

Structures of ACOS from *Physcomitrium patens*, arabidopsis, *Oryza sativa* (PpACOS6, AtACOS5,

OsACOS12, respectively) and that of *P. patens* TKPR (PpTKPR1) were modeled with SWISS-

MODEL (https://swissmodel.expasy.org/) and AlphaFold (https://alphafold.ebi.ac.uk/). The

modeled structures of ACOSs were compared to the X-ray crystal structure of tobacco 4CL2

(Nt4CL2; PDB id, 5BST) and their active site sequences were aligned using Clustal Omega to

reveal enzyme-specific residues. Likewise, the modeled structure of PpTKPR1 was compared to

crystal structures of *Medicago truncatula* CCR (MrCCR; PDB id, 4R1U), *Panicum virgatum*

dihydroflavonol 4-reductase (PvDFR; PDB id, 8FEM) and *Sorghum bicolor* anthocyanidin

reductase (SbANR; PDB id, 8FIP) and their active site sequences were aligned using Clustal

Omega.

Synteny analysis

Coordinates for genes of the polyketide pathway in the taxa discussed and total scaffold lengths of

their genomes were obtained from Phytozome 13. The coordinates were used to calculate the

lengths of DNA spanning chosen pairs of syntenic genes (from the beginning of one gene to the

end of the other). Total genomic scaffold lengths were assumed to represent total genomic

euchromatin. The ratio of these values was viewed as the probability by chance alone of one gene

being present within the spanned DNA on a particular chromosome. The probability of n genes

being located in this span of DNA was calculated as the ratio to the power, n. The uncorrected

probabilities for *Arabidopsis thaliana* were calculated in this way using data from the arabidopsis

TAIR 10 genome sequence in Phytozome 13. The values corrected for heterochromatin utilised

data from the recently sequenced arabidopsis Col-XJTU assembly (Wang *et al*., 2022): the length

of CEN1 (3.8 Mb) was subtracted from the length of DNA spanned by the five syntenic genes of

the polyketide pathway (26 Mb) located on Chr 1 and the combined length of the five CENs (21.7

Mb) and two NORs (0.64 Mb) was subtracted from the genome size (134 Mb). This gave a length

for euchromatin spanned by five syntenic genes of the pathway on Chr 1 of 22.2 Mb and for total

genomic euchromatin of 112 Mb. Corrected probabilities of the observed synteny on Chr 1 and

Chr 4 occurring by chance alone were then obtained by entering these calculated values in the

equation given in the main text.

RESULTS AND DISCUSSION

The embryophyte-specific polyketide pathway and an expanded definition of sporopollenin

It had been suggested previously that ACOS and TKPR are embryophyte-specific (de Azevedo

Souza *et al*., 2008; Grienenberger *et al*., 2010). However, information about algal homologues of

these enzymes was lacking. Taking advantage of newly determined genome sequences, we

expanded earlier studies and searched for ACOS and TKPR in the genomes of thirty-five

embryophytes representing all major lineages (Supplementary Data Tables S1, S2). Putative

ACOS sequences collated from BLASTp searches were identified based on phylogenetic

clustering with known enzymes (Supplementary Data Figs S1, S2). In parallel, for additional

support for enzyme identification, we searched for amino acid residues that are conserved in ACOS

but not in its paralogues. ACOS and 4CL, both of which belong to the 4CL family, catalyze

formation of CoA esters. Both enzymes use ATP and CoA as co-substrates, but they differ in their

acid substrates. While ACOS accepts hydroxyfatty acids (provided by CYP703A and CYP704B

in the polyketide pathway, Fig 1) as substrate, 4CL accepts 4-coumaric acid derivatives. Crystal

structures of several 4CLs pinpointed active site residues responsible for substrate specificity of

the enzyme (Hu *et al*., 2010). Sequence alignment of representative enzymes of the 4CL family

(Supplementary Data Fig. S3) and comparison of 4CL crystal structures (e.g., tobacco 4CL2) with

ACOS structure models (Fig. 2A, B) indicated that Gly^{241} and $S^{339}C(V/I)Tx(S/T)$ (numbering of

AtACOS5) are uniquely conserved in ACOS and could serve as a determinant of ACOS.

Reflecting the different substrate specificity of ASCL and 4CL, corresponding residues in Nt4CL2

are Ser^{243} and $G^{339}PVLxM$ (Fig. 2B). We then performed a BLASTp search of all algal genomes

in the PhycoCosm database for algal homologues of ACOS (Supplementary Data Table S1) and

constructed a phylogenetic tree with representative embryophyte enzymes of the 4CL family. As

shown in Fig. S1, none of the algal homologues, including 13 from charophytes, is found in the

ACOS clade and none of them possesses the above-mentioned ACOS signature residues

(Supplementary Data Fig. S3). We concluded that none of the algae examined has an ACOS.

Similarly, sequences and modeled structures of TKPRs were compared with those of

paralogous DFR, ANR and CCR to identify two substrate binding site residues Arg^{137} and

$(Ser/Thr)^{197}$ (numbering of PpTKPR1) as determinants of TKPR (Supplementary Data Fig. S4;

Fig. 2C, D). BLASTp searches yielded a total of 29 algal homologues, including four from

charophytes (Supplementary Data Table S2). None of them resides in the TKPR clade

(Supplementary Data Fig. S2) and, furthermore, no charophyte homologue has the TKPR signature

residues, Arg and (Ser/Thr). The cytochrome P450 family had already been extensively studied

and meticulously curated in the P450 database (https://drnelson.uthsc.edu/, Nelson, 2009).

CYP703A is present only in embryophytes, while CYP704B is found in embryophytes, protists and red algae, but, importantly, not in green algae (Morant *et al*., 2007; Li *et al*., 2010; Hansen *et al*., 2021). These findings led us to conclude that all of the five enzymes in the polyketide pathway are present in embryophytes but absent in green algae, and produce polyketide precursors of genuine SP. Therefore, we have expanded our definition of SP as follows:

"Sporopollenin is a chemically resistant complex heteropolymer present in the outer walls of spores and pollen grains and is composed partly of hydroxylated polyketides derived from the conserved polyketide pathway, which typically involves CYP703A, CYP704B, ACOS, ASCL and TKPR."

Evolutionary origin of the polyketide pathway

Next we searched the literature for studies on the evolutionary origins of the enzymes in the polyketide pathway. It is generally accepted that enzymes of secondary metabolism have been derived from pre-existing enzymes in primary metabolism via gene duplication and neofunctionalisation (Firn and Jones, 2000).

(1) ACOS

The 4CL family belongs to the adenylate-forming enzyme superfamily present in all organisms. The superfamily shares a common reaction of the formation of an adenylated intermediate as in fatty acid activation by fatty acyl-CoA synthetase. 4CL produces CoA esters of hydroxycinnamic acids which are intermediates in the biosynthesis of a vast array of phenolic secondary metabolites, including monolignols and flavonoids (Hahlbrock and Scheel, 1989). Based on kinetic and phylogenetic analysis, it has been suggested that the members of the superfamily share a common

ancestor similar to extant acyl-CoA synthetase in primary fatty acid metabolism (Fig. 1) (Linne *et*

al., 2007; de Azevedo Souza *et al*., 2008).

(2) ASCL

ASCL belongs to the family of type III polyketide synthases (PKS). Type III PKSs condense a

starter CoA substrate with a number of (mostly three) malonyl-CoA molecules and produce a

variety of polyketide secondary metabolites with different biological functions (Abe and Morita,

2010). There are more than twenty type III PKSs in plants and the most well-known is chalcone

synthase (CHS) that catalyzes the first committed step in flavonoid biosynthesis. Type III PKSs,

along with several condensing enzymes in fatty acid metabolism, belong to the thiolase

superfamily. An earlier phylogenetic study showed that type III PKSs have evolved, along with β-

ketoacyl synthases in fatty acid synthesis, from an ancestral thiolase-like condensing enzyme in

lipid metabolism (Fig. 1) (Jiang et al. 2008).

(3) TKPR

TKPR and its paralogues DFR and CCR all belong to the short-chain dehydrogenase/reductase

(SDR) superfamily present in all living organisms. DFR plays a role in the biosynthesis of

anthocyanins while CCR catalyses the first committed step of lignin biosynthesis (Lacombe *et al*.,

1997; Petit *et al*., 2007). Enzymes in the SDR family display similarities to sugar-metabolizing

enzymes in bacteria and it has been suggested that an ancestor of the superfamily could have been

UDP-galactose-4-epimerase in primary sugar metabolism (Fig. 1) (Baker and Blasco, 1992;

Kallberg *et al*., 2002).

(4) CYP703A and CYP704B

The CYP family is one of the largest enzyme families and present in all kingdoms of life. The

family is made of many clans and each clan contains individual CYP enzymes. The CYP 71 and

86 clans, to which CYP 703A and 704B belong respectively, are mainly involved in fatty acid

metabolism. All extant CYPs are monophyletic, and CYP51 (sterol 14α-demethylase) in sterol

biosynthesis is widely accepted as the ancestral progenitor (Yoshida et al. 2000; Lamb et al., 2021).

To summarize, it appears that ACOS, ASCL and TKPR all evolved from ancestral enzymes in

primary metabolism in the common ancestor of embryophytes. It could be more than a coincidence

that their paralogues play important roles in phenylpropanoid pathways (Fig. 1) and it may reflect

critical contributions that SP and phenylpropanoids made during early evolution of land plants.

ACOS, ASCL and TKPR were co-selected during evolution

Earlier, we found that the genome of *Zostera marina* (a seagrass) that produces exine-less pollen

does not contain *ASCL*, which we attributed to secondary gene loss (Suh and Ashton, 2022). The

Z. marina genome also lacks an *ACOS*, and it contains one copy of a *TKPR*-like sequence that

would produce a truncated TKPR-like protein because of nonsense mutation and pseudogenisation.

Also, *Zingiber officinale* produces pollen with a reduced exine with regular discontinuities

(Subbarayudu *et al.*, 2014). A tBLASTn search against the genome of *Z. officinale* Roscoe

(taxid:94328) (https://blast.ncbi.nlm.nih.gov/, accessed in February 2024) failed to find *ACOS*,

ASCL or *TKPR*. Therefore, in line with our new and expanded definition, neither *Z. marina* nor *Z.*

officinale makes genuine SP. Based on these data, we also concluded that the three genes of the

polyketide pathway were co-selected during embryophyte evolution; loss of one of these genes

would have alleviated the selective pressure for retention of the remaining genes in the pathway.

CYP703A and *CYP704B* genes were found in the *Z. marina* and *Z. officinale* genomes unlike the

other three genes in the polyketide pathway. That they did not suffer gene loss is probably because

the CYP enzymes they encode are required for the biosynthesis of cutin monomers in addition to

SP biosynthesis (Li *et al*., 2010).

Gene dosage: Copy number of the genes in the polyketide pathway

We examined whether the copy number of the five genes in the pathway is controlled. The *CYP*

family underwent diversification bursts (Nelson, 2019) and clans 71 and 86 are the two largest

clans in embryophytes. For instance, the genome of *P. patens* has 71 CYPs and 41 of them are in

clan 71. Nonetheless, *CYP703A* is often the only single gene family in the *CYP71* clan (Nelson,

2019). Also, *CYP704* is often maintained as a single copy or at low copy numbers (Hansen *et al*.,

2021). In the 4CL family, while bona fide *4CL* genes have undergone gene family expansion,

ACOS genes are encoded by single-copy genes in most plants (de Azevedo Souza *et al*., 2009).

Out of 35 plants we examined, only the genomes of *P. patens* and *Glycine max* have two copies

of *ACOS* and the rest have a single copy (Supplementary Data Table S1). Likewise, whereas

multiple copies of *CHS* genes (encoding chalcone synthases) are found in most plants, only one

(bryophytes, lycophytes, gymnosperms) or two copies of *ASCL* (angiosperms) were found in plant

species that we examined (Suh and Ashton, 2022). As for *TKPR*, we found two copies of *TKPR* in

most plant species and a single copy in a few species including *Ginkgo biloba* and *Spirodela*

polyrhiza (Supplementary Data Table S2). These data indicate that gene dosage of all five genes

in the pathway has been subjected to strict control during evolution. Indeed, the genomes of

arabidopsis and rice contain one copy each of *CYP703A*, *CYP704B* and *ACOS* and two copies each

of *ASCL* and *TKPR*. The genome of *P. patens* has one copy each of *ACOS* and *ASCL* and two

copies of *TKPR* despite undergoing two whole genome duplications (Rensing *et al.*, 2008; Barker

and Ashton, 2016; Lang *et al.*, 2018). We argue that the evolutionary benefits of gene dosage

control were maintenance of steady metabolic flux and avoidance of the accumulation of

intermediate(s) that could be prematurely incorporated into SP, harming its structural integrity.

Synteny of genes of the polyketide pathway in model plants

Synteny of genes of the polyketide pathway can be observed in some model plants for which

sequenced whole genome assemblies are available. The most obvious example is arabidopsis in

which five genes, *CYP703A2*, *PKSA*, *ACOS*, *TKPR2* and *CYP704B*, are located on Chromosome

1 (Chr 1) and two genes, *PKSB* and *TKPR1*, are present on Chr 4. (*PKSA* and *PKSB* are *ASCL*

orthologues in arabidopsis (Kim *et al*., 2010)). It is unlikely that the synteny of Chr 4 resulted from

a recent segmental duplication since *PKSB* and *TKPR1* on Chr 4 are much closer together (0.23

Mb) than *PKSA* and *TKPR2* on Chr 1 (25.4 Mb) and are not separated by a recognisable *ACOS*

gene or CEN4 (Fig. 3).

We have calculated the probabilities by chance alone of the detected levels of synteny in

arabidopsis, correcting for confounding factors by subtracting the amounts of heterochromatin

(centromeres (CENs) and nucleolar organising regions (NORs)) in each chromosome and in the

whole genome and calculating the actual span of syntenic genes in the euchromatin, as follows:

Prob = (length of euchromatin spanned by syntenic genes in a single chromosome/total

euchromatin in genome)^{n}

Where n = number of syntenic genes in the chromosome

Based on our calculations, the probabilities of the observed synteny arising by chance alone are

low, e.g. 1 in 3,290 for the five genes on Chr 1 (This probability estimate may be higher than the

actual probability since the CEN1 sequence contains a gap and, based on the physical map, CEN1

is approx. 9 Mb (Hosouchi *et al*., 2002)) and 1 in 243,000 for *PKSB* and *TKPR1* on Chr 4. We

conclude that synteny must have provided a selective advantage during the evolution of SP. The most obvious possibility is that it facilitated epigenetic coordinate gene regulation thereby ensuring appropriate spatio-temporal expression. Interestingly, some syntenic genes, e.g. *PKSB* and *TKPR1* that span approx. 0.2 Mb of euchromatin on Chr 4, are close enough together to occupy a single euchromatin loop (0.2–2 Mb) in the chromocenter-loop model of arabidopsis interphase chromosomes (Fransz *et al*., 2002). That said, it is also clear that the number of syntenic genes of the pathway and their degree (closeness) of synteny varies widely between taxa (Fig. 3). Three genes of the polyketide pathway are located on Chr 1 and two more on Chr 7 of *Zea mays*, two are on Chr 8 of rice, two are on Chr 2 of *Physcomitrium patens*, two are on Chr 1 of *Marchantia polymorpha* and two are on Chr 32 of *Ceratopteris richardii*. Interestingly, a chromosome in each of four taxa, Chr 1 of arabidopsis, Chr 1 of *Z. mays*, Chr 8 of rice and Chr 2 of *P. patens* possesses a *CYP703A* or *CYP704B* gene plus either a *TKPR* gene or an *ASCL* orthologue or both. The distances between six gene pairs, four in arabidopsis, one in *P. patens* and one in *M. polymorpha* are such that each gene pair could be accommodated within a single euchromatin loop. However, we conclude other means of achieving coordinate gene expression must be operating as well as epigenetic mechanisms. Also we cannot rule out synteny conferring a selective advantage by an entirely different mechanism. As more sequenced genome level assemblies become available, it will be interesting to discover how widespread this phenomenon is.

Conclusions

All evidence indicates that the polyketide pathway and therefore SP is embryophyte-specific. The addition of the polyketide pathway in the definition of genuine SP will allow SP to be distinguished unequivocally from algaenans and other chemically resistant 'SP-like' algal spore wall substances.

This new, expanded definition of SP implies that SP was a major evolutionary innovation unique to the embryophyte lineage that may have contributed to plant terrestrialisation. This study should also spur research on possible evolutionary relationships between genuine embryophyte SP and other 'SP-like materials' in reproductive bodies including algal zygospores and dinoflagellate cysts.

SUPPLEMENTARY DATA

Supplementary data are available online at https://academic.oup.com/aob and consist of the following.

Table S1: Sequences used for 4CL family tree construction (Fig S1).

Table S2: Sequences used for CCR family tree construction (Fig S2).

Fig. S1: Maximum Likelihood tree of embryophyte ACOS and 4CL sequences and algal homologues.

Fig. S2: Maximum Likelihood tree of TKPR and other enzymes of the CCR family and algal homologues.

Fig. S3: Multiple sequence alignment of ACOSs, 4CLs and algal homologues.

Fig. S4: Multiple sequence alignment of representative enzymes of the embryophyte CCR family and algal homologues.

FUNDING

This work was supported by the Natural Sciences and Engineering Research Council of Canada Discovery grant [RGPIN-2018-04286 to D.-Y.S.). D.K.S. was supported in part by University of Regina Graduate Scholarships.

ACKNOWLDEGEMENTS

We thank William B. Browing (Southern Illinois Univ., USA) for informing us of the absence of *ASCL* in the *Zingiber officinale* genome. D-YS conceived the concept and D-YS, DKS and NWA collected and analysed data. D-YS and NWA wrote the manuscript.

LITERATURE CITED

Abe I, Morita H. 2010. Structure and function of the chalcone synthase superfamily of plant type III polyketide synthases. *Natural Product Reports* **27:** 809–838.

de Azevedo Souza C, Barbazuk B, Ralph SG, Bohlmann J, Hamberger B, Douglas CJ. 2008. Genome-wide analysis of a land plant-specific *acyl:coenzymeA synthetase* (*ACS*) gene family in *Arabidopsis*, poplar, rice and *Physcomitrella*. *New Phytologist* **179**: 987–1003.

de Azevedo Souza C, Kim SS, Koch S, *et al*. **2009**. A novel fatty acyl-CoA synthetase is required for pollen development and sporopollenin biosynthesis in *Arabidopsis*. *Plant Cell* **21**: 507–525.

Baker ME, Blasco R. 1992. Expansion of the mammalian 3β-hydroxysteroid dehydrogenase/plant dihydroflavonol reductase superfamily to include a bacterial cholesterol dehydrogenase, a bacterial UDP-galactose 4-epimerase, and open reading frames in vaccinia virus and fish lymphocystis disease virus. *FEBS Letters* **301**: 89–93.

Barker EI, Ashton NW. 2016. Ancestral and more recently acquired syntenic relationships of MADS-box genes uncovered by the *Physcomitrella patens* pseudochromosomal genome assembly. *Plant Cell Reports* **35**: 505–512.

Delwiche CF, Graham LE, Thomson N. 1989. Lignin-like compounds and sporopollenin in Coleochaete, an algal model for land plant ancestry. *Science* **245**: 399–401.

Feofilova EP. 2010. The fungal cell wall: modern concepts of its composition and biological function. *Microbiology* **79**: 711–720.

Ferrer J-L, Austin MB, Sterwart Jr C, Noel JP. 2008. Structure and function of enzymes involved in the biosynthesis of phenylpropanoids. *Plant Physiology and Biochemistry* **46**: 356–370.

Fransz P, de Jong JH, Lysak M, Castiglione MR, Schubert I. 2002. Interphase chromosomes in *Arabidopsis* are organized as well define chromocenters from which euchromatin loops emanate. *Proceedings of the National Academy of Sciences of the USA* **99**: 14584–14589.

Firn RD, Jones CG. 2000. The evolution of secondary metabolism—a unifying model. *Molecular Microbiology* **37:** 989–994.

Grienenberger E, Kim SS, Lallemand B, *et al*. 2010. Analysis of *TETRAKETIDE α-PYRONE REDUCTASE* function in *Arabidopsis thaliana* reveals a previously unknown, but conserved, biochemical pathway in sporopollenin monomer biosynthesis. *Plant Cell* **22**: 4067–4083.

Grienenberger E, Quilichini TD. 2021. The toughest material in the plant kingdom: An update on sporopollenin. *Frontiers in Plant Science* **12**: 703864.

Hahlbrock K, Scheel D. 1989. Physiology and molecular biology of phenylpropanoid metabolism. *Annual Review of Plant Physiology and Plant Molecular Biology* **40:** 347–369.

Hansen CC, Nelson DR, Møller BL, Werck-Reichhart D. 2021. Plant cytochrome P450 plasticity and evolution. *Molecular Plant* **14:** 1244–1265.

Heslop-Harrison J. 1969. An acetolysis-resistant membrane investing tapetum and sporogenous tissue in the anthers of certain Compositae. *Canadian Journal of Botany* **47:** 541–542.

Hosouchi T, Kumekawa N, Tsuruoka H, Kotani H. 2002. Physical map-based sizes of the centromeric regions of *Arabidopsis thaliana* chromosomes 1, 2 and 3. *DNA Research* 9: 117–121.

Hu Y, Gai Y, Yin L, *et al*. 2010. Crystal structures of a *Populus tomentosa* 4-coumarate:CoA

ligase shed light on its enzymatic mechanisms. *Plant Cell* **22:** 3093–3104.

Jiang C, Kim SY, Suh D-Y. 2008. Divergent evolution of the thiolase superfamily and chalcone

synthase family. *Molecular Phylogenetics and Evolution* **49**: 691–701.

Kallberg U, Oppermann U, Jörnvall H, Persson B. 2002. Short-chain dehydrogenase/reductase

(SDR) relationships: A large family with eight clusters common to human, animal, and plant

genomes. *Protein Science* **11**: 636−641.

Kim SS, Grienenberger E, Lallemand B, *et al*. 2010. *LAP6/POLYKETIDE SYNTHASE A* and

LAP5/POLYKETIDE SYNTHASE B encode hydroxyalkyl α-pyrone synthases required for

pollen development and sporopollenin biosynthesis in *Arabidopsis thaliana*. *Plant Cell* **22**:

4045–4066.

Lacombe E, Hawkins S, Van Doorsselaere J, *et al*. 1997. Cinnamoyl CoA reductase, the first

committed enzyme of the lignin branch biosynthetic pathway: cloning, expression and

phylogenetic relationships. *Plant Journal* **11:** 429–441.

Lamb DC, Hargrove TY, Zhao B, *et al*. 2021. Concerning P450 evolution: Structural analyses

support bacterial origin of sterol 14α-demethylases. *Molecular Biology and Evolution* **38:** 952–

967.

Lang D, Ullrich KK. Murat F, *et al.* 2018. The *Physcomitrella patens* chromosome-scale

assembly reveals moss genome structure and evolution. *Plant Journal* **93**: 515–533.

Li H, Pinot F, Sauveplane V, *et al*. 2010. Cytochrome P450 family member CYP704B2 catalyzes

the ω-hydroxylation of fatty acids and is required for anther cutin biosynthesis and pollen exine

formation in rice. *Plant Cell*, **22**: 173–190.

Linne U, Schäfer A, Stubbs MT, Marahiel MA. 2007. Aminoacyl-coenzyme A synthesis

catalyzed by adenylation domains. *FEBS Letters* **581**: 905–910.

Morant M, Jørgensen K, Schaller H, *et al*. **2007**. CYP703 is an ancient cytochrome P450 in land

plants catalyzing in-chain hydroxylation of lauric acid to provide building blocks for

sporopollenin synthesis in pollen. *Plant Cell* **19**: 1473–1487.

Nelson DR. 2009. The Cytochrome P450 Homepage. *Human Genomics* **4**: 59–65.

Nelson DR. 2019. Cytochrome P450s in the sugarcane *Saccharum spontaneum*. *Tropical Plant*

Biology **12:** 150–157.

Petit P, Granier T, d'Estaintot BL, *et al*. **2007**. Crystal structure of grape dihydroflavonol 4-

reductase, a key enzyme in flavonoid biosynthesis. *Journal of Molecular Biology* **368:** 1345–

1357.

Rensing SA, Lang D, Zimmer AD, *et al***. 2008**. The *Physcomitrella* genome reveals evolutionary

insights into the conquest of land by plants. *Science* **319**: 64–69.

Rowley JR, Southworth D. 1967. Deposition of sporopollenin on lamellae of unit membrane

dimensions. *Nature* **213:** 703–704.

Subbarayudu S, Shankar Naik B, Sunitibala Devi H, Bhau BS, Shaik Sha Valli Khan P. 2014.

Microsporogenesis and pollen formation in *Zingiber officinale* Roscoe. *Plant Systematics and*

Evolution **300:** 619–632.

Shimizu Y, Ogata H, Goto S. 2017. Type III polyketide synthases: Functional classification and

phylogenomics. *Chembiochem* **18:** 50–65.

Suh D-Y, Ashton NW. 2022. A sporopollenin definition for the genomics ages. *New Phytologist*

236: 2009–2013.

de Vries J, Archibald JM. 2018. Plant evolution: landmarks on the path to terrestrial life. *New*

Phytologist **217**: 1428–1434.

Wang B, Yang X, Jia Y, *et al*. **2022**. High-quality *Arabidopsis thaliana* genome assembly with

nanopore and HiFi long reads. *Genomics Proteomics & Bioinformatics* **20**: 4–13.

Xue JS, Qiu S, Jia X-L, *et al*. **2023**. Stepwise changes in flavonoids in spores/pollen contributed

to terrestrial adaptation of plants. *Plant Physiology* **193:** 627–642.

Yoshida Y, Aoyama Y, Noshiro M, Gotoh O. 2000. Sterol 14-demethylase P450 (CYP51)

provides a breakthrough for the discussion on the evolution of cytochrome P450 gene

superfamily. *Biochemical and Biophysical Research Communications* **273:** 799–804.

Figure captions

Fig. 1. The origin and evolution of the polyketide pathway in sporopollenin biosynthesis.

ACOS, ASCL and TKPR in the polyketide pathway are shown with their evolutionarily related

enzymes in the phenylpropanoid pathway. Also shown are the enzymes of primary metabolism

proposed to share the same ancestors with the related enzymes in the polyketide and

phenylpropanoid pathways. Related enzymes are indicated by text in the same colour. For instance,

ACOS and 4CL are thought to have evolved from the ancestral form of acyl-CoA synthetase in

fatty acid metabolism and are presented using blue text. 4CL, 4-coumarate:CoA ligase; ACOS,

acyl-CoA synthetase; ANR, anthocyanidin reductase; ASCL, anther-specific chalcone synthase-

like; CCR, cinnamoyl-CoA reductase; CHS, chalcone synthase; DFR, dihydroflavonol 4-reductase.

For further information about the enzymes in the pathways, readers are referred to Grienenberger

and Quilichini (2021) and Ferrer et al. (2008).

Fig. 2. ACOS- and TKPR-specific residues at their substrate binding sites.

(**A**, **B**) The modeled structure of arabidopsis ACOS (AtACOS5, blue) is superimposed with the crystal structure of tobacco 4CL (Nt4CL2, PDB id, 5BST, orange). Their acid substrate binding pockets (white-boxed) are magnified in **B** to show the uniquely conserved residues in ACOS (Gly^{241}, Ser^{339}, Cys^{340}, Ile^{341}, Thr^{342} and Thr^{344}) and 4CL (Ser^{243}, Gly^{339}, Pro^{340}, Val^{341}, Leu^{342} and Met^{344}). These substrate-determining residues, also shown in Supplementary Data Fig. S3, allow facile identification of ACOS from its paralogues, including 4CL. (**C**, **D**) The modeled structure of *P. patens TKPR* (PpTKPR1, blue) is superimposed with the crystal structures of *Medicago truncatula* CCR (MtCCR; PDB id, 4R1U, orange), *Panicum virgatum* DFR (PvDFR; PDB id, 8FEM, cyan) and *Sorghum bicolor* ANR (SbANR; PDB id, 8FIP, purple). Their substrate binding pockets (white-boxed) are magnified in **D** and the TKPR-specific Arg^{137} and Ser^{197} in PpTKPR1 are substituted with Tyr^{123} and Val^{180} in MtCCR, Asn^{148} and Val^{199} in PvDFR, and Tyr^{138} and Pro^{206} in SbANR (Supplementary Data Fig. S4), conferring their different substrate specificity.

Fig. 3. Synteny of genes of the polyketide pathway in model plant genomes

The ideograms are not drawn to scale. Genes are colour-coded as follows: *CYP703A* and *CYP704B*

(black), *ASCL* (red), *ACOS* (blue), *TKPR* (purple). (*PKSA* and *PKSB* are *ASCL* orthologues in

arabidopsis). Numbers in brackets refer to DNA in Mb spanning pairs of genes. CEN1 (3.8 Mb)

is approximately 15 Mb from the left end of Chr 1 of arabidopsis and separates *PKSA* and *ACOS*.

Distances within or close to the range (0.2–2.0 Mb) for a euchromatin loop in the arabidopsis

chromocenter-loop model of interphase chromosomes are indicated by an asterisk. Probabilities

by chance alone of the observed degrees of synteny were calculated (*Prob = (length of DNA*

spanned by syntenic genes of the pathway in a single chromosome/total genomic scaffold length)^{n}.

Where n = number of syntenic genes in the chromosome) and are shown as fractions given to three

significant figures. The values shown in brackets for arabidopsis have been corrected for

heterochromatin. Data used for this figure were obtained from Phytozome 13 and also, in the case

of arabidopsis, from Wang *et al*. (2022).