Extensive longevity and DNA virus-driven adaptation in nearctic *Myotis* bats

Juan M Vazquez[1,2,20], M. Elise Lauterbur[3,4,20], Saba Mottaghinia[5], Léa Gaucherand[6], Sarah Maesen[5], Michael Singer[7], Sarah Villa[7], Melanie Bucci[3], Devaughn Fraser[8,9], Genavieve Gray-Sandoval[3], Zeinab R Haidar[10,111], Melissa Han[12], William Kohler[12], Tanya M. Lama[13], Amandine Le Corf[5], Clara Loyer[5], Dakota McMillan[1,14], Stacy Li[1,15], Johnathan Lo[15], Carine Rey[5], Samantha LR Capel[8], Kathleen Slocum[16], Melissa Sui[7], William Thomas[17], Janet Debelak Tyburec[18], Rachel Brem[19], Richard Miller[12], Michael Buchalski[8], Jose Pablo Vazquez-Medina[1], Sébastien Pfeffer[6], Lucie Etienne[6,21]*, David Enard[3,21,22]**, Peter H Sudmant[1,14,21,22,23]***

Affiliations:

1. Department of Integrative Biology, University of California, Berkeley, Berkeley, CA USA
2. Current affiliation: Department of Biology, Pennsylvania State University, State College, PA USA
3. Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ USA
4. Current affiliation: Department of Biology, University of Vermont, Burlington, VT USA
5. Centre International de Recherche en Infectiologie (CIRI), Inserm U1111, UCBL1, CNRS UMR5308, Ecole Normale Supérieure ENS de Lyon, Université de Lyon, Lyon, France
6. Université de Strasbourg, Architecture et Réactivité de l'ARN, Institut de Biologie Moléculaire et Cellulaire du CNRS, Strasbourg, France.
7. Department of Molecular and Cellular Biology, University of California, Berkeley, Berkeley, CA USA
8. Wildlife Genetics Research Unit, Wildlife Health Laboratory, California Department of Fish and Wildlife, Sacramento, CA, USA
9. Current affiliation: Wildlife Diversity Program, Wildlife Division, Connecticut Department of Energy and Environmental Protection, Burlington, CT, USA
10. Department of Biology, California State Polytechnic University, Humboldt, Arcata, CA USA
11. Current affiliation: Western EcoSystems Technology Inc, Cheyenne, WY USA
12. Department of Pathology and Clinical Laboratories, University of Michigan, Ann Arbor, MI USA
13. Department of Biological Sciences, Smith College, Northampton, MA USA
14. Department of Science and Biotechnology, Berkeley City College, Berkeley, CA USA
15. Center for Computational Biology, University of California, Berkeley, Berkeley, CA USA
16. Bat Conservation International, Austin, TX USA
17. Department of Ecology and Evolution, Stony Brook University, Stony Brook NY USA
18. Bat Survey Solutions, LLC, Tucson, AZ USA
19. Department of Plant and Microbial Biology, University of California, Berkeley, Berkeley, CA 94720, USA
20. These authors contributed equally
21. Senior author
22. These authors contributed equally
23. Lead contact

* Correspondence: lucie.etienne@ens-lyon.fr
** Correspondence: denard@arizona.edu

40   *** Correspondence: psudmant@berkeley.edu

Summary Paragraph

The genus *Myotis* is one of the largest clades of bats, and exhibits some of the most extreme variation in lifespans among mammals alongside unique adaptations to viral tolerance and immune defense[1–5]. To study the evolution of longevity-associated traits and response to infectious disease, we generated cell lines and near-complete genome assemblies for 8 closely related species of *Myotis*. Using genome-wide screens of positive selection, analyses of structural variation, and functional experiments in primary cells, we identify new patterns of adaptation contributing to longevity, cancer resistance, and viral interactions in bats. We demonstrate distinct modes of adaptation to DNA and RNA viruses in bats, with bats demonstrating genome-wide overrepresentation of positive selection for DNA virus-interacting proteins, and significant rates of copy number variation for RNA virus-interacting proteins, in contrast to other mammals. We show that the recurrent evolution of longevity seen in *Myotis* leads to pervasive positive selection in cancer pathways, and demonstrate a unique response to DNA damage in primary cells of the long-lived *M. lucifugus*. Together, our results suggest that bats' remarkable longevity and immunity are linked through pleiotropic adaptations against viruses and aging-related disease.

Keywords

Aging, Bats, Cancer, Evolutionary Biology, Functional Genomics, Immunity, Infectious Disease

Introduction

Bats (order *Chiroptera*) represent approximately 20% of all known mammalian species and are one of the most phenotypically diverse clades of mammals[6,7]. Since their emergence 60 million years ago[8–10], many bat lineages have independently evolved a wide variety of life history strategies and phenotypic traits, including exceptional changes in longevity, viral tolerance, and immune defense[1,4,11–14]. Such systems, in which shared traits have evolved *de novo* multiple times, are powerful resources for dissecting the genetic basis of phenotypes. Rigorous approaches to studying these traits, however, depend on high-quality, well-annotated genomes to test evolutionary and genomic hypotheses, and on experimental functional systems to validate these hypotheses.

The largest genus of bats - *Myotis* - is estimated to have emerged approximately 33 million years ago[15,16], and encompasses over 139 described species spanning six continents and a wide range of ecological niches[6,15–17]. *Myotis* species demonstrate some of the most extreme variation in lifespan amongst mammals[1,2,18–20], including a six-fold difference in lifespan between the longest-lived species (*M. brandtii*, 42 years[18,21], **Fig. 1A**) and the shortest-lived species (*M. nigricans*, 7 yrs[18,22]) which diverged approximately 10.6 million years ago[10,17,23,24]. In addition, *Myotis* species have been used as systems for investigating virus tolerance and other pathogen resistance[25–27] associated with the expansion and contraction of antiviral defenses[28–31], which have contributed to bats' ecological role as zoonotic reservoirs[3,4,13,32–34].

While studies on the genetic basis of longevity in short-lived model organisms have been crucial for identifying and dissecting several key aging pathways, comparative studies of exceptionally long-lived

organisms have uncovered novel genes and pleiotropic effects governing lifespan[35–48]. Similarly, studies of
infectious disease response are common and powerful in model organisms, but the lack of diversity and use
of inbred lines limits their scope. Bats in particular present an important case study in, and opportunity to
study, variation in virus adaptation strategies due to bats' role as zoonotic reservoirs and their specific
resistance to viruses[44,49].

Here we combined comparative and functional approaches in *Myotis* to uncover genomic and
functional evidence of adaptation to both aging-related and infectious diseases. Using near-complete
genome assemblies, we identified significant structural variation encompassing stress response, immunity,
and inflammation genes, including a trans-species copy number polymorphism of Protein kinase R (PKR).
Additionally, we quantify the rates of evolution in body size and lifespan across mammals and find that both
ancestral and extant *Myotis* species are overrepresented at the extreme of increased lifespan. Consistent
with this observation, we identified pervasive selection of genes in longevity- and cancer-related processes
throughout *Myotis*. We observed some of the strongest signatures of selection at immune-related genes,
including strong evidence of adaptation in virus-interacting proteins in *Myotis* and other bats. In contrast to
humans and other primates, in which virus adaptation is driven by interactions with RNA viruses[50], we find
that virus adaptation in bats is driven by interactions with DNA viruses, implying that bats' exceptional
immune adaptations are more protective against DNA viruses. This also highlights a fundamental
discordance between the fitness impacts of viruses on humans vs bats. Together, our results suggest that
pleiotropy and co-evolution of traits in *Myotis* has played a key role in the evolution of exceptional longevity
and infectious disease resistance.

Results

High quality chromosome-level assemblies of 8 *Myotis* bat species

To study how lifespan and viral response have evolved in *Myotis*, we collected skin punches and
derived primary cell lines from several North American ("nearctic")[23] species (**Fig. 1A,C),** including from one
of the longest-lived mammals, *Myotis lucifugus[18]*. Using these cell lines and flash frozen tissues we generated
de novo haplotype-resolved, chromosome-scale genome assemblies for eight species (**Fig. 1A, Extended
Data Fig. 1**) using a combination of long-read PacBio HiFi sequencing and HiC scaffolding. These genomes
are highly contiguous, with an average of 98.6% of nucleotides assembled into 22-23 syntenic chromosome-
scale scaffolds corresponding to the published karyotype[51] with an average QV of 66 (Extended Data Fig. 1;
Extended Data Table 1). These genomes have among the highest contig NG50 and lowest auNG scores of
any *Chiroptera* genome published to date (**Fig. 1A, E**; **Extended Data Fig. 1**; **Extended Data Table 1**). We
identified an average of 20,869 protein coding genes with mammalian homologs per genome, with
BUSCO[52,53] scores ranging from 98.2% to 98.5% (**Fig. 1D**). Using these annotations, we examined the
conflicting phylogenies within nearctic *Myotis[15,17,54–57]* by using 1-to-1 orthologs of protein-coding genes
across 536 mammalian genomes (**Extended Data Table 2**) to build a time-calibrated, maximum likelihood
tree of *Eutheria*. Our results conclusively resolve the relationship between sister taxa (*M. lucifugus* and *M.
occultus*; *M. yumanensis and M. velifer*; and *M. evotis* and *M. thysanodes*) with both 100% bootstrap support
and 100% Local Posterior Probability (LPP) scores at all nodes within nearctic *Myotis*. However, the root of
nearctic *Myotis* remains unresolved with 0% LPP support despite 100% bootstrapping support, although our
placement of the nearctic common ancestor as sister to *Myotis brandtii* is supported by geography and recent
work in the literature[56–58]. (**Fig. 1B**; **Extended Data Fig. 2A**; **Extended Data Table 1**)

Across all 8 genomes, each autosome has been completely assembled telomere-to-telomere (T2T)
in at least one species (**Fig. 1E**). Within assemblies, 29%-70% of chromosomes are fully assembled with an
average of less than one gap per chromosome (**Extended Data Table 1**). Overall, these fully annotated
genomes represent some of the most contiguous mammalian assemblies to date.

Evolution of structural variation within constrained karyotypes

With our near-complete assemblies, we first sought to study the landscape of structural variation
within the tightly-conserved karyotype seen in *Myotis*. With only six exceptions across over 60 studied
species[59–63], all *Myotis* species have a conserved 2n=42 karyotype - a remarkable phenomenon for a genus
spread across six continents and 33 million years of divergence[6,7,16,51]. The conserved *Myotis* karyotype
consists of three large autosomes and one small metacentric autosome; 17 small telocentric autosomes; and
metacentric X and Y chromosomes[51,64]. Consistent with this broad cytological conservation, we find large
scale synteny across the nearctic *Myotis* in this study (**Fig. 2A**). However, structural variants (SVs) including
inversions, duplications, and translocations are relatively common within chromosomes, especially adjacent
to putative centromeric regions (**Fig. 2A, B**). This includes a ~20Mb block at the sub-telomeric end of Chr
V15 that displays frequent and recurrent inversions and translocations across the nearctic *Myotis*, and which
contains a 10Mb locus that was recently identified as a potential target of recent selection by adaptive
introgression[57] (**Extended Data Fig. 3D**).

Using SyRI[65], we identified between 6,813 - 8,013 SVs per nearctic *Myotis* genome relative to their
outgroup, *M. myotis*; 97 - 99% of these events were under 10Kb. In the three large autosomes, which
constitute ~30% of the genome, we cataloged an average of 509 SVs (**Extended Data Table 3**). In contrast,
in the small autosomes, constituting ~65% of the genome, we observed an average of 316 events,
highlighting the distinct structural evolution between these chromosome types with SVs found at 2-fold higher
density in larger autosomes (**Extended Data Table 3**). However, large (≥10Kb) duplications, large inverted
duplications, and large inverted translocations were more common on small autosomes compared to the
large autosomes (**Extended Data Table 3**). We also quantified the distribution of transposable elements
(TEs) across chromosomes (**Fig. 2B,C**). Consistent with prior observations in other Chiropteran genomes[66],
LINE elements appeared to be enriched around the predicted centromere (**Fig. 2B**) with simulations showing
that this trend is primarily limited to the long, metacentric autosomes (**Extended Data Fig. 3A**; **Extended
Data Table 3**). The concentration of segmental duplications was also significantly correlated with TE density
in each species (linear regression, p < 0.001; **Fig. 2B,D; Extended Data Fig. 3B**) highlighting the importance
of TEs in facilitating structural evolution. Together these results highlight high levels of structural variation
coincident with mobile element activity occurring in the context of a highly constrained karyotype.

An actively segregating, trans-species copy number polymorphism of the antiviral factor Protein Kinase R, PKR

Among the structural variants identified in nearctic *Myotis*, the gene Protein Kinase R (*PKR*/*EIF2AK2*)
stood out as an interferon-stimulated gene with antiviral activity against both DNA and RNA viruses. While
the functionality of this *Myotis*-specific duplication had been previously characterized[30], the absence of long-
read sequencing data for *Myotis* has prevented a more detailed characterization of copy number and
evolutionary history among Nearctic *Myotis.* Using our near-complete genome assemblies, we fully resolved
the sequence and structure of the two previously known haplotypes: H1, containing a single copy of *PKR*

(*PKR2*); and H2, containing two tandemly duplicated copies of *PKR* (*PKR1* and *PKR2*; **Fig. 3A**). We also identified a third haplotype - H3 - with three tandem duplicates of *PKR* (*PKR1*, *PKR2*, and a third copy) present only in *M. californicus*. Remarkably, while 7 out of 9 *Myotis* species carried duplicated haplotypes, 5 of these cases were heterozygous for the duplicated haplotype (i.e. H1/H2 or H1/H3; **Fig. 3B**). Two *Myotis* individuals (*lucifugus* and *evotis*) only carried non-duplicated (i.e. H1/H1; **Fig. 3B**). To determine the evolutionary history of the duplicates, we used GeneRax[67] to construct a tree from alignments of all *PKR* gene copies across Neartic *Myotis*, using *Pipistrellus pygmaeus* as a non-*Myotis* outgroup (**Fig. 3C**). Our results suggest that *PKR2* is the ancestral copy of *PKR*, and that *PKR1* originated from a single duplication event at the root of *Myotis*. These results highlight that both the duplicated and unduplicated haplotypes have likely been segregating for tens of millions of years, representing an ancient trans-species polymorphism.

PKR is a stress response and innate immune factor that becomes active upon detection of dsRNA through sensing, auto-phosphorylation and dimerization[68,69]. Active PKR shuts down protein translation and restricts viral replication[68,69]. Although PKR1 and PKR2 are both expressed in basal or interferon-stimulated *Myotis velifer* cells, only their independent functional impacts have previously been investigated[30]. Given the co-expression and the complex haplotype diversity we identified in *Myotis*, we set out to determine whether the two *PKR* copies act additively, synergistically, or as dominant negatives in several essential functions. We therefore investigated the functional impact of the duplicates' co-expression on total PKR protein expression levels, cell viability, inhibition of translation, antiviral restriction and homo/hetero-dimer formation (**Fig. 3D-H**). To pinpoint the contribution of PKRs, we used a heterologous cell system[30] with PKR-KO Hela cells transfected with either an empty pSG5 plasmid (control), or pSG5 encoding PKR1, PKR2, or PKR1+PKR2 (1:1 ratio) from *Myotis myotis* (myoMyo) or *Myotis velifer* (myoVel). In this system, PKR transfection leads to its autoactivation[30]. First, we investigated PKR protein expression and found that, although *Myotis* Flag-PKR1 was expressed at lower levels than Flag-PKR2, PKR1+PKR2 co-expression did not affect the overall PKRs' steady-state protein expression levels (**Extended Data Fig. 4A**).

Second, we performed cell viability assays at two doses of transfected PKRs and found that cell viability was only affected at high dose (**Extended Data Fig. 4D**). Third, using a luciferase reporter translational assay, we found that PKR1+PKR2 coexpression led to an intermediate translation shutdown, showing neither synergistic nor dominant negative effect of the PKR duplicates (**Fig. 3D**). Fourth, we tested PKR1+PKR2 co-expression effects on infections by two different model RNA viruses: VSV-GFP (Vesicular stomatitis virus encoding a GFP reporter[70], a representative of *Rhabdoviridae* family that is common in bats, including *Myotis*) and SINV-GFP (Sindbis virus, a representative of *Togaviridae*) (**Fig. 3E-F**, **Extended Data Fig. 4C,E**). We found that PKR1+PKR2 restricted viral infections to a similar, though slightly higher, extent as PKR1 or PKR2 alone (**Fig. 3E-F; Extended Data Fig. 4C-D**, $p>0.05$). While the recently duplicated PKR1s homodimerized, PKR1 and PKR2 did not form heterodimers (**Fig. 3G-H**, **Extended Data Fig. 4B** in SINV condition). This suggests that, since their duplication, divergence between *Myotis* PKR copies has been particularly strong at the protein-protein interface hindering their capacity to form heterodimers.

Together, these results suggest that PKR1 and PKR2 are neither dominant negative nor synergistic with each other, but rather act in a mostly additive form for their main cellular effector functions. Despite their capacity to more broadly escape some DNA and RNA viral antagonists[30], increased cell toxicity at high PKR dose suggests a possible tradeoff upon PKR duplication, which might participate in also explaining why PKR is rarely duplicated in mammals. Consistent with this, PKR1, which is generally more potent in our experiments, is expressed at significantly lower levels in *Myotis velifer* cells than PKR2[30]. Altogether, these genomic and functional results unveil gene duplicates still segregating across several *Myotis* species, possibly shaped by recent viral infections and cell toxicity tradeoff.

Genome-wide adaptation to DNA viruses

To further explore how viruses may have shaped the genomes of *Myotis* bats we tested for adaptive signatures in virus-interacting proteins (VIPs) in *Myotis* and other bats. VIPs are host proteins that physically interact with viral proteins (e.g. *CD45*, **Fig. 4A**), and can be proviral (contributing to viral infection, e.g. viral receptors), antiviral (protective against viral infection, e.g. interferons), or both depending on infection stage and virus type. VIPs commonly have major roles in basic host function, and as a result most host cell biological pathways include VIPs. Previous studies investigating positive selection across mammals have found an enrichment for adaptation among a set of 5,527 manually curated VIPs, defined as host proteins that have at least one experimentally verified physical interaction with a viral protein, RNA, or DNA[71].

By calculating an enrichment score from the ratio of positive selection in VIPs compared to their matched control genes using BUSTED-MH[72], we found that, like other mammals, *Myotis* show an enrichment for adaptation at VIPs (**Fig. 4D; Extended Data Table 5**). Physical host-virus interactions may not always result in fitness effects in the host. We therefore repeated our analysis using a gene set restricted to VIPs with experimental evidence of specific pro- or anti-viral effects, and thus with a stronger expectation of fitness effects. We expect that if viruses were not the primary driver of adaptation in VIPs, then there would be no difference in the enrichment of selection in VIPs with vs without demonstrated pro- or anti-viral effects. We observed an even stronger significant elevation in the ratio of positive selection in these proviral and antiviral VIPs (**Fig. 4B; Extended Data Table 5**), but no elevation in this ratio in VIPs without experimental evidence of specific pro- or anti-viral effects (**Fig. 4C; Extended Data Table 5**). This is consistent with the expectation of viral interaction as the cause of enrichment of positive selection in VIPs in bats[73]. We repeated this analysis using a dataset of 47 publicly-available non-*Myotis* bat genomes, and confirmed these same patterns across bats more broadly, even when excluding *Myotis* genomes (**Extended Data Fig. 5A**).

Previous work has suggested that bats may have different physiological responses to DNA and RNA viruses[74]. To determine if this was reflected in genomic VIP adaptation, we compared the enrichment of positive selection in VIPs that interact only with DNA viruses (DNA-only VIPs) to those that interact only with RNA viruses (RNA-only VIPs). Remarkably, we found that VIP adaptation in *Myotis* and other bats is driven by selection in DNA VIPs (**Fig. 4E, Extended Data Fig. 5B**) in marked contrast to the observed pattern in RNA VIPs, which show no evidence of genome-wide enrichment in adaptation (**Fig. 4F, Extended Data Fig. 5C**).

Adaptation to viral pathogens can also occur via gene copy number changes[75,76,77,78]. We tested whether VIPs were enriched among genes that were recently gained or lost throughout *Myotis* using the method of Huang *et al*. (2023) to test the cumulative per gene, per branch copy number changes and birth-death rates of VIPs compared to other non-VIP genes.[42] While the birth-death rate of VIP genes overall was no different than non-VIP genes (p = 0.071, **Extended Data Fig. 5D**), VIP genes were significantly more likely to have undergone expansions and/or contractions on at least one branch of the *Myotis* family (p < 0.001; **Fig. 4G**). Furthermore, we find that this pattern is driven exclusively by copy number changes in RNA-only VIPs, and not by changes in copy number of DNA-only VIPs (**Fig. 4H-I**). This suggests that there is variation in gene family birth rates across species, but that VIPs are more dynamic across the nearctic *Myotis* as a whole than other types of genes highlighting the importance of the viral immune response in recent *Myotis* evolution.

In contrast to what we observe in bats, VIP adaptation in humans is driven by positive selection in RNA - and not DNA - VIPs[71,79]. To investigate if DNA VIP-driven adaptation in bats is exceptional among

mammals, we replicated these analyses across four other large mammalian orders that are well represented
among publicly-available mammalian genomes: *Primates*, *Glires*, *Euungulata*, and *Carnivora*. We found that
while other mammalian orders show a mix of adaptation enrichments in both RNA and DNA VIPs, none show
an absence of genome-wide enrichment of adaptation in RNA VIPs as observed in bats (**Extended Data**
Fig. 5A-C). These results highlight that bats, including *Myotis*, may have faced greater selective pressures
from DNA viruses than from RNA viruses, in contrast to all other mammals.

Evolution of body size & lifespan across nearctic *Myotis*

Beyond their remarkable immune adaptations, bats are exceptional as the longest-lived clade of
mammal after correcting for their body size[2,20,80], demonstrating a ~11x range of lifespans within a ~650x
range of body sizes[1,80,81]; *Myotis* species in particular exhibit the full dynamic range of bat lifespans within a
narrow range of body sizes. While bats have been noted as an exception to the strong allometric scaling
(positive correlation with body size) of lifespan otherwise seen across mammals and other metazoans, this
exception has not been tested using phylogenetically corrected statistics leveraging well-resolved
phylogenies.

To test the hypothesis of non-allometric scaling of lifespan in bats, we modeled the evolution of body
size and lifespan independently across a supertree of over 1000 mammals[55] (**Fig. 5A-B; Extended Data**
Fig. 6A-B; **Extended Data Table 6**). We generally observed agreement between evolutionary patterns of
body size and lifespan in the two simulations, such as in whales (*Cetacea*)[82,83], elephantids
(*Proboscidea*)[38,84,85], and primates (**Fig. 5A-B**). However, while only minor changes were observed in their
body size, we observed some of the largest and most-rapid changes in lifespan across mammals in bats
(**Fig. 5A-B; Extended Data Fig. 6A-B**; **Extended Data Table 6**). These changes are largely independent
between species and genera, consistent with the theory of multiple independent increases in lifespan across
bats. We used phylogenetically-corrected generalized linear models and ANCOVA to quantify the
relationship between body size and lifespan across mammals. While we find that non-bat mammals
experience an average 0.159% increase in lifespan per 1% increase in body size, bats experience a 0.223%
increase in lifespan years per 1% increase in body size. After phylogenetic correction, however, these rates
were not significantly different (**Extended Data Fig. 6C-D**; pANCOVA: p=0.29). In *Myotis*, we saw many of
the fastest increases in lifespan relative to their most recent ancestor, including for *Myotis grisescens* (4.15x
increase, 100th percentile), *Myotis brandtii* (2.25x increase, 100th percentile), *Myotis lucifugus* (1.56x, 98th
percentile), *Myotis myotis* (1.1x increase, 79nd percentile), and the *Myotis* common ancestor (1.26x increase,
92rd percentile) (**Fig. 5B, Extended Data Fig. 6B**). Together these results demonstrate that *Myotis* bats and
their recent ancestors have evolved some of the most extreme increases in lifespan among mammals,
despite similar allometric lifespan scaling of bats and mammals after phylogenetic correction.

Evolutionary signatures of cancer resistance and pleiotropy with DNA virus adaptations in *Myotis*

Rapid changes in body size and lifespan can have major implications for the evolution of cancer risk
and resistance across mammals. The multistage model of carcinogenesis[86–89] posits that cancers arise from
mutations accumulated during development and over an organism's life; and that lifetime *cancer risk* scales
linearly with body size, and with lifespan as a power-law with exponent 6[88–91]. While these positive
correlations have been well-established *within species*[92–97], when comparing *between* species and

accounting for phylogeny and other life history traits, there is only a weak correlation between body size and
cancer risk,– and no correlation between lifespan and cancer risk[91,96–99]. This observation, which has been
coined *Peto's Paradox*, suggests that species with more cells or longer lifespans have adapted to reduce
their cancer risks (i.e. increased cancer resistance) (**Fig. 5C**).

We hypothesized that the extreme changes in lifespan observed throughout the *Myotis* phylogeny
should exert a selective pressure on genes associated with cancer processes. Using aBSREL[100] to test for
selection at terminal and internal branches within *Myotis*, we found that at FDR<=5%, an average of 5.46%
of protein-coding genes had at least one region under positive selection per node (omega >1); in contrast,
an average of 20.8% of protein-coding genes per node had all regions with an omega less than 1, suggesting
negative selection (**Extended Data Table 7**). Genes under either positive or negative selection were enriched
for several pathways in immunity, cancer, and aging, with many sitting at the intersection of multiple
processes, suggesting possible pleiotropic selective pressures (**Extended Data Table 7**). Confirming our
methods, many of the genes and pathways we identify as under selection in *Myotis* have been previously
identified in the literature, such as members of the Cluster of Differentiation (CD) family[26,101], Serpin family[44],
insulin signaling pathway[39,102], redox repair[103], and iron storage[104,105] (**Extended Data Fig. 6E; Extended**
Data Table 7).

We quantified the contribution of genes under positive selection to pathways associated with the
hallmark of cancer[106–108] by measuring the proportion of cancer-associated pathways overrepresented
among genes under positive selection throughout the phylogeny (**Fig. 5E**). We observed that most nodes
within nearctic *Myotis* were enriched for cancer hallmark pathways among genes under positive selection,
especially at the recent ancestors of the longest-lived species (e.g. *M. lucifugus, M. occultus*; **Fig. 5E**).
Furthermore, when considering the union of all genes under positive selection in the evolutionary history of
each species since their common ancestor, we observed significant enrichments in the representation of
cancer-associated pathways in many of the lineages with the greatest cumulative increases in lifespan (*M.*
lucifugus, M. occultus, M. evotis, M. yumanensis; **Fig. 5F, Extended Data Fig. 6F**). This suggests that
selection in genes involved in cancer-related processes occurred concurrently with the recurrent increases
in lifespan seen in this clade.

The longest-lived bat in our study, *M. lucifugus*, had an overrepresentation of pathways specifically
associated with DNA double-strand break (DSB) repair when looking at both lineage-wide and node-specific
enrichments in positive selection using the Reactome database[109] (**Fig. 5G; Extended Data Table 7**). This
includes 35 out of 65 genes in the high-fidelity Homologous Recombination Repair pathway, and 21/37
members of the Homology-Directed Repair via Single Strand Annealing (**Fig. 5G; Extended Data Fig. 6E;**
Extended Data Table 7). These results suggest that *M. lucifugus* might have an enhanced response to DNA
DSBs relative to other bats. To test this hypothesis, we assessed the tolerance of *M. lucifugus* to
neocarzinostatin (NCS), a potent radiomimetic agent that induces DNA double-strand breaks[110–112] (**Fig. 5H**),
compared to *M. evotis*, three non-*Myotis* bats (*Eidolon helvum, Pteropus rodrigensis,* and *Rousettus*
lanosus). At low doses of neocarzinostatin, *M. lucifugus* was the only species tested showing sensitivity to
neocarzinostatin after 24 hours, with a drop in viability and concomitant increase in apoptosis. At high doses,
M. lucifugus had the highest level of apoptosis and the greatest drop in viability of all the bats tested. This is
consistent with other long-lived species, including elephants[36,37,113], naked mole rats[46], and bowhead
whales[40,114], where longevity are associated with an increased ability to clear out damaged cells.

To better understand the genes that may be driving this difference, we examined RNA expression in
M. lucifugus cells after 6 and 18 hours of treatment with 100nM NCS. As expected, at 6 hours we observe

up-regulation of genes associated with cell cycle arrest (e.g. *MDM2*, *CDKN1A*) and cell death (e.g. *BAX*), and down-regulation of genes in pathways associated with cell division and growth at 6 hours. At 18 hours, we observe up-regulation of genes in pathways associated with cellular stress response and continued down-regulation of the cell cycle (**Fig. 6A-D, Extended Data Fig. 7A, Extended Data Table 9**). At 6 hours, however, we also observed down-regulation of genes associated with DNA damage repair and synthesis, which together with the up-regulation of *BAX* is consistent with the observed sensitivity to NCS in *M. lucifugus*.

Unexpectedly, however, we observed that the strongest pathway enrichment in our DNA-damage RNA-seq data were genes associated with the innate immune response, including genes in the PI3K/AKT pathway (**Fig. 6C-D, Extended Data Table 9**). We hypothesized that the observed relationship between the unique DNA damage response of *M. lucifugus* may be due to pleiotropy with DNA VIPs and selection in these genes. Indeed, we observed that there was a highly significant enrichment for genes that are differentially expressed after NCS treatment and DNA-only VIPs (p=3.34e-5, **Fig. 6E**; α=0.0245, **Extended Data Fig. 7B**). The genes at the intersection of DNA-only VIPs and NCS response are overrepresented in pathways associated with the cell cycle, senescence, transcription, and DNA damage repair (**Fig. 6F**). To test how selection may have impacted these pleiotropic signatures, we repeated these analyses only considering genes under selection in *Myotis lucifugus*. The intersection between DNA-only VIPs and NCS-DE genes remained highly significant (p=1.88e-3, **Extended Data Fig. 7C**) with only three pathways remaining enriched at FDR<=0.05: Cell Cycle, DNA Damage Repair, and Deubiquination (**Fig. 6G**); DNA-only VIP genes which are not NCS-responsive did not show these signatures (**Extended Data Fig. 7D-G).** Together these results suggest that selection on the innate immune response may drive *agonistic* pleiotropy in aging-associated traits like DNA damage response and cell cycle regulation.

Discussion

Bats are widely known for their long lifespan, and viral tolerance[1,4,13,44,80,97,115–117]. As highly complex and pleiotropic processes, the genes and mechanisms underlying these phenotypes can be challenging to identify, especially for rapidly-evolving phenotypes such as host-pathogen interactions. Here we outline an approach that enables functional comparative biology by generating cell lines from wing punches of wild caught bats for genome assembly, comparative genomics, and functional follow up. Cell lines are generated from minimally-invasive biopsies collected in the field thus avoiding disturbing natural populations. Using this approach, we identify new patterns of adaptation contributing to longevity, cancer resistance, and viral interactions in bats, and demonstrate a unique DNA damage response in primary cells of the long-lived *M. lucifugus.* Given the high density of bat species concentrated at single locations world-wide[118,119] it is feasible to collect wing punches from a large number of individuals across a wide phylogenetic range; these wing punches can be used to generate cell lines and sequencing libraries for reference genomes in a matter of weeks. Furthermore, we find that primary-cell-line-derived genomes are of higher quality than those derived from primary tissues (**Figure S1**). As this approach can generate genomic resources from minimal material gathered via non-lethal sampling, it is also well-suited for conservation genomics and the study of rare or endangered species for which acquiring sufficient amounts of material can be challenging[120–122].

The evolution of body size and lifespan across mammals - and in particular, the rapid evolution of lifespan such as in Yangochiroptera - has major implications for the co-evolution of cancer risk and resistance. While models of body size evolution are well-studied in mammals[11,38,84,123] the evolution of

lifespan is less well understood. We find that many extant and ancestral *Myotis* species have experienced some of the fastest rates of change in lifespan seen in mammals, despite minor changes in body size. This altered scaling of longevity in *Myotis* is predicted to have serious consequences for their intrinsic, per-cell cancer risk. Similar to other systems where the evolution of cancer resistance has been largely driven by rapid changes in body size[36–38,45,124,125], the rapid and repeated changes in lifespan across an order of magnitude observed in *Myotis* are hypothesized to result in strong selective pressures on lifetime cancer suppression to avoid premature death by malignancies[80,86,91,126]. This is supported both by recent meta-analyses on cancer risk in vertebrates[96,97,127] demonstrating little to no correlation between neoplasias and either species body size or lifespan; as well as by studies showing weak or no correlation between mutation rates and either species body size or lifespan, respectively[128–130]. For bats, however, there are very limited records for neoplasias in the literature - and only for Yinpterochiropterans[96,97]. The difficulties of maintaining captive colonies of *Myotis* and other Yangochriopterans[131], coupled with their exceptional longevity, necessitates the use of proxy measurements for cancer resistance in these species, such as measuring somatic mutation rates across lifespan and the use of *in vitro* models of carcinogenesis to experimentally test the cancer resistance of long-lived bats[117].

Bats are also important models for understanding infectious disease response and immune adaptation because of their unique resistance to viruses and role as zoonotic reservoirs[3,74,132]. Here, we show that while bats have adapted to both DNA and RNA viruses, they demonstrate a remarkable dominance in adaptations to DNA viruses. This is in contrast to humans and other primates, in which adaptation to RNA viruses dominates[50,79]. While bats are known to mount adaptive immune responses to some RNA viruses, this suggests that adaptations in bats' innate and adaptive immune systems, such expansions in *MHC-I* and *APOBEC3*[44], or the loss of the *PYHIN* gene family[49], may be sufficient as a first line of defense to protect against the fitness impacts of some RNA viruses, but not against DNA viruses. This is consistent with previous observations that most zoonoses, including those hosted by bats, are RNA viruses[13] and that bats are more likely than other mammals to asymptomatically harbor RNA viruses, while being more susceptible themselves to other pathogens such as fungi[74]. We also recapitulate and expand on previous results related to positive selection in immune genes and immune gene family expansion, contraction, and loss. In addition, we demonstrate complex patterns of structural variation in immune genes, including a segregating duplication of the Protein Kinase R gene (*PKR*), a major protein involved in the antiviral innate immune system, that has functional relevance in its activity against both DNA and RNA viruses[30]. By resolving individual haplotypes in nine *Myotis* species, we were able to confirm a single duplication event at the origin of *Myotis PKR1* and *PKR2*. We further demonstrate the functional implications of copy number variation in *PKR*, including differences in copy expression and an additive effect on cellular effector functions.

Multiple hypotheses have been proposed to connect the unique physiology and ecology of bats with the evolution of remarkable adaptations such as viral infection tolerance, stress tolerance, and exceptional longevity[115]. Hypothesized drivers of disease resistance and longevity evolution in bats include the evolution of flight (e.g. "flight as fever" hypothesis[133], though this hypothesis has recently been critiqued[134]), the disposable soma hypothesis[135]; metabolic state[136]; torpor[1]; and other adaptations to specific environments[101,137–139]. Additionally, many studies have highlighted the intersection of these traits, including a relationship between hibernation and both longevity[1,132] and infectious disease resistance[74,132]. Our results are consistent with an *agonistic* pleiotropy hypothesis, wherein genetic adaptations for many specific traits (e.g. DNA virus innate immunity) may prove beneficial to other seemingly-unrelated traits (e.g. cancer resistance, cellular homeostasis, longevity). Importantly, bats are also the only mammal to harbor active DNA

transposons[140,141], the suppression of which may also contribute to signatures of selection in DNA-interacting
VIPs.

Consistent with the agonistic pleiotropy hypothesis, many of the genes and pathways highlighted in
this study have been found to play vital roles across physiological traits in bats and other species. As a core
part of the integrated stress response PKR is not only associated with the innate immune response, but also
has been identified as a longevity-associated gene in multiple studies[142–144]. Additionally, two genes under
selection in nearctic *Myotis - FTH1* and *IGFN1* - have been implicated in functional studies as key hibernation
genes[145–147], viral interacting proteins[148–151], and as pro-longevity genes[152–154]. Similarly, many DNA VIPs
such as *BRCA1/2* and *POLG* are core DNA maintenance genes essential for cancer resistance and
longevity[46,155–161]. Such essential genes are generally under increased constraint; however, it is possible that
internal pressures, such as DNA transposon activity, and external pressures such as viral infections and
other environmental stressors, are shaping the fitness landscape of these essential genes such that novel
optimizations become accessible to evolutionary processes[140]. Changes in structural genomic variation also
play a crucial role here, such as our findings related to PKR and RNA VIPs in *Myotis*. Many of the overarching
pathways we observed under selection, such as inflammation, senescence, and ferroptosis, lie directly at the
intersection of aging-related immune processes[44,49,147,152,162–169]. More broadly, our findings on pervasive
selection on DNA-only VIPs and the extreme evolution of longevity and cancer risk in other bats support a
more general hypothesis that traits such as cancer risk, cellular homeostasis, and antiviral response have
evolved in tandem due to pleiotropic selection at coinciding points in bats' evolutionary history. Further work
leveraging model systems such as *Myotis* for functional evolutionary genomics is necessary to disambiguate
the underlying selective pressures driving these functional genetic changes in bats and beyond.

431 Acknowledgements

PacBio HiFi sequencing was done by the DNA Technologies and Expression Analysis Cores at the UC Davis
Genome Center (NIH SIG 1S10OD010786-01). Analyses were performed using the High Performance
Computing (HPC) resources at the University of Arizona (supported by the University of Arizona TRIF, UITS,
and Research, Innovation, and Impact (RII); maintained by the UArizona Research Technologies
department); the Savio computational cluster resource provided by the Berkeley Research Computing
program at the University of California, Berkeley (supported by the UC Berkeley Chancellor, Vice Chancellor
for Research, and Chief Information Officer); and the Vermont Advanced Computing Core at . We thank the
members of the LP2L team (CIRI) for helpful discussions.

MEL was funded by NSF PRFB #2010884, and a Dovetail Tree of Life Grant. JMV was funded by NSF PRFB
#2109915, NIH NIA T32AG000266, and NIA 1K99AG088361. DE was funded by NIH NIGMS MIRA grant
5R35GM142677. PHS was funded by NIH NIGMS R35GM142916 and by the Vallee Scholars Award. LE
was funded by a grant from the Agence Nationale de la Recherche (ANR, #ANR-202-CE15-0020-01 to LE)
and CNRS MITI These Internationale (to M.S. and L.E.). SP is funded by the ANR (ANR-21-CE35-0018-01)
and by the Interdisciplinary Thematic Institute IMCbio+ supported by IdEx Unistra (ANR-10-IDEX-0002), by
SFRI-STRAT'US project (ANR-20-SFRI-0012) and EUR IMCBio (IMCBio ANR-17-EURE-0023) under the
framework of the French Investments for the Future Program. LE and DE were further funded by a grant
from the Joint Call for Proposals between the CNRS and the University of Arizona (International Research
Center, IRC, 2021-2024). LE, DE, PHS and SP were further funded by the International Research Project
(IRP) RAPIDvBAT from the CNRS, the University of Arizona and the University of California, Berkeley. LG

was funded by a postdoctoral fellowship from the Fondation pour la Recherche Médicale (FRM,
#SPF202209015746).

Image attributions for the bat photos featured in Fig. 1: J. Scott Altenbach (*M. thysanodes*); M. Elise Lauterbur
(*M. occultus, M. velifer*); U.S. National Park Service (*M. auriculus*); Frank Carey (*M. californicus*); Michael
Durham (*M. evotis*); J. N. Stuart (*M. volans*); and SMBishop & the Wikimedia Foundation (*M. lucifugus*).

Author Contributions:

Conceptualization, J.M.V., M.E.L., D.E., P.H.S., L.E., D.F., M.B.; Methodology, J.M.V., M.E.L., D.E.,
P.H.S., L.E., G.G.S.; Software, J.M.V., M.E.L., D.E., P.H.S., L.E., T.M.L.; Validation, J.M.V., M.E.L.,
D.E., P.H.S., L.E., J.P.V.M.; Formal Analysis, J.M.V., M.E.L., D.E., P.H.S., L.E., S.M., A.L.C., D.M.,
S.V.; Investigation, J.M.V., M.E.L., S.M., M.B., D.F., G.G.S., L.G., Z.R.H., M.H., W.K., T.M.L.,
A.L.C., C.L. S.M., D.M., S.L., J.L., C.R., S.L.R.C., M.S., K.S., W.T., J.D.T., S.V., R.M., M.B.,
J.P.V.M., S.P., L.E., D.E., P.H.S.; Resources, J.M.V., M.E.L., D.E., P.H.S., L.E., M.B., D.F., G.G.S.,
L.G., Z.R.H., M.H., W.K., T.M.L., A.L.C., S.M., D.M., S.L., J.L., C.R., S.L.R.C., M.S., K.S., W.T.,
J.D.T., S.V., R.M., M.B.; Data Curation, J.M.V., M.E.L., S.M., M.B., D.F., G.G.S., L.G., Z.R.H., M.H.,
W.K., T.M.L., A.L.C., S.M., D.M., S.L., J.L., C.R., S.L.R.C., M.S., K.S., W.T., J.D.T., S.V., R.M.,
M.B., J.P.V.M., S.P., L.E., D.E., P.H.S.; Writing - Original Draft, J.M.V., M.E.L., D.E., P.H.S., L.E.,
S.M., J.L.; Writing - Review & Editing, J.M.V., M.E.L., S.M., M.B., D.F., G.G.S., L.G., Z.R.H., M.H.,
W.K., T.M.L., A.L.C., S.M., D.M., S.L., J.L., C.R., S.L.R.C., M.S., K.S., W.T., J.D.T., S.V., R.M.,
M.B., J.P.V.M., S.P., L.E., D.E., P.H.S.; Visualization, J.M.V., M.E.L., D.E., P.H.S., L.E., S.M.,
A.L.C.; Supervision, J.M.V., M.E.L., D.E., P.H.S., L.E.; Project Administration, J.M.V., M.E.L., D.E.,
P.H.S., L.E.; Funding Acquisition, J.M.V., M.E.L., D.E., P.H.S., L.E.

Declaration of Interests:

473 The authors declare no competing interests.

References

1. Wilkinson, G.S., and Adams, D.M. (2019). Recurrent Evolution of Extreme Longevity in Bats.
Biol. Lett. *15*, 20180860. https://doi.org/10.1098/rsbl.2018.0860.

2. Austad, S.N., and Fischer, K.E. (1991). Mammalian Aging, Metabolism, and Ecology: Evidence
from the Bats and Marsupials. J. Gerontol. *46*, B47–B53.
https://doi.org/10.1093/geronj/46.2.b47.

3. Guth, S., Mollentze, N., Renault, K., Streicker, D.G., Visher, E., Boots, M., and Brook, C.E.
(2022). Bats host the most virulent-but not the most dangerous-zoonotic viruses. Proc. Natl.
Acad. Sci. U. S. A. *119*, e2113628119. https://doi.org/10.1073/pnas.2113628119.

4. Irving, A.T., Ahn, M., Goh, G., Anderson, D.E., and Wang, L.-F. (2021). Lessons from the host
defences of bats, a unique viral reservoir. Nature *589*, 363–370.
https://doi.org/10.1038/s41586-020-03128-0.

5.   Cooper, L.N., Ansari, M.Y., Capshaw, G., Galazyuk, A., Lauer, A.M., Moss, C.F., Sears, K.E.,
Stewart, M., Teeling, E.C., Wilkinson, G.S., et al. (2024). Bats as instructive animal models for
studying longevity and aging. Ann. N. Y. Acad. Sci. *1541*, 10–23.
https://doi.org/10.1111/nyas.15233.

6.   Wilson, D.E., and Reeder, D.M. (2005). Mammal Species of the World: A Taxonomic and
Geographic Reference (JHU Press).

7.   Simmons, N.B., and Cirranello, A.L. Bat Species of the World: A Taxonomic and Geographic
Database. https://batnames.org/.

8.   Teeling, E.C., Hedges, S.B., and Kumar, S. (2009). Bats (Chiroptera). The timetree of life, 499–
503.

9.   Rietbergen, T.B., Ostende, L.W. van den H., Aase, A., Jones, M.F., Medeiros, E.D., and
Simmons, N.B. (2023). The Oldest Known Bat Skeletons and Their Implications for Eocene
Chiropteran Diversification. PLoS One *18*, e0283505.
https://doi.org/10.1371/journal.pone.0283505.

10. Kumar, S., Stecher, G., Suleski, M., and Hedges, S.B. (2017). TimeTree: A Resource for
Timelines, Timetrees, and Divergence Times. Mol. Biol. Evol. *34*, 1812–1819.
https://doi.org/10.1093/molbev/msx116.

11. Moyers Arévalo, R.L., Amador, L.I., Almeida, F.C., and Giannini, N.P. (2020). Evolution of Body
Mass in Bats: Insights from a Large Supermatrix Phylogeny. J. Mamm. Evol. *27*, 123–138.
https://doi.org/10.1007/s10914-018-9447-8.

12. Datzmann, T., von Helversen, O., and Mayer, F. (2010). Evolution of Nectarivory in
Phyllostomid Bats (Phyllostomidae Gray, 1825, Chiroptera: Mammalia). BMC Evol. Biol. *10*,
165. https://doi.org/10.1186/1471-2148-10-165.

13. Hayman, D.T.S., Bowen, R.A., Cryan, P.M., McCracken, G.F., O'Shea, T.J., Peel, A.J., Gilbert,
A., Webb, C.T., and Wood, J.L.N. (2013). Ecology of zoonotic infectious diseases in bats:
current knowledge and future directions. Zoonoses Public Health *60*, 2–21.
https://doi.org/10.1111/zph.12000.

14. Camacho, J., Bernal-Rivera, A., Peña, V., Morales-Sosa, P., Robb, S.M.C., Russell, J., Yi, K.,
Wang, Y., Tsuchiya, D., Murillo-García, O.E., et al. (2023). Sugar assimilation underlying
dietary evolution of Neotropical bats. Nat. Ecol. Evol. *8*, 1735–1750.
https://doi.org/10.1038/s41559-024-02485-7.

15. Morales, A.E., Ruedi, M., Field, K., and Carstens, B.C. (2019). Diversification rates have no
effect on the convergent evolution of foraging strategies in the most speciose genus of bats,
Myotis. Evolution *73*, 2263–2280. https://doi.org/10.1111/evo.13849.

16. Gunnell, G.F., Smith, R., and Smith, T. (2017). 33 Million Year Old Myotis (Chiroptera,
Vespertilionidae) and the Rapid Global Radiation of Modern Bats. PLoS One *12*, e0172621.
https://doi.org/10.1371/journal.pone.0172621.

17. Ruedi, M., Stadelmann, B., Gager, Y., Douzery, E.J.P., Francis, C.M., Lin, L.-K., Guillén-
Servent, A., and Cibois, A. (2013). Molecular Phylogenetic Reconstructions Identify East Asia
as the Cradle for the Evolution of the Cosmopolitan Genus Myotis (Mammalia, Chiroptera).
Mol. Phylogenet. Evol. *69*, 437–449. https://doi.org/10.1016/j.ympev.2013.08.011.

18. Tacutu, R., Craig, T., Budovsky, A., Wuttke, D., Lehmann, G., Taranukha, D., Costa, J.,
Fraifeld, V.E., and de Magalhães, J.P. (2013). Human Ageing Genomic Resources: Integrated
Databases and Tools for the Biology and Genetics of Ageing. Nucleic Acids Res. *41*, D1027–
D1033. https://doi.org/10.1093/nar/gks1155.

19. Jones, K.E., Bielby, J., Cardillo, M., Fritz, S.A., O'Dell, J., Orme, C.D.L., Safi, K., Sechrest, W.,
Boakes, E.H., Carbone, C., et al. (2009). PanTHERIA: A Species-Level Database of Life
History, Ecology, and Geography of Extant and Recently Extinct Mammals. Ecology *90*, 2648–
2648. https://doi.org/10.1890/08-1494.1.

20. Austad, S.N. (2010). Methusaleh's Zoo: How Nature Provides Us with Clues for Extending
Human Health Span. J. Comp. Pathol. *142*, S10–S21.
https://doi.org/10.1016/j.jcpa.2009.10.024.

21. Podlutsky, A.J., Khritankov, A.M., Ovodov, N.D., and Austad, S.N. (2005). A new field record
for bat longevity. J. Gerontol. A Biol. Sci. Med. Sci. *60*, 1366–1368.
https://doi.org/10.1093/gerona/60.11.1366.

22. Wilson, D.E., and Tyson, E.L. (1970). Longevity Records for Artibeus Jamaicensis and Myotis
Nigricans. J. Mammal. *51*, 203. https://doi.org/10.2307/1378570.

23. Stadelmann, B., Lin, L.-K., Kunz, T.H., and Ruedi, M. (2007). Molecular Phylogeny of New
World \emphMyotis (Chiroptera, Vespertilionidae) Inferred from Mitochondrial and Nuclear DNA
Genes. Mol. Phylogenet. Evol. *43*, 32–48. https://doi.org/10.1016/j.ympev.2006.06.019.

24. Agnarsson, I., Zambrana-Torrelio, C.M., Flores-Saldana, N.P., and May-Collado, L.J. (2011). A
Time-Calibrated Species-Level Phylogeny of Bats (Chiroptera, Mammalia). PLoS Curr. *3*,
RRN1212. https://doi.org/10.1371/currents.RRN1212.

25. Armero, A., Li, R., Bienes, K.M., Chen, X., Li, J., Xu, S., Chen, Y., Hughes, A.C., Berthet, N.,
and Wong, G. (2022). Myotis Fimbriatus Virome, a Window to Virus Diversity and Evolution in
the Genus Myotis. Viruses *14*, 1899. https://doi.org/10.3390/v14091899.

26. He, X., Korytář, T., Zhu, Y., Pikula, J., Bandouchova, H., Zukal, J., and Köllner, B. (2014).
Establishment of Myotis Myotis Cell Lines - Model for Investigation of Host-Pathogen
Interaction in a Natural Host for Emerging Viruses. PLoS One *9*, e109795.
https://doi.org/10.1371/journal.pone.0109795.

27. Seltmann, A., Troxell, S.A., Schad, J., Fritze, M., Bailey, L.D., Voigt, C.C., and Czirják, G.Á.
(2022). Differences in Acute Phase Response to Bacterial, Fungal and Viral Antigens in
Greater Mouse-Eared Bats (Myotis Myotis). Sci. Rep. *12*, 15259.
https://doi.org/10.1038/s41598-022-18240-6.

28. Hayward, J.A., Tachedjian, M., Johnson, A., Irving, A.T., Gordon, T.B., Cui, J., Nicolas, A.,
Smith, I., Boyd, V., Marsh, G.A., et al. (2022). Unique Evolution of Antiviral Tetherin in Bats. J.
Virol. *96*, e0115222. https://doi.org/10.1128/jvi.01152-22.

29. Fernandes, A.P., Águeda-Pinto, A., Pinheiro, A., Rebelo, H., and Esteves, P.J. (2022).
Evolution of TRIM5 and TRIM22 in Bats Reveals a Complex Duplication Process. Viruses *14*.
https://doi.org/10.3390/v14020345.

30. Jacquet, S., Culbertson, M., Zhang, C., El Filali, A., De La Myre Mory, C., Pons, J.-B., Filippi-
Codaccioni, O., Lauterbur, M.E., Ngoubangoye, B., Duhayer, J., et al. (2022). Adaptive
duplication and genetic diversification of protein kinase R contribute to the specificity of bat-

virus interactions. Sci Adv *8*, eadd7540. https://doi.org/10.1126/sciadv.add7540.

31. Jacquet, S., Pontier, D., and Etienne, L. (2020). Rapid Evolution of HERC6 and Duplication of
a Chimeric HERC5/6 Gene in Rodents and Bats Suggest an Overlooked Role of HERCs in
Mammalian Immunity. Front. Immunol. *11*, 605270.
https://doi.org/10.3389/fimmu.2020.605270.

32. Chomel, B., Stuckey, M., Boulouis, H., and Aguilar Setién, Á. (2014). Bat-Related Zoonoses.
Zoonoses - Infections Affecting Humans and Animals, 697–714. https://doi.org/10.1007/978-
94-017-9457-2_28.

33. Williams, E.P., Spruill-Harrell, B.M., Taylor, M.K., Lee, J., Nywening, A.V., Yang, Z., Nichols,
J.H., Camp, J.V., Owen, R.D., and Jonsson, C.B. (2021). Common themes in zoonotic spillover
and disease emergence: Lessons learned from bat- and rodent-borne RNA viruses. Viruses
13, 1509. https://doi.org/10.3390/v13081509.

34. Mollentze, N., and Streicker, D.G. (2020). Viral zoonotic risk is homogenous among taxonomic
orders of mammalian and avian reservoir hosts. Proc. Natl. Acad. Sci. U. S. A. *117*, 9423–
9430. https://doi.org/10.1073/pnas.1919176117.

35. Kolora, S.R.R., Owens, G.L., Vazquez, J.M., Stubbs, A., Chatla, K., Jainese, C., Seeto, K.,
McCrea, M., Sandel, M.W., Vianna, J.A., et al. (2021). Origins and Evolution of Extreme Life
Span in Pacific Ocean Rockfishes. Science *374*, 842–847.
https://doi.org/10.1126/science.abg5332.

36. Sulak, M., Fong, L., Mika, K., Chigurupati, S., Yon, L., Mongan, N.P., Emes, R.D., and Lynch,
V.J. (2016). Correction: TP53 copy number expansion is associated with the evolution of
increased body size and an enhanced DNA damage response in elephants. Elife *5*.
https://doi.org/10.7554/eLife.24307.

37. Vazquez, J.M., Sulak, M., Chigurupati, S., and Lynch, V.J. (2018). A Zombie LIF Gene in
Elephants Is Upregulated by TP53 to Induce Apoptosis in Response to DNA Damage. Cell
Rep. *24*, 1765–1776. https://doi.org/10.1016/j.celrep.2018.07.042.

38. Vazquez, J.M., and Lynch, V.J. (2021). Pervasive duplication of tumor suppressors in
Afrotherians during the evolution of large bodies and reduced cancer risk. Elife *10*, e65041.
https://doi.org/10.7554/eLife.65041.

39. Davies, K.T.J., Tsagkogeorga, G., Bennett, N.C., Dávalos, L.M., Faulkes, C.G., and Rossiter,
S.J. (2014). Molecular Evolution of Growth Hormone and Insulin-like Growth Factor 1
Receptors in Long-Lived, Small-Bodied Mammals. Gene *549*, 228–236.
https://doi.org/10.1016/j.gene.2014.07.061.

40. Vazquez, J.M., Kraft, M., and Lynch, V.J. (2022). A CDKN2C retroduplication in Bowhead
whales is associated with the evolution of extremely long lifespans and alerted cell cycle
dynamics. bioRxiv, 2022.09.07.506958. https://doi.org/10.1101/2022.09.07.506958.

41. Foote, A.D., Liu, Y., Thomas, G.W.C., Vinař, T., Alföldi, J., Deng, J., Dugan, S., van Elk, C.E.,
Hunter, M.E., Joshi, V., et al. (2015). Convergent evolution of the genomes of marine
mammals. Nat. Genet. *47*, 272–275. https://doi.org/10.1038/ng.3198.

42. Huang, Z., Jiang, C., Gu, J., Uvizl, M., Power, S., Douglas, D., and Kacprzyk, J. (2023).
Duplications of Human Longevity-Associated Genes Across Placental Mammals. Genome Biol.
Evol. *15*. https://doi.org/10.1093/gbe/evad186.

43. Li, S., Vazquez, J.M., and Sudmant, P.H. (2023). The evolution of aging and lifespan. Trends
Genet. *39*, 830–843. https://doi.org/10.1016/j.tig.2023.08.005.

44. Jebb, D., Huang, Z., Pippel, M., Hughes, G.M., Lavrichenko, K., Devanna, P., Winkler, S.,
Jermiin, L.S., Skirmuntt, E.C., Katzourakis, A., et al. (2020). Six reference-quality genomes
reveal evolution of bat adaptations. Nature *583*, 578–584. https://doi.org/10.1038/s41586-020-
2486-3.

45. Glaberman, S., Bulls, S.E., Vazquez, J.M., Chiari, Y., and Lynch, V.J. (2021). Concurrent
evolution of antiaging gene duplications and cellular phenotypes in long-lived turtles. Genome
Biol. Evol. *13*, evab244. https://doi.org/10.1093/gbe/evab244.

46. MacRae, S.L., Croken, M.M., Calder, R.B., Aliper, A., Milholland, B., White, R.R., Zhavoronkov,
A., Gladyshev, V.N., Seluanov, A., Gorbunova, V., et al. (2015). DNA Repair in Species with
Extreme Lifespan Differences. Aging *7*, 1171–1184. https://doi.org/10.18632/aging.100866.

47. Baines, C., Meitern, R., Kreitsberg, R., and Sepp, T. (2022). Comparative study of the evolution
of cancer gene duplications across fish. Evol. Appl. *15*, 1834–1845.
https://doi.org/10.1111/eva.13481.

48. López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., and Kroemer, G. (2013). The
Hallmarks of Aging. Cell *153*, 1194–1217. https://doi.org/10.1016/j.cell.2013.05.039.

49. Moreno Santillán, D.D., Lama, T.M., Gutierrez Guerrero, Y.T., Brown, A.M., Donat, P., Zhao,
H., Rossiter, S.J., Yohe, L.R., Potter, J.H., Teeling, E.C., et al. (2021). Large-scale genome
sampling reveals unique immunity and metabolic adaptations in bats. Mol. Ecol. *30*, 6449–
6467. https://doi.org/10.1111/mec.16027.

50. Enard, D., and Petrov, D.A. (2020). Ancient RNA virus epidemics through the lens of recent
adaptation in human genomes. Philos. Trans. R. Soc. Lond. B Biol. Sci. *375*, 20190575.
https://doi.org/10.1098/rstb.2019.0575.

51. Sotero-Caio, C.G., Baker, R.J., and Volleth, M. (2017). Chromosomal evolution in Chiroptera.
Genes (Basel) *8*. https://doi.org/10.3390/genes8100272.

52. Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V., and Zdobnov, E.M. (2015).
BUSCO: assessing genome assembly and annotation completeness with single-copy
orthologs. Bioinformatics *31*, 3210–3212. https://doi.org/10.1093/bioinformatics/btv351.

53. Manni, M., Berkeley, M.R., Seppey, M., and Zdobnov, E.M. (2021). BUSCO: assessing
genomic data quality and beyond. Current Protocols *1*.

54. Amador, L.I., Moyers Arévalo, R.L., Almeida, F.C., Catalano, S.A., and Giannini, N.P. (2018).
Bat Systematics in the Light of Unconstrained Analyses of a Comprehensive Molecular
Supermatrix. J. Mamm. Evol. *25*, 37–70. https://doi.org/10.1007/s10914-016-9363-8.

55. Upham, N.S., Esselstyn, J.A., and Jetz, W. (2019). Inferring the mammal tree: Species-level
sets of phylogenies for questions in ecology, evolution, and conservation. PLoS Biol. *17*,
e3000494. https://doi.org/10.1371/journal.pbio.3000494.

56. Korstian, J.M., Paulat, N.S., Platt, R.N., Stevens, R.D., and Ray, D.A. (2022). SINE-Based
Phylogenomics Reveal Extensive Introgression and Incomplete Lineage Sorting in Myotis.
Genes *13*, 399. https://doi.org/10.3390/genes13030399.

57. Foley, N.M., Harris, A.J., Bredemeyer, K.R., Ruedi, M., Puechmaille, S.J., Teeling, E.C.,
Criscitiello, M.F., and Murphy, W.J. (2024). Karyotypic stasis and swarming influenced the
evolution of viral tolerance in a species-rich bat radiation. Cell Genom *4*, 100482.
https://doi.org/10.1016/j.xgen.2023.100482.

58. Foley, N.M., Mason, V.C., Harris, A.J., Bredemeyer, K.R., Damas, J., Lewin, H.A., Eizirik, E.,
Gatesy, J., Karlsson, E.K., Lindblad-Toh, K., et al. (2023). A genomic timescale for placental
mammal evolution. Science *380*, eabl8189. https://doi.org/10.1126/science.abl8189.

59. Volleth, M. (2012). Variations on a theme: Karyotype comparison in Eurasian Myotis species
and implications for phylogeny. Vespertilio *16*, 329–350.

60. McBee, K. (1986). Standard karyology of nine species of vespertilionid bats (Chiroptera:
Vespertilionidae) from Thailand. Annals of Carnegie Museum *55*, 95–116.

61. Bickham, J.W., McBee, K., and Schlitter, D.A. (1986). Chromosomal variation among seven
species of Myotis (Chiroptera: Vespertilionidae). J. Mammal. *67*, 746–750.
https://doi.org/10.2307/1381139.

62. Zhang, W.D. (1984). A study on karyotype of Myotis chinensis and M. laniger Peter. J Anhui
Normal Univ *7*, 42–47.

63. Zima, J. (1985). Synopsis of karyotypes of vespertilionid bats (Mammalia: Chiroptera). Acta
Univ. Carol. Biol *1981*, 311–329.

64. O'Brien, S.J., Menninger, J.C., and Nash, W.G. (2006). Atlas of Mammalian Chromosomes
(John Wiley & Sons).

65. Goel, M., Sun, H., Jiao, W.-B., and Schneeberger, K. (2019). SyRI: finding genomic
rearrangements and local sequence differences from whole-genome assemblies. Genome Biol.
20, 277. https://doi.org/10.1186/s13059-019-1911-0.

66. de Sotero-Caio, C.G., Cabral-de-Mello, D.C., Calixto, M. da S., Valente, G.T., Martins, C.,
Loreto, V., de Souza, M.J., and Santos, N. (2017). Centromeric enrichment of LINE-1
retrotransposons and its significance for the chromosome evolution of Phyllostomid bats.
Chromosome Res. *25*, 313–325. https://doi.org/10.1007/s10577-017-9565-9.

67. Morel, B., Kozlov, A.M., Stamatakis, A., and Szöllősi, G.J. (2020). GeneRax: A Tool for
Species-Tree-Aware Maximum Likelihood-Based Gene Family Tree Inference under Gene
Duplication, Transfer, and Loss. Mol. Biol. Evol. *37*, 2763–2774.
https://doi.org/10.1093/molbev/msaa141.

68. Kaufman, R.J. (1999). Double-stranded RNA-activated protein kinase mediates virus-induced
apoptosis: a new role for an old actor. Proc. Natl. Acad. Sci. U. S. A. *96*, 11693–11695.
https://doi.org/10.1073/pnas.96.21.11693.

69. García, M.A., Gil, J., Ventoso, I., Guerra, S., Domingo, E., Rivas, C., and Esteban, M. (2006).
Impact of protein kinase PKR in cell biology: from antiviral to antiproliferative action. Microbiol.
Mol. Biol. Rev. *70*, 1032–1060. https://doi.org/10.1128/MMBR.00027-06.

70. Ostertag, D., Hoblitzell-Ostertag, T.M., and Perrault, J. (2007). Overproduction of double-
stranded RNA in vesicular stomatitis virus-infected cells activates a constitutive cell-type-
specific antiviral response. J. Virol. *81*, 503–513. https://doi.org/10.1128/JVI.01218-06.

71. Enard, D., Cai, L., Gwennap, C., and Petrov, D.A. (2016). Viruses are a dominant driver of
protein adaptation in mammals. Elife *5*. https://doi.org/10.7554/eLife.12469.

72. Murrell, B., Weaver, S., Smith, M.D., Wertheim, J.O., Murrell, S., Aylward, A., Eren, K., Pollner,
T., Martin, D.P., Smith, D.M., et al. (2015). Gene-wide identification of episodic selection. Mol.
Biol. Evol. *32*, 1365–1371. https://doi.org/10.1093/molbev/msv035.

73. Souilmi, Y., Lauterbur, M.E., Tobler, R., Huber, C.D., Johar, A.S., Moradi, S.V., Johnston,
W.A., Krogan, N.J., Alexandrov, K., and Enard, D. (2021). An ancient viral epidemic involving
host coronavirus interacting genes more than 20,000 years ago in East Asia. Curr. Biol. *31*,
3704. https://doi.org/10.1016/j.cub.2021.07.052.

74. Brook, C.E., and Dobson, A.P. (2015). Bats as "special" reservoirs for emerging zoonotic
pathogens. Trends Microbiol. *23*, 172–180. https://doi.org/10.1016/j.tim.2014.12.004.

75. Kondrashov, F.A. (2011). Gene Dosage and Duplication. In Evolution after Gene Duplication,
K. Dittmar and D. Liberles, eds. (John Wiley & Sons), pp. 57–76.

76. Kondrashov, F.A., Rogozin, I.B., Wolf, Y.I., and Koonin, E.V. (2002). Selection in the evolution
of gene duplications. Genome Biol. *3*, RESEARCH0008. https://doi.org/10.1186/gb-2002-3-2-
research0008.

77. Rastogi, S., and Liberles, D.A. (2005). Subfunctionalization of duplicated genes as a transition
state to neofunctionalization. BMC Evol. Biol. *5*, 28. https://doi.org/10.1186/1471-2148-5-28.

78. Assis, R., and Bachtrog, D. (2013). Neofunctionalization of young duplicate genes in
Drosophila. Proc. Natl. Acad. Sci. U. S. A. *110*, 17409–17414.
https://doi.org/10.1073/pnas.1313759110.

79. Enard, D., and Petrov, D.A. (2018). Evidence that RNA Viruses Drove Adaptive Introgression
between Neanderthals and Modern Humans. Cell *175*, 360–371.e13.
https://doi.org/10.1016/j.cell.2018.08.034.

80. Brunet-Rossinni, A.K., and Austad, S.N. (2004). Ageing studies on bats: a review.
Biogerontology *5*, 211–222. https://doi.org/10.1023/B:BGEN.0000038022.65024.d8.

81. Wilkinson, G.S., and South, J.M. (2002). Life History, Ecology and Longevity in Bats. Aging
Cell *1*, 124–131. https://doi.org/10.1046/j.1474-9728.2002.00020.x.

82. Pyenson, N.D., and Sponberg, S.N. (2011). Reconstructing body size in extinct crown Cetacea
(neoceti) using allometry, phylogenetic methods and tests from the fossil record. J. Mamm.
Evol. *18*, 269–288. https://doi.org/10.1007/s10914-011-9170-1.

83. Montgomery, S.H., Geisler, J.H., McGowen, M.R., Fox, C., Marino, L., and Gatesy, J. (2013).
The evolutionary history of cetacean brain and body size: Cetacean brain evolution. Evolution
67, 3339–3353. https://doi.org/10.1111/evo.12197.

84. Puttick, M.N., and Thomas, G.H. (2015). Fossils and living taxa agree on patterns of body
mass evolution: a case study with Afrotheria. Proc. Biol. Sci. *282*, 20152023.
https://doi.org/10.1098/rspb.2015.2023.

85. Sulak, M., Fong, L., Mika, K., Chigurupati, S., Yon, L., Mongan, N.P., Emes, R.D., and Lynch,
V.J. (2016). TP53 copy number expansion is associated with the evolution of increased body
size and an enhanced DNA damage response in elephants. Elife *5*.

731  https://doi.org/10.7554/elife.11994.

86. Nunney, L. (2016). Commentary: The multistage model of carcinogenesis, Peto's paradox and evolution. Int. J. Epidemiol. *45*, 649–653. https://doi.org/10.1093/ije/dyv201.

87. Nunney, L. (1999). Lineage selection and the evolution of multistage carcinogenesis. Proc. Biol. Sci. *266*, 493–498. https://doi.org/10.1098/rspb.1999.0664.

88. Armitage, P. (1985). Multistage Models of Carcinogenesis. Environ. Health Perspect. *63*, 195–201. https://doi.org/10.1289/ehp.8563195.

89. Armitage, P., and Doll, R. (2004). The Age Distribution of Cancer and a Multi-Stage Theory of Carcinogenesis. Br. J. Cancer *91*, 6602297. https://doi.org/10.1038/sj.bjc.6602297.

90. Peto, R., Roe, F.J., Lee, P.N., Levy, L., and Clack, J. (1975). Cancer and ageing in mice and men. Br. J. Cancer *32*, 411–426. https://doi.org/10.1038/bjc.1975.242.

91. Peto, R. (2015). Quantitative Implications of the Approximate Irrelevance of Mammalian Body Size and Lifespan to Lifelong Cancer Risk. Philos. Trans. R. Soc. Lond. B Biol. Sci. *370*, 20150198. https://doi.org/10.1098/rstb.2015.0198.

92. Siegel, R.L., Miller, K.D., and Jemal, A. (2018). Cancer Statistics, 2018. CA Cancer J. Clin. *68*, 7–30. https://doi.org/10.3322/caac.21442.

93. Garner, W.B., Smith, B.D., Ludmir, E.B., Wakefield, D.V., Shabason, J., Williams, G.R., Martin, M.Y., Wang, Y., Ballo, M.T., and VanderWalde, N.A. (2023). Predicting Future Cancer Incidence by Age, Race, Ethnicity, and Sex. J. Geriatr. Oncol. *14*, 101393. https://doi.org/10.1016/j.jgo.2022.10.008.

94. Nunney, L. (2024). The effect of body size and inbreeding on cancer mortality in breeds of the domestic dog: a test of the multi-stage model of carcinogenesis. R. Soc. Open Sci. *11*, 231356. https://doi.org/10.1098/rsos.231356.

95. Kent, M.S., Karchemskiy, S., Culp, W.T.N., Lejeune, A.T., Pesavento, P.A., Toedebusch, C., Brady, R., and Rebhun, R. (2022). Longevity and mortality in cats: A single institution necropsy study of 3108 cases (1989-2019). PLoS One *17*, e0278199. https://doi.org/10.1371/journal.pone.0278199.

96. Compton, Z.T., Harris, V., Mellon, W., Rupp, S., Mallo, D., Kapsetaki, S.E., Wilmot, M., Kennington, R., Noble, K., Baciu, C., et al. (2023). Cancer prevalence across vertebrates. Res. Sq. https://doi.org/10.21203/rs.3.rs-3117313/v1.

97. Vincze, O., Colchero, F., Lemaître, J.-F., Conde, D.A., Pavard, S., Bieuville, M., Urrutia, A.O., Ujvari, B., Boddy, A.M., Maley, C.C., et al. (2022). Cancer risk across mammals. Nature *601*, 263–267. https://doi.org/10.1038/s41586-021-04224-5.

98. Nunney, L. (2018). Size Matters: Height, Cell Number and a Person's Risk of Cancer. Proc. R. Soc. B *285*, 20181743. https://doi.org/10.1098/rspb.2018.1743.

99. Caulin, A.F., and Maley, C.C. (2011). Peto's Paradox: evolution's prescription for cancer prevention. Trends Ecol. Evol. *26*, 175–182. https://doi.org/10.1016/j.tree.2011.01.002.

100. Smith, M.D., Wertheim, J.O., Weaver, S., Murrell, B., Scheffler, K., and Kosakovsky Pond, S.L. (2015). Less is more: an adaptive branch-site random effects model for efficient detection of

episodic diversifying selection. Mol. Biol. Evol. *32*, 1342–1353.
https://doi.org/10.1093/molbev/msv022.

101.Pei, G., Balkema-Buschmann, A., and Dorhoi, A. (2024). Disease tolerance as immune
defense strategy in bats: One size fits all? PLoS Pathog. *20*, e1012471.
https://doi.org/10.1371/journal.ppat.1012471.

102.Seim, I., Fang, X., Xiong, Z., Lobanov, A.V., Huang, Z., Ma, S., Feng, Y., Turanov, A.A., Zhu,
Y., Lenz, T.L., et al. (2013). Genome Analysis Reveals Insights into Physiology and Longevity
of the Brandt's Bat Myotis Brandtii. Nat. Commun. *4*, 2212.
https://doi.org/10.1038/ncomms3212.

103.Matsuda, Y., and Makino, T. (2024). Comparative genomics reveals convergent signals
associated with the high metabolism and longevity in birds and bats. Proc. Biol. Sci. *291*,
20241068. https://doi.org/10.1098/rspb.2024.1068.

104.Blumer, M., Brown, T., Freitas, M.B., Destro, A.L., Oliveira, J.A., Morales, A.E., Schell, T.,
Greve, C., Pippel, M., Jebb, D., et al. (2022). Gene losses in the common vampire bat
illuminate molecular adaptations to blood feeding. Sci. Adv. *8*, eabm6494.
https://doi.org/10.1126/sciadv.abm6494.

105.Morales, A.E., Burbrink, F.T., Segall, M., Meza, M., Munegowda, C., Webala, P.W., Patterson,
B.D., Thong, V.D., Ruedi, M., Hiller, M., et al. (2024). Distinct genes with similar functions
underlie convergent evolution in Myotis bat ecomorphs. Mol. Biol. Evol. *41*.
https://doi.org/10.1093/molbev/msae165.

106.Hanahan, D., and Weinberg, R.A. (2011). Hallmarks of Cancer: The Next Generation. Cell *144*,
646–674. https://doi.org/10.1016/j.cell.2011.02.013.

107.Hanahan, D., and Weinberg, R.A. (2000). The Hallmarks of Cancer. Cell *100*, 57–70.
https://doi.org/10.1016/S0092-8674(00)81683-9.

108.Hanahan, D. (2022). Hallmarks of Cancer: New Dimensions. Cancer Discov. *12*, 31–46.
https://doi.org/10.1158/2159-8290.CD-21-1059.

109.Milacic, M., Beavers, D., Conley, P., Gong, C., Gillespie, M., Griss, J., Haw, R., Jassal, B.,
Matthews, L., May, B., et al. (2024). The reactome pathway knowledgebase 2024. Nucleic
Acids Res. *52*, D672–D678. https://doi.org/10.1093/nar/gkad1025.

110.Ohtsuki, K., and Ono, Y. (1997). The biochemical mechanisms involved in the biological effects
induced by neocarzinostatin (NCS) and the NCS chromophore. In Neocarzinostatin (Springer
Japan), pp. 129–154. https://doi.org/10.1007/978-4-431-66914-2_7.

111.Bañuelos, A., Reyes, E., Ocadiz, R., Alvarez, E., Moreno, M., Monroy, A., and Gariglio, P.
(2003). Neocarzinostatin induces an effective p53-dependent response in human
papillomavirus-positive cervical cancer cells. J. Pharmacol. Exp. Ther. *306*, 671–680.
https://doi.org/10.1124/jpet.103.051557.

112.Ishida, N., Miyazaki, K., Kumagai, K., and Rikimaru, M. (1965). Neocarzinostatin, an Antitumor
Antibiotic of High Molecular Weight Isolation, Physicochemical Properties and Biological
Activities. The Journal of Antibiotics, Series A *18*, 68–76.
https://doi.org/10.11554/antibioticsa.18.2_68.

113.Abegglen, L.M., Caulin, A.F., Chan, A., Lee, K., Robinson, R., Campbell, M.S., Kiso, W.K.,

Schmitt, D.L., Waddell, P.J., Bhaskara, S., et al. (2015). Potential Mechanisms for Cancer
Resistance in Elephants and Comparative Cellular Response to DNA Damage in Humans.
JAMA *314*, 1850–1860. https://doi.org/10.1001/jama.2015.13134.

114. Firsanov, D., Zacher, M., Tian, X., Zhao, Y., George, J.C., Sformo, T.L., Tombline, G., Biashad,
S.A., Gilman, A., Hamilton, N., et al. (2023). DNA Repair and Anti-Cancer Mechanisms in the
Longest-Living Mammal: The Bowhead Whale. bioRxiv, 2023.05.07.539748.
https://doi.org/10.1101/2023.05.07.539748.

115. Lagunas-Rangel, F.A. (2020). Why do bats live so long?—Possible molecular mechanisms.
Biogerontology *21*, 1–11. https://doi.org/10.1007/s10522-019-09840-3.

116. Chionh, Y.T., Cui, J., Koh, J., Mendenhall, I.H., Ng, J.H.J., Low, D., Itahana, K., Irving, A.T.,
and Wang, L.-F. (2019). High basal heat-shock protein expression in bats confers resistance to
cellular heat/oxidative stress. Cell Stress Chaperones *24*, 835–849.
https://doi.org/10.1007/s12192-019-01013-y.

117. Hua, R., Ma, Y.-S., Yang, L., Hao, J.-J., Hua, Q.-Y., Shi, L.-Y., Yao, X.-Q., Zhi, H.-Y., and Liu,
Z. (2024). Experimental evidence for cancer resistance in a bat species. Nat. Commun. *15*,
1401. https://doi.org/10.1038/s41467-024-45767-1.

118. Peixoto, F.P., Braga, P.H.P., and Mendes, P. (2018). A synthesis of ecological and
evolutionary determinants of bat diversity across spatial scales. BMC Ecol. *18*, 18.
https://doi.org/10.1186/s12898-018-0174-z.

119. USGS (2022). North American Bat Ranges.
http://dds.cr.usgs.gov/pub/data/nationalatlas/bat000p010g_nt00373.tar.gz
http://dds.cr.usgs.gov/pub/data/nationalatlas/bat000p010g_nt00373.tar.gz.

120. Rhie, A., McCarthy, S.A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M.,
Chow, W., Fungtammasan, A., Kim, J., et al. (2021). Towards complete and error-free genome
assemblies of all vertebrate species. Nature *592*, 737–746. https://doi.org/10.1038/s41586-
021-03451-0.

121. The Darwin Tree of Life Project Consortium (2022). Sequence Locally, Think Globally: The
Darwin Tree of Life Project. Proceedings of the National Academy of Sciences *119*,
e2115642118. https://doi.org/10.1073/pnas.2115642118.

122. Lewin, H.A., Richards, S., Lieberman Aiden, E., Allende, M.L., Archibald, J.M., Bálint, M.,
Barker, K.B., Baumgartner, B., Belov, K., Bertorelle, G., et al. (2022). The Earth BioGenome
Project 2020: Starting the Clock. Proceedings of the National Academy of Sciences *119*,
e2115635118. https://doi.org/10.1073/pnas.2115635118.

123. Slater, G.J., Goldbogen, J.A., and Pyenson, N.D. (2017). Independent evolution of baleen
whale gigantism linked to Plio-Pleistocene ocean dynamics. Proc. Biol. Sci. *284*.
https://doi.org/10.1098/rspb.2017.0546.

124. Tollis, M., Robbins, J., Webb, A., Kuderna, L.F.K., Caulin, A.F., Garcia, J.D., Bérubé, M.,
Pourmand, N., Marquès-Bonet, T., O'Connell, M., et al. (2019). Return to the sea, get huge,
beat cancer: An analysis of cetacean genomes including an assembly for the humpback whale
(Megaptera novaeangliae). Mol. Biol. Evol. *36*, 1746–1763.
https://doi.org/10.1093/molbev/msz099.

125. Nair, N.U., Cheng, K., Naddaf, L., Sharon, E., Pal, L.R., Rajagopal, P.S., Unterman, I., Aldape,

K., Hannenhalli, S., Day, C.-P., et al. (2022). Cross-species identification of cancer resistance-associated genes that may mediate human cancer risk. Sci. Adv. *8*, eabj7176. https://doi.org/10.1126/sciadv.abj7176.

126. Caulin, A.F., Graham, T.A., Wang, L.-S., and Maley, C.C. (2015). Solutions to Peto's paradox revealed by mathematical modelling and cross-species cancer gene analysis. Philos. Trans. R. Soc. Lond. B Biol. Sci. *370*, 20140222. https://doi.org/10.1098/rstb.2014.0222.

127. Vazquez, J.M., Pena, M.T., Muhammad, B., Kraft, M., Adams, L.B., and Lynch, V.J. (2022). Parallel evolution of reduced cancer risk and tumor suppressor duplications in Xenarthra. bioRxiv, e82558. https://doi.org/10.1101/2022.08.04.502824.

128. Beichman, A.C., Zhu, L., and Harris, K. (2024). The evolutionary interplay of somatic and germline mutation rates. Annu. Rev. Biomed. Data Sci. *7*, 83–105. https://doi.org/10.1146/annurev-biodatasci-102523-104225.

129. Cagan, A., Baez-Ortega, A., Brzozowska, N., Abascal, F., Coorens, T.H.H., Sanders, M.A., Lawson, A.R.J., Harvey, L.M.R., Bhosle, S., Jones, D., et al. (2022). Somatic mutation rates scale with lifespan across mammals. Nature *604*, 517–524. https://doi.org/10.1038/s41586-022-04618-z.

130. Bergeron, L.A., Besenbacher, S., Zheng, J., Li, P., Bertelsen, M.F., Quintard, B., Hoffman, J.I., Li, Z., St Leger, J., Shao, C., et al. (2023). Evolution of the germline mutation rate across vertebrates. Nature *615*, 285–291. https://doi.org/10.1038/s41586-023-05752-y.

131. Lollar, A. (2018). The Rehabilitation and Captive Care of Insectivorous Bats. Weatherford International: Houston, TX, USA.

132. Morales, A.E., Dong, Y., Brown, T., Baid, K., Kontopoulos, D.-G., Gonzalez, V., Huang, Z., Ahmed, A.-W., Bhuinya, A., Hilgers, L., et al. (2025). Bat genomes illuminate adaptations to viral tolerance and disease resistance. Nature *638*, 449–458. https://doi.org/10.1038/s41586-024-08471-0.

133. O'Shea, T.J., Cryan, P.M., Cunningham, A.A., Fooks, A.R., Hayman, D.T.S., Luis, A.D., Peel, A.J., Plowright, R.K., and Wood, J.L.N. (2014). Bat flight and zoonotic viruses. Emerg. Infect. Dis. *20*, 741–745. https://doi.org/10.3201/eid2005.130539.

134. Levesque, D.L., Boyles, J.G., Downs, C.J., and Breit, A.M. (2021). High Body Temperature is an Unlikely Cause of High Viral Tolerance in Bats. J. Wildl. Dis. *57*, 238–241. https://doi.org/10.7589/JWD-D-20-00079.

135. Kirkwood, T.B.L. (2017). The disposable soma theory. The evolution of senescence in the tree of life, 23–39.

136. Toshkova, N., Zhelyzkova, V., Reyes-Ruiz, A., Haerens, E., de Castro Deus, M., Lacombe, R.V., Lecerf, M., Gonzalez, G., Jouvenet, N., Planchais, C., et al. (2024). Temperature sensitivity of bat antibodies links metabolic state of bats with antigen-recognition diversity. Nat. Commun. *15*, 5878. https://doi.org/10.1038/s41467-024-50316-x.

137. Gutiérrez-Guerrero, Y.T., Ibarra-Laclette, E., Martínez Del Río, C., Barrera-Redondo, J., Rebollar, E.A., Ortega, J., León-Paniagua, L., Urrutia, A., Aguirre-Planter, E., and Eguiarte, L.E. (2020). Genomic consequences of dietary diversification and parallel evolution due to nectarivory in leaf-nosed bats. Gigascience *9*. https://doi.org/10.1093/gigascience/giaa059.

138.Camacho, J., Bernal-Rivera, A., Peña, V., Morales-Sosa, P., Robb, S., Russell, J., Yi, K.,
Wang, Y., Tsuchiya, D., Murillo-García, O.E., et al. (2023). Sugar assimilation underlying
dietary evolution of Neotropical bats. bioRxiv, 2023.07.02.547432.
https://doi.org/10.1101/2023.07.02.547432.

139.Mandl, J.N., Schneider, C., Schneider, D.S., and Baker, M.L. (2018). Going to bat(s) for studies
of disease tolerance. Front. Immunol. 9, 2112. https://doi.org/10.3389/fimmu.2018.02112.

140.Ray, D.A., Feschotte, C., Pagan, H.J.T., Smith, J.D., Pritham, E.J., Arensburger, P., Atkinson,
P.W., and Craig, N.L. (2008). Multiple Waves of Recent DNA Transposon Activity in the Bat,
Myotis Lucifugus. Genome Res. 18, 717–728. https://doi.org/10.1101/gr.071886.107.

141.Thomas, J., Phillips, C.D., Baker, R.J., and Pritham, E.J. (2014). Rolling-circle transposons
catalyze genomic innovation in a mammalian lineage. Genome Biol. Evol. 6, 2595–2610.
https://doi.org/10.1093/gbe/evu204.

142.Yashin, A.I., Wu, D., Arbeev, K., Yashkin, A.P., Akushevich, I., Bagley, O., Duan, M., and
Ukraintseva, S. (2021). Roles of interacting stress-related genes in lifespan regulation: insights
for translating experimental findings to humans. J. Transl. Genet. Genom. 5, 357–379.
https://doi.org/10.20517/jtgg.2021.26.

143.Wu, L., Zhu, X., Liu, Y., Zhao, D., Yu, B.C., Wei, Z., Lin, X., and Qi, L.S. (2025). Identification of
replicative aging and inflammatory aging signatures via whole-genome CRISPRi screens.
Genome Biol. 26, 233. https://doi.org/10.1186/s13059-025-03683-7.

144.Bullido, M.J., Martínez-García, A., Tenorio, R., Sastre, I., Muñoz, D.G., Frank, A., and
Valdivieso, F. (2008). Double stranded RNA activated EIF2 alpha kinase (EIF2AK2; PKR) is
associated with Alzheimer's disease. Neurobiol. Aging 29, 1160–1166.
https://doi.org/10.1016/j.neurobiolaging.2007.02.023.

145.Vermillion, K.L., Anderson, K.J., Hampton, M., and Andrews, M.T. (2015). Gene expression
changes controlling distinct adaptations in the heart and skeletal muscle of a hibernating
mammal. Physiol. Genomics 47, 58–74. https://doi.org/10.1152/physiolgenomics.00108.2014.

146.Lam, B., Kajderowicz, K.M., Keys, H.R., Roessler, J.M., Frenkel, E.M., Kirkland, A., Bisht, P.,
El-Brolosy, M.A., Jaenisch, R., Bell, G.W., et al. (2024). Multi-species genome-wide CRISPR
screens identify GPX4 as a conserved suppressor of cold-induced cell death. bioRxivorg,
2024.07.25.605098. https://doi.org/10.1101/2024.07.25.605098.

147.Sone, M., and Yamaguchi, Y. (2024). Cold resistance of mammalian hibernators ~ a matter of
ferroptosis? Front. Physiol. 15, 1377986. https://doi.org/10.3389/fphys.2024.1377986.

148.Kaelber, J.T., Demogines, A., Harbison, C.E., Allison, A.B., Goodman, L.B., Ortega, A.N.,
Sawyer, S.L., and Parrish, C.R. (2012). Evolutionary reconstructions of the transferrin receptor
of Caniforms supports canine parvovirus being a re-emerged and not a novel pathogen in
dogs. PLoS Pathog. 8, e1002666. https://doi.org/10.1371/journal.ppat.1002666.

149.Demogines, A., Abraham, J., Choe, H., Farzan, M., and Sawyer, S.L. (2013). Dual host-virus
arms races shape an essential housekeeping protein. PLoS Biol. 11, e1001571.
https://doi.org/10.1371/journal.pbio.1001571.

150.Kerr, S.A., Jackson, E.L., Lungu, O.I., Meyer, A.G., Demogines, A., Ellington, A.D., Georgiou,
G., Wilke, C.O., and Sawyer, S.L. (2015). Computational and functional analysis of the virus-
receptor interface reveals host range trade-offs in New World arenaviruses. J. Virol. 89,

936 11643–11653. https://doi.org/10.1128/JVI.01408-15.

151. Kaur, H., Kalayjian, R., Wu, K., Tassiopoulos, K., Palella, F., Taiwo, B., Bush, W., Hileman, C., Bedimo, R., Koletar, S., et al. (2022). Associations of L-Ferritin and Tim-1 with frailty measures in people with HIV: a cross-sectional and longitudinal study. Lancet Healthy Longev. *3*, S5. https://doi.org/10.1016/s2666-7568(22)00066-6.

152. Kim, J., Jo, Y., Cho, D., and Ryu, D. (2022). L-threonine promotes healthspan by expediting ferritin-dependent ferroptosis inhibition in C. elegans. Nat. Commun. *13*, 6554. https://doi.org/10.1038/s41467-022-34265-x.

153. Daghlas, I., and Gill, D. (2021). Genetically predicted iron status and life expectancy. Clin. Nutr. *40*, 2456–2459. https://doi.org/10.1016/j.clnu.2020.06.025.

154. Perez, K., Ciotlos, S., McGirr, J., Limbad, C., Doi, R., Nederveen, J.P., Nilsson, M.I., Winer, D.A., Evans, W., Tarnopolsky, M., et al. (2022). Single nuclei profiling identifies cell specific markers of skeletal muscle aging, frailty, and senescence. Aging (Albany NY) *14*, 9393–9422. https://doi.org/10.18632/aging.204435.

155. Trifunovic, A., Wredenberg, A., Falkenberg, M., Spelbrink, J.N., Rovio, A.T., Bruder, C.E., Bohlooly-Y, M., Gidlöf, S., Oldfors, A., Wibom, R., et al. (2004). Premature ageing in mice expressing defective mitochondrial DNA polymerase. Nature *429*, 417–423. https://doi.org/10.1038/nature02517.

156. Van Goethem, G., Dermaut, B., Löfgren, A., Martin, J.J., and Van Broeckhoven, C. (2001). Mutation of POLG is associated with progressive external ophthalmoplegia characterized by mtDNA deletions. Nat. Genet. *28*, 211–212. https://doi.org/10.1038/90034.

157. Cao, L., Li, W., Kim, S., Brodie, S.G., and Deng, C.-X. (2003). Senescence, aging, and malignant transformation mediated by p53 in mice lacking the Brca1 full-length isoform. Genes Dev. *17*, 201–213. https://doi.org/10.1101/gad.1050003.

158. Vijg, J., Perls, T., Franceschi, C., and van Orsouw, N.J. (2001). BRCA1 gene sequence variation in centenarians. Ann. N. Y. Acad. Sci. *928*, 85–96. https://doi.org/10.1111/j.1749-6632.2001.tb05639.x.

159. Fearon, E.R. (1997). Human cancer syndromes: clues to the origin and nature of cancer. Science *278*, 1043–1050. https://doi.org/10.1126/science.278.5340.1043.

160. Donoho, G., Brenneman, M.A., Cui, T.X., Donoviel, D., Vogel, H., Goodwin, E.H., Chen, D.J., and Hasty, P. (2003). Deletion of Brca2 exon 27 causes hypersensitivity to DNA crosslinks, chromosomal instability, and reduced life span in mice. Genes Chromosomes Cancer *36*, 317–331. https://doi.org/10.1002/gcc.10148.

161. Wooster, R., Bignell, G., Lancaster, J., Swift, S., Seal, S., Mangion, J., Collins, N., Gregory, S., Gumbs, C., and Micklem, G. (1995). Identification of the breast cancer susceptibility gene BRCA2. Nature *378*, 789–792. https://doi.org/10.1038/378789a0.

162. Sotgia, S., Zinellu, A., Mangoni, A.A., Serra, R., Pintus, G., Caruso, C., Deiana, L., and Carru, C. (2017). Cellular immune activation in Sardinian middle-aged, older adults and centenarians. Exp. Gerontol. *99*, 133–137. https://doi.org/10.1016/j.exger.2017.10.005.

163. Lee, K.-A., Flores, R.R., Jang, I.H., Saathoff, A., and Robbins, P.D. (2022). Immune senescence, immunosenescence and aging. Front. Aging *3*, 900028.

977   https://doi.org/10.3389/fragi.2022.900028.

164. Yousefzadeh, M.J., Flores, R.R., Zhu, Y., Schmiechen, Z.C., Brooks, R.W., Trussoni, C.E., Cui, Y., Angelini, L., Lee, K.-A., McGowan, S.J., et al. (2021). An aged immune system drives senescence and ageing of solid organs. Nature *594*, 100–105. https://doi.org/10.1038/s41586-021-03547-7.

165. Lee, C.-S., Chang, C.-H., Chen, C.-Y., Shih, C.-Y., Peng, J.-K., Huang, H.-L., Chen, P.-Y., Huang, T.-L., Chen, C.-Y., and Tsai, J.-S. (2022). Upregulation of cluster of differentiation 36 mRNA expression in peripheral blood mononuclear cells correlates with frailty severity in older adults. J. Cachexia Sarcopenia Muscle *13*, 1948–1955. https://doi.org/10.1002/jcsm.13003.

166. Kimmel, J.C., Yi, N., Roy, M., Hendrickson, D.G., and Kelley, D.R. (2021). Differentiation reveals latent features of aging and an energy barrier in murine myogenesis. Cell Rep. *35*, 109046. https://doi.org/10.1016/j.celrep.2021.109046.

167. Tian, Y., Lu, J., Hao, X., Li, H., Zhang, G., Liu, X., Li, X., Zhao, C., Kuang, W., Chen, D., et al. (2020). FTH1 inhibits ferroptosis through ferritinophagy in the 6-OHDA model of Parkinson's disease. Neurotherapeutics *17*, 1796–1812. https://doi.org/10.1007/s13311-020-00929-z.

168. Ricklefs, R.E. (2010). Life-history connections to rates of aging in terrestrial vertebrates. Proc. Natl. Acad. Sci. U. S. A. *107*, 10314–10319. https://doi.org/10.1073/pnas.1005862107.

169. Tian, S., Zeng, J., Jiao, H., Zhang, D., Zhang, L., Lei, C.-Q., Rossiter, S.J., and Zhao, H. (2023). Comparative analyses of bat genomes identify distinct evolution of immunity in Old World fruit bats. Sci Adv *9*, eadd0141. https://doi.org/10.1126/sciadv.add0141.

170. Jonathan Sleeman, Center Director, USGS National Wildlife Health Center (2020). NWHC Operations During the COVID-19 Pandemic and Information About Coronaviruses in Wildlife.

171. White-Nose Syndrome Disease Management Working Group National White-Nose Syndrome Decontamination Protocol.

172. Wu, N., Nguyen, X.-N., Wang, L., Appourchaux, R., Zhang, C., Panthu, B., Gruffat, H., Journo, C., Alais, S., Qin, J., et al. (2019). The interferon stimulated gene 20 protein (ISG20) is an innate defense antiviral factor that discriminates self versus non-self translation. PLoS Pathog. *15*. https://doi.org/10.1371/journal.ppat.1008093.

173. Legrand, A., Dahoui, C., De La Myre Mory, C., Noy, K., Guiguettaz, L., Versapuech, M., Loyer, C., Pillon, M., Wcislo, M., Guéguen, L., et al. (2024). SAMD9L acts as an antiviral factor against HIV-1 and primate lentiviruses by restricting viral and cellular translation. PLoS Biol. *22*, e3002696. https://doi.org/10.1371/journal.pbio.3002696.

174. Išerić, H., Alkan, C., Hach, F., and Numanagić, I. (2022). Fast characterization of segmental duplication structure in multiple genome assemblies. Algorithms Mol. Biol.

175. Brown, M., González De la Rosa, P.M., and Mark, B. (2023). A Telomere Identification Toolkit https://doi.org/10.5281/zenodo.10091385.

176. Li, H. (2021). New strategies to improve minimap2 alignment accuracy. Bioinformatics.

177. Goel, M., and Schneeberger, K. (2022). plotsr: visualizing structural similarities and rearrangements between multiple genomes. Bioinformatics *38*, 2922–2926. https://doi.org/10.1093/bioinformatics/btac196.

178. Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A.,
and Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. Mol. Biol. Evol. *37*, 1530–1534.
https://doi.org/10.1093/molbev/msaa015.

179. Zhang, C., and Mirarab, S. (2022). Weighting by gene tree uncertainty improves accuracy of
quartet-based species trees. Mol. Biol. Evol. *39*. https://doi.org/10.1093/molbev/msac215.

180. Romiguier, J., Ranwez, V., Delsuc, F., Galtier, N., and Douzery, E.J.P. (2013). Less is more in
mammalian phylogenomics: AT-rich genes minimize tree conflicts and unravel the root of
placental mammals. Mol. Biol. Evol. *30*, 2134–2144. https://doi.org/10.1093/molbev/mst116.

181. Dos Reis, M., and Yang, Z. (2019). Bayesian Molecular Clock Dating Using Genome-Scale
Datasets. Methods Mol. Biol. *1910*, 309–330. https://doi.org/10.1007/978-1-4939-9074-0_10.

182. Yang, Z. (2007). PAML 4: phylogenetic analysis by maximum likelihood. Mol. Biol. Evol. *24*,
1586–1591. https://doi.org/10.1093/molbev/msm088.

183. Phillips, M.J. (2016). Geomolecular dating and the origin of placental mammals. Syst. Biol. *65*,
546–557. https://doi.org/10.1093/sysbio/syv115.

184. Gunnell, G.F., and Simmons, N.B. (2005). Fossil evidence and the origin of bats. J. Mamm.
Evol. *12*, 209–246. https://doi.org/10.1007/s10914-005-6945-2.

185. Eiting, T.P., and Gunnell, G.F. (2009). Global Completeness of the Bat Fossil Record. J.
Mamm. Evol. *16*, 151–173. https://doi.org/10.1007/s10914-009-9118-x.

186. Storch, G., Sigé, B., and Habersetzer, J. (2002). Tachypteron franzeni n. gen., n. sp., earliest
emballonurid bat from the Middle Eocene of Messel (Mammalia, Chiroptera). Palaontol. Z. *76*,
189–199. https://doi.org/10.1007/bf02989856.

187. Ravel, A., Marivaux, L., Tabuce, R., Ben Haj Ali, M., Essid, E.L.M., and Vianey-Liaud, M.
(2012). A new large philisid (Mammalia, Chiroptera, Vespertilionoidea) from the late Early
Eocene of Chambi, Tunisia: LARGE BAT FROM CHAMBI. Palaeontology *55*, 1035–1041.
https://doi.org/10.1111/j.1475-4983.2012.01160.x.

188. Lim, B.K. (2009). Review of the origins and biogeography of bats in South America. Chiroptera
Neotropical *15*, 391–410.

189. Morgan, G.S., and Czaplewski, N.J. (2003). A New Bat (Chiroptera: Natalidae) from the Early
Miocene of Florida, with Comments on Natalid Phylogeny. J Mammal *84*, 729–752.
https://doi.org/10.1644/1545-1542(2003)084<0729:ANBCNF>2.0.CO;2.

190. Elliot, M.G., and Mooers, A.Ø. (2014). Inferring ancestral states without assuming neutrality or
gradualism using a stable model of continuous character evolution. BMC Evol. Biol. *14*, 226.
https://doi.org/10.1186/s12862-014-0226-8.

191. Kosakovsky Pond, S.L., Poon, A.F.Y., Velazquez, R., Weaver, S., Hepler, N.L., Murrell, B.,
Shank, S.D., Magalis, B.R., Bouvier, D., Nekrutenko, A., et al. (2020). HyPhy 2.5-A
customizable platform for evolutionary hypothesis testing using PHYlogenies. Mol. Biol. Evol.
37, 295–299. https://doi.org/10.1093/molbev/msz197.

192. Mendes, F.K., Vanderpool, D., Fulton, B., and Hahn, M.W. (2021). CAFE 5 models variation in
evolutionary rates among gene families. Bioinformatics *36*, 5516–5518.

https://doi.org/10.1093/bioinformatics/btaa1022.

Materials and Methods

Data availability

All sequencing data and genomes generated in this study are available on NCBI under Bioprojects PRJNA973719, PRJNA1035541, and PRJNA1295526. Annotations generated in this study are available at https://github.com/docmanny/myotis-gene-annotations. All other code is available at https://github.com/sudmantlab/MyotisGenomeAssembly. All other data are archived on DataDryad at DOI: 10.5061/dryad.02v6wwqfh.

Samples and materials

All bats sampled for this study were wild caught under scientific collection permits for California and Arizona (see **Extended Data Table 1**). Bats were sampled using standard mist-netting procedures, including taking standard body measurements, following USGS recommendations for White-Nose Syndrome and COVID-19 prevention[170,171]. For *M. lucifugus*, the donor individual was field-caught in California and transported to the Genetics Laboratory of the California Department of Fish and Wildlife, where they were euthanized via isofluorane. The *M. velifer* individual was caught in Arizona and euthanized in the field via isoflurane. For both *M. lucifugus* and *M. velifer*, tissues were collected and preserved via flash-freezing in liquid nitrogen. All other genomes were generated from primary cell lines. Additional information can be found in the *Extended Methods* and in **Extended Data Table 1**.

All PKR experiments were performed using HeLa PKR-KO cells (kindly provided by A. Geballe, Fred Hutchinson Cancer Center, Seattle WA). The cells were maintained at 37°C with 5% CO_2 and cultured in DMEM supplemented with 5% fetal bovine serum (FBS), 1% penicillin/ streptomycin mix and 1 µg/mL puromycin (Sigma-Aldrich). All transfections were performed 24 hours after seeding, using 3 µL of TransIT-LT1 Transfection Reagent (Mirus Bio) per 1 µg of DNA and Opti-MEM media. We used previously-generated pSG5-FLAGx2 vectors encoding either *M. myotis* PKR1 (GenBank OP006550), *M. myotis* PKR2 (GenBank OP006559), *M. velifer* PKR1 (GenBank OP006558), or *M. velifer* PKR2 (GenBank OP006557)[30]. Plasmids encoding the interferon-stimulated gene ISG20[172] and a constitutively active variant of the sterile alpha motif domain-containing protein 9-like SAMD9L-F886Lfs*11 (here, SAMD9L[173]) were used as controls in viral infections and cell translation experiments, respectively.

Near-complete genome assembly and annotation

Details on DNA extraction, preparation of PacBio HiFi and Omni-C (Dovetail Genomics) libraries, and genome assembly and manual curation can be found in the *Extended Methods*.

Gene & repetitive element annotation and alignments

1089 Details describing our procedure for identifying and annotation repetitive elements and protein-
1090 coding gene annotations can be found in the *Extended Methods.*

RNA sequencing for gene annotations and differential expression

1092 Details on RNA extraction, library preparation, sequencing, and can be found in the *Extended Methods* and
1093 **Extended Data Table 9**.

Structural variation

To understand the distribution of structural variants, including segmental duplication events, we used
SyRI (Synteny and Rearrangement Identifier[65]) and BISER (Brisk Inference of Segmental duplication
Evolutionary stRucture[174]). We first masked telomere regions using TIDK (Telomere Identification toolKit[175]),
and mapped the primary 22 scaffolds of the nearctic *Myotis* genomes to each other with minimap2[176]. The
scaffold corresponding to the X chromosome was omitted because there is no corresponding scaffold in the
M. yumanensis assembly. To verify homologous chromosomes and fix strand orientation, we used *fixchr*
from the SyRI package and manually renamed scaffolds accordingly, then re-mapped them with minimap2.
We ran SyRI on the resulting files and plotted the results with plotsr[177]. We ran BISER on the primary 22
scaffolds of the nearctic *Myotis* genomes with –keep-contigs and default settings to generate bed files with
the inferred segmental duplication regions.

Gene Trees & Phylogeny

A phylogeny of all 536 mammals in our alignments was generated using IQTREE[178] (version 2.3.1)
using all gene alignments with the settings "-B 1000 -m GTR+F3x4+R6." To generate gene trees, we first
filtered our gene alignments to exclude alignments with over 50% gaps in the sequence and less than 4
species. With the remaining alignments, we used IQTREE to find the best-fitting substitution model and tree
using settings "--wbtl --bnni --alrt 1000 -B 1000 --safe". The best substitution models for each gene were
saved as a NEXUS file. As the bootstrap values for this tree were unanimously 100%, we further confirmed
the species tree by using ASTRAL-IV[179] (v1.24.4.7) using the settings "-t 54 -u 2 -C", and confirmed that our
phylogeny agreed with other previously-published *Eutherian* phylogenies, including phylogenies of 39
mammals[180], of 241 placental mammals[58], and of 11 nearctic *Myotis* species[56].

Time-calibration of 59 bat genomes

Using our codon alignments of 59 bat genomes, we generated a time-calibrated phylogeny using
mcmctree[181] and PAML[182] (v. 4.10.0) using an approximate likelihood method. Using the pan-bat codon
alignments and our phylogeny as input, with fossil calibrations based on previously published work[9,44,58,183–
189], we ran *mcmctree* twice to generate the Hessian matrix and confirm convergence. This was followed by
10 independent chains using the "out.BV" file from the first run. Finally, the output files of all 10 chains were
combined to compute final divergence time estimates (see Extended Data Table 2).

Ancestral Body Size, Lifespan, and Cancer Risk reconstruction

1122

To explore how body size and lifespan have evolved over time in mammals, we used a super-
phylogeny of mammal species[55] subsampled to only contain species with extant body size and lifespan data
collected from AnAge[18] and PanTHERIA[19]. Ancestral body sizes and lifespans were simulated separately
using StableTraits[190] and the settings "*--iterations 100000 --skip 1000 --chains 8*". Estimates were further
validated by comparing our initial results to a second run of StableTraits using the same settings. To test the
robustness of RICR against phylogenetic uncertainty, we repeated these simulations using 5,182/10,000 of
the alternative topologies for the mammalian tree published in Upham *et al*. (2019).

Selection Scans & Evolutionary Rates

1130

aBSREL

1131

To conservatively test for branch-specific selection, we used aBSREL[100,191] (version 2.5.48) to test
for selection at each branch within the nearctic *Myotis* clade for 15,734 gene alignments spanning 536
mammals. These genes were identified as 1:1 orthologs across the full alignment, with no more than 50%
sequence gaps and at least 4 species present in the alignment. For each gene alignment, the full phylogeny
was trimmed to match the species present, and the HYPHY script "label-tree.bf" was used to highlight all
nodes within nearctic *Myotis* as foreground branches. aBSREL was then run using the pruned and
highlighted tree and gene alignments on the command line with the options "--code Universal --branches
Foreground". We defined a gene as under selection if one or more regions within the gene demonstrated a
selective pressure with an FDR-corrected p-value ≤ 0.05; a gene was specifically identified as under positive
selection if at least one of these significant regions had an $\omega > 1$.

BUSTED

1142

To quantify the total amount of positive selection across the *Myotis* tree or the different species trees
used in this manuscript, we used an improved version of the BUSTED[72,191] test called BUSTED-MH (see
Extended Methods for details on BUSTED vs BUSTED-MH). We applied BUSTED-MH to 19,646 *Myotis*
orthologous CDS alignments with at least five orthologs. These orthologs are cases where the Orthofinder
gene trees coincide with the species tree. This effectively removes issues regarding whether we should use
the gene or the species tree, at the cost of removing 2,110 genes from the *Myotis* selection analysis. Similarly,
we applied BUSTED-MH to 17,469 non-*Myotis* bat alignments with at least five orthologs. The species in
these two sets (*Myotis* bats and non-*Myotis* bats) are mutually exclusive. This includes a subset of 14,091
alignments with orthologs present in two thirds of the non-*Myotis* bat species, regardless of the genes'
presence/absence in the *Myotis*-only complement. This is specifically used to show that patterns of virus-
driven adaptation are representative of all bats. We also tested 17,890 primate alignments with at least five
orthologs with BUSTED-MH, as well as 19,311 glire, 18,000 carnivora and 18,504 ungulate alignments.

Enrichment scores for VIPs

1155

We expanded on the set of VIPs reported in Souilmi *et al*. (2021)[73] (5,291 VIPs) by searching the
literature for more recent publications reporting VIPs, with a final dataset of 5,527 VIPs. These VIPs ("All

VIPs") were further sub-categorized into 4 groups based on their interactions with either DNA viruses and/or
RNA viruses: "DNA VIPs", "DNA-only VIPs", "RNA VIPs", and "RNA-only VIPs" (**Extended Data Table 5**).
For each VIP set, we selected sets of matching control genes by controlling for 16 different criteria. (Extended
Data Table 5). Sets of control genes were resampled in a bootstrap procedure to generate 95% confidence
intervals for sets of genes across a range of p-values; additional details can be found in the *Extended*
Methods.

Gene Duplications and CAFE analysis

To quantify patterns of gene duplication and loss, we quantified the copy number of genes with human
orthologs from our gene annotations for each nearctic *Myotis* genome. To calculate per-gene expansion and
loss rates and their statistical significance, we ran CAFE[192] v5 on the previously described set of copy number
counts using our time-calibrated species tree pruned to include only the nine nearctic *Myotis* species. CAFE
uses a birth-death model of gene family evolution to investigate changes in gene family size accounting for
evolutionary history using the species phylogeny. Evolutionary branch length is incorporated into the model
expectations, thus accounting for potential bias resulting from differences in branch lengths. We ran CAFE
on the subset of genes with two or more copies in at least one species using a Poisson distribution for the
root frequency (*-p*), first generating an error model to correct for genome assembly and annotation error (-
e). We compared the base model (each gene family belongs to the same evolutionary rate category) to
gamma models (each gene family can belong to one of k evolutionary rate categories) with different values
of k from 2 to 13. A final gamma model with k=9 was chosen to balance model log likelihood with the number
of gene families for which the optimizer failed. The model was run three separate times to ensure
convergence.

To understand if genes in these pathways have higher birth rates or are more likely to have significant
changes in gene copy number than expected relative to other genes, we evaluated the gene copy birth rate
λ and number of genes that have significantly expanded or contracted in copy number on at least one branch
within our nearctic *Myotis* phylogeny using a bootstrap procedure. Following Huang et al.[42], we tested if VIP
genes in particular underwent significant copy number changes or had significantly different birth/death rates
than non-VIP genes. For each category of VIP genes (all VIPs, DNA VIPs, DNA only VIPs, RNA VIPs, and
RNA only VIPs), we generated 100 bootstrap sets of control non-VIP genes with the same number of genes
as the corresponding VIP gene set. We ran CAFE on each set of VIP genes and the corresponding control
non-VIP genes to infer per-gene birth-death rates and per-gene, per-branch expansion/loss events.

Assessment of DNA Double-Strand Break Tolerance

We assessed each species' tolerance to DNA double strand breaks using a by measuring viability,
cytotoxicty, and apoptosis across a range of doses of Neocarzinostatin, a radiomimetic drug. We measured
dose response curves in wing-derived primary dermal fibroblasts across 5 bat species (*Myotis lucifugus*, n=8;
Myotis evotis, n=8; *Rousettus langosus*, n=2; *Eidolon helvum*, n=2; *Pteropus rodrigensis*, n=2) using the
multiplexed ApoTox-Glo assay (Promega). Using 96-well plates, two individuals and 11 doses were assessed
simultaneously with four technical replicates. Results were normalized to treatment controls for each
individual in R as previously described[36,37,40,45].

PKR1/2 Characterization in PKR-KO HeLa cells

Western blot

We assessed for the steady state protein expression of *M. myotis* Flag-PKRs after transfection of 350 ng or 700 ng total of DNA plasmids encoding: PKR1 alone, PKR2 alone or both PKR1 and PKR2 (175 ng of each and 350 ng of each, respectively). Briefly, cells were re-suspended and lysed in ice-cold RIPA buffer (50 mM Tris pH8, 150 mM NaCl, 2 mM EDTA, 0.5% NP40) with protease inhibitor cocktail (Roche) and sonicated. 20 μL of the clarified fraction was denatured with 5 μL of 6x Laemmli buffer at 95°C for 5 min and loaded into 4-20% BioRad Criterion TGX Stain-Free precast gel. The wet transfer into a PVDF membrane was executed overnight at 4°C. The membranes were blocked in a 1xTBS-T buffer (Tris HCl 50 mM pH8, NaCl 30 mM, 0.05% of Tween 20) containing 10 % powder milk, and were incubated for 1h. The membranes were incubated with primary mouse anti-Flag (Sigma F3165) and anti-Tubulin (Sigma T5168) antibodies and secondary anti-Mouse IgG-Peroxidase conjugated (Sigma A9044). Detection was made using the Chemidoc Imaging System (BioRad) with SuperSignal West Pico Chemiluminescent Substrate (ThermoFisher Scientific).

Luciferase reporter assays

Luciferase reporter assays were carried out to investigate whether the two PKR paralogs have synergistic, additive or dominant negative effect in translation shutdown. 25,000 PKR-KO cells were seeded in 24-well plates. The next day, cells were transfected with 350 ng of empty pSG5, 350 ng of PKR1, 350 ng of PKR2, or 175 ng of PKR1+ 175 ng of PKR2, with additional 50 ng of FFLuc firefly luciferase reporter plasmid per well. Sterile alpha motif domain-containing proteins 9L (SAMD9L gain-of-function mutant) was used as a positive control of translational repression[173]. 24 h post transfection, cells were washed with PBS, lysed by a 5× reporter lysis buffer (Promega) and incubated overnight at -20°C. Cells were then collected and 100 μl of the luciferase substrate (Promega) was added to 20 μl of the lysis supernatant. Alternatively, cells were lysed using BrightGlow Lysis Reagent (Promega E2620). The relative luminescence units (RLUs) proportional to the translation of the luciferase reporter were immediately quantified with LUMIstar Omega microplate reader optima (BMG Labtech). All luciferase assays were conducted in technical duplicates in at least five independent experiments. Fold change results were normalized to the empty pSG5 condition within each independent experiment.

PKR co-immunoprecipitation

Hela PKR-KO cells were transfected with 1.25μg plenti6 HA-tagged *M. myotis* PKR1 plasmid per million cells and 1.25μg of either plenti6 myc empty vector, myc-tagged *M. myotis* PKR1 or myc-tagged *M. myotis* PKR2 plasmid using TransIT-LTI transfection reagent (Mirus Bio). The next day, some wells were infected with Sindbis virus expressing GFP (SINV-GFP) at MOI 2 for 24 hours. Cells were then scraped with cold PBS and pelleted. For the IP, cells were lysed in 500μl IP buffer (50mM Tris HCl pH7.5, 140mM NaCl, 6 mM MgCl2, 0.1% NP40) supplemented with RNase (RiboLock, Fisher Scientific) and protease (complete EDTA-free protease inhibitor cocktail, Sigma) inhibitors for 10 minutes on ice, then centrifuged at 12,000 *xg* for 10 min at 4°C. 5% of the volume was kept for input, while the rest was incubated with 40μl μMACS anti-c-myc MicroBeads (Miltenyi Biotec) for 1h at 4°C with constant rotation. Samples were then loaded onto μMACS columns placed in the magnetic field of a μMACS Separator (Miltenyi Biotec), washed 4 times with cold IP buffer, and eluted with the μMACS denaturing elution buffer. Proteins were denatured in elution buffer for 5

min at 95°C, then loaded onto a 4-20% BioRad Criterion TGX Stain-Free precast gel and transferred onto an Amersham Protran nitrocellulose membrane (Sigma) for 1h. Membranes were blocked for 1h in 5% milk in PBS (Euromedex) supplemented with 0.2% tween (Fisher) and incubated with mouse anti-myc monoclonal antibodies (Abcam 9E10, cat# ab32) then secondary anti-mouse IgG antibodies conjugated with HRP (Sigma, cat# A4416), or with rat anti-HA antibodies conjugated with HRP (Roche, Sigma, cat# 12013819001). Images were taken on a Fusion FX imager (Vilber) with SuperSignal West Femto Chemiluminescent Substrate (ThermoFisher Scientific).

Cell viability assay

Hela PKR-KO cells were transfected 24h after plating in 96 well plates (n=10,000 cells/well), with 100 or 200 ng of pSG5 plasmid: empty or coding for *M. myotis* or *M. velifer* PKR1, PKR2 or PKR1+2 equal mix (50%-50%). 24 hours post-transfection, positive control cells were treated with an apoptosis-inducing drug, Etoposide, at different doses (250, 200 or 100 µM). 48 hours post transfection, cells were harvested and lysed to quantify luminescent signal according to CellTiter-Glo® Luminescent Cell Viability Assay (Promega) kit protocol.

VSV and SINV infections

VSV infections. 200,000 cells were plated in 12-well and transfected 24 h after with 350 ng of pSG5 plasmid: empty, or encoding *M. myotis* or *M. velifer* PKR1, PKR2, or equal input of PKR1 and PKR2 (175 ng per plasmid), or a plasmid encoding interferon-stimulated exonuclease gene 20 (ISG20), due to its known antiviral activity against VSV as positive control[172]. Cells were infected 24 h post transfection with replicative VSV-GFP virus[70] at a MOI of 3. Cells were fixed with 4% paraformaldehyde 16-18 hours post infection. VSV infection was quantified by measuring the percentage of GFP positive cell populations with BD FACSCanto II Flow Cytometer (SFR BioSciences). Fold change results were normalized to the empty pSG5 condition across at least three independent experiments.

SINV infections. Hela PKR-KO cells were transfected with 5µg pSG5 empty vector, *M. myotis* PKR1, *M. myotis* PKR2 or 2.5µg *M. myotis* PKR1 + 2.5µg *M. myotis* PKR2 per million cells using TransIT-LTI transfection reagent (Mirus Bio). The next day, some wells were infected with SINV-GFP at MOI 0.2. Cells were then placed into a CellCyte X live cell imaging system (Cytena) and pictures of every well were taken every 2h for 48h. The fraction of GFP+ cells over the total cell area was measured and averaged from six photos of 2 individual wells per condition, and repeated for a total of three independent experiments.

Fig. Legends

Fig. 1: 8 near-complete reference assemblies for North American (nearctic) *Myotis*. A) Phylogeny of 38 bat genomes with 3 outgroup species: cow *(bosTau9)*, mouse *(mm39)*; and human *(T2T-CHM13v2.0)*. Bars at the tips of the phylogeny indicate the Area under the NGx curve (AuNG) score of each genome, a metric of genome contiguity independent of genome size. The dotted line represents the AuNG score for the T2T-CHM13v2.0 assembly. **B)** The time-calibrated phylogeny of 9 nearctic and two representative Palaearctic *Myotis* species based on orthologous codon alignments. Blue bars represent age uncertainties. The branch highlighted in red (*M. volans* as outgroup to the other nearctic *Myotis*) has 100% bootstrap support, but 0% local posterior probability support. **C)** Map of capture sites for new cell lines with representative images for the species sequenced in this study (see "Acknowledgements" for attributions). **D)** Mammalian BUSCO scores for annotations generated for the 8 new *Myotis* genomes. **E)** Ideogram bar plot indicating completion status of each chromosome in assembly. Pie graphs indicate completion status of all chromosomes in assembly. Percentages above ideogram indicate the proportion of T2T assembled chromosomes across species. All chromosomes were positively identified based on size, synteny, and homology to human chromosomes[51]. "Complete (T2T)" status indicates that a chromosome is fully assembled telomere-to-telomere without gaps; "Draft (T2T, gaps)" status indicates that a chromosome is fully scaffolded with both telomeres, but has one or more gaps in the assembly; "Incomplete" status indicates that a chromosome was positively identified, but was not scaffolded from telomere to telomere (only contains one telomere).

Fig. 2: A varied structural variation landscape across 9 nearctic *Myotis* species. A) Synteny between chromosomes of 9 *Myotis* species showing syntenic regions (grey), inversions (orange), translocations (green), and duplications (blue). **B)** Distribution of transposable elements (top) and segmental duplications (red heatmap, bottom) in *M. velifer*. Putative locations of centromeres are denoted by the dotted lines. **C)** Pie chart of overall genomic proportions of TEs in *M. velifer*. **D)** Histogram of segmental duplication size distributions genome-wide in *M. velifer*.

Fig. 3: Evolutionary history and function of an actively segregating copy number polymorphism of *PKR* in *Myotis*. A) Structural comparison of the main PKR haplotypes in two species. Orthologous regions between the two haplotypes are indicated by grey bands, while syntenic duplications are indicated in green. Exons are annotated with black marks. **B)** Cartoon of the PKR locus in the two phased haplotype assemblies of each species. While *PKR2* is present across all haplotypes, *PKR1* and *PKR copy 3* are polymorphic within and across species. Each number indicates the number of exons per gene. **C)** Reconciled gene tree for *PKRs* across all haplotypes and species shown in B. Haplotype corresponding to the reference (A) and alternate (B) haplotype for each species are represented by upper- and lower-diagonal triangles, respectively. **D)** Effect of Myotis PKR expression on luciferase reporter translation in PKR-KO cells, measured in Relative Light Units (RLU) and normalized to the empty pSG5 control. Human SAMD9L-GoF is a positive control of translation inhibition[173]. **E-F)** Effect of PKR on viral infection by VSV-GFP (E, and S3B) and SINV-GFP (F 34h post-infection, and S3C), measured by flow cytometry as the % of GFP+ cells, normalized to the pSG5 control. Although all conditions restricted VSV and SINV, the expression of both PKR1+PKR2 was not beneficial against these viral infections. ISG20 served as a positive control of VSV-GFP restriction[172]. **G)** Co-immunoprecipitation (IP) of PKR-KO cells transfected with *Myotis* HA-PKR1 and either *Myotis* myc-PKR1, *Myotis* myc-PKR2 or a myc-empty vector control. Proteins were pulled down with anti-myc beads and lysates from 5% input or IP samples were run on a western blot and stained for HA and myc. Representative of 3 independent experiments. Extended Data Fig. 3E for SINV conditions. H, Quantification of the three independent experiments (mock and SINV): HA left and myc right. For **D-F**, error bars indicate the means ± SEM for at least three independent experiments. Statistics, unpaired t-test (p<0.05). mPKR, Myotis myotis PKR; vPKR, Myotis velifer PKR. For gel source data, see Supplementary Figure 1.

Fig. 4: Adaptation to DNA viruses, but not RNA viruses, is enriched in *Myotis* and other bats. A) Diagram of an example VIP, host cell transmembrane receptor CD45, showing strength of adaptation at each codon and the locations of direct interactions with the human adenovirus protein sec49K. **B-F)** Enrichment plots showing the ratio of positive selection in only *Myotis* bats in VIPs versus matched sets of control genes at different p-value thresholds. The solid line shows the median ratio; the color of the line, and the number above each point, represents the number of VIPs with significant BUSTED-MH p-values at the given threshold; the grey band represents the 95% confidence interval generated by bootstrapping sets of matched control genes. The number above each point indicates the number of VIP genes and sampled control genes used for each ratio calculation. The 95% confidence interval widens at lower p-values because there are fewer VIPs with significant p-values as the alpha threshold decreases, reducing power. **G-I)** The same as above, across all bats in this study excluding *Myotis*. **J-L)** The number of VIP genes inferred by CAFE to have significantly expanded or contracted in copy number on at least one branch of the nearctic *Myotis* phylogeny (blue line) compared to the distribution across 100 randomized sets of non-VIP control genes (black line). Vertical dashed lines indicated the 95% highest density interval for the control gene distribution.

Fig. 5: Evolution of cancer risk and resistance in *Myotis* and other mammals. A, B) Cophylo plot of the evolution of body size (**A**) and lifespan (**B**) across *Eutheria*. Branch lengths in **A-B** are scaled proportional to the rate of change of the trait over time. **C)** The phylogeny of nearctic *Myotis* species with inset pie charts representing the proportion of the top 100 Reactome pathways overrepresented among genes under selection at each node that are associated with cancer-related processes. Below, pie chart indicates expected proportion of pathways are cancer-associated after 1000 random samples of 100 pathways from the full Reactome database. Asterisks represent nodes with proportions greater than the expected value at p≤0.05 using Fisher's exact test. **D)** Proportion of the top 100 Reactome pathways overrepresented among genes under selection across all nodes in a species' evolutionary history that are associated with cancer-related processes. **E)** Volcano plot of overrepresented pathways in Reactome among the union set of genes under selection across all nodes in the evolutionary history for *M. lucifugus*. **F)** Viability and Apoptosis fold-change in 5 bat species in response to different doses of neocarzinostatin (NCS), a potent inducer of DNA double-strand breaks. Points represent individual replicates normalized to each species' control, while bars represent mean ± 95% confidence intervals.

Fig. 6: **Pleiotropy among genes under selection in *M. lucifugus* between DNA-damage responsive genes and DNA-only VIPs. A-B)** Volcano plots of genes that are differentially expressed in *M. lucifugus* after 6 hours (**A**) and 18 hours (**B**) of treatment with 100nM neocarzinostatin (NCS). **C-D)** Pathway-level Gene Set Enrichment Analysis (GSEA) of genes differentially expressed at 6 and 18 hours post-treatment. **E)** Upset plot between genes that are differentially expressed after NCS treatment and DNA-only VIPs; the hypergeometric P-value of the intersection is shown. **F)** Pathway-level overrepresentation analysis of genes that are DNA-only VIPs and are also differentially expressed in response to NCS treatment. **G)** Pathway-level overrepresentation analysis of genes identified as under selection in *M. lucifugus* via ABSREL that are DNA-only VIPs and are also differentially expressed in response to NCS treatment.

Extended Data

Extended Data Fig. 1-7:
1. **Contiguity statistics for 8 novel *Myotis* genomes**
2. **Estimated node divergence times for *Chiroptera* and ASTRAL-IV scores.**
3. **Correlations of TEs and structural variation in *Myotis* genomes**
4. **Additional functional characterization of *Myotis* PKR duplicates**
5. **VIP selection enrichments in bats and other mammals**
6. **Evolution of Body Size and Lifespan across *Myotis***
7. **Genes differentially expressed after neocarzinostatin treatment and their overlap with DNA-only VIPs**

Extended Data Table 1. Genome Statistics
Extended Data Table 2. Accessions for additional genomes used in this study
Extended Data Table 3. SyRI-identified structural variants (SVs)
Extended Data Table 4. Experimental data for PKR experiments
Extended Data Table 5. List of VIPs and VIP subclasses
Extended Data Table 6. Ancestral State Reconstructions for body size and lifespan in mammals
Extended Data Table 7. aBSREL significant gene lists and Reactome enrichments
Extended Data Table 8. Experimental data for Neocarzinostatin experiments
Extended Data Table 9. RNA-seq QC statistics and GSEA results

A

Reference Chr

V1, V3, V5, V7, V10, V13, V16, V20, V23, V8, V11, V14, V18, V21, V24, V9, V12, V15, V19, V22, V25

Chromosome Position (Mbp)

Genomes
- *M. occultus*
- *M. lucifugus*
- *M. yumanensis*
- *M. velifer*
- *M. californicus*
- *M. auriculus*
- *M. evotis*
- *M. thysanodes*
- *M. volans*

Annotations
- Syntenic
- Inversion
- Translocation
- Duplication

B

Proportion of Region

V1, V3, V5, V7, V8, V9, V10, V11

V12, V13, V14, V15, V16, V18, V19, V20, V21, V22, V23, V24, V25, X

Chromosome Position in mMyoVel1 (Mb)

Transposable Element
- DNA
- LTR
- Retroposon
- Unknown/Other
- LINE
- RC
- SINE
- non-TE

Proportion of Region (SD)
0.75 0.50 0.25

C

12.7% 1.4%
1.7%
3.5%
2.1%
78.7%

Genome Proportions (mMyoVel1)

D

Scaled Density
1.00
0.50
0.00

1,000 10,000 100,000 1,000,000

SD length (bp) genome−wide

A

Body Size (log(g))

5   10   15

B

Lifespan (log(yrs))

0   1   2   3   4   5

Whales

Bats

Primates

Elephants

122   31

C

* M.lucifugus
* M.occultus
* M.yumanensis
M.velifer
M.evotis
* M.thysanodes
* M.auriculus
M.californicus
M.volans

14%

86%

Reactome Pathways

Cancer
Not Cancer

D

Enriched Pathways
Associated with
Cancer (%)

0%   5%   10% 15%

E

100 300 500 648

Extracellular matrix
organization

Collagen formation

Resolution of D-Loop
Structures
(Through Holiday
Junction Intermediates)

HDR through
Homologous
Recombination (HRR)

Homologous DNA Pairing
and Strand Exchange

Diseases associated
with the TLR signaling
cascade

MyD88 deficiency
(TLR2/4)

Sensing of
DNA DSBs

Immune System

-log(FDR)

log$_2$(Enrichment Ratio)

F

Species

Myotis lucifugus (n=8)
Myotis evotis (n=9)
Eidolon helvum (n=2)
Pteropus rodrigensis (n=2)
Rousettus lanosus (n=2)

Apoptosis

*** ***   *   *** n.s. n.s. n.s.

Viability

*** n.s.   ***   *** ***

Fold-Change

NCS [nM]   0   10   500

*** $p_{adj} \leq 5e$-4   ** $p_{adj} \leq 5e$-3   * $p_{adj} \leq 0.05$ n.s. $p_{adj} > 0.05$