Extensive longevity and DNA virus-driven adaptation in nearctic *Myotis* bats

Juan M Vazquez[1,2,20], M. Elise Lauterbur[3,4,20], Saba Mottaghinia[5], Léa Gaucherand[6], Sarah Maesen[5], Michael Singer[7], Sarah Villa[7], Melanie Bucci[3], Devaughn Fraser[8,9], Genavieve Gray-Sandoval[3], Zeinab R Haidar[10,111], Melissa Han[12], William Kohler[12], Tanya M. Lama[13], Amandine Le Corf[5], Clara Loyer[5], Dakota McMillan[1,14], Stacy Li[1,15], Johnathan Lo[15], Carine Rey[5], Samantha LR Capel[8], Kathleen Slocum[16], Melissa Sui[7], William Thomas[17], Janet Debelak Tyburec[18], Rachel Brem[19], Richard Miller[12], Michael Buchalski[8], Jose Pablo Vazquez-Medina[1], Sébastien Pfeffer[6], Lucie Etienne[6,21]*, David Enard[3,21,22]**, Peter H Sudmant[1,14,21,22,23]***

Affiliations:

1. Department of Integrative Biology, University of California, Berkeley, Berkeley, CA USA
2. Current affiliation: Department of Biology, Pennsylvania State University, State College, PA USA
3. Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ USA
4. Current affiliation: Department of Biology, University of Vermont, Burlington, VT USA
5. Centre International de Recherche en Infectiologie (CIRI), Inserm U1111, UCBL1, CNRS UMR5308, Ecole Normale Supérieure ENS de Lyon, Université de Lyon, Lyon, France
6. Université de Strasbourg, Architecture et Réactivité de l'ARN, Institut de Biologie Moléculaire et Cellulaire du CNRS, Strasbourg, France.
7. Department of Molecular and Cellular Biology, University of California, Berkeley, Berkeley, CA USA
8. Wildlife Genetics Research Unit, Wildlife Health Laboratory, California Department of Fish and Wildlife, Sacramento, CA, USA
9. Current affiliation: Wildlife Diversity Program, Wildlife Division, Connecticut Department of Energy and Environmental Protection, Burlington, CT, USA
10. Department of Biology, California State Polytechnic University, Humboldt, Arcata, CA USA
11. Current affiliation: Western EcoSystems Technology Inc, Cheyenne, WY USA
12. Department of Pathology and Clinical Laboratories, University of Michigan, Ann Arbor, MI USA
13. Department of Biological Sciences, Smith College, Northampton, MA USA
14. Department of Science and Biotechnology, Berkeley City College, Berkeley, CA USA
15. Center for Computational Biology, University of California, Berkeley, Berkeley, CA USA
16. Bat Conservation International, Austin, TX USA
17. Department of Ecology and Evolution, Stony Brook University, Stony Brook NY USA
18. Bat Survey Solutions, LLC, Tucson, AZ USA
19. Department of Plant and Microbial Biology, University of California, Berkeley, Berkeley, CA 94720, USA
20. These authors contributed equally
21. Senior author
22. These authors contributed equally
23. Lead contact

* Correspondence: lucie.etienne@ens-lyon.fr
** Correspondence: denard@arizona.edu
*** Correspondence: psudmant@berkeley.edu

Abstract

The genus *Myotis* is one of the largest clades of bats, and exhibits some of the most extreme variation in lifespans among mammals alongside unique adaptations to viral tolerance and immune defense. To study the evolution of longevity-associated traits and infectious disease, we generated cell lines and near-complete genome assemblies for 8 closely related species of *Myotis*. Using genome-wide screens of positive selection, analyses of structural variation, and functional experiments in primary cells, we identify new patterns of adaptation contributing to longevity, cancer resistance, and viral interactions in bats. We show that the recurrent evolution of longevity seen in *Myotis* leads to some of the highest predicted increases in cancer risk across mammals and demonstrate a unique DNA damage response in primary cells of the long-lived *M. lucifugus*. We also find evidence of abundant adaptation in response to DNA viruses - but not RNA viruses - in *Myotis* and other bats in sharp contrast with other mammals, potentially contributing to the role of bats as reservoirs of zoonoses. Together, our results demonstrate how genomics and primary cells derived from diverse taxa uncover the molecular bases of extreme adaptations in non-model organisms.

Keywords

Aging, Bats, Cancer, Evolutionary Biology, Functional Genomics, Immunity, Infectious Disease

Introduction

Bats (order *Chiroptera*) represent approximately 20% of all known mammalian species and are one of the most phenotypically diverse clades of mammals[1,2]. Since their emergence 60 million years ago[3–5], many bat lineages have independently evolved a wide variety of life history strategies and phenotypic traits, including exceptional changes in longevity, viral tolerance, and immune defense[6–11]. Such systems, in which shared traits have evolved *de novo* multiple times, are powerful resources for dissecting the genetic basis of phenotypes. Rigorous approaches to studying these traits, however, depend on high-quality, well-annotated genomes to test evolutionary and genomic hypotheses, and on experimental functional systems to validate these hypotheses.

The largest genus of bats - *Myotis* - is estimated to have emerged approximately 33 million years ago[12,13], and encompasses over 139 described species spanning six continents and a wide range of ecological niches[1,12–14]. *Myotis* species demonstrate some of the most extreme variation in lifespan amongst mammals[6,15–18], including a six-fold difference in lifespan between the longest-lived species (*M. brandtii*, 42 years[15,19], **Fig. 1A**) and the shortest-lived species (*M. nigricans*, 7 yrs[15,20]) which diverged approximately 10.6 million years ago[5,14,21,22]. In addition, *Myotis* species have been used as systems for investigating virus tolerance and other pathogen resistance[23–25] associated with the expansion and contraction of antiviral defenses[26–29], which have contributed to bats' ecological role as zoonotic reservoirs[10,11,30–33].

While studies on the genetic basis of longevity in short-lived model organisms have been crucial for identifying and dissecting several key aging pathways, comparative studies of exceptionally long-lived organisms have uncovered novel genes and pleiotropic effects governing lifespan[34–47]. Similarly, studies of infectious disease response are common and powerful in model organisms, but the lack of diversity and use of inbred lines limits their scope. Bats in particular present an important case study in, and opportunity to study, variation in virus adaptation strategies due to bats' role as zoonotic reservoirs and their specific resistance to viruses[43,48].

Here we combined comparative and functional approaches in *Myotis* to uncover genomic and functional evidence of adaptation to both aging-related and infectious diseases. Using near-complete genome assemblies, we identified significant structural variation encompassing stress response, immunity, and inflammation genes, including a trans-species copy number polymorphism of Protein kinase R (PKR). Additionally, we quantify relative intrinsic cancer risk across mammals and find that both ancestral taxa and extant species in *Myotis* are overrepresented at the extreme of increased relative cancer risk. Consistent with this observation, we identified pervasive selection of genes in longevity- and cancer-related processes throughout *Myotis*. We observed some of the strongest signatures of selection at immune-related genes, including strong evidence of adaptation in virus-interacting proteins in *Myotis* and other bats. In contrast to humans and other primates, in which virus adaptation is driven by interactions with RNA viruses[49], we find that virus adaptation in bats is driven by interactions with DNA viruses, implying that bats' exceptional immune adaptations are more protective against DNA viruses. This also highlights a fundamental discordance between the fitness impacts of viruses on humans vs bats. Together, our results suggest that pleiotropy and co-evolution of traits in *Myotis* has played a key role in the evolution of exceptional longevity and infectious disease resistance.

Results

High quality chromosome-level assemblies of 8 *Myotis* bat species

To study how lifespan and viral response have evolved in *Myotis*, we collected skin punches and derived primary cell lines from several North American ("Nearctic")[21] species (**Fig. 1A,C),** including from one of the longest-lived mammals, *Myotis lucifugus*[15]. Using these cell lines and flash frozen tissues we generated *de novo* haplotype-resolved, chromosome-scale genome assemblies for eight species (**Fig. 1A, Extended Data Fig. 1**) using a combination of long-read PacBio HiFi sequencing and HiC scaffolding. These genomes are highly contiguous, with an average of 98.6% of nucleotides assembled into 22-23 syntenic chromosome-scale scaffolds corresponding to the published karyotype[50] with an average QV of 66 (Extended Data Fig. 1; **Table S1**). These genomes have among the highest contig NG50 and lowest auNG scores of any *Chiroptera* genome published to date (**Fig. 1A, E**; **Extended Data Fig. 1**; **Table S1**). We identified an average of 27,536 protein coding genes per genome, with BUSCO[51,52] scores ranging from 98.2% to 98.5% (**Fig. 1D**). Using these annotations, we resolved the conflicting phylogenies of Nearctic *Myotis*[14,53–56] by using 1-to-1 orthologs of protein-coding genes across 534 mammalian genomes (**Table S2**) to build a time-calibrated, maximum likelihood tree of *Eutheria*. Our results conclusively resolve the relationship between known sister species (*M. lucifugus* and *M. occultus*; *M. yumanensis and M. velifer*; and *M. evotis* and *M. thysanodes*) with 100% bootstrap support at all nodes in the Nearctic *Myotis*. (**Fig. 1B**; **Extended Data Fig. 2A**; **Table S1**)

Across all 8 genomes, each autosome has been completely assembled telomere-to-telomere (T2T) in at least one species (**Fig. 1E**). Within assemblies, 29%-70% of chromosomes are fully assembled with an average of less than one gap per chromosome (**Table S1**). Overall, these fully annotated genomes represent some of the most contiguous mammalian assemblies to date.

Evolution of structural variation within constrained karyotypes

With our near-complete assemblies, we first sought to study the landscape of structural variation within the tightly-conserved karyotype seen in *Myotis*. With only six exceptions across over 60 studied species[57–61], all *Myotis* species have a conserved 2n=42 karyotype - a remarkable phenomenon for a genus spread across six

continents and 20 million years of divergence[1,2]. The conserved *Myotis* karyotype consists of three large autosomes and one small metacentric autosome; 17 small telocentric autosomes; and metacentric X and Y chromosomes[50,62]. Consistent with this broad cytological conservation, we find large scale synteny across the Nearctic *Myotis* in this study (**Fig. 2A**). However, structural variants (SVs) including inversions, duplications, and translocations are relatively common within chromosomes, especially adjacent to putative centromeric regions (**Fig. 2A, B**). This includes a ~20Mb block at the sub-telomeric end of Chr V15 that displays frequent and recurrent inversions and translocations across the Nearctic *Myotis*, and which contains a 10Mb locus that was recently identified as a potential target of recent selection by adaptive introgression[56] (**Extended Data Fig. 3D**).

Using SyRI[63], we identified between 6,813 - 8,013 SVs per Nearctic *Myotis* genome relative to their outgroup, *M. myotis*; 97 - 99% of these events were under 10Kb. In the three large autosomes, which constitute ~30% of the genome, we cataloged an average of 509 SVs (**Table S3**). In contrast, in the small autosomes, constituting ~65% of the genome, we observed an average of 316 events, highlighting the distinct structural evolution between these chromosome types with SVs found at 2-fold higher density in larger autosomes (**Table S3**). However, large (≥10Kb) duplications, large inverted duplications, and large inverted translocations were more common on small autosomes compared to the large autosomes (**Table S3**). We also quantified the distribution of transposable elements (TEs) across chromosomes (**Fig. 2B,C**). Consistent with prior observations in other Chiropteran genomes[64], LINE elements appeared to be enriched around the predicted centromere (**Fig. 2B**) with simulations showing that this trend is primarily limited to the long, metacentric autosomes (**Extended Data Fig. 3A**; **Table S3**). The concentration of segmental duplications was also significantly correlated with TE density in each species (linear regression, p < 0.001; **Fig. 2B,D; Extended Data Fig. 3B**) highlighting the importance of TEs in facilitating structural evolution. Together these results highlight high levels of structural variation coincident with mobile element activity occurring in the context of a highly constrained karyotype.

An actively segregating, trans-species copy number polymorphism of the antiviral factor Protein Kinase R, PKR

Among the duplicated loci identified in Nearctic *Myotis*, the gene Protein Kinase R (*PKR*/*EIF2AK2*) stood out as an interferon-stimulated gene with antiviral activity against both DNA and RNA viruses. While the functionality of this *Myotis*-specific duplication had been previously characterized[28], the absence of long-read sequencing data for *Myotis* has prevented a more detailed characterization of copy number and evolutionary history among Neartic *Myotis.* Using our near-complete genome assemblies, we fully resolved the sequence and structure of the two previously known haplotypes: H1, containing a single copy of *PKR* (*PKR2*); and H2, containing two tandemly duplicated copies of *PKR* (*PKR1* and *PKR2*; **Fig. 3A**). We also identified a third haplotype - H3 - with three tandem duplicates of *PKR* (*PKR1*, *PKR2*, and a third copy) present only in *M. californicus*. Remarkably, while 7 out of 9 *Myotis* species carried duplicated haplotypes, 5 of these cases were heterozygous for the duplicated haplotype (i.e. H1/H2 or H1/H3; **Fig. 3B**). Two *Myotis* individuals (*lucifugus* and *evotis*) only carried non-duplicated (i.e. H1/H1; **Fig. 3B**). To determine the evolutionary history of the duplicates, we used GeneRax[65] to construct a tree from alignments of all *PKR* gene copies across Nearctic *Myotis*, using *Pipistrellus pygmaeus* as a non-*Myotis* outgroup (**Fig. 3C**). Our results suggest that *PKR2* is the ancestral copy of *PKR*, and that *PKR1* originated from a single duplication event at the root of *Myotis*. These results highlight that both the duplicated and unduplicated haplotypes have likely been segregating for tens of millions of years, representing an ancient trans-species polymorphism.

PKR is a stress response and innate immune factor that becomes active upon detection of dsRNA through sensing, auto-phosphorylation and dimerization[66,67]. Active PKR shuts down protein translation and restricts viral replication[66,67]. Although PKR1 and PKR2 are both expressed in basal or interferon-stimulated *Myotis velifer* cells, only their independent functional impacts have previously been investigated[28]. Given the co-expression and the complex haplotype diversity we identified in *Myotis*, we set out to determine whether the two *PKR* copies act additively, synergistically, or as dominant negatives in several essential functions. We therefore investigated the functional impact of the duplicates' co-expression on total PKR protein expression levels, cell viability, inhibition of translation, antiviral restriction and homo/hetero-dimer formation (**Fig. 3D-H**). To pinpoint the contribution of PKRs, we used a heterologous cell system[28] with PKR-KO Hela cells transfected with either an empty pSG5 plasmid (control), or pSG5 encoding PKR1, PKR2, or PKR1+PKR2 from *Myotis myotis* (myoMyo) or *Myotis velifer* (myoVel). In this system, PKR transfection leads to its autoactivation[28]. First, we investigated PKR protein expression and found that, although *Myotis* Flag-PKR1 was expressed at lower levels than Flag-PKR2, PKR1+PKR2 co-expression did not affect the overall PKRs' steady-state protein expression levels (**Extended Data Fig. 4A**).

Second, we performed cell viability assays at two doses of transfected PKRs and found that cell viability was only affected at high dose (**Extended Data Fig. 4D**). Third, using a luciferase reporter translational assay, we found that PKR1+PKR2 coexpression led to an intermediate translation shutdown, showing neither synergistic nor dominant negative effect of the PKR duplicates (**Fig. 3D**). Fourth, we tested PKR1+PKR2 co-expression effects on infections by two different model RNA viruses: VSV-GFP (Vesicular stomatitis virus encoding a GFP reporter[68], a representative of *Rhabdoviridae* family that is common in bats, including *Myotis*) and SINV-GFP (Sindbis virus, a representative of *Togaviridae*) (**Fig. 3E-F**, **Extended Data Fig. 4C,E**). We found that PKR1+PKR2 restricted viral infections to a similar, though slightly higher, extent as PKR1 or PKR2 alone (**Fig. 3E-F; Extended Data Fig. 4C-D**, p>0.05).   e recently duplicated PKR1s homodimerized, PKR1 and PKR2 did not form heterodimers (**Fig. 3G-H** and **Extended Data Fig. 4B** in SINV condition). This suggests that, since their duplication, divergence between *Myotis* PKR copies has been particularly strong at the protein-protein interface hindering their capacity to form heterodimers.

Together, these results suggest that PKR1 and PKR2 are neither dominant negative nor synergistic with each other, but rather act in a mostly additive form for their main cellular effector functions. Despite their capacity to more broadly escape some DNA and RNA viral antagonists[28], increased cell toxicity at high PKR dose suggests a possible tradeoff upon PKR duplication, which might participate in also explaining why PKR is rarely duplicated in mammals. Consistent with this, PKR1 is known to be expressed at significantly lower levels in *Myotis velifer* cells than PKR2 [28]. Altogether, these genomic and functional results unveil gene duplicates still segregating across several *Myotis* species, possibly shaped by recent viral infections and cell toxicity tradeoff.

Genome-wide adaptation to DNA viruses

To further explore how viruses may have shaped the genomes of *Myotis* bats we tested for adaptive signatures in virus-interacting proteins (VIPs) in *Myotis* and other bats. VIPs are host proteins that physically interact with viral proteins (e.g. *CD45*, **Fig. 4A**), and can be proviral (contributing to viral infection, e.g. viral receptors), antiviral (protective against viral infection, e.g. interferons), or both depending on infection stage and virus type. VIPs commonly have major roles in basic host function, and as a result most host cell biological pathways include VIPs. Previous studies investigating positive selection across mammals have found an enrichment for adaptation among a set of 5,527 manually curated VIPs, defined as host proteins that have at least one experimentally verified physical interaction with a viral protein, RNA, or DNA[69].

By calculating an enrichment score from the ratio of positive selection in VIPs compared to their matched control genes using BUSTED-MH[70], we found that, like other mammals, *Myotis* show an enrichment for adaptation at VIPs (**Fig. 4D; Table S5**). Physical host-virus interactions may not always result in fitness effects in the host. We therefore repeated our analysis using a gene set restricted to VIPs with experimental evidence of specific pro- or anti-viral effects, and thus with a stronger expectation of fitness effects. We observed an even stronger significant elevation in the ratio of positive selection in these proviral and antiviral VIPs (**Fig. 4B; Table S5**), but no elevation in this ratio in VIPs without experimental evidence of specific pro- or anti-viral effects (**Fig. 4C; Table S5**). This is consistent with the expectation of viral interaction as the cause of enrichment of positive selection in VIPs in bats[71]. We repeated this analysis using a dataset of 47 publicly-available non-*Myotis* bat genomes, and confirmed these same patterns across bats more broadly, even when excluding *Myotis* genomes (**Extended Data Fig. 5A**).

Previous work has suggested that bats may have different physiological responses to DNA and RNA viruses[72]. To determine if this was reflected in genomic VIP adaptation, we compared the enrichment of positive selection in VIPs that interact only with DNA viruses (DNA-only VIPs) to those that interact only with RNA viruses (RNA-only VIPs). Remarkably, we found that VIP adaptation in *Myotis* and other bats is driven by selection in DNA VIPs (**Fig. 4E, Extended Data Fig. 5B**) in marked contrast to the observed pattern in RNA VIPs, which show no evidence of genome-wide enrichment in adaptation (**Fig. 4F, Extended Data Fig. 5C**).

Adaptation to viral pathogens can also occur via gene copy number changes[73,74,75,76]. We tested whether VIPs were enriched among genes that were recently gained or lost throughout *Myotis* using the method of Huang *et al*. (2023) to test the cumulative per gene, per branch copy number changes and birth-death rates of VIPs compared to other non-VIP genes.[41] While the birth-death rate of VIP genes overall was no different than non-VIP genes (p = 0.071, **Extended Data Fig. 5D**), VIP genes were significantly more likely to have undergone expansions and/or contractions on at least one branch of the *Myotis* family (p < 0.001; **Fig. 4G**). Furthermore, we find that this pattern is driven exclusively by copy number changes in RNA-only VIPs, and not by changes in copy number of DNA-only VIPs (**Fig. 4H-I**). This suggests that there is variation in gene family birth rates across species, but that VIPs are more dynamic across the Nearctic *Myotis* as a whole than other types of genes highlighting the importance of the viral immune response in recent *Myotis* evolution.

In contrast to what we observe in bats, VIP adaptation in humans is driven by positive selection in RNA - and not DNA - VIPs[69,77]. To investigate if DNA VIP-driven adaptation in bats is exceptional among mammals, we replicated these analyses across four other large mammalian orders that are well represented among publicly-available mammalian genomes: *Primates*, *Glires*, *Euungulata*, and *Carnivora*. We found that while other mammalian orders show a mix of adaptation enrichments in both RNA and DNA VIPs, none show an absence of genome-wide enrichment of adaptation in RNA VIPs as observed in bats (**Extended Data Fig. 5A-C**). These results highlight that bats, including *Myotis*, may have faced greater selective pressures from DNA viruses than from RNA viruses, in contrast to all other mammals.

Evolution of Lifespan & Peto's Paradox across Nearctic *Myotis*

Beyond their remarkable immune adaptations, bats are exceptional as the longest-lived clade of mammal after correcting for their body size[17,18,78], demonstrating a 1.5-fold range of lifespans within a 3-fold range of body sizes[6,78,79]; *Myotis* species in particular exhibit the full dynamic range of bat lifespans within an even more narrow range of body sizes. While bats have been noted as an exception to the strong allometric scaling (positive

correlation with body size) of lifespan otherwise seen across mammals and other metazoans, this exception has not been tested using phylogenetically corrected statistics leveraging well-resolved phylogenies.

 To test the hypothesis of non-allometric scaling of lifespan in bats, we modeled the evolution of body size and lifespan independently across a supertree of over 1000 placental mammals (*Eutheria*)[54] (**Fig. 5A-B; Extended Data Fig. 6A-B**; **Table S6**). We generally observed agreement between evolutionary patterns of body size and lifespan in the two simulations, such as in whales (*Cetacea*)[80,81], elephantids (*Proboscidea*)[37,82,83], and primates (**Fig. 5A-B**). However, while only minor changes were observed in their body size, we observed some of the largest and most-rapid changes in lifespan across mammals in bats (**Fig. 5A-B; Extended Data Fig. 6A-B**; **Table S6**). These changes are largely independent between species and genera, consistent with the theory of multiple independent increases in lifespan across bats. This is especially true in *Myotis*, where we saw many of the fastest increases in lifespan relative to their most recent ancestor, including for *Myotis grisescens* (4.15x increase, 100th percentile), *Myotis brandtii* (2.25x increase, 100th percentile), *Myotis lucifugus* (1.56x, 98th percentile), *Myotis myotis* (1.1x increase, 79nd percentile), and the *Myotis* common ancestor (1.26x increase, 92rd percentile) (**Fig. 5B, Extended Data Fig. 6B**).

 We used phylogenetically-corrected generalized linear models and ANCOVA to quantify the relationship between body size and lifespan across mammals. We find that while non-bat mammals experience an average 0.159% increase in lifespan per 1% increase in body size, bats experience a 0.223% increase in lifespan years per 1% increase in body size, though these rates were not significantly different after phylogenetic correction (**Extended Data Fig. 6C-D**; pANCOVA: p=0.29). Together these results demonstrate that *Myotis* bats and their recent ancestors have evolved some of the most extreme increases in lifespan among mammals, despite similar allometric lifespan scaling of bats and mammals after phylogenetic correction.

 Rapid changes in body size and lifespan can have major implications for the evolution of cancer risk and resistance across mammals. Within a species, lifetime *cancer risk* scales linearly with body size, and with lifespan by a power-law of exponent 6[84–87]. In contrast to a strong *within*-species correlation, when comparing *between*-species, there is only a weak correlation between body size and cancer risk - and no correlation between lifespan and cancer risk[87–91] - a phenomenon known as Peto's Paradox. This observation suggests that species with more cells or longer lifespans have adapted to reduce their cancer risks (i.e. increased cancer resistance) (**Fig. 5C**). We hypothesized that the very rapid evolution of increased lifespan in *Myotis* would thus result in a dramatic increase in their expected cancer risk compared to other mammals. Cancer risk can be explicitly measured as the prevalence of neoplasias in a species; however, there are very limited records for neoplasias in the literature, and none available for *Myotis* and other bats in family *Vespertilliodinae*[90,91].

 An alternate approach is to quantify the Relative Intrinsic Cancer Risk (RICR) between extant mammals and their direct ancestors, calculated as the $\log_2$ ratio of cancer risk between the two nodes (**Fig. 5C,D**)[37,87]. In a scenario where a species has increased in either body size or lifespan relative to its direct ancestor, we would observe an increase in RICR corresponding to an increase in their potential cancer risk; this increase, in turn, is predicted to impart a proportional selective pressure on genes in cancer-related pathways to increase cancer resistance. We used our evolutionary estimates of body size and lifespan across *Eutheria* to quantify RICR across the placental mammals (**Fig. 5D**)[37], and found that bats were overrepresented in the top 10% of RICR (odds ratio 1.15), highlighting the impact of rapid lifespan evolution on cancer risk. This is consistent with recent pan-mammal and pan-vertebrate studies of cancer risk[90,91], which found that bats were present among the 20 species with the lowest rates of neoplasia, and not at all among the 20 species with the highest rates of neoplasia[91]. The longest-lived *Myotis* (*M. grisescens*: 39 yrs, 1st percentile; *M. brandtii*: 42 yrs, 2nd percentile; *M. lucifugus*: 36 yrs, 4th pct) and their most recent common ancestors (*lucifugus-occultus*: ~26 yrs, 8th pct;

Myotis common ancestor: ~22 yrs, 14th pct) demonstrated some of the most pronounced increases in RICR among mammals (**Fig. 5D; Table S6**). Similar to other extreme cases of body size and lifespan in vertebrates[37,39,44,46,92–96], the pronounced changes in RICR seen in *Myotis* imply strong selective pressure to evolve cancer resistance mechanisms at multiple points across *Chiroptera* in general, and within *Myotis* in particular.

Evolutionary signatures of cancer resistance and pleiotropy with DNA virus adaptations in *Myotis*

We hypothesized that the extreme changes in RICR observed throughout the *Myotis* phylogeny should exert a selective pressure on genes associated with cancer processes. Using aBSREL[97] to test for selection at terminal and internal branches within *Myotis*, we found that at FDR<=5%, an average of 5.46% of protein-coding genes had at least one region under positive selection per node (omega >1); in contrast, an average of 20.8% of protein-coding genes per node had all regions with an omega less than 1, suggesting negative selection (**Table S7**). Genes under selection were enriched for several pathways involved in immunity, cancer, and aging (**Table S7**), with many sitting at the intersection of multiple processes, including members of the Cluster of Differentiation (CD) family, Serpin family, insulin signaling pathway, redox repair, and iron storage (**Extended Data Fig. 6E; Table S7**), suggesting possible pleiotropic influences on genes under selection.

We quantified the contribution of genes under selection to pathways associated with the hallmark of cancer[98–100] by measuring the proportion of cancer-associated pathways overrepresented among genes under selection throughout the phylogeny (**Fig. 5E**). Consistent with the observation that most nodes within *Myotis* demonstrate increases in RICR, we observed that most nodes within nearctic *Myotis* were enriched for cancer hallmark pathways among genes under selection, especially at the recent ancestors of the longest-lived species (e.g. *M. lucifugus, M. occultus*; **Fig. 5E**). Furthermore, when considering the union of all genes under selection in the evolutionary history of each species since their common ancestor, we observed significant enrichments in the representation of cancer-associated pathways in many of the lineages with the greatest increases in RICR (*M. lucifugus, M. occultus, M. evotis, M. yumanensis;* **Fig. 5F, Extended Data Fig. 6F**). This suggests that selection in genes involved in cancer-related processes occurred concurrently with the recurrent increases in lifespan and RICR seen in this clade.

The longest-lived bat in our study, *M. lucifugus*, had an overrepresentation of pathways specifically associated with DNA double-strand break (DSB) repair when looking at both lineage-wide and node-specific enrichments in positive selection using the Reactome database[101] (**Fig. 5G; Table S7**). This includes 35 out of 65 genes in the high-fidelity Homologous Recombination Repair pathway, and 21/37 members of the Homology-Directed Repair via Single Strand Annealing (**Fig. 5G; Extended Data Fig. 6E; Table S7**). These results suggest that *M. lucifugus* might have an enhanced response to DNA DSBs relative to other bats. To test this hypothesis, we assessed the tolerance of *M. lucifugus* to neocarzinostatin, a potent radiomimetic agent that induces DNA double-strand breaks (**Fig. 5H**), compared to *M. evotis*, three non-*Myotis* bats (*Eidolon helvum, Pteropus rodrigensis,* and *Rousettus lanosus*). At low doses of neocarzinostatin, *M. lucifugus* was the only species tested showing sensitivity to neocarzinostatin after 24 hours, with a drop in viability and concomitant increase in apoptosis. At high doses, *M. lucifugus* had the highest level of apoptosis and the greatest drop in viability of all the bats tested. This is consistent with other long-lived species, including elephants[35,36,102], naked mole rats[45], and bowhead whales[39,103], where longevity and RICR are associated with an increased ability to clear out damaged cells.

To better understand the genes that may be driving this difference, we examined RNA expression in *M. lucifugus* cells after 6 and 18 hours of treatment with 100uM NCS. As expected, at 6 hours we observe up-regulation of genes associated with cell cycle arrest (e.g. *MDM2, CDKN1A*) and cell death (e.g. *BAX*),and down-regulation of genes in pathways associated with cell division and growth at 6 hours. At 18 hours, we observe up-regulation of genes in pathways associated with cellular stress response and continued down-regulation of the cell cycle (**Fig. 6A-D, Extended Data Fig. 7A, Table S9**). At 6 hours, however, we also observed down-regulation of genes associated with DNA damage repair and synthesis, which together with the up-regulation of *BAX* is consistent with the observed sensitivity to NCS in *M. lucifugus*.

Unexpectedly, however, we observed that the strongest pathway enrichment in our DNA-damage RNA-seq data were genes associated with the innate immune response, including genes in the PI3K/AKT pathway (**Fig. 6C-D, Table S9**). We hypothesized that the observed relationship between the unique DNA damage response of *M. lucifugus* may be due to pleiotropy with DNA VIPs and selection in these genes. Indeed, we observed that there was a highly significant enrichment for genes that are differentially expressed after NCS treatment and DNA-only VIPs (p=3.34e-5, **Fig. 6E**; α=0.0245, **Extended Data Fig. 7B**). The genes at the intersection of DNA-only VIPs and NCS response are overrepresented in pathways associated with the cell cycle, senescence, transcription, and DNA damage repair (**Fig. 6F**). To test how selection may have impacted these pleiotropic signatures, we repeated these analyses only considering genes under selection in *Myotis lucifugus*. The intersection between DNA-only VIPs and NCS-DE genes remained highly significant (p=1.88e-3, **Extended Data Fig. 7C**) with only three pathways remaining enriched at FDR<=0.05: Cell Cycle, DNA Damage Repair, and Deubiquination (**Fig. 6G**); DNA-only VIP genes which are not NCS-responsive did not show these signatures (**Extended Data Fig. 7D-G).** Together these results suggest that selection on the innate immune response may drive *agonistic* pleiotropy in aging-associated traits like DNA damage response and cell cycle regulation.

Discussion

Bats are widely known for their long lifespan, cancer resistance, and viral tolerance[6,10,11,43,78,90,104–106]. As highly complex and pleiotropic processes, the genes and mechanisms underlying these phenotypes can be challenging to identify, especially for rapidly-evolving phenotypes such as host-pathogen interactions. Here we outline an approach that enables functional comparative biology by generating cell lines from wing punches of wild caught bats for genome assembly, comparative genomics, and functional follow up. Cell lines are generated from minimally-invasive biopsies collected in the field thus avoiding disturbing natural populations. Using this approach, we identify new patterns of adaptation contributing to longevity, cancer resistance, and viral interactions in bats, and demonstrate a unique DNA damage response in primary cells of the long-lived *M. lucifugus.* Given the high density of bat species concentrated at single locations world-wide[107,108] it is feasible to collect wing punches from a large number of individuals across a wide phylogenetic range; these wing punches can be used to generate cell lines and sequencing libraries for reference genomes in a matter of weeks. This approach is useful to expand genetic resources across the tree of life[109–111] and for conservation genomics. As this approach can generate genomic resources from minimal material gathered via non-lethal sampling, it is also well-suited for the study of rare or endangered species for which acquiring sufficient amounts of material can be challenging.

The evolution of body size and lifespan across mammals - and the rapid evolution of lifespan in *Yinpterochiroptera* in particular - has major implications for the co-evolution of cancer risk and resistance. While

models of body size evolution are well-studied in mammals[7,37,82,112] the evolution of lifespan is less well understood. We find that *Myotis* demonstrates an increased rate of change in lifespan given body size compared to other mammals. This altered scaling of longevity in *Myotis* has dramatic consequences for their intrinsic, per-cell cancer risk and for the evolution of tumor-suppressor genes and pathways. While cancer risk scales linearly with body size, it scales with age as a power law of $6^{84,87,88}$. Meanwhile, while mammalian body sizes span a 10^6 range of masses, they only span a 10^2 range of lifespans[16,113]. Unlike other systems where the evolution of cancer resistance has been driven by rapid changes in body size[35–37,44,92,96], the body size of *Myotis* has not significantly changed since their common ancestor. Instead, the rapid and repeated changes in lifespan across an order of magnitude in *Myotis* lead to some of the most significant changes in intrinsic cancer risk seen across mammals.

Bats are also important models for understanding infectious disease response and immune adaptation because of their unique resistance to viruses and role as zoonotic reservoirs[31,72,114]. Here, we show that while bats have adapted to both DNA and RNA viruses, they demonstrate a remarkable dominance in adaptations to DNA viruses. This is in contrast to humans and other primates, in which adaptation to RNA viruses dominates[49,77]. While bats are known to mount adaptive immune responses to some RNA viruses, this suggests that adaptations in bats' innate and adaptive immune systems, such expansions in *MHC-I* and *APOBEC3*, or the loss of the *PYHIN* gene family[48], may be sufficient as a first line of defense to protect against the fitness impacts of some RNA viruses, but not against DNA viruses. This is consistent with previous observations that most zoonoses, including those hosted by bats, are RNA viruses[10] and that bats are more likely than other mammals to asymptomatically harbor RNA viruses, while being more susceptible themselves to other pathogens such as fungi[72]. We also recapitulate and expand on previous results related to positive selection in immune genes and immune gene family expansion, contraction, and loss. In addition, we demonstrate complex patterns of structural variation in immune genes, including a segregating duplication of the Protein Kinase R gene (*PKR*), a major protein involved in the antiviral innate immune system, that has functional relevance in its activity against both DNA and RNA viruses[28]. By resolving individual haplotypes in nine *Myotis* species, we were able to confirm a single duplication event at the origin of *Myotis PKR1* and *PKR2*. We further demonstrate the functional implications of copy number variation in *PKR*, including differences in copy expression and an additive effect on cellular effector functions.

Multiple hypotheses have been proposed to connect the unique physiology and ecology of bats with the evolution of remarkable adaptations such as viral infection tolerance, stress tolerance, and exceptional longevity[104]. Hypothesized drivers of disease resistance and longevity evolution in bats include the evolution of flight (e.g. "flight as fever" hypothesis[115], though this hypothesis has recently been critiqued[116]), the disposable soma hypothesis[117]; metabolic state[118]; torpor[6]; and other adaptations to specific environments[9,119–121]. Additionally, many studies have highlighted the intersection of these traits, including a relationship between hibernation and both longevity[6,114] and infectious disease resistance[72,114]. Our results are consistent with an *agonistic* pleiotropy hypothesis, wherein genetic adaptations for many specific traits (e.g. DNA virus innate immunity) may prove beneficial to other seemingly-unrelated traits (e.g. cancer resistance, cellular homeostasis, longevity). Importantly, bats are also the only mammal to harbor active DNA transposons, the suppression of which may also contribute to signatures of selection in DNA-interacting VIPs.

Consistent with the agonistic pleiotropy hypothesis, many of the genes and pathways highlighted in this study have been found to play vital roles across physiological traits in bats and other species. For example, two genes under selection in nearctic *Myotis* - *FTH1* and *IGFN1* - have been implicated in functional studies as key hibernation genes[122–124], viral interacting proteins[125–128], and as pro-longevity genes[129–131]. Similarly, many DNA VIPs such as *BRCA1/2* and *POLG* are core DNA maintenance genes essential for cancer resistance and

longevity[45,132–138]. Such essential genes are generally under increased constraint; however, it is possible that external pressures such as viral infection and active DNA transposable elements, are shaping the fitness landscape of these essential genes such that novel optimizations become accessible to evolutionary processes[139]. Beyond individual genes, many of the overarching pathways under selection in *Myotis*, such as those associated with inflammation, senescence, and ferroptosis lie directly at the intersection of aging-related immune processes[43,48,124,129,140–147]. While together our results suggest the possibility that traits such as cancer risk, cellular homeostasis, and antiviral response have evolved in tandem due to pleiotropic selection at overlapping points in bats' evolutionary histories, further functional validation will be required to disentangle the functional impacts of these genetic changes and disambiguate the drivers of selection.

Acknowledgements

PacBio HiFi sequencing was done by the DNA Technologies and Expression Analysis Cores at the UC Davis Genome Center (NIH SIG 1S10OD010786-01). Analyses were performed using the High Performance Computing (HPC) resources at the University of Arizona (supported by the University of Arizona TRIF, UITS, and Research, Innovation, and Impact (RII); maintained by the UArizona Research Technologies department); the Savio computational cluster resource provided by the Berkeley Research Computing program at the University of California, Berkeley (supported by the UC Berkeley Chancellor, Vice Chancellor for Research, and Chief Information Officer); and the Vermont Advanced Computing Core at . We thank the members of the LP2L team (CIRI) for helpful discussions.

MEL was funded by NSF PRFB #2010884, and a Dovetail Tree of Life Grant. JMV was funded by NSF PRFB #2109915, NIH NIA T32AG000266, and NIA 1K99AG088361. DE was funded by NIH NIGMS MIRA grant 5R35GM142677. PHS was funded by NIH NIGMS R35GM142916 and by the Vallee Scholars Award. LE was funded by a grant from the Agence Nationale de la Recherche (ANR, #ANR-202-CE15-0020-01 to LE) and CNRS MITI These Internationale (to M.S. and L.E.). SP is funded by the ANR (ANR-21-CE35-0018-01) and by the Interdisciplinary Thematic Institute IMCbio+ supported by IdEx Unistra (ANR-10-IDEX-0002), by SFRI-STRAT'US project (ANR-20-SFRI-0012) and EUR IMCBio (IMCBio ANR-17-EURE-0023) under the framework of the French Investments for the Future Program. LE and DE were further funded by a grant from the Joint Call for Proposals between the CNRS and the University of Arizona (International Research Center, IRC, 2021-2024). LE, DE, PHS and SP were further funded by the International Research Project (IRP) RAPIDvBAT from the CNRS, the University of Arizona and the University of California, Berkeley. LG was funded by a postdoctoral fellowship from the Fondation pour la Recherche Médicale (FRM, #SPF202209015746).

Image attributions for the bat photos featured in Fig. 1: J. Scott Altenbach (*M. thysanodes*); M. Elise Lauterbur (*M. occultus, M. velifer*); U.S. National Park Service (*M. auriculus*); Frank Carey (*M. californicus*); Michael Durham (*M. evotis*); J. N. Stuart (*M. volans*); and SMBishop & the Wikimedia Foundation (*M. lucifugus*).

Author Contributions:

Conceptualization, J.M.V., M.E.L., D.E., P.H.S., L.E., D.F., M.B.; Methodology, J.M.V., M.E.L., D.E., P.H.S., L.E., G.G.S.; Software, J.M.V., M.E.L., D.E., P.H.S., L.E., T.M.L.; Validation, J.M.V., M.E.L., D.E., P.H.S., L.E., J.P.V.M.; Formal Analysis, J.M.V., M.E.L., D.E., P.H.S., L.E., S.M., A.L.C., D.M., S.V.; Investigation, J.M.V., M.E.L., S.M., M.B., D.F., G.G.S., L.G., Z.R.H., M.H., W.K., T.M.L., A.L.C., C.L. S.M., D.M., S.L., J.L., C.R., S.L.R.C., M.S., K.S., W.T., J.D.T., S.V., R.M., M.B., J.P.V.M., S.P.,

L.E., D.E., P.H.S.; Resources, J.M.V., M.E.L., D.E., P.H.S., L.E., M.B., D.F., G.G.S., L.G., Z.R.H., M.H., W.K., T.M.L., A.L.C., S.M., D.M., S.L., J.L., C.R., S.L.R.C., M.S., K.S., W.T., J.D.T., S.V., R.M., M.B.; Data Curation, J.M.V., M.E.L., S.M., M.B., D.F., G.G.S., L.G., Z.R.H., M.H., W.K., T.M.L., A.L.C., S.M., D.M., S.L., J.L., C.R., S.L.R.C., M.S., K.S., W.T., J.D.T., S.V., R.M., M.B., J.P.V.M., S.P., L.E., D.E., P.H.S.; Writing - Original Draft, J.M.V., M.E.L., D.E., P.H.S., L.E., S.M., J.L.; Writing - Review & Editing, J.M.V., M.E.L., S.M., M.B., D.F., G.G.S., L.G., Z.R.H., M.H., W.K., T.M.L., A.L.C., S.M., D.M., S.L., J.L., C.R., S.L.R.C., M.S., K.S., W.T., J.D.T., S.V., R.M., M.B., J.P.V.M., S.P., L.E., D.E., P.H.S.; Visualization, J.M.V., M.E.L., D.E., P.H.S., L.E., S.M., A.L.C.; Supervision, J.M.V., M.E.L., D.E., P.H.S., L.E.; Project Administration, J.M.V., M.E.L., D.E., P.H.S., L.E.; Funding Acquisition, J.M.V., M.E.L., D.E., P.H.S., L.E.

Declaration of Interests:

The authors declare no competing interests.

References

1. Wilson, D. E. & Reeder, D. M. *Mammal Species of the World: A Taxonomic and Geographic Reference*. (JHU Press, 2005).

2. Simmons, N. B. & Cirranello, A. L. Bat Species of the World: A Taxonomic and Geographic Database. https://batnames.org/.

3. Teeling, E. C., Hedges, S. B. & Kumar, S. Bats (Chiroptera). *The timetree of life* 499–503 (2009).

4. Rietbergen, T. B. *et al.* The Oldest Known Bat Skeletons and Their Implications for Eocene Chiropteran Diversification. *PLoS One* **18**, e0283505 (2023).

5. Kumar, S., Stecher, G., Suleski, M. & Hedges, S. B. TimeTree: A Resource for Timelines, Timetrees, and Divergence Times. *Mol. Biol. Evol.* **34**, 1812–1819 (2017).

6. Wilkinson, G. S. & Adams, D. M. Recurrent Evolution of Extreme Longevity in Bats. *Biol. Lett.* **15**, 20180860 (2019).

7. Moyers Arévalo, R. L., Amador, L. I., Almeida, F. C. & Giannini, N. P. Evolution of Body Mass in Bats: Insights from a Large Supermatrix Phylogeny. *J. Mamm. Evol.* **27**, 123–138 (2020).

8. Datzmann, T., von Helversen, O. & Mayer, F. Evolution of Nectarivory in Phyllostomid Bats (Phyllostomidae Gray, 1825, Chiroptera: Mammalia). *BMC Evol. Biol.* **10**, 165 (2010).

9. Camacho, J. *et al.* Sugar assimilation underlying dietary evolution of Neotropical bats. *bioRxiv*

2023.07.02.547432 (2023) doi:10.1101/2023.07.02.547432.

10. Hayman, D. T. S. *et al.* Ecology of zoonotic infectious diseases in bats: current knowledge and

future directions. *Zoonoses Public Health* **60**, 2–21 (2013).

11. Irving, A. T., Ahn, M., Goh, G., Anderson, D. E. & Wang, L.-F. Lessons from the host defences of

bats, a unique viral reservoir. *Nature* **589**, 363–370 (2021).

12. Morales, A. E., Ruedi, M., Field, K. & Carstens, B. C. Diversification rates have no effect on the

convergent evolution of foraging strategies in the most speciose genus of bats, Myotis. *Evolution*

73, 2263–2280 (2019).

13. Gunnell, G. F., Smith, R. & Smith, T. 33 Million Year Old Myotis (Chiroptera, Vespertilionidae) and

the Rapid Global Radiation of Modern Bats. *PLoS One* **12**, e0172621 (2017).

14. Ruedi, M. *et al.* Molecular Phylogenetic Reconstructions Identify East Asia as the Cradle for the

Evolution of the Cosmopolitan Genus Myotis (Mammalia, Chiroptera). *Mol. Phylogenet. Evol.* **69**,

437–449 (2013).

15. Tacutu, R. *et al.* Human Ageing Genomic Resources: Integrated Databases and Tools for the

Biology and Genetics of Ageing. *Nucleic Acids Res.* **41**, D1027–D1033 (2013).

16. Jones, K. E. *et al.* PanTHERIA: A Species-Level Database of Life History, Ecology, and

Geography of Extant and Recently Extinct Mammals. *Ecology* **90**, 2648–2648 (2009).

17. Austad, S. N. Methusaleh's Zoo: How Nature Provides Us with Clues for Extending Human Health

Span. *J. Comp. Pathol.* **142**, S10–S21 (2010).

18. Austad, S. N. & Fischer, K. E. Mammalian Aging, Metabolism, and Ecology: Evidence from the

Bats and Marsupials. *J. Gerontol.* **46**, B47–53 (1991).

19. Podlutsky, A. J., Khritankov, A. M., Ovodov, N. D. & Austad, S. N. A new field record for bat

longevity. *J. Gerontol. A Biol. Sci. Med. Sci.* **60**, 1366–1368 (2005).

20. Wilson, D. E. & Tyson, E. L. Longevity Records for Artibeus Jamaicensis and Myotis Nigricans. *J.*

Mammal. **51**, 203 (1970).

21. Stadelmann, B., Lin, L.-K., Kunz, T. H. & Ruedi, M. Molecular Phylogeny of New World

Myotis (Chiroptera, Vespertilionidae) Inferred from Mitochondrial and Nuclear DNA Genes.

Mol. Phylogenet. Evol. **43**, 32–48 (2007).

22. Agnarsson, I., Zambrana-Torrelio, C. M., Flores-Saldana, N. P. & May-Collado, L. J. A Time-

Calibrated Species-Level Phylogeny of Bats (Chiroptera, Mammalia). _PLoS Curr._ **3**, RRN1212

(2011).

23. Armero, A. _et al._ Myotis Fimbriatus Virome, a Window to Virus Diversity and Evolution in the Genus

Myotis. _Viruses_ **14**, 1899 (2022).

24. He, X. _et al._ Establishment of Myotis Myotis Cell Lines - Model for Investigation of Host-Pathogen

Interaction in a Natural Host for Emerging Viruses. _PLoS One_ **9**, e109795 (2014).

25. Seltmann, A. _et al._ Differences in Acute Phase Response to Bacterial, Fungal and Viral Antigens in

Greater Mouse-Eared Bats (Myotis Myotis). _Sci. Rep._ **12**, 15259 (2022).

26. Hayward, J. A. _et al._ Unique Evolution of Antiviral Tetherin in Bats. _J. Virol._ **96**, e0115222 (2022).

27. Fernandes, A. P., Águeda-Pinto, A., Pinheiro, A., Rebelo, H. & Esteves, P. J. Evolution of TRIM5

and TRIM22 in Bats Reveals a Complex Duplication Process. _Viruses_ **14**, (2022).

28. Jacquet, S. _et al._ Adaptive duplication and genetic diversification of protein kinase R contribute to

the specificity of bat-virus interactions. _Sci Adv_ **8**, eadd7540 (2022).

29. Jacquet, S., Pontier, D. & Etienne, L. Rapid Evolution of HERC6 and Duplication of a Chimeric

HERC5/6 Gene in Rodents and Bats Suggest an Overlooked Role of HERCs in Mammalian

Immunity. _Front. Immunol._ **11**, 605270 (2020).

30. Chomel, B., Stuckey, M., Boulouis, H. & Aguilar Setién, Á. Bat-Related Zoonoses. _Zoonoses -_

Infections Affecting Humans and Animals 697–714 (2014) doi:10.1007/978-94-017-9457-2_28.

31. Guth, S. _et al._ Bats host the most virulent-but not the most dangerous-zoonotic viruses. _Proc. Natl._

Acad. Sci. U. S. A. **119**, e2113628119 (2022).

32. Williams, E. P. _et al._ Common themes in zoonotic spillover and disease emergence: Lessons

learned from bat- and rodent-borne RNA viruses. _Viruses_ **13**, 1509 (2021).

33. Mollentze, N. & Streicker, D. G. Viral zoonotic risk is homogenous among taxonomic orders of

mammalian and avian reservoir hosts. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 9423–9430 (2020).

34. Kolora, S. R. R. *et al.* Origins and Evolution of Extreme Life Span in Pacific Ocean Rockfishes.

Science **374**, 842–847 (2021).

35. Sulak, M. *et al.* Correction: TP53 copy number expansion is associated with the evolution of

increased body size and an enhanced DNA damage response in elephants. *Elife* **5**, (2016).

36. Vazquez, J. M., Sulak, M., Chigurupati, S. & Lynch, V. J. A Zombie LIF Gene in Elephants Is

Upregulated by TP53 to Induce Apoptosis in Response to DNA Damage. *Cell Rep.* **24**, 1765–1776

(2018).

37. Vazquez, J. M. & Lynch, V. J. Pervasive duplication of tumor suppressors in Afrotherians during

the evolution of large bodies and reduced cancer risk. *Elife* **10**, e65041 (2021).

38. Davies, K. T. J. *et al.* Molecular Evolution of Growth Hormone and Insulin-like Growth Factor 1

Receptors in Long-Lived, Small-Bodied Mammals. *Gene* **549**, 228–236 (2014).

39. Vazquez, J. M., Kraft, M. & Lynch, V. J. A CDKN2C retroduplication in Bowhead whales is

associated with the evolution of extremely long lifespans and alerted cell cycle dynamics. *bioRxiv*

2022.09.07.506958 (2022) doi:10.1101/2022.09.07.506958.

40. Foote, A. D. *et al.* Convergent evolution of the genomes of marine mammals. *Nat. Genet.* **47**, 272–

275 (2015).

41. Huang, Z. *et al.* Duplications of Human Longevity-Associated Genes Across Placental Mammals.

Genome Biol. Evol. **15**, (2023).

42. Li, S., Vazquez, J. M. & Sudmant, P. H. The evolution of aging and lifespan. *Trends Genet.* **39**,

830–843 (2023).

43. Jebb, D. *et al.* Six reference-quality genomes reveal evolution of bat adaptations. *Nature* **583**, 578–

584 (2020).

44. Glaberman, S., Bulls, S. E., Vazquez, J. M., Chiari, Y. & Lynch, V. J. Concurrent evolution of

antiaging gene duplications and cellular phenotypes in long-lived turtles. *Genome Biol. Evol.* **13**,

evab244 (2021).

45. MacRae, S. L. *et al.* DNA Repair in Species with Extreme Lifespan Differences. *Aging* **7**, 1171–1184 (2015).

46. Baines, C., Meitern, R., Kreitsberg, R. & Sepp, T. Comparative study of the evolution of cancer gene duplications across fish. *Evol. Appl.* **15**, 1834–1845 (2022).

47. López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The Hallmarks of Aging. *Cell* **153**, 1194–1217 (2013).

48. Moreno Santillán, D. D. *et al.* Large-scale genome sampling reveals unique immunity and metabolic adaptations in bats. *Mol. Ecol.* **30**, 6449–6467 (2021).

49. Enard, D. & Petrov, D. A. Ancient RNA virus epidemics through the lens of recent adaptation in human genomes. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **375**, 20190575 (2020).

50. Sotero-Caio, C. G., Baker, R. J. & Volleth, M. Chromosomal evolution in Chiroptera. *Genes (Basel)* **8**, (2017).

51. Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**, 3210–3212 (2015).

52. Manni, M., Berkeley, M. R., Seppey, M. & Zdobnov, E. M. BUSCO: assessing genomic data quality and beyond. *Current Protocols* **1**, (2021).

53. Amador, L. I., Moyers Arévalo, R. L., Almeida, F. C., Catalano, S. A. & Giannini, N. P. Bat Systematics in the Light of Unconstrained Analyses of a Comprehensive Molecular Supermatrix. *J. Mamm. Evol.* **25**, 37–70 (2018).

54. Upham, N. S., Esselstyn, J. A. & Jetz, W. Inferring the mammal tree: Species-level sets of phylogenies for questions in ecology, evolution, and conservation. *PLoS Biol.* **17**, e3000494 (2019).

55. Korstian, J. M., Paulat, N. S., Platt, R. N., Stevens, R. D. & Ray, D. A. SINE-Based Phylogenomics Reveal Extensive Introgression and Incomplete Lineage Sorting in Myotis. *Genes* **13**, 399 (2022).

56. Foley, N. M. *et al.* Karyotypic stasis and swarming influenced the evolution of viral tolerance in a

579   species-rich bat radiation. *Cell Genom* **4**, 100482 (2024).

57. Volleth, M. Variations on a theme: Karyotype comparison in Eurasian Myotis species and implications for phylogeny. *Vespertilio* **16**, 329–350 (2012).

58. McBee, K. Standard karyology of nine species of vespertilionid bats (Chiroptera: Vespertilionidae) from Thailand. *Annals of Carnegie Museum* **55**, 95–116 (1986).

59. Bickham, J. W., McBee, K. & Schlitter, D. A. Chromosomal variation among seven species of Myotis (Chiroptera: Vespertilionidae). *J. Mammal.* **67**, 746–750 (1986).

60. Zhang, W. D. A study on karyotype of Myotis chinensis and M. laniger Peter. *J Anhui Normal Univ* **7**, 42–47 (1984).

61. Zima, J. Synopsis of karyotypes of vespertilionid bats (Mammalia: Chiroptera). *Acta Univ. Carol. Biol* **1981**, 311–329 (1985).

62. O'Brien, S. J., Menninger, J. C. & Nash, W. G. *Atlas of Mammalian Chromosomes*. (John Wiley & Sons, 2006).

63. Goel, M., Sun, H., Jiao, W.-B. & Schneeberger, K. SyRI: finding genomic rearrangements and local sequence differences from whole-genome assemblies. *Genome Biol.* **20**, 277 (2019).

64. de Sotero-Caio, C. G. *et al.* Centromeric enrichment of LINE-1 retrotransposons and its significance for the chromosome evolution of Phyllostomid bats. *Chromosome Res.* **25**, 313–325 (2017).

65. Morel, B., Kozlov, A. M., Stamatakis, A. & Szöllősi, G. J. GeneRax: A Tool for Species-Tree-Aware Maximum Likelihood-Based Gene Family Tree Inference under Gene Duplication, Transfer, and Loss. *Mol. Biol. Evol.* **37**, 2763–2774 (2020).

66. Kaufman, R. J. Double-stranded RNA-activated protein kinase mediates virus-induced apoptosis: a new role for an old actor. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 11693–11695 (1999).

67. García, M. A. *et al.* Impact of protein kinase PKR in cell biology: from antiviral to antiproliferative action. *Microbiol. Mol. Biol. Rev.* **70**, 1032–1060 (2006).

68. Ostertag, D., Hoblitzell-Ostertag, T. M. & Perrault, J. Overproduction of double-stranded RNA in

vesicular stomatitis virus-infected cells activates a constitutive cell-type-specific antiviral response. *J. Virol.* **81**, 503–513 (2007).

69. Enard, D., Cai, L., Gwennap, C. & Petrov, D. A. Viruses are a dominant driver of protein adaptation in mammals. *Elife* **5**, (2016).

70. Murrell, B. *et al.* Gene-wide identification of episodic selection. *Mol. Biol. Evol.* **32**, 1365–1371 (2015).

71. Souilmi, Y. *et al.* An ancient viral epidemic involving host coronavirus interacting genes more than 20,000 years ago in East Asia. *Curr. Biol.* **31**, 3704 (2021).

72. Brook, C. E. & Dobson, A. P. Bats as 'special' reservoirs for emerging zoonotic pathogens. *Trends Microbiol.* **23**, 172–180 (2015).

73. Kondrashov, F. A. Gene Dosage and Duplication. in *Evolution after Gene Duplication* (eds. Dittmar, K. & Liberles, D.) 57–76 (John Wiley & Sons, 2011).

74. Kondrashov, F. A., Rogozin, I. B., Wolf, Y. I. & Koonin, E. V. Selection in the evolution of gene duplications. *Genome Biol.* **3**, RESEARCH0008 (2002).

75. Rastogi, S. & Liberles, D. A. Subfunctionalization of duplicated genes as a transition state to neofunctionalization. *BMC Evol. Biol.* **5**, 28 (2005).

76. Assis, R. & Bachtrog, D. Neofunctionalization of young duplicate genes in Drosophila. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 17409–17414 (2013).

77. Enard, D. & Petrov, D. A. Evidence that RNA Viruses Drove Adaptive Introgression between Neanderthals and Modern Humans. *Cell* **175**, 360–371.e13 (2018).

78. Brunet-Rossinni, A. K. & Austad, S. N. Ageing studies on bats: a review. *Biogerontology* **5**, 211–222 (2004).

79. Wilkinson, G. S. & South, J. M. Life History, Ecology and Longevity in Bats. *Aging Cell* **1**, 124–131 (2002).

80. Pyenson, N. D. & Sponberg, S. N. Reconstructing body size in extinct crown Cetacea (neoceti) using allometry, phylogenetic methods and tests from the fossil record. *J. Mamm. Evol.* **18**, 269–

288 (2011).

81. Montgomery, S. H. *et al.* The evolutionary history of cetacean brain and body size: Cetacean brain evolution. *Evolution* **67**, 3339–3353 (2013).

82. Puttick, M. N. & Thomas, G. H. Fossils and living taxa agree on patterns of body mass evolution: a case study with Afrotheria. *Proc. Biol. Sci.* **282**, 20152023 (2015).

83. Sulak, M. *et al.* TP53 copy number expansion is associated with the evolution of increased body size and an enhanced DNA damage response in elephants. *Elife* **5**, (2016).

84. Armitage, P. Multistage Models of Carcinogenesis. *Environ. Health Perspect.* **63**, 195–201 (1985).

85. Armitage, P. & Doll, R. The Age Distribution of Cancer and a Multi-Stage Theory of Carcinogenesis. *Br. J. Cancer* **91**, 6602297 (2004).

86. Peto, R., Roe, F. J., Lee, P. N., Levy, L. & Clack, J. Cancer and ageing in mice and men. *Br. J. Cancer* **32**, 411–426 (1975).

87. Peto, R. Quantitative Implications of the Approximate Irrelevance of Mammalian Body Size and Lifespan to Lifelong Cancer Risk. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **370**, 20150198 (2015).

88. Nunney, L. Size Matters: Height, Cell Number and a Person's Risk of Cancer. *Proc. R. Soc. B* **285**, 20181743 (2018).

89. Caulin, A. F. & Maley, C. C. Peto's Paradox: evolution's prescription for cancer prevention. *Trends Ecol. Evol.* **26**, 175–182 (2011).

90. Vincze, O. *et al.* Cancer risk across mammals. *Nature* **601**, 263–267 (2022).

91. Compton, Z. T. *et al.* Cancer prevalence across vertebrates. *Res. Sq.* (2023) doi:10.21203/rs.3.rs-3117313/v1.

92. Tollis, M. *et al.* Return to the sea, get huge, beat cancer: An analysis of cetacean genomes including an assembly for the humpback whale (Megaptera novaengliae). *Mol. Biol. Evol.* **36**, 1746–1763 (2019).

93. Tollis, M., Schneider-Utaka, A. K. & Maley, C. C. The evolution of human cancer gene duplications across mammals. *Mol. Biol. Evol.* **37**, 2875–2886 (2020).

94. Vazquez, J. M. *et al.* Parallel evolution of reduced cancer risk and tumor suppressor duplications in Xenarthra. *Elife* **11**, e82558 (2022).

95. Caulin, A. F., Graham, T. A., Wang, L.-S. & Maley, C. C. Solutions to Peto's paradox revealed by mathematical modelling and cross-species cancer gene analysis. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **370**, 20140222 (2015).

96. Nair, N. U. *et al.* Cross-species identification of cancer resistance-associated genes that may mediate human cancer risk. *Sci. Adv.* **8**, eabj7176 (2022).

97. Smith, M. D. *et al.* Less is more: an adaptive branch-site random effects model for efficient detection of episodic diversifying selection. *Mol. Biol. Evol.* **32**, 1342–1353 (2015).

98. Hanahan, D. & Weinberg, R. A. Hallmarks of Cancer: The Next Generation. *Cell* **144**, 646–674 (2011).

99. Hanahan, D. & Weinberg, R. A. The Hallmarks of Cancer. *Cell* **100**, 57–70 (2000).

100. Hanahan, D. Hallmarks of Cancer: New Dimensions. *Cancer Discov.* **12**, 31–46 (2022).

101. Milacic, M. *et al.* The reactome pathway knowledgebase 2024. *Nucleic Acids Res.* **52**, D672–D678 (2024).

102. Abegglen, L. M. *et al.* Potential Mechanisms for Cancer Resistance in Elephants and Comparative Cellular Response to DNA Damage in Humans. *JAMA* **314**, 1850–1860 (2015).

103. Firsanov, D. *et al.* DNA Repair and Anti-Cancer Mechanisms in the Longest-Living Mammal: The Bowhead Whale. *bioRxiv* 2023.05.07.539748 (2023) doi:10.1101/2023.05.07.539748.

104. Lagunas-Rangel, F. A. Why do bats live so long?—Possible molecular mechanisms. *Biogerontology* **21**, 1–11 (2020).

105. Chionh, Y. T. *et al.* High basal heat-shock protein expression in bats confers resistance to cellular heat/oxidative stress. *Cell Stress Chaperones* **24**, 835–849 (2019).

106. Hua, R. *et al.* Experimental evidence for cancer resistance in a bat species. *Nat. Commun.* **15**, 1401 (2024).

107. Peixoto, F. P., Braga, P. H. P. & Mendes, P. A synthesis of ecological and evolutionary

determinants of bat diversity across spatial scales. *BMC Ecol.* **18**, 18 (2018).

108.USGS. North American Bat Ranges. (2022).

109.Rhie, A. *et al.* Towards complete and error-free genome assemblies of all vertebrate species.

Nature **592**, 737–746 (2021).

110.The Darwin Tree of Life Project Consortium. Sequence Locally, Think Globally: The Darwin Tree of

Life Project. *Proceedings of the National Academy of Sciences* **119**, e2115642118 (2022).

111.Lewin, H. A. *et al.* The Earth BioGenome Project 2020: Starting the Clock. *Proceedings of the*

National Academy of Sciences **119**, e2115635118 (2022).

112.Slater, G. J., Goldbogen, J. A. & Pyenson, N. D. Independent evolution of baleen whale gigantism

linked to Plio-Pleistocene ocean dynamics. *Proc. Biol. Sci.* **284**, (2017).

113.Tacutu, R. *et al.* Human Ageing Genomic Resources: new and updated databases. *Nucleic Acids*

Res. **46**, D1083–D1090 (2018).

114.Morales, A. E. *et al.* Bat genomes illuminate adaptations to viral tolerance and disease resistance.

Nature **638**, 449–458 (2025).

115.O'Shea, T. J. *et al.* Bat flight and zoonotic viruses. *Emerg. Infect. Dis.* **20**, 741–745 (2014).

116.Levesque, D. L., Boyles, J. G., Downs, C. J. & Breit, A. M. High Body Temperature is an Unlikely

Cause of High Viral Tolerance in Bats. *J. Wildl. Dis.* **57**, 238–241 (2021).

117.Kirkwood, T. B. L. The disposable soma theory. *The evolution of senescence in the tree of life* 23–

39 (2017).

118.Toshkova, N. *et al.* Temperature sensitivity of bat antibodies links metabolic state of bats with

antigen-recognition diversity. *Nat. Commun.* **15**, 5878 (2024).

119.Gutiérrez-Guerrero, Y. T. *et al.* Genomic consequences of dietary diversification and parallel

evolution due to nectarivory in leaf-nosed bats. *Gigascience* **9**, (2020).

120.Mandl, J. N., Schneider, C., Schneider, D. S. & Baker, M. L. Going to bat(s) for studies of disease

tolerance. *Front. Immunol.* **9**, 2112 (2018).

121.Pei, G., Balkema-Buschmann, A. & Dorhoi, A. Disease tolerance as immune defense strategy in

bats: One size fits all? *PLoS Pathog.* **20**, e1012471 (2024).

122. Vermillion, K. L., Anderson, K. J., Hampton, M. & Andrews, M. T. Gene expression changes controlling distinct adaptations in the heart and skeletal muscle of a hibernating mammal. *Physiol. Genomics* **47**, 58–74 (2015).

123. Lam, B. *et al.* Multi-species genome-wide CRISPR screens identify GPX4 as a conserved suppressor of cold-induced cell death. *bioRxivorg* 2024.07.25.605098 (2024) doi:10.1101/2024.07.25.605098.

124. Sone, M. & Yamaguchi, Y. Cold resistance of mammalian hibernators ~ a matter of ferroptosis? *Front. Physiol.* **15**, 1377986 (2024).

125. Kaelber, J. T. *et al.* Evolutionary reconstructions of the transferrin receptor of Caniforms supports canine parvovirus being a re-emerged and not a novel pathogen in dogs. *PLoS Pathog.* **8**, e1002666 (2012).

126. Demogines, A., Abraham, J., Choe, H., Farzan, M. & Sawyer, S. L. Dual host-virus arms races shape an essential housekeeping protein. *PLoS Biol.* **11**, e1001571 (2013).

127. Kerr, S. A. *et al.* Computational and functional analysis of the virus-receptor interface reveals host range trade-offs in New World arenaviruses. *J. Virol.* **89**, 11643–11653 (2015).

128. Kaur, H. *et al.* Associations of L-Ferritin and Tim-1 with frailty measures in people with HIV: a cross-sectional and longitudinal study. *Lancet Healthy Longev.* **3**, S5 (2022).

129. Kim, J., Jo, Y., Cho, D. & Ryu, D. L-threonine promotes healthspan by expediting ferritin-dependent ferroptosis inhibition in C. elegans. *Nat. Commun.* **13**, 6554 (2022).

130. Daghlas, I. & Gill, D. Genetically predicted iron status and life expectancy. *Clin. Nutr.* **40**, 2456–2459 (2021).

131. Perez, K. *et al.* Single nuclei profiling identifies cell specific markers of skeletal muscle aging, frailty, and senescence. *Aging (Albany NY)* **14**, 9393–9422 (2022).

132. Trifunovic, A. *et al.* Premature ageing in mice expressing defective mitochondrial DNA polymerase. *Nature* **429**, 417–423 (2004).

133. Van Goethem, G., Dermaut, B., Löfgren, A., Martin, J. J. & Van Broeckhoven, C. Mutation of POLG

is associated with progressive external ophthalmoplegia characterized by mtDNA deletions. *Nat.*

Genet. **28**, 211–212 (2001).

134. Cao, L., Li, W., Kim, S., Brodie, S. G. & Deng, C.-X. Senescence, aging, and malignant

transformation mediated by p53 in mice lacking the Brca1 full-length isoform. *Genes Dev.* **17**, 201–

213 (2003).

135. Vijg, J., Perls, T., Franceschi, C. & van Orsouw, N. J. BRCA1 gene sequence variation in

centenarians. *Ann. N. Y. Acad. Sci.* **928**, 85–96 (2001).

136. Fearon, E. R. Human cancer syndromes: clues to the origin and nature of cancer. *Science* **278**,

1043–1050 (1997).

137. Donoho, G. *et al.* Deletion of Brca2 exon 27 causes hypersensitivity to DNA crosslinks,

chromosomal instability, and reduced life span in mice. *Genes Chromosomes Cancer* **36**, 317–331

(2003).

138. Wooster, R. *et al.* Identification of the breast cancer susceptibility gene BRCA2. *Nature* **378**, 789–

792 (1995).

139. Ray, D. A. *et al.* Multiple Waves of Recent DNA Transposon Activity in the Bat, Myotis Lucifugus.

Genome Res. **18**, 717–728 (2008).

140. Sotgia, S. *et al.* Cellular immune activation in Sardinian middle-aged, older adults and

centenarians. *Exp. Gerontol.* **99**, 133–137 (2017).

141. Lee, K.-A., Flores, R. R., Jang, I. H., Saathoff, A. & Robbins, P. D. Immune senescence,

immunosenescence and aging. *Front. Aging* **3**, 900028 (2022).

142. Yousefzadeh, M. J. *et al.* An aged immune system drives senescence and ageing of solid organs.

Nature **594**, 100–105 (2021).

143. Lee, C.-S. *et al.* Upregulation of cluster of differentiation 36 mRNA expression in peripheral blood

mononuclear cells correlates with frailty severity in older adults. *J. Cachexia Sarcopenia Muscle*

13, 1948–1955 (2022).

144. Kimmel, J. C., Yi, N., Roy, M., Hendrickson, D. G. & Kelley, D. R. Differentiation reveals latent

features of aging and an energy barrier in murine myogenesis. *Cell Rep.* **35**, 109046 (2021).

145. Tian, Y. *et al.* FTH1 inhibits ferroptosis through ferritinophagy in the 6-OHDA model of Parkinson's

disease. *Neurotherapeutics* **17**, 1796–1812 (2020).

146. Ricklefs, R. E. Life-history connections to rates of aging in terrestrial vertebrates. *Proc. Natl. Acad.*

Sci. U. S. A. **107**, 10314–10319 (2010).

147. Tian, S. *et al.* Comparative analyses of bat genomes identify distinct evolution of immunity in Old

World fruit bats. *Sci Adv* **9**, eadd0141 (2023).

148. Jonathan Sleeman, Center Director, USGS National Wildlife Health Center. NWHC Operations

During the COVID-19 Pandemic and Information About Coronaviruses in Wildlife. (2020).

149. White-Nose Syndrome Disease Management Working Group. *National White-Nose Syndrome*

Decontamination Protocol. www.WhiteNoseSyndrome.org.

150. Wu, N. *et al.* The interferon stimulated gene 20 protein (ISG20) is an innate defense antiviral factor

that discriminates self versus non-self translation. *PLoS Pathog.* **15**, (2019).

151. Legrand, A. *et al.* SAMD9L acts as an antiviral factor against HIV-1 and primate lentiviruses by

restricting viral and cellular translation. *PLoS Biol.* **22**, e3002696 (2024).

152. Išerić, H., Alkan, C., Hach, F. & Numanagić, I. Fast characterization of segmental duplication

structure in multiple genome assemblies. *Algorithms Mol. Biol.* (2022).

153. Brown, M., González De la Rosa, P. M. & Mark, B. *A Telomere Identification Toolkit*. (2023).

doi:10.5281/zenodo.10091385.

154. Li, H. New strategies to improve minimap2 alignment accuracy. *Bioinformatics* (2021).

155. Goel, M. & Schneeberger, K. plotsr: visualizing structural similarities and rearrangements between

multiple genomes. *Bioinformatics* **38**, 2922–2926 (2022).

156. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the

Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

157. Zhang, C. & Mirarab, S. Weighting by gene tree uncertainty improves accuracy of quartet-based

species trees. *Mol. Biol. Evol.* **39**, (2022).

158. Romiguier, J., Ranwez, V., Delsuc, F., Galtier, N. & Douzery, E. J. P. Less is more in mammalian phylogenomics: AT-rich genes minimize tree conflicts and unravel the root of placental mammals. *Mol. Biol. Evol.* **30**, 2134–2144 (2013).

159. Foley, N. M. *et al.* A genomic timescale for placental mammal evolution. *Science* **380**, eabl8189 (2023).

160. Dos Reis, M. & Yang, Z. Bayesian Molecular Clock Dating Using Genome-Scale Datasets. *Methods Mol. Biol.* **1910**, 309–330 (2019).

161. Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**, 1586–1591 (2007).

162. Phillips, M. J. Geomolecular dating and the origin of placental mammals. *Syst. Biol.* **65**, 546–557 (2016).

163. Gunnell, G. F. & Simmons, N. B. Fossil evidence and the origin of bats. *J. Mamm. Evol.* **12**, 209–246 (2005).

164. Eiting, T. P. & Gunnell, G. F. Global Completeness of the Bat Fossil Record. *J. Mamm. Evol.* **16**, 151–173 (2009).

165. Storch, G., Sigé, B. & Habersetzer, J. Tachypteron franzeni n. gen., n. sp., earliest emballonurid bat from the Middle Eocene of Messel (Mammalia, Chiroptera). *Palaontol. Z.* **76**, 189–199 (2002).

166. Ravel, A. *et al.* A new large philisid (Mammalia, Chiroptera, Vespertilionoidea) from the late Early Eocene of Chambi, Tunisia: LARGE BAT FROM CHAMBI. *Palaeontology* **55**, 1035–1041 (2012).

167. Lim, B. K. Review of the origins and biogeography of bats in South America. *Chiroptera Neotropical* **15**, 391–410 (2009).

168. Morgan, G. S. & Czaplewski, N. J. A New Bat (Chiroptera: Natalidae) from the Early Miocene of Florida, with Comments on Natalid Phylogeny. *J Mammal* **84**, 729–752 (2003).

169. Elliot, M. G. & Mooers, A. Ø. Inferring ancestral states without assuming neutrality or gradualism using a stable model of continuous character evolution. *BMC Evol. Biol.* **14**, 226 (2014).

813 170. Kosakovsky Pond, S. L. *et al.* HyPhy 2.5-A customizable platform for evolutionary hypothesis

814   testing using PHYlogenies. *Mol. Biol. Evol.* **37**, 295–299 (2020).

815 171. Mendes, F. K., Vanderpool, D., Fulton, B. & Hahn, M. W. CAFE 5 models variation in evolutionary

816   rates among gene families. *Bioinformatics* **36**, 5516–5518 (2021).

Materials and Methods

Data availability

819   All sequencing data and genomes generated in this study are available on NCBI under Bioprojects
820 PRJNA973719, PRJNA1035541, and PRJNA1295526. Annotations generated in this study are available at
821 https://github.com/docmanny/myotis-gene-annotations.   All   other   code   is   available   at
822 https://github.com/sudmantlab/MyotisGenomeAssembly. All other data are archived on DataDryad at DOI:
823 10.5061/dryad.02v6wwqfh.

Samples and materials

All bats sampled for this study were wild caught under scientific collection permits for California and
Arizona (see Supplemental Table 1). Bats were sampled using standard mist-netting procedures, including
taking standard body measurements, following USGS recommendations for White-Nose Syndrome and COVID-
19 prevention[148,149]. For *M. lucifugus*, the donor individual was field-caught in California and transported to the
Genetics Laboratory of the California Department of Fish and Wildlife, where they were euthanized via
isofluorane. The *M. velifer* individual was caught in Arizona and euthanized in the field via isoflurane. For both
M. lucifugus and *M. velifer*, tissues were collected and preserved via flash-freezing in liquid nitrogen. All other
genomes were generated from primary cell lines. Additional information can be found in the *Extended Methods*
and in **Table S1**.

All PKR experiments were performed using HeLa PKR-KO cells (kindly provided by A. Geballe, Fred
Hutchinson Cancer Center, Seattle WA). The cells were maintained at 37°C with 5% CO_2 and cultured in DMEM
supplemented with 5% fetal bovine serum (FBS), 1% penicillin/ streptomycin mix and 1 µg/mL puromycin (Sigma-
Aldrich). All transfections were performed 24 hours after seeding, using 3 µL of TransIT-LT1 Transfection
Reagent (Mirus Bio) per 1 µg of DNA and Opti-MEM media. We used previously-generated pSG5-FLAGx2
vectors encoding either *M. myotis* PKR1 (GenBank OP006550), *M. myotis* PKR2 (GenBank OP006559), *M.*
velifer PKR1 (GenBank OP006558), or *M. velifer* PKR2 (GenBank OP006557)[28]. Plasmids encoding the
interferon-stimulated gene ISG20[150] and a constitutively active variant of the sterile alpha motif domain-
containing protein 9-like SAMD9L-F886Lfs*11 (here, SAMD9L[151]) were used as controls in viral infections and
cell translation experiments, respectively.

Near-complete genome assembly and annotation

Details on DNA extraction, preparation of PacBio HiFi and Omni-C (Dovetail Genomics) libraries, and genome assembly and manual curation can be found in the *Extended Methods*.

Gene & repetitive element annotation and alignments

Details describing our procedure for identifying and annotation repetitive elements and protein-coding gene annotations can be found in the *Extended Methods.*

RNA sequencing for gene annotations and differential expression

Details on RNA extraction, library preparation, sequencing, and can be found in the *Extended Methods* and **Table S9**.

Structural variation

To understand the distribution of structural variants, including segmental duplication events, we used SyRI (Synteny and Rearrangement Identifier[63]) and BISER (Brisk Inference of Segmental duplication Evolutionary stRucture[152]). We first masked telomere regions using TIDK (Telomere Identification toolKit[153]), and mapped the primary 22 scaffolds of the nearctic *Myotis* genomes to each other with minimap2[154]. The scaffold corresponding to the X chromosome was omitted because there is no corresponding scaffold in the *M. yumanensis* assembly. To verify homologous chromosomes and fix strand orientation, we used *fixchr* from the SyRI package and manually renamed scaffolds accordingly, then re-mapped them with minimap2. We ran SyRI on the resulting files and plotted the results with plotsr[155]. We ran BISER on the primary 22 scaffolds of the nearctic *Myotis* genomes with –keep-contigs and default settings to generate bed files with the inferred segmental duplication regions.

Gene Trees & Phylogeny

A phylogeny of all 534 mammals in our alignments was generated using IQTREE[156] (version 2.3.1) using all gene alignments with the settings "-B 1000 -m GTR+F3x4+R6." To generate gene trees, we first filtered our gene alignments to exclude alignments with over 50% gaps in the sequence and less than 4 species. With the remaining alignments, we used IQTREE to find the best-fitting substitution model and tree using settings "--wbtl --bnni --alrt 1000 -B 1000 --safe". The best substitution models for each gene were saved as a NEXUS file. As the bootstrap values for this tree were unanimously 100%, we further confirmed the species tree by using ASTRAL-IV[157], and confirmed that our phylogeny agreed with other previously-published *Eutherian* phylogenies, including phylogenies of 39 mammals[158], of 241 placental mammals[159], and of 11 nearctic *Myotis* species[55].

Time-calibration of 59 bat genomes

Using our codon alignments of 59 bat genomes, we generated a time-calibrated phylogeny using mcmctree[160] and PAML[161] (v. 4.10.0) using an approximate likelihood method. Using the pan-bat codon alignments and our phylogeny as input, with fossil calibrations based on previously published work[4,43,159,162–168], we ran *mcmctree* twice to generate the Hessian matrix and confirm convergence. This was followed by 10 independent chains using the "out.BV" file from the first run. Finally, the output files of all 10 chains were combined to compute final divergence time estimates (see Table S2).

Ancestral Body Size, Lifespan, and Cancer Risk reconstruction

To explore how body size and lifespan have evolved over time in mammals, we used a super-phylogeny of mammal species[54] subsampled to only contain species with extant body size and lifespan data collected from AnAge[15] and PanTHERIA[16]. Ancestral body sizes and lifespans were simulated separately using StableTraits[169] and the settings "*--iterations 100000 --skip 1000 --chains 8*". Estimates were further validated by comparing our initial results to a second run of StableTraits using the same settings.

The Relative Incidence of Cancer Risk (RICR) was calculated across our mammalian phylogeny following the method of Vazquez and Lynch (2021)[37]. The cancer risk K for each node was calculated using the log of its median predicted body size and lifespan. An organism's lifetime risk of cancer K is proportional to Dt^6, where D is the body size and t is the maximum lifespan. RICR is then calculated as the $\log_2$ of the ratio between the cancer risk of a given node and its direct ancestor.

Selection Scans & Evolutionary Rates

aBSREL

To conservatively test for branch-specific selection, we used aBSREL[97,170] (version 2.5.48) to test for selection at each branch within the Nearctic *Myotis* clade for 15,734 gene alignments spanning 534 mammals. These genes were identified as 1:1 orthologs across the full alignment, with no more than 50% sequence gaps and at least 4 species present in the alignment. For each gene alignment, the full phylogeny was trimmed to match the species present, and the HYPHY script "label-tree.bf" was used to highlight all nodes within nearctic *Myotis* as foreground branches. aBSREL was then run using the pruned and highlighted tree and gene alignments on the command line with the options "--code Universal --branches Foreground". We defined a gene as under selection if one or more regions within the gene demonstrated a selective pressure with an FDR-corrected p-value ≤ 0.05; a gene was specifically identified as under positive selection if at least one of these significant regions had an $\omega > 1$.

BUSTED

To quantify the total amount of positive selection across the *Myotis* tree or the different species trees used in this manuscript, we used an improved version of the BUSTED[70,170] test called BUSTED-MH (see *Extended Methods* for details on BUSTED vs BUSTED-MH). We applied BUSTED-MH to 19,646 *Myotis* orthologous CDS alignments with at least five orthologs. These orthologs are cases where the Orthofinder gene

trees coincide with the species tree. This effectively removes issues regarding whether we should use the gene or the species tree, at the cost of removing 2,110 genes from the *Myotis* selection analysis. Similarly, we applied BUSTED-MH to 17,469 non-*Myotis* bat alignments with at least five orthologs. The species in these two sets (*Myotis* bats and non-*Myotis* bats) are mutually exclusive. This includes a subset of 14,091 alignments with orthologs present in two thirds of the non-*Myotis* bat species, regardless of the genes' presence/absence in the *Myotis*-only complement. This is specifically used to show that patterns of virus-driven adaptation are representative of all bats. We also tested 17,890 primate alignments with at least five orthologs with BUSTED-MH, as well as 19,311 glire, 18,000 carnivora and 18,504 ungulate alignments.

Enrichment scores for VIPs

We expanded on the set of VIPs reported in Souilmi *et al*. (2021)[71] (5,291 VIPs) by searching the literature for more recent publications reporting VIPs, with a final dataset of 5,527 VIPs. These VIPs ("All VIPs") were further sub-categorized into 4 groups based on their interactions with either DNA viruses and/or RNA viruses: "DNA VIPs", "DNA-only VIPs", "RNA VIPs", and "RNA-only VIPs" (**Table S5**). For each VIP set, we selected sets of matching control genes by controlling for 16 different criteria. (Table S5). Sets of control genes were resampled in a bootstrap procedure to generate 95% confidence intervals for sets of genes across a range of p-values; additional details can be found in the *Extended Methods*.

Gene Duplications and CAFE analysis

To quantify patterns of gene duplication and loss, we quantified the copy number of genes with human orthologs from our gene annotations for each nearctic *Myotis* genome. To calculate per-gene expansion and loss rates and their statistical significance, we ran CAFE[171] v5 on the previously described set of copy number counts using our time-calibrated species tree pruned to include only the nine nearctic *Myotis* species. CAFE uses a birth-death model of gene family evolution to investigate changes in gene family size accounting for evolutionary history using the species phylogeny. Evolutionary branch length is incorporated into the model expectations, thus accounting for potential bias resulting from differences in branch lengths. We ran CAFE on the subset of genes with two or more copies in at least one species using a Poisson distribution for the root frequency (-*p*), first generating an error model to correct for genome assembly and annotation error (-*e*). We compared the base model (each gene family belongs to the same evolutionary rate category) to gamma models (each gene family can belong to one of *k* evolutionary rate categories) with different values of *k* from 2 to 13. A final gamma model with *k*=9 was chosen to balance model log likelihood with the number of gene families for which the optimizer failed. The model was run three separate times to ensure convergence.

To understand if genes in these pathways have higher birth rates or are more likely to have significant changes in gene copy number than expected relative to other genes, we evaluated the gene copy birth rate λ and number of genes that have significantly expanded or contracted in copy number on at least one branch within our nearctic *Myotis* phylogeny using a bootstrap procedure. Following Huang et al.[41], we tested if VIP genes in particular underwent significant copy number changes or had significantly different birth/death rates than non-VIP genes. For each category of VIP genes (all VIPs, DNA VIPs, DNA only VIPs, RNA VIPs, and RNA only VIPs), we generated 100 bootstrap sets of control non-VIP genes with the same number of genes as the corresponding VIP gene set. We ran CAFE on each set of VIP genes and the corresponding control non-VIP genes to infer per-gene birth-death rates and per-gene, per-branch expansion/loss events.

Assessment of DNA Double-Strand Break Tolerance

We assessed each species' tolerance to DNA double strand breaks using a by measuring viability, cytotoxicty, and apoptosis across a range of doses of Neocarzinostatin, a radiomimetic drug. We measured dose response curves in wing-derived primary dermal fibroblasts across 5 bat species (*Myotis lucifugus*, n=8; *Myotis evotis*, n=8; *Rousettus langosus*, n=2; *Eidolon helvum*, n=2; *Pteropus rodrigensis*, n=2) using the multiplexed ApoTox-Glo assay (Promega). Using 96-well plates, two individuals and 11 doses were assessed simultaneously with four technical replicates. Results were normalized to treatment controls for each individual in R as previously described[35,36,39,44].

PKR1/2 Characterization in PKR-KO HeLa cells

Western blot

We assessed for the steady state protein expression of *M. myotis* Flag-PKRs after transfection of 350 ng or 700 ng total of DNA plasmids encoding: PKR1 alone, PKR2 alone or both PKR1 and PKR2 (175 ng of each and 350 ng of each, respectively). Briefly, cells were re-suspended and lysed in ice-cold RIPA buffer (50 mM Tris pH8, 150 mM NaCl, 2 mM EDTA, 0.5% NP40) with protease inhibitor cocktail (Roche) and sonicated. 20 µL of the clarified fraction was denatured with 5 µL of 6x Laemmli buffer at 95°C for 5 min and loaded into 4-20% BioRad Criterion TGX Stain-Free precast gel. The wet transfer into a PVDF membrane was executed overnight at 4°C. The membranes were blocked in a 1xTBS-T buffer (Tris HCl 50 mM pH8, NaCl 30 mM, 0.05% of Tween 20) containing 10 % powder milk, and were incubated for 1h. The membranes were incubated with primary mouse anti-Flag (Sigma F3165) and anti-Tubulin (Sigma T5168) antibodies and secondary anti-Mouse IgG-Peroxidase conjugated (Sigma A9044). Detection was made using the Chemidoc Imaging System (BioRad) with SuperSignal West Pico Chemiluminescent Substrate (ThermoFisher Scientific).

Luciferase reporter assays

Luciferase reporter assays were carried out to investigate whether the two PKR paralogs have synergistic, additive or dominant negative effect in translation shutdown. 25,000 PKR-KO cells were seeded in 24-well plates. The next day, cells were transfected with 350 ng of empty pSG5, 350 ng of PKR1, 350 ng of PKR2, or 175 ng of PKR1+ 175 ng of PKR2, with additional 50 ng of FFLuc firefly luciferase reporter plasmid per well. Sterile alpha motif domain-containing proteins 9L (SAMD9L gain-of-function mutant) was used as a positive control of translational repression[151]. 24 h post transfection, cells were washed with PBS, lysed by a 5× reporter lysis buffer (Promega) and incubated overnight at -20°C. Cells were then collected and 100 µl of the luciferase substrate (Promega) was added to 20 µl of the lysis supernatant. Alternatively, cells were lysed using BrightGlow Lysis Reagent (Promega E2620). The relative luminescence units (RLUs) proportional to the translation of the luciferase reporter were immediately quantified with LUMIstar Omega microplate reader optima (BMG Labtech). All luciferase assays were conducted in technical duplicates in at least five independent experiments. Fold change results were normalized to the empty pSG5 condition within each independent experiment.

PKR co-immunoprecipitation

Hela PKR-KO cells were transfected with 1.25µg plenti6 HA-tagged *M. myotis* PKR1 plasmid per million cells and 1.25µg of either plenti6 myc empty vector, myc-tagged *M. myotis* PKR1 or myc-tagged *M. myotis* PKR2 plasmid using TransIT-LTI transfection reagent (Mirus Bio). The next day, some wells were infected with Sindbis

virus expressing GFP (SINV-GFP) at MOI 2 for 24 hours. Cells were then scraped with cold PBS and pelleted. For the IP, cells were lysed in 500µl IP buffer (50mM Tris HCl pH7.5, 140mM NaCl, 6 mM MgCl2, 0.1% NP40) supplemented with RNase (RiboLock, Fisher Scientific) and protease (complete EDTA-free protease inhibitor cocktail, Sigma) inhibitors for 10 minutes on ice, then centrifuged at 12,000 *xg* for 10 min at 4°C. 5% of the volume was kept for input, while the rest was incubated with 40µl µMACS anti-c-myc MicroBeads (Miltenyi Biotec) for 1h at 4°C with constant rotation. Samples were then loaded onto µMACS columns placed in the magnetic field of a µMACS Separator (Miltenyi Biotec), washed 4 times with cold IP buffer, and eluted with the µMACS denaturing elution buffer. Proteins were denatured in elution buffer for 5 min at 95°C, then loaded onto a 4-20% BioRad Criterion TGX Stain-Free precast gel and transferred onto an Amersham Protran nitrocellulose membrane (Sigma) for 1h. Membranes were blocked for 1h in 5% milk in PBS (Euromedex) supplemented with 0.2% tween (Fisher) and incubated with mouse anti-myc monoclonal antibodies (Abcam 9E10, cat# ab32) then secondary anti-mouse IgG antibodies conjugated with HRP (Sigma, cat# A4416), or with rat anti-HA antibodies conjugated with HRP (Roche, Sigma, cat# 12013819001). Images were taken on a Fusion FX imager (Vilber) with SuperSignal West Femto Chemiluminescent Substrate (ThermoFisher Scientific).

Cell viability assay

Hela PKR-KO cells were transfected 24h after plating in 96 well plates (n=10,000 cells/well), with 100 or 200 ng of pSG5 plasmid: empty or coding for *M. myotis* or *M. velifer* PKR1, PKR2 or PKR1+2 equal mix (50%-50%). 24 hours post-transfection, positive control cells were treated with an apoptosis-inducing drug, Etoposide, at different doses (250, 200 or 100 µM). 48 hours post transfection, cells were harvested and lysed to quantify luminescent signal according to CellTiter-Glo® Luminescent Cell Viability Assay (Promega) kit protocol.

VSV and SINV infections

VSV infections. 200,000 cells were plated in 12-well and transfected 24 h after with 350 ng of pSG5 plasmid: empty, or encoding *M. myotis* or *M. velifer* PKR1, PKR2, or equal input of PKR1 and PKR2 (175 ng per plasmid), or a plasmid encoding interferon-stimulated exonuclease gene 20 (ISG20), due to its known antiviral activity against VSV as positive control[150]. Cells were infected 24 h post transfection with replicative VSV-GFP virus[68] at a MOI of 3. Cells were fixed with 4% paraformaldehyde 16-18 hours post infection. VSV infection was quantified by measuring the percentage of GFP positive cell populations with BD FACSCanto II Flow Cytometer (SFR BioSciences). Fold change results were normalized to the empty pSG5 condition across at least three independent experiments.

SINV infections. Hela PKR-KO cells were transfected with 5µg pSG5 empty vector, *M. myotis* PKR1, *M. myotis* PKR2 or 2.5µg *M. myotis* PKR1 + 2.5µg *M. myotis* PKR2 per million cells using TransIT-LTI transfection reagent (Mirus Bio). The next day, some wells were infected with SINV-GFP at MOI 0.2. Cells were then placed into a CellCyte X live cell imaging system (Cytena) and pictures of every well were taken every 2h for 48h. The fraction of GFP+ cells over the total cell area was measured and averaged from six photos of 2 individual wells per condition, and repeated for a total of three independent experiments.

 # Fig. Legends

Fig. 1: 8 near-complete reference assemblies for North American (Nearctic) *Myotis*. A) Phylogeny of 38 bat genomes with 3 outgroup species: cow *(bosTau9)*, mouse *(mm39)*; and human *(T2T-CHM13v2.0)*. Bars at the tips of the phylogeny indicate the Area under the NGx curve (AuNG) score of each genome, a metric of genome contiguity independent of genome size. The dotted line represents the AuNG score for the T2T-CHM13v2.0 assembly. **B)** The time-calibrated phylogeny of 9 Nearctic and two representative Palaearctic *Myotis* species based on orthologous codon alignments. Blue bars represent age uncertainties. **C)** Map of capture sites for new cell lines with representative images for the species sequenced in this study (see "Acknowledgements" for attributions). **D)** Mammalian BUSCO scores for annotations generated for the 8 new *Myotis* genomes. **E)** Ideogram bar plot indicating completion status of each chromosome in assembly. Pie graphs indicate completion status of all chromosomes in assembly. Percentages above ideogram indicate the proportion of T2T assembled chromosomes across species. All chromosomes were positively identified based on size, synteny, and homology to human chromosomes[50]. "Complete (T2T)" status indicates that a chromosome is fully assembled telomere-to-telomere without gaps; "Draft (T2T, gaps)" status indicates that a chromosome is fully scaffolded with both telomeres, but has one or more gaps in the assembly; "Incomplete" status indicates that a chromosome was positively identified, but was not scaffolded from telomere to telomere (only contains one telomere).

Fig. 2: A varied structural variation landscape across 9 nearctic *Myotis* species. A) Synteny between chromosomes of 9 *Myotis* species showing syntenic regions (grey), inversions (orange), translocations (green), and duplications (blue). **B)** Distribution of transposable elements (top) and segmental duplications (red heatmap, bottom) in *M. velifer*. Putative locations of centromeres are denoted by the dotted lines. **C)** Pie chart of overall genomic proportions of TEs in *M. velifer*. **D)** Histogram of segmental duplication size distributions genome-wide in *M. velifer*.

Fig. 3: Evolutionary history and function of an actively segregating copy number polymorphism of *PKR* in *Myotis*. A) Structural comparison of the main PKR haplotypes in two species. Orthologous regions between the two haplotypes are indicated by grey bands, while syntenic duplications are indicated in green. Exons are annotated with black marks. **B)** Cartoon of the PKR locus in the two phased haplotype assemblies of each species. While *PKR2* is present across all haplotypes, *PKR1* and *PKR copy 3* are polymorphic within and across species. Each number indicates the number of exons per gene. **C)** Reconciled gene tree for *PKRs* across all haplotypes and species shown in B. Haplotype corresponding to the reference (A) and alternate (B) haplotype for each species are represented by upper- and lower-diagonal triangles, respectively. **D)** Effect of Myotis PKR expression on luciferase reporter translation in PKR-KO cells, measured in Relative Light Units (RLU) and normalized to the empty pSG5 control. Human SAMD9L-GoF is a positive control of translation inhibition[151]. **E-F)** Effect of PKR on viral infection by VSV-GFP (E, and S3B) and SINV-GFP (F 34h post-infection, and S3C), measured by flow cytometry as the % of GFP+ cells, normalized to the pSG5 control. Although all conditions restricted VSV and SINV, the expression of both PKR1+PKR2 was not beneficial against these viral infections. ISG20 served as a positive control of VSV-GFP restriction[150]. **G)** Co-immunoprecipitation (IP) of PKR-KO cells transfected with *Myotis* HA-PKR1 and either *Myotis* myc-PKR1, *Myotis* myc-PKR2 or a myc-empty vector control. Proteins were pulled down with anti-myc beads and lysates from 5% input or IP samples were run on a western blot and stained for HA and myc. Representative of 3 independent experiments. Extended Data Fig. 3E for SINV conditions. H, Quantification of the three independent experiments (mock and SINV): HA left and myc right. For **D-F**, error bars indicate the means ± SEM for at least three independent experiments. Statistics, unpaired t-test (p<0.05). mPKR, Myotis myotis PKR; vPKR, Myotis velifer PKR.

Fig. 4: Adaptation to DNA viruses, but not RNA viruses, is enriched in *Myotis* and other bats. A) Diagram of an example VIP, host cell transmembrane receptor CD45, showing strength of adaptation at each codon and the locations of direct interactions with the human adenovirus protein sec49K. **B-F)** Enrichment plots showing the ratio

of positive selection in only *Myotis* bats in VIPs versus matched sets of control genes at different p-value thresholds.
The solid line shows the median ratio; the color of the line, and the number above each point, represents the number
of VIPs with significant BUSTED-MH p-values at the given threshold; the grey band represents the 95% confidence
interval generated by bootstrapping sets of matched control genes. The 95% confidence interval widens at lower
p-values because there are fewer VIPs with significant p-values as the alpha threshold decreases, reducing power.
G-I) The same as above, across all bats in this study excluding *Myotis*. **J-L)** The number of VIP genes inferred by
CAFE to have significantly expanded or contracted in copy number on at least one branch of the nearctic *Myotis*
phylogeny (blue line) compared to the distribution across 100 randomized sets of non-VIP control genes (black
line). Vertical dashed lines indicated the 95% highest density interval for the control gene distribution.

Fig. 5: Evolution of cancer risk and resistance in *Myotis* and other mammals. A, B) Cophylo plot of the
evolution of body size (**A**) and lifespan (**B**) across *Eutheria*. Branch lengths in **A-B** are scaled proportional to the
rate of change of the trait over time. **C)** Diagram illustrating the relationship between changes in body size and
lifespan with changes in cancer risk and resistance. **D)** Relative Intrinsic Cancer Risk (RICR) for every node in
Eutheria, ranked from greatest reduction in cancer risk to greatest increase in cancer risk. RICR relative to the most
recent ancestor of select nodes are highlighted, as well as the average RICR across for all nodes within select
clades. **E)** The phylogeny of Nearctic *Myotis* species with inset pie charts representing the proportion of the top 100
Reactome pathways overrepresented among genes under selection at each node that are associated with cancer-
related processes. Below, pie chart indicates expected proportion of pathways are cancer-associated after 1000
random samples of 100 pathways from the full Reactome database. Asterisks represent nodes with proportions
greater than the expected value at p≤0.05 using Fisher's exact test. **F)** Proportion of the top 100 Reactome pathways
overrepresented among genes under selection across all nodes in a species' evolutionary history that are
associated with cancer-related processes. **G)** Volcano plot of overrepresented pathways in Reactome among the
union set of genes under selection across all nodes in the evolutionary history for *M. lucifugus*. **H)** Viability and
Apoptosis fold-change in 5 bat species in response to different doses of neocarzinostatin (NCS), a potent inducer
of DNA double-strand breaks. Points represent individual replicates normalized to each species' control, while bars
represent mean ± 95% confidence intervals.

Fig. 6: **Pleiotropy among genes under selection in *M. lucifugus* between DNA-damage responsive genes
and DNA-only VIPs. A-B)** Volcano plots of genes that are differentially expressed in *M. lucifugus* after 6 hours (**A**)
and 18 hours (**B**) of treatment with 100uM neocarzinostatin (NCS). **C-D)** Pathway-level Gene Set Enrichment
Analysis (GSEA) of genes differentially expressed at 6 and 18 hours post-treatment. **E)** Upset plot between genes
that are differentially expressed after NCS treatment and DNA-only VIPs; the hypergeometric P-value of the
intersection is shown. **F)** Pathway-level overrepresentation analysis of genes that are DNA-only VIPs and are also
differentially expressed in response to NCS treatment. **G)** Pathway-level overrepresentation analysis of genes
identified as under selection in *M. lucifugus* via ABSREL that are DNA-only VIPs and are also differentially
expressed in response to NCS treatment.

Supplemental Information

Extended Data Fig. 1-7:

1. **Contiguity statistics for 8 novel *Myotis* genomes**
2. **Estimated node divergence times for *Chiroptera* and ASTRAL-IV scores.**
3. **Correlations of TEs and structural variation in *Myotis* genomes**
4. **Additional functional characterization of *Myotis* PKR duplicates**
5. **VIP selection enrichments in bats and other mammals**
6. **Evolution of Cancer Risk and Resistance across *Myotis***
7. **Genes differentially expressed after neocarzinostatin treatment and their overlap with DNA-only VIPs**

Table S1. Genome Statistics

Table S2. Accessions for additional genomes used in this study

Table S3. SyRI-identified structural variants (SVs)

Table S4. Experimental data for PKR experiments

Table S5. List of VIPs and VIP subclasses

Table S6. Ancestral State Reconstructions for body size, lifespan, and RICR in mammals

Table S7. aBSREL significant gene lists and Reactome enrichments

Table S8. Experimental data for Neocarzinostatin experiments

Table S9. RNA-seq QC statistics and GSEA results

A

Reference Chr

Chromosome Position (Mbp)

Genomes
- *M. occultus*
- *M. lucifugus*
- *M. yumanensis*
- *M. velifer*
- *M. californicus*
- *M. auriculus*
- *M. evotis*
- *M. thysanodes*
- *M. volans*

Annotations
- Syntenic
- Inversion
- Translocation
- Duplication

B

Proportion of Region

Chromosome Position in mMyoVel1 (Mb)

Transposable Element
- DNA
- LTR
- Retroposon
- Unknown/Other
- LINE
- RC
- SINE
- non-TE

Proportion of Region (SD)
0.75 0.50 0.25

C

12.7%   1.4%
1.7%
3.5%
2.1%

78.7%

Genome Proportions (mMyoVel1)

D

Scaled Density

SD length (bp) genome-wide

A

B

Body Size (log(g))
5   10   15

Lifespan (log(yrs))
0   1   2   3   4   5

Whales

Bats

Primates

Elephants

C

Cancer Risk & Resistance

Reduced Risk { Cells ↓ Lifespan ↓ } Relaxed Resistance

Increased Risk { Cells ↑ Lifespan ↑ } Augmented Resistance

D

Avg. Vespertilionidae
Avg. Myotis
Avg. Phyllostomidae
Avg. Yinpterochiroptera
Evo-Thy
Myotis californicus
Yinpterochiroptera
Myotis evotis
Myotis volans
Phyllostomidae
Myotis yumanensis
Evo-Luc
Myotis myotis
Vespertilionidae
Yum-Vel
Myotis
Myotis lucifugus
Myotis grisescens
Myotis brandtii

RICR (log₂)

-10

Yangochiroptera
Myotis thysanodes
Myotis velifer
Avg. Yangochiroptera
Avg. Molossidae
Molossidae

Species

Myotis lucifugus (n=8)
Myotis evotis (n=9)
Eidolon helvum (n=2)
Pteropus rodrigensis (n=2)
Rousettus lanosus (n=2)

E

* M.lucifugus
* M.occultus
* M.yumanensis
* M.velifer
* M.evotis
* M.thysanodes
* M.auriculus
* M.californicus
M.volans

14%
86%

Cancer
Not Cancer

Reactome Pathways

F

Enriched Pathways Associated with Cancer (%)

0%   5%   10%   15%

G

100   300   500   648

-log(FDR)

7.0
6.5

4.0

3.5

3.0

2.5

2.0

1.5

1.0

0.5

0.0

Extracellular matrix organization

Collagen formation

Resolution of D-Loop Structures (Through Holiday Junction Intermediates)

HDR through Homologous Recombination (HRR)

Homologous DNA Pairing and Strand Exchange

Diseases associated with the TLR signaling cascade

MyD88 deficiency (TLR2/4)

Sensing of DNA DSBs

Immune System

log₂(Enrichment Ratio)

0.0   0.5   1.0   1.5   2.0

H

Apoptosis

Fold-Change

2.5
2.0
1.5
1.0
0.5

*** *** * n.s. *** n.s. n.s.

Viability

1.5
1.0
0.5
0.0

n.s. *** *** *** ***

NCS [uM]
0   10   500

*** p_{adj}≤5e-4   ** p_{adj}≤5e-3   * p_{adj}≤0.05   n.s. p_{adj}> 0.05