The dinucleotide structure of NAD enables specific reduction on mineral surfaces

Delfina P. Henriques Pereira*[a], Xiulan Xie[b], Tuğçe Beyazay[c], Nicole Paczia [d], Zainab Subrati[b], Jürgen Belz[e], Kerstin Volz[e], Harun Tüysüz[c], Martina Preiner*[a]

[a]   Delfina P. Henriques Pereira
        Dr. Martina Preiner
        Microcosm Earth Center
        Max-Planck-Institute for Terrestrial Microbiology
        Hans-Meerwein-Str. 4; 35032 Marburg
        E-mail: martina.preiner@mpi-marburg.mpg.de; delfina.pereira@mpi-marburg.mpg.de
[b]   Dr. Xiulan Xie
        Zainab Subrati
        Department of Chemistry
        Philipps University Marburg
        Hans-Meerwein-Str. 4; 35032 Marburg
[c]   PD Dr. Harun Tüysüz
        Dr. Tuğçe Beyazay
        Heterogeneous Catalysis
        Max-Planck-Institut für Kohlenforschung
        Kaiser-Wilhelm-Platz 1; 45470 Mülheim an der Ruhr
[d]   Dr. Nicole Paczia
        Metabolomics and small molecule mass spectrometry
        Max-Planck-Institute for Terrestrial Microbiology
        Karl-von-Frisch-Str. 10; 35043 Marburg
[e]   Prof. Dr. Kerstin Volz
        Dr. Jürgen Belz
        Department of Physics
        Philipps University Marburg
        Hans-Meerwein-Straße 6; 35032 Marburg

        Supporting information for this article is given via a link at the end of the document

Abstract: Nucleotide-derived cofactors could function as a missing link between the informational and the metabolic part at life's emergence. One well-known example is nicotinamide dinucleotide (NAD), one of the evolutionarily most conserved redox cofactors found in metabolism. Here, we propose that the role of these cofactors could even extend to missing links between geo- and biochemistry. We show NAD^+ can be reduced under close-to nature conditions with nickel-iron-alloys found in water-rock-interaction settings rich in hydrogen (serpentinizing systems) and that nicotinamide mononucleotide (NMN), a precursor molecule to NAD, has different properties regarding reduction specificity and sensitivity than NAD. The additional adenosine monophosphate (AMP) "tail" of the dinucleotide, a shared trait between many organic cofactors, seems to play a crucial mechanistic role in preventing overreduction of the nicotinamide-bearing nucleotide. This specificity is also connected to the used transition metals. While the combination of nickel and iron promotes the reduction of NAD^+ to 1,4-NADH most efficiently, in the case of NMN, the presence of nickel leads to the accumulation of overreduction products. Testing the reducing abilities of both NADH and NMNH under abiotic conditions showed that both molecules act as equally effective, soluble hydride donors in non-enzymatic, proto-metabolic stages of life's emergence.

Introduction

To form connections between undirected geochemical reactions and the first, directed, potentially even autocatalytic chemical reactions ultimately leading to metabolism, organic cofactors have dropped in and out of focus of research for decades.[1–6,6–10] These cofactors are a common denominator between "information first"[1,11] and "metabolism first" emergence of life hypotheses and thus might help to actively bridge this dated temporal separation of life's most central components. The prebiotic synthesis of several organic cofactors was described or hypothesized via numerous synthetic ways.[3,9,11]

Many central cofactors are structurally related, mostly via an adenosine-derived "tail" added to a chemically active group (s. Fig.1). The chemical properties and impact of this tail to the molecular dynamics of the molecule remain largely underexplored, with only a few exceptions.[5] Even when cofactors do not exhibit an obvious molecular connection such as adenosine, many (e.g. pterins or folates) derive from nucleotides in their biosynthesis.[8,12–14]

Figure 1. A selection of central organic cofactors with an adenosine-based tail (blue) connected to a functional part that determines the role of these cofactors in metabolism (black): Electron/hydride transfer, methyl transfer or acetyl-transfer. In grey, the function-associated half of nicotinamide dinucleotide (NAD) is highlighted: nicotinamide

Hence, many organic cofactors such as nicotinamide dinucleotide are often described as "RNA-derived"[10,15,16]. But we are proposing that RNA (and metabolism) could be considered "cofactor-derived". Based on theoretical work[2] cofactors could have preceded enzymatic reactions. They would constitute central parts of self-sustaining non-enzymatic reaction networks that intertwined with the synthesis of information-bearing molecules such as RNA and influenced the evolution/selection of enzymatic cofactor binding sites.[17,18]

An important aspect of evaluating the role of common nucleotide structures in organic cofactors as the adenosine-derived "tail" is to investigate its function in a non-biochemical setting.[5] So to take one of the most central organic cofactors, NAD, as an example: how much are the hydride transfers onto and from the nicotinamide influenced by the non-redox active adenosine part? Getting insight into the mechanisms in question is one of the main goals of this study.

NAD is a central redox cofactor in metabolism and is presumed to trace back to the Last Universal Common Ancestor (LUCA) of Archaea and Bacteria – and possibly beyond.[2,19,20] In a previous study we demonstrated that NAD[+] can react with H_2 and metal powder (Ni, Co, Fe) to specifically form the biologically

relevant form of reduced NAD (1,4-NADH) under hydrothermal conditions.[21] Another study achieved NAD[+] reduction without H_2 – but lower yields – with iron sulfides.[22] These publications introduced organic cofactors as possible transition points between geo- and biochemistry.

Across life's domains, the de novo synthesis of NAD implicates the synthesis of the deaminated form of NMN – nicotic acid mononucleotide (NaMN) – to which AMP is transferred from adenosine triphosphate (ATP) to form nicotinic acid adenine dinucleotide (NAAD). The ladder is then aminated to NAD[+]. All known de novo pathways are considered aerobic and are thought to have commonly emerged from an ancestral anaerobic pathway.[23,24] Anaerobic archaea require L-aspartate as a substrate for NaMN synthesis, and alternatively use fumarate as an electron acceptor, instead of oxygen.[25] Adenylylation (or AMPylation) is a common last step in the de novo pathway of coenzymes, usually from an ATP molecule.[26–29] In many organisms, but particularly in archaea and methanogens, an alternative salvage pathway directly adds ribose-5-phosphate to nicotinamide (Nam), producing nicotinamide mononucleotide (NMN), avoiding two extra steps of deamidation and amidation.[30]

When it comes to NAD's prebiotic synthesis pathways, several have been proposed e.g. via mineral-assisted synthesis under hydrothermal conditions[9,15] or achieved in parts via nitrile- and amino acid derived pathways.[11,31,32] Approaching prebiotic redox reactions, one has to think about a possible primary electron source in the geochemical setting. Here, we are mainly investigating the options of water-rock-interaction systems, where protons of water are being reduced to hydrogen (H_2) gas by electrons of Fe(II) containing minerals (serpentinizing systems).[33,34] Minerals found in such systems are known to promote hydrogen activation and electron transfer for CO_2 fixation.[35–37]

Apart from water-rock interactions constituting a rather simple, possibly wide-spread geological scenario on a mostly unknown early Earth, H_2 as an ancient electron source makes a lot of sense from a biological perspective as well. Looking at the reconstructed physiology of LUCA, it is likely that these first cells lived as anaerobic autotrophs, so were fixing carbon dioxide (CO_2) with the electrons provided by H_2[20,38–40] – a metabolic pathway still employed by many autotrophic prokaryotes today.[41] In order to access the electrons of H_2, these organisms possess enzymes called hydrogenases, which, depending on the exact organism, employ either Fe or Fe and Ni in their active centers.[42]

Incidentally, serpentinizing systems are rich in Fe and, depending on the system, can also be abundant in Ni as well. Hotter systems tend to feature Ni-Fe-alloys with higher nickel composition, while cooler ones are richer in iron.[43] Nickel-containing intermetallic compounds such as awaruite (Ni_3Fe) or taenite ($NiFe_3$ to Ni_2Fe) are products of the reaction of H_2 with Ni(II) compounds in serpentinizing systems[43–45] and are also found in meteorites.[46] Native metals are also not impossible to find, if the conditions on site are reducing enough.[47,48]

In this study, we first tested various Ni- and Fe-containing compounds as "proto-hydrogenases" for NAD^+ reduction and then compared NAD to NMN under hydrothermal conditions, to assess the relevancy of the adenosine-derived tail in pre-enzymatic reduction. We investigated the stability, reducibility and specificity of these organic cofactors and reduction products.

Results and Discussion

SCREENING NATURALLY OCCURRING IRON AND NICKEL CONTAINING MINERALS

Ni-Fe containing minerals found in hydrothermal settings were tested for the reduction of NAD^+ under conditions comparable to those found in mild serpentinizing hydrothermal systems (40 °C, 0.133 M phosphate buffer pH 8.5, 5 bar H_2, SI Scheme 1). These minerals were synthesized scratch via the nano-casting method by using tea leaves as template (s. SI methods) and were previously characterized.[36,49] Additionally, we conducted pre- and post-reaction characterization with Scanning Transmission Electron Microscopy (STEM) (SI Fig. 55–58). The metal content in these reactions is equivalent to that of the cofactor (1 metal atom per cofactor).

The resulting H_2 concentration at our conditions is 3.6 mM (using Henry's law, s. SI Eq. 1 and references [19,33]), which is comparable to the H_2 concentrations found in the effluent of serpentinizing systems.[50,51] The buffer was bubbled with N_2 for 1 h and handled inside a glove box to approximate the anoxic conditions on early Earth. Several controls were implemented, including controls without metal and H_2, respectively. The liquid and solid phase were analysed by ^{1}H-NMR and STEM, respectively (SI Fig. 25 and 26).

After 4 h, most of the dissolved NAD^+ was still preserved in solution under all reaction conditions except $NiFe_3$ which reacted faster (Fig. 2). Under H_2, 1,4- and 1,6-NADH formed in all samples at different yields (SI Table 5 and SI Fig. 25). The distribution of these two products is similar regardless of the mineral used, accumulating about 30% 1,6-NADH relative to the amount of NADH (SI Table 7 and SI Fig. 27). To ensure 1,6-NADH is a reduction product rather than a product of rearrangement, controls starting from 1,4-NADH over metal were conducted (SI Scheme 2 and SI Table 8). The majority of the 1,6-NADH yield can accordingly be attributed to reduction processes. Samples under Ar also produced reduced NAD when a Fe-rich mineral was used ($NiFe_3$, Fe; SI Table 6 and SI Fig. 26). In addition to transferring hydrides from H_2 to NAD^+, iron can oxidize, donating its own electrons either by producing nascent H_2 gas, ultimately reducing NAD^+ or by direct electron transfer to NAD^+. This process can also be used as a proxy for the constant H_2-production in serpentinizing systems[21]. Ni by itself is H_2-dependent in the promotion of NAD^+ reduction.

STEM imaging before and after the reactions (the latter including a washing and dilution step to assure true surface alteration) confirms that Fe, both under Ar and H_2, gets associated with phosphate ions in ratio that suggests the formation of iron phosphate, while Ni does not associate with phosphate, suggesting it stays in its native form (SI Fig. 57 and 58).[21]

Figure 2. NAD^+ reduction at 40 °C over 4 h with equimolar amounts (1 metal atom per NAD molecule) of several Ni-Fe-alloys and 5 bar of H_2 (or Ar). A) Segment of the ^{1}H-NMR spectra where the chemical shift of the hydrogen on the 2^{nd} nicotinamide carbon is visible upon reduction. 1,4-NADH features a characteristic peak at $\delta = 6.9$ ppm and 1,6-NADH at $\delta = 7.1$ ppm (SI Table 4). The ratio of products is similar among all minerals (3.5:1). B) Yield of 1,4-NADH obtained for several metals after 4 h under H_2 and Ar. Reduction under Ar is detected only with minerals whose metal content is ≥ 75% Fe. With 5 bar of H_2, all metals can facilitate 1,4-NADH synthesis. Mixed alloys are more efficient than the pure metals.

The Fe in Ni-rich minerals is expected to slowly reduce NAD under Ar conditions. However, it is likely that the resulting products do not reach the detection threshold within 4 h. Overall, bimetallic minerals are significantly more efficient than the individual transition metals when hydrogen is available. Introducing one Ni atom to a Fe atom increases the yield by 300% (NiFe vs Fe). Their properties, already observed in a previous study,[21] seem to complement each other for the reduction of NAD^+ with H_2: Fe being mostly an electron donor, while Ni promotes hydride transfer from H_2.

THE "UNIVERSAL" ADENINE NUCLEOTIDE IN ORGANIC COFACTORS

Many central cofactors share an adenosine-derived tail (Fig. 1) attached to a "functional" part. In the case of NAD, we define NMN as the functional part (containing the hydride-transferring nicotinamide) and adenosine monophosphate (AMP) as the adenosine-derived tail. NAD is stable in water, with its pH range depending on the reduction state of the nicotinamide: NADH is more stable at pH > 7, while NAD^+ is more stable under acidic conditions.[21] To investigate the role of the AMP-tail in a prebiotic context, several experiments were designed to compare NAD and NMN. 2D ^{1}H-NMR enabled the assignment of peaks for 1,4-NMNH and NMN in accordance with literature and in comparison to the pair NAD/NADH (SI Tables 2 and 3, and SI Fig. 1–19)).[52] We initially focussed on $NiFe_3$, the most efficient of the Ni-Fe minerals in the above described NAD experiment (s. Fig. 2). All other reaction conditions (buffer, pH, temperature, metal to cofactor ratio) were maintained (SI Scheme 3 and 4).

Figure 3. The reduction of NMN and NAD+ promoted with equimolar quantities of NiFe$_3$ nanopowder (normalized to the number of metal atoms) and 5 bar of H$_2$ (40 °C) was monitored for 4 h (SI Scheme 4 and 5). Both reactions were carried out in oxygen-free, aqueous solutions at a pH of 8.5 (0.133 M phosphate buffer). A) ^{1}H-NMR (600MHz) spectra monitoring NMN reduction over time. Dashed lines mark the peaks used for identifying and quantifying reduced NMN species previously identified via 2D NMR. Due to the complexity of the mixture not all peaks could be assigned (SI Figs. 38–51) B) Proposed mechanism for NMN reduction with H$_2$ and Ni-Fe minerals. The reduction site and additional protons are highlighted in red. Full arrows represent proposed reaction mechanisms supported by the data obtained, while the dashed arrow is an additional reaction that could not be excluded. Grey circles indicate the proton providing the NMR signal for quantification. C) Time course of NAD+ reduction. Reduced NAD species in a relative percentage to the starting concentration of NAD+ – all time points represent the mean and SD of at least triplicates of the same reaction. The ring chart represents the distribution of products after 4 h, percentage in the center indicates assigned products. D) Time course of NMN reduction. Reduced NMN species presented in a relative percentage to the starting concentration of NMN – all time points represent the mean and SD of at least triplicates of the same reaction. The ring chart represents the distribution of products after 4 h, percentage in the center indicates assigned products.

Reduction products were characterized through different 2D-NMR correlation spectra (SI Fig. 38–51) and Liquid Chromatography Mass Spectroscopy (LC-MS, SI Fig. 52–54. Products were quantified via 600 MHz ^{1}H-NMR spectroscopy using DSS as an internal standard (SI Table 9). Also controls with Ar instead of H$_2$ and without metal were conducted (SI Fig. 31Without metals, NMN does not react and remains stable (SI Fig. 28 and 30).Under Ar, NMN still gets reduced due to the abundant iron in the mineral compound, but more slowly than under H$_2$ (SI Tables 10 and 11).

An additional 2 h experiment with NAD+ under H$_2$ solidifies the hypothesis that the increase of 1,4- and 1,6-NADH is steady and inversely proportional to the decrease of NAD+ in solution (Fig. 3C, SI Scheme 4). After 4 h, on average 55% of NAD+ was reduced with 19% being oxidized. The remaining 25% can in part be attributed to nicotinamide formation in phosphate buffer and other degradation reactions (SI Table 12 and SI Fig. 32).[53]

NMN, however, shows a completely different reaction profile. After only 1 h, 69% of the starting NMN had been consumed, and a variety of products was observed in 1D ^{1}H-NMR (SI Scheme 5, SI Table 13 and SI Fig. 33). 2D-NMR spectroscopy facilitated the identification of the overreduction of NMN's nicotinamide ring with two and three hydrogenation sites, so 1,4,6-NMNH$_3$, and 1,2,4,6-NMNH$_5$ respectively (Fig. 3A, B and D; SI Fig. 38–51).

While the fully reduced species 1,2,4,6-NMNH$_5$ formed quickly and its concentration remained constant over time, the concentration of twice reduced 1,4,6-NMNH$_3$ increased with once reduced 1,4-NMNH decreasing. This indicates that not all reductions are a step-wise process (s. Fig. 3B).

Under Ar, 1,2,4,6-NMNH$_5$ did not form at all, demonstrating that H$_2$ is necessary for the full hydrogenation of the nicotinamide ring. 1,4,6-NMNH$_3$, however, was also forming under Ar, albeit in far lower yields (3%) than under H$_2$ (25%) after 4h (SI Figs. 31). The yield of 1,4 NMNH was comparable in both atmospheres (7% under Ar; 9% under H$_2$). Transferring these observations to environmental conditions suggests that less reducing conditions could be favourable for specific NMN reduction.

The above mentioned products accumulated in different amounts, 1,4-NMNH being the main product after 1 h under H$_2$ (Fig. 3A and D). Other side products formed at a comparable rate, rapidly depleting the reagent NMN. Consequently, the production of 1,4-NMNH came to halt halfway through the experiment (2 h) and subsequently began to decrease in concentration. The concentration of 1,4,6-NMNH$_3$ continuously increased over time. Even though the concentration of 1,4-NMNH decreased from 35.3% to 8.7% in 2 h, the total amount of reduced NMN remained

relatively stable, exceeding 39%. This suggests that 1,4-NMNH is the first and main product of NMN reduction, which can subsequently undergo further reduction to other species, mainly 1,4,6-NMNH$_3$.

We were able to exclude two products commonly found in NAD$^+$ reduction, where the 2nd or 6th carbon of the nicotinamide ring is reduced.[54–56] Subsequently, reduction products presumably starting with these two one-time reduced products could be excluded (SI Scheme 8 and SI Fig. 40–41). In the case of NADH, its 1,2-reduced form is known to be unstable, so it is likely this is the case with 1,2-NMNH as well, leading to its absence in our reaction (SI Scheme 9).[57]

In the case of 1,4-NMNH loss over time, we saw several routes: i) mainly the further reduction to 1,4,6-NMNH$_3$, ii) 1,4-NMNH getting hydrolysed at it 5th or 6th position, and iii) 1,4-NMNH engaging in a Diels-Alder type reaction with 1,6-NMNH (SI Scheme 10). Via LC-MS we were able to exclude Diels-Alder products and confirm it to be a hydration product, most likely OH$^-$ being added to 1,4-NMNH at the 6th position (Fig. 4B, SI Fig. 52–54) Ultimately we could assign the hydration product NMNH$_2$OH to be the peak at 7.35 ppm in the ^{1}H-NMR (SI Fig. 46 and 47). Both 1,4,6-NMNH$_3$ and NMNH$_2$OH have several conformational isomers, which result in several peaks within the ^{1}H-NMR spectrum as indicated in Fig. 4A.

An often reported side-product of NAD reduction (e.g. via cyclic voltammetry) is a 4,4'-linked NAD dimer,[58] which also qualifies as a possible side reaction of NMN reduction, too. Here, after careful interpretation of our 2D NMRs of the 1 h and 4 h reaction with NMN and the 4h reaction with NAD, we can exclude the presence of such dimers (s. SI Fig. 51; no peak at 40 to 50 ppm in ^{13}C of a bridging methine corresponding to the linkage). This was also confirmed via LC-MS (no double charged molecules were detected). As these dimers are a direct result of radical-forming 1e$^-$ transfers onto NAD,[59] we can draw the conclusion that direct hydride or 2e$^-$ transfer is the present mechanism in our reactions.

After quantification of all identified species, we can account for at least 68% of transformed NMN for all reactions, often more. Unidentified species encountered in lower yields can also stem from the differently reduced versions of the degradation product nicotinamide.[60] It is furthermore possible, that some NMN was lost due to strong interaction with the mineral surface.

In summary, we demonstrated that NMN is much more reactive than NAD$^+$. In 1 h, more NMN is consumed than NAD$^+$ in 4 h, under the same experimental conditions. The first and main product of both reactions seems to be 1,4-NADH/NMNH. However, while 1,4-NADH remains stable in solution, 1,4-NMNH quickly undergoes further reduction to form increasingly reduced products. Even when the experimental conditions are more reducing or using a higher metal to cofactor ratio, 1,4-NADH does not get overreduced and is stable.[21]

Where does this specificity for 1,4-NADH come from? It is known and well-described that NAD in aqueous solution alternates between a folded and open conformation,[61–64] could shield the nicotinamide ring from excessive overreduction – and possibly also from side reactions such as hydrolysis and Diels-Alder reactions described earlier (SI Scheme 10).

Figure 4. Selectivity and efficiency of NMN reduction with H$_2$ gas is influenced by the type of metal prevalent. The reduction of NMN assisted by NiFe alloys (NMN 1:1 nanopowder) yields products that directly correlate with the products obtained when utilizing the metals separately (NMN 1:200 micropowder). Ni^0 promotes the accumulation of 1,4,6-NMNH$_3$ and 1,2,4,6-NMNH$_5$ (green) and other side products. 1,4-NMNH (white) seems to not react further when using Fe^0, accumulating over time.

We deduce from these results that – in a mineral-dependent prebiotic context – the adenosine-derived tail can be essential for the targeted reduction of the 1,4-position, as well as for the stability of this specific reduction product. Nevertheless, we wanted to put to the test how much the choice of catalyst influences the outcome of the reactions.

TO EACH ITS OWN METAL AND MECHANISM

Starting from the observation that during NMN reduction, it's biologically active form 1,4-NMNH is a main product decreasing with the length of the reaction, we deducted that less efficient catalysts (e.g. less surface area) might help to reduce NMN more specifically than the previously used rather efficient nanoparticular NiFe-alloys.

As both Ni and Fe visibly worked less efficiently for NAD$^+$ reduction but Fe promotes reduction in higher yields, we decided to work with the same nanoparticular iron powder used for NAD$^+$ reduction (Fig. 2 and SI Scheme 3). While reducing NMN steadily, the Fe powder by itself (1:1 ratio to the cofactor) does not promote the formation of three times reduced 1,2,4,6-NMNH$_5$, while twice reduced 1,4,6-NMNH$_3$ is only produced in almost untraceable amounts (s. SI Table 10 and SI Fig. 28). The hydration product visible as a peak at 7.35 ppm still accumulates over the 4h reaction time. Other possible reduction products shown in SI Scheme 8 are not detected.

In order to explore whether this observation is a mere metal-dependent effect, we repeated experiments with NMN and H$_2$ over commercially available Fe and Ni micropowder (200:1 ratio to the cofactor; 50:1 in SI Scheme 7 and related figures), both metals pretreated under 5 bar of H$_2$ over night at 50 °C before adding the buffer solution containing the cofactor.. The results (Fig. 4) show a remarkable trend of overreduction with Ni micropowder (particle size: 3–7 μm), while Fe micropowder (particle size: <150 μm) mostly displays two main products: 1,4-NMNH and NMNH$_2$OH (SI Table 16). Comparing these spectra of

Ni only and Fe only promoted NMN reduction with those of NiFe nanopowder (Fig. 4; SI Fig. 37), it becomes apparent, that the distinct product patterns of each metal prevail within the mixture (note that this comparison is a qualitative one).

Nickel has long been recognized as a hydrogenation catalyst,[65] – but why does it, when not combined with Fe, only reluctantly reduce NAD (Fig. 2 and [21]) and yet overreduce NMN, not leaving any traceable amount of single-reduced species? The answer probably lies again in the structural differences between NAD and NMN. NMN can be more easily absorbed to a hydrogenated Ni surface, possibly over the entirety of its nicotinamide ring (s. SI Scheme 12). This could also explain the fast formation of its fully reduced product $NMNH_5$ shown in Fig. 4C. NAD in a staggered formation could only absorb partly on the surface, avoiding overreduction.[66]

If this is true, why does Fe not overreduce NMN as readily as Ni? Here, we can reflect on the mechanisms postulated by us in Pereira et al.,[21] that Fe both serves as a (less effective) hydrogenation catalyst and a strong electron donor, either via direct electron transfer to the nicotinamide cofactor or the formation of nascent hydride groups on its surface.

Assuming that Fe predominantly reduces NMN through direct electron transfer, the reduction process prioritizes the species with the most favourable redox potential first—namely, 1,4-NMNH (and 1,4-NADH, respectively). This hypothesis was substantiated by cyclic voltammetry measurements, which revealed that among all the reduction products obtained from NMN, 1,4-NMNH exhibits the highest oxidation potential (SI Fig. 20–24, 34 and 35 and SI 14).

Another possible explanation could be that the Ni catalyst does not alter as much as the Fe surface, meaning there would be a constant supply of hydrides available. For Fe, the previously described association with phosphate could block active centers which further prevent over reduction.

While the combination of nickel's hydrogenation strengths and iron's electron donation increases the yield of 1,4-NADH immensely compared to only Fe or only Ni (SI Fig. 36 and SI Table 5 and 15), the directed reduction of NMN to 1,4-NMNH does not occur in higher yields. In fact, adding nickel to the mix increases overreduction in such a way, that one could argue that NMN by itself could only have played a role as prebiotic hydride acceptor and donor in an environment depleted of Ni.

Thus, environmental conditions such as metal availability could have influenced the prebiotic selection process of redox cofactors. This study has shown that the adenosine-derived tail of NAD stabilizes the functional nicotinamide part of the molecule. Like that, the reduction and oxidation properties of NAD could have been maintained within a broader variety of environmental conditions than without the tail. At the same time, NMN by itself could have been a better hydride acceptor under nickel deprived, catalytically less efficient or less reducing conditions. Regardless, reducing NAD and NMN with the help of H_2 and metal catalysts is just one part of these cofactors' role in the prebiotic path towards the first functioning cells.

THE REDUCTION CAPABILITY OF NMNH AND NADH

Recently, it was shown, that Fe^{3+} ions (among other metal ions and also minerals) can promote the reaction of 1,4-NADH with pyruvate to lactate abiotically (SI Scheme 11).[10] Here, we show that both 1,4-NMNH and 1,4-NADH can reduce pyruvate to comparable amounts under aqueous conditions with Fe^{3+} in 17 h at 40°C, based on recently published experiments by Mayer and Moran.[10] These results underline that the adenosine nucleotide tail does not (strongly) influence the efficiency of the catalysed hydride transfer (Fig. 5; SI Figure 59–61)

Figure 5. Reduction of pyruvate to lactate with 1,4-NMNH and 1,4-NADH. The method of the reaction was performed according to Mayer and Moran (2023)[16] with the help of Fe(III) chloride as a catalyst in 17 h. Both nicotinamides perform equally well in this reaction. In both cases, lactate forms in comparable amounts – 2 repetitions per reaction. (for original NMR spectra s. SI Fig. 59 and 60).

Investigating the redox potential of both 1,4-NADH and 1,4-NMNH standards via cyclic voltammetry helped comparing their oxidation potential with that of the reaction mixtures of NiFe-assisted reduction of NAD^+ and NMN with H_2 (s. SI Fig. 20–24, 34 and 35 and SI Table 14). In the case of NiFe-assisted NAD^+ reduction, the resulting mixture shows only the oxidation potential of 1,4-NADH, while in the case of NMN reduction, the oxidation potential of both 1,4-NMNH and that of a second reduced species (most likely the species with the 2nd highest concentration 1,4,6-$NMNH_3$) is measured. As the second signal has a lower oxidation potential (meaning is harder to oxidize), we postulate that in the case of NMN, as for NAD, the 1,4-species is the most relevant reductant, not only in a biological but also in a prebiotic context.[67–69] One could argue that it is possible that the 1,4-position of an overreduced species would show a similar oxidation potential as a single reduced 1,4-species. However, as the oxidation of the latter leads to the aromatization of the nicotinamide ring, this reaction would be energetically favourable.

This ultimately leads to the conclusion that both 1,4-NMNH and 1,4-NADH are equally good hydride donors and could have both played a role in prebiotic hydride transfer. As the synthesis of both from their oxidized species is highly influenced by effectiveness and mechanism of the hydrogenation catalysts, it is possible that the environment had an important influence on the prebiotic selection of nicotinamides.

Conclusion

In this study, we have shown that both nicotinamide mono- and dinucleotide can be reduced under conditions found in serpentinizing systems, i.e. with H_2 gas promoted by Fe and Ni containing minerals. We further observed that the distinct structure of the molecules influences the reduction product spectrum associated with the nicotinamide ring.

There is a significant difference in how both molecules react to the addition of nickel to iron-containing catalysts – while most of the overreduction of NMN seems to be due to the contribution of nickel, the specific reduction of NAD^+ to 1,4-NADH and smaller amounts of 1,6-NADH with Ni-Fe-alloys is much faster than with iron or nickel alone. The overreduction of NMN, only consisting of nicotinamide and a phosphorylated ribose part, inevitably would lead to the molecule being a less effective hydride donor. The rather quick overreduction could be due to NMN being able to interact more directly with the mineral/metal surface than NAD. The additional AMP part attributes NAD its well-described staggered position in solution [61–64] which could prevent it from associating to the mineral surface the same way NMN does (SI Scheme 12). Further (theoretical) investigation will be necessary to get a definite answer.

Looking at the reverse reaction, only oxidation at the 1,4 position of the nicotinamide ring is observed in a biological context. Prebiotically, a 1,6-species could also be relevant for reduction, but this, to our knowledge, has not yet been observed in an experimental setup. However, it has been shown that the 1,4-species can participate in non-enzymatic reduction of further electron acceptors.[5,10] Here, we additionally confirmed that both 1,4-NMNH and 1,4-NADH act equally well as a hydride source in a non-enzymatic context. A next step would aim for the self-sustaining combination of both reduction and oxidation of these redox cofactors.

When it comes to the concrete question of prebiotic selection of organic cofactors, our study suggests that the ability to provide hydrides might not be the bottleneck, but rather the setup of the environment in which these reactions would have taken place. Overall, in the case of NAD and NMN, the dinucleotide would be more adapt to a variety of environments while the mononucleotide might be advantageous in an environment with less efficient catalysts (i.e. easier to reduce).

Lastly, our and other's observations of a stabilizing function of adenosine-derived tails could be applied other cofactors such as FAD, CoA or SAM. It seems feasible that also there, the extended structure could have been of merit in a prebiotic setting prior to a biological function.[70]

Supporting Information

The authors have cited additional references within the Supporting Information.[71–77]

Data Availability Statement

The data that support the findings of this study are available in the Supporting Information of this article. Original analysis files (LC-MS, NMR) will be provided by the corresponding authors upon reasonable request.

Acknowledgements

D:P:H.P an M.P. thank Bill Martin for discussions and support. D.P.H.P and M.P thank Alicia Casitas and team for providing access and help to cyclovoltammetry measurements. D.P.H.P. and M.P. thank Kendra Belthle for exploratory surface adsorption measurements and discussions. X.X. thanks Armin Geyer for discussions and the DFG funding for NMR spectrometer NEO600 (Forschungsgroßgeräte project number 508097909). M.P. thanks the Max Planck Society and the International Max Planck Research School 'Principles of microbial life' for funding. H.T. thanks MPG, the Volkswagen Foundation (96_742) and Deutsche Forschungsgemeinschaft (TU 315/8-1 / TU 315/8-3)This project was supported by the European Regional Development Fund (ERDF) and the Recovery Assistance for Cohesion and the Territories of Europe (REACT-EU).

Author contributions

Conceptualization: M.P. & D.P.H.P. Methodology: M.P. & D.P.H.P. Investigation: D.P.H.P., M.P., Z.S, T.B. Validation: D.P.H.P., Z.S. & X.X. Formal analysis: X.X., J.B., N.P., D.P.H.P. & M.P. Resources: H.T., K.V. Writing Original Draft: M.P. & D.P.H.P. Writing Review & Editing: all authors. Visualization: M.P., D.P.H.P. & X.X. Supervision: M.P.

Keywords: nucleotide cofactors• hydrogen • emergence of life • mineral catalysis • 2D NMR

References
[1]   A. D. Goldman, B. Kacar, *Journal of Molecular Evolution* **2021**, *89*, 127–133.
[2]   J. C. Xavier, W. Hordijk, S. Kauffman, M. Steel, W. F. Martin, *Proceedings of the Royal Society B: Biological Sciences* **2020**, *287*, 20192377.
[3]   A. Kirschning, *Natural Product Reports* **2021**, *38*, 993–1010.
[4]   S. Basak, S. Nader, S. S. Mansy, *JACS Au* **2021**, *1*, 371–374.
[5]   L. Sebastianelli, H. Kaur, Z. Chen, R. Krishnamurthy, S. S. Mansy, *Chemistry A European J* **2024**, e202400411.
[6]   H. B. White, *Journal of Molecular Evolution* **1976**, *7*, 101–104.
[7]   E. Werner, S. Pinna, R. J. Mayer, J. Moran, *Journal of American Chemical Society* **2023**, *145*, 21630–21637.
[8]   F. L. Sousa, W. F. Martin, *Biochimica et Biophysica Acta - Bioenergetics* **2014**, *1837*, 964–981.
[9]   J. C. Fontecilla-Camps, *Angewandte Chemie - International Edition* **2019**, *58*, 42–48.
[10]   R. J. Mayer, J. Moran, *Chem* **2024**, *10*, DOI 10.1016/j.chempr.2024.05.007.
[11]   H. J. Kim, S. A. Benner, *Chemistry - A European Journal* **2018**, *24*, 581–584.
[12]   D. E. Graham, R. H. White, *Nat. Prod. Rep.* **2002**, *19*, 133–147.
[13]   V. D. Crécy-Lagard, G. Phillips, L. L. Grochowski, B. E. Yacoubi, F. Jenney, M. W. W. Adams, A. G. Murzin, R. H. White, *ACS Chem. Biol.* **2012**, *7*, 1807–1816.

[14] G. Schwarz, R. R. Mendel, *Annu. Rev. Plant Biol.* **2006**, *57*, 623–647.

[15] S. D. Copley, E. Smith, H. J. Morowitz, *Bioorganic Chemistry* **2007**, *35*, 430–443.

[16] H. J. Cleaves, S. L. Miller, *Journal of Molecular Evolution* **2001**, *52*, 73–77.

[17] S. D. Fried, K. Fujishima, M. Makarov, I. Cherepashuk, K. Hlouchova, *Interface* **2022**, *19*, 20210641.

[18] A. C. Sanchez-Rocha, M. Makarov, L. Pravda, M. Novotný, K. Hlouchová, *eLife* **2024**, DOI 10.7554/eLife.94174.1.

[19] J. L. E. Wimmer, N. Vieira, J. C. Xavier, K. Kleinermanns, W. F. Martin, M. Preiner, *Microorganisms* **2021**, *9*, 458.

[20] J. L. E. Wimmer, J. C. Xavier, A. d. N. Vieira, D. P. H. Pereira, J. Leidner, F. L. Sousa, K. Kleinermanns, M. Preiner, W. F. Martin, *Frontiers in Microbiology* **2021**, *12*, 793664.

[21] D. P. Henriques Pereira, J. Leethaus, T. Beyazay, A. do Nascimento Vieira, K. Kleinermanns, H. Tüysüz, W. F. Martin, M. Preiner, *The FEBS Journal* **2022**, 16329.

[22] J. M. Weber, B. L. Henderson, D. E. LaRowe, A. D. Goldman, S. M. Perl, K. Billings, L. M. Barge, *Astrobiology* **2021**, *22*, 1–10.

[23] T. I. Gossmann, M. Ziegler, P. Puntervoll, L. F. De Figueiredo, S. Schuster, I. Heiland, *The FEBS Journal* **2012**, *279*, 3355–3363.

[24] A. Kirschning, *Nat. Prod. Rep.* **2022**, *39*, 2175–2199.

[25] H. Sakuraba, T. Satomura, R. Kawakami, S. Yamamoto, Y. Kawarabayasi, H. Kikuchi, T. Ohshima, *Extremophiles* **2002**, *6*, 275–281.

[26] T. P. Begley, C. Kinsland, E. Strauss, in *Vitamins & Hormones*, Elsevier, **2001**, pp. 157–171.

[27] E. Strauss, C. Kinsland, Y. Ge, F. W. McLafferty, T. P. Begley, *Journal of Biological Chemistry* **2001**, *276*, 13513–13516.

[28] A. Moreno, V. Taleb, M. Sebastián, E. Anoz-Carbonell, M. Martínez-Júlvez, M. Medina, *IUBMB Life* **2022**, *74*, 655–671.

[29] I. Saint-Girons, C. Parsot, M. M. Zakin, O. Barzu, G. N. Cohen, H. Weissbach, *Critical Reviews in Biochemistry* **1988**, *23*, S1–S42.

[30] S. Sharma, Y.-C. Hsieh, J. Dietze, M. Bockwoldt, Ø. Strømland, M. Ziegler, I. Heiland, *Metabolites* **2022**, *12*, 569.

[31] J. Yi, H. Kaur, W. Kazöne, S. A. Rauscher, L. Gravillier, K. B. Muchowska, J. Moran, *Angewandte Chemie* **2022**, *134*, DOI 10.1002/ange.202117211.

[32] M. Bechtel, N. Kurrle, O. Trapp, *Chemistry A European J* **2024**, e202402055.

[33] L. Schwander, M. Brabender, N. Mrnjavac, J. L. E. Wimmer, M. Preiner, W. F. Martin, *Front. Microbiol.* **2023**, *14*, 1257597.

[34] D. E. Allen, W. E. Seyfried, *Geochimica et Cosmochimica Acta* **2004**, *68*, 1347–1354.

[35] M. Preiner, K. Igarashi, K. Muchowska, M. Yu, S. J. Varma, K. Kleinermanns, M. K. Nobu, Y. Kamagata, H. Tüysüz, J. Moran, W. F. Martin, *Nature Ecology & Evolution* **2020**, *4*, 534–542.

[36] T. Beyazay, C. Ochoa-Hernández, Y. Song, K. S. Belthle, W. F. Martin, H. Tüysüz, *Angewandte Chemie - International Edition* **2023**, *62*, e202218189.

[37] T. Beyazay, K. S. Belthle, C. Farès, M. Preiner, J. Moran, W. F. Martin, H. Tüysüz, *Nature Communications* **2023**, *14*, 570.

[38] M. C. Weiss, F. L. Sousa, N. Mrnjavac, S. Neukirchen, M. Roettger, S. Nelson-Sathi, W. F. Martin, *Nature Microbiology* **2016**, *1*, 16116.

[39] E. R. R. Moody, S. Álvarez-Carretero, T. A. Mahendrarajah, J. W. Clark, H. C. Betts, N. Dombrowski, L. L. Szánthó, R. A. Boyle, S. Daines, X. Chen, N. Lane, Z. Yang, G. A. Shields, G. J. Szöllősi, A. Spang, D. Pisani, T. A. Williams, T. M. Lenton, P. C. J. Donoghue, *Nat Ecol Evol* **2024**, DOI 10.1038/s41559-024-02461-1.

[40] M. Brabender, D. P. Henriques Pereira, N. Mrnjavac, M. L. Schlikker, Z.-I. Kimura, J. Sucharitakul, K. Kleinermanns, H. Tüysüz, W. Buckel, M. Preiner, W. F. Martin, *Proc. Natl. Acad. Sci. U.S.A.* **2024**, *121*, e2318969121.

[41] G. Fuchs, *Annual Review of Microbiology* **2011**, *65*, 631–658.

[42] R. K. Thauer, *European Journal of Inorganic Chemistry* **2011**, *7*, 919–921.

[43] R. Frost, *Journal of Petrology* **1985**, *26*, 31–63.

[44] F. Klein, W. Bach, *Journal of Petrology* **2009**, *50*, 37–59.

[45] D. I. Foustoukos, M. Bizimis, C. Frisby, S. B. Shirey, *Geochimica et Cosmochimica Acta* **2015**, *150*, 11–25.

[46] J. A. P. M. Devienne, T. A. Berndt, W. Williams, L. Nagy, *Geophysical Research Letters* **2023**, *50*, e2022GL102602.

[47] V. Dekov, *J. Geophys. Res.* **2006**, *111*, 2005JB003955.

[48] C. Kanellopoulos, E. Valsami-Jones, P. Voudouris, C. Stouraiti, R. Moritz, C. Mavrogonatos, P. Mitropoulos, *Geosciences* **2018**, *8*, 287.

[49] M. Yu, G. Moon, E. Bill, H. Tüysüz, *Applied Energy Materials* **2019**, *2*, 1199–1209.

[50] J. L. Charlou, J. P. Donval, Y. Fouquet, P. Jean-Baptiste, N. Holm, *Chemical Geology* **2002**, *191*, 345–359.

[51] C. Konn, J. P. Donval, V. Guyader, Y. Germain, A. S. Alix, E. Roussel, O. Rouxel, *Deep-Sea Research Part I: Oceanographic Research Papers* **2022**, *179*, 103630.

[52] R. Zapata-Pérez, A. Tammaro, B. V. Schomakers, A. M. L. Scantlebery, S. Denis, H. L. Elfrink, J. Giroud-Gerbetant, C. Cantó, C. López-Leonardo, R. L. McIntyre, M. Van Weeghel, Á. Sánchez-Ferrer, R. H. Houtkooper, *The FASEB Journal* **2021**, *35*, DOI 10.1096/fj.202001826R.

[53] L. Jr. Rover, J. C. B. Fernandes, G. Neto, de Oliveira, L. T. Kubota, E. Katekawa, S. H. P. Serrano, *Analytical Biochemistry* **1998**, *55*, 50–55.

[54] T. Saba, J. Li, J. W. H. Burnett, R. F. Howe, P. N. Kechagiopoulos, X. Wang, *ACS Catalysis* **2021**, *11*, 283–289.

[55] T. Saba, J. W. H. Burnett, J. Li, P. N. Kechagiopoulos, X. Wang, *Chemical Communications* **2020**, *56*, 1231–1234.

[56] S. E. Godtfredsen, M. Ottesen, N. R. Andersen, *Carlsberg Research Communications* **1979**, *44*, 65–75.

[57] S. Chaykin, L. Meissner, *Biochemical and Biophysical Research Communications* **1964**, *14*, 233–240.

[58] I. Ali, T. Khan, S. Omanovic, *Journal of Molecular Catalysis A: Chemical* **2014**, *387*, 86–91.

[59] N. H. A. Besisa, K.-S. Yoon, M. Yamauchi, *Chem. Sci.* **2024**, *15*, 3240–3248.

[60] A. Zarei, L. Khazdooz, M. Enayati, S. Madarshahian, T. J. Wooster, G. Ufheil, A. Abbaspourrad, *RSC Adv.* **2021**, *11*, 21036–21047.

[61] R. V. Hull, P. S. Conger, R. J. Hoobler, *Biophysical Chemistry* **2001**, *90*, 9–16.

[62] P. E. Smith, J. J. Tanner, *Journal of Molecular Recognition* **2000**, *13*, 27–34.

[63] S. K. Hoffmann, J. Goslar, S. Lijewski, *Appl Magn Reson* **2013**, *44*, 817–826.

[64] R. Parthasarathy, S. M. Fridey, *Science* **1984**, *226*, 969–971.

[65] H. Adkins, H. I. Cramer, *J. Am. Chem. Soc.* **1930**, *52*, 4349–4358.

[66] A. Damian, S. Omanovic, *Journal of Molecular Catalysis A: Chemical* **2006**, *253*, 222–233.

[67] R. A. Weusthuis, P. L. Folch, A. Pozo-Rodríguez, C. E. Paul, *iScience* **2020**, *23*, 101471.

[68] R. D. Braun, K. S. V. Santhanam, P. J. Elving, *J. Am. Chem. Soc.* **1975**, *97*, 2591–2598.

[69] P. J. Elving, C. O. Schmakel, K. S. V. Santhanam, P. Zuman, *C R C Critical Reviews in Analytical Chemistry* **1976**, *6*, 1–67.

[70] L. M. Longo, J. Jablonska, P. Vyas, M. Kanade, R. Kolodny, N. Ben-Tal, D. S. Tawfik, **2020**, *9*, e64415.

[71] S. J. Varma, K. B. Muchowska, P. Chatelain, J. Moran, *Nature Ecology and Evolution* **2018**, *2*, 1019–1024.

[72] S. Berger, S. Braun, *200 and More NMR Experiments, a Practical Course*, WILEY-VCH Verlag, Weinheim, Germany, **2004**.

[73] L. Patiny, A. Borel, *J. Chem. Inf. Model.* **2013**, *53*, 1223–1228.

[74] R. R. Ernst, G. Bodenhausen, *Principles of Nuclear Magnetic Resonance in One and Two Dimensions, Oxford University Press*, Oxford University Press, Oxford, **1997**.

[75] K. Wüthrich, *NMR of Proteins and Nucleic Acids*, John Wiley & Sons, Ltd, New York, **1986**.

[76] J. N. S. Evans, *Biomolecular NMR Spectroscopy*, Oxford University Press, Oxford, **1996**.

[77] H.-O. Kalinowski, S. Berger, S. Braun, *[13]C-NMR-Spektroskopie*, Georg Thieme Verlag, Stuttgart, **1988**.