The polyadenylase PAPI is required for virulence plasmid maintenance in pathogenic bacteria

Katherine Schubert[1], Micah Braly[2], Jessica Zhang[3], Michele E. Muscolo[4], Hanh N. Lam[2*], Karen Hug[2], Henry Moore[5], Joshua W. McCausland[3,6,7], Derfel Terciano[2], Todd Lowe[5], Cammie F. Lesser[4,8,9], Christine Jacobs-Wagner[3,6,7,10], Helen Wang[11], Victoria Auerbuch[2#]

[1]Department of Molecular, Cell, and Developmental Biology, UC Santa Cruz, Santa Cruz, CA 95064, United States

[2]Department of Microbiology and Environmental Toxicology, UC Santa Cruz, Santa Cruz, CA 95064, United States

[3]Department of Biology, Stanford University, Stanford, CA 94305, United States

[4]Center for Bacterial Pathogenesis, Division of Infectious Diseases, Department of Medicine, Massachusetts General Hospital, MA 02115, United States

[5]Department of Biomolecular Engineering, UC Santa Cruz, Santa Cruz, CA 95064, United States

[6]Sarafan ChEM-H Institute, Stanford University, Stanford, CA 94305, United States

[7]Howard Hughes Medical Institute, Stanford University, Stanford, CA 94305, USA

[8]Department of Microbiology, Blavatnik Institute, Harvard Medical School, Boston, MA 02115, United States

[9]Broad Institute of MIT and Harvard, Cambridge, MA 02142, United States

[10]Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA 94305, United States

[11]Department of Medical Biochemistry and Microbiology, Uppsala University, Sweden

* Current address: Department of Biological Sciences, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, United States

Abstract

Many species of pathogenic bacteria harbor critical plasmid-encoded virulence factors, and yet

the regulation of plasmid replication is often poorly understood despite playing a critical role in

plasmid-encoded gene expression. Human pathogenic *Yersinia*, including the plague agent *Y.*

pestis and its close relative *Y. pseudotuberculosis*, require the type III secretion system (T3SS)

virulence factor to subvert host defense mechanisms and colonize host tissues. The *Yersinia*

T3SS is encoded on the IncFII plasmid for *Yersinia* virulence (pYV). Several layers of gene

regulation enables a large increase in expression of *Yersinia* T3SS genes at mammalian body

temperature. Surprisingly, T3SS expression is also controlled at the level of gene dosage. The

number of pYV molecules relative to the number of chromosomes per cell, referred to as

plasmid copy number, increases with temperature. The ability to increase and maintain elevated

pYV plasmid copy number, and therefore T3SS gene dosage, at 37°C is important for *Yersinia*

virulence. In addition, pYV is highly stable in *Yersinia* at all temperatures, despite being

dispensable for growth outside the host. Yet how *Yersinia* reinforces elevated plasmid

replication and plasmid stability remains unclear. In this study, we show that the chromosomal

gene *pcnB* encoding the polyadenylase PAP I is required for regulation of pYV plasmid copy

number (PCN), maintenance of pYV in the bacterial population outside the host, robust T3SS

activity, and *Yersinia* virulence in a mouse infection model. Likewise, *pcnB*/PAP I is also

required for robust expression of the *Shigella flexneri* virulence plasmid-encoded T3SS.

Furthermore, *Yersinia* and *Shigella pcnB*/PAP I is required for maintaining normal PCN of model

antimicrobial resistance (AMR) plasmids whose replication is regulated by sRNA, thereby

increasing antibiotic resistance by ten-fold. These data suggest that *pcnB*/PAP I contributes to

the spread and stabilization of virulence and AMR plasmids in bacterial pathogens, and is

essential in maintaining the gene dosage required to mediate plasmid-encoded traits.

Importantly *pcnB*/PAP I has been bioinformatically identified in many species of bacteria despite

being studied in only a few species to date. Our work highlights the potential importance of

pcnB/PAP I in antibiotic resistance, and shows for the first time that *pcnB*/PAP I reinforces PCN

and virulence plasmid stability in natural pathogenic hosts with a direct impact on bacterial

virulence.

Author Summary

Many pathogens carry extrachromosomal DNA elements known as plasmids, which encode

genes that confer bacterial virulence or antimicrobial resistance (AMR). Acquisition of these

plasmids by bacteria can lead to the emergence of new pathogenic traits and the spread of

AMR, yet the mechanisms by which plasmids are retained in bacterial populations particularly in

the absence of selective pressure remain incompletely understood. Here we show that the

major bacterial polyadenylase enzyme PAP I, encoded by the *pcnB* gene, is critical for the

human pathogen *Yersinia pseudotuberculosis* to maintain its native virulence plasmid as well as

AMR plasmids. Very little is known about the process of polyadenylation in bacteria, or the post-

transcriptional addition of adenosine residues to the 3' end of transcripts. This study represents

the first demonstration that PAP I-mediated polyadenylation contributes to bacterial

pathogenesis.

Introduction

Many species of pathogenic bacteria harbor plasmids that encode genes important for virulence

or antimicrobial resistance (AMR). Virulence plasmids are typically characterized by their

relatively large size (>40kb) and low plasmid copy number (PCN), which refers to the average

number of plasmid copies per cell or chromosome equivalent [1]. In addition to virulence or

AMR genes, these plasmids typically contain additional genes important for plasmid replication,

PCN control, and plasmid maintenance. Virulence plasmids are commonly found in members of

the Enterobacteriaceae family, which includes several species of Gram-negative pathogens. For

example, *Yersinia pseudotuberculosis* and the related plague agent *Yersinia pestis* naturally

harbor a ~70 kb low copy number virulence plasmid known as the *plasmid for Yersinia virulence*

(pYV), also called pCD1 in *Y. pestis* [2-4]. Given that *Y. pestis* has killed over 200 million people

since antiquity [5], pYV has played a grim but important role in human history.

The pYV plasmid encodes several factors important for *Yersinia* virulence, including a major

virulence factor known as the type III secretion system (T3SS) [6-8]. The T3SS is used to inject

a series of effector proteins known as *Yersinia outer proteins* (Yops) directly into the cytosol of

target host cells during infection. Yops dampen the host immune response and promote

Yersinia pathogenesis, making pYV essential for *Yersinia* virulence [3, 9-13]. Expression and

activity of the T3SS exerts a metabolic burden on the bacteria. Therefore, *Yersinia* repress

expression of the T3SS at low environmental temperatures when the T3SS is dispensable, but

induce T3SS expression at mammalian body temperature, 37°C [14-19]. Several mechanisms

contribute to this temperature dependency. For example, the master regulator of the T3SS is

only translated at 37°C due to control by an RNA thermometer [18, 20]. More recently, it was

reported that temperature-dependent changes in T3SS gene expression is also regulated at the

level of gene dosage via changes in pYV PCN [21]. At environmental temperatures such as

26°C, pYV PCN has been reported to be low (~1-2 copies per chromosome). Upon exposure to

mammalian temperatures of 37°C, pYV PCN has been shown to be rapidly upregulated to 3+

copies per chromosome. Importantly, this ability to upregulate pYV PCN and T3SS gene

dosage is essential for normal T3SS activity and *Y. pseudotuberculosis* virulence [21, 22].

Dynamic changes in pYV PCN and retention of the pYV plasmid during these rapid changes are

therefore required for the *Y. pseudotuberculosis* pathogenic lifestyle.

The pYV plasmid encodes several systems important for plasmid regulation and maintenance,

including a *cop-rep* locus that regulates plasmid replication, a ParAB partitioning system that

ensures each daughter cell inherits the plasmid, and a ParDE toxin-antitoxin (TA) system that

promotes plasmid maintenance in the bacterial population [1, 23]. Together, these systems are

predicted to synergistically enforce plasmid stability, or the percentage of cells in the population

that maintain the plasmid. The highly conserved *cop-rep* locus places pYV in the IncFII group of

plasmids [1, 24, 25]. The *cop-rep* locus encodes the essential <u>*replication initiation factor*</u> RepA

and two negative regulators of RepA expression: the CopA small RNA (sRNA) and the CopB

protein. The functions of RepA, CopA, and CopB have been elucidated using plasmid R1, an

IncFII plasmid originally isolated from a clinical strain of *S. enterica* that has been extensively

studied in *Escherichia coli* model systems [25-27]. These studies support the notion that CopA

and CopB regulate the rate of plasmid replication to minimize fluctuations above or below the

average PCN set-point [25, 28].

Using *Y. pseudotuberculosis* as a model, we identified the chromosomal *pcnB* gene as a

previously unknown regulator of pYV replication and stability. The *pcnB* gene was named for

<u>p</u>lasmid <u>c</u>opy <u>n</u>umber <u>B</u> since loss of *pcnB* was found to decrease PCN for several model

plasmids in *E. coli* [29, 30]. The *pcnB* gene encodes the poly A polymerase (PAP I), the major

bacterial polyadenylase ubiquitous in β- and γ-proteobacteria [29-36]. Polyadenylation, or the

addition of adenosine residues to the 3' OH of an RNA transcript, is poorly understood in

bacteria compared to eukaryotes. PAP I-mediated polyadenylation promotes RNA degradation

as part of a housekeeping mechanism [37], but is also implicated in certain gene regulatory

networks [38-43]. PAP I-dependent polyadenylation has been shown to destabilize RNAs that

negatively regulate replication of certain plasmid types in *E. coli*, including CopA from plasmid

R1 and RNAI from ColE1 plasmids [29, 44-47]. However, PAP I has not been studied in the

context of a naturally occurring virulence or AMR plasmid within its native host, and has never

been linked to bacterial virulence.

Here, we report that *Y. pseudotuberculosis pcnB* mutants display reduced PCN and decreased

stability of pYV. This defect is associated with reduced T3SS activity in bacterial culture and

impaired virulence in a mouse infection model. Furthermore, we show that *pcnB* is required for

Y. pseudotuberculosis to maintain model AMR plasmids belonging to various plasmid families,

highlighting its crucial role in bacterial pathogenicity and antibiotic resistance.

Results

Y. pseudotuberculosis toggles pYV plasmid copy number in response to temperature

To determine the average PCN per chromosome equivalents at the bacterial population level,

we used droplet digital (dd)PCR to measure the number of pYV molecules relative to

chromosomes in wildtype *Y. pseudotuberculosis* IP2666pIB1 [22]. Cells were either grown at

26°C to represent growth in environmental conditions outside of the host or at 37°C in low

calcium media to represent host cell contact during mammalian infection. The pYV-encoded

T3SS is expressed at 37°C and becomes active for secretion following host cell membrane

contact or calcium chelation [48-51]. Consistent with previous studies [21, 22], we observed an

increase in pYV PCN from an average of ~1.5 pYV copies per chromosome equivalents at 26°C

to ~3 copies at 37°C/low calcium (Fig 1A). In addition, using a YPIII/pIBX strain of *Y.*

pseudotuberculosis harboring a pYV-encoded luciferase reporter, we found that pYV PCN

relative to cell density increases one hour after shifting the cultures to 37°C/low calcium

conditions (Fig 1B).

To examine pYV PCN at the single-cell level, we employed a microscopy-based approach. To

visualize and quantify pYV puncta in living *Y. pseudotuberculosis* cells, we generated a

YPIII/pIBX *Y. pseudotuberculosis* strain that harbors a *parB-msfGFP* translational fusion in the

native *parB* pYV locus. This approach has been used to visualize plasmids in other bacterial

species [52, 53], as ParB binds to plasmid-encoded *parS* sequences but has never been used

to study pYV in *Y. pseudotuberculosis*. At 26°C, cells were found to have an average of about

four pYV puncta per cell (Fig 1C-D). This number increased to an average of about seven

puncta per cell by 1.5 hours after the shift to 37°C (Fig 1C-D). The higher number of pYV per

cell than per chromosome equivalents (Figs 1D vs. 1A) is consistent with the expectation that

Yersinia has more than one chromosome per cell on average under nutrient-rich (fast) growth

conditions, similar to *E. coli* [54, 55]. Regardless of the normalization method (per cell or

chromosome), *Yersinia* pYV PCN consistently increased by ~1.5-fold upon shifting bacterial

cultures from 26°C to 37°C/low calcium conditions. The pYV plasmid was highly stable in *Y.*

pseudotuberculosis, with the vast majority of cells or clones retaining pYV (Fig 1D).

Disruption of the *pcnB* gene encoding the polyadenylase PAP I decreases pYV PCN and

T3SS expression

Y. pseudotuberculosis with a functional T3SS, but lacking T3SS Yop effectors (Δyop6), induces

activation of the transcription factor NFκB in mammalian cells [56]. The LPS metabolite ADP

heptose is delivered to the host cell cytoplasm through the T3SS, stimulating Alpk1/TIFA-

dependent NFκB activation [57], although YopJ inhibits NFκB signaling [58, 59]. We used

human embryonic fibroblast cells, HEK293T, expressing an NFκB luciferase reporter to identify

a *Y. pseudotuberculosis* IP2666pIB1 Δyop6 transposon (Tn) mutant that induced elevated

bioluminescence compared to the parental strain, (Δyop6/*pil*::Tn; Fig S1A) [56]. This mutant was

found to harbor a Tn insertion in a previously unannotated locus adjacent to the highly

conserved pYV *cop-rep* locus. We will refer to this locus as the plasmid inhibitory locus (*pil*) (Fig

S1B). Given the location of the *pil* locus relative to *cop-rep*, we hypothesized that the *pil*::Tn

mutation affects the pYV PCN. To test this, we introduced the *pil::Tn* insertion mutation into a

clean wildtype *Y. pseudotuberculosis* IP2666pIB1 strain background with intact T3SS effector

Yop-encoding genes (referred to as *pil*::Tn). Consistent with our hypothesis, the *pil::Tn* mutant

displayed elevated secretion of the T3SS cargo proteins YopE, YopD, and YopH compared to

wildtype *Yersinia* (Fig S1C). In addition, we constructed a mutant in the YPIII/pIBX background

with a plasmid insertion into the same locus in *pil* (*pil*::pNQ). As expected, both the *pil*::Tn and

pil::pNQ strains had elevated pYV PCN at 26°C and early after shifting to 37°C relative to the

parental strain (Fig S1DE), which is likely the underlying cause of the increased T3SS activity .

Insertion into the *pil* locus was associated with poor growth compared to the parental strain

under T3SS-inducing conditions, and suppressor mutants could be readily isolated (Fig S1F).

We hypothesized that constitutively elevated pYV PCN contributed to this growth defect,

resulting in selective pressure favoring suppressor mutations that would alleviate high pYV

PCN. Therefore, we used the *pil* insertion mutation as a tool to identify genes important for pYV

PCN homeostasis. We carried out whole genome sequencing of a subset of *pil*::pNQ

suppressor mutants and identified several categories of mutations affecting growth, pYV PCN,

and the T3SS (Table S1). Two clones harbored a single point mutation that was associated with

reduced pYV PCN and T3SS activity relative to the *pil*::pNQ parental strain. This mutation was

mapped to the *pcnB* gene encoding the polyadenylase PAP I, and was predicted to confer a

leucine to arginine change at position 291 (L291R) in the PAP I neck domain linked to RNA

binding (Fig 2A). The L291 residue is conserved in PAP I from *E. coli*, suggesting that the

residue might be important for PAP I structure and/or function (Fig 2A). We will use the term

pcnB when referring to the gene, and PAP I when referring to the *pcnB*-encoded protein.

To confirm that the *pcnB* gene encoding PAP I affects *Y. pseudotuberculosis* pYV PCN, we

generated a full in-frame deletion of the *pcnB* gene (Δ*pcnB*) in the wildtype and *pil*::Tn

IP2666pIB1 strain backgrounds. We used ddPCR to measure relative pYV PCN and found that

compared to wildtype, the Δ*pcnB* mutant had lower pYV PCN following growth at 26°C and

37°C/low calcium (Fig 2B). In keeping with our previous findings, the *pil*::Tn mutant displayed

significantly higher pYV PCN at 26°C compared to wildtype. Furthermore, loss of PAP I in the

pil::Tn*/*Δ*pcnB* double mutant significantly reduced pYV PCN at 26°C compared to the *pil*::Tn

parental strain. These results show that loss of PAP I leads to decreased pYV PCN. Despite

this, loss of *pcnB* did not prevent *Yersinia* from increasing pYV PCN upon shift from 26°C to

37°C, although not to the same high PCN as in wildtype (Fig 2B). Consistent with this reduction

in T3SS gene dosage, the Δ*pcnB* mutant displayed two-fold less expression and secretion of

the T3SS effector protein YopE compared to wildtype *Y. pseudotuberculosis* (Fig 2C). Similarly,

pcnB deletion reduced YopE expression and secretion in the *pil*::Tn strain background (Fig 2C).

Taken together, these data suggest that *pcnB* is essential for maintaining pYV PCN, and that

this function is important for enabling robust T3SS expression under T3SS-inducing conditions.

To determine how the *pcnB* gene and the *pil*::Tn disruption broadly impact expression of pYV-

encoded genes, we carried out RNA-seq analyses on wildtype, Δ*pcnB*, *pil*::Tn, and

pil::Tn*/*Δ*pcnB Y. pseudotuberculosis* IP2666pIB1 grown at 26°C and 37°C/low calcium

(Supplementary Dataset 1). Most pYV-encoded gene transcripts were present at higher steady-

state level in the *pil*::Tn mutant compared to wildtype *Y. pseudotuberculosis* at both

temperatures (Fig S2), consistent with an increase in pYV gene dosage. However, T3SS genes

are not translated at 26°C due to the action of several RNA thermometers in *Yersinia* [48], and

the *pil*::Tn strain does not display T3SS activity at 26˚C despite elevated T3SS gene transcripts

at this temperature (Fig S3). Interestingly, the *pil*::Tn strain displayed altered the RNA-Seq read

pattern of the region containing *pil*, *copB*, and *repA* at both temperatures (Fig S4). We observed

no difference in pYV PCN or T3SS expression when we generated a full in-frame deletion of the

predicted ORF found in the *pil* locus (Fig S1C,E). Therefore, we propose that Tn or pNQ

insertion into *pil* leads to derepressed transcription of the *cop-rep* locus leading to elevated pYV

replication through an as-yet unknown mechanism. In contrast, deletion of *pcnB* resulted in

fewer transcripts of most pYV-encoded genes compared to wildtype *Yersinia* at both

temperatures (Fig S2), consistent with a decrease in pYV gene dosage. In addition, *pcnB*

deletion in the *pil*::Tn background also led to a decrease in the steady-state levels of most T3SS

genes compared to the *pil*::Tn parental strain (Fig S2). We conclude that loss of *pcnB* leads to a

decrease in pYV PCN in both the wildtype and *pil*::Tn backgrounds, leading to decreased

transcription of pYV-encoded genes and mitigation of the T3SS activity-induced growth arrest.

PAP I is required for pYV stability

When working with the ∆*pcnB* strain, we noted that cells lost the pYV plasmid at a higher rate

than observed with wildtype *Y. pseudotuberculosis* (Table 1). To determine whether PAP I

affects pYV stability in addition to pYV PCN, we generated a ∆*pcnB* strain in the ParB-msfGFP

YPIII/pIBX *Y. pseudotuberculosis* background and observed cells lacking clear fluorescent foci

under both non-T3SS-inducing (26˚C) and T3SS-inducing conditions (37˚C, low calcium media.

We observed ∆*pcnB* cells with lower cellular GFP fluorescence, consistent with plasmid loss

given that the msfGFP-labeled ParB is encoded by pYV (Fig 3A). The presence of these

plasmid-devoid cells in the ΔpcnB strain resulted in a clear bimodal distribution of cellular

fluorescence (Fig 3B), showing that ~34% of ΔpcnB cells lacked pYV (low cellular fluorescence)

compared to only ~3% for the wild-type population at the time of imaging at 26ºC (p<10^{-100},

Kolmogorov-Smirnov (KS) test; n = 7033 cells per strain). The ΔpcnB mutant also displayed a

high percentage (24%) of cells lacking pYV under the 37°C/low calcium condition based on the

fraction of cell population with low cellular fluorescence (Fig 3B). By analyzing only those cells

that retained at least one copy of pYV, we found that the ΔpcnB mutant showed the expected

increase in ParB-msfGFP puncta per cell from 26°C to 37°C/low calcium (Fig 3B-C). However,

the distributions were significantly shifted toward lower values relative to the wildtype strain

(p<10^{-100}; Fig 3D-E). While the wildtype strain had, on average, 4.5 and 7.9 pYV spots per cell

at 26°C and 37°C/low calcium, respectively, the ΔpcnB strain had 3.9 and 6.3. Thus, the pcnB

gene does not affect the ability of the cells to respond to the T3SS-inducing environmental shift,

but its loss leads to lower average PCN and increased frequency of plasmidless Yersinia cells.

To further examine the extent of the pYV plasmid stability defect in the ΔpcnB mutant, we

passaged bacteria on solid media for several days at 26°C and then assessed pYV retention for

each clone by patching onto low calcium media containing the dye Congo red, and incubating at

37°C. Congo red only binds to bacteria that are expressing a functional T3SS [60]. Wildtype

cells retained the pYV plasmid at 100% while the ΔpcnB mutant cells only retained pYV in 82%

of cells following incubation at 26°C for 16 hours during passaging (Table 1). When cells were

exposed to T3SS-inducing conditions at 37°C/low calcium and returned to 26°C to mimic the Y.

pseudotuberculosis facultative lifecycle, we found that only 58% of ΔpcnB cells retained pYV

compared to 99% for wildtype (Table 1). This observed increase in pYV loss likely stems from

the growth arrest of pYV⁺ clones specifically under T3SS-inducing conditions at 37°C [15, 49,

50], which provides a selective growth advantage for ∆pcnB clones that have lost pYV.

Altogether, these data support a role for PAP I in regulation of pYV PCN and plasmid stability

during the *Y. pseudotuberculosis* facultative lifecycle.

Table 1. PAP I is required for pYV maintenance in *Y. pseudotuberculosis*

Strain:	Percent Retention[ŧ] of pYV after growth at 26°C only:	Percent Retention[ŧ] of pYV after 16 hr exposure to 37°C:
Wildtype	100.0 ±0	99.1 ±1.2
∆pcnB	81.6 ±17.3	57.7 ±10.7
pil::Tn	100.0 ±0	99.1 ±1.3
pil::Tn ∆pcnB	98.1 ±1.8	75.9 ±13.1

251
[ŧ]Percent retention of the pYV plasmid was averaged from three independent experiments where ±
represents the standard deviation between experiments.

The PAP I[L291R] mutation affects PAP I protein levels.

The original *pcnB* suppressor mutation identified in our *pil*::Tn suppressor screen was a single

point mutation that changed leucine in position 291 to an arginine (L291R) (Fig 2A). Although

the corresponding L291 residue is conserved in *E. coli*, it has not yet been identified as a critical

PAP I residue. Thus, as a first step in investigating the role of this residue, we assessed

whether the L291R mutation affects PAP I expression. We introduced the L291R missense

mutation into the native *pcnB* locus of both *Y. pseudotuberculosis* IP2666pIB1 (with or without a

PAP I C-terminal FLAG tag) and YPIII/pIBX to generate PAP I[L291R] strains. We found that the

PAP I[L291R] protein was poorly expressed, particularly at 37°C/low calcium (Fig. 4A). These data

suggest that the PAP I[L291R] allele may be unstable, leading to low PAP I expression at least at

37°C/low calcium. Interestingly, a strain engineered with a more conservative leucine to alanine

mutation at position 291, PAP I^{L291A}, did not significantly affect PAP I expression levels (Fig 4A).

Consistent with these data, the PAP I^{L291R} mutant showed enhanced loss of pYV compared to

the wildtype strain or the PAP I^{L291A} mutant, as indicated by the emergence of white, fast-

growing colonies following growth under 37°C/low calcium conditions on Congo red agar (Fig

4B). Interestingly, the PAP I^{L291R} strain displayed a greater loss of pYV following long passage at

37°C/low calcium compared to when bacteria were passaged at 26°C (Table 2), reflective of the

relative stability of the PAP I^{L291R} protein at 26°C. Lastly, the PAP I^{L291R} strain had significantly

lower average pYV PCN compared to the wildtype and PAP I^{L291A} strains at 37°C/low calcium

(Fig 4C), and this correlated with lower relative T3SS activity (Fig 4D). Taken together, these

data suggest that the PAP I^{L291R} mutation disrupts PAP I protein levels, explaining why we

identified this mutation in our suppressor screen.

Mutation of residues predicted to affect polyadenylase activity decreases pYV PCN,

plasmid maintenance, and T3SS activity

PAP I D80 and D82 have been shown to be essential for PAP I polyadenylase catalytic activity

in *E. coli* [61, 62]. These residues are also present in *Y. pseudotuberculosis* PAP I (Fig. 2A). To

determine whether the PAP I polyadenylase activity is responsible for PAP I's roles in pYV PCN

and maintenance, we mutated the two corresponding aspartic acid residues (D114 and D116) to

alanine to produce a PAP I^{D2A} catalytic mutant. These mutations did not affect PAP I steady

state protein levels (Fig 4A). However, we found that the PAP I^{D2A} mutant displayed reduced

pYV PCN at 26°C and 37°C/low calcium compared to wildtype *Y. pseudotuberculosis*, to the

same degree as observed in the Δ*pcnB* strain which lacks PAP I completely (Fig 4C). Spot test

serial dilution analyses found that PAP I^{D2A} grows similarly to wildtype and Δ*pcnB* at 26°C (Fig

4B). Furthermore, at 37°C/low calcium, PAP I^{D2A} and Δ*pcnB* exhibited about 10-fold more

growth compared to wildtype (Fig 4B), consistent with a reduction in growth arrest due to

decreased T3SS activity. Similarly, the strain expressing PAP I^{D2A}, like the $\Delta pcnB$ mutant,

displayed an increased number of plasmidless, rapidly-growing colonies (Fig. 4B) and reduced

pYV stability (Table 2). Lastly, the PAP I^{D2A} allele did not support normal T3SS activity (Fig 4D).

Taken together, these data suggest that *Yersinia* PAP I functions as a polyadenylase that

maintains pYV PCN, plasmid stability, and T3SS gene expression and activity.

Table 2. Important PAP I residues are required for stability of the pYV plasmid

Strain:	Percent Retention[†] of pYV after growth at 26°C only:	Percent Retention[†] of pYV after exposure to 37°C:
Wildtype	100.0 ±0	97.3 ±3.8
$\Delta pcnB$	80.2 ±7.8	47.7 ±14.0
PAPI^{D2A}	94.0 ±1.7	21.0 ±7.0
PAPI^{L291R}	98.6 ±2.5	68.6 ±7.9
PAPI^{L291A}	100.0 ±0	100 ±0

298
[†]Percent retention of the pYV plasmid was averaged from three independent experiments where ±
represents the standard deviation between experiments.

***Yersinia* PAP I is required for plasmid stability and plasmid-mediated antibiotic**

resistance from a broad range of plasmid families

To determine if *Yersinia* PAP I contributes to regulation of PCN and retention of other plasmid

types, we introduced a ColE1-like pGFP-uv plasmid into wildtype and $\Delta pcnB$ *Y.*

pseudotuberculosis. This plasmid lacks active partitioning systems such as ParABS, but can be

maintained within the population by antibiotic resistance selection [63]. We observed a decrease

in fluorescence intensity in $\Delta pcnB$ colonies compared to wildtype when grown on carbenicillin-

containing media, suggesting that lowering of pGFP-uv PCN in the absence of PAP I in *Y.*

pseudotuberculosis leads to less robust antibiotic resistance (Fig. 5A). To determine if the

decreased PCN of the pGFP-uv plasmid in the Δ*pcnB* strain leads to plasmid loss, we

measured kinetics of plasmid retention over 40 hours after removing antibiotic selection. We

found that over 70% of Δ*pcnB* cells lose the pGFP-uv plasmid within 10 hours of removing

carbenicillin and the plasmid is completely lost from the population by ~36 hours. In contrast,

100% of wildtype cells retained pGFP-uv over the same 36 hour time course (Fig. 5B).

To ensure that the growth defect seen for Δ*pcnB* harboring pGFP-uv was due to loss of PAP I

specifically, we next constructed a PAP I over-expression plasmid in the pTrc99 vector

backbone. Importantly, pTrc99 is also a ColE1-like plasmid that lacks partitioning systems [64].

We found that Δ*pcnB* harboring pTrc99::empty displays a 10-fold growth defect when spotted

onto LB-carbenicillin but not LB-only plates (Fig. 5C), consistent with what was previously

observed with the pGFP-uv plasmid. Importantly, this growth defect was nearly completely

abolished by expression of PAP I from the pTrc99 plasmid (Fig. 5C). In addition to ColE1

plasmids, we found that PAP I is also required for robust antibiotic resistance conferred by

plasmids with p15A-like origins of replication (pACYC184, pBAD33) (Fig S5). Interestingly,

replication of plasmids containing ColE-like and p15A-like origins of replication, as well as IncFII

plasmids like pYV, is regulated by plasmid-specific sRNA [65-67], suggesting a general role for

PAP I in regulating plasmids with replicons regulated by a sRNA. To test if loss of *pcnB* would

affect antibiotic resistance conferred by a plasmid whose replication is not known to be

regulated by a sRNA, we introduced the mini-F plasmid pNF06::*ccdAB* into wildtype and Δ*pcnB*

Y. pseudotuberculosis [30, 44, 68, 69]. The Δ*pcnB* strain grew equally well to the wildtype strain

in the presence of antibiotic selection, suggesting that pNF06::*ccdAB* maintenance and copy

number is likely not affected by PAP I (Fig. 5D). These data suggest that PAP I is important for

Yersinia to maintain normal PCN and stability of plasmids whose replication is regulated by a

sRNA.

PAP I is required for *Y. pseudotuberculosis* virulence in mice

It is well established that the pYV plasmid and T3SS are critical for *Y. pseudotuberculosis*

virulence in mouse infection models [6, 70]. To determine if loss of PAP I contributes to *Yersinia*

virulence, we infected mice via the intraperitoneal (IP) route and monitored colonization of the

spleen and liver. Four days post-inoculation, significantly fewer Δ*pcnB*-infected mice exhibited

robust colonization of the spleen and liver compared to wildtype-infected mice (Fig 6A and B).

Interestingly, Δ*pcnB*-infected mice formed two distinct populations: one population where the

spleen and liver were colonized to the same levels as wildtype-infected mice, and a second

population that had completely cleared the infection (Fig. 6A and B). We isolated several Δ*pcnB*

clones from the livers of colonized mice and found that their pYV stability and T3SS activity

phenotypes were similar to the inoculum strain (Fig S6). In addition, we sequenced the

genomes of these isolates and could not identify any significant mutations, suggesting that host

tissue colonization by the Δ*pcnB* mutant in a subset of mice does not stem from suppressor

mutations that stabilize pYV. Taken together, these data suggest that PAP I increases the

likelihood of *Yersinia* colonization of host tissues.

PAP I is required for maintenance of plasmid-encoded T3SS expression and antibiotic

resistance in *Shigella flexneri*

The human pathogen *Shigella flexneri* harbors a virulence plasmid, pINV, with a RepFIIA origin

similar to the one found on *Yersinia* pYV [1]. To test whether PAP I also regulates *Shigella*

pINV, we deleted the *pcnB* gene in the *S. flexneri* M90T Δ*ipaH2.5* (Δ*ipaH2.5::tet^R*) strain. The

resulting Δ*ipaH2.5*Δ*pcnB* mutant was defective in Congo red binding compared to its parental

strain (Fig 7A), suggesting a T3SS defect. Indeed, the Δ*ipaH2.5*Δ*pcnB* strain exhibited a stark

defect in expression and secretion of the T3SS translocon proteins IpaB, IpaC, and IpaD

compared to the parental strain (Fig 7BC). Lastly, we introduced pTRC99 that confers IPTG-

inducible expression of *Yersinia* PAP I into the Δ*ipaH2.5* and Δ*ipaH2.5*Δ*pcnB* strains. Consistent

with our previous data using *Y. pseudotuberculosis* strains (Fig 5C), deletion of *pcnB* led to a

>10-fold loss of antibiotic resistance that could be complemented by PAP I expression (Fig 7D).

Taken together, these data suggest that PAP I is essential for maintaining robust plasmid

replication, resulting in increased expression of plasmid-encoded virulence and antibiotic

resistance genes in pathogenic bacteria.

Discussion

Y. pseudotuberculosis modulates the PCN of its pYV virulence plasmid to control the dosage of

pYV-encoded genes in response to temperature [21, 22]. This ability of *Y. pseudotuberculosis* to

upregulate pYV PCN at mammalian body temperature is critical for its ability to cause disease.

Here, we show that in addition to dynamic regulation of pYV PCN, *Y. pseudotuberculosis*

requires the chromosomal gene *pcnB*, encoding the PAP I polyadenylase, to maintain pYV PCN

homeostasis and prevent pYV loss from the bacterial population. The role of the *pcnB* gene in

promoting virulence plasmid-encoded T3SS expression is conserved in both *Y.*

pseudotuberculosis and *S. flexneri*. Although *pcnB*/PAP I has been shown to regulate plasmid

replication of model plasmids in *E. coli*, this is the first study to link this gene to bacterial

pathogenesis. In addition, the requirement of *Yersinia* and *Shigella pcnB*/PAP I for robust

plasmid-encoded antibiotic resistance suggests that PAP I represents a potential new drug

target.

Putative PAP I orthologs have been bioinformatically identified in many bacterial species hailing

from diverse phyla, and yet studies on *pcnB*/PAP I have historically been confined to a small

number of non-pathogenic bacterial species [36]. *E. coli* PAP I (EcPAP) has been shown to

promote RNA degradation of target transcripts. Degradation of RNA is generally initiated by the

endonuclease RNase E and, in the absence of 3' secondary structures, the RNA fragments

produced by RNase E are then digested by the 3'-5' exonulceases PNPase and RNase II [71].

However, RNase II- and PNPase-mediated RNA degradation is significantly impaired for

transcripts that have terminal stem-loop structures, such as mRNAs containing intrinsic

terminators at their 3' ends and structured sRNAs [71]. PAP I has been shown to add poly(A)

tails to the 3' ends of transcripts containing terminal stem-loops structures to promote their

decay [72-74] and it is thought that PAP I-dependent polyadenylation of these transcripts

provides a "toe-hold" for PNPase and RNase II to bind and digest the RNA [74-77]. Components

of the RNA degradosome such as PNPase are known to be important for T3SS function in *Y.*

pseudotuberculosis [78-80] and PAP I has been shown to copurify with PNPase in *E. coli* [81],

suggesting that PAP I may represent a novel component of the *Yersinia* RNA degradosome.

Importantly, many of the major domains and catalytic residues are conserved between *Yersinia*

and *Shigella* PAP I and EcPAP. EcPAP polyadenylase activity is abolished when the two

catalytic residues Asp80 and Asp82 are mutated to alanine residues in *E. coli* [61, 62]. To test if

the phenotypes of the Δ*pcnB* strain was due to loss of PAP I polyadenylase activity, we mutated

the corresponding catalytic residues (Asp114 and Asp116) in PAP I from *Y. pseudotuberculosis*

to generate a PAPI^{D2A} strain with both catalytic residues replaced by alanine residues. The

PAPI^{D2A} mutant recapitulated the phenotypes observed for the Δ*pcnB* strain, including reduced

pYV PCN, plasmid stability, and T3SS activity. This supports that PAP I acts as a polyadenylase

to regulate plasmid maintenance and T3SS activity in *Y. pseudotuberculosis*, and our data

suggests that this may also be the case in *S. flexneri.*

EcPAP has been shown to polyadenylate both chromosomally- and plasmid-derived transcripts

and was first identified due to its regulation of a sRNA that regulates plasmid replication [29, 30].

EcPAP has been shown to regulate PCN for the IncFII family plasmid R1 by polyadenylating

and targeting CopA regulatory sRNA for degradation, leading to de-repression of R1 plasmid

replication and increasing R1 PCN (46). This is consistent with our finding that PAP I is required

for normal PCN and maintenance of the IncFII family *Yersinia* pYV plasmid, and for normal

expression of the *Shigella* T3SS encoded on the related pINV virulence plasmid [1]. Taken

together, these data point to the possibility that PAP I targets CopA sRNA from the *Yersinia* pYV

and *Shigella* pINV *cop-rep* loci for polyadenylation. Bacteria are thought to maintain large

extrachromosomal elements such as IncF virulence plasmids within a tight PCN range near a

specific "set point" that is optimal for that species lifestyle [25]. This is particularly complicated in

the case of pYV since this plasmid seems to have a different optimal PCN "set point" depending

on temperature. Given that the Δ*pcnB* mutant displays a PCN defect at both 26°C and 37°C,

this suggests that pYV plasmid replication rates are unusually low in the absence of PAP I in

general, possibly due to the buildup of CopA sRNA. Reduced plasmid replication rates can also

lead to plasmid stability defects when the plasmid replication rate drops low enough that cells

can no longer reliably pass a copy of the plasmid to daughter cells during cell division [82]. Both

reduced PCN and reduced plasmid stability can theoretically drive compounding disturbances in

plasmid maintenance systems due to reduced gene dosage of stability systems such as

partitioning and toxin-antitoxin (TA) systems. In keeping with this idea, our RNA-seq analysis

supports the notion that the decrease in pYV-encoded gene dosage in the absence of PAP I

leads to decreased expression of the ParDE TA system in Δ*pcnB* cells that may be responsible

for killing plasmidless cells (Supplementary Dataset 1)[23]. Altogether, this supports the claim

that PAP I is required for maintenance of the pYV PCN "set point" at both 26°C and 37°C,

possibly through modulation of CopA sRNA levels. This, in turn, may decrease the frequency at

which plasmidless daughter cell arise. Alternatively, PAP I may also regulate stability of

additional transcripts that control plasmid partitioning and/or post-segregational killing. Future

studies will focus on identification of polyadenylated transcripts in *Yersinia* and *Shigella*.

While there is an overall reduction in pYV PCN in the absence of *pcnB*/PAP I, Δ*pcnB* cells that

retain the pYV plasmid still increase pYV PCN upon shifting from 26°C to 37°C. In addition, we

have found that the *pcnB* gene is transcribed at 26°C and 37°C (Supplementary Dataset 1) and

the PAP I protein is also produced at both temperatures (Fig 4A). Taken together, this suggests

that PAP I is not required for the temperature-dependent shift in pYV PCN but rather for the

maintenance of the pYV plasmid in general as it undergoes dynamic shifts in PCN during the

Yersinia facultative lifestyle. We hypothesize that PAP I polyadenylates and promotes

degradation of pYV-encoded CopA sRNA to regulate cellular CopA levels at both 26°C and

37°C. This would help maintain CopA levels within a range that promotes maintenance of pYV

PCN near its optimal "set point" and ensuring that pYV-encoded gene dosage remains high

enough to support replication and maintenance of the plasmid under conditions where pYV PCN

is normally kept low. In the absence of PAP I, we hypothesize that pYV plasmid replication rates

decrease due to abnormally high CopA sRNA levels, leading to reduced pYV PCN, plasmid

stability, T3SS activity, and host tissue colonization.

We found that PAP I is required for normal *Y. pseudotuberculosis* virulence in a mouse

intraperitoneal infection model, consistent with its role in promoting high pYV-encoded gene

dosage and pYV stability, and therefore T3SS activity. Interestingly, $\Delta pcnB$-infected mice fell

into two groups: one in which the spleen and liver were colonized with *Yersinia* to wildtype-

infected levels, and another in which no $\Delta pcnB$ cells could be recovered from spleens and livers

four days post-inoculation. Overall, this led to a significant decrease in spleen and liver

colonization by $\Delta pcnB$ compared to wildtype *Y. pseudotuberculosis*. Consistent with our finding

that deletion of *pcnB* leads to enhanced pYV loss from the bacterial population *in vitro*, wildtype

cultures used for mouse inoculation had 100% retention of pYV whereas the $\Delta pcnB$ inoculum

had an average retention of 81% at the time of inoculation. This 19% decrease in pYV+ $\Delta pcnB$

cells in the inoculum are not likely to explain the difference in virulence, as wildtype inoculums of

10^3 to $3x10^3$ yielded similar spleen and liver colonization levels. These data suggest that

pcnB/PAP I may aid *Yersinia* surviving a colonization bottleneck during infection. To our

knowledge, this is the first time that PAP I has been directly linked to virulence of a bacterial

pathogen and highlights the importance of understanding how PAP I and virulence plasmid

gene dosage contribute to bacterial pathogenesis. All $\Delta pcnB$ cells recovered from the spleen

and livers of infected mice retained at least one copy of the pYV plasmid despite the plasmid

stability defect observed for this strain *in vitro*. These data suggest a selective pressure to retain

pYV during growth in mouse tissues, consistent with the importance of the T3SS to *Yersinia*

virulence. This contrasts with what we observe *in vitro* when pYV and T3SS expression is a

metabolic burden and not a benefit. These data suggest that *pcnB*/PAP I is required for

maintenance of the pYV plasmid when pYV is dispensable to survival. This is necessary to

ensure that enough of *the Y. pseudotuberculosis* population is pYV$^+$ to be able to colonize a

new host and survive in host tissues.

In addition to pYV and pINV, we found that *pcnB* is important for robust replication of sRNA-

regulated plasmids encoding antimicrobial resistance (AMR) genes in *Yersinia* and *Shigella*.

When *Y. pseudotuberculosis* or *S. flexneri* Δ*pcnB* cells are grown in the presence of the

plasmid-selective antibiotic, we observed a ten-fold decrease in growth (Fig 5 and 7), consistent

with previous studies in *E. coli* [29]. Similarly, we found a ten-fold decrease in GFP fluorescence

in the *Y. pseudotuberculosis* Δ*pcnB* strain carrying the ColE1 pGFPuv plasmid compared to

wildtype, suggesting a ten-fold decrease in plasmid-encoded gene expression in the absence of

pcnB. In addition, time course experiments found that the majority of the Δ*pcnB* population loses

the pGFPuv plasmid within just 10 hours of removing the selective antibiotic. These data

support that ColE1 PCN and plasmid stability are reduced in the absence of PAP I, rendering

cells more antibiotic susceptible. Interestingly, the rate at which pYV is lost from *Y.*

pseudotuberculosis in vitro under non-T3SS inducing conditions was far slower than for

pGFPuv. We passaged wildtype and Δ*pcnB* strains in LB medium at 26°C and while 100% of

wildtype retained pYV, only ~60% of Δ*pcnB* cells retained pYV by 13 days with a still increasing

rate of loss. As pGFPuv has no active plasmid partitioning or maintenance systems but pYV has

both ParABS and ParDE systems, we hypothesize that the rate of plasmid loss in cells lacking

PAP I activity will depend on the presence and type of plasmid-encoded mechanisms that

promote plasmid stability.

Through our RNA-seq analysis we identified several differentially regulated transcripts in the *Y.*

pseudotuberculosis Δ*pcnB* mutant that are involved in responding to several types of stress,

including the universal stress protein UspB, the *Yersinia* urease operon, and several cold-shock

proteins (Supplemental Dataset 1). Importantly, these differentially regulated genes are

encoded on the *Y. pseudotuberculosis* chromosome and so should not be significantly affected

by the pYV PCN defect in the Δ*pcnB* mutant. In addition, EcPAP from *E. coli* has been

previously reported to play a critical role in regulating mRNA transcripts important for

responding to stress where loss of PAP I leads to stabilization of stress response transcripts

[43]. Deletion of *pcnB* in *E. coli* results in increased tolerance to several stresses, including

osmotic stress, acid stress, heat shock, and cold shock [83]. Interestingly, we found that loss of

PAP I does not affect *Y. pseudotuberculosis* viability following heat shock, growth at 4°C, acid

shock, or osmotic shock (Fig S7). Our RNA-seq analysis also found that some stress response

genes such as CspE are downregulated in ΔpcnB cells, suggesting that PAP I normally

promotes stability of at least some stress-related transcripts in *Y. pseudotuberculosis*. These

data suggest that while PAP I stabilizes sRNA-regulated plasmids in multiple bacterial species,

its effect on stress responses is species-specific.

While *E. coli* EcPAP and PAP I from *Y. pseudotuberculosis* and *S. flexneri* are highly conserved

in general, some regions of the proteins are less conserved and might account for their different

impacts on stress tolerance. Interestingly, while the PAP I N-terminal region following the initial

propeptide sequence is predicted to be disordered, this is also the region that displays the least

amount of conservation between the three proteins. In contrast, this N-terminal region is 100%

conserved between *Y. pseudotuberculosis* and *Y. pestis* (Fig S8). This disordered region is also

significantly expanded in length for PAP I from *Y. pseudotuberculosis* and *Y. pestis* compared to

E. coli and *Shigella* PAP I (Fig 2A). Given that disordered regions of RNA-binding proteins have

been proposed to mediate interactions with target RNAs and/or other proteins [84], it is possible

that *Yersinia* PAP I has evolved to target different RNA transcripts and/or interacts with one or

more unique protein partners compared to EcPAP. *Yersinia* PAP I complements the plasmid-

encoded antibiotic resistance defect of the *Shigella* ΔpcnB mutant, suggesting that PAP I plays

a conserved role in regulating plasmid-derived sRNAs. However, given the different role of PAP

I in stress resistance in *Yersinia* and *E. coli*, PAP I may be adapted to regulate chromosomally-

derived mRNA transcripts in a more species-specific way.

In this study, we show for the first time that *pcnB*/PAP I is required for maintenance of a large

IncFII virulence plasmid (pYV) that undergoes dynamic yet specific changes in PCN to regulate

virulence gene dosage in *Y. pseudotuberculosis*. PAP I was also found to be required for normal

Y. pseudotuberculosis virulence, providing the first direct evidence that PAP I regulates

virulence of a Gram-negative pathogen. PAP I is required for maintenance of plasmids from

multiple families in *Y. pseudotuberculosis* and *S. flexneri* (Fig 5 and 7; Fig S5), including families

that commonly include virulence and antimicrobial resistance plasmids. PAP I homologs can be

found in many species of Gram-negative pathogenic bacteria, many of which harbor

extrachromosomal plasmids important for their pathogenic lifestyles, and genomic studies show

that the *pcnB* gene is highly subject to horizontal gene transfer [36]. Our data support the idea

that bacterial PAP I has a conserved role in promoting the stability of many sRNA-regulated

plasmids and may be critical for the establishment of new plasmids in a bacterial population

following horizontal transfer by promoting plasmid replication. However, given the different

effects of PAP I loss to stress resistance in *E. coli* and *Yersinia*, PAP I-mediated polyadenylation

of transcripts that are chromosomally-encoded likely varies across species. Future studies will

explore the possibility of targeting PAP I activity as a novel antibiotic and/or plasmid-curing

strategy as well as the contribution of PAPI-mediated polyadenylation to regulation of gene

expression in different pathogenic bacteria.

Materials and Methods

Bacterial strains, plasmids, and growth conditions

All strains used in this study are listed in Table S2. *Y. pseudotuberculosis* strains were grown in

LB or 2xYT media (pH=7) at 26°C or 37°C shaking at 250rpm, unless otherwise indicated. *S.*

flexneri were grown in tryptic soy broth media at 30°C or 37°C on a roller.

Construction of *Y. pseudotuberculosis* and *S. flexneri* mutant strains

All primers and plasmids used to generate *Y. pseudotuberculosis* strains for this study are listed

in Tables S3 and S4, respectively. Inserts were amplified by PCR using primers designed for

Gibson Assembly and final recombinant plasmids were all made using the pCVD442 suicide

vector backbone (λpir-dependent replicon, carbenicillin resistant [Carb^R], *sacB* gene conferring

sucrose sensitivity) (Addgene) following digestion with NdeI and BbsI. All ligations were

performed using NEBuilder HiFi DNA Assembly kit (New England Biolabs). Recombinant

pCVD442 plasmids were transformed into *E. coli* S17-1 λpir competent cells and introduced into

Y. pseudotuberculosis IP2666pIB1 or YPIII/pIBX via conjugation. Carb^R and irgansan-resistant

(*Yersinia* selective antibiotic) clones were grown in the absence of antibiotics and plated onto

media containing 10% sucrose to induce SacB-mediate toxicity for pCVD442 loss. Carb

sensitive, sucrose-resistant, and Congo red-positive (pYV+) clones were screened by PCR and

sequence verified by Sangar sequencing. <u>*pil* insertion strains:</u> A *pil::Tn Y. pseudotuberculosis*

IP2666pIB1 Δyop6 strain was originally isolated from a Tn*Himar1* library screened for altered

NFκB induction compared to the parental strain (52). The Tn insertion was then transferred to

the wildtype IP2666pIB1 strain using the pSR47s suicide plasmid, as described above. The

pil::pNQ Y. pseudotuberculosis YPIII/pIBX mutant was generated using the synthesized

pNQ705 suicide plasmid carrying a truncated fragment of *pil* (Genescript)[85]. <u>Δ*pcnB*:</u> ~500 bp

upstream of *pcnB* and ~500 bp downstream of *pcnB* was PCR amplified using genomic DNA

(gDNA) from *Y. pseudotuberculosis* IP2666pIB1. Amplified fragments were cloned into

pCVD442 to produce pCVD442::Δ*pcnB*. <u>ParB-GFP:</u> To make ParB-GFP YPIII/pIBX strains,

three fragments were amplified for insertion into pCVD442: (1) the last ~500 bp of the *parB*

gene excluding the stop codon was amplified from *Y. pseudotuberculosis* gDNA, (2) msfGFP

was amplified from pDK13 and (3) 500 bp downstream of the *parB* gene was amplified from *Y.*

pseudotuberculosis gDNA. The resulting fragments were cloned into pCVD442 to generate a

parB-msfGFP construct that includes 500 bp of downstream homology after msfGFP. <u>PAPI-C-</u>

<u>6xHis and PAPI-C-3xFLAG:</u> To make PAPI-6xHis strains, *pcnB* was amplified from *Y.*

pseudotuberculosis gDNA and cloned into pET28 that was previously digested with XhoI and

NcoI to produce PAPI-C-6xHis. PAPI-C-6xHis was then amplified from pET28 and 500bp

downstream of *pcnB* was amplified from gDNA for insertion into pCVD442 to make

pCVD442::PAPI-c-6xHis containing 500 bp of homology flanking the 6xHis tag. We

subsequently found that the 6xHis tag was not sufficient to visualize the PAPI protein, and so

alternative PAPI-C-3xFLAG strains were made in order to carry out western blot analyses. To

make PAPI-3xFLAG, three fragments were amplified for insertion into pCVD442: (1) the *pcnB*

gene was amplified from *Y. pseudotuberculosis* gDNA (2) 3xFLAG was amplified from pET28

and (3) 300 bp downstream of *pcnB* was amplified from *Y. pseudotuberculosis* gDNA. The

fragments were then cloned into pCVD442 to make pCVD442::PAPI-c-3xFLAG containing 300

bp of downstream homology following the 3xFLAG tag. <u>PAP 1 point mutations:</u> To make PAP

I^{D2A}, PAP I^{L291R}, and PAP I^{L291A} mutants, we used the Q5 site-directed mutagenesis kit (New

England Biolabs) to introduce base pair changes at the corresponding residues using the

pCVD442::PAPI-c-6xHis or pCVD442::PAPI-c-3xFLAG recombinant plasmids as templates.

<u>*ΔipaH2.5ΔpcnB Shigella*:</u> A fragment containing 200 bp of homology upstream and 240bp of

downstream *pcnB* flanking sequence was amplified from ECK0142 (*ΔpcnB E. coli* K-12

BW25113) (PMID 16738554) using pcnb_F and pcnb_R. This fragment was introduced into the

chromosome of *ΔipaH2.5::tetR Shigella* (PMID: 25378474) using lambda red recombination

(PMID: 10829079). The insertion was confirmed using pcnB_R_op and k2_wanner.

In vitro suppressor screen

LB cultures of the *pil::*pNQ YPIII/pIBX mutant strain were grown at 26ºC shaking overnight

before diluting in fresh media and spotting for single colonies onto low calcium plates (LB

supplemented with 20 mM sodium oxalate and 20 mM MgCl$_2$). Plates were incubated at 37ºC

for ~16 hours to induce T3SS expression and selective pressure for growth of suppressor

mutants. Clones that broke growth arrest and thus formed large colonies were selected and

screened for changes in pYV PCN using the luciferase assay (see below). Suppressor mutants

were also screened for changes in T3SS activity using an *in vitro* secretion assay (see below).

Identification of mutations arising from *pil::*pNQ suppressor screen

A subset of *pil::*pNQ YPIII/pIBX suppressor mutants were selected for whole genome

sequencing. In brief, cells were grown shaking at 26ºC overnight and total genomic DNA was

extracted using the Blood & Tissue DNeasy kit (Qiagen) according to manufacturer

recommendations. Qubit quantification, library preparation, and Illumina paired-end sequencing

with 150 bp read length was carried out by the UC Davis DNA Technologies Core. Adapters and

low-quality reads were trimmed by Trimmomatic (Version 0.39)[86]. The trimmed sequences

from each sample were aligned to the reference genome using BWA [87] The reference

sequence was composed of the YPIII chromosome (CP009792.1) [88] and its plasmid

(CP032567.1)[89]. Samtools [90] was used to sort and index the aligned reads. Duplicate reads

originated from a single DNA fragment during sample preparation were located and tagged by

MarkDuplicates in Picard (http://broadinstitute.github.io/picard/). Single nucleotide

polymorphisms were detected and joined by GATK (Version 4.1.6) [91].

Droplet digital PCR (ddPCR)

ddPCR to determine pYV plasmid copy number was carried out as previously described [92].

Total DNA was extracted from bacterial cells using GeneJet Genomic DNA Purification kit

(Thermo Scientific) according to the manufacturer's recommendations for Gram-negative

bacteria. Each genomic DNA sample was split into two reactions and the primer pair

F_chrom_ddPCR/R_chrom_ddPCR was used for to amplify chromosomal DNA and

F_pYV_ddPCR/R_pYV_ddPCR was used to amplify pYV DNA. Individual ddPCR samples were

prepared as follows: 10.5 µL of total genomic DNA was combined with 12.5 µL of 2xEVA-Green

master mix for unlabeled probes (Bio-Rad), 1 µL of HindIII-HF (NEB), and 1 µL of respective

primer pairs (200 nM final concentration) in 8-strip PCR tubes. PCR tubes were incubated at

room temperature for 10 minutes to allow HindIII to digest genomic DNA. Droplet generation

was carried out using an Automated Droplet Generator (Bio-Rad) and droplets were

subsequently transferred to a deep-dish qPCR plate (Bio-Rad) for PCR amplification. PCR was

carried out with hot-start activation at 95°C for 5 mins following by denaturation at 94°C for 30s,

amplification at 58°C for 1 min over 40 cycles, and signal stabilization at 4°C for 10 min and

90°C for 5 min. A ramp rate of 2C/s was used for all PCR steps. Droplets were analyzed using a

QX200 droplet reader (Bio-Rad) and data was analyzed using QuantaSoft Analysis Standard

Edition (version 1.2) (Bio-Rad). Statistical significance was calculated using a one-way ANOVA

with Tukey's multiple-comparisons test (**** $P<0.0001$, *** $P<0.001$, ** $P<0.01$, * $P<0.05$). All

graphs and statistical analyses were generated using GraphPad Prism (10.2.0).

Luciferase pYV plasmid copy number assay

Overnight cultures for strains in the YPIII/pIBX background were grown shaking at 26°C before

subculturing to an OD_{600} of 0.2 in 2xYT low calcium media (containing 20mM sodium oxalate

and 20mM $MgCl_2$). Cultures were then grown shaking for 1.5h at 26°C before shifting to 37°C.

Luminescence and OD_{600} were measured after 1.5h at 26°C just before shifting to 37°C and

every hour after shifting to 37°C. Samples were collected at each timepoint in white walled,

clear bottom 96 well plates and luminescence (using lum aperture for 96 well plate with

measurement time=0.1s) and OD_{600} (using Photometric 600 excitation filter) was measured

using a PerkinElmer Multimode Plate Reader EnVision system. Relative plasmid copy number

was determined by dividing luminescence by OD_{600} for the same sample. Statistical analysis

was determined using a one-way ANOVA with Dunnett's multiple comparisons test (****

$P<0.0001$, *** $P<0.001$, ** $P<0.01$, * $P<0.05$) where applicable. All graphs were made using

GraphPad Prism (10.2.0).

ParB-msfGFP microscopy and image analysis

Wildtype and Δ*pcnB* YPIII/pIBX cells expressing ParB-msfGFP from LB plates supplemented

with kanamycin were inoculated into EZ Rich Defined Medium (EZRDM) (Teknova) without

antibiotic supplementation. EZRDM medium was chosen to reduce medium autofluorescence

for downstream imaging purposes. Cultures were grown shaking at 200 rpm overnight at 26ºC

to a final OD_{600} of ~2-4. A secretion analysis was performed to confirm that *Y.*

pseudotuberculosis displays normal T3SS activity when grown in EZRDM under T3SS-inducing

conditions (37°C, supplemented with 20 mM sodium oxalate and 20 mM $MgCl_2$), referred to as

low calcium EZRDM (Fig S9). For non-T3SS inducing conditions, overnight cultures were

diluted to an OD_{600} of ~0.2 in fresh EZRDM and grown shaking for 3 hours at 26ºC. For T3SS-

inducing conditions, overnight cultures were diluted to an OD600 of ~0.2 in low calcium EZRDM

and grown shaking for 1.5 hours at 26ºC before shifting to 37ºC for an additional 1.5 hours.

Cells were prepared for imaging by spotting onto a 1% agarose-low calcium EZRDM pad and

covering with a no. 1.5 coverslip. Imaging was carried out using a Nikon Eclipse Ti microscope

equipped with a 100X Plan Apo 1.45 NA phase-contrast oil objective, a Hamamatsu Orca-

Flash4.0 V2 CMOS camera, and CoolLED pE-400 (Lumencor). The microscopes were

controlled by the Nikon Elements software. The following Chroma filter cubes were used to

acquire the fluorescence images: GFP: excitation ET470/40x, dichroic T495lpxr, emission

ET525/50 m. Images were acquired with 50 ms of exposure (50% power) for GFP and 100 ms

for phase contrast. For each strain and condition, three independent biological replicates

(inoculated with different colonies) were grown and imaged.

For image analysis, cells were segmented and cell outlines were generated using the open-

source Oufti software [93] using the following parameters: Edgemode, 1; Dilate, 1; openNum, 0;

InvertImage, 0; ThreshFactorM, 0.86254; ThreshMinLevel, 0.3; EdgeSigmaL, 0.5; LogThresh,

0. Cell meshes were further curated using the SVMCuratedCellDetection.m script to identify and

remove misdetected cells [94]. To quantify the number of pYV copies per cell, detection of pYV

spots was performed using the ImageJ plug-in package ThunderSTORM [95, 96]. For of cells

grown at 26ºC, image filtering used a wavelet filter (B-spline) with a scale of 2.0 and order of

3.0. Under non-T3SS-inducing conditions, a local maximum using 5.0*std(Wave.F1) and a

connectivity of 8-neighborhood was used to localize molecules based on peak intensity

thresholds, and sub-pixel localization used a Gaussian point spread function (PSF) with a fitting

radius of four pixels and an initial sigma of one pixel. For the 37ºC/low calcium condition, the

parameters were adjusted where the local maximum instead used 2.0*std (Wave.F1) and a

connectivity of 8-neighborhood, and sub-pixel localization used a Gaussian PSF with a fitting

radius of three pixels and an initial sigma of one pixel. Once parameter optimization was

completed, the custom-made ImageJ macro BatchProcess_ThunderSTORM,ijm was used to

iterate ThunderSTORM spot detection over all image files, applying the same parameters for a

given biological replicate. Detected spots were linked with cell meshes using the custom

MATLAB code analyze_ensemble_spot_intensity.m, detailed further below. Code developed as

part of this study has been deposited in the publicly accessible Github code repository

(https://github.com/JacobsWagnerLab/published/tree/master/Schubert_et_al_2024). Three

independent experiments were performed and analyzed per strain and condition.

Elaborating on the MATLAB script analyze_ensemble_spot_intensity.m, each cell was

quantified by averaging whole cell intensity, recording the number of ThunderSTORM-detected

spots, and noting the cell length. Only cells with length of at least 1µm were kept to rule out

incorrect cell segmentations. Each cell was identified and measured from its original Oufti mesh.

There, the fluorescent GFP images were background-subtracted, then cell contours were used

to obtain the boundaries of the cell area. Cellular fluorescence was determined by Oufti [93]. To

link ThunderSTORM spots, (x,y) localizations, to Oufti meshes, the cell meshes were first

converted to MATLAB polygons using the function *polyshape*. The function *inpolygon* was then

used to feed the (x,y) localizations and polygon shape to find if any spots occurred inside the

cell. For Fig 3 C-E, cells with zero spot were excluded from the analysis.

To assess what fraction of cells were devoid of GFP signal, strains were grouped by condition

(inducing vs. non-inducing as 37ºC/low calcium vs. 26ºC, respectively). Each strain in the

condition was subsetted by random sampling without replacement to match the lowest N value

(7,033 cells in 26ºC and 10,003 cells in 37ºC/low calcium) using the function *datasample*. These

data were then bootstrapped 100 times to recalculate histograms and determine the error for

each bin (using the custom function *Hist_Errors*). Since the $\Delta pcnB$ conditions had clear bimodal

distributions for low/high fluorescent states, an Otsu threshold was calculated using the function

otsuthresh to find the split between them [97]. Cell with intensity values below this threshold

were deemed cells without plasmid content. To compare these two distribution shapes

statistically, a Kolmogorov-Smirnov (KS) test was used via the function *kstest2*.

Images were visualized using NIS Elements and Fiji. Image pixel intensity values were auto-

scaled for display using Fiji, and all images were converted to RGB format for display. Data

were plotted using GraphPad Prism 10.2.0, and Adobe Illustrator was used to generate figures.

***In vitro* pYV plasmid stability assay**

Cells were struck for single colonies onto LB plates and grown at 26°C for 1.5-2 days. To test

plasmid stability after 26°C-only passage, single colonies were then patched onto LB plates for

~16 hours at 26°C before heavily streaking for single colonies onto fresh LB plates at 26°C for

1.5-2 days. To test plasmid stability after exposure to T3SS-inducing conditions (37˚C), the

same colonies were also patched onto LB plates supplemented with 20 mM sodium oxalate to

induce T3SS activity for ~16 hours at 37ºC before streaking for single colonies onto LB plates

and returning to 26ºC to grow for 1.5-2 days. Final plasmid stability was determined for both

conditions by patching single colonies from the final 26°C streak on LB plates supplemented

with 20 mM sodium oxalate and containing 1% Congo red. The percentage of cells that retained

the pYV plasmid was calculated by dividing the number of red (pYV+) CFU by the total number

of CFU. Single colony PCR was used on a subset of patches to validate that red CFU retain

pYV and while white CFU do not.

***In vitro* T3SS activity assay**

Quantification of secreted *Yersinia* T3SS cargo was carried out as previously described [98]. *Y.*

pseudotuberculosis cultures were grown shaking overnight at 26°C before subculturing in LB

low calcium media (LB containing 20 mM sodium oxalate and 20 mM $MgCl_2$). Cultures were

grown shaking at 26°C for 1.5 hours before shifting to 37°C/shaking for an additional 1.5 hours.

Cultures were normalized to the same number of cells by normalizing OD_{600} values and pelleted

at 13,200 rpm for 15 min at room temperature. Supernatants were transferred to new tubes and

proteins precipitated with 10% trichloroacetic acid (TCA). A bovine serum albumin (BSA) protein

control was spiked into each sample immediately before the addition of TCA. Samples were

incubated in TCA on ice for at least 1h before pelleting at 13,200 rpm for 15 min at 4°C. Pellets

were then washed twice with ice-cold 100% acetone and resuspended in final sample buffer

(FSB) containing 0.2M dithiothreitol (DTT). Samples were boiled for 10 min at 95°C before

loading and separating on a 12.5% SDS-PAGE gel. Gels were stained with Coomassie blue to

visualize proteins and imaged using Bio-Rad Image Lab Software Quantity and Analysis tools.

Shigella **T3S secretion assay**

S. flexneri cultures were grown with shaking overnight in tryptic soy broth (TCS) media with 15

µg/mL tetracycline and 50 µg/mL kanamycin at 37°C. In the AM, the cultures were sub-cultured

in TCS and grown for 2.5 hours with shaking at 37°C to reach OD_{600} ~1. Cultures were

normalized to the same number of cells based on the OD_{600} and pelleted at 13,000 rpm for 1

min before resuspending in PBS and 10 µM Congo red and incubated for 30 min shaking at

37°C. Pellets and supernatants were separated by centrifugation and supernatants were

transferred to new tubes and precipitated with 10% TCA. Supernatant samples were incubated

on ice and subsequently pelleted at 15000 rpm for 15 min at 4°C. Pellets were then washed with

ice-cold 100% ethanol and resuspended in 1x loading dye buffer with 5% beta-mercaptoethanol.

Samples were boiled for 10 min at 95°C before loading on a 12% SDS-PAGE gel for western

blot analysis.

RNA isolation and RNA-seq analysis

Y. pseudotuberculosis IP2666pIB1 strains were grown in 3 mL LB overnight at 26°C and diluted

the following morning to an OD_{600} of 0.2 in 5 mL fresh LB. Each strain was diluted into two

separate cultures (one for each temperature). For 26°C samples, cultures were diluted in plain

LB media and grown shaking at 26°C for 3h before extracting RNA. For 37°C samples, cultures

were diluted into LB low calcium media (containing 20mM sodium oxalate and 20mM $MgCl_2$)

and grown shaking at 26°C for 1.5 hours before shifting to 37°C to growing shaking for an

additional 1.5 hours. After incubating, 1-2mL of each culture was collected and pelleted for 5

min at 4,000 rpm. Supernatants were removed and cell pellets were resuspended in 1mL RNA

Protect (Qiagen) and total RNA was isolated using the RNeasy minikit (Qiagen) according to the

manufacturer's protocol. DNA was removed from samples using the TURBO DNA-free kit (Life

Technologies/Thermo Fischer) and RNA quality was determined using an Agilent 2000

Bioanalyzer. Samples were DNase treated with Invitrogen DNase. Library preparation was

performed using Illumina's Stranded Total RNA Prep Ligation with Ribo-Zero Plus kit and 10 bp

IDT for Illumina indices. Sequencing was done on a NovaSeq 6000 giving 2x51bp reads.

Demultiplexing, quality control, and adapter trimming was performed with bcl-convert (v4.0.3).

Read mapping was performed with HISAT2 [99]. Read quantification was performed using

Subread's featureCounts functionality [100]. Read counts were normalized using edgeR's

Trimmed Mean of M values (TMM) algorithm (Supplementary Dataset 1)[101]. Subsequent

values were then converted to counts per million (cpm). Differential expression analysis was

performed using edgeR's exact test for differences between two groups of negative-binomial

counts with an estimated dispersion value of 0.1. Genes for which the |logFC| > 1 and p < 0.05

were considered differentially expressed (Supplementary Dataset 1).

Heatmaps for pYV-encoded genes were generated using average TMM values via ClustVis

[102]. RNA-Seq data are available for viewing, and the track hub data can be found

at http://lowelab.ucsc.edu/hubs/yersinia_pcnB_pil_expression/hub.txt on the UCSC Microbial

Genome Browser. For these data, chromosomal reads were normalized separately from pYV-

encoded reads due to differences in pYV PCN among samples. For this analysis, raw paired

reads were aligned to the *Y. pseudotuberculosis* IP2666pIB1 genome (ASM381434v1) using

Bowtie2 (v.2.3.5.1) with default options. Bam files were further processed to remove PCR

duplicates using samtools (v1.15.1) as follows, samtools sort -n {input_bam} | samtools fixmate

-m - - | samtools sort - | samtools markdup -r - - | samtools view -bq 10 -f 0x1 - > {output_bam}.

Finally, bamCoverage-deeptools (v3.5.4) was used to normalize the chromosomal read counts

using the following options: --normalizeUsing CPM --ignoreForNormalization NZ_CP032567.1 -

-outFileFormat bedgraph --binSize 1 --region NZ_CP032566.1. For the non-normalized pYV

reads we used bamCoverage-deeptools with options --ignoreForNormalization

NZ_CP032566.1 --outFileFormat bedgraph --binSize 1 --region NZ_CP032567.1. Finally, the

separate bedgraph files were concatenated and converted to bigwig format using

bedGraphToBigWig (v4) with default options.

Western blot analysis

Cells were grown at the indicated temperature before preparing samples for western blotting.

Samples were normalized to the same number of cells using OD_{600} at the time of sample

collection. Bacterial pellets were then resuspended in FSB containing 0.2M DTT and boiled at

95°C for 15 minutes. Secreted proteins were precipitated using TCA in the same way as

described for the *in vitro* secretion assay above. The same volume of each sample was

separated on a 12.5% SDS-PAGE gel and proteins were transferred to a blotting membrane

(Immunobilon-P) using a wet mini trans-blot cell (Bio-Rad) at 4°C. Blots were blocked for one

hour in Tris-buffered saline containing Tween 20 and 5% milk. Blots were probed with anti-RpoA

(gift from Melanie Marketon), goat anti-YopE (Santa Cruz Biotech), mouse M2 anti-FLAG

(Sigma), and horseradish peroxidase-conjugated secondary antibodies (Santa Cruz Biotech).

Bands were visualized using a BioRad ChemiDoc and quantification analysis was performed

using Image Lab software (Bio-Rad).

For *S. flexneri* strains, samples were separated by a 12% SDS-PAGE gel and transferred to a

nitrocellulose membrane. Blots were probed with anti-IpaB, anti-IpaC, and anti-IpaD antibodies,

and anti-rabbit secondary antibodies. Bands were visualized using SuperSignal West Pico

PLUS Chemiluminescent Substrate on an Azure 300 Imager.

Mouse Infections

C57BL/6 mice were purchased from Jackson Laboratory. Six- to eight-week-old C57Bl/6 mice

were infected with $1\text{-}3\times10^3$ *Y. pseudotuberculosis* via intraperitoneal (IP) injection as previously

described [103]. In brief, LB overnight cultures of *Y. pseudotuberculosis* were grown shaking at

26ºC before spinning down at 5000 rpm and resuspending in sterile PBS (pH 7.4). CFU used for

the inoculum cultures were confirmed to contain pYV prior to infection using single colony PCR

with primers specific to the pYV plasmid. Samples were then diluted in PBS such that 100 μL

contained $1\text{-}3\times10^3$ cells. Inoculums were plated to confirm the inoculation dose and to estimate

the percentage of cells that were pYV+ by patching onto LB plates supplemented with 20 mM

sodium oxalate and containing 1% Congo red dye. The plasmid retention of the starting

inoculums was found to be 100% for wild type and 81% on average for $\Delta pcnB$. Mice were

injected with 100 μL of inoculum and spleens and livers harvested four days post-inoculation.

Organs were homogenized and serial dilutions of the homogenate plated to determine CFU per

gram (CFU/g) of tissue.

Construction of PAP I expression vector

The *pcnB* ORF was PCR amplified from *Y. pseudotuberculosis* genomic DNA using the primer

pair FpcnB_pTrc/RpcnB_pTrc and ligated into pTrc99a previously digested with NcoI and XbaI

using NEBuilder HiFi DNA Assembly kit (New England Biolabs, Inc.). Constructs were

transformed into DH5α *E. coli* for propagation and sequence verified via Sangar sequencing.

The resulting pTrc99a::PAPI recombinant expression vector was introduced to wildtype and

$\Delta pcnB$ *Y. pseudotuberculosis* by electroporation and selection on LB plates containing

carbenicillin. Expression of PAPI was induced by plating onto LB plates containing 0.2 mM

IPTG.

Spot tests to analyze antibiotic resistance conferred by pGFPuv, pTrc99, and pNF06

Wildtype and Δ*pcnB Y. pseudotuberculosis* IP2666pIB1 were electroporated with 1 μL of

pGFPuv [Carb^R] (Gifted by Dr. Manel Camps), pTrc99a, or pNF06::ccdAB [Kan^R] (Gifted by Dr.

Laurence Van Melderen) and single colonies were selected by plating on LB plates containing

carbenicillin (pGFPuv, pTrc99a) or kanamycin (pNF06). For spot test dilution series analyses,

overnight cultures were grown at 26°C in the presence of antibiotics before diluting in PBS (pH

7.4) and making a series of 1:10 dilutions. Serial dilutions were spotted onto LB plates

containing no antibiotic or LB plates containing the selective antibiotic and plates were

visualized after 1-1.5 days of growth at 26°C.

Δ*ipa2.5::tet^R* and Δ*ipa2.5*Δ*pcnB S. flexneri* M90Twere electroporated with pTrc99a, and single

colonies were selected by plating on LB plates containing ampicillin. For spot tests, overnight

cultures were grown at 30°C in the presence of antibiotics before diluting in PBS (pH 7.4) and

making a series of 1:10 dilutions. Serial dilutions were spotted onto LB plates containing no

antibiotic or LB plates containing the selective antibiotic and plates were visualized after 1 day

of growth at 30°C.

pGFPuv plasmid loss curve

Wildtype and Δ*pcnB Y. pseudotuberculosis* IP2666pIB1 lacking pYV (pYV-) were electroporated

with pGFPuv as described above. Overnight cultures were grown at 26°C in the presence of

carbenicillin. Overnight cultures were diluted 1:1000 into LB media without antibiotics to allow

loss of the plasmid. To assess plasmid retention at each timepoint, aliquots of the cultures were

plated onto LB agar and incubated at 26°C. Fifty single colonies from each sample were

patched onto LB agar containing carbenicillin. Percentage of plasmid retention was calculated

as the number of patched colonies able to grow in the presence of carbenicillin relative to the

total number of patched colonies.

Acknowledgments

We would like to acknowledge Dr. Manel Camps (UC Santa Cruz) for providing the pGFPuv

plasmid as well as general advice and guidance on this work. We would also like to

acknowledge Dr. Laurence Van Melderen (Université Libre de Bruxelles) for providing us with

the pNF06::empty and pNF06::*ccdAB* constructs. Finally, we would like to thank Dr. Petra

Dersch (Universität Münster) for providing general advice and technical support for this project.

Funding: This study was supported by National Institutes of Health (www.NIH.gov) grant

R01AI141511 (to VA), R01AI128360 (to CFL), and the Swedish Research Council 2018-02376

and 2022-00741 (to HW). CJW is an investigator of the Howard Hughes Medical Institute. HNL

was supported by NIH K99AI139281. KS was supported by an NIH T32 GM008646 training

grant. The funders had no role in study design, data collection and analysis, decision to publish,

or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Figure Legends

Figure 1. *Y. pseudotuberculosis* pYV plasmid copy number (PCN) dynamically changes

in response to temperature. (A) pYV plasmid copy number per chromosome was determined

for wildtype *Y. pseudotuberculosis* IP2666pIB1 grown at 26°C or 37°C/low calcium using droplet

digital PCR (dd)PCR. **(B)** Relative pYV plasmid copy number of wildtype *Y. pseudotuberculosis*

YPIII/pIBX was determined following growth at 26°C or 37°C/low calcium using a luciferase

reporter stably incorporated in the pIBX plasmid, normalized to bacterial cell density (OD_{600}). (A-

B) Averages of three independent experiments are shown ± standard deviation (Student t-test,

*** $p<0.001$, ** $p<0.01$, * $p<0.05$). **(C)** Representative phase contrast (left) and fluorescence

(right) images of *Y. pseudotuberculosis* YPIII/pIBX cells where the pYV plasmid is visualized

using a ParB-msfGFP fluorescent marker. Images are shown for cells grown at 26°C or under

the 37°C/low calcium condition. **(D)** Histogram depicting the number of ParB-msfGFP-labeled

puncta of pYV quantified from data illustrated in panel C. Three biological replicates per

condition were imaged and analyzed; for each strain and condition, between 790 to 4428 cells

were analyzed per replicate.

Figure 2. Disruption of the *pcnB* gene encoding poly(A) polymerase PAP I decreases pYV

PCN as well as T3SS expression and activity. (A) PAP I from *Y. pseudotuberculosis*

IP2666pIB1, *E. coli* K12, and *S. flexneri* M90T are highly conserved. Known PAP I domains

include the head domain (blue), the neck domain (green), the body domain (purple), and the leg

domain (yellow) [62]. Residues in boxes represent known catalytic residues from EcPAP and

the residue indicated with an arrow represents the residue that was mutated in the L291R *pcnB*

suppressor mutants. **(B)** pYV plasmid copy number per chromosome was determined for *Y.*

pseudotuberculosis IP2666pIB1 strains grown at 26°C or 37°C/low calcium using ddPCR.

Averages of three independent experiments are shown ± standard error of the mean. Statistical

significance was calculated using a one-way ANOVA with Tukey's multiple comparisons test

(**** p<0.0001, *** p<0.001, ** p<0.01, * p<0.05). **(C)** *Y. pseudotuberculosis* IP2666pIB1 strains

were grown in low calcium media at 37°C before preparing samples for western blotting.

Bacterial pellets were collected and secreted proteins (Yops) were precipitated from the

supernatant (sup). Images shown represent a single blot that was independently probed for

YopE and RpoA, which serves as a loading control.

Figure 3. The *pcnB* gene is required for maintenance of the pYV plasmid. (A)

Representative phase contrast (left) and fluorescence (right) images of *Y. pseudotuberculosis*

YPIII/pIBX Δ*pcnB* cells where the pYV plasmid is visualized using a ParB-msfGFP fluorescent

marker. Images are shown for cells grown at 26°C or under the 37°C/low calcium condition.

White arrowheads indicate plasmidless cells. **(B)** Histograms comparing wildtype vs. Δ*pcnB*

cells under 26°C or 37°C/low calcium conditions. Mean fluorescence intensity values are plotted

on the x-axis (log_{10}). Shaded error bars represent the mean ± standard error of the mean from

bootstrapping the data 100 times. Both histograms were randomly sampled without replacement

to match the lowest n value of either condition (7,003 and 10,033 for the 26°C and 37°C/low

calcium conditions, respectively). Both distributions are significantly different ($p < 10^{-100}$) by

Kolmogorov-Smirnov tests. A.U. indicates arbitrary units. **(C)** Histograms depicting the number

of ParB-msfGFP-labeled puncta of pYV quantified from data illustrated in panel A. Only cells

with at least one detected pYV spot were considered. Three biological replicates were imaged

and analyzed per condition. For each strain and condition, between 2063 to 14,479 cells were

analyzed per replicate. **(D)** Same as (C) except that the comparison is between wildtype and

Δ*pcnB* cells grown at 26°C. For each strain and condition, between 760 to 4179 cells were

analyzed per replicate. **(E)** Same as (D) except for the 37°C/low calcium condition. For each

strain and condition, between 2046 to 14,479 cells were analyzed per replicate. Distributions in

panels D and E are significantly different ($p < 10^{-80}$) by Kolmogorov-Smirnov tests.

Figure 4. PAP I point mutations lead to a decrease in pYV PCN, plasmid stability, and T3SS activity. (A) *Y. pseudotuberculosis* IP2666pIB1 strains harboring various PAP I[FLAG] alleles grown at 26°C or 37°C/low calcium were pelleted and prepared for western blot analyses. Images shown represent a single blot that was cut after transferring and independently probed for expression of PAP I[FLAG] (top) and RpoA (loading control, bottom). Data shown is representative of three biological replicates. **(B)** *Y. pseudotuberculosis* IP2666pIB1 strains were grown at 26°C before diluting and spotting onto either LB or low calcium plates containing Congo red dye and incubated at 26°C or 37°C, respectively, prior to imaging. **(C)** Relative pYV plasmid copy number was estimated for strains in the *Y. pseudotuberculosis* pIBX/YPIII background. For each timepoint, luminescence was measured and normalized to cell density (OD_{600}). Averages of three independent experiments are shown ± standard error of the mean. Each strain per condition was compared to the wildtype (WT) strain in a one-way ANOVA with Dunnett's multiple comparisons test (**** $P<0.0001$, *** $P<0.001$, ** $P<0.01$, * $P<0.05$). **(D)** *Y. pseudotuberculosis* IP2666pIB1 strains were grown at 37°C/low calcium and secreted proteins in the supernatant fractions were precipitated and visualized after separation on as SDS-PAGE gel by Coomassie blue staining. Bovine serum albumin (BSA) was used as a loading and protein precipitation control. The YopE T3SS effector protein band from four independent experiments was visualized, quantified, and normalized to BSA from the same samples. Bars represent the mean fold change in YopE/BSA relative to wildtype and error bars represent the standard error of the mean between different experiments. Statistical significance was determined using a one-way ANOVA with Dunnett's multiple comparisons test (**** $P<0.0001$, *** $P<0.001$, ** $P<0.01$, * $P<0.05$).

Figure 5. The *pcnB* gene is required for plasmid-mediated antibiotic resistance in

***Y. pseudotuberculosis.* (A)** Wildtype and Δ*pcnB Y. pseudotuberculosis* IP2666pIB1 harboring

pGFP-uv were spotted simultaneously onto either plain LB plates or LB plates containing

carbenicillin to assess retention of pGFP-uv. Plates were incubated at 26°C for ~16 hours

before imaging spots under a GFP filter. **(B)** Retention of the pGFP-uv plasmid by wildtype and

Δ*pcnB Y. pseudotuberculosis* IP2666pIB1 was assessed at 26°C over 40 hours following the

removal of carbenicillin. **(C)** Wildtype and Δ*pcnB Y. pseudotuberculosis* IP2666pIB1 harboring

either no vector, a pTrc99::empty vector control, or a pTrc99::PAP I recombinant plasmid were

spotted onto either plain LB plates or LB plates containing carbenicillin to select for pTrc99 and

0.2 mM IPTG to induce PAP I expression. Plates were incubated at 26°C for ~16 hours prior to

imaging. **(D)** Wildtype and Δ*pcnB Y. pseudotuberculosis* IP2666pIB1 harboring either no vector

or a pNF06 mini-F plasmid with a *ccdAB* TA system (pNF06::*ccdAB*) were spotted onto either

plain LB plates or LB plates containing 25 µg/mL kanamycin to select for pNF06. Plates were

incubated at 26°C for ~16 hours prior to imaging.

Figure 6. The *pcnB* gene is required for normal *Y. pseudotuberculosis* virulence in a

mouse IP infection model. Six- to eight-week-old C57Bl/6 mice were infected with 1-3x10^3 *Y.*

pseudotuberculosis IP2666pIB1 via intraperitoneal (IP) injection. Tissues were collected,

homogenized, and plated to determine CFU per gram (CFU/g) tissue 4 days post-inoculation.

The $\log_{10}$ CFU/g tissue from the spleens **(A)** and livers **(B)** of infected mice are shown. Each

circle represents one mouse organ; lines represent geometric means.

Figure 7. The *pcnB* gene is required for plasmid-encoded expression of T3SS and

antibiotic resistance genes in *Shigella flexneri*. (A) *S. flexneri* M90T Δ*ipaH2.5* and

ΔipaH2.5ΔpcnB, as well as a virulence plasmid cured (VP⁻) BS103 strain, were incubated on

Congo red media at 37˚C. **(B)** *S. flexneri* M90T *ΔipaH2.5* and *ΔipaH2.5ΔpcnB* carrying pTRC99

with and without IPTG-inducible *Yersinia* PAP I were incubated at 37˚C, and secreted T3SS

effector proteins were visualized after separation on as SDS-PAGE gel using anti-IpaB, anti-

IpaC, and anti-IpaD antibodies. **(C)** *S. flexneri* M90T *ΔipaH2.5* and *ΔipaH2.5ΔpcnB* carrying

pTRC99 with and without IPTG-inducible *Yersinia* PAP I were grown on either plain TCS media

or TCS containing ampicillin and IPTG. Shown is one of two independent experiments.

1002

1003

1004

1005 **Supporting Information**

1006

Figure S1. A transposon insertion into *pil* leads to a constitutively high pYV PCN,

elevated T3SS expression, and a severe growth defect under T3SS-inducing conditions.

(A) HEK293T cells expressing an NFκB luciferase reporter were left uninfected (uninf) or were

infected with an effectorless *Δyop6* strain (*ΔyopHEMOJ*), a *ΔyscNU* T3SS-deficient strain, or

pil::Tn in the *Y. pseudotuberculosis* IP2666pIB1 background. Averages of three independent

replicates ± standard error the mean are shown (Student t-test, $p < 0.04$). **(B)** Schematic

representation of the location of *pil*. The *pil* locus is located ~123bp away from the *copB* gene.

(C) Wildtype, *pil::Tn*, and *Δpil* IP2666pIB1 were grown at 37°C/low calcium and secreted T3SS

cargo proteins (Yops) were precipitated and visualized by Coomassie blue staining. Bovine

serum albumin (BSA) serves as a loading and precipitation control. Data shown is

representative of three independent replicates. **(D-E)** Relative pYV PCN was estimated for

strains in the YPIII/pIBX background of *Y. pseudotuberculosis* using a luciferase plasmid copy

number assay. For each timepoint, luminescence was measured and normalized to cell density

(OD$_{600}$). Representative data from three independent experiments are shown. **(E)** Serial

dilutions for wildtype YPIII/pIBX and the *pil*::pNQ congenic *Y. pseudotuberculosis* strains were

spotted onto low calcium agar plates for ~16 hours at 37°C before imaging. The yellow arrow

indicates an example of a large colony in the *pil*::pNQ mutant background that represents a

candidate suppressor mutant.

1025

1026   **Figure S2. T3SS gene mRNA levels are higher in the *pil*::Tn mutant and lower in the**

1027   **Δ*pcnB* mutant compared to wildtype.** Heat maps displaying relative expression of pYV-

1028   encoded genes from RNA-seq analysis of *Y. pseudotuberculosis* IP2666pIB1 strains grown at

1029   26°C or 37°C/low calcium.

1030

1031   **Figure S3. The *pil*::Tn strain exhibits normal temperature and calcium regulation of T3SS**

1032   **activity.** Strains were grown in regular LB or low calcium LB media, at either 26°C or at 37°C.

1033   Secreted proteins in the supernatant were precipitated and visualized by SDS-PAGE and

1034   Coomassie blue staining. Bovine serum albumin (BSA) was added as a loading and

1035   precipitation control. Data shown is representative data of three biological replicates.

1036

1037   **Figure S4. Transposon insertion in the *pil* locus alters RNA-Seq read patterns at the *cop-***

1038   ***rep* locus.** Reads from RNA-seq analysis mapped to the *cop-rep* locus and adjacent *pil* locus

1039   from wildtype and *pil*::Tn (*pil*) *Y. pseudotuberculosis* IP2666pIB1 grown at 26°C or 37°C/low

1040   calcium. The putative *pil* locus is indicated by a black box and the approximate location of the

1041   Tn insertion is indicated with an arrow.

1042

Figure S5. The *pcnB* gene is required for plasmid-mediated antibiotic resistance in

Y. pseudotuberculosis. Wildtype and Δ*pcnB Y. pseudotuberculosis* IP2666pIB1 harboring

either pBAD33 or pACYC184 were grown overnight in LB supplemented with 20 µg/mL

chloramphenicol, and the overnight cultures spotted onto LB agar lacking or containing

chloramphenicol to assess plasmid retention. Plates were incubated at 26°C for ~16 hours

before imaging. Each experiment was carried out for a total of two independent experiments.

Figure S6. *Y. pseudotuberculosis* isolated from infected livers demonstrate similar

plasmid loss and T3SS activity phenotypes as their parental strains. Independent colony

forming units from infected mouse livers were isolated and assessed for their ability to retain

pYV and carry out T3SS activity. WT-3 and WT-5 are wildtype IP2666pIB1-derived isolates from

two different mice. Δ*pcnB*-2-9, Δ*pcnB*-2-10, Δ*pcnB*-3-1, and Δ*pcnB*-3-2 are Δ*pcnB*-derived

isolates from two different mice. **(A)** Liver isolates and their parental strains were incubated

overnight at 26°C in LB or 37°C in LB/low calcium media. Colonies were patched onto low

calcium media containing Congo red to induce T3SS activity and assess the percentage of

colonies that retained pYV (i.e.-red colonies). **(B)** Liver isolates and their parental strains were

incubated in LB low calcium media at 37°C, and secreted proteins visualized by Coomassie blue

staining. Bovine serum albumin (BSA) was added as a loading and precipitation control. Data

shown is representative of two independent experiments.

Figure S7. Loss of PAP I does not affect *Y. pseudotuberculosis* growth following stress

exposure. (A) Wildtype and Δ*pcnB Y. pseudotuberculosis* IP2666pIB1 cultures were grown to

mid-log phase at 26°C before shifting to 42°C for 1 hour to induce heat shock. Spot dilutions

were plated onto LB plates and incubated at 26°C for ~16 hours before imaging. **(B)** Strains in

the IP2666pIB1 background were grown at 26°C overnight (stationary phase) before diluting

and spotting into LB plates. Plates were incubated at 26°C for 4 hours before transferring to 4°C

and allowing growth for ~2 days prior to imaging. **(C)** Wildtype and Δ*pcnB Y.*

pseudotuberculosis IP2666pIB1 were grown at 37°C (low calcium) to mid-log phase before

inducing stress. Growth was assayed by plating spot dilutions following stress exposure. Stress

was induced by adjusting media to pH=3.0 to induce acid shock or by addition of 800Mm $NaCl_2$

to induce osmotic shock.

1074

1075   **Figure S8. PAP I from *Y. pseudotuberculosis* and *Y. pestis* are 100% conserved.** Predicted

1076   PAP I sequences from *Y. pseudotuberculosis* IP2666pIB1 and *Y. pestis* KIM5.

1077

Figure S9. *Y. pseudotuberculosis* displays normal T3SS activity in EZ-RDM low calcium

media. *Y. pseudotuberculosis* IP2666pIB1 wildtype, Δ*pcnB*, and pYV- strains were grown in

either low calcium LB or low calcium EZ-RDM media. The pYV- strain cured of pYV was used

as a negative control [103]. Secreted proteins were visualized by Coomassie blue staining.

Bovine serum albumin (BSA) was added as a loading and precipitation control. Data shown is

representative of two independent experiments.

1084

1085   **Table S1. Phenotypic categories and mutations identified in suppressor screen.**

1086

1087   **Table S2. *Y. pseudotuberculosis* strains used in this study.**

1088

1089   **Table S3. Primers used in this study.**

1090

1091   **Table S4. Plasmids used in this study.**

1092

Dataset S1. Transcriptome analysis of *Y. pseudotuberculosis* IP2666pIB1, *pil*::Tn, Δ*pcnB*,

and *pil*::Tn/Δ*pcnB* congenic strains.

References

1.   Pilla G, Tang CM. Going around in circles: virulence plasmids in enteric pathogens. Nat

Rev Microbiol. 2018;16(8):484-95. doi: 10.1038/s41579-018-0031-2. PubMed PMID: 29855597.

2.   Gemski P, Lazere JR, Casey T, Wohlhieter JA. Presence of a virulence-associated

plasmid in *Yersinia pseudotuberculosis*. Infect Immun. 1980;28(3):1044-7. doi:

10.1128/iai.28.3.1044-1047.1980. PubMed PMID: 6249747; PubMed Central PMCID:

PMCPMC551057.

3.   Cornelis GR. The *Yersinia* Yop virulon, a bacterial system to subvert cells of the primary

host defense. Folia Microbiol (Praha). 1998;43(3):253-61. doi: 10.1007/BF02818610. PubMed

PMID: 9717252.

4.   Sun YC, Jarrett CO, Bosio CF, Hinnebusch BJ. Retracing the evolutionary path that led

to flea-borne transmission of *Yersinia pestis*. Cell Host Microbe. 2014;15(5):578-86. doi:

10.1016/j.chom.2014.04.003. PubMed PMID: 24832452; PubMed Central PMCID:

PMCPMC4084870.

5.   Glatter KA, Finkelman P. History of the Plague: An Ancient Pandemic for the Age of

COVID-19. Am J Med. 2021;134(2):176-81. Epub 20200924. doi:

10.1016/j.amjmed.2020.08.019. PubMed PMID: 32979306; PubMed Central PMCID:

PMCPMC7513766.

6.   Cornelis GR, Boland A, Boyd AP, Geuijen C, Iriarte M, Neyt C, et al. The virulence

plasmid of *Yersinia*, an antihost genome. Microbiol Mol Biol Rev. 1998;62(4):1315-52. doi:

10.1128/MMBR.62.4.1315-1352.1998. PubMed PMID: 9841674; PubMed Central PMCID:

PMCPMC98948.

7.   Portnoy DA, Wolf-Watz H, Bolin I, Beeder AB, Falkow S. Characterization of common

virulence plasmids in *Yersinia* species and their role in the expression of outer membrane

proteins. Infect Immun. 1984;43(1):108-14. doi: 10.1128/iai.43.1.108-114.1984. PubMed PMID:

6317562; PubMed Central PMCID: PMCPMC263395.

8.   Demeure C, Dussurget O, Fiol GM, Le Guern AS, Savin C, Pizarro-Cerda J. *Yersinia*

pestis and plague: an updated view on evolution, virulence determinants, immune subversion,

vaccination and diagnostics. Microbes Infect. 2019;21(5-6):202-12. Epub 20190625. doi:

10.1016/j.micinf.2019.06.007. PubMed PMID: 31252217.

9.   Cornelis GR. The type III secretion injectisome. Nat Rev Microbiol. 2006;4(11):811-25.

doi: 10.1038/nrmicro1526. PubMed PMID: 17041629.

10.   Coburn B, Sekirov I, Finlay BB. Type III secretion systems and disease. Clin Microbiol

Rev. 2007;20(4):535-49. doi: 10.1128/CMR.00013-07. PubMed PMID: 17934073; PubMed

Central PMCID: PMCPMC2176049.

11.   Bliska JB, Wang X, Viboud GI, Brodsky IE. Modulation of innate immune responses by

Yersinia type III secretion system translocators and effectors. Cell Microbiol. 2013;15(10):1622-

31. Epub 20130729. doi: 10.1111/cmi.12164. PubMed PMID: 23834311; PubMed Central

PMCID: PMCPMC3788085.

12.   Deng W, Marshall NC, Rowland JL, McCoy JM, Worrall LJ, Santos AS, et al. Assembly,

structure, function and regulation of type III secretion systems. Nat Rev Microbiol.

2017;15(6):323-37. Epub 20170410. doi: 10.1038/nrmicro.2017.20. PubMed PMID: 28392566.

13.   Schubert KA, Xu Y, Shao F, Auerbuch V. The *Yersinia* Type III Secretion System as a

Tool for Studying Cytosolic Innate Immune Surveillance. Annu Rev Microbiol. 2020;74:221-45.

Epub 20200713. doi: 10.1146/annurev-micro-020518-120221. PubMed PMID: 32660389.

14.   Brubaker RR. The Vwa+ virulence factor of yersiniae: the molecular basis of the

attendant nutritional requirement for Ca++. Rev Infect Dis. 1983;5 Suppl 4:S748-58. doi:

10.1093/clinids/5.supplement_4.s748. PubMed PMID: 6195719.

15.   Sturm A, Heinemann M, Arnoldini M, Benecke A, Ackermann M, Benz M, et al. The cost

of virulence: retarded growth of *Salmonella* Typhimurium cells expressing type III secretion

system 1. PLoS Pathog. 2011;7(7):e1002143. Epub 20110728. doi:

10.1371/journal.ppat.1002143. PubMed PMID: 21829349; PubMed Central PMCID:

PMCPMC3145796.

16.   Milne-Davies B, Helbig C, Wimmi S, Cheng DWC, Paczia N, Diepold A. Life After

Secretion-*Yersinia enterocolitica* Rapidly Toggles Effector Secretion and Can Resume Cell

Division in Response to Changing External Conditions. Front Microbiol. 2019;10:2128. Epub

20190913. doi: 10.3389/fmicb.2019.02128. PubMed PMID: 31572334; PubMed Central PMCID:

PMCPMC6753693.

17.   Bolin I, Portnoy DA, Wolf-Watz H. Expression of the temperature-inducible outer

membrane proteins of yersiniae. Infect Immun. 1985;48(1):234-40. doi: 10.1128/iai.48.1.234-

240.1985. PubMed PMID: 3980086; PubMed Central PMCID: PMCPMC261940.

18.   Bohme K, Steinmann R, Kortmann J, Seekircher S, Heroven AK, Berger E, et al.

Concerted actions of a thermo-labile regulator and a unique intergenic RNA thermosensor

control *Yersinia* virulence. PLoS Pathog. 2012;8(2):e1002518. Epub 20120216. doi:

10.1371/journal.ppat.1002518. PubMed PMID: 22359501; PubMed Central PMCID:

PMCPMC3280987.

19.   Dewoody RS, Merritt PM, Marketon MM. Regulation of the *Yersinia* type III secretion

system: traffic control. Front Cell Infect Microbiol. 2013;3:4. Epub 20130206. doi:

10.3389/fcimb.2013.00004. PubMed PMID: 23390616; PubMed Central PMCID:

PMCPMC3565153.

20.   Hoe NP, Goguen JD. Temperature sensing in *Yersinia pestis*: translation of the LcrF

activator protein is thermally regulated. J Bacteriol. 1993;175(24):7901-9. doi:

10.1128/jb.175.24.7901-7909.1993. PubMed PMID: 7504666; PubMed Central PMCID:

PMCPMC206968.

21.   Wang H, Avican K, Fahlgren A, Erttmann SF, Nuss AM, Dersch P, et al. Increased

plasmid copy number is essential for *Yersinia* T3SS function and virulence. Science.

2016;353(6298):492-5. Epub 20160630. doi: 10.1126/science.aaf7501. PubMed PMID:

27365311.

22. Schneiders S, Hechard T, Edgren T, Avican K, Fallman M, Fahlgren A, et al.

Spatiotemporal Variations in Growth Rate and Virulence Plasmid Copy Number during *Yersinia*

pseudotuberculosis Infection. Infect Immun. 2021;89(4). Epub 20210317. doi:

10.1128/IAI.00710-20. PubMed PMID: 33495272; PubMed Central PMCID: PMCPMC8090943.

23. Schott S, Scheuer R, Ermoli F, Glatter T, Evguenieva-Hackenberg E, Diepold A. A

ParDE toxin-antitoxin system is responsible for the maintenance of the *Yersinia* virulence

plasmid but not for type III secretion-associated growth inhibition. Front Cell Infect Microbiol.

2023;13:1166077. Epub 20230509. doi: 10.3389/fcimb.2023.1166077. PubMed PMID:

37228670; PubMed Central PMCID: PMCPMC10203498.

24. Villa L, Garcia-Fernandez A, Fortini D, Carattoli A. Replicon sequence typing of IncF

plasmids carrying virulence and resistance determinants. J Antimicrob Chemother.

2010;65(12):2518-29. Epub 20101008. doi: 10.1093/jac/dkq347. PubMed PMID: 20935300.

25. Nordstrom K. Plasmid R1--replication and its control. Plasmid. 2006;55(1):1-26. Epub

20050930. doi: 10.1016/j.plasmid.2005.07.002. PubMed PMID: 16199086.

26. Meynell E, Datta N. The relation of resistance transfer factors to the F-factor (sex-factor)

of *Escherichia coli* K12. Genet Res. 1966;7(1):134-40. doi: 10.1017/s0016672300009538.

PubMed PMID: 5324663.

27. Datta N, Kontomichalou P. Penicillinase synthesis controlled by infectious R factors in

Enterobacteriaceae. Nature. 1965;208(5007):239-41. doi: 10.1038/208239a0. PubMed PMID:

5326330.

28. Olsson JA, Paulsson J, Nordstrom K. Effect of the CopB auxiliary replication control

system on stability of maintenance of Par(+) plasmid R1. J Bacteriol. 2004;186(1):207-11. doi:

10.1128/JB.186.1.207-211.2004. PubMed PMID: 14679240; PubMed Central PMCID:

PMCPMC303431.

29.   Lopilato J, Bortner S, Beckwith J. Mutations in a new chromosomal gene of *Escherichia*

coli K-12, *pcnB*, reduce plasmid copy number of pBR322 and its derivatives. Mol Gen Genet.

1986;205(2):285-90. doi: 10.1007/BF00430440. PubMed PMID: 3100913.

30.   Liu JD, Parkinson JS. Genetics and sequence analysis of the *pcnB* locus, an

Escherichia coli gene involved in plasmid copy number control. J Bacteriol. 1989;171(3):1254-

61. doi: 10.1128/jb.171.3.1254-1261.1989. PubMed PMID: 2537812; PubMed Central PMCID:

PMCPMC209738.

31.   Sippel AE. Purification and characterization of adenosine triphosphate: ribonucleic acid

adenyltransferase from *Escherichia coli*. Eur J Biochem. 1973;37(1):31-40. doi: 10.1111/j.1432-

1033.1973.tb02953.x. PubMed PMID: 4580885.

32.   Cao GJ, Sarkar N. Identification of the gene for an *Escherichia coli* poly(A) polymerase.

Proc Natl Acad Sci U S A. 1992;89(21):10380-4. doi: 10.1073/pnas.89.21.10380. PubMed

PMID: 1438224; PubMed Central PMCID: PMCPMC50342.

33.   Martin G, Keller W. Mutational analysis of mammalian poly(A) polymerase identifies a

region for primer binding and catalytic domain, homologous to the family X polymerases, and to

other nucleotidyltransferases. EMBO J. 1996;15(10):2593-603. PubMed PMID: 8665867;

PubMed Central PMCID: PMCPMC450192.

34.   Masters M, March JB, Oliver IR, Collins JF. A possible role for the *pcnB* gene product of

Escherichia coli in modulating RNA: RNA interactions. Mol Gen Genet. 1990;220(2):341-4. doi:

10.1007/BF00260507. PubMed PMID: 1691435.

35.   Yue D, Maizels N, Weiner AM. CCA-adding enzymes and poly(A) polymerases are all

members of the same nucleotidyltransferase superfamily: characterization of the CCA-adding

enzyme from the archaeal hyperthermophile *Sulfolobus shibatae*. RNA. 1996;2(9):895-908.

PubMed PMID: 8809016; PubMed Central PMCID: PMCPMC1369424.

36.   Jones GH. Acquisition of pcnB [poly(A) polymerase I] genes via horizontal transfer from

the beta, gamma-Proteobacteria. Microb Genom. 2021;7(2). doi: 10.1099/mgen.0.000508.

PubMed PMID: 33502308; PubMed Central PMCID: PMCPMC8208693.

37.   Maes A, Gracia C, Innocenti N, Zhang K, Aurell E, Hajnsdorf E. Landscape of RNA

polyadenylation in *E. coli*. Nucleic Acids Res. 2017;45(5):2746-56. doi: 10.1093/nar/gkw894.

PubMed PMID: 28426097; PubMed Central PMCID: PMCPMC5389530.

38.   Joanny G, Le Derout J, Brechemier-Baey D, Labas V, Vinh J, Regnier P, et al.

Polyadenylation of a functional mRNA controls gene expression in *Escherichia coli*. Nucleic

Acids Res. 2007;35(8):2494-502. Epub 20070329. doi: 10.1093/nar/gkm120. PubMed PMID:

17395638; PubMed Central PMCID: PMCPMC1885654.

39.   Reichenbach B, Maes A, Kalamorz F, Hajnsdorf E, Gorke B. The small RNA GlmY acts

upstream of the sRNA GlmZ in the activation of glmS expression and is subject to regulation by

polyadenylation in *Escherichia coli*. Nucleic Acids Res. 2008;36(8):2570-80. Epub 20080311.

doi: 10.1093/nar/gkn091. PubMed PMID: 18334534; PubMed Central PMCID:

PMCPMC2377431.

40.   Mohanty BK, Maples VF, Kushner SR. Polyadenylation helps regulate functional tRNA

levels in *Escherichia coli*. Nucleic Acids Res. 2012;40(10):4589-603. Epub 20120128. doi:

10.1093/nar/gks006. PubMed PMID: 22287637; PubMed Central PMCID: PMCPMC3378859.

41.   Holmqvist E, Wagner EGH. Impact of bacterial sRNAs in stress responses. Biochem

Soc Trans. 2017;45(6):1203-12. Epub 20171103. doi: 10.1042/BST20160363. PubMed PMID:

29101308; PubMed Central PMCID: PMCPMC5730939.

42.   Sinha D, Matz LM, Cameron TA, De Lay NR. Poly(A) polymerase is required for RyhB

sRNA stability and function in *Escherichia coli*. RNA. 2018;24(11):1496-511. Epub 20180730.

doi: 10.1261/rna.067181.118. PubMed PMID: 30061117; PubMed Central PMCID:

PMCPMC6191717.

43. Francis N, Laishram RS. Transgenesis of mammalian PABP reveals mRNA

polyadenylation as a general stress response mechanism in bacteria. iScience.

2021;24(10):103119. Epub 20210911. doi: 10.1016/j.isci.2021.103119. PubMed PMID:

34646982; PubMed Central PMCID: PMCPMC8496165.

44. Masters M, Colloms MD, Oliver IR, He L, Macnaughton EJ, Charters Y. The *pcnB* gene

of *Escherichia coli*, which is required for ColE1 copy number maintenance, is dispensable. J

Bacteriol. 1993;175(14):4405-13. doi: 10.1128/jb.175.14.4405-4413.1993. PubMed PMID:

8331071; PubMed Central PMCID: PMCPMC204881.

45. Xu F, Lin-Chao S, Cohen SN. The *Escherichia coli pcnB* gene promotes adenylylation of

antisense RNAI of ColE1-type plasmids in vivo and degradation of RNAI decay intermediates.

Proc Natl Acad Sci U S A. 1993;90(14):6756-60. doi: 10.1073/pnas.90.14.6756. PubMed PMID:

7688127; PubMed Central PMCID: PMCPMC47011.

46. He L, Soderbom F, Wagner EG, Binnie U, Binns N, Masters M. PcnB is required for the

rapid degradation of RNAI, the antisense RNA that controls the copy number of ColE1-related

plasmids. Mol Microbiol. 1993;9(6):1131-42. doi: 10.1111/j.1365-2958.1993.tb01243.x. PubMed

PMID: 7523833.

47. Soderbom F, Binnie U, Masters M, Wagner EG. Regulation of plasmid R1 replication:

PcnB and RNase E expedite the decay of the antisense RNA, CopA. Mol Microbiol.

1997;26(3):493-504. doi: 10.1046/j.1365-2958.1997.5871953.x. PubMed PMID: 9402020.

48. Pienkoss S, Javadi S, Chaoprasid P, Holler M, Rossmanith J, Dersch P, et al. RNA

Thermometer-coordinated Assembly of the *Yersinia* Injectisome. J Mol Biol.

2022;434(18):167667. Epub 20220603. doi: 10.1016/j.jmb.2022.167667. PubMed PMID:

35667470.

49. Kupferberg LL, Higuchi K. Role of calcium ions in the stimulation of growth of virulent

strains of *Pasteurella pestis*. J Bacteriol. 1958;76(1):120-1. doi: 10.1128/jb.76.1.120-121.1958.

PubMed PMID: 13563403; PubMed Central PMCID: PMCPMC290166.

50. Higuchi K, Kupferberg LL, Smith JL. Studies on the nutrition and physiology of

Pasteurella pestis. III. Effects of calcium ions on the growth of virulent and avirulent strains of

Pasteurella pestis. J Bacteriol. 1959;77(3):317-21. doi: 10.1128/jb.77.3.317-321.1959. PubMed

PMID: 13641190; PubMed Central PMCID: PMCPMC290369.

51. Rosqvist R, Magnusson KE, Wolf-Watz H. Target cell contact triggers expression and

polarized transfer of *Yersinia* YopE cytotoxin into mammalian cells. EMBO J. 1994;13(4):964-

72. doi: 10.1002/j.1460-2075.1994.tb06341.x. PubMed PMID: 8112310; PubMed Central

PMCID: PMCPMC394898.

52. Takacs CN, Wachter J, Xiang Y, Ren Z, Karaboja X, Scott M, et al. Polyploidy, regular

patterning of genome copies, and unusual control of DNA partitioning in the Lyme disease

spirochete. Nat Commun. 2022;13(1):7173. Epub 20221122. doi: 10.1038/s41467-022-34876-4.

PubMed PMID: 36450725; PubMed Central PMCID: PMCPMC9712426.

53. Erdmann N, Petroff T, Funnell BE. Intracellular localization of P1 ParB protein depends

on ParA and parS. Proc Natl Acad Sci U S A. 1999;96(26):14905-10. doi:

10.1073/pnas.96.26.14905. PubMed PMID: 10611311; PubMed Central PMCID:

PMCPMC24746.

54. Wang JD, Levin PA. Metabolism, cell growth and the bacterial cell cycle. Nat Rev

Microbiol. 2009;7(11):822-7. Epub 20091006. doi: 10.1038/nrmicro2202. PubMed PMID:

19806155; PubMed Central PMCID: PMCPMC2887316.

55. Cooper S, Helmstetter CE. Chromosome replication and the division cycle of

Escherichia coli B/r. J Mol Biol. 1968;31(3):519-40. doi: 10.1016/0022-2836(68)90425-7.

PubMed PMID: 4866337.

56. Miller HK, Kwuan L, Schwiesow L, Bernick DL, Mettert E, Ramirez HA, et al. IscR is

essential for *Yersinia pseudotuberculosis* type III secretion and virulence. PLoS Pathog.

2014;10(6):e1004194. Epub 20140612. doi: 10.1371/journal.ppat.1004194. PubMed PMID:

24945271; PubMed Central PMCID: PMCPMC4055776.

57.   Zhou P, She Y, Dong N, Li P, He H, Borio A, et al. Alpha-kinase 1 is a cytosolic innate

immune receptor for bacterial ADP-heptose. Nature. 2018;561(7721):122-6. Epub 20180815.

doi: 10.1038/s41586-018-0433-3. PubMed PMID: 30111836.

58.   Orth K, Palmer LE, Bao ZQ, Stewart S, Rudolph AE, Bliska JB, et al. Inhibition of the

mitogen-activated protein kinase kinase superfamily by a *Yersinia* effector. Science.

1999;285(5435):1920-3. doi: 10.1126/science.285.5435.1920. PubMed PMID: 10489373.

59.   Orth K, Xu Z, Mudgett MB, Bao ZQ, Palmer LE, Bliska JB, et al. Disruption of signaling

by *Yersinia* effector YopJ, a ubiquitin-like protein protease. Science. 2000;290(5496):1594-7.

doi: 10.1126/science.290.5496.1594. PubMed PMID: 11090361.

60.   Riley G, Toma S. Detection of pathogenic *Yersinia enterocolitica* by using congo red-

magnesium oxalate agar medium. J Clin Microbiol. 1989;27(1):213-4. doi:

10.1128/jcm.27.1.213-214.1989. PubMed PMID: 2913030; PubMed Central PMCID:

PMCPMC267269.

61.   Raynal LC, Carpousis AJ. Poly(A) polymerase I of *Escherichia coli*: characterization of

the catalytic domain, an RNA binding site and regions for the interaction with proteins involved

in mRNA degradation. Mol Microbiol. 1999;32(4):765-75. doi: 10.1046/j.1365-

2958.1999.01394.x. PubMed PMID: 10361280.

62.   Toh Y, Takeshita D, Nagaike T, Numata T, Tomita K. Mechanism for the alteration of the

substrate specificities of template-independent RNA polymerases. Structure. 2011;19(2):232-

43. doi: 10.1016/j.str.2010.12.006. PubMed PMID: 21300291.

63.   Million-Weaver S, Alexander DL, Allen JM, Camps M. Quantifying plasmid copy number

to investigate plasmid dosage effects associated with directed protein evolution. Methods Mol

Biol. 2012;834:33-48. doi: 10.1007/978-1-61779-483-4_3. PubMed PMID: 22144351; PubMed

Central PMCID: PMCPMC3804865.

64.   Amann E, Ochs B, Abel KJ. Tightly regulated tac promoter vectors useful for the

expression of unfused and fused proteins in *Escherichia coli.* Gene. 1988;69(2):301-15. doi:

10.1016/0378-1119(88)90440-4. PubMed PMID: 3069586.

65.   Lacatena RM, Cesareni G. Base pairing of RNA I with its complementary sequence in

the primer precursor inhibits ColE1 replication. Nature. 1981;294(5842):623-6. doi:

10.1038/294623a0. PubMed PMID: 6171736.

66.   Brantl S. Plasmid Replication Control by Antisense RNAs. Microbiol Spectr.

2014;2(4):PLAS-0001-2013. doi: 10.1128/microbiolspec.PLAS-0001-2013. PubMed PMID:

26104196.

67.   Selzer G, Som T, Itoh T, Tomizawa J. The origin of replication of plasmid p15A and

comparative studies on the nucleotide sequences around the origin of related plasmids. Cell.

1983;32(1):119-29. doi: 10.1016/0092-8674(83)90502-0. PubMed PMID: 6186390.

68.   Uga H, Matsunaga F, Wada C. Regulation of DNA replication by iterons: an interaction

between the ori2 and incC regions mediated by RepE-bound iterons inhibits DNA replication of

mini-F plasmid in *Escherichia* coli. EMBO J. 1999;18(13):3856-67. doi:

10.1093/emboj/18.13.3856. PubMed PMID: 10393200; PubMed Central PMCID:

PMCPMC1171462.

69.   Jurenas D, Fraikin N, Goormaghtigh F, De Bruyn P, Vandervelde A, Zedek S, et al.

Bistable Expression of a Toxin-Antitoxin System Located in a Cryptic Prophage of *Escherichia*

coli O157:H7. mBio. 2021;12(6):e0294721. Epub 20211130. doi: 10.1128/mBio.02947-21.

PubMed PMID: 34844426; PubMed Central PMCID: PMCPMC8630535.

70.   Mecsas J, Bilis I, Falkow S. Identification of attenuated *Yersinia pseudotuberculosis*

strains and characterization of an orogastric infection in BALB/c mice on day 5 postinfection by

signature-tagged mutagenesis. Infect Immun. 2001;69(5):2779-87. doi: 10.1128/IAI.67.5.2779-

2787.2001. PubMed PMID: 11292689; PubMed Central PMCID: PMCPMC98225.

71.   Spickler C, Mackie GA. Action of RNase II and polynucleotide phosphorylase against

RNAs containing stem-loops of defined structure. J Bacteriol. 2000;182(9):2422-7. doi:

10.1128/JB.182.9.2422-2427.2000. PubMed PMID: 10762241; PubMed Central PMCID:

PMCPMC111303.

72.   O'Hara EB, Chekanova JA, Ingle CA, Kushner ZR, Peters E, Kushner SR.

Polyadenylylation helps regulate mRNA decay in *Escherichia coli*. Proc Natl Acad Sci U S A.

1995;92(6):1807-11. doi: 10.1073/pnas.92.6.1807. PubMed PMID: 7534403; PubMed Central

PMCID: PMCPMC42371.

73.   Hajnsdorf E, Braun F, Haugel-Nielsen J, Regnier P. Polyadenylylation destabilizes the

rpsO mRNA of *Escherichia coli*. Proc Natl Acad Sci U S A. 1995;92(9):3973-7. doi:

10.1073/pnas.92.9.3973. PubMed PMID: 7732015; PubMed Central PMCID: PMCPMC42084.

74.   Xu F, Cohen SN. RNA degradation in *Escherichia coli* regulated by 3' adenylation and 5'

phosphorylation. Nature. 1995;374(6518):180-3. doi: 10.1038/374180a0. PubMed PMID:

7533264.

75.   Blum E, Carpousis AJ, Higgins CF. Polyadenylation promotes degradation of 3'-

structured RNA by the *Escherichia coli* mRNA degradosome in vitro. J Biol Chem.

1999;274(7):4009-16. doi: 10.1074/jbc.274.7.4009. PubMed PMID: 9933592.

76.   Coburn GA, Miao X, Briant DJ, Mackie GA. Reconstitution of a minimal RNA

degradosome demonstrates functional coordination between a 3' exonuclease and a DEAD-box

RNA helicase. Genes Dev. 1999;13(19):2594-603. doi: 10.1101/gad.13.19.2594. PubMed

PMID: 10521403; PubMed Central PMCID: PMCPMC317069.

77.   Marujo PE, Hajnsdorf E, Le Derout J, Andrade R, Arraiano CM, Regnier P. RNase II

removes the oligo(A) tails that destabilize the rpsO mRNA of *Escherichia coli*. RNA.

2000;6(8):1185-93. doi: 10.1017/s135583820000073x. PubMed PMID: 10943897; PubMed

Central PMCID: PMCPMC1369992.

78.   Rosenzweig JA, Chromy B, Echeverry A, Yang J, Adkins B, Plano GV, et al.

Polynucleotide phosphorylase independently controls virulence factor expression levels and

export in *Yersinia* spp. FEMS Microbiol Lett. 2007;270(2):255-64. Epub 20070328. doi:

10.1111/j.1574-6968.2007.00689.x. PubMed PMID: 17391372.

79.   Rosenzweig JA, Weltman G, Plano GV, Schesser K. Modulation of *Yersinia* type three

secretion system by the S1 domain of polynucleotide phosphorylase. J Biol Chem.

2005;280(1):156-63. Epub 20041026. doi: 10.1074/jbc.M405662200. PubMed PMID: 15509583.

80.   Knittel V, Vollmer I, Volk M, Dersch P. Discovering RNA-Based Regulatory Systems for

Yersinia Virulence. Front Cell Infect Microbiol. 2018;8:378. Epub 20181025. doi:

10.3389/fcimb.2018.00378. PubMed PMID: 30460205; PubMed Central PMCID:

PMCPMC6232918.

81.   Mohanty BK, Maples VF, Kushner SR. The Sm-like protein Hfq regulates

polyadenylation dependent mRNA decay in Escherichia coli. Mol Microbiol. 2004;54(4):905-20.

doi: 10.1111/j.1365-2958.2004.04337.x. PubMed PMID: 15522076.

82.   Werbowy O, Boratynski R, Dekowska A, Kaczorowski T. Genetic analysis of

maintenance of pEC156, a naturally occurring *Escherichia coli* plasmid that carries genes of the

EcoVIII restriction-modification system. Plasmid. 2015;77:39-50. Epub 20141208. doi:

10.1016/j.plasmid.2014.12.002. PubMed PMID: 25500017.

83.   Francis N, Behera MR, Natarajan K, Laishram RS. Tyrosine phosphorylation controlled

poly(A) polymerase I activity regulates general stress response in bacteria. Life Sci Alliance.

2023;6(3). Epub 20221219. doi: 10.26508/lsa.202101148. PubMed PMID: 36535710; PubMed

Central PMCID: PMCPMC9764084.

84.   Ottoz DSM, Berchowitz LE. The role of disorder in RNA binding affinity and specificity.

Open Biol. 2020;10(12):200328. Epub 20201223. doi: 10.1098/rsob.200328. PubMed PMID:

33352065; PubMed Central PMCID: PMCPMC7776568.

85.   Milton DL, Norqvist A, Wolf-Watz H. Cloning of a metalloprotease gene involved in the

virulence mechanism of *Vibrio anguillarum*. J Bacteriol. 1992;174(22):7235-44. doi:

10.1128/jb.174.22.7235-7244.1992. PubMed PMID: 1429449; PubMed Central PMCID:

PMCPMC207417.

86.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics. 2014;30(15):2114-20. Epub 20140401. doi: 10.1093/bioinformatics/btu170.

PubMed PMID: 24695404; PubMed Central PMCID: PMCPMC4103590.

87.   Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009;25(14):1754-60. Epub 20090518. doi: 10.1093/bioinformatics/btp324.

PubMed PMID: 19451168; PubMed Central PMCID: PMCPMC2705234.

88.   Johnson SL, Daligault HE, Davenport KW, Jaissle J, Frey KG, Ladner JT, et al. Thirty-

Two Complete Genome Assemblies of Nine *Yersinia* Species, Including *Y. pestis*, *Y.*

pseudotuberculosis, and *Y. enterocolitica*. Genome Announc. 2015;3(2). Epub 20150430. doi:

10.1128/genomeA.00148-15. PubMed PMID: 25931590; PubMed Central PMCID:

PMCPMC4417686.

89.   Zoubeidi A, Schwiesow L, Auerbuch V, Lam HN. Complete Genome Assembly of

Yersinia pseudotuberculosis IP2666pIB1. Microbiol Resour Announc. 2019;8(7). Epub

20190214. doi: 10.1128/MRA.01592-18. PubMed PMID: 30801065; PubMed Central PMCID:

PMCPMC6376424.

90.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9. Epub 20090608. doi:

10.1093/bioinformatics/btp352. PubMed PMID: 19505943; PubMed Central PMCID:

PMCPMC2723002.

91.   Auwera Gvd, O'Connor BD. Genomics in the cloud : using Docker, GATK, and WDL in

Terra. First edition. ed. Sebastopol, CA: O'Reilly Media; 2020. xxiv, 467 pages p.

92.   Hechard T, Wang H. Determination of Growth Rate and Virulence Plasmid Copy

Number During *Yersinia pseudotuberculosis* Infection Using Droplet Digital PCR. Methods Mol

Biol. 2023;2674:101-15. doi: 10.1007/978-1-0716-3243-7_7. PubMed PMID: 37258963.

93.   Paintdakhi A, Parry B, Campos M, Irnov I, Elf J, Surovtsev I, et al. Oufti: an integrated

software package for high-accuracy, high-throughput quantitative microscopy analysis. Mol

Microbiol. 2016;99(4):767-77. Epub 20151218. doi: 10.1111/mmi.13264. PubMed PMID:

26538279; PubMed Central PMCID: PMCPMC4752901.

94.   Campos M, Govers SK, Irnov I, Dobihal GS, Cornet F, Jacobs-Wagner C. Genomewide

phenotypic analysis of growth, cell morphogenesis, and cell cycle events in *Escherichia coli*. Mol

Syst Biol. 2018;14(6):e7573. Epub 20180625. doi: 10.15252/msb.20177573. PubMed PMID:

29941428; PubMed Central PMCID: PMCPMC6018989.

95.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an

open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82. Epub

20120628. doi: 10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central PMCID:

PMCPMC3855844.

96.   Ovesny M, Krizek P, Borkovec J, Svindrych Z, Hagen GM. ThunderSTORM: a

comprehensive ImageJ plug-in for PALM and STORM data analysis and super-resolution

imaging. Bioinformatics. 2014;30(16):2389-90. Epub 20140425. doi:

10.1093/bioinformatics/btu202. PubMed PMID: 24771516; PubMed Central PMCID:

PMCPMC4207427.

97.   Otsu N. A Threshold Selection Method from Gray-Level Histograms. IEEE Trans Syst

Man Cybern. 1979;9(1):62-6.

98.   Kwuan L, Adams W, Auerbuch V. Impact of host membrane pore formation by the

Yersinia pseudotuberculosis type III secretion system on the macrophage innate immune

response. Infect Immun. 2013;81(3):905-14. Epub 20130107. doi: 10.1128/IAI.01014-12.

PubMed PMID: 23297383; PubMed Central PMCID: PMCPMC3584892.

99.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and

genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol. 2019;37(8):907-15. Epub

20190802. doi: 10.1038/s41587-019-0201-4. PubMed PMID: 31375807; PubMed Central

PMCID: PMCPMC7605509.

100.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics. 2014;30(7):923-30. Epub

20131113. doi: 10.1093/bioinformatics/btt656. PubMed PMID: 24227677.

101.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. Bioinformatics. 2010;26(1):139-40. Epub

20091111. doi: 10.1093/bioinformatics/btp616. PubMed PMID: 19910308; PubMed Central

PMCID: PMCPMC2796818.

102.   Metsalu T, Vilo J. ClustVis: a web tool for visualizing clustering of multivariate data using

Principal Component Analysis and heatmap. Nucleic Acids Res. 2015;43(W1):W566-70. Epub

20150512. doi: 10.1093/nar/gkv468. PubMed PMID: 25969447; PubMed Central PMCID:

PMCPMC4489295.

103.   Auerbuch V, Golenbock DT, Isberg RR. Innate immune recognition of *Yersinia*

pseudotuberculosis type III secretion. PLoS Pathog. 2009;5(12):e1000686. Epub 20091204.

doi: 10.1371/journal.ppat.1000686. PubMed PMID: 19997504; PubMed Central PMCID:

PMCPMC2779593.

1467

Figure 1

Figure 2

Figure 4

Figure 5

A

Δ2.5ΔpcnB

VP⁻   Δ2.5

B   Δ2.5   Δ2.5ΔpcnB

pTRC99   pTRC99   pTRC99   pTRC99
empty   PAP I   empty   PAP I

IpaB →

IpaC →
IpaD →

Pellets

C   Δ2.5   Δ2.5ΔpcnB

pTRC99   pTRC99   pTRC99   pTRC99
empty   PAP I   empty   PAP I

Supernatants

D

No Selection
(No ampicillin)

+ Selection for pTRC99 and IPTG
(+ampicillin)

Δ2.5 pTrc99::empty

Δ2.5ΔpcnB pTrc99::empty

Δ2.5 pTrc99::PAPI

Δ2.5ΔpcnB pTrc99::PAPI