<u>**Title**</u>

Astrocyte regional specialization is shaped by postnatal development

<u>**Authors**</u>

Margaret E. Schroeder[1,2], Dana M. McCormack[1], Lukas R. Metzner[1], Jinyoung Kang[1,2], Katelyn X. Li[1], Eunah Yu[1],

Lisa Melamed[1], Kirsten M. Levandowski[1,4], Heather Zaniewski[1], Qiangge Zhang[1,4], Edward S. Boyden[1,2,3,5,6,7,8,9],

Fenna M. Krienen[10,*], and Guoping Feng[1,2,3,4*]

<u>**Affiliations**</u>

1. McGovern Institute for Brain Research, MIT, Cambridge, MA, USA.
2. Department of Brain and Cognitive Sciences, MIT, Cambridge, MA, USA.
3. Yang Tan Collective, MIT, Cambridge, MA, USA.
4. The Broad Institute of MIT and Harvard, Cambridge, MA, USA.
5. Center for Neurobiological Engineering and K. Lisa Yang Center for Bionics, MIT, Cambridge, MA, USA.
6. Department of Biological Engineering, MIT, Cambridge, MA, USA.
7. Koch Institute, MIT, Cambridge, MA, USA.
8. Howard Hughes Medical Institute, Cambridge, MA, USA.
9. Media Arts and Sciences, MIT, Cambridge, MA, USA.
10. Princeton Neuroscience Institute, Princeton University, Princeton, NJ, USA

*Corresponding author

<u>**Abstract**</u>

Astrocytes are an abundant class of glial cells with critical roles in neural circuit assembly and function. Though

many studies have uncovered significant molecular distinctions between astrocytes from different brain regions,

how this regionalization unfolds over development is not fully understood. We used single-nucleus RNA sequencing

to characterize the molecular diversity of brain cells across six developmental stages and four brain regions in the

mouse and marmoset brain. Our analysis of over 170,000 single astrocyte nuclei revealed striking regional

heterogeneity among astrocytes, particularly between telencephalic and diencephalic regions, at all developmental

time points surveyed in both species. At the stages sampled, most of the region patterning was private to astrocytes

and not shared with neurons or other glial types. Though astrocytes were already regionally patterned in late

embryonic stages, this region-specific astrocyte gene expression signature changed dramatically over postnatal

development, and its composition suggests that regional astrocytes further specialize postnatally to support their

local neuronal circuits. Across mouse and marmoset, we found hundreds of species differentially expressed genes,

as well as divergence in the expression of astrocytic region- and age-differentially expressed genes and the timing

of astrocyte maturation relative to birth between the species. Finally, we used expansion microscopy to show that

astrocyte morphology is also regionally specialized across cortex, striatum, and thalamus in the mouse.

<u>**Introduction**</u>

The mammalian brain is composed of thousands of heterogeneous molecularly-defined cell types[1,2]. This

heterogeneity is prominent between cells from different anatomical regions that arise from distinct developmental

compartments. This regional specialization is critical for circuit formation and proper brain function. In recent years,

this heterogeneity has been cataloged through large-scale single-cell and single-nucleus RNA sequencing

(scRNAseq and snRNAseq, respectively), which enables molecular profiling in unprecedented detail and scale[3].

The past decade has seen the publication of multiple brain cell type transcriptomic atlases, including of the entire

adult mouse brain[1,2], the adult human brain[4], the developing mouse[5] and human[6] brains, and the adult marmoset

brain[7,8]. Most, but not all of these atlases have focused primarily on characterizing neurons, long considered the

brain's principal cell type. Indeed, several studies used cell sorting methods to enrich for neurons[9].

Astrocytes, an abundant class of glia, play critical roles in neuronal circuit assembly and function in healthy and

pathological states[10–14]. While their morphological heterogeneity has long been appreciated[15,16], their molecular

heterogeneity, particularly across brain regions, was only revealed more recently by microarray[17] and bulk RNA
sequencing studies[18–20], and later by single-cell RNA sequencing studies[8,21–25]. Early lineage tracing studies in the
mouse spinal cord and brain revealed that astrocyte precursors from different embryonic domains are molecularly
distinct[26,27]. Adult mouse astrocytes maintain epigenetic marks from their region-restricted radial glia ancestors[28],
which may contribute to the significant heterogeneity of adult astrocyte populations. There is also abundant
evidence supporting the role of extrinsic cues in astrocyte regionalization, including the formation of various cortical
morphological subtypes from a shared astrocyte progenitor[29], the up- or down-regulation of ion channels,
transporters, receptors in response to neuronal inputs[30], and the molecular and morphological adaptation of distinct
developmentally-patterned septal astrocyte subtypes after cross-region heterotopic transplant[31].

As has been found with neurons, it is likely that transcriptionally-defined astrocyte populations are developmentally
influenced by their respective microenvironments and perform distinct functions. Yet, the developmental time course
of astrocyte regional patterning, the composition of astrocyte subtypes over development, and the conservation of
these features between rodents and primates remain unclear[32]. To address this knowledge gap, we applied
snRNAseq to characterize astrocyte molecular diversity across six developmental stages and four brain regions in
mouse and marmoset. To complement the transcriptomic studies, we characterized complex astrocyte morphology
and protein localization at high resolution across brain regions using expansion microscopy.

We used single nucleus sequencing to generate a dataset of 1.4 million brain cell nuclei across multiple stages and
brain regions in mouse and marmoset. A unified study, with data generated from a single lab using highly consistent
methodology, has the advantage of reduced technical variation compared to datasets integrated across research
groups, nuclei isolation protocols, and sequencing platforms, which is difficult to remove *in silico*[33,34].

Our analysis shows that astrocytes are regionally patterned before birth and at all subsequent time points.
Importantly, we found dramatic changes in the transcriptional signatures underlying astrocyte regional identity
between birth and early adolescence in both species, highlighting the importance of postnatal regional cues in
shaping astrocyte identity. We explored the functional implications of genes differentially expressed between
astrocytes from different brain regions, and between astrocytes at different developmental time points. Furthermore,
we identified both region-shared and region-divergent developmental transcriptional signatures in astrocytes.

Many of the region-, age-, and species-differentially expressed genes in astrocytes implicated morphogenesis
pathways. Indeed, astrocyte morphology, which is highly ramified and complex, including sub-micron scale
processes that contact synapses and blood vessels, is essential for their many functions[35,36]. Therefore, to assess
whether astrocyte morphology is also regionally specialized, we used a new variant of expansion microscopy, ExR[37]
to characterize virally-labeled astrocyte morphology and nanoscale protein expression with enhanced resolution.
We found that gray matter thalamic astrocytes in mice were significantly smaller and less complex than their striatal
and cortical counterparts, alongside differences in protein expression.

Results
A multi-region transcriptomic atlas of the developing mouse and marmoset brain
To create the cross-region, cross-species, cross-development snRNAseq atlas, we dissected prefrontal cortex
(PFC), motor cortex (MO), striatum, and thalamus from freshly harvested mouse and marmoset brains at late
embryonic, neonatal, early adolescent, late adolescent, young adult, and aged timepoints and snap-froze the tissue
(**Fig. 1A**). We collected tissues from 2 marmoset donors, one male and one female, at gestational day (GD)135,
neonate, 7 months, 14 months, 30 months (4 donors, previously collected data in the lab) and 11+ years). For
mouse, we collected tissue from 3 mouse biological replicates, at least one female, at E18.5, P4, P14, P32, P90,
and 90 weeks (see **Table S1** for mapping of donors to biological replicates, see **Methods**). We generated single-
nuclei suspensions from the snap-frozen tissue without enriching for any particular cell type, and generated single-
nucleus transcriptomes using 10x Genomics Chromium v3.1 chemistry (see **Fig. S1** for sequencing coverage
statistics). Though the adult (4 donors aged 29-32 months, together labeled 30 months) marmoset snRNAseq data
was generated using a different nuclei isolation protocol and reference genome[7], the data integrated very well

across studies (**Fig. S2A**). The data were also well integrated across biological sex (**Fig. S2-3A**). Quantitative
measures of integration quality[38,39] suggest our integration is well-mixed across biological replicates while
preserving true biological variability (donor mixing = 0.8981 for mouse and 0.9155 for marmoset; neighbor
consistency= 0.6680 for mouse and 0.4694 for marmoset; and average silhouette width = 0.6437 for mouse and
0.6634 for marmoset; see **Methods**).
108
After rigorous quality control, including removal of ambient RNA, low-quality nuclei, and doublets (see **Methods**),
we obtained 597,668 mouse nuclei and 881,832 marmoset nuclei, which were composed of 12 broad cell classes
(**Fig. 1B(i), Fig. 1C(i); Fig. S2-3**). We annotated more granular cell type (Leiden[40]-determined) clusters within each
cell class (**Fig. S4**, **Tables S2-4**). We employed the Allen Brain Cell Atlas's (ABCA) MapMyCells[41] portal[41] to refine
our annotations of neuronal subtypes and to help correct for modest cross-region contamination resulting from
dissection error (**Fig. S5-6**, see **Methods**). Throughout the paper, "dissected" brain region refers to the original
region label for the sample in which the nucleus was processed, while "region" or "assigned region" refers to the
brain region assigned post-hoc in the case of cells informatically predicted to arise from neighboring structures.
117
The total neuron-to-astrocyte ratio (across regions and developmental time points) was modestly higher in mouse
(6.28) than marmoset (5.24). To more quantitatively assess cell type composition differences across dissected
region (cortex, striatum, or thalamus), age, and sex, we used single-cell compositional data analysis (scCODA)[42],
which implements a Bayesian model of cell type counts to address the issue of low sample sizes in snRNAseq data.
scCODA confirmed many significant differences in cell type proportion between regions and ages for each species,
including the expected low numbers of excitatory neurons in striatum, increasing oligodendrocyte abundance with
age in both species, and minimal sex differences in cell type composition (**Fig. S2-3C**, **Tables S5-8**).
125
We found several cell type clusters that were enriched or depleted in developing (late embryonic or neonate) brains
(**Fig. S4C,F**). For example, in both species, there were immature cortical excitatory neuron, microglia, and astrocyte
clusters composed mostly of nuclei from late embryonic and fetal donors. The committed oligodendrocyte precursor
(COP) and newly formed oligodendrocyte (NFOL) cluster was primarily composed of nuclei from the neonate time
point in marmoset, with some nuclei even coming from late embryonic donors, but was primarily composed of nuclei
from early adolescent (P14) donors in mouse, with no COP/NFOLs coming from E18.5 mouse, indicating earlier
oligodendrocyte maturation in marmoset.
133
Astrocyte regional heterogeneity is embryonically patterned and unfolds over postnatal development
We observed striking regional heterogeneity among astrocytes at all developmental time points sampled in both
species, particularly between astrocytes of diencephalic (thalamus) and telencephalic (cortex and striatum) origin
(**Fig. 2A, Fig. 3A**). This is in line with multiple studies demonstrating embryonic regional patterning of
astrocytes[6,8,27]. These regional populations further divided into an immature population, primarily composed of
nuclei from late embryonic and neonatal time points, and a mature population, composed of nuclei from late
adolescent timepoints onward. These separate populations suggest embryonically-patterned regional astrocyte
populations undergo significant changes from the time of birth (neonate or P4) to early adolescence (7 months or
P14). Notably, abundant populations of immature astrocytes remained present in the mouse striatum through
adulthood (P90).
144
Marmoset. Transcription factors and morphogen gradients set up initial boundaries between developmental
compartments such as the telencephalon and diencephalon[43,44]. It could be that such early influences are present
only transiently at initial astrocyte specification, or that later stages retain initial molecular distinctions and
accumulate others over development. A combination of the two is also possible, where some genes follow one
pattern (developmentally transient expression) or the other (sustained expression throughout the lifespan). We
calculated **r**egional **d**ifferentially **e**xpressed **g**enes (rDEGs) at each developmental time point from metacells, 1-
dimensional vectors of averaged normalized expression across all cells in a given grouping (see **Methods**), of each
region. For marmoset, where each donor (biological replicate) was represented in each brain region, we calculated
rDEGs separately for each donor and required that rDEGs be above threshold (minimum expression and log fold-

change) requirements in both donors. We found 70 rDEGs whose expression differed between fetal cortical and
thalamic astrocytes, 142 of such rDEGs by early adolescence (7 months), and 134 in adulthood (30 months, see
Fig. 2B(i-iii) for the expression pattern of the union of these rDEGs at each timepoint). Focusing on cortex-thalamus
rDEGs (which were most numerous, **Table S9**), we found that relatively few persisted across all developmental
timepoints (**Fig. 2C**). 50% were shared between fetal and neonate, and 51% between late adolescent and aged,
but only 4% continued to act as regional patterning signatures throughout the lifespan. We found that many more
rDEGs were shared between late adolescent, young adult, and aged time points (61) than between fetal, neonate,
and 7-month time points (23) and neonate, 7-month, and 14-month time points (25, **Fig. 2C**). These data suggest
that regional astrocyte gene expression signatures emerge in the embryonic brain, change drastically over the
course of early postnatal development and stabilize during adolescence into adulthood.
164
If early telencephalic and diencephalic patterning persists in astrocytes, cortex and striatum should retain common
rDEGs compared to thalamus. To assess the degree of pairwise astrocyte rDEGs sharing across the 3 brain
structures, we correlated the log fold-change difference in astrocyte regional gene expression between different
region pairs (e.g., cortex-striatum vs. cortex-thalamus) for both rDEGs (log fold-change > 0.5) and non-rDEGs at
fetal, early adolescent, and adult timepoints. We found that cortex-striatum vs. cortex-thalamus fold-changes
exhibited a high degree of correlation in fetal marmoset (Pearson's r = 0.78), which decreased over developmental
time (Pearson's r = 0.47 in adult marmoset, **Fig. S7A-C(i)**). At all 3 time points, striatum-thalamus vs. striatum-
cortex fold-changes were negatively or uncorrelated (r = -0.43, -0.26, and 0.04 at GD135, 7 months, and 30 months
respectively, **Fig. S7A-C(ii)**). Finally, thalamus-striatum vs. thalamus-cortex fold-changes were highly correlated
(r > 0.80) at all 3 timepoints (**Fig. S7A-C(iii)**). Together, these results suggest that cortical and striatal astrocytes
share transcriptional divergence from thalamic astrocytes at all ages, but become more transcriptionally similar later
in development. At the same time, cortical and thalamic astrocytes both diverge from striatal astrocytes, but in
distinct ways, as indicated by the negative correlation between striatum-thalamus and striatum-cortex fold-changes
at GD135 and 7 months, and the lack of correlation at 30 months.
179
Dorsal radial glia populate the neocortex in a stereotyped progression, giving rise first to glutamatergic neurons,
then to astrocytes, and finally to oligodendrocytes[45–47]. In the developing thalamus, radial glial progenitors likely
follow the same cell type sequence[46,48]. As we showed previously for variable genes across the adult neocortex[7],
far more adult cortex-thalamus rDEGs are private to astrocytes than are shared with neurons or OPCs, despite their
shared lineage[47] (**Fig. 2D**). Surprisingly, this remained true even at the earliest stages we sampled (GD135 and
neonate). Together these observations suggest that astrocytes gain regional identity early in their maturation, but
that their continued regional identity is facilitated by distinct genes across their lifespan (**Fig. 2C**).
187
Many rDEGs nominate core cellular functions that may be further regionally specialized in astrocytes. For example,
ephrins such as *EFNB2* (identified as rDEG in 30 month marmoset astrocytes) and *EFNA5* (7, 14, and 30 months
and aged) are up-regulated in cortical astrocytes and ephrin receptor *EPHB1* (7, 14, and 30 months and aged) is
upregulated in cortical and striatal astrocytes. Neuron-astrocyte signaling via ephrin ligands and receptors regulates
axon guidance and synaptogenesis[49]. Thus, neuron-astrocyte ephrin signaling may be specialized in the
telencephalon. Cyclic-AMP-related signaling molecules *ADCY1* (30 months and aged) and *ADCY8* (identified at all
ages) are upregulated in thalamic astrocytes. As in neurons, astrocytic cAMP is an important second messenger
following GPCR activation[50], and modulates synaptic plasticity[51]. *ITPR1* (7, 14, and 30 months and aged), a calcium
channel that controls calcium release from the endoplasmic reticulum, an important source of intracellular calcium
during astrocyte signaling[52], is also upregulated in thalamic astrocytes. These rDEGs suggest that thalamic
astrocytes may have developed specialized pathways for calcium and cAMP signaling, potentially in response to
the release of upstream GPCR ligands by thalamus-projecting and thalamic neurons. Additionally, astrocyte rDEGs
included ion channels (e.g. *TRPM3* (GD135, 7, 14, and 30 months, and aged), a non-selective Ca^{2+} permeable ion
channel and thalamic rDEG); synapse-related proteins (e.g., *SPARC, a thalamic rDEG at* 7, 14, and 30 months and
aged, which regulates synaptogenesis[53]); neurotransmitter transporters and receptors (e.g., *SLC6A11*/GAT3 (7, 14,
and 30 months and aged), a thalamic rDEG and GABA transporter, *SLC1A3*/GLAST (at GD135 and neonate), a
glutamate transporter higher in the cortex and commonly used astrocyte marker gene, and *GRM3*/mGluR3 (all ages

except GD135), a telencephalic rDEG and metabotropic glutamate receptor); and a thyroid hormone receptor
(*SLCO1C1* (GD135, neonate, and 30 months), a cortical and later striatal rDEG). These rDEGS point more directly
to astrocyte adaptation to the local synaptic and neuronal niche.
208
To characterize astrocyte rDEG pathways in a more unbiased manner, we used WebGestalt 2024[54,55] over-
representation analysis to test for enrichment of cortex-thalamus rDEGs (bidirectionally, i.e. upregulated either in
cortex or thalamus) in GO Biological Process and KEGG pathways. Enriched pathways implicated oxytocin and
calcium signaling and neuronal projection development for GD135 astrocyte rDEGs; ephrin signaling, synaptic
transmission, and calcium ion homeostasis for 7-month astrocyte rDEGs; and glutamatergic synaptic transmission,
oxytocin signaling, and cGMP-PKG signaling for adult marmoset astrocyte rDEGs (**Fig. 2E**). A summary of
WebGestalt results for cortex-thalamus astrocyte rDEGs each age is provided in **Table S10**. To complement this
pathway analysis and facilitate exploration of rDEG functions, we queried UniProt[56] for each rDEG (see **Methods**)
to return its full protein name, GO Cellular Compartment, GO Molecular Function, and GO Biological Process
annotation(s). These annotations are included in **Table S9** for marmoset and **Table S10** for mouse. Together, these
results suggest that astrocytes are regionally specialized with varied physiological adaptations necessary to support
neuronal transmission and activity in their local environment.
221
Compared to cortex-thalamus expression differences, there were many fewer cortex-striatum rDEGs (0 in fetal
marmoset, 12 in neonate, 8 in 7-month, 9 in 14-month, 25 in 30-month, and 12 in aged). At neonate, 7- and 14-
month time points, at least half of these cortex-striatum rDEGs overlapped with cortex-thalamus rDEGs. At 14
months, these overlapped genes included *MYO16*, an unconventional myosin protein implicated in
neurodevelopment[57] (higher in striatum and thalamus); *UNC5C*, a netrin receptor family member involved in axon
guidance[58,59] (higher in cortex); *MAPK10*, a mitogen-activated protein kinase (higher in striatum and thalamus);
STXBP6, a syntaxin binding protein which is part of the SNARE complex in neurons (higher in striatum and
thalamus); *GRIK2*, a kainate-type ionotropic glutamate receptor subunit (higher in striatum and thalamus); *EYA2*, a
transcriptional coactivator and phosphatase (higher in striatum and thalamus); *DYNC1I1*, a member of the
cytoplasmic dynein 1 complex involved in intracellular transport (higher in striatum and thalamus); and *PTPRE*, a
protein tyrosine phosphatase family member involved in cell signaling with various downstream consequences
(higher in cortex). Each of these genes points to a biological process, such as glutamate sensing, phosphorylation,
and exocytosis, for which striatal and thalamic astrocytes may be differentially invested.
235
Mouse. We calculated mouse astrocyte rDEGs using the previously described metacell method on expression data
pooled across all biological replicates (see **Methods**). As with marmoset, mouse astrocyte gene expression varied
across developmental timepoints, and most astrocyte rDEGs were not shared with other cell types (**Fig. 3A-D,**
Table S11). E18.5 astrocyte cortex-thalamus rDEGs (79 total) included *Cacna2d1*, *Cntn5*, *Nrxn1*, *Creb5*, *Slco1c1*,
and *Slc6a11*, and together were enriched for biological processes including cell-cell adhesion, axon guidance, and
postsynaptic organization (**Fig. 3E(i), Table S12**). P14 astrocyte cortex-thalamus rDEGs included 26 of the rDEGs
present at E18.5 (19% of total P14 rDEGs), in addition to rDEGs that only emerged at P14. These included voltage-
gated calcium channel subunit *Cacna1a*, the glutamate-gated kainate receptor *Grik4*, the N-glycoprotein *Thsd7a*,
the cholesterol transporter *Gramd1b*, and the inward-rectifying potassium channel *Kcnj6*. Together, P14 cortex-
thalamus astrocyte rDEGs were enriched in glutamatergic synapse, hormone transport, and postsynaptic
organization pathways (**Fig. 3E(ii), Table S12**). There were 124 cortex-thalamus astrocyte rDEGs at P90, which
included many of the rDEGs present at earlier time points (17% of P90 rDEGs were present at E18.5 and 52% were
present at P14), and were enriched in neuron migration, axon guidance, calcium signaling, and cell junction
pathways (**Fig. 3E(iii), Table S12**). Compared to marmoset, mouse astrocytes had more cortex-striatum astrocyte
rDEGs throughout development, especially at P14 (13 at E18.5, 24 at P4, 63 at P14, 15 at P90, and 33 at 90
weeks). However, at most 15 of these (at P14) overlapped with cortex-thalamus rDEGs, suggesting a more distinct
transcriptional niche for mouse striatal astrocytes compared to cortex, as explored below.
253
As we did with marmoset, we assessed in mouse the degree to which regional imprinting of astrocyte gene
expression persists across development by correlating log-fold change gene expression differences between region

pairs at E18.5, P14, and P90. We found that cortex-striatum vs. cortex-thalamus fold-changes exhibited a high
degree of correlation at E18.5 (Pearson's r = 0.85), which decreased dramatically at P14 (r = 0.26) and increased
again at P90 (r = 0.40, **Fig. S7D-F(i)**). Compared to marmoset, striatum-cortex vs. striatum-thalamus rDEGs were
more positively correlated at juvenile and adult stages (P14 (r = 0.48) and P90 (r = 0.19, **Fig. S7D-F(ii)**)). Consistent
with marmoset, thalamus-striatum vs. thalamus-cortex fold-changes were highly correlated (r > 0.70) at all 3
timepoints (**Fig. S7D-F(iii)**). Together, these results suggest that while cortical and striatal astrocytes are similarly
divergent from thalamic astrocytes (likely reflecting the their distinct telencephalic-diencephalic origins), mouse
striatal astrocytes develop and maintain a unique transcriptional signature distinct from cortex and thalamus.
264
To validate the existence of these regional astrocyte populations and the differential expression of selected rDEGs
in situ, we conducted multiplexed RNA fluorescence *in situ* hybridization (FISH) using the RNAscope HiPlex or
RNAscope Multiplex Fluorescent v2 kit (Advanced Cell Diagnostics) in neonate and adult animals of both species.
We used CellProfiler 4.2.5[60,61] to quantify the fraction of astrocytes positive for each target gene in each region and
the fraction of each astrocyte nuclei covered by the probe for each target gene in each region (referred to as mean
intensity, see **Methods**). Most rDEGs followed the expected regional and developmental expression pattern in
marmoset astrocytes, including *SPARC*, which was enriched in diencephalic astrocytes (more so in adulthood),
FOXG1, which marked telencephalic astrocytes, *GFAP,* which was elevated in thalamus in adult but not neonate,
and *KCNH7*, which was a telencephalic rDEG in neonate but not adult (**Fig. S8-9**, see **Methods** and
Supplementary Note 1 for a discussion of the few rDEGs whose in situ expression differed from snRNAseq
predictions).
276
Similarly, we found that most mouse astrocyte rDEGs followed the expected regional and developmental expression
pattern in P4 and P90 mouse astrocytes, including *Clmn, Slco1c1, Csmd1*, *Sparc* (in adult mouse), and *Kcnd2*
(**Figs. S10-11,** see **Table S13** for source data and statistics). For additional validation, we analyzed the differential
expression of our selected mouse rDEGs in the Allen Mouse Brain Cell Atlas whole-brain MERSCOPE v1 dataset[1],
and found it to be largely consistent with our snRNAseq data (**Fig. S11B**, see **Methods**). This was also true for the
whole list P90 mouse rDEGs, which also showed differential expression across a wider selection of brain regions
(**Fig. S11C**).
284
To assess the extent of astrocyte intra-regional heterogeneity, we performed subclustering on cortical, striatal, and
thalamic astrocytes from all developmental time points separately for each species (**Fig. S12-13**, see **Methods**).
We found at least 4 astrocyte subclusters within each region, which primarily distinguished protoplasmic and
fibrous/interlaminar subtypes (the latter being identified by *GFAP*, *AQP4*, and/or *ID3* expression[8,23,62,63]) and
immature and mature astrocytes. In both species, the majority of astrocytes in the cortex and striatum were
protoplasmic. In the marmoset thalamus, a larger proportion of astrocytes were *GFAP+*, *AQP4+*, or *ID3+* (**Fig.**
S12C, Table S14), suggesting a higher proportion of fibrous astrocytes, consistent with the greater abundance of
white matter in this region (**Fig. S2B-C**). Nevertheless, it is unclear the extent to which the definitions of
protoplasmic, fibrous, and intralaminar apply outside of the cortex. The mouse striatum had the most intra-regional
heterogeneity, with 12 subclusters (**Fig. S13B**), in large part due to immature populations including *Top2a+* rostral
migratory stream progenitors. As in previous studies, we found that *CRYM/Crym* marks a subset of striatal
astrocytes[19] and *SPARC/Sparc* marks thalamic astrocytes in both species[18]. Several of our subclusters mapped
specifically (>70% of cells in the subcluster) to a single adult mouse Allen Brain Cell Atlas (ABCA) cluster, though
many mapped to several ABCA clusters, especially immature and mixed fibrous/protoplasmic subclusters (**Table**
S15). Taken together with our FISH data, which was obtained in gray matter regions, our subclustering analysis
suggests that most of the astrocytes in our study, and therefore likely most of the resulting rDEGs, arise from
protoplasmic or gray matter astrocytes.
302
Shared and subtype-specific predicted mechanisms of neuron-astrocyte communication
Many of the astrocyte rDEGs implicated neuron-astrocyte communication, suggesting that the regional molecular
identity of astrocytes may arise in part from customized interactions with the vast diversity of specialized neuronal
types across the mammalian brain[1,4,64]. Our previous analysis showed that rDEGs are not substantially shared

across neurons and glia (**Figs. 2-3D**), which rules out the influence of pan-cell type regional patterning. Neurons
and astrocytes communicate via myriad signaling pathways. We assessed whether neuron and astrocyte cluster
pairs sampled from the same region were over-enriched for known ligand-receptor (L-R) interactions using
CellPhoneDB[65]. To increase the specificity of our predicted L-R results, we restricted CellPhoneDB analysis to
neurons and astrocytes only (see **Methods**).
312
Across most brain regions and ages in the marmoset, we found neurexin and neuroligin (NRXN/NLGN) family
members, contactin (CNTN) family members, fibroblast growth factor and receptor (FGF/FGFR) family members,
and neural cell adhesion molecule (NCAM) family members to be the most enriched predicted neuron-astrocyte
and astrocyte-neuron L-R molecules (**Table S16**). Despite the commonality of these L-R pairs between astrocytes
and all neuronal subtypes, for each neuronal subtype in a given region, we found unique or near-unique L-R and
R-L pairs with astrocytes. For example, in the fetal marmoset thalamus, *SLT3→ROBO2* is specific to midbrain-
derived GRIK1+ thalamic inhibitory neurons and immature thalamic astrocytes, while *AFDN→EPHA7* signaling is
specific to immature astrocytes and TRN GABAergic neurons, compared to the other neuronal subtypes examined
(**Fig. 4A**). Later in development, at 14 months, many of the same neuron-astrocyte and astrocyte-neuron L-R
combinations were present, while some new pairs, such as *EFNA5→EPHB1* for thalamic astrocytes to *GRIK1+*
midbrain-derived GABAergic neurons, emerged (**Fig. 4B**, see **Table S17** for quantification of the fraction of L-R
pairs shared across 3 or more ages for all neuronal clusters within a region for both species).
325
To summarize the shared and divergent expression of predicted L-R pairs underlying neuron-astrocyte
communication across regions, we examined the overlap of these pairs for the most abundant neuronal cluster and
the most abundant astrocyte cluster in cortex, striatum, and thalamus (**Fig. 4C**). We found that many L-R pairs were
shared across regions at both GD135 and 14 months (43% and 29% of total L-R pairs, respectively), while the
thalamus (at GD135), and later striatum (at 14 months) had the most L-R pairs not shared with other regions for the
clusters examined. To exclude the possibility that the region-specificity of neuron-astrocyte L-R pairs is due solely
to neuronal heterogeneity, we performed analyses examining the magnitude and specificity of L-R pairs between
local (non-projecting) neurons and regional astrocyte populations, the overlap of neuron-astrocyte and neuron-OPC
L-R pairs within a region, and the proportion of astrocyte subtype DEGs overlapping with ligand (when source) or
receptor (when target) lists compared to neurons. We found that the region-specificity of neuron-astrocyte L-R pairs
is not solely explained by neuronal heterogeneity, reflecting a contribution of astrocyte regional heterogeneity. For
detailed results, please see the Methods section and associated notebooks in our GitHub repository. In several
cases, different members of the same family were used as region-specific neuron-astrocyte/astrocyte-neuron L-R
pairs in different regions. For example, in 14-month marmoset, *EFNA5→EPHA5* was unique to cortical astrocytes
→ cortical L2/3IT glutamatergic neurons, while *EFNA5→EPHA7* was unique to striatal astrocytes → *DRD1*+ medium
spiny neurons, and *EFNA5→EPHA6* was shared across all three regional A→N subtype pairs.
342
To assess how the expression of L-R pairs underlying neuron-astrocyte communication changes over the course
of development, we examined the concordance of L-R pairs of a single neuronal cluster (cortical glutamatergic
L2/3IT neuron) and cortical astrocyte at different developmental time points. In contrast to the expression of rDEGs,
a larger proportion (20/90, 22%) of L-R pairs (all L-R pairs meeting minimum expression criteria, including pairs
shared with other neuronal and astrocyte clusters, were included in this analysis) were shared between all time
points (**Fig. 4D**), suggesting that these putative mediators of neuron-astrocyte communication emerge early and
are maintained throughout development. However, 15/90 (17%) L-R pairs emerged at 7 months and were
maintained throughout adulthood. At later developmental time points, many more L-R pairs were shared between
ages than were unique (**Fig. 4D**). This, along with the increased proportion of rDEGs shared across later time points
(**Fig. 2C**), suggests that the expression of molecules underlying neuron-astrocyte communication stabilizes
postnatally in marmosets at some point between 0 and 7 months.
354

Overall, patterns of predicted neuron-astrocyte and astrocyte-neuron communication in mouse were similar to
marmoset, including implication of neurexin and neuroligin, contactin, fibroblast growth factor and receptor, and
neural cell adhesion molecular families (**Fig. S14A-B**, **Table S16**). As in marmoset, mouse thalamus had more
unique neuron-astrocyte L-R pairs than cortex or striatum at P4 and P90 (**Fig. S14C**). Unlike in marmoset, mice
had more age-specific L-R pairs (from cortical L2/3IT glutamatergic neurons to cortical astrocytes) at earlier time
points (particularly at P4, 17/84 or 20% of all L-R pairs unique at this time point) before stabilizing with more shared
L-R pairs at later time points. This suggests that mediators of mature neuron-astrocyte interactions emerge
relatively later in mouse (**Fig. S14D).** Only 11% of L2/3IT→astrocyte L-R pairs were shared across the lifespan.

Taken together, these results suggest that many L-R pairs potentially underlying neuron-astrocyte communication

are shared across developmental time points and regions in both species. However, more neuron-astrocyte

predicted L-R pairs emerged later in development.

366
Age-dependent refinement of astrocyte identity
In mouse, initiation of gliogenesis in the diencephalon precedes that in the telencephalon by approximately 1
gestational day (E13.5 vs E14.5[66,67]). To determine whether relative immaturity of telencephalic glia compared to
diencephalic glia could explain the robust regional expression differences we observed at each sampled time point
(**Figs. 2, 3**), we examined the developmental trajectory of astrocytes in pseudotime, a prediction of position along
a low-dimensional developmental trajectory based on RNA expression only, using Palantir[68]. Palantir recovered the
known developmental trajectory of the oligodendrocyte lineage in both species (**Fig. S15A-B**).[69] Furthermore, it
underscored the precocious myelination in the marmoset brain compared to mouse, as evidenced by a faster rate
of pseudotime progression towards maturity and a larger proportion of newly-formed and myelinating
oligodendrocytes at earlier time points in marmoset (**Fig. S15C-D**).

377
In astrocytes from both species, pseudotime analysis with separate terminal states for mature telencephalic ("AST-
TE" branch) and diencephalic ("AST-DI" branch) astrocytes revealed a transcriptional developmental trajectory
within astrocytes that aligned with actual age and annotation of mature and immature Leiden clusters (**Fig. 5A-B**,
Fig. S15E-F). Pseudotime values were slightly higher in mature diencephalic versus mature telencephalic
astrocytes, and higher in cortical astrocytes than striatal astrocytes, in both species at mature time points,
suggesting arrival at distinct terminal states, though these need not be more or less mature than one another (**Fig.
5A-B, Fig. S15E-F**). The rate of maturation (that is, distribution of pseudotime values during development relative
to those in adulthood) also differed slightly across regions. To identify genes potentially driving pseudotime
transitions, we used Mellon[70] to calculate gene change scores, a measure of expression change in regions of low
cell-state density (see **Methods**), for each pseudotime trajectory branch. Many of the top 25 change-scoring genes
in both branches were cortex-thalamus rDEGs (**Fig. S15C-D**, **Table S18**), suggesting that the expression of region-
enriched genes is correlated with, and potentially drives, astrocyte maturation.

390
Next, for each region we binned marmoset astrocytes by pseudotime quintile in the appropriate trajectory branch
rather than by actual age and recomputed rDEGs. As with rDEGs grouped by actual age, the number of rDEGs
increased from pseudotime bin 1 (PT1) to pseudotime bin 5 (PT5): 54 rDEGs at PT1, 108 at PT2, 117 at PT3, 177
at PT4, and 181 at PT5. We found that matching by predicted maturational stage largely recapitulated the original
rDEGs calculated from actual age: 56% of PT1 cortex-thalamus rDEGs overlapped with GD135 rDEGs, 81% of
PT2 rDEGs overlapped with neonate rDEGs, 39% of PT3 rDEGs overlapped with 7-month rDEGs, 85% of PT4
rDEGs overlapped with 14-month rDEGs, 81% of PT5 rDEGs overlapped with adult rDEGs, and 85% of PT5 rDEGs
overlapped with aged rDEGs. This suggests that regional imprinting of astrocytes is not simply driven by relative
differences in the birth timing of cells across the different brain structures.

400
We next sought to determine the sequence of molecular changes that unfold in astrocytes within a given region
over time. We calculated **a**ge **d**ifferentially **e**xpressed **g**enes (aDEGs) within each brain region from metacells of
each age (see **Methods**). In each brain region, there were over 100 unique aDEGs (unique after pooling pairwise

age combinations). In marmoset, the largest fraction of aDEGs distinguished GD135/neonate from 7-month and
older astrocytes (**Fig. 5C**). aDEGs enriched in the 30-month dataset could conceivably arise from the different
sample preparation and reference genome used in our previous study[7]; for this reason we used the 14-month time
point to further assess age-related changes across regions.
408
Examining the overlap of marmoset astrocyte GD135 vs. 14-month aDEGs between brain regions (409 aDEGs in
total), we found that ~19% were shared between cortex, striatum, and thalamus (**Fig. 5D**). The striatum had a
modest number of GD135 vs. 14-month aDEGs not shared with other regions (21/409), while the cortex had 3-fold
more (63/409), and the thalamus had the most (156/409), as expected given the stark regional heterogeneity
between telencephalon and diencephalon (**Fig. 2**). Additionally, we found that few to no GD135 vs. 14-month
aDEGs were shared between astrocytes, OPCs, and excitatory neurons or astrocytes, OPCs, and GABAergic
neurons in the cortex (**Fig. 5E**).
416
We found similar results in mice, where we calculated P4 vs. P90 rDEGs, as mouse E18.5 astrocytes were
transcriptionally immature relative to marmoset GD135 astrocytes and GD135-P4 timepoints appear to have better
correspondence, as discussed in the next section). One notable difference from marmoset was that early adolescent
(P14) astrocytes in mice expressed many aDEGs shared with embryonic and neonate timepoints, particularly in
striatum (**Fig. S16A**). As in marmoset, thalamic astrocytes had more unique aDEGs than their cortical and striatal
counterparts (**Fig. S16B**), and most astrocyte aDEGs were not shared with other radial glia-derived cell types (**Fig.
S16C**).
424
We found very few (3 or less) astrocyte aDEGs that were cell type-agnostic and region-specific (i.e., that were also
aDEGs in neurons and OPCs for a given brain region, see **Methods**). In contrast, there were 74 (marmoset GD135
vs. 14-month) and 56 (mouse P4 vs. P90) aDEGs that were astrocyte-specific and region-agnostic, reflecting more
universal aspects of astrocyte transcriptional maturation, regardless of brain region. We found a similar number of
astrocyte-specific, region-specific aDEGs (20 in striatum, 51 in cortex, and 135 in thalamus for GD135 vs. 14-month
marmoset), which reflect the brain region's influence on the maturation of astrocytes *only* in a given brain region. In
both species, the developmental pattern of selected astrocyte-specific, cortex-specific aDEGs was similar but not
identical in the striatum, and more dissimilar with the thalamus (**Fig. 5F**, **Fig. S16D**).
433
Conservation and divergence of astrocyte patterning in mouse and marmoset
Hundreds of differentially expressed genes distinguish adult human and mouse astrocytes[71], and engrafting human
glial progenitors into mouse brain results in mature astrocytes that retain certain human-specific astrocyte
characteristics[72]. This suggests that aspects of an astrocyte's developmental program are cell intrinsic and are
shaped by its species-specific genomic features. We therefore aimed to compare transcriptional signatures of
telencephalic and diencephalic regional astrocyte populations between marmoset and mouse. We integrated a
randomly downsampled subset (100,000 nuclei each) of mouse and marmoset nuclei (all cell types included). To
do so, we used 547 highly variable 1:1 ortholog genes selected from top differentially expressed genes of
superclusters (related groups of Leiden clusters) shared across species and employed the semi-supervised
variational auto-encoder scANVI[73] (see **Methods**). The resulting integrated UMAP plot showed broad conservation
of superclusters between mouse and marmoset, despite differences in cell type proportions across development
(**Fig. S17A-B**). Another method called SATURN[74], which avoids 1:1 mapping of genes, had largely concordant
cross-species integration results (**Fig. S17C-D**).
447
Species-integrated astrocytes partitioned into three superclusters that segregated by developmental stage and by
brain structure (diencephalon vs telencephalon) (**Fig. 6A**). This indicates that at the level of broad cephalic domains,
region patterning is conserved between the two species. Mature telencephalic astrocytes showed better species
integration than diencephalic or immature astrocytes, and immature mouse astrocytes composed a distinct cluster
(**Fig. 6A**). This finding implies that at birth, mouse astrocyte maturity lags behind that of marmoset, in line with our
findings about oligodendrocyte maturation (**Fig. S13A-B, G-H**). Interestingly, marmoset astrocyte aDEGs had more
discrete expression boundaries across time (**Fig. 5C**), while mouse astrocyte aDEGs had more continuous temporal

expression, especially in the striatum (**Fig. S16**). Additionally, marmoset rDEGs shared across ages were largely
divided into younger (GD135, neonate) and older (7 months and older) groups (**Fig. 2C**). In contrast, temporally
overlapping mouse rDEGs were more evenly distributed across individual ages and smaller groups of ages (**Fig.
3C**), suggesting that developmental changes occur more slowly over the sampled timepoints in mouse.
459
In mouse but not marmoset, we observed immature astrocyte clusters composed of nuclei from all time points,
suggesting continued generation of new astrocytes throughout the lifespan. These included the *Top2a+* immature
astrocyte population seen in the neurogenic subventricular zone throughout the lifespan (**Fig. S18**), which forms
part of the rostral migratory stream.[75] All marmoset astrocytes and cortical and thalamic mouse astrocytes exhibited
separate embryonic and neonatal subclusters from early adolescent and older cells (**Fig. S12-13A, C**). In contrast,
in the mouse striatum, there were several more immature clusters (9 total), including some composed of astrocytes
from mature timepoints (**Fig. S13B**, "Str_Ast2", "Str_Ast6", and "Str_Ast12"). However, the relative immaturity of
mouse astrocytes after adolescence is not driven solely by the persistence of *Top2a+* cells in the SVZ, as we found
an immature population of *Top2a-* striatal astrocytes in mouse but not marmoset that may reside outside of the SVZ
(**Fig. S13B,** portions of "Str_Ast2" and "Str_Ast12"). We note that the presence of this immature cluster in the
striatum does not suggest that all mouse astrocytes are less mature than their marmoset counterparts in
adolescence and adulthood.
472
We next tested whether genes that best distinguished astrocytes from a given brain region in one species were
more likely than chance to be rDEGs in the other species. Focusing on astrocyte cortex-thalamus rDEGs at each
developmental time point, we found that the majority of rDEGs were not shared across species, and that the
proportion of shared rDEGs decreased only slightly from fetal to early adolescence time points, from ~14-16% to
~13-14% in both species (**Fig. 6B**, full list of species-overlapping and species-unique cortex-thalamus astrocyte
rDEGs at each developmental time point in **Table S21**). The proportion of overlapping rDEGs was not greater than
chance (see **Methods**, p-value from a Fisher's exact test > 0.05 at all developmental time points).
480
Similarly, the majority of astrocyte aDEGs in cortex and thalamus were not shared between species. 51 cortical
aDEGs (22% of mouse cortical P4-P90 aDEGs and 25% of marmoset GD135-14-month aDEGs) and 61 thalamic
aDEGs (23% of mouse thalamic P4-P90 aDEGs and 21% of marmoset thalamic GD135-14-month aDEGs) were
shared between species (**Fig. 6C**, p-value from a Fisher's exact test on the proportion overlapped = 0.025 for
cortex and 0.454 for marmoset). Lists of species-shared and species unique aDEGs by region are provided in Table
S22.
487
Next, we directly tested for differential expression of 1:1 orthologs between species within shared superclusters.
We calculated **s**pecies **d**ifferentially **e**xpressed **g**enes (sDEGs) based on 1:1 orthologs between species within each
integrated supercluster using our metacell method (see **Methods**). We found hundreds of sDEGs in both
telencephalic (464 total) and diencephalic (579 total) astrocytes whose expression could clearly distinguish between
marmoset- and mouse-derived populations (**Fig. 6D**). Astrocyte sDEGs encoded both cytosolic and membrane-
bound proteins with varied cellular functions (**Table S24**). For example, telencephalic astrocyte sDEGs higher in
marmoset included *NALCN/Nalcn*, a non-selected sodium leak channel; the RNA-binding protein *RBFOX2/Rbfox2*;
KCNT2/*Kcnt2*, a sodium-activated potassium channel subunit; *FABP7/Fabp7*, a fatty acid binding protein with
established roles in neurogenesis; and *DNM3/Dnm3*, a multi-domain GTPase involved in membrane remodeling.
Even this short list of genes suggests that important cellular functions such as ion buffering, RNA processing, fatty
acid binding, and membrane remodeling may differ between astrocytes of different species. Furthermore, 63/286
telencephalic and 71/395 diencephalic sDEGs were SFARI[76] 3.0 Autism Spectrum Disorder (ASD)-related genes
(see **Methods**). These overlaps are significantly higher than chance (assuming 20,000 protein-coding genes in the
human genome, p-values from Fisher's exact tests $< 10^{-15}$ for both telencephalic and diencephalic astrocyte
sDEGs). Complete lists of sDEGs for each supercluster analyzed, and GO annotations for Cellular Compartment,
Biological Process, and Molecular Function are provided in **Table S23.**
504

We validated the differential expression of 2 astrocyte sDEGs, *NRG3/Nrg3* (present in neurons in both species but
higher in marmoset astrocytes) and *LAMA2/Lama2* (higher in mouse astrocytes) *in situ* using RNAscope (**Fig. 6E**).
50% of the genes that distinguish diencephalic astrocytes between species were shared with telencephalic
astrocytes (**Table S24**). These telencephalic-diencephalic astrocyte shared sDEGs made up a larger fraction (62%)
of telencephalic astrocyte sDEGs. This suggests that evolution has acted on the astrocyte class as a whole, while
also shaping divergent regional astrocyte programs between species. Additionally, as with astrocyte rDEGs and
aDEGs, we found that most astrocyte sDEGs were not shared with other cell types (superclusters), including OPCs,
cortical MGE-derived *PVALB*+ interneurons, GABAergic TRN neurons, striatal MSNs, and microglia (**Fig. 6F**). This
result underscores that evolutionary divergence of a cell type's transcriptome unfolds at different rates across cell
types[62,77,78]. Taken together, these findings support both conservation and divergence of postnatal astrocyte
regional specialization in mouse and marmoset.
516
Astrocytes have regionally divergent morphology and protein expression
Many of the genes we found to vary in astrocytes by region, age, and species implicate processes involved in
morphological specification. Indeed, astrocyte morphology, which is highly ramified and complex, is essential for
their specialized functions: end feet contact blood vessels to help form the blood-brain barrier and shuttle water and
nutrients, while terminal processes closely appose synapses to uptake ions and neurotransmitters[35] and regulate
synapse development and function[79]. Because many of these morphological features exist at the sub-micron scale,
conventional light microscopy is not sufficient to visualize the full morphological complexity of astrocytes[36]. We
wondered whether nanoscale astrocyte morphology might also be regionally specialized between gray matter
regions in cortex, striatum, and thalamus, as recently demonstrated for several CNS regions using diffraction-limited
approaches[25]. Thus, we used expansion revealing (ExR), a new variant of protein decrowding expansion
microscopy[37], to visualize astrocyte morphology with enhanced resolution and compare morphological properties
in the PFC, striatum, and thalamus, as these regional populations represented the major molecular subpopulations
in our snRNAseq data (**Fig. 2, 3**).
530
We used a viral approach to label astrocytes for expansion (**Fig. 7A-B**, see **Methods**) and created 3D binary
segmentations to quantitatively assess morphological differences across regions (**Fig. 7D**, Movies **S19-36**). We
calculated the volume, surface area, equivalent diameter (measures of size), surface area to volume ratio (S:V, a
measure of shape, inversely proportional to size), aspect ratio (a measure of elongation), fractal dimension (FD)[80]
(a measure of complexity and self-similarity), and branching complexity via Sholl analysis[81], most of which have
been used to characterize astrocyte morphology in prior studies[36,82].
537
After correcting for multiple comparisons across the 6 univariate measures, we found significant differences in size
(volume, surface area, and equivalent diameter), shape (surface area to volume ratio), and morphological
complexity (FD) between astrocytes from different brain regions, particularly between cortical/striatal and thalamic
astrocytes (**Fig. 7E,** n = 52-60 astrocytes from 3 female and 5 male mice for each region, with statistical significance
determined using a linear mixed effects model with "animal" as the random effect group variable, see **Table S25**).
Specifically, thalamic astrocytes were smaller and less complex compared with cortical and striatal astrocytes, while
striatal astrocytes had smaller surface area and surface area to volume ratios compared to cortical astrocytes. To
exclude the possibility that proximity to under-digested blood vessels or incomplete capture in the axial dimension
impacted these differences, we repeated this analysis on a subset of astrocytes meeting additional criteria and
found largely similar results (**Fig. S19A**, see **Methods**). Similarly, Sholl analysis revealed fewer intersections at
radii larger than ~25 μm for thalamic compared to cortical and striatal astrocytes (**Fig. 7E(vi)**). Taken together,
these results suggest that thalamic astrocytes are smaller and less morphologically complex than their cortical and
striatal counterparts, in line with prior work using conventional microscopy[25].
551
We next probed rDEG protein product expression level and localization in astrocytes at the nanoscale using ExR.
We processed tissue from 2 of the virally-labeled adult C57 Bl/6J mice used for morphology analysis and proceeded
with staining for Glast (encoded by the telencephalic rDEG *Slc1a3*) and Gat3 (encoded by the thalamic rDEG
Slc6a11) alongside GFP (astrocyte processes) and Cav2.1 (a presynaptic protein) in gels prepared from PFC,

striatum, and thalamus. We found that the expression of Glast and Gat3 at the protein level in GFP-labelled
astrocytes agreed with snRNAseq predictions: Glast expression was highest in the cortex, lower in the striatum,
and lowest in the thalamus, while Gat3 expression was higher in the thalamus than cortex or striatum (**Fig. 7F**, **Fig.**
S19B). Both proteins localized to astrocyte processes near synapses. Taken together, these results support the
differential expression of rDEG protein products between telencephalic and diencephalic astrocytes, and reveal the
localization of rDEG protein products on and near astrocyte processes, in close proximity to neuronal synapses.
562
Discussion
Astrocytes are a ubiquitous, versatile brain cell type with increasingly appreciated roles in health and disease. While
their regional molecular heterogeneity has been evident for some time[83,84], the source of this regional heterogeneity,
in particular, the relative contributions of embryonic patterning versus response to environmental cues after birth, is
not well understood[85]. To help bridge this knowledge gap, we generated a unified, multi-region, postnatal
developmental snRNAseq atlas of mouse and marmoset brain cells. Because our dataset contains all brain cell
types, we anticipate this atlas will be a valuable resource for the field. As such, we have made both raw and
processed data publicly available on NeMO and the Broad Single Cell Portal, respectively (see **Data and Code**
Availability). The latter is useful for exploring cell type clusters and querying the expression pattern of genes of
interest across ages and regions, and does not require coding expertise.
573
We found that astrocytes were regionally patterned before birth in both species, a discovery that was not
unexpected given the prevalence of homeobox patterning genes among astrocyte regionally differentially expressed
genes[8], evidence from a recent study showing regionally patterned glioblasts in the first-trimester human brain[6],
and older lineage tracing showing regional allocation of astrocytes based on the region of their originating radial
glia[27]. Less predictably, we discovered dramatic changes in astrocyte regional identity between birth and early
adolescence, in line with their maturation during this period. This period also coincides with peak synaptogenesis,
pruning, and myelination[86], consistent with the notion that astrocyte specialization depends on the activity of
neighboring cells[87].
582
The functions of embryonically-patterned and postnatally-acquired astrocyte rDEGs were varied, but implicate
known astrocyte processes, including supporting synaptic transmission, ion transport, neurotransmitter uptake, cell-
cell adhesion, and morphological specification. The function of some rDEGs, including *SLC6A11* (GAT-3) and
SPARC, has been studied in astrocytes, and shown to be important in modulating the effects of brain injury[88] and
controlling synaptogenesis[53], respectively. We anticipate future mechanistic studies of other astrocyte rDEGs will
reveal yet more essential functions.
589
We found that neuron-astrocyte and astrocyte-neuron predicted ligand-receptor pairs, many of which were
specialized for distinct neuronal subtypes, were upregulated during postnatal development into adulthood, again
supporting the hypothesis that astrocytes specialize in postnatal development to meet the needs of local neurons.
Despite the striking regional heterogeneity of astrocytes, many predicted neuron-astrocyte ligand-receptor pairs
were shared across regions. Even those not shared across regions were functionally similar, suggesting neurons
and astrocytes have developed a common language of molecular communication across the forebrain. Indeed,
some of our rDEGs were members of the same protein family or functional class, pointing to variations on a common
theme of neuron-astrocyte crosstalk across brain regions. Many of the top predicted neuron-astrocyte ligand-
receptor pairs, such as neurexins and neuroligins, are more traditionally associated with neuron-neuron contact at
the synapse[89]. However, adhesion molecules such as ephrins, neurexins/neuroligins, and NrCAMs have been
shown to play important roles in neuron-astrocyte communication[49].
601
In both species, we found hundreds of age differentially expressed genes (aDEGs), many of which astrocyte-
specific but region-agnostic, some of which were astrocyte-specific and region-specific, and very few of which were
cell type-agnostic but region-specific. The thalamus had the most unique astrocyte developmental gene expression
signature of the three brain regions, suggesting that thalamic astrocytes undergo distinct developmental changes
from their telencephalic counterparts. Our astrocyte-specific, region-agnostic aDEGs can be interpreted as a core

forebrain astrocyte developmental program, and were more likely to be shared across the species. For example, in
marmoset, this included *NTRK2*, which encodes the BDNF receptor TrkB, the short isoform of which has been
shown to be essential for astrocyte morphogenesis[90]. Perhaps unsurprisingly, several of our region-specific aDEGs
were also rDEGs, and/or had a high degree of functional overlap with rDEGs. While pseudotime approaches have
limitations[91] and may not fully capture how maturational states differ across brain regions, they can provide
information about the progression of change along a trajectory that is correlated with actual age. For this reason,
we used pseudotime to compare relative maturation differences across regions. This analysis suggested that
intrinsic maturation rates are relatively low drivers of regional differences in gene expression.

In both species, most astrocyte rDEGs, aDEGs, and sDEGs were not shared with OPCs or neurons, suggesting
that astrocyte region- and age- specializations are unique, rather than general to all radial-glia derived cell types in
the same region, developmental time point, or species. This suggests either that regional gene expression
signatures change throughout neuro- and glio-genesis, or that the downstream transcriptional effects of this early
regional patterning depend on the daughter cell's fate. Evidence for both exists in the cortex[92]. Why neurons and
astrocytes exhibit stark regional patterning in adulthood, albeit in different ways, while the oligodendrocyte lineage
does not, is an outstanding question for future study.

The present study characterized two mammalian neuroscience model species, mouse and marmoset. While mice
and humans have a high degree of genetic conservation[93], mice have certain limitations as a model for studying
the human brain including lack of a well-developed prefrontal cortex and complex social behaviors, and poor visual
acuity. In light of these limitations, non-human primates, with whom we share much closer genetic ancestors, are
considered as more translationally-relevant models of brain function and dysfunction. The common marmoset has
become an increasingly popular non-human primate model in neuroscience studies due to its faster generation time
for genetic engineering, shorter lifespan than other larger primates for developmental and late-onset disease
studies, and complex social behaviors[94].

Our data suggest that the development of astrocyte regional heterogeneity, marked by embryonic regional
patterning along cephalic boundaries followed by dramatic postnatal specialization, is broadly conserved between
mouse and marmoset. However, the expression of many rDEGs and aDEGs differs across species, and we
identified hundreds of species differentially expressed genes within both telencephalic and diencephalic astrocytes
(**Fig. 5**). These sDEGs encode proteins involved in key cellular functions which may have undergone evolutionary
selection, and a significant portion have been associated with ASD. Future studies exploring the function of these
sDEGs within astrocytes may reveal how primate astrocytes have evolved to suit the unique anatomy and
physiology of the primate brain. Taken together, these findings suggest that each species may have evolved by
recruiting different sets of genes that facilitate postnatal regional specialization of astrocytes. We found that many
cell types in the marmoset brain are transcriptionally more mature at time of birth than their mouse counterparts, in
line with previously documented precocious development in early postnatal marmosets[95]. This species divergence
in transcriptional maturity at time of birth suggests that researchers should use caution when comparing early
postnatal time points between rodents and NHPs, especially in light of differences in developmental tempo between
species[96].

We used expansion revealing (ExR) combined with a viral astrocyte labeling approach to circumvent the diffraction
limit of light microscopy and the limitations of immunostaining, respectively, to visualize astrocyte processes with
enhanced resolution. Our quantitative profiling of astrocyte morphology in the mouse brain shows that astrocyte
size, shape and complexity do vary across brain regions, and are most distinct between thalamus and cortex. Prior
studies have also found appreciable morphological differences in mouse astrocytes across brain regions[19,25,97].
Although other higher resolution approaches such as electron microscopy may reveal additional differences in
aspects of astrocyte morphology, we demonstrate that expansion microscopy, particularly ExR, offers an
inexpensive and accessible alternative to other super-resolution approaches for characterizing astrocyte
morphology with enhanced resolution. We anticipate that this approach could also be used to study morphological
changes in astrocytes after manipulation and/or in disease contexts.

658
There are several notable limitations to the current study, only some of which we discuss here. The first is the
reliance on 10x Chromium snRNAseq, which is subject to dropout and 3' bias, and produces short reads that cannot
be used to map splice variants or many single-nucleotide polymorphisms that may differ between cell types and
species. Additionally, the use of nuclei instead of whole cells prevents our detection of RNAs in the cytoplasm,
including those locally translated in distant processes of both neurons[98] and astrocytes[99], which are likely relevant
in establishing cell type and state identity. However, others have found similar cell type discrimination capabilities
for single-cell and single-nucleus RNAseq in the mouse cortex, despite lower RNA content (20-50% of total cellular
mRNA) in single nuclei.[100] The second significant limitation is the relatively small sample size, especially for
marmosets due to practical limitations including cost, which limits our ability to compare between sexes.
668
Third, we relied in part on pathway analysis to summarize patterns and deduce functional implications arising from
sets of rDEGs. Our use of WebGestalt did not incorporate any fold-change or p-value information for genes, treating
each DEG equally regardless of its differential expression level, which may skew results. Furthermore, pathway
analysis is only as accurate as the underlying annotations, which can be lacking for glial biology. Finally, many
genes are involved in several pathways. For these reasons, we encourage interested readers to directly examine
our DEG lists provided in the **Supplementary Tables**.
675
Fourth, there are significant challenges in integrating snRNAseq data across species. Even before data analysis,
read alignment will differ across species, varying with the quality and content of reference genome annotation (for
example, we used an optimized version of the mm10 reference genome[101], for which no analogous version exists
for the marmoset). During data analysis, approaches requiring direct merging of the cell x gene count matrices (as
in **Fig. 6**) results in the loss of biological information, because only roughly 50-60% of total genes detected in either
species were mapped as one-to-one orthologs. For this reason, we also integrated the data across species with an
orthogonal approach that does not rely on one-to-one ortholog mapping (**Fig. S17C**). Both approaches are further
limited by the need for *a priori* cell type annotation, which may bias towards or against integration of shared and
unshared superclusters, respectively. Therefore, we are most confident in sDEGs, which are all one-to-one
orthologs and calculated within shared superclusters, as a measure of species divergence.
686
Fifth, our approach for labeling astrocytes for morphological analysis relies on viral infection and manual
identification of astrocytes meeting a minimum brightness level for imaging and segmentation, which may be biased
towards a certain astrocyte subtype. Finally, we relied on RNAscope HiPlex to assess rDEG mRNA levels *in situ*.
Any multiplexing technique that involves repeated stripping and restaining suffers from some level of reduced
fluorescence intensity in later rounds, as well as some amount of registration error. Therefore, any researcher
interested in following up on a gene or protein of interest that was imaged in a later round should perform additional
confirmatory studies with a single round of imaging.
694
Taken together, our data support a model of astrocyte regional specialization that includes both embryonic
patterning and postnatal specialization in response to local environmental cues, including synapse formation and
neuronal activity, as has been previously suggested[25,84]. To determine whether or not early transcriptional patterning
is required for proper postnatal astrocyte specialization for such a role, a cross-region astrocyte heterotopic
transplant would be illuminating. That is, would a thalamic-born astrocyte be able to acquire the transcriptional and
morphological profile of a cortical astrocyte if transplanted in early postnatal life? Evidence from such an experiment
in septal astrocyte populations suggests the answer is yes[31]. Alternatively, but not mutually exclusively, early
developmental regional patterning may "prime" astrocytes to receive and react appropriately to the signals they
receive in their local niches later in development, as a recent study has shown in the context of GABA-induced
morphogenesis[102]. We anticipate the current study will be a useful starting point for hypotheses such as these.
705
Data and Code Availability
Raw data (sequencing reads in fastq format) for all 10x Chromium snRNAseq samples and CellBender-cleaned
aligned counts matrices (in .h5 format) are publicly available for download on the Neuroscience Multi-omic Data

Archive (NeMO) at: https://data.nemoarchive.org/biccn/grant/u01_feng/feng/transcriptome/sncell/10x_v3.1/. Pre-
processed, clustered, and annotated data (in .h5ad format) is available for download, exploration, and gene search
on the Broad Single Cell Portal at: https://singlecell.broadinstitute.org/single_cell/study/SCP2719/a-multi-region-
transcriptomic-atlas-of-developmental-cell-type-diversity-in-mouse-brain (mouse) and
https://singlecell.broadinstitute.org/single_cell/study/SCP2706/a-multi-region-transcriptomic-atlas-of-
developmental-cell-type-diversity-in-marmoset-brain#study-summary (marmoset). Registered RNAscope
HiPlex/v2 FISH image stacks, raw and background-subtracted expansion microscopy image volumes and binarized
3D tracings (both in .tif format) are available for download on BossDB at https://bossdb.org/project/schroeder2025.
Custom scripts used to analyze data are available at https://github.com/Feng-Lab-MIT/AstrocyteHeterogeneity.
718

Author Contributions
M.E.S., K.X.L, G.F., and F.M.K. conceptualized the project. M.E.S. harvested and microdissected mouse brain
tissue and microdissected marmoset brain tissue with assistance from K.X.L. and D.M.M. Q.Z. oversaw marmoset
breeding and donor selection and performed brain tissue harvest. M.E.S., L.M., and K.X.L. conducted single-
nucleus RNA sequencing experiments. M.E.S., D.M.M., and K.X.L. analyzed the single-nucleus RNA sequencing
data with guidance from F.M.K. M.E.S. and L.M. performed mouse colony management. L.M. performed stereotaxic
surgery for AAV injection in adult mice and performed transcardial perfusions alongside D.M.M. E.Y. and J.K.
optimized collagenase treatment for blood vessel preservation and made ExR gels. M.E.S. dissected, pre-stained,
stained and imaged low-expansion ExR gels and analyzed images. J.K. stained and imaged high-expansion ExR
gels. H.Z. and K.M.L. imaged marmoset tissue on the TissueFAXS microscope and provided guidance with
marmoset tissue sectioning and staining. F.M.K., G.F., and E.S.B. supervised the project. M.E.S., F.M.K., E.S.B.,
and G.F. acquired funding. M.E.S. wrote and revised the manuscript with D.M.M., F.M.K., G.F., and E.S.B. An
author contributions matrix, inspired by a recent *Nature* news article (https://www.nature.com/nature-
index/news/researchers-embracing-visual-tools-contribution-matrix-give-fair-credit-authors-scientific-papers) is
included below.
734
735

	MES	DMM	LRM	JK	KXL	EY	LM	QZ	HZ	KL	ESB	FMK	GF
Conceptualization	█				█						█	█	█
Investigation	█	█	█	█	░	█	░	░	░	░		░	
Data curation	█	█											
Analysis	█	█			░							█	
Supervision	█										█	█	█
Funding acquisition	█										█	█	█
Project administration	█							░					
Writing - original draft	█	█								░		█	█
Writing - revisions	█	█					░					█	█
MAJOR													
SUPPORT													

736
Acknowledgements

Pictograms were generated using BioRender.com. MES was supported by the MathWorks Science Fellowship at
MIT, the Collamore-Rogers Fellowship at MIT, the NSF Graduate Research Fellowship Program #1745302, and
NIH 1F31MH133329-01. GF was supported by the National Institute of Mental Health and BRAIN Initiative
(U01MH114819), Hock E. Tan and K. Lisa Yang Center for Autism Research of the Yang Tan Collective at MIT,
the Poitras Center for Psychiatric Disorders Research at MIT, Stanley Center for Psychiatric Research at Broad
Institute of MIT and Harvard. This work was supported by Brain Initiative grant UM1MH130981 to GF and FMK.
ESB was supported by HHMI, Lisa Yang, NIH R01AG087374, NIH 1R01EB024261, Good Ventures, Tom Stocky
and Avni Shah, Kathleen Octavio, NIH 1R01AG070831, European Research Council (ERC) SYNERGY Grant No
835102. We thank Eric Nyase, Andrew Harrahill, and In-Hye Kang for AAV packaging and preparation, Ken Chan
and Ben Deverman for the BI103 plasmid and tissue from an injected animal co-stained with Sox9 to help assess
expression in astrocytes, Michael Debarbardine for generating mitochondrial gene annotations for
mCalJa1.2.Pat.X, Yanay Rosen for generating the marmoset protein embedding for SATURN, Menglong Zeng for
guidance particularly on molecular cloning and for sharing many reagents, Morgan Fleishman for animal colony
maintenance and support, Victoria Beja-Glasser for guidance and the gift of several wild-type mice, Ryan Kast for
guidance particularly on RNA FISH and sharing many reagents, Nik Jorstad for guidance on analysis, Samia Silva
de Castro, Jitendra Sharma, and Yefei Chen for performing necropsies and harvesting marmoset tissue, Alyssa
Lutservitz and Xian Adikonis for tips on the 10x snRNAseq protocol, 10x Genomics Technical Support for answering
many questions and providing replacement reagents when required, Jennifer Shih and Mriganka Sur for the gift of
Aldh1l1-Cre mice, Lizzy Joo for the Cre genotyping primer sequences (originally from the Brueckner Lab at Yale),
and Stephen Yu for assistance with marmoset PCR-based sex genotyping. We thank all members of the Feng and
Boyden labs for insight and suggestions throughout the project. We gratefully acknowledge the MIT BioMicroCenter,
the Broad Institute Genomics Platform, and MIT Department of Comparative Medicine for their support and
assistance of this work.
761
762 **Competing Interests**
763 J.K. and E.S.B. are co-inventors on a patent application for ExR (US 2020/0271556 A1). E.S.B. co-founded a
764 company to explore clinical applications of expansion microscopy technologies. The other authors declare no
765 competing interests.

    Figures and Legends

Figure 1. A multi-region transcriptomic atlas of brain cell diversity across postnatal development in marmoset and mouse. A, Cross-development, cross-region sampling strategy in marmoset (top row) and mouse (bottom row). **(i)** Developmental time points profiled (some approximate, see **Methods** and **Table S1**) GD, gestational day; E, embryonic day; P, postnatal day. **(ii)** Brain regions profiled, including prefrontal cortex (PFC, red dashed boxes), motor cortex (MO, blue dashed boxes), striatum (yellow dashed boxes), and thalamus (green dashed boxes), shown in either sagittal (left) or coronal slices (right, for subcortical regions only). Schematics generated using BioRender.com. **B-C**, Integrated UMAP embedding of marmoset (**B**, 881,832 nuclei) or mouse (**C**, 597,668 nuclei) nuclei from PFC, MO, striatum, and thalamus across all developmental time points assayed and a

776   randomly downsampled portion of adult nuclei from our previous study[7] colored by **(i)** assigned cell type **(ii)**
777   dissected brain region, or **(iii)** developmental time point. Legend for **B-C(i)** is shared.
778
779

Figure 2. Developmental changes and cell-type specificity of astrocyte regional heterogeneity over postnatal development in the marmoset. A, Integrated UMAP embeddings of 103,009 marmoset astrocytes colored by **(i)** assigned brain region (one-hot color encoded in **(iv)**), **(ii)** developmental time point (one-hot color encoded in **(v)**, and **(iii)** Leiden cluster assignment. **B,** Expression heatmaps (rows are cells, columns are genes) of regional differentially expressed genes (rDEGs) between astrocytes from cortex, striatum, and thalamus at **(i)** GD135, **(ii)** 7 month, and **(iii)** 30 month-old marmosets in log counts per million (logCPM). The raster plots beneath each heatmap indicate the time point (GD135, 7 months, or 30 months) and region(s) of upregulation (cortex, striatum, and/or thalamus) for each rDEG. Genes are ordered first by the time point at which they are an rDEG,

then by the regions in which they are most highly expressed, and are plotted more than once if they are present at
more than one time point. The same set of genes are plotted in the same order in **(i-iii)**. Striatal astrocytes are
ordered by subtype identity (see **Methods**). **C,** UpSet plot showing the number of unique and overlapping cortex-
thalamus rDEGs between developmental time points. The colored dots below each vertical bar indicate which
age(s) share that set of rDEGs, while the colored horizontal bars indicate the total number of cortex-thalamus rDEGs
for each age. Overlap categories with 0 rDEGs are not shown. **D,** UpSet plot (as in **(C)**) showing the number of
overlapping cortex-thalamus rDEGs between OPCs (light blue), astrocytes (dark blue), excitatory neurons (pink),
and inhibitory neurons (gray) for **(i)** neonate and **(ii)** adult marmoset. **E,** Gene ontology (GO) and pathway analysis
on cortex-thalamus (enriched in either region) astrocyte rDEGs via WebGestalt 2024 in **(i)** GD135, **(ii)** 7 month, and
(iii) 30 month marmoset astrocytes. Lollipop plots show the enrichment ratio of GO Biological Process and KEGG
pathways from an over-representation analysis with weighted set cover redundancy reduction, with tip size inversely
proportional to the false discovery rate (FDR).
801
802

Figure 3. Developmental changes and cell-type specificity of astrocyte regional heterogeneity over postnatal development in the mouse. A, Integrated UMAP embeddings of 68,485 mouse astrocytes colored by **(i)** brain region (one-hot color encoded in **(iv)**), **(ii)** developmental time point (one-hot color encoded in **(v)**), and **(iii)** Leiden cluster assignment. **B,** Expression heatmap (rows are cells, columns are genes) of regional differentially expressed genes (rDEGs) between astrocytes from cortex, striatum, and thalamus at **(i)** E18.5, **(ii)** P14, and **(iii)** P90 mice in log counts per million (logCPM). The raster plots beneath each heatmap indicate the time point (GD135, 7 months, or 30 months) and region(s) of upregulation (cortex, striatum, and/or thalamus) for each rDEG. Genes are ordered first by the time point at which they are an rDEG, then by the regions in which they are most highly

expressed, and are plotted more than once if they are present at more than one time point. The same set of genes
are plotted in the same order in **(i-iii)**. Striatal astrocytes are ordered by subtype identity (see **Methods**). **C,** UpSet
plot showing the number of overlapping cortex-thalamus rDEGs between developmental time points. The colored
dots below each vertical bar indicate which age(s) share that set of rDEGs, while the colored horizontal bars indicate
the total number of cortex-thalamus rDEGs for each age. Overlap categories with 0 rDEGs are not shown. **D,** UpSet
plot (as in **(C)**) showing the number of overlapping cortex-thalamus rDEGs between OPCs (light blue), astrocytes
(dark blue), inhibitory neurons (gray), and excitatory neurons (pink) for **(i)** P4 and **(ii)** P90 mouse. Overlap categories
with 0 rDEGs are not shown. **E,** Gene ontology (GO) and pathway analysis on cortex-thalamus (enriched in either
region) astrocyte rDEGs via WebGestalt 2024 in **(i)** E18.5, **(ii)** P32, and **(iii)** P90 month mouse astrocytes. Lollipop
plots show the enrichment ratio of GO Biological Process and KEGG pathways from an over-representation analysis
with weighted set cover redundancy reduction, with tip size inversely proportional to the false discovery rate (FDR).
823
824

Figure 4. Cell-cell communication analysis for neuron-astrocyte and astrocyte-neuron predicted ligand-receptor pairs across regions and developmental time points in marmoset. A, Dot plot showing magnitude and specificity of the top 25 near-unique (shared with at most one other neuronal cluster) CellPhoneDB-predicted **(i)** neuron-astrocyte and **(ii)** astrocyte-neuron ligand receptor pairs for the most abundant astrocyte and neuronal Leiden clusters in the fetal marmoset thalamus. The source cell (top of the plot) expresses the ligand (left side of arrow on the row labels), while the target cell (bottom of the plot) expresses the receptor (right side of arrow on the row labels). The color of the dot indicates ligand-receptor expression magnitude ("Lr_means", calculated as the average of the mean expression of the ligand in the source group and the mean expression of the receptor in the target group), while the size of the dot is inversely related to the p-value on ligand-receptor expression sensitivity (-

log10(p)). **B,** Same as **(A)**, for the 14-month marmoset thalamus. **C,** UpSet plot showing the number of overlapping neuron-astrocyte predicted ligand-receptor pairs between regions, from the most abundant neuronal and astrocyte subtypes in each region for **(i)** fetal and **(ii)** late adolescent marmoset. For cortex (purple), glutamatergic L2/3IT neurons to cortical astrocytes; striatum (yellow), *DRD1+* medium spiny neurons to striatal telencephalic astrocytes; and thalamus (green) thalamic glutamatergic neurons to thalamic astrocytes. The colored dots below each vertical bar indicate which regional neuron-astrocyte (N-A) subtypes share that set of L-R pairs, while the colored horizontal bars indicate the total number of L-R pairs for each N-A subtype. Unlike in panel **A**, all L-R pairs meeting minimum expression criteria, including pairs shared with other neuronal and astrocytic clusters, were included in this analysis and that for panel **(D)**. **D,** UpSet plot (as in **(C)**) showing the number of overlapping cortical glutamatergic L2/3IT neuron to cortical astrocyte predicted ligand-receptor pairs between ages.

Figure 5. The postnatal developmental specification of marmoset astrocytes within and across brain regions. A, Integrated UMAP embeddings of 103,009 marmoset astrocytes colored by **(i-ii)** Palantir-predicted pseudotime, **(iii)** developmental time point, **(iv)** brain region (assigned), and **(v)** Leiden cluster assignment. Trajectory path (black lines and arrows) are overlaid for **(i)** telencephalic astrocyte (AST-TE) and **(ii)** diencephalic astrocyte (AST-DI) branches. **B,** Violin plot (scanpy's default) showing the estimated distribution of pseudotime values for the astrocytes in **(A)** grouped by region within each developmental time point (color code as in **A(i)**). Vertical dashed lines indicate separation between time points. **C,** Heatmaps (rows corresponding to nuclei and columns to gene) showing expression in logCPM of astrocyte age differentially expressed genes (aDEGs) in astrocytes from **(i)** cortex, **(ii)** striatum, and **(iii)** thalamus, grouped by developmental time point as indicated on the left of the heatmap. The strategy for calculating aDEGs is schematized on the left. **D,** UpSet plot showing the number of overlapping GD135 vs. 14-month astrocyte aDEGs between cortex, striatum, and thalamus. The colored dots below each vertical bar indicate which region(s) share that set of aDEGs, while the colored horizontal bars indicate the total number of cortex-thalamus aDEGs for each region. Overlap categories with 0 aDEGs are not shown. **E,** UpSet plot (as in **(D)**) showing the number of overlapping GD135 vs. 14-month cortical astrocyte aDEGs between OPCs (light blue), astrocytes (dark blue), excitatory neurons (pink), and inhibitory neurons (gray). **F,** Matrix plot showing mean expression of selected cortex group astrocyte-specific, region-specific (AS-RS) aDEGs (rows) in marmoset astrocytes grouped by region and developmental time point (columns, blocked by region first and then

864 by increasing age within each region block). Expression units of mean logCPM are standardized between 0 and 1
865 by subtracting the minimum and dividing by the maximum for each trait.
866
867

868
Figure 6. Conservation and divergence of the development of astrocyte heterogeneity in mouse and
marmoset. A, scANVI-integrated UMAP embeddings of marmoset and mouse astrocytes, colored by **(i)**
supercluster, **(ii)** species, **(iii)** age, and **(iv)** region. **B,** Venn diagrams showing regional differentially expressed
genes (rDEGs) between cortex and thalamus astrocytes shared across mouse and marmoset at **(i)** fetal, **(ii)** early
adolescent, and **(iii)** young adult time points. **C,** Venn diagram showing age differentially expressed genes (aDEGs)
shared between mouse and marmoset astrocytes within the **(i)** cortex and **(ii)** thalamus. **D,** Heatmaps showing
expression in logCPM of species differentially expressed genes (sDEGs) between marmoset and mouse within **(i)**
telencephalic astrocytes and **(ii)** diencephalic astrocytes. **E,** *In situ* validation of selected sDEGs in marmoset and
mouse tissue with the RNAscope v2 assay. **(i)** Heatmap (rows are nuclei, columns are genes) of sDEGs *NRG3*/*Nrg3*
(higher in marmoset) and *LAMA2*/*Lama2* (higher in mouse) expression in marmoset and mouse astrocytes in
logCPM. **(ii)** Top row: single-channel and composite maximum intensity projections of cropped fields of view in the

marmoset (top row) and mouse (bottom row) thalamus stained via RNAscope v2 FISH for astrocyte marker
SLC1A3/Slc1a3, *NRG3/Nrg3*, and mouse *LAMA2/Lama2*. Scale bar, 50μm. Bottom row: high-magnification images
of the boxed astrocytes in the top row. Scale bar, 5μm. **(iii)** CellProfiler quantification of sDEG abundance in both
species from RNAscope v2 data (n = 2 donors per species, see **Methods**). From left to right: Fraction of probe
positive astrocytes for *LAMA2/Lama2*, fraction of probe positive for *NRG3/Nrg3*, normalized mean intensity of
LAMA2/Lama2 (mean intensity in expanded astrocyte nuclei divided by mean intensity in all nuclei, including
astrocyte nuclei), and normalized mean intensity of *NRG3/Nrg3*. Data points are from individual biological replicates,
with 2 slices averaged for mouse, and the horizontal black line denotes the median. **F,** UpSet plot showing shared
sDEGs across superclusters, comparing telencephalic astrocytes, diencephalic astrocytes, oligodendrocyte
precursor cells (OPCs) medial ganglionic eminence-derived and *PVALB+* cortical GABAergic neurons (Ctx Gad
MGE-PVALB), thalamic reticular nucleus (TRN) GABAergic neurons, striatal medium spiny neurons (MSNs), and
microglia. The colored dots below each vertical bar indicate which supercluster(s) share that set of sDEGs, while
the colored horizontal bars indicate the total number of mouse-marmoset sDEGs for each supercluster. For
simplicity, only supercluster combinations with 10 or more shared sDEGs are shown.
894
895

Figure 7. Expansion microscopy of virally-labeled astrocytes in the mouse and marmoset brain. A, Viral labeling approach for mouse astrocytes (see **Methods** for details). Created using BioRender.com. **B,** Brain slice hemispheres containing regions of interest in **(i)** PFC **(ii)** striatum and **(iii)** thalamus after pre-expansion staining for GFP, as visible through the eyepiece on a dissecting microscope under blue light illumination. **C-E, (i)** Single z-slices of background-subtracted images of ~3.5x expanded astrocytes in the **(i)** prefrontal cortex, **(ii)** striatum, and **(iii)** thalamus, co-stained with GFP, GFAP, and the blood vessel marker Lectin. Shown are 6 examples of medium to high GFP-expressing astrocytes from 6 separate mice. Scale bar, 10µm in biological units. Contrast was adjusted to 35% saturation (Fiji's "auto") in most cases and was further increased for dimmer astrocytes to aid visibility; therefore, astrocytes are not equally contrast adjusted. See Movies S1-36 for 3D-projected image volumes and their corresponding 3D segmentations. **D, (i)** Single z-slice of one of the thalamic astrocytes in **C(iii)** and **(ii)** its corresponding binary segmentation used as input for morphology analysis. **E,** Bar plots showing quantified morphological properties for mouse astrocytes from PFC, striatum, and thalamus (n = 52-60 astrocytes from 3

female and 5 male mice for each region, with statistical significance determined using a linear mixed effects model
with "animal" as the random effect group variable, see **Table S25**): **(i)** volume, **(ii)** surface area, **(iii)** surface area /
volume, **(iv)** equivalent diameter, **(v)** aspect ratio, **(vi)** Sholl analysis (number of intersections with concentric shells
as a function of radius), and **(vii)** fractal dimension by the box counting method (see **Methods**). ns, non significant,

*p≤0.05, **p≤0.01, ***p≤0.001, ****p≤0.0001. **F,** Maximum intensity projection composites of cropped (in x and y
and across 51 z-slices) regions of GFP-labeled astrocyte processes co-stained with either **(i)** Glast (telencephalic
rDEG) or **(iii)** Gat3 (thalamic rDEG) and the synaptic protein Cav2.1 alongside GFP generated using the ExR
protocol (~18x expansion factor). Shown are processes from astrocytes in the PFC (top row), striatum (middle row),
and thalamus (bottom row) of the same mouse (though **(i)** and **(iii)** are from different mice). Scale bar, 0.5μm.
Contrast was manually adjusted by setting equal minimum and maximum intensity values for Cav2.1 and rDEG
targets and using Fiji's "Auto" for GFP. **(ii, iv)** Bar plots showing quantified mean intensity of either Glast or Gat3
within masked GFP+ regions (astrocytes) in the whole ExR image volume (n = 10-16 fields of view from 2 mice,
with statistical significance determined using a linear mixed effect model as described in the **Methods, see Table**
S26 for numerical results). In panels **E** and **F**, error bars indicate standard error of the mean.
923
924

<u>Supplementary Figures and Movies</u>
Supplementary Figure 1. Summary of sequencing coverage for the cross-region developmental
snRNAseq atlas.
Supplementary Figure 2. Distribution of nuclei across cell types, regions, ages, sexes, and study for
marmoset single-nucleus RNAseq data.
Supplementary Figure 3. Distribution of nuclei across cell types, regions, ages, sexes, and study for
mouse single-nucleus RNAseq data.
Supplementary Figure 4. Leiden cluster annotations and proportions for the cross-region developmental
snRNAseq atlas.
Supplementary Figure 5. Dissection strategies for marmoset brains and region reassignment to mitigate
cross-region contamination for developing samples
Supplementary Figure 6. Region reassignment for cross-contaminant nuclei in mouse samples.
Supplementary Figure 7. Correlation of pairwise astrocyte rDEG log-fold change between region pairs
across development in mouse and marmoset.
Supplementary Figure 8. Validation of marmoset astrocyte rDEG expression in situ using multiplexed
FISH.
Supplementary Figure 9. Quantification of selected rDEG and astrocyte subtype marker expression in situ
in adult and neonate marmoset.
Supplementary Figure 10. Validation of mouse astrocyte rDEG expression in situ using multiplexed FISH.
Supplementary Figure 11. Quantification of selected rDEG and astrocyte subtype marker expression in
situ in adult and neonate mouse.
Supplementary Figure 12. Astrocyte sub-clustering captures intra-regional heterogeneity in marmoset.
Supplementary Figure 13. Astrocyte sub-clustering captures intra-regional heterogeneity in mouse.
Supplementary Figure 14. Cell-cell communication analysis for neuron-astrocyte and astrocyte-neuron
predicted ligand-receptor pairs across regions and developmental time points in mouse.
Supplementary Figure 15. Pseudotime inference in mouse and marmoset oligodendrocyte lineage and
mouse astrocytes.
Supplementary Figure 16. Gene expression signatures underlying the postnatal developmental
specification of mouse astrocytes within and across brain regions.
Supplementary Figure 17. Cell type composition across development in both species and cross-species
integration with SATURN.
Supplementary Figure 18. Expression of Top2a in astrocytes of the mouse subventricular zone via
multiplexed FISH.
Supplementary Figure 19. Additional detail on astrocyte morphology and rDEG protein expression
differences across brain regions in mouse.
Supplementary Figure 20. High-concentration collagenase treatment preserves blood vessel morphology
in ExR samples.
Supplementary Movies 1-36. 3D-projected image volumes of ~4x expanded mouse astrocytes.
963
<u>Supplementary Tables</u>
Supplementary Table 1. Biological donor information.
Supplementary Table 2. Abundance of cell types and Leiden clusters and descriptions of Leiden clusters
in mouse and marmoset.
Supplementary Table 3. Proportional breakdown of each Leiden cluster by age, assigned region, and sex
for mouse and marmoset.
Supplementary Table 4. Proportional breakdown of MapMyCells-derived Allen Brain Cell Atlas subclass
assignments by Leiden cluster for mouse and marmoset.
Supplementary Table 5. Summary of significant compositional differences in cell type for marmoset as
found by scCODA.
Supplementary Table 6. Summary of significant compositional differences in cell type for mouse as found
by scCODA.

Supplementary Table 7. Summary of significant compositional differences in Leiden cluster for marmoset
as found by scCODA.
Supplementary Table 8. Summary of significant compositional differences in Leiden cluster for mouse as
found by scCODA.
Supplementary Table 9. Marmoset rDEGs shared across both replicates for each age.
Supplementary Table 10. WebGestalt enrichment results for marmoset rDEGs by age.
Supplementary Table 11. Mouse rDEGs for each age.
Supplementary Table 12. WebGestalt enrichment results for mouse rDEGs by age.
Supplementary Table 13. Image-level quantification of each rDEG probe for adult and neonate marmoset
and mouse and results of statistical tests for mouse.
Supplementary Table 14. Proportion astrocytes positive for one or more fibrous marker(s) in each time
point-region combination for both species.
Supplementary Table 15. Top ABCA MapMyCells cluster assignments for astrocyte subclusters.
Supplementary Table 16. CellPhoneDB-generated cell-cell communication results for each region-age
combination in mouse and marmoset.
Supplementary Table 17. Fraction of CellPhoneDB-generated ligand-receptor pairs shared across 3 or more
ages for all neuronal clusters within a region for both species.
Supplementary Table 18. Genes with the highest Mellon change scores over pseudotime for mouse and
marmoset telencephalic and diencephalic astrocytes.
Supplementary Table 19. Marmoset astrocyte, OPC, GABAergic, and glutamatergic neuron aDEGs for each
brain region.
Supplementary Table 20. Mouse astrocyte, OPC, GABAergic, and glutamatergic neuron aDEGs for each
brain region.
Supplementary Table 21. Shared and species-unique cortex-thalamus astrocyte rDEGs for each
developmental timepoint in mouse and marmoset.
Supplementary Table 22. Shared and species-unique fetal-late adolescent astrocyte aDEGs for each region
in mouse and marmoset.
Supplementary Table 23. sDEGs for each supercluster.
Supplementary Table 24. Telencephalic-specific, diencephalic-specific, and shared astrocyte sDEGs.
Supplementary Table 25. Summary statistics for quantification of mouse astrocyte morphology across
brain regions.
Supplementary Table 26. Summary statistics for quantification of Glast and Gat3 expression in ~18x ExR
images from mouse PFC, striatum, and thalamus.
Supplementary Table 27. Blocking strategy for marmoset region dissection based on custom brain matrix.
Supplementary Table 289. Sequencing coverage statistics for all 10x Chromium reactions.
Supplementary Table 30. Table used to convert between mouse, marmoset, and human gene IDs for 1:1
orthologs.
Supplementary Table 31. List of reagents, including vendor and product information, used for experiments.
Supplementary Table 32. Expansion factor measurements for mouse ExR samples.
Supplementary Table 33. Notes on each ~4x expanded mouse astrocyte imaged (Fig. 7 and Fig. S19).
016
017

References

1. Yao, Z. *et al.* A high-resolution transcriptomic and spatial atlas of cell types in the whole mouse brain. *Nature* **624**, 317–332 (2023).
2. Langlieb, J. *et al.* The molecular cytoarchitecture of the adult mouse brain. *Nature* **624**, 333–342 (2023).
3. Aldridge, S. & Teichmann, S. A. Single cell transcriptomics comes of age. *Nat. Commun.* **11**, 4307 (2020).
4. Siletti, K. *et al.* Transcriptomic diversity of cell types across the adult human brain. *Science* **382**, eadd7046 (2023).
5. La Manno, G. *et al.* Molecular architecture of the developing mouse brain. *Nature* **596**, 92–96 (2021).
6. Braun, E. *et al.* Comprehensive cell atlas of the first-trimester developing human brain. *Science* **382**, eadf1226 (2023).
7. Krienen, F. M. *et al.* A marmoset brain cell census reveals regional specialization of cellular identities. *Sci. Adv.* **9**, eadk3986 (2023).
8. Lin, J.-P. *et al.* Transcriptomic architecture of nuclei in the marmoset CNS. *Nat. Commun.* **13**, 5531 (2022).
9. Yao, Z. *et al.* A taxonomy of transcriptomic cell types across the isocortex and hippocampal formation. *Cell* **184**, 3222-3241.e26 (2021).
10. Allen, N. J. & Barres, B. A. Glia — more than just brain glue. *Nature* **457**, 675–677 (2009).
11. Ben Haim, L. & Rowitch, D. H. Functional diversity of astrocytes in neural circuit regulation. *Nat. Rev. Neurosci.* **18**, 31–41 (2016).
12. Oliveira, J. F. & Araque, A. Astrocyte regulation of neural circuit activity and network states. *Glia* **70**, 1455–1466 (2022).
13. Lee, H. G., Wheeler, M. A. & Quintana, F. J. Function and therapeutic value of astrocytes in neurological diseases. *Nat. Rev. Drug Discov. 2022 215* **21**, 339–358 (2022).
14. Verkhratsky, A. & Nedergaard, M. Physiology of Astroglia. *Physiol. Rev.* **98**, 239–389 (2018).
15. Ramón Y Cajal, S., Swanson, N. & Swanson, L. *Histology of the Nervous System of Man and Vertebrates*. (Oxford University Press, Oxford, New York, 1995).
16. Oberheim, N. A., Goldman, S. A. & Nedergaard, M. Heterogeneity of astrocytic form and function. *Methods Mol. Biol.* **814**, 23–45 (2012).
17. Bachoo, R. M. *et al.* Molecular diversity of astrocytes with implications for neurological disorders. *Proc. Natl. Acad. Sci.* **101**, 8384–8389 (2004).
18. Morel, L. *et al.* Molecular and Functional Properties of Regional Astrocytes in the Adult Brain. *J. Neurosci.* **37**, 8706–8717 (2017).
19. Chai, H. *et al.* Neural Circuit-Specialized Astrocytes: Transcriptomic, Proteomic, Morphological, and Functional Evidence. *Neuron* **95**, 531-549.e9 (2017).
20. John Lin, C. C. *et al.* Identification of diverse astrocyte populations and their malignant analogs. *Nat. Neurosci. 2017 203* **20**, 396–405 (2017).
21. Saunders, A. *et al.* Molecular Diversity and Specializations among the Cells of the Adult Mouse Brain. *Cell* **174**, 1015-1030.e16 (2018).
22. Batiuk, M. Y. *et al.* Identification of region-specific astrocyte subtypes at single cell resolution. *Nat. Commun.* **11**, 1220 (2020).
23. Bayraktar, O. A. *et al.* Astrocyte layers in the mammalian cerebral cortex revealed by a single-cell in situ transcriptomic map. *Nat. Neurosci.* **23**, 500–509 (2020).
24. Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* **174**, 999-1014.e22 (2018).
25. Endo, F. *et al.* Molecular basis of astrocyte diversity and morphology across the CNS in health and disease. *Science* **378**, (2022).
26. Hochstim, C., Deneen, B., Lukaszewicz, A., Zhou, Q. & Anderson, D. J. Identification of Positionally Distinct Astrocyte Subtypes whose Identities Are Specified by a Homeodomain Code. *Cell* **133**, 510–522 (2008).
27. Tsai, H. H. *et al.* Regional astrocyte allocation regulates CNS synaptogenesis and repair. *Science* **337**, 358–362 (2012).

28. Welle, A. *et al.* Epigenetic control of region-specific transcriptional programs in mouse cerebellar
and cortical astrocytes. *Glia* **69**, 2160–2177 (2021).
29. Clavreul, S. *et al.* Cortical astrocytes develop in a plastic manner at both clonal and cellular levels.
Nat. Commun. **10**, (2019).
30. Farmer, W. T. *et al.* Neurons diversify astrocytes in the adult brain through sonic hedgehog
signaling. *Science* **351**, 849–854 (2016).
31. Xie, Y. *et al.* Developmental origin and local signals cooperate to determine septal astrocyte
identity. 2023.10.08.561428 Preprint at https://doi.org/10.1101/2023.10.08.561428 (2023).
32. Molofsky, A. V. & Deneen, B. Astrocyte development: A Guide for the Perplexed. *Glia* **63**, 1320–
1329 (2015).
33. Tung, P. Y. *et al.* Batch effects and the effective design of single-cell gene expression studies.
Sci. Rep. 2017 71 **7**, 1–15 (2017).
34. Tran, H. T. N. *et al.* A benchmark of batch-effect correction methods for single-cell RNA
sequencing data. *Genome Biol.* **21**, 1–32 (2020).
35. Zhou, B., Zuo, Y.-X. & Jiang, R.-T. Astrocyte morphology: Diversity, plasticity, and role in
neurological diseases. *CNS Neurosci. Ther.* **25**, 665–673 (2019).
36. Baldwin, K. T., Murai, K. K. & Khakh, B. S. Astrocyte morphology. *Trends Cell Biol.* **34**, 547–565
(2024).
37. Sarkar, D. *et al.* Revealing nanostructures in brain tissue via protein decrowding by iterative
expansion microscopy. *Nat. Biomed. Eng.* **6**, 1057–1073 (2022).
38. Cao, Z.-J. & Gao, G. Multi-omics single-cell data integration and regulatory inference with graph-
linked embedding. *Nat. Biotechnol.* **40**, 1458–1466 (2022).
39. Luecken, M. D. *et al.* Benchmarking atlas-level data integration in single-cell genomics. *Nat.*
Methods 2021 191 **19**, 41–50 (2021).
40. Traag, V., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-connected
communities. *Sci. Rep.* **9**, (2018).
41. Allen Institute for Brain Science. MapMyCells.
42. Büttner, M., Ostner, J., Müller, C. L., Theis, F. J. & Schubert, B. scCODA is a Bayesian model for
compositional single-cell data analysis. *Nat. Commun.* **12**, 6876 (2021).
43. Rubenstein, J. L. R., Shimamura, K., Martinez, S. & Puelles, L. REGIONALIZATION OF THE
PROSENCEPHALIC NEURAL PLATE. *Annu. Rev. Neurosci.* **21**, 445–477 (1998).
44. Sasai, Y. & De Robertis, E. M. Ectodermal Patterning in Vertebrate Embryos. *Dev. Biol.* **182**, 5–20
(1997).
45. Di Bella, D. J. *et al.* Molecular logic of cellular diversification in the mouse cerebral cortex. *Nature*
595, 554–559 (2021).
46. Anthony, T. E., Klein, C., Fishell, G. & Heintz, N. Radial Glia Serve as Neuronal Progenitors in All
Regions of the Central Nervous System. *Neuron* **41**, 881–890 (2004).
47. Rowitch, D. H. & Kriegstein, A. R. Developmental genetics of vertebrate glial–cell specification.
Nature **468**, 214–222 (2010).
48. Frassoni, C., Amadeo, A., Ortino, B., Jaranowska, A. & Spreafico, R. Organization of radial and
non-radial glia in the developing rat thalamus. *J. Comp. Neurol.* **428**, 527–542 (2000).
49. Tan, C. X. & Eroglu, C. Cell adhesion molecules regulating astrocyte–neuron interactions. *Curr.*
Opin. Neurobiol. **69**, 170–177 (2021).
50. Sobolczyk, M. & Boczek, T. Astrocytic Calcium and cAMP in Neurodegenerative Diseases. *Front.*
Cell. Neurosci. **16**, (2022).
51. Zhou, Z. *et al.* Astrocytic cAMP modulates memory via synaptic plasticity. *Proc. Natl. Acad. Sci.*
U. S. A. **118**, e2016584118 (2021).
52. Bazargani, N. & Attwell, D. Astrocyte calcium signaling: The third wave. *Nat. Neurosci.* **19**, 182–
189 (2016).
53. Kucukdereli, H. *et al.* Control of excitatory CNS synaptogenesis by astrocyte-secreted proteins
hevin and SPARC. *Proc. Natl. Acad. Sci. U. S. A.* **108**, (2011).
54. Elizarraras, J. M. *et al.* WebGestalt 2024: faster gene set analysis and new support for
metabolomics and multi-omics. *Nucleic Acids Res.* **52**, W415–W421 (2024).
55. Zhang, B., Kirov, S. & Snoddy, J. WebGestalt: an integrated system for exploring gene sets in

various biological contexts. *Nucleic Acids Res.* **33**, W741–W748 (2005).

56. The UniProt Consortium. UniProt: the Universal Protein Knowledgebase in 2025. *Nucleic Acids Res.* **53**, D609–D617 (2025).

57. Liu, Y. F. *et al.* Autism and Intellectual Disability-Associated KIRREL3 Interacts with Neuronal Proteins MAP1B and MYO16 with Potential Roles in Neurodevelopment. *PLOS ONE* **10**, e0123106 (2015).

58. Kim, D. & Ackerman, S. L. The UNC5C Netrin Receptor Regulates Dorsal Guidance of Mouse Hindbrain Axons. *J. Neurosci.* **31**, 2167–2179 (2011).

59. Wetzel-Smith, M. K. *et al.* A rare mutation in UNC5C predisposes to late-onset Alzheimer's disease and increases neuronal cell death. *Nat. Med.* **20**, 1452–1457 (2014).

60. Carpenter, A. E. *et al.* CellProfiler: image analysis software for identifying and quantifying cell phenotypes. *Genome Biol.* **7**, R100 (2006).

61. Stirling, D. R. *et al.* CellProfiler 4: improvements in speed, utility and usability. *BMC Bioinformatics* **22**, 433 (2021).

62. Bakken, T. E. *et al.* Comparative cellular analysis of motor cortex in human, marmoset and mouse. *Nat. 2021 5987879* **598**, 111–119 (2021).

63. Hodge, R. D. *et al.* Conserved cell types with divergent features in human versus mouse cortex. *Nat. 2019 5737772* **573**, 61–68 (2019).

64. Fang, R. *et al.* Conservation and divergence of cortical cell organization in human and mouse revealed by MERFISH. *Science* **377**, 56–62 (2022).

65. Efremova, M., Vento-Tormo, M., Teichmann, S. A. & Vento-Tormo, R. CellPhoneDB: inferring cell–cell communication from combined expression of multi-subunit ligand–receptor complexes. *Nat. Protoc.* **15**, 1484–1506 (2020).

66. Delaunay, D. *et al.* Early Neuronal and Glial Fate Restriction of Embryonic Neural Stem Cells. *J. Neurosci.* **28**, 2551–2562 (2008).

67. Hong, W. *et al.* Temporal-spatial Generation of Astrocytes in the Developing Diencephalon. *Neurosci. Bull.* **40**, 1–16 (2024).

68. Setty, M. *et al.* Characterization of cell fate probabilities in single-cell data with Palantir. *Nat. Biotechnol.* **37**, 451–460 (2019).

69. Marques, S. *et al.* Oligodendrocyte heterogeneity in the mouse juvenile and adult central nervous system. *Science* **352**, 1326–1329 (2016).

70. Otto, D. J., Jordan, C., Dury, B., Dien, C. & Setty, M. Quantifying cell-state densities in single-cell phenotypic landscapes using Mellon. *Nat. Methods* **21**, 1185–1195 (2024).

71. Zhang, Y. *et al.* Purification and Characterization of Progenitor and Mature Human Astrocytes Reveals Transcriptional and Functional Differences with Mouse. *Neuron* **89**, 37–53 (2016).

72. Han, X. *et al.* Forebrain Engraftment by Human Glial Progenitor Cells Enhances Synaptic Plasticity and Learning in Adult Mice. *Cell Stem Cell* **12**, 342–353 (2013).

73. Xu, C. *et al.* Probabilistic harmonization and annotation of single-cell transcriptomics data with deep generative models. *Mol. Syst. Biol.* **17**, e9620 (2021).

74. Rosen, Y. *et al.* Toward universal cell embeddings: integrating single-cell RNA-seq datasets across species with SATURN. *Nat. Methods* 1–9 (2024) doi:10.1038/s41592-024-02191-z.

75. Sohn, J. *et al.* The Subventricular Zone Continues to Generate Corpus Callosum and Rostral Migratory Stream Astroglia in Normal Adult Mice. *J. Neurosci.* **35**, 3756–3763 (2015).

76. Abrahams, B. S. *et al.* SFARI Gene 2.0: a community-driven knowledgebase for the autism spectrum disorders (ASDs). *Mol. Autism* **4**, 36 (2013).

77. Jorstad, N. L. *et al.* Comparative transcriptomics reveals human-specific cortical features. *Science* **382**, eade9516 (2023).

78. Krienen, F. M. *et al.* Innovations present in the primate interneuron repertoire. *Nature* **586**, 262–269 (2020).

79. Chung, W.-S., Baldwin, K. T. & Allen, N. J. Astrocyte Regulation of Synapse Formation, Maturation, and Elimination. *Cold Spring Harb. Perspect. Biol.* **16**, a041352 (2024).

80. Liebovitch, L. S. & Toth, T. A fast algorithm to determine fractal dimensions by box counting. *Phys. Lett. A* **141**, 386–390 (1989).

81. Ferreira, T. A. *et al.* Neuronal morphometry directly from bitmap images. *Nat. Methods* **11**, 982–

984 (2014).

82. Soto, J. S. *et al.* Astrocyte Gi-GPCR signaling corrects compulsive-like grooming and anxiety-
related behaviors in Sapap3 knockout mice. *Neuron* **0**, (2024).

83. Zhang, Y. & Barres, B. A. Astrocyte heterogeneity: an underappreciated topic in neurobiology.
Curr. Opin. Neurobiol. **20**, 588–594 (2010).

84. Bayraktar, O. A., Fuentealba, L. C., Alvarez-Buylla, A. & Rowitch, D. H. Astrocyte Development
and Heterogeneity. *Cold Spring Harb. Perspect. Biol.* **7**, a020362 (2015).

85. Schober, A. L., Wicki-Stordeur, L. E., Murai, K. K. & Swayne, L. A. Foundations and implications
of astrocyte heterogeneity during brain development and disease. *Trends Neurosci.* (2022)
doi:10.1016/J.TINS.2022.06.009.

86. Semple, B. D., Blomgren, K., Gimlin, K., Ferriero, D. M. & Noble-Haeusslein, L. J. Brain
development in rodents and humans: Identifying benchmarks of maturation and vulnerability to
injury across species. *Prog. Neurobiol.* **106–107**, 1–16 (2013).

87. Freeman, M. R. Specification and Morphogenesis of Astrocytes. *Science* **330**, 774–778 (2010).

88. Cho, F. S. *et al.* Enhancing GAT-3 in thalamic astrocytes promotes resilience to brain injury in
rodents. *Sci. Transl. Med.* **14**, 4310 (2022).

89. Craig, A. M. & Kang, Y. Neurexin–neuroligin signaling in synapse development. *Curr. Opin.*
Neurobiol. **17**, 43–52 (2007).

90. Holt, L. M. *et al.* Astrocyte morphogenesis is dependent on BDNF signaling via astrocytic TrkB.T1.
eLife **8**, e44667 (2019).

91. Deconinck, L., Cannoodt, R., Saelens, W., Deplancke, B. & Saeys, Y. Recent advances in
trajectory inference from single-cell omics data. *Curr. Opin. Syst. Biol.* **27**, 100344 (2021).

92. Nowakowski, T. J. *et al.* Spatiotemporal gene expression trajectories reveal developmental
hierarchies of the human cortex. *Science* **358**, 1318–1323 (2017).

93. Breschi, A., Gingeras, T. R. & Guigó, R. Comparative transcriptomics in human and mouse. *Nat.*
Rev. Genet. **18**, 425–440 (2017).

94. Okano, H., Hikishima, K., Iriki, A. & Sasaki, E. The common marmoset as a novel animal model
system for biomedical and neuroscience research applications. *Semin. Fetal. Neonatal Med.* **17**,
336–340 (2012).

95. Abbott, D. H., Barnett, D. K., Colman, R. J., Yamamoto, M. E. & Schultz-Darken, N. J. Aspects of
Common Marmoset Basic Biology and Life History Important for Biomedical Research. *Comp.*
Med. **53**, 339–350 (2003).

96. Lindhout, F. W., Krienen, F. M., Pollard, K. S. & Lancaster, M. A. A molecular and cellular
perspective on human brain evolution and tempo. *Nature* **630**, 596–608 (2024).

97. Khakh, B. S. & Deneen, B. The Emerging Nature of Astrocyte Diversity. *Annu. Rev. Neurosci.* **42**,
187–207 (2019).

98. Holt, C. E. & Schuman, E. M. The Central Dogma Decentralized: New Perspectives on RNA
Function and Local Translation in Neurons. *Neuron* **80**, 648–657 (2013).

99. Sakers, K. *et al.* Astrocytes locally translate transcripts in their peripheral processes. *Proc. Natl.*
Acad. Sci. **114**, E3830–E3838 (2017).

100. Bakken, T. E. *et al.* Single-nucleus and single-cell transcriptomes compared in matched cortical
cell types. *PLoS ONE* **13**, (2018).

101. Pool, A.-H., Poldsam, H., Chen, S., Thomson, M. & Oka, Y. Recovery of missing single-cell RNA-
sequencing data with optimized transcriptomic references. *Nat. Methods* **20**, 1506–1515 (2023).

102. Cheng, Y.-T. *et al.* Inhibitory input directs astrocyte morphogenesis through glial GABABR. *Nature*
617, 369–376 (2023).

103. Paxinos, G., Watson, C., Petrides, M., Rosa, M. & Tokuno, H. *The Marmoset Brain in Stereotaxic*
Coordinates. (Elsevier, 2012).

104. Fleming, S. J. *et al.* Unsupervised removal of systematic background noise from droplet-based
single-cell experiments using CellBender. *Nat. Methods* **20**, 1323–1335 (2023).

105. Virshup, I. *et al.* The scverse project provides a computational ecosystem for single-cell omics
data analysis. *Nat. Biotechnol.* **41**, 604–606 (2023).

106. Takabayashi, S. & Katoh, H. Sex Identification Using the ZFX and ZFY Genes in Common
Marmosets (Callithrix jacchus). *Exp. Anim.* **60**, 417–420 (2011).

107. Zargari, M. *et al.* Fetal Sex Determination using Non-Invasive Method of Cell-free Fetal DNA in
Maternal Plasma of Pregnant Women During 6th– 10th Weeks of Gestation. *Avicenna J. Med.*
Biotechnol. **3**, 201–206 (2011).
108. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data
analysis. *Genome Biol.* **19**, 15 (2018).
109. Lopez, R., Regier, J., Cole, M. B., Jordan, M. I. & Yosef, N. Deep generative modeling for single-
cell transcriptomics. *Nat. Methods* **15**, 1053–1058 (2018).
110. Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP. *Nat.*
Biotechnol. **37**, 38–44 (2019).
111. Bernstein, N. J. *et al.* Solo: Doublet Identification in Single-Cell RNA-Seq via Semi-Supervised
Deep Learning. *Cell Syst.* **11**, 95-101.e5 (2020).
112. Heumos, L. *et al.* Best practices for single-cell analysis across modalities. *Nat. Rev. Genet.* **24**,
550–572 (2023).
113. Kolberg, L. *et al.* g:Profiler—interoperable web service for functional enrichment analysis and
gene identifier mapping (2023 update). *Nucleic Acids Res.* **51**, W207–W212 (2023).
114. Cunningham, F. *et al.* Ensembl 2019. *Nucleic Acids Res.* **47**, D745–D751 (2019).
115. van Dijk, D. *et al.* Recovering Gene Interactions from Single-Cell Data Using Data Diffusion. *Cell*
174, 716-729.e27 (2018).
116. Lex, A., Gehlenborg, N., Strobelt, H., Vuillemot, R. & Pfister, H. UpSet: Visualization of
Intersecting Sets. *IEEE Trans. Vis. Comput. Graph.* **20**, 1983–1992 (2014).
117. Ashburner, M. *et al.* Gene Ontology: tool for the unification of biology. *Nat. Genet.* **25**, 25–29
(2000).
118. The Gene Ontology Consortium *et al.* The Gene Ontology knowledgebase in 2023. *Genetics* **224**,
iyad031 (2023).
119. Kanehisa, M. & Goto, S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res.*
28, 27–30 (2000).
120. Lin, Z. *et al.* Evolutionary-scale prediction of atomic-level protein structure with a language model.
Science **379**, 1123–1130 (2023).
121. Geirsdottir, L. *et al.* Cross-Species Single-Cell Analysis Reveals Divergence of the Primate
Microglia Program. *Cell* **179**, 1609-1622.e16 (2019).
122. Dimitrov, D. *et al.* Comparison of methods and resources for cell-cell communication inference
from single-cell RNA-Seq data. *Nat. Commun.* **13**, 3224 (2022).
123. Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. Generalizing RNA velocity to transient
cell states through dynamical modeling. *Nat. Biotechnol.* **38**, 1408–1414 (2020).
124. La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494–498 (2018).
125. Aivazidis, A. *et al.* Model-based inference of RNA velocity modules improves cell fate prediction.
2023.08.03.551650 Preprint at https://doi.org/10.1101/2023.08.03.551650 (2023).
126. Dijk, D. van *et al.* MAGIC: A diffusion-based imputation method reveals gene-gene interactions in
single-cell RNA-sequencing data. 111591 Preprint at https://doi.org/10.1101/111591 (2017).
127. Wang, H. *et al.* Region-specific distribution of Olig2-expressing astrocytes in adult mouse brain
and spinal cord. *Mol. Brain* **14**, 36 (2021).
128. Martynoga, B., Morrison, H., Price, D. J. & Mason, J. O. Foxg1 is required for specification of
ventral telencephalon and region-specific regulation of dorsal telencephalic precursor proliferation
and apoptosis. *Dev. Biol.* **283**, 113–127 (2005).
129. Kita, Y. *et al.* Cellular-resolution gene expression profiling in the neonatal marmoset brain reveals
dynamic species- and region-specific differences. *Proc. Natl. Acad. Sci.* **118**, e2020125118
(2021).
130. Shimogori, T. *et al.* Digital gene atlas of neonate common marmoset brain. *Neurosci. Res.* **128**,
1–13 (2018).
131. Gleichman, A. J., Kawaguchi, R., Sofroniew, M. V. & Carmichael, S. T. A toolbox of astrocyte-
specific, serotype-independent adeno-associated viral vectors using microRNA targeting
sequences. *Nat. Commun.* **14**, 7426 (2023).
132. Chen, F., Tillberg, P. W. & Boyden, E. S. Expansion microscopy. *Science* **347**, 543–548 (2015).
133. Challis, R. C. *et al.* Systemic AAV vectors for widespread and targeted gene delivery in rodents.

Nat. Protoc. **14**, 379–414 (2019).

134. Lock, M., Alvira, M. R., Chen, S.-J. & Wilson, J. M. Absolute Determination of Single-Stranded
and Self-Complementary Adeno-Associated Viral Vector Genome Titers by Droplet Digital PCR.
Hum. Gene Ther. Methods **25**, 115–125 (2014).

135. Valdes, P. A. *et al.* Improved immunostaining of nanostructures and cells in human brain
specimens through expansion-mediated protein decrowding. *Sci. Transl. Med.* **16**, eabo0049
(2024).

136. Kang, J. *et al.* Multiplexed expansion revealing for imaging multiprotein nanostructures in healthy
and diseased brain. *Nat. Commun.* **15**, 9722 (2024).

137. Arshadi, C., Günther, U., Eddison, M., Harrington, K. I. S. & Ferreira, T. A. SNT: a unifying toolbox
for quantification of neuronal anatomy. *Nat. Methods* **18**, 374–377 (2021).