Adaptive challenges of past and future invasion of *Drosophila suzukii*: insights from novel genomic resources and statistical methods combining individual and pool sequencing data

Louise Camus[1,*], Nicolas Rode[1], Svitlana Serga[1,2], Anne Loiseau[1], Xiao Chen[3], Carole Iampietro[4], Marc Kenis[5], William Marande[6], Julián Mensch[7], Hugues Parinello[8], Marija Savić Veselinović[9], Sophie Valière[4], Jinping Zhang[10], Arnaud Estoup[1], Simon Boitard[1,†,*] and Mathieu Gautier[1,†,*]

[1] CBGP, Univ Montpellier, CIRAD, INRAE, IRD, Institut Agro Montpellier, Montpellier, France

[2] National Antarctic Scientific Center of Ukraine, Taras Shevchenko blv. 16, Kyiv, 01601, Ukraine

[3] College of Plant Protection, Yunnan Agricultural University, Kunming, 650201 China

[4] INRAE, GeT-PlaGe, Genotoul, 31326 Castanet-Tolosan, France

[5] CABI, Delémont, Switzerland

[6] French Plant Genomic Resource Center, CNRGV, INRAE, Castanet Tolosan, France

[7] Departamento de Ecología, Genética y Evolución. Facultad de Ciencias Exactas y Naturales. Universidad de Buenos Aires, Ciudad Universitaria, Pabellón II (C1428 EHA), Buenos Aires, Argentina

[8] MGX-Montpellier GenomiX, Univ Montpellier, CNRS, INSERM, Montpellier France

[9] University of Belgrade, Faculty of Biology, Studentski trg 16, 11000 Belgrade, Serbia

[10] MARA-CABI Joint Laboratory for Bio-safety, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, China

[†]These authors are joint senior authors on this work

[*]Corresponding authors: louise.camus@inrae.fr, simon.boitard@inrae.fr and mathieu.gautier@inrae.fr

Keywords: *D. suzukii*, Biological invasions, Genomic Offset, Local Adaptation, GEA, BayPass

Running title: The genomics of past and future invasion in *D. suzukii*

Abstract

Climate change and globalization are accelerating biological invasions, making it crucial to understand how species adapt in new environments to improve management strategies. Genomic data can provide valuable insights into adaptation through Genotype-Environment Association (GEA) studies, which identify genes and biological processes tied to invasion success, and through the geometric Genomic Offset statistics (gGO), which estimate genetic (mal)adaptation to new environments. In this study, we investigate genetic adaptation in the invasive pest *Drosophila suzukii* using novel genomic resources and statistical methods. We used a newly developed chromosome-level genome assembly and data from 37 populations, combining publicly available and newly generated pooled and individual sequencing data, analyzed with an enhanced version of BAYPASS software, tailored for such hybrid datasets. Our findings reveal multiple genomic regions related to invasion success, suggesting a polygenic basis for the shared adaptive traits involved in invasion success. Through a GEA incorporating 27 environmental covariates, we estimate gGO between source environments and invaded areas, shedding light on the potential adaptive challenges *D. suzukii* faced during previous invasions. In addition, we estimate gGO for geographical areas not yet invaded to predict future invasion risks and identify regions from which preadapted populations may originate. Our results suggest that *D. suzukii* populations faced limited adaptive challenges in their major invasion range, and that certain uninvaded regions remain at high risk of future invasion. Our study offers significant insights into *D. suzukii* adaptation and provides a practical population genomics framework to predict biological invasions across diverse species.

Introduction

Invasive species can have significant negative impacts on biodiversity (Mollot *et al.*, 2017), economy (Haubrock *et al.*, 2021; Bradshaw *et al.*, 2016), human health (Mazza *et al.*, 2014), and food security (Bruce, 2010; Paini *et al.*, 2016), with the total reported costs recently estimated at more than a trillion US dollars over the past few decades (Diagne *et al.*, 2021). From an evolutionary perspective, invasive species raise questions about how populations can rapidly colonize new environments and the extent to which their genetic makeup is critical. One hypothesis suggests that invasive success may be due to one or more *de novo* advantageous mutations (Colautti and Barrett, 2013), although the short evolutionary timescale of biological invasions makes this hypothesis less likely (Barrett and Schluter, 2008). Alternatively, invasive success may be due to various forms of pre-adaptation, enabled by standing genetic variation in the native range that allows the exploitation of a wide range of environments, including anthropized ones (Estoup *et al.*, 2016; Hufbauer *et al.*, 2012).

With the advent of next-generation genotyping and sequencing approaches, genome-environment association (GEA) studies have proven informative in unraveling the genetic mechanisms involved in local adaptation, including to invaded environments. In fact, by identifying genomic regions whose allele frequencies covary with environmental factors (Yang *et al.*, 2022; Hofmeister *et al.*, 2021; Ma *et al.*, 2020) or phenotypic traits (Turner *et al.*, 2021; Olazcuaga *et al.*, 2020; Pfenninger *et al.*, 2021), they can provide insights into the biological processes that enable genetic adaptation. More generally, since GEA models summarize the relationship between the (adaptive) genomic composition of populations and their environment, they can be used to predict maladaptation to novel environments. Fitzpatrick and Keller (2015) introduced the Genomic Offset statistic to quantify the gap between the optimal genetic configuration for a new environment (as predicted by the GEA model) and the actual genetic makeup of a population. Genomic Offset can therefore be interpreted as a possible risk of maladaptation (Capblancq *et al.*, 2020). Several studies have confirmed the correlation between Genomic Offset and average population fitness, using both computer simulations and common garden experiments (Gain *et al.*, 2023; Láruson *et al.*, 2020; Lotterhos, 2023; Rhoné *et al.*, 2020; Fitzpatrick *et al.*, 2021). However, Genomic Offset remains underutilized in invasion biology, despite its promise in predicting the success of invasive populations, as shown by a recent simulation study linking it to their probability of establishment (Camus *et al.*, 2024). Moreover, the concept of Genomic Offset, which implies that populations adapt to new environments through preexisting variants, is consistent with the hypothesis of (genetic) pre-adaptation to invaded environments. Following this idea, calculating the Genomic Offset between the source and invaded environments can reveal which geographic areas have posed or would pose the greatest adaptive challenges to any source population.

Using the fruit fly *Drosophila suzukii* as a biological model, the present study will illustrate the contribution of genome scan and GEA approaches to understanding the adaptation of invasive species.

Among these, Genomic Offset methods will provide insights into the past and future adaptive challenges of this emblematic pest. Native to East Asia, *Drosophila suzukii* was first recorded in North America (California) and southern Europe (Spain and Italy) in 2008 (Asplen *et al.*, 2015), with studies indicating independent invasion events on these continents (Fraimout *et al.*, 2017; Adrion *et al.*, 2014). By 2013, it had spread to South America (Deprá *et al.*, 2014) and is now widespread in the Americas and Europe. It has also been detected in various parts of Africa, such as Algeria (Aouari *et al.*, 2022), Morocco (Boughdad *et al.*, 2021), Kenya (Kwadha *et al.*, 2021), and most recently in South Africa (IPCC, 2024). By laying eggs in ripening fruits, *D. suzukii* has caused significant damage to crops, particularly berry crops (e.g. cherries or strawberries) in the various areas where it has invaded, making it a major agricultural pest (Walsh *et al.*, 2011).

To date, few studies have comprehensively investigated the adaptation of *D. suzukii* to invaded environments. To our knowledge, only one study by Feng *et al.* (2024) examined *D. suzukii* environmental adaptation using GEA, identifying genes associated with environmental variables such as temperature and precipitation. Another notable study by Olazcuaga *et al.* (2020) examined the broad genetic basis of *D. suzukii* invasion success and found evidence for only a few genes associated with population invasion status. Both studies were limited by their reliance on a limited number of sequenced pooled population samples (Pool-Seq) and a fragmented genome assembly derived from the sequencing of a strain of North American origin (Paris *et al.*, 2020). As *D. suzukii* research expands, large-scale whole genome sequencing (WGS) efforts include sequenced individual samples (Ind-Seq; e.g. Lewald *et al.*, 2021) in addition to the aforementioned Pool-Seq samples. However, the heterogeneity of the types and origins of the data still present practical challenges for their combined analysis.

To deepen our understanding of the genetic basis of past *D. suzukii* invasion and to further identify geographic regions at risk of future invasion, we present in this study the combined analysis of 27 Pool-Seq and 82 Ind-Seq samples, together representing a total of 37 populations. These data are primarily sourced from carefully curated publicly available datasets (Gautier, 2023), with the addition of newly sequenced samples from four previously unsampled areas. To enable efficient integration of these heterogeneous data for GEA and other genome scan approaches, we extended the Bayesian hierarchical models implemented in BAYPASS software (Gautier, 2015) for the joint analysis of hybrid data sets consisting here of read counts from Pool-Seq samples and genotype likelihoods (GL) for individuals subjected to Ind-Seq, for every marker. In addition, to improve the precision of genomic analyzes, we rely on a newly developed chromosome-level assembly for the genome of *D. suzukii* that we derived from an isofemale strain of Japanese origin (i.e., representative of the native range). Based on these new datasets, genomic resources, and methodological developments, we first searched for key genes and biological processes that may have been involved in the adaptive process during invasion by extending the approach described in Olazcuaga *et al.* (2020). To further

characterize the *D. suzukii* invasion and provide a global picture of its genetic underpinnings, we use Genomic Offset statistics based on a GEA with 27 covariables characterizing the environment of the 37 studied populations. We more specifically addressed the three following questions : i) which parts of the invaded areas of Europe and North America may have posed the greatest adaptive challenge to *D. suzukii*? ii) can we predict which parts of the geographic areas that have not yet been (fully) invaded (e.g. in South America and Africa) are at a higher risk of being invaded in the future ? and iii) can we anticipate which region could host the highest-risk source population for a putative invasive *D. suzukii* population establishing in Australia where this pest is currently absent ? These latter applications aims at illustrating how Genomic Offset statistics may represent an additional potentially valuable tool to be considered in Integrated Pest Management (IPM) practice. By providing a comprehensive view of the adaptive history of *D. suzukii*, we seek to describe a generic and operational population genomics framework to study and predict biological invasions between species.

Materials and Methods

Construction and annotation of a new *de novo* genome assembly

We generated PacBio HiFi and high-throughput chromosome conformation capture (HiC) sequencing data from individuals belonging to an inbred strains of Japanese origin (Ohashi *et al.*, 1991) to construct a male chromosome-level assembly, hereafter referred to as *dsu_isojap1.0*. Briefly, the primary whole-genome assembly was generated with `hifiasm` (v0.19.8; Cheng *et al.*, 2021), followed by contaminant removal with `Kraken2` (v2.1.2; Wood *et al.*, 2019) and self-alignment curation using `minimap2` (v2.20-r1061; Li, 2021). For more details on the data generation and assembly procedure, see Supplementary Note N1. The assembly was also visually compared with other available assemblies for closely related species (e.g. *D. biarmipes*) using dot plots generated with `dgenies` (v1.5.0; Cabanettes and Klopp, 2018), and its general properties were summarized with snail plots using the `BlobToolKit` viewer (v4.3.5; Challis *et al.*, 2020). These include the representation of contiguity and completeness which was evaluated with `BUSCO` on the `diptera_odb10` dataset (v5.4.7; Manni *et al.*, 2021). We finally annotated repetitive elements using `RepeatMasker` (v4.1.4; Smit *et al.*, 2023) run with options `-s -no_is -cutoff 200 -frag 20000`, the `wublast` search engine (option `-e wublast`) and the curated and annotated TE database developed by (Mérel *et al.*, 2021) for the *D. suzukii* genome. The assembly *dsu_isojap1.0* was then submitted to the NCBI repository for gene annotation with the NCBI Eukaryotic Genome Annotation Pipeline (Thibaud-Nissen *et al.*, 2013).

D. suzukii population samples and whole genome sequencing data

Whole-genome sequencing (WGS) data representative of 32 worldwide population samples, derived from either Pool-Seq (n=24 pools; Table S1) or Ind-Seq (n=73 individuals representative of 8 populations; Table S2) were obtained from the NCBI SRA repository (https://www.ncbi.nlm.nih.gov/sra/). These data were the result of a careful curation of publicly available WGS data from three recent studies (Olazcuaga *et al.*, 2020; Lewald *et al.*, 2021; Feng *et al.*, 2024), aimed at removing contaminated samples (Gautier, 2023) and excluding (or merging) geographically redundant samples (see Supplementary Note N2 for details). To further improve the worldwide representation of the samples, we newly sequenced four DNA pools representative of a single South American (AR-Gol from Argentina, n=100) and three European (CH-Del from Switzerland, n=58; DE-Jen from Germany, n=100; and RS-Zaj from Serbia, n=100) populations, as well as nine native individuals collected in Chengdu (China) in 2014. Note that DE-Jen individuals are from the same sampling used in Olazcuaga *et al.* (2020) and Feng *et al.* (2024), but with all non-*D. suzukii* individuals discarded (Gautier, 2023). Briefly, for sequencing these new samples, we extracted DNA from whole bodies (or pools of thoraxes) using the DNeasy Blood and Tissue Kit (Qiagen, USA) according to the manufacturer's recommendations and constructed libraries using the NEBNext Ultra II DNA Library prep kit for Illumina (New England Biolabs). These were paired-end (PE) sequenced (2x150 bp or 2x250 bp) on a NovaSeq6000 Illumina sequencer (Tables S1 and S2) on the Get Plage platform (INRAE, Toulouse).

WGS data processing, SNP calling and genotyping

WGS data were first filtered using `fastp` (v0.23.2; Chen *et al.*, 2018) run with default options to remove contaminating adapter sequences and trim poor quality bases (i.e., Q<15). Read pairs with either one read containing more than 40% low-quality bases or more than 5 N bases were removed. Filtered reads were then mapped onto the new *dsu_isojap1.0* assembly, complemented with the complete mitochondrial sequence of *D. suzukii* (GenBank Accession ID: *KU588141*) and the complete sequence for the Wolbachia endosymbiont wRI strain of *D. simulans* (NCBI ID: *GCA_000022285*), which is closely related to that of *D. suzukii* (Turelli *et al.*, 2018), using `bwa-mem2` with default options (v2.2.1; Vasimuddin *et al.*, 2019). PCR duplicates were then removed using the `view` and `markdup` programs of `SAMtools` (v1.14; Li *et al.*, 2009). The mean realized coverage was estimated using `mosdepth` (v0.3.6; Pedersen and Quinlan, 2017) with the option *-Q 20* to ignore reads with low mapping quality. As detailed in Tables S1 and S2, on autosomes (i.e. scaffolds 2L, 2R, 3 and 4) the coverages varied from 42.94 (US-CAOx) to 105.6 (JP-Kan) with a median of 61.3 for Pool-Seq data; and from 1.47 to 13.7 with a median of 6.98 for Ind-Seq data.

Variant calling was performed using the Haplotype Caller implemented in `FreeBayes` (v1.3.6; Garrison and Marth, 2012), which does not require realignment around indels and base recalibration,

jointly considering all 27 Pool-Seq and 82 Ind-Seq alignment files from *D. suzukii* samples, with the following options *-K -C 1 -F 0.01 -G 5 -E -1 –limit-coverage 500 -n 4 -m 30 -q 20*. For X-linked contigs, we also specified the ploidy of the 5 male individuals with the *-A* option. Variant calling was performed in parallel on a computer grid for the 1,179 nonoverlapping 250 kb windows covering the whole genome assembly. The resulting vcf files were reordered and concatenated per chromosome type using the `view` and `concat` programs of `bcftools` (v1.20; Danecek *et al.*, 2021), and filtered using a custom `awk` script to keep only bi-allelic variants (including the reference allele) with `QUAL>20, NS>100, 100<DP<5000, SAF>0, SAR>0, RPR>1, RPL>1, EPP>0, SRP>0`, and a global MAF computed from the overall read counts > 0.01. This yielded 6,814,770 autosomal bi-allelic variants consisting of 6,732,348 SNPs, 71 MNPs (Multi-Nucleotide Polymorphisms), 52,124 indels and 8,239 so-called complex variants. Likewise, 1,226,417 X-linked bi-allelic variants were obtained, including 1,209,454 SNPs, 22 MNPs, 15,343 indels and 1,598 complex variants. To provide a global visualization of the genetic structuring of the populations, we performed a random allele PCA using the `randomallele.pca` function of the R package `poolfstat` (v2.2.0 Gautier *et al.*, 2022) on the entire *vcf* combining Pool-Seq and Ind-Seq data parsed with the `vcf2pooldata` function.

Genome scan analyses using combined Pool-Seq and Ind-Seq data

We investigated the genetic basis of *D. suzukii* invasive success using the BAYPASS C_2 statistic (Olazcuaga *et al.*, 2020), which identifies overly differentiated markers between two population groups while accounting for population structure. All genome-wide scans were performed using an upgraded version (v3.0) of BAYPASS (Gautier, 2015; Olazcuaga *et al.*, 2020). As detailed in Supplementary Note N3, this new version extends all the previously described models to the analysis of individual genotyping data (coded as genotype likelihood, i.e. GL), which allows one to properly deal with low-to-mid-coverage Ind-Seq data (e.g., <10-15X) by integrating over genotype uncertainty. In addition, it now allows analyzing hybrid data consisting of sample allele and/or read counts and/or GL data under all the previously described models (Gautier, 2015; Olazcuaga *et al.*, 2020). Following the BAYPASS manual recommendation (see also Gautier *et al.*, 2018), the complete dataset was first split into subdata sets of 68,150 markers and 61,320 markers for autosomes and X-chromosome, respectively. The program was then run on each of these with the default option to both estimate the scaled covariance matrix of population allelic frequencies (Ω), which allows accounting for the neutral population structure in the different analyses, and three C_2 statistics (Olazcuaga *et al.*, 2020) contrasting native versus i) all worldwide (C_2^{WW}); ii) European (C_2^{EU}); or iii) American (C_2^{AM}) invasive populations. As observed in previous similar analyses (e.g., Gautier *et al.*, 2018; Olazcuaga *et al.*, 2020), we found a fairly high correlation of the posterior mean estimates of Ω entries over subdata sets (from the same chromosome type), and thus only considered those obtained from a single randomly chosen subdata set for representation and further analysis

under the `covmcmc` mode (see GO estimation, below). Likewise, C_2 estimates were highly consistent across three independent analyses (with different seeds) of each subdata set, so only results from the first analysis were reported.

To identify and delineate significant windows associated with invasive status based on the three different contrasts, i.e. with an excess of high C_2 values, we relied on the local-score approach described in Fariello *et al.* (2017) and (re)implemented in the R function `compute.local.scores` available in the BAYPASS software package (v3.0). More precisely, individual SNP scores were computed as $-\log 10(p) - \xi$, where p is the p-value associated with a given C_2 and ξ determines the a threshold parameter (Fariello *et al.*, 2017) that we set to $\xi = 2$ (i.e., $p < 10^{-2}$ were cumulated and others were scored 0 in the underlying Lindley process). As recommended in the BAYPASS manual, to avoid penalizing the approach, we discarded low-polymorphic SNPs (with an estimated overall allele frequency $\hat{\pi} < 0.2$ or $\hat{\pi} > 0.8$) from the local–score calculation.

For each significant window, the closest annotated gene located within 5 kb upstream or downstream of the C_2 peak was identified, and the *D. melanogaster* gene orthologs were retrieved using the `datasets` NCBI command-line tool (v16.40.0). Gene Ontology (GO) enrichment analysis for Biological Process was performed on the resulting list of candidate genes (from all three C_2) using the R package *clusterProfiler* (v4.12.0; Wu *et al.*, 2021) with default settings. GO terms were obtained from the `org.Dm.eg.db` annotation R package.

Estimation of the geometric Genomic Offset (gGO)

We first performed a GEA under the BAYPASS standard covariate model (Gautier, 2015) with option `-covmcmc`, using the matrix Ω estimated above, to estimate the effect (regression coefficients) of environmental covariables on the (scaled) population allele frequencies. This allows modeling the relationship between the genomic diversity and environmental variation of (all) the studied populations. The environment was here characterized with two data types. First, we considered the 19 commonly used bioclimatic variables, along with the mean monthly climate moisture index following Winkler *et al.* (2020) who showed the importance of humidity on fitness-related traits in *D. suzukii*, and mean monthly-near surface wind speed, suggested as one of the main driver for adaptation in *D. melanogaster* European populations (Bogaerts-Márquez *et al.*, 2021). Data for these 21 covariables, averaged from 1981 to 2010, were sourced from the Chelsa database (Karger *et al.*, 2017), and extracted using the R packages *dismo* (v1.3.5; Hijmans *et al.*, 2023) and *raster* (v3.5.15; Hijmans, 2022). We also included six 2017 land-use covariables from the ESRI Land Cover 2020 database (Karra *et al.*, 2021) processed as percentages for six of nine classes (crops, built area, trees, water, bareground and rangeland) with a resolution comparable to that of the Chelsa data. The remaining classes (ice/snow, clouds and flooded vegetation) were removed as uninformative (i.e no variation across the considered locations). Finally, to address variable interdependence, which can

reduce Genomic Offset's predictive performance of establishment probability (Camus *et al.*, 2024),

we performed a PCA on all the 27 covariables and retained the first 11 PCs (explaining $> 95\%$ of

the variance) as environmental covariables in the GEA analysis.

We used the estimated regression coefficients estimated with BAYPASS for both autosomes and X-

chromosome and computed the geometric GO (gGO; Gain *et al.* (2023)) using the `compute_genetic_offset`

function of BAYPASS v3.0 (Camus *et al.*, 2024; Gautier *et al.*, 2024). Note that no marker pre-

selection was performed, as suggested by recent studies (Gain *et al.*, 2023; Lind and Lotterhos, 2024;

Camus *et al.*, 2024). For graphical representation, following an approach similar to Lachmuth *et al.*

(2023), all gGO maps were uniformly scaled by the highest gGO value between all pairs of localities

represented by the 37 studied populations (observed for the IT-Tre/IR-Dub sample pair), i.e., the

maximum gGO value of established populations in the current worldwide *D. suzukii* range.

Results

A new male chromosome-level genome assembly

242 The combination of PacBio HiFi long reads and HiC sequencing data allowed the generation of a

243 highly contiguous assembly consisting of 54 scaffolds totaling 282,295,391 bp (N50=32.7 Mb) and

244 including 99.06% (98.48% complete and single) of the 3,285 BUSCO `diptera_odb10` dataset genes

245 (Manni *et al.*, 2021, see Table 1 and Figure S1A).

NCBI Assembly Name	CBGP_Dsuzu_IsoJpt1.0	Dsuz_RU_1.0	LBDM_Dsuz_2.1.pri
RefSeq ID	GCA_043229965.1	GCA_037355615.1	GCA_013340165.1
Reference	*this study*	unpublished	Paris *et al.* (2020)
Sample Origin	Japan	USA	USA
	males (IsoJpt strain)	females (WT10 strain)	females (WT3-2.0 strain)
Size in bp	282,295,391 (n=54)	178,754,567 (n=29)	268,012,156 (n=546)
N50 in bp (nb. seq.)	32,723,399 (n=3)	26,747,965 (n=4)	2,609,782 (n=16)
N90 in bp (nb. seq.)	2,973,411 (n=12)	3,161,696 (n=9)	228,192 (n=136)
nb. gaps (%N)	56 (0.00%)	0 (0.00%)	0 (0.00%)
Nb. (%) of BUSCO genes (diptera_odb10, n=3,285)			
Complete	3,254 (99.06%)	3,220 (98.02%)	3,200 (97.41%)
Compl. & Single	3,235 (98.48%)	3,204 (97.53%)	3,106 (94.55%)
Compl. & Duplicated	19 (0.578%)	16 (0.487%)	94 (2.86%)
Fragmented	12 (0.365%)	17 (0.518%)	49 (1.49%)
Missing	19 (0.578%)	48 (1.46%)	36 (1.10%)

Table 1: Descriptive statistics of the new *dsu_isojap1.0* assembly for the *D. suzukii* genome and the two other (partial) chromosome-level assemblies available from the NCBI repository as of March 2025. Aligning *dsu_isojap1.0* with the `RU_1.0` assembly (Figure S2) allowed the identification of 40 sequence similarity blocks > 1 Mb totaling 171.5 Mb (i.e. 96.1% of the `RU_1.0` assembly) with only 1.33% nucleotide divergence. Likewise, compared to the more fragmented `WT3-2.0` assembly, 35 blocks of sequence similarity > 1 Mb and totaling 153.6 Mb were found with a nucleotide divergence of 0.813%.

246 The alignment of the *dsu_isojap1.0* assembly with `Dbia2.1`, the chromosome-level reference assem-

247 bly of the closely related species *Drosophila biarmipes* (Figure S1B), showed that 6 large scaffolds

completely covered the 5 *D. biarmipes* autosome arms (2L, 2R, 3L, 3R and 4) and the X chromosome. Then, these *dsu_isojap1.0* scaffolds were named following the nomenclature of *D. biarmipes* (and *D. melanogaster*), as also justified by cytogenetic studies (Drosopoulou *et al.*, 2019). Likewise, as detailed in Table S3, a total of 3,197 genes of the `diptera_odb10` dataset were found complete and in single copy in the *dsu_isojap1.0* and the *Dmel6* assembly for the *D. melanogaster* genome (Hoskins *et al.*, 2015). The overlap in BUSCO gene content between the syntenic chromosome arms of the two species was almost complete with only a few discrepancies, aligning with the very close but not perfect conservation of Muller elements among *Drosophila* species (Bhutkar *et al.*, 2008). More specifically, an overlap of 99.4%, 99.9% and 100% was found for chromosomes X, 3 and 4 respectively. The overlap was a bit lower for chromosome arms 2L and 2R (95.8% and 96.0%) respectively, suggesting a less complete sequence. However, an additional large scaffold (40.07 Mb) containing only 19 BUSCO genes, 8 mapping to the 2L and 11 to the 2R *D. melanogaster* chromosome arms, likely corresponded to the (extended) centromeric regions of chromosome 2 (as confirmed by their repeat content, see below and Figure S1) and was thus named *2c*. Similarly, the X scaffold (32.72 Mb in length) actually included seven of the 8 contigs assigned to the X chromosome (Supplementary Note N1), the remaining eighth (6.316 Mb) was renamed *scf_Xc* as it was probably more centromeric (see below and Figure S1C). In contrast, none of the three contigs (total 16.74 Mb) assigned to the Y chromosome could be scaffolded. The largest (13.21 Mb) was then named Y, while the other two were manually joined into an 3.338 Mb Y-linked scaffold (*scf_Y1*) based on alignment of the Occludin-Related Y (*Ory*) *D. melanogaster* gene. Note that no BUSCO genes in common with the *Dmel6* assembly mapped to the Y or `scf_Xc` contigs (Table S3).

In agreement with previous studies on the WT3_v2.0 assembly (Paris *et al.*, 2020; Mérel *et al.*, 2021, e.g.), we found that *dsu_isojap1.0* was characterized by a very high TE content (53.24% of the assembly). The count distribution of the different TE orders over the scaffolds highlighted distinct chromosomal regions (Figure S1C). In particular, the center of chr3, one of the two extremities of 2L, X and 2R, and the entire chromosomes 4, Y and scf_Y1, scf_2c (93.7%) and scf_Xc scaffolds lacked simple repeats and were enriched in Long Terminal Repeat retrotransposons and Rolling Circle transposons. This suggests that such TE signature specifies centromeric regions in the *D. suzukii* genome, as shown by Chang *et al.* (2019) in *D. melanogaster*, although different TE superfamilies are involved. Finally, the gene annotation pipeline identified a total of 17,936 genes (including 15,749 coding and 2,219 noncoding) and 266 nontranscribed pseudogenes. Most annotated genes were mapped to the identified autosomal scaffolds, with 75.6% located on chromosomes 2L, 2R, 3, and 4, increasing to 79.3% when including scf_2c (Figure S3A). Similarly, 14.4% of genes were mapped to the X-linked scaffold (chrX), rising to 14.5% with the addition of scf_Xc. A total of 404 genes (2.25%) were annotated on the Y-linked scaffolds (chrY and scf_Y1), while 712 genes (3.97%) were assigned to the remaining small scaffolds. Furthermore, along the main chromosomes, the gene

density was lower in the centromeric regions (e.g., middle of chr3), which were identified based on their TE content, as well as in the other scaffolds, including scf_2c and scf_Xc (Figure S3B). Overall, our new annotated chromosome-level *dsu_isojap1.0* assembly represented a clear improvement compared to previous assemblies, both in terms of contiguity and completeness, which we could rely on to perform further population genomics analyzes.

A combined Ind-Seq and Pool-Seq curated dataset representing the *D. suzukii* worldwide genetic diversity

To visualize the global genetic structuring of the 37 studied populations, we first performed a random allele PCA focusing on autosomal variants. As shown in Figure 1B and in agreement with previous studies (e.g. Fraimout *et al.*, 2017; Lewald *et al.*, 2021), the first factorial plan grouped the populations into four main clusters: i) the native area; ii) European invaded area; iii) populations from the Northwest US and Hawaii; and iv) all other populations from the American invaded area (including the newly generated AR-Gol from Argentina). A more detailed inspection showed that within the native area cluster (i.e., within Asia), Japanese populations tended to cluster together, as did Chinese populations.

Within the European cluster, which includes FR-Run from La Réunion Island in the Indian Ocean due to its recent European origin (Fraimout *et al.*, 2017), PC1 could only loosely be associated with a north-south geographical gradient. In fact, the northernmost IR-Dub (sampled in Ireland) was closer to the Mediterranean populations (ES-Bar, IT-Tre and FR-Cor), while PC1 (and PC2) isolated the newly generated sample from Serbia (RS-Zaj), possibly due to its eastern position or more recent sampling. In contrast, the proximity of CH-Del to DE-Dos matched its geographic origin. Finally, FR-Lez (from southern France) was more closely related to German (DE-Jen and DE-Dos) than to Mediterranean populations. In general, the positioning of the populations of the invaded European area did not clearly reflect geographical or temporal origins, suggesting migration across Europe and possible heterogeneous introgression with propagules from the North American continent (Fraimout *et al.*, 2017).

For the American invaded area, differentiation between eastern and northwestern populations matched previous findings (Lewald *et al.*, 2021). The proximity of US-Haw, US-Wa, and US-Sok, which form a distinct cluster, is consistent with evidence of gene flow from Hawaii to Northwestern America (Fraimout *et al.*, 2017; Gautier *et al.*, 2022). Five eastern (US-Vir, US-Me, US-Ga, US-Nca, US-Ny) and three central US populations (US-Mi, US-Col, US-Wis) clustered together. South Californian and South American populations also clustered with this group, but more closely with each other, in agreement with the previously inferred origin of the South American populations (Gautier *et al.*, 2022). Interestingly, US-Wat was positioned between the Northwestern and Eastern/Central/Southwestern American clusters, consistent with its intermediate geographic origin.

Figure 1: Description of the 37 studied *D. suzukii* population samples and structuring of genetic diversity on autosomal variants. A) Geographical location of the 37 *D. suzukii* populations. Populations from the native area are displayed in green, while invasive American and European populations are displayed in red and blue, respectively. B) Random allele PCA representation of the population samples. Pool-Seq samples are represented by black dots, while Ind-Seq samples are indicated by crosses colored according to their population of origin (with 95% ellipses summarizing the overall dispersion of the underlying individuals around their population center of gravity). C) Correlation plot of the scaled covariance matrix Ω estimated from the combined analysis of Pool-Seq and Ind-Seq data with the new version of BAYPASS software (v3.0; Gautier, 2015) on autosomal variants. The name of the populations represented by Ind-Seq samples are highlighted in bold.

The correlation plot of the Ω matrix estimated with BAYPASS showed a similar structuring into four groups (Figure 1C). Reassuringly, this Ω representation did not cluster samples by sequencing type (Pool-Seq or Ind-Seq), suggesting that the statistical models implemented in the new BAYPASS version perform as expected on hybrid datasets.

Both the random allele PCA and the Ω matrix estimation were independently repeated with X–linked variants, yielding results similar to autosomal variants (Figure S4). Despite minor differences (see Figure S4 legend), the overall clustering patterns remain consistent and robust, providing a clear description of population structure over the whole set of curated Pool-Seq and Ind-Seq samples.

Genome scan for association with invasive status

Cumulating association signal of markers with a C_2 p-value below 0.01 using a local-score, we could identify 231 significant windows for C_2^{WW}, 208 windows for C_2^{EU}, and 18 windows for C_2^{AM} (Figure 2), predominantly located on chromosomes 3 (n=195) and 2R (n=108), followed by the X chromosome (n=74), chromosome 2L (n=69), and chromosome 4 (n=11) (see Tables S4 to S8 for the detail of all the windows for each chromosome).

Figure 2: Manhattan plot of the C_2 p-values based local-score contrasting native versus A) all worldwide (C_2^{WW}); B) European (C_2^{EU}); and or C) American (C_2^{AM}) invasive populations. The dashed lines indicate the chromosome-specific 1% significance threshold for the local-score. The number of significant local-score windows is given in the main legend of each plot. Details about all the windows (separated by chromosome) can be found in Tables S4 to S8, and the top 45 genes associated with the highest local-score values are presented in Table 2.

The size of the identified windows ranged from 155 bp to 476,822 bp and included 7 to 202 SNPs (Tables S4 to S8). Merging the windows that overlapped between different C_2 tests resulted in a final set of 457 candidate regions: 171 specific to C_2^{WW} specific; 154 specific to C_2^{EU}; and 10 specific to C_2^{AM}. The windows detected only with C_2^{EU} or C_2^{AM} likely reflect continent-specific signatures, otherwise they would also have been detected by the C_2^{WW} test, its larger sample size being expected to increase power. Conversely, our ability to assess convergence between adaptation signatures in Europe and America may be limited by the lower power of the C_2^{EU} and even more the C_2^{AM} tests, which both detected fewer significant regions than C_2^{WW}. Hence, the 171 C_2^{WW}–specific regions might result from a combination of signals that are consistent between continents but too weak to be detected in any of them. However, 52 windows were shared by C_2^{WW} and C_2^{EU}; 6 by C_2^{WW} and

C_2^{AM}; and only 2 by the three contrasts, which remains higher than the expected number (n = 0.54) if all tests were completely independent (see Supplementary Note N4).

Gene annotation of the 457 significant windows highlighted 396 positional candidate genes among which 322 could be associated with a *D. melanogaster* ortholog (Tables S4 to S8). The GO enrichment analysis for Biological Process (for all candidate genes with orthologs) showed several terms that were significantly enriched in our candidate genes dataset compared to the rest of the genome (Figure S6). The GO terms with the most significant enrichment were "cell adhesion", "epithelial morphogenesis", and "dicarboxylic acid transport". Further down the list, significant terms were mostly related to morphogenesis but also included cognition, chemosensory and olfactory behavior, and memory.

More specifically, the window at the 2L centromeric end associated with the *Histone H1-like* candidate gene clearly stood out, since it was among the two windows shared by the three contrasts and displayed the highest local-score in each test. The three corresponding windows were also unusually large (476,822 bp, 440,361 bp and 356,304 bp for C_2^{WW}, C_2^{EU} and C_2^{AM} respectively), although inflated by an extended region of extremely low SNP density (4 SNPs spanning ca. 300 Mb) beyond the local-score peak (Figure S7). Nevertheless, the local-score peak was located approximately just 1kb upstream of the *Histone H1-like* gene, confirming a strong signal associated with it. Histones play a crucial role in regulating chromatin structure and gene expression, and Histone H1, the least conserved histone, is essential for heterochromatin silencing and genome stability in *D. melanogaster* (Bayona-Feliu *et al.*, 2016). Although its study has gained traction recently, its full range of functions remains unclear and is likely to include multiple variants with distinct functions (Prendergast and Reinberg, 2021). It should also be noted that we here identified an "H1-like" gene, rather than a canonical Histone H1, which may thus deserve further confirmatory functional characterization in *D. suzukii*.

Among our 45 top candidates (mainly related to C_2^{EU} and C_2^{WW}, Table 2), we also identified a diverse set of genes related to key physiological and behavioral traits, including resistance to stress and detoxification. For instance, in *D. melanogaster*, *CG30015* is likely involved in heavy metal susceptibility (Zhou *et al.*, 2017), *Olf413* has been identified as a candidate for the toxicological response to monoterpene pesticides (Sabio *et al.*, 2024), and *Prosalpha2* likely plays a role in a genetic resistance network to pesticides (Zhang and Zhang, 2019). Additionally, both *ChChd2* and *ClpX* may contribute to the protection against oxidative stress (Meng *et al.*, 2017; Hill *et al.*, 2018). Together, these genes may support adaptive responses to environmental stressors and anthropogenic pressures, including exposure to pesticides, potentially facilitating the establishment of *D. suzukii* in agricultural landscapes.

Several genes related to dispersal and movement were also among the top 45 candidates, such as i) *CG34215* found associated with flight performance in *D. melanogaster* (Spierer *et al.*, 2021); ii)

Gene	Chr	Position in bp	Ortholog	max. Lscore $(-\log_{10} \mathrm{P}_{C2})$	Best C_2	Other C_2: Lscore $(-\log_{10} \mathrm{P}_{C2})$
histone H1-like	2L	25,717,391-25,718,121	-	88.9 (10.1)	C_2^{EU}	C_2^{WW}: 56.6 (5.81) C_2^{AM}: 22.2 (4.16)
Myo81F	3	48,614,974-51,239,613	*Myo81F*	38.0 (6.48)	C_2^{EU}	C_2^{WW}: 17.9 (4.65)
LOC108009451	2R	7,413,548-7,455,825	*CG30015*	34.6 (5.83)	C_2^{EU}	C_2^{WW}: 16.1 (5.58) C_2^{EU}: 23.5 (5.30)
Fancm	3	66,612,689-66,617,774	*Fancm*	27.4 (5.18)	C_2^{WW}	C_2^{EU}: 16.5 (4.20)
Tsp42Ec	2R	17,579,181-17,581,562	*Tsp42Ec*	26.7 (6.45)	C_2^{EU}	-
LOC108011780	3	66,148,636-66,152,754	-	26.7 (5.08)	C_2^{WW}	-
olf413	3	25,100,604-25,210,799	*olf413*	23.9 (7.71)	C_2^{EU}	C_2^{WW}: 9.20 (4.31)
ppk11	2L	18,914,671-18,917,502	*ppk11*	22.9 (5.64)	C_2^{WW}	C_2^{EU}: 6.61 (3.97)
LOC108018145	2R	17,284,668-17,409,796	*CG34215*	21.5 (4.63)	C_2^{EU}	C_2^{EU}: 21.0 (3.99) C_2^{EU}: 15.3 (4.74)
meigo	3	67,030,298-67,031,912	*meigo*	20.8 (5.75)	C_2^{WW}	-
LOC118876994	2R	22,793,692-22,795,240	-	20.7 (8.47)	C_2^{WW}	C_2^{EU}: 19.1 (8.76)
LOC108012509	3	5,481,114-5,699,654	-	20.2 (5.10)	C_2^{EU}	C_2^{WW}: 11.8 (4.52)
CheB42c	2R	17,520,645-17,521,480	*CheB42c*	20.1 (5.74)	C_2^{EU}	-
stum	2R	22,603,541-22,616,256	*stum*	20.1 (5.14)	C_2^{EU}	C_2^{WW}: 6.63 (3.86)
zfh2	4	1,323,039-1,382,835	*zfh2*	19.6 (5.92)	C_2^{EU}	-
Chchd2	X	21,071,650-21,073,530	*Chchd2*	19.3 (6.41)	C_2^{EU}	-
LOC108017765	2R	7,370,854-7,374,343	-	18.5 (8.81)	C_2^{WW}	C_2^{EU}: 10.3 (4.33) C_2^{AM}: 7.06 (4.26)
CLIP-190	2L	12,713,010-12,740,766	*CLIP-190*	18.3 (5.60)	C_2^{WW}	-
dpy	2L	7,226,343-7,361,390	*dpy*	18.0 (4.46)	C_2^{EU}	C_2^{EU}: 6.97 (3.43)
Hmgcr	3	87,507,123-87,527,598	*Hmgcr*	17.9 (3.88)	C_2^{EU}	-
Oamb	3	70,820,820-70,851,333	*Oamb*	17.8 (4.72)	C_2^{WW}	C_2^{AM}: 16.3 (4.37)
PolQ	3	67,647,198-67,654,301	*PolQ*	17.5 (7.10)	C_2^{WW}	C_2^{AM}: 7.54 (4.48)
LOC136117410	X	30,121,354-30,126,274	-	17.4 (4.77)	C_2^{WW}	-
Prosalpha2	3	67,376,995-67,378,199	*Prosalpha2*	17.2 (5.45)	C_2^{WW}	-
RpL26	3	23,059,597-23,225,527	*RpL26*	16.7 (5.98)	C_2^{WW}	C_2^{EU}: 12.6 (5.05)
LOC108009111	2R	23,856,122-23,858,356	*CG15861*	16.4 (4.78)	C_2^{WW}	-
smash	3	62,145,253-62,208,229	*smash*	16.4 (5.32)	C_2^{WW}	-
psq	2R	7,697,594-7,753,927	*psq*	16.3 (5.09)	C_2^{EU}	-
LOC108021082	2L	6,713,709-6,813,712	-	16.0 (5.63)	C_2^{WW}	C_2^{AM}: 6.55(4.40)
LOC108006723	3	63,351,954-63,357,438	*CG31551*	15.6 (4.06)	C_2^{EU}	-
ClpX	3	69,004,325-69,009,580	*ClpX*	15.5 (7.76)	C_2^{EU}	C_2^{AM}: 7.45 (4.93)
LOC108005385	X	20,678,509-20,728,641	*CG13003*	15.3 (4.78)	C_2^{WW}	-
LOC108010813	2L	1,423,888-1,424,584	-	15.2 (6.37)	C_2^{EU}	-
LOC108017397	3	77,334,337-77,338,474	-	15.2 (4.47)	C_2^{WW}	C_2^{EU}: 13.9 (4.68)
SNF4Aγ	3	66,954,451-67,026,359	*SNF4Aγ*	15.0 (6.95)	C_2^{WW}	C_2^{EU}: 12.9 (6.85) C_2^{WW}: 9.61 (6.12) C_2^{EU}: 7.24 (5.45)
Smyd4-3	2R	8,962,627-8,965,112	*Smyd4-3*	14.9 (5.26)	C_2^{WW}	-
LOC108010319	2R	15,504,095-15,509,232	*CG15673*	14.3 (5.45)	C_2^{EU}	C_2^{WW}: 11.7 (4.71)
Ptr	2R	2,129,406-2,145,743	*Ptr*	14.2 (5.02)	C_2^{WW}	C_2^{EU}: 6.66 (4.09)
LOC139352732	3	31,534,547-32,058,847	*CG40470*	14.1 (3.72)	C_2^{EU}	C_2^{WW}: 8.82 (4.22)
LOC108009708	2R	14,107,953-14,113,451	*CG4294*	14.1 (5.77)	C_2^{EU}	C_2^{WW}: 7.00 (4.66)
Obp56h	2R	12,909,224-13,006,256	*Obp56h*	13.6 (5.73)	C_2^{WW}	C_2^{WW}: 9.26 (5.37)
LOC108004869	3	63,498,822-63,528,654	*CG2082*	13.5 (4.33)	C_2^{WW}	-
LOC108007632	3	81,317,724-81,338,568	*CG33203*	13.4 (4.26)	C_2^{WW}	-
Prx2	X	20,378,533-20,380,802	*Prx2*	13.4 (6.07)	C_2^{EU}	-
Gr57a	2R	8,937,103-8,939,349	*Gr57a*	13.3 (5.33)	C_2^{EU}	-

Table 2: Description of the top 45 *D. suzukii* windows with the highest C_2 local-score among the 396 (out of 457) with annotated positional candidate genes. The table gives for each gene i) its position in the new *dsu_isojap1.0* assembly of the *D. suzukii* genome; ii) its corresponding *D. melanogaster* ortholog; (iii) the highest maximum Lindley score (Lscore) and the most significant C_2 P-values (in $-\log_{10}$ scale) for the associated window; iv) the C_2 statistic corresponding to this highest Lscore; and v) other C_2 statistics with which the gene was associated, along with the maximum Lscore and most significant C_2 P-values (in $-\log_{10}$ scale) for the corresponding window.

Oamb which influences locomotion, climbing ability, and exploratory behaviors (El-Kholy *et al.*, 2021); iii) *Hmgcr* involved in sexual dimorphism of locomotor behavior (Belgacem and Martin, 2007); and iv) *Ptr* involved in neuromuscular function during larval development and likely adult lifespan regulation (Parada and Prieto, 2024). Likewise, some of the top candidate genes involved in chemosensation and feeding behavior, such as i) *Cheb42c* which is implicated in pheromone perception (Xu *et al.*, 2002); ii) *Gr57a*, a gustatory receptor found associated with variations in olfactory perception in *D. melanogaster* (Arya *et al.*, 2015); iii) *Obp56h*, which is linked to mating and feeding behavior (Shorter *et al.*, 2016; Delclos *et al.*, 2024; Swarup *et al.*, 2014) and is differentially expressed between summer and winter morphs in *D. suzukii*, suggesting a role in seasonal feeding adjustments (Schwanitz *et al.*, 2022); and iv) *Ppk11* that influences salt taste perception (Liu *et al.*, 2003).

We also identified candidate genes related to immune response, such as i) *CG34215* found to be differentially expressed in *D. suzukii* following infection by the entomopathogenic nematode *Steinernema carpocapsae* (Garriga *et al.*, 2024); and ii) *Psq*, identified as a potential target of *Wolbachia*-induced cytoplasmic incompatibility in *D. melanogaster* (Zheng *et al.*, 2019). Finally, the *CG30015* and *Oamb* candidate genes have been found involved in social behaviors associated with aggression in *D. melanogaster* (Edwards *et al.*, 2006; Luo *et al.*, 2014).

It is worth stressing that most of the genes mentioned above have been found in candidate regions where there are not any or very few other genes. Therefore, there is good statistical evidence that these genes have been the drivers of the selection signal that has been detected in these regions.

Insights into past invasions from gGO

To determine which parts of the invaded areas of Europe and North America posed the greatest adaptive challenge to *D. suzukii*, we computed gGO maps over these two continents by estimating, for each grid cell, the gGO values relative to a reference environment representative of the source populations. For the European continent, for which a single original invasive source population has been previously inferred (Fraimout *et al.*, 2017), we relied on F_3 statistics (Patterson *et al.*, 2012; Gautier *et al.*, 2022) to identify CN-Lia as the closest source population proxy from our data (see Figure S8 for details). The CN-Lia environment was then used as a reference environment to compute the gGO map over Europe (Figure 3A), which generally displays very low gGO, except in a few areas, such as the Alps and the cold regions of north-west Scotland and western Norway, with high or extreme values.

For North America, the reference source environment is more difficult to define, due to the admixed origins of the primary invading populations (Fraimout *et al.*, 2017; Gautier *et al.*, 2022). Interpreting gGO as an environmental distance weighted by the adaptive genome composition (Gain *et al.*, 2023), we proposed to calculate weighted gGO values by combining the two gGO maps obtained using the environments of CN-Nin and US-Haw, respectively, as references (Figure S9), since these

populations serve as the main contributing source proxies for admixture in the invading population (Fraimout *et al.*, 2017). The weights were 75% for the CN-Nin-based and 25% for the US-Haw-based gGO maps, reflecting the estimated proportions of admixture of the primo-invading US population (Fraimout *et al.*, 2017). The resulting estimated gGOs were more heterogeneous across North America (Figure 3B) than in Europe, with high values in northern Canada (colder regions) and medium values in the arid regions of the southwest US (e.g., Arizona, Nevada, Utah, New Mexico). Extreme values were found mainly in the northern cold regions and the central Sonoran Desert. The lowest gGO values were mostly found along the east coast, parts of the west coast, extreme south-east US, and in Central America. However, note that the weighting of gGO may have strongly impacted the results, since the CN-Nin-based and US-Haw-based gGO maps were different (Figure S9), the latter displaying in particular lower values in Central America and western USA.

Together, these results suggest that the adaptive challenge for *D. suzukii* populations to invade Europe and North America was rather limited.

Figure 3: gGO values over European (A) and North American (B) invaded areas. For Europe, the reference environment corresponded to the one of the CN-Lia population, inferred as the closest proxy (see Figure S8). For North America, to account for the admixed origin of the North America invasive populations (Fraimout *et al.*, 2017; Gautier *et al.*, 2022), we computed weighted gGO values by combining the two gGO maps (Figure S9) based on the environments of CN-Nin and US-Haw as references. The combined map used weights of 75% for CN-Nin and 25% for US-Haw, corresponding to their respective estimated admixture proportions in the primo-invading US population (Fraimout *et al.*, 2017). Lowest (highest) gGO values correspond to lowest (strongest) adaptive challenges. Gray pixels represent outlying gGO values (as defined in M&M).

Predicting future invasions from gGO

Following an approach similar to that above, we relied on gGO to predict which uninvaded geographical regions in Africa, South America, and Australia are at higher risk of invasion in the future, by deriving gGO maps over these three continents with respect to a given source environment. For Africa and South America, we selected source environments from representative sites where *D. suzukii* populations have recently established, and thus represent potential sources capable of spreading across these continents. These correspond to Longonot Farm (Nakuru county, Kenya; Kwadha *et al.*, 2021) and Nova Veneza (Santa Catarina, Brazil; Deprá *et al.*, 2014), for Africa

and South America, respectively. For Australia, where *D. suzukii* is not established, we selected the environment of our US-Wa sample as a putative source reference, based on fresh fruit trade statistics (International Trade Center, 2024).

Figure 4: gGO estimates for the geographical regions in Africa (A), South America (B) and Australia (C) that have not been (fully) invaded to date. For Africa and South America, the black arrow indicates the (chosen) source reference environment where *D. suzukii* was first recorded and recently described as established in these continents (i.e. the environment of Longonot Farm - Nakuru county, Kenya - for Africa and of Nova Veneza - Santa Catarina, Brazil - for South America). For Australia, where *D. suzukii* has not yet been found as established, the reference environment associated with our Northwestern American sample US-Wa was used as a (putative) source reference based on trade statistics for fresh fruit exchanges (International Trade Center, 2024). Gray pixels represent outlying gGO values (as defined in M&M).

The gGO map of Africa displayed contrasting patterns (Figure 4A), with arid regions (e.g., the Sahara and Namib deserts) predominantly showing outlying or very high values at their borders, and other regions showing moderate values (e.g., sub-Saharian area). In contrast, some parts of West Africa (e.g., Ivory Coast), Central Africa (e.g., Democratic Republic of the Congo), and East Africa (e.g., the Great Lakes region) displayed low gGO values, i.e., higher invasion risk. Interestingly, the very northern part of the continent, especially in northwestern Morocco, where *D. suzukii* infestation has recently been reported in several berry farms (Boughdad *et al.*, 2021), was among the regions with the lowest values. The north-eastern region of Algeria, including the pomegranate agroecosystem where *D. suzukii* has been detected (Aouari *et al.*, 2022), also showed low gGO values, but with a more heterogeneous pattern. In South America, most of the subcontinent (e.g., the Amazon Basin, Brazilian plateau, northeastern Argentina, and Bolivia) showed very low to moderate gGO values. As expected, the high or extreme outlying values were mainly located in regions with extreme climates, such as the Andes mountains or the west coast and southern parts of Patagonia (Figure 4B). It should be noted that the Chubut region in central Patagonia (Argentina) where *D. suzukii* has only recently been detected and considered established, and from which the AR-Gol sample originates, showed moderate to high gGO values. Finally, the gGO map of Australia displayed a moderate variability (Figure 4C). The highest GO values were located in arid areas (e.g Simpson Desert), while the lowest gGO values were concentrated in the southern regions, particularly along the South

East Coast, and in Tasmania.

For continents that remain uninvaded, such as Australia, estimating gGO can also serve as a useful tool to identify potential source areas posing the highest risk of contributing optimally preadapted invasive *D. suzukii* populations. To illustrate this idea, we note that the two largest exporters of fresh berries to Australia in 2023, after New Zealand and Vietnam (where *D. suzukii* is considered absent), were the USA and Thailand (International Trade Center, 2024). Using Sydney Harbor as a reference environment, since it is an obvious potential sensitive introduction point for pest species in Australia, we derived gGO maps for North America (Figure 5A) and Southeast Asia (Figure 5B). Thailand consistently showed moderate gGO values, with no area of very high or extreme values, and was globally characterized by lower gGO values than North America. Interestingly, in North America, the region with the lowest gGO values was mainly in northeast US (e.g., Great Lake region), which does not produce the *D. suzukii* host plant fruits imported into Australia.

Figure 5: gGO map to assess the areas most at risk as sources of preadapted *D. suzukii* propagules with respect to Australia. gGO values were estimated over areas corresponding to that of the two largest exporters of fresh berries to Australia in 2023 (after New Zealand and Vietnam where *D. suzukii* is absent): North America (A) and Thailand (B, with national borders shown in black). For Australia, the reference environment is that of Sidney's harbor, an obvious potential sensitive introduction point of pest species in Australia. Gray pixels represent outlying gGO values (as defined in M&M).

Discussion

The primary goal of this study was to describe and empirically evaluate a population genomics framework to assess genomic adaptation during biological invasions, using the invasive species *D. suzukii* as a biological model. Leveraging recent studies, we expanded the global representation of *D. suzukii* populations by incorporating publicly available Ind-Seq (Lewald *et al.*, 2021) and Pool-Seq data (Olazcuaga *et al.*, 2020; Feng *et al.*, 2024), along with some newly sequenced samples.

Joint analysis of such heterogeneous data sets raises statistical challenges and can lead to strong technical biases. The random allele PCA originally proposed by Skoglund and Jakobsson (2011) to

deal with ancient DNA sequencing data and implemented in `poolfstat` (Gautier *et al.*, 2022), has proven effective in accounting for sample heterogeneity when characterizing the global structuring of genetic diversity. However, the loss of information resulting from the use of a single read in each (Pool-Seq or Ind-Seq) sample at each genomic position makes it suboptimal for detecting local genomic effects related to adaptive differentiation or association with environmental covariates. To the best of our knowledge, no current population genomics method fully exploits heterogeneous datasets —comprising both Ind-Seq and Pool-Seq data— which become more available across species with the growing use of sequencing technologies and the encouraging shift toward open science. Using Bayesian hierarchical modeling, we developed an extension to the BayPass software (Gautier, 2015) that enables the joint analysis of Ind-Seq and Pool-Seq data and further allows modeling Ind-Seq data using either allele counts or genotype likelihoods (GLs). Although the former approach is sufficient for high-coverage data (e.g., >20X) where individual genotypes can be reliably called beforehand, the latter properly integrates over genotype uncertainty associated with low to medium coverage data (ie, $< 15X$), which is recommended in a variety of population genomics methods (e.g., Kim *et al.*, 2011; Skotte *et al.*, 2013; Korneliussen *et al.*, 2014; Druet and Gautier, 2017; Meisner and Albrechtsen, 2018).

Using our expanded and carefully curated hybrid dataset, we could then search for key genomic regions that may be associated with the invasive success of *D. suzukii* populations using the C_2 contrast statistic (Olazcuaga *et al.*, 2020). In order to improve the accuracy of our genomic analyses, we initially generated a new chromosome-level and annotated assembly for the *D. suzukii* genome from males (to sequence the Y chromosome) of a Japanese strain established in the 1980's (Ohashi *et al.*, 1991), i.e., before the *D. suzukii* invasion outside Asia. This contrasts with previous publicly available assemblies, which were obtained from females of isofemale lines representative of the European or North American invaded areas (e.g. Ometto *et al.*, 2013; Chiu *et al.*, 2013; Paris *et al.*, 2020). The new *dsu_isojap1.0* assembly can thus be considered as more representative of a *D. suzukii* "ancestral" genome for the native area (disregarding subsequent genetic changes that have occurred in the strain) to identify *de novo* mutation in invasive populations, especially regarding structural variation (including TE content), which we did not examine here. Besides, it provides a critical improvement over available assemblies, and may thus represent a valuable reference for other genomic studies on *D. suzukii*.

For our purposes, the close to optimal contiguity of the *dsu_isojap1.0* assembly enabled a better use of the local-score approach of Fariello *et al.* (2017), which accounts for LD information by combining C_2 statistics at neighboring SNPs at a whole chromosome scale. This both increased the support and facilitated the delineation of genomic regions associated with the invasive status of populations. In total, we identified 457 significant local-score windows, which were generally small (that is, mainly < 20 kb), in agreement with the low level of SNP signal clustering already observed in Olazcuaga *et al.*

(2020). Furthermore, in accordance with the notably distinct subsets of SNPs identified as associated based on C_2^{EU} and C_2^{AM} in Olazcuaga *et al.* (2020), the resulting continent-specific windows exhibited a negligible degree of overlap. Likewise, several highly significant C_2^{EU}–specific windows (especially on chromosome 2R) disappeared with C_2^{WW}, i.e., when adding American samples to the test. This suggests a low degree of convergence between the adaptation processes observed in the two (almost) independently invaded areas by *D. suzukii* (i.e. Europe and America).

Surprisingly, the annotation of the genomic windows identified in the present study did not reveal any candidate genes in common with those identified by Olazcuaga *et al.* (2020). This discrepancy may be related to the increased power and robustness provided by our larger dataset and the local-score method, which led us to detect signals not captured by Olazcuaga *et al.* (2020) or, conversely, to discard genes from that previous study that were supported by only a few variants. In addition, we here relied on a much more accurate genome assembly along with an improved variant calling (and genotyping) procedure implemented in `FreeBayes` (Garrison and Marth, 2012), a haplotype caller that relies on local re-alignment of reads. In particular, this may explain why we could not confirm the strong association signal of the three neighboring SNPs upstream of the *cpo* gene, as these were actually located in a low complexity region of the 3R chromosome that was poorly assembled and encompassed several highly complex variants in the previous reference assembly in Paris *et al.* (2020). Although *cpo*, which seemed at first sight particularly attractive as a functional candidate due to its role in the control of diapause (Olazcuaga *et al.*, 2020) was not confirmed, we identified several strong and even more conclusive signals encompassing genes potentially involved in other interesting functions underlying invasion success. Among these functions are detoxification, stress response, and pesticide resistance, which can be advantageous traits. For instance, they may enhance tolerance to pesticides, a factor to which *D. suzukii* is regularly exposed and can rapidly develop resistance, as evidenced by laboratory assessments (Haye *et al.*, 2016; Deans and Hutchison, 2022). In addition, we identified genes related to gustatory and olfactory behaviors, potentially underpinning *D. suzukii*'s generalist feeding strategies by increasing its ability to detect and utilize a variety of host fruits in the environments it has invaded. We also discovered genes associated with immune responses that may be important, as interaction with new pathogens in newly invaded environments can impede or promote invasion success, depending on whether the allocation of resources to immune function detracts from other vital traits or improves adaptive survival in these new habitats. However, association analyses such as the one performed here based on the C_2 are expected to provide a very partial view of the genetic basis of the invasive success of a species, as the underlying traits are expected to be multiple and highly polygenic. Association analyses thus primarily aim at identifying a few important genes, such as the few candidates discussed above, that may deserve special interest for future validation studies to better characterize the traits they control.

To provide a global picture of the (possible) role of genomic adaptation in the invasive success of *D. suzukii*, we considered an alternative approach based on modeling the relationship between whole-genome genomic variation and environmental variables among populations, building on recent work on estimating population maladaptation using the Genomic Offset statistic (e.g. Capblancq *et al.*, 2020), here in its geometric gGO definition (Gain *et al.*, 2023). Under the assumptions that i) the considered environmental variables (at least for some of them) are among the main drivers of invasion success; ii) populations adapt partly genetically to these variables; and iii) genomic information is available from a set of presumably locally adapted populations representative of the species niches in both the native and invaded areas, then a low (resp. high) gGO value between two environments can be interpreted as a relative measure of (genetic) preadaptation (resp. maladaptation) for a population that would be moved from one environment to another. Accordingly, in a simulation study, Camus *et al.* (2024) recently showed that gGO can be related to the probability of invasive population establishment and can be efficiently estimated from the GEA regression model implemented in BayPass. From a practical point of view, the new BayPass extension to the analysis of hybrid datasets allowed estimation of gGO from our combined Pool-Seq and Ind-Seq data, which was not possible (or highly suboptimal) with other available methods.

The worldwide invasion of *D. suzukii* provided us with a valuable case study to illustrate how gGO-based analyzes can provide insights into the level of (genomic) adaptive challenge that poses the environment of a new invaded area to a population adapted to a given region of the native area, thus allowing one to better understand the past dynamics of biological invasion. Interestingly, despite evidence that some key genes were associated with adaptation in invaded areas, gGO indicated that genetic (pre)adaptation may have played a limited role in *D. suzukii*'s invasion success. Indeed, gGO values across North America and Europe were generally low and homogeneous, particularly in Europe, suggesting that most of these invaded areas did not pose a significant adaptive challenge to propagules originating from populations adapted to the environments of the presumed native area. This aligns with the rapid invasion of *D. suzukii* and suggests that once it arrived on these continents, it faced minimal adaptive challenges because some forms of preadaptation to invaded areas existed in native source populations. While not contradicting the findings of Feng *et al.* (2024), who identified many candidate SNPs associated with environmental variables, our results suggest that the underlying genomic regions were likely not crucial in the invasion process. Our conclusions are also in line with a previous study using species distribution modeling in *D. suzukii*, which indicated the absence of an obvious climatic niche shift between Asia and the first observation points in Europe and the US (Fraimout and Monnet, 2018).

Following reverse logic, the gGO maps that we derived for Africa and South America illustrate how gGO can be used to identify the most vulnerable regions in large areas that have not yet been (fully) invaded. According to our results, relatively low gGO values (that is, high risk) were found

in large parts of these continents (for example, the Amazon basin in South America or the Great Lakes region in Africa) where *D. suzukii* is currently absent, but could potentially establish rapidly. Some post hoc surveillance of these regions would be required to assess the extent to which these gGO-based predictions are informative. Indeed, while we report low gGO values in some recently invaded regions (e.g., South Africa), these were more moderate in others (e.g., Patagonia, Chile). In addition, the choice of the reference source environment from which the (preadapted) invading population would originate can be critical. For simplicity, for both Africa and South America, we assumed that the reference source environment is the one where populations have been established for at least two consecutive years. Consideration of other source environments may result in slightly to substantially different gGO maps. For example, in South America, although the first record of *D. suzukii* dates back to 2013 (Deprá *et al.*, 2014), it was reported just one year later, in 2014, in northern Patagonia (Lavagnino *et al.*, 2018), approximately 500 km from the AR-Gol population studied here and sampled in 2023. Given the rapid expansion of *D. suzukii* in Patagonia, AR-Gol could also represent a suitable reference environment to compute a gGO map of invasion across South America. This could potentially lead to even more pessimistic predictions than those based on the Nova Veneza (Brazil) reference environment used in Figure 4B, meaning that an even greater number of regions might be predicted to have low GO values and thus be at higher risk of invasion.. The gGO can also be used to identify the region of origin of a (locally adapted) population that would be the most (genomically) pre-adapted to a specific target area (see also Gautier *et al.*, 2024, for another recent application). Such information can be useful for IPM practice to increase vigilance on imports from the highlighted high risk regions (i.e., those with low gGO values and where *D. suzukii* is established). As here illustrated for the uninvaded Australian continent, the computation of gGO maps using a reference environment associated with the northwestern American US-Wa population (a key fresh fruit import region) supports a high overall risk of invasion from there. Densely populated areas of southern Australia that are likely targets of significant food trade exhibited low gGO values, suggesting that such a US-Wa-like *D. suzukii* population could easily establish there and gradually spread throughout the country. Previous research, which did not incorporate genomic data, similarly identified a high risk of *D. suzukii* invasion, particularly through passive human-mediated pathways (Maino *et al.*, 2021). Additionally, our gGO analysis suggested that invasions from Thailand present a greater (adaptive) risk than those from the US, implying that Australian management should also focus on monitoring imports from Thailand. However, this analysis serves more as an illustration of gGO-based IPM applications than as a source of actionable information for Australia, which already enforces strict control measures on most goods at risk of *D. suzukii* infestation. In a more general perspective, the sequencing of newly established populations could serve as an empirical method to validate gGO predictions in biological invasions. By comparing the most likely sources of invasion inferred from genetic data only, with those expected through gGO

computations accounting for environmental variables (e.g., Gautier *et al.*, 2024), we could assess the relevance of gGO-based predictions in real-world scenarios.

In general, approaches that rely on Genomic Offset statistics appear promising for understanding and predicting *D. suzukii* colonization in different environments. From a biological perspective, gGO estimates made sense globally, as regions with high estimated gGO values corresponded to environmental characteristics known to be physiologically challenging for *D. suzukii*. In particular, high to extreme gGO values were observed in cold regions, which are inhospitable to *D. suzukii* due to its "chill-susceptibility" (Kimura, 2004; Enriquez and Colinet, 2017; Ryan *et al.*, 2016; Winkler *et al.*, 2020), although *D. suzukii* can survive cold winters using natural or anthropogenic shelters (Jakobs *et al.*, 2015; Stockton *et al.*, 2019a; Rossi-Stacconi *et al.*, 2016). Similarly, elevated gGO values have been found in arid and semi-arid desert regions, which are also not suitable for *D. suzukii* due to its low tolerance to drought and high temperatures (Winkler *et al.*, 2020; Fanning *et al.*, 2019; Eben *et al.*, 2018). However, it should be noted that predictions of potential future distributions based on Ecological Niche Models (ENM) show low to no suitability for some areas that were associated with moderate or low gGO values in our analyzes, namely tropical regions such as the Amazon basin or Central Africa (Nair and Peterson, 2023; Ørsted and Ørsted, 2019). This discrepancy likely arises from fundamental differences between ENM and Genomic Offset approaches, which do not rely on the same data to model the relationship between populations and their environments. On the one hand, because ENMs rely on occurrence data, they can underestimate environmental suitability in regions where sampling is less intensive (Dubos *et al.*, 2022), which is likely the case of these tropical zones, overlooking the fact that the climatic characteristics of these regions are not necessarily limiting for the species, as indicated by gGO. On the other hand, the Genomic Offset based method focuses on a limited set of sequenced populations, which may limit its ability to capture the full ecological niche of the species. Moreover, it is important to keep in mind that current Genomic Offset metrics do not provide an absolute quantitative measure of invasion risk or adaptation difficulty. As a result, gGO values can only be interpreted relatively, i.e., by ranking environments according to their susceptibility to invasion by source populations that are (genetically) adapted to a given reference region. The exact conditions under which this relative interpretation is theoretically valid also need to be further explored and could, for example, be challenged if a given target environment is particularly suitable for a species, regardless of the genetic makeup of the population (Lotterhos, 2024).

In addition, other factors not included in the models underlying gGO calculations may be critical for successful invasion and must also be considered to explain or predict the establishment of invasive species. First, some pivotal environmental variables could have been ignored in the GEA or their variation may be imperfectly captured with the analyzed PCs, which may have a decisive impact on the predictive performance of gGO (Camus *et al.*, 2024). For example, we did not (directly)

incorporate fruit-related variables other than land use due to data availability limitations on the considered continental scales. Given that small fruits availability is an important component of $D.$ $suzukii$ fitness, its inclusion in future gGO analyzes could greatly improve the precision of invasion predictions by taking into account additional factors that influence the success of $D.$ $suzukii$ establishment. Note also that many of the top significant candidate genes detected by our C_2 genome scan are related to functions that may be crucial for the species invasion success (e.g., pesticide resistance, immune response, dispersal, or feeding behavior) but are not considered in the gGO analysis. Integrating environmental descriptors that are directly linked to these functions, for instance, pesticide levels, might provide a different picture of the actual adaptive challenge encountered during the invasion(s), with higher and more heterogeneous gGO values. Second, innate characteristics of the species, including phenotypic plasticity, can play an important role for invasion success and may be difficult (if not impossible) to include in gGO computations. Hence, $D.$ $suzukii$ has several characteristics that may have facilitated its successful invasion, such as its ability to i) tolerate cold environments through mechanisms such as overwintering or diapause (Little et $al.$, 2020); ii) exploit a broad trophic niche with host plants that span several families and many fruit species (Olazcuaga et $al.$, 2023); and iii) even subsist on a diet without fruits (Poyet et $al.$, 2015; Stockton et $al.$, 2019b). Finally, contingent and extrinsic factors are also critical to successfully invade a new area. At the very least, enough individuals (of both sexes for sexual species) must first be able to reach it. In the case of $D.$ $suzukii$, its widespread distribution is likely driven by human-mediated passive dispersal rather than active movement (Adrion et $al.$, 2014; Calabria et $al.$, 2012; Maino et $al.$, 2021). This form of assisted migration can significantly affect the dynamics of invasion, influencing which areas are invaded first, and the spread of the species in a newly invaded area. In addition, the number of introduced individuals (propagule size) and their arrival patterns (number and spatio-temporal distribution of introduction events), which together define the so-called "propagule pressure", have also been shown to be a crucial factor in the establishment of invasive populations (Simberloff, 2009; Estoup et $al.$, 2016). In fact, high propagule pressure can buffer adaptive challenges, as shown in our recent simulation study, where the performance of Genomic Offset-based analyzes in predicting the probability of invasive populations being established was found to be reduced for scenarios involving the single introduction of larger numbers of individuals (Camus et $al.$, 2024). For example, even though some populations from Thailand may be more preadapted to Australia, populations from America may be more likely to become established in Australia due to greater fruit infestation and/or higher fruit imports and consequently greater propagule pressure.

Despite these limitations and the avenue for further refinement, we believe that our findings provide valuable insights into the invasion success of $D.$ $suzukii$ and more generally illustrate and highlight the effectiveness and promise of GEA and gGO-based analyzes to elucidate the factors underpinning the dynamic of past and future biological invasions. Our framework for population genomic analysis

and our recent development of methodological tools are inherently generic. As such, they can be applied to any other invasive species in order to help guide management decisions.

Data availability

All data were deposited in the SRA repository (project PRJNA1032894). These included Pacbio HiFi raw reads (Run ID: SRR29552229) and the HiC library paired-end sequences (Run ID: SRR29552230) used for the de novo assembly of the Japanese strain, and all new Pool-Seq and Ind-Seq data (see accession details in Tables S1 and S2). The annotated *dsu_isojap1.0* whole genome assembly is publicly available from the NCBI repository under the ID GCA_043229665.1 (`https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_043229965.1/`). The newly developed version (v3.0) of BayPass software is publicly available from `https://forgemia.inra.fr/mathieu.gautier/baypass_public`

Acknowledgments

We wish to thank Toshiro Aigaki (TMU, Tokyo, Japan), Nicolas Gompel (Bonn University, Germany) and Benjamin Prud'homme (IBDML, Université Aix-Marseille, France) for providing and having maintained individuals from the Japanese isofemale line used for the genome assembly. We are also grateful to Benjamin Prud'homme for providing *D. subpulchrella* individuals, and Dr Marina Stamenković-Radak, Dr Mihailo Jelić, and Mina Rakić for their assistance in collecting the sample from Serbia. Mathieu Gautier, Simon Boitard and Arnaud Estoup have benefited from State aid managed by the Agence Nationale de la Recherche under the France 2030 programme, reference ANR-23-EXMA-0002 ("AgroStat" project of the Programme et Equipements Prioritaires de Recherche "MathsVives"). Svitlana Serga was supported by the PAUSE-ANR Ukraine Program. This work was performed in collaboration with the GeT core facility, Toulouse, France (GeT, `https://doi.org/10.15454/1.5572370921303193E12`). MGX and GeT core facility acknowledge financial support from France Génomique National infrastructure, funded as part of "Investissement d'avenir" program managed by Agence Nationale pour la Recherche (contract ANR-10-INBS-09). We are also grateful to the genotoul bioinformatics platform Toulouse Occitanie (Bioinfo Genotoul, `https://doi.org/10.15454/1.5572369328961167E12`) for providing computing resources. We finally thank the DrosEU consortium for helpful discussions during stimulating meetings.

Bibliography

Adrion, J. R., Kousathanas, A., Pascual, M., Burrack, H. J., Haddad, N. M., Bergland, A. O., Machado, H., Sackton, T. B., Schlenke, T. A., Watada, M., Wegmann, D., and Singh, N. D. 2014. Drosophila suzukii: The Genetic Footprint of a Recent, Worldwide Invasion. *Molecular Biology and Evolution*, 31(12): 3148–3163.

Aouari, I., Barech, G., and Khaldi, M. 2022. First record of the agricultural pest *Drosophila suzukii* (Matsumura, 1931) (Diptera: Drosophilidae) in Algeria. *EPPO Bulletin*, 52(2): 471–478.

Arya, G. H., Magwire, M. M., Huang, W., Serrano-Negron, Y. L., Mackay, T. F., and Anholt, R. R. 2015. The genetic basis for variation in olfactory behavior in drosophila melanogaster. *Chemical senses*, 40(4): 233–243.

Asplen, M. K., Anfora, G., Biondi, A., Choi, D. S., Chu, D., Daane, K. M., Gibert, P., Gutierrez, A. P., Hoelmer, K. A., Hutchison, W. D., Isaacs, R., Jiang, Z. L., Kárpáti, Z., Kimura, M. T., Pascual, M., Philips, C. R., Plantamp, C., Ponti, L., Vétek, G., Vogt, H., Walton, V. M., Yu, Y., Zappalà, L., and Desneux, N. 2015. Invasion biology of spotted wing Drosophila (Drosophila suzukii): a global perspective and future priorities. *Journal of Pest Science*, 88(3): 469–494.

Barrett, R. D. H. and Schluter, D. 2008. Adaptation from standing genetic variation. *Trends in Ecology & Evolution*, 23(1): 38–44.

Baudry, L., Guiglielmoni, N., Marie-Nelly, H., Cormier, A., Marbouty, M., Avia, K., Mie, Y. L., Godfroy, O., Sterck, L., Cock, J. M., Zimmer, C., Coelho, S. M., and Koszul, R. 2020. in-staGRAAL: chromosome-level quality scaffolding of genomes using a proximity ligation-based scaffolder. *Genome Biol*, 21(1): 148.

Bayona-Feliu, A., Casas-Lamesa, A., Carbonell, A., Climent-Cantó, P., Tatarski, M., Pérez-Montero, S., Azorín, F., and Bernués, J. 2016. Histone h1: lessons from drosophila. *Biochimica et Biophysica Acta (BBA)-Gene Regulatory Mechanisms*, 1859(3): 526–532.

Belgacem, Y. H. and Martin, J.-R. 2007. Hmgcr in the corpus allatum controls sexual dimorphism of locomotor activity and body size via the insulin pathway in drosophila. *PLoS One*, 2(1): e187.

Bhutkar, A., Schaeffer, S. W., Russo, S. M., Xu, M., Smith, T. F., and Gelbart, W. M. 2008. Chromosomal rearrangement inferred from comparisons of 12 Drosophila genomes. *Genetics*, 179(3): 1657–1680.

Bogaerts-Márquez, M., Guirao-Rico, S., Gautier, M., and González, J. 2021. Temperature, rainfall and wind variables underlie environmental adaptation in natural populations of Drosophila melanogaster. *Molecular Ecology*, 30(4): 938–954.

Boughdad, A., Haddi, K., El Bouazzati, A., Nassiri, A., Tahiri, A., El Anbri, C., Eddaya, T., Zaid, A., and Biondi, A. 2021. First record of the invasive spotted wing Drosophila infesting berry crops in Africa. *Journal of Pest Science*, 94(2): 261–271.

Bradshaw, C. J. A., Leroy, B., Bellard, C., Roiz, D., Albert, C., Fournier, A., Barbet-Massin, M., Salles, J.-M., Simard, F., and Courchamp, F. 2016. Massive yet grossly underestimated global costs of invasive insects. *Nature Communications*, 7(1): 12986.

Bruce, T. J. A. 2010. Tackling the threat to food security caused by crop pests in the new millennium. *Food Security*, 2(2): 133–141.

Cabanettes, F. and Klopp, C. 2018. D-GENIES : Dot plot large GENomes in an interactive, efficient and simple way. *PeerJ Preprints*, 6: e26567v1.

Calabria, G., Máca, J., Bächli, G., Serra, L., and Pascual, M. 2012. First records of the potential pest species drosophila suzukii (diptera: Drosophilidae) in europe. *Journal of Applied entomology*, 136(1-2): 139–147.

Camus, L., Gautier, M., and Boitard, S. 2024. Predicting species invasiveness with genomic data: Is genomic offset related to establishment probability? *Evolutionary Applications*, 17(6): e13709.

Capblancq, T., Fitzpatrick, M. C., Bay, R. A., Exposito-Alonso, M., and Keller, S. R. 2020. Genomic Prediction of (Mal)Adaptation Across Current and Future Climatic Landscapes. *Annual Review of Ecology, Evolution, and Systematics*, 51(1): 245–269.

Challis, R., Richards, E., Rajan, J., Cochrane, G., and Blaxter, M. 2020. BlobToolKit – Interactive Quality Assessment of Genome Assemblies. *G3*, 10(4): 1361–1374.

Chang, C.-H., Chavan, A., Palladino, J., Wei, X., Martins, N. M. C., Santinello, B., Chen, C.-C., Erceg, J., Beliveau, B. J., Wu, C.-T., Larracuente, A. M., and Mellone, B. G. 2019. Islands of retroelements are major components of drosophila centromeres. *PLOS Biology*, 17(5): 1–40.

Chen, S., Zhou, Y., Chen, Y., and Gu, J. 2018. fastp: an ultra-fast all-in-one fastq preprocessor. *Bioinformatics*, 34(17): i884–i890.

Cheng, H., Concepcion, G. T., Feng, X., Zhang, H., and Li, H. 2021. Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. *Nature Methods*, 18(2): 170–175.

Chiu, J. C., Jiang, X., Zhao, L., Hamm, C. A., Cridland, J. M., Saelao, P., Hamby, K. A., Lee, E. K., Kwok, R. S., Zhang, G., Zalom, F. G., Walton, V. M., and Begun, D. J. 2013. Genome of Drosophila suzukii, the spotted wing drosophila. *G3*, 3(12): 2257–71.

Colautti, R. I. and Barrett, S. C. H. 2013. Rapid Adaptation to Climate Facilitates Range Expansion of an Invasive Plant. *Science*, 342(6156): 364–366.

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., and Li, H. 2021. Twelve years of SAMtools and BCFtools. *GigaScience*, 10(2): giab008.

Deans, C. and Hutchison, W. D. 2022. Propensity for resistance development in the invasive berry pest, spotted-wing drosophila (drosophila suzukii), under laboratory selection. *Pest management science*, 78(12): 5203–5212.

Delclos, P. J., Adhikari, K., Mai, A. B., Hassan, O., Oderhowho, A. A., Sriskantharajah, V., Trinh, T., and Meisel, R. 2024. Trans regulation of an odorant binding protein by a proto-y chromosome affects male courtship in house fly. *Elife*, 13: e90349.

Deprá, M., Poppe, J., Schmitz, H., De Toni, D., and Valente, V. 2014. The first records of the invasive pest Drosophila suzukii in the South American continent. *J Pest Sci*, 87.

Diagne, C., Leroy, B., Vaissière, A.-C., Gozlan, R. E., Roiz, D., Jarić, I., Salles, J.-M., Bradshaw, C. J. A., and Courchamp, F. 2021. High and rising economic costs of biological invasions worldwide. *Nature*, 592(7855): 571–576.

Drosopoulou, E., Gariou-Papalexiou, A., Karamoustou, E., Gouvi, G., Augustinos, A. A., Bourtzis, K., and Zacharopoulou, A. 2019. The chromosomes of Drosophila suzukii (Diptera: Drosophilidae): detailed photographic polytene chromosomal maps and in situ hybridization data. *Molecular Genetics and Genomics*, 294(6): 1535–1546.

Druet, T. and Gautier, M. 2017. A model-based approach to characterize individual inbreeding at both global and local genomic scales. *Molecular Ecology*, 26(20): 5820–5841.

Dubos, N., Préau, C., Lenormand, M., Papuga, G., Monsarrat, S., Denelle, P., Le Louarn, M., Heremans, S., May, R., Roche, P., *et al.* 2022. Assessing the effect of sample bias correction in species distribution models. *Ecological Indicators*, 145: 109487.

Eben, A., Reifenrath, M., Briem, F., Pink, S., and Vogt, H. 2018. Response of d rosophila suzukii (d iptera: D rosophilidae) to extreme heat and dryness. *Agricultural and Forest Entomology*, 20(1): 113–121.

Edwards, A. C., Rollmann, S. M., Morgan, T. J., and Mackay, T. F. C. 2006. Quantitative genomics of aggressive behavior in drosophila melanogaster. *PLoS genetics*, 2(9): e154.

El-Kholy, S. E., Afifi, B., El-Husseiny, I., and Seif, A. 2021. Octopamine signaling via oamb is essential for a well-orchestrated climbing performance of adult drosophila melanogaster.

Enriquez, T. and Colinet, H. 2017. Basal tolerance to heat and cold exposure of the spotted wing drosophila, drosophila suzukii. *PeerJ*, 5: e3112.

Estoup, A., Ravigné, V., Hufbauer, R., Vitalis, R., Gautier, M., and Facon, B. 2016. Is There a Genetic Paradox of Biological Invasion? *Annual Review of Ecology, Evolution, and Systematics*, 47(Volume 47, 2016): 51–72.

Fanning, P. D., Johnson, A. E., Luttinen, B. E., Espeland, E. M., Jahn, N. T., and Isaacs, R. 2019. Behavioral and physiological resistance to desiccation in spotted wing drosophila (diptera: Drosophilidae). *Environmental entomology*, 48(4): 792–798.

Fariello, M. I., Boitard, S., Mercier, S., Robelin, D., Faraut, T., Arnould, C., Recoquillay, J., Bouchez, O., Salin, G., Dehais, P., Gourichon, D., Leroux, S., Pitel, F., Leterrier, C., and San-Cristobal, M. 2017. Accounting for linkage disequilibrium in genome scans for selection without individual genotypes: The local score approach. *Molecular Ecology*, 26(14): 3700–3714.

Feng, S., DeGrey, S. P., Guédot, C., Schoville, S. D., and Pool, J. E. 2024. Genomic Diversity Illuminates the Environmental Adaptation of Drosophila suzukii. *Genome Biology and Evolution*, 16(9): evae195.

Fitzpatrick, M. C. and Keller, S. R. 2015. Ecological genomics meets community-level modelling of biodiversity: mapping the genomic landscape of current and future environmental adaptation. *Ecology Letters*, 18(1): 1–16.

Fitzpatrick, M. C., Chhatre, V. E., Soolanayakanahally, R. Y., and Keller, S. R. 2021. Experimental support for genomic prediction of climate maladaptation using the machine learning approach Gradient Forests. *Molecular Ecology Resources*, 21(8): 2749–2765.

Fraimout, A. and Monnet, A.-C. 2018. Accounting for intraspecific variation to quantify niche dynamics along the invasion routes of drosophila suzukii. *Biological Invasions*, 20(10): 2963–2979.

Fraimout, A., Debat, V., Fellous, S., Hufbauer, R. A., Foucaud, J., Pudlo, P., Marin, J.-M., Price, D. K., Cattel, J., Chen, X., Deprá, M., François Duyck, P., Guedot, C., Kenis, M., Kimura, M. T., Loeb, G., Loiseau, A., Martinez-Sañudo, I., Pascual, M., Polihronakis Richmond, M., Shearer, P., Singh, N., Tamura, K., Xuéreb, A., Zhang, J., and Estoup, A. 2017. Deciphering the Routes of invasion of Drosophila suzukii by Means of ABC Random Forest. *Molecular Biology and Evolution*, 34(4): 980–996.

Fukasawa, Y., Ermini, L., Wang, H., Carty, K., and Cheung, M.-S. 2020. LongQC: A Quality Control Tool for Third Generation Sequencing Long Read Data. *G3*, 10(4): 1193–1196.

Gain, C., Rhoné, B., Cubry, P., Salazar, I., Forbes, F., Vigouroux, Y., Jay, F., and François, O. 2023. A Quantitative Theory for Genomic Offset Statistics. *Molecular Biology and Evolution*, 40(6): msad140.

Garriga, A., Toubarro, D., Morton, A., Simões, N., and García-del Pino, F. 2024. Analysis of the immune transcriptome of the invasive pest spotted wing drosophila infected by steinernema carpocapsae. *Bulletin of Entomological Research*, 114(5): 622–630.

Garrison, E. and Marth, G. 2012. Haplotype-based variant detection from short-read sequencing. *arXiv*, 1207.3907.

Gautier, M. 2015. Genome-Wide Scan for Adaptive Divergence and Association with Population-Specific Covariates. *Genetics*, 201(4): 1555–1579.

Gautier, M. 2023. Efficient *k-mer* based curation of raw sequence data: application in *Drosophila suzukii*. *Peer Community Journal*, 3: e79.

Gautier, M., Yamaguchi, J., Foucaud, J., Loiseau, A., Ausset, A., Facon, B., Gschloessl, B., Lagnel, J., Loire, E., Parrinello, H., Severac, D., Lopez-Roques, C., Donnadieu, C., Manno, M., Berges, H., Gharbi, K., Lawson-Handley, L., Zang, L.-S., Vogel, H., Estoup, A., and Prud'homme, B. 2018. The Genomic Basis of Color Pattern Polymorphism in the Harlequin Ladybird. *Current Biology*, 28(20): 3296–3302.e7.

Gautier, M., Vitalis, R., Flori, L., and Estoup, A. 2022. f-Statistics estimation and admixture graph construction with Pool-Seq or allele count data using the R package poolfstat. *Molecular Ecology Resources*, 22(4): 1394–1416.

Gautier, M., Micol, T., Camus, L., Moazami-Goudarzi, K., Naves, M., Guéret, E., Engelen, S., Lemainque, A., Colas, F., Flori, L., and Druet, T. 2024. Genomic Reconstruction of the Successful Establishment of a Feralized Bovine Population on the Subantarctic Island of Amsterdam. *Molecular Biology and Evolution*, 41(7): msae121.

Haubrock, P. J., Turbelin, A. J., Cuthbert, R. N., Novoa, A., Taylor, N. G., Angulo, E., Ballesteros-Mejia, L., Bodey, T. W., Capinha, C., Diagne, C., Essl, F., Golivets, M., Kirichenko, N., Kourantidou, M., Leroy, B., Renault, D., Verbrugge, L., and Courchamp, F. 2021. Economic costs of invasive alien species across Europe. *NeoBiota*, 67: 153–190.

Haye, T., Girod, P., Cuthbertson, A., Wang, X., Daane, K., Hoelmer, K., Baroffio, C., Zhang, J., and Desneux, N. 2016. Current swd ipm tactics and their practical implementation in fruit crops across different regions around the world. *Journal of Pest Science*, 89: 643–651.

Hijmans, R. J. 2022. *raster: Geographic Data Analysis and Modeling*. R package version 3.5-15.

Hijmans, R. J., Phillips, S., Leathwick, J., and Elith, J. 2023. *dismo: Species Distribution Modeling*. R package version 1.3-14.

Hill, V. M., O'Connor, R. M., Sissoko, G. B., Irobunda, I. S., Leong, S., Canman, J. C., Stavropoulos, N., and Shirasu-Hiza, M. 2018. A bidirectional relationship between sleep and oxidative stress in drosophila. *PLoS biology*, 16(7): e2005206.

Hofmeister, N. R., Werner, S. J., and Lovette, I. J. 2021. Environmental correlates of genetic variation in the invasive European starling in North America. *Molecular Ecology*, 30(5): 1251–1263.

Hoskins, R. A., Carlson, J. W., Wan, K. H., Park, S., Mendez, I., Galle, S. E., Booth, B. W., Pfeiffer, B. D., George, R. A., Svirskas, R., Krzywinski, M., Schein, J., Accardo, M. C., Damia, E., Messina, G., Méndez-Lago, M., de Pablos, B., Demakova, O. V., Andreyeva, E. N., Boldyreva, L. V., Marra, M., Carvalho, A. B., Dimitri, P., Villasante, A., Zhimulev, I. F., Rubin, G. M., Karpen, G. H., and Celniker, S. E. 2015. The release 6 reference sequence of the drosophila melanogaster genome. *Genome Research*, 25(3): 445–58.

Hufbauer, R. A., Facon, B., Ravigné, V., Turgeon, J., Foucaud, J., Lee, C. E., Rey, O., and Estoup, A. 2012. Anthropogenically induced adaptation to invade (AIAI): contemporary adaptation to human-altered habitats within the native range can promote invasions. *Evolutionary Applications*, 5(1): 89–101.

International Trade Center 2024. Trade Map. Accessed: 2024-07-22.

IPCC 2024. International Plant Protection Convention country report: Notification of the detection of Drosophila suzukii, the Spotted Wing Drosophila in the Republic of South Africa. Technical Report ZAF-58/2, National Plant Protection Organization of South Africa.

Jakobs, R., Gariepy, T. D., and Sinclair, B. J. 2015. Adult plasticity of cold tolerance in a continental-temperate population of drosophila suzukii. *Journal of Insect Physiology*, 79: 1–9.

Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder, H. P., and Kessler, M. 2017. Climatologies at high resolution for the earth's land surface areas. *Scientific Data*, 4(1): 170122.

Karra, K., Kontgis, C., Statman-Weil, Z., Mazzariello, J. C., Mathis, M., and Brumby, S. P. 2021. Global land use/land cover with Sentinel 2 and deep learning. *2021 IEEE International Geoscience and Remote Sensing Symposium IGARSS*, pages 4704–4707.

Kim, S. Y., Lohmueller, K. E., Albrechtsen, A., Li, Y., Korneliussen, T., Tian, G., Grarup, N., Jiang, T., Andersen, G., Witte, D., Jorgensen, T., Hansen, T., Pedersen, O., Wang, J., and Nielsen, R. 2011. Estimation of allele frequency and association mapping using next-generation sequencing data. *BMC Bioinformatics*, 12(1): 231.

Kimura, M. T. 2004. Cold and heat tolerance of drosophilid flies with reference to their latitudinal distributions. *Oecologia*, 140: 442–449.

Korneliussen, T. S., Albrechtsen, A., and Nielsen, R. 2014. Angsd: Analysis of next generation sequencing data. *BMC Bioinformatics*, 15(1): 356.

Kwadha, C. A., Okwaro, L. A., Kleman, I., Rehermann, G., Revadi, S., Ndlela, S., Khamis, F. M., Nderitu, P. W., Kasina, M., George, M. K., Kithusi, G. G., Mohamed, S. A., Lattorff, H. M. G., and Becher, P. G. 2021. Detection of the spotted wing drosophila, Drosophila suzukii, in continental sub-Saharan Africa. *Journal of Pest Science*, 94(2): 251–259.

Lachmuth, S., Capblancq, T., Prakash, A., Keller, S. R., and Fitzpatrick, M. C. 2023. Novel genomic offset metrics integrate local adaptation into habitat suitability forecasts and inform assisted migration. *Ecological Monographs*, 94(1): e1593.

Láruson, A. J., Yeaman, S., and Lotterhos, K. E. 2020. The Importance of Genetic Redundancy in Evolution. *Trends in Ecology & Evolution*, 35(9): 809–822.

Lavagnino, N. J., Cichón, L. I., De La Vega, G. J., Garrido, S. A., Fanara, J. J., *et al.* 2018. New records of the invasive pest Drosophila suzukii (Matsumura)(Diptera: Drosophilidae) in the South American continent. *Revista de la Sociedad Entomológica Argentina*, 77(1): 27.

Lewald, K. M., Abrieux, A., Wilson, D. A., Lee, Y., Conner, W. R., Andreazza, F., Beers, E. H., Burrack, H. J., Daane, K. M., Diepenbrock, L., Drummond, F. A., Fanning, P. D., Gaffney, M. T., Hesler, S. P., Ioriatti, C., Isaacs, R., Little, B. A., Loeb, G. M., Miller, B., Nava, D. E., Rendon, D., Sial, A. A., Bezerra da Silva, C. S., Stockton, D. G., Van Timmeren, S., Wallingford, A., Walton, V. M., Wang, X., Zhao, B., Zalom, F. G., and Chiu, J. C. 2021. Population genomics of Drosophila suzukii reveal longitudinal population structure and signals of migrations in and out of the continental United States. *G3*, 11(12): jkab343.

Li, H. 2021. New strategies to improve minimap2 alignment accuracy. *Bioinformatics*, 37(23): 4572–4574.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R. 2009. The sequence alignment/map format and samtools. *Bioinformatics*, 25(16): 2078–9.

Lind, B. M. and Lotterhos, K. E. 2024. The limits of predicting maladaptation to future environments with genomic data. *bioRxiv*, 2024.01.30.577973.

Little, C. M., Chapman, T. W., and Hillier, N. K. 2020. Plasticity is key to success of drosophila suzukii (diptera: Drosophilidae) invasion. *Journal of Insect Science*, 20(3): 5.

Liu, L., Leonard, A. S., Motto, D. G., Feller, M. A., Price, M. P., Johnson, W. A., and Welsh, M. J. 2003. Contribution of drosophila deg/enac genes to salt taste. *Neuron*, 39(1): 133–146.

Lotterhos, K. E. 2023. The paradox of adaptive trait clines with nonclinal patterns in the underlying genes. *Proceedings of the National Academy of Sciences*, 120(12): e2220313120.

Lotterhos, K. E. 2024. Interpretation issues with "genomic vulnerability" arise from conceptual issues in local adaptation and maladaptation. *Evolution Letters*, 8(3): 331–339.

Luo, J., Lushchak, O. V., Goergen, P., Williams, M. J., and Nässel, D. R. 2014. Drosophila insulin-producing cells are differentially modulated by serotonin and octopamine receptors and affect social behavior. *PloS one*, 9(6): e99732.

Ma, L., Cao, L.-J., Hoffmann, A. A., Gong, Y.-J., Chen, J.-C., Chen, H.-S., Wang, X.-B., Zeng, A.-P., Wei, S.-J., and Zhou, Z.-S. 2020. Rapid and strong population genetic differentiation and genomic signatures of climatic adaptation in an invasive mealybug. *Diversity and Distributions*, 26(5): 610–622.

Maino, J. L., Schouten, R., and Umina, P. 2021. Predicting the global invasion of drosophila suzukii to improve australian biosecurity preparedness. *Journal of Applied Ecology*, 58(4): 789–800.

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., and Zdobnov, E. M. 2021. BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology and Evolution*, 38(10): 4647–4654.

Matthey-Doret, C., Baudry, L., Bignaud, A., Cournac, A., Remi-Montagne, Guiglielmoni, N., Foutel-Rodier, T., and Scolari, V. F. 2020. hicstuff: Simple library/pipeline to generate and handle hi-c data.

Mazza, G., Tricarico, E., Genovesi, P., and Gherardi, F. 2014. Biological invaders are threats to human health: an overview. *Ethology Ecology & Evolution*.

Meisner, J. and Albrechtsen, A. 2018. Inferring Population Structure and Admixture Proportions in Low-Depth NGS Data. *Genetics*, 210(2): 719–731.

Meng, H., Yamashita, C., Shiba-Fukushima, K., Inoshita, T., Funayama, M., Sato, S., Hatta, T., Natsume, T., Umitsu, M., Takagi, J., *et al.* 2017. Loss of parkinson's disease-associated protein chchd2 affects mitochondrial crista structure and destabilizes cytochrome c. *Nature communications*, 8(1): 15500.

Mérel, V., Gibert, P., Buch, I., Rodriguez Rada, V., Estoup, A., Gautier, M., Fablet, M., Boulesteix, M., and Vieira, C. 2021. The Worldwide Invasion of Drosophila suzukii Is Accompanied by a Large

Increase of Transposable Element Load and a Small Number of Putatively Adaptive Insertions. *Molecular Biology and Evolution*, 38(10): 4252–4267.

Mollot, G., Pantel, J. H., and Romanuk, T. N. 2017. Chapter Two - The Effects of Invasive Species on the Decline in Species Richness: A Global Meta-Analysis. In D. A. Bohan, A. J. Dumbrell, and F. Massol, editors, *Advances in Ecological Research*, volume 56 of *Networks of Invasion: A Synthesis of Concepts*, pages 61–83. Academic Press.

Nair, R. R. and Peterson, A. T. 2023. Mapping the global distribution of invasive pest drosophila suzukii and parasitoid leptopilina japonica: implications for biological control. *PeerJ*, 11: e15222.

Ohashi, Y. Y., Haino-Fukushima, K., and Fuyama, Y. 1991. Purification and charachterization of an ovulation stimulating substance from the male accessory glands of drosophila suzukii. *Insect Biochemistry*, 21(4): 413–419.

Olazcuaga, L., Loiseau, A., Parrinello, H., Paris, M., Fraimout, A., Guedot, C., Diepenbrock, L. M., Kenis, M., Zhang, J., Chen, X., Borowiec, N., Facon, B., Vogt, H., Price, D. K., Vogel, H., Prud'homme, B., Estoup, A., and Gautier, M. 2020. A Whole-Genome Scan for Association with Invasion Success in the Fruit Fly Drosophila suzukii Using Contrasts of Allele Frequencies Corrected for Population Structure. *Molecular Biology and Evolution*, 37(8): 2369–2385.

Olazcuaga, L., Baltenweck, R., Leménager, N., Maia-Grondard, A., Claudel, P., Hugueney, P., and Foucaud, J. 2023. Metabolic consequences of various fruit-based diets in a generalist insect species. *eLife*, 12: e84370.

Ometto, L., Cestaro, A., Ramasamy, S., Grassi, A., Revadi, S., Siozios, S., Moretto, M., Fontana, P., Varotto, C., Pisani, D., Dekker, T., Wrobel, N., Viola, R., Pertot, I., Cavalieri, D., Blaxter, M., Anfora, G., and Rota-Stabelli, O. 2013. Linking Genomics and Ecology to Investigate the Complex Evolution of an Invasive Drosophila Pest. *Genome Biology and Evolution*, 5(4): 745–757.

Ørsted, I. V. and Ørsted, M. 2019. Species distribution models of the spotted wing drosophila (drosophila suzukii, diptera: Drosophilidae) in its native and invasive range reveal an ecological niche shift. *Journal of applied ecology*, 56(2): 423–435.

Paini, D. R., Sheppard, A. W., Cook, D. C., De Barro, P. J., Worner, S. P., and Thomas, M. B. 2016. Global threat to agriculture from invasive species. *Proceedings of the National Academy of Sciences*, 113(27): 7575–7579.

Parada, C. and Prieto, D. 2024. Survival, movement, and lifespan: Decoding the roles of patched-related (ptr) in drosophila melanogaster. *bioRxiv*, pages 2024–05.

Paris, M., Boyer, R., Jaenichen, R., Wolf, J., Karageorgi, M., Green, J., Cagnon, M., Parinello, H., Estoup, A., Gautier, M., Gompel, N., and Prud'homme, B. 2020. Near-chromosome level genome

assembly of the fruit pest Drosophila suzukii using long-read sequencing. *Scientific Reports*, 10(1): 11227.

Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschoreck, T., Webster, T., and Reich, D. 2012. Ancient Admixture in Human History. *Genetics*, 192(3): 1065–1093.

Pedersen, B. S. and Quinlan, A. R. 2017. Mosdepth: quick coverage calculation for genomes and exomes. *Bioinformatics*, 34(5): 867–868.

Pfenninger, M., Reuss, F., Kiebler, A., Schönnenbeck, P., Caliendo, C., Gerber, S., Cocchiararo, B., Reuter, S., Blüthgen, N., Mody, K., Mishra, B., Bálint, M., Thines, M., and Feldmeyer, B. 2021. Genomic basis for drought resistance in European beech forests threatened by climate change. *eLife*, 10: e65532.

Poyet, M., Le Roux, V., Gibert, P., Meirland, A., Prevost, G., Eslin, P., and Chabrerie, O. 2015. The wide potential trophic niche of the asiatic fruit fly drosophila suzukii: the key of its invasion success in temperate europe? *PloS one*, 10(11): e0142785.

Prendergast, L. and Reinberg, D. 2021. The missing linker: emerging trends for h1 variant-specific functions. *Genes & development*, 35(1-2): 40–58.

Rhoné, B., Defrance, D., Berthouly-Salazar, C., Mariac, C., Cubry, P., Couderc, M., Dequincey, A., Assoumanne, A., Kane, N. A., Sultan, B., Barnaud, A., and Vigouroux, Y. 2020. Pearl millet genomic vulnerability to climate change in West Africa highlights the need for regional collaboration. *Nature Communications*, 11(1): 5274.

Rossi-Stacconi, M. V., Kaur, R., Mazzoni, V., Ometto, L., Grassi, A., Gottardello, A., Rota-Stabelli, O., and Anfora, G. 2016. Multiple lines of evidence for reproductive winter diapause in the invasive pest drosophila suzukii: useful clues for control strategies. *Journal of pest science*, 89: 689–700.

Ryan, G. D., Emiljanowicz, L., Wilkinson, F., Kornya, M., and Newman, J. A. 2016. Thermal tolerances of the spotted-wing drosophila drosophila suzukii (diptera: Drosophilidae). *Journal of Economic Entomology*, 109(2): 746–752.

Sabio, M. C., Alzogaray, R., and Fanara, J. J. 2024. Genetic architecture of the toxicological response to eucalyptol and citronellal in drosophila melanogaster. *Pesticide Biochemistry and Physiology*, 202: 105938.

Sario, S., Marques, J. P., Farelo, L., Afonso, S., Santos, C., and Melo-Ferreira, J. 2024. Dissecting the invasion history of spotted-wing drosophila (drosophila suzukii) in portugal using genomic data. *BMC Genomics*, 25(1): 813.

Schwanitz, T. W., Polashock, J. J., Stockton, D. G., Rodriguez-Saona, C., Sotomayor, D., Loeb, G., and Hawkings, C. 2022. Molecular and behavioral studies reveal differences in olfaction between winter and summer morphs of drosophila suzukii. *PeerJ*, 10: e13825.

Shorter, J. R., Dembeck, L. M., Everett, L. J., Morozova, T. V., Arya, G. H., Turlapati, L., St. Armour, G. E., Schal, C., Mackay, T. F., and Anholt, R. R. 2016. Obp56h modulates mating behavior in drosophila melanogaster. *G3: Genes, Genomes, Genetics*, 6(10): 3335–3342.

Simberloff, D. 2009. The role of propagule pressure in biological invasions. *Annual review of ecology, evolution, and systematics*, 40(1): 81–102.

Skoglund, P. and Jakobsson, M. 2011. Archaic human ancestry in east asia. *Proceedings of the National Academy of Sciences*, 108(45): 18301–18306.

Skotte, L., Korneliussen, T. S., and Albrechtsen, A. 2013. Estimating individual admixture proportions from next generation sequencing data. *Genetics*, 195(3): 693–702.

Smit, A., Hubley, R., and Green, P. 2023. *RepeatMasker Open-4.0*. Institute for Systems Biology.

Spierer, A. N., Mossman, J. A., Smith, S. P., Crawford, L., Ramachandran, S., and Rand, D. M. 2021. Natural variation in the regulation of neurodevelopmental genes modifies flight performance in drosophila. *PLoS Genetics*, 17(3): e1008887.

Stockton, D., Wallingford, A., Rendon, D., Fanning, P., Green, C. K., Diepenbrock, L., Ballman, E., Walton, V. M., Isaacs, R., Leach, H., *et al.* 2019a. Interactions between biotic and abiotic factors affect survival in overwintering drosophila suzukii (diptera: Drosophilidae). *Environmental Entomology*, 48(2): 454–464.

Stockton, D. G., Brown, R., and Loeb, G. M. 2019b. Not berry hungry? discovering the hidden food sources of a small fruit specialist, drosophila suzukii. *Ecological Entomology*, 44(6): 810–822.

Swarup, S., Morozova, T. V., Sridhar, S., Nokes, M., and Anholt, R. R. 2014. Modulation of feeding behavior by odorant-binding proteins in drosophila melanogaster. *Chemical senses*, 39(2): 125–132.

Thibaud-Nissen, F., Souvorov, A., Murphy, T., DiCuccio, M., and Kitts, P. 2013. Eukaryotic genome annotation pipeline. In *The NCBI Handbook*. National Center for Biotechnology Information (US). https://www.ncbi.nlm.nih.gov/books/NBK169439/.

Turelli, M., Cooper, B. S., Richardson, K. M., Ginsberg, P. S., Peckenpaugh, B., Antelope, C. X., Kim, K. J., May, M. R., Abrieux, A., Wilson, D. A., Bronski, M. J., Moore, B. R., Gao, J.-J., Eisen, M. B., Chiu, J. C., Conner, W. R., and Hoffmann, A. A. 2018. Rapid global spread of wri-like wolbachia across multiple drosophila. *Current Biology*, 28(6): 963–971.e8.

Turner, K. G., Ostevik, K. L., Grassa, C. J., and Rieseberg, L. H. 2021. Genomic Analyses of Phenotypic Differences Between Native and Invasive Populations of Diffuse Knapweed (Centaurea diffusa). *Frontiers in Ecology and Evolution*, 8.

Vasimuddin, M., Misra, S., Li, H., and Aluru, S. 2019. Efficient architecture-aware acceleration of bwa-mem for multicore systems. In *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS)*, pages 314–324.

Walsh, D. B., Bolda, M. P., Goodhue, R. E., Dreves, A. J., Lee, J., Bruck, D. J., Walton, V. M., O'Neal, S. D., and Zalom, F. G. 2011. Drosophila suzukii (Diptera: Drosophilidae): Invasive Pest of Ripening Soft Fruit Expanding its Geographic Range and Damage Potential. *Journal of Integrated Pest Management*, 2(1): G1–G7.

Winkler, A., Jung, J., Kleinhenz, B., and Racca, P. 2020. A review on temperature and humidity effects on Drosophila suzukii population dynamics. *Agricultural and Forest Entomology*, 22(3): 179–192.

Wood, D. E., Lu, J., and Langmead, B. 2019. Improved metagenomic analysis with kraken 2. *Genome Biology*, 20(1): 257.

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., Fu, X., Liu, S., Bo, X., and Yu, G. 2021. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *The Innovation*, 2(3): 100141.

Xu, A., Park, S.-K., D'Mello, S., Kim, E., Wang, Q., and Pikielny, C. 2002. Novel genes expressed in subsets of chemosensory sensilla on the front legs of male drosophila melanogaster. *Cell and tissue research*, 307: 381–392.

Yang, F., Crossley, M. S., Schrader, L., Dubovskiy, I. M., Wei, S.-J., and Zhang, R. 2022. Polygenic adaptation contributes to the invasive success of the Colorado potato beetle. *Molecular Ecology*, 31(21): 5568–5580.

Zhang, G. and Zhang, W. 2019. Protein–protein interaction network analysis of insecticide resistance molecular mechanism in drosophila melanogaster. *Archives of Insect Biochemistry and Physiology*, 100(1): e21523.

Zheng, Y., Shen, W., Bi, J., Chen, M.-Y., Wang, R.-F., Ai, H., and Wang, Y.-F. 2019. Small rna analysis provides new insights into cytoplasmic incompatibility in drosophila melanogaster induced by wolbachia. *Journal of Insect Physiology*, 118: 103938.

Zhou, C., McCarthy, S. A., and Durbin, R. 2022. YaHS: yet another Hi-C scaffolding tool. *Bioinformatics*, 39(1): btac808.

1103 Zhou, S., Luoma, S. E., St. Armour, G. E., Thakkar, E., Mackay, T. F., and Anholt, R. R. 2017. A

1104 drosophila model for toxicogenomics: Genetic variation in susceptibility to heavy metal exposure.

1105 *PLoS genetics*, 13(7): e1006907.