Transient proliferation by reversible YAP-mediated increase of the cyclin D1/p27 ratio

Katherine R. Ferrick[1,2], Yilin Fan[1,2,3], Nalin Ratnayeke[1,2,4], Mary N. Teruel[5], Tobias Meyer[1,2,6, *]

[1]Department of Cell and Developmental Biology, Weill Cornell Medicine, New York, NY, USA

[2]Department of Chemical and Systems Biology, Stanford Medicine, Stanford, CA, USA

[3]Current: Department of Pathology and Center for Cancer Research, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA

[4]Current: Cancer Biology and Genetics Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[5]Department of Biochemistry, Weill Cornell Medicine, New York, NY, USA

[6]Lead contact

[*]Correspondence: tom4003@med.cornell.edu

HIGHLIGHTS

- YAP signaling controls cell cycle entry and exit by up- and down-regulating the cyclin D1/p27 ratio above and below a conserved threshold
- YAP-induced proliferation is intrinsically transient since contact inhibition of YAP suppresses EGFR signaling after a delay to reduce this cyclin D1/p27 ratio
- YAP can still be robustly inhibited after Merlin/NF2 ablation but only at higher local cell density
- The YAP-regulated cyclin D1/p27 ratio is primarily controlled by MEK-ERK rather than mTOR activity

SUMMARY

Hippo-YAP signaling orchestrates epithelial tissue repair and is therefore an attractive target in regenerative medicine. Yet it is unresolved how YAP controls the underlying transient proliferative response. Here we show that YAP-TEAD activation increases the nuclear cyclin D1/p27 protein ratio in G1 phase, towards a threshold level that dictates whether individual cells enter or exit the cell cycle. YAP increases this ratio indirectly, by increasing EGFR and other receptor activities that signal primarily through ERK. Conversely, contact inhibition suppresses YAP activity which gradually downregulates mitogen signaling and the cyclin D1/p27 ratio. Increasing YAP activity by ablating the suppressor Merlin/NF2 reveals a robust balancing mechanism in which YAP can still be inhibited after cell division further increases local cell density. Thus, critical for tissue repair, the proliferation response is intrinsically transient since the YAP-induced and mitogen-mediated increase in the cyclin D1/p27 ratio is reliably reversed through delayed contact inhibition of YAP.

INTRODUCTION

Acute chemical and physical injury inactivates the Hippo pathway, which activates the transcriptional cofactor YAP to increase proliferation, along with inducing metabolic changes and matrix remodeling[1,2]. YAP activation initiates transcription by binding TEAD and other nuclear transcription factors[3]. While YAP and its paralog TAZ are dispensable for the basal proliferation of several tissues, they are crucial for regeneration and repair following injury in tissues such as skin, liver, and kidney[1]. Inhibitors of Hippo pathway proteins are therefore being developed for potential clinical use in tissue repair[4–6].

Hippo signaling is kept active in epithelial sheets and other tissues by cell-cell contacts, such as adherens and tight junctions, and is inactivated by mechanical stress during tissue repair[1,7,8]. Adaptor proteins like Merlin (NF2) contribute to contact inhibition signaling to activate LATS1/2, which is the principal kinase of the Hippo pathway suppressing YAP-TEAD activity[9,10]. Consequently, loss of LATS activity leads to dramatic increases in the size of tissues and organs, and tumorigenesis[9,11,12]. Despite the general finding that YAP activation promotes proliferation, the mechanism how YAP initiates cell cycle entry is unresolved.

Different mechanisms have been described through which YAP signaling could increase proliferation. Constitutive YAP activation increases expression of a broad array of cell cycle genes, including DNA replication factors and regulators of G2/M progression[13–15]. YAP also promotes secretion of mitogens[16,17] and cooperates with AP-1 transcription factors in inducing

target genes[15,18,19]. Likewise, cell cycle entry is regulated by both receptor signaling and contact

inhibition, which may signal independently of YAP[20,21]. Nevertheless, cell cycle entry is

ultimately controlled by cyclin-dependent kinase (CDK) activity in G1 phase which must

phosphorylate and inactivate retinoblastoma protein (Rb), the suppressor of the cell cycle

transcription factor E2F. How YAP controls Rb phosphorylation in G1 is not established: YAP

could induce cell cycle entry by directly regulating cyclin D, indirectly through regulating mitogen

signaling, or by reprogramming the cell state.

As persistent YAP activation induces organ hyperplasia and cancer, YAP-TEAD must

also be robustly inactivated after the tissue repair is completed. Hippo-dependent and Hippo-

independent pathways have been described that can inactivate YAP through phosphorylation,

cytoplasmic sequestration, and degradation[3]. The complexity of both the proliferation initiation

and termination, together with additional roles of YAP in cell differentiation and cell metabolism,

has made it challenging to use animal models to determine how Hippo-YAP signaling

coordinates a proliferation response that must be transient, generating only the cells needed to

recover the original tissue size during repair.

Here we use multiplexed, single-cell immunofluorescence analysis to examine YAP-

regulated proliferation in thousands of epithelial cells in a monolayer. We show that over a wide

range of stimuli, proliferation is controlled through a balance mechanism whereby mitogen

signaling is opposed by contact inhibition suppressing YAP activity. YAP activation drives

proliferation by both increasing the nuclear ratio of the protein level of cyclin D1 and decreasing

the level of the CDK inhibitor p27. Strikingly, YAP-activated cells only enter the cell cycle if the

cyclin D1/p27 ratio exceeds a critical threshold level during the G1 phase, whereas cells arrest

the cell cycle below this threshold. YAP activation increases this cyclin D1/p27 ratio indirectly by

increasing the activity of EGFR and other surface receptors, which in turn signal primarily

through MEK-ERK, rather than the often-implicated mTOR pathway. Ensuring a strong

proliferation response, this inactivation of YAP downregulated EGFR signaling to lower the

cyclin D1/p27 ratio, but only after a delay. Ensuring robust YAP signal termination, even a

strong increase in YAP activity generated by ablation of the upstream Hippo component Merlin

could still be reversed by contact inhibition-mediated YAP inactivation, but only at higher local

cell density. Together, our study identifies a balancing mechanism centered on an increase and

delayed decrease in the cyclin D1/p27 ratio that makes YAP-induced proliferation intrinsically

transient, which is essential for generating a commensurate number of new cells during repair.

RESULTS

Contact inhibition of YAP opposes mitogen stimulation in a balance mechanism controlling cell proliferation

93
94
Even within epithelial monolayers subject to the same mitogen stimuli, there is considerable
heterogeneity in cell cycle signaling[22–24] . We therefore used single cell analysis to distinguish
between various plausible mechanisms of how YAP activation might be linked to contact
inhibition and mitogen signaling to control cell cycle entry. While YAP and TAZ both promote
proliferation, we focus here primarily on YAP as it is generally the limiting component in
regulating proliferation[13,14]. We used hTERT-immortalized human retinal pigmented epithelial
cells (RPE1) that reversibly arrest the cell cycle with serum starvation or contact inhibition[22]. We
use the terms "contact inhibition" and "local cell density" interchangeably since the local cell
density in monolayers closely predicts whether proliferation is suppressed in individual RPE1
cells[22]. We used different concentrations of serum (fetal bovine) to understand the role of
mitogens in the process.
We quantified local cell density by subdividing segmented nuclei for each image into a
10 x 10 grid (~75 x 75 µm) and counting the cells within each block (Figure 1A). Each cell was
then assigned its local density based on the number of cells in its block (Figure S1A, see also
STAR Methods). To assess changes in proliferation, we measured both phosphorylation of
retinoblastoma protein (p-Rb, S807/811), which is a bimodal marker of cyclin-CDK activity[25] that
is low in arrested cells and remains high in cycling cells from G1 through mitosis, and EdU
incorporation into DNA, which occurs during DNA replication in S phase (Figure 1B, S1B).
We first confirmed that increasing the mitogen stimulus from low (0.1% serum) and
intermediate (1% serum) to high (10% serum) levels increased the percentage of proliferating
cells using both metrics of proliferation, p-Rb (Figure 1C) and EdU (Figure S1C). The proportion
of proliferating cells progressively decreased as local cell density increased for high and low
serum levels, supporting that mitogen signaling and contact inhibition oppose each other in
regulating proliferation. This conclusion was also supported by an alternative metric of local cell
density that was based on the area of neighboring cell nuclei[22] within a radius of similar scaling
(Figure S1D). This inverse regulation represents an intuitive balancing mechanism: the inhibition
of proliferation at a given cell density can be overcome by increasing mitogen signaling, and the
increase in proliferation after increasing mitogen signaling can be antagonized once cell division
has increased local cell density.
We next determined how YAP activity changes this balance between mitogen signaling
and contact inhibition by stably expressing a doxycycline (DOX)-inducible, constitutively active

YAP variant (5 regulatory serines mutated to alanine[26], CA-YAP). As leaky expression of CA-

YAP (without DOX induction) was sufficient to increase proliferation (at a similar expression

level compared to the endogenous YAP, Figure S1E), we did not use DOX induction unless

otherwise stated. Even with this weaker YAP expression, the cell density dependence of

proliferation was abolished in CA-YAP cells (Figure 1C), demonstrating that contact inhibition of

proliferation requires contact inhibition of YAP activity. Moreover, CA-YAP cells retained

mitogen sensitivity independent of contact inhibition since increasing serum levels further

increased the percentage of proliferating cells (Figure 1D, S1F).

We next tested whether contact inhibition and mitogen signaling regulate endogenous

YAP activity. As the ratio of the nuclear to cytoplasmic (n/c) YAP and the absolute nuclear level

of YAP are both relative measures of YAP activity[27,28], we measured both parameters as a

function of local cell density (Figure 1E, see also STAR Methods). At 0.1% serum, we observed

the expected inhibition of YAP at increasing cells density when monitoring the YAP n/c ratio or

the nuclear YAP level. At 1% and 10% serum, the YAP n/c ratio was diminished at higher cell

densities, but the nuclear level of YAP was higher than for low serum conditions.

We used this single-cell data to analyze the percent of cells with phosphorylated Rb as a

function of the relative YAP activity. Markedly, using both metrics, increasing serum levels

increased proliferation for cells with identical YAP levels (Figure 1F). Thus, YAP activation and

mitogen stimuli synergistically increase proliferation, and exogenously added mitogens increase

proliferation directly without requiring increased YAP.

Finally, we determined how acute increases and decreases in YAP activity regulate Rb

phosphorylation. Using recently developed inhibitors, we inhibited YAP-TEAD signaling with a

pan-TEAD inhibitor[29], and activated endogenous YAP using a LATS1/2 inhibitor[5,30]. We

confirmed that TEAD inhibition promoted a strong cytoplasmic localization of endogenous YAP

while LATS inhibition triggered its nuclear enrichment (Figure 1G). TEAD inhibition greatly

reduced the fraction of proliferating cells for low but not for high serum stimuli, suggesting that

YAP-TEAD activity is particularly critical for inducing proliferation for low mitogen stimulation

conditions (Figure 1H, S1G). Conversely, LATS inhibition greatly boosted proliferation at low

serum stimuli and had a smaller relative effect at high serum stimuli due to the already high

percentage of proliferating cells (Figure S1H). Finally, this analysis showed that high mitogens

can induce proliferation independently of YAP-TEAD activity since TEAD inhibition could

reverse the portion of proliferation induced by YAP but not a YAP-independent component of

mitogen-induced proliferation (Figure 1H).

Together, these experiments strongly support a model that contact inhibition of YAP

activity and mitogen signaling compete to respectively suppress or promote proliferation over a

wide range of cell densities and mitogen stimuli. However, this balance can be broken in two

ways: by high mitogen stimuli inducing proliferation in cells lacking YAP-TEAD activity and

persistent activation of YAP abolishing contact inhibition of proliferation.

YAP-regulated cell cycle entry and exit is dictated by a cyclin D1/p27 threshold

We next used the quantitative analysis of the balance between contact inhibition and mitogen

signaling to identify the mechanism of how YAP regulation controls cell-cycle entry and exit.

Cells enter the cell cycle during the G1 phase by activating cyclin-CDK complexes to

hyperphosphorylate Rb and derepress the E2F cell cycle transcription program[25]. In contrast, in

cells that exit the cell cycle or stay arrested, cyclin-CDKs stay inactive and Rb remains

dephosphorylated. In monolayers of RPE1 and other epithelial cells, this bimodal Rb

phosphorylation is thought to be primarily controlled by the CDK4/6 activator cyclin D1 and the

cyclin-CDK2/4/6 inhibitor p27[22], and cyclin D1 is generally thought to contribute to the YAP-

mediated proliferation regulation[3]. We included p27 in our analysis since both cyclin D1[31] and

p27 are key regulators of cell proliferation in vivo, with p27 deletion resulting in multi-organ

hyperplasia[32].

Using multiplexed immunofluorescence[33], we computationally gated for G1 cells by their

diploid 2N DNA content and lack of incorporation of EdU (Figure S2A). We then measured the

levels of nuclear localized cyclin D1 and p27 and whether Rb was phosphorylated in each G1-

phase cell. Notably, p27 protein levels are often used as a marker of non-proliferating cells, but

we observed substantial heterogeneity in the nuclear levels of both p27 and cyclin D1 across

individual cells irrespective of Rb phosphorylation (Figure S2B). Increasing the mitogen stimulus

(from 0.1% to 10% serum) resulted in the cell population shifting towards higher average cyclin

D1 levels and lower average p27 levels.

We next determined how local cell density and YAP activity regulate the cyclin D1 and

p27 levels. As local density increased, the median cyclin D1 level of WT cells gradually

decreased, while the median p27 level strongly increased (Figure 2A). This trend was opposite

to the effects of increasing serum, which both globally increased cyclin D1 levels and

suppressed the density-dependent increase in p27 protein. Markedly, the CA-YAP cells lost the

serum and density dependence of both, keeping the levels of cyclin D1 high and p27 low in G1

phase, arguing that YAP activity is not only upregulating the level of cyclin D1 but also

downregulating the level of p27 (Figure 2B).

Explaining why the level of cyclin D1 or p27 alone cannot predict whether Rb is
phosphorylated, only the level of cyclin D1 relative to p27 was tightly correlated with Rb
phosphorylation in individual cells. Only cells with relatively more cyclin D1 than p27 had Rb
phosphorylated (Figure 2C-D, S2C). Thus, even for low mitogen levels, a small fraction of cells
had sufficiently high cyclin D1 and low p27 to have Rb phosphorylated. Conversely, at high
mitogen stimulation, more cells populated the high-cyclin D1/low-p27 state and had Rb
phosphorylated, demonstrating a critical threshold cyclin D1/p27 ratio for proliferation.
Strikingly, in CA-YAP cells with persistently active YAP, Rb phosphorylation remained
closely predicted by the relative nuclear cyclin D1 and p27 levels (Figure 2E, S2D-E), implying
that YAP is signaling through both cyclin D1 and p27 to control proliferation. Notably, we
identified some CA-YAP cells with high cyclin D1 that did not have Rb phosphorylated, but
these cells had increased protein levels of another CDK inhibitor, p21, which we demonstrated
in separate experiments (Figure 2F). Since p21 levels were only weakly regulated by serum
starvation or contact inhibition (Figure S2F-G), p27 is functioning as the primary proliferation
suppressor relevant for contact inhibition and YAP signaling[34,35].
The fundamental role of the cyclin D1/p27 ratio in YAP-regulated proliferation is further
seen in a comparison of different perturbations that change the cyclin D1/p27 ratio and
correspondingly change Rb phosphorylation (Figure 2G). The siRNA-mediated knockdown of
cyclin D1 strongly reduced the proportion of cells with phosphorylated Rb by shifting most of the
CA-YAP cells to a high-p27 and quiescent state, but again preserved the control of Rb
phosphorylation by the cyclin D1/p27 ratio (Figure 2H, S2H). Moreover, knocking down p27 in
WT cells resulted in more cells ending up below the dashed line but still preserved the control of
Rb phosphorylation by the cyclin D1/p27 ratio (Figure 2G). Conversely, in cells where p27 levels
were increased by CRISPRa induction, most cells were above the dashed line and proliferation
was completely suppressed (Figure 2I). However, LATS inhibitor-mediated activation of YAP in
these cells lowered p27 and increased cyclin D1 to push some cells below the threshold cyclin
D1/p27 ratio where they could again proliferate (Figure 2I).
We conclude that the crucial parameter for the YAP-regulated control of cell cycle entry
is the nuclear cyclin D1/p27 ratio in G1 phase, with a threshold dictating whether individual cells
enter or exit the cell cycle (Figure 2J). Moreover, YAP activity and mitogen signaling both
control proliferation through the same downstream cyclin D1/p27 ratio. Notably, we were only
able to identify this molecular mechanism of how YAP controls the cell cycle entry and exit
decision by measuring nuclear p27, cyclin D1 and p-Rb levels within the same, G1-gated single
cell.

YAP regulation gradually changes the nuclear cyclin D1/p27 ratio

We next used RNA sequencing (RNAseq) to distinguish between different potential signaling pathways through which YAP activation may regulate cyclin D1 and p27 levels over time. Previous studies suggested various regulatory mechanisms, such as direct induction of cyclin D1[14], indirect induction of mitogen signaling, or reprogramming the cell state[15,36,37]. We measured gene expression changes up to 36-hours following CA-YAP induction in low serum cells (Figure 3A). Critically, we inhibited CDK4/6 activity in these cells (palbociclib) to keep cells in a state with minimal E2F activation to suppress the otherwise dominating effects of the E2F cell cycle transcription program that may obscure the YAP-specific transcription program.

We confirmed that the conserved YAP target gene signature[38] was significantly enriched after 12-36 hours of induction (Figure 3B), including the canonical YAP target genes *CYR61* and *ANKRD1* that were significantly induced within 6h (Figure 3C, S3B). We also confirmed previously-characterized induction of G1-related target genes *CDK6*[39] and *SKP2*[40] (Figure S3B), proliferation-supporting transcription factors (*MYBL1, ETS1, MYC,* Figure S3C)[15], as well as TEAD and AP1 target genes[18] (*TEAD1/2*, *FOSL1/2, JUN*, Figure S3D). Compared with the fast induction of the YAP target gene *CYR61*, we observed *CCND1* (cyclin D1) expression increasing only after 12-36 hours and only 1.4-fold. In comparison, the previously-reported suppressed gene *DDIT4*[41] rapidly dropped whereas the expression of p27 (*CDKN1B*) and p21 (*CDKN1A*) changed only minimally over time (Figure 3D).

Since there was only a small fold change in cyclin D1 and p27 expression, we next tested whether cyclin D1 and p27 are more strongly regulated at the protein level[42–44] following acute YAP-TEAD activation. LATS inhibitor treatment of serum-starved WT cells increased the nuclear/cytoplasmic YAP ratio and we confirmed the protein-level increase of select YAP targets (CDK6 and SKP2, Figure S3E). LATS inhibition strongly increased cyclin D1 protein levels but also resulted in a small increase in p27 and transient decrease in p21 (Figure 3E, S3F). Over 3-12 hours following LATS stimulation, the increase in the cyclin D1/p27 ratio was paralleled by increasing Rb phosphorylation (Figure 3F-G), and cells progressed to S-phase within 12-24 hours (Figure S3G). Supporting that LATS signals through YAP-TEADs, TEAD inhibition reversed the LATSi-induced changes of these protein levels and reversed the percent of proliferating cells back to control levels (Figure 3F, S3G).

We next inhibited TEAD in proliferating cells to determine how YAP-TEAD inhibition regulates cyclin D1 and p27 levels and Rb phosphorylation in G1 phase to terminate proliferation. We first confirmed that TEAD inhibition of CA-YAP cells, or of LATSi pre-treated

WT cells, rapidly reduced the YAP ratio (Figure S3H). Markedly, however, TEAD inhibition

resulted in only a delayed reduction of the nuclear level of cyclin D1 protein in G1 phase (Figure

3H). Moreover, TEAD inhibition caused a strong and slow increase of the nuclear protein levels

of p27 and a relatively smaller increase in p21 levels across the population. Concordantly, the

mean cyclin D1/p27 ratio and the G1 phase Rb phosphorylation both decreased but only after a

delay with most cells arresting by 24-36 hours (Figure 3I, S3I). Moreover, single-cell analysis

confirmed that the cyclin D1/p27 ratio still predicted whether Rb was phosphorylated in G1 as

individual cells gradually reduced their cyclin D1 and increased their p27 level (Figure 3J).

Thus, while acute YAP activation increases the cyclin D1/p27 ratio and triggers Rb

phosphorylation relatively quickly, YAP inactivation in proliferating cells lowers the cyclin D1/p27

ratio and Rb phosphorylation only after a delay of 12 to 24 hours. This result raises the question

of how YAP activity can induce such a prolonged increase in the cyclin D1/p27 ratio that keeps

Rb hyperphosphorylated in G1-phase cells long after YAP has been inhibited.

YAP activation induces and prolongs proliferation by increasing EGFR signaling

Our RNAseq time course analysis also supported earlier data that YAP activation increases

multiple factors related to receptor signaling[16,45–47] (Figure 4A, S4A). These factors included

receptors and secreted ligands known to regulate cell proliferation including *ERBB3* and *AXL*

receptors[3], along with ligands like *FGF1/2* and the EGFR ligand precursor *HBEGF* (paralog of

AREG)[48,49], providing a potential cell autonomous activation of EGFR and other receptors. We

were intrigued by this possibility since the slow kinetics of the expression changes matched the

delay we observed in the changes of the cyclin D1/p27 ratio in the G1 phase after TEAD

inhibition.

Since expression levels of *AXL* (a direct YAP target gene)[15,50,51], *EGFR,* and *HRAS* were

all gradually increased by YAP activation, along with the EGFR-activating ligand *HBEGF* (Figure

4B), we first considered whether the residual proliferation observed in serum starved cells might

be induced by YAP through activation of EGFR signaling. Indeed, the small portion of WT cells

proliferated in the absence of mitogens could be inhibited by titration of the EGFR inhibitor

gefitinib in the absence of exogenous EGF (Figure 4C). The EGFR inhibition also mirrored the

TEAD inhibition, supporting that YAP-TEAD activity may signal through EGFR at low stimulation

conditions to promote proliferation (Figure S1G). However, EGFR inhibition in CA-YAP cells

could only partially suppress Rb phosphorylation (Figure 4E, Figure S4B), suggesting that

additional receptor signaling pathways are contributing to the YAP-induced proliferation.

We next determined how rapidly TEAD inhibition changes the protein level of EGFR and

AXL. Western blot analysis revealed that their total levels gradually decreased after TEAD

inhibition in CA-YAP cells over a similar time scale as the loss of proliferation (Figure 4D). Since

EGFR levels were YAP dependent, we further tested whether YAP activation increases EGF

sensitivity. Indeed, while EGF stimulation caused only a small increase in Rb phosphorylation in

WT cells, Rb phosphorylation was much more strongly increased in CA-YAP cells (Figure 4E).

Furthermore, while both WT and CA-YAP cells had similarly low basal AKT and ERK

phosphorylation, the CA-YAP cells kept their phosphorylated AKT and ERK levels higher for at

least 10 hours after EGF stimulation compared to WT cells, again consistent with sensitized

EGF signaling (Figure 4F). As a control, we also confirmed that TEAD inhibition of CA-YAP cells

prevented this prolonged AKT and ERK signaling.

Finally, we tested whether there is an extended period of enhanced EGF signaling also

after transient LATS inactivation upstream of YAP. We used WT cells pre-treated with LATSi

and then either maintained (sustained) or washed out the LATSi prior to a 14-hour addition of

EGF (Figure 4G). While LATSi washout reduced the nuclear/cytoplasmic YAP ratio and nuclear

YAP levels back to control conditions (Figure 4G, S4C), cells retained sensitivity to EGF for at

least 14 hours, as evidenced by the increased p-Rb (Figure 4H, Figure S4D), S6

phosphorylation (p-S6, a readout of PI3K/mTOR activity, Figure 4I, S4E), and nuclear FRA1

levels (a readout of integrated MEK-ERK activity[52], Figure 4J). In support of a general

prolongation of proliferation signaling, we also found that transient LATSi treatment sustained

the fraction of G1 cells with phosphorylated Rb after HGF and LPA stimulation (Figure S4F).

We conclude that YAP activity increases the activity of EGFR in the absence of added

EGF in an autocrine manner[16,48], while also increasing the sensitivity of cells to exogenous EGF

stimulation. This sensitization extends to other receptors since gefitinib could only suppress a

fraction of the YAP-induced proliferation response. Notably, the increased EGFR signaling

persisted for over 14 hours after YAP inactivation, suggesting that YAP is inducing a prolonged

proliferation response since YAP inactivation only terminates proliferation and signaling after a

delay.

YAP activation signals primarily through MEK-ERK to increase the cyclin D1/p27 ratio

We next examined which signaling pathways downstream of the YAP-induced receptor

activation control the cyclin D1/p27 ratio. We focused on PI3K-AKT-mTOR and MEK-ERK

signaling[45,53], since various functions of YAP activation are mediated through mTOR

activation,[54,55] and both mTOR and MEK-ERK can regulate cyclin D1 and p27 levels[20,34,56]

(Figure 5A). We measured phosphorylation of AKT and S6 protein (p-S6) to monitor PI3K

pathway activity and protein levels of the transcription factor Fra1 to monitor the ERK pathway.

Serum-starved CA-YAP cells displayed both elevated p-AKT(S473) and p-S6(240/244) as well

as Fra1 levels (Figure 5B-C), which similarly depended on YAP-TEAD activity, as TEAD

inhibition reduced their phosphorylation and protein levels in CA-YAP cells (Figure 5D).

To assess the functional importance of PI3K-AKT-mTOR signaling, we inhibited PI3K

(LY-294002) or AKT (MK2206), which had only small inhibitory effects on Rb phosphorylation in

CA-YAP cells (Figure 5E), despite reducing the p-AKT(S473) and p-S6(240/244 and 235/236)

levels (Figure S5A-B). Nevertheless, inhibiting mTORC1/2 (mTORi, Torin2) caused an

approximately 50% reduction in proliferation (Figure 5E). In contrast, inhibition of the obligate

upstream regulator of ERK, MEK (MEKi, PD0325901), completely suppressed Rb

phosphorylation and readouts of ERK activity (Figure 5F, S5C-D).

In support of YAP regulating cyclin D1 protein levels in G1 phase through mTOR and

MEK/ERK, rather than doing so directly, MEK inhibitor alone (Figure S5D) or the combined

inhibition of MEK and mTOR fully reduced the cyclin D1 level in CA-YAP cells to the level of WT

cells (Figure 5G, S5E). A time course analysis showed that the drop in cyclin D1 levels is rapid

after mTOR inhibition but slower and stronger after MEK inhibition even though both inhibitors

rapidly suppressed p-S6 and p-ERK, respectively (Figure 5H-I). As mTOR can rapidly control

translation of cyclin D1 mRNA[57], this fast regulation was anticipated. In contrast, mTOR

inhibition did not alter p27 levels but MEK inhibition not only reduced cyclin D1 levels but also

slowly and strongly increased p27 levels, explaining why MEK inhibition suppresses Rb

phosphorylation more completely than the regulation of mTOR. Thus, transient YAP-mediated

proliferation responses result from an increase and delayed decrease of the activity of EGFR

and other receptors, which in turn transiently increases the cyclin D1/p27 ratio through

reversibly regulating MEK-ERK and, to a lesser degree, mTOR signaling pathways.

Partial activation of YAP from ablating Merlin can be reversed by increased contact

inhibition

To evaluate the robustness of the termination of YAP activation in monolayers, we increased

YAP activity by ablating Merlin, an adaptor protein and tumor suppressor that indirectly activates

LATS1/2[58–60]. We generated clonal Merlin knockout RPE1 cells (Merlin KO) using

CRISPR/Cas9. As expected, these cells had increased nuclear localization of YAP that could be

suppressed by TEAD inhibition (Figure S6A-B).

However, unlike CA-YAP, Merlin KO increased proliferation without abolishing contact

inhibition of proliferation. Instead, a higher cell density was required to inhibit proliferation

compared to WT, both at low and high serum levels (Figure 6A, S6C). Moreover, inhibiting

TEAD in Merlin KO cell suppressed Rb phosphorylation in both cases to levels similar as TEAD-

inhibited WT cells (Figure S1G), supporting a model whereby Merlin regulates proliferation

primarily through YAP-TEAD activity[9]. Importantly, we found that YAP activity in Merlin KO cells

could be as effectively inhibited by contact inhibition as in WT cells but required a higher local

cell density to reach the same YAP inhibition (both at low and high serum levels, Figure 6B).

We next examined the role of EGFR in the Merlin KO cells, as a possible mechanism

mediating serum-independent proliferation downstream of the higher YAP signaling. Like CA-

YAP cells, Merlin KO cells displayed activated EGFR signaling in the absence of added EGF,

since the EGFR inhibitor gefitinib reduced the percentage of proliferating cells (Figure 6C, S6D).

Similar to the transiently LATS-inhibited cells, the Merlin KO cells had increased sensitivity to

EGF while still retaining a strong local cell density dependence (Figure 6E). In addition, we

found increased proliferation in response to several other receptor tyrosine kinase ligands

(Figure S6D), again consistent with YAP inducing a broad upregulation of receptor signals not

limited to EGFR.

Finally, we found that Merlin KO cells still suppressed cyclin D1 and increased p27

levels at higher local cell density (Figure 6E), demonstrating that these cells suppress

proliferation through the same mechanism as WT cells: inhibiting YAP activity and

downregulating mitogen signaling and the cyclin D1/p27 ratio (Figure 6F). However, Merlin KO

cells employ alternative mechanisms from inactivating Merlin[61,62] to contact inhibit YAP.

We conclude that even though constitutive YAP activation abolished contact inhibition of

proliferation, cells can robustly contact inhibit YAP-TEAD even if Hippo signaling is reduced and

YAP activation increased. However, cells that have such an induced YAP activity now require

higher local cell density to suppress YAP activity and proliferation.

DISCUSSION

Our study shows that contact inhibition of YAP and mitogen signaling control the cyclin D1/p27

ratio and cell cycle entry in G1 phase through a balance mechanism that shapes an intrinsically

transient YAP-induced proliferation response (Figure 6G). Our study supports a model where

YAP activation controls proliferation by gradually increasing the expression of HB-EGF and

other RTK ligands, as well as various receptors and known regulators of mitogen signaling[3,54,63].

This regulation explains how YAP activation is increasing EGF and other mitogen signals in a

cell autonomous manner (Figure 4C), while simultaneously sensitizing cells to exogenous

stimulation by EGF and other mitogens (Figure S4F, S6E), which is likely relevant for the role of

YAP activation in the inflammatory injury response[64,65]. We further show that YAP activation

increases downstream signaling through MEK-ERK signaling and, to an unexpectedly lesser

degree, mTOR signaling, to control the nuclear protein levels of cyclin D1 and p27 (Figure 5I).

Most importantly, our study revealed a predominant role of both cyclin D1 and p27 in YAP-

induced proliferation. We showed that YAP-stimulated cells decide in the G1 phase whether to

enter the cell cycle by increasing the nuclear cyclin D1/p27 ratio above a critical activity

threshold, or to remain quiescent or exit the cell cycle by staying below this threshold (Figure

2C-E).

Moreover, as the YAP-induced cell division increases the local cell density, we show that

the ensuing contact inhibition of YAP-TEAD causes a delayed inhibition of mitogen signaling,

which lowers the cyclin D1/p27 ratio to terminate the proliferation response. Thus, there are two

delay mechanisms ensuring a prolonged proliferation response: cells wait to inhibit YAP until

cell division has increased the local cell density, then the resulting contact inhibition of YAP

lowers the cyclin D1/p27 ratio and arrests proliferation after a delay (Figure 6H). YAP signals

often fluctuate[66–68], and the slow integration of YAP signals further ensures that the proliferation

response is robustly started and ended. Slow integration provides a filter mechanism which

prevents short-lived YAP activity changes from triggering unwanted cell cycle entry or

premature cell cycle exit.

We also evaluated the robustness of YAP inactivation by using ablation of Merlin/NF2, a

suppressor of Hippo signaling to YAP[69–71]. While Merlin has been shown to have additional

signaling functions at the plasma membrane, our study shows that the increased EGFR

signaling and proliferation in Merlin KO cells can be reversed by TEAD inhibition (Figure 6A),

suggesting that Merlin primarily signals through YAP-TEAD to control EGFR signaling, the

cyclin D1/p27 ratio, and proliferation. In support that the termination of YAP signaling and

proliferation is robust, our study shows that contact inhibition is merely shifted but not abolished

when YAP activity is increased by ablation of Merlin, and cells can still effectively terminate

proliferation but at higher local cell density. Moreover, since the loss of Merlin can boost

proliferation without a loss of contact inhibition, this result also suggests that Merlin has a

unique advantage as a drug target, since a potential Merlin inhibitor would trigger a self-limiting

proliferation response. This could offer a safer alternative strategy to LATS inhibition, which was

recently shown to cause a widespread and uncontrolled proliferation response and de-

differentiation in mice[6].

We also found that inhibiting MEK-ERK, rather than mTOR, in CA-YAP cells greatly
reduces the cyclin D/p27 ratio and stops proliferation. This crucial role of MEK-ERK in YAP-
induced proliferation was surprising, since oncogenic YAP mutants can drive resistance to
KRAS inhibition and KRAS primarily regulates proliferation through MEK-ERK and mTOR
activation[29,45,72]. The resistance to KRAS inhibitor may therefore by mediated by YAP-mediated
direct regulation of cell cycle entry. We confirmed that YAP activation can also directly induce
cyclin D1 as well as SKP2, which can degrade p27[73,74]. Indeed, we found that this upregulation
may be particularly important at high levels of YAP activation[75,76], since DOX induction of CA-
YAP in high serum conditions can still increase cyclin D1 and proliferation when MEK is
inhibited (Figure S6F-G).
Our study also supports that YAP regulates proliferation through additional modulatory
pathways described in previous studies. YAP gradually induces a number of Hippo pathway
components such as LATS, Merlin and Angiomotin-like genes[77,78] (Figure S6H-I), which likely
contribute to the delayed downregulation of YAP along with the contact inhibition of YAP.
Moreover, our finding that persistent activation of YAP suppresses contact inhibition of
proliferation is supported by our own and previous RNAseq data[15,19] showing that YAP
activation not only increases mitogen signaling regulators but also reduces the expression of
contact inhibition regulators at the plasma membrane such as ephrins[49] and protocadherins[79,80]
(Figure S6J-K), which may mediate cell density sensing. Finally, we showed the stress-
regulated p21 CDK inhibitor can be rate-limiting in suppressing proliferation in some YAP-
activated cells (Figure 2F), supporting a redundant role of both p27 and p21 inhibitors in
promoting cell cycle exit.
451

Concluding remarks
YAP activation must be transient to drive the epithelial regenerative response. Our study
identifies a self-limiting control mechanism that allows YAP to trigger a commensurate
proliferation response. We show that transient proliferation responses are initiated by YAP
activation increasing EGF and mitogen signaling to increase the cyclin D1/p27 ratio above a
critical threshold. The delayed contact inhibition-mediated inactivation of YAP then terminates
proliferation by gradually lowering mitogen signaling and the cyclin D1/p27 ratio below this
critical threshold (Figure 6H). In contrast, we demonstrate that persistent inactivation of
LATS1/2 or activation of YAP activity abolishes contact inhibition and promotes persistent
mitogen signaling that keeps the cyclin D1/p27 ratio high, explaining how oncogenic mutations
that constitutively inactivate LATS1/2 or activate YAP elicit unrestricted proliferation.

Limitations

We identified cyclin D1 and p27 as the primary regulators of Rb phosphorylation for RPE1 cells
in response to contact inhibition and serum starvation with an added contribution from p21. In
other cell types and cancer models, additional phosphorylation, degradation, and translation-
mediated control steps may further contribute to the timing and overall level of proliferation. We
also studied EGFR signaling as a representative effector of YAP, but our data suggest a
pleiotropic effect on additional receptors and cell adhesion pathways, which likely affect the
activity of Merlin and other adaptor proteins at the plasma membrane and feed into the
dynamics of different mitogen responses. Furthermore, as our mechanistic studies required in
vitro, single-cell experiments, how these control mechanisms of the proliferation response apply
to various animal models of epithelial repair remains to be determined. Finally, in addition to
Merlin, other adaptor proteins like angiomotin and cytoskeleton regulators like RhoA, can
regulate YAP activity in response to cell density changes. Future work is also needed to
understand specific contributions from such regulators and when they engage to restrict or
promote YAP-TEAD activity.

Acknowledgments

We thank Nathaniel Kastan and James Hudspeth for early access to a previous iteration of the
LATS1/2 inhibitor (TRULI); Alexander Pfannenstein, Marielle Koeberlin, and Sanjeev Sharma
for technical support; the Stanford Shared FACS Facility and Weill Cornell FACS Facility for cell
sorting; the Weill Cornell Genomics Core for library prep and RNA-sequencing; Marielle
Koeberlin for critical reading of the manuscript; and James Ferrell, Alexander Dunn, Daniel
Jarosz, Ovijit Chaudhuri, Elizabeth Kiddie, Members of the Meyer and Teruel laboratories for
technical feedback and helpful discussions. This work was funded by an NIH T32 training grant
(5 T32 GM 113854-3), NIH grants T.M. (R35 GM12702601) and M.N.T. (R01 DK131432); K.F.
was supported by an NSF Graduate Research Fellowship (DGE-1656518).

Author contributions

492 Conceptualization, K.F., Y.F., and T.M.; Methodology, K.F., N.R., Y.F., and T.M.; Experiments,
493 visualization, and analysis, K.F.; Manuscript preparation, K.F., M.N.T., and T.M.; Funding
494 acquisition, T.M., and M.N.T.

Declaration of interests

497   The authors declare no competing interests.

Resource availability

500   Immunofluorescence protocols, and cell lines are available upon request to the lead contact.

501   Plasmids generated in this study will be deposited to Addgene.

Data and code availability

504   MATLAB image-processing scripts and iterative imaging analysis pipeline are previously

505   published[22,81] and publicly available on Github (https://github.com/MeyerLab/image-analysis-

506   [ratnayeke-2022](https://github.com/MeyerLab/image-analysis-ratnayeke-2022)).

507   Additional information is available upon request from the lead contact.

Supplemental information titles and legends

510   Figures S1-S7 and Tables S1-S2

STAR METHODS

Cell culture

All experiments were performed with human retinal pigmented epithelial cells (RPE1-hTERT

immortalized, ATCC Cat# CRL-4000) unless specified otherwise. RPE1 were cultured in

DMEM/F12 growth media with HEPES (Thermo Scientific Cat#11039047), supplemented with

10% fetal bovine serum (Millipore Sigma Cat#F4135-500ML) and passaged using trypsin-EDTA

(0.05%, Gibco Cat# 25300054). Human mammary epithelial cells (MCF10A, ATCC Cat# CRL-

10317, RRID:CVCL_0598) were cultured in DMEM/F12 growth media with HEPES

supplemented with 5% horse serum (Gibco Cat# 16050122), 20 ng/mL EGF (PeproTech Cat#

C8052), 0.5 µg/mL hydrocortisone (Sigma Cat#H0888), 100 ng/mL cholera toxin (Sigma Cat#

C8052), and 10 µg/mL insulin (Sigma Cat#I1882). For serum starvation experiments, cells were

first washed with serum-free DMEM/F12 media, then maintained in serum-free (0%) or 0.1%

serum-containing DMEM/F12 (serum-starvation conditions). 0.1% serum-containing media had

similar proliferation as 0% serum-containing media, but cells were slightly more adherent to

glass-bottom plates for subsequent immunofluorescence protocols and therefore used for a

subset of experiments. For mitogen-release experiments, mitogens and growth factors (EGF,

HGF, and FGF at 25 ng/mL and LPA at 1 µM) were diluted in starvation media. HEK-293T cells

used for lentivirus production were cultured in DMEM growth media (Gibco Cat# 11995065)

supplemented with 10% fetal bovine serum. Cells were cultured at 37 °C and 5% CO_2.

Plasmid generation

pSpCas9(BB)-2A-Puro (PX459) V2.0 was a gift from Feng Zhang (Addgene # 62988,

RRID:Addgene_62988). Phosphorylated and annealed sgRNA oligos against *NF2* (see Cell line

generation below) were cloned into restriction enzyme-digested PX459[82].

pQCXIH-Myc-YAP-5SA was a gift from Kun-Liang Guan (Addgene #33093, RRID:

Addgene_33093). Doxycycline-inducible, myc-epitope-tagged CA-YAP (YAP5SA) was

generated using Gibson assembly of PCR-amplified myc-YAP5SA with restriction enzyme-

digested plasmid backbone derived from a bicistronic vector with a TetOn promoter driving

target gene expression and PGK-driven rtTA expression, pCW-Cas9 (Addgene #50661, RRID:

Addgene_50661, a gift from Eric Lander and David Sabatini).

Cell line generation

Merlin knockout cells were generated by transient transfection of parental RPE1 cells with

pX459 (2 ng/uL, containing gRNA scaffolds targeting Merlin/Nf2) and Lipofectamine 2000

(1:500, Thermo Cat# 11668019) in Opti-MEM (Fisher Scientific Cat#31-985-070) and growth

media overnight, then exchanged for growth media with puromycin selection (10 μg/mL,

Invivogen Cat#ant-pr-1) for 72h. After a recovery period in selection-free media, cells were

diluted to 0.5 cells/100 μL and seeded to 96-well plates for single-cell clonal expansion. 24

single-cell clones were further expanded for preliminary knockout assessment by western blot

for loss of Merlin protein expression. 2 clones (Merlin KO-clone 1 forward:

CACCGCACAGTGGCCTGGCTCAAAA, reverse: AAACTTTTGAGCCAGGCCACTGTGC ;

Merlin KO-clone 2 forward: CACCGTGATTTGGTGTGCCGGACTC, reverse:

AAACGAGTCCGGCACACCAAATCAC) were selected for validation by genomic DNA

purification (Qiagen CAT#69504), PCR of the Merlin locus and sequencing. After initial

characterizations confirming Merlin KO-clone 1 and Merlin KO-clone 2 were similarly

proliferative (data not shown), we proceeded to use KO-clone 1 for all subsequent experiments

in this study.

Lenti-EF1a-dCas9-VPR-Puro (Addgene # 99373, RRID: Addgene_99373) was a gift

from Kristen Brennard and stably expressed in RPE1 cells using puromycin selection (10

ug/mL, Invivogen Cat#ant-pr-1). CA-YAP and dCas9-expressing cells were generated using

third-generation lentiviral transduction. Lentivirus was generated in HEK-293T cells by

transfection with packaging and envelope plasmids pMDLg/pRRE (Addgene # 12251, RRID:

Addgene_12251), pRSV-rev (Addgene #1225, RRID: Addgene_1225), and pCMV-VSV-G

(Addgene #8454, RRID: Addgene_8454) in Lipofectamine 2000 and Opti-MEM. After 72h

transfection, supernatant was passed through a 0.22 uM filter (Millipore Cat# SCGP00525) and

concentrated using 100 kDa centrifugal filters (Millipore Cat# UFC910024). Lentivirus was then

frozen or directly transduced into cells. 72h after lentiviral transduction, cells were split into

antibiotic-selection media or, if expressing a fluorescent construct, expanded prior to sorting by

a BD Influx cell sorter (Stanford Shared FACS Facility, Weill Cornell FACS Facility).

siRNA, sgRNA, and plasmid transfection

All oligo sequences are listed in Table S1. Pooled siRNA (ON-TARGETplus, Horizon Discovery

Biosci.) were used for siCtrl (Non-targeting Control, Horizon Cat#D-001810-10-05), siCycD1

(siCCND1, Horizon Cat# L-003210-00-0005), and sip27 (siCDKN1B, Horizon Cat# L-040178-

00-0005). For CRISPRa-mediated overexpression of genes through targeting of endogenous

promoters, pooled sgRNA (Horizon) were used for sgCtrl (non-targeting control, Horizon Cat#

U-009500-10-05), sgCycD1 (sgCCND1, Horizon Cat# P-003210-01-0005), and sgp27

(sgCDKN1B, Horizon Cat# P-003472-01-0005) with tracRNA (Edit-R tracrRNA, Cat# U-002005-

05). RPE1 cells were transfected with siRNA and sgRNA using DharmaFECT 1 (1:500 in Opti-

MEM, Dharmacon Cat# T-2001-03) according to manufacturer protocol using 20 nM siRNA, or

sgRNA and tracrRNA (in 1:1 ratio, according to manufacturer protocol). Cells were incubated in

transfection mixture for 6-12h in growth media, then replaced with media appropriate for the

experiment conditions and assessed 24-48h after transfection.

Drugs and mitogens

Human growth factors and mitogens were dissolved in sterile, ultrapure water (Fisher Scientific

Cat#10977023) and used at the following concentrations unless otherwise stated: 25 ng/mL

recombinant EGF (PeproTech cat#AF-100-15), 25 ng/mL recombinant HGF (Life Technologies

Cat#PHG0254), 25 ng/mL recombinant FGF basic (R&D Systems Cat#233-FB-025), 1 uM LPA

(oleoyl-L-alpha-lysophosphatidic acid, Sigma-Aldrich Cat#L7260-1MG).

Drugs were dissolved in DMSO (Sigma-Aldrich Cat# D2650) and used at the following

concentrations unless otherwise stated: 0.5 µM TDI-011536 (Selleck Chemicals Cat#E1314,

CAS#2687970-96-1, LATSi), 5 µM TRULI (a gift from James Hudspeth, CAS#1424635-83-5), 1

µM GNE-7887 (ChemieTek Cat#CT-GNE7883, CAS# 2648450-42-2,TEADi), 10 µM gefitinib

(Selleck Chemicals Cat#S1025, CAS#184475-35-2, EGFRi), 10 µM LY294002 (Cayman

Chemical Cat#70920, CAS#154447-36-6 , PI3Ki), 200 nM MK2206 (Cayman Chemical

Cat#11593, CAS#1032350-13-2, AKTi), 100 nM Torin2 (Cayman Chemical Cat#14185,

CAS#1223001-51-1, mTORi), 100 nM PD0325901 (Selleck Chemicals Cat#S1036,

CAS#391210-10-9, MEKi), 1 µM palbociclib (PD-0332991, Selleck Chemicals cat#S1116,

CAS#827022-32-2, CDK4/6i), 0.5 ug/mL doxycycline hyclate (Sigma-Aldrich cat#D9891,

CAS#24390-14-5, DOX), 10 µM 5-ethynyl-2'-deoxyuridine (EdU, Cayman Chemical Cat#20518,

CAS# 61135-33-9). For drug treatments and time courses, control conditions were treated with

DMSO, except mitogen release controls (mock), which were dissolved in water. For all EdU-

incorporation, cells received an EdU spike-in (10 µM, in serum-free media) for 12-15 min prior to

fixation.

Western blotting

Cells grown in 6-well plates were washed once in cold, sterile PBS and lysed in chilled, 1x RIPA

buffer (Boston BioProducts Cat#BP-115-5X) or non-denaturing lysis buffer (20 mM HEPES, 150

mM NaCl, 2 mM EDTA, 0.1% IGEPAL-CA630) containing HALT protease and phosphatase

inhibitor cocktail (1:100, Thermo Fisher Cat#78439). Lysates were harvested on ice using cell

scrapers, sheared by syringe (25G) and centrifuged at 15,000 RCF for 15 min at 4C.

Supernatant was stored at -20C. Protein concentration was calculated by Pierce BCA Protein

Assay (Thermo Fisher Cat#23225) according to manufacturer protocol and a BSA standard

curve.

10-30 ug protein was loaded onto 7.5%, 12%, or 4-20% Mini-PROTEAN TGX gels for

resolution by SDS-PAGE in 1x Laemmli sample buffer (Fisher Scientific, Cat#AAJ60660AC, or

Bio-Rad Cat#1610737) alongside Page RulerPlus (Thermo Scientific Cat#26620) molecular

ladders in Tris/Glycine/SDS running buffer (Bio-Rad Cat#161-0772). Gels were run at 150V for

1h then transferred to 0.2 um PVDF membranes by semi-dry transfer (Bio-Rad Trans-Blot SD,

Cat# 1703940) in Tris/Glycine buffer (Bio-Rad Cat# 1610734) with 10% methanol, or semi-dry

transfer (Bio-Rad Trans-Blot Turbo Cat# 1704150) with Trans-Blot Turbo Transfer Packs (Bio-

Rad Cat# 1704156). Membranes were washed in TBST (20 mM Tris, pH 7.5, 150 mM NaCl,

0.1% Tween-20), blocked for 30 min in 5% milk/TBST (VWR Cat#10128-602), and incubated at

RT for 3h or overnight at 4C in primary antibody dilutions in 5% BSA/TBST + 0.01% NaN3.

Primary antibodies used were rabbit anti-AXL (1:1000, Cell Signaling Technology Cat# 8661,

RRID:AB_11217435), rabbit anti-EGFR (1:1000, Cell Signaling Technology Cat# 4267,

RRID:AB_2246311), rabbit anti-GAPDH (1:1000, Cell Signaling Technology Cat# 5174,

RRID:AB_10622025), mouse anti-Vinculin (1:1000, Thermo Fisher Scientific Cat# 14-9777-80,

RRID:AB_2573027), rabbit anti-Merlin (1:1000, Cell Signaling Technology Cat# 12888,

RRID:AB_2650551) (see also Table S2).

After 2x washes in TBST, membranes were incubated with anti-rabbit or anti-mouse
HRP-conjugated secondary antibodies (1:5000, CST#7074, RRID: AB_2099233 or CST#7076,
RRID: AB_330924) or fluorophore-conjugated secondary antibodies (1:10,000, LI-COR
Biosciences, IRDye 680 LT CAT#925-68020, or IRDye 800CW, CAT#925-32210) in 5%
BSA/TBST for 1h at RT. Membranes treated with enhanced chemiluminescence substrate
(Thermo Fisher Cat#PI34080) were detected by film (Thomas Sci. Cat#EK-5130) or Licor
Odyssey Fc. Membranes treated with fluorophore-conjugated secondary antibodies were
imaged on a Licor Odyssey Fc. Normalized protein level changes were calculated using FIJI [83]
by measuring the integrated intensity of a set box size encompassing the protein band. The
background signal was measured by applying the same box to an adjacent region of the blot
and subtracted from the median protein intensity. An equivalent box size was used for
reference/loading control protein bands as the protein of interest.
645
Immunofluorescence
Cells were plated on bovine fibronectin-coated (1:75 in PBS, Sigma-Aldrich F1141,
CAT#F1141) or collagen-coated (1:100 in PBS, Advanced Biomatrix Cat#5005-B), 96-well,
glass-bottom plates (Cellvis CAT# P96-1.5h-N). Cells were fixed in 4% PFA (in PBS, Fisher
Scientific Cat#AA433689) for 15 min at room temperature (RT), washed 3x with PBS,
permeabilized in 0.2% Triton X-100 for 15 min, and blocked in blocking solution (10% FBS, 5%
BSA, 0.1% Triton X-100, 0.01% NaN3 in PBS) for 1h at RT. For conditions with EdU
incorporation (10 µM in basal medium, 12-15 min at 37C, Cayman Chemical Cat#20518) prior
to fixation, cells were washed 1x with PBS after the blocking step and treated with a click
chemistry solution (2 mM CuSO4, 20 mg/mL sodium ascorbate, 3 µM AFDye picolyl azide (488,
568, or 647, Click Chemistry Tools Cat# 1276, 1292, or 1300 respectively) in TBS (50 mM Tris,
150 mM NaCl, pH 8.3) for 25 min. Following a 1x PBS wash, cells were incubated with primary
antibody for 3h at room temperature or overnight at 4C (see also Table S2) in blocking solution,
then washed 3x in PBS. Fluorophore-conjugated secondary antibodies (Thermo Fisher
Scientific, goat anti-mouse, goat anti-rabbit, 1:1000 in blocking solution) and Hoechst 333342
(Thermo Fisher Scientific Cat#H3570, 1:10,000) were incubated on cells for 1h at RT, then
washed 3x in PBS. Wells were imaged in 1x PBS or 1x N-acetylcysteine (NAC, 700 mM in
ddH2O, pH 7.4, Sigma-Aldrich A7250) imaging buffer for iterative immunofluorescence
protocol[33] (4i).
For the 4i protocol, wells were then washed 1x in dH2O, treated with elution buffer (0.5M
L-glycine (Sigma-Aldrich Cat#G2879), 3M urea (Sigma-Aldrich Cat#U5378), 3M guanidine

hydrochloride (Sigma-Aldrich Cat# G3272), 70 mM TCEP-HCl (Goldbio Cat# TCEP50) in

ddH2O, pH 2.5) for 20 min on an orbital shaker. After 1x PBS wash, elution was confirmed by

fluorescence microscopy. Cells were then blocked with 4i blocking solution (1% BSA + 150 mM

maleimide (Sigma-Aldrich Cat# 129585) in PBS) for 30 min, then incubated with the second

round of primaries (in blocking solution) for at least 3h RT or overnight at 4C. Secondary

staining, imaging in NAC, and subsequent elution and 4i blocking steps were repeated up to 5

times. Control wells with secondary antibody treatment alone were used to confirm proper

elution of previous antibody rounds. Antibody list and dilutions used for immunofluorescence are

listed below and in Table S2:

Immunofluorescence Antibodies

The following primary antibodies were used exclusively for round 1 of immunofluorescence

staining: rabbit anti-CDK6 (1:1000, Abcam Cat#ab124821, RRID: AB_10999714), rabbit anti-

YAP (1:400, Cell Signaling Technology Cat# 14074, RRID:AB_2650491), mouse anti-p27

(1:1000, Cell Signaling Technology Cat#3698, RRID:AB_2077832), mouse anti-p21 (1:500, BD

Bioscience Cat#556430, RRID:AB_396414), rabbit phospho-AKT(S473)(1:400, Cell Signaling

Technology Cat#4060, RRID:AB_2315049), rabbit phospho-MAPK1/2(Thr202/Tyr204)(1:400,

Cell Signaling Technology Cat#4370, RRID:AB_2315112), mouse phospho-S6(235/236)(1:800,

Cell Signaling Technology Cat#62016, RRID:AB_2799618).

686   The following primary antibodies were used for any round of immunofluorescence:

mouse anti-cyclin D1 (1:200, Santa Cruz Biotech Cat#sc-8396, RRID:AB_627344), mouse anti-

FRA1 (1:200, Santa Cruz Biotech Cat#sc-28310, RRID: AB_ AB_627632), rabbit anti-Myc

(1:800, Cell Signaling Technology Cat#5605, RRID:AB_1903938), mouse anti-Myc tag (1:8,000,

Cell Signaling Technology Cat#2276, RRID:AB_331783), mouse anti-N-cadherin (1:200, Santa

Cruz Biotech Cat#sc-393933, RRID: AB_2832921), rabbit anti-p27 (1:1600, Cell Signaling

Technology Cat#3686, RRID:AB_2077850), rabbit anti-p21 (1:2500, Cell Signaling Technology

Cat#2947, RRID:AB_823586), rabbit anti-phospho-Rb(Ser807/811)(1:2500, Cell Signaling

Technology Cat#8516, RRID:AB_1117658), rabbit anti-phospho-S6(240/244)(1:2500, Cell

Signaling Technology Cat#5364, RRID:AB_10694233), rabbit anti-Skp2 (1:800, Cell Signaling

Technology Cat#2652, RRID:AB_11178941), mouse anti-YAP (1:200, Santa Cruz Biotech.

Cat#sc-101199,RRID: AB_1131430). Antibodies were previously validated in RPE1[22] and for

4i[84], and in this study by drug treatment (including AKT, mTORC1/2, MEK, and LATS1/2

inhibitors).

The following secondaries were used in variable combination (488, 514 or 568, and
647), at 1:1000: goat anti-mouse IgG Alexa Fluor 488 (Thermo Fisher Scientific Cat#A11029,
RRID:AB_2534088), goat anti-mouse IgG Alexa Fluor 514 (Thermo Fisher Scientific
Cat#A31558, RRID:AB_1037789), goat anti-mouse IgG Alexa Fluor 568 (Thermo Fisher
Scientific Cat#A-11031 RRID:AB_144696), goat anti-rabbit IgG Alexa Fluor 568 (Thermo Fisher
Scientific Cat#A11036, RRID:AB_1056366), goat anti-rabbit IgG Alexa Fluor 647 (Thermo
Fisher Scientific Cat#A21245, RRID:AB_23813).
707
Microscopy
For automated epifluorescence microscopy, cells were imaged on a Ti2-E inverted microscope
(Nikon) using penta-band (DAPI/CFP/YFP/TRITC/Cy5, Chroma: 89405) Sedat filter sets with an
LED light source (Lumencor Spectra X or Spectra III), Hamamatsu ORCA-fusionBT or ORCA-
Flash4.0 V3 sCMOS camera, and 20x objective (Nikon CFI Plan Apo Lambda, 0.75 NA) in a
humidified chamber at 37C. 25-30 tiled images were acquired per well in 16-bit mode with 2x2
binning. Fixed cells were imaged in 1x N-acetylcysteine imaging buffer. Fluorophores for
secondary antibodies were previously optimized to minimize bleed-through[81].
716
RNA sequencing experimental conditions
WT and doxycycline-inducible YAP5SA cells were seeded at 38,000 cells per well of 96-well,
optical-grade plates (Sigma-Aldrich Cat#CLS3904) in 1.0% FBS. After ~2h to allow time for
adherence to the tissue culture surface, cells were washed with 0.1% FBS then pre-arrested
with CDK4/6 inhibitor (palbociclib, 1 uM, SelleckChem CAT#S1116) in either 1.0% FBS or 0.1%
FBS for 28-36h prior to doxycycline (500 ng/mL, Sigma-Aldrich CAT#D9891) induction or
DMSO control treatment for 6, 12, 24, or 36h in the presence of CDK4/6 inhibitor to maintain
cells in G0 and prevent additional transcriptional variation due to cell cycle progression. Each
experiment (repeated on three separate occasions) was performed in technical duplicate (with 6
replicate wells per technical duplicate), reserving one replicate for immunofluorescence staining
to confirm quality of cell cycle arrest (by DNA content and EdU incorporation) and the other for
harvesting RNA.
729
RNA-seq and analysis
Prior to harvesting, 96-well plates were washed with PBS, aspirated fully and transferred
immediately to -80C. RNA was later harvested from the plates by briefly thawing prior to using
extraction by RNeasy Mini kit (Qiagen Cat# 74106) and QIAshredder (Qiagen Cat#79654), with

pooling of 6 technical replicate wells/condition directly following the lysis step prior to following

the manufacturer protocol, including the optional DNase treatment (Qiagen Cat# 79254).

Samples were eluted in ultraPure ddH2O and assessed for quality using an Agilent Bioanalyzer

TapeStation. Illumina Stranded mRNA library preparation with polyA capture and sequencing

was performed on a NovaSeq6000 system (paired-end, 2 x 50) at the Weill Cornell Genome

Resources Core Facility over two rounds to get sufficient read depth.

Raw reads were checked for quality using FASTQC then aligned to the human genome

assembly (GRCh38) with HISAT2 [85] and assigned to genomic features with featureCounts[86].

Differential expression analysis was conducted with DESeq2 [87].

Image quantifications and analysis

Illumination bias correction, nuclear segmentation, image cropping and alignment across

iterative rounds of immunofluorescence, and analysis were performed with a custom MATLAB

pipeline (previously published[81] and deposited on Github: https://github.com/MeyerLab/image-

analysis-ratnayeke-2021) incorporating local density measurements[22]. For illumination bias

correction, images of background autofluorescence from immunofluorescence blocking solution

in cell-free wells were taken for each experiment and used to flatten raw images. Cell nuclei

were automatically segmented using Hoechst staining, a Gaussian filter, and a curvature-based

object splitting algorithm. A segmentation mask from the first round of imaging was used to align

subsequent imaging rounds and crop the images for the overlapping regions. Image jitter

between 4i rounds was typically less than 500 nm.

For nuclear-localized protein stains (EdU, cyclin D1, p27, phospho-Rb, c-Myc, Fra1,

SKP2, CDK6, p21), global background subtraction was performed automatically by expanding a

7.8 μm ring from the nuclear mask and subtracting the 25th percentile of pixel signal as

estimated background. For protein stains with cytoplasmic and nuclear localization (YAP,

phospho-S6, phospho-AKT, phospho-ERK, N-cadherin), median background signal was

estimated during analysis by manual averaging of background pixels of representative cell-free

regions and subtracting as a flat threshold. For quantification of immunofluorescence signals,

DNA content was calculated as the total nuclear Hoechst intensity (median intensity x nuclear

area); nuclear fluorescence signals were calculated as the median nuclear intensity;

cytoplasmic fluorescence signals were calculated as the median cytoplasmic intensity

(expanding the nuclear mask with a ring of inner radius 0.65 um and outer radius 3.23 um to

minimize the chance of sampling cytoplasm of neighboring cells). To compare experimental

replicates, fold-change in immunofluorescence staining was used to account for the variation in

staining and real fluorescence unit values.

Cell gating for G1-phase cells was performed by distinguishing G0/G1 and G2 cell

populations by 2N or 4N DNA context, then selecting cells below the EdU-incorporation

threshold (see sample EdU histograms, Figure S1B) with 2N DNA. Otherwise, for calculating

percent EdU-positive or percent Rb phosphorylated (all-cell gating), cells across cell cycle

phases were analyzed together. Rb phosphorylation was defined by histogram analysis of the

bifurcated phospho-Rb stain and calculating the fraction of cells above the threshold cutoff (see

sample p-Rb histograms, Figure S1B).

Local cell density analysis

Local cell density analysis was adapted from Fan et al., 2021. Briefly, local density was

measured by dividing imaging sites (25-30 tiled images/well) into a 3D histogram based on the

centroid coordinates of segmented nuclei (10 x 10 grid, MATLAB hist3). The xy location

coordinates of each cell centroid were rounded and the number of cells per block was

automatically counted, then scaled back to match the original image size (giving numerical

output rather than integer values)[22]. The local cell density value assigned for each cell was

based on the scaled number of cells in its assigned local density block.

Quantification corrections

While cells were counted prior to seeding, final cell plating density varied across experiments,

and absolute proliferation levels varied per independent experiment. We therefore assessed

both absolute (% EdU, % p-Rb) and fold-change proliferation metrics for assessing drug and

condition effect sizes.

During the iterative steps of multiplexed immunofluorescence, some cells washed off

during later rounds of imaging. As a proxy for null cells, cells that were segmented from the first

round of imaging were gated out of the final analysis if their Hoechst value was less than 100

RFU for a subsequent imaging round.

Statistical Analysis

Statistical analyses were performed using Student's one-sample or paired (MATLAB ttest) and

two-sample (MATLAB ttest2) t tests. Linear regressions for data binned by local cell density

were performed using fitlm (MATLAB). 95% confidence intervals and SEM are plotted as

indicated for representative experiments and summary data.

REFERENCES

1. Moya, I.M., and Halder, G. (2019). Hippo-YAP/TAZ signalling in organ regeneration and regenerative medicine. Nat. Rev. Mol. Cell Biol. *20*, 211–226. 10.1038/s41580-018-0086-y.

2. Driskill, J.H., and Pan, D. (2023). Control of stem cell renewal and fate by YAP and TAZ. Nat. Rev. Mol. Cell Biol. *24*, 895–911. 10.1038/s41580-023-00644-5.

3. Totaro, A., Panciera, T., and Piccolo, S. (2018). YAP/TAZ upstream signals and downstream responses. Nat. Cell Biol. *20*, 888–899. 10.1038/s41556-018-0142-z.

4. Fan, F., He, Z., Kong, L.-L., Chen, Q., Yuan, Q., Zhang, S., Ye, J., Liu, H., Sun, X., Geng, J., et al. (2016). Pharmacological targeting of kinases MST1 and MST2 augments tissue repair and regeneration. Sci. Transl. Med. *8*, 352ra108. 10.1126/scitranslmed.aaf2304.

5. Kastan, N., Gnedeva, K., Alisch, T., Petelski, A.A., Huggins, D.J., Chiaravalli, J., Aharanov, A., Shakked, A., Tzahor, E., Nagiel, A., et al. (2021). Small-molecule inhibition of Lats kinases may promote Yap-dependent proliferation in postmitotic mammalian tissues. Nat. Commun. *12*, 3100. 10.1038/s41467-021-23395-3.

6. Namoto, K., Baader, C., Orsini, V., Landshammer, A., Breuer, E., Dinh, K.T., Ungricht, R., Pikiolek, M., Laurent, S., Lu, B., et al. (2024). NIBR-LTSi is a selective LATS kinase inhibitor activating YAP signaling and expanding tissue stem cells in vitro and in vivo. Cell Stem Cell *31*, 554-569.e17. 10.1016/j.stem.2024.03.003.

7. Meng, Z., Moroishi, T., and Guan, K.-L. (2016). Mechanisms of Hippo pathway regulation. Genes Dev. *30*, 1–17. 10.1101/gad.274027.115.

8. Gumbiner, B.M., and Kim, N.-G. (2014). The Hippo-YAP signaling pathway and contact inhibition of growth. J. Cell Sci. *127*, 709–717. 10.1242/jcs.140103.

9. Zhang, N., Bai, H., David, K.K., Dong, J., Zheng, Y., Cai, J., Giovannini, M., Liu, P., Anders, R.A., and Pan, D. (2010). The Merlin/NF2 tumor suppressor functions through the YAP oncoprotein to regulate tissue homeostasis in mammals. Dev. Cell *19*, 27–38. 10.1016/j.devcel.2010.06.015.

10. Huang, J., Wu, S., Barrera, J., Matthews, K., and Pan, D. (2005). The Hippo signaling pathway coordinately regulates cell proliferation and apoptosis by inactivating Yorkie, the Drosophila Homolog of YAP. Cell *122*, 421–434. 10.1016/j.cell.2005.06.007.

11. Benhamouche, S., Curto, M., Saotome, I., Gladden, A.B., Liu, C.-H., Giovannini, M., and McClatchey, A.I. (2010). Nf2/Merlin controls progenitor homeostasis and tumorigenesis in the liver. Genes Dev. *24*, 1718–1730. 10.1101/gad.1938710.

835   12.   Yu, F.-X., Zhao, B., and Guan, K.-L. (2015). Hippo pathway in organ size control, tissue
836   homeostasis, and cancer. Cell *163*, 811–828. 10.1016/j.cell.2015.10.044.

837   13.   Zhao, B., Ye, X., Yu, J., Li, L., Li, W., Li, S., Yu, J., Lin, J.D., Wang, C.-Y., Chinnaiyan,
838   A.M., et al. (2008). TEAD mediates YAP-dependent gene induction and growth control.
839   Genes Dev. *22*, 1962–1971. 10.1101/gad.1664408.

840   14.   Mizuno, T., Murakami, H., Fujii, M., Ishiguro, F., Tanaka, I., Kondo, Y., Akatsuka, S.,
841   Toyokuni, S., Yokoi, K., Osada, H., et al. (2012). YAP induces malignant mesothelioma
842   cell proliferation by upregulating transcription of cell cycle-promoting genes. Oncogene
843   *31*, 5117–5122. 10.1038/onc.2012.5.

844   15.   Zanconato, F., Forcato, M., Battilana, G., Azzolin, L., Quaranta, E., Bodega, B., Rosato,
845   A., Bicciato, S., Cordenonsi, M., and Piccolo, S. (2015). Genome-wide association
846   between YAP/TAZ/TEAD and AP-1 at enhancers drives oncogenic growth. Nat. Cell
847   Biol. *17*, 1218–1227. 10.1038/ncb3216.

848   16.   Zhang, J., Ji, J.-Y., Yu, M., Overholtzer, M., Smolen, G.A., Wang, R., Brugge, J.S.,
849   Dyson, N.J., and Haber, D.A. (2009). YAP-dependent induction of amphiregulin
850   identifies a non-cell-autonomous component of the Hippo pathway. Nat. Cell Biol. *11*,
851   1444–1450. 10.1038/ncb1993.

852   17.   Zhang, H., Pasolli, H.A., and Fuchs, E. (2011). Yes-associated protein (YAP)
853   transcriptional coactivator functions in balancing growth and differentiation in skin. Proc
854   Natl Acad Sci USA *108*, 2270–2275. 10.1073/pnas.1019603108.

855   18.   Koo, J.H., Plouffe, S.W., Meng, Z., Lee, D.-H., Yang, D., Lim, D.-S., Wang, C.-Y., and
856   Guan, K.-L. (2020). Induction of AP-1 by YAP/TAZ contributes to cell proliferation and
857   organ growth. Genes Dev. *34*, 72–86. 10.1101/gad.331546.119.

858   19.   Liu, X., Li, H., Rajurkar, M., Li, Q., Cotton, J.L., Ou, J., Zhu, L.J., Goel, H.L., Mercurio,
859   A.M., Park, J.-S., et al. (2016). Tead and AP1 coordinate transcription and motility. Cell
860   Rep. *14*, 1169–1180. 10.1016/j.celrep.2015.12.104.

861   20.   Lavoie, H., Gagnon, J., and Therrien, M. (2020). ERK signalling: a master regulator of
862   cell behaviour, life and fate. Nat. Rev. Mol. Cell Biol. *21*, 607–632. 10.1038/s41580-020-
863   0255-7.

864   21.   Chiasson-MacKenzie, C., and McClatchey, A.I. (2018). Cell-Cell Contact and Receptor
865   Tyrosine Kinase Signaling. Cold Spring Harb. Perspect. Biol. *10*.
866   10.1101/cshperspect.a029215.

867   22.   Fan, Y., and Meyer, T. (2021). Molecular control of cell density-mediated exit to
868   quiescence. Cell Rep. *36*, 109436. 10.1016/j.celrep.2021.109436.

23. Yang, H.W., Chung, M., Kudo, T., and Meyer, T. (2017). Competing memories of
mitogen and p53 signalling control cell-cycle entry. Nature *549*, 404–408.
10.1038/nature23880.

24. Aoki, K., Kumagai, Y., Sakurai, A., Komatsu, N., Fujita, Y., Shionyu, C., and Matsuda,
M. (2013). Stochastic ERK activation induced by noise and cell-to-cell propagation
regulates cell density-dependent proliferation. Mol. Cell *52*, 529–540.
10.1016/j.molcel.2013.09.015.

25. Rubin, S.M., Sage, J., and Skotheim, J.M. (2020). Integrating old and new paradigms of
G1/S control. Mol. Cell *80*, 183–192. 10.1016/j.molcel.2020.08.020.

26. Zhao, B., Wei, X., Li, W., Udan, R.S., Yang, Q., Kim, J., Xie, J., Ikenoue, T., Yu, J., Li,
L., et al. (2007). Inactivation of YAP oncoprotein by the Hippo pathway is involved in
cell contact inhibition and tissue growth control. Genes Dev. *21*, 2747–2761.
10.1101/gad.1602907.

27. Cai, J., Zhang, N., Zheng, Y., de Wilde, R.F., Maitra, A., and Pan, D. (2010). The Hippo
signaling pathway restricts the oncogenic potential of an intestinal regeneration program.
Genes Dev. *24*, 2383–2388. 10.1101/gad.1978810.

28. Lee, M.-J., Byun, M.R., Furutani-Seiki, M., Hong, J.-H., and Jung, H.-S. (2014). YAP
and TAZ regulate skin wound healing. J. Invest. Dermatol. *134*, 518–525.
10.1038/jid.2013.339.

29. Hagenbeek, T.J., Zbieg, J.R., Hafner, M., Mroue, R., Lacap, J.A., Sodir, N.M., Noland,
C.L., Afghani, S., Kishore, A., Bhat, K.P., et al. (2023). An allosteric pan-TEAD inhibitor
blocks oncogenic YAP/TAZ signaling and overcomes KRAS G12C inhibitor resistance.
Nat. Cancer *4*, 812–828. 10.1038/s43018-023-00577-0.

30. Kastan, N.R., Oak, S., Liang, R., Baxt, L., Myers, R.W., Ginn, J., Liverton, N., Huggins,
D.J., Pichardo, J., Paul, M., et al. (2022). Development of an improved inhibitor of Lats
kinases to promote regeneration of mammalian organs. Proc Natl Acad Sci USA *119*,
e2206113119. 10.1073/pnas.2206113119.

31. Bienvenu, F., Jirawatnotai, S., Elias, J.E., Meyer, C.A., Mizeracka, K., Marson, A.,
Frampton, G.M., Cole, M.F., Odom, D.T., Odajima, J., et al. (2010). Transcriptional role
of cyclin D1 in development revealed by a genetic-proteomic screen. Nature *463*, 374–
378. 10.1038/nature08684.

32. Koff, A. (2006). How to decrease p27Kip1 levels during tumor development. Cancer Cell
9, 75–76. 10.1016/j.ccr.2006.01.020.

33. Gut, G., Herrmann, M.D., and Pelkmans, L. (2018). Multiplexed protein maps link
subcellular organization to cellular states. Science *361*. 10.1126/science.aar7042.

34.   Cheng, M., Sexl, V., Sherr, C.J., and Roussel, M.F. (1998). Assembly of cyclin D-
dependent kinase and titration of p27Kip1 regulated by mitogen-activated protein kinase
kinase (MEK1). Proc Natl Acad Sci USA *95*, 1091–1096. 10.1073/pnas.95.3.1091.

35.   Polyak, K., Lee, M.H., Erdjument-Bromage, H., Koff, A., Roberts, J.M., Tempst, P., and
Massagué, J. (1994). Cloning of p27Kip1, a cyclin-dependent kinase inhibitor and a
potential mediator of extracellular antimitogenic signals. Cell *78*, 59–66. 10.1016/0092-
8674(94)90572-x.

36.   Croci, O., De Fazio, S., Biagioni, F., Donato, E., Caganova, M., Curti, L., Doni, M.,
Sberna, S., Aldeghi, D., Biancotto, C., et al. (2017). Transcriptional integration of
mitogenic and mechanical signals by Myc and YAP. Genes Dev. *31*, 2017–2022.
10.1101/gad.301184.117.

37.   Monroe, T.O., Hill, M.C., Morikawa, Y., Leach, J.P., Heallen, T., Cao, S., Krijger,
P.H.L., de Laat, W., Wehrens, X.H.T., Rodney, G.G., et al. (2019). YAP partially
reprograms chromatin accessibility to directly induce adult cardiogenesis in vivo. Dev.
Cell *48*, 765-779.e7. 10.1016/j.devcel.2019.01.017.

38.   Cordenonsi, M., Zanconato, F., Azzolin, L., Forcato, M., Rosato, A., Frasson, C., Inui,
M., Montagner, M., Parenti, A.R., Poletti, A., et al. (2011). The Hippo transducer TAZ
confers cancer stem cell-related traits on breast cancer cells. Cell *147*, 759–772.
10.1016/j.cell.2011.09.048.

39.   Xie, Q., Chen, J., Feng, H., Peng, S., Adams, U., Bai, Y., Huang, L., Li, J., Huang, J.,
Meng, S., et al. (2013). YAP/TEAD-mediated transcription controls cellular senescence.
Cancer Res. *73*, 3615–3624. 10.1158/0008-5472.CAN-12-3793.

40.   Zhang, S., Chen, Q., Liu, Q., Li, Y., Sun, X., Hong, L., Ji, S., Liu, C., Geng, J., Zhang,
W., et al. (2017). Hippo Signaling Suppresses Cell Ploidy and Tumorigenesis through
Skp2. Cancer Cell *31*, 669-684.e7. 10.1016/j.ccell.2017.04.004.

41.   Kim, M., Kim, T., Johnson, R.L., and Lim, D.-S. (2015). Transcriptional co-repressor
function of the hippo pathway transducers YAP and TAZ. Cell Rep. *11*, 270–282.
10.1016/j.celrep.2015.03.015.

42.   Diehl, J.A., Cheng, M., Roussel, M.F., and Sherr, C.J. (1998). Glycogen synthase kinase-
3beta regulates cyclin D1 proteolysis and subcellular localization. Genes Dev. *12*, 3499–
3511. 10.1101/gad.12.22.3499.

43.   Nakayama, K., Nagahama, H., Minamishima, Y.A., Miyake, S., Ishida, N., Hatakeyama,
S., Kitagawa, M., Iemura, S., Natsume, T., and Nakayama, K.I. (2004). Skp2-mediated
degradation of p27 regulates progression into mitosis. Dev. Cell *6*, 661–672.
10.1016/s1534-5807(04)00131-5.

44.   Malek, N.P., Sundberg, H., McGrew, S., Nakayama, K., Kyriakides, T.R., and Roberts,
940   J.M. (2001). A mouse knock-in model exposes sequential proteolytic pathways that
941   regulate p27Kip1 in G1 and S phase. Nature *413*, 323–327. 10.1038/35095083.

942   45.   Shao, D.D., Xue, W., Krall, E.B., Bhutkar, A., Piccioni, F., Wang, X., Schinzel, A.C.,
943   Sood, S., Rosenbluh, J., Kim, J.W., et al. (2014). KRAS and YAP1 converge to regulate
944   EMT and tumor survival. Cell *158*, 171–184. 10.1016/j.cell.2014.06.004.

945   46.   Wang, X., Freire Valls, A., Schermann, G., Shen, Y., Moya, I.M., Castro, L., Urban, S.,
946   Solecki, G.M., Winkler, F., Riedemann, L., et al. (2017). YAP/TAZ Orchestrate VEGF
947   Signaling during Developmental Angiogenesis. Dev. Cell *42*, 462-478.e7.
948   10.1016/j.devcel.2017.08.002.

949   47.   Komuro, A., Nagai, M., Navin, N.E., and Sudol, M. (2003). WW domain-containing
950   protein YAP associates with ErbB-4 and acts as a co-transcriptional activator for the
951   carboxyl-terminal fragment of ErbB-4 that translocates to the nucleus. J. Biol. Chem. *278*,
952   33334–33341. 10.1074/jbc.M305597200.

953   48.   He, C., Lv, X., Hua, G., Lele, S.M., Remmenga, S., Dong, J., Davis, J.S., and Wang, C.
954   (2015). YAP forms autocrine loops with the ERBB pathway to regulate ovarian cancer
955   initiation and progression. Oncogene *34*, 6040–6054. 10.1038/onc.2015.52.

956   49.   Gregorieff, A., Liu, Y., Inanlou, M.R., Khomchuk, Y., and Wrana, J.L. (2015). Yap-
957   dependent reprogramming of Lgr5(+) stem cells drives intestinal regeneration and cancer.
958   Nature *526*, 715–718. 10.1038/nature15382.

959   50.   Yuan, Y., Park, J., Feng, A., Awasthi, P., Wang, Z., Chen, Q., and Iglesias-Bartolome, R.
960   (2020). YAP1/TAZ-TEAD transcriptional networks maintain skin homeostasis by
961   regulating cell proliferation and limiting KLF4 activity. Nat. Commun. *11*, 1472.
962   10.1038/s41467-020-15301-0.

963   51.   King, B., Araki, J., Palm, W., and Thompson, C.B. (2020). Yap/Taz promote the
964   scavenging of extracellular nutrients through macropinocytosis. Genes Dev. *34*, 1345–
965   1358. 10.1101/gad.340661.120.

966   52.   Gillies, T.E., Pargett, M., Minguet, M., Davies, A.E., and Albeck, J.G. (2017). Linear
967   Integration of ERK Activity Predominates over Persistence Detection in Fra-1
968   Regulation. Cell Syst. *5*, 549-563.e5. 10.1016/j.cels.2017.10.019.

969   53.   Edwards, A.C., Stalnecker, C.A., Jean Morales, A., Taylor, K.E., Klomp, J.E., Klomp,
970   J.A., Waters, A.M., Sudhakar, N., Hallin, J., Tang, T.T., et al. (2023). TEAD Inhibition
971   Overcomes YAP1/TAZ-Driven Primary and Acquired Resistance to KRASG12C
972   Inhibitors. Cancer Res. *83*, 4112–4129. 10.1158/0008-5472.CAN-23-2994.

973   54.   Vaidyanathan, S., Salmi, T.M., Sathiqu, R.M., McConville, M.J., Cox, A.G., and Brown,
974   K.K. (2022). YAP regulates an SGK1/mTORC1/SREBP-dependent lipogenic program to

support proliferation and tissue growth. Dev. Cell *57*, 719-731.e8.
10.1016/j.devcel.2022.02.004.

55.   Tumaneng, K., Schlegelmilch, K., Russell, R.C., Yimlamai, D., Basnet, H., Mahadevan,
N., Fitamant, J., Bardeesy, N., Camargo, F.D., and Guan, K.-L. (2012). YAP mediates
crosstalk between the Hippo and PI(3)K–TOR pathways by suppressing PTEN via miR-
29. Nat. Cell Biol. *14*, 1322–1329. 10.1038/ncb2615.

56.   Zoncu, R., Efeyan, A., and Sabatini, D.M. (2011). mTOR: from growth signal integration
to cancer, diabetes and ageing. Nat. Rev. Mol. Cell Biol. *12*, 21–35. 10.1038/nrm3025.

57.   Averous, J., Fonseca, B.D., and Proud, C.G. (2008). Regulation of cyclin D1 expression
by mTORC1 signaling requires eukaryotic initiation factor 4E-binding protein 1.
Oncogene *27*, 1106–1113. 10.1038/sj.onc.1210715.

58.   Hamaratoglu, F., Willecke, M., Kango-Singh, M., Nolo, R., Hyun, E., Tao, C., Jafar-
Nejad, H., and Halder, G. (2006). The tumour-suppressor genes NF2/Merlin and
Expanded act through Hippo signalling to regulate cell proliferation and apoptosis. Nat.
Cell Biol. *8*, 27–36. 10.1038/ncb1339.

59.   Striedinger, K., VandenBerg, S.R., Baia, G.S., McDermott, M.W., Gutmann, D.H., and
Lal, A. (2008). The neurofibromatosis 2 tumor suppressor gene product, merlin, regulates
human meningioma cell growth by signaling through YAP. Neoplasia *10*, 1204–1212.
10.1593/neo.08642.

60.   Yin, F., Yu, J., Zheng, Y., Chen, Q., Zhang, N., and Pan, D. (2013). Spatial organization
of Hippo signaling at the plasma membrane mediated by the tumor suppressor
Merlin/NF2. Cell *154*, 1342–1355. 10.1016/j.cell.2013.08.025.

61.   Wada, K.-I., Itoga, K., Okano, T., Yonemura, S., and Sasaki, H. (2011). Hippo pathway
regulation by cell morphology and stress fibers. Development *138*, 3907–3914.
10.1242/dev.070987.

62.   Aragona, M., Panciera, T., Manfrin, A., Giulitti, S., Michielin, F., Elvassore, N., Dupont,
S., and Piccolo, S. (2013). A mechanical checkpoint controls multicellular growth
through YAP/TAZ regulation by actin-processing factors. Cell *154*, 1047–1059.
10.1016/j.cell.2013.07.042.

63.   Ong, Y.T., Andrade, J., Armbruster, M., Shi, C., Castro, M., Costa, A.S.H., Sugino, T.,
Eelen, G., Zimmermann, B., Wilhelm, K., et al. (2022). A YAP/TAZ-TEAD signalling
module links endothelial nutrient acquisition to angiogenic growth. Nat. Metab. *4*, 672–
682. 10.1038/s42255-022-00584-y.

64.   Su, T., Bondar, T., Zhou, X., Zhang, C., He, H., and Medzhitov, R. (2015). Two-signal
requirement for growth-promoting function of Yap in hepatocytes. eLife *4*.
10.7554/eLife.02948.

65.   Chen, Q., Zhang, N., Xie, R., Wang, W., Cai, J., Choi, K.-S., David, K.K., Huang, B.,
1012   Yabuta, N., Nojima, H., et al. (2015). Homeostatic control of Hippo signaling activity
1013   revealed by an endogenous activating mutation in YAP. Genes Dev. *29*, 1285–1297.
1014   10.1101/gad.264234.115.

1015   66.   Meyer, K., Lammers, N.C., Bugaj, L.J., Garcia, H.G., and Weiner, O.D. (2023).
1016   Optogenetic control of YAP reveals a dynamic communication code for stem cell fate and
1017   proliferation. Nat. Commun. *14*, 6929. 10.1038/s41467-023-42643-2.

1018   67.   Franklin, J.M., Ghosh, R.P., Shi, Q., Reddick, M.P., and Liphardt, J.T. (2020). Concerted
1019   localization-resets precede YAP-dependent transcription. Nat. Commun. *11*, 4581.
1020   10.1038/s41467-020-18368-x.

1021   68.   Manning, S.A., Dent, L.G., Kondo, S., Zhao, Z.W., Plachta, N., and Harvey, K.F. (2018).
1022   Dynamic fluctuations in subcellular localization of the hippo pathway effector yorkie
1023   in vivo. Curr. Biol. *28*, 1651-1660.e4. 10.1016/j.cub.2018.04.018.

1024   69.   Curto, M., Cole, B.K., Lallemand, D., Liu, C.-H., and McClatchey, A.I. (2007). Contact-
1025   dependent inhibition of EGFR signaling by Nf2/Merlin. J. Cell Biol. *177*, 893–903.
1026   10.1083/jcb.200703010.

1027   70.   Moleirinho, S., Hoxha, S., Mandati, V., Curtale, G., Troutman, S., Ehmer, U., and Kissil,
1028   J.L. (2017). Regulation of localization and function of the transcriptional co-activator
1029   YAP by angiomotin. eLife *6*. 10.7554/eLife.23966.

1030   71.   Yi, C., Troutman, S., Fera, D., Stemmer-Rachamimov, A., Avila, J.L., Christian, N.,
1031   Persson, N.L., Shimono, A., Speicher, D.W., Marmorstein, R., et al. (2011). A tight
1032   junction-associated Merlin-angiomotin complex mediates Merlin's regulation of
1033   mitogenic signaling and tumor suppressive functions. Cancer Cell *19*, 527–540.
1034   10.1016/j.ccr.2011.02.017.

1035   72.   Kapoor, A., Yao, W., Ying, H., Hua, S., Liewen, A., Wang, Q., Zhong, Y., Wu, C.-J.,
1036   Sadanandam, A., Hu, B., et al. (2014). Yap1 activation enables bypass of oncogenic Kras
1037   addiction in pancreatic cancer. Cell *158*, 185–197. 10.1016/j.cell.2014.06.003.

1038   73.   Jang, W., Kim, T., Koo, J.S., Kim, S.-K., and Lim, D.-S. (2017). Mechanical cue-induced
1039   YAP instructs Skp2-dependent cell cycle exit and oncogenic signaling. EMBO J. *36*,
1040   2510–2528. 10.15252/embj.201696089.

1041   74.   Kossatz, U., Dietrich, N., Zender, L., Buer, J., Manns, M.P., and Malek, N.P. (2004).
1042   Skp2-dependent degradation of p27kip1 is essential for cell cycle progression. Genes
1043   Dev. *18*, 2602–2607. 10.1101/gad.321004.

1044   75.   Adachi, Y., Kimura, R., Hirade, K., Yanase, S., Nishioka, Y., Kasuga, N., Yamaguchi,
1045   R., and Ebi, H. (2023). Scribble mis-localization induces adaptive resistance to KRAS

G12C inhibitors through feedback activation of MAPK signaling mediated by YAP-
induced MRAS. Nat. Cancer *4*, 829–843. 10.1038/s43018-023-00575-2.

76. Lin, L., Sabnis, A.J., Chan, E., Olivas, V., Cade, L., Pazarentzos, E., Asthana, S., Neel,
D., Yan, J.J., Lu, X., et al. (2015). The Hippo effector YAP promotes resistance to RAF-
and MEK-targeted cancer therapies. Nat. Genet. *47*, 250–256. 10.1038/ng.3218.

77. Dai, X., Liu, H., Shen, S., Guo, X., Yan, H., Ji, X., Li, L., Huang, J., Feng, X.-H., and
Zhao, B. (2015). YAP activates the Hippo pathway in a negative feedback loop. Cell Res.
25, 1175–1178. 10.1038/cr.2015.101.

78. Moroishi, T., Park, H.W., Qin, B., Chen, Q., Meng, Z., Plouffe, S.W., Taniguchi, K., Yu,
F.-X., Karin, M., Pan, D., et al. (2015). A YAP/TAZ-induced feedback mechanism
regulates Hippo pathway homeostasis. Genes Dev. *29*, 1271–1284.
10.1101/gad.262816.115.

79. Nardone, G., Oliver-De La Cruz, J., Vrbsky, J., Martini, C., Pribyl, J., Skládal, P., Pešl,
M., Caluori, G., Pagliari, S., Martino, F., et al. (2017). YAP regulates cell mechanics by
controlling focal adhesion assembly. Nat. Commun. *8*, 15321. 10.1038/ncomms15321.

80. Pastushenko, I., Mauri, F., Song, Y., de Cock, F., Meeusen, B., Swedlund, B., Impens, F.,
Van Haver, D., Opitz, M., Thery, M., et al. (2021). Fat1 deletion promotes hybrid EMT
state, tumour stemness and metastasis. Nature *589*, 448–455. 10.1038/s41586-020-03046-
1.

81. Ratnayeke, N., Baris, Y., Chung, M., Yeeles, J.T.P., and Meyer, T. (2023). CDT1 inhibits
CMG helicase in early S phase to separate origin licensing from DNA synthesis. Mol.
Cell *83*, 26-42.e13. 10.1016/j.molcel.2022.12.004.

82. Ran, F.A., Hsu, P.D., Wright, J., Agarwala, V., Scott, D.A., and Zhang, F. (2013).
Genome engineering using the CRISPR-Cas9 system. Nat. Protoc. *8*, 2281–2308.
10.1038/nprot.2013.143.

83. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat. Methods *9*, 676–682. 10.1038/nmeth.2019.

84. Stallaert, W., Kedziora, K.M., Taylor, C.D., Zikry, T.M., Ranek, J.S., Sobon, H.K.,
Taylor, S.R., Young, C.L., Cook, J.G., and Purvis, J.E. (2022). The structure of the
human cell cycle. Cell Syst. *13*, 230-240.e3. 10.1016/j.cels.2021.10.007.

85. Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based
genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat. Biotechnol.
37, 907–915. 10.1038/s41587-019-0201-4.

86.   Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose
1081   program for assigning sequence reads to genomic features. Bioinformatics *30*, 923–930.
1082   10.1093/bioinformatics/btt656.

1083   87.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
1084   dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550. 10.1186/s13059-014-
1085   0550-8.

Figure 1. Contact inhibition of YAP opposes mitogen stimulation in a balance mechanism that controls proliferation

Figure 1. Contact inhibition of YAP opposes mitogen stimulation in a balance mechanism that controls proliferation

A) Local cell density metric by subdividing images into 10 x 10 grids and counting the cells (by centroid of segmented nucleus) for each block (see also STAR Methods, scale bar = 100 μm).
B) <u>Left</u>: Diagram of cell cycle phases: EdU incorporation marks S-phase, Rb hyperphosphorylation (p-Rb, S807/811) cycling cells. <u>Right</u>: sample images of wildtype (WT) RPE1 cells at increasing local cell density: Hoechst for DNA (white), EdU for S-phase cells (green), p-Rb for cycling cells (blue), and YAP (grey). Magenta outline is the automated nuclear mask (scale bar = 10 μm).
C) Local cell density (relative units, R.U.) was measured by binning cell numbers per local block area. The percent of Rb-phosphorylated cells was calculated per local density bin for WT and CA-YAP cells maintained in increasing serum levels (0.1%, 1%, or 10%) for 36h, n > 20,000 cells/condition. See also **Figure S1**.
D) Mean percent p-Rb for cells treated as in (C). n = 3 independent experiments. Student's t-test: $p=6.86 \times 10^{-6}$ (0.1% serum), $p=2.50 \times 10^{-5}$ (1% serum), $p = 5.90 \times 10^{-4}$ (10% serum).
E) Median nuclear/cytoplasmic (n/c) YAP ratio (left) and nuclear YAP levels (right, real fluorescent units, R.F.U.) per local density bin for WT cells treated as in (C), n > 20,000 cells/condition. Shaded error bars are 95% confidence intervals.
F) Percent p-Rb for WT cells binned by their YAP ratio (left) or nuclear YAP levels (right).
G) Representative images of YAP staining for WT cells treated with TEAD inhibitor (TEADi, 0.5 μM GNE-7883), LATS inhibitor (LATSi, 0.5 μM TDI-011536) or in combination in 10% serum for 24h.
H) Fold-change in percent p-Rb for WT cells treated with TEADi, LATSi, or in combination, normalized to DMSO in 0.1% (left), 1% (middle), or 10% serum (right). N = 3 independent experiments.
All binned plots have n>100 cells/bin and >20,000 cells/condition.

Figure 2. YAP-regulated cell cycle entry and exit is dictated by a cyclin D1/p27 threshold

Figure 2. YAP-regulated cell cycle entry and exit is dictated by a cyclin D1/p27 threshold

A-B) Median levels (real fluorescent units (RFU)) of cyclin D1 (left) and p27 (right) per local cell density bin for WT (A) and CA-YAP (B) cells at 0.1%, 1%, and 10% serum. Binned data only include G1-gated cells, n > 100 cells/bin. Shaded error bars are 95% confidence intervals, n > 15,000 cells per condition. Data are representative of 2 independent experiments.

C) Representative images of multiplexed immunofluorescence staining comparing cyclin D1 (green) and p27 levels (magenta) with aligned p-Rb stain (cyan with gray nuclear mask overlay) in WT or CA-YAP cells. Scale bar = 50 µm.

D-E) Single-cell scatterplots of co-stained cyclin D1 and p27 levels (log2-transformed), colored by p-Rb (purple, p-Rb-negative; orange, p-Rb-positive) in WT (D) and CA-YAP (E). Left, 0.1% serum; right, 10% serum. Mean ± SD of 2 independent experiments.

F) Representative single-cell scatterplot of serum-starved CA-YAP cells co-stained for cyclin D1 and p27 protein levels, colored by p21 protein levels (top: parula color scale) or p-Rb status (bottom: purple, p-Rb-negative; orange, p-Rb-positive).

G) Top: experimental conditions for siRNA transfection followed by 48h serum starvation. Bottom: mean cyclin D1/p27 ratio (left) and p-Rb (right) for WT and CA-YAP cells transfected with siRNA against cyclin D1 and p27, normalized to non-targeting control (siCtrl). N = 3 independent experiments.

H) Single-cell scatterplots of co-stained cyclin D1 and p27 levels colored by p-Rb for CA-YAP cells treated as in (G).

I) Top: experimental conditions for sgRNA-mediated overexpression (CRISPRa) of p27 in dCas9-expressing WT cells treated with DMSO or LATSi (0.5 µM TDI-011536). Bottom: Single-cell scatterplots of co-stained cyclin D1 and p27 levels colored by p-Rb for p27-overexpressing cells treated with DMSO (left) or LATSi (right).

J) Schematic of mitogen signaling and contact inhibition of YAP-TEAD activity converging on the cyclin D1/p27 ratio to control whether cells enter or exit the cell cycle.

Dashed lines approximate separation of p-Rb-positive and p-Rb-negative cell populations. Single-cell data are representative of 2 independent experiments, with n = 8,000 random cells/scatter plot and gated for G1-phase (defined as 2N DNA/EdU-negative status). See also **Figure S2**.

Figure 3. Delayed YAP-mediated control of the nuclear cyclin D1/p27 ratio and cell cycle arrest

Figure 3. Delayed YAP-mediated control of the cyclin D1/p27 ratio and cell cycle arrest

A) Schematic for CA-YAP expression in RPE1 cells. RNA-sequencing performed in confluent cells pre-conditioned at 0.1% and palbociclib-arrested (CDK4/6i, 1 µM), prior to doxycycline-induction over 36h (0.5 µg/mL DOX). See also **Figure S3** and STAR METHODS.

B) Up-regulated genes (q<0.05, Log2 fold-change >1) for 12, 24, and 36h YAP induction showed the Cordenonsi YAP signature (using ToppGene)[38].

C-D) Log2 fold-change (Log2FC) induction (C) of *CYR61* (canonical YAP target gene, grey) compared with *CCND1* (cyclin D1, red) and (D) regulation of DDIT4 (suppressed YAP target gene, grey), *CDKN1A* (p21, dark blue), and *CDKN1B* (p27, blue) at times since overexpression. Adjusted p-values (NS at all other time points): *CCND1*, 3.17×10^{-4} (12h), 4.45×10^{-5} (24h), 3.96×10^{-12} (36h); *CDKN1A*, 8.65×10^{-7} (12h), 9.5×10^{-4} (24h); *CDKN1B*, 0.0012 (24h), 0.0312 (36h).

E) <u>Top</u>: Experimental conditions for LATSi treatment (0.5 µM TDI-011536) of serum-starved WT cells for 0-24h. <u>Bottom</u>: Histogram distribution of cyclin D1, p27, and p21 protein levels over LATSi time course.

F) Mean cyclin D1/p27 ratio (left) and percent p-Rb (right) for cells treated as in (E). n = 3 independent experiments and 3 well-replicates/experiment, as indicated by matched colors. Student's t-test: cyclin D1/p27, p = 0.0062 (6h), 0.0022 (12h), 2.63×10^{-4} (24h); p-Rb, p = 0.0217 (6h), 4.12×10^{-4} (12h), 0.0087 (24h).

G) Single-cell scatterplots of cyclin D1 and p27 protein levels colored by p-Rb status (purple, p-Rb-negative; orange, p-Rb-positive).

H) <u>Top</u>: experimental conditions for TEADi treatment (1 µM GNE-7883) of serum-starved CA-YAP cells. <u>Bottom</u>: Histogram distribution of log2-transformed cyclin D1, p27, and p21 protein levels over TEADi time course.

I) Mean cyclin D1/p27 ratio (left) and percent p-Rb (right) for cells treated as in (H). n = 3 biological replicates, where well-replicates are indicated by matched colors. Student's t-test: cyclin D1/p27 ratio, p = 0.027 (24h), p = 0.0015 (36h); % p-Rb, p = 0.0188 (3h), p = 0.0011 (6h), 0.0119 (12h), p = 0.0481 (24h), p = 1.98×10^{-5} (36h).

J) Single-cell scatterplots of log2-transformed cyclin D1 and p27 protein levels colored by p-Rb status for cells treated as in (H).

All histogram and single-cell data (n = 8,000 cells/scatterplot) are representative of 3 independent experiments.

Figure 4. YAP activation induces proliferation by increasing EGFR signaling

Figure 4. YAP activation induces proliferation by increasing EGFR signaling

A) Heatmap representation of differential expression (row z-score) of select up-regulated ligand and receptor genes, grouped by 3 independent replicates and time points.

B) Log2 fold-change (Log2FC) of select receptor signaling-related genes, *AXL* (direct YAP target grey), *EGFR* (blue), *HBEGF* (green), and *HRAS* (red). Adjusted p-values: *AXL,* 6.20×10^{-4} (6h), 9.11×10^{-12} (12h), 2.33×10^{-11} (24h), 2.19×10^{-21} (36h); *EGFR,* 4.97×10^{-16} (6h), 1.07×10^{-14} (12h), 1.07×10^{-8} (24h), 1.23×10^{-5} (36h); *HBEGF*, 3.90×10^{-4} (12h), 8.57×10^{-12} (24h), 5.24×10^{-18} (36h); *HRAS*, 4.10×10^{-4} (24h), 6.05×10^{-5} (36h).

C) Percent p-Rb binned by local cell density for WT (left) and CA-YAP (right) cells serum-starved and treated with gefitinib (EGFRi, 0-20 µM) for 24h. See also **Figure S4.**

D) Western blot analysis of EGFR and AXL from whole-cell lysates of serum-starved CA-YAP cells treated with TEADi for 0-36h, with GAPDH and Vinculin loading controls. Values indicate fold-change difference relative to 0h control (after normalization to loading control). Data are representative of 2 independent experiments.

E) Percent p-Rb binned by local density for serum-starved WT and CA-YAP cells after 14h (25 ng/mL). Data are representative of 2 independent experiments, n > 25,000 cells/condition.

F) Western blot analysis of AKT and ERK phosphorylation levels in whole-cell lysates of serum-starved WT (left), CA-YAP (middle), and TEADi pre-treated CA-YAP cells (right) after 0-10h EGF treatment (25 ng/mL). Values normalized to loading control and fold-change relative to 0h control.

G) Top: experimental conditions for serum-starvation of WT cells, pre-treatment with LATSi, and sustained or washout of LATSi during 14h EGF treatment (25 ng/mL). Bottom: Mean YAP ratio (n/c) for treatment conditions. N = 3 independent experiments, matched colors indicate conditions from the same experimental replicate. Student's t-test, YAP ratio: LATSi sustained, p= 0.0278 (CTRL), 0.027 (EGF); LATSi washout, not significant (NS).

H) Percent p-Rb binned by local cell density for cells with LATSi sustained (left) or washout (right) with or without EGF treatment (n > 20,000 cells/condition).

I-J) Histogram distribution of phosphorylated-S6 (p-S6 240/244, I) and Fra1 protein distributions for DMSO-treated (top) or LATSi-pretreated cells (bottom, LATSi sustained and washout). Data are representative of 3 independent experiments.

All single-cell data represent values from G1-gated cells.

Figure 5. YAP activation signals through ERK and mTOR to increase the cyclin D1/p27 ratio

1212 **Figure 5. YAP activation signals through ERK and mTOR to increase the cyclin D1/p27**
1213 **ratio**
1214
A) Schematic representation of proposed indirect regulation of cyclin D1/p27 ratio by
YAP/TEAD via EGFR-activated signaling pathways.
B-C) <u>Top</u>: experimental conditions for serum-starvation of WT and CA-YAP cells followed by
24h treatment with TEAD inhibitor for 24h in starvation media. <u>Bottom</u>: Mean fold-change (FC)
in protein levels of p-AKT(473, left) and p-S6(240/244, right) (B) and Fra1 (C) for serum-starved
WT and CA-YAP cells treated with TEADi for 24h (normalized to WT/DMSO condition). N = 2
(p-AKT) or 3 (p-S6, Fra1) independent experiments.
D) Representative multiplexed images for Fra1 and p-S6(240/244) staining in CA-YAP cells
treated with TEADi as in (B). Scale bar = 50 μm.
E-F) Mean percentage p-Rb for WT and CA-YAP cells treated with PI3K (LY294002, 10 μM),
AKT (MK2206, 200 nM), or mTORC1/2 (Torin2, 100 nM) inhibitors (E) or MEK (PD0325901,
100 nM) inhibitor (F) as in (B). N = 3 independent experiments. Student's t-test: WT/mTORi, p =
0.0372; CA-YAP/mTORi, p = 0.0103 (E). CA-YAP/MEKi, p = 0.0028 (F). See also **Figure S5**.
G) Mean FC in cyclin D1 protein levels for WT and CA-YAP cells treated with combined mTORi
and MEKi relative to WT/DMSO condition as in (B). N = 2 independent experiments.
H) <u>Top</u>: experiment conditions for mTORi or MEKi time course treatment of serum-starved CA-
YAP cells. <u>Bottom</u>: fold-change in p-ERK (MEKi-treated, dark grey) and p-S6(240/244, mTORi-
treated, light grey) following drug treatment; mean ± SEM (n = 2 independent experiments, 2-3
replicate wells/experiment).
I) Fold-change in protein levels of cyclin D1, p27, and p-Rb for MEKi and mTORi for CA-YAP
cells treated as in (H). Mean ± SEM.
1236

Figure 6. Partial activation of YAP by Merlin knockout can be reversed by increased contact inhibition

Figure 6. Partial activation of YAP by Merlin knockout can be reversed by increased contact inhibition

A) TEAD-dependent hyperproliferation of Merlin KO cells. Percent p-Rb binned by local density for Merlin knockout cells (Merlin KO, yellow) in 0.1% (left) or 10% serum (right) treated with DMSO or TEADi (0.5 M) compared with WT cells (grey).

B) Median YAP ratio (n/c) binned by local cell density for Merlin KO cells in 0.1% or 10% serum, compared with WT cells (same experiment as in **Figure 1E**). Data are representative of 3 independent experiments. Shaded error are 95% confidence intervals.

C) Percent p-Rb binned by local density for serum-starved Merlin KO cells treated with gefitinib (EGFRi) as in Figure 4C. Data are representative of 2 independent experiments.

D) Percent p-Rb binned by local cell density for serum-starved Merlin KO cells with control (CTRL) or EGF (25 ng/mL) treatment for 12h. Data are representative of 2 independent experiments.

E) Median cyclin D1 (left) or p27 (right) protein levels for Merlin KO cells in titrated serum. Shaded error are 95% confidence intervals. Data are representative of 2 independent experiments.

F) Median cyclin D1/p27 ratio binned by local cell density for WT (grey), Merlin KO (yellow), and CA-YAP cells (red) in 0.1% (left) or 10% serum (right). Shaded error are 95% confidence intervals.

G) Model for Hippo-pathway inhibition of YAP-TEAD activity, which controls signaling through and downstream of EGFR to the cyclin D1/p27 ratio.

H) Model proposing transitions between low and high YAP-TEAD activity as contact inhibition is temporarily reduced then recovered after a delay during epithelial sheet regeneration.

For all representative binned plots, data are pooled from at least 2 wells, n > 100 cells/bin, >10,000 cells/plot. All quantifications are for G1-gated cells. See also **Figure S6**.

Figure S1

Figure S1, related to Figure 1

A) Representative local density estimation in an imaging site, where circles represent the location of the centroids of cell nuclei. Cell location coordinates were divided into a 10 x 10, equally-spaced grid (local density blocks). The z-scale color indicates the number of cell centroids within a block. Cells were assigned local densities based on the number of cells in their block, scaled by the image size to give a relative density unit. See also *STAR METHODS*.
B) Histogram plots of binary distribution of EdU (<u>top</u>) and Rb phosphorylation (phospho-Rb 807/811, <u>bottom</u>) in WT RPE1 cells with 10%, 1%, or 0.1% serum. Dashed line indicates cutoff values used to designed whether cells are EdU-positive or have phosphorylated Rb. Data are representative of 3 independent experiments (same experiment as in Figure 1C).
C) Percent EdU-positive cells per local density bin for WT cells in titrated serum levels. Data are representative of 3 independent experiments.
D) Percent p-Rb binned by alternate local density metric (area of neighboring cell nuclear pixels within an 80-pixel (~51.6 µm) radius) for WT cells in titrated serum, showing similar density dependence as the primary local density metric.
E) Mean nuclear YAP protein levels in serum-titrated WT and CA-YAP cells, normalized (norm.) to WT/0.1% serum condition for each experiment, n = 3 independent experiments.
F) Mean percent EdU-positive in WT and CA-YAP across serum conditions, n = 3 independent experiments. Student's t-test: $p = 1.01 \times 10^{-5}$ (0.1% serum), $p = 8.13 \times 10^{-5}$ (1% serum), $p = 7.13 \times 10^{-4}$ (10% serum).
G-H) Percent p-Rb (left) and median YAP ratio (nuclear/cytoplasmic, n/c, right) binned by local density for WT and CA-YAP cells in 0.1% and 10% serum. Cells were treated with DMSO and TEADi (1 µM GNE-7883, G, dashed lines) or LATSi (0.5 µM TDI-011536, H, orange lines) for 24h. Shaded error are 95% confidence intervals, n > 15,000 cells/condition. Arrows highlight drug-mediated changes for matched serum conditions.
I) Percent p-Rb per local density bin for WT (grey) and CA-YAP cells treated with DMSO (red) or TEADi (0.5 µM, red, dashed line) in titrated serum.
All binned and histogram data are from one representative replicate of 3 independent experiments, with >100 cells/bin for binned data (n >15,000 cells/condition).

Figure S2

A sample image-based cytometry

B WT cyclin D1 — 0.1% serum / 10% serum

p27

log2(protein (RFU) (G1)

C WT — 10% serum / 1% serum / 0.1% serum

cyclin D1/p27 ratio

D 0.1% serum cyclin D1 — WT (as in**B**) / CA-YAP

p27

p-Rb+

log2(protein (RFU) (G1)

E CA-YAP — 10% serum / 1% serum / 0.1% serum

cyclin D1/p27 ratio

F WT, 0% serum — log2(p21)

log2(Cyclin D1) (G1)

pRb- / pRb+

G CA-YAP / WT — Local Density

0% FBS — WT / CA-YAP

H WT, si-Ctrl — p-Rb- / p-Rb+

WT, si-Cyclin D1

WT, si-p27

log2(Cyclin D1) (G1)

1295 **Figure S2, related to Figure 2**
1296

A) Density scatterplot of DNA content (Hoechst staining, real fluorescence units (RFU)) and
EdU-incorporation for representative G0/G1 gating of cells. G1 population cells were defined as
2N DNA/EdU-negative, S-cells as EdU-positive, and G2 as 4N DNA/EdU-negative. All-cell
gating reflects cells within the outer red boundary (n = 8,000 random cells).
B) Histogram distribution of cyclin D1 protein (top) and p27 protein (bottom) stain in WT cells in
0.1% or 10% FBS. Data are representative of n = 2 independent experiments
C) Percent p-Rb binned by cyclin D1/p27 ratio for serum titration in WT cells (n > 15,000
cells/condition).
D) Histogram distribution of cyclin D1 (top), p27 (middle), and p-Rb (bottom) stain comparing
CA-YAP cells with WT in 0.1% FBS (same WT data as S1B), n > 20,000 cells/condition. Dashed
line indicates p-Rb cutoff.
E) Percent p-Rb binned by cyclin D1/p27 ratio for serum titration in CA-YAP cells (n > 15,000
cells/condition).
F) Density scatterplot of cyclin D1 stain against p27 stain, colored by binary p-Rb status (left) or
p21 stain (right, parula scale) in starved WT cell.
G) Mean p21 (left) and p27 (right) protein levels (RFU) binned by local density for starved WT
(grey), and CA-YAP (red) cells (n > 20,000 cells/condition). Shaded error are 95% confidence
intervals.
H) Density scatterplots of cyclin D1 and p27 protein levels (RFU) colored by p-Rb status for
serum-starved WT cells transfected with siRNA against control (si-Ctrl, left), cyclin D1 (si-Cyclin
D1, middle), or p27 (si-p27, right). Dashed, diagonal line visually approximates where p-Rb
bifurcates in CA-YAP conditions (matching Figure 2H).
All data representative of 2 independent experiments and gated for G1-phase. All single-cell
scatterplots show n = 8,000 cells.
1321

Figure S3

A 0.1% FBS CA-YAP — % EdU vs DOX (h): -, 6, 12, 24, 36

B Log2FC vs DOX (h), 0.1% FBS — ANKRD1, SKP2, CDK6

C Log2FC vs DOX (h), 0.1% FBS — MYBL1, ETS1, MYC

D TEAD/AP-1-associated (positive feedback), row z-score — JUN, FOSL2, MYC, TEAD1, FOSL1, TEAD4; DOX (h), 0.1% serum: 0, 6, 12, 24, 36

E WT — YAP ratio (n/c), CDK6 (G1), SKP2 (G1); LATSi (h): -, 3, 6, 12, 24; TEADi (h): 24

F WT — Cyclin D1 (G1), p27 (G1), p21 (G1); LATSi (h): -, 3, 6, 12, 24; TEADi (h): 24

G WT — % EdU-positive; LATSi (h): -, 3, 6, 12, 24; TEADi (h): 24

H YAP ratio (n/c) vs TEADi (h), 0% serum — WT/LATSi, CA-YAP

I Cyclin D1/p27 ratio (G1), % p-Rb (G1), % EdU-positive vs TEADi (h), 0% serum — WT/LATSi, CA-YAP

Figure S3, related to Figure 3

A) Mean percent EdU for matched replicates of Figure 3A reserved for immunofluorescence staining for CA-YAP cells in 0.1% FBS and arrested with palbociclib (1μM) during treatment with DMSO (-) or doxycycline (DOX) for 0-36h. N = 3 independent experiments.

B-C) Log2 fold-change (Log2FC) induction of ANKRD1 (canonical YAP target gene, grey) compared with *CDK6* (green) and *SKP2* (purple) (B) and transcription factor targets *MYC* (red) compared with *MYBL1* (orange) and *ETS1* (yellow) at times since overexpression (C). Adjusted p-values: *CDK6*, 0.0451 (6h), 4.22×10^{-22} (12h), 3.08×10^{-52} (24h), 3.26×10^{-58} (36h); *SKP2*, 3.78×10^{-13} (12h), 3.49×10^{-65} (24h), 1.86×10^{-70} (36h); *MYC*, 2.18×10^{-6} (12h), 2.39×10^{-8} (24h), 1.50×10^{-13} (36h); *MYBL1*, 7.77×10^{-13} (12h), 2.76×10^{-10} (24h), 3.48×10^{-7} (36h); *ETS1*, 1.85×10^{-13} (6h), 4.55×10^{-16} (12h), 6.24×10^{-37} (24h), 1.23×10^{-57} (36h).

D) Heatmap of TEAD/AP-1-associated positive feedback genes in DOX-treated, CA-YAP cells. Color scale indicates row z-score.

E-G) Mean YAP ratio (E, left) and fold-change in protein levels of CDK6 (E, middle), SKP2 (E, right), cyclin D1 (F, left), p27 (F, middle), p21 (F, right), and percent EdU (G) for serum starved, WT cells treated with LATSi (0.5 μM TDI-011536) for 0-24h or co-inhibited with TEADi (grey outline, 1 μM GNE-7883). N = 2 (CDK6) or 3 independent experiments (all other proteins). Student's one-sample t-test: SKP2, p = 0.0159 (3h), 0.0178 (6h), 0.0468 (12h); cyclin D1, p = 0.0168 (6h), 0.0271 (24h); EdU, p = 0.0285 (12h), 6.65×10^{-4} (24h).

H-I) Mean YAP ratio (H), cyclin D1/p27 ratio (I, left), percent p-Rb (I, middle) and EdU-positive (I, right) cells following TEADi treatment of starved WT cells treated with LATSi (dashed line) or CA-YAP (solid line, same data as Figure 3I). Mean ± SEM (n = 3 independent experiments). Student's two-sample t-test, CA-YAP: cyclin D1/p27 ratio, p = 0.0354 (24h, 36h); %p-Rb, p = 0.0377 (24h), 0.0328 (36h); EdU, 0.0038 (24h), 0.0012 (36h).

All data are G1-gated. Colored rings indicate replicate wells within matched experimental replicates.

Figure S4

Figure S4, related to Figure 4

A) <u>Top</u>: GO terms enriched for significantly up-regulated genes after 24-hour YAP induction (q < 0.05, Log2FC > 1.0) relative to control (0.1 % FBS) (Figure 5A). <u>Bottom</u>: GO process terms enriched for significantly down-regulated genes (q < 0.05, Log2FC < -1.0).

B) Mean percent p-Rb for serum-starved WT and CA-YAP cells treated with gefinitib (EGFR inhibitor) at indicated doses (µM) for 24h (n = 2 independent experiments).

C) Mean fold-change in nuclear YAP levels for WT cells treated with DMSO or LATSi ± washout during EGF treatment (0.5 µM TDI-011536, 25 ng/mL EGF) as in Figure 4G (n = 3 independent experiments).

D) Percent p-Rb binned by local density for serum-starved WT cells pre-treated with DMSO prior to 14h EGF (25 ng/mL) or control (CTRL) treatment, as in Figure 4G. Data are representative of 3 independent experiments (n > 25,000 cells/condition).

E) Mean fold-change in cytoplasmic p-S6 (240/244, <u>left</u>) and cyclin D1 (<u>right</u>) for cells treated as in (E), G1-gated (n = 3 independent experiments, where matched colors indicate values from same experiment).

F) Mean percent p-Rb for serum-starved WT cells pre-treated with DMSO or LATSi (5 µM, TRULI) for 24h prior to washout and 14h mitogen treatment (25 ng/mL EGF and HGF, 1 µM LPA). N = 2 independent experiments, where matched colors indicate well-replicates/experiment.

All single-cell data are G1-gated cells.

 Figure S5, related to Figure 5

 A-B) Fold-change in protein levels of p-AKT(S473) and cyclin D1 (A), and p-S6(B, <u>top</u>: 235/236;

 <u>bottom</u>: 240/244) for serum-starved WT and CA-YAP cells treated with PI3Ki, AKTi, or mTORi,

 as in Figure 5E. N = 2 (p-AKT, p-S6-240/244) or 3 (cyclin D1, p-S6-235/236) independent

 experiments, where matched colors indicate values from the same experiment.

 C-D) Mean fold-change in protein levels of p-ERK and Fra1 (C), cyclin D1 and p27 (D) for

 serum-starved WT and CA-YAP cells treated with MEKi as in Figure 5F. N = 2 (p-ERK, Fra1,

 p27) or 3 (cyclin D1) independent experiments, matched colors indicate values from the same

 experiment.

 E) Mean percent p-Rb in serum-starved WT and CA-YAP cells treated with DMSO or combined

 mTOR+MEKi (100 nM Torin2, 100 nM PD0325901). N = 2 independent experiments, matched

 colors indicate values from the same experiment.

 All data are from cells gated for G1-phase.

Figure S6

Figure S6, related to Figure 6

A) Representative immunofluorescence images of YAP staining in WT, Merlin KO, and TEADi-treated Merlin KO (0.5 µM GNE-7883) in 10% serum. Scale bar = 50 µm.

B) Mean fold-change in YAP ratio and nuclear YAP levels in Merlin KO treated with DMSO or TEADi relative to WT cells, as in (A). N = 3 independent experiments.

C) Percent p-Rb per local cell density bin for Merlin KO cells maintained in titrated serum levels (gating for all cells, as in Figure 1C). N > 20,000 cells/condition.

D) Mean percent p-Rb for serum-starved Merlin KO cells treated with gefitinib (EGFR inhibitor) at indicated doses (µM) for 24h. N = 2 independent experiments.

E) Top: starvation-release protocol for WT or Merlin KO cells with control (CTRL), EGF, HGF, or FGF (all 25 ng/mL). Bottom: percent p-Rb comparing WT (left) and Merlin KO (right) treated with CTRL or mitogens (n>12,000 cells/condition). Data are representative of 2 independent experiments.

F) Mean fold-change in p-ERK (top) and cyclin D1 (bottom) for WT and DOX-induced CA-YAP cells treated as in Figure 5F in starvation (left) and 10% serum conditions (right). N = 2 independent experiments, where matched colors indicate values from the same experiment.

G) Percent p-Rb binned by local density for WT and DOX-induced CA-YAP cells starved (left) or in 10% serum (right) treated with DMSO or MEKi as in (F). Starved: n > 5,000 cells (WT), n > 20,000 cells/condition (CA-YAP); 10% serum: n > 10,000 cells (WT), n > 20,000 cells/condition. Data are representative of 2 independent experiments.

H-I) Heatmap of DOX-induced gene expression changes for a subset of YAP/Hippo pathway genes (H) and log2-fold change (Log2FC) in expression of *LATS2* (orange), *NF2* (Merlin, yellow), and *AMOTL2* (Angiomotin-like 2, red) over time of induction (I). Z-scale is row-count normalized z-score (H), n = 3 independent experiments.

Adjusted p-values (I): *LATS2*, 0.0085 (6h), 1.09×10^{-17} (12h), 2.75×10^{-54} (24h), 1.56×10^{-59} (36h); *NF2*, 2.12×10^{-8} (6h), 1.18×10^{-25} (12h), 1.24×10^{-11} (24h), 9.16×10^{-15} (36h); *AMOTL2*, 0.0078 (6h), 2.18×10^{-21} (12h), 9.00×10^{-46} (24h), 3.44×10^{-48} (36h).

J-K) Heatmap of differential expression (row z-score) of select down-regulated inhibitory and cell-contact related genes (J) and protocadherins (K) from RNAseq, grouped by 3 independent replicates and time points.

All binned data are G1-gated unless stated otherwise, with n > 100 cells/bin.