Pre-fusion AAA+ remodeling of target-SNARE protein complexes enables synaptic transmission

K. Ian White[1,2,3,4,5,**], Yousuf A. Khan[1,2,3,4,5], Kangqiang Qiu[6], Ashwin Balaji[7], Sergio Couoh-Cardel[1,2,3,4,5], Luis Esquivies[1,2,3,4,5], Richard A. Pfuetzner[1,2,3,4,5], Jiajie Diao[6], Axel T. Brunger[1,2,3,4,5,8,*]

[1]Department of Molecular and Cellular Physiology, Stanford University, Stanford, California, United States
[2]Department of Neurology and Neurological Sciences, Stanford University, Stanford, California, United States
[3]Department of Structural Biology, Stanford University, Stanford, California, United States
[4]Department of Photon Science, Stanford University, Stanford, California, United States
[5]Howard Hughes Medical Institute, Stanford University, Stanford, California, United States
[6]Department of Cancer Biology, University of Cincinnati College of Medicine, Cincinnati, United States
[7]Department of Chemistry, Stanford University, Stanford, California, United States
[8]Lead contact

[*]Correspondence: brunger@stanford.edu
[**]Correspondence: kiwhite@stanford.edu

Keywords

Membrane vesicle trafficking, AAA+ protein; neurotransmitter release; synapse; nanocluster; active zone; syntaxin; SNAP-25; NSF; ATP; hydrolysis

Summary

Membrane fusion is driven by SNARE complex formation across cellular contexts, including vesicle fusion during synaptic transmission. Multiple proteins organize trans-SNARE complex assembly and priming, leading to fusion. One target membrane SNARE, syntaxin, forms nanodomains at the active zone, and another, SNAP-25, enters non-fusogenic complexes with it. Here, we show that the AAA+ protein NSF (N-ethylmaleimide sensitive factor) and SNAP (soluble NSF attachment protein) must act prior to fusion. We show that syntaxin clusters are conserved, that NSF colocalizes with them, and characterize SNARE populations within and near these clusters using cryo-EM. Supercomplexes of NSF, α-SNAP, and either a syntaxin tetramer or two binary complexes of syntaxin—SNAP-25 reveal atomic details of SNARE processing and show how sequential ATP hydrolysis drives disassembly. These results suggest a functional role for syntaxin clusters as reservoirs and a corresponding role for NSF in syntaxin liberation and SNARE protein quality control preceding fusion.

Introduction

The membrane-anchored SNARE (soluble N-ethylmaleimide-sensitive factor acceptor protein receptor) proteins provide the energy to drive membrane fusion in various cellular contexts, such as intracellular trafficking and exocytosis. One critical exocytic process is synaptic transmission, in which a synaptic vesicle fuses with the presynaptic plasma membrane in the active zone of a neuron to release neurotransmitter into the synaptic cleft in response to the arrival of Ca^{2+} in the presynapse. While many proteins precisely act in concert to ensure synchronized and efficient Ca^{2+}-triggered fusion[1–5], this process is fundamentally mediated by the "zippering" of four SNARE motifs typically contributed by three to four SNARE proteins in the neuronal context— syntaxin-1a and SNAP-25 on the target plasma membrane (*t*-SNAREs), and synaptobrevin-2 on the vesicle (*v*-SNARE)—which come together to form a highly stable parallel four-helix bundle during Ca^{2+}-triggered membrane fusion. This *cis*-SNARE complex must then be disassembled in an ATP-dependent manner by the AAA+ protein NSF in concert with the adaptor SNAP proteins[6,7].

While the molecular mechanisms of this core fusion cycle and the accessory proteins required to ensure Ca^{2+}-dependent neurotransmitter release have been extensively studied[4], the organization of the various components in the context of the crowded cellular milieu remains uncertain. Numerous studies suggest nanoscale organization within the presynaptic active zone, where proteins and lipids[8–11] such as cholesterol and PIP_2 colocalize or even phase separate into discrete clusters with implications for organization within the active zone and even across the synapse[12].

The SNARE protein syntaxin shows such organization. Syntaxin is conserved from yeast to mammals[13,14]; a short C-terminal transmembrane domain, a SNARE domain, and a C-terminal

Habc domain characterize it[15,16]. In the neuronal context, the majority (~ 67%) of syntaxin

molecules form nanodomains of non-uniform density[17,18], often near the active zone[8,17–21], and

these nanodomains frequently overlap with other essential active zone proteins and even docked

granules[19,20,22–25]. Morphologically, they are dense, around 50–60 nm in diameter, and harbor as

many as 70 syntaxin molecules each[17–19,22,26]. Increased presynaptic activity drives a decrease in

syntaxin cluster size[27]. These clusters depend on the syntaxin SNARE domain (H3), which

drives oligomerization[28].

Despite extensive study, the functional purpose of these clusters has remained unclear. Moreover,

NSF, α-SNAP, and syntaxin appear to interact *in vitro*[29], and α-SNAP associates with liposomes

presenting reconstituted full-length syntaxin[30]. Binary (syntaxin—SNAP-25) complexes of

different stoichiometries may also form [31–37], and NSF disassembles these complexes *in*

vitro[31,38–40]. Furthermore, continuous ATP hydrolysis by NSF on plasma-membrane mimicking

vesicles with reconstituted syntaxin and SNAP-25 produces a primed state for the most efficient

and synchronous synaptic vesicle fusion[3]. These observations support a critical role for NSF not

only in recycling SNARE proteins following fusion, but also in regulating the pool of fusogenic

syntaxin available to enter the vesicle fusion cycle.

Here, we show that spatially organized homo- or hetero-oligomers of syntaxin serve as

conserved molecular protein stores, and that NSF disassembles syntaxin nanodomains while

acting as a quality control factor to disassemble off-pathway binary complexes of syntaxin and

SNAP-25—a prerequisite for synaptic vesicle priming and fusion. We also describe the

molecular mechanism of disassembly and present a sequential ATP hydrolysis mechanism by

which NSF couples γ-phosphate release to SNARE substrate remodeling, with broad

implications for the AAA+ protein family.

Results

Syntaxin nanodomains are conserved hotspots of NSF activity

First, we hypothesized that NSF is enriched at syntaxin clusters in the active zone, where it must

act to liberate fusogenic syntaxin molecules. To test this, we monitored the colocalization of

syntaxin and NSF in PC12 cells, a model system for neuronal secretion, using structured

illumination microscopy (SIM). All cells were transfected with GFP-syntaxin and as well NSF

with either a short C-terminal linker followed by mScarlet-I (NSF-mScarlet-I; Fig. 1A), a mutant

form of NSF (NSF(ΔN)-mScarlet-I; Fig. 1B,), or mScarlet-I alone (Fig. 1C). NSF-mScarlet-I is

hexameric and, together with α-SNAP, retains the capacity to bind and disassemble the ternary

SNARE complex (Fig. S1E). Clear, punctate co-clustering between NSF and syntaxin is visually

apparent upon inspection of the micrographs (Fig. 1A–C); Pearson's colocalization coefficient

(PCC) was calculated between the green and red channels to quantitate this (Fig. 1D, Methods).

Cells transfected with both GFP-syntaxin and NSF-mScarlet-I showed significant and

reproducible GFP-syntaxin-NSF-mScarlet-I colocalization (Fig. 1A, PCC = 60% ± 2%; p <

0.001 vs. mScarlet-I only;), suggesting an interaction between NSF and syntaxin molecules

within the nanodomains. As a control, mScarlet-I alone did not colocalize with syntaxin, showing

a nearly three-fold reduction in colocalization (Fig 1C; PCC = 22% ± 2%). To verify that

colocalization is consistent with the binding mode of NSF, in which NSF N-domains engage

multiple α-SNAPs, which in turn coat the SNARE complex[6,40,41], we examined colocalization of

GFP-syntaxin with a mutant form of NSF with the N-domain required for α-SNAP binding—and

thus SNARE complex binding—removed (NSF(ΔN)-mScarlet-I)[42]. As expected, this reduced

colocalization to a level nearly indistinguishable from the mScarlet-I control (Fig. 1B, PCC =

28% ± 3%; p = 0.1776 vs. mScarlet-I only). Two previously characterized[43] point mutations to

the D2 domain also disrupted colocalization between syntaxin and NSF (Fig. S1A–B). These

colocalization studies suggest that NSF associates with some multimeric species involving

syntaxin molecules *in vivo.* While these results showed clustering of syntaxin in the context of

synaptic transmission, we found that clustering also occurs in the case of the conserved syntaxin

homolog in yeast, Sso1p, as well (Fig 1F). During cell division, the mother cell transfers cargo to

the budding daughter cell via secretory vesicles; Sso1p clusters are enriched in the budding cell

in areas undergoing active fusion.

Discovery of a syntaxin tetramer complex with NSF—α-SNAP

To gain insight into the contents of the syntaxin nanodomains as well as the molecular basis of

the interaction between NSF and syntaxin, we purified the complex consisting of NSF, α-SNAP,

and soluble syntaxin (residues 2–253, transmembrane region omitted) by mixing the individual

purified proteins and performing size exclusion chromatography (SEC), Fig. S2A). We then

performed cryo-EM single-particle analysis (SPA) of this complex. Following 3D classification,

5% of the particles formed a class with an overall resolution of 3.75 Å, revealing a complex

(referred to as sx20S) between hexameric NSF, four α-SNAP molecules, and a parallel four-helix

bundle of syntaxin H3 SNARE domains (sx20S; Fig. 2A, Fig. S3, Tables S1–2). The remaining

particles showed no evidence of SNARE substrate. The existence of a syntaxin tetramer is

consistent with crosslinking mass spectrometry analysis of purified soluble syntaxin[36]. The

sx20S complex further supports the notion that NSF and α-SNAP recognize parallel, four-helix

SNARE complexes of variable composition[6,7].

Similar to structures of hexameric NSF and α-SNAP bound to the neuronal ternary SNARE

complex (referred to as 20S complex) under non-hydrolyzing conditions[6,41], the sx20S complex

forms a three-tiered architecture (Fig. 2A). NSF comprises three structured domains connected

by flexible linkers—the adaptor N-domain followed by the D1 and D2 AAA+ domains. Six NSF

protomers come together to form the active form of the enzyme, a homo-hexamer. In the top tier,

or spire, the NSF hexamer's multiple N domains bind four α-SNAP molecules and a tetramer of

syntaxin H3 SNARE domains. This spire structure positions the N-terminal end of the syntaxin

tetramer near the pore of the hexameric D1 ATPase ring of NSF; this middle tier is responsible

for ATP hydrolysis and SNARE complex disassembly. The bottom tier consists of the inactive

NSF D2 ATPase ring responsible for the oligomerization of the complex.

The spire structure is dynamic and rocks about relative to the D1 pore, but three of the six N

domains are relatively well-ordered and engaged with the α-SNAP—syntaxin sub-complex. The

D1 ring is asymmetric and split; protomers A–F are defined from the bottom of the split to the

top. All six D1 domains are ordered and engage the N-terminal linker preceding the H3 domain

of one of the four syntaxin molecules via conserved pore loops, which position Y294 such that it

intercalates between every other syntaxin substrate side chain passing through the pore (Fig. 2B).

The nucleotide states of the NSF ATPase domains are consistent with a pre-hydrolysis

conformation; D1 protomer A is ADP bound, while D1 protomers B–F are ATP bound with no

evidence for divalent cation as expected (Table S2). All D2 protomers are ATP bound and lack

divalent cation.

While the spire in the complete sx20S complex reconstruction was ordered enough to distinguish

the various components, continuous disorder limited the local resolution of the spire

reconstruction. To overcome this, we performed signal subtraction to remove the D1 and D2

rings, followed by local alignment of the α-SNAP—syntaxin sub-complex structure alone (Fig.

S3, Methods). This approach improved the resolution of the reconstruction of the α-SNAP—

syntaxin sub-complex to 4.25 Å, revealing a parallel, four-helix bundle of syntaxin SNARE

motifs with a density of sufficient quality to assign characteristic side chains (Fig. 2C). All four

helices are aligned in the same register, with a unique hydrophobic layer (-3 layer) composed of

four F216 side chains present; remarkably, all fit within the core of the bundle (Fig. 2D, Fig.

S9A) and induce little overall deformation of the individual H3 domains (Fig. S9B-D, bottom

row). The parallel syntaxin tetramer bundle observed in our structure would allow membrane

incorporation of the C-terminal transmembrane domains of all four syntaxin molecules (Fig. 2A).

To determine whether NSF and α-SNAP act to disassemble the syntaxin tetramer, we devised an

in vitro, gel-based disassembly assay (Fig. 1E). Wild-type NSF, α-SNAP, and soluble syntaxin

(residues 2–253) were purified independently, mixed in the absence of Mg^{2+} (*i.e.*, non-

hydrolyzing conditions), and the complex of all three was purified. The subsequent addition of

Mg^{2+} to the purified complex resulted in a significant loss of α-SNAP—syntaxin following

filtration relative to the non-hydrolyzing control, consistent with the disassembly of the syntaxin

tetramer and concurrent disassociation of α-SNAP (Fig. 1E, 21 ± 1% loss; Fig. S1D).

Structure of a 2:1 syntaxin—SNAP-25 complex with NSF—α-SNAP

In addition to the syntaxin homo-tetramer, hetero-oligomers with SNAP-25 are expected to be

present near or even within syntaxin nanodomains. Indeed, SNAP-25 colocalizes with syntaxin

in neurons[19,22,24] and readily forms binary complexes with syntaxin *in vitro*[31–35,44,45]. Although

SNAP-25 is required for fusion, assembly of a fusogenic ternary *trans*-SNARE complex likely

proceeds first through a syntaxin—synaptobrevin intermediate templated by Munc18-1, also

colocalized to syntaxin clusters[24], and by Munc13-1[31,46,47]. Disassembly of the binary complex is

critical, as genetic deletion of NSF is lethal[48]. However, an NSF mutation that differentially

blocks ternary, but not binary, SNARE complex disassembly causes only intermediate

sensorineural defects[38]. Thus, NSF and α-SNAP must first disassemble binary complexes of

syntaxin and SNAP-25 to set the stage for the action of Munc18 and Munc13, ultimately

resulting in synchronous and efficient Ca^{2+}-triggered fusion[3,31,32,46].

To gain insight into the structural nature of binary complex formation and disassembly by NSF,

we prepared a complex of NSF, α-SNAP, and the binary *t*-SNARE complex (21bin20S)

composed of full-length soluble SNAP-25 and soluble syntaxin-1a (1–267) in the absence of

divalent cation (*i.e.*, non-hydrolyzing conditions) and performed single-particle cryo-EM (Tables

S1–2, Fig. S2B, Figs. S4–5). First, models of the NSF D1 and D2 domains with substrate bound

in the D1 pore were built into two high-resolution (2.88 Å, 2.92 Å) classes from a single dataset,

differing only in the state of F protomer engagement with substrate (Fig. S4). Next, the dataset

from which they were derived was combined with two additional datasets to enable the 3D

reconstruction of weakly populated states with ordered density for the α-SNAP—SNARE spire

(Fig. S5). Classes with evidence of NSF D1 domains engaging ordered α-SNAP—SNARE sub-

complexes were carried forward; eight 21bin20S models were built into reconstructions ranging

from 3.39–4.01 Å in resolution, differing mainly in the orientation of N domains and a sub-

complex of three α-SNAP molecules and four SNARE motifs relative to the D1 ATPase ring

(Fig. S10A). All classes revealed substrate-bound NSF without Mg^{2+} needed for ATP hydrolysis

and disassembly (Table S2), and share the same overall architecture, with a spire composed of

NSF N-domains and the α-SNAP—SNARE subcomplex on top, the NSF D1 ring in the middle

tier, and the NSF D2 ring forming the bottom tier (Fig. 3A). In all cases, the NSF D1 ATPase

ring engages SNARE substrate, with one syntaxin molecule bound to the D1 ring pore in most

structures (Fig. 3B, Table S2).

Like the sx20S complex, continuous variability of the α-SNAP—SNARE subcomplex was

present in the 21bin20S reconstructions. We again performed signal subtraction and local

alignment on α-SNAP—SNARE sub-complexes from all five classes with ordered spire

structures (Fig. S5; see Methods). This procedure produced a single α-SNAP—SNARE sub-

complex reconstruction with an overall resolution of 3.74 Å (Fig. 3C), sufficient to assign

characteristic side chains, identify the SNAREs forming the complex and their directionality

(Fig. 3D, Fig. S9A). A model composed of the ternary neuronal SNARE complex[49] (PDB ID

1N7S) with synaptobrevin removed was docked into the density, preserving a distinctive kink in

the peptide linker leading into the SN2 SNARE motif of SNAP-25. In place of synaptobrevin, a

second syntaxin molecule was placed to complete the four-helix bundle with little deviation from

its usual backbone configuration of the ternary complex (Fig. S9B). As such, the stoichiometry

of the syntaxin—SNAP-25 complex observed is 2:1, the dominant species in solution[33,36,44,45].

Sidechains comprising essential layers repack within the core of the complex (Fig. 3D) compared

to the ternary SNARE complex. For example, the ionic central layer (0 layer) is reorganized;

Arg56 of synaptobrevin is replaced with a second Gln226 from the additional syntaxin molecule,

which adopts an extended rotamer to mimic the arginine it replaces. The SNAP-25 linker, which

joins the two SNARE motifs, is largely disordered but blocks binding by a fourth α-SNAP

molecule, as observed for the neuronal SNARE complex[6].

Docking the 2:1 binary complex structure back into the seven individual 21bin20S

reconstructions reveals an apparent preference for syntaxin, with six of eight reconstructions

featuring the N-terminus of syntaxin H3 bound within the pore of the D1 ring (Fig. 3B). As in the

case of the 20S with ternary SNARE complex, this observation is consistent with previous work,

in which disassembly was not contingent on specific SNARE N-terminus identity[6,40]. Taken

together, we observe that the NSF D1 ring can bind to at least two different SNAREs, syntaxin,

and SNAP-25, in various configurations.

Structure of a 2:2 syntaxin—SNAP-25 complex with NSF—α-SNAP

To gain insight into the disassembly of the binary SNARE complex, we repeated the binary 20S

preparation but added Mg^{2+} and an ATP regeneration system[50] (*i.e.*, hydrolyzing conditions) to

the purified complex ~1 min before vitrification for single particle cryo-EM (Tables S1–2, Fig.

S2C, Fig. S6). The addition of the regeneration system more closely mimics the metabolic state

of the cell, and it allows multiple cycles of SNARE disassembly and reassembly to proceed as

observed under far more dilute conditions[32]. Following 3D classification, 41% of particles

formed a complete complex, all of which have ordered α-SNAP—SNARE spire density and

differ largely by spire pose (Fig. S10A). Interestingly, classes with poorly defined SNARE

substrate, as observed under the non-hydrolyzing condition, were no longer observed. The

remaining 59% of particles were SNARE-substrate free and grouped into three classes with split

D1 and D2 rings; presumably, this is a post-disassembly, substrate-released state, and it is not

observed under non-hydrolyzing conditions.

Substrate-bound reconstructions reveal a flattened D1 ring in closer proximity to the D2 ring and

a strikingly different α-SNAP—SNARE spire structure than observed in the 21bin20S complex

under non-hydrolyzing conditions (Fig. 4A, Fig. S11A). Again, syntaxin is bound within the pore

of the D1 ring (Fig. 4B, Table S2). We observed a nearly symmetric α-SNAP—SNARE

subcomplex comprising four α-SNAP molecules coating a four-helix bundle. Even without

further processing, this structure was recognizable as a 2:2 syntaxin—SNAP-25 sub-complex

formed by two syntaxin H3 SNARE domains and two SNAP-25 SN1 SNARE domains; this

binary 2:2 configuration was previously observed in crystal structures of syntaxin-1 H3 and

SNAP-25 SN1 domains[34] and was unexpected as we prepared the complex with full-length

proteins, implying that two independent SNAP-25 molecules contribute their SN1 SNARE

domains to this complex. Presumably, disassembly of the binary 2:1 complex (Fig. 3C) had

occurred and was followed by reassembly to form the 2:2 binary complex (Fig. 4C) before

vitrification.

To improve the resolution of the spire reconstruction, signal subtraction, and local alignment

were performed as before (Fig. S6). This approach improved the reconstruction of the spire

proteins to 4.22 Å, enabling docking of a crystal structure of the 2:2 syntaxin—SNAP-25 SN1

complex[34] into the density (Fig. 4C), which was of sufficient quality to assign characteristic side

chains, to identify the SNARE domains and their directionality (Fig. 4D, Fig. S9A). The 2:2

binary complex was fit back into the full reconstructions, enabling modeling of the full

22bin20S. Furthermore, we were able to reproduce this result by reconstituting the complex with

only the SNARE domains of syntaxin and SNAP-25 under non-hydrolyzing conditions

(min22bin20S, Fig. S7A–D), yielding the same spire configuration but to a higher resolution of

3.86 Å following local reconstruction (Fig. S7E, Fig. S9A). This reconstruction also reveals the

interface between an N-domain of NSF and an α-SNAP molecule in detail; the interface is

characterized by a core hydrophobic interaction surrounded by charge complementation (Fig.

S7F).

Insights into SNARE substrate engagement and processing by NSF

Collectively, we obtained many datasets with NSF, α-SNAP, and several SNARE substrates

under non-hydrolyzing and hydrolyzing conditions (Tables S1–2), presenting an opportunity to

investigate the molecular mechanism of SNARE substrate processing and chemomechanical

energy transduction in NSF. Together, the resulting high-resolution structures derived from the

different classes from these datasets capture the hydrolysis cycle of NSF and reveal important

details about its reaction coordinate (Fig. 5–6, Figs. S11–13, Table S2).

For datasets with particles collected under non-hydrolyzing conditions, NSF is found in two

general conformations, primarily differentiated by their F protomer state (Fig. 5A, left column).

Two high-resolution 21bin20S classes from the 2:1 binary SNARE complex dataset reveal large-

scale conformational displacement connected to D1 F promoter nucleotide state and SNARE

substrate engagement. The first class is ATP-bound and fully SNARE substrate-engaged, while

the second has a nucleotide-free and SNARE substrate-disengaged F protomer (Fig 5A, left, top,

and bottom, respectively; Fig. S4). The engagement of F at the top of the ring accompanies a

relaxation of A into the split. Otherwise, the classes are nearly identical; protomers B–E D1

domains also bind ATP, while protomer A, at the bottom of the D1 split, is ADP-bound in both

cases. Comparison of the two classes reveals limited differences outside of the protomers A and

F positions, as the change in protomer F position allows additional relaxation of protomer A into

the split.

In contrast to the non-hydrolyzing datasets, particles in the hydrolyzing dataset fall into two

categories—substrate-bound 22bin20S (Fig. 5A, middle column) or substrate-free NSF (post-

hydrolysis; Fig. 5A, right column), of which there are ten and three classes, respectively (Table

S2, Fig. S6). The particles show a population shift towards a flatter D1 ring that is closer to D2

relative to the non-hydrolyzing class, particularly in the case of the SNARE-substrate free classes

(Fig. S11). Most strikingly, no SNARE substrate-bound classes show an F protomer as fully

engaged as those in non-hydrolyzing classes; in all but one class, F is ADP bound, SNARE

substrate disengaged, and sits between engaged A and E protomers. However, a subclass with an

ATP-bound F protomer was found using a subdivision of a large class with 3D Variability

Analysis[51] (3DVA; Fig. S6, Fig. 12C–D) for which the interprotomer interface with the

neighboring D1 protomer E is nearly—but not completely—formed. Critically, the pore loop

approaches the substrate-engaged state found in the non-hydrolyzing classes; together, this

reveals a trajectory of protomer A dissociation, nucleotide release and rebinding, and, finally,

SNARE substrate rebinding (Fig. 5D). Furthermore, classes with ordered syntaxin substrate are

gripped by the tyrosine residues in the pore of the D1 ring around eight residues further into the

primary sequence under hydrolyzing conditions (Fig. S10B).

Sequential hydrolysis by NSF drives conformational cycling

As noted above, the nucleotide states of the D1 protomers help explain the pattern of engagement

with the SNARE substrate (Fig. 5A–C, Table S2). How does nucleotide hydrolysis drive this

process? Unlike the non-hydrolyzing 21bin20S reconstructions, hydrolyzing 22bin20S

reconstructions are only ATP-bound in protomer E, indicative of a sequential hydrolysis

mechanism (Fig. 5A, middle column). Except for the rebinding F protomer described above, all

other D1 protomers are ADP bound with variability in the presence of Mg^{2+} and inorganic

phosphate (P_i); in most cases, protomers C and D have ADP•P_i and Mg^{2+}, protomer B is ADP-

bound, and protomer A is typically ADP-bound only. Finally, substrate-free classes are always

ADP bound, even in protomer E; they are flat, have some N-domains packed between the D1 and

D2 rings, and are characterized by a split in both the D1 and D2 rings between protomers A and

F (Fig. 5A, right column); this is consistent with a SNARE substrate side-loading mechanism

proposed for the NSF ortholog in yeast, Sec18[52]. In both hydrolyzing and non-hydrolyzing

datasets, all D2 domains are ATP-bound, with Mg^{2+} present in the hydrolyzing condition.

Together, these classes represent key steps in the hydrolysis cycle, revealing the binding of Mg^{2+}, hydrolysis of ATP to ADP•P_i at protomer E, and subsequent P_i release down-ring; these states are accompanied by conformational change in the Walker A, Walker B, and Sensor 1 active site residues in *cis* and the arginine fingers of the down-ring protomer in *trans* (Fig. 5B), and are connected in turn to larger scale conformational change within the D1 domain (Fig. 5C). Critically, P_i release following hydrolysis leads to arginine finger disengagement (Fig. 5B, right two panels) and subsequent relaxation and solvation of the interprotomer interface, enabling the protomer A transition to protomer F and subsequent substrate rebinding.

What triggers hydrolysis at E, and what conformational changes follow? While hydrolysis could proceed spontaneously, this would be inefficient in the absence of an ATP-bound protomer F up-ring to capture accompanying substrate translocation. We hypothesized that some allosteric mechanism must regulate this process. To identify such a mechanism in an unbiased manner, we performed principal components analysis (PCA) of Cα coordinates of 132 D1 protomers across all datasets followed by hierarchical density-based clustering (HDBSCAN) in high-dimensional space, revealing the conformational cycling of the NSF D1 protomers in a low-dimensional coordinate as well as a corresponding allosteric mechanism for modulating hydrolysis by protomers E (Fig. 6; see also Methods).

Overall, we find protomers divided into five interpretable clusters connected to ATP hydrolysis, P_i release, and protomer cycling at the D1 split during SNARE disengagement, nucleotide rebinding, and SNARE reengagement (Fig. 6A). Here, we focus on the unique features of the protomers E under hydrolyzing conditions in two dimensions of the 11-dimensional latent space, where they cluster far from other protomers (Fig. 6A, transition 1). As noted above, protomers E are the only ATP-Mg^{2+} bound D1 protomers under hydrolyzing conditions (Fig. 6B), and they are

uniquely characterized by an extreme large-small subdomain angle (Fig. 6C, Fig. S11).

Examination of PCA and clustering results reveal conformational states unique to these protomers (Fig. 6E), notably a discrete Walker A (α_3) helix position ("down" state) relative to the rest of the D1 large subdomain that uniquely positions the nucleotide near key catalytic residues (N374, R388; Fig. S13), a unique conformation of a highly conserved[56] segment of the N-D1 linker connected to N-domain engagement ("up A" state), and the absence of a fully engaged "latch loop" (residues 457–467) contributed by the up-ring protomer F in *trans* ("intermediate A" state; Fig. S12A). All three of these features appear structurally coupled (Fig. S12B) and comprise an allosteric integration mechanism by which the position of nucleotide in the active site is modulated at a distance by both N-domain engagement to SNARE substrate via the N-D1 linker conformation and by the presence of an up-ring protomer F via latch loop engagement.

To test the importance of the latch loop interaction in *trans*, a mutant form of NSF was generated in which a portion of the latch loop (residues 460–466) is replaced with a short glycine-serine linker (SGS). An NSF activity assay based on fluorescence dequenching (see Methods) shows a highly significant, ~72% reduction in the rate of SNARE complex disassembly ($p = 0.0005$; Fig. 6D). On the other hand, the functional importance of the N-D1 linker has already been established. We note that a mutation to NSF in the most conserved region N-D1 linker, I209N, has a direct behavioral effect in zebrafish; NSF(I209N) is defective in the disassembly of ternary SNARE complexes but not binary SNARE complexes, explaining this phenotype[38,56]. This isoleucine side chain directly contacts the engaged latch loop and is adjacent to S207, a putative phosphorylation site[53], raising the possibility of post-translational modification in regulating NSF activity. Together, these observations support an allosteric mechanism critical in both effecting and modulating NSF function.

Discussion

Our observation of NSF colocalization to syntaxin clusters *in vivo*, coupled with direct structural

evidence of disassembly of both syntaxin tetramers and syntaxin—SNAP-25 complexes,

explains a previously overlooked but essential aspect of the synaptic vesicle fusion cycle, with

broad implications for membrane trafficking (Fig. 7A). While NSF is well known for its role in

ternary *trans*-SNARE complex disassembly following fusion, this work, combined with

functional studies[3,31,32,46], also establishes NSF as a critical factor for synaptic vesicle priming

before fusion. NSF is enriched at and near syntaxin clusters, which are, in turn, found at the

presynapse. The observation here of a parallel, four-helix syntaxin-1a bundle suggests that these

clusters must be structured, at least in part, and that NSF and α-SNAP act to disassemble the

syntaxin tetramers on the perimeter of a given cluster, increasing syntaxin mobility at the

synapse[27]. Individual syntaxin molecules liberated from clusters by NSF can then be captured by

the SM protein Munc18 and ultimately be used to assemble fusogenic *trans*-SNARE complexes.

However, a high local concentration of SNAP-25 is present near the syntaxin clusters as well, so

free syntaxin may also enter a non-productive binary SNARE complex with SNAP-25 with a

stoichiometry dependent upon local concentrations and configurations of either protein. In this

work, we observe binary complexes with two stoichiometries—2:1 and 2:2 binary complexes of

syntaxin and SNAP-25, with the latter re-forming following NSF-mediated disassembly of the

former under hydrolyzing conditions. The 2:1 configuration dominates in the non-hydrolyzing

condition—and in other *in vitro* studies[33,45]—but is derived from co-expressed proteins that were

extensively purified without NSF action. The 2:2 configuration is only found following NSF-

mediated disassembly of the 2:1 configuration. This effect may be due to the dissociation of the

SNAP-25 SN1 and SN2 domains during disassembly by NSF, increasing the likelihood of

binding to other SN1 domains in *trans*. The structural plasticity of the SNARE complexes is also

of note; taken together, the repeated observation of a twisted, parallel, four-helix bundle with

minimal backbone distortion (Fig S9B) emphasizes the functional conservation latent in the

SNARE domain sequences studied thus far, as well as the importance of additional SNARE

protein accessory domains in differentiating SNARE protein function[52].

Regardless, NSF and α-SNAP must thus constantly act to disassemble both syntaxin tetramers

and binary SNARE complexes to maintain a fusogenic population of *t*-SNAREs[3], which

ultimately enter the *trans*-SNARE complex with the *v*-SNARE synaptobrevin-2 and set the stage

for Ca^{2+}-triggered fusion in cooperation with synaptotagmin and complexin[54,55]. Indeed, the

energy released by SNARE protein zippering during vesicle fusion should be related to the

hydrolysis of ATP by NSF, which effectively maintains them in a high-energy state. Thus,

syntaxin nanodomains could effectively act as high-density storage sites for syntaxin, and NSF

could tap these stores under periods of high-frequency vesicle release. Modulation of NSF

activity by post-translational modification could thus be one way to regulate this process[38].

Finally, given the observation of clustered Sso1p in yeast, and the high degree of structural and

functional conservation between NSF and Sec18[52], this seems likely to be a conserved

organizing principle across fusion contexts.

At the atomic level, how does NSF accomplish the energy transfer needed to disassemble various

SNARE complexes? The structural details of the hydrolysis cycle (Fig. 5) suggest sequential

hydrolysis by the six D1 ATPase domains of NSF. Hydrolysis drives conformational change,

which is coupled to the application of force to one member of the SNARE complex, destabilizing

it and promoting disassembly (Fig. 7B, Movie S1). Rather than relying on a non-physiological

nucleotide state promoted by chelation of divalent cations or non-hydrolyzable nucleotide

analogs (*e.g.*, ATPγS), we established a reaction condition more closely approximating cellular

metabolism by using a physiological ATP regeneration system. Critically, hydrolysis proceeds

sequentially around the asymmetric NSF D1 ring, occurring near the "top" of the ring at

protomer E. Down-ring protomers reveal evidence for a transient post-hydrolysis reaction state

in which ADP•P_i is generally observed before P_i release from protomers C or B and an associated

disruption and solvation of the interprotomer interface. This mechanism is likely conserved in

other members of the AAA+ family, such as p97 (Supplemental Discussion).

Furthermore, the continuous transition from protomer A at the bottom of the ring to protomer F

at the top follows ADP release and ATP rebinding (Fig. 7B, Movie S1). This cycling appears

coupled to the triggering of hydrolysis at protomer E in a process allosterically mediated by

several key structural elements. Protomers E in the hydrolyzing condition position ATP in the

active site via a unique pre-hydrolysis conformation of the Walker A helix. Presumably, full

protomer F rebinding triggers hydrolysis, likely driven by an allosteric shift associated with full

latch engagement with the N-D1 linker region and further formation of the interprotomer

interface (Fig. 6, Fig. S12).

Protomer F rebinding is also connected to the mechanism by which engagement of the SNARE

substrate is maintained during processing. In the absence of a completely protomer F-rebound

state in the hydrolyzing condition, however, we cannot distinguish between SNARE substrate

rebinding before or after protomer E hydrolysis; the former case (ATP in fully rebound F and E)

represents a classical "hand-over-hand" mechanism, while the latter case (ATP in fully rebound

F, ATP ADP•Pi in E), a ratchet mechanism in which F re-engages substrate during or

immediately following a translocation event driven by protomer E hydrolysis, locking in any

substrate translocation that took place[57]. Given the highly flattened D1 ring state of the SNARE-

substrate free classes with ADP•Pi in protomers E (Fig. S11), the latter case is perhaps more

likely; flattening upon hydrolysis at protomer E could exert a pulling force on the substrate, and

translocation would then effectively take place with substrate engagement by a new protomer F

and release of the old protomer A.

Together, our findings expand the role of NSF in the membrane fusion cycle while further

demonstrating its contribution as a model system for understanding energy transduction and

protein remodeling in AAA+ proteins. NSF acts as a mobile engine of the presynapse—working

together with the SNAP adaptor proteins to convert the chemical energy latent in ATP into the

physical work needed to organize, maintain, and drive synaptic function through disassembly of

a variety of SNARE proteins.

Acknowledgments

We thank Richard Held, Jeremy Leitz, Rama Ranganathan, Subu Subramanian, Benjamin White,

Susan Wu, and Minglei Zhao for stimulating discussions; E. Montabana and C. Zhang for

assistance with the electron microscopy data collection at the Stanford University cEMc; and

Paul D. Adams, Pavel Afonine, and Nigel Moriarty for assistance with PHENIX refinement.

Funding was provided by the US National Institutes of Health (NIH; RO1MH63105 to A.T.B.)

and the Helen Hay Whitney Foundation/HHMI (to K.I.W.). Molecular graphics and analyses

performed with *UCSF ChimeraX*, developed by the Resource for Biocomputing, Visualization,

and Informatics at the University of California, San Francisco, with support from National

Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure and Computational

Biology, National Institute of Allergy and Infectious Diseases. This article is subject to HHMI's

Open Access to Publications policy. HHMI laboratory heads have previously granted a non-

Author contributions

K.I.W. and A.T.B. developed the research plan and experimental strategy. K.Q. and J.D.

performed the cellular imaging experiments in PC12 cells and analyzed the data. Y.A.K. and

A.B. performed the cellular imaging experiments in yeast and analyzed the data. K.I.W., Y.A.K.

S.C., L.E., and R.P. purified all proteins. K.I.W., S.C., and Y.A.K. performed the disassembly

experiments. K.I.W. performed the cryo-EM experiments and analyzed the data. K.I.W. and

A.T.B. wrote the manuscript with input from the other authors.

Declaration of interests

459 All authors declare no competing interests.

Supplemental information

Associated Supplemental information file: Supplemental Discussion, Supplemental Figures 1–13

Supplemental Tables 1–2.

Supplemental Movie 1

Figures

Figure 1 | NSF colocalizes to syntaxin nanodomains in an N-domain dependent manner and
disassembles syntaxin oligomers

The interaction between NSF and syntaxin nanodomains in living cells. NSF-mScarlet-I and

syntaxin-GFP were transfected into PC12 cells, and their conjugated fluorescent proteins were

observed by SIM. Their interaction was assessed by measuring Pearson's colocalization

coefficient (PCC) between the channels using *ImageJ*. Clustering of the syntaxin homolog Sso1p

also occurs in yeast.

A. Colocalization between syntaxin-GFP nanodomains and wild-type NSF-mScarlet-I. NSF

often forms punctate clusters that overlap with syntaxin (PCC = 60% ± 2%, mean ± SEM).

B. Colocalization between syntaxin-GFP nanodomains and NSF(ΔN)-mScarlet-I, in which the

N-domain of NSF that mediates the interaction with the α-SNAP—SNARE subcomplex has

been removed. Disruption of this interaction is evident and quantifiably reduces NSF

colocalization to syntaxin nanodomains (PCC = 28% $\pm$ 3%, mean $\pm$ SEM).

C. Control showing colocalization between syntaxin-GFP and mScarlet-I alone. No specific

interaction between these proteins is apparent (PCC = 22% $\pm$ 2%, mean $\pm$ SEM).

D. Comparison of wild-type NSF-mScarlet-I, NSF(ΔN)-mScarlet-I, and mScarlet-I clustering to

syntaxin-GFP nanodomains (M $\pm$ SEM, n = 10). Statistical significance was established by two-

tailed Student's t-tests, with $p < 0.05$ considered indicative of statistical significance.

Colocalization between wild-type NSF-mScarlet-I and syntaxin-GFP is significantly different

from both colocalization between NSF(ΔN)-mScarlet-I and syntaxin-GFP and colocalization

between mScarlet-I and syntaxin-GFP ($p < 0.0001$). Colocalization between NSF(ΔN)-mScarlet-

I and syntaxin-GFP and colocalization between mScarlet-I and syntaxin-GFP was not

significantly different (p = 0.1776).

E. NSF and α-SNAP disassemble oligomeric syntaxin as assessed by a gel-based de-enrichment

assay. sx20S was purified and split into two pools. EDTA was added to the first pool as a non-

hydrolyzing control, while Mg^{2+} was added to the second to initiate syntaxin complex

disassembly. Each pool was then concentrated using a 100,000 MWCO filtration device,

retaining 20S complex or α-SNAP—syntaxin subcomplex but allowing individual syntaxin and

α-SNAP molecules to pass, de-enriching them. Under hydrolyzing conditions, α-SNAP and

syntaxin were significantly de-enriched by around 21 $\pm$ 1% (p = 0.0148), consistent with the

disassembly of oligomeric syntaxin by NSF.

F. Sso1p forms dense clusters in the yeast plasma membrane during cell division (white

triangles). Sso1p-GFP was expressed in *S. cerevisiae* and imaged by SIM. Sso1p clusters were

enriched in the budding daughter cell (left); discrete clusters typically approached sizes near the

Nyquist limit of 160 nm (Fig S1C).

Figure 2 | NSF and α-SNAP engage a tetrameric SNARE complex composed of four
syntaxin H3 domains

The structures of the sx20S (class #1, 3.75 Å) and the α-SNAP—syntaxin tetramer subcomplex

(class #2, 4.07 Å).

A. Hexameric NSF engages the α-SNAP—syntaxin tetramer subcomplex via its N-domains

(salmon), several of which engage four α-SNAP molecules (gold) surrounding the soluble,

tetrameric syntaxin H3 complex (dark red) within the catalytic D1 ring of NSF (light blue). The

structural D2 ring of NSF (purple) does not engage the SNARE substrate. A hypothetical

membrane is illustrated in grey. For the SNARE proteins, domains visible in the reconstruction

of the SNARE complex are indicated with an asterisk; the syntaxin Habc domains were not

observed.

B. NSF engages one of the four syntaxin molecules in the tetramer via a series of pore loop

interactions; a tyrosine side chain (Y294) on each of the six protomers intercalates in between

every other side chain of a portion of syntaxin N-terminal to its H3 domain. The corresponding

sharpened map is contoured at 4 σ.

C. The α-SNAP—syntaxin tetramer subcomplex model, with two of the four α-SNAP models

hidden for clarity. The syntaxin H3 SNARE complex forms a twisted, parallel α-helical bundle

with a series of key SNARE complex layers preserved. See Fig. S9 for additional detail.

D. Slices through the α-SNAP—syntaxin tetramer subcomplex model and sharpened map at the -

4–0 layers, with key residues lightened for emphasis. Each layer is characterized by nearly planar

interactions between four identical side chains within the helical bundle. The primary

interactions between α-SNAP and the syntaxin H3 complex are observed near the -2 and -3

layers, where several functionally essential[58] α-SNAP side chains (L197, Y200) are positioned by

a loop that buries them within the groove of the bundle. No symmetry was imposed during

reconstruction.

Figure 3 | NSF and α-SNAP engage a 2:1 binary SNARE complex of syntaxin H3 and SNAP-25 SN1 and SN2 SNARE domains under non-hydrolyzing conditions

A representative 21bin20S model (class #5, 3.39 Å) and the α-SNAP—2:1 binary SNARE subcomplex model (class #13, 3.74 Å).

A. Hexameric NSF engages the α-SNAP—2:1 binary SNARE subcomplex via multiple N-domains (salmon), which engage three α-SNAP molecules (gold) surrounding the 2:1 binary SNARE complex composed of two syntaxin H3 domains (dark red) and the SN1 and SN2 SNARE domains of a single SNAP-25 molecule (green). Part of the linker N-terminal to one of the syntaxin H3 domains is found within the catalytic D1 ring of NSF (light blue). The structural D2 ring of NSF (purple) does not engage the substrate. A hypothetical membrane is illustrated in grey. For the SNARE proteins, domains visible in the reconstruction of the SNARE complex are indicated with an asterisk.

B. NSF engages one of the two syntaxin molecules in the 2:1 binary SNARE complex via a

series of pore loop interactions involving Y294. A sharpened map is shown, contoured at 5 σ. Of

the eight 21bin20S classes with the 2:1 binary SNARE complex from the dataset collected under

non-hydrolyzing conditions, six have syntaxin molecules, and two have SNAP-25 molecules

bound to the pore of the D1 ring.

C. The α-SNAP—2:1 binary SNARE subcomplex model. Three α-SNAP molecules bind the 2:1

binary SNARE complex; a flexible linker between SNAP-25 SN1 and SN2 blocks access by a

fourth. The 2:1 binary SNARE complex forms a twisted, parallel α-helical bundle with a series

of characteristic SNARE complex layers preserved. This configuration also places the N-

terminal portion of the linker, which bears cysteines often palmitoylated *in vivo*, near the

membrane[59]. See Fig. S9 for additional detail.

D. Layer slices through the α-SNAP—2:1 binary SNARE subcomplex model and sharpened

map. Each layer is characterized by nearly planar interactions between four side chains within

the helical bundle. The primary interactions between α-SNAP and the syntaxin H3 complex are

found near the -2 and -3 layers, where several functionally essential[58] α-SNAP side chains (L197,

Y200) are positioned by a loop that buries them within the bundle groove.

Figure 4 | NSF and α-SNAP engage a 2:2 binary SNARE complex of syntaxin H3 and SNAP-25 SN1 SNARE domains under hydrolyzing conditions

A representative 22bin20S model (class #19, 3.56 Å) and the α-SNAP—2:2 binary SNARE subcomplex model (class #23, 4.14 Å).

A. Hexameric NSF engages the α-SNAP—2:2 binary SNARE subcomplex via multiple N-domains (salmon), which engage three α-SNAP molecules (gold) surrounding the soluble, 2:2 binary SNARE complex composed of two syntaxin H3 domains (dark red) and two SNAP-25 SN1 domains contributed by two separate full-length SNAP-25 molecules (green). A portion of the linker N-terminal to one of the syntaxin H3 domains is observed within the catalytic D1 ring of NSF (light blue). The structural D2 ring of NSF (purple) does not contain density for a substrate. A hypothetical membrane is illustrated in grey. For the SNARE proteins, domains

visible in the reconstruction of the SNARE complex are indicated with an asterisk. See Fig. S9

for additional detail.

B. NSF engages one of the two syntaxin molecules in the 2:2 binary SNARE complex via a

series of pore loop interactions involving Y294. A sharpened map is shown, contoured at 6 σ. Of

the five 22bin20S classes with the 2:2 binary SNARE complex, all are found with syntaxin in the

D1 pore, while two are observed with the N-terminus of the SNAP-25 SN1 domain in the pore.

C. The α-SNAP—2:2 binary SNARE subcomplex model. Four α-SNAP molecules bind the 2:2

binary SNARE complex, with two SNAP-25 SN1 SNARE domains and two syntaxin H3

domains. This configuration would place two independent N-terminal portions of the SNAP-25

linker bearing cysteine sidechains often palmitoylated *in vivo* near the membrane[59].

D. Slices through the α-SNAP—syntaxin tetramer subcomplex at the -4–0 layers. Each layer is

characterized by nearly planar interactions between four side chains within the helical bundle.

The primary interactions between α-SNAP and the syntaxin H3 complex are found near the -2

and -3 layers, where several functionally essential[58] hydrophobic α-SNAP side chains, L197 and

Y200, are positioned by a loop that buries them within the bundle groove. Symmetry was not

enforced during local reconstruction. Note that the near-twofold symmetry of the bundle leaves

some sidechain densities poorly defined, likely resulting from the superposition of the syntaxin

and SNAP-25 residues in a given layer.

Figure 5 | Comparison of NSF structures under non-hydrolyzing and hydrolyzing conditions reveals interplay between sequential AAA+ hydrolysis and substrate rebinding

A. Key conformational states of the NSF D1 ATPase ring. Two primary states dominate in the absence of Mg^{2+} (left); the top class (#4) has a SNARE substrate disengaged, nucleotide-free protomer F, while the bottom class (#3) has a fully SNARE substrate-engaged, ATP-bound F. With the addition of Mg^{2+} (middle), E is ATP- and Mg^{2+}-bound. Protomers D, C, and occasionally B retain P_i. and Mg^{2+}. All A protomers are ADP-bound. 3DVA-derived subclasses reveal a coordinate for nucleotide exchange and protomer F engagement. The top subclass (#18) has a substrate-disengaged, ADP-bound F. The bottom subclass (#17) has a nearly substrate-engaged, ATP-bound F. Finally, substrate-free, post-hydrolysis classes such as #24 (right) show an ATP-free D1 ring. For clarity, the spire, SNARE substrates, and the D2 ring are omitted in the figures. See Tables S1–2 for more information on specific classes.

B. Atomic details of nucleotide hydrolysis. The P-loop binds nucleotide. Arginine fingers R385 and R388 from the down-ring protomer coordinate nucleotide phosphates. Without Mg^{2+}, a water molecule sits between the nucleotide phosphates and Walker B residue side chains of D328 and E329. Sensor 1 residue N374 is oriented towards the γ-phosphate of ATP. In the hydrolyzing condition, Mg^{2+} appears in the active site, tightly coordinated by the β- and γ-phosphate moieties of ATP and the side chain of Walker A motif T267. Following hydrolysis, ADP, P_i, and Mg^{2+} remain; the side chains of Walker B residue E329 and Sensor 1 residue N374 tightly coordinate the P_i as it disassociates. Subsequent P_i release down-ring leads to arginine finger disassociation, disrupting the interprotomer interface. Active sites shown are derived (from left to right) from D1 class protomers #4-E, #16-E, #16-D, and #16-B.

C. Alignment of 130 D1 protomers from 23 models derived from all four datasets. Structural alignment was performed on the D1 large subdomains. Only Cα atoms of nucleotide-bound protomers and corresponding Pβ atoms are shown for clarity. The inset (right) shows a key

region of the protein that shows changes in the N-D1 linker and N-terminus of the D1 domain

and the Walker A helix. The presence of the latch loop contributed by the up-ring protomer also

influences the conformational state of this region. In contrast, the loop is omitted for clarity, and

the binding location is indicated with a star.

D. Alignment of tyrosine residues in the pore of the D1 ring of substrate-bound models reveals

protomer F as the site of nucleotide exchange and substrate re-engagement. ADP-bound F is

disengaged from the substrate (blue). The release of ADP (nucleotide-free, grey) and ATP-

binding (red) are accompanied by a continuous transition to substrate engagement. The lowest

tyrosine corresponding to an ATP-bound F shown (red triangle) is from the 3DVA-derived model

shown in **A** (middle column, bottom).

Figure 6 | Principal component analysis of NSF D1 protomer coordinates reveals further details related to sequential hydrolysis

A. 132 D1 domain models were aligned by large subdomain (residues 210–392), PCA was

performed on Cα coordinates, and HDBSCAN was used to cluster the models in 11-dimensional

space with a minimum cluster size of 3 models. Protomers cluster into discrete groups based on

conformation; these are easily distinguished upon projection into a 2D space defined by the first

and fifth modes. A proposed hydrolysis cycle connects these clusters as indicated by arrows

between clusters and descriptions below. Superimposed models corresponding to these

transitions are shown on the right. The cluster corresponding to pre-hydrolysis protomers E is

circled with a dashed line.

B. Protomer nucleotide state, mapped to the 2D PCA coordinate from **A**. The ATP + Mg^{2+} state is

unique to protomers E (top left, dashed line).

C. The angle (º) between helices α_3 and α_8 (a proxy for the relative angle between the D1 large

and small subdomains). Protomers E have the largest angle due (top left, dashed line).

D. A mutant form of NSF, NSF(Δlatch), was generated in which part of the latch loop (residues

460–466) was replaced with a short glycine-serine linker (SGS) and assayed for its ability to

disassemble the ternary SNARE complex composed of syntaxin, SNAP-25, and synaptobrevin,

relative to wild-type NSF as described previously[6]. Traces show the mean ± SEM of eight

replicates for each form of NSF, with and without adding $MgCl_2$ to initiate the reaction.

Disassembly rates were calculated by linear regression to each trace in the linear region of the

time course, from ~95–350 s, depending on the trace. NSF(Δlatch) disassembled SNARE

complex at 28 ± 1% the rate of wild type.

E. Three conformational features that distinguish the pre-hydrolysis protomer E cluster (top left,

dashed line), mapped to the 2D PCA coordinate from **A**. These features are in contact and

structurally coupled. The position of nucleotide in the active site is modulated by the position of

the Walker A helix (α_3), which occupies three discrete states; pre-hydrolysis protomers E occupy

the "down" state (left). The position of the helix is allosterically modulated by at least two

inputs. First, the N-D1 linker conformation (center) is connected to the state of N-domain

binding to α-SNAP, and it interacts with the latch loop of the up-ring protomer upon rebinding.

Second, the latch loop of the up-ring protomer contacts this region in *trans* and is also

structurally coupled (right). These two features push against the C-terminal end of the Walker A

helix and ultimately modulate the position of the ATP γ-phosphate relative to key catalytic

residues.

Figure 7 | An expanded model for NSF action across cellular and atomic scales

A. A model for disassembling non-fusogenic SNARE species by NSF and α-SNAP at the

presynapse and progression to priming and fusion. Syntaxin nanodomains comprised at least in

part of tetrameric syntaxin SNARE complexes serve as a reservoir for the liberation of

individual, fusogenic syntaxin molecules by NSF upon disassembly. These syntaxin molecules

may also enter non-productive binary complexes with SNAP-25, with 2:1 or 2:2 stoichiometry;

such complexes must also be disassembled by NSF to ensure *trans*-SNARE complex formation

with synaptobrevin on incoming synaptic vesicles in concert with other factors, such as Munc18,

Munc13, complexin, and synaptotagmin. Where possible, full-length structures (structures

reported here and PDB IDs 1N7S, 4JEU, 5W5C) were supplemented with AlphaFold2[60]

structures to provide additional context, and ISOLDE[61] was used with secondary structure

restraints to reposition flexible regions.

B. A sequential hydrolysis model for SNARE complex disassembly by NSF is inferred from

comparing different classes from non-hydrolyzing and hydrolyzing conditions. Following

hydrolysis-independent substrate side loading, the D1 ring is found in a pre-hydrolysis state with

ATP in protomer E of D1. Protomer F nucleotide exchange then enables ATP exchange and

rebinding to both SNARE substrate and the formation of the interprotomer interface with

protomer E, triggering ATP hydrolysis in protomer E and substrate translocation. This is

accompanied by P_i and Mg^{2+} release from down-ring protomers; in many structures, the release

occurs between protomers C and B. The cycle then begins anew. Blue corresponds to ADP-

bound, red to ATP-bound, purple to ADP•P_i bound, and grey to nucleotide-free. See Movie S1 for

an animated rendition.

Methods

PC12 transfection and imaging

The PC12 cells, generously provided by Prof. Tobias Meyer from Stanford University, were

cultured on glass-bottomed dishes (MatTek; P35G-1.5-14-C) and allowed to adhere for 24 hr. To

introduce genetic material into the cells, transfection was performed using the TurboFect™

Transfection Reagent (#R0532, Thermo Fisher Scientific, USA) according to the manufacturer's

instructions. In brief, the plasmids (NSF-mScarlet-I plasmid: 1.75 μg; GFP-Syntaxin plasmid:

0.5 µg) were combined with 6.75 µL of TurboFect in 225 µL of DMEM medium (without FBS)

and incubated for 20 min. The DNA-TurboFect mixtures were carefully added drop-wise to the

cell cultures and incubated for 6 hr. Following the incubation, the cell cultures were replaced

with complete culture medium. After 24 hr post-transfection, the cells underwent three washes

with PBS. Subsequently, the cells were imaged using a structured illumination Nikon N-SIM

system (version AR5.11.00 64bit, Tokyo, Japan) equipped with an Apochromat 100×/1.49

numerical aperture oil-immersion objective lens. Using a 488 nm laser, the samples were excited

to acquire green SIM images, which captured emission within the 500–550 nm bandwidth,

specifically for GFP. Additionally, red SIM images were obtained by exciting the samples with a

561 nm laser to capture emission within the 570–640 nm bandwidth, corresponding to mScarlet-

I. To determine the degree of colocalization between syntaxin and NSF, Pearson's correlation

coefficient was calculated between channels *ImageJ*[62].

Yeast imaging

Super-resolution microscopy of *S. cerevisiae* expressing GFP-tagged Sso1 was performed using

structured illumination microscopy (SIM) on the OMX-Blaze system (Applied Precision,

Issaquah, WA). Glass-bottom dishes (1.5H, Cellvis) were pretreated with 0.01% poly-l-lysine by

adding 600–1000 µL to the center and incubated overnight. Following incubation, the dishes

were washed with deionized water to remove excess poly-l-lysine. Yeast cultures were grown

overnight, washed twice with phosphate-buffered saline (PBS), and resuspended to an optical

density (OD) of 0.4. A 10–20 µL volume of the yeast suspension was deposited onto the treated

dishes, followed by the addition of Diamond Antifade (Thermo Fisher Scientific) to prevent

photobleaching during imaging. The glass-bottom dishes were placed in a metal holder with 1–2

drops of immersion oil (refractive index 1.512) added before placement on the microscope stage.

SIM imaging was conducted using a laser appropriate for GFP excitation, and images were

captured with a 100× objective for a *z*-slice thickness of 0.125 μm, spanning a total thickness of

2.6 μm. Widefield images were also collected at 0.2 μm slice thickness when necessary.

Protein expression and purification

All NSF, the wild-type and mutant NSF-mScarlet-I, and NSF(Δ460–466) proteins were derived

from NSF from the Chinese hamster *C. griseus*, expressed in BL21(DE3)-RIL *E. coli*, and

purified as described previously[6], with final reassembly of hexameric NSF in reassembly buffer

(50 mM Tris pH 8.0, 150 mM NaCl, 1 mM EDTA, 1 mM ATP, and 1 mM TCEP). α-SNAP from

rat *R. norvegicus* was expressed in *E. coli* and purified as described previously[63]. Soluble rat

neuronal SNARE complex for the fluorescence-based disassembly assay (wild-type SNAP-25A,

syntaxin-1A 1–265, and 6×His-synaptobrevin-2 1–96) was purified by co-expression in C41 *E.*

coli and purified as described previously[63] Purified soluble rat neuronal SNARE complex was

fluorescently labeled with Oregon Green 488 maleimide dye as described previously[6].

Syntaxin-1A fragment 2-253 (Stx[2-253]) was N-terminal His$_6$-tagged using the plasmid pTEV5.

Overexpression in *E. coli* BL21(DE3) was carried out in 8 L of auto-inducible media at 37ºC for

4 hr and then 30ºC overnight. Cells were collected by centrifugation; half of the pellet was

resuspended into 200 mL of 50 mM TRIS pH 8, 500 mM NaCl, and 10 mM imidazole and

supplemented with DNAse, lysozyme, 1 mM MgCl$_2$, and 2 eComplete Protease Inhibitor

Cocktail tablets (Roche). Cells were lysed by sonication (1 s on/3 s off, total 20 min, 55%

amplitude, 4ºC). Debris were removed by centrifugation at 43,000 rpm for 30 min at 4ºC using a

Ti45 rotor. The supernatant was then mixed with 5 mL of nickel-NTA agarose resin (Qiagen)

equilibrated with the lysis buffer. The suspension was gently stirred for 1 hr at 4ºC. The resin was

subsequently recovered by centrifugation at 4,200 rpm for 10 min at 4ºC, poured into a column,

and attached to an AKTA Start (Cytiva). An extensive wash was performed using 20 CV of 50

mM TRIS pH 8, 500 mM NaCl, and 40 mM imidazole until the 280 nm signal was stable.

Protein was eluted in 1 mL fractions in 50 mM TRIS pH8, 200 mM NaCl, and 400 mM

imidazole. Syntaxin-enriched fractions were identified by SDS-PAGE and pooled. To remove the

His-tag, the protein was mixed with an excess of TEV protease and dialyzed against 20 mM

HEPES pH 7.5, 50 mM NaCl, 1 mM DTT, overnight at 4ºC. The sample was centrifuged at

4,000 rpm for 10 min at 4ºC, filtered, and loaded onto a Mono-Q 5/50 column attached to an

AKTA Pure (Cytiva). The resin was extensively washed with 20 mM HEPES pH 7.5, 50 mM

NaCl, 0.2 mM EDTA until 280 nm signal was stable. Elution was performed in 120 CV over 2 hr

from 0.050–1 M NaCl. 2 mL fractions within the major peaks were run on SDS-PAGE, and

fractions containing Stx^{2-253} were concentrated using a 10 kDa cutoff centrifugal filtration device.

The concentrate was filtered and further purified using a Superdex Increase 75 16/60 column

(Cytiva) equilibrated with 20 mM TRIS pH 8, 150 mM NaCl. Fractions were run on an SDS-

PAGE, and fractions containing Stx^{2-253} were concentrated using a 10 kDa MWCO centrifugal

filtration device. A final concentration of 745 mM was determined by using the extinction

coefficient 5,960 M^{-1} cm^{-1}. Stx^{2-253} was aliquoted, flash frozen, and stored at -80ºC until further

use. An independent sample purified from the same batch of cells was analyzed on MALDI-TOF,

and a monomeric molecular weight of 29,208.50 g/mol was confirmed.

Binary neuronal SNARE complex for cryo-EM imaging studies (SNAP-25A with all cysteines

mutated to serine, syntaxin-1A 1–267) was purified following co-expression of SNARE proteins

from *E. coli* BL21 STAR (Invitrogen). Cells were grown overnight at 30°C using an auto-

inducing LB medium[64]. After harvesting the cells by centrifugation, the pellet was resuspended

in lysis buffer (50 mM NaPi pH 8.0, 300 mM NaCl, 20 mM imidazole, 0.5 mM TCEP)

supplemented with lysozyme, DNase I, and antiprotease tablets (Roche) and subjected to

sonication (3 s on, 9 s off, 60% power for 5 min, repeated 3×) and centrifugation (10,000 rpm for

10 min followed by 43,000 rpm for 40 min). The clarified lysate was bound to Ni-

NTA agarose beads (QIAGEN) equilibrated in a lysis buffer. Beads were harvested by

centrifugation (2200 rpm for 10 min), poured into a gravity column, and washed with lysis

buffer, urea buffer (50 mM NaP$_i$ pH 8.0, 300 mM NaCl, 30 mM imidazole, 0.5 mM TCEP, 7.5 M

urea), and wash buffer (50 mM NaPi pH 8.0, 300 mM NaCl, 30 mM imidazole, 0.5 mM TCEP).

The SNARE complex was then eluted with wash buffer supplemented with an additional

320 mM imidazole. The fresh eluent of the Ni-NTA affinity purified SNARE complex was

supplemented with tobacco etch virus (TEV) protease and dialyzed against buffer A1 (50 mM

Tris-HCl, pH 8.0, 250 mM NaCl, 0.5mM TCEP) overnight at 4°C. After dialysis, the sample was

diluted to a final NaCl concentration of 50 mM using buffer A1 without NaCl, and the His-tag-

free complex was subjected to anion exchange chromatography (Buffer A2: 50 mM Tris-HCl, pH

8.0, 50 mM NaCl, 0.5 mM TCEP; Buffer B1: 50 mM Tris-HCl, pH 8.0, 500 mM NaCl, 0.5 mM

TCEP) using a MonoQ 4.6/100 column (Cytiva) and a linear gradient of NaCl starting at 50 mM

and ending at 500 mM. The peaks containing the purified complex were pooled and loaded onto

a Superdex 75 16/60 column to normalize NaCl concentration (50 mM Tris-HCl pH 8.0, 50 mM

NaCl, 0.5 mM TCEP).

The sx20S supercomplex was formed by mixing ATP-bound NSF, α-SNAP, and syntaxin to a

ratio of 1:5:30 in reassembly buffer and purifying it by size-exclusion chromatography using a

Superose 6 10/300 GL column; the first peak to elute was collected and concentrated to around

15 mg mL^{-1} for vitrification and cryo-EM imaging. The 21bin20S supercomplex was purified

similarly but with a 1:5:30 ratio of NSF:binary SNARE complex:α-SNAP. In the case of

preparing 21bin20S supercomplex for imaging under hydrolyzing conditions, EDTA was omitted

from the final SEC buffer.

Gel-based syntaxin disassembly assay

Soluble syntaxin (2–253) and α-SNAP were mixed at concentrations of 100 μM and 150 μM,

respectively, in 50 mM Tris pH 8.0, 150 mM NaCl, 1 mM TCEP, and 1 mM EDTA (SEC buffer)

and incubated for 20 min. NSF was subsequently added to a final concentration of 2.5 μM,

bringing the total reaction volume to 500 μL. The mixture was then injected into a Superdex 200

16/60 column (Cytiva) and complex peak fractions were pooled and concentrated using a

100,000 MWCO filtration device. For each assay condition, 5 μg of complex was used. Two

experimental conditions were performed in triplicate; either EDTA or MgCl$_2$ was added to a final

concentration of 4 mM. Samples were incubated for 20 min at room temperature and re-

concentrated using 100k MWCO filters before SDS-PAGE. Re-concentration effectively de-

enriches liberated syntaxin and α-SNAP molecules. Densitometry was performed using

ImageJ[62], and statistical significance was established between the conditions using a two-tailed

Student's *t*-test.

Gel-based ternary SNARE complex disassembly assay

To verify that NSF activity was not hampered by the C-terminal addition of mScarlet-I, SNARE

complex disassembly was evaluated by gel electrophoresis. NSF, NSF-mScarlet-I, α-SNAP, and

the neuronal SNARE complex were dialyzed into 25 mM TRIS pH 8, 150 mM NaCl, 1 mM ATP.

Then, components were combined into either EDTA buffer (25 mM TRIS pH 8, 2 mM ATP, 1

mM EDTA) or Mg^{2+} buffer (25 mM TRIS pH 8, 2 mM ATP, 4 mM MgCl$_2$). The reaction volume

was 10 μL, and contained 500 nM NSF or NSF-mScarlet-I and 50 nM α-SNAP. Assays were

started by adding 10 μM SNARE complex, and samples were then incubated at 37ºC for 10 min

in a thermocycler. 2 uL of 5×-Laemli loading buffer was then added, and samples were

incubated at 95ºC for 5 min. Finally, 5 µL of each reaction was loaded onto an AnykD TGX gel

(BioRad) and run at 200 V for 45 min. The gel was stained with Coomassie R-250 (BioRad). As

internal controls, all single components were diluted at a similar concentration as the reactions

using Mg^{2+} buffer and treated in parallel. Note that the additional control marked as "SNARE

complex unboiled" corresponds to the SNARE complex incubated at 37ºC but not at 95ºC after

adding loading buffer. For reference, monomeric NSF is ~83 kDa while NSF-mScarlet-I is ~112

kDa.

Fluorescence dequenching-based neuronal SNARE complex disassembly assay

The details of the fluorescence dequenching assay have been previously described[6,32,38,40,41,63].

Fluorescence dequenching assays were carried out using a FlexStation II 384-well plate reader

(Molecular Devices) with a final reaction volume of 60 µL in 20 mM Tris pH 8, 100 mM NaCl, 2

mM ATP, and 0.5 mM TCEP. The assay plate was first prepared by adding 50 µL of 60 nM wild-

type or Δ460–466 mutant NSF, 2.4 µM α-SNAP, 480 nM OG-labeled neuronal SNARE complex

in reaction buffer to each well. Then, 100 µL of reaction buffer (as a control) or ATP hydrolysis

initiation buffer (reaction buffer supplemented with 24 µM $MgCl_2$) was added to appropriate

wells of the compound plate, and fluorescence dequenching was monitored. Excitation was

performed at 485 nm, and emission was monitored at 525 nm with a dichroic mirror at 515 nm.

First, fluorescence intensity was monitored before reaction initiation for 60 s. Then, 10 µL of

compound plate solution was transferred to each assay plate well (diluting all protein 5/6-fold

and $MgCl_2$ concentrations six-fold) and triturated, initiating the disassembly reaction. The final

protein concentrations following mixing were 50 nM NSF, 2.0 µM α-SNAP, and 400 nM

SNARE complex. The final $MgCl_2$ concentration was 4 mM. Dequenching was then monitored

for 30 min at 20 ºC. Linear regression over the first several minutes was performed to estimate

relative disassembly rates. The statistical significance of changes to disassembly rates was

assessed using the two-tailed Student's *t*-test.

Sample vitrification for cryo-EM SPA

The samples were prepared immediately before vitrification (Fig. S2). Quantifoil R1.2/1.3 200

mesh gold grids (Quantifoil Micro Tools GmbH, Germany) were cleaned with chloroform, dried

for several hours, and subjected to glow discharge using a PELCO easiGlow Glow Discharge

Cleaning System (Ted Pella) under vacuum at 15 mA for 45 s with a 15 s hold time. Immediately

following 21bin20S or sx20S supercomplex purification, proteins were concentrated to ~15 mg

mL^{-1} in 17 µL. For the hydrolyzing condition, 2 µL of 10 × ATP Regeneration Solution (50 mM

Tris pH 8.0, 150 mM NaCl, 1 mM ATP, 160 mM creatine phosphate, and 0.3 mg/mL rabbit

muscle creatine kinase (Roche)) was added, and the sample was incubated on ice for 2 min; in

contrast, 2 µL of reassembly buffer was added in the case of non-hydrolyzing conditions. Then,

for the hydrolyzing condition, 1 µL of 20 × Reaction Initiation Solution (50 mM Tris pH 8.0, 150

mM NaCl, 1 mM ATP, 20 mM MgCl2, 1% Nonidet P-40) was added, otherwise 1 µL of 1%

Nonidet P-40 was added in the case of non-hydrolyzing conditions. Using a Vitrobot Mark IV

(Thermo Fisher Scientific), 4 µL of a given sample was applied to a grid at room temperature

and 100% humidity and, blotted for 4 s with a blot force of 3, and subsequently plunge-frozen in

liquid ethane.

Cryo-EM SPA data collection

Grids were imaged using a Titan Krios electron microscope (Thermo Fisher Scientific) operated

at 300 kV and equipped with a K3 camera (Gatan). Data collection was performed using

SerialEM[65]. All datasets were collected with a nominal magnification of 22,500× for a physical

pixel size of 1.096 Å and a super-resolution pixel size of 0.548 Å. Dose rate was either ~10 or

~20 e$^-$/px/s depending on whether correlated double sampling (CDS) was enabled on the K3, for

a total dose of around 30 e$^-$/Å^2. Nine grid holes were imaged for each stage movement using

multiple record setup in *SerialEM*, with one collection per hole. At each hole, either a 2 s or 4 s

exposure fractionated into either 40 or 80 frames was collected, depending on whether CDS was

enabled. Defocus values ranged from -1–-3 μm. See Table S1 for details.

Cryo-EM SPA image processing

All cryo-EM datasets were processed using a software pipeline consisting of *MotionCorr2*[66],

CTFFIND4[67], *RELION 3*[68,69], and *cryoSPARC*[51,70]. Early-stage processing was carried out in

RELION 3, with final 3D refinement and variability analysis carried out with *CryoSPARC*.

Dataset-specific processing schemes are presented in Figs. S3–7. Generally, all processing

proceeded as follows. Processing began in *RELION 3*, with per-micrograph optics groups

defined for a given dataset based on the nine beam shift positions using the custom script

update_relion_tilt_class_og.py. For example, the hydrolyzing condition micrographs were

collected in two sessions, so 18 total optics groups were defined. The raw image stacks were

dose-weighted and motion-corrected using the *RELION* implementation of *MotionCorr2*, with

seven patches along the *x* direction and five patches along *y*. CTF parameters were estimated for

each micrograph using *CTFFIND4* with a resolution range of 3–30 Å, an FFT box size of 1024

px, and otherwise default parameters. Micrographs were then selected based on maximum

resolution and the CTF figure of merit.

Generally, an initial round of manual picking, extraction, and 2D classification were used to

assess data set quality and provide particles to reconstruct an initial 3D model. Good particles

from manual picking were then used for automated particle picking in *RELION*. Particles were

then extracted with a box size of 640 px, binned 8× (4.384 Å Å pixel size) to a box size of 80 px,

and subjected to at least one round of 2D classification to eliminate obviously bad particles.

Next, multiple rounds of 3D classification with a 280 Å mask diameter and a regularization

parameter $T = 4$ were performed over many iterations to determine the proper number of discrete

states in a given dataset, and to ensure that particles were classified properly. Convergence of

classification was monitored with a custom script, *startle.py*. Several classes slightly larger than

the number of discrete states were used, and classes that appeared to lie along a continuous

conformational coordinate were subsequently re-merged into a single class. In the case of the

hydrolyzing dataset, SNARE substrate-bound and SNARE substrate-free particles were split into

separate groups of particles, re-classified, and processed independently.

Following classification, particles were re-centered and re-extracted with a box size of 588 px

and binned 2× (1.096 Å pixel size) with a box size of 294 px. Following an initial round of 3D

refinement and post-processing, CTF refinement was performed. Typically, defocus,

astigmatism, B-factor, tilt, and trefoil were refined on a per-particle or per-micrograph basis,

depending on the specific class. Particles were then subjected to an additional round of 3D

refinement and post-processing, and optimal parameters for particle polishing were determined

by training on several classes, yielding different parameter sets. Particle polishing was then

performed with these sets, and 3D refinement and post-processing were used to assess the

results. Polished particles were then subjected to a final round of 3D refinement, post-processing,

and CTF refinement, during which anisotropic magnification was estimated. At this stage,

particles and 3D reconstructions were imported into *CryoSPARC* for a final round of non-

uniform refinement, which typically improved the quality of reconstructions.

In the case of one class observed under hydrolyzing conditions (c4-7-16-20-24), a large degree of

continuous heterogeneity was observed. 3D variability analysis (3DVA) revealed a trajectory of

protomer F nucleotide and rebinding, so particles in the first mode were split using a custom

script, *split_by_3dva.py*, as current tools to do so natively were unavailable at the time. This

script fits a normal distribution to the eigenvector magnitude distribution and allows slicing with

different cutoffs. For the 3DVA subclasses (slices O and P), particles in the $\pm1.5\ \sigma$ tails of the

distribution were used for reconstruction.

These reconstructions were considered final for the overall 20S architecture, but further

processing was required to generate local reconstructions of the various α-SNAP—SNARE

subcomplexes. In the case of the sx20S dataset, only one class was observed with ordered spire

density, so local refinement was relatively simple. *CryoSPARC* signal subtraction

(*particle_subtract*; masks generated using *Segger* 2.5.3[71] in *Chimera* 1.15[72]) was used to remove

the signal from NSF N, D1, and D2 domains, leaving only the α-SNAP—syntaxin H3 SNARE

domain tetramer subcomplex. Local refinement (*new_local_refine*) was then performed on these

particles with a 40°/20 Å search extent and a custom fulcrum at the N-terminal end of the

SNARE bundle near the point of interaction with the F protomer of the D1 ring. The resulting

volumes were re-centered on the center of mass (*volume_alignment_tools*), and particle

alignments were updated as well (*align_3D*). Then, a final round of local refinement

(*new_local_refine*) was performed with the map center as the fulcrum (the default setting), a

smaller search extent of 10°/5 Å, and a maximum alignment resolution of 0.1°. Per-particle scales

were also reset, and minimization was performed over a per-particle scale. This produced the

final reconstruction of this subcomplex for subsequent modeling. In the case of the α-SNAP—

2:1 and α-SNAP—2:2 binary SNARE subcomplexes, this procedure was followed for each class

meeting certain criteria (number of particles, ordered spire, resolution), yielding five local

reconstructions for the α-SNAP—2:1 binary SNARE subcomplex and six local reconstructions

for the α-SNAP—2:2 binary SNARE subcomplex. In each case, these reconstructions were re-

centered (*volume_alignment_tools*), aligned to one another with the center of mass at the origin

(*align_3D)*, and a final round of local refinement was run to produce a final higher-resolution

reconstruction incorporating data from multiple classes. This produced the final reconstruction of

these subcomplexes for subsequent modeling. For more details about all data sets, see Figs. S3–

7, Tables S1–2, and Fig. S8.

Model building and refinement

All manual model building and refinements were carried out using *Coot*[73]. Semi-manual

refinement into low-resolution features was performed with *UCSF ChimeraX*[74] and the

ISOLDE[61] plugin. Automated refinement was performed using *phenix.real_space_refine*[75,76].

First, models of the three α-SNAP—SNARE subcomplexes were built by rigid body fitting of

the best possible starting models into their corresponding densities using *UCSF ChimeraX*. In

each case, 3–4 copies of a high-quality α-SNAP model from the previous best 20S model (PDB

ID 6MDM[6]) were rigid body fit into the density. In the case of the syntaxin H3 domain tetramer

complex, four copies of the syntaxin chain from the high-resolution crystal structure (PDB ID

1N7S[49]) of the neuronal SNARE complex were docked into the density, with several

characteristic residues as fiducials. The quality of the local reconstructions of the SNARE—α-

SNAP spires allowed unique assignment of the identity of the SNARE domains, their

directionality, and register (Figs. 2D, 3D, 4D, Fig. S7DE, Fig. S9A). In the case of the 2:1 binary

SNARE complex, the syntaxin and SNAP-25 components of the same crystal structure were

docked into the density, but with the SNARE domain of synaptobrevin replaced with an

additional syntaxin H3 SNARE domain; key side chain densities as well as a distinctive kink

leading into the SNAP-25 SN2 domain guided placement (Fig. 3C, Fig. S9A). In the case of the

2:2 binary SNARE complex, a previous crystal structure (PDB ID 1JTH[34]) was docked into the

density without modification. In all cases, individual components were trimmed, or residues

were added where supported by the density. A round of *phenix.real_space_refine* was then

performed with only rigid-body refinement enabled for each component. Subsequent cycles of

manual model building were performed in *Coot* and *ISOLDE* followed by 1–7 macrocycles of

phenix.real_space_refine with global minimization and local grid search enabled. Ramachandran

and secondary structure restraints were used throughout.

Models of NSF or the different 20S supercomplexes were then built. Gold-standard models were

first built into two high-resolution NSF classes from a single non-hydrolyzing dataset with near-

atomic resolutions using the previous best model of NSF (PDB ID 6MDM[6]) as a starting model.

The starting model was first docked into cryo-EM density using ChimeraX. Next, a round of

phenix.real_space_refine was performed with only rigid-body refinement enabled; each large

and small subdomain of each protomer was treated as a single rigid body (twelve total).

Nucleotides and water molecules were then added where appropriate. Subsequent cycles of

manual model building were performed in *Coot* and *ISOLDE* followed by 1–7 macrocycles of

phenix.real_space_refine with global minimization and local grid search enabled. Ramachandran

and secondary structure restraints were used throughout.

The previously described models were then used as starting models in building the sx20S and

bin20S supercomplexes. In sx20S and bin20S reconstructions with well-defined N-domains (i.e.,

a recognizable two-lobed density), one or more instances of the re-processed model derived from

the crystal structure (PDB ID 1QCS[77]) as re-processed by *PDB_REDO*[78] were rigid body fit into

density before refinement. Next, the appropriate α-SNAP—SNARE subcomplex was docked

into the density, completing the spire structure. A round of *phenix.real_space_refine* was

performed with only rigid-body refinement enabled; each large and small subdomain of each

protomer, each N domain, and each α-SNAP—SNARE subcomplex was treated as a rigid body

(for a 20S particle with four engaged N-domains, this would be 17 total rigid body groups).

Nucleotides, Mg^{2+}, P_i, and water molecules were then added where appropriate. Subsequent

cycles of manual model building were performed in *Coot* and *ISOLDE* followed by 1–7

macrocycles of *phenix.real_space_refine* with global minimization and local grid search enabled.

Ramachandran and secondary structure restraints were used throughout.

All model figures were prepared with *UCSF ChimeraX*, in many cases with custom CXC scripts

generated by the Python scripts described below.

Automated NSF model analysis and coordinate PCA

Several custom Python scripts (*preprocess_models.py, analyze_models.py*) using the *Bio.PDB*

module[79] of the Biopython project[80] as well as *pandas*[81], *numpy*[82], *scipy*[83], *scikit-learn*[84], and

matplotlib[85] were used to standardize all 25 NSF, sx20S, or bin20S models for further analysis,

such as to perform principle components analysis (PCA) and HDBSCAN clustering, to measure

NSF D1 large-small subdomain angles, to perform planarity fits for NSF domains and

subdomains, to measure distances between key catalytic residues and nucleotides, and to map

various values to structures for visualization in *UCSF ChimeraX*.

For PCA, 132 NSF D1 domains from 22 models over all datasets were split into large and small

subdomains, which were then aligned to one another (large, residues 224–373; small, residues

400–486) using *Bio.PDB.Superimposer*. Splitting removes the largest source of variance in the

data, the rigid body motions of the two subdomains relative to each other, which is considered

trivial for the purposes of this analysis. Next, a Cα coordinate matrix $\mathbf{C}$ with dimensions $n_{models} \times$

$n_{residues} \times 3$ was constructed for residues shared by all models and flattened into two dimensions,

creating $\mathbf{C_{flat}}$ with n_{models} rows and $n_{residues} \times 3$ columns. $\mathbf{C_{flat}}$ was then rescaled using

sklearn.preprocessing.StandardScaler and PCA was performed on $\mathbf{C_{flat}}$ using

sklearn.decomposition.PCA. 13 components were found to contribute 99% of the variance, so a

transformed matrix $\mathbf{C_{trans}}$ with dimensions $n_{models} \times 13$, with eigenvectors sorted along the second

axis by eigenvalue, was calculated. The cluster analysis of models was then performed using

sklearn.cluster.HDBSCAN with a minimum cluster size of three, producing 11 clusters

accounting for 119/132 total models. This analysis serves as the basis for Fig. 6A.

For NSF D1 and D2 plane fits, centers of mass for all subdomains were calculated, and the

planes for different combinations of these points were fit using a least squares approach (Fig.

S11). For NSF D1 large-small subdomain angle calculations, the primary axes of two α-helices

in the large (residues 350–358) and small subdomains (residues 442–452) were calculated by

singular value decomposition using *np.linalg.svd*, with implementation inspired by

helix_analysis.py from the MDAnalysis package[86]. Vectors corresponding to these helical axes u

and v were normalized and centered at the coordinate origin, and the angle between them was

calculated as $\theta = 2 \times \tan^{-1}\left(\sqrt{\sum_i \vec{u}_i^2}\ \sqrt{\sum_i \vec{v}_i^2}\right)$.

Distances between key active site atoms over all models (Fig. S13) were calculated using

np.linalg.norm, where the norm of the difference vector of the coordinates of each atom is the

distance between them in Å. Significant changes in protomer E active site residues relative to the

other five protomers were assessed by pairwise application of the Welch's t-test assuming non-

equal variances as implemented in *scipy.stats.ttest_ind*. Distance distributions were considered

statistically different if $p < 0.05$.

All these values, as well as manually curated, qualitative data labels, were plotted in various

ways using *matplotlib* or incorporated into automatically generated UCSF ChimeraX CXC

scripts.

References

1. Rizo, J., and Südhof, T.C. (2012). The membrane fusion enigma: SNAREs, Sec1/Munc18
proteins, and their accomplices--guilty as charged? Annu. Rev. Cell Dev. Biol. *28*, 279–308.
https://doi.org/10.1146/annurev-cellbio-101011-155818.

2. Brunger, A.T., Choi, U.B., Lai, Y., Leitz, J., and Zhou, Q. (2018). Molecular Mechanisms of
Fast Neurotransmitter Release. Annu. Rev. Biophys. *47*, 469–497.
https://doi.org/10.1146/annurev-biophys-070816-034117.

3. Leitz, J., Wang, C., Esquivies, L., Pfuetzner, R.A., Peters, J.J., Couoh-Cardel, S., Wang, A.L.,
and Brunger, A.T. (2024). Beyond the MUN domain, Munc13 controls priming and
depriming of synaptic vesicles. Cell Rep. *43*. https://doi.org/10.1016/j.celrep.2024.114026.

4. Brunger, A.T., Choi, U.B., Lai, Y., Leitz, J., White, K.I., and Zhou, Q. (2019). The pre-
synaptic fusion machinery. Curr. Opin. Struct. Biol. *54*, 179–188.
https://doi.org/10.1016/j.sbi.2019.03.007.

5. Rizo, J. (2022). Molecular Mechanisms Underlying Neurotransmitter Release. Annu. Rev.
Biophys. *51*, 377–408. https://doi.org/10.1146/annurev-biophys-111821-104732.

6. White, K.I., Zhao, M., Choi, U.B., Pfuetzner, R.A., and Brunger, A.T. (2018). Structural
principles of SNARE complex recognition by the AAA+ protein NSF. eLife *7*, e38888.
https://doi.org/10.7554/eLife.38888.

7. Zhao, M., Wu, S., Zhou, Q., Vivona, S., Cipriano, D.J., Cheng, Y., and Brunger, A.T. (2015).
Mechanistic insights into the recycling machine of the SNARE complex. Nature *518*, 61–67.
https://doi.org/10.1038/nature14148.

8. Rickman, C., Medine, C.N., Dun, A.R., Moulton, D.J., Mandula, O., Halemani, N.D.,
Rizzoli, S.O., Chamberlain, L.H., and Duncan, R.R. (2010). t-SNARE Protein
Conformations Patterned by the Lipid Microenvironment. J. Biol. Chem. *285*, 13535–13541.
https://doi.org/10.1074/jbc.M109.091058.

9. Murray, D.H., and Tamm, L.K. (2009). Clustering of Syntaxin-1A in Model Membranes is
Modulated by Phosphatidylinositol-4,5-bisphosphate and Cholesterol. Biochemistry *48*,
4617–4625. https://doi.org/10.1021/bi9003217.

10. Aoyagi, K., Sugaya, T., Umeda, M., Yamamoto, S., Terakawa, S., and Takahashi, M. (2005).
The activation of exocytotic sites by the formation of phosphatidylinositol 4,5-bisphosphate
microdomains at syntaxin clusters. J. Biol. Chem. *280*, 17346–17352.
https://doi.org/10.1074/jbc.M413307200.

11. Lang, T., Bruns, D., Wenzel, D., Riedel, D., Holroyd, P., Thiele, C., and Jahn, R. (2001).
SNAREs are concentrated in cholesterol-dependent clusters that define docking and fusion
sites for exocytosis. EMBO J. *20*, 2202–2213. https://doi.org/10.1093/emboj/20.9.2202.

12. Emperador-Melero, J., and Kaeser, P.S. (2020). Assembly of the presynaptic active zone.
Curr. Opin. Neurobiol. *63*, 95–103. https://doi.org/10.1016/j.conb.2020.03.008.

13. Weimbs, T., Low, S.H., Chapin, S.J., Mostov, K.E., Bucher, P., and Hofmann, K. (1997). A
conserved domain is present in different families of vesicular fusion proteins:
A new superfamily. Proc. Natl. Acad. Sci. U. S. A. *94*, 3046–3051.

14. Bennett, M.K., Calakos, N., and Scheller, R.H. (1992). Syntaxin: A Synaptic Protein
Implicated in Docking of Synaptic Vesicles at Presynaptic Active Zones. Science *257*, 255–
259. https://doi.org/10.1126/science.1321498.

15. Sutton, R.B., Fasshauer, D., Jahn, R., and Brunger, A.T. (1998). Crystal structure of a
SNARE complex involved in synaptic exocytosis at 2.4 Å resolution. Nature *395*, 347–353.
https://doi.org/10.1038/26412.

16. Fernandez, I., Ubach, J., Dulubova, I., Zhang, X., Südhof, T.C., and Rizo, J. (1998). Three-
Dimensional Structure of an Evolutionarily Conserved N-Terminal Domain of Syntaxin 1A.
Cell *94*, 841–849. https://doi.org/10.1016/s0092-8674(00)81742-0.

17. Sieber, J.J., Willig, K.I., Kutzner, C., Gerding-Reimers, C., Harke, B., Donnert, G.,
Rammner, B., Eggeling, C., Hell, S.W., Grubmüller, H., et al. (2007). Anatomy and
Dynamics of a Supramolecular Membrane Protein Cluster. Science *317*, 1072–1076.
https://doi.org/10.1126/science.1141727.

18. Bar-On, D., Wolter, S., Van De Linde, S., Heilemann, M., Nudelman, G., Nachliel, E.,
Gutman, M., Sauer, M., and Ashery, U. (2012). Super-resolution Imaging Reveals the
Internal Architecture of Nano-sized Syntaxin Clusters. J. Biol. Chem. *287*, 27158–27167.
https://doi.org/10.1074/jbc.M112.353250.

19. Knowles, M.K., Barg, S., Wan, L., Midorikawa, M., Chen, X., and Almers, W. (2010). Single
secretory granules of live cells recruit syntaxin-1 and synaptosomal associated protein 25
(SNAP-25) in large copy numbers. Proc. Natl. Acad. Sci. *107*, 20810–20815.
https://doi.org/10.1073/pnas.1014840107.

20. Wan, L., Chen, X., and Almers, W. (2021). Munc18-dependent and -independent clustering
of syntaxin in the plasma membrane of cultured endocrine cells. Proc. Natl. Acad. Sci. *118*,
e2025748118. https://doi.org/10.1073/pnas.2025748118.

21. Ullrich, A., Böhme, M.A., Schöneberg, J., Depner, H., Sigrist, S.J., and Noé, F. (2015).
Dynamical Organization of Syntaxin-1A at the Presynaptic Active Zone. PLOS Comput.
Biol. *11*, e1004407. https://doi.org/10.1371/journal.pcbi.1004407.

22. Barg, S., Knowles, M.K., Chen, X., Midorikawa, M., and Almers, W. (2010). Syntaxin
clusters assemble reversibly at sites of secretory granules in live cells. Proc. Natl. Acad. Sci.
107, 20804–20809. https://doi.org/10.1073/pnas.1014823107.

23. Sakamoto, H., Ariyoshi, T., Kimpara, N., Sugao, K., Taiko, I., Takikawa, K., Asanuma, D.,
Namiki, S., and Hirose, K. (2018). Synaptic weight set by Munc13-1 supramolecular
assemblies. Nat. Neurosci. *21*, 41–49. https://doi.org/10.1038/s41593-017-0041-9.

24. Pertsinidis, A., Mukherjee, K., Sharma, M., Pang, Z.P., Park, S.R., Zhang, Y., Brunger, A.T.,
Südhof, T.C., and Chu, S. (2013). Ultrahigh-resolution imaging reveals formation of
neuronal SNARE/Munc18 complexes in situ. Proc. Natl. Acad. Sci. *110*, E2812–E2820.
https://doi.org/10.1073/pnas.1310654110.

25. Yin, P., Gandasi, N.R., Arora, S., Omar-Hmeadi, M., Saras, J., and Barg, S. (2018). Syntaxin
clusters at secretory granules in a munc18-bound conformation. Mol. Biol. Cell *29*, 2700–
2708. https://doi.org/10.1091/mbc.E17-09-0541.

26. Lang, T. (2007). SNARE proteins and 'membrane rafts.' J. Physiol. *585*, 693–698.
https://doi.org/10.1113/jphysiol.2007.134346.

27. Bademosi, A.T., Lauwers, E., Padmanabhan, P., Odierna, L., Chai, Y.J., Papadopulos, A.,
Goodhill, G.J., Verstreken, P., van Swinderen, B., and Meunier, F.A. (2016). In vivo single-
molecule imaging of syntaxin1A reveals polyphosphoinositide- and activity-dependent
trapping in presynaptic nanoclusters. Nat. Commun. *7*, 13660.
https://doi.org/10.1038/ncomms13660.

28. Sieber, J.J., Willig, K.I., Heintzmann, R., Hell, S.W., and Lang, T. (2006). The SNARE Motif
Is Essential for the Formation of Syntaxin Clusters in the Plasma Membrane. Biophys. J. *90*,
2843–2851. https://doi.org/10.1529/biophysj.105.079574.

29. Hanson, P.I., Otto, H., Barton, N., and Jahn, R. (1995). The N-Ethylmaleimide-sensitive
Fusion Protein and α-SNAP Induce a Conformational Change in Syntaxin (*). J. Biol. Chem.
270, 16955–16961. https://doi.org/10.1074/jbc.270.28.16955.

30. Stepien, K.P., Prinslow, E.A., and Rizo, J. (2019). Munc18-1 is crucial to overcome the
inhibition of synaptic vesicle fusion by αSNAP. Nat. Commun. *10*, 4326.
https://doi.org/10.1038/s41467-019-12188-4.

31. Ma, C., Su, L., Seven, A.B., Xu, Y., and Rizo, J. (2013). Reconstitution of the vital functions
of Munc18 and Munc13 in neurotransmitter release. Science *339*, 421–425.
https://doi.org/10.1126/science.1230473.

32. Choi, U.B., Zhao, M., White, K.I., Pfuetzner, R.A., Esquivies, L., Zhou, Q., and Brunger,
A.T. (2018). NSF-mediated disassembly of on- and off-pathway SNARE complexes and
inhibition by complexin. eLife *7*, e36497. https://doi.org/10.7554/eLife.36497.

33. Fasshauer, D., Bruns, D., Shen, B., Jahn, R., and Brunger, A.T. (1997). A Structural Change
Occurs upon Binding of Syntaxin to SNAP-25. J. Biol. Chem. *272*, 4582–4590.
https://doi.org/10.1074/jbc.272.7.4582.

34. Misura, K.M.S., Gonzalez, L.C., May, A.P., Scheller, R.H., and Weis, W.I. (2001). Crystal
Structure and Biophysical Properties of a Complex between the N-terminal SNARE Region
of SNAP25 and Syntaxin 1a *. J. Biol. Chem. *276*, 41301–41309.
https://doi.org/10.1074/jbc.M106853200.

35. Weninger, K., Bowen, M.E., Choi, U.B., Chu, S., and Brunger, A.T. (2008). Accessory
Proteins Stabilize the Acceptor Complex for Synaptobrevin, the 1:1 Syntaxin/SNAP-25
Complex. Structure *16*, 308–320. https://doi.org/10.1016/j.str.2007.12.010.

36. Hesselbarth, J., and Schmidt, C. (2023). Mass spectrometry uncovers intermediates and off-
pathway complexes for SNARE complex assembly. Commun. Biol. *6*, 1–15.
https://doi.org/10.1038/s42003-023-04548-0.

37. Hayashi, T., McMahon, H., Yamasaki, S., Binz, T., Hata, Y., Südhof, T.C., and Niemann, H.
(1994). Synaptic vesicle membrane fusion complex: action of clostridial neurotoxins on
assembly. EMBO J. *13*, 5051–5061. https://doi.org/10.1002/j.1460-2075.1994.tb06834.x.

38. Gao, Y., Khan, Y.A., Mo, W., White, K.I., Perkins, M., Pfuetzner, R.A., Trapani, J.G.,
Brunger, A.T., and Nicolson, T. (2023). Sensory deficit screen identifies nsf mutation that
differentially affects SNARE recycling and quality control. Cell Rep. *42*.
https://doi.org/10.1016/j.celrep.2023.112345.

39. Hayashi, T., Yamasaki, S., Nauenburg, S., Binz, T., and Niemann, H. (1995). Disassembly of
the reconstituted synaptic vesicle membrane fusion complex in vitro. EMBO J. *14*, 2317–
2325. https://doi.org/10.1002/j.1460-2075.1995.tb07226.x.

40. Vivona, S., Cipriano, D.J., O'Leary, S., Li, Y.H., Fenn, T.D., and Brunger, A.T. (2013).
Disassembly of All SNARE Complexes by N-Ethylmaleimide-sensitive Factor (NSF) Is
Initiated by a Conserved 1:1 Interaction between α-Soluble NSF Attachment Protein (SNAP)
and SNARE Complex. J. Biol. Chem. *288*, 24984–24991.
https://doi.org/10.1074/jbc.M113.489807.

41. Zhao, M., Wu, S., Zhou, Q., Vivona, S., Cipriano, D.J., Cheng, Y., and Brunger, A.T. (2015).
Mechanistic insights into the recycling machine of the SNARE complex. Nature *518*, 61–67.
https://doi.org/10.1038/nature14148.

42. Nagiec, E.E., Bernstein, A., and Whiteheart, S.W. (1995). Each Domain of the N-
Ethylmaleimide-sensitive Fusion Protein Contributes to Its Transport Activity (*). J. Biol.
Chem. *270*, 29182–29188. https://doi.org/10.1074/jbc.270.49.29182.

43. Matveeva, E.A., He, P., and Whiteheart, S.W. (1997). N-Ethylmaleimide-sensitive Fusion
Protein Contains High and Low Affinity ATP-binding Sites That Are Functionally Distinct *.
J. Biol. Chem. *272*, 26413–26418. https://doi.org/10.1074/jbc.272.42.26413.

44. Xiao, W., Poirier, M.A., Bennett, M.K., and Shin, Y.-K. (2001). The neuronal t-SNARE
complex is a parallel four-helix bundle. Nat. Struct. Biol. *8*, 308–311.
https://doi.org/10.1038/86174.

45. Margittai, M., Fasshauer, D., Pabst, S., Jahn, R., and Langen, R. (2001). Homo- and
Heterooligomeric SNARE Complexes Studied by Site-directed Spin Labeling *. J. Biol.
Chem. *276*, 13169–13177. https://doi.org/10.1074/jbc.M010653200.

46. Lai, Y., Choi, U.B., Leitz, J., Rhee, H.J., Lee, C., Altas, B., Zhao, M., Pfuetzner, R.A., Wang,
A.L., Brose, N., et al. (2017). Molecular Mechanisms of Synaptic Vesicle Priming by
Munc13 and Munc18. Neuron *95*, 591-607.e10.
https://doi.org/10.1016/j.neuron.2017.07.004.

47. Stepien, K.P., Xu, J., Zhang, X., Bai, X.-C., and Rizo, J. (2022). SNARE assembly
enlightened by cryo-EM structures of a synaptobrevin–Munc18-1–syntaxin-1 complex. Sci.
Adv. *8*, eabo5272. https://doi.org/10.1126/sciadv.abo5272.

48. Sanyal, S., and Krishnan, K.S. (2001). Lethal comatose mutation in Drosophila reveals
possible role for NSF in neurogenesis. NeuroReport *12*, 1363.

49. Ernst, J.A., and Brunger, A.T. (2003). High Resolution Structure, Stability, and
Synaptotagmin Binding of a Truncated Neuronal SNARE Complex *. J. Biol. Chem. *278*,
8630–8636. https://doi.org/10.1074/jbc.M211889200.

50. de la Peña, A.H., Goodall, E.A., Gates, S.N., Lander, G.C., and Martin, A. (2018). Substrate-
engaged 26S proteasome structures reveal mechanisms for ATP-hydrolysis–driven
translocation. Science *362*, eaav0725. https://doi.org/10.1126/science.aav0725.

51. Punjani, A., and Fleet, D.J. (2021). 3D variability analysis: Resolving continuous flexibility
and discrete heterogeneity from single particle cryo-EM. J. Struct. Biol. *213*, 107702.
https://doi.org/10.1016/j.jsb.2021.107702.

52. Khan, Y.A., White, K.I., Pfuetzner, R.A., Singal, B., Esquivies, L., Mckenzie, G., Liu, F.,
DeLong, K., Choi, U.B., Montabana, E., et al. (2024). Sec18 side-loading is essential for
universal SNARE recycling across cellular contexts. Preprint at bioRxiv,
https://doi.org/10.1101/2024.08.30.610324 https://doi.org/10.1101/2024.08.30.610324.

53. Zhou, H., Di Palma, S., Preisinger, C., Peng, M., Polat, A.N., Heck, A.J.R., and Mohammed,
S. (2013). Toward a Comprehensive Characterization of a Human Cancer Cell
Phosphoproteome. J. Proteome Res. *12*, 260–271. https://doi.org/10.1021/pr300630k.

54. Zhou, Q., Zhou, P., Wang, A.L., Wu, D., Zhao, M., Südhof, T.C., and Brunger, A.T. (2017).
The primed SNARE–complexin–synaptotagmin complex for neuronal exocytosis. Nature
548, 420–425. https://doi.org/10.1038/nature23484.

55. Zhou, Q., Lai, Y., Bacaj, T., Zhao, M., Lyubimov, A.Y., Uervirojnangkoorn, M., Zeldin, O.B.,
Brewster, A.S., Sauter, N.K., Cohen, A.E., et al. (2015). Architecture of the synaptotagmin–
SNARE machinery for neuronal exocytosis. Nature *525*, 62–67.
https://doi.org/10.1038/nature14975.

56. Smith, G.R., Contreras-Moreira, B., Zhang, X., and Bates, P.A. (2004). A link between
sequence conservation and domain motion within the AAA+ family. J. Struct. Biol. *146*,
189–204. https://doi.org/10.1016/j.jsb.2003.11.022.

57. Khan, Y.A., White, K.I., and Brunger, A.T. (2022). The AAA+ superfamily: a review of the
structural and mechanistic principles of these molecular machines. Crit. Rev. Biochem. Mol.
Biol. *57*, 156–187. https://doi.org/10.1080/10409238.2021.1979460.

58. Huang, X., Sun, S., Wang, X., Fan, F., Zhou, Q., Lu, S., Cao, Y., Wang, Q.-W., Dong, M.-Q.,
Yao, J., et al. (2019). Mechanistic insights into the SNARE complex disassembly. Sci. Adv.
5, eaau8164. https://doi.org/10.1126/sciadv.aau8164.

59. Greaves, J., Gorleku, O.A., Salaun, C., and Chamberlain, L.H. (2010). Palmitoylation of the
SNAP25 Protein Family. J. Biol. Chem. *285*, 24629–24638.
https://doi.org/10.1074/jbc.M110.119289.

60. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate
protein structure prediction with AlphaFold. Nature *596*, 583–589.
https://doi.org/10.1038/s41586-021-03819-2.

61. Croll, T.I. (2018). ISOLDE: a physically realistic environment for model building into low-
resolution electron-density maps. Acta Crystallogr. Sect. Struct. Biol. *74*, 519–530.
https://doi.org/10.1107/S2059798318002425.

62. Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years
of image analysis. Nat. Methods *9*, 671–675. https://doi.org/10.1038/nmeth.2089.

63. Cipriano, D.J., Jung, J., Vivona, S., Fenn, T.D., Brunger, A.T., and Bryant, Z. (2013).
Processive ATP-driven Substrate Disassembly by the N-Ethylmaleimide-sensitive Factor
(NSF) Molecular Machine♦. J. Biol. Chem. *288*, 23436–23445.
https://doi.org/10.1074/jbc.M113.476705.

64. Studier, F.W. (2005). Protein production by auto-induction in high-density shaking cultures.
Protein Expr. Purif. *41*, 207–234. https://doi.org/10.1016/j.pep.2005.01.016.

65. Mastronarde, D.N. (2005). Automated electron microscope tomography using robust
prediction of specimen movements. J. Struct. Biol. *152*, 36–51.
https://doi.org/10.1016/j.jsb.2005.07.007.

66. Zheng, S.Q., Palovcak, E., Armache, J.-P., Verba, K.A., Cheng, Y., and Agard, D.A. (2017).
MotionCor2 - anisotropic correction of beam-induced motion for improved cryo-electron
microscopy. Nat. Methods *14*, 331–332. https://doi.org/10.1038/nmeth.4193.

67. Rohou, A., and Grigorieff, N. (2015). CTFFIND4: Fast and accurate defocus estimation from
electron micrographs. J. Struct. Biol. *192*, 216–221.
https://doi.org/10.1016/j.jsb.2015.08.008.

68. Scheres, S.H.W. (2012). RELION: Implementation of a Bayesian approach to cryo-EM
structure determination. J. Struct. Biol. *180*, 519–530.
https://doi.org/10.1016/j.jsb.2012.09.006.

69. Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E., and Scheres,
S.H. (2018). New tools for automated high-resolution cryo-EM structure determination in
RELION-3. eLife *7*, e42166. https://doi.org/10.7554/eLife.42166.

70. Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017). cryoSPARC:
algorithms for rapid unsupervised cryo-EM structure determination. Nat. Methods *14*, 290–
296. https://doi.org/10.1038/nmeth.4169.

71. Pintilie, G.D., Zhang, J., Goddard, T.D., Chiu, W., and Gossard, D.C. (2010). Quantitative
analysis of cryo-EM density map segmentation by watershed and scale-space filtering, and
fitting of structures by alignment to regions. J. Struct. Biol. *170*, 427–438.
https://doi.org/10.1016/j.jsb.2010.03.007.

72. Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and
Ferrin, T.E. (2004). UCSF Chimera—A visualization system for exploratory research and
analysis. J. Comput. Chem. *25*, 1605–1612. https://doi.org/10.1002/jcc.20084.

73. Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and development of
Coot. Acta Crystallogr. D Biol. Crystallogr. *66*, 486–501.
https://doi.org/10.1107/S0907444910007493.

74. Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H., and
Ferrin, T.E. (2018). UCSF ChimeraX: Meeting modern challenges in visualization and
analysis. Protein Sci. *27*, 14–25. https://doi.org/10.1002/pro.3235.

75. Adams, P.D., Afonine, P.V., Bunkóczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J.,
Hung, L.-W., Kapral, G.J., Grosse-Kunstleve, R.W., et al. (2010). PHENIX: a comprehensive
Python-based system for macromolecular structure solution. Acta Crystallogr. D Biol.
Crystallogr. *66*, 213–221. https://doi.org/10.1107/S0907444909052925.

76. Afonine, P.V., Poon, B.K., Read, R.J., Sobolev, O.V., Terwilliger, T.C., Urzhumtsev, A., and
Adams, P.D. (2018). Real-space refinement in PHENIX for cryo-EM and crystallography.
Acta Crystallogr. Sect. Struct. Biol. *74*, 531–544.
https://doi.org/10.1107/S2059798318006551.

77. Yu, R.C., Jahn, R., and Brunger, A.T. (1999). NSF N-Terminal Domain Crystal Structure:
Models of NSF Function. Mol. Cell *4*, 97–107. https://doi.org/10.1016/S1097-
2765(00)80191-4.

78. Joosten, R.P., Long, F., Murshudov, G.N., and Perrakis, A. (2014). The PDB_REDO server
for macromolecular structure model optimization. IUCrJ *1*, 213–220.
https://doi.org/10.1107/S2052252514009324.

79. Hamelryck, T., and Manderick, B. (2003). PDB file parser and structure class implemented
in Python. Bioinformatics *19*, 2308–2310. https://doi.org/10.1093/bioinformatics/btg299.

80. Cock, P.J.A., Antao, T., Chang, J.T., Chapman, B.A., Cox, C.J., Dalke, A., Friedberg, I.,
Hamelryck, T., Kauff, F., Wilczynski, B., et al. (2009). Biopython: freely available Python
tools for computational molecular biology and bioinformatics. Bioinformatics *25*, 1422–
1423. https://doi.org/10.1093/bioinformatics/btp163.

81. McKinney, W. (2010). Data Structures for Statistical Computing in Python. In Proceedings
of the 9th Python in Science Confference, pp. 56–61. https://doi.org/10.25080/Majora-
92bf1922-00a.

82. Harris, C.R., Millman, K.J., van der Walt, S.J., Gommers, R., Virtanen, P., Cournapeau, D.,
Wieser, E., Taylor, J., Berg, S., Smith, N.J., et al. (2020). Array programming with NumPy.
Nature *585*, 357–362. https://doi.org/10.1038/s41586-020-2649-2.

83. Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D.,
Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0: fundamental
algorithms for scientific computing in Python. Nat. Methods *17*, 261–272.
https://doi.org/10.1038/s41592-019-0686-2.

84. Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Prettenhofer, P., Weiss, R., Dubourg, V., et al. (2011). Scikit-learn: Machine Learning in
Python. J. Mach. Learn. Res. *12*, 2825–2830.

85. Hunter, J.D. (2007). Matplotlib: A 2D Graphics Environment. Comput. Sci. Eng. *9*, 90–95.
https://doi.org/10.1109/MCSE.2007.55.

86. Gowers, R.J., Linke, M., Barnoud, J., Reddy, T.J.E., Melo, M.N., Seyler, S.L., DomaÅ„ski,
J., Dotson, D.L., Buchoux, S., Kenney, I.M., et al. (2016). MDAnalysis: A Python Package
for the Rapid Analysis of Molecular Dynamics Simulations. In Proceedings of the 15th
Python in Science Conference, S. Benthall and S. Rostrup, eds., pp. 98–105.
https://doi.org/10.25080/Majora-629e541a-00e.

1285
1286