Structural remodeling of target-SNARE protein complexes by NSF enables synaptic transmission

K. Ian White[1,2,3,4,5,**], Yousuf A. Khan[1,2,3,4,5], Kangqiang Qiu[6], Ashwin Balaji[7], Sergio Couoh-Cardel[1,2,3,4,5], Luis Esquivies[1,2,3,4,5], Richard A. Pfuetzner[1,2,3,4,5], Jiajie Diao[6], Axel T. Brunger[1,2,3,4,5 ,*]

[1]Department of Molecular and Cellular Physiology, Stanford University, Stanford, California, United States
[2]Department of Neurology and Neurological Sciences, Stanford University, Stanford, California, United States
[3]Department of Structural Biology, Stanford University, Stanford, California, United States
[4]Department of Photon Science, Stanford University, Stanford, California, United States
[5]Howard Hughes Medical Institute, Stanford University, Stanford, California, United States
[6]Department of Cancer Biology, University of Cincinnati College of Medicine, Cincinnati, United States
[7]Department of Chemistry, Stanford University, Stanford, California, United States

[*]Correspondence: brunger@stanford.edu
[**]Correspondence: kiwhite@stanford.edu

Keywords

Membrane vesicle trafficking, AAA+ protein; neurotransmitter release; synapse; nanocluster; active zone; syntaxin; SNAP-25; NSF; ATP; hydrolysis

Abstract

Synaptic vesicles containing neurotransmitters fuse with the plasma membrane upon the arrival of an action potential at the active zone. Multiple proteins organize trans-SNARE complex assembly and priming, leading to fusion. One target membrane SNARE, syntaxin, forms nanodomains at the active zone, and another, SNAP-25, enters non-fusogenic complexes with it. Here, we reveal mechanistic details of AAA+ protein NSF (N-ethylmaleimide sensitive factor) and SNAP (soluble NSF attachment protein) action before fusion. We show that syntaxin clusters are conserved, that NSF colocalizes with them, and characterize SNARE populations that may exist within or near them using cryo-EM. Supercomplexes of NSF, α-SNAP, and either a syntaxin tetramer or one of two binary complexes of syntaxin—SNAP-25 reveal atomic details of SNARE processing and show how sequential ATP hydrolysis drives disassembly. These results suggest a functional role for syntaxin clusters as reservoirs and a corresponding role for NSF in syntaxin liberation and SNARE protein quality control preceding fusion.

Introduction

The membrane-anchored SNARE (soluble N-ethylmaleimide-sensitive factor acceptor protein receptor) proteins provide the energy to drive membrane fusion in various cellular contexts, such as intracellular trafficking and exocytosis. One critical exocytic process is synaptic transmission, in which a synaptic vesicle fuses with the presynaptic plasma membrane in the active zone of a neuron to release neurotransmitter molecules into the synaptic cleft in response to the arrival of Ca^{2+} in the presynapse. While many proteins precisely act in concert to ensure synchronized and efficient Ca^{2+}-triggered fusion[1–5], this process is fundamentally mediated by the "zippering" of four SNARE motifs typically contributed by three to four SNARE proteins in the neuronal context—syntaxin-1a and SNAP-25 on the target plasma membrane (*t*-SNAREs), and synaptobrevin-2 on the vesicle (*v*-SNARE)—which come together to form a highly stable parallel four-helix bundle during Ca^{2+}-triggered membrane fusion. This *cis*-SNARE complex is then disassembled in an ATP-dependent manner by the AAA+ protein NSF in concert with the adaptor SNAP proteins[6,7].

While the molecular mechanisms of this core fusion cycle and the accessory proteins required to ensure Ca^{2+}-dependent neurotransmitter release have been extensively studied[4], the organization of the various components in the crowded cellular milieu remains uncertain. Numerous studies suggest nanoscale organization within the presynaptic active zone, where proteins and lipids[8–11] such as cholesterol and PIP_2 colocalize or even phase separate into discrete clusters with implications for organization within the active zone and even across the synapse[12].

The SNARE protein syntaxin shows such an organization[11,13]. Syntaxin is conserved from yeast to mammals[14,15]; it comprises a short C-terminal transmembrane domain, a SNARE domain, and

an N-terminal Habc domain [16,17]. In the neuronal context, the majority (~ 67%) of syntaxin

molecules form nanodomains of non-uniform density[13,18], often near the active zone[8,13,18–21], and

these nanodomains frequently overlap with other essential active zone proteins and even docked

granules[11,19,20,22–25]. Morphologically, they are dense, around 50–60 nm in diameter, and harbor

as many as 70 syntaxin molecules each[13,18,19,22,26]. Increased presynaptic activity drives a

decrease in syntaxin cluster size[27]. These clusters depend on the syntaxin SNARE domain (H3),

which drives oligomerization[28].

Despite extensive study, the functional purpose of these clusters has remained unclear. Moreover,

NSF, α-SNAP, and syntaxin appear to interact *in vitro*[29], and α-SNAP associates with liposomes

presenting reconstituted full-length syntaxin[30]. Binary (syntaxin—SNAP-25) complexes of

different stoichiometries may also form [31–37], and NSF disassembles these complexes *in*

vitro[31,38–40]. Furthermore, continuous ATP hydrolysis by NSF on plasma-membrane mimicking

vesicles with reconstituted syntaxin and SNAP-25 produces a primed state for the most efficient

and synchronous synaptic vesicle fusion[3]. These observations support a critical role for NSF not

only in recycling SNARE proteins following fusion, but also in regulating the pool of fusogenic

syntaxin available to enter the vesicle fusion cycle.

Here, we show that spatially organized nanodomains of syntaxin may serve as conserved

molecular protein stores, where NSF likely disassembles syntaxin homo- and hetero-oligomers.

Thus, NSF may act as a quality control factor to disassemble off-pathway binary complexes of

syntaxin and SNAP-25—a prerequisite for synaptic vesicle priming and fusion. We also describe

the molecular disassembly mechanism and present a sequential ATP hydrolysis mechanism by

which NSF couples γ-phosphate release to SNARE substrate remodeling, with broad

implications for the AAA+ protein family.

Results

Syntaxin nanodomains are conserved hotspots of NSF activity

First, we hypothesized that NSF is enriched at syntaxin clusters in the active zone, where it could

act to liberate fusogenic syntaxin molecules. To test this, we monitored the colocalization of

syntaxin and NSF in PC12 cells, a model system for neuronal secretion, using structured

illumination microscopy (SIM). All cells were transfected with syntaxin-GFP and as well NSF

with either a short C-terminal linker followed by mScarlet-I (NSF-mScarlet-I; Fig. 1A), a mutant

form of NSF (NSF(ΔN)-mScarlet-I; Fig. 1B), or mScarlet-I alone (Fig. 1C), or several

catalytically impaired mutants (Supplementary Fig. 1A). NSF-mScarlet-I is hexameric and,

together with α-SNAP, retains the capacity to bind and disassemble the ternary SNARE complex

(Supplementary Fig. 2A). Clear, punctate co-clustering between NSF and syntaxin is visually

apparent upon inspection of the micrographs (Fig. 1A–C); Pearson's colocalization coefficient

(PCC) was calculated between the green and red channels to quantitate this (Fig. 1D, Methods).

Cells transfected with both syntaxin-GFP and NSF-mScarlet-I showed significant and

reproducible syntaxin-GFP—NSF-mScarlet-I colocalization (Fig. 1A, PCC = 60% ± 2%; p <

0.001 vs. mScarlet-I only;), suggesting an interaction between NSF and syntaxin molecules

within the nanodomains. As a control, mScarlet-I alone did not colocalize with syntaxin, showing

a nearly three-fold reduction in colocalization (Fig 1C; PCC = 22% ± 2%). To verify that

colocalization is consistent with the binding mode of NSF, in which NSF N-domains engage

multiple α-SNAPs, which in turn coat the SNARE complex[6,40,41], we examined colocalization of

GFP-syntaxin with a mutant form of NSF with the N-domain required for α-SNAP binding—and

thus SNARE complex binding—removed (NSF(ΔN)-mScarlet-I)[42]. As expected, this reduced

colocalization to a level nearly indistinguishable from the mScarlet-I control (Fig. 1B, PCC =

28% ± 3%; $p = 0.1776$ vs. mScarlet-I only). Two previously characterized[43] point mutations to

the D2 domain also disrupted colocalization between syntaxin and NSF (Supplementary Fig.

1A–B). These colocalization studies suggest that NSF is associated with some clustered,

multimeric species involving syntaxin molecules *in vivo*, and that the interaction is dependent

upon the recognition of a left-handed, four-helix SNARE complex. While these results showed

clustering of syntaxin in the context of synaptic transmission, we found that clustering also

occurs in the case of the conserved syntaxin homolog in yeast, Sso1p, as well (Fig. 1F,

Supplementary Fig. 1C). During cell division, the mother cell transfers plasma membrane and

cargo to the budding daughter cell via secretory vesicles, and tethering and fusion take place

across daughter cell plasma membrane[44]; we observe Sso1p clusters enriched throughout the

budding cell in areas likely undergoing active fusion.

Discovery of a syntaxin tetramer complex with NSF—α-SNAP

To gain insight into the contents of the syntaxin nanodomains as well as the molecular basis of

the interaction between NSF and syntaxin, we purified the complex consisting of NSF, α-SNAP,

and soluble syntaxin (residues 2–253, transmembrane region omitted) by mixing the individual

purified proteins and performing size exclusion chromatography (SEC), Supplementary Fig. 3A).

We note that soluble syntaxin exists as a mixture of monomeric and oligomeric species[36]. We

then performed cryo-EM single-particle analysis (SPA) of this complex. Following 3D

classification, 5% of the particles formed a class with an overall resolution of 3.75 Å, revealing a

complex (referred to as sx20S) between hexameric NSF, four α-SNAP molecules, and a parallel

four-helix bundle of syntaxin H3 SNARE domains (sx20S; Fig. 2A, Supplementary Fig. 4,

Supplementary Tables 1–2, Supplementary Discussion). The remaining particles showed no

evidence of SNARE substrate. The finding of a syntaxin tetramer is consistent with crosslinking

mass spectrometry analysis of purified soluble syntaxin[36], and corroborates the specificity of the

system for parallel, left-handed, four-helix SNARE bundles as no other species identified by

mass spectrometry were found in complex with NSF and α-SNAP. The sx20S complex thus

further supports the notion that NSF and α-SNAP recognize parallel, four-helix SNARE

complexes of variable composition[6,7].

Similar to structures of hexameric NSF and α-SNAP bound to the neuronal ternary SNARE

complex (referred to as 20S complex) under non-hydrolyzing conditions[6,41], the sx20S complex

forms a three-tiered architecture (Fig. 2A). NSF comprises three structured domains connected

by flexible linkers—the adaptor N-domain followed by the D1 and D2 AAA+ domains. Six NSF

protomers come together to form the active form of the enzyme, a homo-hexamer. In the top tier,

or spire, the NSF hexamer's multiple N domains bind four α-SNAP molecules and a tetramer of

syntaxin H3 SNARE domains. This spire structure positions the N-terminal end of the syntaxin

tetramer near the pore of the hexameric D1 ATPase ring of NSF; this middle tier is responsible

for ATP hydrolysis and SNARE complex disassembly. The bottom tier consists of the non-

hydrolytic NSF D2 ATPase ring responsible for the oligomerization of the complex. We note

weak, poorly defined density on the outside of the D2 pore, suggestive of a highly mobile Habc

domain; this is consistent with observations of the H_{abc} domain in the yeast 20S complex of

Sec17, Sec18, and the yeast SNARE complex (including the syntaxin-1a homolog Sso1p), in

which the Sso1p Habc domain is found outside of the D2 pore, packed against the surface of the

ring[45]. Presumably, the remaining Habc domains extend outwards from the space between the

spire and the D1 ring, although they are not resolved in our structures of the neuronal 20S

complex.

The spire structure appears dynamic and rocks about relative to the D1 pore, but three of the six

N domains are relatively well-ordered and engaged with the α-SNAP—syntaxin sub-complex.

The D1 ring is asymmetric and split; protomers A–F are defined from the bottom of the split to

the top. All six D1 domains are ordered and engage the N-terminal linker preceding the H3

domain of one of the four syntaxin molecules via conserved pore loops, which position Y294

such that it intercalates between every other syntaxin substrate side chain passing through the

pore (Fig. 2B). The nucleotide states of the NSF ATPase domains are consistent with a pre-

hydrolysis conformation; D1 protomer A is ADP bound, while D1 protomers B–F are ATP bound

with no evidence for divalent cation as expected (Supplementary Table 2). All D2 protomers are

ATP bound and lack divalent cation.

While the spire in the complete sx20S complex reconstruction was ordered enough to distinguish

the various components, continuous disorder limited the local resolution of the spire

reconstruction. To overcome this, we performed signal subtraction to remove the D1 and D2

rings, followed by local alignment of the α-SNAP—syntaxin sub-complex structure alone

(Supplementary Fig. 4, Methods). This approach improved the resolution of the reconstruction of

the α-SNAP—syntaxin sub-complex to 4.25 Å, revealing a parallel, four-helix bundle of

syntaxin SNARE motifs with a density of sufficient quality to assign characteristic side chains

(Fig. 2C). All four helices are aligned in the same register, with a unique hydrophobic layer (-3

layer) composed of four F216 side chains present; remarkably, all fit within the core of the

bundle (Fig. 2D, Fig. 5A) and induce little overall deformation of the individual H3 domains

(Fig. 5B–D, bottom row). The parallel syntaxin tetramer bundle observed in our structure would

allow membrane incorporation of the C-terminal transmembrane domains of all four syntaxin

molecules (Fig. 2A).

To determine whether NSF and α-SNAP act to disassemble the syntaxin tetramer, we devised an

in vitro, gel-based disassembly assay (Supplementary Fig. 2B–C). Wild-type NSF, α-SNAP, and

soluble syntaxin (residues 2–253) were purified independently, mixed in the absence of Mg^{2+}

(*i.e.*, non-hydrolyzing conditions), and the sx20S complex was purified. The subsequent addition

of Mg^{2+} to the purified complex resulted in a significant loss of α-SNAP—syntaxin following

filtration relative to the non-hydrolyzing control, consistent with the disassembly of the syntaxin

tetramer and concurrent disassociation of α-SNAP (Supplementary Fig. 2C, $21 \pm 1\%$ loss). This

activity is also shown in the context of the Munc18 capture assay described below (Fig. 1E).

183 NSF-catalyzed disassembly of the tetrameric syntaxin SNARE complex
184 is required for efficient syntaxin capture by Munc18

NSF and α-SNAP constantly disassemble SNARE complexes at the synapse, and individual

syntaxin molecules can then be captured by Munc18 for efficient priming, and, ultimately, for

fusion to occur[3,46,47]. The capture of free syntaxin molecules, either from clusters or otherwise,

accompanies the formation of "on-granule" clusters of syntaxin—Munc18 complex[20,22,25], with

implications for animal behavior[48]. Our co-localization results (Fig. 1A–D) suggest that syntaxin

is at least partially sequestered in stable SNARE complexes that are a substrate for NSF. This

notion is further supported by the finding that α-SNAP must be present for NSF colocalization to

occur, as inferred from the co-localization experiments comparing NSF with and without the N-

domain of NSF, the deletion of which precludes α-SNAP binding (Fig. 1A–B). Finally, we note

that α-SNAP itself inhibits fusion by competing with Munc18 to bind syntaxin[30].

Having shown that NSF and α-SNAP disassemble the tetrameric syntaxin SNARE complex

(Supplementary Fig. 2B–C), we next performed a related experiment to test the hypothesis that

disassembly precedes Munc18 capture by monitoring the formation of the syntaxin—Munc18

complex. Again, wild-type NSF, α-SNAP, and soluble syntaxin (residues 2–253) were purified

independently, mixed in the absence of Mg^{2+} (*i.e.*, non-hydrolyzing conditions), and the sx20S

complex was purified. The pool was split in half, and Munc18 was then added to each in excess,

followed by the addition of either EDTA (non-hydrolyzing control) or Mg^{2+} (hydrolyzing

conditions) to initiate the disassembly reaction. Following incubation at room temperature for 10

min, size-exclusion chromatography (Fig. 1E) and SDS-PAGE (Supplementary Fig. 2D–E) were

performed to assess syntaxin oligomer disassembly and formation of the syntaxin—Munc18

complex[30].

Under non-hydrolyzing conditions, a supercomplex of sx20S—Munc18 formed, with some

syntaxin captured by Munc18. Strikingly, the addition of Mg^{2+} triggered nearly complete

disassembly of the tetrameric syntaxin SNARE complex, leaving free NSF, a 2-fold enriched

syntaxin—Munc18 and nearly two-fold enriched free α-SNAP as determined by integrated area

(Fig 1E). Together, these results support the notion that NSF and α-SNAP disassemble the

tetrameric syntaxin SNARE complex and show that complete capture by Munc18 is dependent

upon this disassembly process, consistent with previous observations of the inhibitory nature of

α-SNAP. The finding of a sx20S—Munc18 complex is not surprising, as the N-termini of the

syntaxin molecules remain accessible, enabling a weak syntaxin—Munc18 interaction[49].

Structure of a 2:1 syntaxin—SNAP-25 complex with NSF—α-SNAP

In addition to the syntaxin homo-tetramer, hetero-oligomers with SNAP-25 are expected to be

present near or even within syntaxin nanodomains. Indeed, SNAP-25 forms distinct clusters near

syntaxin clusters in neurons[19,22,24] and readily forms binary complexes with syntaxin *in vitro*[31–]

[35,50,51]. Although SNAP-25 is required for fusion, assembly of a fusogenic ternary *trans*-SNARE

complex likely proceeds first through a syntaxin—synaptobrevin intermediate templated by

Munc18-1, also found near syntaxin clusters[20,24,25], and by Munc13-1[31,46,52]. Disassembly of the

binary complex is critical, as genetic deletion of NSF is lethal[53]. However, an NSF mutation that

differentially blocks ternary, but not binary, SNARE complex disassembly causes only

intermediate sensorineural defects[38]. Thus, NSF and α-SNAP likely disassemble binary

complexes of syntaxin and SNAP-25 *in vivo* to set the stage for the action of Munc18 and

Munc13, ultimately resulting in synchronous and efficient Ca^{2+}-triggered fusion[3,31,32,52].

To gain insight into the structural nature of binary complex formation and disassembly by NSF,

we prepared a complex of NSF, α-SNAP, and the binary *t*-SNARE complex (21bin20S)

composed of full-length soluble SNAP-25 and soluble syntaxin-1a (1–267) in the absence of

divalent cation (*i.e.*, non-hydrolyzing conditions) and performed single-particle cryo-EM

(Supplementary Tables 1–2, Supplementary Fig. 3B, Supplementary Figs. 5–6). First, models of

the NSF D1 and D2 domains with substrate bound in the D1 pore were built into two high-

resolution (2.88 Å, 2.92 Å) classes from a single dataset, differing only in the state of F protomer

engagement with substrate (Supplementary Fig. 5). Next, the dataset from which they were

derived was combined with two additional datasets to enable the 3D reconstruction of weakly

populated states with ordered density for the α-SNAP—SNARE spire (Supplementary Fig. 6).

Classes with evidence of NSF D1 domains engaging ordered α-SNAP—SNARE sub-complexes

were carried forward; eight 21bin20S models were built into reconstructions ranging from 3.39–

4.01 Å in resolution, differing mainly in the orientation of N domains and a sub-complex of three

α-SNAP molecules and four SNARE motifs relative to the D1 ATPase ring (Fig. 6A). All classes

revealed substrate-bound NSF without Mg^{2+} needed for ATP hydrolysis and disassembly

(Supplementary Table 2), and share the same overall architecture, with a spire composed of NSF

N-domains and the α-SNAP—SNARE subcomplex on top, the NSF D1 ring in the middle tier,

and the NSF D2 ring forming the bottom tier (Fig. 3A). In all cases, the NSF D1 ATPase ring

engages SNARE substrate, with one syntaxin molecule bound to the D1 ring pore in most

structures (Fig. 3B, Supplementary Table 2).

Like the sx20S complex, continuous variability of the α-SNAP—SNARE subcomplex was

present in the 21bin20S reconstructions. We again performed signal subtraction and local

alignment on α-SNAP—SNARE sub-complexes from all five classes with ordered spire

structures (Supplementary Fig. 6; see Methods). This procedure produced a single α-SNAP—

SNARE sub-complex reconstruction with an overall resolution of 3.74 Å (Fig. 3C), sufficient to

assign characteristic side chains, identify the SNAREs forming the complex and their

directionality (Fig. 3D, Fig. 5A). A model composed of the ternary neuronal SNARE complex[54]

with synaptobrevin removed was docked into the density, preserving a distinctive kink in the

peptide linker leading into the SN2 SNARE motif of SNAP-25. In place of synaptobrevin, a

second syntaxin molecule was placed to complete the four-helix bundle with little deviation from

its usual backbone configuration of the ternary complex (Fig. 5B). As such, the stoichiometry of

the syntaxin—SNAP-25 complex observed is 2:1, the dominant species in solution[33,36,50,51,55].

Sidechains comprising essential layers repack within the core of the complex (Fig. 3D) compared

to the ternary SNARE complex. For example, the ionic central layer (0 layer) is reorganized;

Arg56 of synaptobrevin is replaced with a second Gln226 from the additional syntaxin molecule,

which adopts an extended rotamer to mimic the arginine it replaces. The SNAP-25 linker, which

joins the two SNARE motifs, is largely disordered but blocks binding by a fourth α-SNAP

molecule, as observed for the neuronal SNARE complex[6].

Docking the 2:1 binary complex structure back into the seven individual 21bin20S

reconstructions reveals an apparent preference for syntaxin, with six of eight reconstructions

featuring the N-terminus of syntaxin H3 bound within the pore of the D1 ring (Fig. 3B). As in the

case of the 20S with ternary SNARE complex, this observation is consistent with previous work,

in which disassembly was not contingent on specific SNARE N-terminus identity[6,40]. Taken

together, we observe that the NSF D1 ring can bind to at least two different SNAREs, syntaxin,

and SNAP-25, in various configurations.

Structure of a 2:2 syntaxin—SNAP-25 complex with NSF—α-SNAP

To gain insight into the disassembly of the binary SNARE complex, we repeated the binary 20S

preparation but added Mg^{2+} and an ATP regeneration system[56] (*i.e.*, hydrolyzing conditions) to

the purified complex ~1 min before vitrification for single particle cryo-EM (Supplementary

Tables 1–2, Supplementary Fig. 3C, Supplementary Fig. 7). The addition of the regeneration

system more closely mimics the metabolic state of the cell, and it allows multiple cycles of

SNARE disassembly and reassembly to proceed as observed under far more dilute conditions[32].

Following 3D classification, 41% of particles formed a complete complex, all of which have

ordered α-SNAP—SNARE spire density and differ largely by spire pose (Fig. 6A). Interestingly,

classes with poorly defined SNARE substrate, as observed under the non-hydrolyzing condition,

were no longer observed. The remaining 59% of particles were SNARE-substrate free and

grouped into three classes with split D1 and D2 rings; presumably, this is a post-disassembly,

substrate-released state, and it is not observed under non-hydrolyzing conditions.

Substrate-bound reconstructions reveal a flattened D1 ring in closer proximity to the D2 ring and

a strikingly different α-SNAP—SNARE spire structure than observed in the 21bin20S complex

under non-hydrolyzing conditions (Fig. 4A, Supplementary Fig. 8A). Again, syntaxin is bound

within the pore of the D1 ring (Fig. 4B, Supplementary Table 2). We observed a nearly

symmetric α-SNAP—SNARE subcomplex comprising four α-SNAP molecules coating a four-

helix bundle. Even without further processing, this structure was recognizable as a 2:2

syntaxin—SNAP-25 sub-complex formed by two syntaxin H3 SNARE domains and two SNAP-

25 SN1 SNARE domains; this binary 2:2 configuration was previously observed in crystal

structures of syntaxin-1 H3 and SNAP-25 SN1 domains[34] and was unexpected as we prepared

the complex with full-length proteins, implying that two independent SNAP-25 molecules

contribute their SN1 SNARE domains to this complex. Presumably, disassembly of the binary

2:1 complex (Fig. 3C) had occurred and was followed by reassembly to form the 2:2 binary

complex (Fig. 4C) before vitrification.

To improve the resolution of the spire reconstruction, signal subtraction, and local alignment

were performed as before (Supplementary Fig. 7). This approach improved the reconstruction of

the spire proteins to 4.22 Å, enabling docking of a crystal structure of the 2:2 syntaxin—SNAP-

25 SN1 complex[34] into the density (Fig. 4C), which was of sufficient quality to assign

characteristic side chains, to identify the SNARE domains and their directionality (Fig. 4D, Fig.

5A). The 2:2 binary complex was fit back into the full reconstructions, enabling modeling of the

full 22bin20S. Furthermore, we were able to reproduce this result by reconstituting the complex

with only the SNARE domains of syntaxin and SNAP-25 under non-hydrolyzing conditions

(min22bin20S, Supplementary Fig. 9A–D), yielding the same spire configuration but to a higher

resolution of 3.86 Å following local reconstruction (Supplementary Fig. 9E, Fig. 5A). This

reconstruction also reveals the interface between an N-domain of NSF and an α-SNAP molecule

in detail; the interface is characterized by a core hydrophobic interaction surrounded by charge

complementation (Supplementary Fig. 9F).

Insights into SNARE substrate engagement and processing by NSF

Hydrolysis by NSF is required for efficient priming and, ultimately, fusion. Collectively, we

obtained many datasets with NSF, α-SNAP, and several SNARE substrates under non-

hydrolyzing and hydrolyzing conditions (Supplementary Tables 1–2), presenting an opportunity

to investigate the molecular mechanism of SNARE substrate processing and chemomechanical

energy transduction in NSF. Together, the resulting high-resolution structures derived from the

different classes from these datasets capture the hydrolysis cycle of NSF and reveal important

details about its reaction coordinate.

For datasets with particles collected under non-hydrolyzing conditions, NSF is found in two

general conformations, primarily differentiated by their F protomer state (Fig. 7A, left column).

Two high-resolution 21bin20S classes from the 2:1 binary SNARE complex dataset reveal large-

scale conformational displacement connected to D1 F promoter nucleotide state and SNARE

substrate engagement. The first class is ATP-bound and fully SNARE substrate-engaged, while

the second has a nucleotide-free and SNARE substrate-disengaged F protomer (Fig 5A, left,

bottom, and top, respectively; Supplementary Fig. 5). The engagement of F at the top of the ring

accompanies a relaxation of A into the split. Otherwise, the classes are nearly identical;

protomers B–E D1 domains also bind ATP, while protomer A, at the bottom of the D1 split, is

ADP-bound in both cases. Comparison of the two classes reveals limited differences outside of

the protomers A and F positions, as the change in protomer F position allows additional

relaxation of protomer A into the split.

In contrast to the non-hydrolyzing datasets, particles in the hydrolyzing dataset fall into two categories—substrate-bound 22bin20S (Fig. 7A, middle column) or substrate-free NSF (post-hydrolysis; Fig. 7A, right column), of which there are ten and three classes, respectively (Supplementary Table 2, Supplementary Fig. 7). The particles show a population shift towards a flatter D1 ring that is closer to D2 relative to the non-hydrolyzing class, particularly in the case of the SNARE-substrate free classes, which also show an expanded pore (Supplementary Fig. 8A–B). Most strikingly, no SNARE substrate-bound classes show an F protomer as fully engaged as those in non-hydrolyzing classes; in all but one class, F is ADP bound, SNARE substrate disengaged, and sits between engaged A and E protomers. However, a subclass with an ATP-bound F protomer was found using a subdivision of a large class with 3D Variability Analysis[57] (3DVA; Supplementary Fig. 10C–D) for which the interprotomer interface with the neighboring D1 protomer E is nearly—but not completely—formed. Critically, the pore loop approaches the substrate-engaged state found in the non-hydrolyzing classes; together, this reveals a trajectory of protomer A dissociation, nucleotide release and rebinding, and, finally, SNARE substrate rebinding (Fig. 7D). Furthermore, in classes with ordered syntaxin substrate, the unstructured region of syntaxin N-terminal to the H3 domain is engaged by the D1 pore loops around eight residues further into the primary sequence under hydrolyzing conditions (Fig. 6B).

Sequential hydrolysis by NSF drives conformational cycling

As noted above, the nucleotide states of the D1 protomers help explain the pattern of engagement with the SNARE substrate (Fig. 7A–C, Supplementary Table 2). How does nucleotide hydrolysis drive this process? Unlike the non-hydrolyzing 21bin20S reconstructions, hydrolyzing 22bin20S reconstructions are only ATP-bound in protomer E, consistent with a sequential hydrolysis mechanism (Fig. 7A, middle column). Except for the rebinding F protomer described above, all

other D1 protomers are ADP bound with variability in the presence of Mg^{2+} and inorganic

phosphate (P_i); in most cases, protomers C and D have ADP•P_i and Mg^{2+}, protomer B is ADP-

bound, and protomer A is typically ADP-bound only. Finally, substrate-free classes are always

ADP bound, even in protomer E; they are flat, have some N-domains packed between the D1 and

D2 rings, and are characterized by a split in both the D1 and D2 rings between protomers A and

F (Fig. 7A, right column); this is consistent with a SNARE substrate side-loading mechanism

proposed for the NSF ortholog in yeast, Sec18[45]. In both hydrolyzing and non-hydrolyzing

datasets, all D2 domains are ATP-bound, with Mg^{2+} present in the hydrolyzing condition.

Together, these classes represent key steps in the hydrolysis cycle, revealing the binding of Mg^{2+},

hydrolysis of ATP to ADP•P_i at protomer E, and subsequent P_i release down-ring; these states are

accompanied by a conformational change in the Walker A, Walker B, and Sensor 1 active site

residues in *cis* and the arginine fingers of the down-ring protomer in *trans* (Fig. 7B), and are

connected in turn to larger scale conformational change within the D1 domain (Fig. 7C).

Critically, P_i release following hydrolysis leads to arginine finger disengagement (Fig. 7B, right

two panels) and subsequent relaxation and solvation of the interprotomer interface, enabling the

protomer A transition to protomer F and subsequent substrate rebinding.

What triggers hydrolysis at E, and what conformational changes follow? While hydrolysis could

proceed spontaneously, this would be inefficient in the absence of an ATP-bound protomer F up-

ring to capture accompanying substrate translocation. We hypothesized that some allosteric

mechanism likely regulates this process. To identify such a mechanism in an unbiased manner,

we performed principal components analysis (PCA) of Cα coordinates of 132 D1 protomers

across all datasets followed by hierarchical density-based clustering (HDBSCAN) in high-

dimensional space, revealing the conformational cycling of the NSF D1 protomers in a low-

dimensional coordinate as well as a corresponding allosteric mechanism for modulating

hydrolysis by protomers E (Figs. 8–9).

Overall, we find protomers divided into five interpretable clusters connected to ATP hydrolysis,

P_i release, and protomer cycling at the D1 split during SNARE disengagement, nucleotide

rebinding, and SNARE re-engagement (Fig. 8A). Here, we focus on the unique features of the

protomers E under hydrolyzing conditions in two dimensions of the 11-dimensional latent space,

where they cluster far from other protomers (Fig. 8A, transition 1). As noted above, protomers E

are the only ATP-Mg^{2+} bound D1 protomers under hydrolyzing conditions (Fig. 8B), and they are

uniquely characterized by an extreme large-small subdomain angle (Fig. 8C, Supplementary Fig.

8). Examination of PCA and clustering results reveal conformational states unique to these

protomers (Supplementary Fig. 10A, Fig. 8), notably a discrete Walker A (α_3) helix position

("down" state) relative to the rest of the D1 large subdomain that uniquely positions the

nucleotide near two key catalytic residues (Sensor I, N374 and Arginine finger 2, R388; Fig. 8E

left, Fig. 9), a unique conformation of a highly conserved[58] segment of the N-D1 linker

connected to N-domain engagement ("up A" state, Fig. 8E, middle), and the absence of a fully

engaged "latch loop" (residues 457–467) contributed by the up-ring protomer F in *trans*

("intermediate A" state; Fig. 8E, right). All three of these features appear structurally coupled

(Supplementary Fig. 10B) and comprise an allosteric integration mechanism by which the

position of nucleotide in the active site is modulated at a distance by both N-domain engagement

to SNARE substrate via the N-D1 linker conformation and by the presence of an up-ring

protomer F via latch loop engagement. This modulation in turn influences the distance between

key catalytic active site residues and nucleotide (Fig. 9).

To test the importance of the latch loop interaction in *trans*, a mutant form of NSF was generated

in which a portion of the latch loop (residues 460–466) is replaced with a short glycine-serine

linker (SGS). An NSF activity assay based on fluorescence dequenching (see Methods) shows a

highly significant, ~72% reduction in the rate of SNARE complex disassembly ($p = 0.0005$; Fig.

8D). On the other hand, the functional importance of the N-D1 linker has already been

established. A mutation to NSF in the most conserved region N-D1 linker, I209N, has a direct

behavioral effect in zebrafish; NSF(I209N) is defective in the disassembly of ternary SNARE

complexes but not binary SNARE complexes, explaining this phenotype[38,58]. This isoleucine side

chain directly contacts the engaged latch loop and is adjacent to S207, a putative phosphorylation

site[59], further raising the possibility of post-translational modification in regulating NSF

activity[60]. Together, these observations support an allosteric mechanism critical in modulating

NSF function.

411 Discussion

Using structural and biophysical studies, we have discovered a key organizational principle at the

active zone in the presynapse that likely involves the remodeling of syntaxin clusters. Synaptic

vesicle fusion and neurotransmitter release are among the most central processes of synaptic

transmission, and understanding the organization and the molecular mechanisms of these

processes in the cellular context is essential. Our work helps to answer a key unanswered

question—what is the purpose of phase-separated syntaxin at the presynapse?

Our observation of NSF colocalization to syntaxin clusters *in vivo*, coupled with direct evidence

of disassembly of both syntaxin tetramers and syntaxin—SNAP-25 complexes, explains a

previously overlooked but essential aspect of the synaptic vesicle fusion cycle with broad

implications for membrane trafficking (Fig. 10A). While NSF is well known for its role in

ternary *trans*-SNARE complex disassembly following fusion, this work, combined with

functional studies[3,31,32,38,52], further establishes NSF as a critical factor for synaptic vesicle

priming before fusion. NSF is enriched at and near syntaxin clusters, which are, in turn, found at

the presynapse. This enrichment is highly dependent upon the presence of the NSF N-domain,

which binds α-SNAP (Supplementary Figure 2E–F), which in turn binds parallel, four-helix

SNARE complexes through a stereotyped binding interface[6,7,61]. The observation here of a

parallel, four-helix syntaxin-1a bundle suggests that these clusters are structured, at least in part,

and that NSF and α-SNAP act to disassemble the syntaxin tetramers within or on the perimeter of

a given cluster, increasing syntaxin mobility at the synapse[27]. Individual syntaxin molecules

liberated from clusters by NSF can then be captured by the SM protein Munc18 and ultimately

be used to assemble fusogenic *trans*-SNARE complexes.

However, a high local concentration of SNAP-25 is present near the syntaxin clusters as

well[11,24,62], so free syntaxin may also enter a non-productive binary SNARE complex with

SNAP-25 with a stoichiometry dependent upon local concentrations and configurations of either

protein. NSF disassembly of these species prior to fusion is critical *in vivo*[38]. In this work, we

observe binary complexes with two stoichiometries—2:1 and 2:2 binary complexes of syntaxin

and SNAP-25, with the latter re-forming following NSF-mediated disassembly of the former

under hydrolyzing conditions. The 2:1 configuration dominates in the non-hydrolyzing

condition—and in other *in vitro* studies[33,51]—but is derived from co-expressed proteins that were

extensively purified without NSF action. The 2:2 configuration is only found following NSF-

mediated disassembly of the 2:1 configuration. This striking observation is likely related to

differences in the effective concentration of syntaxin dimers preceding complex formation during

expression in and purification from *E. coli* versus following disassembly *in vitro*[55]. Regardless,

these structures are consistent with the notion that SN1 is a critical initial mediator of SNARE

assembly[63]. The structural plasticity of the SNARE complexes is also of note; taken together, the

repeated observation of a twisted, parallel, four-helix bundle with minimal backbone distortion

(Fig. 5) emphasizes the functional conservation latent in the SNARE domain sequences studied

thus far, as well as the importance of additional SNARE protein accessory domains in

differentiating SNARE protein function[45]. Thus far, only parallel SNARE bundles have been

observed the context of various 20S complexes in this work and elsewhere[6,7,32,61]. While NSF

and alpha-SNAP rapidly disassemble anti-parallel SNARE complexes[32], such complexes may be

too rare or heterogeneous to be observable by cryo-EM.

We note that several recent studies[20,48] also highlight the critical connection between syntaxin

clustering, neurotransmission, and animal behavior and emphasize the role of the protein

Munc18 in capturing free syntaxin from clusters for subsequent priming. Our disassembly results

show that syntaxin capture by Munc18 depends on its liberation from SNARE complexes (Fig.

1E) on the timescales of synaptic transmission. This is also consistent with previous observations

of increasing syntaxin mobility at the synapse with NSF action[27]. As such, a complete

mechanistic picture of the system involves NSF, α-SNAP, and Munc18; Munc18 can directly

bind any monomeric syntaxin at or near the syntaxin clusters, but it likely cannot actively

remove syntaxin from oligomeric complexes within the clusters, especially on the timescales of

synaptic transmission. Our model also suggests a mechanism for the transit of syntaxin from off-

pathway, Munc18-free clusters to on-granule clusters of syntaxin-1a—Munc18[20,22] and further

explains the inhibition of Munc18 binding of syntaxin by α-SNAP[30].

Regardless, NSF and α-SNAP likely disassemble both syntaxin tetramers and binary SNARE

complexes to maintain a fusogenic population of *t*-SNAREs[3,38], which ultimately enter the *trans*-

SNARE complex with the *v*-SNARE synaptobrevin-2 and set the stage for Ca^{2+}-triggered fusion

in cooperation with synaptotagmin and complexin[64,65]. Indeed, the energy released by SNARE

protein zippering during vesicle fusion can be directly traced to the hydrolysis of ATP by NSF,

which effectively maintains them in a high-energy state. Thus, syntaxin nanodomains could

effectively act as high-density storage sites for syntaxin, and NSF could tap these stores under

periods of high-frequency vesicle release. Modulation of NSF activity by post-translational

modification could thus be one way to regulate this process[38,60]. Finally, given the observation of

clustered Sso1p in yeast, and the high degree of structural and functional conservation between

NSF and Sec18[45], this seems likely to be a conserved organizing principle across fusion contexts.

At the atomic level, how does NSF accomplish the energy transfer needed to disassemble various

SNARE complexes? The structural details of the hydrolysis cycle (Fig. 7) are consistent with

sequential hydrolysis by the six D1 ATPase domains of NSF. Hydrolysis drives conformational

change, which is coupled to the application of force to one member of the SNARE complex,

destabilizing it and promoting disassembly over multiple hydrolysis events (Fig. 10B,

Supplementary Movie 1). Rather than relying on a non-physiological nucleotide state promoted

by chelation of divalent cations or non-hydrolyzable nucleotide analogs (*e.g.*, ATPγS), we

established a reaction condition more closely approximating cellular metabolism by using a

physiological ATP regeneration system. Our sequential hydrolysis model suggests that hydrolysis

proceeds around the asymmetric NSF D1 ring, occurring near the "top" of the ring at protomer E.

Down-ring protomers reveal evidence for a transient post-hydrolysis reaction state in which

ADP•P_i is generally observed before P_i release from protomers C or B and an associated

disruption and solvation of the interprotomer interface. This mechanism is likely conserved in

other members of the AAA+ family, such as p97 (Supplementary Discussion).

Furthermore, the continuous transition from protomer A at the bottom of the ring to protomer F

at the top follows ADP release and ATP rebinding (Fig. 10B, Supplementary Fig. 10D–E,

Supplementary Movie 1). This cycling appears coupled to the triggering of hydrolysis at

protomer E in a process allosterically mediated by several key structural elements (Figs. 8–9);

presumably, full protomer F rebinding triggers hydrolysis, likely driven by an allosteric shift

associated with full latch engagement with the N-D1 linker region, further formation of the

interprotomer interface, and mediation of the distance between nucleotide and key catalytic

residues such as Sensor I and the second Arginine finger (N374 and R388, respectively; Fig. 9).

Protomer F rebinding is also connected to the mechanism by which engagement of the SNARE

substrate is maintained during processing. In the absence of a completely protomer F-rebound

state in the hydrolyzing condition, however, we cannot distinguish between SNARE substrate

rebinding before or after protomer E hydrolysis; the former case (ATP in fully rebound F and E)

represents a classical "hand-over-hand" mechanism, while the latter case (ATP in fully rebound

F, ATP ADP•Pi in E), a ratchet mechanism in which F re-engages substrate during or

immediately following a translocation event driven by protomer E hydrolysis, locking in

substrate translocation[66]. The flattened D1 rings of the SNARE-substrate-free classes with

ADP•Pi in protomers E (Supplementary Fig. 8) are suggestive of the latter; flattening upon

hydrolysis at protomer E could exert a pulling force on the substrate, and translocation would

then take place with substrate engagement by a new protomer F and release of the old protomer

A. Multiple rounds of translocation are expected to ultimately lead to melting of the engaged

SNARE domain and subsequent disruption of the SNARE complex. These large-scale

conformational changes may also apply a force via the NSF N-domains, which could aid in the

disassembly process[6,7,61].

Together, our findings expand the role of NSF in synaptic transmission and the membrane fusion

cycle while further demonstrating its contribution as a model system for understanding energy

transduction and protein remodeling in AAA+ proteins. NSF acts as a mobile engine of the

presynapse—working together with the SNAP adaptor proteins and factors such as Munc18 to

convert the chemical energy latent in ATP into the physical work needed to organize, maintain,

and drive synaptic function through the disassembly of a variety of SNARE proteins.

Methods

PC12 transfection and imaging

The PC12 cells, generously provided by Prof. Tobias Meyer from Stanford University, were

cultured on glass-bottomed dishes (MatTek; P35G-1.5-14-C) and allowed to adhere for 24 hr. To

introduce genetic material into the cells, transfection was performed using the TurboFect™

Transfection Reagent (#R0532, Thermo Fisher Scientific, USA) according to the manufacturer's

instructions. In brief, the plasmids (NSF-GGGSGGGS-mScarlet-I in pTwist CMV Puro, 1.75

μg; syntaxin1a-eGFP plasmid, 0.5 μg) were combined with 6.75 μL of TurboFect in 225 μL of

DMEM medium (without FBS) and incubated for 20 min. The DNA-TurboFect mixtures were

carefully added drop-wise to the cell cultures and incubated for 6 hr. Following the incubation,

the cell cultures were replaced with complete culture medium. After 24 hr post-transfection, the

cells underwent three washes with PBS. Subsequently, the cell membranes were imaged using a

structured illumination Nikon N-SIM system (version AR5.11.00 64bit, Tokyo, Japan) equipped

with an Apochromat 100×/1.49 numerical aperture oil-immersion objective lens. Using a 488 nm

laser, the samples were excited to acquire green SIM images, which captured emission within the

500–550 nm bandwidth, specifically for GFP. Additionally, red SIM images were obtained by

exciting the samples with a 561 nm laser to capture emission within the 570–640 nm bandwidth,

corresponding to mScarlet-I. To determine the degree of colocalization between syntaxin and

NSF, Pearson's correlation coefficient was calculated between channels *ImageJ*[67].

Yeast imaging

The Sso1-GFP strain was created previously[68] and provided by the lab of Daniel Jarosz; the yeast

harbors a plasmid with Sso1-GFP similar to that used in the PC12 experiments, in which GFP is

found outside of the cell while Sso1p itself is in the plasma membrane and cytoplasm. BY4741 is

the parental strain; N-SWAT Sso1 GFP is the protein, and **MATa his3Δ1 leu2Δ0 met15Δ0**

ura3Δ0 is the genotype. Expression was carried out using the native yeast promoter NOP1.

Super-resolution microscopy of *S. cerevisiae* expressing Sso1-GFP was performed using

structured illumination microscopy (SIM) on the OMX-Blaze system (Applied Precision,

Issaquah, WA). Glass-bottom dishes (1.5H, Cellvis) were pretreated with 0.01% poly-l-lysine by

adding 600–1000 μL to the center and incubated overnight. Following incubation, the dishes

were washed with deionized water to remove excess poly-l-lysine. Yeast cultures were grown

overnight, washed twice with phosphate-buffered saline (PBS), and resuspended to an optical

density (OD) of 0.4. A 10–20 μL volume of the yeast suspension was deposited onto the treated

dishes, followed by the addition of Diamond Antifade (Thermo Fisher Scientific) to prevent

photobleaching during imaging. The glass-bottom dishes were placed in a metal holder with 1–2

drops of immersion oil (refractive index 1.512) added before placement on the microscope stage.

SIM imaging was conducted using a laser appropriate for GFP excitation, and images were

captured with a 100× objective for a *z*-slice thickness of 0.125 μm, spanning a total thickness of

2.6 μm. Widefield images were also collected at 0.2 μm slice thickness when necessary.

Protein expression and purification

All NSF, the wild-type and mutant NSF-mScarlet-I, and NSF(Δ460–466) proteins were derived

from NSF from the Chinese hamster *C. griseus*, expressed in BL21(DE3)-RIL *E. coli*, and

purified as described previously[6], with final reassembly of hexameric NSF in reassembly buffer

(50 mM Tris pH 8.0, 150 mM NaCl, 1 mM EDTA, 1 mM ATP, and 1 mM TCEP). α-SNAP from

rat *R. norvegicus* was expressed in *E. coli* and purified as described previously[69]. Soluble rat

neuronal SNARE complex for the fluorescence-based disassembly assay (wild-type SNAP-25A,

syntaxin-1A 1–265, and 6×His-synaptobrevin-2 1–96) was purified by co-expression in C41 *E.*

coli and purified as described previously[69] Purified soluble rat neuronal SNARE complex was

fluorescently labeled with Oregon Green 488 maleimide dye as described previously[6].

Syntaxin-1A fragment 2-253 (Stx$^{2\text{-}253}$) was N-terminal His$_6$-tagged using the plasmid pTEV5.

Overexpression in *E. coli* BL21(DE3) was carried out in 8 L of auto-inducible media at 37ºC for

4 hr and then 30ºC overnight. Cells were collected by centrifugation; half of the pellet was

resuspended into 200 mL of 50 mM TRIS pH 8, 500 mM NaCl, and 10 mM imidazole and

supplemented with DNAse, lysozyme, 1 mM MgCl$_2$, and 2 eComplete Protease Inhibitor

Cocktail tablets (Roche). Cells were lysed by sonication (1 s on/3 s off, total 20 min, 55%

amplitude, 4ºC). Debris were removed by centrifugation at 43,000 rpm for 30 min at 4ºC using a

Ti45 rotor. The supernatant was then mixed with 5 mL of nickel-NTA agarose resin (Qiagen)

equilibrated with the lysis buffer. The suspension was gently stirred for 1 hr at 4ºC. The resin was

subsequently recovered by centrifugation at 4,200 rpm for 10 min at 4ºC, poured into a column,

and attached to an AKTA Start (Cytiva). An extensive wash was performed using 20 CV of 50

mM TRIS pH 8, 500 mM NaCl, and 40 mM imidazole until the 280 nm signal was stable.

Protein was eluted in 1 mL fractions in 50 mM TRIS pH8, 200 mM NaCl, and 400 mM

imidazole. Syntaxin-enriched fractions were identified by SDS-PAGE and pooled. To remove the

His-tag, the protein was mixed with an excess of TEV protease and dialyzed against 20 mM

HEPES pH 7.5, 50 mM NaCl, 1 mM DTT, overnight at 4ºC. The sample was centrifuged at

4,000 rpm for 10 min at 4ºC, filtered, and loaded onto a Mono-Q 5/50 column attached to an

AKTA Pure (Cytiva). The resin was extensively washed with 20 mM HEPES pH 7.5, 50 mM

NaCl, 0.2 mM EDTA until 280 nm signal was stable. Elution was performed in 120 CV over 2 hr

from 0.050–1 M NaCl. 2 mL fractions within the major peaks were run on SDS-PAGE, and

fractions containing Stx^{2-253} were concentrated using a 10 kDa cutoff centrifugal filtration device.

The concentrate was filtered and further purified using a Superdex Increase 75 16/60 column

(Cytiva) equilibrated with 20 mM TRIS pH 8, 150 mM NaCl. Fractions were run on an SDS-

PAGE, and fractions containing Stx^{2-253} were concentrated using a 10 kDa MWCO centrifugal

filtration device. A final concentration of 745 mM was determined by using the extinction

coefficient 5,960 M^{-1} cm^{-1}. Stx^{2-253} was aliquoted, flash frozen, and stored at -80ºC until further

use. An independent sample purified from the same batch of cells was analyzed on MALDI-TOF,

and a monomeric molecular weight of 29,208.50 g/mol was confirmed.

Binary neuronal SNARE complex for cryo-EM imaging studies (SNAP-25A with all cysteines

mutated to serine, syntaxin-1A 1–267) was purified following co-expression of SNARE proteins

from *E. coli* BL21 STAR (Invitrogen). Cells were grown overnight at 30°C using an auto-

inducing LB medium[70]. After harvesting the cells by centrifugation, the pellet was resuspended

in lysis buffer (50 mM NaPi pH 8.0, 300 mM NaCl, 20 mM imidazole, 0.5 mM TCEP)

supplemented with lysozyme, DNase I, and antiprotease tablets (Roche) and subjected to

sonication (3 s on, 9 s off, 60% power for 5 min, repeated 3×) and centrifugation (10,000 rpm for

10 min followed by 43,000 rpm for 40 min). The clarified lysate was bound to Ni-

NTA agarose beads (QIAGEN) equilibrated in a lysis buffer. Beads were harvested by

centrifugation (2200 rpm for 10 min), poured into a gravity column, and washed with lysis

buffer, urea buffer (50 mM NaP_i pH 8.0, 300 mM NaCl, 30 mM imidazole, 0.5 mM TCEP, 7.5 M

urea), and wash buffer (50 mM NaPi pH 8.0, 300 mM NaCl, 30 mM imidazole, 0.5 mM TCEP).

The SNARE complex was then eluted with wash buffer supplemented with an additional

320 mM imidazole. The fresh eluent of the Ni-NTA affinity purified SNARE complex was

supplemented with tobacco etch virus (TEV) protease and dialyzed against buffer A1 (50 mM

Tris-HCl, pH 8.0, 250 mM NaCl, 0.5mM TCEP) overnight at 4°C. After dialysis, the sample was

diluted to a final NaCl concentration of 50 mM using buffer A1 without NaCl, and the His-tag-

free complex was subjected to anion exchange chromatography (Buffer A2: 50 mM Tris-HCl, pH

8.0, 50 mM NaCl, 0.5 mM TCEP; Buffer B1: 50 mM Tris-HCl, pH 8.0, 500 mM NaCl, 0.5 mM

TCEP) using a MonoQ 4.6/100 column (Cytiva) and a linear gradient of NaCl starting at 50 mM

and ending at 500 mM. The peaks containing the purified complex were pooled and loaded onto

a Superdex 75 16/60 column to normalize NaCl concentration (50 mM Tris-HCl pH 8.0, 50 mM

NaCl, 0.5 mM TCEP).

The sx20S supercomplex was formed by mixing ATP-bound NSF, syntaxin, and α-SNAP to a

ratio of 1:5:30 in reassembly buffer and purifying it by size-exclusion chromatography using a

Superose 6 10/300 GL column; the first peak to elute was collected and concentrated to around

15 mg mL^{-1} for vitrification and cryo-EM imaging. The 21bin20S supercomplex was purified

similarly but with a 1:5:30 ratio of NSF, binary SNARE complex, and α-SNAP. In the case of

preparing 21bin20S supercomplex for imaging under hydrolyzing conditions, EDTA was omitted

from the final SEC buffer.

Size exclusion chromatography (SEC)-based syntaxin disassembly and Munc18 capture
assay

The sx20S complex was prepared as described in the "Protein expression and purification"

Methods Section without EDTA in the reassembly buffer and with a 1:50:30 ratio of ATP-bound

NSF, syntaxin, and α-SNAP. The NSF and α-SNAP concentrations were informed by the fraction

of sx20S to NSF in the cryo-EM experiment (Supplementary Fig. 4) and resulted in an increase

of the fraction of 20S with the tetrameric syntaxin SNARE complex relative to free NSF.

Following concentration to 50 mg mL^{-1} (or around 66 µM assuming full sx20S formation,

although the actual concentration is likely lower), the sx20S sample was split into two 35 µL

pools and resuspended to 500 µL in 464 µL of 43 µM Munc18, for a final concentration of 5 µM

sx20S (likely lower) and 40 µM Munc18. Either 1 µL of 500 mM EDTA or 1 µL of 500 mM

MgCl$_2$ was then added to the pool, in the latter case initiating the disassembly reaction.

Following incubation for 10 min at room temperature, the pools were independently subjected to

size exclusion chromatography using a Superdex 200 Increase 10/300 GL column (Cytiva), and

fractions were analyzed by SDS-PAGE. Absorbance data for each run were processed using the

script *elution.py*. Data parsed using *pandas*[71] and *numpy*[72], smoothed using Savitzky-Golay

filtering as implemented in *scipy*[73] with a window of 1000 points and a polynomial order of 2,

and peaks with absorbance amplitudes greater than 10 mAU, distances greater than 1 mL, and

prominences greater than 0.5 mL were found using *find_peaks* in *scipy*. Peaks were integrated

using the *trapz* function as implemented in *numpy*. Data were plotted using *matplotlib*[74].

Filtration-based syntaxin disassembly assay

Soluble syntaxin (2–253) and α-SNAP were mixed at concentrations of 100 µM and 150 µM,

respectively, in 50 mM Tris pH 8.0, 150 mM NaCl, 1 mM TCEP, and 1 mM EDTA (SEC buffer)

and incubated for 20 min. NSF was subsequently added to a final concentration of 2.5 µM,

bringing the total reaction volume to 500 µL. The mixture was then injected into a Superdex 200

16/60 column (Cytiva) and complex peak fractions were pooled and concentrated using a

100,000 MWCO filtration device. For each assay condition, 5 µg of complex was used. Two

experimental conditions were performed in triplicate; either EDTA or MgCl$_2$ was added to a final

concentration of 4 mM. Samples were incubated for 20 min at room temperature and re-

concentrated using 100k MWCO filters before SDS-PAGE. Re-concentration effectively de-

enriches liberated syntaxin and α-SNAP molecules. Densitometry was performed using

ImageJ[67], and statistical significance was established between the conditions using a two-tailed

Student's *t*-test.

Gel-based ternary SNARE complex disassembly assay

To verify that NSF activity was not hampered by the C-terminal addition of mScarlet-I, SNARE

complex disassembly was evaluated by gel electrophoresis. Note that the ternary neuronal

SNARE complex runs as a single band unless boiled in SDS buffer, or disassembled by NSF, in

which case the individual SNARE components are enriched as the complex band disappears.

NSF, NSF-mScarlet-I, α-SNAP, and the neuronal SNARE complex were dialyzed into 25 mM

TRIS pH 8, 150 mM NaCl, 1 mM ATP. Then, components were combined into either EDTA

buffer (25 mM TRIS pH 8, 2 mM ATP, 1 mM EDTA) or Mg^{2+} buffer (25 mM TRIS pH 8, 2 mM

ATP, 4 mM MgCl$_2$). The reaction volume was 10 µL, and contained 500 nM NSF or NSF-

mScarlet-I and 50 nM α-SNAP. Assays were started by adding 10 µM SNARE complex, and

samples were then incubated at 37ºC for 10 min in a thermocycler. 2 uL of 5×-Laemli loading

buffer was then added, and samples were incubated at 95ºC for 5 min. Finally, 5 µL of each

reaction was loaded onto an AnykD TGX gel (BioRad) and run at 200 V for 45 min. The gel was

stained with Coomassie R-250 (BioRad). As internal controls, all single components were diluted

at a similar concentration as the reactions using Mg^{2+} buffer and treated in parallel. Note that the

additional control marked as "SNARE complex unboiled" corresponds to the SNARE complex

incubated at 37ºC but not at 95ºC after adding loading buffer. For reference, monomeric NSF is

~83 kDa while NSF-mScarlet-I is ~112 kDa.

Fluorescence dequenching-based neuronal SNARE complex disassembly assay

The details of the fluorescence dequenching assay have been previously described[6,32,38,40,41,69].

Fluorescence dequenching assays were carried out using a FlexStation II 384-well plate reader

(Molecular Devices) with a final reaction volume of 60 µL in 20 mM Tris pH 8, 100 mM NaCl, 2

mM ATP, and 0.5 mM TCEP. The assay plate was first prepared by adding 50 µL of 60 nM wild-

type or Δ460–466 mutant NSF, 2.4 µM α-SNAP, 480 nM OG-labeled neuronal SNARE complex

in reaction buffer to each well. Then, 10 µL of reaction buffer (as a control) or ATP hydrolysis

initiation buffer (reaction buffer supplemented with 24 µM $MgCl_2$) was added to appropriate

wells of the compound plate, and fluorescence dequenching was monitored. Excitation was

performed at 485 nm, and emission was monitored at 525 nm with a dichroic mirror at 515 nm.

First, fluorescence intensity was monitored before reaction initiation for 60 s. Then, 10 µL of

compound plate solution was transferred to each assay plate well (diluting all protein 5/6-fold

and $MgCl_2$ concentrations six-fold) and triturated, initiating the disassembly reaction. The final

protein concentrations following mixing were 50 nM NSF, 2.0 µM α-SNAP, and 400 nM

SNARE complex. The final $MgCl_2$ concentration was 4 mM. Dequenching was then monitored

for 30 min at 20 ºC. Linear regression over the first several minutes was performed to estimate

relative disassembly rates. The statistical significance of changes to disassembly rates was

assessed using the two-tailed Student's *t*-test.

Sample vitrification for cryo-EM SPA

The samples were prepared immediately before vitrification (Supplementary Fig. 3). Quantifoil

R1.2/1.3 200 mesh gold grids (Quantifoil Micro Tools GmbH, Germany) were cleaned with

chloroform, dried for several hours, and subjected to glow discharge using a PELCO easiGlow

Glow Discharge Cleaning System (Ted Pella) under vacuum at 15 mA for 45 s with a 15 s hold

time. Immediately following 21bin20S or sx20S supercomplex purification, proteins were

concentrated to ~15 mg mL^{-1} in 17 µL. For the hydrolyzing condition, 2 µL of 10 × ATP

Regeneration Solution (50 mM Tris pH 8.0, 150 mM NaCl, 1 mM ATP, 160 mM creatine

phosphate, and 0.3 mg/mL rabbit muscle creatine kinase (Roche)) was added, and the sample

was incubated on ice for 2 min; in contrast, 2 µL of reassembly buffer was added in the case of

non-hydrolyzing conditions. Then, for the hydrolyzing condition, 1 µL of 20 × Reaction

Initiation Solution (50 mM Tris pH 8.0, 150 mM NaCl, 1 mM ATP, 20 mM MgCl$_2$, 1% Nonidet

P-40) was added, otherwise 1 µL of 1% Nonidet P-40 was added in the case of non-hydrolyzing

conditions. Using a Vitrobot Mark IV (Thermo Fisher Scientific), 4 µL of a given sample was

applied to a grid at room temperature and 100% humidity and, blotted for 4 s with a blot force of

3, and subsequently plunge-frozen in liquid ethane.

Cryo-EM SPA data collection

Grids were imaged using a Titan Krios electron microscope (Thermo Fisher Scientific) operated

at 300 kV and equipped with a K3 camera (Gatan). Data collection was performed using

SerialEM[75]. All datasets were collected with a nominal magnification of 22,500× for a physical

pixel size of 1.096 Å and a super-resolution pixel size of 0.548 Å. The dose rate was either ~10

or ~20 e⁻/px/s depending on whether correlated double sampling (CDS) was enabled on the K3,

for a total dose of around 30 e⁻/Å². Nine grid holes were imaged for each stage movement using

the multiple record setup in *SerialEM*, with one collection per hole. At each hole, either a 2 s or 4

s exposure fractionated into either 40 or 80 frames was collected, depending on whether CDS

was enabled. Defocus values ranged from -1–-3 μm. See Table 1 for details.

Cryo-EM SPA image processing

All cryo-EM datasets were processed using a software pipeline consisting of *MotionCorr2*[76],

CTFFIND4[77], *RELION 3*[78,79], and *cryoSPARC*[57,80]. Early-stage processing was carried out in

RELION 3, with final 3D refinement and variability analysis carried out with *CryoSPARC*.

Dataset-specific processing schemes are presented in Supplementary Figs. 4–7 and

Supplementary Fig. 9. Generally, all processing proceeded as follows. Processing began in

RELION 3, with per-micrograph optics groups defined for a given dataset based on the nine

beam shift positions using the custom script *update_relion_tilt_class_og.py*. For example, the

hydrolyzing condition micrographs were collected in two sessions, so 18 total optics groups were

defined. The raw image stacks were dose-weighted and motion-corrected using the *RELION*

implementation of *MotionCorr2*, with seven patches along the *x* direction and five patches along

y. CTF parameters were estimated for each micrograph using *CTFFIND4* with a resolution range

of 3–30 Å, an FFT box size of 1024 px, and otherwise default parameters. Micrographs were

then selected based on maximum resolution and the CTF figure of merit.

Generally, an initial round of manual picking, extraction, and 2D classification were used to

assess data set quality and provide particles to reconstruct an initial 3D model. Good particles

from manual picking were then used for automated particle picking in *RELION*. Particles were

then extracted with a box size of 640 px, binned 8× (4.384 Å Å pixel size) to a box size of 80 px,

and subjected to at least one round of 2D classification to eliminate obviously bad particles.

Next, multiple rounds of 3D classification with a 280 Å mask diameter and a regularization

parameter $T = 4$ were performed over many iterations to determine the proper number of discrete

states in a given dataset, and to ensure that particles were classified properly. Convergence of

classification was monitored with a custom script, *startle.py*. Several classes slightly larger than

the number of discrete states were used, and classes that appeared to lie along a continuous

conformational coordinate were subsequently re-merged into a single class. In the case of the

hydrolyzing dataset, SNARE substrate-bound and SNARE substrate-free particles were split into

separate groups of particles, re-classified, and processed independently.

Following classification, particles were re-centered and re-extracted with a box size of 588 px

and binned 2× (1.096 Å pixel size) with a box size of 294 px. Following an initial round of 3D

refinement and post-processing, CTF refinement was performed. Typically, defocus,

astigmatism, B-factor, tilt, and trefoil were refined on a per-particle or per-micrograph basis,

depending on the specific class. Particles were then subjected to an additional round of 3D

refinement and post-processing, and optimal parameters for particle polishing were determined

by training on several classes, yielding different parameter sets. Particle polishing was then

performed with these sets, and 3D refinement and post-processing were used to assess the

results. Polished particles were then subjected to a final round of 3D refinement, post-processing,

and CTF refinement, during which anisotropic magnification was estimated. At this stage,

particles and 3D reconstructions were imported into *CryoSPARC* for a final round of non-

uniform refinement, which typically improved the quality of reconstructions.

In the case of one class observed under hydrolyzing conditions (c4-7-16-20-24), a large degree of

continuous heterogeneity was observed. 3D variability analysis (3DVA) revealed a trajectory of

protomer F nucleotide and rebinding, so particles in the first mode were split using a custom

script, *split_by_3dva.py*, as current tools to do so natively were unavailable at the time. This

script fits a normal distribution to the eigenvector magnitude distribution and allows slicing with

different cutoffs. For the 3DVA subclasses (slices O and P), particles in the ±1.5 σ tails of the

distribution were used for reconstruction.

These reconstructions were considered final for the overall 20S architecture, but further

processing was required to generate local reconstructions of the various α-SNAP—SNARE

subcomplexes. In the case of the sx20S dataset, only one class was observed with ordered spire

density, so local refinement was relatively simple. *CryoSPARC* signal subtraction

(*particle_subtract*; masks generated using *Segger* 2.5.3[81] in *Chimera* 1.15[82]) was used to remove

the signal from NSF N, D1, and D2 domains, leaving only the α-SNAP—syntaxin H3 SNARE

domain tetramer subcomplex. Local refinement (*new_local_refine*) was then performed on these

particles with a 40°/20 Å search extent and a custom fulcrum at the N-terminal end of the

SNARE bundle near the point of interaction with the F protomer of the D1 ring. The resulting

volumes were re-centered on the center of mass (*volume_alignment_tools*), and particle

alignments were updated as well (*align_3D*). Then, a final round of local refinement

(*new_local_refine*) was performed with the map center as the fulcrum (the default setting), a

smaller search extent of 10°/5 Å, and a maximum alignment resolution of 0.1°. Per-particle scales

were also reset, and minimization was performed over a per-particle scale. This produced the

final reconstruction of this subcomplex for subsequent modeling. In the case of the α-SNAP—

2:1 and α-SNAP—2:2 binary SNARE subcomplexes, this procedure was followed for each class

meeting certain criteria (number of particles, ordered spire, resolution), yielding five local

reconstructions for the α-SNAP—2:1 binary SNARE subcomplex and six local reconstructions

for the α-SNAP—2:2 binary SNARE subcomplex. In each case, these reconstructions were re-

centered (*volume_alignment_tools*), aligned to one another with the center of mass at the origin

(*align_3D)*, and a final round of local refinement was run to produce a final higher-resolution

reconstruction incorporating data from multiple classes. This produced the final reconstruction of

these subcomplexes for subsequent modeling. For more details about all data sets, see

Supplementary Figs. 4–7, Supplementary Fig. 9, Supplementary Tables 1–2, and Supplementary

Fig. 11.

Model building and refinement

All manual model building and refinements were carried out using *Coot*[83]. Semi-manual

refinement into low-resolution features was performed with *UCSF ChimeraX*[84] and the

ISOLDE[85] plugin. Automated refinement was performed using *phenix.real_space_refine*[86,87]. No

composite maps were used at any stage of model building or refinement.

First, models of the three α-SNAP—SNARE subcomplexes were built by rigid body fitting of

the best possible starting models into their corresponding densities using *UCSF ChimeraX*. In

each case, 3–4 copies of a high-quality α-SNAP model from the previous best 20S model (PDB

ID 6MDM[6]) were rigid body fit into the density. In the case of the syntaxin H3 domain tetramer

complex, four copies of the syntaxin chain from the high-resolution crystal structure (PDB ID

1N7S[54]) of the neuronal SNARE complex were docked into the density, with several

characteristic residues as fiducials. The quality of the local reconstructions of the SNARE—α-

SNAP spires allowed unique assignment of the identity of the SNARE domains, their

directionality, and register. In the case of the 2:1 binary SNARE complex, the syntaxin and

SNAP-25 components of the same crystal structure were docked into the density, but with the

SNARE domain of synaptobrevin replaced with an additional syntaxin H3 SNARE domain; key

side chain densities as well as a distinctive kink leading into the SNAP-25 SN2 domain guided

placement (Fig. 3C, Fig. 5A). In the case of the 2:2 binary SNARE complex, a previous crystal

structure (PDB ID 1JTH[34]) was docked into the density without modification. In all cases,

individual components were trimmed, or residues were added where supported by the density. A

round of *phenix.real_space_refine* was then performed with only rigid-body refinement enabled

for each component. Subsequent cycles of manual model building were performed in *Coot* and

ISOLDE followed by 1–7 macrocycles of *phenix.real_space_refine* with global minimization and

local grid search enabled. Ramachandran and secondary structure restraints were used

throughout. Note that while sequence information was included upon deposition for the full-

length construct used in the case of each SNARE, some regions (e.g., the syntaxin Habc domain,

the SNAP-25 SN2 domain in the case of the 2:2 binary SNARE complex) were not visible in the

reconstructions and were not modeled as a result.

Reference models of NSF were then built. Gold-standard models were first built into two high-

resolution NSF classes from a single non-hydrolyzing dataset with near-atomic resolutions using

the previous best model of NSF (PDB ID 6MDM[6]) as a starting model. The starting model was

first docked into cryo-EM density using ChimeraX. Next, a round of *phenix.real_space_refine*

was performed with only rigid-body refinement enabled; each large and small subdomain of each

protomer was treated as a single rigid body (twelve total). Nucleotides and water molecules were

then added where appropriate. Subsequent cycles of manual model building were performed in

Coot and *ISOLDE* followed by 1–7 macrocycles of *phenix.real_space_refine* with global

minimization and local grid search enabled. Ramachandran and secondary structure restraints

were used throughout.

The previously described reference models were then used as starting models in building the

sx20S and bin20S supercomplexes. In sx20S and bin20S reconstructions with well-defined N-

domains (i.e., a recognizable two-lobed density), one or more instances of the re-processed

model derived from the crystal structure (PDB ID 1QCS[88]) as re-processed by *PDB_REDO*[89]

were rigid body fit into density before refinement. Next, the appropriate α-SNAP—SNARE

subcomplex was docked into the density, completing the spire structure. Spires in the global 20S

reconstructions were generally heterogeneous; although secondary structure could often be

distinguished, side chain density could not, leading to poor map-model correlations in these

regions. A round of *phenix.real_space_refine* was performed with only rigid-body refinement

enabled; each large and small subdomain of each protomer, each N domain, and each α-SNAP—

SNARE subcomplex was treated as a rigid body (for a 20S particle with four engaged N-

domains, this would be 17 total rigid body groups). Nucleotides, Mg^{2+}, P_i, and water molecules

were then added where appropriate. Subsequent cycles of manual model building were

performed in *Coot* and *ISOLDE* followed by 1–7 macrocycles of *phenix.real_space_refine* with

global minimization and local grid search enabled. Ramachandran and secondary structure

restraints were used throughout.

All model figures were prepared with *UCSF ChimeraX*, in many cases with custom CXC scripts

generated by the Python scripts described below.

Automated NSF model analysis and coordinate PCA

Several custom Python scripts (*preprocess_models.py, analyze_models.py*) using the *Bio.PDB*

module[90] of the Biopython project[91] as well as *pandas*[71], *numpy*[72], *scipy*[73], *scikit-learn*[92], and

matplotlib[74] were used to standardize all 25 NSF, sx20S, or bin20S models for further analysis,

such as to perform principle components analysis (PCA) and HDBSCAN clustering, to measure

NSF D1 large-small subdomain angles, to perform planarity fits for NSF domains and

subdomains, to measure distances between key catalytic residues and nucleotides, and to map

various values to structures for visualization in *UCSF ChimeraX*.

For PCA, 132 NSF D1 domains from 22 models over all datasets were split into large and small

subdomains, which were then aligned to one another (large, residues 224–373; small, residues

400–486) using *Bio.PDB.Superimposer*. Splitting removes the largest source of variance in the

data, the rigid body motions of the two subdomains relative to each other, which is considered

trivial for the purposes of this analysis. Next, a Cα coordinate matrix $\mathbf{C}$ with dimensions n_{models} ×

$n_{residues}$ × 3 was constructed for residues shared by all models and flattened into two dimensions,

creating $\mathbf{C_{flat}}$ with n_{models} rows and $n_{residues}$ × 3 columns. $\mathbf{C_{flat}}$ was then rescaled using

sklearn.preprocessing.StandardScaler and PCA was performed on $\mathbf{C_{flat}}$ using

sklearn.decomposition.PCA. 13 components were found to contribute 99% of the variance, so a

transformed matrix $\mathbf{C_{trans}}$ with dimensions n_{models} × 13, with eigenvectors sorted along the second

axis by eigenvalue, was calculated. The cluster analysis of models was then performed using

sklearn.cluster.HDBSCAN with a minimum cluster size of three, producing 11 clusters

accounting for 119/132 total models. This analysis serves as the basis for Fig. 8A.

For NSF D1 and D2 plane fits, centers of mass for all subdomains were calculated, and the

planes for different combinations of these points were fit using a least squares approach

(Supplementary Fig. 8). For NSF D1 large-small subdomain angle calculations, the primary axes

of two α-helices in the large (residues 350–358) and small subdomains (residues 442–452) were

calculated by singular value decomposition using *np.linalg.svd*, with implementation inspired by

helix_analysis.py from the MDAnalysis package[93]. Vectors corresponding to these helical axes *u*

and *v* were normalized and centered at the coordinate origin, and the angle between them was

calculated as $\theta = 2 \times \tan^{-1}\left(\sqrt{\sum_i \vec{u}_i^2}\sqrt{\sum_i \vec{v}_i^2}\right)$.

Distances between key active site atoms over all models (Fig. 9) were calculated using

np.linalg.norm, where the norm of the difference vector of the coordinates of each atom is the

distance between them in Å. Significant changes in protomer E active site residues relative to the

other five protomers were assessed by pairwise application of the Welch's t-test assuming non-

equal variances as implemented in *scipy.stats.ttest_ind*. Distance distributions were considered

statistically different if $p < 0.05$.

All these values, as well as manually curated, qualitative data labels, were plotted in various

ways using *matplotlib* or incorporated into automatically generated UCSF *ChimeraX* CXC

scripts.

Data availability

The atomic coordinates have been deposited in the Protein Data Bank (PDB) under accession

codes 9OJ2 [http://doi.org/10.2210/pdb9oj2/pdb] (sx20S complex (NSF-alphaSNAP-syntaxin-

1a), non-hydrolyzing, class 1); 9OJJ [http://doi.org/10.2210/pdb9ojj/pdb] (sx20S complex (NSF-

alphaSNAP-syntaxin-1a), alphaSNAP-syntaxin-1a subcomplex local refinement, non-

hydrolyzing, class 1); 9OJR [http://doi.org/10.2210/pdb9ojr/pdb] (21bin20S complex (NSF-

alphaSNAP-2:1 syntaxin-1a:SNAP-25), non-hydrolyzing, class 3); 9OJU

[http://doi.org/10.2210/pdb9oju/pdb] (21bin20S complex (NSF-alphaSNAP-2:1 syntaxin-

1a:SNAP-25), non-hydrolyzing, class 4); 9OJZ [http://doi.org/10.2210/pdb9ojz/pdb] (21bin20S

complex (NSF-alphaSNAP-2:1 syntaxin-1a:SNAP-25), non-hydrolyzing, class 5); 9OK3

[http://doi.org/10.2210/pdb9ok3/pdb] (21bin20S complex (NSF-alphaSNAP-2:1 syntaxin-

1a:SNAP-25), 3:2:1 alphaSNAP-syntaxin-1a-SNAP-25 subcomplex local refinement, non-

hydrolyzing, class 13); 9OK5 [http://doi.org/10.2210/pdb9ok5/pdb] (22bin20S complex (NSF-

alphaSNAP-2:2 syntaxin-1a:SNAP-25), hydrolyzing, class 16); 9OKC

[http://doi.org/10.2210/pdb9okc/pdb] (22bin20S complex (NSF-alphaSNAP-2:2 syntaxin-

1a:SNAP-25), hydrolyzing, class 17); 9OLJ [http://doi.org/10.2210/pdb9olj/pdb] (22bin20S

complex (NSF-alphaSNAP-2:2 syntaxin-1a:SNAP-25), hydrolyzing, class 18); 9OLO

[http://doi.org/10.2210/pdb9olo/pdb] (22bin20S complex (NSF-alphaSNAP-2:2 syntaxin-

1a:SNAP-25), hydrolyzing, class 19); 9OM6 [http://doi.org/10.2210/pdb9om6/pdb] (22bin20S

complex (NSF-alphaSNAP-2:2 syntaxin-1a:SNAP-25), 4:2:2 alphaSNAP-syntaxin-1a-SNAP-25

subcomplex local refinement, hydrolyzing, class 23); 9OMQ

[http://doi.org/10.2210/pdb9omq/pdb] (NSF, substrate free, hydrolyzing, class 24); 9PAF

[http://doi.org/10.2210/pdb9paf/pdb] (21bin20S complex (NSF-alphaSNAP-2:1 syntaxin-

1a:SNAP-25), non-hydrolyzing, class 6); 9PAG [http://doi.org/10.2210/pdb9pag/pdb] (21bin20S

complex (NSF-alphaSNAP-2:1 syntaxin-1a:SNAP-25), non-hydrolyzing, class 7); 9PB9

[http://doi.org/10.2210/pdb9pb9/pdb] (21bin20S complex (NSF-alphaSNAP-2:1 syntaxin-

1a:SNAP-25), non-hydrolyzing, class 8); 9PBA [http://doi.org/10.2210/pdb9pba/pdb] (21bin20S

complex (NSF-alphaSNAP-2:1 syntaxin-1a:SNAP-25), non-hydrolyzing, class 9); 9PBF

[http://doi.org/10.2210/pdb9pbf/pdb] (21bin20S complex (NSF-alphaSNAP-2:1 syntaxin-

1a:SNAP-25), non-hydrolyzing, class 10); 9PBV [http://doi.org/10.2210/pdb9pbv/pdb]

(21bin20S complex (NSF-alphaSNAP-2:1 syntaxin-1a:SNAP-25), non-hydrolyzing, class 11);

9PC3 [http://doi.org/10.2210/pdb9pc3/pdb] (21bin20S complex (NSF-alphaSNAP-2:1 syntaxin-

1a:SNAP-25), non-hydrolyzing, class 12); 9PCX [http://doi.org/10.2210/pdb9pcx/pdb]

(22bin20S complex (NSF-alphaSNAP-2:2 syntaxin-1a:SNAP-25), hydrolyzing, class 14); 9PCZ

[http://doi.org/10.2210/pdb9pcz/pdb] (22bin20S complex (NSF-alphaSNAP-2:2 syntaxin-

1a:SNAP-25), hydrolyzing, class 15); 9PD1 [http://doi.org/10.2210/pdb9pd1/pdb] (22bin20S

complex (NSF-alphaSNAP-2:2 syntaxin-1a:SNAP-25), hydrolyzing, class 20); 9PD8

[http://doi.org/10.2210/pdb9pd8/pdb] (22bin20S complex (NSF-alphaSNAP-2:2 syntaxin-

1a:SNAP-25), hydrolyzing, class 21); 9PDB [http://doi.org/10.2210/pdb9pdb/pdb] (22bin20S

complex (NSF-alphaSNAP-2:2 syntaxin-1a:SNAP-25), hydrolyzing, class 22); 9PDD

[http://doi.org/10.2210/pdb9pdd/pdb] (22bin20S complex (NSF-alphaSNAP-2:2 syntaxin-

1a:SNAP-25), hydrolyzing, class 29); 9PF2 [http://doi.org/10.2210/pdb9pf2/pdb] (NSF, substrate

free, hydrolyzing, class 25); 9PFC [http://doi.org/10.2210/pdb9pfc/pdb] (NSF, substrate free,

hydrolyzing, class 26); 9PFF [http://doi.org/10.2210/pdb9pff/pdb] (min22bin20S complex (NSF-

alphaSNAP-2:2 syntaxin-1a H3:SNAP-25 SN1), non-hydrolyzing, class 27); and 9PFG

[http://doi.org/10.2210/pdb9pfg/pdb] (min22bin20S complex (NSF-alphaSNAP-2:2 syntaxin-1a

H3:SNAP-25 SN1), 4:2:2 alphaSNAP-syntaxin-1a H3-SNAP-25 SN1 subcomplex local

refinement, non-hydrolyzing, class 28). Previously published models used in this work include

6MDM [http://doi.org/10.2210/pdb6mdm/pdb] (The 20S supercomplex engaging the SNAP-25

N-terminus (class 1)); 1N7S [http://doi.org/10.2210/pdb1n7s/pdb] (High Resolution Structure of

a Truncated Neuronal SNARE Complex); 1QCS [http://doi.org/10.2210/pdb1qcs/pdb] (N-

TERMINAL DOMAIN OF N-ETHYLMALEIMIDE SENSITIVE FACTOR (NSF)); 1JTH

[http://doi.org/10.2210/pdb1jth/pdb] (Crystal structure and biophysical properties of a complex

between the N-terminal region of SNAP25 and the SNARE region of syntaxin 1a); and 1HVV

[http://doi.org/10.2210/pdb1hvv/pdb] (SELF-ASSOCIATION OF THE H3 REGION OF

SYNTAXIN 1A).

Code Availability

We used *AlphaFold2*[94], *ChimeraX 1.6–1.8*[84], *Coot 0.9*[83], *CryoSPARC*[57,80], *CTFFIND4*[77],

ImageJ[67], *ISOLDE*[85], *MolProbity*[95], *MotionCorr2*[76], *PHENIX 1.2.X*[86,87,96], *Prism* (GraphPad

Software), *RELION 3*[78], *Segger*[81], and *SerialEM 4*[75]. Custom scripts were written in *Python 3.1*

with packages including *numpy*[72], *scipy*[73], *scikit-learn*[92], *biopython*[90,91], *pandas*[71], and

matplotlib[74] and are available at https://github.com/brungerlab/kiw_scripts.

References

1. Rizo, J., and Südhof, T.C. (2012). The membrane fusion enigma: SNAREs, Sec1/Munc18
proteins, and their accomplices--guilty as charged? Annu. Rev. Cell Dev. Biol. *28*, 279–308.
https://doi.org/10.1146/annurev-cellbio-101011-155818.

2. Brunger, A.T., Choi, U.B., Lai, Y., Leitz, J., and Zhou, Q. (2018). Molecular Mechanisms of
Fast Neurotransmitter Release. Annu. Rev. Biophys. *47*, 469–497.
https://doi.org/10.1146/annurev-biophys-070816-034117.

3. Leitz, J., Wang, C., Esquivies, L., Pfuetzner, R.A., Peters, J.J., Couoh-Cardel, S., Wang,
A.L., and Brunger, A.T. (2024). Beyond the MUN domain, Munc13 controls priming and
depriming of synaptic vesicles. Cell Rep. *43*. https://doi.org/10.1016/j.celrep.2024.114026.

4. Brunger, A.T., Choi, U.B., Lai, Y., Leitz, J., White, K.I., and Zhou, Q. (2019). The pre-
synaptic fusion machinery. Curr. Opin. Struct. Biol. *54*, 179–188.
https://doi.org/10.1016/j.sbi.2019.03.007.

5. Rizo, J. (2022). Molecular Mechanisms Underlying Neurotransmitter Release. Annu. Rev.
Biophys. *51*, 377–408. https://doi.org/10.1146/annurev-biophys-111821-104732.

6. White, K.I., Zhao, M., Choi, U.B., Pfuetzner, R.A., and Brunger, A.T. (2018). Structural
principles of SNARE complex recognition by the AAA+ protein NSF. eLife *7*, e38888.
https://doi.org/10.7554/eLife.38888.

7. Zhao, M., Wu, S., Zhou, Q., Vivona, S., Cipriano, D.J., Cheng, Y., and Brunger, A.T.
(2015). Mechanistic insights into the recycling machine of the SNARE complex. Nature *518*,
61–67. https://doi.org/10.1038/nature14148.

8. Rickman, C., Medine, C.N., Dun, A.R., Moulton, D.J., Mandula, O., Halemani, N.D.,
Rizzoli, S.O., Chamberlain, L.H., and Duncan, R.R. (2010). t-SNARE Protein
Conformations Patterned by the Lipid Microenvironment. J. Biol. Chem. *285*, 13535–13541.
https://doi.org/10.1074/jbc.M109.091058.

9. Murray, D.H., and Tamm, L.K. (2009). Clustering of Syntaxin-1A in Model Membranes is
Modulated by Phosphatidylinositol-4,5-bisphosphate and Cholesterol. Biochemistry *48*,
4617–4625. https://doi.org/10.1021/bi9003217.

10. Aoyagi, K., Sugaya, T., Umeda, M., Yamamoto, S., Terakawa, S., and Takahashi, M. (2005).
The activation of exocytotic sites by the formation of phosphatidylinositol 4,5-bisphosphate
microdomains at syntaxin clusters. J. Biol. Chem. *280*, 17346–17352.
https://doi.org/10.1074/jbc.M413307200.

11. Lang, T., Bruns, D., Wenzel, D., Riedel, D., Holroyd, P., Thiele, C., and Jahn, R. (2001).
SNAREs are concentrated in cholesterol-dependent clusters that define docking and fusion
sites for exocytosis. EMBO J. *20*, 2202–2213. https://doi.org/10.1093/emboj/20.9.2202.

12. Emperador-Melero, J., and Kaeser, P.S. (2020). Assembly of the presynaptic active zone.
Curr. Opin. Neurobiol. *63*, 95–103. https://doi.org/10.1016/j.conb.2020.03.008.

13. Sieber, J.J., Willig, K.I., Kutzner, C., Gerding-Reimers, C., Harke, B., Donnert, G.,
Rammner, B., Eggeling, C., Hell, S.W., Grubmüller, H., et al. (2007). Anatomy and
Dynamics of a Supramolecular Membrane Protein Cluster. Science *317*, 1072–1076.
https://doi.org/10.1126/science.1141727.

14. Weimbs, T., Low, S.H., Chapin, S.J., Mostov, K.E., Bucher, P., and Hofmann, K. (1997). A
conserved domain is present in different families of vesicular fusion proteins:
A new superfamily. Proc. Natl. Acad. Sci. U. S. A. *94*, 3046–3051.

15. Bennett, M.K., Calakos, N., and Scheller, R.H. (1992). Syntaxin: A Synaptic Protein
Implicated in Docking of Synaptic Vesicles at Presynaptic Active Zones. Science *257*, 255–
259. https://doi.org/10.1126/science.1321498.

16. Sutton, R.B., Fasshauer, D., Jahn, R., and Brunger, A.T. (1998). Crystal structure of a
SNARE complex involved in synaptic exocytosis at 2.4 Å resolution. Nature *395*, 347–353.
https://doi.org/10.1038/26412.

17. Fernandez, I., Ubach, J., Dulubova, I., Zhang, X., Südhof, T.C., and Rizo, J. (1998). Three-
Dimensional Structure of an Evolutionarily Conserved N-Terminal Domain of Syntaxin 1A.
Cell *94*, 841–849. https://doi.org/10.1016/s0092-8674(00)81742-0.

18. Bar-On, D., Wolter, S., Van De Linde, S., Heilemann, M., Nudelman, G., Nachliel, E.,
Gutman, M., Sauer, M., and Ashery, U. (2012). Super-resolution Imaging Reveals the
Internal Architecture of Nano-sized Syntaxin Clusters. J. Biol. Chem. *287*, 27158–27167.
https://doi.org/10.1074/jbc.M112.353250.

19. Knowles, M.K., Barg, S., Wan, L., Midorikawa, M., Chen, X., and Almers, W. (2010).
Single secretory granules of live cells recruit syntaxin-1 and synaptosomal associated protein
25 (SNAP-25) in large copy numbers. Proc. Natl. Acad. Sci. *107*, 20810–20815.
https://doi.org/10.1073/pnas.1014840107.

20. Wan, L., Chen, X., and Almers, W. (2021). Munc18-dependent and -independent clustering
of syntaxin in the plasma membrane of cultured endocrine cells. Proc. Natl. Acad. Sci. *118*,
e2025748118. https://doi.org/10.1073/pnas.2025748118.

21. Ullrich, A., Böhme, M.A., Schöneberg, J., Depner, H., Sigrist, S.J., and Noé, F. (2015).
Dynamical Organization of Syntaxin-1A at the Presynaptic Active Zone. PLOS Comput.
Biol. *11*, e1004407. https://doi.org/10.1371/journal.pcbi.1004407.

22. Barg, S., Knowles, M.K., Chen, X., Midorikawa, M., and Almers, W. (2010). Syntaxin
clusters assemble reversibly at sites of secretory granules in live cells. Proc. Natl. Acad. Sci.
107, 20804–20809. https://doi.org/10.1073/pnas.1014823107.

23. Sakamoto, H., Ariyoshi, T., Kimpara, N., Sugao, K., Taiko, I., Takikawa, K., Asanuma, D.,
Namiki, S., and Hirose, K. (2018). Synaptic weight set by Munc13-1 supramolecular
assemblies. Nat. Neurosci. *21*, 41–49. https://doi.org/10.1038/s41593-017-0041-9.

24. Pertsinidis, A., Mukherjee, K., Sharma, M., Pang, Z.P., Park, S.R., Zhang, Y., Brunger, A.T.,
Südhof, T.C., and Chu, S. (2013). Ultrahigh-resolution imaging reveals formation of
neuronal SNARE/Munc18 complexes in situ. Proc. Natl. Acad. Sci. *110*, E2812–E2820.
https://doi.org/10.1073/pnas.1310654110.

25. Yin, P., Gandasi, N.R., Arora, S., Omar-Hmeadi, M., Saras, J., and Barg, S. (2018). Syntaxin
clusters at secretory granules in a munc18-bound conformation. Mol. Biol. Cell *29*, 2700–
2708. https://doi.org/10.1091/mbc.E17-09-0541.

26. Lang, T. (2007). SNARE proteins and 'membrane rafts.' J. Physiol. *585*, 693–698.
https://doi.org/10.1113/jphysiol.2007.134346.

27. Bademosi, A.T., Lauwers, E., Padmanabhan, P., Odierna, L., Chai, Y.J., Papadopulos, A.,
Goodhill, G.J., Verstreken, P., van Swinderen, B., and Meunier, F.A. (2016). In vivo single-
molecule imaging of syntaxin1A reveals polyphosphoinositide- and activity-dependent
trapping in presynaptic nanoclusters. Nat. Commun. *7*, 13660.
https://doi.org/10.1038/ncomms13660.

28. Sieber, J.J., Willig, K.I., Heintzmann, R., Hell, S.W., and Lang, T. (2006). The SNARE
Motif Is Essential for the Formation of Syntaxin Clusters in the Plasma Membrane. Biophys.
J. *90*, 2843–2851. https://doi.org/10.1529/biophysj.105.079574.

29. Hanson, P.I., Otto, H., Barton, N., and Jahn, R. (1995). The N-Ethylmaleimide-sensitive
Fusion Protein and α-SNAP Induce a Conformational Change in Syntaxin (*). J. Biol. Chem.
270, 16955–16961. https://doi.org/10.1074/jbc.270.28.16955.

30. Stepien, K.P., Prinslow, E.A., and Rizo, J. (2019). Munc18-1 is crucial to overcome the
inhibition of synaptic vesicle fusion by αSNAP. Nat. Commun. *10*, 4326.
https://doi.org/10.1038/s41467-019-12188-4.

31. Ma, C., Su, L., Seven, A.B., Xu, Y., and Rizo, J. (2013). Reconstitution of the vital functions
of Munc18 and Munc13 in neurotransmitter release. Science *339*, 421–425.
https://doi.org/10.1126/science.1230473.

32. Choi, U.B., Zhao, M., White, K.I., Pfuetzner, R.A., Esquivies, L., Zhou, Q., and Brunger,
A.T. (2018). NSF-mediated disassembly of on- and off-pathway SNARE complexes and
inhibition by complexin. eLife *7*, e36497. https://doi.org/10.7554/eLife.36497.

33. Fasshauer, D., Bruns, D., Shen, B., Jahn, R., and Brunger, A.T. (1997). A Structural Change
Occurs upon Binding of Syntaxin to SNAP-25. J. Biol. Chem. *272*, 4582–4590.
https://doi.org/10.1074/jbc.272.7.4582.

34. Misura, K.M.S., Gonzalez, L.C., May, A.P., Scheller, R.H., and Weis, W.I. (2001). Crystal
Structure and Biophysical Properties of a Complex between the N-terminal SNARE Region
of SNAP25 and Syntaxin 1a *. J. Biol. Chem. *276*, 41301–41309.
https://doi.org/10.1074/jbc.M106853200.

35. Weninger, K., Bowen, M.E., Choi, U.B., Chu, S., and Brunger, A.T. (2008). Accessory
Proteins Stabilize the Acceptor Complex for Synaptobrevin, the 1:1 Syntaxin/SNAP-25
Complex. Structure *16*, 308–320. https://doi.org/10.1016/j.str.2007.12.010.

36. Hesselbarth, J., and Schmidt, C. (2023). Mass spectrometry uncovers intermediates and off-
pathway complexes for SNARE complex assembly. Commun. Biol. *6*, 1–15.
https://doi.org/10.1038/s42003-023-04548-0.

37. Hayashi, T., McMahon, H., Yamasaki, S., Binz, T., Hata, Y., Südhof, T.C., and Niemann, H.
(1994). Synaptic vesicle membrane fusion complex: action of clostridial neurotoxins on
assembly. EMBO J. *13*, 5051–5061. https://doi.org/10.1002/j.1460-2075.1994.tb06834.x.

38. Gao, Y., Khan, Y.A., Mo, W., White, K.I., Perkins, M., Pfuetzner, R.A., Trapani, J.G.,
Brunger, A.T., and Nicolson, T. (2023). Sensory deficit screen identifies nsf mutation that
differentially affects SNARE recycling and quality control. Cell Rep. *42*.
https://doi.org/10.1016/j.celrep.2023.112345.

39. Hayashi, T., Yamasaki, S., Nauenburg, S., Binz, T., and Niemann, H. (1995). Disassembly of
the reconstituted synaptic vesicle membrane fusion complex in vitro. EMBO J. *14*, 2317–
2325. https://doi.org/10.1002/j.1460-2075.1995.tb07226.x.

40. Vivona, S., Cipriano, D.J., O'Leary, S., Li, Y.H., Fenn, T.D., and Brunger, A.T. (2013).
Disassembly of All SNARE Complexes by N-Ethylmaleimide-sensitive Factor (NSF) Is
Initiated by a Conserved 1:1 Interaction between α-Soluble NSF Attachment Protein (SNAP)
and SNARE Complex. J. Biol. Chem. *288*, 24984–24991.
https://doi.org/10.1074/jbc.M113.489807.

42. Nagiec, E.E., Bernstein, A., and Whiteheart, S.W. (1995). Each Domain of the N-
Ethylmaleimide-sensitive Fusion Protein Contributes to Its Transport Activity (*). J. Biol.
Chem. *270*, 29182–29188. https://doi.org/10.1074/jbc.270.49.29182.

43. Matveeva, E.A., He, P., and Whiteheart, S.W. (1997). N-Ethylmaleimide-sensitive Fusion
Protein Contains High and Low Affinity ATP-binding Sites That Are Functionally Distinct
*. J. Biol. Chem. *272*, 26413–26418. https://doi.org/10.1074/jbc.272.42.26413.

44. Donovan, K.W., and Bretscher, A. (2015). Tracking individual secretory vesicles during
exocytosis reveals an ordered and regulated process. J. Cell Biol. *210*, 181–189.
https://doi.org/10.1083/jcb.201501118.

45. Khan, Y.A., White, K.I., Pfuetzner, R.A., Singal, B., Esquivies, L., Mckenzie, G., Liu, F.,
DeLong, K., Choi, U.B., Montabana, E., et al. (2025). SNARE disassembly requires
Sec18/NSF side loading. Nat. Struct. Mol. Biol., 1–13. https://doi.org/10.1038/s41594-025-
01590-w.

46. Stepien, K.P., Xu, J., Zhang, X., Bai, X.-C., and Rizo, J. (2022). SNARE assembly
enlightened by cryo-EM structures of a synaptobrevin–Munc18-1–syntaxin-1 complex. Sci.
Adv. *8*, eabo5272. https://doi.org/10.1126/sciadv.abo5272.

47. Dawidowski, D., and Cafiso, D.S. (2013). Allosteric Control of Syntaxin 1a by Munc18-1:
Characterization of the Open and Closed Conformations of Syntaxin. Biophys. J. *104*, 1585.
https://doi.org/10.1016/j.bpj.2013.02.004.

48. Pei, Q., Chen, Q., Tian, Z., Zhu, L., Chen, Y., Gong, J., Wang, S., Xiang, Y., Khamo, J., Fan,
J., et al. (2024). Munc18 modulates syntaxin phase separation for exocytosis.

49. Meijer, M., Burkhardt, P., de Wit, H., Toonen, R.F., Fasshauer, D., and Verhage, M. (2012).
Munc18-1 mutations that strongly impair SNARE-complex binding support normal synaptic
transmission. EMBO J. *31*, 2156–2168. https://doi.org/10.1038/emboj.2012.72.

50. Xiao, W., Poirier, M.A., Bennett, M.K., and Shin, Y.-K. (2001). The neuronal t-SNARE
complex is a parallel four-helix bundle. Nat. Struct. Biol. *8*, 308–311.
https://doi.org/10.1038/86174.

51. Margittai, M., Fasshauer, D., Pabst, S., Jahn, R., and Langen, R. (2001). Homo- and
Heterooligomeric SNARE Complexes Studied by Site-directed Spin Labeling *. J. Biol.
Chem. *276*, 13169–13177. https://doi.org/10.1074/jbc.M010653200.

52. Lai, Y., Choi, U.B., Leitz, J., Rhee, H.J., Lee, C., Altas, B., Zhao, M., Pfuetzner, R.A.,
Wang, A.L., Brose, N., et al. (2017). Molecular Mechanisms of Synaptic Vesicle Priming by
Munc13 and Munc18. Neuron *95*, 591-607.e10.
https://doi.org/10.1016/j.neuron.2017.07.004.

53. Sanyal, S., and Krishnan, K.S. (2001). Lethal comatose mutation in Drosophila reveals
possible role for NSF in neurogenesis. NeuroReport *12*, 1363.

54. Ernst, J.A., and Brunger, A.T. (2003). High Resolution Structure, Stability, and
Synaptotagmin Binding of a Truncated Neuronal SNARE Complex *. J. Biol. Chem. *278*,
8630–8636. https://doi.org/10.1074/jbc.M211889200.

55. Pribicevic, S., Graham, A.C., Cafiso, D.S., Pérez-Lara, Á., and Jahn, R. (2024). Intermediate
steps in the formation of neuronal SNARE complexes. J. Biol. Chem. *300*, 107591.
https://doi.org/10.1016/j.jbc.2024.107591.

56. de la Peña, A.H., Goodall, E.A., Gates, S.N., Lander, G.C., and Martin, A. (2018). Substrate-
engaged 26S proteasome structures reveal mechanisms for ATP-hydrolysis–driven
translocation. Science *362*, eaav0725. https://doi.org/10.1126/science.aav0725.

57. Punjani, A., and Fleet, D.J. (2021). 3D variability analysis: Resolving continuous flexibility
and discrete heterogeneity from single particle cryo-EM. J. Struct. Biol. *213*, 107702.
https://doi.org/10.1016/j.jsb.2021.107702.

58. Smith, G.R., Contreras-Moreira, B., Zhang, X., and Bates, P.A. (2004). A link between
sequence conservation and domain motion within the AAA+ family. J. Struct. Biol. *146*,
189–204. https://doi.org/10.1016/j.jsb.2003.11.022.

59. Zhou, H., Di Palma, S., Preisinger, C., Peng, M., Polat, A.N., Heck, A.J.R., and Mohammed,
S. (2013). Toward a Comprehensive Characterization of a Human Cancer Cell
Phosphoproteome. J. Proteome Res. *12*, 260–271. https://doi.org/10.1021/pr300630k.

60. Matveeva, E.A., Whiteheart, S.W., Vanaman, T.C., and Slevin, J.T. (2001). Phosphorylation
of the N-Ethylmaleimide-sensitive Factor Is Associated with Depolarization-dependent
Neurotransmitter Release from Synaptosomes. J. Biol. Chem. *276*, 12174–12181.
https://doi.org/10.1074/jbc.m007394200.

61. Huang, X., Sun, S., Wang, X., Fan, F., Zhou, Q., Lu, S., Cao, Y., Wang, Q.-W., Dong, M.-
Q., Yao, J., et al. (2019). Mechanistic insights into the SNARE complex disassembly. Sci.
Adv. *5*, eaau8164. https://doi.org/10.1126/sciadv.aau8164.

62. Mertins, J., Finke, J., Sies, R., Rink, K.M., Hasenauer, J., and Lang, T. (2021). The
mesoscale organization of syntaxin 1A and SNAP25 is determined by SNARE–SNARE
interactions. eLife *10*, e69236. https://doi.org/10.7554/eLife.69236.

63. Kraichely, K.N., Sandall, C.R., Liang, B., Kiessling, V., and Tamm, L.K. (2025).
Functionally distinct SNARE motifs of SNAP25 cooperate in SNARE assembly and
membrane fusion. Biophys. J. *124*, 637–650. https://doi.org/10.1016/j.bpj.2024.12.034.

64. Zhou, Q., Zhou, P., Wang, A.L., Wu, D., Zhao, M., Südhof, T.C., and Brunger, A.T. (2017).
The primed SNARE–complexin–synaptotagmin complex for neuronal exocytosis. Nature
548, 420–425. https://doi.org/10.1038/nature23484.

65. Zhou, Q., Lai, Y., Bacaj, T., Zhao, M., Lyubimov, A.Y., Uervirojnangkoorn, M., Zeldin,
O.B., Brewster, A.S., Sauter, N.K., Cohen, A.E., et al. (2015). Architecture of the
synaptotagmin–SNARE machinery for neuronal exocytosis. Nature *525*, 62–67.
https://doi.org/10.1038/nature14975.

66. Khan, Y.A., White, K.I., and Brunger, A.T. (2022). The AAA+ superfamily: a review of the
structural and mechanistic principles of these molecular machines. Crit. Rev. Biochem. Mol.
Biol. *57*, 156–187. https://doi.org/10.1080/10409238.2021.1979460.

67. Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years
of image analysis. Nat. Methods *9*, 671–675. https://doi.org/10.1038/nmeth.2089.

68. Yofe, I., Weill, U., Meurer, M., Chuartzman, S., Zalckvar, E., Goldman, O., Ben-Dor, S.,
Schütze, C., Wiedemann, N., Knop, M., et al. (2016). One library to make them all:
streamlining the creation of yeast libraries via a SWAp-Tag strategy. Nat. Methods *13*, 371–
378. https://doi.org/10.1038/nmeth.3795.

69. Cipriano, D.J., Jung, J., Vivona, S., Fenn, T.D., Brunger, A.T., and Bryant, Z. (2013).
Processive ATP-driven Substrate Disassembly by the N-Ethylmaleimide-sensitive Factor
(NSF) Molecular Machine♦. J. Biol. Chem. *288*, 23436–23445.
https://doi.org/10.1074/jbc.M113.476705.

70. Studier, F.W. (2005). Protein production by auto-induction in high-density shaking cultures.
Protein Expr. Purif. *41*, 207–234. https://doi.org/10.1016/j.pep.2005.01.016.

71. McKinney, W. (2010). Data Structures for Statistical Computing in Python. In Proceedings
of the 9th Python in Science Confference, pp. 56–61. https://doi.org/10.25080/Majora-
92bf1922-00a.

72. Harris, C.R., Millman, K.J., van der Walt, S.J., Gommers, R., Virtanen, P., Cournapeau, D.,
Wieser, E., Taylor, J., Berg, S., Smith, N.J., et al. (2020). Array programming with NumPy.
Nature *585*, 357–362. https://doi.org/10.1038/s41586-020-2649-2.

73. Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D.,
Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0: fundamental
algorithms for scientific computing in Python. Nat. Methods *17*, 261–272.
https://doi.org/10.1038/s41592-019-0686-2.

74. Hunter, J.D. (2007). Matplotlib: A 2D Graphics Environment. Comput. Sci. Eng. *9*, 90–95.
https://doi.org/10.1109/MCSE.2007.55.

75. Mastronarde, D.N. (2005). Automated electron microscope tomography using robust
prediction of specimen movements. J. Struct. Biol. *152*, 36–51.
https://doi.org/10.1016/j.jsb.2005.07.007.

76. Zheng, S.Q., Palovcak, E., Armache, J.-P., Verba, K.A., Cheng, Y., and Agard, D.A. (2017).
MotionCor2 - anisotropic correction of beam-induced motion for improved cryo-electron
microscopy. Nat. Methods *14*, 331–332. https://doi.org/10.1038/nmeth.4193.

77. Rohou, A., and Grigorieff, N. (2015). CTFFIND4: Fast and accurate defocus estimation from
electron micrographs. J. Struct. Biol. *192*, 216–221.
https://doi.org/10.1016/j.jsb.2015.08.008.

78. Scheres, S.H.W. (2012). RELION: Implementation of a Bayesian approach to cryo-EM
structure determination. J. Struct. Biol. *180*, 519–530.
https://doi.org/10.1016/j.jsb.2012.09.006.

79. Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E., and
Scheres, S.H. (2018). New tools for automated high-resolution cryo-EM structure
determination in RELION-3. eLife *7*, e42166. https://doi.org/10.7554/eLife.42166.

80. Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017). cryoSPARC:
algorithms for rapid unsupervised cryo-EM structure determination. Nat. Methods *14*, 290–
296. https://doi.org/10.1038/nmeth.4169.

81. Pintilie, G.D., Zhang, J., Goddard, T.D., Chiu, W., and Gossard, D.C. (2010). Quantitative
analysis of cryo-EM density map segmentation by watershed and scale-space filtering, and
fitting of structures by alignment to regions. J. Struct. Biol. *170*, 427–438.
https://doi.org/10.1016/j.jsb.2010.03.007.

82. Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C.,
and Ferrin, T.E. (2004). UCSF Chimera—A visualization system for exploratory research
and analysis. J. Comput. Chem. *25*, 1605–1612. https://doi.org/10.1002/jcc.20084.

83. Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and development of
Coot. Acta Crystallogr. D Biol. Crystallogr. *66*, 486–501.
https://doi.org/10.1107/S0907444910007493.

84. Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H., and
Ferrin, T.E. (2018). UCSF ChimeraX: Meeting modern challenges in visualization and
analysis. Protein Sci. *27*, 14–25. https://doi.org/10.1002/pro.3235.

85. Croll, T.I. (2018). ISOLDE: a physically realistic environment for model building into low-
resolution electron-density maps. Acta Crystallogr. Sect. Struct. Biol. *74*, 519–530.
https://doi.org/10.1107/S2059798318002425.

86. Adams, P.D., Afonine, P.V., Bunkóczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J.,
Hung, L.-W., Kapral, G.J., Grosse-Kunstleve, R.W., et al. (2010). PHENIX: a
comprehensive Python-based system for macromolecular structure solution. Acta
Crystallogr. D Biol. Crystallogr. *66*, 213–221. https://doi.org/10.1107/S0907444909052925.

87. Afonine, P.V., Poon, B.K., Read, R.J., Sobolev, O.V., Terwilliger, T.C., Urzhumtsev, A., and
Adams, P.D. (2018). Real-space refinement in PHENIX for cryo-EM and crystallography.
Acta Crystallogr. Sect. Struct. Biol. *74*, 531–544.
https://doi.org/10.1107/S2059798318006551.

88. Yu, R.C., Jahn, R., and Brunger, A.T. (1999). NSF N-Terminal Domain Crystal Structure:
Models of NSF Function. Mol. Cell *4*, 97–107. https://doi.org/10.1016/S1097-
2765(00)80191-4.

89. Joosten, R.P., Long, F., Murshudov, G.N., and Perrakis, A. (2014). The PDB_REDO server
for macromolecular structure model optimization. IUCrJ *1*, 213–220.
https://doi.org/10.1107/S2052252514009324.

90. Hamelryck, T., and Manderick, B. (2003). PDB file parser and structure class implemented
in Python. Bioinformatics *19*, 2308–2310. https://doi.org/10.1093/bioinformatics/btg299.

91. Cock, P.J.A., Antao, T., Chang, J.T., Chapman, B.A., Cox, C.J., Dalke, A., Friedberg, I.,
Hamelryck, T., Kauff, F., Wilczynski, B., et al. (2009). Biopython: freely available Python
tools for computational molecular biology and bioinformatics. Bioinformatics *25*, 1422–
1423. https://doi.org/10.1093/bioinformatics/btp163.

92. Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Prettenhofer, P., Weiss, R., Dubourg, V., et al. (2011). Scikit-learn: Machine Learning in
Python. J. Mach. Learn. Res. *12*, 2825–2830.

93. Gowers, R.J., Linke, M., Barnoud, J., Reddy, T.J.E., Melo, M.N., Seyler, S.L., DomaÅ„ski,
J., Dotson, D.L., Buchoux, S., Kenney, I.M., et al. (2016). MDAnalysis: A Python Package
for the Rapid Analysis of Molecular Dynamics Simulations. In Proceedings of the 15th
Python in Science Conference, S. Benthall and S. Rostrup, eds., pp. 98–105.
https://doi.org/10.25080/Majora-629e541a-00e.

94. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate
protein structure prediction with AlphaFold. Nature *596*, 583–589.
https://doi.org/10.1038/s41586-021-03819-2.

95. Davis, I.W., Leaver-Fay, A., Chen, V.B., Block, J.N., Kapral, G.J., Wang, X., Murray, L.W.,
Arendall, W.B., Snoeyink, J., Richardson, J.S., et al. (2007). MolProbity: all-atom contacts

and structure validation for proteins and nucleic acids. Nucleic Acids Res. *35*, W375–W383.
https://doi.org/10.1093/nar/gkm216.

96. Liebschner, D., Afonine, P.V., Baker, M.L., Bunkóczi, G., Chen, V.B., Croll, T.I., Hintze,
B., Hung, L.-W., Jain, S., McCoy, A.J., et al. (2019). Macromolecular structure
determination using X-rays, neutrons and electrons: recent developments in Phenix. Acta
Crystallogr. Sect. Struct. Biol. *75*, 861–877. https://doi.org/10.1107/S2059798319011471.

97. Greaves, J., Gorleku, O.A., Salaun, C., and Chamberlain, L.H. (2010). Palmitoylation of the
SNAP25 Protein Family. J. Biol. Chem. *285*, 24629–24638.
https://doi.org/10.1074/jbc.M110.119289.

1256

Acknowledgments

We thank Richard Held, Jeremy Leitz, Rama Ranganathan, Subu Subramanian, Benjamin White,

Susan Wu, and Minglei Zhao for stimulating discussions; E. Montabana and C. Zhang for

assistance with the electron microscopy data collection at the Stanford University cEMc; and

Paul D. Adams, Pavel Afonine, and Nigel Moriarty for assistance with PHENIX refinement.

Funding was provided by the US National Institutes of Health (NIH; RO1MH63105 to A.T.B.)

and the Helen Hay Whitney Foundation/HHMI (to K.I.W.). Molecular graphics and analyses

performed with *UCSF ChimeraX*, developed by the Resource for Biocomputing, Visualization,

and Informatics at the University of California, San Francisco, with support from National

Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure and Computational

Biology, National Institute of Allergy and Infectious Diseases. This article is subject to HHMI's

Open Access to Publications policy. HHMI laboratory heads have previously granted a non-

exclusive CC BY 4.0 license to the public and a sublicensable license to HHMI in their research

articles. Pursuant to those licenses, the author-accepted manuscript of this article can be made

freely available under a CC BY 4.0 license immediately upon publication.

Author contributions

1272

K.I.W. and A.T.B. developed the research plan and experimental strategy. K.Q. and J.D.

performed the cellular imaging experiments in PC12 cells and analyzed the data. Y.A.K. and

A.B. performed the cellular imaging experiments in yeast and analyzed the data. K.I.W., Y.A.K.

S.C., L.E., and R.P. purified all proteins. K.I.W., S.C., and Y.A.K. performed the disassembly

experiments. K.I.W. performed the cryo-EM experiments and analyzed the data. K.I.W. and

A.T.B. wrote the manuscript with input from the other authors.

Competing Interest Statement

1279

1280   All authors declare no competing interests.

Figures

Figure 1 | NSF colocalizes to syntaxin nanodomains in an N-domain dependent manner and disassembles syntaxin oligomers

A. Colocalization between syntaxin-GFP nanodomains and NSF-mScarlet-I. NSF often forms clusters overlapping with syntaxin (PCC = 60% ± 2%, mean ± SEM).

B. Colocalization between syntaxin-GFP nanodomains and NSF(ΔN)-mScarlet-I; disruption of colocalization is evident (PCC = 28% ± 3%, mean ± SEM).

C. Colocalization between syntaxin-GFP and mScarlet-I alone. No interaction is apparent (PCC = 22% ± 2%, mean ± SEM).

D. Comparison of NSF-mScarlet-I, NSF(ΔN)-mScarlet-I, and mScarlet-I clustering with syntaxin-GFP nanodomains (M ± SEM, $n = 10$ images). Statistical significance was established by two-tailed Student's t-tests. Colocalization between NSF-mScarlet-I and syntaxin-GFP is significantly different than between NSF(ΔN)-mScarlet-I and syntaxin-GFP ($p = 8.9002 \times 10^{-8}$) or between mScarlet-I and syntaxin-GFP ($p = 5.2106 \times 10^{-10}$). Colocalization between NSF(ΔN)-mScarlet-I and syntaxin-GFP and colocalization between mScarlet-I and syntaxin-GFP were not significantly different ($p = 0.1776$).

E. To demonstrate disassembly tetrameric syntaxin-1a SNARE complexes by NSF and α-SNAP, and to show the insufficiency of Munc18 alone in disassembly, sx20S was prepared from purified components, split into two pools, resuspended in with a ~20-fold excess of Munc18. Following addition of EDTA or Mg^{2+}, pools were subjected to size-exclusion chromatography (SEC). Raw and filtered SEC profiles of the non-hydrolyzing (EDTA, blue) and hydrolyzing (Mg^{2+}, red) chromatograms at 290 nm are shown. The first peak corresponds to a complex of sx20S-Munc18 under non-hydrolyzing conditions (9.6 mL), or free NSF (10.2 mL) under hydrolyzing conditions. The second peak is consistent with an intermediate species of all proteins (11.6 mL). The third peak corresponds to a complex of syntaxin—Munc18 (13.6 mL), enriched twofold as assessed by integrated area under the curve (IAUC) under hydrolyzing conditions (12.8–14.8 mL; EDTA, 537.5 mAU^2; Mg^{2+}, 1058.5 mAU^2). The fourth peak is dominated by free α-SNAP, enriched twofold under hydrolyzing conditions (14.8–16.8 mL; EDTA, 224.4 mAU^2; Mg^{2+}, 442.7 mAU^2).

F. Sso1p forms dense clusters in the *S. cerevisiae* plasma membrane during cell division as assessed by SIM. Sso1p clusters were enriched in the budding daughter cell (left); discrete clusters typically approached sizes near the Nyquist limit of 160 nm (n = 46 cell images).

1314

1315

Figure 2 | NSF and α-SNAP engage a tetrameric SNARE complex composed of four syntaxin H3 domains

A. The structures of the sx20S (class #1, 3.75 Å) and the α-SNAP—syntaxin tetramer subcomplex (class #2, 4.07 Å). Hexameric NSF engages the α-SNAP—syntaxin tetramer subcomplex via its N-domains (salmon), several of which engage four α-SNAP molecules (gold) surrounding the soluble, tetrameric syntaxin H3 complex (dark red) within the catalytic D1 ring of NSF (light blue). The D2 ring of NSF (purple) does not engage the SNARE substrate. A hypothetical membrane is illustrated in grey. In the protein domain legend, ordered syntaxin domains are lightly shaded and those visible in the reconstruction of the SNARE complex are denoted with an asterisk; the syntaxin Habc domains were not observed.

B. NSF (light blue) engages one of the four syntaxin molecules (dark red) in the tetramer via a

series of pore loop interactions (protomers A–F, indicated); a tyrosine side chain (Y294) on each

of the six protomers intercalates in between every other side chain of a portion of syntaxin N-

terminal to its H3 domain. The corresponding sharpened map is contoured at 4 σ.

C. The α-SNAP—syntaxin tetramer subcomplex model, with two of the four α-SNAP models

hidden for clarity. The syntaxin H3 SNARE complex forms a twisted, parallel α-helical bundle

with a series of key SNARE complex layers preserved.

D. Slices through the α-SNAP—syntaxin tetramer subcomplex model and sharpened map at the -

4–0 layers, with key residues lightened for emphasis. Each layer is characterized by nearly planar

interactions between four identical side chains within the helical bundle. The primary

interactions between α-SNAP and the syntaxin H3 complex are observed near the -2 and -3

layers, where several functionally essential α-SNAP side chains (L197, Y200) are positioned by

a loop that buries them within the groove of the bundle. No symmetry was imposed during

1340

Figure 3 | NSF and α-SNAP engage a 2:1 binary SNARE complex of syntaxin H3 and
SNAP-25 SN1 and SN2 SNARE domains under non-hydrolyzing conditions

A. A representative 21bin20S model (class #5, 3.39 Å) and α-SNAP—2:1 binary SNARE

subcomplex model (class #13, 3.74 Å). Hexameric NSF engages the α-SNAP—2:1 binary

SNARE subcomplex via multiple N-domains (salmon), which engage three α-SNAP molecules

(gold) surrounding the 2:1 binary SNARE complex composed of two syntaxin H3 domains (dark

red) and the SN1 and SN2 SNARE domains of a single SNAP-25 molecule (green). A linker N-

terminal to one of the syntaxin H3 domains is found within the catalytic D1 ring of NSF (light

blue). The D2 ring of NSF (purple) does not engage substrate. A hypothetical membrane is

illustrated in grey. In the protein domain legend, ordered syntaxin and SNAP-25 domains are

shaded and those visible in the reconstruction of the SNARE complex are denoted with an

asterisk; syntaxin Habc domains were not observed.

B. NSF (light blue) engages one of the two syntaxin molecules in the 2:1 binary SNARE

complex (dark red) via pore loop interactions involving Y294 (protomers A–F, indicated). A

sharpened map is shown, contoured at 5 σ. Of the eight 21bin20S classes with the 2:1 binary

SNARE complex from the dataset collected under non-hydrolyzing conditions, six engage

syntaxin and two engage SNAP-25 within the pore of the D1 ring.

C. The α-SNAP—2:1 binary SNARE subcomplex model. Three α-SNAP molecules bind the 2:1

binary SNARE complex; a flexible linker between SNAP-25 SN1 and SN2 blocks access by a

fourth. The 2:1 binary SNARE complex forms a twisted, parallel α-helical bundle with a series

of characteristic SNARE complex layers preserved. This configuration also places the N-

terminal portion of the linker, bearing cysteines often palmitoylated *in vivo*, near the

membrane[97].

D. Layer slices through the α-SNAP—2:1 binary SNARE subcomplex model and sharpened

map. Each layer is characterized by nearly planar interactions between four side chains within

the helical bundle. The primary interactions between α-SNAP and the syntaxin H3 complex are

found near the -2 and -3 layers, where several functionally essential[61] α-SNAP side chains (L197,

Y200) are positioned by a loop that buries them within the bundle groove.

1369

1370

1371 **Figure 4 | NSF and α-SNAP engage a 2:2 binary SNARE complex of syntaxin H3 and**
1372 **SNAP-25 SN1 SNARE domains under hydrolyzing conditions**

1373 **A.** A representative 22bin20S model (class #19, 3.56 Å) and α-SNAP—2:2 binary SNARE

1374 subcomplex model (class #23, 4.14 Å). Hexameric NSF engages the α-SNAP—2:2 binary

1375 SNARE subcomplex via N-domains (salmon), which engage three α-SNAP molecules (gold)

1376 surrounding the soluble, 2:2 binary SNARE complex composed of two syntaxin H3 domains

1377 (dark red) and two SNAP-25 SN1 domains contributed by two separate full-length SNAP-25

1378 molecules (green). A portion of a linker N-terminal to one syntaxin H3 domain is observed

1379 within the catalytic D1 ring of NSF (light blue). The D2 ring of NSF (purple) shows no substrate

1380 density. A hypothetical membrane is illustrated in grey. In the protein domain legend, ordered

1381 syntaxin and SNAP-25 domains are lightly shaded and those visible in the reconstruction of the

SNARE complex are denoted with an asterisk; syntaxin Habc domains and SNAP-25 SN2

domains were not observed.

B. NSF (light blue) engages one of the two syntaxin molecules (dark red) in the 2:2 binary

SNARE complex via pore loop interactions involving Y294 (protomers A–F, indicated). A

sharpened map is shown, contoured at 6 σ. Of the five 22bin20S classes with the 2:2 binary

SNARE complex, all are found with syntaxin in the D1 pore, while two are observed with the N-

terminus of the SNAP-25 SN1 domain in the pore.

C. The α-SNAP—2:2 binary SNARE subcomplex model. Four α-SNAP molecules bind the 2:2

binary SNARE complex, with two SNAP-25 SN1 SNARE domains and two syntaxin H3

domains. This configuration locates two independent N-terminal portions of the SNAP-25 linker

bearing cysteine sidechains often palmitoylated *in vivo* near the membrane[97].

D. Slices through the α-SNAP—syntaxin tetramer subcomplex at the -4–0 layers. Each layer is

characterized by nearly planar interactions between four side chains within the helical bundle.

The primary interactions between α-SNAP and the syntaxin H3 complex are found near the -2

and -3 layers, where several functionally essential[61] hydrophobic α-SNAP side chains, L197 and

Y200, are buried within the bundle groove. Symmetry was not enforced during local

reconstruction. Note that the near-twofold symmetry of the bundle leaves some sidechain

densities poorly defined.

1400

1401

Figure 5 | Reconstruction and comparison of SNARE complexes and sub-complexes

A. Signal subtraction and local refinement in *CryoSPARC*[80] produced reconstructions of four

different α-SNAP—SNARE subcomplexes. In each case, the density of the SNARE proteins in

the core of the subcomplex was sufficiently well-defined to assign the SNARE domain identity, directionality, and register.

B. Crystal structures of the neuronal ternary SNARE complex (PDB ID 1N7S[54]) and the 2:2 syntaxin—SNAP-25 complex (PDB ID 1JTH[34]) were superimposed using UCSF ChimeraX[84] with cryo-EM structures from this work of the tetrameric syntaxin H3 sub-complex, the 2:1 syntaxin—SNAP-25 sub-complex, and the 2:2 syntaxin—SNAP-25 sub-complex within the bin20S complex. These structures show similar quaternary structure, maintaining the same register and similar twist in the central SNARE layers. The SNARE complexes and sub-complexes within their respective 20S complexes are somewhat untwisted at the N- and C-terminal ends relative to the crystal structures of the SNARE complexes alone, likely due to a combination of the absence of crystal packing as well as interactions with α-SNAPs and NSF.

C. Quantitative comparison of individual SNARE domain ternary structure. Each pixel in a pair of matrices represents the mean Cα RMSD (Å, left) and corresponding standard deviation (right) between two instances of a given SNARE domain in two different structures. Only Cα atoms shared by all structures were considered. The same color scale is preserved across all comparisons. The mean Cα RMSD generally varies between 1–2 Å depending on the specific pair.

D. Individual Cα RMSDs mapped to their corresponding SNARE domain structures, aligned. Core regions of the complexes are structurally similar, but unwinding near the SNARE domain termini leads to larger deviations, particularly near the N-termini where NSF engages one of the N-terminal segments leading into a SNARE domain (SNAP-25 or syntaxin).

1427

Figure 6 | Multimodal engagement of SNARE substrate by NSF

A. The α-SNAP—SNARE subcomplex is engaged by the NSF N domains. The orientation of the

subcomplex relative to the NSF D1 ring split, the tilt of the subcomplex relative to the D1 ring,

and the specific NSF N-domains engaged to the subcomplex are major determinants of particle

class identity in different datasets. Three distinctive "grips" are observed in the case of the

classes with the 2:1 syntaxin—SNAP-25 complex identified under non-hydrolyzing conditions

(top), and two are observed in the case of the classes with the 2:2 syntaxin—SNAP-25 SNARE

complex identified under hydrolyzing conditions (bottom). Pairs of N-domains typically bind a

given α-SNAP near its C-terminus; arrows indicate the most well-ordered N-domains as assessed

by the presence of two-lobed density. The two cases are not directly comparable, as the 2:1

syntaxin—SNAP-25 sub-complex is bound by three α-SNAP molecules, while the 2:2

syntaxin—SNAP-25 sub-complex is bound by four α-SNAP molecules. This difference is likely

due to the flexible linker that runs along the 2:1 syntaxin—SNAP-25 sub-complex surface

between the SNAP-25 SN1 and SN2 SNARE domains. **B.** The SNARE N-termini are engaged

by the NSF D1 pore. Variable syntaxin engagement by NSF between non-hydrolyzing and

hydrolyzing classes is observed. Alignment of structures with ordered syntaxin substrates in the

pore of the D1 ring under non-hydrolyzing (red) and hydrolyzing (blue) conditions reveals

disruption of the syntaxin H3 domain up to around residues 197 and 205, respectively.

1446

1447

Figure 7 | Comparison of NSF structures under non-hydrolyzing and hydrolyzing conditions reveals the interplay between sequential AAA+ hydrolysis and substrate rebinding

A. Key conformational states of the NSF D1 ATPase ring. Two states dominate in the absence of Mg^{2+} (left); the top class (#4) has SNARE substrate disengaged, nucleotide-free protomer F, while the bottom class (#3) has SNARE substrate-engaged, ATP-bound F. With the addition of Mg^{2+} (middle), E is Mg^{2+}-bound. Protomers D, C, and occasionally B retain P$_i$ and Mg^{2+}. Protomers A are ADP-bound. The top subclass (#18) has substrate-disengaged, ADP-bound F. The bottom subclass (#17) has nearly substrate-engaged, ATP-bound F. Substrate-free, post-hydrolysis classes, e.g., #24 (right) show an ATP-free D1 ring. Spire, SNARE substrates, and the D2 ring are mostly hidden here.

B. Details of hydrolysis. The P-loop binds nucleotide. Arginine fingers R385 and R388 from the down-ring protomer coordinate nucleotide phosphates. Without Mg^{2+}, a water molecule sits between the γ-phosphate and Walker B residue side chains of D328 and E329. Sensor 1 residue N374 is oriented towards the γ-phosphate of ATP. Under hydrolyzing conditions, Mg^{2+} is coordinated by the β- and γ-phosphate moieties of ATP and the side chain of Walker A motif T267. Following hydrolysis, ADP, P$_i$, and Mg^{2+} remain; Walker B E329 and Sensor 1 N374 side chains coordinate the P$_i$ as it disassociates. P$_i$ release down-ring leads to arginine finger disassociation, disrupting the interprotomer interface. Models are derived from D1 protomers #4-E, #16-E, #16-D, and #16-B.

C. Alignment of 130 D1 protomer large subdomains from 23 models derived from all datasets. Only Cα and Pβ atoms are shown. The inset shows a key region of the protein that shows coupled changes in the N-D1 linker and the Walker A helix. The latch loop contributed by the up-ring protomer is also coupled to the conformational state of this region.

1472   **D.** Alignment of tyrosine residues in the pore of the D1 ring of substrate-bound models reveals

1473   protomer F as the site of nucleotide exchange and substrate re-engagement. ADP-bound F is

1474   disengaged from the substrate (blue). The release of ADP (nucleotide-free, grey) and ATP-

1475   binding (red) accompany substrate (re)engagement. The lowest tyrosine corresponding to an

1476   ATP-bound F shown (red triangle) is from the model shown in **A** (middle column, bottom).

1477

Figure 8 | Principal component analysis of NSF D1 protomer coordinates reveals quantitative mechanistic details of sequential hydrolysis

A. 132 D1 domain models were aligned by large subdomain, subjected to Cα coordinate PCA, and clustered with HDBSCAN in 11-dimensional space (left). Protomers cluster into discrete

groups based on conformation, visible upon projection into a plane defined by the first and fifth

modes. A proposed hydrolysis cycle is indicated. Superimposed models corresponding to these

transitions are shown (right). The cluster corresponding to pre-hydrolysis protomers E is circled.

B. Protomer nucleotide state, mapped to the 2D PCA coordinate from **A**. The ATP + Mg^{2+} state is

unique to protomers E (circled).

C. The angle (º) between helices α_3 and α_8 (a proxy for the relative angle between the D1 large

and small subdomains). Protomers E have the largest angle (circled).

D. A mutant form of NSF, NSF(Δlatch), in which part of the latch loop (residues 460–466) was

replaced with a short glycine-serine linker (SGS), was assayed for its ability to disassemble the

ternary SNARE complex composed of syntaxin, SNAP-25, and synaptobrevin[6]. Traces show the

mean ± SEM of eight replicates for each form of NSF, with and without adding $MgCl_2$.

Disassembly rates were calculated by linear regression to each trace in the linear region of the

time course, from ~95–350 s. NSF(Δlatch) disassembled SNARE complex at 28 ± 1% the rate of

wild type.

E. Three conformational features distinguish the pre-hydrolysis protomer E cluster (circled).

These features are in contact and structurally coupled. The position of nucleotide in the active

site is modulated by the position of the Walker A helix (α_3), which occupies three discrete states;

pre-hydrolysis protomers E occupy the "down" state (left). The position of the α_3 is allosterically

modulated by at least two inputs. The N-D1 linker conformation (center) is connected to the state

of N-domain binding to α-SNAP, and it interacts with the latch loop of the up-ring protomer

upon rebinding. The latch loop of the up-ring protomer contacts this region in *trans* and is also

structurally coupled (right). These two elements push against the C-terminal end of α_3 and

ultimately modulate the position of the ATP γ-phosphate relative to key catalytic residues.

1506

1507

Figure 9 | The pre-hydrolysis state of protomers E has several unique active site features.

A. Distance distributions between the P262 Cα and a given active site residue. P262 Cα is a

relatively invariant reference atom for measuring distances, as it is well-constrained by local

geometry and the presence or absence of Pγ, Pi, or Mg^{2+} in the active site. Protomers E (orange

bars) have a significantly closer Sensor I N374 distance to P262 (4.8 ± 0.5 Å vs. 5.7 ± 0.9

overall, $p < 0.005$ for all pairs) than any other protomer (black bars). The terminal carbon of the

second arginine finger, R388 Cζ, was also significantly closer in protomers E than in any other

protomer (7.5 ± 0.4 Å vs. 13.7 ± 9.6 Å, $p < 0.05$ for all pairs). While not a specific observation to

protomer E, the P262 Cα-Walker B D328 Cγ distance distribution is bimodal, with ADP-bound

protomers generally having a smaller distance between the two atoms. Welch's t-test, assuming

unequal variances, was used to assess significance in all cases.

B. Distance distributions between the Pβ atom of either ATP or ADP of a given active site

residue. Pβ was used as it is found in all nucleotide-bound protomers. The pattern of significant

differences between protomers E and protomers A–D and F described for P262 Cα is mostly

repeated here, with the Sensor I N374 average distance to Pβ found to be highly significant (6.6

± 0.4 Å vs. 8.6 ± 1.3, $p < 0.005$ for all pairs). As in **A**, the terminal carbon of the second arginine

finger, R388 Cζ, was also significantly closer to Pβ in protomers E than in any other protomer

(6.3 ± 0.6 Å vs. 11.6 ± 8.8, $p < 0.05$ for all pairs). Some differences are present in other cases.

Aside from specific concerns related to protomers E, many of these distributions also show a

bimodal pattern consistent with different nucleotide states. Welch's *t*-test, assuming unequal

variances, was used to assess significance in all cases.

C. Distances from the histograms in **B** mapped to the coordinate determined by PCA.

1530

1531

Figure 10 | An expanded model for NSF action across cellular and atomic scales

A. A model for disassembling non-fusogenic SNARE species by NSF and α-SNAP at the

presynapse and progression to priming and fusion. Syntaxin nanodomains comprised at least in

part of tetrameric syntaxin SNARE complexes serve as a reservoir for the liberation of

individual, fusogenic syntaxin molecules by NSF upon disassembly. These syntaxin molecules

may also enter non-productive binary complexes with SNAP-25, with 2:1 or 2:2 stoichiometry;

such complexes would be disassembled by NSF to ensure *trans*-SNARE complex formation with

synaptobrevin on incoming synaptic vesicles in concert with other factors, such as Munc18,

Munc13, complexin, and synaptotagmin. Where possible, full-length structures (structures

reported here and PDB IDs 1N7S, 4JEU, 5W5C) were supplemented with AlphaFold2[94]

structures to provide additional context, and ISOLDE[85] was used with secondary structure

restraints to reposition flexible regions.

B. A sequential hydrolysis model for SNARE complex disassembly by NSF is inferred from

comparing different classes from non-hydrolyzing and hydrolyzing conditions. Following

hydrolysis-independent substrate side loading, the D1 ring is found in a pre-hydrolysis state with

ATP in protomer E of D1. Protomer F nucleotide exchange then enables ATP exchange and

rebinding to both SNARE substrate and the formation of the interprotomer interface with

protomer E, triggering ATP hydrolysis in protomer E and substrate translocation. This is

accompanied by P_i and Mg^{2+} release from down-ring protomers; in many structures, the release

occurs between protomers C and B. The cycle then begins anew. Blue corresponds to ADP-

bound, red to ATP-bound, purple to ADP•P_i bound, and grey to nucleotide-free. See Movie S1 for

an animated rendition.

Statistics and reproducibility

1555   No statistical method was used to predetermine the sample size, but experiments described in

1556   this study were performed with at least three samples for each group in the experiments shown

1557   throughout.

Reporting summary

1559   Further information on research design is available in the Nature Portfolio Reporting Summary

1560   linked to this article.

1561