Human brain changes after first psilocybin use

Lyons T[1], Spriggs M[1], Kerkelä L[2], Rosas FE[1,3,4,5,6], Roseman L[1], Mediano PAM[5,7,8],

Timmermann C[1], Oestreich L[9], Pagni BA[13], Zeifman RJ[1,13], Hampshire A[16], Trender W[15],

Douglass HM[1], Girn M[11,12], Godfrey K[1], Kettner H [1,12], Sharif F[1], Espasiano L[1], Gazzaley A[14],

Wall MB[10], Erritzoe D[1], Nutt DJ[1], *Carhart-Harris RL[1,12]

[1] Centre for Psychedelic Research, Division of Psychiatry, Department of Brain Sciences, Imperial
College London, London, UK.

[2] UCL Great Ormond Street Institute of Child Health, University College London, London, UK.

[3] Department of Informatics, University of Sussex, Brighton, UK.

[4] Data Science Institute, Imperial College London, London, UK.

[5] Centre for Complexity Science, Imperial College London, London, UK.

[6] Centre for Eudaimonia and Human Flourishing, University of Oxford, Oxford, UK.

[7] Department of Psychology, University of Cambridge, Cambridge, UK.

[8] Department of Psychology, Queen Mary University of London, London, UK.

[9] School of Psychology and Australian Institute for Bioengineering and Nanotechnology (AIBN), University
of Queensland, Australia.

[10] Invicro London, Hammersmith Hospital, Faculty of Medicine, Imperial College London, UK.

[11] Department of Neurology and Neurosurgery, Montreal Neurological Institute, McGill University,
Montreal, QC, Canada.

[12] Carhart-Harris Lab, Department of Neurology, University of California San Francisco, San Francisco,
CA, USA.

[13] NYU Langone Center for Psychedelic Medicine, Department of Psychiatry, NYU Grossman School of
Medicine, NY, US.

[14] Neuroscape, Department of Neurology, University of California San Francisco, San Francisco, CA,
USA.

[15] Department of Brain Sciences, Imperial College London, UK.

[16] Department of Neuroimaging, Institute of Psychiatry, Psychology & Neuroscience, King's College
London, UK.

* robin.carhart-harris@ucsf.edu

ABSTRACT (147 words)

Psychedelics have robust effects on acute brain function and long-term behavior but whether they also cause enduring functional and anatomical brain changes is unknown. In a placebo-controlled, within-subjects, electroencephalography, and magnetic resonance imaging study in 28 healthy, entirely psychedelic-naive participants, anatomical and functional brain changes were detected from one-hour to one-month after a single high-dose (25 mg) of psilocybin. Increases in cognitive flexibility, psychological insight, and well-being were seen at one-month. Diffusion imaging done before and one-month after 25mg psilocybin revealed decreased axial diffusivity bilaterally in prefrontal-subcortical tracts that correlated with decreased brain network modularity over the same time period. Decreased modularity also correlated with improved well-being. Increased cortical signal entropy at 1- and 2-hours post-dosing predicted improved psychological well-being at one-month. Next-day psychological insight mediated the entropy to well-being relationship. All effects were exclusive to 25mg psilocybin; no effects occurred with a 1mg psilocybin 'placebo' dose.

KEYWORDS

Neuroplasticity; plasticity; psychedelic; psilocybin; serotonin; 5-HT$_{2A}$; neuroimaging; well-being; mental health; insight

INTRODUCTION

Psilocybin (4-PO-HO-DMT) is the precursor of psilocin (4-HO-DMT), a serotonin receptor agonist. Converging evidence supports a role for serotonin 2A receptor (5-HT2AR) agonism in eliciting the characteristic brain and behavioural effects of this and related psychedelics in humans [1,2]. Pre-clinical research supports an association between 5-HT2AR agonism and increased 'neuroplasticity' [3-7], although see [8].

Clinical trials have assessed psilocybin-therapy for psychopathology [9-14]. Human research has found enduring psychological changes after single high-dose psilocybin [15], including increased cognitive flexibility in depressed patients [16] (although see [17] with LSD) and improved well-being in diverse samples [18,19]. Several human neuroimaging studies have observed changes in acute brain function with psychedelics [20-24]. Long-term brain changes are less well characterized, but see [25,26] in healthy volunteers and [16,27-30] in depression, plus Table S1 for a list of relevant studies.

In the present work, we sought to address important knowledge gaps regarding human brain changes with psilocybin. Electroencephalography (EEG), functional magnetic resonance imaging (fMRI) and diffusion tensor imaging (DTI) recordings were done in healthy human volunteers receiving their first-ever high-dose of a psychedelic (25mg psilocybin). Brain (fMRI/DTI) and behavioural outcomes were assessed at baseline and one-month post-dosing. EEG was done during dosing. Due to prior evidence of enduring behavioural changes after psilocybin [16,26,27,29], a fixed-order, repeated measures design was used; participants received an initial 'placebo' dose of psilocybin (1mg) one-month prior to a subsequent 25mg dose. Dosing was oral. See Figures S2-3 for study design details.

RESULTS

Demographics

Twenty-eight healthy volunteers participated, mean age = 41 (SD=8.7), 43% female. See Figure S4 for a demographics table.

Neuroimaging outcomes

Acute brain effects

 Time-dependent changes in LZc and spectral power (EEG)

Eyes-closed, task-free 'resting-state' brain activity during each of the two psilocybin dosing sessions (i.e., 1mg and 25mg, one-month apart) was investigated using EEG performed at pre-dose baseline and 1-, 2-, and 4.5-hours post-dosing. Robust pre vs acute (i.e., 'on-drug') brain changes were observed after the 25mg dose, as revealed by *Dose* (1mg vs 25mg) x *Time* (0-4.5h) linear mixed-effects modelling. There were significant ($p<0.001$) increases in the informational entropy of spontaneous scalp potentials (Lempel-Ziv complexity, LZc) at 1- and 2-hours post 25mg, and significant decreases in alpha power (Figure 1A and B). Additionally, power increases in the gamma band at 2h and decreases in the theta band at 1 and 2h were observed. No changes in EEG recorded activity were observed under 1mg psilocybin (See Table S5 and Figure S6 for these data).

Figure 1. **Acute EEG. Significant effects of 25mg (vs 1mg) on brain entropy (LZc) and alpha power.** Increases in LZc (A) and decreases in alpha power (B) were maximal at the two-hour timepoint, coinciding with maximum subjective intensity. The red (25mg) and gray (1mg, placebo) timelines correspond to means and standard errors (shaded areas) at 1, 2, and 4.5h post-dose. Vertical bars index contrasts, *** = $p < 0.001$. See Figure S7 for single-subject data.

Long-term brain changes

Anatomical changes (MRI)

Diffusion tensor imaging revealed a significant interaction between '*Timepoint*' (i.e., one factor, three-levels, 1) baseline, 2) 1-month post-1mg, 3) 1-month post-25mg) and mean axial diffusivity (AD) - the diffusion coefficient along the principal diffusion direction, see Figure 2A. ANOVA revealed significant changes in AD for two tracts after 25mg psilocybin, a bilateral prefrontal cortex - striatum tract (PFC-STR; $F(2, 48) = 10.30$, $p = 0.005$) and a PFC - thalamus

tract (PFC-THA; F(2, 48) = 9.51, p = 0.008). See Figure S18 for all relevant tracts. Post-hoc t-
tests found significant decreases in AD one-month post-25mg vs. one-month post-1mg for the
PFC-STR (t(24) = -3.72, p = 0.006) and PFC-THA tracts (t(24) = -3.85, p = 0.005). Separating
by hemispheres for post-hoc t-tests, decreases were apparent in both hemispheres; however,
with Bonferroni correction, only decreases in the left hemisphere remained significant (see
Figure S19). No significant differences in AD were observed following the 1mg control. With
free-water correction (see Figure S21 and section 5.5. of the supplement), consistent AD effects
were seen post-25mg psilocybin, and fractional anisotropy (FA) changes also became
statistically significant (Figure S21). A tract-based spatial statistics (TBSS) method was also
done on request (see Methods) but yielded no significant results.

Brain response to emotional faces
Whole-brain analyses of the Blood Oxygen Level Dependent (BOLD) fMRI response to
emotional face stimuli (happy, fearful, neutral) yielded a large effect (95% *CI* [0.33, 1.34], *d*=0.9)
for the salient contrast of one-month post-25mg vs one-month post-1mg psilocybin for fearful
faces vs fixation cross, Figures 2C & 2D (see Section 4 of the supplementary file, Figures S8 &
S9, for results for happy and neutral faces – and for the post-25mg vs pre-dose baseline
contrast; 95% *CI* [0.04, 0.98], *d*=0.3). However, an inclusive (i.e., all factors, all levels) whole-
brain analysis, failed to find a significant interaction between emotional face type and *Timepoint*
on BOLD response. Explorative amygdala region of interest (ROI) results can be found in the
supplement (Figures S10 & S11), along with generalized psychophysiological interaction (gPPI)
functional connectivity (FC) findings. Amygdala effects were observed for the left hemisphere
and gPPI effects were observed that survived FDR correction for exploring both hemispheres
(See Table S12 & Figure S13).

Resting-state FC (RSFC)
Explorative 'resting-state' (RS) functional connectivity (RSFC) analyses were applied to data
from a single 8-minutes eyes-closed, task-free, RS fMRI run. Based on prior hypotheses, four
regions were initially chosen for seed-based RSFC analyses: (i) bilateral parahippocampus
(PH), (ii) bilateral amygdala, (iii) ventromedial prefrontal cortex (vmPFC), and (iv) subgenual
anterior cingulate cortex (sgACC). Two further seeds were examined after peer review: (v) the
hippocampus and (vi) the dorsal anterior cingulate cortex (dACC). In ANOVAs across all
timepoints, three of the initial four seed did not yield significant results and neither did the further
two seeds. However, an increase in amygdala RSFC was evident one-month post-25mg vs

one-month post-1mg psilocybin (M_{diff}=3.68, SE=0.64, $p < 0.0001$; 95% *CI* [2.00, 5.25], d=1.2) as

well as when post-25mg was contrasted against pre-dose baseline (M_{diff}=1.49, SE=0.56,

p=0.046; 95% *CI* [0.02, 2.96], d=0.6, Bonferroni-corrected). See Figure S14 for the relevant

maps and S15 for single subject data. The robustness of these results was highlighted by an

inclusive ANOVA across all timepoints (Figure S16) that found a significant interaction between

Timepoint and *amygdala RSFC* in consistent regions as those shown in Figure S14 for the key

contrast.

Network modularity (RSFC)

Previous work highlighted a relationship between improved depressive symptoms and

decreased brain network modularity after psilocybin-therapy for depression [27]. Unlike in this

previous work, statistically significant group-average decreases in modularity were not observed

here, and neither did we find changes in mass univariate network analyses. More specifically,

we found no statistically significant changes in *within* - or *between* network RSFC. However,

controlling for *Well-being* before the 25mg session, a correlation was found between *modularity*

changes (decreases) one-month post-25mg and contemporaneous changes (improvements) in

Well-being (r= -0.40, p=0.04, two-tailed; Figure 2E and F; see also Figure 4B). This relationship

is consistent with prior work in depression that found decreased network modularity post-

psilocybin correlating with improved depressive symptom severity in two separate trials [27].

Figure 2. One-month post-25mg psilocybin anatomical and functional brain changes. **A**)
DTI. Combined bilateral prefrontal-subcortical white-matter tracts where significant decreases in
axial diffusivity (AD) were observed one-month post-25mg, Bonferroni corrected. **B**) Violin plot
shows decreases in AD with dose; * = $p < 0.05$, ** = $p < 0.01$. See supplement for consistent
results with free-water correction (Figure S21) as well as single subject data (Figure S20). **C**)
BOLD responses to fearful faces (vs fixation cross) for the key contrast one-month post-25mg
vs one-month post-1mg. See Figures S8-9 for baseline and single-subject values. Blue depicts
locations of reduced response to fearful faces one-month post-25mg (Z = 2.3, $p < 0.05$, cluster-
corrected). **D**) Plots show Z values from the significant clusters shown in C. Note, the delta
values for this specific contrast contributes to the correlation matrix shown in Figure 4B; i.e., the

variable is labeled 'Emo-fMRI' in 4B. **E**) *Brain modularity* at the three timepoints, NS = no

contrast was significantly different. **F**) Controlling for *Well-being* prior to 25mg, there was a

significant negative correlation between *Modularity Change* and *Well-being Change* in the key

contrast of one-month post-25mg vs one-month post-1mg (r = -0.4, p = 0.04, 2-tailed), i.e.,

decreased modularity correlated with improved well-being (see also Figure 4B). Note that

plotting the residuals in 2F does not highlight the robust improvements in well-being that were

seen (note, these changes are evident in 3C). For further DTI and fMRI results, including single

subject data, see the supplementary file. Modularity and AD (PFC-STR and PFC-THA tracts

merged) change values for the salient one-month post-25mg vs one-month post-1mg contrast

also contribute to Figure 4B.

194
195
196 **Psychological outcomes**

197
Acute subjective intensity

An interaction between *Dose* (1mg vs 25mg) and *Time* (0-6h) on the *Intensity* of subjective drug

effects was seen ($F_{(8,197)}$=57.73, p < 0.0001). Greater subjective intensity was reported in

relation to 25mg psilocybin compared with both time-zero baseline (all p < 0.0001 from 1-5h; p <

0.001 at 6h) and the 1mg control (all p < 0.0001 from 1-5h; p = 0.001 at 6h; Figure 3A). No

significant elevation in subjective intensity was observed with 1mg. All, except one, of the

participants (94%) rated the 25mg experience as the "*single most unusual state of*

consciousness" of their entire life. The remaining person ranked it "*among the top five most*

unusual experiences" of their entire life. In contrast, most participants rated their 1mg

experience as "*no more unusual than an everyday state of consciousness*" (see Figure S26)

reinforcing the impression from the EEG data that the 1mg 'placebo' dose was functionally

inactive.

Figure 3. Behavioral analyses. (A) Subjective intensity. A significant interaction between
Dose (1mg vs 25mg) and *Time* (0-6h) ($F_{(8,197)}$ = 57.73, p < 0.0001) on *subjective intensity* rated
in-session. Post-hoc t-tests, 25mg vs 1mg, *** = p < 0.001, **** = p < 0.0001. **(B)
Psychological insight scale (PIS).** A significant interaction between *Dose* (1mg vs 25mg) and
Timepoint (one-day, two-weeks & one-month post-dosing) on *psychological insight* ($F_{(2,44)}$ =
14.05, p < 0.0001); insight scores (PIS) were significantly greater at all timepoints post-25mg vs
1mg (**** = p < 0.0001). **(C) Warwick-Edinburgh Mental Well-Being Scale (WEMWBS).** A
significant interaction was observed between *Dose* (1mg vs 25mg) and *Timepoint* (pre-dose
baseline, two-weeks & one-month post-dose) on *Well-being* ($F_{(2,44)}$ = 11.92, p < 0.0001); *Well-
being* (WEMWBS) increases were significant at two-weeks (M_{diff} = 5.83, SE = 1.54, p < 0.01;
95% *CI* [-10.64, -1.02]*, d* = 0.8) and one-month (M_{diff} = 4.70, SE = 1.47, p < 0.05; 95% *CI* [-9.29,

-0.10], *d* = 0.6) post-25 mg psilocybin. Post-hoc *t*-tests, 25mg vs 1mg, * = $p < 0.05$, ** = $p <$

0.01. **(D) Intra- and extra-dimensional attentional set-shifting (IDED) & Extradimensional**

Shift Errors (EDS, Cognitive Flexibility). Analyses revealed a significant interaction between

Task Performance and *Timepoint* ($F_{(7.29,167.67)}$ = 2.47, p = 0.018; see S23A). Post-hoc tests

showed significantly fewer EDS errors in the key contrast of one-month post-25mg vs one-

month post-1mg (M_{diff} = 0.06, *SE* = 0.02, p = 0.016*; 95% *CI* [0.00, 0.12], *d* = 0.6), outlier

corrected, see Figure S23C for details. Fewer EDS errors reflects faster (correct) detection of

rule shifts. Data are expressed as mean ±SEM. T-tests, * = $p < 0.05$, ** = $p < 0.01$. No

significant effects of 1mg were observed on any of the statistical tests.

Long-term psychological changes

Psychological insight

Psychological insight was assessed via the 'psychological insight scale' (PIS) [31]. Analyses

revealed a robust interaction between *Dose* (1mg vs 25mg) and sub-acute *Timepoints* on

Insight ($F_{(2,44)}$ = 14.05, $p < 0.0001$; Figure 3B). Insight scores were significantly greater after

25mg versus 1mg psilocybin at all sub-acute timepoints— i.e. one-day (M_{diff} = 50.90, *SE* = 5.51,

$p < 0.0001$; 95% *CI* [39.50, 62.29], *d* = 1.9), two-weeks (M_{diff} = 29.83, *SE* = 4.72, $p < 0.0001$;

95% *CI* [20.08, 39.57], *d* = 1.3) and one-month post-dosing (M_{diff} = 27.86, *SE* = 4.37, $p < 0.0001$;

95% *CI* [18.90, 36.82], *d* = 1.2), Bonferroni-corrected.

Psychological well-being

Psychological well-being was assessed using the Warwick-Edinburgh Mental Wellbeing Scale

(WEMWBS), a single dimension, self-report measure of well-being rated with reference to a

two-week period [32]. Analyses revealed a significant interaction between *Dose* (1mg vs 25mg)

and sub-acute *Timepoints* on *Well-being* ($F_{(2,44)}$ = 11.92, $p < 0.0001$). Bonferroni-corrected *post-*

hoc comparisons revealed significantly greater increases in well-being at two-weeks (M_{diff} =

5.83, *SE* = 1.54, $p < 0.01$; 95% *CI* [-10.64, -1.02], *d* = 0.8) and one-month (M_{diff} = 4.70, *SE* =

1.47, $p < 0.05$; 95% *CI* [-9.29, -0.10], *d* = 0.6) post-25mg psilocybin (vs 1mg), Figure 3C. There

were no significant changes post-1mg dose vs (pre any intervention) baseline.

Cognitive flexibility

Cognitive flexibility was measured via an intra-dimensional/extra-dimensional (IDED) task,

optimised for internet-based delivery [33,34]. Extradimensional shift (EDS) is an index of

participants' ability to correctly identify rule shifts, and thus, *Cognitive Flexibility*. Participants

completed the task at baseline and one-month post-dosing. Greenhouse-Geisser correction was

applied due to a violation of sphericity for the interaction term (Epsilon ε = 0.456). Analyses revealed a significant interaction between *Task Discrimination Stage* and *Timepoint* on *Cognitive Flexibility* ($F_{(7.29,167.67)}$ = 2.47, *p* = 0.018). FDR-correction of post-hoc comparisons (i.e., for the 9 IDED task phases, Figure 22) plus Bonferroni correction within the salient EDS phase, showed a significant decrease in EDS errors one-month post-25mg vs one-month post-1mg (Figure 3D), indicative of greater cognitive flexibility (M_{diff} = 0.06, *SE* = 0.02, *p = 0.016*; 95% *CI* [0.00, 0.12], *d*=0.5). Contextualized by no global difference in performance or at IDS, IDR or EDR stages, the EDS change pertains to flexible attentional set as opposed to perseveration or more general problem with processing discriminations / maintaining rules. No IDED changes were observed after 1mg psilocybin. See S22-25 of the supplement for more detailed IDED results.

Correlations between long-term change outcomes

The relationship between various contemporaneous (i.e., synchronous in time) changes in brain and behavioral variables observed in the key contrast of one-month post-1mg vs one-month post-25mg was assessed via a correlation matrix that included the relevant contrast values for: 1) *EDS errors* (Cog-flex); 2) *Amygdala-RSFC* (amg-RSFC); 3) *Well-being*; 4) *BOLD responses to emotional faces* (Emo-fMRI); 5) *Brain Network Modularity*; 6) *DTI-measured axial diffusivity* in the merged PFC-THA and PFC-STR tracts (DTI), and 7) *Psychological Insight* (one-month post 1mg vs one-month post 25mg); Figure 4B. Values represent Pearson correlation coefficients, calculated via pairwise linear regressions. Significance was assessed via non-parametric cluster statistics, appropriate for non-independent tests. A significant cluster was found (*p* = 0.006, cluster corrected, coloured cells), implying a nonspurious correlational structure. All of the significant relationships are directionally consistent with the direction of the changes in each outcome for the key contrast of one-month post-1mg vs one-month post-25mg psilocybin.

Acute brain entropy predicts increased psychological insight and improved well-being

In two data driven analyses, the informational entropy of spontaneous cortical activity (*LZc*, EEG) under 25mg psilocybin predicted both next-day psychological insight (Figure 4A) and improvements in psychological well-being at one-month post-25mg psilocybin (Figure 4C). In the former case, LZc increases within posterior electrodes (P3, P4, and O1) recorded during 2-hours post-25mg, predicted psychological insight scored one-day later (r_s = 0.59, p = 0.006, Figure 4A). In the latter case, LZc predicted improved psychological well-being scores (one-month post-25mg versus one-month post-1mg) directly from most sensors at all three post-dose

timepoints (i.e., at 1, 2, and 4.5 hours; r_s = 0.66, p = 0.006, Figure 4C). No such predictive

relationships were found for the 1mg condition. Note that regressions were done in a data-

driven way to identify— via cross-validation — the most sensitive sensors for the relationship

i.e., those with the greatest predictive capacity. Repeating these analyses using all sensors

contributing equally produced consistent results (see 3.4. in the supplement).

A final mediation analysis (Figure 4D) was performed using linear models that included key

variables at three timepoints: (i) whole-brain LZc 2h post-25mg psilocybin (i.e., averaged signal

across all sensors at this timepoint), (ii) *Insight* measured one-day post-25mg, and (iii) *Well-*

being changes one-month post-25mg. This analysis revealed that acute *LZc* was a significant

predictor of next-day *Insight* (β = 197, SE = 67, p = 0.008, normalised β = 0.55). Both *LZc* (β =

41.8, SE = 14.2, p = 0.008, normalised β = 0.54) and *Insight* (β = 0.11, SE = 0.04, p = 0.01,

normalised β = 0.50) were significant predictors of increased *Well-being* at one-month; however,

the effect of *LZc* on *Well-being* became statistically insignificant if *Insight* was added to the

model (β = 29.6, SE = 17.2, p = 0.10, normalised β = 0.38). These results satisfy the criteria for

statistical mediation by [35] – as well as more recent recommendations [36] – implying that the

relationship between acute LZc and improved well-being at one-month post-25mg

was significantly mediated by sub-acute psychological insight.

Figure 4. Correlational and predictive modelling. A) A cross-validated predictive model revealed a significant positive correlation between acute Lempel-Ziv complexity (LZc, or 'brain entropy') for P3, P4, and O1 electrodes, and psychological insight, measured one-day post-25mg. Inset shows electrode-level relationship between LZc (brain entropy) at 2h post-dose (when subjective effects were intense) and next-day insight (see also 3.4 in the supplement for a validation of this result using data from all sensors). **B)** Correlation matrix examining relationships between functional and anatomical MRI metric values and contemporaneous psychological outcomes. Coloured cells represent significant Pearson correlations, corrected for multiple comparisons (non-parametric cluster test, p = 0.006). Red/pink cells reflect significant positive correlations, blue cells reflect significant negative correlations. All relationships are directionally consistent with the original direction of changes. All values derive from the key one-month post-1mg versus one-month post-25mg contrast. **C)** Model predicting one-month well-being increases directly from acute increases in brain entropy at all three post-dosing timepoints. Plot shows electrode-level relationship with (average) data from the 3 post-dose timepoints, i.e., 1-, 2-, and 4.5-hours post-dose (see also 3.4 in the supplement for a validation of the result using data from all sensors). **D**) Model shows that increased well-being at one-

month post-25mg can be explained by increased brain entropy (LZc) across all sensors, but also that the strongest path entails brain entropy first predicting next-day *Insight*, which then, in-turn, predicts improvements in *Well-being* (1-month). The values between the arrows in 4D represent normalized regression coefficients. Relationships were statistically significant at p < 0.05.

DISCUSSION (1188 words)

The present work sheds important new light on human brain changes after first-time high-dose psilocybin. The high-dose session constituted the first-ever psychedelic experience for all of the study participants. All except one participant rated their high-dose experience with psilocybin as the *single most unusual conscious state* of their entire lives, and the single exception rated it within their *top-five most unusual*. Significant changes in brain function and anatomy were evident from one-hour to one-month after high-dose psilocybin. No acute or long-term changes were apparent after a 1mg 'placebo' dose. Empirical modeling highlighted the predictive power of increased brain entropy under high-dose psilocybin, implying its role in manifesting not just the characteristic 'psychedelic' effects of psilocybin, but also its longer-term mental-health impact. More specifically, improved well-being could be predicted directly from acute increases in brain entropy as early as one-hour post dosing. Prediction could also be done through a sequence where increased brain entropy first predicted next day psychological insight, which then predicted the one-month improvements in well-being. If non-pharmacological variables are standardized as 'psychologically supportive' [37], these results imply that human brain changes as early as one-hour into a 25mg psilocybin experience— and that seem closely related to the subjective 'trip'— can predict mental health improvements one-month later.

Decreased axial diffusivity in prefrontal-subcortical tracts one-month after participants' first-ever high-dose of a psychedelic could be construed as potential anatomical 'neuroplasticity', echoing earlier in-vivo animal studies following single-dose psilocybin that reported increases in synaptic spine formation in female mice[5] and synaptic density in pigs[6]. However, interpretation of the axial diffusivity changes is complex, not least due to crossing fiber-related confounds[38]. The diffusion-weighted signal can change due to neurofibril or glial growth, altered myelination, axon density, membrane permeability or extracellular fluid [39]. Decreases in axial diffusivity have been observed with meditation[40], healthy neurodevelopment[41,42] and learning[43,44], but also with axonal injury[45,46], ageing and related pathology[47].

5-HT2AR agonism has been associated with *in vivo* dendro-architectural changes in adult mouse brain [5], axonal development in embryonic mouse brain [48], and oligodendrocyte changes in rodent brain tissue *in vitro* [49]. The high density of 5-HT2ARs in the human prefrontal cortex [50] and greater number of cortical compared with subcortical terminals[51] suggest a PFC 5-HT2A receptor locus of action for the observed diffusivity changes. The DTI results were robust to free-water correction, and the AD decreases correlated with decreased modularity at one-month post-25mg psilocybin, lending support to the inference that functionally relevant microstructural changes had occurred. Further research using multi-shell sequences is needed to disambiguate the current findings and inform on their robustness and replicability. Until then, serious caution is advised when interpreting the biological basis of these DTI findings. If the effects *are* microstructural, however, then decreases in AD and FA exclusively after the 25mg dose might tentatively relate to two types of change: 1) the pruning of weak or redundant connections; 2) neurogenesis with under-myelinated axons.

Decreased EEG alpha power, a well-replicated effect of psychedelics, is linked with cortical disinhibition [52] and increased brain entropy under psychedelics appears to be dose-dependent [53] and has been associated with acute and sub-acute psychological changes [54,55]. Decreased cortical alpha power and increased signal entropy are robust and reliable markers of the acute action of psychedelics in humans [20,21,55-57]. Future research in larger samples could test whether acute EEG changes, induced by 5-HT2AR agonist psychedelics, can predict downstream anatomical neuroplasticity [3] or therapeutically-relevant functional brain changes [27]. However, we failed to find a relationship between acute increases in entropic brain activity and the downstream DTI changes.

Compared with other outcomes in this study, the fMRI-measured brain changes assessed one-month post 25mg were relatively modest; for example, explorative ROI and network-based RSFC analyses (see supplement and methods), including network modularity, yielded largely non-significant results, albeit with some exceptions, including brain responsivity to facial expressions, amygdala RSFC, and a statistically significant *modularity change* to *well-being improvement* correlation (Figure 2C,D,F and Figures S8-17 in the supplement) as well as *modularity change* to *axial diffusivity change* (Figure 4B). Previous work assessing post psilocybin therapy changes in brain function in individuals with depression have revealed more robust changes [16,27-29]. For example, in two separate trials of psilocybin for depression,

decreased brain network modularity was seen that correlated with or predicted improvements in
symptom severity post-treatment [27] (see also [58]). Doss et al. found similar effects with
psilocybin-therapy for depression that they labelled increased "neural flexibility" [16]. In the
present study, changes in network modularity failed to reach statistical significance (Figure 2E)
but we did see a *network modularity change* vs *mental health change* relationship (Figure 2F,
4B) that was directionally consistent with previous work, i.e., decreased modularity correlated
with improved mental health [27].

Reductions in amygdala and salience-network responses to fearful emotional stimuli post-25mg
psilocybin were observed in the present study (Figure 2C&D and S8-11) and these are broadly
consistent with prior work in healthy volunteers [26]. Psychophysiological interaction results were
observed for the bilateral amygdala within the face paradigm (see S12-13). Since these
analyses were included on request, we refrain from a detailed discussion here but point the
reader to a relevant discussion in the supplementary file. Briefly, we note some similarities and
differences between a prior study examining amygdala PPI responses to emotional faces post
psilocybin-therapy for depression [59]. Also note, S10&11 (happy > neutral and fear) and two
prior psilocybin studies in depression [28,60] for evidence of *augmented* brain reactivity to
emotional face stimuli post-psilocybin-therapy, most reliably in the non-negative emotional
domains – i.e., happy and neutral faces. These latter results could be interpreted as consistent
with the re-opening critical periods for social learning [61]. However, some have questioned the
reliability of brain responsiveness to emotional faces as a biomarker of therapeutic action [62].

The present study's fMRI results suggest that enduring functional brain changes post psilocybin
are less robust and reliable in healthy versus mentally unwell populations e.g., [27]. If this principle
is reliable, it could imply that greater baseline atypicality in the clinical populations, primes for a
more robust remediating change via psilocybin. However, given the reliability of enduring
psychological changes post psilocybin in healthy samples [18,19], it remains plausible— if not
likely— that functional brain changes *do* occur in healthy populations but their detection is
dependent on— or sensitive to— experimental modality, metric and parameter choices [27,29,63]. In
short, we may have not yet found a sufficiently sensitive assay to detect true functional brain
changes.

The present multi-modal neuroimaging study in healthy participants sheds new light on the brain
effects of first-time high-dose psychedelic use and the therapeutic action of psilocybin-therapy,

suggesting, for the first time, how therapeutically relevant effects (i.e., improved well-being) can
be forecast via an *acute* human brain action, i.e., entropic brain activity, that is known to relate
to the subjective 'trip' [55]. Recent evidence suggests that this characteristic effect of psychedelics
is dose-dependent and somewhat exclusive to this category of drug versus psychoactive
stimulants and cannabinoids [53]. Long-term improvements in well-being were predicted by the
acute increases in entropic brain activity, temporally coinciding with the psilocybin's acute
subjective effects. Results support a role for psychological insight in mediating the causal
association between increased entropic brain activity and potentially enduring improvements in
well-being. These and prior results highlight psychological insight [31] and entropic brain effects
[64,65] as important to the action of psychedelic therapy. Possible white-matter changes, as well as
improvements in well-being and cognition were also observed in this study. All acute and
enduring psychological and neurobiological effects were exclusive to the high-dose psilocybin
condition.

Justified by prior evidence of enduring psychological changes post-psychedelics [9,10], this study
used a fixed-order cross-over design. Pairwise contrasts on all the study's main outcomes
showed no evidence of acute brain or behavioural effects or changes one-month post-1mg;
however, order confounds cannot be entirely discounted. Between-subject confirmatory studies
are now required to examine the reliability of these novel findings.

ACKNOWLEDGEMENTS

Funded by philanthropic donations to the Centre for Psychedelic Research, Imperial College London,
founded in April 2019. TL received an MRC-DTP Studentship, HMD received an Imperial College London
President's PhD Scholarships. RCH was supported by the Alex Mosley Charitable Trust between 2015
and April 2019. From April 2019 to July 2021, RCH was supported by donations to the Centre for
Psychedelic Research at Imperial College London, upon its creation; and from July 2021 to present, RCH
has been supported by The Ralph Metzner Distinguished Professorship at University of California San
Francisco. Some further support for this study was provided by The Beckley Foundation. Infrastructure
support was provided by the NIHR Imperial Biomedical Research Centre and the NIHR Imperial Clinical
Research Facility. The views expressed are those of the author(s) and not necessarily those of the NHS,
the NIHR or the Department of Health and Social Care.

We thank Manca Peskar, Victoria Nygart, Hannes Kettner and Mellissa Shukuroglou for their invaluable
assistance on study visits; Dr Roberta Murphy and Dr Jonny Martell for clinical oversight; Dr Rosalind
Watts for advice, training and support on guiding and integration procedures; Dr Robin Tyacke for

controlled drug oversight; Joe Peill, Gregory Cooper, Katie Trinci and Pablo Mallaroni for their support
with data analysis; and Bruna Giribaldi for advice and support.
468
AUTHOR CONTRIBUTIONS
RCH designed the study and wrote the original and revised manuscripts. RCH and DJN provided
oversight. TL coordinated the study, led the study visits, collected the data, created charts and plots, and
worked with RCH to compile the paper and revisions. MS coordinated the study to completion, leading
study visits and collecting data for the final set of participants. LE, LR, CT and FR supervised dosing
sessions and performed data collection. LE and DE provided medical cover and assisted with study visits.
TL analyzed the behavioral data and conducted regression and correlation analyses. FR and PMM ran
the clustering analyses of long-term changes, completed the mediation analysis, and created the
corresponding figures (Figures 4A-D). PMM and FR also created Figure 1A-B. LK analyzed the diffusion
MR data. LR, BAP, CT, MG, and MBW analyzed the functional MR data. LK created the diffusion images
and RCH organized them into plots. FR, PMM, FS, CT, MS, and HD analyzed the EEG data. AH and WT
guided our use of the IDED, which HD analyzed, and TL created charts for. BAP and RJZ led the PPI
analyses and created the relevant charts. All authors were given the opportunity to provide edits of the
manuscript and accepted the final draft.
483
DECLARATION OF INTERESTS
RC-H is a scientific advisor to Entheos Labs, Journey Collab, Otsuka, and Mindstate and has received
fees or share from these companies. **DJN** is Chief Research Officer for Awakn Life Sciences and, within
the past 2 years, has been a scientific advisor to COMPASS Pathways, Neural therapeutics, Algernon
pharmaceuticals and Alvarius. **RJZ** and **BAP** are postdoctoral fellows in the NYU Langone Psychedelic
Medicine Research Training program funded by MindMed. **MBW's** primary employer is Invicro LLC., a
contract research organization which provides services to the pharmaceutical and biotechnology
industries. **DE** has, within the last 2 years, received fees for scientific advisory work from the following
(novel psychedelic) companies: Mydecine, Field Trip Health, SmallPharma Ltd, Aya Biosciences,
Clerkenwell Health, and Mindstate Design Lab, and has received an honorarium fee from each of
Compass Pathways and Lundbeck for a talk about psychedelic science.
495
496
497

REFERENCES

Becker, A. M. *et al.* Ketanserin reverses the acute response to LSD in a randomized, double-blind, placebo-controlled, crossover study in healthy subjects. *Int J Neuropsychopharmacol* (2022). https://doi.org/10.1093/ijnp/pyac075

Vollenweider, F. X., Vollenweider-Scherpenhuyzen, M. F., Babler, A., Vogel, H. & Hell, D. Psilocybin induces schizophrenia-like psychosis in humans via a serotonin-2 agonist action. *Neuroreport* **9**, 3897-3902 (1998).

Ly, C. *et al.* Psychedelics Promote Structural and Functional Neural Plasticity. *Cell Rep* **23**, 3170-3182 (2018). https://doi.org/10.1016/j.celrep.2018.05.022

Vaidya, V. A., Marek, G. J., Aghajanian, G. K. & Duman, R. S. 5-HT2A receptor-mediated regulation of brain-derived neurotrophic factor mRNA in the hippocampus and the neocortex. *J Neurosci* **17**, 2785-2795 (1997).

Shao, L. X. *et al.* Psilocybin induces rapid and persistent growth of dendritic spines in frontal cortex in vivo. *Neuron* **109**, 2535-2544 e2534 (2021). https://doi.org/10.1016/j.neuron.2021.06.008

Raval, N. R. *et al.* A Single Dose of Psilocybin Increases Synaptic Density and Decreases 5-HT2A Receptor Density in the Pig Brain. *Int J Mol Sci* **22** (2021). https://doi.org/10.3390/ijms22020835

Vargas, M. V. *et al.* Psychedelics promote neuroplasticity through the activation of intracellular 5-HT2A receptors. *Science* **379**, 700-706 (2023). https://doi.org/10.1126/science.adf0435

Moliner, R. *et al.* Psychedelics promote plasticity by directly binding to BDNF receptor TrkB. *Nat Neurosci* **26**, 1032-1041 (2023). https://doi.org/10.1038/s41593-023-01316-5

Griffiths, R. R. *et al.* Psilocybin produces substantial and sustained decreases in depression and anxiety in patients with life-threatening cancer: A randomized double-blind trial. *J Psychopharmacol* **30**, 1181-1197 (2016). https://doi.org/10.1177/0269881116675513

Ross, S. *et al.* Rapid and sustained symptom reduction following psilocybin treatment for anxiety and depression in patients with life-threatening cancer: a randomized controlled trial. *J Psychopharmacol* **30**, 1165-1180 (2016). https://doi.org/10.1177/0269881116675512

Carhart-Harris, R. *et al.* Trial of Psilocybin versus Escitalopram for Depression. *N Engl J Med* **384**, 1402-1411 (2021). https://doi.org/10.1056/NEJMoa2032994

Carhart-Harris, R. L. *et al.* Psilocybin with psychological support for treatment-resistant depression: six-month follow-up. *Psychopharmacology (Berl)* **235**, 399-408 (2018). https://doi.org/10.1007/s00213-017-4771-x

Goodwin, G. M. *et al.* Single-Dose Psilocybin for a Treatment-Resistant Episode of Major Depression. *N Engl J Med* **387**, 1637-1648 (2022). https://doi.org/10.1056/NEJMoa2206443

Bogenschutz, M. P. *et al.* Percentage of Heavy Drinking Days Following Psilocybin-Assisted Psychotherapy vs Placebo in the Treatment of Adult Patients With Alcohol Use Disorder: A Randomized Clinical Trial. *JAMA Psychiatry* (2022). https://doi.org/10.1001/jamapsychiatry.2022.2096

Griffiths, R., Richards, W., Johnson, M., McCann, U. & Jesse, R. Mystical-type experiences occasioned by psilocybin mediate the attribution of personal meaning and

544		spiritual significance 14 months later. *Journal of psychopharmacology (Oxford, England)*
545		**22**, 621-632 (2008). https://doi.org/10.1177/0269881108094300
546	16	Doss, M. K. *et al.* Psilocybin therapy increases cognitive and neural flexibility in patients
547		with major depressive disorder. *Transl Psychiatry* **11**, 574 (2021).
548		https://doi.org/10.1038/s41398-021-01706-y
549	17	Wiessner, I. *et al.* LSD, afterglow and hangover: Increased episodic memory and verbal
550		fluency, decreased cognitive flexibility. *Eur Neuropsychopharmacol* **58**, 7-19 (2022).
551		https://doi.org/10.1016/j.euroneuro.2022.01.114
552	18	Mans, K. *et al.* Sustained, Multifaceted Improvements in Mental Well-Being Following
553		Psychedelic Experiences in a Prospective Opportunity Sample. *Front Psychiatry* **12**,
554		647909 (2021). https://doi.org/10.3389/fpsyt.2021.647909
555	19	Mason, N. L., Mischler, E., Uthaug, M. V. & Kuypers, K. P. C. Sub-Acute Effects of
556		Psilocybin on Empathy, Creative Thinking, and Subjective Well-Being. *J Psychoactive*
557		*Drugs* **51**, 123-134 (2019). https://doi.org/10.1080/02791072.2019.1580804
558	20	Kometer, M., Schmidt, A., Jancke, L. & Vollenweider, F. X. Activation of serotonin 2A
559		receptors underlies the psilocybin-induced effects on alpha oscillations, N170 visual-
560		evoked potentials, and visual hallucinations. *The Journal of neuroscience : the official*
561		*journal of the Society for Neuroscience* **33**, 10544-10551 (2013).
562		https://doi.org/10.1523/jneurosci.3007-12.2013
563	21	Valle, M. *et al.* Inhibition of alpha oscillations through serotonin-2A receptor activation
564		underlies the visual effects of ayahuasca in humans. *Eur Neuropsychopharmacol* **26**,
565		1161-1175 (2016). https://doi.org/10.1016/j.euroneuro.2016.03.012
566	22	Kringelbach, M. L. *et al.* Dynamic coupling of whole-brain neuronal and
567		neurotransmitter systems. *Proc Natl Acad Sci U S A* **117**, 9566-9576 (2020).
568		https://doi.org/10.1073/pnas.1921475117
569	23	Madsen, M. K. *et al.* Psychedelic effects of psilocybin correlate with serotonin 2A
570		receptor occupancy and plasma psilocin levels. *Neuropsychopharmacology* **44**, 1328-
571		1334 (2019). https://doi.org/10.1038/s41386-019-0324-9
572	24	Carhart-Harris, R. L. *et al.* Neural correlates of the LSD experience revealed by
573		multimodal neuroimaging. *Proceedings of the National Academy of Sciences of the*
574		*United States of America* **113**, 4853-4858 (2016).
575		https://doi.org/10.1073/pnas.1518377113
576	25	McCulloch, D. E. *et al.* Lasting effects of a single psilocybin dose on resting-state
577		functional connectivity in healthy individuals. *J Psychopharmacol* **36**, 74-84 (2022).
578		https://doi.org/10.1177/02698811211026454
579	26	Barrett, F. S., Doss, M. K., Sepeda, N. D., Pekar, J. J. & Griffiths, R. R. Emotions and
580		brain function are altered up to one month after a single high dose of psilocybin. *Sci Rep*
581		**10**, 2214 (2020). https://doi.org/10.1038/s41598-020-59282-y
582	27	Daws, R. E. *et al.* Increased global integration in the brain after psilocybin therapy for
583		depression. *Nat Med* **28**, 844-851 (2022). https://doi.org/10.1038/s41591-022-01744-z
584	28	Roseman, L., Demetriou, L., Wall, M. B., Nutt, D. J. & Carhart-Harris, R. L. Increased
585		amygdala responses to emotional faces after psilocybin for treatment-resistant depression.
586		*Neuropharmacology* **142**, 263-269 (2018).
587		https://doi.org/10.1016/j.neuropharm.2017.12.041
588	29	Carhart-Harris, R. L. *et al.* Psilocybin for treatment-resistant depression: fMRI-measured
589		brain mechanisms. *Sci Rep* **7**, 13187 (2017). https://doi.org/10.1038/s41598-017-13282-7

30   Pasquini, L., Palhano-Fontes, F. & Araujo, D. B. Subacute effects of the psychedelic
ayahuasca on the salience and default mode networks. *J Psychopharmacol* **34**, 623-635
(2020). https://doi.org/10.1177/0269881120909409
31   Peill, J. M. *et al.* Validation of the Psychological Insight Scale: A new scale to assess
psychological insight following a psychedelic experience. *Journal of*
psychopharmacology (Oxford, England) **36**, 31-45 (2022).
https://doi.org/10.1177/02698811211066709
32   Tennant, R. *et al.* The Warwick-Edinburgh Mental Well-being Scale (WEMWBS):
development and UK validation. *Health Qual Life Outcomes* **5**, 63 (2007).
https://doi.org/10.1186/1477-7525-5-63
33   Roberts, A. C., Robbins, T. W. & Everitt, B. J. The effects of intradimensional and
extradimensional shifts on visual discrimination learning in humans and non-human
primates. *Q J Exp Psychol B* **40**, 321-341 (1988).
34   Owen, A. M., Roberts, A. C., Polkey, C. E., Sahakian, B. J. & Robbins, T. W. Extra-
dimensional versus intra-dimensional set shifting performance following frontal lobe
excisions, temporal lobe excisions or amygdalo-hippocampectomy in man.
Neuropsychologia **29**, 993-1006 (1991). https://doi.org/10.1016/0028-3932(91)90063-e
35   Baron, R. M. & Kenny, D. A. The moderator-mediator variable distinction in social
psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc*
Psychol **51**, 1173-1182 (1986). https://doi.org/10.1037//0022-3514.51.6.1173
36   Hayes, A. F. & Rockwood, N. J. Regression-based statistical mediation and moderation
analysis in clinical research: Observations, recommendations, and implementation. *Behav*
Res Ther **98**, 39-57 (2017). https://doi.org/10.1016/j.brat.2016.11.001
37   Carhart-Harris, R. L. *et al.* Psychedelics and the essential importance of context. *J*
Psychopharmacol **32**, 725-731 (2018). https://doi.org/10.1177/0269881118754710
38   Figley, C. R. *et al.* Potential Pitfalls of Using Fractional Anisotropy, Axial Diffusivity,
and Radial Diffusivity as Biomarkers of Cerebral White Matter Microstructure. *Front*
Neurosci **15**, 799576 (2021). https://doi.org/10.3389/fnins.2021.799576
39   Li, X., Bai, Y., Liao, Y. & Xin, S. X. Assessment of the effects of mimicking tissue
microstructural properties on apparent diffusion coefficient and apparent exchange rate in
diffusion MRI via a series of specially designed phantoms. *Magn Reson Med* **87**, 292-301
(2022). https://doi.org/10.1002/mrm.28990
40   Tang, Y. Y., Lu, Q., Fan, M., Yang, Y. & Posner, M. I. Mechanisms of white matter
changes induced by meditation. *Proc Natl Acad Sci U S A* **109**, 10570-10574 (2012).
https://doi.org/10.1073/pnas.1207817109
41   Cascio, C. J., Gerig, G. & Piven, J. Diffusion tensor imaging: Application to the study of
the developing brain. *J Am Acad Child Adolesc Psychiatry* **46**, 213-223 (2007).
https://doi.org/10.1097/01.chi.0000246064.93200.e8
42   Kumar, R., Nguyen, H. D., Macey, P. M., Woo, M. A. & Harper, R. M. Regional brain
axial and radial diffusivity changes during development. *J Neurosci Res* **90**, 346-355
(2012). https://doi.org/10.1002/jnr.22757
43   Sagi, Y. *et al.* Learning in the fast lane: new insights into neuroplasticity. *Neuron* **73**,
1195-1203 (2012). https://doi.org/10.1016/j.neuron.2012.01.025
44   Huber, E., Donnelly, P. M., Rokem, A. & Yeatman, J. D. Rapid and widespread white
matter plasticity during an intensive reading intervention. *Nat Commun* **9**, 2260 (2018).
https://doi.org/10.1038/s41467-018-04627-5

45   Budde, M. D., Xie, M., Cross, A. H. & Song, S. K. Axial diffusivity is the primary
637   correlate of axonal injury in the experimental autoimmune encephalomyelitis spinal cord:
638   a quantitative pixelwise analysis. *J Neurosci* **29**, 2805-2813 (2009).
639   https://doi.org/10.1523/JNEUROSCI.4605-08.2009
640   46   Carhart-Harris, R. L. & Nutt, D. J. Serotonin and brain function: a tale of two receptors. *J*
641   *Psychopharmacol* **31**, 1091-1120 (2017). https://doi.org/10.1177/0269881117725915
642   47   Sexton, C. E. *et al.* Accelerated changes in white matter microstructure during aging: a
643   longitudinal diffusion tensor imaging study. *J Neurosci* **34**, 15425-15436 (2014).
644   https://doi.org/10.1523/JNEUROSCI.0203-14.2014
645   48   Vicenzi, S., Foa, L. & Gasperini, R. J. Serotonin functions as a bidirectional guidance
646   molecule regulating growth cone motility. *Cell Mol Life Sci* **78**, 2247-2262 (2021).
647   https://doi.org/10.1007/s00018-020-03628-2
648   49   Fan, L. W. *et al.* Exposure to serotonin adversely affects oligodendrocyte development
649   and myelination in vitro. *J Neurochem* **133**, 532-543 (2015).
650   https://doi.org/10.1111/jnc.12988
651   50   Beliveau, V. *et al.* A High-Resolution In Vivo Atlas of the Human Brain's Serotonin
652   System. *The Journal of neuroscience : the official journal of the Society for Neuroscience*
653   **37**, 120-128 (2017). https://doi.org/10.1523/jneurosci.2830-16.2016
654   51   Sherman, S. M. & Guillery, R. W. On the actions that one nerve cell can have on another:
655   distinguishing "drivers" from "modulators". *Proc Natl Acad Sci U S A* **95**, 7121-7126
656   (1998). https://doi.org/10.1073/pnas.95.12.7121
657   52   Jensen, O. & Mazaheri, A. Shaping functional architecture by oscillatory alpha activity:
658   gating by inhibition. *Front Hum Neurosci* **4**, 186 (2010).
659   https://doi.org/10.3389/fnhum.2010.00186
660   53   Murray, C. H. *et al.* Neural complexity is increased after low doses of LSD, but not
661   moderate to high doses of oral THC or methamphetamine. *Neuropsychopharmacology*
662   (2024). https://doi.org/10.1038/s41386-024-01809-2
663   54   Lebedev, A. V. *et al.* LSD-induced entropic brain activity predicts subsequent personality
664   change. *Hum Brain Mapp* **37**, 3203-3213 (2016). https://doi.org/10.1002/hbm.23234
665   55   Timmermann, C. *et al.* Human brain effects of DMT assessed via EEG-fMRI. *Proc Natl*
666   *Acad Sci U S A* **120**, e2218949120 (2023). https://doi.org/10.1073/pnas.2218949120
667   56   Timmermann, C. *et al.* Neural correlates of the DMT experience assessed with
668   multivariate EEG. *Sci Rep* **9**, 16324 (2019). https://doi.org/10.1038/s41598-019-51974-4
669   57   Schartner, M. M., Carhart-Harris, R. L., Barrett, A. B., Seth, A. K. &
670   Muthukumaraswamy, S. D. Increased spontaneous MEG signal diversity for
671   psychoactive doses of ketamine, LSD and psilocybin. *Sci Rep* **7**, 46421 (2017).
672   https://doi.org/10.1038/srep46421
673   58   Deco, G. *et al.* Different hierarchical reconfigurations in the brain by psilocybin and
674   escitalopram for depression. *Nature Mental Health* **2**, 1096-1110 (2024).
675   https://doi.org/10.1038/s44220-024-00298-y
676   59   Mertens, L. J. *et al.* Therapeutic mechanisms of psilocybin: Changes in amygdala and
677   prefrontal functional connectivity during emotional processing after psilocybin for
678   treatment-resistant depression. *J Psychopharmacol* **34**, 167-180 (2020).
679   https://doi.org/10.1177/0269881119895520

60   Wall, M. B. *et al.* Reduced brain responsiveness to emotional stimuli with escitalopram
but not psilocybin therapy for depression. *medRxiv*, 2023.2005.2029.23290667 (2023).
https://doi.org/10.1101/2023.05.29.23290667
61   Nardou, R. *et al.* Psychedelics reopen the social reward learning critical period. *Nature*
618, 790-798 (2023). https://doi.org/10.1038/s41586-023-06204-3
62   Nord, C. L., Gray, A., Charpentier, C. J., Robinson, O. J. & Roiser, J. P. Unreliability of
putative fMRI biomarkers during emotional face processing. *Neuroimage* **156**, 119-127
(2017). https://doi.org/10.1016/j.neuroimage.2017.05.024
63   Carhart-Harris, R. L. *et al.* Canalization and plasticity in psychopathology.
Neuropharmacology **226**, 109398 (2022).
https://doi.org/10.1016/j.neuropharm.2022.109398
64   Carhart-Harris, R. L. *et al.* The entropic brain: a theory of conscious states informed by
neuroimaging research with psychedelic drugs. *Front Hum Neurosci* **8**, 20 (2014).
https://doi.org/10.3389/fnhum.2014.00020
65   Carhart-Harris, R. L. The entropic brain - revisited. *Neuropharmacology* **142**, 167-178
(2018). https://doi.org/10.1016/j.neuropharm.2018.03.010
696