Structured stabilization in recurrent neural circuits through inhibitory synaptic plasticity

Dylan Festa   Claudia Cusseddu   Julijana Gjorgjieva

School of Life Sciences, Technical University of Munich, Freising, Germany

Inhibitory interneurons play a dual role in recurrently connected biological circuits: they regulate global neural activity to prevent runaway excitation, and contribute to complex neural computations. While the first role can be achieved through unstructured connections tuned for homeostatic rate stabilization, computational tasks often require structured excitatory-inhibitory (E/I) connectivity. Here, we consider a broad class of pairwise inhibitory spike-timing dependent plasticity (iSTDP) rules, demonstrating how inhibitory synapses can self-organize to both stabilize excitation and generate functionally relevant connectivity structures — a process we call "structured stabilization". We show that in both E/I circuit motifs and large spiking recurrent neural networks the choice of iSTDP rule can lead to either mutually connected E/I pairs, or to lateral inhibition, where an inhibitory neuron connects to an excitatory neuron that does not directly connect back to it. In a one-dimensional ring network, if two inhibitory populations follow these distinct forms of iSTDP, the effective connectivity within the excitatory population self-organizes into a Mexican-hat-like profile with excitatory influence in the center and inhibitory influence away from the center. This leads to emergent dynamical properties such as surround suppression and modular spontaneous activity. Our theoretical work introduces a family of rules that retains the broad applicability and simplicity of spike-timing-based plasticity, while promoting structured, self-organized stabilization. These findings highlight the rich interplay between iSTDP rules, circuit structure, and neuronal dynamics, offering a framework for understanding how inhibitory plasticity shapes network function.

1   Introduction

In cortical circuits, GABA-ergic interneurons constitute approximately 10% to 20% of the total neural population, and can be further divided into subtypes that exhibit distinct characteristics in connectivity, plasticity and tuning properties (Jiang et al. 2015; Tremblay et al. 2016). However, fully understanding how this diverse neural population influences neural circuit function remains a significant challenge.

At a fundamental level, a global dynamical balance between excitation and inhibition can explain the irregular, Poisson-like firing patterns observed *in vivo* in cortical circuits (Van Vreeswijk and Sompolinsky 1998; Brunel 2000). This balanced regime enhances the circuit's sensitivity to input, allowing for rapid processing of information (Tsodyks and Sejnowski 1995). It further supports and regulates other key computational tasks, such as the tuning of receptive fields (Wallenstein and Hasselmo 1997; Froemke et al. 2007; Dorrn et al. 2010; Kleberg et al. 2014; Rubin et al. 2017), the formation of neural assemblies (Sadeh and Clopath 2021), and the storage of precise firing sequences (Maes et al. 2021; Riquelme et al. 2023; Gong and Brunel 2024).

Dynamical excitatory-inhibitory (E/I) balance can be achieved without any specific inhibitory connectivity structures. Often circuit models consider inhibition simply as a "blanket" that regulates excitation, and assimilate it into a global normalization term or a correction to the synaptic weights between excitatory neurons (Jin 2002; Ravid Tannenbaum and Burak 2016; Montangie et al. 2020). While this simplified approach captures fundamental properties of neural dynamics, it overlooks the possible advantages of more specific E/I connectivity patterns. Incorporating structured inhibitory connections can further refine computations and mediate neural functions such as E/I co-tuning (Clopath et al. 2016; Mackwood et al. 2021; Miehl and Gjorgjieva 2022; Eckmann et al. 2024), memory formation (Bergoin et al. 2023; Miehl et al. 2023; Agnes and Vogels 2024), predictive coding (Schulz et al. 2021), and shaping of neural dynamics (Mongillo et al. 2018; Dahmen et al. 2020; Sadeh and Clopath 2020; Hafizi et al. 2022).

Experimental findings demonstrate that inhibitory neurons form non-random connectivity structures not explainable by connectivity rules that purely depend on distance or morphology. These structures vary across brain areas and depend on interneuron subtypes. (Peng et al. 2021; Bollmann et al. 2023; Reimann et al. 2023; Wu et al. 2023; Kuan et al. 2024; Znamenskiy et al. 2024). In the neonatal cortex, even before the onset of external

sensory stimuli, structured inhibition shapes circuit development (Bollmann et al. 2023). For instance, effective connectivity in the form of a Mexican-hat with short-range excitatory effects and longer-range inhibition results in an interesting signature of developmental dynamics: widespread modular activity which propagates throughout the sensory cortex (Smith et al. 2018; Mulholland et al. 2021).

Our knowledge of the mechanisms and principles that enable the formation of inhibitory connectivity structures, and their persistence over time, is still lacking. Genetic mechanisms play a role, especially during developmental stages (Duan et al. 2020; Hanganu-Opatz et al. 2021; Wu et al. 2023). A second, equally important mechanism behind the establishment of specific connectivity patterns is synaptic plasticity triggered by neural activity (Sreenivasan et al. 2022). This mechanism offers the advantage of a closed-loop regulation of neural dynamics: connections can be determined and tuned based on activity, with higher precision in compact cortical circuits and flexibility in the face of change.

Although some theoretical studies considered the impact of different forms of iSTDP in feedforward circuit architectures with a single postsynaptic neuron (Kleberg et al. 2014; Luz and Shamir 2014), there is still no general understanding about the effects of pairwise iSTDP on structure formation and at the circuit level. Theoretical work proposed a iSTDP rule that guarantees the homeostatic regulation of neural rates (Vogels et al. 2011), becoming the "canonical" choice for plastic circuits. However, this rule is not fully concerned with structure and tends to over-regulate rates, producing narrow rate distributions in the excitatory population. In other work, inhibitory rules are based on experimental measurements (Lagzi et al. 2021; Bergoin et al. 2024), derived normatively from computation objectives (Mackwood et al. 2021), or are found through numerical optimization on a large parameter space (Confavreux et al. 2024). Here, we propose a unifying approach by analyzing a broad family of pairwise iSTDP rules, dissecting the principles and the mechanisms that contribute to excitatory regulation and structure formation, and testing them on large scale recurrent neuronal networks (RNNs) of spiking neurons.

We first derive analytically how the specific choice of iSTDP rule can determine the relevant activity components driving the formation of inhibitory connections. On one extreme, rules dominated by firing rates are weakly influenced by structure, and pursue rigidly the goal of homeostatic rate regulation. These rules promote "blanket" inhibition. At the opposite end of the spectrum, we find covariance-dominated rules, strongly influenced by the pairwise spiking statistics that, in the absence of external inputs, arise from the underlying circuit connectivity. In this case, while stabilizing the circuit, inhibition also forms specific motifs, such as mutually connected E/I pairs or lateral inhibition, an effect that we call "structured stabilization". We confirm these results through numerical simulations in large RNNs with random connectivity. We further explore the functional implications of these iSTDP rules by building a RNN where the excitatory connectivity follows a one-dimensional ring structure and demonstrate that inhibition can self-organize, leading to a Mexican-hat effective connectivity. This leads to emergent properties such as surround suppression and long-range "modular" correlations of spontaneous activity. In sum, our work explores how families of inhibitory plasticity rules shape different connectivity motifs involving inhibition, resulting in large-scale connectivity structures and emergent computations.

We initially considered a general iSTDP rule that depends exclusively on local spike times applied to a synapse between an inhibitory presynaptic and an excitatory postsynaptic neuron (inh-to-exc) (Kempter et al. 1999; Luz and Shamir 2014; Akil et al. 2021). The rule contains both fixed components triggered by each pre- and postsynaptic firing events, denoted as α_{pre} and α_{post}, and a kernel function, $L(\tau)$ triggered by the co-occurrence of pre-post ($\tau > 0$) or post-pre ($\tau < 0$) spiking events (see Methods, Section 4.1). Under these rules, the mean-field equation for the dynamics of an average synaptic weight is:

$$\frac{d\langle W \rangle}{dt} = \alpha_{\text{pre}} r_{\text{pre}} + \alpha_{\text{post}} r_{\text{post}} + B \, r_{\text{pre}} \, r_{\text{post}} + Q \quad,$$

$$\text{with} \quad B := \int_{-\infty}^{+\infty} d\tau \, L(\tau) \quad \text{and} \quad Q := \int_{-\infty}^{+\infty} d\tau \, L(\tau) \, C(\tau) \; ; \tag{1}$$

where r_{pre} and r_{post} are the pre- and post-synaptic firing rates, B is the net area of the iSTDP kernel $L(\tau)$, and Q captures the effect of the iSTDP kernel on the pre-post spike train pairwise interactions, conveyed by the covariance density curve $C(\tau)$ (see Supplementary Material, or Kempter et al. 1999; Luz and Shamir 2014; Lagzi and Fairhall 2024).

2 Results

Figure 1: Effects of plasticity for E/I motifs in a reduced model. **A.** Schematics of the reduced circuit model. Two Poisson units, one excitatory and one inhibitory, fire with varying probability, each incoming spike is convolved with an exponential synaptic kernel that either increases or decreases the instantaneous firing probability. Only the inhibitory synapse (dashed line) is plastic. **B,C.** The circuit is simulated in two configurations: unidirectional (U), with $w_{\mathrm{exc}} = 0$ and mutual (M), with $w_{\mathrm{exc}} = 0.5$. The plots show the numerical covariance density between excitatory and inhibitory neuron before plasticity is activated. **D,E,F.** Pairwise interaction kernels of the three iSTDP rules tested (Eq. 1). We used the following numerical parameters. **D.** $B = 1$, $\alpha_{\mathrm{pre}} = -25$, $\alpha_{\mathrm{post}} = 0$, $\tau_+ = 50$ ms. **E.** $B = 0$, $\alpha_{\mathrm{pre}} = -0.12$, $\alpha_{\mathrm{post}} = 1.45$, $\tau_+ = 50$ ms, $\tau_- = 500$ ms. **F.** $B = 0$, $\alpha_{\mathrm{pre}} = -0.14$, $\alpha_{\mathrm{post}} = 7.95$, $\tau_+ = 30$ ms, $\tau_- = 30$ ms. **G,H,I.** Change in inhibitory synaptic weight over time for the M (brown) and U (cyan) circuit configurations, under the rules above each column. Dashed lines are analytic predictions for the fixed point (Methods, Eq. 10). **J,K,L.** Change in excitatory firing rate over time for the M (dark blue) and U (dark green) configurations, under the rules above. Dashed horizontal lines are the analytic predictions for the fixed-point (Methods, Eq. 10).

2.1 The influence of inhibitory plasticity on circuit structure

To understand how inhibitory plasticity influences circuit structure, first, we developed an analytical framework, defining a family of iSTDP rules. Using these rules, we aimed to derive the closed-form solution for a reduced E/I motif, and demonstrate how the shape of iSTDP affects the learned inhibitory connections.

Several terms in Eq. 1 depend solely on pre- and post-synaptic firing rates. Furthermore, in the general category of rules where B is high, the contribution of the $B\,r_{\mathrm{pre}}\,r_{\mathrm{post}}$ term can easily be an order of magnitude above Q, even in the presence of strong pre-post correlations (see Methods, Section 4.3). Therefore, when

$Q \ll B\, r_{\mathrm{pre}}\, r_{\mathrm{post}}$, the plasticity dynamics of Eq. 1 are strongly dominated by the firing rates. We call this plasticity regime "rate-dominated". The previously established and well-studied rate-homeostatic iSTDP rule (Vogels et al. 2011) is a rate-dominated rule with $\alpha_{\mathrm{post}} = 0$. This rule drives the inhibitory weight to adapt continuously so that the postsynaptic rate stays close to the target value $\tilde{r}_{\mathrm{post}} \approx -\alpha_{\mathrm{pre}}/B$, corresponding to a stable fixed point solution in Eq. 1.

While a large value of B simplifies the analysis by allowing us to neglect the covariance-based terms, this only captures a small fraction of the possible plasticity rules. We next used our framework to explore a broader range of plasticity rules, including those where the iSTDP kernel has zero or near-zero net area ($B \approx 0$). This amplifies the contribution of the second-order spike-timing correlations between pre- and postsynaptic neuron in the Q term. We refer to this class of rules as "covariance-dominated". Covariance-dominated rules include both synaptic potentiation and depression in their iSTDP kernel, so that their net area is zero. Examples are symmetric kernels with strong short-range potentiation and weak long-range depression (as in Fig. 1E), or antisymmetric kernels where pre-post and post-pre interactions are opposite in sign and balance each other (Fig. 1F).

Analyzing the full dynamics of Eq. 1 is challenging because the covariance $C(\tau)$ generally depends on the synaptic weights themselves. We therefore considered a minimal two-neuron E/I circuit motif (Fig. 1A) with fixed exc-to-inh connection weight (w_{exc}) but plastic inh-to-exc synapse (w_{inh}). We modeled the neurons as Poisson processes, where each incoming spike changed the instantaneous spiking probability through an excitatory or inhibitory post-synaptic potential (EPSP/IPSP) kernel (internal traces shown in Fig. 1A, see also Methods Section 4.4). The inhibitory neuron's firing rate was kept constant by an external homeostatic mechanism. The excitatory neuron received an excitatory fixed external input in addition to the inhibitory input. Using this simplified model, we derived analytically a closed-form expression for the fixed-point of the iSTDP dynamics (Methods, Eq. 10; Supplementary Material).

With this analytical framework in hand, we investigated how different iSTDP rules affect the resulting inhibitory synaptic strength and the excitatory postsynaptic activity. We considered two configurations within our E/I circuit motif: (1) a unidirectional configuration (U, Fig. 1B), where the weight from the excitatory to the inhibitory neuron, w_{exc} was set to zero, resulting in initial uncorrelated activity between the two neurons; and (2) a mutual configuration (M, Fig. 1C), where a strong weight from the excitatory to the inhibitory neuron, w_{exc} introduced a positive post-pre correlation in the spiking activity (compare covariance density plots in Fig. 1B and C). We considered the behavior of three iSTDP rules under both U and M configurations: a rate-dominated rule with a symmetric interaction kernel (Fig. 1D), a covariance-dominated rule also with a symmetric kernel (Fig. 1E), and a covariance-dominated rule with an antisymmetric kernel (Fig. 1F). As expected, the rate-dominated rule exhibited minimal sensitivity to the presence of the mutual connection. In both the U and the M configurations, the inhibitory synapse regulated the excitatory firing rate towards the target rate dictated by the rate-dependent components of the rule $\tilde{r}_{\mathrm{exc}} \approx -\alpha_{\mathrm{pre}}/B$ (Fig. 1G). Minor deviations from this target arose from the non-zero contribution of the Q term.

In contrast, the covariance-dominated rules exhibited strong dependence on the initial connectivity structure and on the choice of iSTDP kernel. With the symmetric kernel, the final inhibitory weight was significantly higher in the M configuration compared to the U configuration (Fig. 1H, compare brown and cyan). The effect was reversed for the antisymmetric kernel, where the final inhibitory weight was much lower in the M configuration (Fig. 1I). Notably, these rules can regulate the excitatory rate, but with high selectivity to structure. We call this effect "structured stabilization". The emergence of structured inhibition arises from the interplay between the iSTDP kernel and the second-order statistics of the pre-post neuron pair. In the M configuration, the presence of a reciprocal excitatory-to-inhibitory connection generated an increase in correlated post-pre events at short timescales (Fig. 1C). For symmetric iSTDP, these correlated events fell within the positive left branch of the pairwise interaction kernel, leading to synaptic potentiation (compare Fig. 1C and E). Conversely, with the antisymmetric iSTDP rule, the same correlated events fell within the negative region of the kernel, resulting in synaptic depression (compare Fig. 1C and F). The fixed components of the iSTDP rule, α_{pre} and α_{post}, were also crucial for ensuring that the excitatory firing rate converged to the desired target rate (set to $25\mathrm{Hz}$) in the preferred motif configuration.

In sum, we provide a general framework for investigating how different iSTDP rules shape both inhibitory strength and circuit structure.

Figure 2: Parametric study of different shapes of iSTDP kernel in the reduced 2D model. **A**. Representation of four possible shapes of iSTDP kernel, labeled by their effects in the covariance-dominated regime (i.e. with near-zero rate-dependent terms). **B**. Parametric study based on the analytic solution for each of the four iSTDP kernels, varying α_{post} and the net area under the iSTDP kernel, B (see Eq. 1). The color map represents the level of inhibition, where 1 indicates that the excitatory neuron is completely silenced, and 0 indicates no inhibition, thus the excitatory firing rate matches the input current **C-E**. Final inhibitory weight and final excitatory rate as a function of the incoming excitatory weight for three different iSTDP rules. Dots are numerical simulations, gray lines are analytic results (Methods, Eq. 10), color shades indicate different input levels. **C**. Results for the symmetric, self-stabilizing rule. The green dots represent the same conditions shown in Fig. 1H,K. **D**. Results for the antisymmetric, self-stabilizing rule. The green dots represent the same conditions shown in Fig. 1I,L. **E**. Results for the symmetric rule in a rate-dominated regime.

2.2 The shape of inhibitory STDP affects inhibitory structure

Having established that covariance-dominated rules can achieve structured stabilization, we next investigated how the specific form of the iSTDP kernel shapes the learned inhibitory connectivity. We considered four kernel shapes: symmetric, antisymmetric, and their "inverted" counterparts, simply obtained by reversing the sign of the kernel (Fig. 2A). We analyzed the behavior of these kernels in the reduced two-neuron E/I motif. To explore the parameter space, we varied the rate-dependent parameters B and α_{pre}, while keeping $\alpha_{post} = 0$. With this parametrization, the covariance-dominated regime corresponds to values of B and α_{pre} close to zero.

Our analysis confirmed that both symmetric and antisymmetric kernels yield stable fixed-points in the covariance-dominated regime (Fig. 2A,B, top row, and Supplementary Fig. S1C,D). These kernels, which we term "self-

stabilizing" exhibit robustness to parameter fluctuations and reliably converge to a stable inhibitory weight. In contrast, the inverted antisymmetric and the inverted symmetric rule proved unstable in the covariance-dominated regime, with inhibitory weights either decaying to zero or saturating depending on the initial conditions. We categorize these as "saturating" kernels (Fig. 2A, bottom panels, and Fig. S1E,F in Supplementary Material). Saturating kernels can still reach stability with strong rate-dependent terms, which reduce the overall influence of the kernel shape on the final weight.

This difference in stability arises from how the kernels interact with changes in pre- and postsynaptic correlations. Consider a synapse operating under a covariance-dominated plasticity rule at its fixed point. At equilibrium, potentiation and depression events balance exactly. A perturbation that strengthens the inhibitory weight leads to a reduction in pre-post firing events due to the increased inhibition. For self-stabilizing kernels, which have a positive right branch (corresponding to pre-post events), this reduction in correlation reduces the occurrence of synaptic potentiation events, shifting the balance in favor of synaptic depression. The synapse therefore depresses, eventually compensating the initial perturbation from equilibrium. Conversely, saturating kernels show the opposite effect: since their kernel is negative in the pre-post region, a reduction of pre-post correlation results in fewer synaptic depression events, so the initial balance is further shifted in favor of synaptic potentiation, amplifying the perturbation and leading the synapse towards saturation. Finally, in rate-dominated rules, the specific shape of iSTDP kernel becomes less relevant. Weight fluctuations impact the dynamics through the $B\, r_{\text{pre}}\, r_{\text{post}}$ term rather than the correlation-dependent Q term. Consequently, these rules consistently converge to the classic rate-homeostatic solution $\tilde{r}_{\text{exc}} \approx -\alpha_{\text{pre}}/B$ (Vogels et al. 2011).

To further characterize the self-stabilizing rules, we extended our analysis beyond the U and M circuit configurations presented in Fig. 1, exploring a broader range of excitatory-to-inhibitory weights w_{exc} and input currents to the excitatory neuron. For the symmetric kernel, the learned inhibitory weight increased with w_{exc}, while for the antisymmetric kernel it decreased (Fig. 2C and D). Increasing the excitatory external input led to stronger inhibitory weights in both cases (Fig. 2C and D, top panels), however the rate of the excitatory neuron still scaled linearly with the input strength (Fig. 2C and D, bottom panels). In contrast, the rate-homeostatic rule exhibited a minimal dependence on w_{exc} and a very tight control on the excitatory rate (Fig. 2E), which restricts the dynamic range of the excitatory neuron, limiting its ability to respond to changes in input.

In summary, our analysis of the reduced E/I motif revealed a spectrum of iSTDP rules with distinct outcomes on circuit structure and dynamics. On the one hand, covariance-dominated rules with self-stabilizing kernels enable structured stabilization. As a consequence, they selectively reinforce connectivity motifs based on pre- and post-synaptic correlations, and allow a broader range of excitatory firing rates. On the other hand, rate-dominated rules prioritize homeostatic control of excitatory activity over the formation of structural motifs. Thus, these rules ensure robust dynamical stability, potentially at the cost of computational flexibility. These results demonstrate a trade-off between the emergence of specific inhibitory structures and the tight regulation of excitatory rates, suggesting that different iSTDP rules may be suited for different computational tasks.

2.3 Structured stabilization in randomly connected RNNs

To validate our analytical results derived on the reduced E/I motifs in larger networks, we simulated a randomly connected RNN of conductance-based leaky integrate-and-fire (LIF) neurons, with 900 excitatory and 100 inhibitory units. The excitatory connections (exc-to-exc and exc-to-inh) were fixed and sparse, while the inh-to-exc connections were dense and plastic (Table 2), allowing them to be shaped by the iSTDP rules.

We first implemented a covariance-dominated plasticity rule with a symmetric kernel (Fig. 3A). Initially, the network produced a pathological high-firing regime. However, inhibitory plasticity quickly stabilized the dynamics, bringing it to the asynchronous irregular firing regime characteristic of balanced networks (Fig. 3B). To analyze the resulting inhibitory connectivity, we categorized inh-to-exc synaptic weights as either "mutual" or "unidirectional". We call an inh-to-exc weight mutual when a reciprocal exc-to-inh connection exists between the same two neurons, reflecting a bidirectional link. Conversely, a weight is unidirectional in the absence of such excitatory reciprocal connection. Following the onset of plasticity, we observed that mutual inhibitory weights increased much more than unidirectional ones (Fig. 3C). Some mutual weights reached saturation, while the unidirectional remained weak. This selectivity was further confirmed by directly visualizing the matrix of synaptic weights after learning: the pattern of fixed exc-to-inh weights (Fig. 3E) was mirrored by the learned inh-to-exc weight structure (Fig. 3F). The same was true for the full weight matrix after convergence (see Supplementary Material for full weight matrix).

Figure 3: Structure emerges in RNNs of conductance-based LIF neurons under different iSTDP rules. **A** *left*. Representation of the random network with ($N_{\text{exc}} = 900$, $N_{\text{inh}} = 100$) spiking neurons. The inh-to-exc weights form all-to-all connections and are plastic, all other weights are fixed. Exc-to-exc and exc-to-inh connections are sparse. **A** *middle*. Kernel of the symmetric, covariance-dominated iSTDP rule used in network 1 (panels B-F). Parameters: $B = 0$, $\alpha_{\text{pre}} = -0.1$, $\alpha_{\text{post}} = 0$, $\tau_+ = 30$ ms, $\tau_- = 600$ ms. **A** *right*. Antisymmetric, covariance-dominated iSTDP rule usedin network 2 (panels G-K) with parameters $B = 0.3$, $\alpha_{\text{pre}} = -0.7$, $\alpha_{\text{post}} = 0.2$, $\tau_+ = 60$ ms, $\tau_- = 30$ ms. **B**. Evolution of population mean firing rates over time, with final distribution of rates over the full populations. **C**. Evolution of inh-to-exc synaptic weights over time and final distribution. The weights are split into a "mutual" (brown) and a "unidirectional" (cyan) group. Only few weights reach the saturating value, set to 80. **D**. Synaptic weights of four random inhibitory neurons after convergence. Each neuron is represented by a different column. The y coordinate represent excitatory neuron indices (1 to 150). Excitatory incoming weights are shown as right horizontal pulses (blue), inhibitory outgoing weights are shown as left pulses (red). Weights lower than 5% of saturation level are not shown. Matching left and right pulses indicate that the inhibitory neuron both receives and sends a connection to the same excitatory neuron. **E-F**. The exc-to-inh weights (**E** fixed and sparse) and the inh-to-exc weights (**F** all-to-all, plastic) after learning. **G-K**. Equivalent plots in a second network with identical network dynamics and initial conditions, but using antisymmetric iSTDP. See Supplementary Material for an example of full weight matrix after weight convergence.

Figure 4: Inhibitory self-organization in a spiking RNN structured as a 1D ring. **A**. Representation of the model. 800 excitatory neurons are arranged to a 1D ring, and connected to each other and to their inhibitory counterpart according to their "tuning distance". 100 "PV" inhibitory neurons connect all-to-all to the excitatory neurons with plasticity following a symmetric iSTDP rule, 100 "SST" neurons connect all-to-all to excitatory neurons with plasticity following an antisymmetric iSTDP. For both rules $B = 0$, $\alpha_{\mathrm{pre}} = -0.1$, $\alpha_{\mathrm{post}} = 0$. Full weight matrix after weight convergence in Supplementary Material. **B**. Evolution of mean firing rate over time. **C**. Profile of the average outgoing weights as a function of tuning distance. The excitatory weights (blue, light blue) are fixed, the inhibitory weights (dark red, orange) are learned through plasticity. For convenience, the profiles are normalized so that the maximum is either 1 or -1. Full, not-normalized, weight matrix after learning in Supplementary Fig. S4. **D**. Average effective interaction profile between the excitatory neurons, as a function of tuning (see Methods). **E**. Pearson correlation during spontaneous activity between an excitatory neuron and its neighbors, ordered by tuning distance and averaged over the entire population. **F**. Mean responses of the excitatory population to stimuli of different sizes, averaged over 100 trials. **G**. Response of the center neuron to stimuli of different sizes.

Next, we investigated the impact on an antisymmetric iSTDP rule within the same random RNN architecture and neural dynamics. As with the symmetric rule, this antisymmetric iSTDP led the network to a balanced dynamical regime. However, it promoted a markedly different connectivity pattern: purely lateral inhibition. In this configuration, inhibitory neurons strengthened their connections onto excitatory neurons lacking reciprocal connections back to them, thus forming exclusively unidirectional connections (Fig. 2H). This resulted in an inhibitory-to-excitatory connectivity structure that was a negative image of the pattern observed with the symmetric kernel: the presence of exc-to-inh synapses was mirrored by a corresponding absence of inh-to-exc connections (Fig. 2J,K). Achieving robust lateral inhibition with the antisymmetric iSTDP rule required the introduction of a small rate-dependent component ($B = 0.3$). This deviation from a purely covariance-dominated regime introduces characteristics of rate-homeostatic plasticity, such as rapid convergence to a narrow distribution of excitatory firing rates (Fig. 2G). This requirement is perhaps unsurprising given the challenges of parameter tuning in large RNNs, where closed-form solutions are unavailable (see Discussion for further considerations).

To sum up, these RNN simulations confirm that our analytical predictions from the reduced E/I model generalize to larger, more complex networks. Symmetric iSTDP rules reinforce mutual inhibition, mirroring pre-existing excitatory structure, whereas antisymmetric kernels favor lateral inhibition.

2.4 Emergent computations from structured stabilization in a ring network

Having shown that covariance-dominated iSTDP shapes E/I connectivity based on local correlations arising from direct connections, we next explored the functional consequences of structured stabilization in a recurrent network with a one-dimensional ring architecture. Ring networks are well-suited to model continuous attractor dynamics, supporting bumps of persistent activity (Amari 1977; Ben-Yishai et al. 1995; Khona and Fiete 2022). This makes them relevant for studying neural computations such as head-direction encoding (Xie et al. 2002; Kim et al. 2019; Petrucco et al. 2023) and orientation selectivity in the primary visual cortex (Goldberg et al.

2004).

We built a ring network of 800 excitatory and 200 inhibitory neurons arranged on a circle, with exc-to-exc
and exc-to-inh connectivity following a profile that decayed exponentially with the angular distance between
neurons (i.e. a rescaled Von Mises, Fig. 4A). We further divided the inhibitory population into two subclasses,
each implementing a distinct iSTDP rule but sharing identical neural dynamics. Recent experimental work has
measured a symmetric iSTDP rule for the plasticity between parvalbumin-expressing (PV) interneurons and ex-
citatory pyramidal neurons in the orbitofrontal cortex (Lagzi et al. 2021). The same work also measured an
antisymmetric iSTDP rule for the plasticity between somatostatic-expressing (SST) interneurons and excitatory
pyramidal neurons (Lagzi et al. 2021). Therefore, we identified our two populations of inhibitory interneurons in
our ring network with "PV" and "SST" and implemented a covariance-dominated symmetric iSTDP rule for the
PV population and a covariance-dominated antisymmetric rule for the SST population. Initially, both PV-to-exc
and SST-to-exc connections were all-to-all, weak, and unbiased (see Fig. 4A for schematics, and Methods Table 3
for numerical parameters).

As the network evolved under the influence of the two plasticity rules, the inhibitory populations developed spe-
cialized roles. PV interneurons, governed by symmetric iSTDP, formed strong mutual connections with excita-
tory neurons, mirroring the distance-dependent excitatory structure (Fig. 4C, red). Conversely, SST interneurons,
driven by antisymmetric iSTDP, established lateral inhibition across the ring (Fig. 4C, orange; full connectivity
matrix after learning in Supplementary Material). This structured stabilization, with both localized and broader
components, raised the possibility of nontrivial effective interactions between excitatory neurons, potentially
mediated by the different roles of the PV and SST interneurons.

To estimate the effective interaction profile between excitatory neurons, we used linear response theory, equiv-
alent to a Taylor expansion of the network dynamics around its operating point (Sadeh and Clopath 2020). We
found a Mexican-hat effective connectivity profile, characterized by short-range excitation directly inherited
from exc-to-exc connectivity and a broader inhibitory surround resulting from the combined action of PV and
SST interneurons (Fig. 4D). This effective connectivity profile raised questions about the resulting network dy-
namics and its response to external stimuli. We examined the emergent spontaneous activity and observed a
complex correlation structure, with periodic oscillations in correlations over distance (Fig. 4E, see Supplemen-
tary Material for full correlation matrix). This emerging dynamics is reminiscent of the correlation maps observed
in the developing cortex of ferrets, and is considered a signature of modular cortical organization (Smith et al.
2018).

Finally, we considered the network's response to external stimuli. The network received inputs of different size
modeled as an instantaneous increase in external currents to a center neuron and a varying number of its neigh-
bors. For example, size five corresponded to a step-signal to the center neuron and to its four neighbors, two on
each side. We presented each stimulus 100 times in randomized trials with a $0.5\,\mathrm{s}$ stimulus presentation followed
by a $0.5\,\mathrm{s}$ blank. Plasticity was deactivated, with connection weights fixed to their value after learning from spon-
taneous activity. We found that larger stimuli elicited lower peak firing rates, with higher response for neurons
with receptive fields closer to the stimulus' edges (Fig. 4F, raster plot in Supplementary Material). Focusing on
the center neuron's response as a function of stimulus size revealed the characteristic surround suppression pro-
file (Fig. 4G), consistent with classic findings on contextual effects (DeAngelis et al. 1994; Cavanaugh et al. 2002).
These results demonstrate how structured stabilization shaped by the interplay of symmetric and antisymmetric
iSTDP rules can give rise to complex dynamics and computations, including surround suppression.

3   Discussion

In this work, we have shown that general iSTDP rules can stabilize excitation at the circuit level, while simultane-
ously driving the formation of specific connectivity patterns, or motifs, among neurons. The shape of the iSTDP
kernel determines which motifs arise, selectively reinforcing structures based on the correlations present in the
network. In the absence of structured external inputs, or training signals, circuits self-organize purely through
its spontaneous internal dynamics.

Analytical methods have been developed to explore the self-organized formation of motifs in large networks
through STDP, but so far they only focused on exc-to-exc plasticity (Ocker et al. 2015; Ravid Tannenbaum and
Burak 2016; Triplett et al. 2018; Ocker and Doiron 2019; Montangie et al. 2020; Manz and Memmesheimer
2023). These studies do not account for the formation of inhibitory structures and E/I motifs, whose inclusion
can significantly increase the complexity of the the analytical expressions, due to the combinatorial interactions

among the multiple neural populations within each motif. Here, we employed an analytical framework originally developed by Kempter et al. (1999) for exc-to-exc plasticity. While there has been substantial progress in theoretical neuroscience involving iSTDP rules, previous work has been largely restricted to feedforward circuits and covariance-dominated plasticity rules (i.e. rules lacking rate-dependent terms) (Luz and Shamir 2014; Akil et al. 2021; Vignoud and Robert 2022; Lagzi and Fairhall 2024). In particular, the work of Luz and Shamir (2014) explored both symmetric and antisymmetric iSTDP kernels, but their scope was limited to feedforward circuits. In contrast, our work introduces a closed-form solution for a simple recurrent excitatory-inhibitory circuit, incorporating the rate-dependent terms of the generalized iSTDP. These terms are essential for achieving stable solutions and regulating neural activity. This approach provides deeper insight into both the regulatory and structural properties of generalized iSTDP rules, expanding our understanding of plasticity in closed-loop circuits. (Akil et al. 2021) proposed a mean-field approach to study how iSTDP shapes recurrent networks of LIF neurons. Although their approach captures more realistic single neuron dynamics than our Poisson neurons and reduced two-dimensional system, it assumes a regime of weak interactions, where internally-generated correlations become negligible as the size of the neural population grows, and plasticity is driven exclusively by externally-induced correlations. In contrast, in our case finite network size, strong interactions and a correlation-dominated rule together give rise to self-organization driven solely by these internal correlations. It remains an open question whether and to what extent external correlations, when present, would dominate the internal ones.

A challenge of numerically simulating large-scale RNNs equipped with plasticity is to determine the parameters that ensure stable solutions. But fine-tuning these parameters can be time-consuming. One potential solution is to employ numerical methods for parameter search based on deep learning, which have already been applied to search for generalized excitatory and inhibitory STDP rules acting on recurrent circuits (Confavreux et al. 2024). Our framework provides a complementary approach by offering analytical insights into the interactions of the iSTDP kernel and other parameters. This may help understand some of the degeneracies encountered in the numerical searches, and simplify the parameter search process. For example, we predict that in rate-dominated rules the evolution of plastic weights is primarily determined by the area under the STPD kernel and the rate-dependent parameters, rather than by the specific shape of the kernel.

Our work reveals that circuit structure can self-organize at the level of E/I neural motifs, even in networks with simple sparse (excitatory) connectivity. Higher order structures, such as dense assemblies of interconnected neurons, amplify correlations among their members and potentially introduce negative correlations across assemblies. Previous studies that report E/I tuning at the assembly level for the classical rate-homeostatic iSTDP rule align with our findings Lagzi et al. (2021) and Lagzi and Fairhall (2024). Specifically, enhanced within-assembly correlations and reduced or negative across-assembly correlations introduce a slow change in the plasticity dynamics, once the excitatory target-rate prescribed by the homeostatic component is reached. By considering a broader class of plasticity rules, we have shown that this selectivity can be enhanced and regulated flexibly, without requiring structured external inputs or large-scale connectivity patterns.

Additionally, we demonstrated the computational advantages of including two distinct inhibitory interneuron populations in a recurrent E/I circuit. This builds on earlier findings showing that a similar configuration can promote the formation of neural assemblies and maintain them over time (Lagzi et al. 2021; Bergoin et al. 2023; Bergoin et al. 2024). Our work extends these insights by showing how the interplay between distinct interneuron types can benefit other functionally relevant circuit configurations, such as ring models, leading to emergent functional properties, such as surround suppression and long-range correlations reminiscent of modular activity observed in the cortex (Smith et al. 2018).

While this study focuses on inhibitory plasticity, the same formalism can be extended to excitatory STDP, generalizing the existing results on self-stabilizing excitatory plasticity (Kempter et al. 1999) to large recurrent circuits with multiple plastic neuron types. Our analysis suggests that excitatory STDP rules with pre-post potentiation saturate in covariance-dominated regimes. Stability can be preserved by incorporating additional, often non-local, regularization mechanism (Litwin-Kumar and Doiron 2014). Alternatively, stronger rate-dependent terms could ensure both rate homeostasis and structure formation — a direction that, with few exceptions (see e.g. Manz and Memmesheimer 2023), remains largely unexplored. Overall, this work underscores the capacity of simple, local plasticity rules to regulate circuit activity and drive the self-organization of structured connectivity in recurrent networks, with broad implications for understanding neural development and learning.

4 Methods

4.1 General mean-field solution

We consider a class of pairwise iSTDP rules triggered by spiking events and purely based on spike-timing difference between pre- and postsynaptic neuron firing, defined by the following expression.

$$\text{on } t_k^{(\text{pre})} \;\rightarrow\; \Delta W^{(\text{pre})} = \alpha_{\text{pre}} + \sum_{k'<k} L^{\Delta_-}\left(t_k^{(\text{pre})} - t_{k'}^{(\text{post})}\right) \;,$$

$$\text{on } t_k^{(\text{post})} \;\rightarrow\; \Delta W^{(\text{post})} = \alpha_{\text{post}} + \sum_{k'<k} L^{\Delta_+}\left(t_k^{(\text{post})} - t_{k'}^{(\text{pre})}\right) \;; \qquad (2)$$

where $t_k^{(\text{pre})}$ and $t_k^{(\text{post})}$ represent pre- and postsynaptic spike events, respectively. Each presynaptic spike triggers a weight change consisting of a fixed component, denoted α_{pre}, and a pairwise interaction term. Similarly, each postsynaptic spike triggers a weight change with a fixed component α_{post} and a pairwise interaction term. The terms of pairwise interaction capture the cumulative influence of joint spiking events, incorporating the contribution of all preceding spikes ($k' < k$) weighted by a kernel $L^{\Delta_\pm}(\cdot)$ that depends on the time differences between the spikes. The two functions $L^{\Delta_-}(\cdot)$ and the $L^{\Delta_+}(\cdot)$ refer to the post-pre and the pre-post pairwise interaction, and are merged in a single function, known as the iSTDP (pairwise interaction) kernel, as follows:

$$L(\tau) := \begin{cases} L^{\Delta_+}(\tau) & \text{if } \tau \geq 0 \\ L^{\Delta_-}(-\tau) & \text{if } \tau < 0 \,. \end{cases} \qquad (3)$$

By convention, $\tau > 0$ (the Δ_+ branch of the curve) represents pre-post interactions (presynaptic spike preceding the postsynaptic spike), while $\tau < 0$ (the Δ_- branch) represents post-pre interactions (postsynaptic spike preceding the presynaptic spike). In our framework, we consider $L(\tau)$ to be independent of α_{pre} and α_{post}. This is in contrast to previous formulations that incorporate these terms into the kernel representation (e.g., Vogels et al. 2011 and Confavreux et al. 2024). We assume that synaptic plasticity operates on a much slower timescale than neural dynamics, allowing us to treat synaptic weights as constant when calculating first- and second-order firing statistics (Kempter et al. 1999). Under these assumptions, we can write the evolution of the mean synaptic weights as:

$$\frac{d\langle W \rangle}{dt} = \alpha_{\text{pre}} r_{\text{pre}} + \alpha_{\text{post}} r_{\text{post}} + B\, r_{\text{pre}}\, r_{\text{post}} + Q \quad,$$

$$\text{with} \quad B := \int_{-\infty}^{+\infty} d\tau\, L(\tau) \quad \text{and} \quad Q := \int_{-\infty}^{+\infty} d\tau\, L(\tau)\, C(\tau) \;; \qquad (4)$$

where B is the integral of the iSTDP kernel, and Q is the only term that depends on the pairwise correlation density between pre and post-synaptic spike trains, here denoted $C(\tau)$. For a complete derivation see Supplementary Material or Kempter et al. (1999) and Luz and Shamir (2014).

4.2 Expressions for the symmetric and antisymmetric exponential kernel

Eq. 4 can be applied to any form of iSTDP kernel, including, e.g., kernels with delayed interactions (Babadi and Abbott 2013). Here, we focus on two specific exponentially decaying shapes: symmetric and antisymmetric, defined as follows:

$$L^{\text{sy}}(\tau) = \frac{1}{2}\left(\frac{1}{\tau_+} A_+\, e^{-|\tau|/\tau_+} - \frac{1}{\tau_-} A_-\, e^{|\tau|/\tau_-}\right) \quad \text{and} \quad L^{\text{an}}(\tau) = \begin{cases} -\frac{A_-}{\tau_-}\, e^{\tau/\tau_-} & \text{if } \tau < 0 \\ \frac{A_+}{\tau_+}\, e^{-\tau/\tau_+} & \text{if } \tau \geq 0 \end{cases} \qquad (5)$$

With this parametrization, $A_+ = A_-$ implies $B = 0$, corresponding to covariance-dominated rules, whereas for $A_- = 0$ we have $B = A_+$, a rate-dominated regime. Such parametrization allows us to keep track of the entire spiking history through activity traces updated at every spike time and decaying in time with the same constants as the iSTDP kernel. These traces will determine the amount of potentiation ($+$) or depression ($-$) triggered by each spike. For the antisymmetric rule, the presynaptic and postsynaptic traces are respectively defined as follows:

$$\frac{dT_{r+}(t)}{dt} = -\frac{T_{r+}(t)}{\tau_+}; \qquad \text{if } t = t^{(\text{pre})}, \quad T_{r+}(t+\varepsilon) = T_{r+}(t) + 1$$

$$\frac{dT_{o-}(t)}{dt} = -\frac{T_{o-}(t)}{\tau_-}; \qquad \text{if } t = t^{(\text{post})}, \quad T_{o-}(t+\varepsilon) = T_{o-}(t) + 1 \qquad (6)$$

The symmetric rule requires, in addition to $T_{r+}(t)$ and $T_{o-}(t)$, the traces $T_{r-}(t)$ and $T_{o+}(t)$, defined analogously.

The implementation of the iSTDP rule is then given by:

$$
\begin{aligned}
\Delta W^{\mathsf{an}}(t) &= \alpha_{\mathsf{pre}} - \frac{A_-}{\tau_-}\, T_{o-}(t) \qquad \text{if} \quad t = t^{(\mathsf{pre})}\,, \\
\Delta W^{\mathsf{an}}(t) &= \alpha_{\mathsf{post}} + \frac{A_+}{\tau_+}\, T_{r+}(t) \qquad \text{if} \quad t = t^{(\mathsf{post})}\,,
\end{aligned}
\tag{7}
$$

for the antisymmetric rule, and

$$
\begin{aligned}
\Delta W^{\mathsf{sy}}(t) &= \alpha_{\mathsf{pre}} + \frac{A_+}{2\tau_+}\, T_{o+}(t) - \frac{A_-}{2\tau_-}\, T_{o-}(t) \qquad \text{if} \quad t = t^{(\mathsf{pre})}\,, \\
\Delta W^{\mathsf{sy}}(t) &= \alpha_{\mathsf{post}} + \frac{A_+}{2\tau_+}\, T_{r+}(t) - \frac{A_-}{2\tau_-}\, T_{r-}(t) \qquad \text{if} \quad t = t^{(\mathsf{post})}\,,
\end{aligned}
\tag{8}
$$

for the symmetric rule.

4.3 Low impact of correlations in rate-dominated rules

To better understand the role of the net area B of the iSTDP kernel in the weight dynamics (Eq. 4), and how it determines whether a rule is rate-dominated or covariance-dominated, we will consider here a simplified scenario, where $L(\tau)$ prescribes a +1 increase in weight for coincident pre-post spikes and no increase otherwise. This choice yields $B = 1$. Consider now a pre and a post-synaptic neuron that fire with Poisson statistics at rates $r_{\mathsf{pre}} = r_{\mathsf{post}} = r = 10\,\mathrm{Hz}$. The term $B\, r_{\mathsf{pre}}\, r_{\mathsf{post}}$ in Eq. 4 takes a value of $100\,\mathrm{s}^{-2}$ regardless of the correlation between the two neurons. But how does it compare to the covariance term Q in the presence of strong correlations between pre and post-synaptic activity? We can consider an regime of (unnaturally) high correlations, such that half of the spikes of the two neurons are synchronous (Pearson correlation 0.5). The term Q, corresponding to the spike-count covariance, is $5\,\mathrm{s}^{-2}$. So even for very strong correlations Q is 20 times smaller than $B\, r_{\mathsf{pre}}\, r_{\mathsf{post}}$. This illustrates that in regimes with strong B, the weight dynamics in Eq. 4 are primarily governed by first-order statistics, with — at best — second-order contributions from correlations in spiking activity.

4.4 The two-dimensional reduced model

Finding a general solution for Eq. 4 can be challenging. For our purposes, we consider a simplified case. First, we model an excitatory and an inhibitory neuron as mutually interacting Poisson processes (Hawkes 1971; Ocker et al. 2017). Each time the presynaptic neuron spikes, the firing probability of the postsynaptic neuron increases (or decreases) proportionally to the excitatory (or inhibitory) post-synaptic potential EPSP (IPSP) kernel:

$$
E_e(\tau) = \frac{1}{\tau_e}\, e^{-\tau/\tau_e} \quad \text{and} \quad E_i(\tau) = -\frac{1}{\tau_i}\, e^{-\tau/\tau_i} \quad \text{with } \tau \geq 0
\tag{9}
$$

We further assume that the input to the inhibitory neuron is regulated homeostatically to keep its rate constant. In these conditions, Eq. 4 has an approximate fixed-point closed-form solution for the inhibitory connection w_{inh}:

$$
\tilde{w}_{\mathsf{inh}} = \frac{\alpha_{\mathsf{pre}}\, r_{\mathsf{inh}} + h_{\mathsf{exc}}\Psi}{r_{\mathsf{inh}}\left(\Psi - M_{ei}^{(10)}\right)} \quad ; \quad \tilde{r}_{\mathsf{exc}} = -\frac{\alpha_{\mathsf{pre}}\, r_{\mathsf{inh}} + M_{ei}^{(10)}\, h_{\mathsf{exc}}}{\Psi - M_{ei}^{(10)}} \quad ;
\tag{10}
$$

$$
\text{with } \Psi = \alpha_{\mathsf{post}} + B\, r_{\mathsf{inh}} + M_{ie}^{(01)}\, w_{\mathsf{exc}}\quad .
$$

The coefficients $M_{ei}^{(10)}$ and $M_{ie}^{(01)}$ are found through the linear response theory (Hawkes 1971; Ocker et al. 2017), and depend on the iSTDP kernel and the EPSP / IPSP parameters (refer to the analytic derivation of the Supplementary Material).

For the symmetric iSTDP:

$$
M_{ei}^{\mathsf{sy},\,(10)} = \frac{A_+}{\tau_+ + \tau_i} - \frac{A_-}{\tau_- + \tau_i} \quad ; \quad M_{ie}^{\mathsf{sy},\,(01)} = \frac{A_+}{\tau_+ + \tau_e} - \frac{A_-}{\tau_- + \tau_e}
\tag{11}
$$

For the antisymmetric iSTDP instead:

$$
M_{ei}^{\mathsf{an},\,(10)} = \frac{A_+}{\tau_+ + \tau_i} \quad ; \quad M_{ie}^{\mathsf{an},\,(01)} = -\frac{A_-}{\tau_- + \tau_e}
\tag{12}
$$

Numerical parameters are in Table 1.

symbol	value	description
N_{exc}	1	number of excitatory neurons
N_{inh}	1	number of inhibitory neurons
τ_e	30 ms	EPSP kernel time constant
τ_i	30 ms	IPSP kernel time constant
h_e	40 Hz	excitatory input current
r_{inh}	80 Hz	inhibitory firing rate
τ_i	30 ms	IPSP kernel time constant
rate-dominated iSTDP rule		
A_0	$5 \cdot 10^{-7}$	global learning rate
B	1	net area under the iSTDP kernel
α_{pre}	-25	constant dependent on presynaptic rate
α_{post}	0	constant dependent on postsynaptic rate
τ_+	50 ms	iSTDP kernel time constant
covariance-dominated symmetric iSTDP rule		
A_0	$3 \cdot 10^{-6}$	global learning rate
B	0	net area under the iSTDP kernel
α_{pre}	-25	constant dependent on presynaptic rate
α_{post}	0	constant dependent on postsynaptic rate
τ_+	50 ms	iSTDP kernel time constant for potentiation
τ_-	500 ms	iSTDP kernel time constant for depression
covariance-dominated antisymmetric iSTDP rule		
A_0	$3 \cdot 10^{-6}$	global learning rate
B	0	net area under the iSTDP kernel
α_{pre}	-25	constant dependent on presynaptic rate
α_{post}	0	constant dependent on postsynaptic rate
τ_+	50 ms	iSTDP kernel time constant for potentiation
τ_-	50 ms	iSTDP kernel time constant for depression

Table 1: Numerical parameters used for Fig. 1. The parameters in Fig. 2 match those of Fig. 1 unless otherwise indicated by the figure caption.

4.5   Randomly connected recurrent neural network

We built a random network of conductance-based leaky integrate-and-fire (LIF) neurons, with 900 excitatory units and 100 inhibitory units. The membrane potential obeyed the following dynamics:

$$\frac{dV(t)}{dt} = \frac{1}{C_{\text{membrane}}} \left(-g_{\text{leak}}(V - V_{\text{L}}) - g_{\text{AMPA}} \left(V - V_{\text{AMPA}} \right) - g_{\text{GABA}} \left(V - V_{\text{GABA}} \right) + \text{input} \right)$$

$$\frac{dg_{\text{AMPA}}(t)}{dt} = -\frac{g_{\text{AMPA}}(t)}{\tau_{\text{AMPA}}} \qquad ; \qquad \frac{dg_{\text{GABA}}(t)}{dt} = -\frac{g_{\text{GABA}}(t)}{\tau_{\text{GABA}}}$$

(13)

With $g_{\text{AMPA}}(t)$ and $g_{\text{GABA}}(t)$ updated proportionally to the excitatory and inhibitory synaptic weights for each incoming signal. We assigned a probability of connection for all non-plastic connections (excitatory to excitatory, excitatory to inhibitory, inhibitory to inhibitory). The inhibitory to excitatory connectivity was all-to-all and plastic. Numerical parameters are in Table 2.

4.6   Ring model

The ring model was implemented with the same neural dynamics and parameters as the random network (Eq. 13). We used 800 excitatory units and 200 inhibitory units, further divided into a PV and an SST subclass. The synaptic weight between an excitatory neuron and its neighbors decayed with distance, following a bell-shaped

symbol	value	description
N_{exc}	900	number of excitatory neurons
N_{inh}	100	number of inhibitory neurons
C_{membrane}	500 pF	membrane capacitance
τ_{AMPA}	5 ms	glutamatergic synaptic time constant
τ_{GABA}	10 ms	GABAergic synaptic time constant
g_{leak}	10	leak conductivity
V_{L}	-60 mV	membrane leak reverse potential
V_{AMPA}	0 mV	AMPA reverse potential
V_{GABA}	-80 mV	GABA reverse potential
$V_{\text{threshold}}$	-50 mV	firing threshold
$p_{\text{e}\leftarrow\text{e}}$	0.2	exc-to-exc connection probability
$p_{\text{i}\leftarrow\text{e}}$	0.1	exc-to-inh connection probability
$p_{\text{e}\leftarrow\text{i}}$	1.0	inh-to-exc connection probability
$p_{\text{i}\leftarrow\text{i}}$	1.0	inh-to-inh connection probability
$w_{\text{e}\leftarrow\text{e}}$	1.0	exc-to-exc synaptic weight
$w_{\text{i}\leftarrow\text{e}}$	1.8	exc-to-inh synaptic weight
$w_{\text{i}\leftarrow\text{i}}$	0.3	inh-to-inh synaptic weight
covariance-dominated symmetric iSTDP rule (panels A-F)		
A_0	$2 \cdot 10^{-2}$	global learning rate
B	0	net integral of the STDP kernel
α_{pre}	-0.1	constant dependent on presynaptic rate
α_{post}	0	constant dependent on postsynaptic rate
τ_+	30 ms	STDP kernel time constant
τ_-	600 ms	STDP kernel time constant
"covariance-dominated" antisymmetric iSTDP rule (panels G-L)		
A_0	$5 \cdot 10^{-3}$	global learning rate
B	0.3	net integral of the iSTDP kernel
α_{pre}	-0.7	constant dependent on presynaptic rate
α_{post}	0.2	constant dependent on postsynaptic rate
τ_+	60 ms	iSTDP kernel time constant for potentiation
τ_-	30 ms	iSTDP kernel time constant for depression

Table 2: Numerical parameters for the random RNN in Fig. 3.

curve modeled as a rescaled von Mises:

$$w_{ij}^{(\bullet\leftarrow\text{e})} := C_{\bullet\leftarrow\text{e}} \left(\exp \left(k_{\bullet\leftarrow\text{e}} \cos \left(\theta_i - \theta_j \right) \right) \right) \; ; \tag{14}$$

where θ_i and θ_j represent the locations of the post- and presynaptic neuron on the ring, with each population equally spaced in the $[0, \, 2\pi)$ interval, $k_{\bullet\leftarrow\text{e}}$ regulates the connection width, finally the constant $C_{\bullet\leftarrow\text{e}}$ is determined so that the discrete sum of incoming weights to either exc, PV or SST population reaches a desired fixed sum. All inh-to-inh connections were fixed and dense. We considered two distinct inhibitory populations with identical dynamics, but different iSTDP rules. The first population had a symmetric iSTDP kernel, the second an antisymmetric one. Numerical parameters are shown in Table 3.

4.7 Software implementation and code availability

All simulations for the reduced models are based on custom code written in Julia (Bezanson et al. 2017, https://julialang.org), also available as separate toolbox. For the large scale simulations we used Brian2 (Stimberg et al. 2019, https://briansimulator.org). Finally some of the analytic integrals were solved using Maple (https://maplesoft.com). The code is fully open-source and publicly available on GitHub under the Creative Commons Attribution 4.0 International (CC BY 4.0) License.

https://github.com/comp-neural-circuits/structured-stabilization-in-recurrent-neural-circuits

symbol	value	description
N_{exc}	800	number of excitatory neurons
N_{PV}	100	number of PV interneurons
N_{SST}	100	number of SST interneurons
sum $w_{\bullet \leftarrow e}$	145	total incoming exc weights to exc, PV, SST
$k_{e \leftarrow e}$	10	width of exc-to-exc connections
$k_{\text{PV} \leftarrow e}$	5	width of exc-to-pv connections
$k_{\text{SST} \leftarrow e}$	10	width of exc-to-sst connections
$p_{\text{PV,SST} \leftarrow \text{PV,SST}}$	0.9	connection probability for PV-PV, SST-SST and PV-SST
$w_{\text{PV} \leftarrow \text{PV}}$	2	PV-to-PV synaptic weight
$w_{\text{SST} \leftarrow \text{PV}}$	0.5	PV-to-SST synaptic weight
$w_{\text{SST} \leftarrow \text{SST}}$	2	SST-to-SST synaptic weight
$w_{\text{PV} \leftarrow \text{SST}}$	0.5	SST-to-PV synaptic weight
symmetric iSTDP rule for the PV population		
A_0	$2 \cdot 10^{-3}$	global learning rate
B	0	net integral of the iSTDP kernel
α_{pre}	-0.2	constant dependent on presynaptic rate
α_{post}	1	constant dependent on postsynaptic rate
τ_+	30 ms	STDP kernel time constant for potentiation
τ_-	600 ms	STDP kernel time constant for depression
antisymmetric iSTDP rule for the SST population		
A_0	$1 \cdot 10^{-3}$	global learning rate
B	0	net integral of the iSTDP kernel
α_{pre}	-0.2	constant dependent on presynaptic rate
α_{post}	1	constant dependent on postsynaptic rate
τ_+	30 ms	iSTDP kernel time constant for potentiation
τ_-	30 ms	iSTDP kernel time constant for depression

Table 3: Numerical parameters for one-dimensional ring model. The parameters that regulate neural dynamics are the same as in Table 2.

440 Acknowledgments

441 This project has received funding from the European Research Council (ERC StG NeuroDevo, Grant agreement
442 No. 804824) and Marie Skłodowska-Curie ITN SmartNets (grant agreement No. 860949) under the European
443 Union's Horizon 2020 research and innovation program, by the International Human Frontier Science Program
444 Organization (RGP0062/2021), and by the Technical University of Munich (TUM). Thanks to the "Computation
445 in Neural Circuits Group" for their useful insights and discussions.

References

Agnes, Everton J. and Tim P. Vogels (Mar. 20, 2024). "Co-Dependent Excitatory and Inhibitory Plasticity Accounts for Quick, Stable and Long-Lasting Memories in Biological Networks". In: *Nature Neuroscience*, pp. 1–11. DOI: 10.1038/s41593-024-01597-4.

Akil, Alan Eric, Robert Rosenbaum, and Krešimir Josić (May 12, 2021). "Balanced Networks under Spike-Time Dependent Plasticity". In: *PLOS Computational Biology* 17.5, e1008958. DOI: 10.1371/journal.pcbi.1008958.

Amari, Shun-ichi (June 1, 1977). "Dynamics of Pattern Formation in Lateral-Inhibition Type Neural Fields". In: *Biological Cybernetics* 27.2, pp. 77–87. DOI: 10.1007/BF00337259.

Babadi, Baktash and L. F. Abbott (Feb. 21, 2013). "Pairwise Analysis Can Account for Network Structures Arising from Spike-Timing Dependent Plasticity". In: *PLOS Computational Biology* 9.2, e1002906. DOI: 10.1371/journal.pcbi.1002906.

Ben-Yishai, R, R L Bar-Or, and H Sompolinsky (Apr. 25, 1995). "Theory of Orientation Tuning in Visual Cortex." In: *Proceedings of the National Academy of Sciences* 92.9, pp. 3844–3848. DOI: 10.1073/pnas.92.9.3844.

Bergoin, Raphaël, Alessandro Torcini, Gustavo Deco, Mathias Quoy, and Gorka Zamora-López (Apr. 28, 2023). "Inhibitory Neurons Control the Consolidation of Neural Assemblies via Adaptation to Selective Stimuli". In: *Scientific Reports* 13.1 (1), p. 6949. DOI: 10.1038/s41598-023-34165-0.

– (July 17, 2024). *Emergence and Long-Term Maintenance of Modularity in Plastic Networks of Spiking Neurons*. DOI: 10.1101/2024.07.15.603496. Pre-published.

Bezanson, Jeff, Alan Edelman, Stefan Karpinski, and Viral B Shah (2017). "Julia: A fresh approach to numerical computing". In: *SIAM review* 59.1, pp. 65–98.

Bollmann, Yannick et al. (Sept. 2023). "Prominent in Vivo Influence of Single Interneurons in the Developing Barrel Cortex". In: *Nature Neuroscience* 26.9 (9), pp. 1555–1565. DOI: 10.1038/s41593-023-01405-5.

Brunel, N. (2000). "Dynamics of Sparsely Connected Networks of Excitatory and Inhibitory Spiking Neurons". In: *Journal of computational neuroscience* 8.3, pp. 183–208. DOI: 10.1023/A:1008925309027.

Cavanaugh, James R., Wyeth Bair, and J. Anthony Movshon (Nov. 2002). "Nature and Interaction of Signals From the Receptive Field Center and Surround in Macaque V1 Neurons". In: *Journal of Neurophysiology* 88.5, pp. 2530–2546. DOI: 10.1152/jn.00692.2001.

Clopath, Claudia, Tim P. Vogels, Robert C. Froemke, and Henning Sprekeler (July 29, 2016). "Receptive Field Formation by Interacting Excitatory and Inhibitory Synaptic Plasticity". In: p. 066589. DOI: 10.1101/066589.

Confavreux, Basile, Everton J. Agnes, Friedemann Zenke, Henning Sprekeler, and Tim P. Vogels (June 17, 2024). *Balancing Complexity, Performance and Plausibility to Meta Learn Plasticity Rules in Recurrent Spiking Networks*. DOI: 10.1101/2024.06.17.599260.

Dahmen, David et al. (Nov. 4, 2020). "Strong Coupling and Local Control of Dimensionality across Brain Areas". In: *bioRxiv*, p. 2020.11.02.365072. DOI: 10.1101/2020.11.02.365072.

DeAngelis, G. C., R. D. Freeman, and I. Ohzawa (Jan. 1994). "Length and Width Tuning of Neurons in the Cat's Primary Visual Cortex". In: *Journal of Neurophysiology* 71.1, pp. 347–374. DOI: 10.1152/jn.1994.71.1.347.

Dorrn, Anja L., Kexin Yuan, Alison J. Barker, Christoph E. Schreiner, and Robert C. Froemke (June 17, 2010). "Developmental Sensory Experience Balances Cortical Excitation and Inhibition". In: *Nature* 465.7300, pp. 932–936. DOI: 10.1038/nature09119.

Duan, Zhe Ran S. et al. (Jan. 8, 2020). "GABAergic Restriction of Network Dynamics Regulates Interneuron Survival in the Developing Cortex". In: *Neuron* 105.1, 75–92.e5. DOI: 10.1016/j.neuron.2019.10.008.

Eckmann, Samuel, Edward James Young, and Julijana Gjorgjieva (June 18, 2024). "Synapse-Type-Specific Competitive Hebbian Learning Forms Functional Recurrent Networks". In: *Proceedings of the National Academy of Sciences* 121.25, e2305326121. DOI: 10.1073/pnas.2305326121.

Froemke, Robert C., Michael M. Merzenich, and Christoph E. Schreiner (Nov. 15, 2007). "A Synaptic Memory Trace for Cortical Receptive Field Plasticity". In: *Nature* 450.7168, pp. 425–429. DOI: 10.1038/nature06289.

Goldberg, Joshua A., Uri Rokni, and Haim Sompolinsky (May 13, 2004). "Patterns of Ongoing Activity and the Functional Architecture of the Primary Visual Cortex". In: *Neuron* 42.3, pp. 489–500. DOI: 10.1016/S0896-6273(04)00197-7.

Gong, Ziyi and Nicolas Brunel (Apr. 10, 2024). *Inhibitory Plasticity Enhances Sequence Storage Capacity and Retrieval Robustness*. DOI: 10.1101/2024.04.08.588573. Pre-published.

Hafizi, Hadi et al. (Aug. 20, 2022). *Inhibition-Dominated Rich-Club Shapes Dynamics in Cortical Microcircuits*. DOI: 10.1101/2021.05.07.443074.

Hanganu-Opatz, Ileana L. et al. (Feb. 3, 2021). "The Logic of Developing Neocortical Circuits in Health and Disease". In: *Journal of Neuroscience* 41.5, pp. 813–822. DOI: 10.1523/JNEUROSCI.1655-20.2020.

Hawkes, Alan G. (1971). "Spectra of Some Self-Exciting and Mutually Exciting Point Processes". In: *Biometrika* 58.1, pp. 83–90. DOI: 10.2307/2334319.

Jiang, Xiaolong et al. (Nov. 27, 2015). "Principles of Connectivity among Morphologically Defined Cell Types in Adult Neocortex". In: *Science* 350.6264, aac9462. DOI: 10.1126/science.aac9462.

Jin, Dezhe (Oct. 2002). "Fast Convergence of Spike Sequences to Periodic Patterns in Recurrent Networks". In: *Physical Review Letters* 89.20. DOI: 10.1103/PhysRevLett.89.208102.

Kempter, Richard, Wulfram Gerstner, and J. Leo van Hemmen (Apr. 1, 1999). "Hebbian Learning and Spiking Neurons". In: *Physical Review E* 59.4, pp. 4498–4514. DOI: 10.1103/PhysRevE.59.4498.

Khona, Mikail and Ila R. Fiete (Nov. 3, 2022). "Attractor and Integrator Networks in the Brain". In: *Nature Reviews Neuroscience*. DOI: 10.1038/s41583-022-00642-0.

Kim, Sung Soo, Ann M. Hermundstad, Sandro Romani, L. F. Abbott, and Vivek Jayaraman (Dec. 2019). "Generation of Stable Heading Representations in Diverse Visual Scenes". In: *Nature* 576.7785 (7785), pp. 126–131. DOI: 10.1038/s41586-019-1767-1.

Kleberg, Florence I., Tomoki Fukai, and Matthieu Gilson (May 7, 2014). "Excitatory and Inhibitory STDP Jointly Tune Feedforward Neural Circuits to Selectively Propagate Correlated Spiking Activity". In: *Frontiers in Computational Neuroscience* 8. DOI: 10.3389/fncom.2014.00053.

Kuan, Aaron T. et al. (Feb. 21, 2024). "Synaptic Wiring Motifs in Posterior Parietal Cortex Support Decision-Making". In: *Nature*, pp. 1–7. DOI: 10.1038/s41586-024-07088-7.

Lagzi, Fereshteh, Martha Canto Bustos, Anne-Marie Oswald, and Brent Doiron (Sept. 7, 2021). "Assembly Formation Is Stabilized by Parvalbumin Neurons and Accelerated by Somatostatin Neurons". In: *bioRxiv*. DOI: 10.1101/2021.09.06.459211.

Lagzi, Fereshteh and Adrienne L. Fairhall (Mar. 20, 2024). "Emergence of Co-Tuning in Inhibitory Neurons as a Network Phenomenon Mediated by Randomness, Correlations, and Homeostatic Plasticity". In: *Science Advances* 10.12, eadi4350. DOI: 10.1126/sciadv.adi4350.

Litwin-Kumar, Ashok and Brent Doiron (Nov. 14, 2014). "Formation and maintenance of neuronal assemblies through synaptic plasticity". In: *Nature Communications* 5, p. 5319. DOI: 10.1038/ncomms6319.

Luz, Yotam and Maoz Shamir (July 7, 2014). "The Effect of STDP Temporal Kernel Structure on the Learning Dynamics of Single Excitatory and Inhibitory Synapses ". In: *PLOS ONE* 9.7, e101109. DOI: 10.1371/journal.pone.0101109.

Mackwood, Owen, Laura B Naumann, and Henning Sprekeler (Apr. 26, 2021). "Learning Excitatory-Inhibitory Neuronal Assemblies in Recurrent Networks". In: *eLife* 10. Ed. by Stephanie E Palmer and Richard B Ivry, e59715. DOI: 10.7554/eLife.59715.

Maes, Amadeus, Mauricio Barahona, and Claudia Clopath (Mar. 25, 2021). "Learning Compositional Sequences with Multiple Time Scales through a Hierarchical Network of Spiking Neurons". In: *PLOS Computational Biology* 17.3, e1008866. DOI: 10.1371/journal.pcbi.1008866.

Manz, Paul and Raoul-Martin Memmesheimer (Apr. 12, 2023). "Purely STDP-based Assembly Dynamics: Stability, Learning, Overlaps, Drift and Aging". In: *PLOS Computational Biology* 19.4, e1011006. DOI: 10.1371/journal.pcbi.1011006.

Miehl, Christoph and Julijana Gjorgjieva (Dec. 2, 2022). "Stability and Learning in Excitatory Synapses by Nonlinear Inhibitory Plasticity". In: *PLOS Computational Biology* 18.12, e1010682. DOI: 10.1371/journal.pcbi.1010682.

Miehl, Christoph, Sebastian Onasch, Dylan Festa, and Julijana Gjorgjieva (2023). "Formation and Computational Implications of Assemblies in Neural Circuits". In: *The Journal of Physiology* 601.15, pp. 3071–3090. DOI: 10.1113/JP282750.

Mongillo, Gianluigi, Simon Rumpel, and Yonatan Loewenstein (Oct. 2018). "Inhibitory Connectivity Defines the Realm of Excitatory Plasticity". In: *Nature Neuroscience* 21.10, pp. 1463–1470. DOI: 10.1038/s41593-018-0226-x.

Montangie, Lisandro, Christoph Miehl, and Julijana Gjorgjieva (May 8, 2020). "Autonomous Emergence of Connectivity Assemblies via Spike Triplet Interactions". In: *PLOS Computational Biology* 16.5, e1007835. DOI: 10.1371/journal.pcbi.1007835.

Mulholland, Haleigh N, Bettina Hein, Matthias Kaschube, and Gordon B Smith (Dec. 8, 2021). "Tightly Coupled Inhibitory and Excitatory Functional Networks in the Developing Primary Visual Cortex". In: *eLife* 10, e72456. DOI: 10.7554/eLife.72456.

Ocker, Gabriel K. et al. (Oct. 1, 2017). "From the Statistics of Connectivity to the Statistics of Spike Times in Neuronal Networks". In: *Current Opinion in Neurobiology*. Computational Neuroscience 46, pp. 109–119. DOI: 10.1016/j.conb.2017.07.011.

Ocker, Gabriel Koch and Brent Doiron (Mar. 1, 2019). "Training and Spontaneous Reinforcement of Neuronal Assemblies by Spike Timing Plasticity". In: *Cerebral Cortex* 29.3, pp. 937–951. DOI: 10.1093/cercor/bhy001.

Ocker, Gabriel Koch, Ashok Litwin-Kumar, and Brent Doiron (Aug. 20, 2015). "Self-Organization of Microcircuits in Networks of Spiking Neurons with Plastic Synapses". In: *PLOS Comput Biol* 11.8, e1004458. DOI: 10.1371/journal.pcbi.1004458.

Peng, Yangfan et al. (June 16, 2021). "Spatially Structured Inhibition Defined by Polarized Parvalbumin Interneuron Axons Promotes Head Direction Tuning". In: *Science Advances* 7.25, eabg4693. DOI: 10.1126/sciadv.abg4693.

Petrucco, Luigi et al. (Apr. 24, 2023). "Neural Dynamics and Architecture of the Heading Direction Circuit in Zebrafish". In: *Nature Neuroscience*, pp. 1–9. DOI: 10.1038/s41593-023-01308-5.

Ravid Tannenbaum, Neta and Yoram Burak (Aug. 12, 2016). "Shaping Neural Circuits by High Order Synaptic Interactions". In: *PLOS Computational Biology* 12.8, e1005056. DOI: 10.1371/journal.pcbi.1005056.

Reimann, Michael W., Daniela Egas Santander, András Ecker, and Eilif B. Muller (Dec. 22, 2023). *Specific Inhibition and Disinhibition in the Higher-Order Structure of a Cortical Connectome*. DOI: 10.1101/2023.12.22.573036.

Riquelme, Juan Luis, Mike Hemberger, Gilles Laurent, and Julijana Gjorgjieva (Feb. 13, 2023). "Single Spikes Drive Sequential Propagation and Routing of Activity in a Cortical Network". In: *eLife* 12. Ed. by Peter Latham, Laura L Colgin, and Abigail Morrison, e79928. DOI: 10.7554/eLife.79928.

Rubin, Ran, L. F. Abbott, and Haim Sompolinsky (Oct. 17, 2017). "Balanced Excitation and Inhibition Are Required for High-Capacity, Noise-Robust Neuronal Selectivity". In: *Proceedings of the National Academy of Sciences*, p. 201705841. DOI: 10.1073/pnas.1705841114.

Sadeh, Sadra and Claudia Clopath (Oct. 27, 2020). "Theory of Neuronal Perturbome in Cortical Networks". In: *Proceedings of the National Academy of Sciences* 117.43, pp. 26966–26976. DOI: 10.1073/pnas.2004568117.

– (Nov. 5, 2021). "Excitatory-Inhibitory Balance Modulates the Formation and Dynamics of Neuronal Assemblies in Cortical Networks". In: *Science Advances* 7.45, eabg8411. DOI: 10.1126/sciadv.abg8411.

Schulz, Auguste, Christoph Miehl, Michael J Berry II, and Julijana Gjorgjieva (Oct. 14, 2021). "The Generation of Cortical Novelty Responses through Inhibitory Plasticity". In: *eLife* 10. Ed. by Maria N Geffen, Joshua I Gold, and Maria N Geffen, e65309. DOI: 10.7554/eLife.65309.

Smith, Gordon B., Bettina Hein, David E. Whitney, David Fitzpatrick, and Matthias Kaschube (Nov. 2018). "Distributed Network Interactions and Their Emergence in Developing Neocortex". In: *Nature Neuroscience* 21.11 (11), pp. 1600–1608. DOI: 10.1038/s41593-018-0247-5.

Sreenivasan, Varun et al. (May 17, 2022). "Input-Specific Control of Interneuron Numbers in Nascent Striatal Networks". In: *Proceedings of the National Academy of Sciences* 119.20, e2118430119. DOI: 10.1073/pnas.2118430119.

Stimberg, Marcel, Romain Brette, and Dan FM Goodman (Aug. 20, 2019). "Brian 2, an Intuitive and Efficient Neural Simulator". In: *eLife* 8. Ed. by Frances K Skinner, Ronald L Calabrese, Frances K Skinner, Fleur Zeldenrust, and Richard C Gerkin, e47314. DOI: 10.7554/eLife.47314.

Tremblay, Robin, Soohyun Lee, and Bernardo Rudy (July 20, 2016). "GABAergic Interneurons in the Neocortex: From Cellular Properties to Circuits". In: *Neuron* 91.2, pp. 260–292. DOI: 10.1016/j.neuron.2016.06.033.

Triplett, Marcus A., Lilach Avitan, and Geoffrey J. Goodhill (Sept. 28, 2018). "Emergence of Spontaneous Assembly Activity in Developing Neural Networks without Afferent Input". In: *PLOS Computational Biology* 14.9, e1006421. DOI: 10.1371/journal.pcbi.1006421.

Tsodyks, M. V. and T. Sejnowski (May 1995). "Rapid State Switching in Balanced Cortical Network Models". In: *Network: Computation in Neural Systems* 6.2, p. 111. DOI: 10.1088/0954-898X/6/2/001.

Vignoud, Gaëtan and Philippe Robert (May 10, 2022). "Spontaneous Dynamics of Synaptic Weights in Stochastic Models with Pair-Based Spike-Timing-Dependent Plasticity". In: *Physical Review E* 105.5, p. 054405. DOI: 10.1103/PhysRevE.105.054405.

Vogels, Tim P., H. Sprekeler, F. Zenke, C. Clopath, and W. Gerstner (Nov. 10, 2011). "Inhibitory Plasticity Balances Excitation and Inhibition in Sensory Pathways and Memory Networks". In: *Science* 334.6062, pp. 1569–1573. DOI: 10.1126/science.1211095.

Van Vreeswijk, C. and H. Sompolinsky (1998). "Chaotic balanced state in a model of cortical circuits". In: *Neural Computation* 10.6, pp. 1321–1371.

Wallenstein, Gene V. and Michael E. Hasselmo (July 1997). "GABAergic Modulation of Hippocampal Population Activity: Sequence Learning, Place Field Development, and the Phase Precession Effect". In: *Journal of Neurophysiology* 78.1, pp. 393–408. DOI: 10.1152/jn.1997.78.1.393.

Wu, Sherry Jingjing et al. (Sept. 6, 2023). "Cortical Somatostatin Interneuron Subtypes Form Cell-Type-Specific Circuits". In: *Neuron* 111.17, 2675–2692.e9. DOI: 10.1016/j.neuron.2023.05.032.

613   Xie, Xiaohui, Richard H. R. Hahnloser, and H. Sebastian Seung (Oct. 9, 2002). "Double-Ring Network Model of
614   the Head-Direction System". In: *Physical Review E* 66.4, p. 041902. DOI: 10.1103/PhysRevE.66.041902.
615   Znamenskiy, Petr et al. (Mar. 20, 2024). "Functional Specificity of Recurrent Inhibition in Visual Cortex". In: *Neuron*
616   112.6, 991–1000.e8. DOI: 10.1016/j.neuron.2023.12.013.

Structured stabilization in recurrent neural circuits through inhibitory synaptic plasticity

S1 Analytic results

S1.1 Mean-field equation for weight dynamics

In this section, we adapt the results introduced by Kempter et al. (1999) to our notation and our specific needs. We first describe a sufficiently general expression for the pairwise STDP rules. Given two neurons, one presynaptic and one postsynaptic, an excitatory or inhibitory STDP rule is a weight update triggered by either a pre- or a postsynaptic spike. The weight update solely depends on the time differences between the previous spike times, relative to the current spike time:

$$\text{on } t = t_k^{(\text{pre})} \ \rightarrow \ \Delta W^{(\text{pre})} = \alpha_{\text{pre}} + \sum_{k' < k} L^{\Delta_-}\left(t - t_{k'}^{(\text{post})}\right)$$

$$\text{on } t = t_k^{(\text{post})} \ \rightarrow \ \Delta W^{(\text{post})} = \alpha_{\text{post}} + \sum_{k' < k} L^{\Delta_+}\left(t - t_{k'}^{(\text{pre})}\right) \ , \tag{S1}$$

where $t_k^{(\text{pre})}$ and $t_k^{(\text{post})}$ represent pre- and postsynaptic spike events, respectively. We can merge the functions L^{Δ_-} and L^{Δ_+} into a single STDP kernel defined as:

$$L(\tau) := \begin{cases} L^{\Delta_+}(\tau) & \text{if } \tau \geq 0 \\ L^{\Delta_-}(-\tau) & \text{if } \tau < 0 \ . \end{cases} \tag{S2}$$

Therefore $\tau > 0$ (the Δ_+ branch) represents pre-post interactions (presynaptic spike preceding the postsynaptic spike), while $\tau < 0$ (the Δ_- branch) represents post-pre interactions (postsynaptic spike preceding the presynaptic spike).

It is useful to express each spike train as a sum of Dirac delta functions centered at each spike time:

$$S^i(t) := \sum_k \delta(t - t_k^i) \tag{S3}$$

where i denotes the neuron index. With this notation, the number of spikes in the $[t_a, t_b]$ interval is:

$$N_{[t_a, t_b]}^i = \int_{t_a}^{t_b} dt \ S^i(t) \tag{S4}$$

Assuming stationary conditions, we can therefore define the firing rate of neuron i as:

$$r_i := \frac{1}{T} \int_0^T dt \ S^i(t) = \langle S^i(t) \rangle_t \ . \tag{S5}$$

Using the definition in Eq. S3, we can then rewrite the pairwise interaction terms in Eq. S1 as:

$$\sum_{k'} L^{\Delta_\pm}(t - t_{k'}^i) = \int_{-\infty}^t d\tau \ S^i(\tau) \ L^{\Delta_\pm}(t - \tau) = \int_0^\infty d\tau^* \ S^i(t - \tau^*) \ L^{\Delta_\pm}(\tau^*) \tag{S6}$$

Where in the last step we performed the change of variable $\tau \leftarrow t - \tau^*$. This leads to the following expression for the instantaneous change in synaptic weights, numerically equivalent to Eq. S1:

$$\text{on } t = t_k^{(\text{pre})} \ \rightarrow \ \Delta W^{(\text{pre})}(t) = \alpha_{\text{pre}} + \int_0^\infty d\tau \ S^{\text{post}}(t - \tau) \ L^{\Delta_-}(\tau)$$

$$\text{on } t = t_k^{(\text{post})} \ \rightarrow \ \Delta W^{(\text{post})}(t) = \alpha_{\text{post}} + \int_0^\infty d\tau \ S^{\text{pre}}(t - \tau) \ L^{\Delta_+}(\tau) \tag{S7}$$

Now, using again the same definition of spike train in Eq. S3, an event-based update can be expressed as an integral over time over the pre- and post-synaptic spiking events. The total weight change occurring in an interval $[t_a, t_b]$ is:

$$\Delta W_{[t_a, t_b]} = \int_{t_a}^{t_b} dt\, S^{\text{pre}}(t)\, \Delta W^{(\text{pre})}(t) + \int_{t_a}^{t_b} S^{\text{post}}(t)\, \Delta W^{(\text{post})}(t)\,. \tag{S8}$$

We can then express the average weight change per second, considering a time interval $[0, T]$ large enough to ensure precise sampling in second-order firing statistics, but still small compared to the rate of changes in ΔW. Under this assumption of separation of timescales between neuronal dynamics and plasticity, we can substitute Eq. S7 into Eq. S8, and write:

$$\langle \Delta W(t) \rangle_t = \frac{1}{T} \int_0^T dt\, S^{\text{pre}}(t)\, \alpha_{\text{pre}} + \frac{1}{T} \int_0^T dt\, S^{\text{post}}(t)\, \alpha_{\text{post}} +$$
$$+ \frac{1}{T} \int_0^T dt \int_0^\infty d\tau\, S^{\text{post}}(t)\, S^{\text{pre}}(t - \tau)\, L^{\Delta_+}(\tau) + \frac{1}{T} \int_0^T dt \int_0^\infty d\tau\, S^{\text{pre}}(t)\, S^{\text{post}}(t - \tau)\, L^{\Delta_-}(\tau). \tag{S9}$$

Since α_{pre} and α_{post} are constants, we can factor them out. The first two terms of the r.h.s. of Eq. S9 become:

$$\frac{1}{T} \int_0^T dt\, S^{\text{pre}}(t) \alpha_{\text{pre}} = r_{\text{pre}}\, \alpha_{\text{pre}} \quad \text{and} \quad \frac{1}{T} \int_0^T dt\, S^{\text{post}}(t) \alpha_{\text{post}} = r_{\text{post}}\, \alpha_{\text{post}} \tag{S10}$$

The pairwise interaction terms instead can be simplified as follows:

$$\frac{1}{T} \int_0^T dt \int_0^\infty d\tau\, S^{\text{post}}(t)\, S^{\text{pre}}(t - \tau)\, L^{\Delta_+}(\tau) = r_{\text{pre}}\, r_{\text{post}} \int_0^\infty d\tau\, L^{\Delta_+}(\tau) + \int_0^\infty d\tau\, C^{\Delta_+}(\tau) L^{\Delta_+}(\tau), \tag{S11}$$

and similarly for the last term in Eq. S9. In the above equation, $C^{\Delta_+}(\tau)$ denotes the covariance density between the pre- and the postsynaptic neuron, following the convention that Δ_+ represents the pre-post order and Δ_- the post-pre order:

$$C^{\Delta_+}(\tau) := \frac{1}{T} \int_0^T \left(S^{\text{post}}(t) - r_{\text{post}} \right) \left(S^{\text{pre}}(t - \tau) - r_{\text{pre}} \right)\,. \tag{S12}$$

An analogous definition holds for $C^{\Delta_-}(\tau)$, with the pre- and post- temporal order inverted. Therefore

$$\langle \Delta W(t) \rangle_t = r_{\text{pre}}\, \alpha_{\text{pre}} + r_{\text{post}}\, \alpha_{\text{post}} + r_{\text{pre}}\, r_{\text{post}} \int_0^\infty d\tau\, \left(L^{\Delta_-}(\tau) + L^{\Delta_+}(\tau) \right) +$$
$$+ \int_0^\infty d\tau\, C^{\Delta_-}(\tau) L^{\Delta_-}(\tau) + \int_0^\infty d\tau\, C^{\Delta_+}(\tau) L^{\Delta_+}(\tau)\,. \tag{S13}$$

Now using Eq. S2, we can write:

$$r_{\text{pre}}\, r_{\text{post}} \int_0^\infty d\tau\, \left(L^{\Delta_-}(\tau) + L^{\Delta_+}(\tau) \right) = r_{\text{pre}}\, r_{\text{post}} \int_{-\infty}^\infty d\tau\, L(\tau)\,; \tag{S14}$$

and given that $C^{\Delta_+}(\tau) = C^{\Delta_-}(-\tau)$ we can perform the following variable substitution:

$$\int_0^\infty d\tau\, C^{\Delta_-}(\tau) L^{\Delta_-}(\tau) = \int_{-\infty}^0 d\tau\, C^{\Delta_+}(\tau) L^{\Delta_+}(\tau) \tag{S15}$$

Finally, substituting Eq. S14 and Eq. S15 in Eq. S13, and defining $C(\tau) := C^{\Delta_+}(\tau)$, we have:

$$\langle \Delta W(t) \rangle_t = \alpha_{\text{pre}} r_{\text{pre}} + \alpha_{\text{post}} r_{\text{post}} + B\, r_{\text{pre}}\, r_{\text{post}} + Q\quad,$$
$$\text{with} \quad B := \int_{-\infty}^{+\infty} d\tau\, L(\tau) \quad \text{and} \quad Q := \int_{-\infty}^{+\infty} d\tau\, L(\tau)\, C(\tau)\,; \tag{S16}$$

Corresponding to Eq. 1 in the main text.

S1.2 Derivation of the closed-form solution for the E/I motif

In this section, we derive a closed-form solution for a reduced model of one inhibitory and one excitatory neuron (Eq. S16). We model the two neurons as mutually interacting Hawkes processes (Main Text, Fig. 1A, Hawkes 1971; Pernice et al. 2011). The excitatory unit receives a constant input current h_{exc}, a plastic inhibitory connection $w_{\text{inh}}(t)$ and it projects a fixed excitatory connection w_{exc} to the inhibitory neuron. For simplicity, we assume that the inhibitory unit is kept at a constant rate r_{inh} by an adaptive external current. In this framework, the average firing rates behave linearly, therefore, the excitatory rate can be written as:

$$r_{\text{exc}}(t) = h_{\text{exc}} - w_{\text{inh}}(t)\, r_{\text{inh}} \, . \tag{S17}$$

As we focus on the connection $w_{\text{inh}}(t)$ from the inhibitory to the excitatory neuron, we rename $r_{\text{inh}} = r_{\text{pre}}$ and $r_{\text{exc}} = r_{\text{post}}$ in Eq. S16 and we get:

$$\left\langle \frac{dw_{\text{inh}}(t)}{dt} \right\rangle = \alpha_{\text{pre}} r_{\text{inh}} + \alpha_{\text{post}} \left(h_{\text{exc}} - w_{\text{inh}} r_{\text{inh}} \right) + B\, r_{\text{inh}}\, \left(h_{\text{exc}} - w_{\text{inh}} r_{\text{inh}} \right) + Q$$
$$= \alpha_{\text{pre}} r_{\text{inh}} - w_{\text{inh}} r_{\text{inh}} \left(\alpha_{\text{post}} + B\, r_{\text{inh}} \right) + h_{\text{exc}} \left(\alpha_{\text{post}} + B\, r_{\text{inh}} \right) + Q. \tag{S18}$$

The only nontrivial quantity to compute is the term Q, as it requires the inhibitory-to-excitatory covariance density curve. To avoid ambiguities, we define the Fourier transform operator acting on a function $f(t)$ and the convolution operator acting on two functions $f(t)$ and $g(t)$ as follows:

$$\hat{f}(\omega) = \int_{-\infty}^{+\infty} dt \;\; f(t)\, e^{-j\,\omega\, t} \quad \text{and} \quad (f * g)(\tau) = \int_{-\infty}^{+\infty} dt \;\; f(t)\, g(\tau - t) \; ; \tag{S19}$$

where j denotes the imaginary unit.

Note that the total integral Q of the product $C(\tau)\, L(\tau)$ (as defined in Eq. S16) is equivalent to its Fourier transform evaluated at $\omega = 0$. But for the inverse convolution theorem, the Fourier transform of the product $C(\tau)\, L(\tau)$ corresponds to the convolution of the two transforms $\hat{C}(\omega)$ and $\hat{L}(\omega)$. In equations:

$$Q = \int_{-\infty}^{\infty} d\tau\; C(\tau)\, L(\tau) = \hat{f}\left(C(\tau)\, L(\tau) \right)(\omega)\Big|_{\omega=0} = \frac{1}{2\pi}\; \left(\hat{C}(\tilde{\omega}) * \hat{L}(\tilde{\omega}) \right)(\omega)\Big|_{\omega=0} =$$
$$= \frac{1}{2\pi}\; \int_{-\infty}^{\infty} d\tilde{\omega}\; \hat{C}(\tilde{\omega})\, \hat{L}(-\tilde{\omega}) \, . \tag{S20}$$

The advantage of re-defining Q in the Fourier space is that, for Hawkes processes, the covariance $\hat{C}(\omega)$ can be expressed in a closed form as a function of network connectivity and neural activity (Hawkes 1971; Pernice et al. 2011; Trousdale et al. 2012; Jovanović et al. 2015). More specifically, given the matrix of post-synaptic potential kernels (Eq. 9 in Methods)

$$\mathbf{E}(\tau) = \begin{pmatrix} \frac{1}{\tau_e}\, e^{-\tau/\tau_e} & 0 \\ 0 & -\frac{1}{\tau_i}\, e^{-\tau/\tau_i} \end{pmatrix} \tag{S21}$$

and the connectivity matrix

$$\mathbf{W} = \begin{pmatrix} 0 & w_{\text{inh}} \\ w_{\text{exc}} & 0 \end{pmatrix} , \tag{S22}$$

we can first write the vector of firing rates $\mathbf{r} = \left[r_{\text{exc}} \, , \, r_{\text{inh}} \right]^{\top}$ as

$$\mathbf{r} = \mathbf{h_0} + \int_{-\infty}^{+\infty} d\tau \;\; \mathbf{G}(\tau)\, S(\tau - t), \tag{S23}$$

where $\mathbf{h_0}$ is the baseline input and $\mathbf{G}(\tau) = \mathbf{E}(\tau)\, \mathbf{W}$ is referred to as the matrix of "interaction kernels". At the equilibrium state,

$$\mathbf{r} = \mathbf{h_0} + \mathbf{G}\, \mathbf{r}, \quad \text{with } \mathbf{G} := \int_{-\infty}^{+\infty} d\tau \;\; \mathbf{G}(\tau), \tag{S24}$$

where the integral in the r.h.s. corresponds to the Fourier transform evaluated at zero, i.e., $\hat{\mathbf{G}}(\omega)|_{\omega=0}$. Equivalently, the rate vector can be written in an explicit form:

$$\mathbf{r} = (\mathbf{I} - \mathbf{G})^{-1}\,\mathbf{h_0}, \tag{S25}$$

where $\mathbf{I}$ represents the identity matrix. Moreover, if we write the covariance matrix (cf. Eq. S12) as

$$\hat{\mathbf{C}}(\tau) = \langle \mathbf{S}(t)\,\mathbf{S}(t-\tau)^T \rangle - \mathbf{r}\,\mathbf{r}^T \tag{S26}$$

then we have the following expression in the Fourier domain, at the equilibrium state:

$$\hat{\mathbf{C}}(\omega) = \left(\mathbf{I} - \hat{\mathbf{G}}(\omega)\right)^{-1}\mathbf{D}\left(\mathbf{I} - \hat{\mathbf{G}}^T(\omega)\right)^{-1} \tag{S27}$$

where $\mathbf{D}$ is a diagonal matrix with the firing rates on its diagonal. Finally, recall that, for a matrix $\mathbf{G}$ with spectral radius < 1,

$$(\mathbf{I} - \mathbf{G})^{-1} = \sum_{\alpha}^{\infty} \mathbf{G}^{\alpha}. \tag{S28}$$

Therefore, the covariance matrix can be written as the following expansion

$$\hat{\mathbf{C}}(\omega) = \sum_{\alpha=0}^{N}\sum_{\beta=0}^{N} \hat{\mathbf{E}}^{\alpha}(\omega)(\hat{\mathbf{E}}^T)^{\beta}(-\omega)\left[(\mathbf{W}^{\alpha})\mathbf{D}(\mathbf{W}^T)^{\beta})\right], \tag{S29}$$

where the indices α,β indicate the number of connections between any spiking neuron in the network to the post- and to the presynaptic neuron, respectively. This captures the influence of each spike propagated in the network through existing synaptic connections on the correlated firing of the pre- and postsynaptic neurons. Within this formalism, in our reduced model we consider only pairwise interactions, that is, $\alpha + \beta = 1$. The inhibitory-to-excitatory covariance is the $(1,2)$ element of $\hat{\mathbf{C}}(\omega)$. We can compute it as:

$$\hat{C}(\omega) = \hat{E}_e(\omega)\,w_{\text{inh}}\,r_{\text{inh}} + \hat{E}_i(-\omega)\,r_{\text{exc}}\,w_{\text{exc}} \tag{S30}$$

By combining Eq. S20 and Eq. S30, we get the following expression for Q:

$$Q = M_{ei}^{(10)}w_{\text{inh}}\,r_{\text{inh}} + M_{ie}^{(01)}w_{\text{exc}}\,r_{\text{exc}}$$
$$\text{with} \quad M_{jk}^{(\alpha\beta)} := \int_{-\infty}^{+\infty} d\omega\,\hat{E}_j^{\alpha}(\omega)\hat{E}_k^{\beta}(-\omega)\hat{L}(-\omega). \tag{S31}$$

The $M_{ei}^{(10)}$, $M_{ie}^{(01)}$ coefficients can be computed analytically depending on the iSTDP rule (see next section). Considering them as constants, we can then write Eq. S18 as:

$$\left\langle \frac{dw_{\text{inh}}(t)}{dt} \right\rangle = \alpha_{\text{pre}}r_{\text{inh}} - w_{\text{inh}}\,r_{\text{inh}}\left(\alpha_{\text{post}} + B\,r_{\text{inh}}\right) + h_{\text{exc}}\left(\alpha_{\text{post}} + B\,r_{\text{inh}}\right) +$$
$$M_{ei}^{(10)}w_{\text{inh}}\,r_{\text{inh}} + M_{ie}^{(01)}w_{\text{exc}}\,r_{\text{exc}} \tag{S32}$$

or equivalently,

$$\left\langle \frac{dw_{\text{inh}}(t)}{dt} \right\rangle = \alpha_{\text{pre}}r_{\text{inh}} - w_{\text{inh}}\,r_{\text{inh}}\left(\Psi - M_{ei}^{(10)}\right) + h_{\text{exc}}\,\Psi \tag{S33}$$

with $\Psi := \alpha_{\text{post}} + Br_{\text{inh}} + M_{ie}^{(01)}w_{\text{exc}}$.

We then find the stationary point (Eq. 10 in Methods)

$$\tilde{w}_{\text{inh}} = \frac{\alpha_{\text{pre}}\,r_{\text{inh}} + h_{\text{exc}}\Psi}{r_{\text{inh}}\left(\Psi - M_{ei}^{(10)}\right)} \tag{S34}$$

The equilibrium point $\tilde{w}_{\text{inh}}$ is a stable fixed point attractor, and positive, that is $\tilde{w}_{\text{inh}} \in [0, w_{\text{max}}]$, if the following conditions hold:

$$\Psi - M_{ei}^{(10)} > 0$$
$$\alpha_{\text{pre}}\,r_{\text{inh}} + \Psi\,h_{\text{exc}} > 0, \tag{S35}$$

If a solution for $\tilde{w}_{\text{inh}}$ exists, we can also express the steady-state postsynaptic rate as follows:

$$\tilde{r}_{\text{exc}} = h_{\text{exc}} - \tilde{w}_{\text{inh}} \, r_{\text{inh}} = -\frac{\alpha_{\text{pre}} r_{\text{inh}} + M_{ei}^{(10)} h_{\text{post}}}{\Psi - M_{ei}^{(10)}} \quad , \tag{S36}$$

The solution is valid only for positive rates. This adds one more condition for a stable and positive solution. The constraints are therefore:

$$\Psi - M_{ei}^{(10)} > 0$$
$$\alpha_{\text{pre}} \, r_{\text{inh}} + \Psi \, h_{\text{exc}} > 0$$
$$r_{\text{inh}} > \frac{M_{ei}^{(10)} \, h_{\text{exc}}}{-\alpha_{\text{pre}}} \quad \text{with } \alpha_{\text{pre}} < 0. \tag{S37}$$

S1.3 Derivation of first-order motif coefficients

To calculate the motif coefficients

$$M_{ei}^{(10)} = \int_{-\infty}^{+\infty} d\omega \, \hat{L}(-\omega) \, \hat{E}_i(\omega) \quad \text{and} \quad M_{ie}^{(01)} = \int_{-\infty}^{+\infty} d\omega \, \hat{L}(-\omega) \, \hat{E}_e(-\omega) \tag{S38}$$

for the connection W, we need to compute the Fourier transform of the postsynaptic potential E_e, E_i and the Fourier transform of the iSTDP kernel.

The Fourier transforms of the excitatory and inhibitory postsynaptic potential (Eq. 9 in Methods) are respectively given by:

$$\hat{E}_e(\omega) = \frac{1}{1 + j\tau_e\omega}, \quad \text{and} \quad \hat{E}_i(\omega) = -\frac{1}{1 + j\tau_i\omega}, \tag{S39}$$

where j is the imaginary unit.

Consider first the Hebbian symmetric STDP kernel, defined via the function

$$L^{\text{sy}}(\tau) = \frac{1}{2} \left(\frac{1}{\tau_+} \, A_+ \, e^{-|\tau|/\tau_+} - \frac{1}{\tau_-} \, A_- \, e^{|\tau|/\tau_-} \right) \tag{S40}$$

with A_+, $A_- > 0$. The Fourier transform is given by

$$\hat{L}^{\text{sy}}(\omega) = \hat{L}^{\text{sy}}(-\omega) = \frac{A_+}{1 + \tau_+^2\omega^2} - \frac{A_-}{1 + \tau_-^2\omega^2}. \tag{S41}$$

By substituting Eq. S39 and Eq. S41 in Eq. S38, we can calculate the first-order motif coefficients $M_{ei}^{\text{sy},(10)}$, $M_{ie}^{\text{sy},(01)}$:

$$M_{ei}^{\text{sy},(10)} = \frac{A_+}{\tau_+ + \tau_i} - \frac{A_-}{\tau_- + \tau_i} \quad ; \quad M_{ie}^{\text{sy},(01)} = \frac{A_+}{\tau_+ + \tau_e} - \frac{A_-}{\tau_- + \tau_e}, \tag{S42}$$

and therefore find an analytical solution for the integral Q in Eq. S31.

Consider now the antisymmetric iSTDP rule, with kernel:

$$L^{\text{an}}(\tau) = \begin{cases} -\frac{A_-}{\tau_-} \, e^{\tau/\tau_-} & \text{if } \tau < 0 \\ \frac{A_+}{\tau_+} \, e^{-\tau/\tau_+} & \text{if } \tau \geq 0 \end{cases} \tag{S43}$$

whose Fourier transform is given by

$$\hat{L}^{\text{an}}(\omega) = \frac{A_+}{1 + j\tau_+\omega} - \frac{A_-}{1 - j\tau_-\omega}. \tag{S44}$$

Again, we can calculate the first order motif coefficients $M_{ei}^{\text{an},(10)}$, $M_{ie}^{\text{an},(01)}$ for the antisymmetric rule by substituting Eq. S39 and Eq. S44 in Eq. S38:

$$M_{ei}^{\text{an},(10)} = \frac{A_+}{\tau_+ + \tau_i} \quad ; \quad M_{ie}^{\text{an},(01)} = -\frac{A_-}{\tau_- + \tau_e}. \tag{S45}$$

and find an analytical solution for Eq. S31.

 # S2 Supplementary figures

Figure S1: Analytic table solutions for different iSTDP interaction kernels (complements Fig. 2 in main text).
A. Representation of the model (see also Fig. 1A). **B-E**. iSTDP rules, represented by their kernel and their additional parameters (left panels) and study of the associated weight dynamics as a one-dimensional differential equation (right panels), black dots represent fixed points. **B** Rate-dominated iSTDP rule. **C-E**. covariance dominated rules presented in the same order as in Fig. 2A, main text. Note that the configurations in **E** and **F** produce unstable fixed points.

Figure S2: Full weight matrix for the random sparse RNN simulation with symmetric iSTDP (see Main Text, Fig. 3B-F).

Figure S3: Full weight matrix for the random sparse RNN simulation with antisymmetric iSTDP (see Main Text, Fig. 3G-K).

Figure S4: Full weight matrix for the recurrent network with 1D ring connectivity structure after weight convergence with two inhibitory populations.

Figure S5: **A**. Full matrix of Pearson correlations between excitatory neurons in the one-dimensional ring model during spontaneous activity, after convergence of plastic weights. **B** Response of the one-dimensional ring model to localized stimuli (see Fig. 4 in main text).

References for Supplementary Material

Hawkes, Alan G. (1971). "Spectra of Some Self-Exciting and Mutually Exciting Point Processes". In: *Biometrika* 58.1, pp. 83–90. DOI: 10.2307/2334319.

Jovanović, Stojan, John Hertz, and Stefan Rotter (Apr. 7, 2015). "Cumulants of Hawkes Point Processes". In: *Physical Review E* 91.4, p. 042802. DOI: 10.1103/PhysRevE.91.042802.

Kempter, Richard, Wulfram Gerstner, and J. Leo van Hemmen (Apr. 1, 1999). "Hebbian Learning and Spiking Neurons". In: *Physical Review E* 59.4, pp. 4498–4514. DOI: 10.1103/PhysRevE.59.4498.

Pernice, Volker, Benjamin Staude, Stefano Cardanobile, and Stefan Rotter (May 19, 2011). "How Structure Determines Correlations in Neuronal Networks". In: *PLOS Computational Biology* 7.5, e1002059. DOI: 10.1371/journal.pcbi.1002059.

Trousdale, James, Yu Hu, Eric Shea-Brown, and Krešimir Josić (Mar. 22, 2012). "Impact of Network Structure and Cellular Response on Spike Time Correlations". In: *PLoS Comput Biol* 8.3, e1002408. DOI: 10.1371/journal.pcbi.1002408.