MeNu Guide
A Metabolite Nutrition Graph to Uncover Interactions with Disease Etiology

Vivian Würf[1,ORCID]   Josch K. Pauling[1,2,*]

[1] LipiTUM, TUM School of Life Sciences, Technical University of Munich, 85354 Freising, Germany

[2] Institute for Clinical Chemistry and Laboratory Medicine, University Hospital and Faculty of Medicine Carl Gustav Carus of the Dresden University of Technology, 01307 Dresden, Germany

[*] Correspondence to `josch.pauling@tum.de`

Abstract

The relationship between diet and disease is well-documented, yet the complex interactions among foods, metabolites, and genetics makes research challenging. This study explores the potential insights offered by a knowledge graph that connects nutrition and diseases on a metabolic level. Ten ontologies and data from six databases were merged, resulting in a graph with over 25 million triple statements, stored in a Turtle file and added to a GraphDB repository. SPARQL queries revealed biases towards specific foods and conditions within the integrated databases. Despite these biases, this knowledge graph serves as a proof-of-concept, demonstrating the feasibility of integrating information from diverse resources to yield valuable insights and enabling the drawing of meaningful conclusions. The graph allows efficient identification of disease-related compounds and their food sources and enables the exploration of changes in metabolite concentrations, such as those occurring during food processing. Researchers could use such a knowledge graph to identify biomarkers, help generate new hypotheses, and improve experimental designs. Expanding the graph with automated text-mining and recipe data would further enhance its utility for nutrition research. Such a resource could advance understanding of the molecular mechanisms behind diet-disease relationships, guiding more targeted interventions.

Keywords Nutrition · Metabolomics · Knowledge Graph · SPARQL · GraphDB

1 Introduction

Nutrition research grapples with multifaceted challenges that render it inherently complex. It is well-established that many diseases have a strong connection to nutrition, including obesity [1], cardiovascular conditions [2], type 2 diabetes mellitus [3] and non-alcoholic fatty liver disease [4]. This highlights the vast scope of how dietary patterns influence health and disease. Certain diets can directly contribute to disease development, such as alcohol consumption leading to alcoholic fatty liver disease [5], while others can be used therapeutically, like exclusive enteral nutrition in the treatment of Crohn's disease [6, 7]. However, the diversity of dietary patterns in different populations makes it challenging to generalize findings across varying cultural, socioeconomic, and geographical contexts. Additionally, conducting controlled experiments in nutrition research is challenging due to the long-term nature of diet-disease relationships.

Despite significant advancements, much remains unknown, especially regarding diet-related disorders and diseases. Conditions like Irritable Bowel Syndrome (IBS) and Inflammatory Bowel Disease (IBD) have reached a concerning level of global prevalence affecting millions of people worldwide [8–13]. However, our understanding of their underlying causes and optimal dietary interventions remains limited. This results in untapped potential for treatment, personalized interventions, supportive therapies, and disease prevention. While there is growing recognition of the role of diet in managing these conditions, tailored dietary strategies that effectively alleviate symptoms and improve the quality of life for patients are still lacking [14–17]. Despite considerable progress, the complexity of human metabolism and the multifactorial nature of disease etiology pose significant challenges to understanding the intricate connections between foods, metabolites, and diseases. Bridging this knowledge gap necessitates adopting innovative approaches to integrate information from heterogeneous data sources.

Multiple initiatives have been launched to compile comprehensive databases of food metabolites to understand their potential roles in human health and disease [18–20]. While the existing databases provide valuable information on the chemical composition of foods and their metabolites, gaps and inconsistencies persist. Additionally, they lack direct links to disease associations. Moreover, not every database utilizes standardized nomenclature and annotation for metabolites, which further complicates integrating data across different sources. As a result, while these databases serve as valuable resources for nutritional research, their limitations underscore the need for continued efforts to improve data quality, coverage, and interoperability to facilitate meaningful insights into the impact of food metabolites on human health.

Knowledge graphs are structured representations of data that capture and store entities and the relationships between them. They have emerged fairly recently as a promising tool for tackling complex biomedical problems. By organizing vast amounts of heterogeneous information from diverse sources such as scientific literature, databases, and electronic health records, knowledge graphs can provide a holistic view of biological systems. Researchers can then use graph analytics to extract meaningful insights and elucidate the intricate relationships between genes, diseases, drugs, and other biomedical entities. The success of knowledge graphs in biomedical research is evidenced by their diverse applications in areas such as drug repurposing, precision medicine, and clinical decision support [21–24]. Leveraging the wealth of interconnected data encapsulated within knowledge graphs enables researchers to navigate the complexities of biological systems, accelerate scientific discovery, and, ultimately, advance human health.

Here, we present the nutrition-disease knowledge graph MeNu GUIDE, which was constructed through a multifaceted approach that harnesses structured data from databases and ontologies (figure 1). Relevant data from various curated databases containing information on nutrition, metabolites, diseases, genes, and pathways were extracted and integrated to populate the knowledge graph. Our nutrition-disease knowledge graph offers a comprehensive and dynamic repository of interconnected information, facilitating easy, fast, and holistic exploration and analysis of the complex interrelationships between diet, metabolism, and health outcomes. By employing graph analytics, its utility extends beyond mere data aggregation, serving as a powerful tool for hypothesis generation and data-driven discovery. The ultimate goal of the nutrition-disease knowledge graph is to unravel the underlying mechanisms linking dietary factors to disease risk and progression, thereby informing evidence-based dietary recommendations and ultimately improving patient health outcomes.

2 Methods

2.1 Database Merging

Information from six different databases was extracted and merged. The utilized databases include the Food Database (FooDB) [18], the Human Metabolome Database (HMDB) [25], MarkerDB [26], Chemical Entities of Biological Interest (ChEBI) [27], Kyoto Encyclopedia of Genes and Genomes (KEGG) [28] and the Virtual Metabolic Human (VMH) database [29]. Three of these databases (FooDB, HMDB and MarkerDB) are curated by the Wishart reasearch group at the University of Alberta in Canada and therefore form an ideal starting point for the database integration process. Beginning with **FooDB**, all relevant data was downloaded as CSV files from the website. Compounds in the Compounds.csv file were matched with their external descriptors in the CompoundExternalDescriptor.csv file to obtain as many identifiers as possible for merging with the other databases. In the next step, the hmdb_metabolites.xml file was downloaded from the **HMDB** website. Relevant information and identifiers were extracted for each compound and saved as a CSV file. The extracted HMDB compounds were then matched with the FooDB compounds based on the lowercase compound names, as it yielded the best results. To extract all compounds present in **MarkerDB** the file all_chemicals.tsv was downloaded from the website. All unique compounds were extracted and the API was queried to retrieve information on each compound. As each entry was associated with an HMDB ID, this information was used to match them with the previously merged FooDB and HMDB compounds.

To integrate the **ChEBI** database, the files containing information on compounds were downloaded from the website. Initially, only the highest confidence (3-star) entries were considered. However, references to 2-star entries in other databases necessitated the inclusion of all possible ChEBI compounds. Information about KEGG and International Chemical Identifier (InChI) identifiers were contained in separate files ('chebiId_inchi.tsv' and 'database_accession.tsv'), which were merged with the 'compounds.tsv' file. Discrepancies between ChEBI IDs in the HMDB XML file required manual verification and correction of duplicated IDs before merging the datasets. To extract all **KEGG** compounds, the provided REST API was utilized. Extracted flat files for each metabolite were parsed using regular expression, and KEGG compounds were matched to the dataframe via KEGG IDs provided by HMDB, FooDB, and ChEBI. **VMH** metabolites were extracted using the provided API. VMH compounds were matched using available VMH IDs, as well as HMDB, FooDB, ChEBI identifiers, and compound names. This resulted in the final table comprising all compounds present in the mentioned databases.

Figure 1: MeNu GUIDE is a knowledge graph containing over 25 million triple statements on the relationships between foods, metabolites and diseases. It was created by combining the information from databases with the knowledge stored in ontologies. MeNu GUIDE is available in the Turtle file format, which can easily be queried using SPARQL.

2.2   Ontology Integration

Ontologies were utilized as a systematic framework for organizing and harmonizing the diverse datasets extracted from various databases. For constructing this knowledge graph, ontologies describing nutrition, diseases, and compounds were integrated using the RDFLib library [30]. The nutrition-relevant ontologies included the Food Ontology (FoodOn) [31], Compositional Dietary Nutrition Ontology (CDNO) [32], Food Interactions with Drugs Evidence Ontology (FIDEO) [33] and Food-Biomarker Ontology (FOBI) [34]. For diseases, Human Disease Ontology (DO) [35] and The Drug Ontology (DrOn) [36] were integrated, and for molecules, ChEBI, ChEBI Integrated Role Ontology (CHIRO) [37], Gene Ontology (GO) [38, 39] and Systems Biology Ontology (SBO) [40] were included. Some ontologies already incorporate others, such as DO including ChEBI, which facilitated the integration process. For ontology alignment, precomposed and curated mappings were extracted from BioPortal [41] using its API. The final merged ontologies were saved as a Turtle file.

2.3 Combining Database and Ontology Information

The data composed from the database extraction was then added to the merged ontologies. To match the compounds in the database to the ontologies, unique Internationalized Resource Identifiers (IRIs) were created for each molecule using a generated compound ID. The compound name was added as label and the available properties such as monoisotopic mass, chemical formula, InChI and SMILES were added using the already available predicates from the ChEBI ontology. For a detailed list of the utilized predicates see table S9. Wherever possible, the compound was connected to corresponding classes in the ChEBI ontology.

The 990 foods and dishes present in the downloaded FooDB data were first matched onto the FoodOn ontology using the common and scientific names, if available. This left 250 entities that could not be matched automatically, but had to be mapped manually. All properties used to connect the food nodes are described in table S10. The next step was to create the relation between foods and metabolites through the content entries provided by FooDB. For this purpose only the content entries were selected, that were associated with a source, unit and with a content greater than zero. The content entries were added as a separate node and then connected to the corresponding food and metabolites via their respective IRIs. All properties associated with the content nodes are shown in table S11.

Similarly, the data on biomarkers from MarkerDB was added in the form of separate measurement nodes to serve as the connection between compounds and diseases. To accomplish this, the conditions from the MarkerDB entries first had to be mapped to the DO ontology. A first automated attempt matched 143 diseases, leaving 150 conditions to be matched manually. A comprehensive overview over the relationships connected to the measurement nodes can be found in table S15.

Finally, a new node was created for each reaction. The substrates and products of a reaction were then linked via their MeNu GUIDE compound IRI. Nodes were also introduced for the corresponding enzymes and genes and connected to the reactions they are involved in. A complete list of all implemented relations can be found in tables S12, S13 and S14 for reactions, enzymes and genes respectively.

The final Turtle file containing all triples was then loaded into GraphDB Version 10.6 [42] to create the final knowledge graph. The content of the graph was then explored using SPARQL Protocol and RDF Query Language (SPARQL) queries.

3 Results and Discussion

3.1 Evaluation of the connection between foods and compounds in the MeNu GUIDE knowledge graph

The final graph consists of 25,413,000 triple statements and contains information on 990 foods, 404,823 compounds, 216 conditions, 223,540 content entries and 2,987 biomarker measurements. To obtain an overview over the relations between the graph contents, highly connected nodes were extracted for each node type. Starting with the food nodes, query 1 was used to count the number of associated content entries. These content entries connect foods and compounds and originate from the FooDB database. One example of such a content entry and its connected nodes is visualized in figure 2. Each of these content entries is always associated with a food and compound. It's properties include the content amount and corresponding unit. Additionally, it also mentions the reference, e.g. database or publication, from which this content information derives. For example, for content_1 the reference is "DTU", which corresponds to the Danish food database Frida [43] provided by the Technical University of Denmark (DTU). If available, the content node also includes a minimum and maximum amount measured. The size of a node in the figure depicts its Resource Description Framework (RDF) rank in the graph, which is a measure of interconnectedness.

Figure 2: *Example of a content node in the MeNu GUIDE knowledge graph connecting foods with the compounds they contain.*

Table 1 displays the top 15 food nodes with the highest number of associated content entries. The top three most highly connected food nodes are cattle, cow milk and pork. The "cattle (beef, veal)" node is connected to 15,473 content nodes. However, querying for the compounds associated with these content nodes reveals that multiple content entries are connected to the same compound. This results in only 92 unique compounds associated with the "cattle (beef, veal)" node

via the 15,473 content entries. Modifying the query to extract the number of distinct compounds associated with a food leads to major changes in the top 15 spots, except for milk, potato and corn, which also appear in table 2.

In general, the highly connected food nodes are indicative for important agricultural industries. It is reasonable to assume that this is due to a research focus on foods in the most influential sectors. This is supported by looking at the Annual Survey of Manufactures compiled by the Bureau of the Census of the U.S. Department of Commerce. Data collected in 2021 shows meat processing to be the largest industry group in food and beverage manufacturing with 26.2% of sales, followed by dairy with 12.8% [44].

Table 1: *Foods with highest associated content count.*

Nr	Food	Content Count
1	cattle (beef, veal)	15473
2	milk (cow)	10667
3	domestic pig	10310
4	breakfast cereal	6422
5	baby food	5002
6	cheese	4685
7	soup	3888
8	sausage	3413
9	pastry	2942
10	biscuit	2917
11	common bean	2599
12	potato	2527
13	pizza	2209
14	corn	2158
15	snack bar	1750

Table 2: *Foods with highest associated compound count.*

Nr	Food	Compound Count
1	milk (cow)	1648
2	carrot	231
3	corn	223
4	sweet orange	186
5	green bell pepper	176
6	beer	175
7	red bell pepper	163
8	potato	161
9	apple	160
10	broccoli	154
11	tea	152
12	black tea	152
13	green tea	152
14	parsley	151
15	mandarin orange	148

Similar trends are visible in the European Union, where about 39.8% of the total agricultural output in 2023 came from animals and animal products, mainly from milk and pigs [45]. This is estimated to value around €213.8 billion, representing an enormous market. Naturally, the contributions differ significantly between countries. In the case of France, cereals make up the largest share of the countries agricultural output in 2023 with 15.1%, shortly followed by wine with 14.3%. On the third and fourth spot are milk and cattle with 13.8% and 10.6% [46]. For Germany, milk made up the largest share of the agricultural output with 21.1%, followed by pigs with 12.6% [47]. Cereal is in the fourth spot with 11.9% and wine only makes up 2% of Germany's agricultural output. Potatoes are listed separately from other vegetables in the country specific fact sheets and make up 6.0% and 7.2% of the France's and Germany's agricultural output. All this matches well with the highly connected food nodes and likely explains why they are so well researched.

Another reason for the overrepresentation of milk in the knowledge graph could be that the Wishart research group, which curates the FooDB and HMDB databases, also hosts the Milk Composition Database (MCDB) [48]. It is reasonable to assume that the MCDB shares data with the other databases in the ecosystem, which might lead to an unintended information focus on milk in this knowledge graph. This suggests that the knowledge graph must be enriched with information from additional sources to mitigate biases introduced during the database selection process. These sources could include more databases as well as the extraction of information from publications. Although this may not completely eliminate the research focus bias, it will help alleviate curation bias.

One advantage of the graph structure is that it allows for easy extraction of compounds that two or more foods have in common. To further illustrate this point, query 3 was used to compare the average content of compounds contained in milk and cheese, which is shown in figure 3. Four observations can be made based on this comparison which validate the knowledge graph content. One aspect is the ten-fold increase in sodium content of cheese which is likely due to the salting during the cheesemaking process. This helps to preserve the cheese and prevent the growth of pathogens [49]. Additionally, there is more lactose in milk compared to cheese. This is to be expected and consistent with the manufacturing process as only minimal amounts of lactose are left in the cheese after the metabolization to lactic acid by the starter culture during curd formation and removal of the whey, which retains about 90% of the lactose present [49]. Another observation is that cheese contains a lot more cholesterol than milk, in this case around 17 times more on average. The reason for this is likely the concentration process of the milk solids during cheese-making by removing water. As so much milk fat is retained in the cheese it is logical that this would also lead to an accumulation of fat-soluble vitamins present in milk. This fits with the increase in retinol and beta-carotene in figure 3. However, it is important to note that the levels of beta-carotene and other carotenoids, such as bixin and norbixin, may be elevated in cheese due to their possible use as food colorants [50]. The increase in compounds such as riboflavin, folate and vitamin B12 could be attributed to production by bacteria during the ripening process depending on which strains were used [51].

Figure 3: *Fold change of compounds contained in milk and cheese found in MeNu GUIDE knowledge graph.*

To further explore the graph contents, query 4 was used to extract which compounds have the most connections to content entries and query 5 to count the unique food connections. The results of both queries are depicted in table 3 and table 4. Overall, the compounds contained in both tables are very similar and represent mainly essential elements and vitamins. This is not surprising as their contents in foods are generally well characterized in national nutritional databases.

Both tables contain the elements sodium, potassium and phosphorus. These are important nutrients that patients on hemodialysis are advised to avoid overconsuming in their diet. The knowledge graph can be queried for foods that are very high and low in these minerals. For example, query 6 was used to extract the average potassium content in foods. The top 15 foods containing the most amount of potassium are depicted in table S16. In the first spot is kombu, which is a kelp belonging to the Laminariaceae family and a popular food in East Asia. Species from this family are known to contain high levels of many minerals, especially potassium [52, 53].

Table 3: Compounds with highest associated contents count.

Nr	Compound	Content Count
1	retinol	8184
2	nicotinic acid	7914
3	sodium	5757
4	potassium	5740
5	iron	5473
6	phosphorus	5180
7	zinc	5119
8	riboflavin	5016
9	folic acid	4878
10	calcium	4873
11	copper	4640
12	pyridoxine	4613
13	pantothenic acid	4567
14	thiamine	4555
15	manganese	4305

Table 4: Compounds with highest associated food counts.

Nr	Compound	Food Count
1	iron	752
2	calcium	745
3	phosphorus	735
4	sodium	734
5	potassium	733
6	riboflavin	722
7	copper	719
8	nicotinic acid	717
9	zinc	715
10	thiamine	698
11	manganese	682
12	pyridoxine	673
13	folic acid	650
14	selenium	641
15	retinol	623

On the second spot in table S16 of foods with the most amount of potassium is the tomato. Fresh tomatoes are known to contain high amounts of potassium, with reported values ranging up to 1783 mg/100g [54]. However, the average value reported in table S16 is even higher with 3839.5 mg/100g and a large standard deviation of 1216.9 mg/100g. This equates to a potassium content of 3.8 % in tomato, which is doubtfully high. One advantage of the chosen graph structure is that each content entry also has a reference associated with it, which can be used to further elucidate this issue. Analysis of the information sources shows that the average value of 3839.5 mg/100g for the node "garden tomato (var.)" originates from two entries in the Dr. Duke's Phytochemical and Ethnobotanical Database (Duke's) [55], which is provided by the U.S. Department of Agriculture (USDA). However, due to the data contained in FooDB there is another node in the knowledge graph named "garden tomato", which is connected to content measurements from the DTU and USDA's FoodData Central [56] which average around 796.5 mg/100g and 472.5 mg/100g respectively, around one fifth of the value from Duke's.

The exact reason for the discrepancy between these measurements is speculative. While the cultivar type of the tomato might play a crucial role, potassium fertilization aimed at improving quality and yield could also lead to higher potassium content [57, 58]. However, most measured values mentioned in literature are closer to the FoodData content than to Duke's. Interestingly, both Duke's and FoodData Central are hosted by the USDA, yet they display significantly different values. This discrepancy underscores the importance of the knowledge graph in easily tracing statements back to their sources for comparison. It also highlights the considerable variability in measurement data, revealing that even for common compounds, information can be inconsistent. This demonstrates the need for continued efforts to enhance the accuracy and consistency of nutrient information.

Looking at top 15 foods with the lowest amount of potassium yields mostly processed foods and alcoholic beverages such as vodka, energy drink and marshmallow. These are not the types of food recommended to patients for a healthy diet. In this case the FoodOn ontology and RDF inference from GraphDB prove useful. By specifying that the returned foods should be of the type vegetables and vegetable products (FOODON_03540358) the adjusted query returns the vegetables with the lowest amounts of potassium, shown in table S17. These kinds of queries can serve as inspiration for healthy meal ingredients and, with future expansions of the knowledge graph to include recipes, enable recommendations for low-potassium dishes. Research indicates that maintaining adherence to optimal diets is still problematic [59–61], so providing a tool that simplifies meal planning would be very beneficial.

3.2 Exploration of the connection between compounds and conditions

Another crucial connection in the knowledge graph is through the measurement nodes that link compounds to conditions (see exemplary figure 4). Each measurement entry is associated with a compound and a condition as well as the sample type in which the compound was measured. Properties of the measurement nodes include the measured amount of the compound, the unit, and the reference, typically a PubMed ID for the publication. If available, additional information such as cohort or sex of the measured subjects is provided.

Figure 4: Example of a measurement node in MeNu GUIDE knowledge graph connecting compound and conditions.

Query 8 was used to extract which metabolites have the most connections to measurements, see table 5. Additionally, the number of connections to conditions via those measurement nodes was collected and is shown in table 6. Eleven of the 15 compounds found in table 5 also appear in table 6, which is much more consistent than the content associations. To further explore the connection between compounds and diseases in the knowledge graph, query 7 was used to find all conditions linked to a specific biomarker. Looking at l-lactic acid for example, the compound description already provides a first hint at the diseases it is involved in. The description states that "lactate measurement in critically ill patients has been traditionally used to stratify patients with poor outcomes", which is likely a reason for why it is associated with so many measurements. Looking at the concentrations in different sample types, the knowledge graph contains measurements for urine, blood and cerebrospinal fluid. Many of the diseases found to be associated with l-lactic acid are inborn-errors of metabolism, such as Leigh disease, fructose-1,6-bisphosphatase deficiency and pyruvate decarboxylase deficiency, which the description already mentions. However, the query provides information on other linked diseases not mentioned in the description such as type 2 diabetes mellitus and epilepsy. At present, the reliance on MarkerDB results in relatively limited measurement connections that focus solely on biomarkers. However, literature mining could be utilized to broaden the scope and introduce additional causal relationships.

Table 5: Compounds with most associated measurements.

Nr	Compound	Measurement Count
1	l-lactic acid	47
2	uric acid	33
3	glycine	28
4	3-hydroxybutyric acid	25
5	l-lysine	24
6	homovanillic acid	21
7	l-arginine	21
8	l-histidine	20
9	pipecolic acid	20
10	l-serine	19
11	cortisol	19
12	homocysteine	18
13	pyruvic acid	17
14	creatinine	17
15	hypoxanthine	17

Table 6: Compounds with most associated conditions.

Nr	Compound	Condition Count
1	l-lactic acid	20
2	uric acid	13
3	l-arginine	13
4	glycine	12
5	homovanillic acid	11
6	3-hydroxybutyric acid	11
7	homocysteine	11
8	pyruvic acid	10
9	xanthine	10
10	bilirubin	10
11	l-serine	9
12	l-lysine	9
13	l-alanine	9
14	creatinine	9
15	dopamine	9

Table 7: *Conditions with most associated measurements.*

Nr	Compound	Measurement Count
1	normal	1623
2	obesity	252
3	Alzheimer's disease	82
4	eosinophilic esophagitis	79
5	uremia	48
6	chronic kidney disease	28
7	type 2 diabetes mellitus	25
8	Lewy body dementia	21
9	meningitis	20
10	schizophrenia	19
11	Cirrhosis	19
12	multiple sclerosis	18
13	Parkinson's disease	18
14	maple syrup urine disease	18
15	celiac disease	17

Table 8: *Conditions with most associated biomarkers.*

Nr	Condition	Biomarker Count
1	normal	877
2	obesity	235
3	eosinophilic esophagitis	79
4	Alzheimer's disease	60
5	uremia	47
6	chronic kidney disease	28
7	type 2 diabetes mellitus	21
8	Lewy body dementia	21
9	meningitis	20
10	schizophrenia	18
11	multiple sclerosis	18
12	Parkinson's disease	18
13	Cirrhosis	18
14	celiac disease	17
15	pre-eclampsia	16

To complete the analysis of the knowledge graph connectivity, query 9 was used to extract the conditions that are connected to the most amount of measurements, seen in table 7. For comparison purposes, the conditions with the highest number of biomarkers associated via these measurement nodes were also extracted and are displayed in table 8. The only difference between these two results is that the condition maple syrup urine disease from table 7 is replaced with pre-eclampsia in table 8.

Table 7 contains two diseases with notable diet associations: celiac disease and maple syrup urine disease. In the case of celiac disease, the critical compound, gluten, is unfortunately absent from the knowledge graph, as it is not part of the compounds extracted from the databases. Thus is cannot be connected to any food or condition. This is another clear knowledge gap that could be filled through text mining of relevant publications. However, for maple syrup urine disease, leucine and isoleucine were identified as biomarkers. Individuals with this disease cannot properly catabolize the branched-chain amino acids leucine, isoleucine, and valine due to a genetic mutation.

This association now can serve as the connection to enable the main feature of the knowledge graph, which is the extraction of foods that contain the compounds connected to a specific disease. The knowledge graph facilitates the direct creation of a connection between the condition, its associated biomarkers, and the foods that contain them, as demonstrated by query 10. This relationship can be visualized using the "CONSTRUCT" keyword in the SPARQL query 11 , with figure 5 showing an exemplary visualization for the top three foods with a link display limit of 50.

Figure 5: *Example of condition-compound-food connection in the MeNu GUIDE knowledge graph showing the top three foods in the knowledge graph containing the most amount of l-leucine, which is a biomarker for maple syrup urine disease.*

Each of the top three foods (eggs, soy bean and other soy product) is connected to the l-leucine node via two content nodes that contain reference measurements from the USDA database. The compound node of l-leucine is then connected to maple syrup urine disease via two measurement nodes, measurement_1331 and measurement_1329. The node properties (figure S1 A) reveal that the source of measurement_1331 is a publication from 2002 that measured an average blood concentration of 655 μM l-leucine levels in newborns with maple syrup urine disease [62]. Analogously, the properties of the measurement_1329 node (figure S1 B) point to a publication from 1980 where a l-leucine level of 892 μM was measured in the cerebrospinal fluid of a two year old child [63].

This example demonstrates the capability of the knowledge graph to reveal critical connections and present them in a visually engaging and easily navigable format. Additionally, by providing the source for each connection within the graph researchers are able to quickly locate publications supporting specific data points or relationships. This not only enables the fast extraction of relevant literature on a certain research question, but also easy verification of the contained data. Most notably though, a single query was utilized to identify compounds associated with a specific disease and to pinpoint which foods contain these compounds. Achieving the same result without the knowledge graph would involve laborious manual searches across multiple databases. This functionality underscores a significant advantage of the knowledge graph, made possible through the successful integration of diverse data sources in an ontology-based framework.

4 Conclusion and Outlook

The MeNu GUIDE knowledge graph serves as a proof of concept, demonstrating the feasibility of linking food and disease at a metabolite-specific level to extract relevant information. For instance, the graph structure facilitates the identification of compounds shared by different foods, allowing for the investigation of metabolic changes that occur during food processing, such as cheese-making. Additionally, the knowledge graph enables users to search for foods with particularly high or low levels of specific compounds, aiding in dietary decisions to either boost essential nutrients or avoid excessive consumption of unfavorable substances. Most notably, a single query can reveal compounds associated with a particular disease and locate foods that contain them. This functionality underscores a significant benefit of the knowledge graph, made possible by the successful integration of various data sources.

The MeNu GUIDE knowledge graph establishes a foundation by offering a standardized, ontology-based framework, which can easily be expanded. The current implementation provides a solid starting point for developing increasingly complex models. The analyses revealed that relying solely on database information often results in significant knowledge gaps. For instance, crucial details such as the role of gluten in celiac disease are missing. Therefore, the primary task moving forward will be to enhance the knowledge graph with additional information. This can include data from other databases as well as insights from scientific literature, which harbors a wealth of untapped knowledge. However, accessing this valuable information is challenging due to its fragmented nature, and manually extracting relevant details would be exceedingly time-consuming. Modern technologies, such as neural networks and machine learning algorithms, present a promising solution to this problem [64–67]. Additionally, incorporating recipes into the database would facilitate the development of a recommendation algorithm tailored to specific nutritional needs. Including information on bioavailability during this process would also be highly beneficial.

The second major objective going forward is validation. Once additional content is integrated, the knowledge graph should be made available as a web application for the scientific community. This will allow researchers and experts to identify and report errors for correction. Given the vast scale of the knowledge graph, containing millions of edges even before incorporating data from literature, it is impractical for a single individual alone to validate the content. Therefore, a collaborative approach is essential. Similar to Wikipedia, experts in relevant fields should have the ability to report issues and make corrections, leveraging the collective intelligence of the community.

By expanding the content, facilitating validation, and providing an intuitive framework for exploration, the knowledge graph has the potential to become an invaluable tool for researchers. It allows to explore the currently available knowledge on a topic and discover gaps that might serve as interesting areas for further investigation. Empowering researchers with a comprehensive knowledge graph has the potential to catalyze breakthroughs in understanding the intricate connections between diet and health. As nutrition research evolves, there is a growing need for a tool that aggregates significant findings and makes them readily accessible and searchable. This pilot project has demonstrated that an ontology-based knowledge graph is well-suited to meet this need.

Data Availability Statement

All data used for this project is publicly available. The Turtle file containing the complete knowledge graph can be downloaded from figshare via `https://doi.org/10.6084/m9.figshare.27216054.v1`.

Code Availability Statement

The MeNu GUIDE source code can be accessed on GitHub via `https://github.com/LipiTUM/MeNuGUIDE`.

Author Contributions

JKP conceived the initial idea, supervised the project and secured the funding. VW and JKP planned and conceptualized the work. VW extracted the database information, merged the ontologies and ran all queries. VW and JKP wrote the manuscript. All authors read, reviewed, and accepted the manuscript in its final form.

Acknowledgments

This project was funded by the Bavarian State Ministry of Science and the Arts in the framework of the Bavarian Research Institute for Digital Transformation (bidt; JKP, VW: Junior Research Group LipiTUM).

Conflicts of interest

The authors declare no conflicts of interest.

References

1. San-Cristobal, R., Navas-Carretero, S., *et al.* Contribution of macronutrients to obesity: implications for precision nutrition. *Nature Reviews Endocrinology* **16,** 305–320 (2020).

2. Violi, F., Pastori, D., *et al.* Nutrition, thrombosis, and cardiovascular disease. *Circulation research* **126,** 1415–1442 (2020).

3. Magkos, F., Hjorth, M. F. & Astrup, A. Diet and exercise in the prevention and treatment of type 2 diabetes mellitus. *Nature Reviews Endocrinology* **16,** 545–555 (2020).

4. Younossi, Z. M., Zelber-Sagi, S., *et al.* Lifestyle interventions in nonalcoholic fatty liver disease. *Nature Reviews Gastroenterology & Hepatology* **20,** 708–722 (2023).

5. Park, S. H., Lee, Y.-S., *et al.* Alcoholic liver disease: a new insight into the pathogenesis of liver disease. *Archives of Pharmacal Research* **45,** 447–459 (2022).

6. Mitrev, N., Huang, H., *et al.* Review of exclusive enteral therapy in adult Crohn's disease. *BMJ open gastroenterology* **8,** e000745 (2021).

7. Kerbiriou, C., Dickson, C., *et al.* Treatment of Active Crohn's Disease With Exclusive Enteral Nutrition Diminishes the Immunostimulatory Potential of Fecal Microbial Products. *Inflammatory Bowel Diseases,* izae124 (2024).

8. Almario, C. V., Sharabi, E., *et al.* Prevalence and Burden of Illness of Rome IV Irritable Bowel Syndrome in the United States: Results from a Nationwide Cross-Sectional Study. *Gastroenterology* **165,** 1475–1487 (2023).

9. Oka, P., Parr, H., *et al.* Global prevalence of irritable bowel syndrome according to Rome III or IV criteria: a systematic review and meta-analysis. *The lancet Gastroenterology & hepatology* **5,** 908–917 (2020).

10. Ng, S. C., Shi, H. Y., *et al.* Worldwide incidence and prevalence of inflammatory bowel disease in the 21st century: a systematic review of population-based studies. *The Lancet* **390,** 2769–2778 (2017).

11. Weisman, M. H., Oleg Stens, *et al.* Inflammatory Bowel Disease Prevalence: Surveillance data from the U.S. National Health and Nutrition Examination Survey. *Preventive Medicine Reports* **33,** 102173. ISSN: 2211-3355. https://www.sciencedirect.com/science/article/pii/S2211335523000645 (2023).

12. Lewis, J. D., Parlett, L. E., *et al.* Incidence, prevalence, and racial and ethnic distribution of inflammatory bowel disease in the United States. *Gastroenterology* **165,** 1197–1205 (2023).

13. Buie, M. J., Quan, J., *et al.* Global hospitalization trends for crohn's disease and ulcerative colitis in the 21st century: a systematic review with temporal analyses. *Clinical gastroenterology and hepatology* **21,** 2211–2221 (2023).

14. Fritscher-Ravens, A., Pflaum, T., *et al.* Many patients with irritable bowel syndrome have atypical food allergies not associated with immunoglobulin E. *Gastroenterology* **157,** 109–118 (2019).

15. Dionne, J., Ford, A. C., *et al.* A systematic review and meta-analysis evaluating the efficacy of a gluten-free diet and a low FODMAPs diet in treating symptoms of irritable bowel syndrome. *Official journal of the American College of Gastroenterology| ACG* **113,** 1290–1300 (2018).

16. Nordin, E., Landberg, R., *et al.* Exploration of differential responses to FODMAPs and gluten in people with irritable bowel syndrome-a double-blind randomized cross-over challenge study. *Metabolomics* **20,** 1–12 (2024).

17. Levine, A., Rhodes, J. M., *et al.* Dietary guidance from the international organization for the study of inflammatory bowel diseases. *Clinical Gastroenterology and Hepatology* **18,** 1381–1392 (2020).

18. *FooDB* https://foodb.ca (2024).

19. Rothwell, J. A., Perez-Jimenez, J., *et al.* Phenol-Explorer 3.0: a major update of the Phenol-Explorer database to incorporate data on the effects of food processing on polyphenol content. *Database* **2013,** bat070. ISSN: 1758-0463. eprint: https://academic.oup.com/database/article-pdf/doi/10.1093/database/bat070/11197871/bat070.pdf. https://doi.org/10.1093/database/bat070 (Oct. 2013).

20. Jarvis, A., Gallo-Franco, J., *et al.* Periodic Table of Food Initiative for generating biomolecular knowledge of edible biodiversity. *Nature food* **5,** 189–193 (2024).

21. Santos, A., Colaço, A. R., *et al.* A knowledge graph to interpret clinical proteomics data. *Nature biotechnology* **40,** 692–702 (2022).

22. Swainston, N., Batista-Navarro, R., *et al.* biochem4j: Integrated and extensible biochemical knowledge through graph databases. *PloS one* **12,** e0179130 (2017).

23. Stebbing, J., Phelan, A., *et al.* COVID-19: combining antiviral and anti-inflammatory treatments. *The Lancet Infectious Diseases* **20,** 400–402 (2020).

24. Chandak, P., Huang, K. & Zitnik, M. Building a knowledge graph to enable precision medicine. *Scientific Data* **10,** 67 (2023).

25. Wishart, D. S., Guo, A., *et al.* HMDB 5.0: the human metabolome database for 2022. *Nucleic acids research* **50,** D622–D631 (2022).

26. Wishart, D. S., Bartok, B., *et al.* MarkerDB: an online database of molecular biomarkers. *Nucleic acids research* **49,** D1259–D1267 (2021).

27. Hastings, J., Owen, G., *et al.* ChEBI in 2016: Improved services and an expanding collection of metabolites. *Nucleic acids research* **44,** D1214–D1219 (2016).

28. Kanehisa, M., Furumichi, M., *et al.* KEGG for taxonomy-based analysis of pathways and genomes. *Nucleic acids research* **51,** D587–D592 (2023).

29. Brunk, E., Sahoo, S., *et al.* Recon3D enables a three-dimensional view of gene variation in human metabolism. *Nature biotechnology* **36,** 272–281 (2018).

30. Krech, D., Grimnes, G. A., *et al. RDFLib* version 7.0.0. Aug. 2023. `https://github.com/RDFLib/rdflib`.

31. Dooley, D. M., Griffiths, E. J., *et al.* FoodOn: a harmonized food ontology to increase global food traceability, quality control and data integration. *npj Science of Food* **2,** 23 (2018).

32. Andrés-Hernández, L., Blumberg, K., *et al.* Establishing a common nutritional vocabulary-from food production to diet. *Frontiers in nutrition* **9,** 928837 (2022).

33. Bordea, G., Nikiema, J., *et al. FIDEO: food interactions with drugs evidence ontology* in *11th International Conference on Biomedical Ontologies* (2020).

34. Castellano-Escuder, P., González-Domínguez, R., *et al.* FOBI: an ontology to represent food intake data and associate it with metabolomic data. *Database* **2020,** baaa033 (2020).

35. Schriml, L. M., Mitraka, E., *et al.* Human Disease Ontology 2018 update: classification, content and workflow expansion. *Nucleic acids research* **47,** D955–D962 (2019).

36. Bona, J. P., Brochhausen, M. & Hogan, W. R. Enhancing the drug ontology with semantically-rich representations of National Drug Codes and RxNorm unique concept identifiers. *BMC bioinformatics* **20,** 1–14 (2019).

37. Vasilevsky, N. & Mungall, C. *chiro: CHEBI Integrated Role Ontology* `https://github.com/obophenotype/chiro` (2024).

38. Ashburner, M., Ball, C. A., *et al.* Gene ontology: tool for the unification of biology. *Nature genetics* **25,** 25–29 (2000).

39. Aleksander, S. A., Balhoff, J., *et al.* The gene ontology knowledgebase in 2023. *Genetics* **224,** iyad031 (2023).

40. Sheriff, R., Karr, J. & Nguyen, T. *SBO: Systems Biology Ontology* `https://github.com/EBI-BioModels/SBO` (2024).

41. Whetzel, P. L., Noy, N. F., *et al.* BioPortal: enhanced functionality via new Web services from the National Center for Biomedical Ontology to access and use ontologies in software applications. *Nucleic acids research* **39,** W541–W545 (2011).

42. Ontotext. *GraphDB* version 10.6. 2024. `https://www.ontotext.com/products/graphdb/?ref=menu`.

43. Christensen, T. & Biltoft-Jensen, A. P. *The New version of Danish food composition database FRIDA including a case study on recipe calculation compared to a chemical analysis* in *39th National Nutrient Databank Conference* (2016). `https://frida.fooddata.dk/?lang=en`.

44. USDA. *Food and Beverage Manufacturing* `https://www.ers.usda.gov/topics/food-markets-prices/processing-marketing/food-and-beverage-manufacturing/` (2024).

45. Eurostat. *Performance of the agricultural sector - Value of agricultural output* `https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Performance_of_the_agricultural_sector` (2024).

46. Agridata. *Country Factsheet France - Agricultural output per sector* `https://agridata.ec.europa.eu/extensions/CountryFactsheets/CountryFactsheets.html?memberstate=France` (2024).

47. Agridata. *Country Factsheet Germany - Agricultural output per sector* `https://agridata.ec.europa.eu/extensions/CountryFactsheets/CountryFactsheets.html?memberstate=Germany` (2024).

48. Foroutan, A., Guo, A. C., *et al.* Chemical composition of commercial cow's milk. *Journal of agricultural and food chemistry* **67,** 4897–4914. `https://www.mcdb.ca` (2019).

49. Fox, P. F., Guinee, T. P., *et al. Fundamentals of cheese science* 391–442 (Springer, 2017).

50. Sharma, P., Segat, A., *et al.* Colorants in cheese manufacture: Production, chemistry, interactions, and regulation. *Comprehensive Reviews in Food Science and Food Safety* **19,** 1220–1242 (2020).

51. Ağagündüz, D., Şahin, T. Ö., *et al.* Lactic acid bacteria as pro-technological, bioprotective and health-promoting cultures in the dairy food industry. *Food Bioscience* **47,** 101617 (2022).

52. Funaki, M., Nishizawa, M., *et al.* Mineral composition in the holdfast of three brown algae of the genus Laminaria. *Fisheries science* **67,** 295–300 (2001).

53. Hwang, J.-H., Kim, N.-G., *et al.* Variation in the chemical composition of Saccharina japonica with harvest area and culture period. *Journal of Aquaculture Research & Development* **5,** 1 (2014).

54. Petek, M., Armanda, A. R., *et al.* Potassium content in tomato and tomato products. *Scientific Papers. Series B. Horticulture* (2021).

55. U.S. Department of Agriculture, A. R. S. *Dr. Duke's Phytochemical and Ethnobotanical Databases* `https://phytochem.nal.usda.gov` (2024).

56. U.S. Department of Agriculture Agricultural Research Service, B. H. N. R. C. *FoodData Central* `https://fdc.nal.usda.gov/` (2024).

57. Daoud, B., Pawelzik, E. & Naumann, M. Different potassium fertilization levels influence water-use efficiency, yield, and fruit quality attributes of cocktail tomato—A comparative study of deficient-to-excessive supply. *Scientia Horticulturae* **272,** 109562 (2020).

58. Sonntag, F., Naumann, M., *et al.* Improvement of cocktail tomato yield and consumer-oriented quality traits by potassium fertilization is driven by the cultivar. *Journal of the Science of Food and Agriculture* **99,** 3350–3358 (2019).

59. Luis, D., Zlatkis, K., *et al.* Dietary quality and adherence to dietary recommendations in patients undergoing hemodialysis. *Journal of Renal Nutrition* **26,** 190–195 (2016).

60. Uribarri, J. *An aspirational diet for dialysis patients: Evidence and theory* in *Seminars in dialysis* **31** (2018), 236–243.

61. Sussman-Dabach, E. J., Joshi, S., *et al.* *Preventing potential pitfalls of a liberalized potassium diet in the hemodialysis population* in *Seminars in Dialysis* **37** (2024), 317–325.

62. Deng, C., Shang, C., *et al.* Rapid diagnosis of phenylketonuria and other aminoacidemias by quantitative analysis of amino acids in neonatal blood spots by gas chromatography–mass spectrometry. *Journal of Chromatography B* **775,** 115–120 (2002).

63. Wendel, U, Becker, K, *et al.* Peritoneal dialysis in maple-syrup-urine disease: studies on branched-chain amino and keto acids. *European Journal of Pediatrics* **134,** 57–63 (1980).

64. Popovski, G., Kochev, S., *et al.* FoodIE: A Rule-based Named-entity Recognition Method for Food Information Extraction. *ICPRAM* **12,** 915 (2019).

65. Lee, J., Yoon, W., *et al.* BioBERT: a pre-trained biomedical language representation model for biomedical text mining. *Bioinformatics* **36,** 1234–1240 (2020).

66. Cenikj, G., Seljak, B. K. & Eftimov, T. *FoodChem: A food-chemical relation extraction model* in *2021 IEEE Symposium Series on Computational Intelligence (SSCI)* (2021), 1–8.

67. Cenikj, G., Eftimov, T. & Seljak, B. K. FooDis: A food-disease relation mining pipeline. *Artificial intelligence in medicine* **142,** 102586 (2023).

List of Abbreviations

CDNO Compositional Dietary Nutrition Ontology

ChEBI Chemical Entities of Biological Interest

CHIRO ChEBI Integrated Role Ontology

DO Human Disease Ontology

DrOn The Drug Ontology

DTU Technical University of Denmark

Duke's Dr. Duke's Phytochemical and Ethnobotanical Database

FIDEO Food Interactions with Drugs Evidence Ontology

FOBI Food-Biomarker Ontology

FooDB Food Database

FoodOn Food Ontology

GO Gene Ontology

HMDB Human Metabolome Database

IBD Inflammatory Bowel Disease

IBS Irritable Bowel Syndrome

InChI International Chemical Identifier

IRI Internationalized Resource Identifier

KEGG Kyoto Encyclopedia of Genes and Genomes

MCDB Milk Composition Database

RDF Resource Description Framework

SBO Systems Biology Ontology

SPARQL SPARQL Protocol and RDF Query Language

USDA U.S. Department of Agriculture

VMH Virtual Metabolic Human