Specific Bacterial Taxa and Their Metabolite, DHPS, Linked to Alzheimer's Disease, Parkinson's Disease, and Amyotrophic Lateral Sclerosis.

Christopher, Courtney Jayde (co-first), Department of Chemistry, University of Tennessee, 1420 Circle Drive, Knoxville, TN 37996. ORCiD: 0000-0002-8387-4779

Morgan, Katherine Hope (co-first), College of Nursing, University of Tennessee, 1412 Circle Drive, Knoxville, TN 37996. ORCiD: 0000-0003-3673-5570.

Tolleson, Christopher Mahone, University of Tennessee Medical Center, Division of Neurology, 1924 Alcoa Hwy, Knoxville, TN 37920. ORCiD: 0009-0003-0416-6729

Trudell, Randall, University of Tennessee Medical Center, Division of Neurology, 1924 Alcoa Hwy, Knoxville, TN 37920. ORCiD: 0009-0002-3051-9578

Fernandez-Romero, Roberto, The Pat Summitt Clinic at The University of Tennessee Medical Center, 1932 Alcoa Hwy., Suite C-150, Knoxville, TN 37920. ORCiD:0009-0007-2306-7215

Rice, Lexis, Department of Biochemistry & Cellular and Molecular Biology, University of Tennessee, 1311 Cumberland Ave., Knoxville, TN 37916. ORCiD: 0009-0004-5800-9128

Abiodun, Blessing A, Department of Chemistry, University of Tennessee, 1420 Circle Drive, Knoxville, TN 37996. ORCiD: 0009-0009-3370-6468

Vickery, Zane, Department of Chemistry, University of Tennessee, 1420 Circle Drive, Knoxville, TN 37996. ORCiD: 0009-0003-5595-4194

Jones, Katarina, Department of Chemistry, University of Tennessee, Department of Chemistry, University of Tennessee, 522 Buehler Hall, 1420 Circle Drive, Knoxville, TN 37996. ORCiD: 0000-0003-3769-8599

Woodall, Brittni Morgan, Department of Chemistry, University of Tennessee, 1420 Circle Drive, Knoxville, TN 37996. ORCiD: 0000-0002-2597-7331

Nagy, Christopher; High Throughput Sequencing Facility; University of North Carolina at Chapel Hill, School of Medicine, 1153 Genome Science Building, SOM CB#7102, Chapel Hill, NC 27599 ORCiD: 0009-0003-3628-9360

Mieczkowski, Piotr Andrzej, Department of Genetics, University of North Carolina at Chapel Hill, School of Medicine, 4256 Genome Science Building, SOM CB#7264, Chapel Hill, NC 27599. ORCiD:0000-0003-2418-0096.

Campagna, Shawn R, Department of Chemistry and Biological and Small Molecule Mass Spectrometry Core, University of Tennessee, 611 Buehler Hall, 1420 Circle Drive, Knoxville, TN 37996-1600. ORCiD 0000-0001-6809-3862

Bowen, Gregory; Integrated Genomics Cores; University of North Carolina at Chapel Hill, 4252 Genome Sciences Building, 250 Bell Tower Dr., Chapel Hill, NC 27514. ORCiD:0009-0008-5687-5507

Ellis, Joseph Christopher,* NetEllis, and Department of Medicine at the UT Graduate School of Medicine, Knoxville, TN 37920. ORCiD:0000-0002-2510-977X; email: chris@netellis.com

Neurodegenerative diseases (NDDs) are multifactorial disorders frequently associated

with gut dysbiosis, oxidative stress, and inflammation; however, the pathophysiological

mechanisms remain poorly understood. We investigated bacterial and metabolic dyshomeostasis

in the gut microbiome associated with early disease stages across three NDDs, amyotrophic

lateral sclerosis (ALS), Alzheimer's Disease (AD), Parkinson's Disease (PD), and healthy

controls (HC) and discovered a previously unrecognized link between a microbial-derived

metabolite with an unknown role in human physiology, 2,3-dihydroxypropane-1-sulfonate

(DHPS), and NDDs. DHPS was downregulated in AD, ALS, and PD, while *Eubacterium* and

Desulfovibrio, capable of metabolizing this metabolite,[1-4] were increased in all disease cohorts.

Additionally, select taxa within the Clostridia class had strong negative correlations to DHPS

suggesting a potential role in DHPS metabolism. Hydrogen sulfide is a catabolic product of

DHPS,[1,5] and hydrogen sulfide promotes inflammation,[6-8] oxidative stress,[9] mitochondrial

damage,[10] and gut dysbiosis,[2,11] known hallmarks of NDD. These findings suggest that cryptic

sulfur metabolism via DHPS is a missing link in our current understanding of NDD onset and

progression. To the best of our knowledge, we are the first to provide evidence of a conserved

gut-brain axis linkage of specific bacterial taxa and their metabolism of DHPS shared by three

neurodegenerative diseases.

Main

The social and economic impact of neurodegenerative diseases (NDDs) such as

Alzheimer's (AD), amyotrophic lateral sclerosis (ALS), and Parkinson's (PD) have devasting

impacts on patients and families alike. In aggregate, neurological disorders cause disability,

reduce lifespan, and are the second leading cause of death worldwide.[12] A growing number of

researchers are exploring the gut-brain axis in Alzheimer's disease using the mouse model.[13]

However, the cause of neurodegenerative disease, broadly, and amyotrophic lateral sclerosis,

Alzheimer's Disease, Parkinson's Disease, specifically, remain poorly understood.

26   The gut is in constant system-wide communication with organs throughout the body,[14]

and dyshomeostasis within the metabolome can contribute to intestinal and systemic

inflammation, oxidative stress, and mitochondrial dysfunction which are involved in the

pathology of NDDs.[15] For example, hydrogen sulfide (H_2S) is a key metabolite involved in sulfur

metabolism with a myriad of beneficial physiological functions and is primarily produced by

members of the *Desulfovibrio* genus.[2,16] However, excess H_2S promotes inflammation,[6-8]

oxidative stress,[9] mitochondrial damage,[10] and gut dysbiosis.[2,11] Dyshomeostasis in sulfur

metabolism has been well-documented in PD,[17-19] and the host-microbe regulation of oxidized

sulfur compounds in the metabolome is gaining recognition in health and disease.[20]

35   In this study, we collected fecal samples and investigated the human gut bacterial

microbiome and metabolome of patients from three NDDs and a healthy controls (HC) cohorts

using a multi-omics approach. Sampling occurred early in disease progression, at the first or

second clinical appointment with a neurologist. These results demonstrate a previously

unrecognized association between microbial-derived metabolite DHPS and NDDs. While the

role of DHPS in human health is unknown, we found DHPS to be significantly downregulated in

the stool metabolome of AD, ALS, and PD cohorts compared to controls. In addition, we

observed that specific taxa responsible for DHPS production (*Eubacterium*) and catabolism

(*Desulfovibrio)* were significantly increased in NDDs. For the first time, we have provided data

directly linking DHPS to NDDs. These findings may suggest that DHPS is a missing link in the

current understanding of sulfur dyshomeostasis in the gut associated with NDDs.

Subjects Were in Early Diagnostic Stages

From summer 2021 to spring 2022, 48 participants enrolled, and 45 completed the study.

Of these, 39 submitted stool samples (AD=5, ALS=11, PD=13, and HC=10). Participants were

non-Hispanic Caucasians living within the 21-counties served by the University of Tennessee

Medical Center (See Supplementary Table 1 for additional demographics) in Knoxville,

Tennessee.

Global Metabolomics

Untargeted metabolomics was used to investigate functional alterations in the gut

metabolome associated with NDDs. We identified 164 metabolites based on exact mass and

retention time and evaluated metabolic profile differences between each NDD and HC using

PLS-DA, which revealed that each NDD had a unique gut metabolic profile compared to HC

(Fig. 1). To determine metabolites altered by NDD, metabolites from each pairwise comparison

with a fold change $> |1.5|$ ($p < 0.1$) or VIP score > 1 were used for further analysis

(Supplementary Table 2). These metabolites were also used to identify unique metabolic

signatures for each NDD. This revealed 14, 20, and 9 unique markers for AD, ALS, and PD,

respectively. Moreover, we identified 19 metabolic markers, which were consistently altered

across all three NDDs in this study (Supplementary Table 3).

AD: Based on the PLS-DA model between the AD and HC metabolomes, the overall

metabolic profile of the gut in AD patients was distinctly different from HC as seen by the

clustering and separation of groups (Fig. 1). Metabolites with the highest VIP scores, thereby

contributing most to global metabolome differences, were UMP (2.8), IMP (2.5), and taurine

(2.4). We observed that 29% of identified metabolites in the gut were altered by AD, and 14 of

these metabolites were unique to AD. Notably, the primary catabolic product of vitamin B_6 (4-

pyridoxate), S-ribosyl-L-homocysteine, 3-methylphenylacetic acid, and metabolites involved in

fatty acid and energy metabolism (carnitine, hydroxylysine, FMN) were unique to AD.

Metabolic profile differences in AD demonstrated altered sulfur metabolism, with

downregulation of cysteine, taurine, methionine, homocysteine, and DHPS, while sulfur

containing B vitamins, biotin and pyridoxine, were increased in the AD cohort compared to HC

(Fig. 2).

ALS: Similarly, the ALS cohort also exhibited global gut metabolome differences

compared to HC, as shown in Figure 1. Metabolites contributing most to differences between

cohorts were N-acetylglucosamine phosphate (VIP = 2.6), cholesterol sulfate (VIP = 2.4), 2-

isopropylmalate (VIP = 2.3), and kynurenic acid (VIP = 2.2) (Fig. 2). Overall, 34% of identified

metabolites were significantly altered in the ALS cohort: of these, histadinol and CDP-choline

were increased, while all other metabolites were decreased. Similar to AD, we observed that ALS

caused alterations in tryptophan metabolism, sulfur metabolism, and acetylated amino acids in

the gut (Fig. 2). There were 19 metabolites unique to ALS, including metabolites involved in

tryptophan metabolism (xanthurenic acid, kynurenic acid), vitamin B_6 (pyridoxine), and energy,

fatty acid, and amino acid metabolism (creatine, creatinine, arginine, CDP-choline, and

glycodeoxycholate). Interestingly, creatine metabolism was unique to the ALS cohort, with no

notable alterations in AD or PD cohorts.

PD: The most dramatic alterations in the gut metabolome were found when comparing

the PD to HC cohort with 35% of identified metabolites being altered by PD (Fig. 1).

Phenyllactic acid (2.6), CMP (2.3), 2-isopropylmalate (2.3), sucralose (2.2), and UMP (2.1) had

the highest VIP scores contributing most to the global metabolome differences between the PD

and HC cohorts in the PLS-DA model. Similar to both AD and ALS, PD caused alterations in

sulfur, tryptophan, vitamin B, and acetylated amino acid metabolism. Other notable differences

in the PD gut metabolome include downregulation of taurine, alanine, nicotinate and

nicotinamide, and cysteine and methionine metabolism (Fig 2). Conversely, biotin, sucralose, 3-

methylthiopropionate, and histidinol were upregulated in PD (Fig 2). There were 10 unique

metabolic markers associated with PD including xylose, lysine, xylitol, and N-acetyl-beta-

alanine. The most dramatic alterations in neurotransmitters (5-Hydroxyindoleacetic acid (5-

HIAA), dopamine, 3,4-dihydroxyphenylacetate (DOPAC)) were observed in the gut metabolome

of PD patients.

Neurodegenerative disease metabolic markers

There were 19 metabolites altered by the three NDDs in the study, serving as metabolic

markers for AD, ALS, and PD (Fig. 2). Many of these metabolites were involved in sulfur and

vitamin B metabolism (DHPS, taurine, 3-methylthiopropionate, cholesterol sulfate, sulfolactate,

and biotin). DHPS is a microbial-derived metabolite, and was on average 9x lower ($p < 0.05$) in

the NDD cohorts compared to the HC cohort (Fig. 2). We also observed that DHPS had strong

correlations to acetylated amino acids and metabolites involved in neurotransmitter, bile acid,

and vitamin B metabolism (Fig. 2).

Sequencing (16s rRNA)

The microbial composition of stool samples from NDD and HC patients were

characterized via 16S amplicon sequencing. Differences in alpha diversity (Fig. 3) were only

observed when comparing the PD and HC cohorts, with PD having higher alpha diversity (p =

0.02). Analysis of OTUs revealed phylum level differences in the abundance of Firmicutes in PD

(53%) compared to HC (67%) stool, while Firmicutes displayed minimal differences in ALS and

AD cohorts (Fig. 3). Phylum Actinobacteria was unchanged in AD but was increased in PD and

ALS compared to HC patients. Differences within the Firmicutes phylum in PD could broadly be

attributed to genus *Eubacterium*. There were six taxa significantly altered across all NDDs

compared to HCs: *Desulfovibrio, Eubacterium, Akkermansia, Paludicola, Sellimonas, and*

Enterococcus. (Fig. 3) All of these taxa were more abundant in the NDD cohorts than HC, except

for *Enterococcus* which was higher in HC.

To assess the relationship between gut microbial composition and function in NDDs, we

identified OTUs correlated with metabolites that were significantly different in NDDs.

Microbial-derived metabolite DHPS was consistently and significantly altered in NDDs and

there were 33, 43, and 15 OTUs significantly correlated with DHPS in PD, AD, and ALS,

respectively, all having negative correlations. Of these, eight OTUs were correlated with DHPS

across PD, AD, and ALS, and all were part of the Firmicutes phylum (class *Clostridia*), except

for one belonging to the Bacteroidetes phylum (Fig. 2).

127   **Discussion**

Cryptic sulfur metabolism in NDDs

All detected metabolites in the transsulfuration pathway were decreased in NDD cohorts,

except for cystathionine. Rapid conversion of homocysteine to cystathionine could suggest

increased fluxes in sulfur metabolism as previous work using longitudinal metabolome data and

constraint-based modeling of gut microbial communities has shown increased flux in the

transsulfuration pathway in PD.[18] This pathway generates glutathione and H_2S, which impact

redox metabolism. Increased fluxes through sulfur metabolism may be indicative of excess H_2S,

which at low concentrations, is neuroprotective and cytoprotective. However, at high levels, H_2S

induces systemic inflammation,[6-8] oxidative stress,[9] mitochondrial damage,[10] and gut

dysbiosis.[2,11] Our findings support the current hypothesis that the increased burden of oxidative

stress causes mitochondrial dysfunction leading to neuroinflammation.

In addition to metabolites in the transsulfuration pathway, other sulfur-containing

metabolites, DHPS and taurine, were significantly altered across NDDs. We found this to be

intriguing as oxidized sulfur-containing small molecules represent an underappreciated class of

unique, biologically relevant metabolites in host-microbiome interactions.[20] It has been

discovered recently that these molecules play a significant role in cellular signaling and impact

disease-related phenotypes thereby bringing oxidized sulfur metabolites to the forefront of

enhancing our understanding of the pathophysiological impact of host-microbiome

interactions.[20-24] Since the role of DHPS in NDDs is unknown, we identified metabolites and

OTUs significantly correlated with DHPS. Five metabolites (taurine, vanillin, histidinol, *N*-

acetylornithine, and sucrose/trehalose), and eight OTUs were significantly (p<0.01) correlated

with DHPS in each NDD cohort. All of the significantly correlated OTUs belonged to phylum

Firmicutes (orders Clostridia and Oscillospirales) except for one, which belonged to the phylum

Bacteroidetes (genus *Odoribacter*). All of these microbes have been directly linked to

metabolizing dietary sulfur to H_2S.[4,25] Together, the metabolites and OTUs correlated with

DHPS suggest that DHPS is involved in cryptic sulfur host-microbiome metabolism in NDDs.

DHPS microbial metabolism in NDDs

Frommeyer and colleagues annotated genes required for a novel bacterial transaldolase

pathway by which select gut microbes (*Enterococcus*, *Clostridium*, *Desulfovibrio*, and

Eubacterium strains) can ferment dietary sulfoquinovose (SQ) to H_2S with DHPS as a transient

intermediate.[2] SQ is obtained through the intake leafy greens and vegetables containing the lipid

sulfoquinovosyl diacylglycerol (SQDG), which is a highly abundant sulfur containing metabolite

common in photosynthetic plants and cyanobacteria. In addition, Hanson and colleagues[2]

reported that DHPS contributes to intestinal H_2S production and that *Eubacterium rectale (E.*

rectale) is the primary DHPS producer, while *Desulfovibrio* and *Bilophila are* the primary DHPS

consumers, in fecal microcosms from human vegetarians. They identified genes necessary for

microbial catabolism of diet-acquired SQ to H_2S and analyzed the expression of these genes

across stool metatranscriptomes of 123 healthy, 28 ulcerative colitis, and 50 Crohn's Disease

patients to identify the overall impact on health.[2] However, they found no significant differences

in expression of these pathways between cohorts with bowel disease and healthy individuals.[2]

Eubacterium rectale plays a significant role in promoting inflammation and damaging

tissue in the gut.[26] We found that the genus *Eubacterium* was increased in PD, ALS, and AD,

and it was one of the eight OTUs strongly correlated with DHPS across all three diseases (Fig.

3). Additionally, sulfate-reducing bacteria (SRB), such as *Desulfovibrio,* generate H_2S, and have

previously been implicated in the development and severity of PD.[17] Here, we observed

Desulfovibrio were significantly altered in PD, ALS, and AD and were more abundant in patients

with NDDs (Fig. 3). Additionally, taurine was strongly correlated with DHPS and significantly

decreased in NDDs and can also be obtained through dietary sources and degraded by

Desulfovibrio and Bilophilia to acetyl-CoA and H_2S. Other studies have also implicated

Bilophilia wadsworthia in PD altering microbial sulfur metabolism, yet the etiology remains

elusive.[18,27-29] Disrupted sulfur metabolism in NDDs could result in increased flux through

DHPS and taurine, promoting overgrowth of SRB and excess H_2S. This may explain why DHPS

producing and consuming microbes are increased in NDDs, while DHPS is decreased.

While we do not have dietary information from participants, it has become fairly

accepted that the gut microbiome composition and function in NDDs are influenced by dietary

habits and nutrition.[30] Since diet-acquired SQDG from leafy greens can be bioconverted to

DHPS by select microbes in the gut, we hypothesize that diet corresponds to DHPS and sulfur

homeostasis in NDDs.

Previous studies have laid the foundation for investigating DHPS in human metabolism

by elucidating the biochemical mechanism of DHPS production and degradation in the human

gut and directly linking dietary sulfonates to H_2S production;[2-4] however, the data did not

demonstrate an association between DHPS, or the genes involved in DHPS metabolism, and

human health outcomes. Recently, Zaparte, Christopher and colleagues[31] provided the first

report of direct DHPS detection in human stool samples and highlighted a compelling connection

between DHPS and gut metabolic dysregulation in people who use e-cigarette and/or

combustible tobacco/marijuana smoke. They observed that DHPS was significantly higher (20x)

in the control cohort compared to the smoking cohorts and noted strong correlations between

DHPS and vitamin B, acetylated amino acids, bile acids, and cholesterol metabolism. This is

consistent with the DHPS findings outlined herein, demonstrating a conserved trend across

independent datasets that DHPS abundance in human stool is associated with overall health

status, supporting the proposal of DHPS as a missing link in the pathophysiology of

inflammation as it may be a key regulator of inflammation and overall human health.

Conclusion

In this study, we investigated bacterial and metabolic dyshomeostasis in the gut

associated with AD, ALS, and PD using an integrative cross-omics approach. Herein, we present

the first linkage of DHPS, a microbial-derived metabolite, to NDDs, and suggest that it

contributes to dyshomeostasis in sulfur metabolism, since DHPS consuming and degrading

microbes are more abundant in NDDs while DHPS is decreased (Fig. 4). We hypothesize that the

decrease in DHPS in NDDs may be a result of increased flux through the recently identified

transaldolase pathway in which dietary sulfonates are metabolized to DHPS and H_2S by select

gut microbes, including *Eubacterium* and *Desulfovibrio*. Excess H_2S, resulting from increased

flux through DHPS, contributes to mitochondrial dysfunction, inflammation, oxidative stress,

and gut dysbiosis in NDDs. These findings suggest that cryptic sulfur metabolism via DHPS is a

missing link in our current understanding of NDD onset and progression.

Limitations

This study lays the foundation for future discoveries on the role of specific bacterial taxa

and their metabolite DHPS however, there are some notable limitations. First, all patients were

sampled in the same geographic location of Eastern Tennessee and the surrounding Appalachian

Mountains. In addition, all the patients were Caucasian. Combined, this limits the

generalizability of the conclusions. Finally, while this study describes a remarkably well

conserved and novel association across different neurodegenerative diseases, this study is limited

by not being able to define these findings as causative given the exploratory nature of untargeted

metabolomics.

Main References

1   Burrichter, A. *et al.* Anaerobic Degradation of the Plant Sugar Sulfoquinovose

Concomitant With H(2)S Production: Escherichia coli K-12 and Desulfovibrio sp. Strain

DF1 as Co-culture Model. *Front Microbiol* **9**, 2792 (2018).

https://doi.org/10.3389/fmicb.2018.02792

2   Hanson, B. T. *et al.* Sulfoquinovose is a select nutrient of prominent bacteria and a source

of hydrogen sulfide in the human gut. *ISME J* **15**, 2779-2791 (2021).

https://doi.org/10.1038/s41396-021-00968-0

3   Liu, J. *et al.* Two radical-dependent mechanisms for anaerobic degradation of the

globally abundant organosulfur compound dihydroxypropanesulfonate. *Proc Natl Acad*

Sci U S A **117**, 15599-15608 (2020). https://doi.org/10.1073/pnas.2003434117

4   Frommeyer, B. *et al.* Environmental and Intestinal Phylum Firmicutes Bacteria

Metabolize the Plant Sugar Sulfoquinovose via a 6-Deoxy-6-sulfofructose Transaldolase

Pathway. *iScience* **23**, 101510 (2020). https://doi.org/10.1016/j.isci.2020.101510

5   Denger, K. *et al.* Sulphoglycolysis in Escherichia coli K-12 closes a gap in the

biogeochemical sulphur cycle. *Nature* **507**, 114-117 (2014).

https://doi.org/10.1038/nature12947

6   Szabo, C. & Papapetropoulos, A. International union of basic and clinical pharmacology.

CII: pharmacological modulation of H2S levels: H2S donors and H2S biosynthesis

inhibitors. *Pharmacological Reviews* **69**, 497-564 (2017).

7   Li, L. *et al.* Hydrogen sulfide is a novel mediator of lipopolysaccharide-induced

inflammation in the mouse. *FASEB J* **19**, 1196-1198 (2005).

https://doi.org/10.1096/fj.04-3583fje

8   Dilek, N., Papapetropoulos, A., Toliver-Kinsky, T. & Szabo, C. Hydrogen sulfide: An

endogenous regulator of the immune system. *Pharmacol Res* **161**, 105119 (2020).

https://doi.org/10.1016/j.phrs.2020.105119

9   Olson, K. R. & Straub, K. D. The Role of Hydrogen Sulfide in Evolution and the

Evolution of Hydrogen Sulfide in Metabolism and Signaling. *Physiology (Bethesda)* **31**,

60-72 (2016). https://doi.org/10.1152/physiol.00024.2015

10   Libiad, M. *et al.* Hydrogen sulfide perturbs mitochondrial bioenergetics and triggers

metabolic reprogramming in colon cells. *J Biol Chem* **294**, 12077-12090 (2019).

https://doi.org/10.1074/jbc.RA119.009442

11   Ijssennagger, N. *et al.* Gut microbiota facilitates dietary heme-induced epithelial

hyperproliferation by opening the mucus barrier in colon. *Proc Natl Acad Sci U S A* **112**,

10038-10043 (2015). https://doi.org/10.1073/pnas.1507645112

12   Global, regional, and national burden of neurological disorders, 1990-2016: a systematic

analysis for the Global Burden of Disease Study 2016. *Lancet Neurol* **18**, 459-480

(2019). https://doi.org/10.1016/s1474-4422(18)30499-x

13   Sun, J. *et al.* Fecal microbiota transplantation alleviated Alzheimer's disease-like

pathogenesis in APP/PS1 transgenic mice. *Translational Psychiatry* **9**, 189 (2019).

https://doi.org/10.1038/s41398-019-0525-3

14   Ahlawat, S., Asha & Sharma, K. Gut–organ axis: a microbial outreach and networking.

Letters in Applied Microbiology **72**, 636-668 (2021).

15   Markovinovic, A., Greig, J., Martín-Guerrero, S. M., Salam, S. & Paillusson, S.

Endoplasmic reticulum-mitochondria signaling in neurons and neurodegenerative

diseases. *J Cell Sci* **135** (2022). https://doi.org/10.1242/jcs.248534

16   Carbonero, F., Benefiel, A., Alizadeh-Ghamsari, A. & Gaskins, H. R. Microbial

pathways in colonic sulfur metabolism and links with health and disease. *Frontiers in*

Physiology **3** (2012). https://doi.org/10.3389/fphys.2012.00448

17   Murros, K. E., Huynh, V. A., Takala, T. M. & Saris, P. E. J. Desulfovibrio Bacteria Are

Associated With Parkinson's Disease. *Front Cell Infect Microbiol* **11**, 652617 (2021).

https://doi.org/10.3389/fcimb.2021.652617

18   Hertel, J. *et al.* Integrated Analyses of Microbiome and Longitudinal Metabolome Data

Reveal Microbial-Host Interactions on Sulfur Metabolism in Parkinson's Disease. *Cell*

Rep **29**, 1767-1777.e1768 (2019). https://doi.org/10.1016/j.celrep.2019.10.035

19   Murros, K. E. Sulfate reducing gut bacteria and Parkinson's disease. *Eur J Neurol* **28**, e21

(2021). https://doi.org/10.1111/ene.14626

20   D'Agostino, G. D., Chaudhari, S. N. & Devlin, A. S. Host–microbiome orchestration of

the sulfated metabolome. *Nature Chemical Biology* **20**, 410-421 (2024).

https://doi.org/10.1038/s41589-023-01526-9

21   Sakurai, T. *et al.* Cholesterol sulfate is a DOCK2 inhibitor that mediates tissue-specific

immune evasion in the eye. *Science Signaling* **11**, eaao4874 (2018).

22   Wang, F., Beck-García, K., Zorzin, C., Schamel, W. W. & Davis, M. M. Inhibition of T

cell receptor signaling by cholesterol sulfate, a naturally occurring derivative of

membrane cholesterol. *Nature Immunology* **17**, 844-850 (2016).

23   Lobel, L., Cao, Y. G., Fenn, K., Glickman, J. N. & Garrett, W. S. Diet posttranslationally

modifies the mouse gut microbial proteome to modulate renal function. *Science* **369**,

1518-1524 (2020).

24   Stewart Campbell, A. *et al.* Safety and target engagement of an oral small-molecule

sequestrant in adolescents with autism spectrum disorder: an open-label phase 1b/2a trial.

Nature Medicine **28**, 528-534 (2022).

25   Nguyen, L. H. *et al.* Association Between Sulfur-Metabolizing Bacterial Communities in

Stool and Risk of Distal Colorectal Cancer in Men. *Gastroenterology* **158**, 1313-1325

(2020). https://doi.org/10.1053/j.gastro.2019.12.029

26   Wang, Y. *et al.* Eubacterium rectale contributes to colorectal cancer initiation via

promoting colitis. *Gut Pathogens* **13**, 2 (2021). https://doi.org/10.1186/s13099-020-

00396-z

27   Baldini, F. *et al.* Parkinson's disease-associated alterations of the gut microbiome predict

disease-relevant changes in metabolic functions. *BMC Biol* **18**, 62 (2020).

https://doi.org/10.1186/s12915-020-00775-7

28   Jo, S. *et al.* Oral and gut dysbiosis leads to functional alterations in Parkinson's disease.

NPJ Parkinson's Disease **8**, 87 (2022). https://doi.org/10.1038/s41531-022-00351-6

29   Murros, K. E. Hydrogen Sulfide Produced by Gut Bacteria May Induce Parkinson's

Disease. *Cells* **11** (2022). https://doi.org/10.3390/cells11060978

30   Kwon, D. *et al.* Diet and the gut microbiome in patients with Parkinson's disease. *NPJ*

Parkinson's Disease **10**, 89 (2024). https://doi.org/10.1038/s41531-024-00681-7

31   Zaparte, A. *et al.* Effects of E-Cigarettes on the Lung and Systemic Metabolome in

People with HIV. *Metabolites* **14**, 434 (2024). https://doi.org/10.3390/metabo14080434

Figure 1. Partial least squares discriminant analysis (PLS-DA) comparing metabolomes

associated with NDD and HC. Individual participants are shown with respective confidence

intervals; HC samples are yellow; AD samples are blue; PD samples are purple; ALS samples are

shown in red. D-F) Variable importance in projection (VIP) scores for each corresponding PLS-

DA model top show the top 25 metabolites contributing most to the differences in metabolic

profiles.

Figure 2. There were 19 metabolic markers for all NDD in this study, as well as unique markers

for each disease. DHPS was one of the 19 markers for NDD as was significantly decreased in

NDD cohorts. DHPS was strongly correlated to acetylated amino acids and taurine. A-C)

volcano plots showing metabolites significantly different between cohorts. The x-axis is the $\log_2$

foldchange and y-axis is the -log p value. The following cutoffs were used: fold change > |1.5| (p

< 0.1). D). Venn diagram showing unique and shared metabolic markers. These metabolites had

fold change > |1.5| (p < 0.1) or VIP score >1. E). Metabolites strongly correlated with DHPS in

PD. F). Eight OTUs were significantly correlated with DHPS across all three NDDs.

OTU136 (d__Bacteria;p__Firmicutes;c__Clostridia;__;__;__)

OUT127 (d__Bacteria;p__Firmicutes;c__Clostridia;o__Clostridia;f__Hungateiclostridiaceae;__)

OTU121

(d__Bacteria;p__Firmicutes;c__Clostridia;o__Oscillospirales;f__Oscillospiraceae;g__Oscillospir

a)

OTU091 (d__Bacteria;p__Firmicutes;c__Clostridia;o__Oscillospirales;f__UCG-010;g__UCG-

010)

OTU045

(d__Bacteria;p__Firmicutes;c__Clostridia;o__Clostridia_vadinBB60_group;f__Clostridia_vadin

BB60_group;g__Clostridia_vadinBB60_group)

OTU060

(d__Bacteria;p__Bacteroidota;c__Bacteroidia;o__Bacteroidales;f__Marinifilaceae;g__Odoribact

er)

OTU050 (d__Bacteria;p__Firmicutes;c__Clostridia;o__Oscillospirales;f__Oscillospiraceae;__)

OTU038

(d__Bacteria;p__Firmicutes;c__Clostridia;o__Oscillospirales;f__[Eubacterium]_coprostanoligen

es_group;g__[Eubacterium]_coprostanoligenes_group)

Figure 3. Gut microbial composition is altered in NDDs A) Phyla level differences in microbial

composition, B) Alpha diversity was significantly greater in PD than HC cohort. There were no

significant differences in the AD and ALS cohort compared to HC, C) log$_2$ fold change of taxa

significantly altered in across AD, ALS, and PD cohorts. Orange represents higher abundance in

disease compared to HC while blue represents lower abundance in disease compared to HC.

Figure 4. DHPS can be obtained through dietary consumption of leafy greens and degraded to H_2S. The abundance of DHPS producing and consuming microbes are increased in NDD while DHPS is decreased in NDD suggesting the rapid degradation of DHPS to yield H_2S. Other microbes (*Enterococcus* and *Clostridium*) can ferment dietary sulfoquinovose (SQ) to H_2S with DHPS as a transient intermediate, as well. Figure was created using biorender.com and adapted from Hanson et al. [2]

Methods

Study Population

This study was approved by the institutional review board of the University of Tennessee Health

Science Center for the protection of human subjects. Participants were recruited during their

first or second visit to a movement disorder specialist for ALS, AD, or PD from the 21-county

regional service area of the University of Tennessee Medical Center, in Knoxville, Tennessee.

Participation was voluntary and only those who were willing and able to complete the study

activities (with a caregiver if needed) were included. Participants were excluded if they had

received antibiotics in the previous six months. Further inclusion criteria for each cohort were as

follows: 1) For AD: 65 years of age or older, met all clinical criteria for mild-to-moderate AD

according to National Institute on Aging-Alzheimer's Association criteria,[1] were otherwise in

good health based on medical history and clinical examination; 2) ALS: age of 30 years or older

at enrollment, and met the El Escorial Criteria[2] for definite, probable, or possible ALS. 3) PD:

age 50 to 85 years at enrollment, diagnosed within the last 5 years, absence of PD medications

with the exception of MAO-B inhibitors, absence of dyskinesia or motor fluctuations, modified

Hoehn and Yahr Stage[3] <2.5 (symptoms ranging from unilateral involvement only [Stage 1] to

mild bilateral involvement with recovery on a pull test [Stage 2.5]); and 4) HC: age 30 or older,

were age matched (+/- 4 years) and sex-matched, in good health based on medical history and

clinical examination. HC age 65 years or older completed the Neuropsychological Battery from

the National Alzheimer's Coordinating Center[4] and the Cognivue® computerized cognitive

assessment prior to enrollment to demonstrate normal cognition.

Sample Collection, Preservation, DNA Extraction

Samples were collected at home by participants within one week of enrollment. Participants were

instructed to place stool from the same specimen into two vials as follows: 1) the DNA

Genotek® gut microbiome DNA collection kit and 2) a 50-mL sterile tube, which was

immediately frozen at home at -20 C.   Stool samples were delivered frozen on ice by same-day

courier service, to the laboratory, where they were aliquoted and stored at -80 C. Nucleic acid

extraction was conducted in batches, using Zymo *Quick*-DNA Fecal/soil Microbe Kit, and

extractions were frozen at -80 C, then shipped on dry ice to the University of North Carolina

(UNC) Chapel Hill High Throughput Sequencing Facility (HTSF) for 16s rRNA/DNA

sequencing.

Metabolomics

Metabolite extractions

All samples were extracted and analyzed at the Biological and Small Molecule Mass

Spectrometry Core at the University of Tennessee Knoxville (RRID: SCR_021368). Stool

samples were kept at 4℃, pre-weighed and homogenized. The exact mass of stool used for

sample was recorded and used for normalization. Roughly 50 mg of stool were aliquoted and

water-soluble metabolites were extracted using an acidic acetonitrile extraction procedure

adapted from Rabinowitz and Kimball[5] using 1.5 mL of 4:4:2 acetonitrile: methanol: water with

0.1 M formic acid. [6,7] All solvents were HPLC grade. The supernatant collected from each

sample were dried under nitrogen, then 300 µL of LC-MS grade water was added to each sample

before mass spectral analysis.

UHPLC-HRMS

A previously described ultra high-performance liquid chromatography high resolution mass spectrometry (UHPLC-HRMS) method was used for untargeted metabolomics analysis[8] using an UltiMate 3000 RS autosampler (Dionex, Sunnyvale, CA, USA), Synergi 2.6 µm Hydro RP column (100 mm $\times$ 2.1 mm, 100 Å; Phenomenex, Torrance, CA, USA), an UltiMate 3000 pump (Dionex), and Exactive Plus Orbitrap mass spectrometer (Thermo Fisher Scientific, Waltham, MA, United States). A previously described 25 min gradient elution, reverse phase ion-paring method with a water:methanol solvent system, and a tributylamine ion pairing reagent was used for chromatographic separation.[9] Metabolites were ionized via negative mode electrospray ionization (ESI) prior to full scan mass spectral analysis as previously described.[8]

Metabolomics data processing

Raw mass spectral files were converted to mzML files using a package from ProteoWizard, msConverter.[10] All mzML files were imported into an open-source software, metabolomics analysis and visualization engine (El-MAVEN) where metabolites were manually identified using an in-house library based on exact mass ($\pm$5 ppm) and retention time ($\pm$2 min).[6,11,12] Metabolite peaks were integrated and raw peak intensities for identified metabolites were exported from El-MAVEN to a csv file. Prior to statistical analysis, raw spectral data were normalized by mass of sample used for extraction.

Statistical Analysis

The normalized data were imported into MetaboAnalyst 5.0 and were filtered via interquartile range (IQR), log transformed, and Pareto scaled [13-15] Partial least squares discriminant analysis (PLS-DA) were performed in MetaboAnalyst 5.0 where variable importance in projection (VIP) scores were assigned to each metabolite to indicate the importance of each metabolite in

contributing to the separation between experimental groups. VIP scores > 1 indicate that a

metabolite significantly contributes to the separation of groups in the PLS-DA model. Venn

diagrams were constructed using metabolites with VIP scores > 1 from pairwise group

comparisons. MetaboAnalyst 5.0 was also used to generate volcano plots to visualize metabolites

significantly different between groups.

16s rRNA sequencing

In this study, we implemented a detailed two-step protocol for 16S rRNA gene amplification,[16]

prioritizing high precision and reproducibility across numerous samples. The initial PCR reaction

was meticulously configured, utilizing specific volumes of Kapa Enhancer, Kapa Buffer A

(KAPA Robust 2G - Roche), forward and reverse Frame Shift Molecular Tags (FSMT) primers

(Supplementary Table 2), and other necessary components, culminating in a total reaction

volume of 50 µL. The FSMT primers, comprising six frames for both 338F and 806R types,

were carefully mixed. To streamline the handling of multiple samples, a 'ready-mix' comprising

essential Kapa components was prepared. This strategy was instrumental in reducing batch

effects, particularly when utilizing reagents from different Kapa kits. The ready-mix was then

allocated into aliquots each adequate for a 96-well plate. The first PCR cycle's conditions were

meticulously programmed to include specific durations and steps for denaturation, annealing,

and extension, set at 95°C for one minute, followed by 10 cycles of 95°C for 15 seconds, 50°C

for 30 seconds, and 72°C for 30 seconds, then 72°C for one minute, and a final indefinite hold at

4°C. Post-PCR, bead purification was performed to selectively exclude DNA fragments shorter

than 300 bp, employing a 0.68:1 bead-to-DNA ratio. The second PCR phase (indexing) followed

a similar methodology but with modifications in the reaction components, such as the Kapa HiFi

Readymix (KAPA HiFi -Roche) and specific Adaptor_primer1 and indexing primers. The

cycling conditions for this phase were set at 95°C for one minute, followed by 22 cycles of 95°C

for 15 seconds, 60°C for 30 seconds, and 72°C for 30 seconds, concluding with 75°C for one

minute and an indefinite hold at 4°C. Finally, for 16S sequencing, we used a series of custom-

designed primers, chosen for their specificity and amplification efficiency. The sequences of

these primers, along with the custom sequencing primer (NextF_Read1_seq), are

comprehensively detailed in the supplementary material. Prepared libraries were pooled and

sequenced on MiSeq Pair End 2x300 v2 flowcell (Supplementary Table 2).

16S data processing and analysis

16S amplicon sequencing was performed using the moving pictures documentation in QIIME2.[17]

Briefly, sequences were imported into QIIME2 (version 2023.2). Paired end sequences were

denoised with DADA2. Trimming values such as p-trim-left was set to 20 and p-trim-len was set

to 220 and sampling depth was set at 120,000 sequences. Finally, an OTU table was produced

for downstream analysis. Traditional microbial ecology analysis such as alpha diversity and beta

diversity were also performed. In addition, taxonomic classification of the reads was performed

in the QIIME2 using Silva 138 99% OTUs full-length sequences.

Relative abundances from operational taxonomic units (OTUs) were analyzed on the phylum,

class, order, family, and genus levels. QIIME taxonomy labels with corresponding abundances

were uploaded to MicrobiomeAnalyst, and the data were scaled by total sum prior to statistical

and meta-analysis of microbiome data.[18] Alpha diversity was accessed at the genus level by

Chao, Shannon, and Simpson indexes with the Mann–Whitney test used to determine if alpha

diversity differed across cohorts. Stacked bar graphs were used to visualize OTU relative

abundances across cohorts. Pairwise comparisons between cohorts were made using the

statistical method EdgeR. OTU correlations were analyzed using either Pearson's r or Spearman's

rho correlation coefficients.

Data Availability Statement

Data will be made available on NCBI via SRA

Metabolomics data files are accessible at MetaboLights (accession No. MTBLS10511).

Methods References

1   Jack, C. R., Jr. *et al.* Introduction to the recommendations from the National Institute on

Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's

disease. *Alzheimers Dement* **7**, 257-262 (2011). https://doi.org:10.1016/j.jalz.2011.03.004

2   Brooks, B. R. El Escorial World Federation of Neurology criteria for the diagnosis of

amyotrophic lateral sclerosis. Subcommittee on Motor Neuron Diseases/Amyotrophic

Lateral Sclerosis of the World Federation of Neurology Research Group on

Neuromuscular Diseases and the El Escorial "Clinical limits of amyotrophic lateral

sclerosis" workshop contributors. *J Neurol Sci* **124 Suppl**, 96-107 (1994).

https://doi.org:10.1016/0022-510x(94)90191-0

3   Goetz, C. G. *et al.* Movement Disorder Society Task Force report on the Hoehn and Yahr

staging scale: Status and recommendations The Movement Disorder Society Task Force

on rating scales for Parkinson's disease. *Movement Disorders* **19**, 1020-1028 (2004).

https://doi.org:https://doi.org/10.1002/mds.20213

4   Besser, L. *et al.* Version 3 of the National Alzheimer's Coordinating Center's Uniform

Data Set. *Alzheimer Dis Assoc Disord* **32**, 351-358 (2018).

https://doi.org:10.1097/wad.0000000000000279

5   Joshua D. Rabinowitz, E. K. Acidic acetonitrile for cellular metabolome extraction from

Escherichia coli. *Anal. Chem* **79**, 6167-6173 (2007).

6   Byerley, L. O. *et al.* Gut Microbiome and Metabolome Variations in Self-Identified

Muscle Builders Who Report Using Protein Supplements. *Nutrients* **14**, 533 (2022).

7   Jones, K. A. *et al.* Cross-Omics Analysis of Fenugreek Supplementation Reveals

492   Beneficial Effects Are Caused by Gut Microbiome Changes Not Mammalian Host

493   Physiology. *International Journal of Molecular Sciences* **23**, 3654 (2022).

494   8   Lu, W. *et al.* Metabolomic analysis via reversed-phase ion-pairing liquid chromatography

495   coupled to a standalone orbitrap mass spectrometer. *Anal Chem* **82**, 3212-3221 (2010).

496   https://doi.org:10.1021/ac902837x

497   9   Bazurto, J. V., Dearth, S. P., Tague, E. D., Campagna, S. R. & Downs, D. M. Untargeted

498   metabolomics confirms and extends the understanding of the impact of aminoimidazole

499   carboxamide ribotide (AICAR) in the metabolic network of Salmonella enterica. *Microb*

500   *Cell* **5**, 74-87 (2017). https://doi.org:10.15698/mic2018.02.613

501   10   Martens, L. *et al.* mzML--a community standard for mass spectrometry data. *Mol Cell*

502   *Proteomics* **10**, R110.000133-R000110.000133 (2011).

503   https://doi.org:10.1074/mcp.R110.000133

504   11   Clasquin, M. F., Melamud, E. & Rabinowitz, J. D. LC-MS data processing with

505   MAVEN: a metabolomic analysis and visualization engine. *Curr Protoc Bioinformatics*

506   **Chapter 14**, Unit14.11 (2012). https://doi.org:10.1002/0471250953.bi1411s37

507   12   Melamud, E., Vastag, L. & Rabinowitz, J. D. Metabolomic Analysis and Visualization

508   Engine for LC−MS Data. *Analytical Chemistry* **82**, 9818-9826 (2010).

509   https://doi.org:10.1021/ac1021166

510   13   Pang, Z. *et al.* Using MetaboAnalyst 5.0 for LC–HRMS spectra processing, multi-omics

511   integration and covariate adjustment of global metabolomics data. *Nature Protocols* **17**,

512   1735-1761 (2022). https://doi.org:10.1038/s41596-022-00710-w

14   Pang, Z. *et al.* MetaboAnalyst 5.0: narrowing the gap between raw spectra and functional

insights. *Nucleic Acids Research* **49**, W388-W396 (2021).

https://doi.org:10.1093/nar/gkab382

15   Chong, J., Wishart, D. S. & Xia, J. Using MetaboAnalyst 4.0 for Comprehensive and

Integrative Metabolomics Data Analysis. *Current Protocols in Bioinformatics* **68**, e86

(2019). https://doi.org:https://doi.org/10.1002/cpbi.86

16   Lundberg, D. S., Yourstone, S., Mieczkowski, P., Jones, C. D. & Dangl, J. L. Practical

innovations for high-throughput amplicon sequencing. *Nat Methods* **10**, 999-1002 (2013).

https://doi.org:10.1038/nmeth.2634

17   Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data

science using QIIME 2. *Nat Biotechnol* **37**, 852-857 (2019).

https://doi.org/10.1038/s41587-019-0209-9

18   Chong, J., Wishart, D. S. & Xia, J. Using MetaboAnalyst 4.0 for Comprehensive and

Integrative Metabolomics Data Analysis. *Current Protocols in Bioinformatics* **68**, e86

(2019). https://doi.org/https://doi.org/10.1002/cpbi.86

Acknowledgements

The authors would like to thank Katherine Havranek and Samia Dutra for stool DNA extractions. We gratefully acknowledge the technical support from the UNC Chapel Hill High Throughput Sequencing Facility (HTSF), a facility that is supported by the University Cancer Research Fund, Comprehensive Cancer Center Core Support grant (P30-CA016086).

Author Contributions

KHM study coordination, recruitment of controls, sample preparation, DNA extractions, funding, and manuscript preparation. CJC collected and analyzed the final metabolomics data, provided bioinformatics for 16S data and cross-omics analysis, and conceptualized results. LR generated a list of microbial derived metabolites from the untargeted analysis, BAA performed stool metabolomics extractions, and ZAV processed the unidentified mass spectral features. PM next generation sequencing. BMW performed a pilot study with a subset of samples in this study. KAJ supervised the untargeted metabolomics analysis. CJC and SRC DHPS identification, characterization, pathway analysis, and conceptualized DHPS models in NDDs. The manuscript was written by CJC and KHM and reviewed by all authors. JCE, originator of MIND study and initial hypothesis, and conceptualization, study design, protocol development, 16S data analysis, manuscript preparation, and funding. RT, CMT, and RR-F screened, recruited, enrolled, and assessed patients and controls and edited the manuscript.

Funding and Conflicts of Interest Statement

Health and Wellbeing Project; and by the Cole Family who support Parkinson's care and research at the Cole Center for Parkinson's Disease and Movement Disorders. The authors have no conflicts of interest to disclose.

Supplementary Information is available for this paper.

Correspondence and requests for materials should be addressed to Christopher Ellis.

Extended Data Figures are available.