Heterosis in crosses between remnant populations of a rare prairie forb: implications for

restoration genetics

Isabelle A. Turner[1,2], Juan Diego Rojas-Gutierrez[1], Bob Easter[3], Christopher G. Oakley*[1]

[1] Department of Botany and Plant Pathology, and the Center for Plant Biology, Purdue

University, West Lafayette, IN 47907, USA

[2] Present address: Department of Biology, Miami University, Oxford, OH 45056, USA

[3] NICHES Land Trust, Lafayette, IN 47905, USA

*Author for correspondence: oakleyc@purdue.edu

Running head: Heterosis in remnant populations of a rare prairie forb

ABSTRACT

- *Premise*

Calls to adopt seed sourcing strategies in biological restoration that preserve local

adaptation while maximizing genetic variation (e.g., regional admixture provenancing)

are increasingly common. Heterosis, the increased fitness of progeny from between-

relative to within-population crosses, could provide an added benefit under such

strategies, but several open questions remain.

- *Methods*

We quantified heterosis in crosses between three small remnant populations of a rare

prairie forb. We measured early fitness components (seed number per fruit, germination,

and juvenile survival) in a greenhouse. Adult fitness components (survival and

reproduction) were quantified over two flowering seasons in two different environments:

a field common garden simulating the initial stages of a restoration, and a greenhouse.

Our approach is unique in estimating heterosis in an environment most relevant to the

early stages of a restoration and in comparing heterosis under field and controlled

conditions.

- *Key Results*

The consequences of between-population crosses for cumulative fitness in the field were

strongly positive in two of the populations (281% and 50% heterosis) and nearly neutral

in a third. Heterosis was generally stronger when measured under field conditions, and

stage specific patterns of heterosis varied among populations.

- *Conclusions*

Regional admixture provenancing in restorations should be beneficial for this species. We

advocate for more research on heterosis in rare species, particularly in restorations. We

also provide a cautionary note on how to calculate and report results for heterosis studies.

Key words: Fragmented populations, Genetic rescue, Heterosis, Prairie, Regional admixture

provenancing, Restoration genetics, Revegetation, *Silene regia*

INTRODUCTION

Limited genetic variation needed to adapt to variable or changing conditions, and the negative fitness consequences of fixed deleterious recessive alleles, are major issues in conservation and restoration genetics (Frankham, 2005; Willi et al., 2006; Bell et al., 2019). One way to probe the relative extent of deleterious recessive alleles fixed within populations in nature is heterosis, the increased fitness in between- relative to within-population crosses (Fenster and Galloway, 2000; Paland and Schmid, 2003; Oakley and Winn, 2012; Lohr and Haag, 2015). Heterosis is common in crosses between natural populations (Fenster, 1991; van Treuren et al., 1993; Ouborg and Van Treuren, 1994; Luijten et al., 2002; Willi and Fischer, 2005; Pickup et al., 2012; Willi, 2013; Weisenberger et al., 2014; Koski et al., 2019; Oakley et al., 2019; Perrier et al., 2020; Söderquist et al., 2020; Thoen et al., 2025). Greater heterosis in small/less diverse compared to large/more diverse populations (Paland and Schmid, 2003; Escobar et al., 2008; Oakley and Winn, 2012; Lohr and Haag, 2015; Spigler et al., 2017) and in selfing compared to outcrossing populations (Busch, 2006; Oakley et al., 2015a) is consistent with a role of drift in fixing partly recessive deleterious alleles (Charlesworth, 2018). This suggests that heterosis may be particularly relevant in conservation and restoration of small remnant populations.

Some outstanding questions in the study of heterosis are how the magnitude depends on experimental assay conditions and to what extent it varies across the life cycle. The former is somewhat surprising given the considerable work on similar questions in inbreeding depression which is also due to partly recessive deleterious alleles. For example, work by Dudash (1990) demonstrated greater inbreeding depression in field compared to greenhouse conditions, and since then the environmental dependence of inbreeding depression has been well documented (reviewed in Fox and Reed, 2010; Cheptou and Donohue, 2011). Heterosis is often estimated in

the greenhouse, though studies estimating heterosis on plants transplanted into the field for at

least part of the life cycle are becoming more common (Fenster, 1991; Ouborg and Van Treuren,

1994; Fenster and Galloway, 2000; Luijten et al., 2002; Wagenius et al., 2010; Maschinski et al.,

2013; Weisenberger et al., 2014; Oakley et al., 2015a; Spigler et al., 2017; Oakley et al., 2019;

Thoen et al., 2025). However, we are unaware of any comparison of heterosis in field vs.

controlled conditions, though there are some comparisons in multiple environments (Fenster and

Galloway, 2000; Willi et al., 2007; Li et al., 2018; Stojanova et al., 2021; Perrier et al., 2022). We

note that heterosis differs from inbreeding depression in that it is caused by alleles fixed within

populations rather than alleles segregating within populations (Whitlock et al., 2000; Whitlock,

2002; Charlesworth, 2018). Also, unlike inbreeding depression which can be comprised of both

strongly deleterious and more mildly deleterious alleles (Charlesworth and Willis, 2009),

heterosis is expected to be mostly due to mildly deleterious alleles (Whitlock et al., 2000;

Glémin et al., 2003; Charlesworth, 2018). Heterosis may therefore be weak for individual fitness

components, particularly early in the life cycle (c.f., Husband and Schemske, 1996; Perrier et al.,

2020). Therefore, obtaining accurate estimates of heterosis may require estimates of fitness

encompassing as many life stages as possible.

One context in which heterosis could be potentially useful is in restoration projects

involving rare species. Restoration practitioners use seeds, or sometimes seedling plugs, to

establish populations (Shaw et al., 2020), often including rare or threatened species that are

dispersal- or seed-limited (Pywell et al., 2002; Seabloom et al., 2003; Clark et al., 2007). The

method of seed sourcing and the genetics of the source populations can greatly influence

establishment success (De Vitis et al., 2022). Traditionally it was thought that seeds should be

sourced as locally as possible to minimize the risk of disrupting adaptation to local

environmental (extrinsic) conditions or outbreeding depression due to (intrinsic) genetic

incompatibilities (McKay et al., 2005; Gann et al., 2019; Török et al., 2024). However, potential

seed sources for restoration are often small remnant populations from which only a small number

of individuals (sometimes only a single individual) are sampled (Godefroid et al., 2011). Thus,

the founding population may harbor fixed deleterious mutations and have limited genetic

variation to adapt to fluctuating or changing conditions, particularly in the face of climate

change. Because of these risks, many authors have called for some version of seed sourcing

strategies that increase genetic variation while also minimizing outbreeding depression and the

loss of local adaptation (Broadhurst et al., 2008; Godefroid et al., 2011; Breed et al., 2013;

Bucharova et al., 2019; St. Clair et al., 2020; Hoffmann et al., 2021; Stojanova et al., 2021;

Jordan et al., 2023). One proposed strategy is "regional admixture provenancing" (Bucharova et

al., 2019), where seed is sourced from multiple populations regionally to increase genetic

variation. These populations would however come from similar habitat in the same eco-region,

likely connected by gene flow prior to fragmentation, to minimize the risks of maladaptation and

genetic incompatibilities. Additionally, heterosis following admixture in a restoration may

provide at least a short term fitness boost that may aid establishment of rare species, analogous

to the "catapult effect" proposed for the role of heterosis in the establishment of invasive species

(Drake, 2006; Li et al., 2018). Despite the potential benefits, there are few empirical tests of the

regional admixture provenancing approach (but see St. Clair et al., 2020) or the potential for

heterosis during a newly established restoration.

Here we used within- and between- population crosses of three small remnant

populations of a rare prairie forb, *Silene regia* Sims., to quantify heterosis for multiple fitness

components in both a greenhouse experiment and a field experiment mimicking a newly

established restoration. We addressed the following questions: 1) What is the magnitude of

heterosis (or conversely, outbreeding depression) under restoration conditions in the field, and

does it vary among populations? 2) Is the magnitude of heterosis greater in a field common

garden compared to controlled greenhouse conditions? and 3) How does the magnitude of

heterosis vary across different life history stages? We expected to observe greater heterosis under

field compared to greenhouse conditions, as well as greater heterosis in estimates of fitness

incorporating later life history stages.

MATERIALS AND METHODS

Study system--

Silene regia (royal catchfly) is a tap-rooted perennial in the Caryophyllaceae family (Dolan,

1994). It is endemic to a dozen states in eastern North America and is thought to have been

broadly distributed across prairies prior to the widespread conversion of this habitat to

agriculture (Menges, 1991). Contemporary populations across much of the native range are small

and isolated and exist as remnants in prairie cemeteries and along railroad rights-of-way and

roadsides (Dolan, 1994). Remnant populations of *S. regia* are potentially at risk of genetic

depletion (Menges and Dolan, 1998). This species does not vegetatively reproduce and achieves

sexual reproduction via hummingbird pollination. It is self-compatible and some geitonogamous

selfing may occur, but it is likely mostly outcrossing as the individual flowers are protandrous

(Dolan, 1994; Menges, 1995).

We selected three remnant focal populations in different counties in western Indiana (Tip

= Tippecanoe; Fou = Fountain; Ver = Vermillion). Exact locations of these populations are not

provided because this species is critically imperiled in Indiana (NatureServe, 2024), but these are

distinct populations, all separated by at least 14 km. These three populations include all known

existing remnants within a 55 km radius, and there may be as few as 10 extant remnant

populations in the entire state of Indiana (B. Easter, pers. obs.). While including more

populations could have broadened our scope of inference, we restricted the sampling to remnants

from the same local area as would be advised under regional admixture provenancing. The

smallest population by census size was Tip (10 flowering individuals) from a ~0.5 ha remnant,

which has likely been small for an extended period (B. Easter, pers. obs.). The largest population

was Ver (~50 flowering individuals) from a ~0.4 ha remnant, and population size reductions are

likely more recent at this site (B. Easter, pers. obs.). The size of Fou, from a ~0.4 ha remnant,

was intermediate to the other populations (~20 flowering individuals). While there is some

variation in census size among these remnants, they are all quite small and are thus all expected

to have limited genetic variation and harbor fixed deleterious mutations.

Generation of seeds for the fitness assays--

Bulk seed was collected from each population in September 2019. Our goal was to characterize

population level heterosis, and the small number of individuals in these populations precluded

experimental designs accounting for distinct maternal lineages. We collected seeds from most of

the reproductive individuals in the two smaller populations and from a haphazard subset of

approximately 20 individuals in the largest population, Ver.

In early October 2019, seeds from each population were surface sterilized in a 10%

bleach solution for 10 minutes, followed by three rinses in deionized-autoclaved water. The

seeds were then placed into moist paper towels in bags and cold stratified in the dark at 4°C for

10, 30, or 60 days. Different stratification periods were used because 60 days would be ideal for

germination, but starting the plants sooner was preferrable due to logistical constraints. Because

of low overall germination we included individual maternal plants arising from all three seed

stratification treatments. While there was a significant effect of seed stratification treatment (for

seeds that produced the parental plants) on seed number per fruit following controlled crosses,

this treatment (involving the generation prior to the experimental assay) did not qualitatively

affect the results for either the effects of cross type or the interaction between maternal

population and cross type (compare Table S1 to Table 1) so we omitted this term from

subsequent analyses.

After stratification, we sowed seeds (5 per cell) into a 72-cell flat insert (cell volume = 70

ml, Greenhouse Megastore, Danville, Illinois, USA), filled with a 3:1 mixture of MetroMix

potting soil (Sun Gro Horticulture, Agawam, Massachusetts, USA) and autoclaved sand. A total

of 30-50 seeds were sown for each population-stratification treatment combination. The seeds

were then germinated in an incubator (Percival scientific, Perry, Iowa, USA) set to a constant

27°C and a 12-hour photoperiod. After approximately 4 weeks, cells were thinned to one

seedling per cell, and the flat was moved to a heated greenhouse (minimum: 20°C) with

supplemental lighting (12-hour photoperiod). Plants were grown there for approximately 6

weeks, watering as needed. Plants were then moved to a large refrigerator (4°C) with lights (10-

hour photoperiod) for approximately one month to synchronize flowering. After the end of this

vernalization period in mid-January 2020, plants were moved back to the greenhouse and

transplanted into 17 cm circular pots (volume = 2.4 l, Greenhouse Megastore, Danville, Illinois,

USA) filled with the same soil mixture. Plants were watered as needed and fertilized monthly

with ¼ strength Hoagland's solution.

Once plants had begun to flower (early March 2020), we performed controlled hand

pollinations within and between populations in all possible combinations and crossing directions.

In mid-March, because of COVID related shutdowns, the plants (19, 13, and 10 individuals from

Fou, Tip, and Ver respectively) were moved to a private residence and grown on light racks with

a 14-hour photoperiod for the duration of the hand pollinations. Most of the pollinations were

performed from mid-March through the end of June 2020. Before each pollination we identified

ovule donor flowers several days prior to anthesis and removed the upper part of the corolla

tube, filaments, and anthers with scissors and forceps. Pollinations were completed when the

stigma lobes of the ovule donor flowers appeared to be receptive (usually 3-4 days later) by

removing stamens from a pollen donor flower with forceps and brushing the dehisced anthers

against the stigma. Each pollination was tracked with a unique color combination of embroidery

floss tied around the pedicel. Pollinations were done haphazardly based on flower availability,

but we attempted to balance the cross types done for a given maternal population on a given day

to minimize confounding treatment effects with age effects. Because we started with bulked seed

collections and were interested in population level estimates of heterosis, and because of

practical constraints of the number of receptive flowers per day on a given plant, we did not

execute a balanced crossing design at the level of individual plants. However, most individual

plants participated as both ovule and pollen donors, with population mean averages of 5-8

flowers as ovule donors and 4-7 flowers as pollen donors for between population crosses per

plant. For within population crosses, each individual plant contributed an average (by

population) of 3-4 flowers as ovule donors and 2-3 flowers as pollen donors.

The number of hand pollinations per maternal population ranged from 39-58 (mean = 46)

for within-population (WIN) crosses and 85-102 (mean = 91) for between-population (BET)

crosses, for a grand total of 411 crosses. The approximately two-fold greater number of hand

pollinations for between-population crosses is because each maternal population was crossed to

two different paternal populations, and we wanted sufficient power to test for interactions

between maternal and paternal populations.

In addition to hand pollinations, we performed control emasculations where flowers were

emasculated but no pollen was applied to the stigma lobes. Of the 69 control emasculations, less

than 3% (2 emasculations) resulted in a fruit, giving us confidence that the vast majority of the

pollinations performed were successful crosses between the intended pollen and ovule donors.

Early fitness components--

We collected fruits of all successful pollinations to be used in subsequent fitness assays. This

species required an extended period between pollination and fruit maturity (mean = 56.8 days,

SD =10.4). There were no patterns in fruit set or the rate of fruit development with respect to

cross type, but fruits of Tip developed on average about 5 days faster than fruits from the other

two populations. For each of 17-39 WIN fruits and 45-71 BET fruits per maternal population

(Table S2), we counted the number of seeds per fruit, excluding small, shriveled seeds presumed

to be inviable.

We selected a subset of the viable seeds for each population and cross type, prioritizing

fruits with enough viable seeds for germination assays and the goal of proportionally balanced

sample sizes. This subset included approximately 300 seeds each of the WIN and the two BET

combinations for populations Tip and Ver and approximately double those numbers for Fou

(Table S2). These seeds were cold stratified as previously described for a period of 7 weeks

starting in mid-November of 2020. Seeds were then sown into 72 cell flat inserts filled with the

same soil mix (5 flats in total) with 7-12 (mean = 9.96, SD = 0.34) seeds per cell in a randomized

block design. The number of cells per flat varied by maternal population with an average of 6

and 12 for WIN and BET crosses respectively from Tip and Ver, and approximately double the

sample size for each cross type from Fou (Table S2). We recorded germination daily for a period

of 24 days starting in mid-January 2021. Most seeds germinated within the first week, and only 5

new germinants were recorded during the second half of the scoring period.

Once seeds had germinated and seedlings had two true leaves, they were transplanted

into individual cells of 72 cell flat inserts and moved to the greenhouse. The seedlings were

transplanted into an incomplete randomized block design with a total of 30 flats (blocks). The

number of cells per flat varied by maternal population, with an average of 5-6 for WIN and 10

for BET crosses from Tip and Ver, and an average of 11 WIN and 24 BET crosses from Fou. Any

seedlings that died within the first week were replaced with another seedling of the same

population-cross type. This mortality was likely due to transplant shock, and these individuals

were omitted from subsequent analyses. Once roots started to poke through the bottom of the

cells, individuals were transplanted first into 6 cm square (volume = 250 ml, Greenhouse

Megastore, Danville, Illinois, USA) pots filled with the same soil mixture. Plants were watered

as needed and fertilized twice with ¼ strength Hoagland's solution. Cumulative juvenile survival

was recorded up to approximately seven weeks after stratification, excluding individuals that

died within one week of the initial transplant. Surviving plants that were established in the 6 cm

pots were then randomly divided into two experiments: one in the greenhouse and one in a field

common garden simulating a new restoration.

Greenhouse experiment--

In late March 2021, seedlings were transplanted into 11 cm square (volume = 1L, Greenhouse

Megastore, Danville, Illinois, USA) pots filled with the same potting mix in an incomplete

randomized block design consisting of 48 flats each containing 7-8 pots. Each maternal

population cross type combination was represented by an average of 1.3-2.6 plants in at least 19

different flats (blocks). The total size of the greenhouse experiment was 354 plants with sample

sizes for between population crosses being approximately double that of within population

crosses, and sample sizes for Fou being approximately double that of the other populations

(Table S2). Plants were grown for approximately 18 months, until mid-October of 2022, which

included two flowering periods. During this time plants were watered as needed and fertilized

once with Osmocote timed release 15-9-12 fertilizer at ½ the recommended dose after the first

flowering period was over.

261   For each plant in the greenhouse experiment we scored adult fitness as the cumulative

number of flowers produced over ~18 months (two flowering seasons) including zeros for plants

that died prior to reproducing or that did not reproduce. We additionally quantified two

components of this composite measure of adult fitness separately: combined cumulative

probability of survival and reproduction (did a plant reproduce or not over the entire course of

the experiment), and cumulative fecundity (total flower number over the entire course of the

experiment) of reproducing plants.

Field experiment--

We established a field common garden experiment to quantify adult fitness components under

more realistic restoration conditions. The site was located on the property of Arbor America LLC

in West Point, Indiana (Tippecanoe County) USA. This location is within the extant range of *S.*

regia, and the field common garden experiment was planted on Coloma Sand (Mixed, mesic

Lamellic Udipsamments), a soil type and series similar to the well-drained sites where *S. regia*

naturally occurs (ISEE Network, 2024). The plot had previously been treated to establish a

prairie strip with an application of Roundup in May 2020 to remove the turf, followed by

controlled burning approximately 1 month later, and another application of Roundup after

approximately 2 more months. Herbicide applications and controlled burns are typical of

establishing grassland restorations and help to promote native forb establishment (Török et al.,

2024). In early March 2021, the plot was hand seeded with a short stature, high-diversity prairie

mix from Spence Restoration Nursery (Muncie, IN, USA).

In early April 2021, seedlings were transplanted into a randomized complete block design

consisting of six blocks containing 120 plants each. In total there were 720 plants, with

approximately double the sample size of the greenhouse experiment for each maternal

population-cross type combination (Table S2). Each block was a rectangular grid of 3x40 plants

with 30 cm spacing between the plants. Plants were hand watered for the first week only, and

plants that died during this first week were considered to have succumbed to transplant shock

and were replaced. In July of both 2021 and 2022, other vegetation within 10 cm of the base of

experimental plants was clipped once to facilitate locating plants for the post-reproductive

censuses. While the conditions of the experimental plot closely mimic the conditions during the

initial phase of a restoration, we acknowledge that this species would likely be sown as seeds in

an actual restoration. Given the extremely low rates of field establishment from seed for this

species (Menges, 1995) and these populations in particular (< 2%, C. Oakley, unpublished data),

and the effort to create enough seeds via experimental crosses, starting these plants in the field as

seedlings was a necessary compromise.

In late October 2021, we recorded whether each plant was still alive and had reproduced,

and for plants that had reproduced we counted the number of fruits. Only about 11% of all plants

had reproduced in the first year. We re-censused the same fitness components in October 2022.

We estimated adult fitness as the total number of fruits (sum over both years) produced per

individual transplanted, including zeros for plants that died before reproducing or surviving

plants that did not reproduce. We additionally calculated the combined cumulative (across both

years) probability of survival and reproduction for each plant and cumulative fecundity (total

number of fruits across both years) for each reproducing plant.

Statistical analyses and estimates of heterosis--

We analyzed each fitness component using ANOVA. We tested for the effects of maternal

population (Fou, Tip, and Ver), paternal population, and their interaction. For all fitness

components except seed number per fruit from the initial crosses (where there was no blocking

possible), we included a term for block in the analysis to account for micro-spatial variation

within the experiment. All effects were treated as fixed effects because the limited number of

populations and blocks preclude treating them as a truly random sample. Inspection of residuals

indicated no major issues with normality or heteroscedasticity for many fitness components, and

for these we used normal error distributions. Juvenile survival and the combined probability of

survival and reproduction in both the greenhouse and field experiments required a generalized

linear model with a binomial error distribution. For each model with a significant interaction

between maternal and paternal populations or significant main effects for both maternal and

paternal population, we used linear contrasts to test *a priori* hypotheses. We first contrasted BET

crosses to WIN crosses (over all populations) to test if there was significant overall heterosis (or

outbreeding depression).

We then performed two different sets of contrasts to test for heterosis (or outbreeding

depression) separately by maternal population. First, we contrasted BET crosses to WIN crosses

for each maternal population. This is the traditional approach to investigating heterosis within

maternal populations. However, this approach may lead to misleading conclusions if progeny

from WIN crosses from different populations have very different mean fitness. To illustrate this

issue, consider a hypothetical example where population A has very low within population

fitness and population B has very high within population fitness. With purely additive effects

(i.e. no dominance complementation or heterosis) the cross between these two populations

should have fitness equal to the mean of the two parents, a.k.a. the midparent value (Lynch and

Walsh, 1998; Demuth and Wade, 2006; Oakley et al., 2015b). Comparing BET crosses to WIN

crosses for population A could lead to the mistaken interpretation of heterosis, and a similar

mistaken interpretation of outbreeding depression could result for the comparison in population

B. To more accurately test for heterosis or outbreeding depression within maternal populations

while accounting for additive differences in mean population fitness we performed contrasts of

BET crosses for a given population to WIN crosses pooled over all populations (because all

populations were crossed in all combinations). In discussing the results, we primarily focus on

these corrected contrasts but point out cases where results for the two methods differ notably. All

statistical analyses were performed in JMP Pro v. 16.1.0 (JMP, 1989-2021).

For each fitness component and composite estimate of fitness we calculated estimates of

heterosis separately for each population using means for each maternal population-cross type

combination averaging over the block means (Table S2). For the traditional approach we

calculated the percent increase in fitness of between-population crosses relative to within-

population crosses ([(BET-WIN)/WIN]*100), where positive values indicate heterosis and

negative values indicate outbreeding depression. To account for differences in mean WIN fitness

among populations, we pooled WIN crosses from all three populations for the denominator

([(BET-WIN_{pool})/WIN_{pool}]*100). We additionally calculated heterosis relative to the midparent

value for each pair of maternal and paternal populations. However, relative fitness of BET

crosses for different paternal populations were always of the same sign (i.e. variation in the

degree of heterosis or outbreeding depression for both populations, but never contrasting

directions of effects; Figure S1). For simplicity, we therefore focus on the estimates pooling

within population crosses across populations but estimates for all three approaches can be found

in Table S3.

For cumulative fitness, calculated separately for the greenhouse and field experiments,

we used the product of four estimates: 1) seed number per fruit, 2) proportion germination, 3)

proportion juvenile survival, and 4) adult fitness. Because this multiplicative measure of

cumulative fitness yields only one estimate per cross type per population, we could not directly

test the effects of population, cross type, and their interaction as with the fitness components.

Other authors have approached this problem by using replicate line level estimates of

multiplicative fitness (see for e.g., Dudash, 1990), but we were unable to use this approach both

because of our initial bulked seed collection and because not all cross types were performed on

all plants.

RESULTS

Early fitness components--

In general, heterosis was weak and/or non-significant for early fitness components (Fig. 1,

Tables 1&S3). For seed number per fruit and germination there were genetically based

differences among maternal populations (Table 1), with Fou having greater overall seed number

per fruit and germination than the other two populations (Fig. S2; Table S2). There was no

overall heterosis for either fitness component (BET_{pool} vs. WIN_{pool} contrast in Table 1). However,

there was suggestive evidence of heterosis in germination for Fou (Table 1). On average,

estimates of population level heterosis (or outbreeding depression) for seed number per fruit

ranged from -14% to 41%, and estimates of heterosis (or outbreeding depression) for

germination ranged from -7% to 23% (Table S3). For juvenile survival there was a significant

interaction between maternal and paternal population (Table 1) and significant overall heterosis

of about 6%. Pairwise contrasts of cross types separately by maternal population indicated

significant heterosis (13%) only for Fou (Fig. 1, Tables 1&S3).

Greenhouse experiment--

For adult fitness (flowers per seedling planted) in the greenhouse experiment, there were no

significant effects for any of the model terms (Fig. 2; Table 2). On average, relative fitness

estimates (Table S3) ranged from modest heterosis (32%) in Ver to weak outbreeding depression

in Fou (-3%) and Tip (-6%). There was limited evidence of significant effects for fitness

components (Fig. 2; Table 2). However, there was a suggestive interaction between maternal and

paternal population for fecundity (Table 2) with about 19% overall heterosis. This was most

evident in Ver with 36% heterosis (Figs. 2&S3; Table S3). For the combined probability of

survival and reproduction there was weak outbreeding depression on average for all populations

-3% to -13% (Table S3). For this experiment, the three WIN crosses had similar mean fitness,

and thus calculations of heterosis were qualitatively similar between the different approaches

(Figs. 2&S3; Table S3).

Field experiment--

For adult fitness (fruits per seedling planted) in the field, there were significant effects of both

maternal and paternal population (Table 3), indicating genetically based differentiation in mean

fitness among populations. Most notably there was four-fold variation in mean fitness of progeny

from WIN crosses, ranging from 2 fruits per seedling planted for Tip, to 3 for Fou, and 8 for Ver

(Fig. S4; Table S2). Additionally, there was a significant interaction between maternal and

paternal populations which could indicate overall heterosis (or outbreeding depression) and/or

variation among populations in the degree of heterosis of BET crosses (Table 3). Indeed,

contrasts revealed significant overall heterosis (BET_{pool} vs. WIN_{pool}) of about 45%. Contrasts of

BET crosses accounting for additive differences between populations (BET vs. WIN_{pool})

indicated significant heterosis only for Fou (Fig. 3; Table 3) of about 78% (Table S3). At least on

average, there was also heterosis for Tip (20%) and Ver (36%) (Tables 3&S3).

The methodology for testing/calculating relative fitness separately for maternal

populations had a considerable effect on adult fitness for some populations (Tables 3&S3). This

is likely due to the extent of population differentiation in mean fitness of progeny from within-

population crosses, which had not been observed for other fitness components thus far. Using the

traditional approach, we would conclude significant heterosis of 160% for Tip (Tip: BET vs.

WIN contrast Table 3) compared to the non-significant 20% heterosis above (Table S3). The

results for Ver do not qualitatively change between approaches but there is a reversal in the sign

of effects between the two approaches, -26% outbreeding depression with the traditional

approach compared to 36% heterosis above (Table S3). Both approaches indicate significant

heterosis for Fou (Table S3).

There was also significant variation among populations and population combinations for

individual fitness components. For the combined probability of survival and reproduction there

were significant effects of both maternal and paternal populations. As with adult fitness, the

probability of surviving and reproducing for WIN crosses in Ver was much greater (about 1.6-

fold) than for the other populations (Fig. S4; Table S2). There was no significant overall

heterosis (about 10% on average) for this fitness component, but there was a suggestive effect of

heterosis (23%) in Fou (Fig. 3; Tables 3&S3). For fecundity, there was a significant effect of

paternal population and a suggestive effect for the interaction between maternal and paternal

populations (Table 3). There was significant overall heterosis for fecundity of about 38% with

heterosis in all populations (29 to 55%), at least on average, significantly so in Fou (Fig. 3;

Tables 3&S3).

As with adult fitness, the methodology for testing/calculating relative fitness for the

fitness components had a strong influence on the results. For probability of survival and

reproduction, the traditional approach would suggest highly significant heterosis for Fou (Fou:

BET vs. WIN contrast Table 3; Table S3) of a much greater magnitude (89% compared to 23%

above) and highly significant outbreeding depression (-34% compared to an average of 11%

heterosis for the corrected comparison) for Ver (Tables 3&S3). For fecundity, both methods

yielded qualitatively similar results, likely because of the more limited variation among maternal

populations in mean fecundity.

Cumulative fitness--

Estimates of cumulative heterosis across all measured life stages were substantial in both

greenhouse and field experiments, but they were generally stronger in the field experiment. For

the greenhouse experiment there was about 35% heterosis overall, and within populations

heterosis was 92% for Fou and 35% for Ver, and there was weak (and non-significant based on

results for fitness components) outbreeding depression (-23%) for Tip (Table S3). Using the

traditional comparison suggested less heterosis for Fou and more for Ver (Table S3). For the field

common garden experiment, heterosis was 112% over all populations. All three populations

exhibited heterosis for cumulative fitness with considerable variation in magnitude among

populations, from weak (6%) in Tip to modest (50%) in Ver to very strong (281%) in Fou (Table

S3). Using the traditional comparisons would cause the false conclusion of very strong heterosis

(131%) in Tip and weak outbreeding depression (-4%) in Ver, but estimates for Fou were similar

for both methods (Table S3).

DISCUSSION

Seed sourcing strategies for restoration such as regional admixture provenancing have been

suggested to help maintain both local adaptation and amounts of genetic variation within

restored populations. Heterosis may provide an added short-term benefit to such strategies. To

our knowledge, our study is the first to quantify heterosis for multiple fitness components in a

newly established restoration. Averaged over all populations, the consequences of between

population crosses for cumulative fitness was strongly positive (112% overall heterosis) in the

field, suggesting that regional admixture provenancing should provide at least a short-term

fitness advantage for this species. Estimates of heterosis from the field were greater than those

from the greenhouse (35% overall heterosis). Additionally, life-stage specific estimates of both

mean population fitness and heterosis varied among populations, particularly in the field

experiment. This suggests that while there is an overall benefit of crosses between populations

under realistic field conditions, relative fitness outcomes can depend on the populations involved

and the environmental conditions of the fitness assay.

Results from the field experiment simulating early restoration conditions is most directly

relevant for restoration practitioners. Over all populations there was significant heterosis for both

fruits per seedling planted (45%) and fecundity (38%), and on average there was consistently

increased relative fitness for these measures (ranging from 20-78%) within populations (Table

S3). Heterosis for the combined probability of survival and reproduction was weaker (10%

overall), and none of the results for individual populations were significant. Thus, admixture of

these populations in field restorations are likely to result in a net positive effect, at least for

reproductive fitness, despite some among population variation in the degree of heterosis.

The among population variation in mean fitness in the field experiment is important for

the calculation and interpretation of heterosis for individual populations, and also has direct

implications for restoration potential. For example, Ver had much greater survival and adult

fitness of WIN crosses compared to other populations (Fig. S4). This highlights the importance

of using the mid-parent value as the frame of reference in calculating heterosis to avoid

confounding additive and non-additive effects (field results for Ver; Table S3). Estimating

heterosis as a proportional deviation from the mid-parental value is commonly used for natural

populations with predominantly selfing mating systems (Lynch and Walsh, 1998; Oakley et al.,

2019; Clo et al., 2021), but it should be broadly applicable to other systems. Alternatively,

studies where different population categories (e.g. small vs. large) are represented by many

populations (Paland and Schmid, 2003; Oakley and Winn, 2012; Willi, 2013) or where each

maternal population is crossed with many paternal populations (Busch, 2006; Weisenberger et

al., 2014; Spigler et al., 2017; Thoen et al., 2025) achieve a similar end. These approaches

minimize the influence of individual populations with very high or very low mean within

population fitness by averaging over many populations. The use of such approaches to minimize

confounding additive and dominance effects can also be important in greenhouse studies if the

parental populations differ in mean fitness (Rojas-Gutierrez et al., 2026). The greater mean

fitness within Ver may be because it is the largest of the three populations and more recently

reduced in size and may therefore harbor fewer fixed deleterious recessive alleles. The low mean

adult fitness and fitness components within Tip for the field experiment (Fig. S4) was also

notable, especially because the origin of this population was geographically closest to the

planting site. This was the smallest population and may therefore suffer the most from fixed

deleterious alleles, representing a situation in which the most "local" population may not always

be the "best" source material for a restoration (Pizza et al., 2023).

Field experiments of heterosis are logistically challenging but are increasingly more

common, and like our study are often based on transplanted seedlings (Fenster, 1991; Ouborg

and Van Treuren, 1994; Wagenius et al., 2010; Weisenberger et al., 2014; Oakley et al., 2015a;

Spigler et al., 2017; Oakley et al., 2019; Thoen et al., 2025). We limit our discussion to studies of

heterosis in rare or recently fragmented species because of the expected context dependency of

heterosis on population size, genetic distance between parents, and spatial scale of the field

experiment. Our overall conclusions agree with an exceptional 14-year field study that found

inter-population variation in heterosis and limited evidence for outbreeding depression of seven

fragmented populations of *Echinacea angustifolia* (Thoen et al., 2025). An 8-year field study in

the same species conducted near the Thoen et al. (2025) study also found no evidence for

outbreeding depression, but curiously did not find any heterosis (Wagenius et al., 2010), which

Thoen et al. (2025) attribute to environment dependent expression of heterosis. Generally

speaking, other field studies in rare taxa also find heterosis and little evidence for outbreeding

depression (Weisenberger et al., 2014), though the magnitude of heterosis seems to depend on

population characteristics (Pickup et al., 2012) and temporal environmental variation

(Maschinski et al., 2013).

Comparisons between field- and greenhouse-based estimates of heterosis are rare in the

literature, though some studies have indicated that heterosis may be stronger under more

stressful conditions (Maschinski et al., 2013; Hahn and Rieseberg, 2017; Stojanova et al., 2021).

Our results indicate that heterosis is greater in field compared to greenhouse conditions for fruits

per seedling planted (adult fitness) and both of its component parts, combined probability of

survival and reproduction and fecundity. We also found stronger average heterosis in the field

compared to greenhouse for cumulative fitness. Stronger heterosis under presumably more

variable and stronger selection pressures in the field compared to the greenhouse is in agreement

with results from many, though not all, similar comparisons for inbreeding depression (Dudash,

1990; Fox and Reed, 2010; Cheptou and Donohue, 2011).

In addition to the comparison between field and greenhouse, we were interested in how

heterosis might change across different life stages. If heterosis is caused by many mildly

deleterious alleles, we would expect to see greater heterosis in later life history stages or in

estimates that integrate multiple fitness components (van Treuren et al., 1993; Willi et al., 2007;

Escobar et al., 2008; Oakley et al., 2015a; Spigler et al., 2017; Perrier et al., 2020; Thoen et al.,

2025) as is seen for inbreeding depression in highly selfing species (Husband and Schemske,

1996). In general, the results for heterosis over all populations follow such a pattern, with no

significant heterosis for the earliest two fitness components, seed number per fruit and

germination, and very modest heterosis for seedling survival. It is important to note that in our

study all early components of fitness were measured in the greenhouse. Similar experiments are

rarely started from seed (but see Luijten et al., 2002), likely due to low field germination rates

(e.g. Thoen et al., 2025) as expected for *S. regia* (Menges, 1995). Strong heterosis for early

fitness components has been reported for some species even in controlled environment

conditions (Busch, 2006; Oakley et al., 2015a; Söderquist et al., 2020), but outbreeding

depression for early fitness components is also possible (Sletvold et al., 2012). For later life

stages in the greenhouse there was stronger overall heterosis for fecundity (19%), though not for

the combined probability of survival and reproduction or adult fitness. Incorporating additional

years of fecundity estimates would likely increase overall heterosis in both the greenhouse and

field environments. Thus, our measures of adult fitness components may underestimate overall

heterosis.

In conclusion, the heterosis reported here for crosses between these small remnant

populations in conditions representative of a newly established restoration strongly suggest that

regional admixture provenancing would be a beneficial strategy for this species. The short-term

fitness boost provided by heterosis is in addition to any potential long-term benefits of increased

genetic variation necessary to adapt to environmental heterogeneity and climate change. We

agree with previous authors that outbreeding depression is unlikely to have serious negative

consequences for restoration and conservation efforts (Frankham, 2015; Ralls et al., 2018; Bell et

al., 2019; Hoffmann et al., 2021; Jordan et al., 2023) if appropriate guidelines are followed

(Frankham et al., 2011; Weeks et al., 2011). Even if there is some outbreeding depression in

recombinant generations in this system, selection is expected to quickly remove deleterious

combinations (Luijten et al., 2002; Erickson and Fenster, 2006; Aitkin and Whitlock, 2013;

Hamilton and Miller, 2016). In future studies of heterosis, we suggest crossing designs and/or

analytical approaches (i.e. comparisons to mid-parent value for paired population designs) to

interpret dominance effects correctly accounting for differences in mean fitness between

populations. Furthermore, we suggest that it is important to report mean values for each cross

type for each population in addition to estimates of heterosis/outbreeding depression. Many open

questions about heterosis remain, and we agree with Menges (2008) that restorations should be

seen as experimental opportunities in addition to their other benefits.

ACKNOWLEDGEMENTS

We thank P. Chidambaram, C. Gallick, G. Lee, M. Lopp, N. Ryan, T. Soto, and J. Will for

assistance with the experiments. We also thank G. Pardillo and Arbor America for permission to

conduct the common garden experiment on their property, and K. Jacobsen for agreeing to house

the plants for crossing for nearly four months during the COVID shutdown. This work was

funded in part by USDA Hatch grant 7000281 to CGO via Purdue College of Agriculture and a

Biodiversity Research grant from the Indiana Native Plant Society to IAT.

AUTHOR CONTRIBUTIONS

I.A.T. and C.G.O. conceived of and designed the study. B.E. provided the seeds and established

the field plot. I.A.T. and C.G.O. conducted the experiment and collected the data. C.G.O.,

J.D.R.G. and I.A.T. analyzed the data. J.D.R.G. produced the figures. I.A.T. and C.G.O. drafted

the manuscript. All authors contributed to revising the manuscript.

LITERATURE CITED

Aitkin, S., and M. C. Whitlock. 2013. Assisted gene flow to facilitate local adaptation to climate change. *Annual Review of Ecology Evolution and Systematics* 44: 367-388.

Bell, D. A., Z. L. Robinson, W. C. Funk, S. W. Fitzpatrick, F. W. Allendorf, D. A. Tallmon, and A. R. Whiteley. 2019. The exciting potential and remaining uncertainties of genetic rescue. *Trends in Ecology & Evolution* 34: 1070-1079.

Breed, M. F., M. G. Stead, K. M. Ottewell, M. G. Gardner, and A. J. Lowe. 2013. Which provenance and where? Seed sourcing strategies for revegetation in a changing environment. *Conservation Genetics* 14: 1-10.

Broadhurst, L. M., A. Lowe, D. J. Coates, S. A. Cunningham, M. Mcdonald, P. A. Vesk, and C. Yates. 2008. Seed supply for broadscale restoration: maximizing evolutionary potential. *Evolutionary Applications* 1: 587-597.

Bucharova, A., O. Bossdorf, N. Hölzel, J. Kollmann, R. Prasse, and W. Durka. 2019. Mix and match: regional admixture provenancing strikes a balance among different seed-sourcing strategies for ecological restoration. *Conservation Genetics* 20: 7-17.

Busch, J. W. 2006. Heterosis in an isolated, effectively small, and self-fertilizing population of the flowering plant *Leavenworthia alabamica*. *Evolution* 60: 184-191.

Charlesworth, B. 2018. Mutational load, inbreeding depression and heterosis in subdivided populations. *Molecular Ecology* 27: 4991-5003.

Charlesworth, D., and J. H. Willis. 2009. The genetics of inbreeding depression. *Nature Reviews Genetics* 10: 783-796.

Cheptou, P.-O., and K. Donohue. 2011. Environment-dependent inbreeding depression: its ecological and evolutionary significance. *New Phytologist* 189: 395-407.

594   Clark, C. J., J. R. Poulsen, D. J. Levey, and C. W. Osenberg. 2007. Are plant populations seed

595   limited? A critique and meta-analysis of seed addition experiments. *American Naturalist*

596   170: 128-142.

597   Clo, J., J. Ronfort, and L. Gay. 2021. Fitness consequences of hybridization in a predominantly

598   selfing species: insights into the role of dominance and eistatic incompatibilities.

599   *Heredity* 127: 393-400.

600   De Vitis, M., K. Havens, R. S. Barak, L. Egerton-Warburton, A. R. Ernst, M. Evans, J. B. Fant,

601   et al. 2022. Why are some plant species missing from restorations? A diagnostic tool for

602   temperate grassland ecosystems. *Frontiers in Conservation Science* 3:

603   10.3389/fcosc.2022.1028295.

604   Demuth, J. P., and M. J. Wade. 2006. Experimental methods for measuring gene interactions.

605   *Annual Review of Ecology, Evolution and Systematics* 37: 289-316.

606   Dolan, R. W. 1994. Patterns of isozyme variation in relation to population size, isolation, and

607   phytogeographic history in royal catchfly. *American Journal of Botany* 81: 965-972.

608   Drake, J. M. 2006. Heterosis, the catapult effect and establishment success of a colonizing bird.

609   *Biology Letters* 2: 304-307.

610   Dudash, M. R. 1990. Relative fitness of selfed and outcrossed progeny in a self-compatible

611   protandrous species, *Sabatia angularis* L. (Gentianaceae): a comparison in three

612   environments. *Evolution* 44: 1129-1139.

613   Erickson, D. L., and C. B. Fenster. 2006. Intraspecific hybridization and the recovery of fitness

614   in the native legume *Chamaecrista fasciculata*. *Evolution* 60: 225-233.

Escobar, J. S., A. Nicot, and D. P. 2008. The different sources of variation in inbreeding

depression, heterosis and outbreeding depression in a metapopulations of *Physa acuta*.

Genetics 180: 1593-1608.

Fenster, C. B. 1991. Gene flow in *Chamaecrista fasciculata* (Leguminosae) II. Gene

establishment. *Evolution* 45: 410-422.

Fenster, C. B., and L. F. Galloway. 2000. Population differentiation in an annual legume: genetic

architecture. *Evolution* 54: 1157-1172.

Fox, C. W., and D. H. Reed. 2010. Inbreeding depression increases with envionmental stress: and

experimental study and meta-analysis. *Evolution* 65: 246-258.

Frankham, R. 2005. Genetics and extinction. *Biological Conservation* 126: 131-140.

Frankham, R. 2015. Genetic rescue of small inbred populations: meta-analysis reveals large and

consistent benefits of gene flow. *Molecular Ecology* 24: 2610-2618.

Frankham, R., J. D. Ballou, M. D. B. Eldridge, R. C. Lacy, K. Ralls, M. R. Dudash, and C. B.

Fenster. 2011. Predicting the probability of outbreeding depression. *Conservation*

Biology 25: 465-475.

Gann, G. D., T. Mcdonald, B. Walder, J. Aronson, C. R. Nelson, J. Jonson, J. G. Hallett, et al.

2019. International principles and standards for the practice of ecological restoration.

Second edition. *Restoration Ecology* 27: S1-S46.

Glémin, S., J. Ronfort, and T. Bataillon. 2003. Patterns of inbreeding depression and architecture

of the load in subdivided populations. *Genetics* 165: 2193-2212.

Godefroid, S., C. Piazza, G. Rossi, S. Buord, A. Stevens, R. Aguraiuja, C. Cowell, et al. 2011.

How successful are plant species reintroductions? *Biological Conservation* 144: 672-682.

Hahn, M. A., and L. H. Rieseberg. 2017. Genetic admixture and heterosis may enhance the

invasiveness of common ragweed. *Evolutionary Applications* 10: 241-250.

Hamilton, J. A., and J. M. Miller. 2016. Adaptive introgression as a resource for management

and genetic conservation in a changing climate. *Conservation Biology* 30: 33-41.

Hoffmann, A. A., A. D. Miller, and A. R. Weeks. 2021. Genetic mixing for population

management: from genetic rescue to provenancing. *Evolutionary Applications*: 634-652.

Husband, B. C., and D. W. Schemske. 1996. Evolution of the magnitude and timing of

inbreeding depression in plants. *Evolution* 50: 54-70.

Isee Network. 2024. Soil Explorer http://SoilExplorer.net.

Jmp. 1989-2021. JMP Version 16. SAS Institute Inc., Cary, NC.

Jordan, R., P. A. Harrison, and M. F. Breed. 2023. The eco-evolutionary risks of not changing

seed provenancing practices in changing environments. *Ecology Letters* 24: e14348.

Koski, M. H., N. C. Layman, C. J. Prior, J. W. Busch, and L. F. Galloway. 2019. Selfing ability

and drift load evolve with range expansion. *Evolution Letters* 3: 500-512.

Li, Y., M. Stift, and M. Van Kleunen. 2018. Admixture increases performance of an invasive

plant beyond first-generation heterosis. *Journal of Ecology* 106: 1595-1606.

Lohr, J. N., and C. R. Haag. 2015. Genetic load, inbreeding depression, and hybrid vigor covary

with population size: An empricial evaluation of theoretical predictions. *Evolution* 69:

3109-3122.

Luijten, S. H., M. Kéry, J. G. B. Oostermeijer, and J. C. M. Den Nijs. 2002. Demographic

consequences of inbreeding and outbreeding in *Arnica montana*: a field experiment.

Journal of Ecology 90: 593-603.

Lynch, M., and B. Walsh. 1998. Genetics and analysis of quantitative traits, 980. Sinauer

Associates, Inc., Sunderland, MA.

Maschinski, J., S. J. Wright, S. Koptur, and E. C. Pinto-Torres. 2013. When is local the best

paradigm? Breeding history influences conservation reintroduction surivval and

population trajectories in times of extreme climate events. *Biological Conservation* 159:

277-284.

Mckay, J. K., C. E. Christian, S. Harrison, and K. J. Rice. 2005. "How local is local?" - A review

of practical and conceptual issues in the genetics of restoration. *Restoration Ecology* 13:

432-440.

Menges, E. S. 1991. Seed germination percentage increases with population size in a fragmented

prairie species. *Conservation Biology* 5: 158-164.

Menges, E. S. 1995. Factors limiting fecundity and germination in small populations of *Silene*

regia (*Caryophyllaceae*), a rare hummingbird-pollinated prairie forb. *American Midland*

Naturalist 133: 242-255.

Menges, E. S. 2008. Restoration demography and genetics of plants: when is a translocation

successful? *Australian Journal of Botany* 56: 187-196.

Menges, E. S., and R. W. Dolan. 1998. Demographic viability of populations of *Silene regia* in

midwestern prairies: relationships with fire management, genetic variation, geographic

location, population size and isolation. *Journal of Ecology* 86: 63-78.

Natureserve. 2024.

https://explorer.natureserve.org/Taxon/ELEMENT_GLOBAL.2.128777/Silene_regia.

Oakley, C. G., and A. A. Winn. 2012. Effects of population size and isolation on heterosis, mean

fitness, and inbreeding depression in a perennial plant. *New Phytologist* 196: 261-270.

682 Oakley, C. G., J. P. Spoelhof, and D. W. Schemske. 2015a. Increased heterosis in selfing

683    populations of a perennial forb. *AoB Plants* 7: plv122.

684 Oakley, C. G., J. Ågren, and D. W. Schemske. 2015b. Heterosis and outbreeding depression in

685    crosses between natural populations of *Arabidopsis thaliana*. *Heredity* 115: 73-82.

686 Oakley, C. G., S. Lundemo, J. Ågren, and D. W. Schemske. 2019. Heterosis is common and

687    inbreeding depression absent in natural populations of *Arabidopsis thaliana*. *Journal of*

688    *Evolutionary Biology* 32: 592-603.

689 Ouborg, N. J., and R. Van Treuren. 1994. The significance of genetic erosion in the process of

690    extinction. 4. Inbreeding load and heterosis in relation to population size in the mint

691    *Salvia pratensis*. *Evolution* 48: 996-1008.

692 Paland, S., and B. Schmid. 2003. Population size and the nature of genetic load in *Gentianella*

693    *germanica*. *Evolution* 57: 2242-2251.

694 Perrier, A., D. Sánchez-Castro, and Y. Willi. 2020. Expressed mutational load increases toward

695    the edge of a species' geographic range. *Evolution* 74: 1711-1723.

696 Perrier, A., D. Sánchez-Castro, and Y. Willi. 2022. Environment dependence of the expression of

697    mutation load and species range limits. *Journal of Evolutionary Biology* 35: 731-741.

698 Pickup, M., D. L. Field, D. M. Rowell, and A. G. Young. 2012. Source population characteristics

699    affect heterosis following genetic rescue of fragmented plant populations. *Proceedings of*

700    *the Royal Society B* 280: 20122058.

701 Pizza, R. B., J. Foster, and L. A. Brudvig. 2023. Where should they come from? Where should

702    they go? Several measures of seed source locality fail to predict plant establishment in

703    early prairie restorations. *Ecological Solutions and Evidence* 4: e12223.

704 Pywell, R. F., J. M. Bullock, A. Hopkins, K. J. Walker, T. H. Sparks, M. J. W. Burke, and S. Peel.

705   2002. Restoration of species-rich grassland on arable land: assessing the limiting

706   processes using a multi-site experiment. *Journal of Applied Ecology* 39: 294-309.

707 Ralls, K., J. D. Ballou, M. R. Dudash, M. D. B. Eldridge, C. B. Fenster, R. C. Lacy, P. Sunnucks,

708   and R. Frankham. 2018. Call for a paradigm shift in the genetic management of

709   fragemented populations. *Conservation Letters* 11: 1-6.

710 Rojas-Gutierrez, J. D., S. J. Mantel, and C. G. Oakley. 2026. Environment-dependent and often

711   antagonistic efects of dominance and epistasis on hetereosis in crosses between natural

712   populations. *BioRxiv*: https://doi.org/10.64898/62026.64802.64810.705147.

713 Seabloom, E. W., E. T. Borer, V. L. Boucher, R. S. Burton, K. L. Cottingham, L. Goldwasser, W.

714   K. Gram, et al. 2003. Competition, seed limitation, disturbance, and reestablishment of

715   California native annual forbs. *Ecological Applications* 13: 575-592.

716 Shaw, N., R. S. Barak, R. E. Campbell, A. Kirmer, S. Pedrini, K. Dixon, and S. Frischie. 2020.

717   Seed use in the field: delivering seeds for restoration success. *Restoration Ecology* 28:

718   S276-S285.

719 Sletvold, N. S., J. M. Grindeland, P. Zu, and J. Ågren. 2012. Strong inbreeding depression and

720   local outbreeding depression in the rewarding orchid *Gymnadenia conopsea*.

721   *Conservation Genetics* 13: 1305-1315.

722 Söderquist, L., A. Broberg, V. Rosenberg, and N. S. Sletvold. 2020. Predicting heterosis and

723   inbreeding depression from population size and density to inform management efforts.

724   *Journal of Applied Ecology* 57: 1459-1468.

725 Spigler, R. B., K. Theodorou, and S. M. Chang. 2017. Inbreeding depression and drift load in

726   small populations at demographic disequilibrium. *Evolution* 71: 81-94.

St. Clair, A. B., P. W. Dunwiddie, J. B. Fant, T. N. Kaye, and A. T. Kramer. 2020. Mixed source

populations increasses genetic diversity of restored rare plant populations. *Restoration*

Ecology 28: 583-593.

Stojanova, B., Z. Münzbergová, and H. Pánková. 2021. Inbreeding depression and heterosis vary

in space and time in the serpentinophyte perennial *Minuartia smejkalii*. *Preslia* 93: 149-

168.

Thoen, R. D., A. Southgate, G. Kiefer, R. G. Shaw, and S. Wagenius. 2025. The conservation

value of small population remnants: Variability in inbreeding depression and heterosis of

a perennial herb, the narrow-leaved purple coneflower (*Echinacea angustifolia*). *Journal*

of Heredity 116: 24-33.

Török, K., O. Valkó, and B. Deák. 2024. Ecosystem restoration with local or broad seed

provenancing: debates and perceptions in science and practice. *Biological Conservation*

293: 110535.

Van Treuren, R., R. Bijlsma, N. J. Ouborg, and W. Van Delden. 1993. The significance of genetic

erosion in the process of extinction. 4. Inbreeding depression and heterosis effects caused

by selfing and outcrossing in *Scabiosa-columbaria*. *Evolution* 47: 1669-1680.

Wagenius, S., H. H. Hangelbrook, C. E. Ridley, and R. G. Shaw. 2010. Biparental inbreeding and

interremnant mating in a perennial prairire plant: Fitness consequences in their first eight

years. *Evolution* 64: 761-771.

Weeks, A. R., C. M. Sgro, A. G. Young, R. Frankham, N. J. Mitchell, K. A. Miller, M. Byrne, et

al. 2011. Assessing the benefits and risks of translocations in changing environments: a

genetic perspective. *Evolutionary Applications* 4: 709-725.

Weisenberger, L. A., S. G. Weller, and A. K. Sakai. 2014. Remnants of populations provide

effective source material for reintroduction of an endangered Hawaiian plant, *Schiedea*

kaalae (Caryophyllaceae). *American Journal of Botany* 101: 1954-1962.

Whitlock, M. C. 2002. Selection, load and inbreeding depression in a large metapopulation.

Genetics 160: 1191-1202.

Whitlock, M. C., P. K. Ingvarsson, and T. Hatfield. 2000. Local drift load and the heterosis of

interconnected populations. *Heredity* 84: 452-457.

Willi, Y. 2013. Mutational meltdown in selfing *Arabidopsis lyrata*. *Evolution* 67: 806-815.

Willi, Y., and M. Fischer. 2005. Genetic rescue in interconnected populations of small and large

size of the self-incompatible *Ranunculus reptans*. *Heredity* 95: 437-443.

Willi, Y., J. Van Buskirk, and A. A. Hoffmann. 2006. Limits to the adaptive potential of small

populations. *Annual Review of Ecology, Evolution and Systematics* 37: 433-458.

Willi, Y., M. Van Kleunen, S. Dietrich, and M. Fischer. 2007. Genetic rescue persists beyond

first-generation outbreeding in small populations of a rare plant. *Proceedings of the Royal*

Society B-Biological Sciences 274: 2357-2364.

Table 1. ANOVA results for early fitness components for maternal and paternal populations and
their interaction and block (where appropriate). Models for both seed number per fruit and
percent germination (per cell) employed normal error distributions (denominator degrees of
freedom in parentheses). The model for juvenile survival employed a binomial error distribution.
Contrasts between cross types within a maternal population are only reported for models with a
P value < 0.1 for either the interaction between maternal population and paternal population, or
both maternal and paternal populations. For the contrasts, WIN = within-population cross, BET
= between-population cross, the subscript pool indicates that the contrast involves all individuals
of a given cross type (pooled over populations). Contrasts that start with a population code
followed by a colon contain at least one cross type specific to that maternal population.
775

	Seed number per fruit (Denom. DF = 229)		Germination (Denom. DF = 346)		Juvenile Survival	
	F	P	F	P	χ^2	P
Maternal population	17.9	**<0.001**	9.1	**<0.001**	5.6	0.060[†]
Paternal population	0.5	0.635	2.4	0.097[†]	6.0	**0.049**
Mat. Pop * Pat. Pop	0.2	0.840	0.9	0.437	14.5	**0.006**
Block	n/a	n/a	4.6	**0.001**	65.3	**<0.001**
Contrast	t	P	t	P	χ^2	P
BET$_{pool}$ vs. WIN$_{pool}$	n/a	n/a	0.3	0.736	5.7	**0.017**
Fou: BET vs. WIN$_{pool}$	n/a	n/a	3.1	**0.002**	21.4	**<0.001**
Tip: BET vs. WIN$_{pool}$	n/a	n/a	-1.0	0.312	0.7	0.408
Ver: BET vs. WIN$_{pool}$	n/a	n/a	-0.8	0.410	0.2	0.672
Fou: BET vs. WIN	n/a	n/a	1.8	0.079[†]	30.6	**<0.001**
Tip: BET vs. WIN	n/a	n/a	-1.3	0.195	0.0	-
Ver: BET vs. WIN	n/a	n/a	0.6	0.565	0.0	-

Table 2. ANOVA results for the greenhouse experiment. The model for the combined probability
of survival and reproduction employed a binomial error distribution, whereas the models for
fecundity (flowers per reproducing plant) and flowers per seedling planted (including zeros)
employed normal error distributions. Contrasts between cross types within a maternal population
are only reported for models with a P value < 0.1 for either the interaction between maternal
population and paternal population, or both maternal and paternal populations. For the contrasts,
WIN = within-population cross, BET = between-population cross, the subscript pool indicates
that the contrast involves all individuals of a given cross type (pooled over populations), and
contrasts that start with a population code followed by a colon contain at least one cross type
specific to that maternal population.

	Survival and reproduction		Fecundity (Denom. DF = 239)		Flowers per seedling planted	
	χ^2	P	F	P	F	P
Maternal population	0.8	0.675	1.4	0.250	1.6	0.194
Paternal population	1.9	0.380	0.8	0.459	1.6	0.208
Mat. Pop * Pat. Pop	3.5	0.483	2.1	0.080[†]	1.1	0.372
Block	45.0	0.554	1.3	0.142	1.2	0.202
Contrast	χ^2	P	t	P	t	P
BET$_{pool}$ vs. WIN$_{pool}$	n/a	n/a	2.5	**0.015**	n/a	n/a
Fou: BET vs. WIN$_{pool}$	n/a	n/a	1.6	0.248	n/a	n/a
Tip: BET vs. WIN$_{pool}$	n/a	n/a	1.2	0.235	n/a	n/a
Ver: BET vs. WIN$_{pool}$	n/a	n/a	3.2	**0.002**	n/a	n/a
Fou: BET vs. WIN	n/a	n/a	1.0	0.300	n/a	n/a
Tip: BET vs. WIN	n/a	n/a	0.5	0.599	n/a	n/a
Ver: BET vs. WIN	n/a	n/a	2.5	**0.014**	n/a	n/a

Table 3. ANOVA results for the field experiment. The model for the combined probability of

survival and reproduction employed a binomial error distribution, whereas the models for

fecundity (flowers per reproducing plant) and fruits per seedling planted (including zeros)

employed normal error distributions. Contrasts between cross types within a maternal population

are only reported for models with a P value < 0.1 for either the interaction between maternal

population and paternal population, or both maternal and paternal populations. For the contrasts,

WIN = within-population cross, BET = between-population cross, the subscript pool indicates

that the contrast involves all individuals of a given cross type (pooled over populations), and

contrasts that start with a population code followed by a colon contain at least one cross type

specific to that maternal population.

	Survival and reproduction		Fecundity (Denom. DF = 258)		Fruits per seedling planted	
	χ^2	P	F	P	F	P
Maternal population	10.9	**0.004**	1.1	0.313	3.3	**0.038**
Paternal population	21.7	**<0.001**	4.4	**0.014**	10.9	**<0.001**
Mat. Pop * Pat. Pop	4.7	0.315	2.1	$0.080^{\dagger}$	2.4	**0.049**
Block	23.5	**<0.001**	1.3	0.267	4.6	**<0.001**
Contrast	χ^2	P	t	P	t	P
BET$_{pool}$ vs. WIN$_{pool}$	1.3	0.263	2.4	**0.019**	2.3	**0.024**
Fou: BET vs. WIN$_{pool}$	3.7	$0.054^{\dagger}$	3.0	**0.003**	3.6	**<0.001**
Tip: BET vs. WIN$_{pool}$	0.0	0.944	1.5	0.123	0.7	0.465
Ver: BET vs. WIN$_{pool}$	0.9	0.338	1.3	0.194	1.3	0.187
Fou: BET vs. WIN	15.3	**<0.001**	1.7	$0.098^{\dagger}$	4.2	**<0.001**
Tip: BET vs. WIN	2.3	0.131	1.9	$0.067^{\dagger}$	2.0	**0.044**
Ver: BET vs. WIN	7.3	**<0.001**	0.7	0.676	-1.4	0.167

Figure 1. Mean fitness components for each maternal population (Fou, Tip, and Ver) and cross type (WIN$_{pool}$ = within-population crosses pooled over populations, BET = between-population cross) for seed number per fruit of the hand pollinations, germination, and juvenile survival. Means and standard errors were calculated as the mean of block means. Significant contrasts of BET vs. WIN$_{pool}$ are indicated where appropriate. ** $P < 0.01$, *** $P < 0.001$ (see Table 1).

Figure 2. Mean fitness components for each maternal population (Fou, Tip, and Ver) and cross type (WIN_{pool} = within-population crosses pooled over populations, BET = between-population cross) for the greenhouse experiment. Means and standard errors were calculated as the mean of block means. Significant contrasts of BET vs. WIN_{pool} are indicated where appropriate. ** $P <$ 0.01 (see Table 2).

Figure 3. Mean fitness components for each maternal population (Fou, Tip, and Ver) and cross type (WIN$_{pool}$ = within-population crosses pooled over populations, BET = between-population cross) for the field common garden experiment. Means and standard errors were calculated as the mean of block means. Significant contrasts of BET vs. WIN$_{pool}$ are indicated where appropriate. † 0.10 > P > 0.05, ** $P < 0.01$, *** $P < 0.001$ (see Table 3).

Figure S1. Mean fitness components for each maternal population (Fou, Tip, and Ver) and cross type (WIN = within-population cross, BET = between-population cross) separately by the two paternal populations for between-population crosses. Means and standard errors were calculated as the mean of block means, except for seed number per fruit which are the raw means and SE.

Figure S2. Mean fitness components for each maternal population (Fou, Tip, and Ver) and cross type (WIN = within-population cross, BET = between-population cross) for seed number per fruit of the hand pollinations, germination, and juvenile survival. Means and standard errors were calculated as the mean of block means, except for seed number per fruit which are the raw means and SE. Within a population, significant contrasts of BET vs. WIN are indicated where appropriate. † 0.10 > P > 0.05, *** $P < 0.001$ (see Table 1).

Figure S3. Mean fitness components for each maternal population (Fou, Tip, and Ver) and cross type (WIN = within-population cross, BET = between-population cross) for the greenhouse experiment. Means and standard errors were calculated as the mean of block means. Within a population, significant contrasts of BET vs. WIN are indicated where appropriate. * *P* < 0.05 (see Table 2).

Figure S4. Mean fitness components for each maternal population (Fou, Tip, and Ver) and cross type (WIN = within-population cross, BET = between-population cross) for the field common garden experiment. Means and standard errors were calculated as the mean of block means. Within a population, significant contrasts of BET vs. WIN are indicated where appropriate. † 0.10 > P > 0.05, *** *P* < 0.001 (see Table 3).

Table S1. ANOVA results for early fitness components for the effects of maternal stratification treatment (10, 30, or 60 days), maternal and paternal populations and their interaction, and block (where appropriate). Models for both seed number per fruit and percent germination (per cell) employed normal error distributions (denominator degrees of freedom in parentheses). The model for juvenile survival employed a binomial error distribution. Contrasts between cross types within a maternal population are only reported for models with a significant interaction between maternal population and paternal population or significant main effects for both of those terms. For the contrasts, WIN = within-population cross, BET = between-population cross, the subscript pool indicates that the contrast involves all individuals of a given cross type (pooled over populations), and contrasts that start with a population code followed by a colon contain at least one cross type specific to that maternal population. Model results were similar to models without incorporating effects of maternal stratification treatment (compare with Table 1) so we proceed with the simpler model throughout the results.

	Seed number per fruit (Denom. DF = 227)		Germination (Denom. DF = 344)		Juvenile Survival	
	F	P	F	P	χ^2	P
Maternal population	13.1	**<0.001**	6.4	**0.002**	5.3	$0.070^{\dagger}$
Paternal population	1.0	0.366	2.7	$0.071^{\dagger}$	6.3	**0.044**
Mat. Pop * Pat. Pop	0.3	0.877	1.1	0.345	14.6	**0.006**
Mat. Strat. Trt.	13.5	**<0.001**	2.0	0.132	0.4	0.804
Block	-	-	4.5	**0.002**	64.9	**<0.001**
Contrast	t	P	t	P	χ^2	P
BET$_{pool}$ vs. WIN$_{pool}$	-	-	0.3	0.758	6.0	**0.015**
Fou: BET vs. WIN$_{pool}$	-	-	2.8	**0.005**	21.5	**<0.001**
Tip: BET vs. WIN$_{pool}$	-	-	-1.0	0.305	0.5	0.458
Ver: BET vs. WIN$_{pool}$	-	-	-0.6	0.520	0.3	0.577
Fou: BET vs. WIN	-	-	1.9	$0.064^{\dagger}$	30.9	**<0.001**
Tip: BET vs. WIN	-	-	-1.5	0.142	0.0	-
Ver: BET vs. WIN	-	-	0.6	0.525	0.0	-

Table S2. Sample sizes and mean values (averaged over block where appropriate) for fitness components for each maternal population and cross type. For seed number per fruit, sample size indicates the number of fruits. For germination, mean is the percent germination per cell (usually starting with 10 seeds) averaged over five flats, and sample size represents the total number of cells over all flats. For juvenile survival, the mean percent survival over the means of the 30 flats is given, and sample size represents the total number of seedlings over all flats. For the greenhouse experiment, sample sizes are the total number of plants at the start of the experiment over all 48 flats, and means are the mean over flat means. For the field experiment, sample sizes are the total number of plants at the start of the experiment over all 6 blocks, and means are the mean over block means. For fecundity in both experiments, sample sizes are the number of plants that reproduced.

| | Fou | | | | Tip | | | | Ver | | | |
| | BET | | WIN | | BET | | WIN | | BET | | WIN | |
	N	Mean	N	Mean	N	Mean	N	Mean	N	Mean	N	Mean
Early												
Seed # / fruit	71	52.3	39	47.8	45	31.7	17	30.9	47	38.6	19	32.4
Germination (%)	120	71.2	60	61.0	60	53.4	30	62.2	59	55.8	30	49.9
Juv. survival (%)	723	86.9	337	73.3	307	78.9	170	78.1	291	77.5	149	79.0
Greenhouse												
Surv. and rep. (%)	118	76.0	59	86.2	59	77.5	30	91.3	59	84.7	29	85.1
Fecundity	112	44.7	55	38.9	53	42.9	28	41.1	55	54.3	26	39.5
Flowers/sdl. planted	118	34.2	59	34.0	59	33.1	30	37.5	59	46.6	29	34.2
Field												
Surv. and rep. (%)	239	44.0	120	23.3	120	34.2	60	23.3	118	39.9	59	60.9
Fecundity	105	16.8	28	12.3	41	14.0	14	8.3	47	14.1	36	11.9
Fruits/sdl. planted	239	7.7	120	3.0	120	5.2	60	2.0	118	5.9	59	8.0

Table S3. Estimates of heterosis (outbreeding depression for negative values) for each maternal population for fitness components and composite fitness estimates.

	Fou_{pool}	Tip_{pool}	Ver_{pool}	Fou	Tip	Ver	Fou*Tip	Fou*Ver	Tip*Fou	Tip*Ver	Ver*Fou	Ver*Tip
Early												
Seed # / fruit	**41**	**-14**	**4**	9	3	19	35	28	-20	1	-9	28
Germination (%)	**23**	**-7**	**-3**	17	-14	12	12	33	-13	-4	-11	16
Juv. survival (%)	**13**	**3**	**1**	19	1	-2	14	16	1	1	-2	-1
Greenhouse												
Surv. and rep. (%)	**-13**	**-11**	**-3**	-12	-15	0	-12	-9	-14	-8	2	-5
Fecundity	**12**	**8**	**36**	15	4	37	7	9	-2	19	31	34
Flowers/sdl. planted	**-3**	**-6**	**32**	1	-12	36	-6	-1	-17	9	31	29
Field												
Surv. and rep. (%)	**23**	**-5**	**11**	89	47	-34	55	23	29	-9	-13	4
Fecundity	**55**	**29**	**30**	37	69	18	39	56	22	46	31	30
Fruits/sdl. planted	**78**	**20**	**36**	157	160	-26	111	85	61	27	22	2
Cumulative												
Greenhouse	**92**	**-23**	**35**	52	-21	78	61	95	-42	8	5	90
Field	**281**	**6**	**50**	289	131	-4	262	265	14	26	-2	50