MANUSCRIPT TITLE

HIV-2 evades restriction by ZAP through adaptations in the U3 LTR region despite increased CpG levels

AUTHORS

Dorota Kmiec[1]*, Rayhane Nchioua[1], Alexander Gabel[2,3], Asimenia Vlachou[1], Sabina Ganskih[2], Sümeyye Erdemci-Evin[1], Stacey Lapp[4], Diane Carnathan[4], Steven Bosinger[4], Ben Berkhout[5], Atze T Das[5], Mathias Munschauer[2,3], Frank Kirchhoff[1]

[1]Institute of Molecular Virology
 Ulm University Medical Center
 89081 Ulm, Germany

[2]Helmholtz Institute for RNA-based Infection Research,
 ´97080 Würzburg, Germany

[3]Institute of Medical Virology
 University Hospital Frankfurt, Goethe University
 60590 Frankfurt am Main, Germany

[4]Department of Pathology and Laboratory Medicine
 School of Medicine
 Emory University
 Atlanta, GA 30322, USA

[5]Amsterdam University Medical Center,
University of Amsterdam,
Medical Microbiology and Infection prevention,
Amsterdam, Netherlands

* To whom correspondence should be addressed. Email: Dorota.Kmiec@uni-ulm.de

ABSTRACT

Simian immunodeficiency viruses infecting sooty mangabeys (SIVsmm) gave rise to nine groups of human immunodeficiency virus type 2 (HIV-2). Two of these (A and B) spread substantially with an estimated 1- 2 million individuals affected. The evolutionary adaptations that facilitated HIV-2's spread in humans are still poorly understood. Here, we report that diverse SIVsmm strains efficiently infect primary human T cells. However, they are more sensitive to interferon than HIV-2, indicating that interferon-stimulated genes (ISGs) pose a

barrier to the successful spread of SIVsmm in humans. One of the best-known antiviral ISGs is the zinc finger antiviral protein (ZAP), which targets CpG dinucleotides in RNA. To evade ZAP-mediated restriction, many viruses, including HIV-1, suppress their CpG content. Unexpectedly, we found that HIV-2 is more resistant to ZAP restriction than HIV-1 and SIVsmm despite having 33% more CpGs. Identification of ZAP binding sites using RNA eCLIP and analyses of chimeric HIV-2/SIVsmm viruses, revealed that the determinants of ZAP resistance map to the U3 region of the long-terminal repeat and promote HIV-2 replication in primary human T cells. Our results indicate that ZAP poses a barrier to SIVsmm infection in humans and that HIV-2 evolved a CpG-independent mechanism to evade it.

INTRODUCTION

Zinc-finger antiviral protein (ZAP), also known as ARTD13, PARP13, and ZC3HAV1, is a broadly expressed cellular RNA-binding protein that inhibits multiple RNA and DNA viruses (1–3). Like other antiviral restriction factors that constitute intrinsic immunity, ZAP evolved under positive selection and its expression is upregulated by interferons (IFNs) (3–6). ZAP binds to CpG dinucleotides in HIV-1 RNAs and recruits the co-factors ubiquitin ligase Tripartite Motif Containing 25 (TRIM25) and endoribonuclease KH and NYN Domain Containing (KHNYN) to degrade it, which in turn reduces viral protein expression and replication (7–10). CpGs are the least abundant dinucleotide type found in the transcriptomes of most vertebrates, including humans (2, 11, 12). ZAP exploits this CpG suppression to distinguish between self and non-self RNAs and target viral transcripts.

The continuous arms race between viruses and their hosts shapes pathogen evolution and enables the evasion of antiviral restriction factors such as ZAP. Many successful viruses mimic the CpG suppression of their host's genomes to evade the recognition by ZAP. For example, HIV-1 suppresses its genomic CpG content by over 80%, which is even lower than CpG suppression of the human transcriptome (60%) (7, 13). However, the CpG content of primate lentiviral genomes varies greatly and does not correlate with ZAP sensitivity (13), suggesting alternative ZAP evasion mechanisms.

Simian immunodeficiency virus (SIV) species naturally infect over 40 African primate species. Of those, SIVs infecting chimpanzees, gorillas and sooty mangabeys successfully crossed the interspecies barrier and gave rise to HIV-1 (SIVcpz, SIVgor) and HIV-2 (SIVsmm). While the

HIV-1 group M originating from SIVcpz has spread globally causing >95% of all HIV infections, HIV-2 infects around 1-2 million people, mostly in West Africa (14). HIV-1 and HIV-2 evolved sophisticated ways to evade or counteract human immune responses and cause persistent infection that, if untreated, eventually leads to a fatal acquired immune deficiency syndrome (AIDS). SIVsmm, the ancestor of HIV-2, does not cause disease in its natural host and is highly prevalent in sooty mangabeys, which are hunted for bushmeat or kept as pets in parts of West Africa (15). SIVsmm strains readily infect human T cells *in vitro* (16–18). Therefore, it is not surprising that SIVsmm crossed the species barrier from monkeys to humans on at least nine occasions. However, only two of these zoonotic transmissions spread among humans, giving rise to epidemic HIV-2 groups A and B. While the overall prevalence of HIV-2 is declining, new infections with HIV-2 groups A, B, and their recombined form (AB) continue to be reported outside of West Africa (19–21). It remains to be defined what type of evolutionary changes the epidemic groups of HIV-2 gained during their human adaptation.

Here, we examined whether ZAP poses a barrier to effective spread of SIVsmm in human cells. We found that SIVsmm strains are more sensitive to IFN than HIV-2 and also significantly more susceptible to inhibition by ZAP in human T cells. Surprisingly, epidemic HIV-2 evolved resistance to ZAP despite acquiring CpG dinucleotides during its adaptation to humans. Through enhanced crosslinking immunoprecipitation (eCLIP) analysis of ZAP-binding sites in viral RNA and exchanging specific genomic regions between HIV-2 and SIVsmm, we show that the factors responsible for HIV-2's evasion of ZAP do not map to the *env* gene coding region, as observed in HIV-1. Instead, they involve changes in the U3 region of the long terminal repeat (LTR), which enhance HIV-2 fitness in primary human T lymphocytes. Our results suggest that CpG-independent changes in the U3 region of the LTR contributed to the evolution of ZAP resistance and epidemic spread of HIV-2 in humans.

METHODS

Cell lines

HEK293T and HeLa cells were obtained from ATCC. HEK293T ZAP KO cells have been previously described (7). THP-1 as well as Jurkat WT and ZAP KO cells were described before (22, 23). TZM-bl cells express CD4, CCR5, and CXCR4 and contain the β-galactosidase genes under the control of the HIV promoter (24, 25).

Adherent cell lines were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 2.5% (during virus production) or 10% (at all other times) heat-inactivated foetal calf serum (FCS), 2 mM L-glutamine, 100 U/ml penicillin, and 100 µg/ml streptomycin. Jurkat cells were cultured in RPMI supplemented with 10% FCS, 2 mM L-glutamine, 100 U/ml penicillin, and 100 µg/ml streptomycin.

Isolation of primary cells

Peripheral blood mononuclear cells (PBMCs) or CD4+ T cells from healthy human donors were isolated using lymphocyte separation medium (Biocoll separating solution; Biochrom) or RosetteSep Human CD4+ T Cell Enrichment Cocktail (Stemcell), stimulated for 3 days with phytohemagglutinin (PHA; 2 µg/ml), and 10 ng/ml interleukin-2 (IL-2) in RPMI 1640 medium containing 10% fetal calf serum. Where indicated, cells were stimulated with IFNα 24h prior to infection.

Expression constructs

The pCG IRES eBFP vector encoding human ZAP-L containing an N-terminal hemagglutinin (HA) tag and empty vector control were described before (13). The infectious molecular clones of HIV-1 NL4-3, HIV-1 NL4-3 CpG high mutant, CH77TF, CH58 TF, THRO, CH167, HIV-2 7312A, ST, KR, GH123, ROD10, MCR, MCN, as well as SIVsmm PGM, SIVsmm PGM nef-IRES eGFP, L1.V1, L1.V2, L2, L3, L4 and L5 were described previously (8, 26–39). HA-ZAP-L zinc-finger 1-4 deletion mutant was generated by Q5 site-directed mutagenesis using pCG HA-ZAP-L IRES BFP as a template. The pCG mCherry-ZAP-L IRES BFP expression construct was generated by Gibson assembly using the XbaI/MluI cloning sites. SIVsmm Env and Nef expression vectors and HIV-2 7312A *env* chimeras were previously reported (16). HIV-2 7312A IRES eGFP constructs carrying premature stop codons in *vpr* and *nef* ORFs were described before (31). SIVsmm Vpr and Nef expression constructs were described before (18, 40). Chimeric HIV-2 and SIVsmm infectious molecular clones as well as CpG mutants thereof were generated using Gibson assembly (NEB) and cloned into pCR XL TOPO or pBR vector using NotI/MluI restriction sites. All constructs were cloned using primers (from biomers.net) listed in Table 1 and their correctness was confirmed by Sanger sequencing (eurofins).

Generation of sooty mangabey ZAP

Whole blood from two sooty mangabeys was collected and PBMCs were purified via Ficoll density gradient centrifugation and RNA from 1 million cells was extracted according to the manufacturer's instructions (RNeasy Mini Kit, Qiagen cat# 74104).

To generate cDNA, the SuperScript™ IV Reverse Transcriptase (cat# 18090010, Thermo Fisher Scientific) kit was used. The cDNA was cleaned using a DNA Clean and Concentrator spin column (Zymo, cat# D4033) and eluted in RNase-free water. ZAP transcript was amplified from cDNA using primer pair binding in the 5' and 3' ZAP-L UTR. Resulting PCR product was ligated into pCR XL TOPO Blunt vector. Sanger sequencing revealed 3 unique ZAP-L alleles. In the next step, ZAP with N-terminal HA tag was cloned into the pCG IRES BFP vector using XbaI/MluI restriction sites. All constructs were cloned using primers listed in Table 1 and their correctness was confirmed by Sanger sequencing (eurofins).

Transduction of Jurkat cells

Vesicular stomatitis virus glycoprotein (VSV-g)-pseudotyped virus stocks were prepared by transfecting HEK293T cells with 1 µg VSV-g and 5 µg proviral construct, followed by a medium change. Supernatants were harvested 48 h later and normalized based on infectivity as measured by the TZM-bl reporter cells. One million Jurkat WT or ZAP KO cells were seeded and treated with IFN-α (200 U/ml) or left unstimulated at the time of transduction. The input virus was removed 24h later, and cells were washed in PBS. Three and six days later, supernatants were harvested and used to infect TZM-bl reporter cells in triplicates. β-Galactosidase activity was measured 2 days later using the Gal-Screen kit (Applied Biosystems) as relative light units per second using a microplate luminometer.

Flow cytometry of infected primary cells

PHA-activated PBMCs were infected with VSV-G-pseudotyped HIV-2 or SIVsmm viruses and harvested three days post-infection for flow cytometric analysis. Cells were fixed and permeabilized (NordicMUBio, Cat#GAS-002) and then stained intracellularly for Gag (FITC conjugated; 6604665; Beckman Coulter) and ZAP (GenTex, cat# GTX120134), followed by secondary anti-rabbit AlexaFluor647 (Invitrogen, cat#A-21245). Rabbit isotype IgG antibody served as a negative control (CellSignaling cat#2729S). The MFI of infected (FITC+) cells were used to determine changes in ZAP protein levels.

Confocal microscopy

Hela cells were seeded onto poly-Lysine coated IBIDI-slides and co-transfected with 125 ng of pCG mCherry-ZAP-L expression plasmid and 125ng of HIV-2 7312A IRES eGFP provirus or empty vector control LT1 transfection reagent. 40 h post-transfection, cells were fixed in PBS containing 2% PFA and 0.05% TritonX and stained for 30min in 1µg/ml DAPI.

Lentiviral sensitivity to ZAP

To assess relative lentiviral sensitivity to ZAP overexpression, HEK293T ZAP KO cells (in 24-well format) were co-transfected using polyethyleneimine Max (PEI Max, Polysciences) transfection reagent with 600 ng of the indicated provirus and increasing concentrations (0, 100, 200, and 400 ng) of pCG HA-ZAP-L IRES BFP expression vector and the amount of DNA was normalized to 1 µg by adding pCG IRES GFP vector. Virus-containing supernatants were harvested 2 days later and used to infect 10.000 TZM-bl reporter cells in triplicates. β-Galactosidase activity was measured 2-3 days later using the Gal-Screen kit (Applied Biosystems) as relative light units per second using a microplate luminometer. Infectious virus yield values of each virus in the presence of ZAP were normalized to the corresponding GFP-only control.

Lentiviral sequence analysis, CpG mapping and RNA structure prediction

Sequences of SIVsmm and HIV-2 isolates were obtained from the Los Alamos HIV sequence database (https://www.hiv.lanl.gov/content/sequence/HIV/mainpage.html). A phylogenetic tree showing the relationship between lentiviruses was generated using full-genome sequences and the ClustalW2 simple phylogeny tool (https://www.ebi.ac.uk/Tools/phylogeny/simple_phylogeny/). CpG frequency was calculated as [number of CpG]/[sequence length], whereas CpG suppression was calculated as [number of CpG * sequence length]/[number of C * number of G]. The RNAfold web server based on the latest ViennaRNA Package (Version 2.6.3; http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi) was used to predict the minimum free energy (MFE) RNA structure, using the dynamic programming algorithm (41).

Western blotting

To examine the viral protein levels under the expression of ZAP, HEK293T ZAP KO cells were co-transfected in 12-well plates with 1.25 µg of indicated provirus and 0.25 µg DNA of pCG HA-ZAP-L IRES BFP vector or empty pCG IRES BFP vector. Two days post-transfection, cells were lysed with coimmunoprecipitation (CO-IP) buffer (150 mM NaCl, 50 mM HEPES, 5 mM EDTA, 0.10% NP-40, 0.5 mM sodium orthovanadate, 0.5 mM NaF, protease inhibitor cocktail from Roche), and cell-free virions were purified by centrifugation of cell culture supernatants through a 20% sucrose cushion at 20,800 × g for 90 min at 4°C and lysed in co-IP lysis buffer. Samples were reduced in the presence of β-mercaptoethanol by boiling at 95°C for 10 min. Proteins were separated in 4% to 12% Bis-Tris gradient acrylamide gels (Invitrogen), blotted onto a polyvinylidene difluoride (PVDF) membrane, and incubated with anti-HIV-1 Env (cat#ADP20421, CFAR), anti-HIV-2 Env (cat#ARP-1410; NIH AIDS Reagent Program), anti-HIV-1 p24 (cat#ab9071; Abcam), anti-SIVmac p27 (cat#ARP-2321; NIH AIDS Reagent Program), anti-GAPDH (Cat# 607902; BioLegend), anti-ZAP (catalogue no. GTX120134; GeneTex), anti-KHNYN (#sc-514168, Santa Cruz Biotechnology), anti-TRIM25 (cat#610570, BD), anti-Hsp90 (cat#sc7947, Santa Cruz Biotechnology) antibodies. Subsequently, blots were probed with IRDye 680RD goat anti-rabbit IgG(H+L) (catalogue no. 926-68071; LI-COR) and IRDye 800CW goat anti-mouse IgG(H+L) (catalogue no. 926-32210; LI-COR) Odyssey antibodies and scanned using a Li-Cor Odyssey reader.

Enhanced cross-linking and immunoprecipitation (eCLIP) sequencing

HEK293T ZAP KO cells (7) (8mln) were co-transfected with 30 µg of pCG HA-ZAP-L IRES BFP and 50 µg of HIV-2 7312A or SIVsmmL1.V1 proviral DNA. After 6h, medium was changed and cells were incubated at 37°C 5%CO2 for another 40h. The culture medium was removed, and cells were washed once with ice-cold PBS followed by cross-linking (254 nm and 0.8 J/cm2 UV light) on ice. Cells were harvested into ice-cold PBS and centrifuged at 400 × g at 4°C for 5 min. The cell pellet was washed once with ice-cold PBS and frozen down at -80°C. Frozen cell pellets were lysed in 50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% (vol/vol) NP-40, 0.5% sodium deoxycholate, 0.25 mM TCEP [Tris(2-carboxyethyl) phosphine hydrochloride], and complete EDTA-free protease inhibitor cocktail (Roche).

Immunoprecipitation and cDNA library preparation was performed as described in the eCLIP method (PMID: 27018577), including the following modifications: immunoprecipitates were

washed two times in 1 ml CLIP lysis buffer and two times in IP wash buffer (50 mM Tris-HCl pH 7.4, 300 mM NaCl, 1 mM EDTA, 1% [vol/vol] NP-40, 0.5% sodium deoxycholate, 0.25 mM TCEP), followed by two washes in 50 mM Tris-HCl pH 7.4, 1 mM EDTA, and 0.5% (vol/vol) NP-40. All other steps were carried out as described in the eCLIP method. Briefly, following cell lysis, unprotected RNA was digested with RNase I, and HA-tagged ZAP proteins were immunoprecipitated using a mouse monoclonal HA-antibody (Abcam, ab49969). Following immunoprecipitation, protein-RNA complexes were separated by SDS-PAGE and transferred onto a nitrocellulose membrane. Bands migrating at the expected size range were excised for immunoprecipitation (IP) and size-matched input (SMI) samples. Crosslinked RNA fragments were released by Proteinase K digestion and recovered RNA fragments were converted into a cDNA library for both IP and SMI samples by following the eCLIP procedure (PMID: 27018577). Sequencing was performed using the Illumina NextSeq 500 platform.

eCLIP data analysis

Resulting paired-end sequencing libraries with read lengths of 2 × 40 nucleotides were adapter- and quality trimmed using cutadapt (v1.18). Reads shorter than 18 nt were discarded. A custom java program was applied to identify and clip the unique molecular identifier (UMI) associated with each read. The trimmed reads were then aligned to the human (hg38, Ensembl release 110) and one of the respective viral genomes (gRNA sequence based on published full genome sequences SIVsmm: GenBank KU182922.1; HIV-2: GenBank KX174311.1) using STAR (v2.7.10a) (42) with the parameters --outFilterScoreMinOverLread 0 --outFilterMatchNminOverLread 0 --outFilterMatchNmin 0 --outFilterType Normal --alignSoftClipAtReferenceEnds No --alignSJoverhangMin 8 --alignSJDBoverhangMin 1 --outFilterMismatchNoverLmax 0.04 --scoreDelOpen −1 --alignIntronMin 20 --alignIntronMax 3000 --alignMatesGapMax 3000 --alignEndsType EndToEnd. PCR duplicates were removed with the UMI-aware deduplication functionality in Picard's MarkDuplicates. Protein binding regions were predicted by using MACS2 (43) which models read coverage enrichment in an IP sample over a paired SMI control under a Poisson distribution. The analysis was performed using the parameters: -s 31 --keep-dup all --nomodel --d-min 25 --call-summits --scale-to small --shift 25 --nolambda --extsize 5 --max-gap 20 --min-length 5. The identified MACS2 peaks were further filtered by applying a one-sided Fisher's exact test. Statistically significant enrichment was determined by calculating the odds-ratio of mapped reads within each peak

against all remaining mapped reads between IP and SMI. Multiple testing correction was applied using the Benjamini-Yekutieli (44) procedure, and only peaks with an adjusted *P* value <0.05 were considered for further analysis. Additionally, the enrichment of regions of interest, such as R, U5, and U3, was calculated by using a one-sided Fisher's exact test, following the same approach as the filtering of MACS2 peaks.

To visualize the eCLIP signal in IP relative to SMI, the relative information content of IP over SMI (45–47)was calculated as $p_i \times \log_2(p_i/q_i)$, where i represents a genomic position, p_i denotes the relative fraction of aligned reads covering that position in IP and q_i denotes the relative fraction of aligned reads covering the same position in SMI. The relative information content was visualized using the Integrative Genome Visualization (IGV) Browser.

Statistical analysis

Statistical analysis was performed using GraphPad Prism software. ANOVA, Mann-Whitney U-test or two-tailed Student's t-test were used to determine statistical significance. Significant differences are indicated as: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. Statistical parameters are specified in the Figure legends.

RESULTS

SIVsmm is highly sensitive to restriction by type I IFN in primary human cells

HIV-2 originated from and is genetically closely related to SIVsmm strains but is highly divergent from HIV-1, the main causative agent of AIDS. To compare the ability of HIV-1, HIV-2 and SIVsmm to evade human intrinsic immune responses, we analysed their replication in the presence of type I IFN in primary human peripheral blood mononuclear cells (PBMCs) containing T lymphocytes (26). To ensure that the results are representative of the lentiviral groups, we selected panels of five infectious molecular clones (IMCs) of HIV-1, including four primary transmitted-founder IMCs of subtypes B and C (CH77, CH58, THRO and CH167), five HIV-2 IMCs including the patient-derived 7312A strain, and five SIVsmm strains representing diverse viral lineages (L1-L5). In the absence of IFNα, all viruses replicated efficiently, with HIV-2 strains reaching a replication peak around day 6 and SIVsmm reaching a peak at day 8-10 (Figure 1A-B). HIV-1 strains replicated very efficiently also in the presence of IFN (2-fold reduction), while HIV-2 replication was inhibited by ~5-fold (Figure 1C). In comparison,

SIVsmm strains were more strongly suppressed by type I IFN (32-fold reduction) and generally replicated poorly in its presence. Thus, SIVsmm exhibits lower replication fitness and immune evasion than HIV-1 and HIV-2, suggesting suboptimal adaptation to type I IFN-induced antiviral restriction factors.

Figure 1   Kmiec et al., 2025

Figure 1. Replication fitness and type I IFN sensitivity of HIV-1, HIV-2 and SIVsmm in primary human cells. *(A) Replication kinetics of indicated lentiviral strain in pre-activated and IFNα-treated (500u/ml; dotted line) or untreated (solid line) human PBMCs up to 10 days post-infection. Infectious virus yield was quantified using a TZM-bl reporter assay. Shown is the mean of 3 individual donors, each tested in triplicates of infection, +/- SEM. (B) Average time needed to reach a replication peak of each virus group based on (A). Mean + SD. (C) Resistance to IFNα treatment based on (A). Mean + SD. *, P < 0.05; **, P < 0.01; calculated by Mann–Whitney test.*

HIV-2 evolved ZAP resistance

ZAP is a broadly acting antiviral restriction factor that forms an antiviral complex with TRIM25 ubiquitin ligase and endoribonuclease KHNYN to degrade CpG-containing viral transcripts. Their expression is upregulated by IFN (Figure S1A) (3, 4), which suggests that ZAP antiviral complex might contribute to the restriction of SIVsmm in human PBMCs.

To determine the contribution of ZAP to IFN susceptibility of HIV-2 and SIVsmm, we used Jurkat CCR5 CRISPR Cas9 ZAP KO cells (22). This cell line was selected due to its resemblance to human T lymphocytes, the natural target cells of HIV. The infection of control and ZAP KO Jurkat cells with HIV-1, HIV-2 and SIVsmm resulted in a detectable virus production of all strains (Figure 2A, S1B). ZAP KO modestly increased infectious HIV-1 production by 50-100% in the absence and presence of IFNα, while HIV-2 was unaffected by ZAP (Figure 2B). In contrast, ZAP KO enhanced SIVsmm infection by 3-fold in the absence of IFNα and up to 5-fold in the presence of IFNα treatment. Thus, ZAP restricts SIVsmm more efficiently than HIV-1 and HIV-2 in human T cells.

To ensure that these results were not biased by relatively low replication of SIVsmm strains in this system, we performed additional overexpression experiments in HEK293T ZAP KO cells (7). For this, we used an expression plasmid encoding the long isoform of ZAP (ZAP-L), which is the most active isoform of ZAP (2, 48, 49). Transfection of SIVsmm IMCs resulted in similar or slightly lower viral protein expression and infectious virus yield than HIV-2 (Figure S1C), in agreement with previous studies data (13, 26). On average, ZAP reduced the p27 and Env protein production of SIVsmm by ~90% but only by ~65% in the case of HIV-2 (Figures 2C-D). HIV-2 strains differed in ZAP sensitivity from being highly resistant (HIV-2 7312A, directly isolated from patient PBMCs) to being as sensitive as SIVsmm (HIV-2 ST, passaged in cell lines) (Figure 2E). In general, SIVsmm virus production was on average 3.5 times more inhibited by ZAP than that of HIV-2 (Figure 2F). Altogether, these results clearly suggest that HIV-2 evolved increased ZAP resistance during its adaptation to humans.

Figure 2. Inhibition of HIV-2 and SIVsmm by ZAP. *(A) HIV-1 (black square), HIV-2 (red circle) and SIVsmm (blue triangle) infectious virus production in parental (CTRL) or ZAP knock out (KO) CCR5-expressing Jurkat cells 6 days after transduction. (B) Infectious virus yield % calculated by dividing virus yield in the absence (KO) and presence (CTRL) of ZAP, in the absence and presence of 500u/ml IFNα treatment. (C) western blot of HEK293T ZAP KO cells and their supernatants following the transfection of empty BFP vector control or HA-tagged ZAP and indicated infectious molecular clone of SIVsmm or HIV-2, and (D) its quantification. Hsp90 serves as loading control.*

*(E) Infectious virus yield of HIV-2 and SIVsmm produced from HEK293T ZAP KO cells co-transfected with indicated infectious molecular clone and human HA-tagged ZAP. Values were normalized to infectious virus produced in the absence of ZAP expression. (F) Average infectious virus produced from HEK293T ZAP KO cells in the presence of ZAP overexpression based on (E). N=3 + SD. *, P < 0.05; **, P < 0.01; calculated using Student's t-test.*

HIV-2 evades ZAP despite higher CpG content in a species-independent manner

CpG suppression plays a key role in the evasion of ZAP restriction of many viruses (1, 2, 7, 50). For the set of HIV-2 and SIVsmm IMCs analysed, however, high CpG numbers did not correlate with increased ZAP sensitivity and even showed the opposite trend (Figure 3A). On average, HIV-2 strains contained about two times more CpGs than HIV-1 (Figure 3B). Extended analysis of the 72 available full viral genome sequences from the Los Alamos HIV database confirmed that epidemic HIV-2 A and B strains show significantly higher CpGs frequencies than SIVsmm (Figure 3C). In contrast, the CpG frequencies in rare HIV-2 F-I did not differ significantly from those of SIVsmm. Thus, epidemic HIV-2 strains evolved ZAP resistance despite an overall increase in CpG content. We have previously shown that the sensitivity of HIV-1 to ZAP is determined by the number of CpGs in the first 600 bp of the *env* ORF rather than the whole genome (13). Thus, we compared the distribution of CpGs across HIV-2 and SIVsmm genomes to identify site-specific differences in CpG content (Figure 3D). HIV-2 contained an increased number of CpGs throughout the genome. The location of CpGs in HIV-2 7312A substantially overlapped with those found in SIVsmmL1 (Figure 3E). Furthermore, HIV-2 did not decrease ZAP expression levels or affect subcellular localization (Figure S2A-C) suggesting that it evades rather than directly counteracts ZAP.

Single-stranded RNA binding domains (RBD) of ZAP orthologues from different vertebrate species exhibit varying degrees of activity and CpG-specificity (10). To determine if differences between human and sooty mangabeys ZAP activity could explain the high sensitivity of SIVsmm, we amplified ZAP from the cDNA of captive sooty mangabeys. We obtained three ZAP variants and cloned them into an expression vector. The otherwise highly conserved RNA binding domain (RBD) of sooty mangabey ZAP differed at three positions from human ZAP (Figure S3A). To determine the potential impact of these mutations on antiviral activity, we co-transfected HEK293T ZAP KO cells with ZAP and HIV-1, HIV-2 or SIVsmm proviruses. Sooty

mangabey ZAP strongly inhibited the control CpG-enriched HIV-1 mutant virus and restricted HIV-1 and SIVsmm as efficiently as human ZAP, while HIV-2 7312A was generally less affected by either orthologue (Figure 3E). Thus, sooty mangabey ZAP is as active as its human orthologue and HIV-2 evolved resistance to both, which might have reduced the selective pressure driving CpG suppression in its RNA genome.

Figure 3

Figure 3. CpG number and distribution in HIV-2 and SIVsmm genomes and the antiviral activity of human and sooty ZAP orthologues. *(A) Correlation between CpG number in HIV-2 (red) and SIVsmm (blue) strain genomes and their resistance to ZAP overexpression (related to Figure 2E). (B) Average CpG number in the tested HIV-1, HIV-2 and SIVsmm strains and (C)*

average CpG frequency in HIV-1, SIVsmm and HIV-2 (groups A-I) full-genome sequences obtained from the Los Alamos HIV sequence database. (D) Left: distribution of CpGs in the aligned HIV-2 7312A (red) and SIVsmm L1 (blue) genomes and overlap in CpG positions (grey). ORF position in the HIV-2 genome is shown for reference above the graph. Right: Quantification of the area under the curve for HIV-2, SIVsmm and overlapping peaks. (E) Infectious virus yield of HIV-1, HIV-2 and SIVsmm co-transfected with 0.4 µg DNA of human and sooty mangabey ZAP in ZAP KO HEK293T cells. N=3 + SD. ***, P < 0.001, calculated using the Mann–Whitney test.

HIV-2 RNA is bound by ZAP but resistant to its antiviral effects due to changes in the 3' half of the genome

RNA binding is essential for the antiviral activity of ZAP (3, 48). To compare ZAP binding events across the HIV-2 and SIVsmm RNA, we performed enhanced cross-linking and immunoprecipitation in combination with RNA sequencing analysis (eCLIP-seq) in HEK293T ZAP KO cells co-transfected with HIV-2 or SIVsmm proviruses and ZAP. We normalized the signal observed in ZAP immunoprecipitations relative to a size-matched input control and observed seven significantly enriched peak regions (FDR <0.05, log 2 fold change > 0.5; Table 2), all of which contained at least one CpG (Figure 4A). In the case of HIV-2 RNA, we identified nine significantly enriched ZAP-binding sites (FDR <0.05, log2 fold change > 0.5; Table 2) of which five overlapped with CpG positions (Figure 4B). In both viruses, most ZAP binding sites localized to the 3' genome half (downstream of the *vpr* gene). To analyse the impact of this region on ZAP resistance, we generated chimeras between the resistant HIV-2 7312A and sensitive SIVsmmL1. To avoid changes within the viral ORFs, we selected the *vpx/vpr* gene border (Figure 4C) to generate IMCs carrying 5' and 3' genome halves of HIV-2 or SIVsmm, respectively. Chimeric virus containing the 3' half of the HIV-2 genome was more resistant to ZAP overexpression than the virus containing the 3' half of SIVsmm (Figures 4D, S4A). Thus, ZAP sensitivity is determined by the 3` viral genome half which contains most of the ZAP binding sites.

Figure 4. Analysis of ZAP binding regions in HIV-2 and SIVsmm RNA and mapping of the determinants of ZAP resistance. *(A) Alignment of HA-ZAP eCLIP data to SIVsmmL1 and (B) HIV-2 7312A genomic RNA. Relative coverage information in IP vs. size-matched input (SMI) is calculated at each position and displayed along the viral genome (blue). ZAP binding sites significantly enriched relative to SMI are shown in dark pink. For each significantly enriched ZAP binding site, crosslinking events are extracted and shown in black (relative information in IP vs. SMI). CpG-dinucleotides are displayed in dark blue. (C) Position of ligation of the half HIV-2/ half SIVsmm virus chimeras and schematic of generated chimeric proviral constructs showing HIV-2*

(red) and SIVsmm (blue) fragments. (D) Sensitivity of those chimeras to ZAP overexpression in HEK293T ZAP KO cells. N=3+ SD.

HIV-2 evades ZAP restriction by an *env*-independent mechanism

The largest ORF found in the 3' half of the HIV-2 and SIVsmm genomes encodes for the Env protein. We have previously shown that CpG content of the *env* encoding region determines ZAP sensitivity of HIV-1 (13). However, CpG levels in the *env* genes of HIV-2 and SIVsmm did not correlate with ZAP restriction (Figure S4B). To determine if the *env* coding region contributed to the difference in ZAP sensitivity of HIV-2 and SIVsmm, we utilized the published HIV-2 7312A recombinants encoding different SIVsmm Env ectodomains (16) (Figure 5A). HIV-2 WT and all chimeras were resistant to ZAP overexpression, while SIVsmmL1 was strongly inhibited (Figure 5B). To additionally determine the contribution of *env* regions encoding for the transmembrane protein domain, we utilized the bicistonic *env* IRES eGFP reporter constructs (Figure 5C). In this system, ZAP targeting of the *env* region in the RNA transcript reduces GFP expression. Comparison of GFP signal of the control HIV-2 7312A-IRES-GFP and SIVsmm PGM-IRES-GFP reporter viruses in the absence and presence ZAP, showed significant inhibition of SIVsmm but not HIV-2 7312A, indicating that the IRES eGFP cassette itself did not change ZAP phenotype (Figure 5D, left). In contrast, ZAP decreased GFP signal of both HIV-2 and SIVsmm *env*-IRES-eGFP reporters, and there was no significant difference between them (Figure 5D, right). These results show that the determinants of ZAP sensitivity of the HIV-2/SIVsmm lineage differ from that of HIV-1 and are *env*-independent.

Figure 5. Impact of ZAP on HIV-2 Env chimeras with SIVsmm Env ectodomains and viral env transcripts. *(A) Schematic of HIV-2 7312A based Env ectodomain chimeras and (B) their sensitivity to ZAP overexpression compared to wild type HIV-2 7312A and SIVsmmL1 strains. (C) Schematic of env IRES eGFP reporter and (D) flow cytometry results showing eGFP mean fluorescence intensity in HEK293T ZAP KO cells co-transfected with ZAP or ZAP mutant without RNA binding domain (ZAP delta RBD; negative control; 100%) and either IRES eGFP reporter viruses or pCG Env IRES eGFP reporter variants. (D) N=3 + SD. *, p < 0.05; **, p < 0.01; ***, p < 0.001, calculated using Student's t-test.*

HIV-2's *nef* encoding region contributes to ZAP resistance

The accessory proteins of HIV are known to antagonize several antiviral restriction factors (51). Therefore, we tested whether the HIV-2 accessory Vpr and Nef proteins, encoded by the 3′ genome half, counteract ZAP. HIV-2 7312A with premature stop codons in *vpr* or *nef* remained resistant to ZAP (Figure 6A), indicating that these proteins do not act as ZAP antagonists.

To examine whether HIV-2 and SIVsmm *vpr* and/or *nef* encoding regions contribute to ZAP sensitivity, we utilized an extended panel of bicistronic IRES eGFP reporters. ZAP reduced eGFP expression from mRNAs containing SIVsmm *vpr* and *nef* genes significantly more efficiently than the corresponding HIV-2 constructs (Figure 6B). The reporter containing SIVsmm *nef* ORF was more efficiently inhibited (85% decrease) by ZAP than that containing *vpr* (45% decrease). Further analyses demonstrated that bicistronic IRES-eGFP constructs containing *nef* alleles from diverse SIVsmm variants are also more sensitive to ZAP restriction than the construct containing HIV-2 7312A *nef* (Figure 6C).

To determine the impact of the *nef* coding region on viral resistance to ZAP, we generated additional chimeric HIV-2 7312A/SIVsmm viruses (Figure 6D). Unfortunately, the SIVsmmL1 chimera containing HIV-2 *nef* genes were poorly infectious precluding meaningful analyses (Fig. S4A), therefore two HIV-2-based constructs were analysed instead, one carrying the *nef* of SIVsmmL1 and the second with the *nef* of SIVsmmL2. Introduction of SIVsmm *nef* region increased HIV-2's sensitivity to ZAP by 30-50% (Figure 6E). These results show that the *nef* ORF is an important determinant of the differential ZAP susceptibility of HIV-2 and SIVsmm.

Figure 6. Mapping of determinants of ZAP resistance using chimeric HIV-2/SIVsmm clones. *(A) Effect of ZAP overexpression (0.3 µg) on HIV-2 7312A wild type, reporter and mutants containing premature stop codons (*) in vpr and nef ORFs in HEK293T ZAP KO cells. (B) eGFP mean fluorescence intensity in HEK293T ZAP KO cells co-transfected with ZAP or ZAP delta RBD (negative control; 100%), and pCG IRES eGFP vectors containing the vpr, env or nef ORF from HIV-2 7312A or SIVsmm pgm or (C) nefs from HIV-2 7312A or SIVsmm L1-L5 (F). (D) Schematic of nef chimeric proviral constructs showing HIV-2 (red) and SIVsmm (blue) fragments and (E) Sensitivity of chimeric viruses (light pink) to ZAP overexpression (0.4 µg) relative to WT HIV-2 (red) and SIVsmm (blue) strains. N=3-5 + SD. *, P < 0.05; **, P < 0.01; ***, P < 0.001, calculated using Student's t-test.*

The *nef*/U3 LTR overlap region determines ZAP resistance of HIV-2

In the SIVsmm/HIV-2 lineage, the *nef* ORF overlaps with the U3 region of 3'LTR by ~350bp (Figure S5). The U3 region is duplicated at 5' and 3' ends in the proviral (DNA) form of the viral genome, but is only found at the 3' end of the genomic and subgenomic RNA (52) (53). To define a potential contribution of this region to ZAP sensitivity, we swapped the entire viral LTRs or its individual U5, R and U3 parts between HIV-2 and SIVsmm L1 (Figure 7A). The exchange of the LTRs completely reversed the ZAP phenotypes of HIV-2 and SIVsmm (Figure 6B). However, it also reduced basal infectious virus production by one to two orders of magnitude even in the absence of ZAP (Figure S6A). Exchange of only the 3' U3 or U5 regions also significantly attenuated HIV-2 7312 by 1.5 and 3.5-fold correspondingly, while mutation of the 3'R region had no significant impact (Figure S6B). The analysis of these 3'LTR chimeras showed that the exchange of the U3 region sensitizes HIV-2 to ZAP (Figure 7C). Thus, the U3 region which partially overlaps with the *nef* ORF is a major determinant of HIV-2's ZAP resistance.

Figure 7. Impact of LTRs on HIV-2's resistance to ZAP. *(A) Schematic of the generated chimeric proviruses in which either both the 5' and 3' LTRs are exchanged, or only the U3, R or U5 region of the 3'LTR. SIVsmm L1 regions are shown in blue while HIV-2 7312A regions are shown in red. (B) Sensitivity of 5'+3' LTR chimeras and (C) 3' LTR U3, R and U5 chimeras to ZAP overexpression in HEK293T ZAP KO cells. N=3 + SD. *, P < 0.05; ***, P < 0.001; calculated using Student's t-test.*

An adaptation in the U3 region of HIV-2 that mediates ZAP resistance promotes viral fitness in primary human T cells

The iCLIP analysis of viral RNA regions bound by ZAP (Figure 4A-B) identified 2 unique binding regions within the SIVsmm L1 U3 (Figure 8A). In comparison, this region was not bound in HIV-2 7312A RNA (Figure 8B). Sequence alignments revealed three CpG sites located in SIVsmmL1 but not in the HIV-2 7312A U3, two of which were found in the proximity of ZAP binding site (Figure 8C). To test whether any of these sites was specifically targeted by ZAP and responsible for differential restriction of both viruses, we generated single mutants that were depleted (SIVsmm) or enriched (HIV-2) in CpGs at the identified sites. The introduction or removal of these CpGs did not have a significant effect on viral infectivity in the absence or presence of ZAP (Figure 8D, S6C).

To map the changes that contribute to resistance to ZAP, we divided the U3 into 3 regions (Figure 8C; region A – green, B – blue and C – pink). Regions A and C contained the ZAP binding sites (ZBR) identified by iCLIP in SIVsmm U3. Region B contained no ZBR and was the most divergent region between HIV-2 and SIVsmms (Figure S7A). Previous SHAPE structure probing of the SIVmac239 RNA (54), a virus closely related and derived from SIVsmm, indicated that region B can fold a stable hairpin structure (Figure S7B), which according to our *in silico* RNA fold predictions does not form in HIV-2 (Figure S7C-D-E). To determine if changes within these regions contribute to ZAP resistance, we generated HIV-2/SIVsmm chimeras with exchanges in the U3 region A, B and C (Figure S8A-B). The introduction of SIVsmm U3 region A into HIV-2 had the most profound effect on ZAP phenotype, leading to a 3-fold decrease in ZAP resistance, while the reverse chimera in SIVsmm backbone remained highly ZAP sensitive (Figure 8C). HIV-2 with U3 region B of SIVsmm also showed a minor loss of ZAP resistance, while the corresponding SIVsmm region B mutant a minor increase in resistance (Figure 8E). In comparison, the exchange of the U3 region C had no effect on either virus. Thus, U3 region A is the main determinant of HIV-2's ZAP resistance.

To determine if the identified U3 region A associated with ZAP resistance contributes to HIV-2 fitness in primary human cells, we examined the replication of the chimeric HIV-2 and SIVsmm viruses in pre-activated human PBMCs (Figure 8F). The exchange of U3 region A reduced the replication of HIV-2 7312A by 3.5-fold, while the exchange of the other two U3 regions had no (region B) or only modest (1.3-fold; region C) effect (Figure 8G). The corresponding SIVsmmL1-based chimeras showed no significant differences in replication.

Together, these results show that ZAP binds CpG sites in SIVsmm U3 LTR and that HIV-2 7312A evolved changes in the U3 region promoting ZAP escape and replication fitness through a mechanism independent of CpG-depletion.

Figure 8. Mapping of U3 region responsible for HIV-2's resistance to ZAP and its impact on viral fitness. *(A) ZAP eCLIP data aligned to SIVsmmL1 and (B) HIV-2 7312A 3' LTR region (based on Figure 4A-B). (C) Alignment of the HIV-2 7312A and SIVsmmL1 U3 sequences showing the location of CpG dinucleotides that are exclusively found in SIVsmmL1 (CpG 1, 2 and 3) and U3 regions A (green), B (blue) and C (pink). (D) Infectious virus yield of HIV-2 and SIVsmm CpG mutants or (E) chimeric viruses carrying exchanges in the 3' U3 region A, B or C, in HEK293T ZAP KO cells transfected with ZAP. N=3-5 +SD. (F) Replication kinetics of chimeric HIV-2 and SIVsmm viruses carrying exchanges in the 3' U3 region A, B or C in human PBMCs over 12 days. (G) Normalized infectious virus yield of indicated U3 mutants as compared to WT HIV-2 7312A or SIVsmmL1 (100%) based on area under the curve analysis of data shown in panel D. Mean of 4 independent donors + SD.*

DISCUSSION

ZAP restricts the replication of many RNA and DNA viruses, often in an IFN-dependent manner (1). The ability of ZAP to preferentially target foreign RNA transcripts relies on its selective binding to CpG dinucleotides (7) which represent the least abundant type of dinucleotide in the vertebrate genomes (55). HIV-1 and many other RNA viruses strongly suppresses their CpG content and thus at least partially escape ZAP's activity (7, 50, 56–59). Here, we found that epidemic HIV-2 evolved resistance to ZAP despite increasing its CpG content by 33% compared to its zoonotic ancestor, SIVsmm. This evasion of ZAP is mediated by changes in the 3' U3 LTR region that promote HIV-2 fitness in primary human T cells. Our results show that ZAP contributes IFN mediated response against SIVsmm in human cells and show the versality of primate lentiviruses in evading ZAP-mediated antiviral immunity. Furthermore, our findings suggest that evolution of an efficient ZAP evasion strategy allows viruses to tolerate increased genomic CpG levels. Thus, our findings demonstrate that ZAP poses a barrier for zoonotic viruses, and that evolution of ZAP evasion strategies can lower the pressure driving CpG suppression in their genomes.

The link between viral CpG suppression and ZAP evasion is well established (7, 13, 50, 56–58, 60–62). However the determinants of ZAP sensitivity are complex and include not only the CpG number, but also their localization, context and local RNA structure (13, 60, 62). We have previously shown that CpG frequency in the 5' end of the *env* gene determines the sensitivity

of primary HIV-1 strains to ZAP (13). However, HIV-2 *env* plays no role in its resistance to ZAP, which underscores the divergent evolutionary paths taken by HIV-1 and HIV-2 in adapting to human immune defences and evading ZAP. In case of SIVsmm, ZAP efficiently decreased both Env and Gag levels, which are expressed from independent, spliced and unspliced viral mRNA, respectively. The broad viral transcript targeting by ZAP can be explained by the fact that all HIV-2 and SIVsmm RNA splicing products contain the untranslated 3' U3 LTR region. Analyses of chimeric HIV-2/SIVsmm recombinants and iCLIP revealed that this RNA region is specifically targeted by ZAP. Furthermore, our findings suggest that while the *nef* ORF contributes to ZAP evasion due to its overlap with 3' U3 LTR, the Nef protein does not act as a direct antagonist of ZAP. While the immunomodulatory functions of HIV-2 and SIVsmm Nefs are highly conserved (16, 63, 64), changes it its coding region can additionally promote viral adaptation to humans and resistance to antiviral restriction factors such as ZAP.

ZAP contributes to restriction of many animal-infecting RNA viruses (65–69) (66, 70, 71),. Here, we found that ZAP contributes to IFN-mediated restriction of SIVsmm in primary human T cells. Therefore, ZAP constitutes a part of the interspecies barrier protecting humans from zoonotic pathogens. ZAP targets the UTRs not only of SIVsmm but also several viral and host mRNAs (7, 72–74). It remains to be established if the UTR region is commonly recognized and targeted by ZAP in other viruses and whether pathogens with increased CpG content other than HIV-2 utilize an CpG-independent ZAP evasion strategy.

While our results identify ZAP as an effective inhibitor of SIVsmm replication in human T cells during IFN-response activation, it is not the only antiviral restriction factor that represents a cross-species barrier to this zoonotic virus. For example, it has been shown that SIVsmm is less efficient at counteracting human APOBEC3F (26), IFI16 (75) and GBP5 than HIV-2 (26, 75–77), suggesting that the latter had to evolve multiple innate immune evasion strategies to spread significantly in humans. Apart from ZAP, type I IFN upregulates the expression of hundreds of other ISGs (78–80), most of which have never been studied in the context of HIV-2 or SIVsmm restriction. Deciphering the contribution of all individual human ISGs to SIVsmm restriction would be an ambitious and time-consuming task that lies beyond the scope of the current study. Nevertheless, the generated chimeric HIV-2/SIVsmm infectious molecular clones will be highly useful for further studies investigating the innate immune mechanisms limiting SIVsmm cross-species transmission and spread in humans.

We found that the U3 region of HIV-2 accumulated changes during its human-specific adaptation that result in changes in the second half of the *nef* ORF, contribute to replicative fitness and mediate ZAP's resistance of HIV-2. The corresponding 5' part of the U3 region of SIVsmm was significantly bound by ZAP, and its introduction into HIV-2 7312A could confer partial ZAP sensitivity. It remains to be clarified how the structure of this RNA region affects the recognition by ZAP and how changes in this region affect the replication and ZAP sensitivity of other groups of HIV-2 and SIVs. Notably, the U3 LTR regions of HIV-2 isolates have extremely high genomic variability (81). Considering the presence of this region in both the unspliced RNA and all major splicing products, it is surprising that so little is known about the physiological role of the U3 region found upstream of the major Sp1 and NF-κB transcription factor binding sites (81). Our finding that this poorly-studied RNA region enables evasion of an IFN-induced antiviral restriction factor uncovers a new role of the HIV-2 U3.

SIVsmm replicates in human cells but is not well equipped to evade IFN-mediated response in humans. The contribution of ZAP to this restriction might at first seem counter-intuitive, as sooty mangabeys express a highly active form of ZAP, and SIVsmm has co-evolved with its host much longer than HIV-2. However, the difference in the antiviral immune response activation can may explain this phenomenon: sooty mangabeys infected by SIVsmm have high viral loads but do not develop the excessive immune activation and inflammation leading to a progressive CD4+ T cell loss and immunodeficiency (17, 39, 82–84). This is likely a result of a long pathogen-host co-evolution and the lack of fully functional TLR4 receptor in sooties (82). In contrast, HIV-2 infection in humans is characterized by high levels of immune activation and IFN production that upregulates the expression of ZAP and its cofactor TRIM25 (2, 9, 66, 68). Our results support a model in which epidemic HIV-2 had to evolve increased fitness and IFN resistance during its adaptation to humans. A part of this human-specific adaptation was mediated by mutations in the 3' U3 LTR region that increase replicative fitness in human T cells and allow evasion of ZAP through a CpG-independent mechanism. This adaptation reduced the selective pressure acting against CpGs in the HIV-2 genome, leading to a substantial accumulation of these rare dinucleotide over the course of its evolution.

In conclusion, we show that ZAP contributes to the IFN-mediated restriction of SIVsmm in human T cells and in part elucidate how HIV-2 evolved to evade it. By demonstrating that HIV-2's resistance to ZAP is mediated by a CpG-independent mechanism involving the 3' LTR U3

region, we identified a novel ZAP evasion mechanism, that requires further molecular characterization. It remains to be clarified whether ZAP evasion contributes to the fitness of HIV-2 *in vivo*, as well as if it removes the evolutionary constraints posed by ZAP on CpG content in lentiviral genomes. It will be interesting to test if untranslated regions of other viruses are also targeted by ZAP and if other successful zoonotic viral pathogens evolved similar evasion mechanisms. Thus, further studies on the interplay between ZAP and viral RNAs will provide insights into the virus-host arms race and the mechanisms shaping viral evolution.

ACKNOWLEDGEMENTS

We thank Regina Burger, Jana Romana Fischer, Daniela Krnavek, Martha Mayer, Birgit Ott, Kerstin Regensburger and Nicola Schrott for laboratory assistance. Prof. Paul Bieniasz and Prof. Molly Ohainle kindly provided the HEK293T ZAP KO and Jurkat ZAP KO cells. TZM-bl cells have been provided by J. C. Kappes and X. Wu and Tranzyme Inc. through the NIH AIDS Reagent Program. The p27 antibody provided by Dr. Karen Kent and Caroline Powell was obtained through the NIH HIV Reagent Program, Division of AIDS, NIAID, NIH and the Centre for AIDS Reagents, NIBSC, UK, supported by EURIPRED (EC FP7 INFRASTRUCTURES-2012 – INFRA-2012-1.1.5: Grant Number 31266). The HIV-2 Env antiserum provided by Dr. Raymond Sweet, SmithKline Beecham Pharmaceuticals was obtained through the same NIH program. We also thank Prof. Dennis Kätzel, Prof. Daniel Sauter and Prof. Konstantin Sparrer for critical reading of the article and helpful discussions.

AUTHOR CONTRIBUTIONS

Dorota Kmiec: Conceptualization, Experimental work, Formal analysis, Methodology, Validation, Writing—original draft. Rayhane Nchioua: Experimental work, Formal analysis, Methodology. Alexander Gabel: Formal analysis, Methodology. Asimenia Vlachou: Experimental work, reagent generation. Sabina Ganskih: Experimental work. Sümeyye Erdemci-Evin: Reagent generation. Stacey Lab: Reagent generation. Diane Carnathan: Reagent generation. Steven Bosinger: Reagent generation. Ben Berkhout: Formal analysis. Atze T Das: Formal analysis, Visualization. Mathias Munschauer: Formal analysis, Visualization. Frank Kirchhoff: Conceptualization, Formal analysis. All authors: writing—review & editing.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR online.

CONFLICT OF INTEREST

The authors report there are no competing interests to declare.

FUNDING

This work was supported by the European Union's Horizon 2020 research and innovation programme under Marie Sklodowska-Curie (grant agreement No. 101062524 to DK) and from Else-Krönung Fresenius Stiftung (2022_EKEA.47 to DK). FK is supported by CRC 1279, an ERC Advanced grant (Traitor viruses) and the BW foundation. Further support comes from the Helmholtz Association under the Helmholtz Young Investigator Group Program, [VH-NG-128 to MM] and the European Research Council (ERC) [ERC-StG COVIDecode, 101040914 to MM].

Funding for open access charge: Ulm University Medical Center and Else-Krönung Fresenius Stiftung.

DATA AVAILABILITY

The data generated by iCLIP has been deposited to GEO repository under the identifier GSE290608 and will be released to the public upon publication of the manuscript. The primary data underlying this article as well as generated resources will be shared on a reasonable request to the corresponding author.

REFERENCES

1. Ficarelli,M., Neil,S.J.D. and Swanson,C.M. (2021) Targeted Restriction of Viral Gene Expression and Replication by the ZAP Antiviral System. *Annu Rev Virol*, **8**, 265–283.

2. Nchioua,R., Kmiec,D., Müller,J.A., Conzelmann,C., Groß,R., Swanson,C.M., Neil,S.J.D., Stenger,S., Sauter,D., Münch,J., *et al.* (2020) SARS-CoV-2 Is Restricted by Zinc Finger Antiviral Protein despite Preadaptation to the Low-CpG Environment in Humans. *mBio*, **11**, e01930-20.

3. Gao,G., Guo,X. and Goff,S.P. (2002) Inhibition of retroviral RNA production by ZAP, a CCCH-type zinc finger protein. *Science*, **297**, 1703–1706.

4. Karki,S., Li,M.M.H., Schoggins,J.W., Tian,S., Rice,C.M. and MacDonald,M.R. (2012) Multiple Interferon Stimulated Genes Synergize with the Zinc Finger Antiviral Protein to Mediate Anti-Alphavirus Activity. *PLoS One*, **7**, e37398.

5. Kerns,J.A., Emerman,M. and Malik,H.S. (2008) Positive selection and increased antiviral activity associated with the PARP-containing isoform of human zinc-finger antiviral protein. *PLoS Genet*, **4**, e21.

6. Wang,N., Dong,Q., Li,J., Jangra,R.K., Fan,M., Brasier,A.R., Lemon,S.M., Pfeffer,L.M. and Li,K. (2010) Viral Induction of the Zinc Finger Antiviral Protein Is IRF3-dependent but NF-κB-independent. *J Biol Chem*, **285**, 6080–6090.

7. Takata,M.A., Gonçalves-Carneiro,D., Zang,T.M., Soll,S.J., York,A., Blanco-Melo,D. and Bieniasz,P.D. (2017) CG dinucleotide suppression enables antiviral defence targeting non-self RNA. *Nature*, **550**, 124–127.

8. Ficarelli,M., Wilson,H., Pedro Galão,R., Mazzon,M., Antzin-Anduetza,I., Marsh,M., Neil,S.J. and Swanson,C.M. (2019) KHNYN is essential for the zinc finger antiviral protein (ZAP) to restrict HIV-1 containing clustered CpG dinucleotides. *Elife*, **8**, e46767.

9. Li,M.M.H., Lau,Z., Cheung,P., Aguilar,E.G., Schneider,W.M., Bozzacco,L., Molina,H., Buehler,E., Takaoka,A., Rice,C.M., *et al.* (2017) TRIM25 Enhances the Antiviral Action of Zinc-Finger Antiviral Protein (ZAP). *PLOS Pathogens*, **13**, e1006145.

10. Gonçalves-Carneiro,D., Takata,M.A., Ong,H., Shilton,A. and Bieniasz,P.D. (2021) Origin and evolution of the zinc finger antiviral protein. *PLoS Pathog*, **17**, e1009545.

11. Stallings,R.L. (1992) CpG suppression in vertebrate genomes does not account for the rarity of (CpG)n microsatellite repeats. *Genomics*, **13**, 890–891.

12. Saxonov,S., Berg,P. and Brutlag,D.L. (2006) A genome-wide analysis of CpG dinucleotides in the human genome distinguishes two distinct classes of promoters. *Proc Natl Acad Sci U S A*, **103**, 1412–1417.

13. Kmiec,D., Nchioua,R., Sherrill-Mix,S., Stürzel,C.M., Heusinger,E., Braun,E., Gondim,M.V.P., Hotter,D., Sparrer,K.M.J., Hahn,B.H., *et al.* (2020) CpG Frequency in the 5' Third of the env Gene Determines Sensitivity of Primary HIV-1 Strains to the Zinc-Finger Antiviral Protein. *mBio*, **11**, e02903-19.

14. Kapoor,A.K. and Padival,S. (2024) HIV-2 Infection. In *StatPearls*. StatPearls Publishing, Treasure Island (FL).

15. Santiago,M.L., Range,F., Keele,B.F., Li,Y., Bailes,E., Bibollet-Ruche,F., Fruteau,C., Noë,R., Peeters,M., Brookfield,J.F.Y., *et al.* (2005) Simian Immunodeficiency Virus Infection in Free-Ranging Sooty Mangabeys (Cercocebus atys atys) from the Taï Forest, Côte d'Ivoire: Implications for the Origin of Epidemic Human Immunodeficiency Virus Type 2. *J Virol*, **79**, 12515–12527.

16. Heusinger,E., Deppe,K., Sette,P., Krapp,C., Kmiec,D., Kluge,S.F., Marx,P.A., Apetrei,C., Kirchhoff,F. and Sauter,D. (2018) Preadaptation of Simian Immunodeficiency Virus SIVsmm Facilitated Env-Mediated Counteraction of Human Tetherin by Human Immunodeficiency Virus Type 2. *Journal of Virology*, **92**, 10.1128/jvi.00276-18.

17. Gautam,R., Carter,A.C., Katz,N., Butler,I.F., Barnes,M., Hasegawa,A., Ratterree,M., Silvestri,G., Marx,P.A., Hirsch,V.M., *et al.* (2007) In vitro characterization of primary SIVsmm isolates belonging to different lineages. In vitro growth on rhesus macaque cells is not predictive for in vivo replication in rhesus macaques. *Virology*, **362**, 257–270.

18. Heigele,A., Kmiec,D., Regensburger,K., Langer,S., Peiffer,L., Stürzel,C.M., Sauter,D., Peeters,M., Pizzato,M., Learn,G.H., *et al.* (2016) The Potency of Nef-Mediated SERINC5 Antagonism Correlates with the Prevalence of Primate Lentiviruses in the Wild. *Cell Host & Microbe*, **20**, 381–391.

19. Ceccarelli,G., Giovanetti,M., Sagnelli,C., Ciccozzi,A., d'Ettorre,G., Angeletti,S., Borsetti,A. and Ciccozzi,M. (2021) Human Immunodeficiency Virus Type 2: The Neglected Threat. *Pathogens*, **10**, 1377.

20. Ibe,S., Yokomaku,Y., Shiino,T., Tanaka,R., Hattori,J., Fujisaki,S., Iwatani,Y., Mamiya,N., Utsumi,M., Kato,S., *et al.* (2010) HIV-2 CRF01_AB: first circulating recombinant form of HIV-2. *J Acquir Immune Defic Syndr*, **54**, 241–247.

21. Guo,H., Hu,H., Zhou,Y., Huan,X., Qiu,T., Fu,G., Lu,J. and Wang,X. (2014) The identification of a novel HIV-1 CRF01_AE/B recombinant using the near full length genome in Jiangsu Province, China. *AIDS Res Hum Retroviruses*, **30**, 1239–1242.

22. Montoya,V.R., Ready,T.M., Felton,A., Fine,S.R., OhAinle,M. and Emerman,M. A Virus-Packageable CRISPR System Identifies Host Dependency Factors Co-Opted by Multiple HIV-1 Strains. *mBio*, **14**, e00009-23.

23. OhAinle,M., Helms,L., Vermeire,J., Roesch,F., Humes,D., Basom,R., Delrow,J.J., Overbaugh,J. and Emerman,M. (2018) A virus-packageable CRISPR screen identifies host factors mediating interferon inhibition of HIV. *eLife*, **7**, e39823.

24. Platt,E.J., Wehrly,K., Kuhmann,S.E., Chesebro,B. and Kabat,D. (1998) Effects of CCR5 and CD4 cell surface concentrations on infections by macrophagetropic isolates of human immunodeficiency virus type 1. *J Virol*, **72**, 2855–2864.

25. Derdeyn,C.A., Decker,J.M., Sfakianos,J.N., Wu,X., O'Brien,W.A., Ratner,L., Kappes,J.C., Shaw,G.M. and Hunter,E. (2000) Sensitivity of human immunodeficiency virus type 1 to the fusion inhibitor T-20 is modulated by coreceptor specificity defined by the V3 loop of gp120. *J Virol*, **74**, 8358–8367.

26. Nchioua,R., Kmiec,D., Gaba,A., Stürzel,C.M., Follack,T., Patrick,S., Kirmaier,A., Johnson,W.E., Hahn,B.H., Chelico,L., *et al.* (2021) APOBEC3F Constitutes a Barrier to Successful Cross-Species Transmission of Simian Immunodeficiency Virus SIVsmm to Humans. *Journal of Virology*, **95**, 10.1128/jvi.00808-21.

27. Ochsenbauer,C., Edmonds,T.G., Ding,H., Keele,B.F., Decker,J., Salazar,M.G., Salazar-Gonzalez,J.F., Shattock,R., Haynes,B.F., Shaw,G.M., *et al.* (2012) Generation of transmitted/founder HIV-1 infectious molecular clones and characterization of their replication capacity in CD4 T lymphocytes and monocyte-derived macrophages. *J Virol*, **86**, 2715–2728.

28. Adachi,A., Gendelman,H.E., Koenig,S., Folks,T., Willey,R., Rabson,A. and Martin,M.A. (1986) Production of acquired immunodeficiency syndrome-associated retrovirus in human and nonhuman cells transfected with an infectious molecular clone. *J Virol*, **59**, 284–291.

29. Parrish,N.F., Gao,F., Li,H., Giorgi,E.E., Barbian,H.J., Parrish,E.H., Zajic,L., Iyer,S.S., Decker,J.M., Kumar,A., *et al.* (2013) Phenotypic properties of transmitted founder HIV-1. *Proc Natl Acad Sci U S A*, **110**, 6626–6633.

30. Fenton-May,A.E., Dibben,O., Emmerich,T., Ding,H., Pfafferott,K., Aasa-Chapman,M.M., Pellegrino,P., Williams,I., Cohen,M.S., Gao,F., *et al.* (2013) Relative resistance of HIV-1 founder viruses to control by interferon-alpha. *Retrovirology*, **10**, 146.

31. Hopfensperger,K., Richard,J., Stürzel,C.M., Bibollet-Ruche,F., Apps,R., Leoz,M., Plantier,J.-C., Hahn,B.H., Finzi,A., Kirchhoff,F., *et al.* (2020) Convergent Evolution of HLA-C Downmodulation in HIV-1 and HIV-2. *mBio*, **11**, e00782-20.

32. Gao,F., Yue,L., Robertson,D.L., Hill,S.C., Hui,H., Biggar,R.J., Neequaye,A.E., Whelan,T.M., Ho,D.D. and Shaw,G.M. (1994) Genetic diversity of human immunodeficiency virus type 2: evidence for distinct sequence subtypes with differences in virus biology. *Journal of Virology*, **68**, 7433–7447.

33. Kumar,P., Hui,H.X., Kappes,J.C., Haggarty,B.S., Hoxie,J.A., Arya,S.K., Shaw,G.M. and Hahn,B.H. (1990) Molecular characterization of an attenuated human immunodeficiency virus type 2 isolate. *J Virol*, **64**, 890–901.

34. Kraus,G., Radaelli,A., Talbott,R., Leavitt,M., Schmidt,A., Badel,P., Bartz,C., Morton,W., Wong-Staal,F. and Looney,D.J. (1998) Characterization of a molecular clone of HIV type 2 infectious for Macaca nemestrina. *AIDS Res Hum Retroviruses*, **14**, 65–77.

35. Ishikawa,K., Tsujimoto,H., Nakai,M., Mingle,J.A.A., Osei-Kwasi,M., Aggrey,S.E., Nettey,V.B.A., Afoakwa,S.N., Fukasawa,M., Kodama,T., *et al.* (1988) Isolation and characterization of HIV-2 from an AIDS patient in Ghana. *AIDS*, **2**, 383.

36. Clavel,F., Guétard,D., Brun-Vézinet,F., Chamaret,S., Rey,M.A., Santos-Ferreira,M.O., Laurent,A.G., Dauguet,C., Katlama,C. and Rozioux,C. (1986) Isolation of a new human retrovirus from West African patients with AIDS. *Science*, **233**, 343–346.

37. Bour,S., Akari,H., Miyagi,E. and Strebel,K. (2003) Naturally occurring amino acid substitutions in the HIV-2 ROD envelope glycoprotein regulate its ability to augment viral particle release. *Virology*, **309**, 85–98.

38. Novembre,F.J., De Rosayro,J., O'Neil,S.P., Anderson,D.C., Klumpp,S.A. and McClure,H.M. (1998) Isolation and Characterization of a Neuropathogenic Simian Immunodeficiency Virus Derived from a Sooty Mangabey. *J Virol*, **72**, 8841–8851.

39. Apetrei,C., Kaur,A., Lerche,N.W., Metzger,M., Pandrea,I., Hardcastle,J., Falkenstein,S., Bohm,R., Koehler,J., Traina-Dorge,V., *et al.* (2005) Molecular epidemiology of simian immunodeficiency virus SIVsm in U.S. primate centers unravels the origin of SIVmac and SIVstm. *J Virol*, **79**, 8991–9005.

40. Hotter,D., Krabbe,T., Reith,E., Gawanbacht,A., Rahm,N., Ayouba,A., Driessche,B.V., Lint,C.V., Peeters,M., Kirchhoff,F., *et al.* (2017) Primate lentiviruses use at least three alternative strategies to suppress NF-κB-mediated immune activation. *PLOS Pathogens*, **13**, e1006598.

41. Zuker,M. and Stiegler,P. (1981) Optimal computer folding of large RNA sequences using thermodynamics and auxiliary information. *Nucleic Acids Research*, **9**, 133–148.

42. Dobin,A., Davis,C.A., Schlesinger,F., Drenkow,J., Zaleski,C., Jha,S., Batut,P., Chaisson,M. and Gingeras,T.R. (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, **29**, 15–21.

43. Zhang,Y., Liu,T., Meyer,C.A., Eeckhoute,J., Johnson,D.S., Bernstein,B.E., Nusbaum,C., Myers,R.M., Brown,M., Li,W., *et al.* (2008) Model-based Analysis of ChIP-Seq (MACS). *Genome Biology*, **9**, R137.

44. Benjamini,Y. and Yekutieli,D. (2001) The control of the false discovery rate in multiple testing under dependency. *The Annals of Statistics*, **29**, 1165–1188.

45. Van Nostrand,E.L., Freese,P., Pratt,G.A., Wang,X., Wei,X., Xiao,R., Blue,S.M., Chen,J.-Y., Cody,N.A.L., Dominguez,D., *et al.* (2020) A large-scale binding and functional map of human RNA-binding proteins. *Nature*, **583**, 711–719.

46. Yee,B.A., Pratt,G.A., Graveley,B.R., Van Nostrand,E.L. and Yeo,G.W. (2019) RBP-Maps enables robust generation of splicing regulatory maps. *RNA*, **25**, 193–204.

47. Van Nostrand,E.L., Pratt,G.A., Yee,B.A., Wheeler,E.C., Blue,S.M., Mueller,J., Park,S.S., Garcia,K.E., Gelboin-Burkhart,C., Nguyen,T.B., *et al.* (2020) Principles of RNA processing from analysis of enhanced CLIP maps for 150 RNA binding proteins. *Genome Biology*, **21**, 90.

48. Kmiec,D., Lista,M.J., Ficarelli,M., Swanson,C.M. and Neil,S.J.D. (2021) S-farnesylation is essential for antiviral activity of the long ZAP isoform against RNA viruses with diverse replication strategies. *PLoS Pathog*, **17**, e1009726.

49. Li,M.M.H., Aguilar,E.G., Michailidis,E., Pabon,J., Park,P., Wu,X., de Jong,Y.P., Schneider,W.M., Molina,H., Rice,C.M., *et al.* (2019) Characterization of Novel Splice Variants of Zinc Finger Antiviral Protein (ZAP). *J Virol*, **93**, e00715-19.

50. Sharp,C., Thompson,B., Nash,T., Diebold,O., Pinto,R., Thorley,L., Lin,Y.-T., Sives,S., Wise,H., Hendry,S.C., *et al.* (2022) CpG dinucleotide enrichment in the influenza A virus genome as a live attenuated vaccine development strateg. 10.1101/2022.04.29.490024.

51. Sauter,D. and Kirchhoff,F. (2018) Multilayered and versatile inhibition of cellular antiviral factors by HIV and SIV accessory proteins. *Cytokine Growth Factor Rev*, **40**, 3–12.

52. Zhang,J. and Crumpacker,C. (2022) HIV UTR, LTR, and Epigenetic Immunity. *Viruses*, **14**, 1084.

53. Emery,A., Zhou,S., Pollom,E. and Swanstrom,R. (2017) Characterizing HIV-1 Splicing by Using Next-Generation Sequencing. *Journal of Virology*, **91**, 10.1128/jvi.02515-16.

54. Pollom,E., Dang,K.K., Potter,E.L., Gorelick,R.J., Burch,C.L., Weeks,K.M. and Swanstrom,R. (2013) Comparison of SIV and HIV-1 genomic RNA structures reveals impact of sequence evolution on conserved and non-conserved structural motifs. *PLoS Pathog*, **9**, e1003294.

55. Bullock,E. and Elton,R.A. (1972) Dipeptide frequencies in proteins and the CpG deficiency in vertebrate DNA. *J Mol Evol*, **1**, 315–325.

56. Lin,Y.-T., Chiweshe,S., McCormick,D., Raper,A., Wickenhagen,A., DeFillipis,V., Gaunt,E., Simmonds,P., Wilson,S.J. and Grey,F. (2020) Human cytomegalovirus evades ZAP detection by suppressing CpG dinucleotides in the major immediate early 1 gene. *PLOS Pathogens*, **16**, e1008844.

57. Nguyen,L.P., Aldana,K.S., Yang,E., Yao,Z. and Li,M.M.H. (2023) Alphavirus Evasion of Zinc Finger Antiviral Protein (ZAP) Correlates with CpG Suppression in a Specific Viral nsP2 Gene Sequence. *Viruses*, **15**, 830.

58. Shao,R., Visser,I., Fros,J.J. and Yin,X. (2024) Versatility of the Zinc-Finger Antiviral Protein (ZAP) As a Modulator of Viral Infections. *International Journal of Biological Sciences*, **20**, 4585–4600.

59. Yang,E., Nguyen,L.P., Wisherop,C.A., Kan,R.L. and Li,M.M.H. (2022) The Role of ZAP and TRIM25 RNA Binding in Restricting Viral Translation. *Front. Cell. Infect. Microbiol.*, **12**.

60. Ficarelli,M., Antzin-Anduetza,I., Hugh-White,R., Firth,A.E., Sertkaya,H., Wilson,H., Neil,S.J.D., Schulz,R. and Swanson,C.M. (2020) CpG Dinucleotides Inhibit HIV-1 Replication through Zinc Finger Antiviral Protein (ZAP)-Dependent and -Independent Mechanisms. *Journal of Virology*, **94**, 10.1128/jvi.01337-19.

61. Zimmer,M.M., Kibe,A., Rand,U., Pekarek,L., Ye,L., Buck,S., Smyth,R.P., Cicin-Sain,L. and Caliskan,N. (2021) The short isoform of the host antiviral protein ZAP acts as an inhibitor of SARS-CoV-2 programmed ribosomal frameshifting. *Nat Commun*, **12**, 7193.

62. Gonçalves-Carneiro,D., Mastrocola,E., Lei,X., DaSilva,J., Chan,Y.F. and Bieniasz,P.D. (2022) Rational attenuation of RNA viruses with zinc finger antiviral protein. *Nat Microbiol*, **7**, 1558–1567.

63. Schmökel,J., Li,H., Bailes,E., Schindler,M., Silvestri,G., Hahn,B.H., Apetrei,C. and Kirchhoff,F. (2009) Conservation of Nef function across highly diverse lineages of SIVsmm. *Retrovirology*, **6**, 36.

64. Schindler,M., Münch,J., Kutsch,O., Li,H., Santiago,M.L., Bibollet-Ruche,F., Müller-Trutwin,M.C., Novembre,F.J., Peeters,M., Courgnaud,V., *et al.* (2006) Nef-Mediated Suppression of T Cell Activation Was Lost in a Lentiviral Lineage that Gave Rise to HIV-1. *Cell*, **125**, 1055–1067.

65. Schwerk,J., Soveg,F.W., Ryan,A.P., Thomas,K.R., Hatfield,L.D., Ozarkar,S., Forero,A., Kell,A.M., Roby,J.A., So,L., *et al.* (2019) RNA-binding protein isoforms ZAP-S and ZAP-L have distinct antiviral and immune resolution functions. *Nat Immunol*, **20**, 1610–1620.

66. MacDonald,M.R., Machlin,E.S., Albin,O.R. and Levy,D.E. (2007) The Zinc Finger Antiviral Protein Acts Synergistically with an Interferon-Induced Factor for Maximal Activity against Alphaviruses. *Journal of Virology*, **81**, 13509–13518.

67. Hayakawa,S., Shiratori,S., Yamato,H., Kameyama,T., Kitatsuji,C., Kashigi,F., Goto,S., Kameoka,S., Fujikura,D., Yamada,T., *et al.* (2011) ZAPS is a potent stimulator of signaling mediated by the RNA helicase RIG-I during antiviral responses. *Nat Immunol*, **12**, 37–44.

68. Buckmaster,M.V. and Goff,S.P. (2022) Riplet Binds the Zinc Finger Antiviral Protein (ZAP) and Augments ZAP-Mediated Restriction of HIV-1. *J Virol*, **96**, e0052622.

69. Lee,H., Komano,J., Saitoh,Y., Yamaoka,S., Kozaki,T., Misawa,T., Takahama,M., Satoh,T., Takeuchi,O., Yamamoto,N., *et al.* (2013) Zinc-finger antiviral protein mediates retinoic acid inducible gene I–like receptor-independent antiviral response to murine leukemia virus. *Proceedings of the National Academy of Sciences*, **110**, 12379–12384.

70. Thibault,P.A., Ryan,A.P., Hung,C., Daugherty,M. and Lee,B. (2019) The long isoform of ZAP widely restricts Paramyxoviruses. *Access Microbiology*, **1**, 854.

71. Zhang,Y., Burke,C.W., Ryman,K.D. and Klimstra,W.B. (2007) Identification and characterization of interferon-induced proteins that inhibit alphavirus replication. *J Virol*, **81**, 11246–11255.

72. Guo,X., Carroll,J.-W.N., MacDonald,M.R., Goff,S.P. and Gao,G. (2004) The Zinc Finger Antiviral Protein Directly Binds to Specific Viral mRNAs through the CCCH Zinc Finger Motifs. *Journal of Virology*, **78**, 12781–12787.

73. Chiu,H.-P., Chiu,H., Yang,C.-F., Lee,Y.-L., Chiu,F.-L., Kuo,H.-C., Lin,R.-J. and Lin,Y.-L. (2018) Inhibition of Japanese encephalitis virus infection by the host zinc-finger antiviral protein. *PLOS Pathogens*, **14**, e1007166.

74. Todorova,T., Bock,F.J. and Chang,P. (2014) PARP13 regulates cellular mRNA post-transcriptionally and functions as a pro-apoptotic factor by destabilizing TRAILR4 transcript. *Nat Commun*, **5**, 5362.

75. Hotter,D., Bosso,M., Jønsson,K.L., Krapp,C., Stürzel,C.M., Das,A., Littwitz-Salomon,E., Berkhout,B., Russ,A., Wittmann,S., *et al.* (2019) IFI16 Targets the Transcription Factor Sp1 to Suppress HIV-1 Transcription and Latency Reactivation. *Cell Host Microbe*, **25**, 858-872.e13.

76. Krapp,C., Hotter,D., Gawanbacht,A., McLaren,P.J., Kluge,S.F., Stürzel,C.M., Mack,K., Reith,E., Engelhart,S., Ciuffi,A., *et al.* (2016) Guanylate Binding Protein (GBP) 5 Is an Interferon-Inducible Inhibitor of HIV-1 Infectivity. *Cell Host Microbe*, **19**, 504–514.

77. Braun,E., Hotter,D., Koepke,L., Zech,F., Groß,R., Sparrer,K.M.J., Müller,J.A., Pfaller,C.K., Heusinger,E., Wombacher,R., *et al.* (2019) Guanylate-Binding Proteins 2 and 5 Exert Broad Antiviral Activity by Inhibiting Furin-Mediated Processing of Viral Envelope Proteins. *Cell Rep*, **27**, 2092-2104.e10.

78. Geiss,G.K., Carter,V.S., He,Y., Kwieciszewski,B.K., Holzman,T., Korth,M.J., Lazaro,C.A., Fausto,N., Bumgarner,R.E. and Katze,M.G. (2003) Gene Expression Profiling of the Cellular Transcriptional Network Regulated by Alpha/Beta Interferon and Its Partial Attenuation by the Hepatitis C Virus Nonstructural 5A Protein. *Journal of Virology*, **77**, 6367–6375.

79. Shaw,A.E., Rihn,S.J., Mollentze,N., Wickenhagen,A., Stewart,D.G., Orton,R.J., Kuchi,S., Bakshi,S., Collados,M.R., Turnbull,M.L., *et al.* (2021) The antiviral state has shaped the CpG composition of the vertebrate interferome to avoid self-targeting. *PLOS Biology*, **19**, e3001352.

80. Samarajiwa,S.A., Forster,S., Auchettl,K. and Hertzog,P.J. (2009) INTERFEROME: the database of interferon regulated genes. *Nucleic Acids Res*, **37**, D852-857.

81. Le Hingrat,Q., Visseaux,B., Bertine,M., Chauveau,L., Schwartz,O., Collin,F., Damond,F., Matheron,S., Descamps,D., Charpentier,C., *et al.* (2020) Genetic Variability of Long Terminal Repeat Region between HIV-2 Groups Impacts Transcriptional Activity. *Journal of Virology*, **94**, 10.1128/jvi.01504-19.

82. Palesch,D., Bosinger,S.E., Tharp,G.K., Vanderford,T.H., Paiardini,M., Chahroudi,A., Johnson,Z.P., Kirchhoff,F., Hahn,B.H., Norgren,R.B., *et al.* (2018) Sooty mangabey genome sequence provides insight into AIDS resistance in a natural SIV host. *Nature*, **553**, 77–81.

83. Apetrei,C., Gautam,R., Sumpter,B., Carter,A.C., Gaufin,T., Staprans,S.I., Else,J., Barnes,M., Cao,R., Garg,S., *et al.* (2007) Virus subtype-specific features of natural simian immunodeficiency virus SIVsmm infection in sooty mangabeys. *J Virol*, **81**, 7913–7923.

84. Silvestri,G., Sodora,D.L., Koup,R.A., Paiardini,M., O'Neil,S.P., McClure,H.M., Staprans,S.I. and Feinberg,M.B. (2003) Nonpathogenic SIV Infection of Sooty Mangabeys Is Characterized by

Limited Bystander Immunopathology Despite Chronic High-Level Viremia. *Immunity*, **18**, 441–452.

TABLES

Table 1. Oligonucleotides used for cloning.

Construct	Primers used	Cloning type
pCG HA-ZAP-L ZnF1-4 deletion IRES BFP	fwd aacctgatggacagaaaggtgc rev gacccgggctcgagtggtgg	SDM
pCG mCherry-ZAP-L IRES BFP	fwd 1 tgtagaagcgcgtaggccttctagatatggtgagcaagggcgagga rev 1 ccgggtccgccttgtacagctcgtccatgcc fwd 2 gctgtacaaggcggacccggaggtgtgctg rev 2 tgtagtactccggggatccgactaactaatcacgcaggctttgtcttcagtatattcaatcacatattgtg	Gibson assembly
pCG sooty mangabey ZAP-L IRES BFP	fwd 1 gccgcggcagagccagcacagccagc rev 1 gaatggttatggcatccttctgacgc fwd 2 tgtagaagcgcgtaggccttctagatatgtacccatacgatgttccagattacgctgcggacccggaggtg rev 2 gtagtactccggggatccgacgcgtctaactaatcacgcaggg	Blunt TOPO cloning and Gibson assembly

5'HIV-2 7312A/3' SIVsmmL1.V1	fwd 1 acgccaagctatttaggtgacgcgtggaagggatttttttatagtgaaagaagacataggatattagatacatattttgag fwd 2 aggcctagcctaaaatgacagaaagacctcc rev 1 tgtcattttaggctaggcctggggggagg rev 2 ctctagatgcatgctcgagcggccgctgctagggattttcctgc		Gibson assembly
5'SIVsmmL1.V1/ 3'HIV-2 7312A	fwd 1 acgccaagctatttaggtgacgcgtggaagggatttattacag fwd 2 aggactagcataatggcagaagcagcctc rev 1 tctgccattatgctagtcctggaggggg rev 2 ctctagatgcatgctcgagcggccgctgctagggattttcctgc		Gibson assembly
HIV-2 7312A + SIVsmm L1.V1 Nef	fwd 1 aagataataggactattataagcttgaataagtattataatttaacaatacattgtaag fwd 2 tacctccaatatgggtggcattacctcc fwd 3 aacaagctgaacaggaaagcagcagcataaag rev 1 tgccacccatattggaggtacgcaaacag rev 2 gctttcctgttcagcttgtttccttcttgttag rev 3 ctctagatgcatgctcgagcccgctgctagggattttc		Gibson assembly
HIV-2 7312A + SIVsmmL2 Nef	fwd 1 aagataataggactattataagcttgaataagtattataatttaacaatacattgtaag fwd 2 tacctccaatatgggtgccagtggctcc		Gibson assembly

	fwd 3 aacaagctgaacaggaaagcagcagcataaag	
	rev 1 tggcacccatattggaggtacgcaaacag	
	rev 2 gctttcctgttcagcttgtttccttcttgtcagc	
	rev 3 ctctagatgcatgctcgagcccgctgctagggattttc	
HIV-2 7312A + SIVsmmL1.V1 LTRs and SIVsmmL1.V1 + HIV-2 7312A LTRs	fwd 1 gccaacctgctagggattttcctgc rev 1 gcaggttggcgcccgaacagggacccgag fwd 2 gactggaagggatttattacagtg rev 2 ccagtccccccttttctttttataaaatgagac fwd 3 gcaggttggcgcccgaacagggac rev 3 caacctgctagggattttcctgc	SOE PCR
HIV-2 7312A + SIVsmm L1.V1 3′ U3	fwd 1 gacttagaagaggctatagaccggttttctcttcccccccc fwd 2 aaggggggactggaagggatttattacagtg fwd 3 gctctgtattcagtcgctctgcggagaggc rev 1 atcccttccagtccccccttttctttttataaaatgagacatg rev 2 agagcgactgaatacagagcgaaatgcagttg rev 3 ctctagatgcatgctcgagcgctgctagggattttcctgcttagg	Gibson assembly
HIV-2 7312A + SIVsmm L1.V1 3′ R	fwd 1 gacttagaagaggctatagaccggttttctcttcccccccc fwd 2 gcattgtattcagtcactctgcggagaggc fwd 3 tttagaagtaagtcaagtgtgtgttcccatc rev 1 agagtgactgaatacaatgcaagaagcgggtacatttatac	Gibson assembly

	rev 2 acacttgacttacttctaaatggcagctttattgaagagg	
	rev 3 ctctagatgcatgctcgagcgctgctagggattttcctgc	
HIV-2 7312A + SIVsmm L1.V1 3' U5	fwd 1 gacttagaagaggctatagaccggtttttctcttcccccccc	Gibson assembly
	fwd 2 attagaagcaagctagtgtgtgttcccatc	
	rev 1 cacactagcttgcttctaattggcagctttattaagagg	
	rev 2 ctctagatgcatgctcgagcgctgctagggattttcctgc	
HIV-2 7312A + SIVsmmL1.V1 U3 fragment A	fwd 1 gacttagaagaggctatagaccggtttttctcttccccc	Gibson assembly
	rev 1 tcactgtaataaatcccttccagtcccc	
	fwd 2 gaagggatttattacagtgaaaggagacataaaatattag	
	rev 2 actcttctgggtacttaacaaatgccttg	
	fwd 3 tgttaagtacccagaagagtttgggtatcagtcaggattac	
	rev 3 ctctagatgcatgctcgagcgcggccgctgctagggat	
HIV-2 7312A + SIVsmmL1.V1 U3 fragment B	fwd 1 gacttagaagaggctatagaccggtttttctcttccccc	Gibson assembly
	rev 1 tgacatactaccaaactcttctgggaacc	
	fwd 2 aagagtttggtagtatgtcaggattgcc	
	rev 2 cagtgtcagcttgtttccttcttgttagc	
	fwd 3 aaggaaacaagctgacactgcagggactttcc	
	rev 3 ctctagatgcatgctcgagcgcggccgctgctagggat	
HIV-2 7312A + SIVsmmL1.V1 U3 fragment C	fwd 1 gacttagaagaggctatagaccggtttttctcttccccc	Gibson assembly
	rev 1 ccctgctgtctcagccagtttcctttcttatg	

	fwd 2 aactggctgagacagcagggactttccac		
	rev 2 actgaatacagagcgaaatgcagttgtatttatac		
	fwd 3 catttcgctctgtattcagtcgctctgcggagag		
	rev 3 ctctagatgcatgctcgagcgcggccgctgctagggat		
SIVsmmL1.V1 + HIV-2 7312A U3 region A	fwd cagtgagaccaagagtccccttaagaaccatgacatacaaattgg	1	Gibson assembly
	rev 1 tcactataaaaaatcccttccagtcccc		
	fwd gaagggattttttatagtgaaagaagacataggatattagatac	2	
	rev 2 ctcttctgggaacctgttgaaggcctcatc		
	fwd 3 tcaacaggttcccagaagagtttggtag		
	fwd 3 gcttaattaagtttaaacgctgctagggattttcctgc		
SIVsmmL1.V1 + HIV-2 7312A U3 region B	fwd cagtgagaccaagagtccccttaagaaccatgacatacaaattg	1	Gibson assembly
	rev 1 tgactgatacccaaactcttctgggtac		
	fwd 2 aagagtttgggtatcagtcaggattacc		
	rev 2 ctgtctcagccagtttcctttcttatgcag		
	fwd 3 aaggaaactggctgagacagcagggactttcc		
	rev 3 gcttaattaagtttaaacgctgctagggattttcctgc		

SIVsmmL1.V1 + HIV-2 7312A U3 region C	fwd 1 cagtgagaccaagagtccccttaagaaccatgacatacaaattg rev 1 ccctgcagtgtcagcttgtttccttcttg fwd 2 aacaagctgacactgcagggactttccag rev 2 actgaatacaatgcaagaagcgggtacatttatac fwd 3 cttcttgcattgtattcagtcactctgc rev 3 gcttaattaagtttaaacgctgctagggattttcctgc		Gibson assembly
HIV-2 7312A + U3 CpG 1	fwd 1 gacttagaagaggctatagaccggttttctcttccccc rev 1 gatgggtctcctcttcctcccgggttac fwd 2 gaagaggagacccatcgtctagtgcacccagcacag rev 2 ctctagatgcatgctcgagcgcggccgctgctagggat		Gibson assembly
SIVsmmL1.V1 - U3 CpG 1	fwd 1 cagtgagaccaagagtccccttaagaaccatgacatacaaattgg rev 1 caatgtgtctcatcttcctgagcttcatctg fwd 2 gaagatgagacacattgcttgatgcatccagcac rev 2 gcttaattaagtttaaacgctgctagggattttcctgc		Gibson assembly
HIV-2 7312A + U3 CpG 2	fwd 1 gacttagaagaggctatagaccggttttctcttcccccccc rev 1 tgcctcttgcgtttagtctagccttccattccttctc fwd 2 tagactaaacgcaagaggcatacctacagag rev 2 ctctagatgcatgctcgagcggccgctgctagggattttcctgcttag		Gibson assembly

SIVsmmL1.V1 - U3 CpG 2	fwd cagtgagaccaagagtcccccttaagaaccatgacatacaaattg	1	Gibson assembly
	rev 1 ggcctcttgctgttagccttctctttacc		
	fwd 2 aaggctaacagcaagaggcctttttaaaaatg		
	rev 2 gcttaattaagtttaaacgcggccgctgctagggattttcctgc		
HIV-2 7312A + U3 CpG 3	fwd 1 gacttagaagaggctatagaccggtttttctcttccccccc	1	Gibson assembly
	rev 1 aatacaatgcgagaagcgggtacatttatac		
	fwd 2 cccgcttctcgcattgtattcagtcgctctg		
	rev ctctagatgcatgctcgagcggccgctgctagggattttcctgcttag	2	
SIVsmmL1.V1 - U3 CpG 3	fwd cagtgagaccaagagtcccccttaagaaccatgacatacaaattg	1	Gibson assembly
	rev 1 aatacagagcaaaatgcagttgtatttatacag		
	fwd 2 actgcattttgctctgtattcagtcactc		
	rev 2 gcttaattaagtttaaacgcggccgctgctagggattttcctgc		

Table 2: Significant peaks identified by eCLIP

SIVsmmL1 ZAP binding peaks												
	start	end	count IP	count SM	noPeak. IP	no Peak. SM	log2FC (IP/SM)	Odds Ratio	p. value	p.adj	score	Log pval
1	416	454	851	42	146869	10870	0.55	1.50	0.004	0.0315	1.50	2.37
2	2707	2723	269	3	147451	10909	2.32	6.63	5.872 e-6	1.145 e-4	3.94	5.23
3	6797	6830	492	20	147228	10892	0.80	1.82	0.002	0.028	1.55	2.53
4	7339	7368	673	29	147047	10883	0.73	1.72	0.001	0.016	1.79	2.85

	start	end	count IP	count SM	noPeak. IP	no Peak. SM	log2FC (IP/SM)	Odds Ratio	p. value	p.adj	score	Log pval
5	8410	8528	2887	143	144833	10769	0.57	1.50	3.660 e-7	1.684 e-5	4.77	6.44
6	9080	9226	2573	125	145147	10787	0.60	1.53	5.761 e-7	1.684 e-5	4.77	6.24
7	9416	9447	755	32	146965	10880	0.76	1.75	5.535 e-4	0.008	2.09	3.26

HIV-2 7312A ZAP binding peaks

	start	end	count IP	count SM	noPeak. IP	no Peak. SM	log2FC (IP/SM)	Odds Ratio	p. value	p.adj	score	Log pval
1	695	724	3117	59	1863543	97117	1.44	2.75	4.771 e-20	1.049 e-18	17.98	19.32
2	762	777	1350	30	1865310	97146	1.18	2.34	8.810 e-8	1.210 e-6	5.92	7.06
3	1318	1334	765	16	1865895	97160	1.23	2.49	2.270 e-5	2.480 e-4	3.61	4.64
4	5801	5857	9830	346	1856830	96830	0.56	1.48	1.495 e-14	2.739 e-13	12.56	13.83
5	6367	6376	1018	32	1865642	97144	0.69	1.66	0.002	0.012	1.90	2.79
6	6798	6899	17477	638	1849183	96538	0.51	1.43	5.384 e-21	1.480 e-19	18.83	20.26
7	7598	7694	17556	408	1849104	96768	1.16	2.25	1.669 e-76	9.177 e-75	74.04	75.78
8	7917	7981	14717	491	1851943	96685	1.44	2.75	4.771 e-20	1.049 e-18	17.98	19.32
9	8774	8821	19858	291	1846802	96885	1.18	2.34	8.810 e-8	1.210 e-6	5.92	7.06