TITLE

Reactive Oxygen Detoxification Contributes to *Mycobacterium abscessus* Antibiotic Survival

AUTHORS

Nicholas A. Bates[1,2], Ronald Rodriguez[3,4], Rama Drwich[1], Abigail Ray[5], Sarah A. Stanley[3],

Bennett H. Penn[1]*

AFFILIATIONS:

1. Department of Internal Medicine, University of California, Davis, Davis, California, United States

of America

2. Graduate Group in Immunology, University of California, Davis, Davis, California, United States

of America

3. Department of Molecular & Cell Biology, University of California, Berkeley, California, USA

4. Department of Plant & Microbial Biology, University of California, Berkeley, California, USA

5. Microbiology Graduate Group, University of California, Davis, Davis, California, United States

of America

*Corresponding author

Email: bhpenn@ucdavis.edu

ABSTRACT

It has long been recognized that when a population of bacteria are exposed to a bactericidal

antibiotic the majority of the population rapidly dies, but that a sub-population of cells, known as

"persister cells" can survive for prolonged periods in a non-growing, but viable, state and can

resume replication upon removal of the antibiotic. Persister cell frequency is dramatically

increased by stress such as nutrient deprivation, but it is unclear how what pathways enable

survival and how the process is regulated and enhanced by environmental stress. Mycobacterial

persisters are particularly resilient, with the ability to endure many weeks of antibiotic exposure.

To identify the genetic determinants of persister cell survival we carried out a transposon

mutagenesis coupled with high-throughput sequencing (Tn-Seq) screen in *Mycobacterium*

abscessus (*Mabs*). We identified a number of pathways contributing to persister cell survival,

including the bacterial proteasome and enzymes needed to detoxify reactive oxygen species

(ROS). We showed that *MAB_2470c*, encoding the KatG catalase/peroxidase was required for

both spontaneous and starvation-induced persister cell formation in response to multiple

antibiotics. Consistent with a role for ROS in antibiotic-mediated cell death we observed an

accumulation of ROS following antibiotic exposure. We tested the role of oxygen in bacterial

death, and found that that hypoxic conditions significantly impaired bacterial killing, and notably,

in the absence of oxygen KatG became dispensable. Taken together, these findings support a

model where the lethality of some antibiotics are enhanced by the accumulation of ROS.

INTRODUCTION

A key goal of antibiotic therapy is eradication of the pathogenic bacteria. While many common

infections respond well to antibiotics, and high cure rates are achieved with 1-2 weeks of therapy[1,

2], there are also infections where bacterial eradication is either very slow, or frequently incomplete.

This challenge is exemplified by mycobacterial infections, where treatment courses extend from

months to years to prevent relapse.

While the ability of mycobacteria to escape antibiotic-mediated killing is multifactorial, the

phenomenon of antibiotic "persistence" is likely to be an important contributor[3-6]. Studies on

penicillin dating from the 1940s noted that while the majority of a susceptible bacterial population

died rapidly over a few hours, a small sub-population of 'persisters' remained viable for over a

week[7]. Importantly, these persister cells had not acquired a mutation conferring heritable antibiotic

resistance, and do not grow in the presence of the antibiotic. Rather, they have entered into a

readily-reversable epigenetic state[8-10] where despite antibiotic-mediated inhibition of critical

processes, they are able to survive. Virtually all bacteria display the ability to form persister cells,

and persister cell development is strongly induced in response to stresses such as nutrient

deprivation or acidic pH[6, 7, 11, 12]. Notably, these same stresses are encountered in the lysosome

of an activated immune cell[13], and studies of pathogens isolated from activated macrophages

indeed show a strong immune-mediated increase in persister cells[5, 14]. Thus, paradoxically, the

immune system may actually impede bacterial eradication by antibiotics.

Persister formation has been studied extensively in model systems such as *Escherichia coli* ,

providing important insights, but also highlighting some of the uncertainties of current models for

this process. Several different pathways have been implicated in persister formation in *E. coli*

including the HipBA toxin-antitoxin system[15] that regulates translation, ppGpp synthesis by

RelA/SpoT enzymes[16, 17], and Lon protease[16]. In each of these models, the postulated effect is to

arrest cell division and render the process targeted by antibiotics non-essential. However,

important questions remain unanswered. It is unclear how the process is regulated and induced

by stress. It is also unclear how some bacterial cells survive when critical processes such as

transcription or translation are arrested by antibiotics.

Even the mechanism of cell death following antibiotic exposure itself remains uncertain, and

somewhat controversial. Historically, antibiotics have been assumed to kill bacteria as a direct

result of inhibition of their target molecules. For example, β-lactam antibiotics directly disrupt cell

wall integrity leading to mechanical cell lysis[18]. However, a number of studies have suggested

that reactive oxygen species (ROS) accumulation triggered by antibiotic stress downstream of

target inhibition might also contribute to cell death[10,19, 20]. However, other studies have found no

such association[21, 22], leaving the role of ROS in antibiotic-mediated cell death unresolved.

Studying persister cell physiology in mycobacterial pathogens offers several advantages.

Mycobacterial persister cells are particularly resilient, as *Mycobacterium smegmatis* (*Msmeg*) and

Mycobacterium tuberculosis (Mtb) persisters can endure many weeks of antibiotic exposure[10, 23,]

[24], and clinically, mycobacterial infections are difficult to eradicate. Fully-susceptible *Mtb* requires

multiple antibiotics for four months or longer[25 26], whereas non-tuberculous mycobacteria typically

require treatment for 12-18 months, and have a relapse rates of roughly 50%[27]. Treatment of

Mycobacterium abscessus (*Mabs*) is particularly challenging. In addition to the possibility of

forming persister cells, it is intrinsically resistant to many classes of antibiotics, leaving few

treatment options. This leads to the use of antibiotics with greater toxicity to patients, and a need

to use these agents for prolonged periods to prevent relapse[28].

Previous genetic screens have studied antibiotic responses in mycobacteria, with some

evaluating heritable resistance, and others investigating persister cell formation. Several studies

of genetic resistance have successfully used either transposon mutagenesis with high-throughput

sequencing of insertion sites (Tn-Seq) or CRISPR-based transcriptional repression with high-

throughput sequencing of guide RNAs (CRISPRi) to identify genes promoting growth in sub-

inhibitory concentrations of antibiotic. These studies have provided insights such as the

importance of cell membrane permeability controlling antibiotic penetration into the cytoplasm[29-]

[31].

The reversable state of persister cell formation has proven challenging to study, likely due to the

effects of widespread bacterial cell death during the experiment, which creates population

bottlenecks that obscure mutant phenotypes[32, 33 ,34.] There has been one highly-effective Tn-Seq

study of rifampin-specific antibiotic tolerance in *Mtb* that successfully isolated over 100 mutants[19],

demonstrating the feasibility of genetic screening in this context. However, whether these

phenotypes seen with rifampin in *Mtb* extend to other organisms and other antibiotics remains to

be determined.

Here, we study the process of persister cell formation in *Mabs*, and describe the results of a

genome-wide genetic screen seeking to identify genes required for both spontaneous and

starvation-induced persister cell formation in *Mabs*. We identified a number of pathways

contributing to persister cell survival and identified the KatG catalase-peroxidase ROS detoxifying

enzyme as conferring the ability to survive for prolonged periods in several antibiotics, providing

support for the idea that reactive oxygen species can indeed contribute to bacterial cell death

from antibiotics.

RESULTS

Starvation-induced persister cell formation in mycobacteria

We first sought to develop conditions suitable for genetic analysis of antibiotic tolerance in

mycobacteria. Genetic screens examining persister cell formation face two inherent obstacles.

First, cell death during the experiment creates population bottlenecks that obscure mutant

phenotypes. In addition, most mycobacterial populations contain spontaneous drug-resistant

mutants that can expand if the population is exposed to a single antibiotic. To overcome these

obstacles, we sought to establish large-scale, high-density culture conditions that still maintained

antibiotic potency.

We began by assessing the feasibility of this approach using wild-type *Msmeg,* exposing the cells

to either a combination of two antibiotics routinely used to treat *Mabs* infections, tigecycline and

linezolid (TIG/LZD)[27, 35], or to the combination of rifampin, isoniazid, and ethambutol

(RIF/INH/EMB) used to treat *Mtb*. We first determined the minimum-inhibitory concentrations

(MICs) and minimum bactericidal concentrations (MBCs) of cultures at the standard density used

in clinical laboratories for mycobacterial drug-susceptibility testing versus the high-density

conditions that would be needed for genetic analysis of antibiotic persister cells. We established

the MIC and MBC at the standard 5E5 colony-forming units (CFU) per ml[36] for wild-type *Msmeg*,

and compared this to the MICs of high-density cultures (3E8 CFU/ml) under fully aerated

conditions. As expected, the MIC and MBC for both combinations were significantly elevated at

high bacterial densities, however, both antibiotic combinations retained sufficient potency to

reduce the bacterial population >1000-fold within 72h (Figure S1A; Figure 1A). We then evaluated

both spontaneous and stress-induced persister cell formation under these conditions in *Msmeg*.

We compared logarithmically growing (mid-log) cultures in rich 7H9 media to cultures starved for

2 days in PBS prior to antibiotic exposure. Consistent with expectations, we found a marked

increase in the frequency of persister cells in starved cultures, with a 100-fold increase in survival

following TIG/LZD exposure and a 10,000-fold increase following RIF/INH/EMB exposure.

We next examined two additional species of pathogenic mycobacteria to assess starvation-

induced persister formation under these high-density conditions. Prior work demonstrated that

multiple stresses, including starvation, induces persister cell formation in both *Mtb* and *Mabs*[37,38,

[39,40]. We again compared cells starved in PBS to mid-log cells growing 7H9, and again found that

cultures of wild-type *Mabs* (ATCC 19977 strain) and *Mtb* (Erdman) also displayed dramatic

increases in the frequency of persister cells in nutrient-deprived cultures (Figure 1B,C). Notably,

for *Mabs* and *Msmeg,* the development of these persister cells required an adaptation period of

several days in nutrient-limited conditions, as formation of persister cells was dramatically

impaired if cells were shifted immediately into starvation conditions with antibiotics, suggesting a

regulated process needed to be completed, and persister cells were not simply slow growing cells

(Figure S2).

Identification of pathways needed for persister formation in *Mabs*

We next used these conditions to carry out a Tn-Seq screen in *Mabs* to identify genes necessary

for forming both spontaneous and starvation-induced persister cells in response to clinically-

relevant antibiotics (TIG/LZD). We conducted the screen using a *Mabs Himar1* Tn library library[29]

comprised of ~55,000 mutations across all 4,992 non-essential *Mabs* genes in strain 19977. We

maintained cells in log-phase growth in 7H9 rich media, or starved cells in PBS as described

above, and then exposed them to TIG/LZD for 6 days, a point at which spontaneous persister

cells comprise the majority of the population and there is minimal further loss in viability of

antibiotic treated mid-log cultures (Figure 1C). Following antibiotic exposure cells were then

washed and placed in antibiotic-free liquid media to recover, genomic DNA was isolated, and Tn

insertion sites sequenced. We then used TRANSIT software[41] to quantify the abundance of each

Tn mutant across different conditions and identify mutants with statistically-significant differences

in distribution in order to identify mutations altering the ability of *Mabs* to form persister cells

(Figure 2A). We identified 277 *Mabs* genes required for surviving TIG/LZD exposure in rich media,

271 genes required for survival during starvation and 362 genes required to survive the combined

exposure to antibiotics and starvation (FDR < 0.05, Log2 fold-change >0.5; Figure 2B-E). Of the

genes required for persister survival, roughly 60% were required in both nutrient-replete and

starvation states, although condition-specific survival determinants were also seen (Figure 2F).

We identified several discrete pathways involved in spontaneous persister cell formation in *Mabs*

when log-phase cultures in rich media were exposed to antibiotics. As expected, we identified

genes with already-established functions in antibiotic responses, including MAB_0984c,

MAB_2752, MAB_2753 which are all homologs of known antibiotic transporters in *Mtb,* and the

tetracycline-responsive transcription factors MAB_4687 and MAB_0314c (Table S1).

Identification of multiple genes known to be involved in bacterial responses to antibiotic

demonstrates an ability of Tn-Seq to identify physiologically relevant pathways in this context.

Interestingly, although cells were exposed to translation-inhibiting antibiotics, and no exogenous

oxidative or nitrosative stress was applied to the cells, we identified a number of factors needed

to combat these stresses and their consequences. This includes *bfrB* (bactoferritin), *aphe*

(peroxiredoxin) and *katG* (catalase/peroxidase) as well as 5 components of the bacterial

proteasome pathway, known to mediate survival in *Mtb* during nitrosative stress. We also

identified members of DNA-damage response pathways including *recF* and *recG* and *uvrA*, *uvrB*

and *uvrC* (Figure 2G). Examining starvation-induced persisters, a number of the same pathways

were again seen. The mutant with the greatest persister defect in this context was *mntH*, a redox-

regulated Mn/Zn transporter implicated in peroxide resistance in other organisms[42, 43] (Table S1).

Taken together, these findings suggest an unexpected scenario whereby translation inhibition

triggers accumulation of reactive oxygen or nitrogen species and damage to macromolecules

such as DNA was occurring.

We next sought to independently confirm the role in antibiotic persistence for a set of genes

identified in the Tn-Seq screen. We selected a set of high-priority genes with strong defects in

persister formation and used oligonucleotide-mediated recombineering (ORBIT)[44] to disrupt their

open reading frames. The initial genes we selected were *pafA* (proteasome pathway), *katG*

(catalase/peroxidase), *recR* (DNA repair), *blaR* (β-lactam sensing), and *MAB1456c*

(cobalamin synthesis). To control for non-specific effects of antibiotic selection during the

recombineering process, we used the same methods to create a parallel strain targeting a non-

coding intergenic region downstream of a redundant tRNA gene (MAB_t5030c). We then

individually screened each of these mutants to determine if they displayed deficits in persister cell

formation. We exposed cells to TIG/LZD, either in rich 7H9 media or under starvation conditions,

as had been done in the as in the pooled Tn-Seq screen. For Δ*katG* we detected clear defects in

persister cell formation following 6 days of antibiotic exposure in either starved or starvation-

induced conditions, and for Δ*pafA* there was a trend toward impairment under starved conditions

(Figure 3A,B).

To confirm the role of *katG* and *pafA* and exclude off-target or polar effects, we performed genetic

complementation analysis by restoring a wild-type copy of each gene into the respective Δ*pafA*

and Δ*katG* mutants. In each case, we integrated a single copy of the wild-type gene, under the

control of its endogenous promoter, into the genome at the L5 *attB* site (hereafter *pafA+, katG+*

strains), and constructed isogenic control strains with empty vector integrated at the same site

(hereafter *pafA-, katG-* strains). We confirmed expression of the re-introduced copy of each gene

by RT-qPCR in the *pafA+*, and *katG+* strains, and found expression within roughly 2-fold of

endogenous wild-type levels (Figure 4A,B). In rich media, the *katG+* strain had a significant, albeit

incomplete restoration of persister survival, with a roughly 50-fold increase relative to the *katG-*

strain following antibiotic exposure. When exposed to antibiotics under starvation conditions,

where the *katG-* defect is more severe, the *katG+* strain showed a near-complete restoration of

survival relative to control cells and over 1000-fold greater than *pafA-* cells (Figure 4C). Analysis

of *pafA* mutants also showed restoration of survival in *pafA+* cells relative to *pafA-* cells although

the overall magnitude of the defect in *pafA-* cells was smaller than in *katG-* cells. We next

evaluated whether the *pafA-*, and *katG-* strains were overall more sensitive to the growth inhibitory

effects of TIG/LZD, or whether they had more specific defects in survival. We performed MIC

determination for TIG and LZD individually for each strain, comparing the *katG+*/*katG-* and

pafA+/*pafA-* strains. We found that that the MICs for each of these strains were unchanged,

demonstrating that these mutants were not more readily inhibited by these antibiotics. Instead,

they have more rapid kinetics of cell death at concentrations above the MIC, consistent with a

specific defect in persister cell formation or survival (Figure 4A,B).

Reactive oxygen contributes to antibiotic lethality in *Mabs*

We next investigated the role of KatG in persister cell formation more broadly. We assessed

whether *katG* conferred protection from other antibiotics with diverse mechanisms of action, and

evaluated a panel of antibiotics are used to treat mycobacterial infections. Because *katG-* mutants

showed the greatest defects in starvation-induced persistence, we analyzed survival of *katG+* and

katG- strains in starvation-adapted cultures and challenged cells with a panel of antibiotics.

Because both TIG and LZD both act by inhibiting translation, we began by exposing cells to either

TIG or LZD alone. As expected, the degree of bacterial killing was significantly less with either

agent alone than when they are added in combination. Upon exposure to either of these antibiotics

the *katG-* cells died more rapidly than *katG+* cells, though we note that the final proportion of

persisters in the population was unchanged in *katG-* cells (Figure 5A). When we exposed cells to

rifabutin (RFB), an RNA polymerase inhibitor, we saw a similar effect, with a 100-fold loss of

viability in *katG-* cells relative to the *katG+* cells. In contrast, when we exposed cultures to either

levofloxacin (gyrase inhibitor) or cefoxitin (β-lactam inhibitor of peptidoglycan cross-linking), *katG*

had little to no effect on cell viability (Figure 5A). Thus, the role of KatG is context-dependent

suggesting that some antibiotics generate oxidative stress that can be ameliorated by KatG

catalase/peroxidase activity while others do not.

We next assessed whether ROS is playing a role in antibiotic-mediated cell-death in *Mabs*. The

identification of *katG* as essential for persister cells to survive exposure to TIG/LZD, suggests that

ROS are present and leading to toxicity. Although TIG/LIN are translation inhibitors that do not

directly generate ROS we evaluated whether they might nonetheless be triggering ROS

accumulation as a secondary effect, and if the decreased survivability of *katG* mutants might be

attributable to ROS. We began by examining ROS levels in *katG+ Mabs* using the ROS indicator

dye cellROX, that is retained in cells when it becomes oxidized[45]. At baseline, during log-phase

growth in rich media < 3% of cells had ROS accumulation (Figure 5B). We saw a modest increase

in ROS accumulation in starved cultures, with roughly 10% of the population cellROX+. However,

when starved cultures were exposed to the TIG/LZD we saw a dramatic increase in ROS

accumulation with over 30% cellROX+ cells. Taken together, these data indicate that translation

inhibition does indeed have unanticipated downstream effects on cellular redox balance, with

ROS accumulation that could be contributing to cell death.

We next sought to test whether ROS were contributing to bacterial cell death by reducing their

generation and assessing the impact on cell survival. Under normal aerobic conditions, a major

source of ROS is oxidative phosphorylation, as superoxide, hydrogen peroxide, and hydroxyl

radicals are natural byproducts of the reduction of O_2 to H_2O. We tested the hypothesis that

oxidative phosphorylation is the source of ROS in antibiotic-treated cells by shifting cultures to

hypoxia, in order to block cellular respiration and associated ROS generation. A well-established

system for studying hypoxia in mycobacteria is the Wayne Model of gradual-onset hypoxia

whereby low-density cultures are inoculated in sealed vessels with minimal headspace. As the

culture slowly grows, the soluble O_2 in the culture is consumed, resulting in a slow onset of hypoxia

over several days, a process that can be monitored by the decolorization of Methylene Blue dye

in the media[46]. Under aerobic conditions in rich media, we observed the expected rapid killing of

Mabs over the first 5 days with the combination of TIG/LZD, with more rapid loss of viability in

KatG- cells (Figure 5C). However under hypoxic conditions, where oxidative phosphorylation is

blocked and ROS production suppressed we saw much slower bacterial killing. Importantly, under

hypoxic conditions *katG-* cells were no longer defective in persister formation relative to *katG+*

cells, consistent with the hypothesis that translation-inhibiting antibiotics also cause secondary

accumulation of lethal ROS in antibiotic treated cells.

We next tested the role of KatG in *Mabs* clinical strains to determine how prevalent the role of

katG in persister formation is amongst different *Mabs* strains. We obtained 2 clinical strains of

Mabs, and used ORBIT to target the *katG* locus, and again targeted the tRNA-proximal intergenic

site as a negative control. We then evaluated the ability of these strains to form both spontaneous

and starvation-induced persister cells upon exposure to TIG/LZD. For clinical strain 1, the Δ*katG*

mutant showed no defects in either spontaneous or stress-induced persister cell formation (Figure

6B). In contrast, for clinical strain 2, *katG* contributed to starvation-induced persistence, as Δ*katG*

displayed dramatic defects in survival (Figure 6C). Interestingly, this phenotype only manifest

during starvation, and was not seen in the absence of stress. Thus, loss of *katG* displays

incomplete penetrance, and the degree of protection conferred by KatG falls along a continuum

among *Mabs* strains, with strains showing no impairment in Δ*katG* cells, some showing defects

in both spontaneous and stress-induced persister survival, and some showing context-dependent

defects. Taken together these results show that antibiotics do trigger ROS accumulation in *Mabs*,

and that detoxification factors like KatG contribute to persister cell survival.

DISCUSSION:

Pathways necessary for persister formation in Mabs. The phenomenon of bacterial persister

cell formation has been recognized since the 1940s, and has been observed in a broad array of

bacterial species. Despite this, a unifying mechanism of persister cell formation has not emerged,

suggesting that different pathways may play a role in different contexts. A large body of work

comes from *E. coli* where toxin-antitoxin systems such as HipBA[15], Lon protease[16, 47] and the

ppGpp synthase RelA all contribute to the formation of persister cells [48-51]. However, even within

this species other mechanisms seem to function, as RelA contributes to persister cell formation

following exposure to β-lactams, but not quinolones[52]. In addition, how exactly these pathways

promoter persister formation remains unclear. While ppGpp produced by RelA seems to act

through Lon protease[16], the critical substrates in this process are note unknown.

It is also unclear how mechanisms identified in one bacterial species may relate to the

mechanisms in other species. *RelA* plays a role in diverse species of bacteria, as it is essential

for persister cell formation in *Pseudomonas aeruginosa*[53]*, Staphylococcus aureus* [54], and *Mtb*[55]

[19]. However, the role of this pathway does not seem to be universal, as deletion of neither *relA*

nor *lon* had an effect on persister formation in *Msmeg* [56]. Conversely, other pathways have been

identified as critical in other organisms, including a reduction ATP synthesis in *S. aureus* [57].

In our dataset we identified a number of pathways contributing to persister cell formation following

exposure to translation-inhibiting antibiotics. This includes multiple members of the bacterial

proteasome, proteins involved in reactive oxygen and nitrogen species, and DNA repair pathways.

Interestingly, we did not identify *relA*, however, a prior study of the *relA Mabs* mutant demonstrates

that this strain still synthesizes ppGpp, suggesting possible genetic redundancy and a need to

disrupt one or more additional genes in a *relA* mutant background in order to study the role of

ppGpp synthesis in *Mabs* persister cell formation.

Mechanisms of antibiotic lethality

The mechanisms of bacterial cell death following antibiotic exposure itself remain somewhat

controversial. The most straightforward explanation is that antibiotics kill by directly disrupting the

function of their target molecule. For example, in this model cell-wall stability inhibitors lead to

direct mechanical lysis, and translation inhibitors act by depleting one or more essential proteins

below a critical threshold[21,22,58]. However, more recently it was postulated that antibiotics kill

through a final common pathway of lethal ROS generation, leading to oxidative damage to DNA

and other macromolecules[59]. Since that time, there have been studies both supporting[19,10,20] and

refuting the role of ROS[21, 22] in antibiotic-mediated cell death in a variety of organisms.

Our analysis of *Mabs* provides new insights into the role of ROS in antibiotic mediated killing.

Consistent with a role for ROS generation contributing to lethality, we find elevated ROS levels in

bacteria exposed to translation inhibitors. In addition, using Tn-Seq, we identified the *katG*

catalase/peroxidase gene as contributing to persister cell viability in both log-phase and nutrient-

starved cells. When oxygen is removed from the cultures, not only is antibiotic killing slowed, but

importantly, the *katG* mutant no longer displays a survival defect. Taken together, these data

demonstrate that in *Mabs*, translation inhibitors such as TIG or LIN do indeed have an indirect

downstream effect triggering ROS accumulation, and that KatG-mediated detoxification of these

ROS is important for survival. These findings are supported by other published studies. In *Mabs*

other groups have also observed reduced antibiotic-mediated killing in hypoxic conditions[37]. In

Mtb, exposure to rifampin also generates ROS[19, 60], and *katG* is found in Tn-Seq datasets

examining rifampin-treated cells[19].

The source of ROS in this setting is uncertain. In principle, any of several derangements could

lead to ROS accumulation. One of the major sources of cellular ROS is oxidative phosphorylation,

as hydrogen peroxide, superoxide, and hydroxyl radicals are natural byproducts of the reduction

of O_2 to H_2O by the electron transport chain. Thus, increased ROS generation by the electron

transport chain is an attractive hypothesis. Alternatively, particularly under starvation conditions,

it is possible that antioxidant and ROS scavenging metabolites may become depleted, thus

accelerating oxidative damage. Our Tn-Seq analysis provides additional insight on this. We noted

a small class of Tn mutants that were better able to survive antibiotic exposure (Figure 2B).

Prominent among this class of mutants were several independent components of the NADH

dehydrogenase complex. Also known as Complex I of the electron transport chain, this complex

is one of the entry points for electrons into the oxidative phosphorylation pathway. The observation

that mutants in this complex are protected suggests that decreasing flux through oxidative

phosphorylation, with a concomitant decrease in ROS generation, may enhance survival during

antibiotic exposure.

However, antibiotic-induced ROS accumulation is not a universal phenomenon. Other antibiotics

we examined in *Mabs* do not trigger ROS accumulation, and loss of *katG* had no effect on survival

in these antibiotics. In addition, ROS detoxification does not always play a protective role, as loss

of catalase and peroxidase activities in E. coli had no effect on persister survival[22]. Taken together,

this suggests a model in which antibiotics cause direct toxicity by acting on their target, but in

addition some antibiotics, notably transcription inhibitors and translation inhibitors, also appear to

have a secondary toxic effect of causing ROS accumulation. How exactly transcription or

translation blockade lead to increased ROS levels is currently unclear and will require further

investigation, although our data suggest that the electron transport chain plays an important role.

Therapeutic implications

Mabs infections are particularly challenging to eradicate, with relapse rates over 50%[28]. Our

results highlight several bacterial processes such as the bacterial proteasome or ROS

detoxification that might targeted therapeutically to reduce the development or survival of persister

cells. Agents targeting these process might not have any antimicrobial activity alone, but might

act to target the unique physiology of persister cells that develop upon antibiotic exposure. This

would represent a new therapeutic class of "persistence inhibitors" that might act cooperatively

with traditional antibiotics to eliminate the subpopulation of persister cells that would otherwise

survive prolonged antibiotic exposure.

Limitations

Tn-Seq screens have inherent drawbacks, including an inability to identify mutants in essential genes, or in cases of genetic redundancy. Thus, there are likely genes needed for spontaneous or starvation-induced persister cell formation in *Mabs* that were not identified in this study. In addition, we studied a exposure to a single class of antibiotic, focusing on the translation inhibitors often used to treat *Mabs* infections, and we studied only spontaneous and starvation-induced persister cells. It is likely that studying other antibiotics, with different mechanisms of action, or different stresses that induce persister cells would identify additional genes contributing to persister formation and would allow better identification of pathways that might be shared amongst different classes of antibiotic.

MATERIALS AND METHODS

Bacterial Strains and Culture conditions:

Mabs ATCC 19977 and clinical *Mabs* strains were grown in BD Middlebrook 7H9 media (liquid) or 7H10 media (solid) supplemented with 0.5% glycerol (Sigma) and 0.2% Tween-80 (Sigma). Sacramento Clinical strains were obtained from the Sacramento County Department of Public health mycobacteriology laboratory. Identity of strains as *Mabs* was confirmed by amplification of the 16s locus and Sanger sequencing. *Msmeg* (MC2 155) was grown in the same media as *Mabs*. *Mtb* Erdman was grown in 7H9 (liquid) or 7H10 (solid) supplemented with 0.5% glycerol, 0.1% Tween-80, and 10% OADC (BD). All cultures were grown at 37C and liquid cultures were shaken. With all liquid cultures, cultures were grown with 90% container headspace or using a gas permeable cap to ensure culture oxygenation. PBS starvation was achieved by washing OD0.5-1.0 *Mabs* 1X in DPBS (-Ca/Mg) and resuspending in DPBS supplemented with 0.1% tyloxapol (Sigma).

Mabs antibiotic killing experiments:

For PBS starvation experiments, freezer stocks of *Mabs* were thawed and grown for 48 hours in

log phase, then PBS starved or passaged in log phase for 48h. OD0.5-1.0 or PBS starved *Mabs*

were then resuspended in antibiotic containing media at OD1.0. For experiments with hypoxia,

Mabs was adjusted to OD0.001 in media with 1.5ug/ml methylene blue and added to a rubber

septum sealed glass vial with 50% headspace. These cultures were grown for 5d, and methylene

blue discoloration was observed at d3. Antibiotics were used at the following concentrations:

tigecycline (Chem-Impex) at 10ug/ml (8 fold above MIC, re-administered every 3 days), linezolid

(Chem-Impex) at 100ug/ml (20 fold above MIC), levofloxacin (Sigma) at 40ug/ml (8 fold above

MIC), cefoxitin (Sigma) at 80ug/ml (8 fold above MIC, re-administered every 3 days), and rifabutin

(Cayman) at 40ug/ml (4 fold above MIC). After antibiotic administration, colony forming units over

time were measured.

Msmeg antibiotic killing experiments:

Freezer stocks of *Msmeg* were streaked to agar plates, grown at 37C until colonies formed (3

days), then stored at room temperature for up to one additional week. Individual colonies were

picked and grown for 48 hours in log phase before being PBS starved or passaged in log phase

for 48h. OD0.5-1.0 or PBS starved *Msmeg* were then resuspended in antibiotic containing media

at OD1.0. Antibiotics were used at the following concentrations: tigecycline (Chem-Impex) at

1.25ug/ml (8 fold above MIC, re-administered every 3 days), linezolid (Chem-Impex) at 2.5ug/ml

(8 fold above MIC), rifampin (Sigma) at 32ug/ml (8 fold above MIC, re-administered every 6 days),

isoniazid (Sigma) at 32ug/ml (8 fold above MIC, re-administered every 6 days), and ethambutol

(Thermo) at 4ug/ml (8 fold above MIC, re-administered every 3 days). After antibiotic

administration, colony forming units over time were measured.

MTB antibiotic killing experiments:

Freezer stocks of *MTB* were thawed and grown for 5-7 days in log phase before being starved for

14d or longer. Non-starved control bugs were thawed such that they were also grown for 5-7 days

in log phase before experimental use. OD0.5-1.0 or PBS starved *MTB* was then resuspended in

antibiotic containing media and adjusted to OD1.0. Antibiotics were used at the following

concentrations: rifampin (Sigma) at 0.1ug/ml (4 fold above MIC, re-administered every 6 days),

isoniazid (Sigma) at 0.1ug/ml (4 fold above MIC, re-administered every 6 days), and ethambutol

(Thermo) at 8ug/ml (4 fold above MIC, re-administered every 6 days). After antibiotic

administration, colony forming units over time were measured.

Transposon insertion sequencing:

The transposon (Tn) library for Mabs was constructed as previously described for other

mycobacterial species[41]. Screening was done by growing a freezer stock of the library for 2.5

days in log phase before 48 hour PBS starvation or continued log phase growth. Samples were

then resuspended in tigecycline/linezolid or DMSO solvent containing media and left for 6 days

(with a re-administration of tigecycline or matching DMSO on day 3). The samples were then

washed 2X in antibiotic free liquid media and grown until OD0.5-1.0. Subsequently, the samples

underwent three more rounds of 100-fold passaging in liquid media before the samples were

collected in trizol (Invitrogen). A sample taken at the time of the commencement of PBS starvation

was collected in trizol and used as the input control. Three independent trials of this experiment

were run to achieve triplicate data. DNA was extracted and submitted to the UC Davis DNA

Technologies Core, where Himar1 transposon containing sequences were amplified and

sequenced on Illumina AVITI. Sequence reads were mapped to the ATCC 19977 genome and

analyzed using TRANSIT software. The resampling method was used with the following

parameters: 0% of N/C termini ignored, 10,000 samples, TTR normalization, LOESS correction,

include sites with all zeros, site restricted resampling. For follow-up analysis, adjusted p value

<0.05 and log2 fold change <-0.5 (essential) or >0.5 (less essential) was used a cutoff for

statistical significance. Mabs gene orthology to Mtb was done with BLASTp using an e-value

cutoff of 0.1. Mabs genes and MTB genes that mapped to each other using independent one-way

BLASTp searches were considered orthologs. For pathway analysis, gene lists (*MTB* orthologs)

were imported into the DAVID knowledgebase and pathway enrichment analysis was done for

biological processes with Gene Ontology and Uniprot key words as well as KEGG pathways.

Duplicated pathways were filtered out and the top pathways by adjusted p value were reported.

Gene Deletion and complementation:

Knockout strains were generated using ORBIT {Murphy, 2018 #325}. Briefly, *Mabs* was

transformed with the kanR ORBIT recombineering plasmid pkm444. 20ml Mab at OD0.5-1.0 was

washed 2X in 10% glycerol and resuspended in 200ul 10% glycerol. 500ng plasmid was added

and electroporated at 2.5kV in 0.2cm cuvettes. The bacteria were allowed to recover overnight

before plating on 150ug/ml kan plates. Clones were selected and regrown in liquid media

supplemented with 150ug/ml kan and 10% OADC (BD) to OD0.5-1.0. The pkm444-*Mabs* was

then diluted to OD0.1 and 200mM glycine (sigma) was added to the media. 16 hours later, 500mM

sucrose (Sigma) and 500ng/ml aTC (Cayman) were added and incubated for an additional 4

hours. Subsequently, the Mab was washed 2X in ice cold 10% glycerol + 500mM sucrose. 200ul

of 10X concentrated *Mabs* was then electroporated with 600ng of the ZeoR ORBIT payload

plasmid pkm496 and 2ug of targeting oligonucleotide (supplementary table 3) at 2.5kV in 0.2cm

cuvettes. The Mab was then allowed to recover overnight in liquid media with 10% OADC and

500ng/ml aTC and plated on 150ug/ml zeocin plates. Mutants were then selected and screened

for gene deletion by PCR amplification and confirmed by sanger sequencing. For genetic

complementation, knockout strains were complemented with PCR amplified WT copies of their

knocked-out gene. Error-free KatG and PafA PCR amplification and vector integration was

confirmed by sanger sequencing. At least 100bp of upstream and downstream sequence was

also included, allowing for retention of the endogenous promoter and terminator. Since *katG* is

known to be transcribed as part of an operon, the upstream gene FurA was also included in the

complementation construct to achieve maximal KatG expression. The mycobacterial integrating

vector pmv306 was used and cloning was done by SLIC. Subsequently, complementation was

confirmed by RNA extraction and RT-qPCR.

MIC plates

Two-fold serial dilutions of antibiotics were prepared in a 96 well plate in 100ul volume. 100ul of

2X bacteria were added (for Mab: used a final OD of 0.001, M. smegmatis: OD0.001, MTB:

OD0.01), making a final volume of 200ul. The plates were incubated until there was visible growth

in the no antibiotic control well. At this time, MICs were assessed visually, and the bacteria were

transferred to a new plate with 20ul of 40% PFA. OD620 measurements were taken in a FilterMax

F3 plate reader (Molecular Devices).

Flow cytometry

OD1 Mab was stained with cellROX green (Thermo) at a final concentration of 5mM for 30min at

37C. The cells were then washed in PBS and resuspended in PBS with 4% PFA and 5ug/ml DAPI.

The samples were run on a LSRII flow cytometer (BD) and analyzed with FlowJo software (BD).

FIGURE LEGENDS

Figure 1. Nutrient starvation induces antibiotic persister cell formation across

mycobacteria. (A) *M. smegmatis* (MC^2 155)*, (B) *M. tuberculosis* (Erdman)*, or (C) *M. abscessus*

(ATCC 19977) were starved in PBS or grown in 7H9 rich media. The indicated antibiotics were

added: Isoniazid (INH), rifampin (RIF), ethambutol (EMB) tigecycline (TIG) and linezolid (LZD).

Surviving colony forming units (CFUs) are shown. Error bars represent SEM, statistical

significance is calculated at each time point using student's t test. ****: p<0.0001, ***: p<0.001,

**: p<0.01, *: p<0.05, ns: p>0.05. Data are combined from 3 independent experiments.

Figure 2. *Mabs* Tn-seq identifies genes essential for persister cell survival. (A)

Experimental design (B-E) Relative abundance of individual genes under the indicted conditions

using Tn-Seq. For B-D comparisons are made to the input log-phase population. The additional

comparison in E is made to PBS starvation. When antibiotics are not added an equal amount of

DMSO solvent is added. Cultures were fully aerated. (F) Distribution of genes essential for each

indicated condition relative to the input population (FDR < 0.05, Log2 fold-change >0.5). (G)

Pathway analysis of the essential genes in each condition, showing significantly enriched

pathways using the NIH DAVID knowledgebase. Screens were run as 3 independent

experiments and data is from the combined results.

Figure 3. Independent gene deletions of *katG* and *pafA* confirm Tn-Seq results. Survival

over time for the indicated genes following exposure to TIG/LZD either in 7H9 rich media or

following starvation for 48h in PBS. ORBIT recombineering mutants were made in ATCC 19977

to disrupt (A) *katG* (Catalase-peroxidase), (B) *pafA* (Pup protein ligase), (C) *MAB1456c/cobA*

(cobalamin synthesis), *blaR* (β-lactam sensing), *recR* (Recombination protein RecR). A-B are

combined data from 4 independent experiments. C-E is representative data from 2 independent

experiments. Error bars represent SEM, statistical significance is calculated at each time point

using student's t test. ****: p<0.0001, ***: p<0.001, **: p<0.01, *: p<0.05, ns: p>0.05.

Figure 4. Complementation analysis *katG* and *pafA* mutants confirms their role in

persister survival. Analysis of *katG- (ΔkatG::pmv306), katG+ (ΔkatG::pmv306 katG), pafA-*

*(ΔpafA::pmv306), pafA+ (pafA::pmv306 pafA)*and control strain (intergenic

site*::zeoR::pmv306).* MIC determination for (A) *katG+/katG-* and (B) *pafA+/pafA-* strains.

Survival over time for (C) *katG+/katG-* and (D) *pafA+/pafA-* strains. Error bars represent

standard deviation, mean expression is displayed above the bar for each group, data are 3

technical replicates from one experiment. ****: p<0.0001, ***: p<0.001, **: p<0.01, *: p<0.05, ns:

p>0.05.

Figure 5. Context-specific accumulation of ROS and role for *katG*. (A) Analysis of

katG+/katG- cells when challenged with a panel of antibiotics. Cells were starved in PBS for 48h

to induce persister cells and then exposed to the indicated antibiotic at 4x MIC. (B) Flow

cytometry of DAPI and cellROX stained *katG$^+$ Mabs* after 72h in each condition are shown.

Percentage of cells in the cellROX+ gate are shown for each plot. (C) Persister survival over

time for aerated and hypoxic and cultures of *Mabs* comparing *katG+/katG-* cells after treatment

with TIG/LIN are shown. Statistical significance is calculated between antibiotic treated

katG+/katG- cells *Mabs.* Error bars represent SEM, statistical significance is calculated at each

time point using student's t test. ****: p<0.0001, ***: p<0.001, **: p<0.01, *: p<0.05, ns: p>0.05.

Figure 6. Variable penetrance of *katG* phenotype among *Mabs* strains Δ*katG* and control

strains were generated in the indicated strain backgrounds using ORBIT recombineering: (A)

ATCC 19977, (B) Sacramento County clinical strain #1 and (C) clinical strain #2. Error bars

represent SEM, statistical significance is calculated at each time point using student's t test.

****: p<0.0001, ***: p<0.001, **: p<0.01, *: p<0.05, ns: p>0.05.

Figure S1. Antibiotic concentration and culture density. (A-E) *Msmeg* was grown to log

phase, concentration adjusted to either 3e7 cells/ml or 5e5 cells/ml, and treated with antibiotics

across the indicated range of concentrations. OD measurements after 36 hours of growth are

shown for (A) Tigecycline, (B) Linezolid, (C) Isoniazid, (D) Rifampin, and (E) Ethambutol. The

antibiotic concentrations used in subsequent experiments are labelled with an arrow.

Figure S2. Adaption to starvation prior to antibiotic administration is required for

persister cell induction. (A) *Msmeg,* (B) *Mtb*, or (C) *Mabs* were starved in PBS or grown in

7H9 rich media. At day 0, all the indicated cultures were changed to PBS with antibiotics and kill

kinetics are shown. Antibiotics in nutrient rich media are included as a control. Error bars

represent SEM, statistical significance was calculated between PBS+abx conditions at each

time point using student's t test. ****: $p<0.0001$, ***: $p<0.001$, **: $p<0.01$, *: $p<0.05$, ns: $p>0.05$.

All graphs are combined from 3 independent experiments.

Table S1. Tn-Seq results

Table S2. DAVID functional pathway analysis

Table S3. Oligonucleotide sequences

REFERENCES

1. Stevens, D.L., A.L. Bisno, H.F. Chambers, E.P. Dellinger, E.J. Goldstein, S.L. Gorbach, J.V. Hirschmann, S.L. Kaplan, J.G. Montoya, and J.C. Wade, *Practice guidelines for the diagnosis and management of skin and soft tissue infections: 2014 update by the Infectious Diseases Society of America.* Clin Infect Dis, 2014. **59**(2): p. e10-52.

2. Metersky, M.L. and A.C. Kalil, *Management of ventilator-associated pneumonia: guidelines.* Clinics in chest medicine, 2018. **39**(4): p. 797-808.

3. Meylan, S., I.W. Andrews, and J.J. Collins, *Targeting Antibiotic Tolerance, Pathogen by Pathogen.* Cell, 2018. **172**(6): p. 1228-1238.

4. Namugenyi, S.B., A.M. Aagesen, S.R. Elliott, and A.D. Tischler, *Mycobacterium tuberculosis PhoY Proteins Promote Persister Formation by Mediating Pst/SenX3-RegX3 Phosphate Sensing.* mBio, 2017. **8**(4): p. e00494-17.

5. Liu, Y., S. Tan, L. Huang, R.B. Abramovitch, K.H. Rohde, M.D. Zimmerman, C. Chen, V. Dartois, B.C. VanderVen, and D.G. Russell, *Immune activation of the host cell induces drug tolerance in Mycobacterium tuberculosis both in vitro and in vivo.* Journal of Experimental Medicine, 2016. **213**(5): p. 809-825.

6. Gold, B. and C. Nathan, *Targeting Phenotypically Tolerant Mycobacterium tuberculosis.* Microbiol Spectr, 2017. **5**(1).

7. Bigger, J.J.T.L., *Treatment of staphylococcal infections with penicillin by intermittent sterilisation.* 1944. **244**(6320): p. 497-500.

8. Ronneau, S., P.W.S. Hill, and S. Helaine, *Antibiotic persistence and tolerance: not just one and the same.* Current opinion in microbiology, 2021. **64**: p. 76-81.

9. Dhar, N. and J.D. McKinney, *Microbial phenotypic heterogeneity and antibiotic tolerance.* Current opinion in microbiology, 2007. **10**(1): p. 30-38.

596   10.   Grant, S.S., B.B. Kaufmann, N.S. Chand, N. Haseley, and D.T. Hung, *Eradication of*

597   *bacterial persisters with antibiotic-generated hydroxyl radicals.* Proceedings of the

598   National Academy of Sciences, 2012. **109**(30): p. 12147-12152.

599   11.   Baker, J.J. and R.B. Abramovitch, *Genetic and metabolic regulation of Mycobacterium*

600   *tuberculosis acid growth arrest.* Scientific Reports, 2018. **8**(1).

601   12.   Saito, K., T. Warrier, S. Somersan-Karakaya, L. Kaminski, J. Mi, X. Jiang, S. Park, K.

602   Shigyo, B. Gold, and J. Roberts, *Rifamycin action on RNA polymerase in antibiotic-*

603   *tolerant Mycobacterium tuberculosis results in differentially detectable populations.*

604   Proceedings of the National Academy of Sciences, 2017. **114**(24): p. E4832-E4840.

605   13.   Hipolito, V.E.B., E. Ospina-Escobar, and R.J. Botelho, *Lysosome remodelling and*

606   *adaptation during phagocyte activation.* Cell Microbiol, 2018. **20**(4).

607   14.   Sukumar, N., S. Tan, B.B. Aldridge, and D.G. Russell, *Exploitation of Mycobacterium*

608   *tuberculosis reporter strains to probe the impact of vaccination at sites of infection.* PLoS

609   Pathog, 2014. **10**(9): p. e1004394.

610   15.   Schumacher, M.A., K.M. Piro, W. Xu, S. Hansen, K. Lewis, and R.G. Brennan, *Molecular*

611   *Mechanisms of HipA-Mediated Multidrug Tolerance and Its Neutralization by HipB.*

612   Science, 2009. **323**(5912): p. 396-401.

613   16.   Harms, A., C. Fino, A. Sørensen Michael, S. Semsey, and K. Gerdes, *Prophages and*

614   *Growth Dynamics Confound Experimental Results with Antibiotic-Tolerant Persister*

615   *Cells.* mBio, 2017. **8**(6): p. 10.1128/mbio.01964-17.

616   17.   Korch, S.B., T.A. Henderson, and T.M. Hill, *Characterization of the hipA7 allele of*

617   *Escherichia coli and evidence that high persistence is governed by (p)ppGpp synthesis.*

618   Mol Microbiol, 2003. **50**(4): p. 1199-213.

619   18.   Wong, F., S. Wilson, R. Helbig, S. Hegde, O. Aftenieva, H. Zheng, C. Liu, T. Pilizota, E.C.

620   Garner, A. Amir, and L.D. Renner, *Understanding Beta-Lactam-Induced Lysis at the*

621   *Single-Cell Level.* Frontiers in Microbiology, 2021. **12**.

622   19.   Saito, K., S. Mishra, T. Warrier, N. Cicchetti, J. Mi, E. Weber, X. Jiang, J. Roberts, A.

623   Gouzy, E. Kaplan, C.D. Brown, B. Gold, and C. Nathan, *Oxidative damage and delayed*

624   *replication allow viable Mycobacterium tuberculosis to go undetected.* Sci Transl Med,

625   2021. **13**(621): p. eabg2612.

626   20.   Vilchèze, C., T. Hartman, B. Weinrick, P. Jain, T.R. Weisbrod, L.W. Leung, J.S.

627   Freundlich, and W.R. Jacobs, Jr., *Enhanced respiration prevents drug tolerance and*

628   *drug resistance in Mycobacterium tuberculosis.* Proc Natl Acad Sci U S A, 2017. **114**(17):

629   p. 4495-4500.

630   21.   Keren, I., S. Minami, E. Rubin, and K. Lewis, *Characterization and transcriptome*

631   *analysis of Mycobacterium tuberculosis persisters.* mBio, 2011. **2**(3): p. e00100-11.

632   22.   Liu, Y. and J.A. Imlay, *Cell death from antibiotics without the involvement of reactive*

633   *oxygen species.* Science, 2013. **339**(6124): p. 1210-3.

634   23.   de Keijzer, J., A. Mulder, J. de Beer, A.H. de Ru, P.A. van Veelen, and D. van Soolingen,

635   *Mechanisms of Phenotypic Rifampicin Tolerance in Mycobacterium tuberculosis Beijing*

636   *Genotype Strain B0/W148 Revealed by Proteomics.* J Proteome Res, 2016. **15**(4): p.

637   1194-204.

638   24.   Gold, B., T. Warrier, and C. Nathan, *A multi-stress model for high throughput screening*

639   *against non-replicating Mycobacterium tuberculosis.* Mycobacteria protocols, 2015: p.

640   293-315.

641   25.   Dorman, S.E., P. Nahid, E.V. Kurbatova, P.P. Phillips, K. Bryant, K.E. Dooley, M. Engle,

642   S.V. Goldberg, H.T. Phan, and J. Hakim, *Four-month rifapentine regimens with or without*

643   *moxifloxacin for tuberculosis.* New England Journal of Medicine, 2021. **384**(18): p. 1705-

644   1718.

645   26.   Nahid, P., S.E. Dorman, N. Alipanah, P.M. Barry, J.L. Brozek, A. Cattamanchi, L.H.

646   Chaisson, R.E. Chaisson, C.L. Daley, and M. Grzemska, *Official American thoracic*

647   *society/centers for disease control and prevention/infectious diseases society of America*

clinical practice guidelines: treatment of drug-susceptible tuberculosis. Clinical infectious

diseases, 2016. **63**(7): p. e147-e195.

27.   Griffith, D.E., T. Aksamit, B.A. Brown-Elliott, A. Catanzaro, C. Daley, F. Gordin, S.M.

Holland, R. Horsburgh, G. Huitt, and M.F. Iademarco, *An official ATS/IDSA statement:*

diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases.

American journal of respiratory and critical care medicine, 2007. **175**(4): p. 367-416.

28.   Griffith, D.E. and C.L. Daley, *Treatment of Mycobacterium abscessus Pulmonary*

Disease. CHEST, 2022. **161**(1): p. 64-75.

29.   Rodriguez, R., N. Campbell-Kruger, J. Gonzalez Camba, J. Berude, R. Fetterman, and

S. Stanley, *MarR-Dependent Transcriptional Regulation of mmpSL5 Induces*

Ethionamide Resistance in Mycobacterium abscessus. Antimicrobial Agents and

Chemotherapy, 2023. **67**(4): p. e01350-22.

30.   Li, S., N.C. Poulton, J.S. Chang, Z.A. Azadian, M.A. DeJesus, N. Ruecker, M.D.

Zimmerman, K.A. Eckartt, B. Bosch, C.A. Engelhart, D.F. Sullivan, M. Gengenbacher,

V.A. Dartois, D. Schnappinger, and J.M. Rock, *CRISPRi chemical genetics and*

comparative genomics identify genes mediating drug potency in Mycobacterium

tuberculosis. Nature microbiology, 2022. **7**(6): p. 766-779.

31.   Carey, A.F., J.M. Rock, I.V. Krieger, M.R. Chase, M. Fernandez-Suarez, S. Gagneux,

J.C. Sacchettini, T.R. Ioerger, and S.M. Fortune, *TnSeq of Mycobacterium tuberculosis*

clinical isolates reveals strain-specific antibiotic liabilities. PLoS pathogens, 2018. **14**(3):

p. e1006939.

32.   Kreutzfeldt, K.M., R.S. Jansen, T.E. Hartman, A. Gouzy, R. Wang, I.V. Krieger, M.D.

Zimmerman, M. Gengenbacher, J.P. Sarathy, M. Xie, V. Dartois, J.C. Sacchettini, K.Y.

Rhee, D. Schnappinger, and S. Ehrt, *CinA mediates multidrug tolerance in*

Mycobacterium tuberculosis. Nat Commun, 2022. **13**(1): p. 2203.

33.   Block, A.M., S.B. Namugenyi, N.P. Palani, A.M. Brokaw, L. Zhang, K.B. Beckman, and

A.D. Tischler, *Mycobacterium tuberculosis Requires the Outer Membrane Lipid*

Phthiocerol Dimycocerosate for Starvation-Induced Antibiotic Tolerance. Msystems,

2023. **8**(1): p. e0069922.

34.   Bellerose, M.M., S.-H. Baek, C.-C. Huang, C.E. Moss, E.-I. Koh, M.K. Proulx, C.M.

Smith, R.E. Baker, J.S. Lee, S. Eum, S.J. Shin, S.-N. Cho, M. Murray, and C.M. Sassetti,

Common Variants in the Glycerol Kinase Gene Reduce Tuberculosis Drug Efficacy.

mBio, 2019. **10**(4): p. e00663-19.

35.   Kumar, K., C.L. Daley, D.E. Griffith, and M.R. Loebinger, *Management of Mycobacterium*

avium complex and Mycobacterium abscessus pulmonary disease: therapeutic

advances and emerging treatments. European respiratory review, 2022. **31**(163).

36.   Woods, G.L., B.A. Brown-Elliott, P.S. Conville, E.P. Desmond, G.S. Hall, G. Lin, G.E.

Pfyffer, J.C. Ridderhof, S.H. Siddiqi, R.J. Wallace, Jr., N.G. Warren, and F.G. Witebsky,

CLSI Standards: Guidelines for Health Care Excellence, in *Susceptibility Testing of*

Mycobacteria, Nocardiae, and Other Aerobic Actinomycetes. 2011, Clinical and

Laboratory Standards Institute

37.   Yam, Y.K., N. Alvarez, M.L. Go, and T. Dick, *Extreme Drug Tolerance of Mycobacterium*
abscessus "Persisters". Front Microbiol, 2020. **11**: p. 359.

38.   Berube, B.J., L. Castro, D. Russell, Y. Ovechkina, and T. Parish, *Novel Screen to Assess*
Bactericidal Activity of Compounds Against Non-replicating Mycobacterium abscessus.
Front Microbiol, 2018. **9**: p. 2417.

39.   Lee, J., N. Ammerman, A. Agarwal, M. Naji, S.Y. Li, and E. Nuermberger, *Differential In*
Vitro Activities of Individual Drugs and Bedaquiline-Rifabutin Combinations against
Actively Multiplying and Nutrient-Starved Mycobacterium abscessus. Antimicrob Agents
Chemother, 2021. **65**(2).

40.   Betts, J.C., P.T. Lukey, L.C. Robb, R.A. McAdam, and K. Duncan, *Evaluation of a*
nutrient starvation model of Mycobacterium tuberculosis persistence by gene and protein
expression profiling. Molecular Microbiology, 2002. **43**(3): p. 717-731.

41.   DeJesus, M.A., C. Ambadipudi, R. Baker, C. Sassetti, and T.R. Ioerger, *TRANSIT-a*
software tool for Himar1 TnSeq analysis. PLoS computational biology, 2015. **11**(10): p.
e1004401.

42.   Shi, H., R. Zhang, L. Lan, Z. Chen, and J. Kan, *Zinc mediates resuscitation of lactic acid-*
injured Escherichia coli by relieving oxidative stress. Journal of Applied Microbiology,
2019. **127**(6): p. 1741-1750.

43.   Hohle, T.H. and M.R. O'Brian, *The mntH gene encodes the major Mn(2+) transporter in*
Bradyrhizobium japonicum and is regulated by manganese via the Fur protein. Mol
Microbiol, 2009. **72**(2): p. 399-409.

44.   Murphy, K.C., S.J. Nelson, S. Nambi, K. Papavinasasundaram, C.E. Baer, and C.M.
Sassetti, *ORBIT: a new paradigm for genetic engineering of mycobacterial*
chromosomes. mBio, 2018. **9**(6): p. e01467-18.

45.   McBee, M.E., Y.H. Chionh, M.L. Sharaf, P. Ho, M.W.L. Cai, and P.C. Dedon, *Production*
of superoxide in bacteria is stress- and cell state-dependent: A gating-optimized flow

cytometry method that minimizes ROS measurement artifacts with fluorescent dyes.

Frontiers in Microbiology, 2017. **8**(MAR).

46.   Wayne, L.G. and L.G. Hayes, *An In Vitro Model for Sequential Study of Shiftdown of*

Mycobacterium tuberculosis through Two Stages of Nonreplicating Persistence. Infection

and Immunity, 1996. **64**(6): p. 2062-2069.

47.   Brazas Michelle, D., B.M. Breidenstein Elena, J. Overhage, and E.W. Hancock Robert,

Role of Lon, an ATP-Dependent Protease Homolog, in Resistance of Pseudomonas

aeruginosa to Ciprofloxacin. Antimicrobial Agents and Chemotherapy, 2007. **51**(12): p.

4276-4283.

48.   Kusser, W. and E.E. Ishiguro, *Involvement of the relA gene in the autolysis of*

Escherichia coli induced by inhibitors of peptidoglycan biosynthesis. Journal of

Bacteriology, 1985. **164**(2): p. 861-865.

49.   Bokinsky, G., E.E.K. Baidoo, S. Akella, H. Burd, D. Weaver, J. Alonso-Gutierrez, H.

García-Martín, T.S. Lee, and J.D. Keasling, *HipA-Triggered Growth Arrest and β-Lactam*

Tolerance in Escherichia coli Are Mediated by RelA-Dependent ppGpp Synthesis.

Journal of Bacteriology, 2013. **195**(14): p. 3173-3182.

50.   Honsa, E.S., V.S. Cooper, M.N. Mhaissen, M. Frank, J. Shaker, A. Iverson, J. Rubnitz,

R.T. Hayden, R.E. Lee, C.O. Rock, E.I. Tuomanen, J. Wolf, and J.W. Rosch, *RelA*

Mutant Enterococcus faecium with Multiantibiotic Tolerance Arising in an

Immunocompromised Host. mBio, 2017. **8**(1): p. e02124-16.

51.   Nguyen, D., A. Joshi-Datar, F. Lepine, E. Bauerle, O. Olakanmi, K. Beer, G. McKay, R.

Siehnel, J. Schafhauser, Y. Wang, B.E. Britigan, and P.K. Singh, *Active starvation*

responses mediate antibiotic tolerance in biofilms and nutrient-limited bacteria. Science,

2011. **334**(6058): p. 982-6.

52.   Kudrin, P., V. Varik, S.R.A. Oliveira, J. Beljantseva, T.D.P. Santos, I. Dzhygyr, D. Rejman,

F. Cava, T. Tenson, and V. Hauryliuk, *Subinhibitory Concentrations of Bacteriostatic*

Antibiotics Induce <i>relA</i>-Dependent and <i>relA</i>-Independent Tolerance to β-Lactams. Antimicrobial Agents and Chemotherapy, 2017. **61**(4): p. 10.1128/aac.02173-16.

53. Viducic, D., T. Ono, K. Murakami, H. Susilowati, S. Kayama, K. Hirota, and Y. Miyake, *Functional analysis of spoT, relA and dksA genes on quinolone tolerance in Pseudomonas aeruginosa under nongrowing condition.* Microbiol Immunol, 2006. **50**(4): p. 349-57.

54. Geiger, T., B. Kästle, F.L. Gratani, C. Goerke, and C. Wolz, *Two small (p)ppGpp synthases in Staphylococcus aureus mediate tolerance against cell envelope stress conditions.* J Bacteriol, 2014. **196**(4): p. 894-902.

55. Dutta, N.K., L.G. Klinkenberg, M.J. Vazquez, D. Segura-Carro, G. Colmenarejo, F. Ramon, B. Rodriguez-Miquel, L. Mata-Cantero, E. Porras-De Francisco, Y.M. Chuang, H. Rubin, J.J. Lee, H. Eoh, J.S. Bader, E. Perez-Herran, A. Mendoza-Losana, and P.C. Karakousis, *Inhibiting the stringent response blocks Mycobacterium tuberculosis entry into quiescence and reduces persistence.* Sci Adv, 2019. **5**(3): p. eaav2104.

56. Bhaskar, A., C. De Piano, E. Gelman, J.D. McKinney, and N. Dhar, *Elucidating the role of (p) ppGpp in mycobacterial persistence against antibiotics.* IUBMB life, 2018. **70**(9): p. 836-844.

57. Conlon, B.P., S.E. Rowe, A.B. Gandt, A.S. Nuxoll, N.P. Donegan, E.A. Zalis, G. Clair, J.N. Adkins, A.L. Cheung, and K. Lewis, *Persister formation in Staphylococcus aureus is associated with ATP depletion.* Nature Microbiology, 2016. **1**(5): p. 16051.

58. Kapoor, G., S. Saigal, and A. Elongavan, *Action and resistance mechanisms of antibiotics: A guide for clinicians.* Journal of Anaesthesiology Clinical Pharmacology, 2017. **33**(3): p. 300-305.

59.   Kohanski, M.A., D.J. Dwyer, B. Hayete, C.A. Lawrence, and J.J. Collins, *A common*

mechanism of cellular death induced by bactericidal antibiotics. Cell, 2007. **130**(5): p.

797-810.

60.   Sebastian, J., S. Swaminath, R.R. Nair, K. Jakkala, A. Pradhan, and P. Ajitkumar, *De*

Novo Emergence of Genetically Resistant Mutants of Mycobacterium tuberculosis from

the Persistence Phase Cells Formed against Antituberculosis Drugs In Vitro. Antimicrob

Agents Chemother, 2017. **61**(2).

FIGURE 1

A. *M. smegmatis*

B. *M. tuberculosis*

C. *M. abscessus*

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

SUPPLEMENTAL FIGURE 1

SUPPLEMENTAL FIGURE 2