B cells imprinted by ancestral SARS-CoV-2 develop pan-sarbecovirus neutralization in immune recalls

Xixian Chen[1,2,#], Ling Li[1,#], Ruiping Du[1,2,#], Zuowei Wang[1,#], Yunjian Li[1,2], Yi Sun[1,2], Rongrong Qin[1,2], Hualong Feng[1], Lin Hu[1], Xuanyi Chen[1], Maosheng Lu[1], Xueyan Huang[1], Liwei Jiang[1,*], Teng Zuo[1,*]

[1]Laboratory of Immunoengineering, Institute of Health and Medical Technology, Hefei Institutes of Physical Science, Chinese Academy of Sciences, Hefei 230031, China

[2]University of Science and Technology of China, Hefei 230026, China

[#]These authors contributed equally to this work

[*]Correspondence: jlw0531@cmpt.ac.cn (L.J.); zuot@cmpt.ac.cn (T.Z.)

Short running title: Evolution of B cells imprinted by ancestral SARS-CoV-2

Summary

Chen et al. demonstrate that B cells imprinted by ancestral SARS-CoV-2 have tremendous potential to develop neutralizing breadth and potency in repeated immune recalls driven by Omicron variants, implicating that ancestral SARS-CoV-2 immune imprinting can be harnessed in developing pan-SARS-CoV-2 and even pan-sarbecovirus vaccines.

Abstract

A key question on ancestral SARS-CoV-2 immune imprinting is to what extent imprinted B cells can develop neutralizing breadth and potency in immune recalls. Here, we longitudinally tracked B cells recognizing wild-type spike in two individuals, who were sequentially infected by Omicron variants after receiving mRNA vaccines. Functional and genetic analysis of 632 monoclonal antibodies (mAbs) from those B cells reveals that mAbs cloned after second infection have dramatically enhanced neutralizing breadth and potency, which is attributed to recall and maturation of pre-existing memory B cells. Among the 11 mAbs that potently neutralize SARS-CoV-2 variants from wild-type to KP.3, 5 mAbs are classified into public clonotypes encoded by IGHV3-53 or IGHV3-66, whereas the rest belong to a rarely reported clonotype encoded by IGHV3-74. Notably, IGHV3-74 mAbs can also broadly neutralize other sarbecoviruses by targeting a novel epitope on receptor-binding domain of spike. These results support that ancestral SARS-CoV-2 immune imprinting can be harnessed in developing pan-SARS-CoV-2 and even pan-sarbecovirus vaccines.

Introduction

Adaptive immune system adopts multiple mechanisms to ensure specific and diverse antibody responses against various pathogens. During B cell development, V(D)J recombination generates a vast diversity of B cell antigen receptors (BCRs) for naïve B cells (Tonegawa, 1983). When infection or vaccination occurs, a small portion of naïve B cell repertoire get activated as their BCRs recognize corresponding antigens by chance (Batista and Harwood, 2009). Activated B cells either differentiate into short-lived plasmablasts (PBs) to produce the initial wave of antibodies, or enter germinal centers (GCs) to undergo repeated rounds of somatic hypermutation and affinity-dependent selection, a process known as antibody affinity maturation (Victora and Nussenzweig, 2022). The evolved GC B cells eventually differentiate into long-lived plasma cells (PCs) to constantly secrete antibodies for months to years, or memory B cells to wait for future encounter with the same or similar antigens (Victora and Nussenzweig, 2022). Compared with primary antigen exposure, secondary antigen exposure typically leads to much faster and stronger antibody responses as recalled memory B cells rapidly differentiate into PB/PCs to secrete high-affinity antibodies (Inoue et al., 2018). In addition, memory B cells can be recruited into newly formed GCs to further mature upon secondary antigen exposure (Inoue et al., 2018). Overall, these mechanisms work in concert to establish antibody-mediated immune protection against infections.

Since the outbreak of SARS-CoV-2, numerous studies on human antibody responses against spike protein of this emerging virus have been conducted to inform the development of vaccines and antibody therapeutics (Lapuente et al., 2024; Qi et al., 2022; Roltgen and Boyd, 2024). Early in the pandemic, extensive isolation and characterization of monoclonal antibodies (mAbs) from convalescents reveals that neutralizing antibodies primarily target epitopes on receptor-binding

domain (RBD) and to a lesser extent on N-terminal domain (NTD) of spike (Ling et al., 2023; Liu and

Wilson, 2022). Furthermore, RBD-specific neutralizing antibodies are categorized into seven major

groups according to their binding modes and fine epitopes (Hastie et al., 2021), whereas NTD-specific

neutralizing antibodies largely target a single supersite (Cerutti et al., 2021; McCallum et al., 2021).

Although spike-specific antibodies from different convalescents are highly diverse, a series of public

antibodies (i.e. antibodies generated by different individuals while sharing common genes and antigen

recognition modes) have been widely reported, including antibodies encoded by IGHV3-53, IGHV3-

66, IGHV1-58 and IGHV1-69 (Robbiani et al., 2020; Wang et al., 2022a; Yan et al., 2022; Yuan et al.,

2020a). In addition, longitudinal profiling of memory B cells from peripheral blood of convalescents

demonstrates that spike- or RBD-specific memory B cells display increased somatic hypermutation,

binding affinity and neutralizing potency over time, reflecting long-lasting GC responses elicited by

primary SARS-CoV-2 infection (Gaebler et al., 2021; Sakharkar et al., 2021; Sokal et al., 2021).

Compared with SARS-CoV-2 infection, two doses of SARS-CoV-2 mRNA vaccines lead to equivalent

plasma neutralizing activity and similar amount of RBD-specific memory B cells (Wang et al., 2021).

Spike- or RBD-specific mAbs cloned from vaccinees and convalescents recognize common epitopes

and share frequently-used antibody genes (Wang et al., 2021). Moreover, analysis of draining lymph

node samples from vaccinees indicates that mRNA vaccines provoke persistent GCs, which last for

at least six months and enable the maturation of spike-specific B cells (Kim et al., 2022; Turner et al.,

2021).

Under the immune pressure established by infection and vaccination, SARS-CoV-2 variants,

represented by Alpha (B.1.1.7), Beta (B.1.351), Gamma (P.1), Delta (B.1.617.2) and Omicron

(B.1.1.529), emerged successively in the first two years of the pandemic. Among them, Omicron

exhibited most striking evasion to neutralizing antibodies induced by ancestral SARS-CoV-2 spike and therefore caused a new wave of infections worldwide (Viana et al., 2022). Omicron infection in individuals administrated with mRNA vaccines predominantly recalls cross-reactive memory B cells elicited by vaccination whereas rarely induces *de novo* Omicron-specific B cells responses, highlighting that ancestral SARS-CoV-2 imprints B cell responses to Omicron (Kaku et al., 2022; Quandt et al., 2022; Wang et al., 2022b; Weber et al., 2023). Consequently, plasma neutralizing activity against ancestral SARS-CoV-2 and Omicron are enhanced synchronously (Kaku et al., 2022; Quandt et al., 2022; Wang et al., 2022b; Weber et al., 2023). Moreover, B cell clones with more mutations and higher affinities towards Omicron progressively remodel spike-specific memory B cell repertoire, suggesting that cross-reactive memory B cells can undergo further affinity maturation in new GCs induced by Omicron breakthrough infection (Kaku et al., 2023; Sokal et al., 2023).

With the rapid emergence of Omicron-derived variants, such as BA.2.75, BA.5, BF.7, BQ.1, XBB.1.5, XBB.1.16, EG.5.1, BA.2.86 and JN.1, repeated infections with these variants have become common and updated vaccines have been applied accordingly in some countries (Huang et al., 2023). In individuals with prior exposures to wild-type (WT) spike, antibody responses boosted by subsequent exposures to emerging Omicron variants, either through infection or vaccination, display biased neutralizing activities to ancestral SARS-CoV-2 and earlier Omicron variants, suggesting that immune imprinting may impede the elicitation of neutralizing antibodies against more recent and future variants (Johnston et al., 2024; Liang et al., 2024; Tortorici et al., 2024; Wang et al., 2023a; Yisimayi et al., 2024). Nonetheless, mAbs from those individuals dramatically surpass mAbs elicited by WT spike in neutralizing breadth and potency, indicating that immune imprinting is potentially beneficial for the development of broadly neutralizing antibodies (bnAbs) (Paciello et al., 2024; Sokal et al., 2023).

To better understand the beneficial effects of ancestral SARS-CoV-2 immune imprinting on antibody

responses to emerging variants, we ask that to what extent imprinted B cells (i.e. memory B cells

elicited by ancestral SARS-CoV-2) can develop neutralizing breadth and potency in immune recalls.

To address this question, we performed a 7-month longitudinal study of B cells recognizing WT spike

in two individuals of mRNA vaccine, from convalescence of breakthrough infection to acute phase of

reinfection. Functional and genetic analysis of 632 mAbs from those B cells reveals that mAbs cloned

after reinfection have dramatically enhanced neutralizing breadth and potency, which is attributed to

recall of pre-existing memory B cells and maturation of those cells either before or after reinfection.

Among the 11 mAbs that potently neutralize all tested SARS-CoV-2 variants from WT to KP.3, 5 mAbs

are classified into public clonotypes encoded by IGHV3-53 or IGHV3-66, whereas the rest belong to

a rarely reported clonotype encoded by IGHV3-74. Notably, IGHV3-74 mAbs can also broadly

neutralize a panel of sarbecoviruses including SARS-CoV-1, Pangolin CoV GD, Bat CoV WIV16 and

RaTG13. Structural and functional analysis shows that IGHV3-74 mAbs can block the binding

between RBD and ACE2 by targeting a novel epitope consisting of E340-R346, V367-S375, N436-

G446 and Q498-V503. Moreover, the best IGHV3-74 mAb, termed KXD355, is highly resilient to

variations on this epitope. Overall, this study demonstrates that both public and rare antibody

clonotypes imprinted by ancestral SARS-CoV-2 have potential to achieve extraordinary neutralizing

breadth and potency in repeated Omicron infections, which implicates that ancestral SARS-CoV-2

immune imprinting can be harnessed in developing pan-SARS-CoV-2 and even pan-sarbecovirus

vaccines.

Results

Reinfection enhances plasma neutralizing antibody responses

In our previous study, we recruited a cohort to study antibody responses elicited by breakthrough infection (Li et al., 2023). Among them, Donor 1 and 2 received two doses of mRNA vaccines in the middle or beginning of 2021 and were infected by BA.5 or BF.7 in the end of 2022 (Figure 1A). With their blood samples collected about two months (T1) after infection, we have identified multiple mAbs with broadly neutralizing capacity (Li et al., 2023). To longitudinally track B cell responses in these two donors, we also collected blood samples about 3 months (T2) and 6 months (T3) after infection. Donor 1 showed symptoms of infection and was confirmed by SARS-CoV-2 antigen test on August 25, 2023, when EG.5.1 caused a new round of infections in China (Wang et al., 2023b). Eleven days later (T4), blood was collected again from Donor 1 to study acute phase antibody responses evoked by reinfection. In parallel, we also collected blood from Donor 2 at T4, though Donor 2 reported no symptoms and negative test results during several waves of infections before this time point.

We measured neutralizing antibody titers of plasma from four time points against a panel of 18 pseudoviruses bearing spikes from ancestral SARS-CoV-2 (WT), Alpha, Beta, Gamma, Delta, BA.1, BA.4/5, BA.2.75, BQ.1, XBB, XBB.1.5, EG.5.1, HK.3, BA.2.86, JN.1, KP.2, KP.3 and SARS-CoV-1 (Figure 1B and Figure S1). Overall, the plasma neutralizing titers (NT50s) of both donors gradually decrease from T1 to T3 with a maximum of 10.7 folds. After reinfection, plasma from Donor 1 at T4 shows substantially increased neutralizing titers against most strains by a maximum of 20.7 folds compared to T3. The titers of plasma from Donor 2 continuously decline or sustain from T3 to T4 for most strains, whereas increase by 1.9 to 2.7 folds for strains including Gamma, BA.1 and XBB. As latest variants of XBB and BA.2.86 lineages, HK.3 and KP.3 are least sensitive to neutralization, highlighting the continuous neutralization escape (Figure 1C). Nonetheless, plasma at T4 from Donor

1 displays relatively stable neutralizing titers across all SARS-CoV-2 variants, indicating that

reinfection enhances the breadth of neutralizing antibody response in Donor 1.

Monoclonal antibodies cloned from WT-spike-specific B cells after reinfection exhibit

improved neutralizing breadth and potency

As B cells imprinted by ancestral SARS-CoV-2 mRNA vaccine (i.e. memory B cells) typically maintain

their specificities to WT spike in recalled responses (Johnston et al., 2024; Kaku et al., 2022; Kaku et

al., 2023; Liang et al., 2024; Quandt et al., 2022; Sokal et al., 2023; Tortorici et al., 2024; Wang et al.,

2023a; Wang et al., 2022b; Weber et al., 2023; Yisimayi et al., 2024), we used WT spike as bait to

track imprinted B cells in peripheral blood mononuclear cells (PBMCs) from these two donors.

Consistent with plasma neutralizing titers, the frequencies of WT-spike-specific B cells in Donor 1

drop sequentially from T1 to T3 and then rise back to initial level (Figure 2A). Unexpectedly, Donor 2

displays similar dynamic changes as Donor 1, suggesting that Donor 2 may have experienced an

asymptomatic infection between T3 and T4.

To examine the binding and neutralizing properties of antibodies from those WT-spike-specific B

cells, we cloned the variable regions of antibodies from those B cells into expression vectors through

single-cell RT-PCR. Then we transfected HEK293T cells with those vectors and harvested

supernatant for ELISA and neutralization assay (Figure 2B). In ELISA, the supernatant was tested

with RBD, NTD, S1, S2 and spike (S) from WT SARS-CoV-2 and spikes from Omicron variants

including BA.4/5, XBB.1.5, EG.5.1 and JN.1. In neutralization assay, the supernatant was tested

against WT, BA.4/5, XBB.1.5, EG.5.1 and JN.1 pseudoviruses.

With published mAbs at T1 included, we identified 448 and 184 WT-spike-specific mAbs from Donor

1 and Donor 2 respectively. The main targets of these mAbs are RBD, NTD and S2, with the

proportions of mAbs to each domain slightly varying from T1 to T4 (Figure 2C). Regarding to cross-

binding with different spikes, the frequencies of RBD-specific mAbs that bind WT and one Omicron

variant, either BA.4/5, XBB.1.5, EG.5.1 or JN.1, decline from T1 to T3 and then ascend (Figure 2D,

Figure S2A). However, similar trend is not observed in NTD- and S2-specific mAbs.

By defining 70% inhibition as cutoff, we identified 127 mAbs that are able to neutralize WT

pseudoviruses (WT-neutralizing mAbs), with 77 from Donor 1 and 50 from Donor 2 (Figure 2E).

Among them, only 5 mAbs target NTD while the others exclusively recognize RBD. Regarding to

cross-neutralizing activity, around 20-30% of mAbs at T4 are able to neutralize all 5 tested

pseudoviruses, whereas only 2 out of 61 mAbs from T1 to T3 have such capacity (Figure 2F).

Moreover, the frequencies of mAbs that cross-neutralize WT and one Omicron variant drop from T1

to T3 and then jump to highest levels at T4 (Figure 2G, Figure S2B).

To further compare the breadth and potency of neutralizing mAbs from different time points, we

purified those mAbs and measured their IC50s against 14 representative pseudoviruses including

WT, Delta, BA.1, BA.4/5, BA.2.75, BQ.1, XBB.1.5, EG.5.1, HK.3, BA.2.86, JN.1, KP.2, KP.3 and

SARS-CoV-1 (Figure 3A). Four neutralizing mAbs screened by supernatant were not included in this

analysis because of low yields. Among all tested mAbs, 4 mAbs from Donor 2 are able to neutralize

all SARS-CoV-2 variants and SARS-CoV-1. In addition, 2 mAbs from Donor 1 and 5 mAbs from Donor

2 can neutralize all SARS-CoV-2 variants from WT to KP.3. Notably, these 11 mAbs are exclusively

identified from PBMCs collected at T4. Along with this finding, differences are observed in the overall

neutralizing breadth of mAbs at T1 to T4 (Figure 3B, Figure S3A). T1 and T4 have higher frequencies

of neutralizing antibody against BA.1, BA.4/5, BA.2.75, BQ.1, XBB.1.5 and BA.2.86 than T2 and T3,

which is consistent with neutralization results of supernatant. Moreover, T4 has higher proportions of

neutralizing antibody against EG.5.1, HK.3, JN.1, KP.2 and KP.3 than the other time points.

Regarding to neutralizing potency, the overall IC50s of mAbs at T4 are lower than that of mAbs at

T2 and T3 against most strains (Figure 3C, D, Figure S3A). In addition, mAbs at T4 also display

superior neutralizing potency than mAbs at T1 against Delta, EG.5.1, HK.3, JN.1 and KP.2. In contrast,

significant differences in IC50s are generally not observed among mAbs at T1 to T3 by statistical

analysis. We also combined mAbs at T1 to T3 together and compared them with mAbs at T4, which

further confirms that mAbs isolated after reinfection have superior neutralizing breadth and potency

than mAbs before reinfection (Figure S3B-D). Taken together, these data suggest that mAbs cloned

from WT-spike-specific B cells after reinfection have enhanced neutralizing breadth and potency.

Reinfection recalls pre-existing memory B cells

To explain the enhanced neutralizing breadth and potency of mAbs cloned after reinfection, we moved

on to analyze the sequences of mAbs from T1 to T4. Binding mAbs from both donors use a wide

range of V genes (Figure 4A, B). The frequencies of each V gene from T1 to T4 are more consistent

for mAbs from Donor 1 than those from Donor 2, likely due to fewer mAbs isolated from Donor 2 at

T2 and T3. The most frequently-used V genes for heavy chain include IGHV3-30, IGHV3-66, IGHV3-

53, IGHV3-33, IGHV1-69 and IGHV1-46. Among them, IGHV3-30 and IGHV3-33 are primarily used

by mAbs targeting S2, whereas IGHV3-66, IGHV3-53 and IGHV1-69 are preferentially used by mAbs

recognizing RBD (Figure S4A). The most frequently-used V genes for light chain include IGKV1-39,

IGKV3-20, IGKV3-15, IGKV3-11, IGKV1-33, IGKV1-5 and IGKV1-9, which are less biased toward

mAbs targeting a certain domain than V genes for heavy chain (Figure S4A). In contrast to binding

mAbs, neutralizing mAbs are predominantly encoded by IGHV3-66, IGHV3-53 and IGHV1-69 (Figure

4A, B). Furthermore, in these neutralizing mAbs, IGHV3-66 and IGHV3-53 typically pair with IGKV1-

33, IGKV1-9 and IGKV3-20, whereas IGHV1-69 commonly pair with IGLV1-40 (Figure S4B). It is

worth mentioning that the pair of IGHV3-74 and IGKV3-20 is used by 10 out of 50 neutralizing mAbs

from Donor 2. Nonetheless, the overall profiles of V gene usage are rather consistent from T1 to T4.

According to V and J genes and CDR3 similarity, we defined 401 and 145 unique clonotypes

respectively from 448 and 184 mAbs (Figure 4C). The expanded clonotypes are primarily found in

mAbs at T4, particular for Donor 2, further indicating that T4 is also in acute phase of reinfection for

Donor 2. As an evidence of recall of pre-existing memory B cells, multiple clonotypes found at T1 to

T3 are extensively expanded at T4, particular for neutralizing clonotypes such as Clonotype 204 and

223 from Donor 1 and Clonotype 88 and 92 from Donor 2 (Figure 4D).

On the other hand, mAbs at T4 have more somatic mutations than those from earlier time points,

which can be either due to recall of pre-existing memory B cells with more mutations, or due to further

maturation of pre-existing memory B cells in new GC reactions (Figure 4E). Taken together, these

data suggest that the enhanced neutralizing breadth and potency of mAbs at T4 is attributed to recall

of pre-existing memory B cells and maturation of those cells either before or after reinfection.

Both public and rare antibody clonotypes can develop extraordinary neutralizing breadth and

potency

After longitudinal analysis of overall B cell responses in these two donors, we moved on to

characterize the 11 mAbs with extraordinary neutralizing breadth (Figure 3A, Figure 5A). Clonotype

analysis reveals that they belong to 5 unique clonotypes including Clonotype 92, 82 and 75 from

Donor 2 and Clonotype 240-2 and 212-1 from Donor 1. Clonotype 92 is encoded by IGHV3-74 and

IGHV3-20, which is rarely reported in RBD-specific antibodies against SARS-CoV-2 (Wang et al.,

2022a). This clonotype contains 10 neutralizing mAbs, with 1 from T2 and 9 from T4. Among them,

KXD350, KXD352, KXD355 and KXD358 can neutralize all tested SARS-CoV-2 variants and SARS-

CoV-1. Further neutralization analysis shows that KXD350 and KXD355 can also neutralize other

sarbecoviruses including Pangolin CoV GD, Bat CoV WIV16 and RaTG13-T372A, whereas KXD352

and KXD358 are escaped by RaTG13-T372A (Figure 5B). According to the phylogenetic tree, mAbs

in this clonotype can be divided into two major clades, suggesting that neutralizing breadth can be

developed through diverse evolution pathways.

The other four clonotypes belong to previously defined IGHV3-53/3-66 public antibody clonotypes

(Wang et al., 2022a). Clonotype 82 is encoded by IGVH3-66 and IGKV1-33 and contains 8

neutralizing mAbs from T4. However, only KXD357 is able to neutralize all SARS-CoV-2 variants, with

IC50s ranging from 0.003 to 0.224 µg/ml. Similar as Clonotype 92, Clonotype 82 also has two major

clades. Although mAbs in both clades are extensively mutated, mAbs in Clade-2 exhibit superior

neutralizing breadth than mAbs in Clade-1. Encoded by IGHV3-53 and IGKV3-15, Clonotype 75

contains previously published KXD02 from T1 and two addition mAbs from T4. In our previous study,

we have shown that KXD02 broadly neutralizes SARS-CoV-2 variants from WT to HK.3 (Li et al.,

2023). However, KXD02 is soon escaped by variants in BA.2.86 lineages including JN.1, KP.2 and

KP.3. In contrast, KXD353 and KXD356 from T4 maintain neutralizing activities against all SARS-

CoV-2 variants, with IC50s ranging from 0.002 to 0.216 µg/ml. In line with increased neutralizing

breadth and potency, KXD353 and KXD356 are also more evolved than KXD02 according to

phylogenetic analysis. As singletons from Donor 1, Clonotype 240-2 (KXD470) is encoded by IGHV3-

66 and IGKV1-9, whereas Clonotype 212-1 (KXD472) is encoded by IGHV3-53 and IGKV1-5.

Although both mAbs are able to neutralize all SARS-CoV-2 variants, they are less potent to KP.2 and

KP.3 than bnAbs from Donor 2.

As all these clonotypes are identified with WT spike, we assume that they are initially primed by

ancestral SARS-CoV-2 mRNA vaccine but not by BA.5 or BF.7 breakthrough infection. To confirm

this assumption, we synthesized the putative germline antibodies of each clonotypes and measured

their binding avidities to WT and BA.4/5 spikes expressed on HEK293T cells (Figure 5C). BF.7 was

not tested since it has only one amino acid different from BA.5 in RBD. All the germlines, except GL-

4, show strong to moderate binding to WT spike. In contrast, they all display weak or even

undetectable binding to BA.4/5 spike. The biased recognition towards WT spike suggest that these

clonotypes are more likely to be activated by mRNA vaccine. Taken together, these data suggest that

both public and rare antibody clonotypes imprinted by ancestral SARS-CoV-2 have potential to

develop ultra-broad and potent neutralizing activities in repeated Omicron infections.

IGHV3-74-encoded bnAbs target a novel epitope and exhibit dramatic resilience to mutations

on the epitope

To evaluate the neutralizing potential of above bnAbs to future SARS-CoV-2 variants, we moved on

to analyze their epitopes. We first performed competition ELISA to roughly map their epitopes with

ACE2, CR3022, S309, P2S-2E9, BD55-5483, BD56-1854 and BD55-1205 as references (Figure 6A).

As reported, CR3022 binds to a cryptic epitope on the inner face of RBD whereas S309 recognizes

N343 proteoglycan epitope on the outer face of RBD (Figure S5A) (Pinto et al., 2020; Yuan et al.,

2020b). P2S-2E9 and BD55-5483 (analogue of BD55-5514) both target the previously defined "mesa"

of RBD, respectively biased to the outer face and inner face (Cao et al., 2022; Hastie et al., 2021; Ju

et al., 2023). As representatives of IGHV3-53/3-66 public antibodies, BD56-1854 and BD55-1205

target receptor-binding-motif (RBM) (Cao et al., 2023).

We tested four mAbs (KXD350, KXD352, KXD355 and KXD358) from Clonotype 92 (IGHV3-74

mAbs) and five mAbs (KXD353, KXD356, KXD357, KXD470 and KXD472) from other clonotypes

(IGHV3-53/3-66 mAbs) (Figure 6A). IGHV3-53/3-66 mAbs compete with BD55-5483, BD56-1854 and

BD55-1205 but not with S309 and P2S-2E9. In contrast, IGHV3-74 mAbs compete with S309, P2S-

2E9 and BD55-5483 while not with BD56-1854 and BD55-1205. Moreover, no competition is observed

between these mAbs and CR3022 except for some minor competition between KXD358 and CR3022.

In addition, they all compete with ACE2, suggesting that they neutralize SARS-CoV-2 through

receptor blockade.

To define the fine epitopes of these mAbs, we performed structure prediction with AlphaFold 3

(Abramson et al., 2024). The predicted structures of IGHV3-53/3-66 mAbs in complex with WT RBD

are similar to the structure of BD55-1205 in complex with XBB.1.5 RBD (Figure S5B). According to

these structures, we defined the common epitope of IGHV3-53/3-66 mAbs, which largely overlap with

the epitope of ACE2 (Figure 6B, C).

On the other hand, AlphaFold 3 generated various structures for the complexes of IGHV3-74 mAbs

and WT RBD. However, only one type of structures is consistent with epitope mapping by competition

ELISA. Here, KXD355 is taken as an example to present the results. Binding to the previously defined

"escarpment" of RBD (Hastie et al., 2021), KXD355 clashes with S309, P2S-2E9 and BD55-5514

(analogue of BD55-5483) but not with BD55-1205 and CR3022 (Figure S5C). As this epitope is

exposed on both up and down RBDs, KXD355 can bind RBDs in both conformations (Figure 6D). In

details, the long CDRH3 of KXD355 extends into a groove formed by three helixes on the "escarpment"

of RBD (Figure 6E). Meanwhile, the other CDRs attach to "mesa" of RBD to provide support for

CDRH3. Accordingly, the epitope of KXD355 includes 22 residues distributed on E340-R346, V367-

S375, W436-G446 and Q498-V503 (Figure 6F).

To confirm the predicted epitopes, we constructed a panel of single-mutated KP.3 spikes and

measured their escape to those mAbs in binding and neutralization (Figure 6G). Compared to KP.3,

most mutants show similar or slightly lower expression levels on HEK293T cells, which is indicated

by an NTD-specific mAb. In contrast, nearly half of these mutants display reduced binding to ACE by

different extent. Regarding to the binding with mAbs, mutants including K403D, D420N, Y421K,

L456V, L456S, K460F, A475V, N487K, Y489A, Y489D, R498A and R498D completely lose binding

to at least one IGHV3-53/3-66 mAb. To a lesser extent, mutants like N343A, N343F, K440D, L441S

and H445D show decreased binding to IGHV3-74 mAbs. Consistent with reduced binding, the

aforementioned mutants also display dramatic escape of neutralization to corresponding mAbs. In

addition, mutants such as T345A, T345F, K444A and K444D are also resistant to certain IGHV3-74

mAbs, although they only partially lose binding to those mAbs.

In contrast to all IGHV3-53/3-66 mAbs and other IGHV3-74 mAbs, KXD352 and KXD355 maintain

low IC50s against all the mutants except L441S, suggesting that they are more resilient to single

mutations on their epitopes (Figure S6). Compared with SARS-CoV-2 variants, SARS-CoV-1 and

other sarbecoviruses have multiple mutations on the epitopes of IGHV3-53/3-66 and IGHV3-74 mAbs

(Figure 6H). However, the best IGHV3-74 mAb, KXD355 can neutralize all the tested sarbecoviruses

with IC50s ranging from 0.003 to 0.151 µg/ml, further highlighting that KXD355 is highly tolerant to

variations on its epitope (Figure 5A, B).

To evaluate the likelihood that these bnAbs are escaped by future circulating SARS-CoV-2 strains,

we collected 25332 spike sequences published between June 1 to October 1 in 2024. With KP.3 as

reference, we analyzed the mutations on the epitopes of these bnAbs. With extremely low frequency,

multiple confirmed escape mutations for IGHV3-53/3-66 mAbs are found in the circulating strains,

including D420N (0.008%), A475V (0.288%), N487K (0.008%) (Figure S7A). In contrast, confirmed

escape mutations for IGHV3-74 mAbs are not found in the circulating strains (Figure S7B). Overall,

these data predict that IGHV3-74 mAbs, particular KXD355, will not be easily escaped by SARS-CoV-

2 in near future.

Discussion

The key scientific question of this study is to what extent B cells imprinted by ancestral SARS-CoV-2

can develop neutralizing breadth and potency in immune recalls. Strategically, we used WT spike as

bait to track those B cells as they typically maintain specificities to WT spike in immune recalls

(Johnston et al., 2024; Kaku et al., 2022; Kaku et al., 2023; Liang et al., 2024; Quandt et al., 2022;

Sokal et al., 2023; Tortorici et al., 2024; Wang et al., 2023a; Wang et al., 2022b; Weber et al., 2023;

Yisimayi et al., 2024) . A recent study pointed out that memory B cells induced by ancestral SARS-

CoV-2 could redirect their specificities to emerging variants in repeated Omicron exposures (Kotaki

et al., 2024). As our focus was memory B cells broadly recognizing SARS-CoV-2 from WT to emerging

variants, we did not take into account those memory B cells with redirected specificities. Regarding

to the samples, we collected blood from 2 individuals of mRNA vaccines at four time points, which are

2 months, 3 months, 6 months and 9 months after BA.5 or BF.7 breakthrough infection. In addition,

Donor 1 experienced a symptomatic reinfection at 11 days before the last time point, while Donor 2

experienced an asymptomatic reinfection shortly before the last time point, as suggested by our data.

Therefore, the timeline of this longitudinal study is ranging from convalescence of breakthrough

infection to acute phase of reinfection.

A major finding of this study is that mAbs cloned after reinfection have superior neutralizing breadth

and potency, which is accompanied with expansion of pre-existing clonotypes and increased mutation

levels. According to the antigen exposure history of these two donors, this finding can be explained

by two possibilities. First, a fraction of pre-existing memory B cells, which have already undergone

extensive maturation and developed superior neutralizing breadth and potency, are recalled by

reinfection. Second, a fraction of pre-existing memory B cells are recalled by reinfection and then

acquire further maturation in new GC responses. Non-mutually exclusive, these two possibilities differ

in whether an addition round of maturation occurs after reinfection. Previous studies suggest that

BA.1 breakthrough infection or booster vaccination can both lead to affinity maturation of pre-existing

memory B cells induced by mRNA vaccines (Alsoussi et al., 2023; Kaku et al., 2023; Sokal et al.,

2023). Here, it remains to be determined whether memory B cells induced by mRNA vaccines can

undergo two rounds of affinity maturation sequentially driven by breakthrough infection and reinfection.

A highlight of this study is the identification of 4 bnAbs against sarbecoviruses. These bnAbs belong

to a clonotype encoded by IGHV3-74, which is rarely used by RBD-specific antibodies (Wang et al.,

2022a). Structural analysis indicates that these IGHV3-74 bnAbs bind to an epitope consisting of

E340-R346, V367-S375, N436-G446 and Q498-V503. In an early study, epitopes on RBD are

classified into seven groups, with RBD-1 to RBD-3 on the top face, RBD-4 and RBD-5 on the outer

face, RBD-6 and RBD-7 on the inner face (Hastie et al., 2021). More recently, a cryptic epitope

recognized by human mAb BIOLS56 and IMCAS74 is defined as RBD-8 (Rao et al., 2023). Referred

to these epitope categories, the epitope of these IGHV3-74 bnAbs stretches across RBD-3 and RBD-

and is therefore a novel epitope. Unexpectedly, this epitope is not conserved among sarbecoviruses although IGHV3-74 bnAbs can broadly neutralize those viruses. According to structures predicted by AlphaFold 3, we speculate that the resilience of IGHV3-74 bnAbs to epitope variations is due to the plasticity of their long CDRH3s, which play a key role in contacting with this epitope.

As a byproduct of defining a novel broadly neutralizing epitope on RBD, our study represents an example of epitope mapping with AlphaFold 3. In our practice with AlphaFold 3, it is straightforward to predict the structures of IGHV3-53/3-66 mAbs, probably because the structures of similar antibodies have been abundantly reported. In contrast, various structures are predicted for IGHV3-74 bnAbs. In this case, traditional epitope mapping through competition ELISA is necessary to pick the best candidates for further confirmation. Overall, our attempt to apply AlphaFold 3 in fine epitope mapping turned out to be successful.

Similar to a study published recently (Jian et al., 2024), we identified multiple IGHV3-53/3-66 public antibodies with ultra-broad neutralizing activities against SARS-CoV-2. In addition, we observed the expansion of clonotypes encoded by IGHV3-53/3-66. According to phylogenetic analysis, the improved neutralizing breadth and potency of IGHV3-53/3-66 public antibodies is somehow not always correlated to the extent of somatic hypermutation, supporting the view that bystander mutations generated in a stochastic manner are critical for the development of bnAbs against SARS-CoV-2 (Bruhn et al., 2024; Korenkov et al., 2023). On the other hand, the neutralizing breadth of IGHV3-53/3-66 public antibodies reported so far is still limited to SARS-CoV-2 variants. It remains to be investigated whether this class of public antibodies can develop pan-sarbecovirus neutralization after further maturation.

Currently, ancestral SARS-CoV-2 immune imprinting has been acknowledged as a great challenge

for the update of SARS-CoV-2 vaccines (Johnston et al., 2024; Liang et al., 2024; Tortorici et al., 2024;

Yisimayi et al., 2024). In one preprint we posted recently, we find that immune imprinting can facilitate

the development of bnAbs against SARS-CoV-2 (Wang et al., 2024 Preprint). Here, we further show

that B cells imprinted by ancestral SARS-CoV-2 can even develop pan-sarbecovirus neutralization.

Taken together, these studies support that ancestral SARS-CoV-2 immune imprinting can be

harnessed in developing pan-SARS-CoV-2 and even pan-sarbecovirus vaccines. From the aspect of

antibody therapeutics, KXD355 exhibits broad and potent neutralizing activity against sarbecoviruses.

More importantly, KXD355 targets a novel epitope and tolerates mutations on this epitope. Therefore,

KXD355 is an ideal candidate for antibody therapeutics against future SARS-CoV-2 variants and

related sarbecoviruses.

Materials and methods

Ethics statement

This study was approved by the Ethics Committee of Hefei Institutes of Physical Science, Chinese

Academy of Sciences (Approval Number: YXLL-2023-47). All donors provided written informed

consent for collection of information, analysis of plasma and PBMCs, and publication of data

generated from their samples.

Human samples

Peripheral blood samples were collected from 2 donors. Plasma and PBMCs were isolated from blood

through Ficoll density gradient centrifugation.

Protein expression and purification

The extracellular domains of spikes from WT, BA.4/5, XBB.1.5, EG.5.1, JN.1 were constructed with a

foldon-trimerization-motif, a His-tag, a tandem Strep-tag II and a FLAG-tag at C-terminal. S1, NTD

and RBD of WT spike were constructed with a Strep-tag II at C-terminal. S2 of WT spike was

constructed with a His-tag at C-terminal. Human ACE2 (huACE2, 1-740) was constructed with a

Strep-tag II at N-terminal. All recombinant proteins were expressed in FreeStyle 293F cells and

purified with Streptactin Agarose Resin 4FF (Yeasen, 20495ES60) or Ni-NTA Agarose (Qiagen,

30210).

Cloning and expression of mAbs

As previously described (Li et al., 2023), PBMCs were first incubated with 200nM WT spike. After

wash, cells were stained with antibodies targeting CD3, CD19, CD27, CD38, human IgM, human IgD,

His-tag and FLAG-tag along with DAPI. WT-spike-specific B cells were gated as DAPI$^-$

CD3$^-$CD19$^+$CD27$^+$IgD$^-$His$^+$FLAG$^+$ and sorted into 96-well PCR plates with one cell per well. After

reverse transcription reaction, variable regions of antibody heavy and light chains from single B cells

were amplified by nested PCR and cloned into human IgG1 expression vectors. Paired heavy and

light chain plasmids were co-transfected into HEK293T cells or FreeStyle 293F cells. Antibodies were

purified with Protein A magnetic beads (GenScript, L00273).

ELISA

Indicated antigens were coated onto 96-well ELISA plates (100 ng/well) at 4˚C overnight. After

blocking with blocking buffer (PBS with 10% FBS), undiluted HEK293T supernatant was added to the

wells and incubated at 37˚C for 1hr. After wash, HRP-conjugated goat anti-human-IgG antibodies

(Zen-bio, 550004) were added and incubated at 37˚C for 1hr. After wash, TMB substrate (Sangon

Biotech, E661007-0100) was added and incubated at room temperature for 5 minutes. Then the

reaction was stopped with stop solution and absorbance at 450 nm (OD450) was measured.

Pseudovirus neutralization assay

To generate pseudoviruses, HEK293T cells were transfected with psPAX2, pLenti-luciferase and

spike-encoding plasmids (WT, Alpha, Beta, Gamma, Delta, BA.1, BA.4/5, BA.2.75, BQ.1, XBB,

XBB.1.5, EG.5.1, HK.3, BA.2.86, JN.1, KP.2, KP.3, SARS-CoV-1, Pangolin CoV GD, Bat CoV WIV16,

Bat CoV RaTG13 and KP.3 mutants) using polyetherimide (PEI). Pseudoviruses supernatants were

collected 48hrs after transfection. 3-fold serially diluted plasma (starting at 1:20), HEK293T

supernatant (starting at 1:4.5), or mAbs (starting at 10 μg/ml) were mixed with pseudoviruses at 37˚C

for 1hr. HEK293T-hACE2 cells (1.5×10^4 per well) were added into the mixture and incubated at 37˚C

for 48hrs. Cells were lysed to measure luciferase activity by Bright-Lite Luciferase Assay System

(Vazyme Biotech, DD1204-02). The percentages of neutralization were determined by comparing with

the virus control. The data were analyzed and plotted with GraphPad Prism.

Competition ELISA

Purified huACE2, CR3022, S309, P2S-2E9, BD55-5483, BD56-1854 and BD55-1205 were labeled

with HRP (ProteinTech, PK20001). WT RBD was coated onto 96-well ELISA plates (50 ng/well). After

blocking with blocking buffer (PBS with 10% FBS), blocking mAbs were added and incubated at 37

˚C for 1 hr. Then HRP-labeled detecting reagents were added in the presence of blocking mAbs and

incubated at 37˚C for 1 hr. The ratio of blocking and detecting reagents was 100:1. After wash, TMB

substrate (Sangon Biotech, E661007-0100) was added and incubated at room temperature for 30

minutes. Then the reaction was stopped with stop solution and absorbance at 450 nm (OD450) was

measured. The percentages of signal reduction caused by blocking mAbs were calculated.

Binding activity measured by FACS

The plasmids encoding spikes of WT, BA.4/5, KP.3 and KP.3 mutants were transfected into HEK293T

cells. After 48hr, cells were harvested. To compare the binding activity of germline and mature

antibodies to WT and BA.4/5 spikes, cells were incubated with 3-fold serially diluted germline or

mature mAbs (starting at 10 µg/ml) for 30 min at 4°C. To compare the binding with KP.3 and KP.3

mutants, cells were incubated with 1 µg/ml mAbs or 10 nM biotinylated ACE2 for 30 min at 4°C. After

wash, cells stained with mAbs were further stained with DAPI and goat anti-human IgG FITC

(Proteintech, SA00003-12), whereas cells stained with ACE2 were further stained with DAPI and

streptavidin APC (Biolegend, 405243). After wash, cells were analyzed by CytoFLEX (BECKMAN

COULTER). FACS data were analyzed with Flowjo V10 to determine the percentages of positive cells.

To compare the binding activity of germline and mature antibodies to WT and BA.4/5 spikes, the

percentages were ploted with GraphPad Prism. To compare the binding with KP.3 and KP.3 mutants,

the percentages for KP.3 mutants were divided by the percentages for KP.3 and the ratios were

recorded.

Sequence analysis

Variable regions of antibody heavy and light chains from single B cells were analyzed with IgBlast

(https://www.ncbi.nlm.nih.gov/igblast/). Using Python 3.12, clonotypes were analyzed with the Bio,

pandas, and numpy packages. Antibodies with same V and J genes and more than 85% similarity in

CDR3 (nucleotide sequence) for both heavy and light chains were defined as clones within the same

clonotype. Spike sequences, uploaded from June 1 2024 to October 1 2024, were obtained from

NCBI Virus. Sequences with lengths outside the range of 1250–1290 or containing 'X' were removed.

Sequence alignment was performed using Muscle5, and statistical analysis was conducted with

Python 3.12.

Statistics and reproducibility

All statistical analyses were performed with GraphPad Prism. The numbers of biological repeats for

each experiment and tests for statistical significance are described in corresponding figure legends.

Supplemental material

Figures S1–S7

Data S1. Excel spreadsheet containing numerical data for figures.

Data Availability Statement

All relevant data are within the manuscript and its supplemental files.

Acknowledgements

The authors thank the donors for providing peripheral blood. The authors thank Professor Linqi Zhang and Peng Chen at Tsinghua University for confirming the pan-sarbecovirus neutralization of IGHV3-74 mAbs, and for providing plasmids encoding spikes of Pangolin CoV GD, Bat CoV WIV16, Bat CoV RaTG13 and plasmids encoding the heavy and light chains of mAb P2S-2E9. This work is supported by the National Key Plan for Scientific Research and Development of China (Grant No. 2022YFC2305800) and the National Natural Science Foundation of China (Grant No. 32200765, Grant No. 32370941).

Author contributions

Conceptualization, L.J. and T.Z.; Validation, X.C., L.L., R.D., Z.W.; Formal Analysis, X.C., L.L., R.D., Z.W. and Y.L.; Investigation, X.C., L.L., R.D., Z.W., Y.L., Y.S., R.Q., H.F., L.H., X.C., M.L., X.H. and L.J.; Writing – Original Draft, X.C., L.L., R.D., Z.W., Y.L., L.J. and T.Z.; Writing – Review & Editing, L.J. and T.Z.; Visualization, X.C., L.L., Z.W. and Y.L.; Supervision, L.J. and T.Z.; Funding Acquisition, L.J. and T.Z..

Competing interests

L.J., T.Z., X.C., L.L., R.D. and Z.W. are inventors on a patent application for antibodies including

KXD350, KXD352, KXD355, KXD358. The other authors have no competing interests.

References

Abramson, J., J. Adler, J. Dunger, R. Evans, T. Green, A. Pritzel, O. Ronneberger, L. Willmore, A.J. Ballard, J.
Bambrick, S.W. Bodenstein, D.A. Evans, C.C. Hung, M. O'Neill, D. Reiman, K. Tunyasuvunakool, Z.
Wu, A. Zemgulyte, E. Arvaniti, C. Beattie, O. Bertolli, A. Bridgland, A. Cherepanov, M. Congreve, A.I.
Cowen-Rivers, A. Cowie, M. Figurnov, F.B. Fuchs, H. Gladman, R. Jain, Y.A. Khan, C.M.R. Low, K.
Perlin, A. Potapenko, P. Savy, S. Singh, A. Stecula, A. Thillaisundaram, C. Tong, S. Yakneen, E.D.
Zhong, M. Zielinski, A. Zidek, V. Bapst, P. Kohli, M. Jaderberg, D. Hassabis, and J.M. Jumper. 2024.
Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* 630:493-500.
Alsoussi, W.B., S.K. Malladi, J.Q. Zhou, Z.M. Liu, B.L. Ying, W. Kim, A.J. Schmitz, T.T. Lei, S.C. Horvath, A.J.
Sturtz, K.M. McIntire, B. Evavold, F.J. Han, S.M. Scheaffer, I.F. Fox, S.F. Mirza, L. Parra-Rodriguez, R.
Nachbagauer, B. Nestorova, S. Chalkias, C.W. Farnsworth, M.K. Klebert, I. Pusic, B.S. Strnad, W.D.
Middleton, S.A. Teefey, S.P.J. Whelan, M.S. Diamond, R. Paris, J.A. O'Halloran, R.M. Presti, J.S. Turner,
and A.H. Ellebedy. 2023. SARS-CoV-2 Omicron boosting induces de novo B cell response in humans.
Nature 617:592-+.
Batista, F.D., and N.E. Harwood. 2009. The who, how and where of antigen presentation to B cells. *Nat Rev
Immunol* 9:15-27.
Bruhn, M., M. Obara, A. Salam, B. Costa, A. Ziegler, I. Waltl, A. Pavlou, M. Hoffmann, T. Graalmann, S.
Pöhlmann, A. Schambach, and U. Kalinke. 2024. Diversification of the VH3-53 immunoglobulin gene
segment by somatic hypermutation results in neutralization of SARS-CoV-2 virus variants. *Eur J
Immunol* 54:
Cao, Y.L., F.C. Jian, J. Wang, Y.L. Yu, W.L. Song, A. Yisimayi, J. Wang, R. An, X.S. Chen, N. Zhang, Y. Wang,
P. Wang, L.J. Zhao, H.Y. Sun, L.L. Yu, S.J. Yang, X. Niu, T.H. Xiao, Q.Q. Gu, F. Shao, X.H. Hao, Y.L.
Xu, R.H. Jin, Z.Y. Shen, Y.C. Wang, and X.S. Xie. 2023. Imprinted SARS-CoV-2 humoral immunity
induces convergent Omicron RBD evolution. *Nature* 614:521-+.
Cao, Y.L., F.C. Jian, Z.Y. Zhang, A. Yisimayi, X.H. Hao, L.L. Bao, F. Yuan, Y.L. Yu, S. Du, J. Wang, T.H. Xiao,
W.L. Song, Y. Zhang, P.L. Liu, R. An, P. Wang, Y. Wang, S.J. Yang, X. Niu, Y.H. Zhang, Q.Q. Gu, F.
Shao, Y.L. Hu, W.D. Yin, A.H. Zheng, Y.C. Wang, C. Qin, R.H. Jin, J.Y. Xiao, and X.L.S. Xie. 2022.
Rational identification of potent and broad sarbecovirus-neutralizing antibody cocktails from SARS
convalescents. *Cell Reports* 41:
Cerutti, G., Y. Guo, T. Zhou, J. Gorman, M. Lee, M. Rapp, E.R. Reddem, J. Yu, F. Bahna, J. Bimela, Y. Huang,
P.S. Katsamba, L. Liu, M.S. Nair, R. Rawi, A.S. Olia, P. Wang, B. Zhang, G.Y. Chuang, D.D. Ho, Z.
Sheng, P.D. Kwong, and L. Shapiro. 2021. Potent SARS-CoV-2 neutralizing antibodies directed against
spike N-terminal domain target a single supersite. *Cell Host Microbe* 29:819-833 e817.
Gaebler, C., Z. Wang, J.C.C. Lorenzi, F. Muecksch, S. Finkin, M. Tokuyama, A. Cho, M. Jankovic, D. Schaefer-
Babajew, T.Y. Oliveira, M. Cipolla, C. Viant, C.O. Barnes, Y. Bram, G. Breton, T. Hagglof, P. Mendoza,
A. Hurley, M. Turroja, K. Gordon, K.G. Millard, V. Ramos, F. Schmidt, Y. Weisblum, D. Jha, M.
Tankelevich, G. Martinez-Delgado, J. Yee, R. Patel, J. Dizon, C. Unson-O'Brien, I. Shimeliovich, D.F.
Robbiani, Z. Zhao, A. Gazumyan, R.E. Schwartz, T. Hatziioannou, P.J. Bjorkman, S. Mehandru, P.D.
Bieniasz, M. Caskey, and M.C. Nussenzweig. 2021. Evolution of antibody immunity to SARS-CoV-2.
Nature 591:639-644.
Hastie, K.M., H. Li, D. Bedinger, S.L. Schendel, S.M. Dennison, K. Li, V. Rayaprolu, X. Yu, C. Mann, M.

Zandonatti, R. Diaz Avalos, D. Zyla, T. Buck, S. Hui, K. Shaffer, C. Hariharan, J. Yin, E. Olmedillas, A.
Enriquez, D. Parekh, M. Abraha, E. Feeney, G.Q. Horn, V.I.C.D.B.t. Co, Y. Aldon, H. Ali, S. Aracic, R.R.
Cobb, R.S. Federman, J.M. Fernandez, J. Glanville, R. Green, G. Grigoryan, A.G. Lujan Hernandez,
D.D. Ho, K.A. Huang, J. Ingraham, W. Jiang, P. Kellam, C. Kim, M. Kim, H.M. Kim, C. Kong, S.J. Krebs,
F. Lan, G. Lang, S. Lee, C.L. Leung, J. Liu, Y. Lu, A. MacCamy, A.T. McGuire, A.L. Palser, T.H. Rabbitts,
Z. Rikhtegaran Tehrani, M.M. Sajadi, R.W. Sanders, A.K. Sato, L. Schweizer, J. Seo, B. Shen, J.L.
Snitselaar, L. Stamatatos, Y. Tan, M.T. Tomic, M.J. van Gils, S. Youssef, J. Yu, T.Z. Yuan, Q. Zhang, B.
Peters, G.D. Tomaras, T. Germann, and E.O. Saphire. 2021. Defining variant-resistant epitopes
targeted by SARS-CoV-2 antibodies: A global consortium study. *Science* 374:472-478.
Huang, C.Q., S. Vishwanath, G.W. Carnell, A.C.Y. Chan, and J.L. Heeney. 2023. Immune imprinting and next-
generation coronavirus vaccines. *Nat Microbiol* 8:1971-1985.
Inoue, T., I. Moran, R. Shinnakasu, T.G. Phan, and T. Kurosaki. 2018. Generation of memory B cells and their
reactivation. *Immunol Rev* 283:138-149.
Jian, F., J. Wang, A. Yisimayi, W. Song, Y. Xu, X. Chen, X. Niu, S. Yang, Y. Yu, P. Wang, H. Sun, L. Yu, J. Wang,
Y. Wang, R. An, W. Wang, M. Ma, T. Xiao, Q. Gu, F. Shao, Y. Wang, Z. Shen, R. Jin, and Y. Cao. 2024.
Evolving antibody response to SARS-CoV-2 antigenic shift from XBB to JN.1. *Nature*
Johnston, T.S., S.H. Li, M.M. Painter, R.K. Atkinson, N.R. Douek, D.B. Reeg, D.C. Douek, E.J. Wherry, and S.E.
Hensley. 2024. Immunological imprinting shapes the specificity of human antibody responses against
SARS-CoV-2 variants. *Immunity* 57:912-925 e914.
Ju, B., Q. Zhang, Z. Wang, Z.Q. Aw, P. Chen, B. Zhou, R. Wang, X. Ge, Q. Lv, L. Cheng, R. Zhang, Y.H. Wong,
H. Chen, H. Wang, S. Shan, X. Liao, X. Shi, L. Liu, J.J.H. Chu, X. Wang, Z. Zhang, and L. Zhang. 2023.
Infection with wild-type SARS-CoV-2 elicits broadly neutralizing and protective antibodies against
omicron subvariants. *Nat Immunol* 24:690-699.
Kaku, C.I., A.J. Bergeron, C. Ahlm, J. Normark, M. Sakharkar, M.N.E. Forsell, and L.M. Walker. 2022. Recall of
preexisting cross-reactive B cell memory after Omicron BA.1 breakthrough infection. *Sci Immunol*
7:eabq3511.
Kaku, C.I., T.N. Starr, P. Zhou, H.L. Dugan, P. Khalife, G. Song, E.R. Champney, D.W. Mielcarz, J.C. Geoghegan,
D.R. Burton, R. Andrabi, J.D. Bloom, and L.M. Walker. 2023. Evolution of antibody immunity following
Omicron BA.1 breakthrough infection. *Nat Commun* 14:2751.
Kim, W., J.Q. Zhou, S.C. Horvath, A.J. Schmitz, A.J. Sturtz, T. Lei, Z. Liu, E. Kalaidina, M. Thapa, W.B. Alsoussi,
A. Haile, M.K. Klebert, T. Suessen, L. Parra-Rodriguez, P.A. Mudd, S.P.J. Whelan, W.D. Middleton, S.A.
Teefey, I. Pusic, J.A. O'Halloran, R.M. Presti, J.S. Turner, and A.H. Ellebedy. 2022. Germinal centre-
driven maturation of B cell response to mRNA vaccination. *Nature* 604:141-145.
Korenkov, M., M. Zehner, H. Cohen-Dvashi, A. Borenstein-Katz, L. Kottege, H. Janicki, K. Vanshylla, T. Weber,
H. Gruell, M. Koch, R. Diskin, C. Kreer, and F. Klein. 2023. Somatic hypermutation introduces bystander
mutations that prepare SARS-CoV-2 antibodies for emerging variants. *Immunity* 56:
Kotaki, R., S. Moriyama, S. Oishi, T. Onodera, Y. Adachi, E. Sasaki, K. Ishino, M. Morikawa, H. Takei, H.
Takahashi, T. Takano, A. Nishiyama, K. Yumoto, K. Terahara, M. Isogawa, T. Matsumura, M. Shinkai,
and Y. Takahashi. 2024. Repeated Omicron exposures redirect SARS-CoV-2-specific memory B cell
evolution toward the latest variants. *Sci Transl Med* 16:eadp9927.
Lapuente, D., T.H. Winkler, and M. Tenbusch. 2024. B-cell and antibody responses to SARS-CoV-2: infection,
vaccination, and hybrid immunity. *Cell Mol Immunol* 21:144-158.
Li, L., X. Chen, Z. Wang, Y. Li, C. Wang, L. Jiang, and T. Zuo. 2023. Breakthrough infection elicits hypermutated
IGHV3-53/3-66 public antibodies with broad and potent neutralizing activity against SARS-CoV-2

variants including the emerging EG.5 lineages. *PLoS Pathog* 19:e1011856.

Liang, C.Y., S. Raju, Z. Liu, Y. Li, G. Asthagiri Arunkumar, J.B. Case, S.M. Scheaffer, S.J. Zost, C.M. Acreman,
M. Gagne, S.F. Andrew, D.C. Carvalho Dos Anjos, K.E. Foulds, J.S. McLellan, J.E. Crowe, Jr., D.C.
Douek, S.P.J. Whelan, S.M. Elbashir, D.K. Edwards, and M.S. Diamond. 2024. Imprinting of serum
neutralizing antibodies by Wuhan-1 mRNA vaccines. *Nature* 630:950-960.

Ling, Z., C. Yi, X. Sun, Z. Yang, and B. Sun. 2023. Broad strategies for neutralizing SARS-CoV-2 and other
human coronaviruses with monoclonal antibodies. *Sci China Life Sci* 66:658-678.

Liu, H., and I.A. Wilson. 2022. Protective neutralizing epitopes in SARS-CoV-2. *Immunol Rev* 310:76-92.

McCallum, M., A. De Marco, F.A. Lempp, M.A. Tortorici, D. Pinto, A.C. Walls, M. Beltramello, A. Chen, Z. Liu, F.
Zatta, S. Zepeda, J. di Iulio, J.E. Bowen, M. Montiel-Ruiz, J. Zhou, L.E. Rosen, S. Bianchi, B. Guarino,
C.S. Fregni, R. Abdelnabi, S.C. Foo, P.W. Rothlauf, L.M. Bloyet, F. Benigni, E. Cameroni, J. Neyts, A.
Riva, G. Snell, A. Telenti, S.P.J. Whelan, H.W. Virgin, D. Corti, M.S. Pizzuto, and D. Veesler. 2021. N-
terminal domain antigenic mapping reveals a site of vulnerability for SARS-CoV-2. *Cell* 184:2332-2347
e2316.

Paciello, I., G. Pierleoni, E. Pantano, G. Antonelli, P. Pileri, G. Maccari, D. Cardamone, G. Realini, F. Perrone,
M.M. Neto, S. Pozzessere, M. Fabbiani, F. Panza, I. Rancan, M. Tumbarello, F. Montagnani, D. Medini,
P. Maes, N. Temperton, E. Simon-Loriere, O. Schwartz, R. Rappuoli, and E. Andreano. 2024. Antigenic
sin and multiple breakthrough infections drive converging evolution of COVID-19 neutralizing responses.
Cell Rep 43:114645.

Pinto, D., Y.J. Park, M. Beltramello, A.C. Walls, M.A. Tortorici, S. Bianchi, S. Jaconi, K. Culap, F. Zatta, A. De
Marco, A. Peter, B. Guarino, R. Spreafico, E. Cameroni, J.B. Case, R.E. Chen, C. Havenar-Daughton,
G. Snell, A. Telenti, H.W. Virgin, A. Lanzavecchia, M.S. Diamond, K. Fink, D. Veesler, and D. Corti. 2020.
Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV antibody. *Nature* 583:290-
295.

Qi, H., B. Liu, X. Wang, and L. Zhang. 2022. The humoral response and antibodies against SARS-CoV-2
infection. *Nat Immunol* 23:1008-1020.

Quandt, J., A. Muik, N. Salisch, B.G. Lui, S. Lutz, K. Kruger, A.K. Wallisch, P. Adams-Quack, M. Bacher, A.
Finlayson, O. Ozhelvaci, I. Vogler, K. Grikscheit, S. Hoehl, U. Goetsch, S. Ciesek, O. Tureci, and U.
Sahin. 2022. Omicron BA.1 breakthrough infection drives cross-variant neutralization and memory B
cell formation against conserved epitopes. *Sci Immunol* 7:eabq2427.

Rao, X., R.C. Zhao, Z. Tong, S.X. Guo, W.Y. Peng, K.F. Liu, S.H. Li, L.L. Wu, J.Y. Tong, Y. Chai, P. Han, F.R.
Wang, P. Jia, Z.H. Li, X. Zhao, D.D. Li, R. Zhang, X. Zhang, W.W. Zou, W.W. Li, Q.H. Wang, G.F. Gao,
Y. Wu, L.P. Dai, and F. Gao. 2023. Defining a de novo non- RBM antibody as RBD-8 and its synergistic
rescue of immune- evaded antibodies to neutralize Omicron SARS- CoV-2. *P Natl Acad Sci USA* 120:

Robbiani, D.F., C. Gaebler, F. Muecksch, J.C.C. Lorenzi, Z. Wang, A. Cho, M. Agudelo, C.O. Barnes, A.
Gazumyan, S. Finkin, T. Hagglof, T.Y. Oliveira, C. Viant, A. Hurley, H.H. Hoffmann, K.G. Millard, R.G.
Kost, M. Cipolla, K. Gordon, F. Bianchini, S.T. Chen, V. Ramos, R. Patel, J. Dizon, I. Shimeliovich, P.
Mendoza, H. Hartweger, L. Nogueira, M. Pack, J. Horowitz, F. Schmidt, Y. Weisblum, E. Michailidis,
A.W. Ashbrook, E. Waltari, J.E. Pak, K.E. Huey-Tubman, N. Koranda, P.R. Hoffman, A.P. West, Jr., C.M.
Rice, T. Hatziioannou, P.J. Bjorkman, P.D. Bieniasz, M. Caskey, and M.C. Nussenzweig. 2020.
Convergent antibody responses to SARS-CoV-2 in convalescent individuals. *Nature* 584:437-442.

Roltgen, K., and S.D. Boyd. 2024. Antibody and B Cell Responses to SARS-CoV-2 Infection and Vaccination:
The End of the Beginning. *Annu Rev Pathol* 19:69-97.

Sakharkar, M., C.G. Rappazzo, W.F. Wieland-Alter, C.L. Hsieh, D. Wrapp, E.S. Esterman, C.I. Kaku, A.Z. Wec,

J.C. Geoghegan, J.S. McLellan, R.I. Connor, P.F. Wright, and L.M. Walker. 2021. Prolonged evolution of the human B cell response to SARS-CoV-2 infection. *Sci Immunol* 6:

Sokal, A., G. Barba-Spaeth, L. Hunault, I. Fernandez, M. Broketa, A. Meola, S. Fourati, I. Azzaoui, A. Vandenberghe, P. Lagouge-Roussey, M. Broutin, A. Roeser, M. Bouvier-Alias, E. Crickx, L. Languille, M. Fournier, M. Michel, B. Godeau, S. Gallien, G. Melica, Y. Nguyen, F. Canoui-Poitrine, F. Pirenne, J. Megret, J.M. Pawlotsky, S. Fillatreau, C.A. Reynaud, J.C. Weill, F.A. Rey, P. Bruhns, M. Mahevas, and P. Chappert. 2023. SARS-CoV-2 Omicron BA.1 breakthrough infection drives late remodeling of the memory B cell repertoire in vaccinated individuals. *Immunity* 56:2137-2151 e2137.

Sokal, A., P. Chappert, G. Barba-Spaeth, A. Roeser, S. Fourati, I. Azzaoui, A. Vandenberghe, I. Fernandez, A. Meola, M. Bouvier-Alias, E. Crickx, A. Beldi-Ferchiou, S. Hue, L. Languille, M. Michel, S. Baloul, F. Noizat-Pirenne, M. Luka, J. Megret, M. Menager, J.M. Pawlotsky, S. Fillatreau, F.A. Rey, J.C. Weill, C.A. Reynaud, and M. Mahevas. 2021. Maturation and persistence of the anti-SARS-CoV-2 memory B cell response. *Cell* 184:1201-1213 e1214.

Tonegawa, S. 1983. Somatic generation of antibody diversity. *Nature* 302:575-581.

Tortorici, M.A., A. Addetia, A.J. Seo, J. Brown, K. Sprouse, J. Logue, E. Clark, N. Franko, H. Chu, and D. Veesler. 2024. Persistent immune imprinting occurs after vaccination with the COVID-19 XBB.1.5 mRNA booster in humans. *Immunity* 57:904-911 e904.

Turner, J.S., J.A. O'Halloran, E. Kalaidina, W. Kim, A.J. Schmitz, J.Q. Zhou, T. Lei, M. Thapa, R.E. Chen, J.B. Case, F. Amanat, A.M. Rauseo, A. Haile, X. Xie, M.K. Klebert, T. Suessen, W.D. Middleton, P.Y. Shi, F. Krammer, S.A. Teefey, M.S. Diamond, R.M. Presti, and A.H. Ellebedy. 2021. SARS-CoV-2 mRNA vaccines induce persistent human germinal centre responses. *Nature* 596:109-113.

Viana, R., S. Moyo, D.G. Amoako, H. Tegally, C. Scheepers, C.L. Althaus, U.J. Anyaneji, P.A. Bester, M.F. Boni, M. Chand, W.T. Choga, R. Colquhoun, M. Davids, K. Deforche, D. Doolabh, L. du Plessis, S. Engelbrecht, J. Everatt, J. Giandhari, M. Giovanetti, D. Hardie, V. Hill, N.Y. Hsiao, A. Iranzadeh, A. Ismail, C. Joseph, R. Joseph, L. Koopile, S.L. Kosakovsky Pond, M.U.G. Kraemer, L. Kuate-Lere, O. Laguda-Akingba, O. Lesetedi-Mafoko, R.J. Lessells, S. Lockman, A.G. Lucaci, A. Maharaj, B. Mahlangu, T. Maponga, K. Mahlakwane, Z. Makatini, G. Marais, D. Maruapula, K. Masupu, M. Matshaba, S. Mayaphi, N. Mbhele, M.B. Mbulawa, A. Mendes, K. Mlisana, A. Mnguni, T. Mohale, M. Moir, K. Moruisi, M. Mosepele, G. Motsatsi, M.S. Motswaledi, T. Mphoyakgosi, N. Msomi, P.N. Mwangi, Y. Naidoo, N. Ntuli, M. Nyaga, L. Olubayo, S. Pillay, B. Radibe, Y. Ramphal, U. Ramphal, J.E. San, L. Scott, R. Shapiro, L. Singh, P. Smith-Lawrence, W. Stevens, A. Strydom, K. Subramoney, N. Tebeila, D. Tshiabuila, J. Tsui, S. van Wyk, S. Weaver, C.K. Wibmer, E. Wilkinson, N. Wolter, A.E. Zarebski, B. Zuze, D. Goedhals, W. Preiser, F. Treurnicht, M. Venter, C. Williamson, O.G. Pybus, J. Bhiman, A. Glass, D.P. Martin, A. Rambaut, S. Gaseitsiwe, A. von Gottberg, and T. de Oliveira. 2022. Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. *Nature* 603:679-686.

Victora, G.D., and M.C. Nussenzweig. 2022. Germinal Centers. *Annu Rev Immunol* 40:413-442.

Wang, Q., Y. Guo, A.R. Tam, R. Valdez, A. Gordon, L. Liu, and D.D. Ho. 2023a. Deep immunological imprinting due to the ancestral spike in the current bivalent COVID-19 vaccine. *Cell Rep Med* 4:101258.

Wang, Q., Y.C. Guo, R.M. Zhang, J.R. Ho, H. Mohri, R. Valdez, D.M. Manthei, A. Gordon, L.H. Liu, and D.V. Ho. 2023b. Antibody neutralisation of emerging SARS-CoV-2 subvariants: EG.5.1 and XBC.1.6. *Lancet Infect Dis* 23:E397-E398.

Wang, Y., M. Yuan, H. Lv, J. Peng, I.A. Wilson, and N.C. Wu. 2022a. A large-scale systematic survey reveals recurring molecular features of public antibody responses to SARS-CoV-2. *Immunity* 55:1105-1117 e1104.

Wang, Z., L. Li, R. Du, X. Chen, Y. Sun, R. Qin, Y. Li, H. Feng, L. Hu, X. Chen, M. Lu, L. Jiang, and T. Zuo. 2024.
Ancestral SARS-CoV-2 immune imprinting persists on RBD but not NTD after sequential Omicron
infections. *bioRxiv*. doi.org/10.1101/2024.05.30.596664 (Preprint posted July 04, 2024)

Wang, Z., F. Schmidt, Y. Weisblum, F. Muecksch, C.O. Barnes, S. Finkin, D. Schaefer-Babajew, M. Cipolla, C.
Gaebler, J.A. Lieberman, T.Y. Oliveira, Z. Yang, M.E. Abernathy, K.E. Huey-Tubman, A. Hurley, M.
Turroja, K.A. West, K. Gordon, K.G. Millard, V. Ramos, J. Da Silva, J. Xu, R.A. Colbert, R. Patel, J.
Dizon, C. Unson-O'Brien, I. Shimeliovich, A. Gazumyan, M. Caskey, P.J. Bjorkman, R. Casellas, T.
Hatziioannou, P.D. Bieniasz, and M.C. Nussenzweig. 2021. mRNA vaccine-elicited antibodies to SARS-
CoV-2 and circulating variants. *Nature* 592:616-622.

Wang, Z., P. Zhou, F. Muecksch, A. Cho, T. Ben Tanfous, M. Canis, L. Witte, B. Johnson, R. Raspe, F. Schmidt,
E. Bednarski, J. Da Silva, V. Ramos, S. Zong, M. Turroja, K.G. Millard, K.H. Yao, I. Shimeliovich, J.
Dizon, A. Kaczynska, M. Jankovic, A. Gazumyan, T.Y. Oliveira, M. Caskey, C. Gaebler, P.D. Bieniasz,
T. Hatziioannou, and M.C. Nussenzweig. 2022b. Memory B cell responses to Omicron subvariants after
SARS-CoV-2 mRNA breakthrough infection in humans. *J Exp Med* 219:

Weber, T., S. Dahling, S. Rose, P. Affeldt, K. Vanshylla, L. Ullrich, L. Gieselmann, F. Teipel, H. Gruell, V. Di
Cristanziano, D.S. Kim, G. Georgiou, M. Koch, C. Kreer, and F. Klein. 2023. Enhanced SARS-CoV-2
humoral immunity following breakthrough infection builds upon the preexisting memory B cell pool. *Sci
Immunol* 8:eadk5845.

Yan, Q., R. Hou, X. Huang, Y. Zhang, P. He, Y. Zhang, B. Liu, Q. Wang, H. Rao, X. Chen, X. Zhao, X. Niu, J.
Zhao, X. Xiong, and L. Chen. 2022. Shared IGHV1-69-encoded neutralizing antibodies contribute to
the emergence of L452R substitution in SARS-CoV-2 variants. *Emerg Microbes Infect* 11:2749-2761.

Yisimayi, A., W. Song, J. Wang, F. Jian, Y. Yu, X. Chen, Y. Xu, S. Yang, X. Niu, T. Xiao, J. Wang, L. Zhao, H.
Sun, R. An, N. Zhang, Y. Wang, P. Wang, L. Yu, Z. Lv, Q. Gu, F. Shao, R. Jin, Z. Shen, X.S. Xie, Y.
Wang, and Y. Cao. 2024. Repeated Omicron exposures override ancestral SARS-CoV-2 immune
imprinting. *Nature* 625:148-156.

Yuan, M., H. Liu, N.C. Wu, C.D. Lee, X. Zhu, F. Zhao, D. Huang, W. Yu, Y. Hua, H. Tien, T.F. Rogers, E. Landais,
D. Sok, J.G. Jardine, D.R. Burton, and I.A. Wilson. 2020a. Structural basis of a shared antibody
response to SARS-CoV-2. *Science* 369:1119-1123.

Yuan, M., N.C. Wu, X. Zhu, C.D. Lee, R.T.Y. So, H. Lv, C.K.P. Mok, and I.A. Wilson. 2020b. A highly conserved
cryptic epitope in the receptor binding domains of SARS-CoV-2 and SARS-CoV. *Science* 368:630-633.

Figures and figure legends

Figure 1. Longitudinal analysis of plasma neutralizing antibody responses in two individuals.

(A) Basic information of donors and samples. (B, C) Summary of plasma neutralizing antibody titers

(NT50). NT50s are calculated with results from three independent experiments, in which duplicates

are performed.

Figure 2. Isolation and characterization of mAbs. (A) FACS plots representing the percentages of

WT-spike-specific B cells in DAPI⁻CD3⁻CD19⁺CD27⁺IgD⁻ B cells. (B) ELISA and neutralization results

of 632 mAbs from two donors. Supernatant of HEK293T is tested against 9 antigens by ELISA, and

tested against 5 pseudoviruses by neutralization assay. For ELISA, supernatant is not diluted. For

neutralization, supernatant is diluted by 4.5 folds. The results are mean values from two independent

experiments, in which duplicates are performed. (C) Frequencies of mAbs targeting each domain

among mAbs binding to WT spike (WT-binding mAbs). (D) Frequencies of cross-binding mAbs among

RBD-specific, NTD-specific, and S2-specific mAbs. (E) Frequencies of mAbs targeting RBD or NTD

among mAbs neutralizing WT pseudoviruses (WT-neutralizing mAbs). (F) Frequencies of mAbs with

714   different neutralizing breadth. Breadth indicates the number of pseudoviruses can be neutralized by

715   mAbs. (G) Frequencies of cross-neutralizing mAbs.

Figure 3. Neutralizing potency of mAbs against SARS-CoV-2 and SARS-CoV-1. (A) IC50s of

mAbs against 14 pseudoviruses. The highest antibody concentration used to determine IC50 is 10

μg/ml. IC50s are calculated with results from two independent experiments, in which duplicates are

performed. The black asterisk indicates mAbs with broadly neutralizing activity against SARS-CoV-2

variants from WT to KP.3. The red asterisk indicates mAbs with broadly neutralizing activity against

SARS-CoV-2 and SARS-CoV-1. (B) Frequencies of neutralizing mAbs against each pseudoviruses

724   among mAbs from T1 to T4. (C) Geometric mean IC50s of mAbs from T1 to T4 against each

725   pseudoviruses. (D) Comparison of IC50s. Statistical analysis is performed by Mann-Whitney test. ns

726   (not significant) $P > 0.05$, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

Figure 4. Sequence analysis. (A, B) Frequencies of V genes used by mAbs binding to WT spike

(WT-binding mAbs) and mAbs neutralizing WT pseudoviruses (WT-neutralizing mAbs) from Donor 1

and Donor 2. (C) Clonotype analysis of WT-binding mAbs from Donor 1 and Donor 2. Values in

parentheses indicate number of clonotypes/number of mAbs. Different clonotypes are distinguished

by alternating colors, and identical clonotypes across different time points are connected by gray lines.

(D) Bubble plot of clonotypes comprising two or more mAbs. Dot size indicates number of mAbs. The

red asterisk indicates clonotypes containing neutralizing mAbs. (E) Somatic hypermutation on

variable regions of heavy and light chains based on DNA sequences. Statistical analysis is performed

by two-tailed unpaired T test. *P<0.05, **P < 0.01, ***P < 0.001.

Figure 5. Clonotypes containing mAbs with extraordinary neutralizing breadth and potency.

(A) Maximum-likelihood phylogenetic trees of each clonotype and IC50s of each mAbs. (B)

Neutralization against sarbecoviruses including Pangolin CoV GD (Pangolin-GD), Bat CoV WIV16

(WIV16) and Bat CoV RaTG13-T372A (RaTG13-T372A). (C) Binding of germline antibodies to WT

and BA.4/5 spikes expressed on HEK293T cells.

Figure 6. Epitope mapping and escape mutations of mAbs with extraordinary neutralizing breadth and potency. (A) Competition ELISA. The blocking indicates signal (OD450) reduced by blocking mAbs. The results are mean values from two independent experiments, in which duplicates are performed. (B) Epitope of IGHV3-53/3-66 mAbs on WT RBD (PDB: 7A94). (C) Epitope of ACE2

on WT RBD (PDB: 7A94). (D) The predicted structure of KXD355 in complex with WT RBD is aligned

to up RBD and down RBD on WT spike (PDB: 7A94). (E) Detailed view of KXD355 in complex with

WT RBD. (F) Epitope of KXD355 on WT RBD (PDB: 7A94). (G) Binding and neutralizing activities

against single-mutated KP.3. The binding activities are measured by FACS. NTD-mAb, a mAb

targeting NTD, is used to evaluate the expression levels of KP.3 mutants. The results of binding

activities are mean values from two independent experiments. For neutralizing activities, IC50s are

calculated with results from two independent experiments, in which duplicates are performed. N/A

(not applicable) indicates that neutralization assay is not done because we are not able to generate

pseudoviruses for the mutant. (H) Alignment of spike sequences from SARS-CoV-2 variants and other

sarbecoviruses tested in neutralization assay. The epitopes of IGHV3-53/3-66 and IGHV3-74 mAbs

are respectively highlighted with blue and red background.

Figure S1. Plasma neutralizing antibody titers. (A) Plasma neutralizing antibody titers against a

panel of pseudoviruses. The data are represented as mean ± SD and they are from three independent

experiments, in which duplicates are performed. The published data for plasma from T1 are not used

here. Instead, we measured the neutralizing tiers of plasma from T1 again, along with plasma from

other time points.

770 **Figure S2. Cross-binding and cross-neutralizing mAbs from Donor 1 and Donor 2.** (A)

771 Frequencies of cross-binding mAbs among RBD-specific, NTD-specific, and S2-specific mAbs from

772 Donor 1 and Donor 2. (B) Frequencies of cross-neutralizing mAbs from Donor 1 and Donor 2.

Figure S3. IC50s of mAbs. (A) Summary of IC50s. The numbers below the line indicate the geometric

mean IC50s and the numbers above the line represent the percentage of mAbs that is able to

777   neutralize the corresponding pseudoviruses (IC50 below 10 μg/ml). (B) Frequencies of neutralizing

778   mAbs against each pseudoviruses among T1-T3 group and T4 group. (C) Geometric mean IC50s of

779   mAbs among T1-T3 group and T4 group. (D) Comparison of IC50s. The numbers below and above

780   indicate the geometric mean IC50s and the percentages of neutralizing mAbs, respectively. Statistical

781   analysis is performed by Mann-Whitney test. ns (not significant) $P > 0.05$, $**P < 0.01$, $****P < 0.0001$.

Figure S4. Analysis of V gene usage. (A) Frequencies of V genes used by mAbs targeting RBD, NTD and S2. (B) Paring of V genes for heavy and light chains in neutralizing mAbs. The number in each box indicates the count of mAbs using the corresponding pair.

Figure S5. Published structures and predicted structures. (A) The published structures of ACE2

(PDB: 7A94), CR3022 (PDB: 6W41), S309 (PDB: 7TLY), P2S-2E9 (PDB: 7XSA), BD55-5514 (PBD:

7Y0W), BD55-1205 (PDB: 8XEA) in complex with RBD. (B) The predicted structures of KXD353,

KXD356, KXD357, KXD470 and KXD472 in complex with WT RBD. (C) The predicted structure of

KXD355 in complex with WT RBD is aligned with published structures of CR3022, S309, P2S-2E9,

BD55-5514 and BD55-1205.

	KXD356	KXD357	KXD353	KXD470	KXD472	KXD352	KXD355	KXD350	KXD358
KP.3	0.006	0.089	0.216	0.763	1.635	0.007	0.008	0.026	0.019
K403D	0.043	>10	>10	>10	>10	0.006	0.009	0.041	0.05
D420N	0.126	2.407	>10	>10	>10	0.005	0.007	0.067	0.016
Y421K	0.08	>10	>10	>10	>10	0.002	0.005	0.043	0.023
L456V	0.038	0.958	1.114	>10	>10	0.022	0.014	0.117	0.023
L456S	0.243	>10	>10	>10	>10	0.012	0.015	0.072	0.028
K460F	0.057	>10	3.384	>10	>10	0.004	0.008	0.053	0.023
A475V	0.148	>10	>10	>10	>10	0.005	0.013	0.039	0.016
K484A	0.003	0.015	2.174	2.529	1.612	0.003	0.005	0.018	0.006
K484D	0.007	0.03	0.631	>10	9.051	0.005	0.009	0.037	0.012
N487K	>10	>10	>10	>10	>10	0.006	0.008	0.044	0.017
Y489A	>10	>10	>10	>10	>10	0.003	0.002	0.006	0.004
Y489D	>10	0.986	>10	>10	>10	0.001	0.002	0.008	0.002
R498A	0.385	0.017	>10	3.665	>10	0.004	0.006	0.029	0.011
R498D					N/A				
P499A	0.004	0.053	0.086	>10	5.506	0.008	0.013	0.042	0.026
P499S	0.008	0.077	0.195	>10	>10	0.016	0.029	0.162	0.024
N343A	0.003	0.029	0.022	>10	3.81	0.021	0.01	>10	>10
N343F	0.004	0.051	0.061	7.918	2.951	0.063	0.273	>10	>10
T345A	0.004	0.035	0.104	6.648	3.247	0.013	0.005	>10	1.66
T345F	0.003	0.049	0.05	3.132	2.047	0.001	0.001	>10	0.484
K440D	0.013	0.074	2.37	>10	>10	0.191	0.55	0.141	>10
L441A	0.003	0.095	0.263	0.361	3.986	0.004	0.004	0.013	0.852
L441S	0.004	0.066	0.162	9.55	1.348	>10	>10	>10	>10
K444A	0.008	0.048	0.181	4.152	5.256	0.023	0.297	5.636	0.907
K444D	0.003	0.011	0.366	>10	>10	0.072	0.382	>10	>10
H445D	0.003	0.095	0.263	0.361	3.986	0.004	0.004	0.013	0.852
R346A	0.001	0.023	0.164	5.175	1.807	0.002	0.01	0.042	0.019
R346D	0.002	0.009	0.358	0.803	2.898	0.006	0.014	0.35	0.037
V367A	0.005	0.026	0.039	4.553	3.388	0.013	0.02	0.064	0.056
V367S	0.004	0.045	0.071	4.755	4.669	0.014	0.02	0.04	0.103
F371A	0.003	0.017	0.373	1.352	0.94	0.004	0.007	0.033	0.009
F371S	0.003	0.007	0.08	0.303	0.241	0.005	0.005	0.021	0.005
F374A	0.008	0.016	0.385	0.985	2.43	0.01	0.011	0.054	0.018
F374S	0.004	0.006	0.1	0.39	0.293	0.01	0.01	0.019	0.016
F375A	0.004	0.073	1.254	1.477	4.554	0.006	0.007	0.063	0.018
F375S	0.002	0.077	1.785	1.608	1.025	0.015	0.017	0.151	0.266
W436A					N/A				
W436S									
H445A	0.008	0.027	0.134	0.773	0.265	0.009	0.021	0.049	0.02
S455A	0.004	0.062	0.101	2.99	7.856	0.008	0.009	0.045	0.013
S455F	0.005	0.012	0.12	0.621	0.102	0.009	0.016	0.047	0.026
K460D					N/A				

797 **Figure S6. IC50s (μg/ml) against KP.3 mutants.**

Figure S7. Variations on the epitopes of IGHV3-53/3-66 and IGHV3-74 mAbs. Frequencies of mutations on the epitopes of IGHV3-53/3-66 mAbs (A) and IGHV3-74 mAbs (B) among a total of 25332 spike sequences with KP.3 as reference. The sequences were published between June 1 to October 1 in 2024.