Human formin FHOD3-mediated actin elongation is required for sarcomere integrity in cardiomyocytes

Dylan A. Valencia[1], Angela N. Koeberlein[1], Haruko Nakano[2,3], Akos Rudas[4], Airi Harui[5], Cassandra Spencer[2], Atsushi Nakano[2,3,6*], and Margot E. Quinlan[1,6*]

[1]*Department of Chemistry and Biochemistry, [2]Department of Molecular, Cell, and Developmental Biology, [3]Eli & Edythe Broad Center of Regenerative Medicine & Stem Cell Research, [4]Department of Computational Medicine, [5]Divison of Pulmonary & Critical Care Medicine, Geffen School of Medicine, [6]Molecular Biology Institute, University of California Los Angeles, Los Angeles, California, 90095*

*To whom correspondence should be addressed: UCLA Department of Chemistry and Biochemistry, 607 Charles E. Young Dr. E., Los Angeles, California 90095. E-mail: anakano@ucla.edu, margot@chem.ucla.edu

Abstract

Contractility and cell motility depend on accurately controlled assembly of the actin cytoskeleton. Formins are a large group of actin assembly proteins that nucleate new actin filaments and act as elongation factors. Some formins may cap filaments, instead of elongating them, and others are known to sever or bundle filaments. The Formin HOmology Domain-containing protein (FHOD)-family of formins is critical to the formation of the fundamental contractile unit in muscle, the sarcomere. Specifically, mammalian FHOD3L plays an essential role in cardiomyocytes. Despite our knowledge of FHOD3L's importance in cardiomyocytes, its biochemical and cellular activities remain poorly understood. It has been proposed that FHOD-family formins act by capping and bundling, as opposed to assembling new filaments. Here, we demonstrate that FHOD3L nucleates actin and rapidly but briefly elongates filaments after temporarily pausing elongation, *in vitro*. We designed function-separating mutants that enabled us to distinguish which biochemical roles are req`uired in the cell. We found that human FHOD3L's elongation activity, but not its nucleation, capping, or bundling activity, is necessary for proper sarcomere formation and contractile function in neonatal rat ventricular myocytes. The results of this work provide new insight into the mechanisms by which formins build specific structures and will contribute to knowledge regarding how cardiomyopathies arise from defects in sarcomere formation and maintenance.

Keywords: Actin; Formin; Formin Homology Domain-Containing Protein 3 (FHOD3); Sarcomere; Cardiomyocyte; Hypertrophic cardiomyopathy (HCM)

Introduction

Higher order actin-based structures, such as the sarcomere, are the foundation for specific cellular functions that demand precise spatial and temporal coordination. Other examples of actin-based structures, including filopodia, stress fibers and the cytokinetic furrow, are also required for a wide range of processes, such as cell migration, adhesion, and division (Blanchoin et al., 2014; Svitkina, 2018). Hundreds of actin-binding proteins, and specific combinations thereof, coordinate to assemble these complex structures by accelerating actin assembly, crosslinking and bundling filaments, and stabilizing and/or disassembling actin filaments (Dominguez and Holmes, 2011; Lappalainen et al., 2022; Pollard, 2016). There are three known classes of actin nucleators that stimulate *de novo* filament assembly. Each class functions by a distinct mechanism, resulting in a unique starting point for the structure it initiates.

One such class of actin nucleators, known as formins, mediates both nucleation and continued
elongation of actin filaments. Formins are defined by their Formin Homology (FH) domains 1
and 2. The FH2 domain is highly conserved and forms a donut-shaped homodimer, which is
sufficient to nucleate new filaments. After nucleation, the FH2 domain remains processively
associated with the faster-growing barbed-end of actin filaments, modifying the rate of
elongation. The FH1 domain is less well conserved but contains proline-rich tracts that bind
profilin-actin to facilitate elongation of actin filaments (Courtemanche, 2018; Paul and Pollard,
2008, 2009). The regulatory domains for most formins are on either side of these FH domains
and an intramolecular interaction leads to autoinhibition (Goode and Eck, 2007; Li and Higgs,
2005; Schonichen and Geyer, 2010). The timing and strength of formin activity differ between
formins, allowing for distinct roles within a given cell (Homa et al., 2021).
62
Multiple classes of formins have been linked to sarcomere structure, including
Dishevelled Associated Activator Of Morphogenesis (Daam1, 2), Diaphanous (Diaph1-3),
Formin HOmology Domain–containing protein (FHOD1, 3), and others (Deng et al., 2021;
Iskratsch et al., 2010; Li et al., 2011; Mi-Mi et al., 2012; Molnár et al., 2014; Rosado et al., 2014;
Sanger et al., 2017; Shwartz et al., 2016; Sundaramurthy et al., 2020; Taniguchi et al., 2009).
There are two mammalian isoforms in the FHOD family of formins, FHOD1 and FHOD3
(Bechtold et al., 2014; Kanaya et al., 2005; Schonichen and Geyer, 2010). FHOD1 is widely
expressed and assembles stress fibers that contribute to adhesion and motility of various cell
types (Gasteier et al., 2003; Iskratsch et al., 2013; Koka et al., 2003; Schulze et al., 2014).
Despite its localization to intercalated discs, structures important for rapid signal transmission
and synchronized contractions, FHOD1 seems to be dispensable in cardiomyocytes (Al Haj et
al., 2015; Dwyer et al., 2014; Sanematsu et al., 2019). In contrast, FHOD3 is implicated in
sarcomere assembly and maintenance (Fujimoto et al., 2016; Iskratsch et al., 2010; Kan-o et
al., 2012a; Taniguchi et al., 2009; Ushijima et al., 2018).
77
There are four isoforms of FHOD3 reported in Uniprot. The major differences are due to
alternative splicing in the N-terminal half. An eight residue insert in the C-terminal half (an acidic
sequence $T(D/E)_5XE$) is present only in Uniprot isoform 4, which is commonly referred to as
FHOD3L (Iskratsch et al., 2010). In this paper, we adopt the commonly accepted nomenclature
in which FHOD3S (short) indicates Uniprot isoform 1 and FHOD3L (long) indicates Uniprot
isoform 4 (Figure 1). FHOD3L is predominantly expressed in striated muscle, whereas FHOD3S
is more widely expressed (Taniguchi et al., 2009). In cardiomyocytes, FHOD3L localizes to
sarcomeres in a striated pattern (Iskratsch et al., 2010; Kan-o et al., 2012b; Taniguchi et al.,
2009). Interestingly, FHOD3L does not localize at the barbed end of actin filaments within
sarcomeres. Instead, a direct interaction between FHOD3L and cardiac myosin-binding protein
C (cMyBP-C) drives localization of FHOD3L to the so-called C zone of sarcomeres (Matsuyama
et al., 2018a). In the absence of cMyBP-C, FHOD3L is diffuse and cardiac function is
compromised, suggesting that a balance between these two proteins is critical for proper heart
function in mice (Matsuyama et al., 2018a).
92
Mutations in the *FHOD3* gene have been deemed causative in at least 1-2% of patients
with hypertrophic cardiomyopathy (HCM), in addition to cases of left ventricular noncompaction
(LVNC), dilated cardiomyopathy (DCM), and progressive high-frequency hearing loss (Arimura
et al., 2013; Boussaty et al., 2023; Myasnikov et al., 2022; Ochoa et al., 2018). Moreover, recent
clinical studies have shown that HCM-causative FHOD3 mutations increase the risk of
cardiovascular death and all-cause death, with the onset of the disease occurring as early as
age 4 and as late as age 63 (Vodnjov et al., 2023; Wu et al., 2021).
100

Despite its known physiological significance, we lack a mechanistic understanding of
FHOD3L's role in cardiac development and function. Early biochemical analysis suggested that
FHOD family formins were an exceptional class of formins. Both purified mammalian isoforms
(FHOD1 and FHOD3) were found to decelerate, rather than hasten, actin assembly *in vitro*
(Schonichen et al., 2013; Taniguchi et al., 2009). In addition, actin bundling was observed for
FHOD1 (Schonichen et al., 2013). Consistently, FHOD1 mediates nuclear movement by
bundling and anchoring actin filaments (Antoku et al., 2023). However, it fails to complete this
function when a highly conserved isoleucine in the FH2 domain is mutated (hereafter, referred
to as the IA mutation) (Kutscheidt et al., 2014). The IA mutation is known to disrupt both
nucleation and elongation in most, if not all, formins, indicating that FHOD1 may enhance actin
assembly in addition to bundling *in vivo* (Patel et al., 2018; Xu et al., 2004). In addition, FHOD
proteins are required to form new sarcomeres in rat cardiomyocytes, human induced
cardiomyocytes, worms and mice (Fenix et al., 2018; Iskratsch et al., 2010; Kan-O et al., 2012;
Sundaramurthy et al., 2020; Taniguchi et al., 2009). FHODs with the IA mutation do not form
new sarcomeres, suggesting that actin assembly is required for sarcomere assembly, in conflict
with the biochemical data (Kan-O et al., 2012; Shwartz et al., 2016; Taniguchi et al., 2009; Xu et
al., 2004). Most recently, in the worm, FHOD protein was shown to cooperate with profilin to
build muscle, providing strong evidence that it functions by elongating actin filaments (Kimmich
et al., 2024). Previously, we found that *Drosophila* Fhod is a potent actin nucleator that can
accelerate actin elongation under certain conditions (Patel et al., 2018). We also showed that
human FHOD1 can nucleate actin but its activity is sensitive to the actin isoform (Patel et al.,
2018). We confirmed that both of these Fhod-family formins lose activity when the IA mutation is
introduced (Patel et al., 2018). Based on these biochemical results and the data regarding
Fhod-family formins in muscle of multiple species, we asked whether human FHOD3L can also
nucleate.
126
Here, we show that purified FHOD3L can accelerate actin assembly by both nucleation
and elongation. We also confirm that FHOD3L potently caps filaments, in the absence of
profilin, and bundles. Using function-separating mutations, we correlate FHOD3L's biochemistry
to the integrity of the sarcomere within cardiomyocytes. We found that reduced nucleation,
capping, and bundling by FHOD3L are well tolerated in neonatal rat ventricular myocytes
(NRVMs), whereas elongation activity is necessary for proper sarcomere formation and
function.
134
135 **Results**
136
137 *Biochemical characterization of human FHOD3*
138
We purified the C-terminal half of human FHOD3L (FHOD3L-CT), encompassing the
FH1 domain, FH2 domain, and C-terminal tail, which is typically sufficient for actin assembly *in*
vitro and *in vivo* (Figure 1, Figure 1—figure supplement 1) (Courtemanche, 2018; Patel et al.,
2018). In contrast to an earlier report, we found that FHOD3L-CT enhances rabbit skeletal
muscle actin (RSA) assembly in bulk pyrene assays (Figure 2A,B, Table 1) (Taniguchi et al.,
2009). (Table 1 summarizes all biochemical measurements.) Consistent with nucleation activity,
when we visualized the products of similar reactions, we observed many more filaments in the
presence of FHOD3L-CT compared to actin alone (Figure 2—figure supplement 1A). We also
asked if the shorter, alternatively spliced FHOD3 isoform, FHOD3S-CT, nucleates actin.
FHOD3S-CT, which is 8 residues shorter than FHOD3L-CT, accelerates actin assembly ~30%
more potently than FHOD3L-CT (Figure 2B, Figure 2—figure supplement 2A). Thus, we
conclude that the C-terminal half of both FHOD3 isoforms (moving forward, we will use S/L

when indicating both the short and long isoforms of FHOD3) are able to nucleate actin
filaments.
153
To further compare FHOD3L-CT to previously characterized formins, we introduced
mutations at conserved residues of the FH2 domain, I1163A and K1309A (Xu et al., 2004). Like
other formins, the I1163A mutant lacked nucleation activity and the K1309A mutant reduced
nucleation by ~90% (Figure 2B). We observed a reduction in the plateau of pyrene traces with
wild-type FHOD3S/L-CT and the mutants. The decrease was more apparent at higher
concentrations of the nucleator and seemed to be dose-dependent (Figure 2A, Figure 2—figure
supplement 2A). Control experiments demonstrated that the fluorescence change does not
reflect quenching due to side binding or bundling (Figure 2—figure supplement 1B). We,
therefore, tested barbed-end binding. We asked whether FHOD3L-CT inhibits filament
elongation by performing bulk seeded-elongation assays. Indeed, we found that FHOD3L-CT
potently slows barbed-end elongation (K_{app} = 0.023 $\pm$ 0.005 nM; Figure 2C,D). Barbed-end
binding by FHOD3S-CT is over 30 times weaker (K_{app} = 0.750 $\pm$ 0.090 nM) (Figure 2D, Figure
2—figure supplement 2B). The IA mutation is generally thought to disrupt binding to both actin
monomers and filament barbed ends. However, FHOD3L-CT I1163A has an affinity of 4.9 $\pm$ 1.4
nM for barbed ends of actin filaments (Figure 2D). While the affinity is >200-fold weaker than
FHOD3L-CT, it still binds barbed ends tightly. Therefore, we interpret the plateau decrease
observed in bulk assembly assays as evidence of barbed-end capping.
171
To directly observe the impact of FHOD3L-CT on growing actin filaments, we used Total
Internal Reflection Fluorescence (TIRF) microscopy in the presence of profilin. Most of the
filaments elongated at rates indistinguishable from actin alone. Initially, we could not confirm
that FHOD3L-CT was bound to these filaments. We, therefore, repeated bulk seeded-elongation
assays in the presence of profilin. Under these conditions, we observed increased rates of actin
assembly at low doses of FHOD3L-CT, confirming that the formin accelerates elongation
(Figure 2E). In contrast, at higher formin concentrations, we found that elongation was actually
slower than actin alone. We attribute the complicated shapes and dose response of these data
to the contributions of both elongation and capping activities. We also note that the
concentrations required for FHOD3L-CT to elicit measurable changes in actin assembly indicate
a marked apparent decrease (500-1000 fold) in affinity for the barbed-end when profilin is
present (Figure 2E).
184
Upon closer observation of individual filaments in the TIRF assay, we detected brief
pauses followed by short "bursts" of fast elongation (Figure 2F,G, Figure 2—figure supplement
1C). A pause lasted ~12 s on average (Figure 2—figure supplement 1D). During the following
bursts, the filaments were dimmer and elongated ~3-fold faster (39 $\pm$ 13 subunits/s), consistent
with formin-mediated elongation (Figure 2F,G, Video 1). The average length of filament
produced during these bursts was 1.10 $\pm$ 0.49 μm (Figure 2—figure supplement 1E), which is
similar to the characteristic run length of *Drosophila* Fhod-A (~2 μm), but very different from
other formins, which often range from 20 – 200 μm (Bremer et al., 2024; Cao et al., 2018;
Moseley and Goode, 2005; Patel et al., 2018; Vizcarra et al., 2014). FHOD3S-CT mediated
similar behavior with indistinguishable pauses, though with slightly, albeit statistically
significantly, shorter bursts (characteristic run length of 0.90 $\pm$ 0.33 μm) (Figure 2—figure
supplement 2C-E). Thus, for brief intervals, FHOD3S/L-CT caps barbed-ends followed by
accelerated elongation, when profilin is present.
198
Next, we investigated bundling by FHOD3L-CT. To do so, we visually examined the
organization of phalloidin-stabilized actin filaments after incubating with FHOD3L-CT. Bundles
were readily apparent (Figure 2—figure supplement 1F). To better characterize bundling

activity, we performed low-speed bundling assays with a fixed concentration of actin filaments
and several concentrations of FHOD3L-CT (Figure 2H,I). We also compared this activity to
bundling by FHOD3S-CT. We found that FHOD3L-CT bundles somewhat more potently than
FHOD3S-CT, which is most apparent at concentrations above 15 nM (Figure 2—figure
supplement 2F,G).
207
Overall, FHOD3S/L-CT nucleates and elongates actin. FHOD3S-CT is a more potent
nucleator, a weaker barbed-end capper, and a slightly weaker bundler compared to FHOD3L-
CT (Table 1). These data suggest that the acidic $T(D/E)_5XE$ insertion of FHOD3L interacts with
different surfaces of actin depending on the biochemical activity. Perhaps the marked increase
in capping by FHOD3L-CT is mediated by binding to an exposed basic region on the filament.
Decreased activity levels of FHOD3L-CT, such as nucleation and bundling, may be due to the
fact that most of the actin monomer surface is acidic.
215
216 **Video 1. TIRF elongation assay**. This video shows elongation assays of actin alone and in the presence
217 of 100 pM pHOD3L-CT. Events – pauses before bursts of acceleration – are indicated with arrows in the
218 last movie. (Conditions as in Fig. 2F.)
219
220 *Biochemical validation of function-separating mutants*
221
We next designed FHOD3L-CT mutants to separate nucleation and elongation. We also
assessed the impact of these mutations on capping and bundling. Previously, Baker et al.
identified mutations in the FH2 domain of Bni1 that diminish nucleation while maintaining
elongation activity (Baker et al., 2015). Based on sequence and structural alignments, FHOD3L
K1193 approximates one of these, Bni1 K1467. FHOD3L-CT K1193L nucleated with less than
30% of the strength seen for wild-type (Figure 3A). The barbed-end affinity of FHOD3L-CT
K1193L was 0.470 ± 0.090 nM, ~12-fold weaker than wild-type (Figure 3B). Importantly,
K1193L-mediated elongation of actin filaments was indistinguishable from wild-type (no
significant difference in elongation rate, run length, or capping duration) (Figure 3C, Figure 3—
figure supplement 1A,B). Low-speed bundling assays showed that K1193L bundling is >2-fold
weaker than wild-type (Figure 3D). In summary, elongation activity is maintained while
nucleation and bundling are strongly reduced by the K1193L mutation.
234
To remove FH1-mediated acceleration of elongation, we substituted glycine-serine
linkers for the polyproline tracts in FHOD3L-CT (Zweifel and Courtemanche, 2020). (The FH2
domain alone was unstable *in vitro*.) Pyrene-based actin assembly assays were confounded by
bundling/aggregation induced by this construct. Although low-speed cosedimentation assays
indicated that the bundling strength of FHOD3L-CT GS-FH1 is slightly weaker than that of wild-
type, images show irregular shapes that may scatter light more, thereby disrupting the pyrene
signal (Figure 3D, Figure 3—figure supplement 1C). Therefore, to compare the nucleation
strength of FHOD3L-CT GS-FH1 with wildtype, we counted filaments using TIRF microscopy.
We observed similar numbers of filaments generated in the presence of low concentrations
(used to minimize bundling) of FHOD3L-CT wild-type or GS-FH1 (Figure 3E, Figure 3—figure
supplement 1D). Barbed-end elongation assays in the absence of profilin demonstrated a
weakened affinity of FHOD3L-CT GS-FH1 ($K_{app} = 0.218 \pm 0.016$ nM) (Figure 3B). This finding
suggests that increasing the stiffness of the FH1 domain impacts barbed end binding but we
cannot say whether the effect is direct or indirect. Interestingly, profilin did not decrease barbed-
end binding of FHOD3L-CT GS-FH1 ($K_{app} = 0.21 \pm 0.13$ nM; Figure 3F). Finally, and most
importantly deceleration was observed at all concentrations tested in assays with profilin,
confirming that elongation is not enhanced by FHOD3L-CT GS-FH1 (Figure 3F). Thus,
nucleation is maintained while acceleration of elongation is no longer detected.

Together, these mutations resulted in a range of nucleation, elongation, capping, and
bundling activities (Table 1). We, therefore, used them to assess the correlation of biochemical
activities with sarcomere formation and function in cardiomyocytes.
257

FHOD3L rescues sarcomere organization and contractility in neonatal rat ventricular myocytes
259

In order to perform structure-function analysis in a cellular context, we established
methods to knock down and rescue FHOD3L in neonatal rat ventricular myocytes (NRVMs)
(similar to (Taniguchi et al., 2009). Briefly, freshly isolated NRVMs were treated with small
interfering RNA (siRNA) targeting FHOD3 using reverse transfection (Figure 4A; see Methods
for details). Adenoviral infection, driving expression of rescue constructs, was initiated two days
after siRNA treatment and cells were examined (fixed or live) two days after infection (Figure
4A). The rescue constructs contained human FHOD3L, which is not targeted by the siRNA used
to remove endogenous rat FHOD3. In all cases, cells were treated with siRNA and adenovirus.
For negative controls, AllStars Negative Control siRNA and/or empty virus were substituted for
the FHOD3-specific reagents. Thus, we refer to the negative control experiment as a mock
knockdown and the knockdown alone as a mock rescue. During assay development, we found
that knockdown reduced FHOD3 mRNA to ~35% of original levels after four days, with either of
two commercially available oligos (Figure 4—figure supplement 1A). At this timepoint, we
detected an ~80% reduction in endogenous FHOD3 protein via Western blot (Figure 4B,C).
274

We analyzed sarcomere structure in fixed samples. To detect sarcomeres, we stained
cells with anti-α-actinin antibodies (Figure 4D). To analyze on a per cell basis, we stained
plasma membranes with WGA (wheat germ agglutinin) and segmented the NRVMs with
CellPose (Figure 4—figure supplement 1B) (Stringer et al., 2021). The DsRed reporter was not
an accurate readout of exogenous 3xHA-FHOD3L (hereafter, FHOD3L) expression levels
(Figure 4—figure supplement 1C). Therefore, we used anti-HA antibodies to visualize and
quantify the expression of exogenous FHOD3 on a per cell basis (Figure 4D, Figure 4—figure
supplement 4G). We found that exogenous FHOD3L expression was ~90% above endogenous
levels at the end of the rescue timeline (Figure 4B,C). We, therefore, examined the impact of
FHOD3L expression level on sarcomere structure. We detected essentially no correlation (R-
squared ranges from 0.01 to 0.16) of sarcomere number, length, or width over a three-fold
change in FHOD3L expression levels (Figure 4—figure supplement 1D-F). The average HA
intensity was within this range for all experiments. Despite the evidence that sarcomere
structure was not a function of FHOD3 expression levels, we decided to exclude NRVMs that
express very highly above endogenous FHOD3 levels. We set an upper cutoff for the
normalized exogenous FHOD3L expression per cell area of 5% and applied that same intensity
level as a cutoff for all other experiments (Figure 4—figure supplement 1G). We also set a lower
cutoff slightly above background HA levels to exclude NRVMs not expressing exogenous
FHOD3L.
294

We confirmed that sarcomeres remained organized upon mock knockdown, counting 12
$\pm$ 13 sarcomeres per cell (Figure 4D,E, Table 2). (Measurements from NRVMs are summarized
in Table 2.) Sarcomeres were largely absent in the mock rescue (3 $\pm$ 7) with α-actinin puncta
and aggregates visible by immunofluorescence (IF) (Figure 4D,E). Expression of wild-type
FHOD3L was sufficient to rescue the loss of sarcomeres and anti-HA staining demonstrated
that exogenous FHOD3L localized to sarcomeres, as expected (Figure 4D). In fact, FHOD3L-
rescued NRVMs formed significantly more sarcomeres than the mock knockdown NRVMs (19 $\pm$
14 vs. 12 $\pm$ 13; Figure 4E). We note that the sarcomere number measurement has a high
standard deviation because even within the expression level cutoffs, some cells lacked

sarcomeres. Such right-tailed distributions are consistent with other reports of sarcomere numbers per cardiomyocyte (Neininger-Castro et al., 2023).

To analyze sarcomere integrity more closely, we measured sarcomere lengths (Z-line to Z-line) and widths (Z-line lengths). In the mock rescue cells, the sarcomeres that remained were both shorter and narrower than those in the mock knockdown cells (Figure 4F, G). When rescued with FHOD3L, sarcomere lengths (1.72 ± 0.18 µm) and sarcomere widths (1.70 ± 0.46 µm) recovered to lengths comparable to the mock knockdown control (Figure 4F, G). To measure thin filament length, we stained NRVMs with phalloidin and anti-HA (Figure 4 H). We observed shorter thin filament lengths in FHOD3L-rescued NRVMs (739 ± 81 nm) compared to mock knockdown NRVMs (925 ± 94 nm) (Figure 4I). The difference in thin filament length could reflect the increased number of sarcomeres in the rescued cells and/or indicate that the sarcomeres have not reached their final steady state after two days of FHOD3L expression. Therefore, FHOD3L-rescued NRVMs form sarcomeres *de novo* that well approximate the sarcomeres in the negative control. We compared further experimental results to FHOD3L rescue cells, due to the slight differences.

We next measured contractile function in FHOD3L-rescued NRVMs. To do so, we used the motionGUI MatLab program that makes use of Digital Image Correlation (DIC) to measure contractility (Huebsch et al., 2015; Nakano et al., 2017). Contraction and relaxation velocities inform us about systolic and diastolic function of the cardiomyocytes, respectively, and thus their ability to promote proper blood flow in organisms (Ferreira-Martins and Leite-Moreira, 2010). As expected, maximal contraction and relaxation velocities were both significantly decreased upon FHOD3 knockdown (Figure 5A,B,D, Figure 5—figure supplement 1A). Both velocities recovered to mock knockdown levels by subsequent expression of FHOD3L (Figure 5A,C,D, Figure 5—figure supplement 1A, Video 2). We also examined expression of FHOD3L in an otherwise untreated background (overexpression). Intensity of exogenous FHOD3L per NRVM was notably lower compared to the wild-type rescue despite infecting with the same titer, suggesting that excel FHOD3L is deleterious and NRVMs attempt to maintain FHOD3L levels below some maximum (Figure 5—figure supplement 2A). Consistent with this idea, contraction and relaxation velocities were reduced upon overexpression of FHOD3L (Figure 5—figure supplement 2B,C). These data further confirm that the expression levels in our rescue experiments, while somewhat higher than endogenous levels, are within an acceptable range.

To assess the impact of FHOD3L depletion on cardiac rhythm, we measured the percentage of beating area in each video that exhibited consistent, rhythmic contractions (defined as no more than 1 beat out of sync in a 10 second period). We observed rhythmic contractions for both the mock knockdown NRVMs and FHOD3L-rescued NRVMs for ~75% of the analyzed videos (Figure 5A,C,E, Video 2). In contrast, primarily arrhythmic contractions were detected in FHOD3-depleted NRVMs, with only ~15% of the videos showing rhythmic beating (Figure 5B,E, Video 2). To assess whether some NRVMs in these conditions were not contracting at all, we estimated the proportion of NRVMs that were contracting per video and found contractions throughout ~95% of the field of view on average for both mock knockdown NRVMs and wild-type FHOD3L-rescued NRVMs (Figure 5—figure supplement 1B). For FHOD3-depleted NRVMs, we observed a reduction in contractile area to ~80% (Figure 5—figure supplement 1B). This value was higher than expected based on other metrics (Figure 5D,E). We attribute much of the movement to neighboring cells pulling each other. We also quantified contractility when we overexpressed FHOD3L. FHOD3L overexpression did not clearly impact the average amount of rhythmic contractions or the area of contraction though variance was increased (Figure 5—figure supplement 2D,E), consistent with the idea that, at high enough levels, FHOD3L activity can be detrimental in NRVMs.

Video 2. Exogenous FHOD3L expression rescues rhythmic contractions in NRVMs. This video shows live NRVMs from control and rescue conditions as in Fig. 5. DIC analysis is indicated by overlaid red and green vectors.

Loss of nucleation but not elongation is tolerated for sarcomere formation and cardiac function

Once we established baseline levels of sarcomere structure and contractility in NRVMs with the wild type FHOD3L rescue, we asked which of FHOD3L's actin organizing activities are important for its cellular function. To this end, we performed rescue experiments with the nucleation-hindering mutant (K1193L) and the elongation-hindering mutant (GS-FH1). Expression of the K1193L mutant resulted in expression levels similar to wild type (Figure 6—figure supplement 1A). Overall, cells expressing FHOD3L K1193L were almost indistinguishable from those expressing FHOD3L. FHOD3L K1193L localization was striated (Figure 6A) and the number of sarcomeres per NRVM was rescued to wild-type levels (15 ± 16) (Figure 6B). Sarcomere lengths and widths were indistinguishable from wild-type (Figure 6C,D). The only difference was thin filament lengths, which were significantly longer in the K1193L-rescued NRVMs (792 ± 79 nm) compared to those in FHOD3L-rescued NRVMs (739 ± 81) (Figure 6E, Figure 6—figure supplement 1B). In agreement with the well-organized, near wild-type appearance of sarcomeres, contraction and relaxation velocities, as well as the proportion of rhythmically contracting NRVMs, were indistinguishable from the FHOD3L rescue (Figure 6F,G, Figure 6—figure supplement 1C,D, Video 3). Thus, ~70% lower nucleation activity, ~20-fold weaker barbed-end capping, and loss of over 50% of bundling activity had no deleterious impact on FHOD3L's ability to function in NRVMs.

In contrast, the elongation deficient mutant (GS-FH1) did not rescue FHOD3L loss in NRVMs. Based on total fluorescence per cell, we determined that GS-FH1 levels were lower than observed for wild-type or K1193L, despite infecting a similar proportion of NRVMs (Figure 6—figure supplement 1A). In a few cells that possibly escaped from RNAi knockdown, we observed a striated pattern of GS-FH1, demonstrating that the protein can fold and localize correctly if sarcomeres are intact (Figure 6—figure supplement 1E). The turnover rate of GS-FH1 may be higher than wild-type intrinsically or because there are few binding sites available. The sarcomere number per cell was only 2 ± 4, similar to the mock rescue (Figure 6A and B). Within the sarcomeres detected, sarcomere lengths and widths were both reduced (Figure 6C and D). We were not able to measure thin filament lengths with confidence, due to the lack of sarcomere organization. Not surprisingly, contractility was nearly abolished after rescuing with GS-FH1 (Figure 6F,G,Figure 6—figure supplement 1C,D, Video 3). Contraction and relaxation velocities were decreased, and contractions were largely arrhythmic (Figure 6F,G, Figure 6—figure supplement 1C,D). We note that barbed-end capping and bundling activities were weaker than wild-type in the GS-FH1 mutant but they were equivalent or greater than the activities found for the K1193L mutant which rescued well (Figure 3B,D). Therefore, we conclude that elongation activity of FHOD3L is the primary activity required for sarcomere formation, whereas reduced nucleation, barbed-end binding, and bundling activities are well tolerated.

Video 3. Expression of FHOD3L function separating mutants in NRVMs. This video shows live NRVMs from rescues with wild type and function separating mutants, corresponding to Fig 6. Two movies for GS-FH1 reflect the range of contractions we observed with this rescue. DIC analysis is indicated by overlaid red and green vectors.

Discussion

406   *FHOD3L elongation is necessary for sarcomeres in NRVMs*
407
Published rates of nucleation and elongation by formins both range over at least an
order of magnitude. It is generally thought that the specific actin assembly properties of a given
formin are set for its function, i.e., the structure this formin will build. To directly test this idea,
Homa et al. replaced Cdc12p, the formin critical to *S. pombe* cytokinesis, with several chimeras
of differing nucleation strength (Homa et al., 2021). Indeed, they found a strong positive
correlation between formins with nucleation strength similar to Cdc12p and their ability to drive
cytokinesis. In a computational model that recapitulates yeast cable structure, it was found that
nucleation and/or elongation could be tuned to build the cables (McInally et al., 2021). Thus,
how formins are used is likely to differ from case to case.
Here we determined that FHOD3L can nucleate *in vitro*, albeit weakly compared to many
formins. Unlike in cytokinesis, we found that this activity is not necessary for its function in
NRVMs. The necessity of actin assembly, nucleation and/or elongation, by Fhod-family formins
has been assumed based on failure of the IA mutation to rescue in multiple species (Kan-O et
al., 2012; Kimmich et al., 2024; Kutscheidt et al., 2014; Shwartz et al., 2016; Taniguchi et al.,
2009). To assess nucleation more specifically, we tested a mutant variant that substantially
reduces the nucleation activity without altering elongation (FHOD3L K1193L). This mutant had
almost no impact on sarcomere formation or function in NRVMs. The only statistically significant
difference we detected was in thin filament length (Figure 6E). A host of proteins are critical to
establishing and maintaining thin filaments at a precise length (Szikora et al., 2022). No other
studies have implicated FHOD3L in this process to date. Thus, we believe that the length
difference is more likely to reflect the state of maturity of the cells (time post infection) and/or the
number of sarcomeres per cell. While we only have three data points at this time, sarcomere
number and thin filament length were highly correlated (inversely) in our experiments, with an R-
squared of 0.90.
432
By rescuing NRVMs with FHOD3L K1193L, we also tested a formin that caps barbed
ends with a K_d that is ~15 times higher than wild type (i.e., ~15x weaker) and bundles filaments
~50% less potently than wild type. The fact that the cardiomyocytes were indistinguishable from
wild-type when expressing this mutant suggests that neither of these activities are required and,
certainly, they are not needed at wild-type levels. Assuming that FHOD3L activities are tuned to
meet its *in vivo* functions, these magnitudes of change in activity would be expected to result in
measurable phenotypes in function and/or morphology of the cell.
440
The fact that K1193L had no impact on the elongation properties of FHOD3L *in vitro* and
still rescues sarcomeres leads to an indirect conclusion that elongation is the actin assembly
activity that is important in NRVMs. Of course, the failure to rescue by FHOD3L GS-FH1, an
elongation incompetent mutant, strongly supports this conclusion (Figure 6). The elongation
properties of FHOD3L are unusual. Its elongation rate is among the highest known, including
mDia1 and Cappuccino (Bor et al., 2012; Kovar et al., 2006). However, processivity is very brief,
resulting in filaments of only ~1 μm. We previously found that the *Drosophila* splice variant
FhodA has a similarly short characteristic run length (Patel et al., 2018). In principle, this
characteristic run length is sufficient to build sarcomeric thin filaments, which are about 1 μm
long in NRVMs. Notably, a splice variant of *Drosophila* Fhod that only differs by having a shorter
tail, FhodB, typically elongates for ~20 μm and the tails of mammalian and *Drosophila* formins
are highly conserved (Bremer et al., 2024). These observations lead us to question whether
processivity of Fhod-family formins is somehow regulated *in vivo* such that it can build short
filaments when needed and longer filaments in other contexts. Interestingly, a computational
model describing Bni1-built actin cables demonstrates that tuning the length of individual

filaments to cell size is sufficient to account for cable differences in cells of different sizes
(McInally et al., 2024).
458
The FH1 domain strongly influences elongation rate *in vitro*. In NRVMs, we found that
replacement of the FH1 domain with a flexible linker (FHOD3L GS-FH1) resulted in a severe
phenotype. These cells had very few sarcomeres. FHOD3L GS-FH1 was indistinguishable from
mock rescue in sarcomere number, structure (shorter and narrower), and function (contraction
and relaxation velocities did not differ). The result could be questioned because the expression
levels were lower than those for the other constructs. We argue against this based on two
observations: First, despite similar metrics in sarcomere structure and function, the rhythmicity
of contraction was markedly diminished, even compared to the mock rescue (Figure 6G).
Second, we do not believe that the low protein levels reflect protein instability. We detect
FHOD3L GS-FH1 bound to residual sarcomeres, suggesting that it folds and is stable when
binding sites within sarcomeres are available (Figure 6—figure supplement 1E). Therefore, we
suspect that low FHOD3L GS-FH1 levels reflect the absence of sarcomeres.
471
Recent work in C. elegans is consistent with our finding. Kimmich et al. removed the
endogenous coding region of the FH1 domain in worm Fhod-1 (the only worm Fhod gene)
(Kimmich et al., 2024). Body wall muscle is severely impacted by this deletion. In addition, they
found that profilin is required for Fhod-1 function. Together these data strongly argue that
elongation is important. Interestingly, the muscle phenotype is more severe in a strain that lacks
the FH2 domain and all of the upstream sequence but not the FH1 domain. Thus, nucleation
activity seems to be necessary in this animal. Ultimately, complete removal of a domain, such
as the GS-FH1 mutant is an important starting point but a blunt tool. Future experiments with
modified elongation (e.g. slower, more processive) will provide further insight into the
biochemical role of FHOD3.
482
483 *Why does FHOD3L elongate in NRVMs?*
484
Given these new data, what could FHOD3L be doing in the cell? In multiple species,
premyofibrils are built but they fail to mature into wild-type myofibrils in the absence of FHOD
(Fenix et al., 2018; Kan-O et al., 2012; Kimmich et al., 2024; Shwartz et al., 2016). It follows that
FHOD3 is not required to build the earliest thin filaments in the sarcomere. Instead, FHOD3
could contribute new thin filaments to expanding, maturing sarcomeres. If this were the case,
we might expect nucleation to be more important than our data indicate. Instead, FHOD3L could
be responsible for elongating filaments that are nucleated by a different protein. For example,
FHOD3L and DAAM could collaborate, or they could contribute to the apparent redundancy and
robustness built in to sarcomerogenesis (Szikora et al., 2022). DAAM knockout mice have
overlapping cardiac phenotypes with those of the FHOD3 knockout (Kan-O et al., 2012; Li et al.,
2011). Furthermore, a lack of sarcomere thickening and loss of sarcomere organization are
common phenotypes in *Drosophila* knockdowns of Daam1 and Fhod (Molnár et al., 2014;
Shwartz et al., 2016). While FHOD3L and DAAM cannot compensate for the loss of one or the
other, they might work together and be able to help each other when one is compromised.
499
Alternatively, FHOD3L could be reinforcing the Z-line structure to facilitate sarcomere
thickening. This idea is consistent with the narrow myofibrils and Z-body disorder observed in
worms lacking functional Fhod-1 (Kimmich et al., 2024; Mi-Mi et al., 2012). However, if capping
and bundling were important to the Z-line integrity, one would expect to find FHOD3L localized
in the Z-line. In mouse heart tissue, FHOD3L appears to bind directly to MyBP-C which localizes
between the Z- and M-lines (Matsuyama et al., 2018b), suggesting that it is not actively
stabilizing the Z-line. FHOD localization varies between species and with development,

suggesting that this role is not conserved, if it exists. If, instead, transient actin assembly were
sufficient for Z-line stabilization, this role would be consistent with our findings that capping and
bundling are not essential in NRVMs.
510
511 **Materials and Methods**
512
Protein expression, purification, and labeling
FHOD3L-CT (residues 963-1622) was cloned into pGEX-6P-2 with an N-terminal
glutathione-S-transferase (GST) tag. The original template, EGFP-Fhod3L, was generously
provided by T. Iskratsch (Queen Mary University of London) (Iskratsch et al., 2010). Point
mutations were generated by site-directed mutagenesis. Truncations were constructed using
FastCloning (Liu and Naismith, 2008). pGEX-FHOD3L-CT GS-FH1, in which the polyproline
tracts were replaced with GS linkers was cloned via Gibson Assembly introducing a gBlock into
pGEX-FHOD3L-CT (not including the FH1 region).
521
The FHOD3L-CT wild-type and mutant constructs were transformed in Rosetta 2 (*E. coli*
DE3) cells (Novagen), which were grown in 1 liter of Terrific Broth supplemented with 100
mg/liter ampicillin and 32 mg/liter chloramphenicol. Expression was induced at an OD of 0.6-0.8
by adding 0.5 mM isopropyl β-D-1-thiogalacto-pyranoside (IPTG) and shaking overnight at
18°C, 210 rpm. The cells were harvested by centrifugation, washed in PBS, and flash frozen in
liquid nitrogen.
528
Cell pellets expressing GST-FHOD3L-CT wild-type and mutants were resuspended in 20
mM HEPES pH 7.5, 150 mM NaCl, 1 mM PMSF, 1 mM DTT, 2 µg/mL DNaseI. All subsequent
steps were performed on ice or at 4°C. The cells were lysed with a microfluidizer, cleared by
centrifugation at 20,000 x g for 20 minutes, and then purified using a HitrapSP-FF cation
exchange column (GE Life Sciences) with a gradient of 0.2 –0.6 M NaCl over 1 column volume
after a 1 column volume wash with 0.2 M NaCl. Peak fractions were dialyzed overnight into 20
mM HEPES pH 8, 200 mM NaCl, 1 mM DTT and Prescission Protease was added to cleave off
the GST-tag. This sample was centrifuged at 4°C for 48,000 rpm for 20 minutes, and further
purified on a MonoS cation exchange column (GE Life Sciences) with a gradient of 0.2–0.95 M
NaCl over 40 column volumes after a 1 column volume wash with 0.2 M NaCl. Peak fractions
were exchanged into storage buffer (10 mM Tris pH 8, 150 mM NaCl, 20% glycerol, 1 mM DTT),
then flash frozen in liquid nitrogen and stored at -80°C.
541
542   Cell pellets expressing GST-FHOD3S-CT were purified and stored similarly to GST-
543 FHOD3L-CT wild-type, except 150 mM NaCl was used throughout the purification with a
544 gradient of 0.15–0.95 M NaCl over 40 column volumes after a 1 column volume wash with 0.15
545 M NaCl on the MonoS cation exchange column (GE Life Sciences).
546
547   Concentrations of C-terminal FHOD3 constructs were determined by running a series of
548 serial dilutions on a Sypro Red-stained quantitative gel using densitometry (ImageJ) with rabbit
549 skeletal actin as the standard. All FHOD3L-CT concentrations are reported as dimer
550 concentrations.
551
552   Human Profilin-1 and *Schizosaccharomyces pombe* profilin were expressed and purified
553 as described for *Drosophila* profilin (Chic) (Bor et al., 2012). Profilin-1 concentration was
554 determined using the extinction coefficient 14,992 M^{-1} cm^{-1}. *S. pombe* profilin concentration was
555 determined using 1.63 OD/mg/ml.(Lu and Pollard, 2001)

556
We used rabbit skeletal muscle actin (RSA) throughout the paper, based on FHOD3L's
role in skeletal and cardiac muscle. Skeletal muscle actin was isolated from rabbit back muscle
acetone powder (Pel-Freez) according to the method described by Spudich and Watt followed
by gel purification (Spudich and Watt, 1971). Skeletal muscle actin was labeled with pyrene
iodoacetamide (Thermo Scientific) or Alexa Fluor 488 NHS-ester (Thermo Scientific) as
described (Sun et al., 2018).
563
564   *Pyrene Assays*
565
Pyrene assays were performed essentially as described (Bor et al., 2012) on an Infinite
200 Pro plate reader (Tecan). FHOD3L-CT was diluted in buffer Z (2 mM Tris pH 8.0, 0.2 mM
ATP, 0.1 mM CaCl$_2$, 0.5 mM TCEP, 0.04% sodium azide) before addition to polymerization
buffer (KMEH: 10mM HEPES, pH 7, 1mM EGTA, 50 mM KCl, 1 mM MgCl$_2$). This mix was
added to Mg^{2+}-actin at a final concentration of 4 µM with 5% pyrene-labeled actin. For bulk
assembly assays, nucleation strengths were calculated from the slope at t$_{1/8}$ through the origin.
The concentration of barbed ends was calculated from the slope (obtained by linear regression
over 90 s) as described in (Pollard, 1986 and Patel et al., 2018). For seeded elongation assays,
actin filaments were sheared by passing three times through a 24-gauge needle and then
aliquoted into each well of a microplate. Proteins were added to the seeds and incubated for 2–
4 min at room temperature. Seeds and additional proteins in KMEH were added to Mg^{2+}-actin,
at a final concentration of 0.5 µM actin with 10% pyrene-labeled actin, to initiate elongation.
Elongation rates were determined by linear regression over the first 90 s and normalized against
the rate of actin alone in each experiment. The affinity of FHOD3-CT for barbed ends was
determined by fitting the data to the quadratic binding equation:
581
582
583
584

$$r = a \,+\, \text{b} * ((\text{[barbed ends]} + \text{[FHOD3} - \text{CT]} + \, K_d) - \sqrt{(\text{[barbed ends]} + \text{[FHOD3} - \text{CT]} + \, K_d)^2 - 4 * \text{[barbed ends]} * \text{[FHOD3} - \text{CT]}}),$$

585
586   where *r* is the normalized elongation rate and *a* and *b* are offset and scaling constants.
587
588   *Low Speed Bundling Assays*
589
10 µM actin was polymerized for 1 hour at room temperature and diluted to 5 µM with
varying amounts of FHOD3-CT constructs in KMEH. Samples were incubated for 30 minutes at
room temperature and then spun for 20 min., 14,000 x g to separate the pellet and supernatant.
The supernatant samples were carefully transferred to new tubes and the pellet samples were
resuspended in an equal volume of 1X sample loading buffer for quantitative comparison.
Samples were boiled in 1X sample loading buffer for 10 minutes and run on 10% SDS-PAGE
gels. The percentage of actin pelleted was determined via densitometry with FIJI (Schindelin et
al., 2012).
598
599   *TIRF Microscopy*
600
601   TIRF microscopy was utilized to measure the elongation rates and run lengths of the
602   FHOD3-CT constructs. Coverslips were rinsed three times in MilliQ water, placed in 2%
603   Hellmanex (Hellma Analytics) at 60–65 °C for 2 h, and rinsed another five times in MilliQ water
604   and allowed to dry.
605

Parallel flow chambers of ~15 μl were assembled on the slide using strips of double-
sided tape. Flow chambers were prepared with the following steps: 1) 20 μl of high-salt buffer
(50 mM Tris pH 7.5, 600 mM NaCl) for 2 min; 2) 12.5 μl of 60 nM NEM-myosin (stock in 10 mM
HEPES pH7, 0.5 M KCl, 10 mM EDTA pH 7.0, 1 mM DTT, 50% glycerol, diluted in high-salt
buffer); 3) 20 μl of high-salt BSA-containing buffer (1% BSA, 50 mM Tris pH 7.5, 600 mM NaCl);
4) 20 μl of low-salt BSA-containing buffer (1% BSA, 50 mM Tris pH 7.5, 150 mM NaCl). 5) 50 μl
of magnesium-actin and additional proteins to be assayed for a final concentration of 1 μM actin
(10% Alexa Fluor 488-labeled) and 5 μM human Profilin-1 in 1X TIRF (KMEH, 0.2 mM ATP, 50
mM DTT, 20 mM glucose, 0.5% methylcellulose (4,000 cP)) buffer supplemented with 250 μg/ml
glucose oxidase and 50 μg/ml catalase.
616
The videos and images were acquired using a Zeiss Axio Observer 7 Basic Marianas
Microscope with Definite Focus 2 equipped with a 3i Vector TIRF System, an Alpha Plan-
Apochromat 63x (1.46 NA) Oil TIRF Objective, and an Andor iXon3 897 512x512 10 MHz
EMCCD Camera, using Slidebook 6 software. Experiments were performed at room
temperature. Images were captured at 2.5-s intervals for 10 min. Bright and dim filaments were
distinguished manually. Filament lengths were quantified with the JFilament plug-in in FIJI
(Schindelin et al., 2012; Smith et al., 2010).
624
625   *Adenoviral generation, purification, and infection*
626
Adenoviruses containing a full-length FHOD3L wild-type construct were generated by
transiently transfecting HEK293 cells using the CalPhos Mammalian Transfection Kit (Catalog
No. 631312, Takara Bio) with pAdenoX-CMV-3xHA-FHOD3L-CMV-DsRed after digesting with
PacI (NEB) in Cutsmart buffer (NEB) for 1 hr at 37 °C. HEK293 cells were cultured in 1X DMEM
(Thermo Fisher, Catalog No. 11965092), 10% FBS (Thermo Fisher, Catalog No. 10082147),
and 1% penicillin/streptomycin/amphotericin B (Thermo Fisher, Catalog No. 15240096). Cells
were lysed 7-10 days later when noticeable cytopathic effect (CPE) was present and centrifuged
at 1.5k x g for 5 min., RT.
635
Additional crude adenovirus encoding full-length FHOD3L was harvested from infected
HEK293. Dilutions and subsequent infections were performed in PBS with magnesium chloride
and calcium chloride (PBS+/+) (Thermo Fisher, Catalog No. 14040117) for 1 hr at room
temperature, rotating the plate every 10-15 minutes. Plaque formation assays were then
performed as in (Baer and Kehn-Hall, 2014). Clone variability was assessed by infecting HEK
293 for each adenoviral clone and observing subsequent DsRed expression over time. Optimal
adenoviral clones were selected for large-scale purification. HEK293 were infected, then
collected 2-3 days later for purification with the Adeno-X Maxi Purification kit (Takara Bio,
Catalog No. 631533).
645
Crude adenoviruses containing pAV-CMV-{3xHA-FHOD3L GS-FH1}:SV40 pA-CMV-
mCherry and pAV-CMV-{3xHA-FHOD3L K1193L}:SV40 pA-CMV-mCherry were commercially
generated and purchased from VectorBuilder. The amount of crude adenovirus was estimated
by infecting HEK293 cells with serially diluted amounts of crude adenovirus. For large scale
purification, HEK293 were infected, then collected 2-3 days later for purification with the Adeno-
X Maxi Purification kit (Takara Bio, Catalog No. 631533).
652

653   Particle titers of purified viruses were quantified by making three separate dilutions of
654 each virus in 0.1% SDS (Thermo Fisher) and vortexing for 5 minutes, spinning down 13,000 x
655 rpm, 5 minutes, and then taking the average of the absorbance readings on the
656 Nanophotometer N50-GO (Implen) to quantify particles/mL as per (Maizel et al., 1968).
657
658 *Neonatal rat ventricular myocyte seeding, siRNA knockdown, and rescue*
659
The 8-well chamber slides (Corning, Catalog No. 354118) were prepared by coating with
10 µg/mL fibronectin (Sigma, Catalog No. F1141) and 20 µg/mL poly-D-lysine hydrobromide
(Sigma, Catalog No. P6407) in PBS overnight at 4°C. Neonatal rat ventricular myocytes
(NRVMs) were seeded at 175,000 NRVMs/well in a mixture of 75% DMEM (Thermo Fisher,
Catalog No. 11965092), 15% Medium-199 (Thermo Fisher, Catalog No. 11150059)
supplemented with 2 mM L-glutamine (Thermo Fisher, Catalog No. A2916801) and 10 mM
HEPES (Thermo Fisher, Catalog No. 15630080). For reverse transfection, the cells were
treated with siRNA targeting FHOD3 (Qiagen, Rn_LOC100360334_2 Flexitube siRNA) or
AllStars Negative Control siRNA (Qiagen, 20 nmol), Lipofectamine RNAiMAX transfection
reagent (Thermo Fisher, Catalog No. 13778075) and Opti-MEM I Reduced Serum Medium
(Thermo Fisher, Catalog No. 31985062) to dilute the siRNA. Media was changed 24 hours later
to one containing penicillin/streptomycin (Thermo Fisher, Catalog No. 15140122). NRVMs were
infected with an optimal amount of adenovirus (determined experimentally to be particle titer
MOI 350 to minimize excessive damage to NRVMs while maintaining sufficient expression) as
above 48 hours after seeding and media containing penicillin/streptomycin was added on top of
the PBS+/+ at the end of the infection. Media was changed 24 hours later and cells were
examined 48 hours after infection.
677
678 *Gene expression analysis by quantitative reverse-transcriptase PCR*
679
RNA was extracted from the NRVMs four days after reverse transfection of siRNA using
the Direct-zol RNA mini prep kit (Zymo Research, Catalog No. R2050). RNA was reverse-
transcribed into complementary DNA using the qScript cDNA synthesis kit (Quanta Biosciences,
Catalog No. 95047-025). Quantitative reverse-transcriptase PCR was performed using
PowerUp SYBR green master mix for qPCR (Applied Biosystems, Catalog No. A25742) on a
Lightcycler 480 (Roche). Each qPCR reaction was repeated three times. Forward and reverse
primer sequences are as follows: GAPDH forward, CCGCATCTTCTTGTGCAGTG; GAPDH
reverse, CGATACGGCCAAATCCGTTC; FHOD3 forward, CAGCCAATCACGGAG; FHOD3
reverse, TGCTGTCCTTGCCCTGA.
689
690 *Western blots*
691
NRVMs were lysed in 100 mM Tris pH 8, 150 mM NaCl, 0.5% Triton-X from 8-well
chamber slides (Corning, Catalog No. 354118) after rescue experiments and samples were
vortexed for 1 minute before centrifuging at 15,000 rpm, 4°C, for 10 minutes. The resulting
supernatant was boiled in sample loading buffer at 100°C for 10 minutes and run on an SDS-
PAGE gel. The gel was transferred to an Immobilon-FL polyvinylidene fluoride membrane
(Millipore, IPFL00010) at 100V for 90 minutes on ice. The membrane was blocked in 4% nonfat
milk in low-salt TBST (20 mM Tris pH 7.6, 150 mM NaCl, 0.05% Tween-20, 0.01% sodium
azide) for 30 minutes at room temperature. It was incubated, rotating at 4°C overnight, with
Fhod3 polyclonal rabbit (Abcam, ab224463) or HA monoclonal rabbit (Cell Signaling

Technologies, 3724S) and GAPDH monoclonal mouse (Santa Cruz Biotechnology, sc-365062)
diluted 1:1000 in low-salt TBST. Membranes were washed 3 times for 5 minutes each the next
day in high-salt TBST (20 mM Tris pH 7.6, 500 mM NaCl, 0.05% Tween-20, 0.01% sodium
azide) and then incubated for 1 hour at room temperature, shaking, with 800CW goat anti-rabbit
IgG secondary antibody (Li-Cor Biosciences, 926-32211) and 680RD goat anti-mouse IgG
secondary antibody (Li-Cor Biosciences, 926-68070) diluted 1:10,000 in high-salt TBST.
Membranes were washed 3 times for 10 minutes each in high-salt TBST and then imaged on a
Li-Cor Odyssey 9120 Infrared Imager (Li-Cor Biosciences).
709
710 *In vitro contractility assay*
711
Contractility assessments were performed by utilizing a video-based technique with the
UCSF Gladstone-developed MATLAB program MotionGUI (Huebsch et al., 2015). Videos for
contractility analysis were acquired using MicroManager software on a Leica SD AF Spinning
Disc system using a HC PL Fluotar 10x (0.3 NA) dry objective lens with an ORCA-Flash4.0 LT
C11440 camera (Hamamatsu) at 30 fps with live NRVMs incubated at 37°C, 5% CO_2 (Tokai
Hit). The videos were converted from ome.tif to a tiff stack with FIJI for analysis with MotionGUI.
A pixel size of 0.538 μm was obtained from the metadata. 8 pixel macroblocks were used for all
assessments. All parameters of the MotionGUI program not specified here were set to their
respective default values. Motion vectors were calculated, and the data were evaluated upon
completion. All videos were subjected to the same post-processing procedures to ensure
consistency during comparative analysis. Each video sample was post-processed using
neighbor-based cleaning with the vector-based cleaning criterion within the program. The
threshold for this post-processing method was set to two for all samples and was adequate for
improving the signal-to-noise ratio enough to identify peaks clearly corresponding to beating
events in most samples.
727
728 *Immunofluorescence and image analysis*
729
For sarcomere integrity analysis, NRVMs were stained before fixation with Wheat Germ
Agglutinin (WGA) TMR (Thermo Fisher, Catalog No. W849) for 10 minutes at 37°C with 5
μg/mL WGA, followed by two PBS washes. NRVMs were then fixed with 4% paraformaldehyde
(Fisher Scientific, Catalog No. AC416780010) in PBS for 15 minutes at 37°C. Cells were
washed 3 times in PBS for 5 minutes each at room temperature. Cells were permeabilized and
blocked with PBS/0.1% Triton-X/10% Goat Serum (Triton-X: Thermo Fisher, Catalog No.
A16046.AP; Goat Serum: Sigma, Catalog No. S26-100ML) for 30 minutes at 37°C. Cells were
incubated with primary antibodies overnight at 4°C. α-actinin (mouse) primary antibody (Sigma,
Catalog No. A7811) was diluted 1:250 and HA-tag (rabbit) primary antibody (Cell Signaling,
Catalog No. 3724) was diluted 1:500 in PBS/0.1% Triton-X/5% Goat Serum. Cells were washed
3 times for 5 minutes each with PBS/0.1% Triton-X/5% Goat Serum at room temperature.
Incubation with secondary antibodies was for 1 hour at 37°C. Alexa Fluor 488 goat anti-mouse
secondary (Thermo Fisher, Catalog No. A-11001) or Alexa Fluor 647 goat anti-rabbit secondary
(Thermo Fisher, Catalog No. A-21244) was diluted 1:500 in PBS/0.1% Triton-X/5% Goat Serum.
Cells were then washed twice in PBS/0.1% Triton-X/5% Goat Serum for 5 minutes each
followed by on PBS wash for 5 minutes before mounting in Vectashield Plus Antifade mounting
media (Vector Laboratories, Catalog No. H-1900-10) with 1 μg/mL DAPI (Fisher Scientific,
Catalog No. EN62248).

For thin filament determination, NRVM's were incubated with sterile-filtered Ringer's relaxation buffer (6 mM potassium phosphate pH 7.0, 100 mM NaCl, 2 mM KCl, 0.1% glucose, 2 mM $MgCl_2$, 1 mM EGTA) for 20 min., room temperature, followed by a brief PBS wash to start the WGA staining. Fixation and blocking were performed as above. Staining with HA-tag (rabbit) primary antibody Alexa Fluor 647 goat anti-rabbit secondary antibody were performed as above. Cells were then washed twice in PBS/0.1% Triton-X/5% Goat Serum for 5 minutes each and 200 nM Alexa Fluor 488 phalloidin (Thermo Fisher, Catalog No. A12379) was added for 1 hr, room temperature in PBS/1% BSA. Cells were washed with PBS twice for 5 minutes before mounting in Prolong Glass Antifade mountant (Thermo Fisher, Catalog No. P36980) with 1 μg/mL DAPI (Fisher Scientific, Catalog No. EN62248).

Twelve 40X magnification images per biological replicate were acquired using a random coordinate generator (https://onlineintegertools.com/generate-integer-pairs), excluding the edges of the wells, for quantification of sarcomeres and thin filaments. Sample size with sufficient power was determined via a small-scale pilot study for sarcomere number, length, and width, as well as thin filaments between mock knockdown, mock rescue, and FHOD3L WT rescue followed by inputting averages of the treatments on https://www.stat.ubc.ca/~rollin/stats/ssize/n1.html. Each well was treated as a 16x16 grid of 40X magnification fields of view. 16-bit images were acquired on an AXIO Imager.D1 fluorescence phase contrast microscope (Zeiss) using an EC Plan-Neofluor 40x M27 (0.75 NA) objective lens with an AxioCam MRm camera (Zeiss). Images of NRVMs in figures were acquired with a Plan-Apochromat oil DIC M27 63x (1.4 NA) objective lens for visualization.

Images showing DsRed reporter fluorescence at the end of the rescue timeline for the NRVMs were taken on a CKX53 inverted phase contrast microscope (Olympus) using a UPLFLN 4x (0.2 NA) objective lens (Olympus) with an ORCA-spark digital CMOS camera (Hamamatsu).

Cells were segmented using Cellpose (Stringer et al., 2021). Two-color images of the WGA membrane stain and nuclear stain were saved as jpeg files on FIJI. These images were then input to the Cellpose environment for batch processing with a cell diameter of 210 pixels, 0.4 cell probability and flow thresholds, 0 stitch threshold, using the cyto2 model for segmentation. The text files of the outlines were saved and overlaid on the respective channels of the images individually using the provided python file from Cellpose as a macro through FIJI (imagej_roi_converter.py). Cellpose segments with no nuclei or 3 or more nuclei, as well as cell segments on the edge of the image were excluded from analysis.

Sarcomere analysis was performed manually in a single-blind manner (for mock knockdown, mock rescue, wild-type rescue, and the K1193L rescue) using blindrename.pl (https://github.com/davalencia0914/sarcApp_Cellpose_Merge) to generate the filenames and a key filename csv file for decoding after analysis. We attempted to blind all other rescue conditions, but they were too easily identified based on expression level and localization differences. Linescans were generated along myofibrils, perpendicular to the Z-lines to make sarcomere length measurements from Z-line peak to Z-line peak. Z-line lengths were measured by visual inspection of the alpha-actinin channel and measured on FIJI with the line tool. Three or more consecutive Z-lines at least 0.70 μm long in a row were analyzed as sarcomeres.

Statistical analysis

To compare two or more groups for the rescue experiments in NRVMs, pair-wise comparisons were performed. In order to reduce Type 1 error rate stemming from multiple comparisons, Bonferroni correction was applied to obtain a corrected alpha. Whether groups were normally distributed or not was determined by the Shapiro-Wilk test. To compare two normally distributed groups, Student's two-sample, unpaired t-test was used. If either group was not normally distributed, the non-parametric Mann-Whitney U test was used. The same analysis was applied to analyze run lengths and capping duration from the TIRF seeded elongation assays.

For the nucleation assay, barbed-end binding assays, and TIRF elongation rate analysis, one-way ANOVA's were performed for 3 treatment comparisons with post-hoc Tukey tests as the variances between all treatments tested were fairly equal.

In all cases when the sample size exceeded 20, we removed outliers from the bottom and upper 2.5% of data to perform all statistical tests, regardless of normality, to avoid introducing bias. However, all plots shown include these outliers.

Data Availability

All data described in the manuscript are contained within the manuscript. Additional information and data are available upon request. All code used can be found on Github (https://github.com/davalencia0914/sarcApp_Cellpose_Merge).

Funding and additional information – This work was supported by NIH grant R01 HL146159 to M.E.Q., and Ruth L. Kirschstein National Research Service Award T32 GM007185 to D.A.V. and T32 AR065972 to A.N.K. Confocal laser scanning microscopy was performed at the Advanced Light Microscopy/Spectroscopy Laboratory and Leica Microsystems Center of Excellence at the California NanoSystems Institute at UCLA (RRID:SCR_022789) with funding support from NIH Shared Instrumentation Grant S10OD025017 and NSF Major Research Instrumentation grant CHE-072251.

Author contributions – D. A. V., A. N. K., A. N., and M. E. Q. designed the experiments. D. A. V. and A. N. K. performed the experiments. A. H. and H. N. provided essential experimental support and advice. D. A. V. and A. R. performed the statistical analysis. D. A. V. and C. S. performed manual sarcomere analysis. D. A. V., H. N., A. N., and M. E. Q. wrote the manuscript with input from all authors.

Acknowledgements – We thank members of the Quinlan lab and Nakano lab for experimental support and scientific feedback, the Reisler lab for valuable discussions and reagents, the Courtemanche lab for NEM-myosin, and T. Iskratsch for the FHOD plasmids. We also thank Dr. Michael D. Roth for access to his lab's BSL II-equipped facilities, enabling the production of adenovirus. Neonatal rat ventricular myocytes (NRVMs) were isolated from post-natal P1-P3 day old Sprague-Dawley rat pups of mixed gender by UCLA Cardiovascular Research Theme Core services. We thank Brian Jeong at California NanoSystems Institute at UCLA for assisting us with video imaging for contractility assays.

References

Al Haj A, Mazur AJ, Radaszkiewicz K, Radaszkiewicz T, Makowiecka A, Stopschinski BE, Schönichen A, Geyer M, Mannherz HG. 2015. Distribution of formins in cardiac muscle: FHOD1 is a component of intercalated discs and costameres. *European Journal of Cell Biology* **94**:101–113. doi:10.1016/j.ejcb.2014.11.003

Antoku S, Schwartz TU, Gundersen GG. 2023. FHODs: Nuclear tethered formins for nuclear mechanotransduction. *Frontiers in Cell and Developmental Biology* **11**.

Arimura T, Takeya R, Ishikawa T, Yamano T, Matsuo A, Tatsumi T, Nomura T, Sumimoto H, Kimura A. 2013. Dilated Cardiomyopathy-Associated *FHOD3* Variant Impairs the Ability to Induce Activation of Transcription Factor Serum Response Factor. https://www.jstage.jst.go.jp/article/circj/77/12/77_CJ-13-0255/_article

Baer A, Kehn-Hall K. 2014. Viral Concentration Determination Through Plaque Assays: Using Traditional and Novel Overlay Systems. *J Vis Exp* 52065. doi:10.3791/52065

Baker JL, Courtemanche N, Parton DL, McCullagh M, Pollard TD, Voth GA. 2015. Electrostatic Interactions between the Bni1p Formin FH2 Domain and Actin Influence Actin Filament Nucleation. *Structure* **23**:68–79. doi:10.1016/j.str.2014.10.014

Bechtold M, Schultz J, Bogdan S. 2014. FHOD proteins in actin dynamics—a formin' class of its own. *Small GTPases* **5**:e973765. doi:10.4161/21541248.2014.973765

Blanchoin L, Boujemaa-Paterski R, Sykes C, Plastino J. 2014. Actin Dynamics, Architecture, and Mechanics in Cell Motility. *Physiological Reviews* **94**:235–263. doi:10.1152/physrev.00018.2013

Bor B, Vizcarra CL, Phillips ML, Quinlan ME. 2012. Autoinhibition of the formin Cappuccino in the absence of canonical autoinhibitory domains. *Mol Biol Cell* **23**:3801–3813. doi:10.1091/mbc.E12-04-0288

Boussaty EC, Ninoyu Y, Andrade L, Li Q, Takeya R, Sumimoto H, Ohyama T, Wahlin KJ, Manor U, Friedman RA. 2023. Altered Fhod3 Expression Involved in Progressive High-Frequency Hearing Loss via Dysregulation of Actin Polymerization Stoichiometry in The Cuticular Plate. *bioRxiv* 2023.07.20.549974. doi:10.1101/2023.07.20.549974

Bremer KV, Wu C, Patel AA, He KL, Grunfeld AM, Chanfreau GF, Quinlan ME. 2024. Formin tails act as a switch, inhibiting or enhancing processive actin elongation. *J Biol Chem* **300**:105557. doi:10.1016/j.jbc.2023.105557

Cao L, Kerleau M, Suzuki EL, Wioland H, Jouet S, Guichard B, Lenz M, Romet-Lemonne G, Jegou A. 2018. Modulation of formin processivity by profilin and mechanical tension. *Elife* **7**:e34176. doi:10.7554/eLife.34176

Courtemanche N. 2018. Mechanisms of formin-mediated actin assembly and dynamics. *Biophys Rev* **10**:1553–1569. doi:10.1007/s12551-018-0468-6

Cutler KJ, Stringer C, Wiggins PA, Mougous JD. 2021. Omnipose: a high-precision morphology-independent solution for bacterial cell segmentation (preprint). Microbiology. doi:10.1101/2021.11.03.467199

Deng S, Silimon RL, Balakrishnan M, Bothe I, Juros D, Soffar DB, Baylies MK. 2021. The actin polymerization factor Diaphanous and the actin severing protein Flightless I collaborate to regulate sarcomere size. *Developmental Biology* **469**:12–25. doi:10.1016/j.ydbio.2020.09.014

Dominguez R, Holmes KC. 2011. Actin Structure and Function. *Annu Rev Biophys* **40**:169–186. doi:10.1146/annurev-biophys-042910-155359

Dwyer J, Pluess M, Iskratsch T, Dos Remedios CG, Ehler E. 2014. The formin FHOD1 in cardiomyocytes. *Anat Rec (Hoboken)* **297**:1560–1570. doi:10.1002/ar.22984

Fenix AM, Neininger AC, Taneja N, Hyde K, Visetsouk MR, Garde RJ, Liu B, Nixon BR, Manalo AE, Becker JR, Crawley SW, Bader DM, Tyska MJ, Liu Q, Gutzman JH, Burnette DT. 2018. Muscle-specific stress fibers give rise to sarcomeres in cardiomyocytes. *eLife* **7**:e42144. doi:10.7554/eLife.42144

Ferreira-Martins J, Leite-Moreira AF. 2010. Physiologic Basis and Pathophysiologic Implications of the Diastolic Properties of the Cardiac Muscle. *J Biomed Biotechnol* **2010**:807084. doi:10.1155/2010/807084

Fujimoto N, Kan-O M, Ushijima T, Kage Y, Tominaga R, Sumimoto H, Takeya R. 2016. Transgenic Expression of the Formin Protein Fhod3 Selectively in the Embryonic Heart: Role of Actin-Binding Activity of Fhod3 and Its Sarcomeric Localization during Myofibrillogenesis. *PLoS ONE* **11**:e0148472. doi:10.1371/journal.pone.0148472

Gasteier JE, Madrid R, Krautkrämer E, Schröder S, Muranyi W, Benichou S, Fackler OT. 2003. Activation of the Rac-binding partner FHOD1 induces actin stress fibers via a ROCK-dependent mechanism. *J Biol Chem* **278**:38902–38912. doi:10.1074/jbc.M306229200

Goode BL, Eck MJ. 2007. Mechanism and Function of Formins in the Control of Actin Assembly. *Annual Review of Biochemistry* **76**:593–627. doi:10.1146/annurev.biochem.75.103004.142647

Homa KE, Zsolnay V, Anderson CA, O'Connell ME, Neidt EM, Voth GA, Bidone TC, Kovar DR. 2021. Formin Cdc12's specific actin assembly properties are tailored for cytokinesis in fission yeast. *Biophysical Journal* **120**:2984–2997. doi:10.1016/j.bpj.2021.06.023

Huebsch N, Loskill P, Mandegar MA, Marks NC, Sheehan AS, Ma Z, Mathur A, Nguyen TN, Yoo JC, Judge LM, Spencer CI, Chukka AC, Russell CR, So P-L, Conklin BR, Healy KE. 2015. Automated Video-Based Analysis of Contractility and Calcium Flux in Human-Induced Pluripotent Stem Cell-Derived Cardiomyocytes Cultured over Different Spatial Scales. *Tissue Engineering Part C: Methods* **21**:467–479. doi:10.1089/ten.tec.2014.0283

Iskratsch T, Lange S, Dwyer J, Kho AL, Remedios C dos, Ehler E. 2010. Formin follows function: a muscle-specific isoform of FHOD3 is regulated by CK2 phosphorylation and promotes myofibril maintenance. *Journal of Cell Biology* **191**:1159–1172. doi:10.1083/jcb.201005060

Iskratsch T, Yu C-H, Mathur A, Liu S, Stévenin V, Dwyer J, Hone J, Ehler E, Sheetz M. 2013. FHOD1 Is Needed for Directed Forces and Adhesion Maturation during Cell Spreading and Migration. *Developmental Cell* **27**:545–559. doi:10.1016/j.devcel.2013.11.003

Kanaya H, Takeya R, Takeuchi K, Watanabe N, Jing N, Sumimoto H. 2005. Fhos2, a novel formin-related actin-organizing protein, probably associates with the nestin intermediate filament. *Genes to Cells* **10**:665–678. doi:10.1111/j.1365-2443.2005.00867.x

Kan-o M, Takeya R, Abe T, Kitajima N, Nishida M, Tominaga R, Kurose H, Sumimoto H. 2012a. Mammalian formin Fhod3 plays an essential role in cardiogenesis by organizing myofibrillogenesis | Biology Open | The Company of Biologists. https://journals.biologists.com/bio/article/1/9/889/19745/Mammalian-formin-Fhod3-plays-an-essential-role-in

Kan-O M, Takeya R, Abe T, Kitajima N, Nishida M, Tominaga R, Kurose H, Sumimoto H. 2012.
Mammalian formin Fhod3 plays an essential role in cardiogenesis by organizing
myofibrillogenesis. *Biology Open* **1**:889–896. doi:10.1242/bio.20121370

Kan-o M, Takeya R, Taniguchi K, Tanoue Y, Tominaga R, Sumimoto H. 2012b. Expression and
Subcellular Localization of Mammalian Formin Fhod3 in the Embryonic and Adult Heart.
PLOS ONE **7**:e34765. doi:10.1371/journal.pone.0034765

Kimmich MJ, Sundaramurthy S, Geary MA, Lesanpezeshki L, Yingling CV, Vanapalli SA, Littlefield
RS, Pruyne D. 2024. FHOD-1/profilin-mediated actin assembly protect sarcomeres
against contraction-induced deformation in C. elegans. doi:10.1101/2024.02.29.582848

Koka S, Neudauer CL, Li X, Lewis RE, McCarthy JB, Westendorf JJ. 2003. The formin-homology-
domain-containing protein FHOD1 enhances cell migration. *Journal of Cell Science*
116:1745–1755. doi:10.1242/jcs.00386

Kovar DR, Harris ES, Mahaffy R, Higgs HN, Pollard TD. 2006. Control of the Assembly of ATP- and
ADP-Actin by Formins and Profilin. *Cell* **124**:423–435. doi:10.1016/j.cell.2005.11.038

Kutscheidt S, Zhu R, Antoku S, Luxton GWG, Stagljar I, Fackler OT, Gundersen GG. 2014. FHOD1
interaction with nesprin-2G mediates TAN line formation and nuclear movement. *Nat*
Cell Biol **16**:708–715. doi:10.1038/ncb2981

Lappalainen P, Kotila T, Jégou A, Romet-Lemonne G. 2022. Biochemical and mechanical
regulation of actin dynamics. *Nat Rev Mol Cell Biol* **23**:836–852. doi:10.1038/s41580-
022-00508-4

Li D, Hallett MA, Zhu W, Rubart M, Liu Y, Yang Z, Chen H, Haneline LS, Chan RJ, Schwartz RJ,
Field LJ, Atkinson SJ, Shou W. 2011. Dishevelled-associated activator of morphogenesis 1
(Daam1) is required for heart morphogenesis. *Development* **138**:303–315.
doi:10.1242/dev.055566

Li F, Higgs HN. 2005. Dissecting Requirements for Auto-inhibition of Actin Nucleation by the
Formin, mDia1. *J Biol Chem* **280**:6986–6992. doi:10.1074/jbc.M411605200

Liu H, Naismith JH. 2008. An efficient one-step site-directed deletion, insertion, single and
multiple-site plasmid mutagenesis protocol. *BMC Biotechnol* **8**:91. doi:10.1186/1472-
6750-8-91

Lu J, Pollard TD. 2001. Profilin Binding to Poly-l-Proline and Actin Monomers along with Ability
to Catalyze Actin Nucleotide Exchange Is Required for Viability of Fission Yeast. *Mol Biol*
Cell **12**:1161–1175. doi:10.1091/mbc.12.4.1161

Maizel JV, White DO, Scharff MD. 1968. The polypeptides of adenovirus: I. Evidence for multiple
protein components in the virion and a comparison of types 2, 7A, and 12. *Virology*
36:115–125. doi:10.1016/0042-6822(68)90121-9

Matsuyama S, Kage Y, Fujimoto N, Ushijima T, Tsuruda T, Kitamura K, Shiose A, Asada Y,
Sumimoto H, Takeya R. 2018a. Interaction between cardiac myosin-binding protein C
and formin Fhod3. *Proceedings of the National Academy of Sciences* **115**:E4386–E4395.
doi:10.1073/pnas.1716498115

Matsuyama S, Kage Y, Fujimoto N, Ushijima T, Tsuruda T, Kitamura K, Shiose A, Asada Y,
Sumimoto H, Takeya R. 2018b. Interaction between cardiac myosin-binding protein C
and formin Fhod3. *Proceedings of the National Academy of Sciences* **115**:E4386–E4395.
doi:10.1073/pnas.1716498115

McInally SG, Kondev J, Goode BL. 2021. Scaling of subcellular actin structures with cell length
through decelerated growth. *eLife* **10**:e68424. doi:10.7554/eLife.68424
McInally SG, Reading AJB, Rosario A, Jelenkovic PR, Goode BL, Kondev J. 2024. Length control
emerges from cytoskeletal network geometry. *Proceedings of the National Academy of*
Sciences **121**:e2401816121. doi:10.1073/pnas.2401816121
Mi-Mi L, Votra S, Kemphues K, Bretscher A, Pruyne D. 2012. Z-line formins promote contractile
lattice growth and maintenance in striated muscles of C. elegans. *J Cell Biol* **198**:87–102.
doi:10.1083/jcb.201202053
Molnár I, Migh E, Szikora S, Kalmár T, Végh AG, Deák F, Barkó S, Bugyi B, Orfanos Z, Kovács J,
Juhász G, Váró G, Nyitrai M, Sparrow J, Mihály J. 2014. DAAM Is Required for Thin
Filament Formation and Sarcomerogenesis during Muscle Development in Drosophila.
PLOS Genetics **10**:e1004166. doi:10.1371/journal.pgen.1004166
Moseley JB, Goode BL. 2005. Differential Activities and Regulation of Saccharomyces cerevisiae
Formin Proteins Bni1 and Bnr1 by Bud6. *Journal of Biological Chemistry* **280**:28023–
28033. doi:10.1074/jbc.M503094200
Myasnikov R, Bukaeva A, Kulikova O, Meshkov A, Kiseleva A, Ershova A, Petukhova A, Divashuk
M, Zotova E, Sotnikova E, Kharlap M, Zharikova A, Vyatkin Y, Ramensky V, Abisheva A,
Muraveva A, Koretskiy S, Kudryavtseva M, Popov S, Utkina M, Mershina E, Sinitsyn V,
Kogan E, Blagova O, Drapkina O. 2022. A Case of Severe Left-Ventricular Noncompaction
Associated with Splicing Altering Variant in the FHOD3 Gene. *Genes* **13**:309.
doi:10.3390/genes13020309
Nakano H, Minami I, Braas D, Pappoe H, Wu X, Sagadevan A, Vergnes L, Fu K, Morselli M,
Dunham C, Ding X, Stieg AZ, Gimzewski JK, Pellegrini M, Clark PM, Reue K, Lusis AJ,
Ribalet B, Kurdistani SK, Christofk H, Nakatsuji N, Nakano A. 2017. Glucose inhibits
cardiac muscle maturation through nucleotide biosynthesis. *eLife* **6**:e29330.
doi:10.7554/eLife.29330
Neininger-Castro AC, Jr JBH, Sanchez ZC, Taneja N, Fenix AM, Moparthi S, Vassilopoulos S,
Burnette DT. 2023. Independent regulation of Z-lines and M-lines during sarcomere
assembly in cardiac myocytes revealed by the automatic image analysis software
sarcApp. *eLife* **12**. doi:10.7554/eLife.87065.2
Ochoa JP, Sabater-Molina M, Garcia-Pinilla JM, Mogensen J, Restrepo-Cordoba A, Palomino-
Doza J, Villacorta E, Martinez-Moreno M, Ramos-Maqueda J, Zorio E, Peña-Peña ML,
García-Granja PE, Rodríguez-Palomares JF, Cárdenas-Reyes IJ, de la Torre-Carpente MM,
Bautista-Pavés A, Akhtar MM, Cicerchia MN, Bilbao-Quesada R, Mogollón-Jimenez MV,
Salazar-Mendiguchía J, Mesa Latorre JM, Arnaez B, Olavarri-Miguel I, Fuentes-Cañamero
ME, Lamounier Jr A, Pérez Ruiz JM, Climent-Payá V, Monserrat L. 2018. Formin
Homology 2 Domain Containing 3 (FHOD3) Is a Genetic Basis for Hypertrophic
Cardiomyopathy - ScienceDirect.
https://www.sciencedirect.com/science/article/pii/S073510971838673X?via%3Dihub
Patel AA, Oztug Durer ZA, van Loon AP, Bremer KV, Quinlan ME. 2018. Drosophila and human
FHOD family formin proteins nucleate actin filaments. *J Biol Chem* **293**:532–540.
doi:10.1074/jbc.M117.800888

Paul A, Pollard T. 2008. The Role of the FH1 Domain and Profilin in Formin-Mediated Actin-
1014   Filament Elongation and Nucleation. *Current Biology* **18**:9–19.
1015   doi:10.1016/j.cub.2007.11.062
1016   Paul AS, Pollard TD. 2009. Review of the mechanism of processive actin filament elongation by
1017   formins. *Cell Motil Cytoskeleton* **66**:606–17. doi:10.1002/cm.20379
1018   Pollard TD. 2016. Actin and Actin-Binding Proteins. *Cold Spring Harb Perspect Biol* a018226.
1019   doi:10.1101/cshperspect.a018226
1020   Rosado M, Barber CF, Berciu C, Feldman S, Birren SJ, Nicastro D, Goode BL. 2014. Critical roles
1021   for multiple formins during cardiac myofibril development and repair. *Mol Biol Cell*
1022   **25**:811–827. doi:10.1091/mbc.E13-08-0443
1023   Sanematsu F, Kanai A, Ushijima T, Shiraishi A, Abe T, Kage Y, Sumimoto H, Takeya R. 2019.
1024   Fhod1, an actin-organizing formin family protein, is dispensable for cardiac development
1025   and function in mice. *Cytoskeleton* cm.21523. doi:10.1002/cm.21523
1026   Sanger JW, Wang J, Fan Y, White J, Mi-Mi L, Dube DK, Sanger JM, Pruyne D. 2017. Assembly and
1027   Maintenance of Myofibrils in Striated Muscle. *Handb Exp Pharmacol* **235**:39–75.
1028   doi:10.1007/164_2016_53
1029   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,
1030   Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P,
1031   Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat*
1032   *Methods* **9**:676–682. doi:10.1038/nmeth.2019
1033   Schonichen A, Geyer M. 2010. Fifteen formins for an actin filament: a molecular view on the
1034   regulation of human formins. *Biochim Biophys Acta* **1803**:152–63.
1035   doi:10.1016/j.bbamcr.2010.01.014
1036   Schonichen A, Mannherz HG, Behrmann E, Mazur AJ, Kuhn S, Silvan U, Schoenenberger C-A,
1037   Fackler OT, Raunser S, Dehmelt L, Geyer M. 2013. FHOD1 is a combined actin filament
1038   capping and bundling factor that selectively associates with actin arcs and stress fibers.
1039   *Journal of Cell Science* **126**:1891–1901. doi:10.1242/jcs.126706
1040   Schulze N, Graessl M, Soares AB, Geyer M, Dehmelt L, Nalbant P. 2014. FHOD1 regulates stress
1041   fiber organization by controlling the dynamics of transverse arcs and dorsal fibers. *J Cell*
1042   *Sci* **127**:1379–1393. doi:10.1242/jcs.134627
1043   Shwartz A, Dhanyasi N, Schejter ED, Shilo B-Z. 2016. The Drosophila formin Fhos is a primary
1044   mediator of sarcomeric thin-filament array assembly. *eLife Sciences* **5**:e16540.
1045   doi:10.7554/eLife.16540
1046   Smith MB, Li H, Shen T, Huang X, Yusuf E, Vavylonis D. 2010. Segmentation and tracking of
1047   cytoskeletal filaments using open active contours. *Cytoskeleton (Hoboken)* **67**:693–705.
1048   doi:10.1002/cm.20481
1049   Spudich JA, Watt S. 1971. The Regulation of Rabbit Skeletal Muscle Contraction. *Journal of*
1050   *Biological Chemistry* **246**:4866–4871. doi:10.1016/S0021-9258(18)62016-2
1051   Stringer C, Wang T, Michaelos M, Pachitariu M. 2021. Cellpose: a generalist algorithm for
1052   cellular segmentation. *Nat Methods* **18**:100–106. doi:10.1038/s41592-020-01018-x
1053   Sun H, Luo Y, Miao Y. 2018. Purification of Globular Actin from Rabbit Muscle and Pyrene
1054   Fluorescent Assays to Investigate Actin Dynamics in vitro. *Bio Protoc* **8**:e3102.
1055   doi:10.21769/BioProtoc.3102

Sundaramurthy S, Votra S, Laszlo A, Davies T, Pruyne D. 2020. FHOD-1 is the only formin in Caenorhabditis elegans that promotes striated muscle growth and Z-line organization in a cell autonomous manner. *Cytoskeleton* **77**:422–441. doi:10.1002/cm.21639

Svitkina T. 2018. The Actin Cytoskeleton and Actin-Based Motility. *Cold Spring Harb Perspect Biol* **10**:a018267. doi:10.1101/cshperspect.a018267

Szikora S, Görög P, Mihály J. 2022. The Mechanisms of Thin Filament Assembly and Length Regulation in Muscles. *International Journal of Molecular Sciences* **23**:5306. doi:10.3390/ijms23105306

Taniguchi K, Takeya R, Suetsugu S, Kan-o M, Narusawa M, Shiose A, Tominaga R, Sumimoto H. 2009. Mammalian formin Fhod3 regulates actin assembly and sarcomere organization in striated muscles. *Journal of Biological Chemistry* **284**:29873–29881. doi:10.1074/jbc.M109.059303

Ushijima T, Fujimoto N, Matsuyama S, Kan-o M, Kiyonari H, Shioi G, Kage Y, Yamasaki S, Takeya R, Sumimoto H. 2018. The actin-organizing formin protein Fhod3 is required for postnatal development and functional maintenance of the adult heart in mice. *J Biol Chem* **293**:148–162. doi:10.1074/jbc.M117.813931

Vizcarra CL, Bor B, Quinlan ME. 2014. The Role of Formin Tails in Actin Nucleation, Processive Elongation, and Filament Bundling. *J Biol Chem* **289**:30602–30613. doi:10.1074/jbc.M114.588368

Vodnjov N, Toplišek J, Maver A, Čuturilo G, Jaklič H, Teran N, Višnjar T, Pušenjak MŠ, Hodžić A, Miljanović O, Peterlin B, Writzl K. 2023. A novel splice-site FHOD3 founder variant is a common cause of hypertrophic cardiomyopathy in the population of the Balkans–A cohort study. *PLOS ONE* **18**:e0294969. doi:10.1371/journal.pone.0294969

Wu G, Ruan J, Liu J, Zhang C, Kang L, Wang J, Zou Y, Song L. 2021. Variant Spectrum of Formin Homology 2 Domain-Containing 3 Gene in Chinese Patients With Hypertrophic Cardiomyopathy. *Journal of the American Heart Association* **10**:e018236. doi:10.1161/JAHA.120.018236

Xu Y, Moseley JB, Sagot I, Poy F, Pellman D, Goode BL, Eck MJ. 2004. Crystal Structures of a Formin Homology-2 Domain Reveal a Tethered Dimer Architecture. *Cell* **116**:711–723. doi:10.1016/S0092-8674(04)00210-7

Zweifel ME, Courtemanche N. 2020. Competition for delivery of profilin–actin to barbed ends limits the rate of formin-mediated actin filament elongation. *J Biol Chem* **295**:4513–4525. doi:10.1074/jbc.RA119.012000

Figure 1

1091
Figure 1.   Domain structure of human FHOD3.   GDB=GTP Binding Domain, DID=Diaphanous
Inhibitory Domain, CC=Coiled-coil, FH=Formin Homology, DAD=Diaphanous Autoregulatory Domain.
Mutations tested in this study are indicated with black arrows. Numbers correspond to FHOD 3L
sequence (Uniprot Isoform 4). The FHOD3L construct spans residues 963–1622 and includes the FH1
domain, FH2 domain, and tail. The FHOD3S construct spans residues 771-1422 (Uniprot Isoform 1
numbering). FHOD3L-specific exons are in gray. The 8-residue $T(D/E)_5XE$ exon that distinguishes
FHOD3L-CT from FHOD3S-CT is circled in red.

1099
1100
1101

Figure 1 Supplement 1

1102
Figure 1—figure supplement 1. Coomassie-stained polyacrylamide gel showing purified FHOD3
constructs. Lower molecular weight contaminants were difficult to remove without significantly diluting the
sample. They were relatively consistent for each construct, and had no detectable impact on FHOD
activity.

Figure 2

1107
Figure 2. Biochemical characterization of human FHOD3L. (**A**) Assembly of 4 µM actin (5% pyrene-
labeled) and the indicated concentrations of FHOD3L-CT from a 2-fold dilution series. The inset shows
the first 200 seconds of data normalized to the plateau of actin alone. (**B**) Relative nucleation activities of
FHOD3L-CT (n=5), FHOD3S-CT (n=4), FHOD3L-CT K1309A (n=3) and I1163A (n=3). Data are means ±
SD. Slopes reported are the average slopes of independent experiments. The asterisks indicate
significance of difference from FHOD3L-CT (see below for details). (**C**) Barbed-end elongation assay.
Final conditions were 0.25 µM F-actin seeds (~0.1 nM barbed ends), 0.5 µM actin (10% pyrene-labeled),
and indicated concentrations FHOD3L-CT. (**D**) Quantification of barbed-end affinity for FHOD3L-CT (from
C), FHOD3S-CT (from Figure 2, figure supplement 2B) and FHOD3L-CT I1163A. Raw data are shown

and the lines are fits to all data points. The K_ds reported are the averages of three independent trials.
(n=3, each; mean ± SD.) (**E**) Barbed-end elongation assay with profilin. Final conditions as in (**C**) plus 1.5
µM *S. pombe* profilin. (**F**) Kymographs of growing filaments from TIRF assays with or without FHOD3L-
CT. Conditions: 1 µM actin (10%-Alexa Fluor 488-labeled), 5 µM Hs profilin-1 ± 0.1 nM FHOD3L-CT. The
red box highlights a pause followed by a burst of elongation. Vertical scale bar, 5 µm. (**G**) Elongation
rates from TIRF assays. Average elongation rates (over 10s of seconds) and formin-mediated elongation
rates (dim) are shown separately. (n=21, profilin-actin; n=20, FHOD3L-CT (avg); n=112, FHOD3L-CT
(dim); 3 channels for all samples; mean ± SD.) Scale bars, 10 µm. (**H**) Coomassie-stained
polyacrylamide gel of pellet fractions from low-speed bundling assays with 5 µM actin and 0-60 nM
FHOD3L-CT. (**I**) Quantification of bundling from (**H**) via densitometry (n=3 each; mean ± SD.) p-values
were determined by one-way ANOVA with post-hoc Tukey test. ** $p < 0.001$, *** $p < 0.0001$.
1128
1129

Figure 2 - supplement 1

1130

Figure 2—figure supplement 1 (A) Actin polymerized in the presence or absence of 12 nM FHOD3L-
CT, stabilized with fluorescent phalloidin and diluted to 5 nM. (n=3, each; mean $\pm$ SD) Scale bars, 10 µm.
(**B**) Pyrene fluorescence readings of 4 µM actin filaments (5% pyrene-labeled) incubated with different
concentrations of FHOD3L-CT I1163A for 5 minutes. (n=2; mean $\pm$ SD.) **(C)** Time-lapse images from
TIRF microscopy. Conditions: 1 µM actin (10%-Alexa Fluor 488-labeled), 5 µM Hs profilin-1 $\pm$ 0.1 nM
FHOD3L-CT. Yellow arrows indicate beginning and end of dim portion of filament elongated by FHOD3L-
CT. 2.5 s / frame. Scale bar, 10 µm. (**D**) Duration of pauses observed in TIRF assays. (**F**) Run lengths
observed in TIRF assays. (For (**D-E**) n=112 (dim); 3 flow channels; mean $\pm$ SD.) **(E)** Epifluorescence
micrographs of phalloidin-stabilized filaments from low-speed bundling assays. Conditions: 5 µM actin
filaments with or without 15 nM FHOD3L-CT, diluted to 5 nM actin after centrifugation. Scale bars, 10 µm.
1141

Figure 2 Supplement 2

1142
Figure 2—figure supplement 2. Comparison of FHOD3S to FHOD3L. (**A**) Assembly of 4 µM actin (5%
pyrene-labeled) and the indicated concentrations of FHOD3S from a 2-fold dilution series. (**B**) Barbed-
end elongation assay. Final conditions: 0.25 µM F-actin seeds (~0.1 nM barbed ends), 0.5 µM G-actin
(10% pyrene-labeled), and 312.5 pM-5 nM FHOD3S-CT. (**C**) Elongation rates from TIRF assays. Average
elongation rates (over 10s of seconds) and formin-mediated elongation rates (dim) are shown separately.
Conditions: 1 µM actin (10%-Alexa Fluor 488-labeled), 5 µM Hs profilin-1 ± 0.1 nM FHOD3L-CT or 1 nM
FHOD3S-CT. (profilin-actin n=21; 3L (avg) n=20; 3S (avg) n=24; 3L (dim) n=112; 3S (dim) n=78; 4 flow
channels for 3S, 3 flow channels for all others; mean ± SD, p-values by one-way ANOVA with post-hoc
Tukey test.) (**D**) Quantification of (dim) run lengths for FHOD3S/L-CT TIRF assays. (FHOD3L-CT n=97;
FHOD3S-CT n=73; 3 flow channels for 3L, 4 flow channels for 3S; mean ± SD, p-values by Mann-
Whitney U test.) (**E**) Duration of capping events by FHOD3L-CT vs FHOD3S-CT observed by TIRF
microscopy. (FHOD3L-CT n=97; FHOD3S-CT n=73; 3 flow channels for 3L, 4 flow channels for 3S; mean
± SD, p-values by Mann-Whitney U test.) (**F**) Coomassie-stained polyacrylamide gel of pellet fractions
from low-speed bundling assays with 5 µM actin and 0-60 nM FHOD3S-CT. (**H**) Comparison of bundling
for FHOD3S/L-CT via densitometry (n=3 each group; mean ± SD.) ns p > 0.05, * p < 0.05, *** p < 0.0001
1158
1159

Figure 3

Figure 3. Biochemical validation of function-separating mutants. (**A**) Relative nucleation activity for FHOD3L-CT (n=5) and FHOD3L-CT K1193L (n=5). Data points are means and error bars are standard deviations. Slopes reported are the average slopes of independent experiments. (**B**) Barbed-end affinity measurements for FHOD3L-CT, FHOD3L-CT GS-FH1, and FHOD3L-CT K1193L. Raw data are shown and the line is a fit to all data points. The K_ds reported are the average of three independent trials. (n=3, each; mean $\pm$ SD.) (**C**) Elongation rates from TIRF assays. Conditions: 1 µM actin (10%-Alexa Fluor 488-labeled), 5 µM Hs profilin-1 $\pm$ 0.1 nM FHOD3L-CT or 1 nM FHOD3L-CT K1193L. Average elongation rates (over 10s of seconds) and formin-mediated elongation rates (dim) are shown separately. (FHOD3L-CT K1193L (avg) n=19; FHOD3L-CT K1193L (dim) n=67; 3 flow channels each; mean $\pm$ SD, p-values by one-way ANOVA with post-hoc Tukey test.) (**D**) Quantification of bundling by FHOD3L-CT GS-FH1 and K1193L (n=3, each; mean $\pm$ SD.) (**E**) Nucleation test. Quantification of number of filaments per FOV for FHOD3L-CT and FHOD3L-CT GS-FH1. Conditions: 4 µM actin with indicated construct. Reaction was stabilized with Alexa488-phalloidin and diluted to 5 nM actin for visualization. 5 images (technical replicates) were taken per independent experiment. (n=15 images each; 3 biological replicates each; mean $\pm$ SD, p-values by one-way ANOVA with post-hoc Tukey test.) Representative images are shown in Figure 3-supplement 1D. (**F**) Barbed-end elongation assay for FHOD3L-CT GS-FH1 in the presence of profilin. Final conditions: 0.25 µM F-actin seeds (~0.1 nM barbed ends), 0.5 µM actin (10% pyrene-labeled), 1.5 µM *S. pombe* profilin, and 1.25 nM-20 nM FHOD3L-CT GS-FH1. (n=2; mean $\pm$ SD.) ns p > 0.05, *** p < 0.0001

Figure 3 Supplement 1

1181
Figure 3—figure supplement 1. (**A**) Quantification of run lengths for FHOD3L-CT and FHOD3L-CT
K1193L. (p>0.05 by Mann-Whitney U test.) (**B**) Duration of capping events by FHOD3L-CT and FHOD3L-
CT K1193L observed by TIRF microscopy. (p>0.05 by Mann-Whitney U test.) (**C**) TIRF micrographs from
nucleation assays using 4 µM actin in the presence or absence of 46 nM FHOD3L-CT GS-FH1, stabilized
with Alexa488-phalloiding and diluted to 5 nM. (**D**) TIRF micrographs from nucleation assays using 4 µM
actin and indicated concentration of FHOD3 construct. Samples were diluted to 5 nM actin in Alexa488
phalloidin for visualization. The number of aggregates was negligible at lower concentrations of FHOD3L-
CT GS-FH1. Quantification shown in Figure 3E.

Figure 4

Figure 4. FHOD3L rescues sarcomere organization in neonatal rat ventricular myocytes.
(**A**) Overview of the rescue experiments. Reverse transfection of siRNA upon plating NRVMs followed by infection with adenovirus to drive exogenous expression. (**B**) Western blot showing depletion of

endogenous FHOD3 after knockdown, and exogenous FHOD3L expression levels after rescue. GAPDH
used as a loading control. (**C**) Quantification of Western blot in (**B**) normalized to GAPDH for each lane
and then normalized to endogenous levels in the mock knockdown. (n=3, each; mean $\pm$ SD.) (**D**)
Sarcomere integrity indicated by immunofluorescent staining of α-actinin (green). Localization of
exogenous HA-FHOD3L is shown in magenta. DAPI (blue) is included in the merged images. WGA is not
shown for clarity. (**E**) Quantification of sarcomere number per NRVM. (n=160 cells, mock KD; n=334 cells,
mock rescue; n=110 cells, FHOD3L rescue.) (**F**) Average sarcomere lengths per NRVM. (mock KD n=100
cells; mock rescue n=71 cells; FHOD3L rescue n=92 cells). (**G**) Average sarcomere widths (Z-line
lengths) per NRVM. (mock KD n=100 cells; mock rescue n=71 cells; FHOD3L rescue n=92 cells) (**H**)
Epifluorescence micrographs showing mock knockdown and FHOD3L-rescued NRVMs stained with
phalloidin (green) to visualize thin filaments and anti-HA (magenta) to show expression of exogenous
FHOD3L. (**I**) Quantification of thin filament lengths for mock knockdown and FHOD3L-rescued NRVMs.
(mock KD n=94 cells; WT rescue n=84 cells) For (**E,F,G,I**) Data from three biological replicates for each
condition are represented by different shades. Mean $\pm$ SD is shown. p-values were calculated with Mann-
Whitney U tests, ns p > 0.05; * p < 0.05; ** p < 0.001.
1209
1210
1211
1212

Figure 4 - Supplement 1

1213
Figure 4—figure supplement 1. Method establishment for FHOD3L rescues in neonatal rat
ventricular myocytes. **(A)** Relative mRNA expression of *FHOD3* in NRVMs after knockdown with
different siRNA oligos compared to *GAPDH* mRNA expression (n=2 wells, 1 biological replicate; mean)**.**
(**B**) Typical image of cellpose segmentation of wild-type FHOD3L-rescued NRVMs, used for per cell
quantification of sarcomeres. ROI's are numbered. Scale bar, 20 µm (**C**) Pairwise correlation of
normalized DsRed vs. HA fluorescence intensity from FHOD3L rescue, plotted per NRVM area.
Correlation is too weak to use DsRed as an expression level reporter (n=347 cells; 1 biological replicate).
(D-F) Pairwise correlation of HA fluorescence intensity vs. (D) sarcomere number, (E) sarcomere length,
(F) sarcomere width from FHOD3L rescue, plotted per NRVM area. (**G**) Normalized and background
corrected 3xHA-FHOD3L fluorescence intensity per NRVM area from a rescue experiment. Magenta line
indicates the upper expression level cutoff (2700 a.u./µm^2) used to select cells for further analysis (n=913
cells; 3 biological replicates; mean $\pm$ SD.)

1226
1227
1228

Figure 5

1229
Figure 5. FHOD3L rescues contractility in neonatal rat ventricular myocytes. (**A-C**) Motion
analysis by digital image correlation. Mock knockdown, mock rescue, and FHOD3L rescue beating
patterns are shown. The first (•) and the second (▲) peak of each duplex represent the contraction and
the relaxation speed, respectively. (**D**) Maximal contraction velocities quantified for mock knockdown,
mock rescue, and FHOD3L wild-type rescue (mock KD n=82 ROIs; mock rescue n=81 ROIs; FHOD3L
rescue n=72 ROIs; 3 biological replicates, each; mean ± SD, p-values by Student's two-sample, unpaired
t-test.) (**E**) Quantification of the percentage of analyzed ROI's from the videos that contained rhythmic
contractions for mock knockdown, mock rescue, and FHOD3L wild-type rescue. (n=3, each; mean ± SD.)
(Note, we did not analyze % rhythmicity because the data rely on percentages. Thus only large changes
are noted.) ns p> 0.05, ** p < 0.001.

Figure 5 Supplement 1

1247
Figure 5—figure supplement 1. (**A**) Maximal relaxation velocities quantified for mock knockdown, mock
rescue, and FHOD3L rescue (mock KD n=82 ROIs; mock rescue n=81 ROIs; FHOD3L rescue n=72
ROIs; 3 biological replicates, each; mean ± SD, p-values by Student's two-sample, unpaired t-test.) (**B**)
Estimate of the percentage of NRVMs contracting in each video for mock knockdown, mock rescue, and
FHOD3L rescue. (n=3, each; mean ± SD.) (Note, we did not statistically analyze the % contracting data
because they are a subjective measurement. Thus, only large changes are noted.) p-values determined
with Student's two-sample, unpaired t-test. * p < 0.05, ** p < 0.001, *** p< 0.0001
1255
1256

Figure 5 Supplement 2

Figure 5—figure supplement 2. (**A**) Comparison of 3xHA-FHOD3L fluorescence intensity per NRVM area from a rescue experiment and an overexpression experiment. Data are normalized and background corrected as in Fig.4-supplemental figure 1G. (**B**) Maximal contraction velocities quantified for mock knockdown and FHOD3L overexpression NRVMs (mock KD n=82 ROIs; FHOD3L overexpression n=79 ROIs; 3 biological replicates, each; mean $\pm$ SD, p-value by Student's two-sample, unpaired t-test.) (**C**) Maximal relaxation velocities quantified for mock knockdown and FHOD3L overexpression NRVMs (mock KD n=82 ROIs; FHOD3L overexpression n=79 ROIs; 3 biological replicates, each; mean $\pm$ SD, p-value by student's two-sample, unpaired t-test.) (**D**) Quantification of rhythmic contractions as in Fig. 5E for mock knockdown and overexpression NRVMs (n=3, each; mean $\pm$ SD.) (**E**) Estimate of the percentage of NRVMs contracting in each video for mock knockdown and overexpression NRVMs (n=3, each; mean $\pm$ SD.) * $p < 0.05$, ** $p < 0.001$, *** $p < 0.0001$

Figure 6

1271
Figure 6. Loss of nucleation but not elongation is tolerated for sarcomere formation and function. (**A**) Images of NRVMs rescued with K1193L or GS-FH1. Sarcomere integrity indicated by immunofluorescent staining of α-actinin (green). Localization of exogenous FHOD3L is shown in magenta. DAPI (blue) is included in the merged images. WGA is not shown for clarity. (**B**) Quantification of sarcomere number per NRVM in the FHOD3L, K1193L, and GS-FH1 rescues. (K1193L n=148 cells; GS-FH1 n=259 cells; 3 biological replicates; mean ± SD, p-values by Mann-Whitney-U test.) (**C**) Average sarcomere lengths per NRVM in the FHOD3L, K1193L, and GS-FH1-rescues. (K1193L n=95 cells; GS-FH1 n=73 cells; 3 biological replicates; mean ± SD, p-value for FHOD3L comparison to GS-FH1 by Student's two-sample, unpaired t-test, all other p-values by Mann-Whitney-U test.) (**D**) Average sarcomere widths (Z-line lengths) per NRVM in the FHOD3L, K1193L, and GS-FH1-rescues. (K1193L n=95 cells; GS-FH1 n=73 cells; 3 biological replicates; mean ± SD, p-values by Mann-Whitney-U test.) (**E**) Quantification of thin filament lengths for FHOD3L and K1193L-rescued NRVMs (K1193L n=99 cells; 3 biological replicates, each; mean ± SD, p-value by Mann-Whitney-U test.) (**G**) Quantification of contracting NRVMs as in Fig. 5D for FHOD3L, K1193L, and GS-FH1-rescued NRVMs. (n=3, each; mean ± SD.) (**H**) Quantification of rhythmic contractions as in Fig. 5E for FHOD3L, K1193L, and GS-FH1-rescued NRVMs. (FHOD3L n=3; K1193L n=3; GS-FH1 n=2; mean ± SD.) ns p > 0.05, ** p < 0.001

Figure 6 - Supplement 1

1288
Figure 6—figure supplement 1. (A) Normalized and background-corrected fluorescence intensity per
NRVM area from the FHOD3L, K1193L, and GS-FH1 rescues. (K1193L n=897 cells; GS-FH1 n=536
cells; 3 biological replicates, each; mean ± SD.) (**B**) Epifluorescence micrographs showing K1193L-
rescued NRVMs stained with phalloidin (green) to visualize thin filaments and anti-HA (magenta) to show
expression of exogenous FHOD3L. (**C**) Maximal relaxation velocities for FHOD3L, K1193L, and GS-FH1-
rescued NRVMs. (K1193L n=81 ROIs; GS-FH1 n=31 ROIs; 3 biological replicates for WT and K1193L; 2
for GS-FH1; mean ± SD, p-values from Mann-Whitney U test.) (**D**) Maximal contraction velocities
quantified for FHOD3L, K1193L, and GS-FH1-rescued NRVMs. (K1193L n=81 ROIs; GS-FH1 n=31 ROIs;
3 biological replicates, WT and K1193L; 2 for GS-FH1; mean ± SD; p-values determined with Mann-
Whitney-U test.) (**E**) Striated HA-FHOD3L demonstrates that the GS-FH1 mutant can localize correctly if
sarcomeres are present. Sarcomere integrity indicated by immunofluorescent staining of α-actinin
(green). Localization of exogenous HA-FHOD3L is shown in magenta. DAPI (blue) is included in the
merged images. WGA is not shown for clarity. ns p > 0.05, ** p < 0.001
1302
1303
1304
1305

1306
1307

	Pyrene-actin-based assays		TIRF-based assays			Cosedimentation
	Nucleation Strength (a.u./s)	Barbed end binding affinity (nM)	Elongation Rate (subunits/s)	Run Length (µm)	Capping duration (s)	Bundling (% actin pelleted at 60 nM)
FHOD3L-CT	0.32 ± 0.04 (n=5)	0.028 ± 0.005	39 ± 13	1.10 ± 0.49	11.8 ± 6.8	81.7 ± 2.9
FHOD3S-CT	0.42 ± 0.04* (n=4)	0.750 ± 0.090*	33 ± 12	0.90 ± 0.33*	12.0 ± 7.8	56 ± 17
FHOD3L-CT I1163A	-0.081 ± 0.002* (n=3)	4.9 ± 1.4*	n.d.	n.d.	n.d.	n.d.
FHOD3L-CT K1309A	0.04 ± 0.02* (n=3)	n.d.	n.d.	n.d.	n.d.	n.d.
FHOD3L-CT K1193L	0.10 ± 0.02* (n=5)	0.470 ± 0.090*	38 ± 12	1.11 ± 0.42	12.7 ± 5.6	30.8 ± 0.7
FHOD3L-CT GS-FH1	Nucleates similarly via TIRF nucleation assay	0.218 ± 0.016*	n.d.	n.d.	n.d.	61.7 ± 0.7

Table 1: Summary of biochemical measurements for FHOD3L-CT and mutants. All data shown are
means ± standard deviation each from 3 independent experiments, unless otherwise indicated. n.d.= no
data. Light grey columns are data acquired with bulk pyrene-actin based assays without profilin added.
Medium grey columns provide data from TIRF analysis of individual filaments (profilin was present). The
dark grey column reports data from cosedimentation assay.
* Statistically different from FHOD3L-CT. Analysis by ANOVA and Tukey post hoc tests, $p < 0.05$. More
details are in the figure legends.
1315

	treatment		Fixed samples				Live samples			
	FHOD3 KD?	FHOD3 AdV?	Avg Sarcomere Number per NRVM	Sarcomere Length (µm)	Z-Line Length (µm)	Thin Filament Length (nm)	Maximal Contraction Velocity (µm/s)	Maximal Relaxation Velocity (µm/s)	Avg % Rhythmic Contractions	Avg % Contracting NRCs in FOV
Mock Knockdown	-	-	12 ± 13	1.71 ± 0.22	1.69 ± 0.43	925 ± 94	7.9 ± 1.6	6.0 ± 1.7	74 ± 9	98.2 ± 3.2
Mock Rescue	+	-	3 ± 7[a]	1.46 ± 0.37[a]	1.38 ± 0.46[a]	n.d.	6.0 ± 1.7[a]	4.4 ± 1.5[a]	14 ± 12	89 ± 11
FHOD3L Rescue	+	+	19 ± 14[a,b]	1.72 ± 0.18[b]	1.70 ± 0.46[b]	739 ± 81[a,b]	7.5 ± 1.5[b]	6.1 ± 1.5[b]	72 ± 20	97.2 ± 4.8
K1193L Rescue	+	+	15 ± 16	1.65 ± 0.22	1.67 ± 0.41	792 ± 79[c]	7.7 ± 1.6	5.9 ± 1.6	88 ± 21	98.2 ± 3.2
GS-FH1 Rescue	+	+	2 ± 4[c]	1.42 ± 0.35[c]	1.23 ± 0.29[c]	n.d.	5.6 ± 1.7[c]	3.6 ± 1.2[c]	3.7 ± 5.2	32 ± 32
FHOD3L Over-expression	-	+	n.d.	n.d.	n.d.	n.d.	6.6 ± 1.6[c]	5.5 ± 1.6[c]	72 ± 21	76 ± 32

Table 2: Summary of FHOD3L and mutant rescue experiments in NRVMs. All data shown are means ± standard deviation each from 3 independent experiments (except for GS-FH1 rescue: maximal contraction and relaxation velocity measurements and average % rhythmic contractions from 2 independent experiments). n.d.= no data. In the lightest columns, "-" indicates treatment with negative control siRNA or empty virus, "+" indicates treatment with FHOD3 siRNA or corresponding FHOD3L adenovirus. Data in the medium-colored columns were acquired from fixed and stained cells. Data in the darkest columns were acquired from live cells. Statistical analyses are described in figure legends.

  a- statistically different from mock KD (mock rescue and Fhod3L rescue)
  b- statistically different from mock rescue (FHOD3L and GS-FH1)
  c- statistically different from FHOD3L rescue (all mutant rescues and overexpression)