Light-guided actin polymerization drives directed motility in protocells

Hideaki T. Matsubayashi[1,2,3†], Shiva Razavi[1,2,4,5], Yuhei O. Tahara[6,7], Oghosa H. Akenuwa[8], T. Willow Rock[1,2], Daichi Nakajima[9], Rina Otsuka-Yamaguchi[3], Hideki Nakamura[1,2,10,11], Daniel A. Kramer[12,13], Tomoaki Matsuura[14], Baoyu Chen[12,13], Christopher T. Lee[8], Makoto Miyata[6,7], Satoshi Murata[9], Shin-ichiro M. Nomura[9], Takanari Inoue[1,2†]

1 Department of Cell Biology, School of Medicine, Johns Hopkins University, 2 Center for Cell Dynamics, Institute of Basic Biomedical Sciences, Johns Hopkins University, 3 Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, 4 Department of Biomedical Engineering, School of Medicine, Johns Hopkins University, 5 Department of Biological Engineering, School of Engineering, Massachusetts Institute of Technology, 6 Graduate School of Science, Osaka Metropolitan University, 7 The OMU Advanced Research Center for Natural Science and Technology, Osaka Metropolitan University, 8 Department of Molecular Biology, University of California San Diego, 9 Department of Robotics, Graduate School of Engineering, Tohoku University, 10 Hakubi Center for Advanced Research, Kyoto University, 11 Department of Synthetic Chemistry and Biological Chemistry, School of Engineering, Kyoto University, 12 Roy J. Carver Department of Biochemistry, Biophysics and Molecular Biology, Iowa State University, 13 Departments of Internal Medicine & Biophysics, University of Texas Southwestern Medical Center, 14 Earth-Life Science Institute, Institute of Science Tokyo

†Corresponding authors. Email: hideaki.matsubayashi.e1@tohoku.ac.jp, jctinoue@jhmi.edu

Abstract

Motility is a hallmark of life's dynamic processes, enabling cells to actively chase prey, repair wounds, and shape organs. Recreating these intricate behaviors using well-defined molecules remains a major challenge at the intersection of biology, physics, and molecular engineering. Although the polymerization force of the actin cytoskeleton is characterized as a primary driver of cell motility, building a minimal platform to test this process in protocellular systems has proven elusive. The difficulty lies in the daunting task of distilling key components from motile cells and integrating them into model membranes in a physiologically relevant manner. To address this, we developed a method to optically control actin polymerization with high spatiotemporal precision within cell-mimetic lipid vesicles known as giant unilamellar vesicles (GUVs). Within these active protocells, the reorganization of actin networks triggered outward membrane extensions as well as the unidirectional movement of GUVs at speeds of up to 0.43 µm/min, within the range of adherent mammalian cells. Notably, our findings reveal the requirements of both branched and linear actin networks for efficient membrane protrusions. This approach offers a powerful platform for unraveling the intricacies of cell migration, designing

39   synthetic cells with active morphodynamics, and advancing bioengineering applications, such as
40   self-propelled delivery systems and autonomous tissue-like materials.

Introduction

Motility is a fundamental property of cells across biological systems, from bacteria to large animals ([1]–[6]). By transcending the randomness and viscosity of the microscopic world, cells move toward their destination by extending membrane protrusions and by clutching onto the extracellular environment. In amoeboid cell migration, these processes are universally galvanized by actin polymerization ([7]–[12]). While hundreds of biomolecules are identified to be required for cell motility ([4], [6], [13]), a minimal list of biomolecular components sufficient for cell motility remains elusive. This primarily reflects the inherent difficulty of eliminating all biomolecules beyond the minimal set in the context of living cells.

Cell migration is driven by the coordination of forward movement, rear contraction, and adhesion/deadhesion of the cell edges ([4], [6], [14]–[16]). Among these processes, actin polymerization at the leading edge is a strong candidate for driving overall forward displacement. For instance, *Dictyostelium*, fish keratocyte, and human neutrophil retain their motility even after myosin contractility is compromised genetically ([17], [18]) or pharmacologically ([19], [20]). Cells also manage to migrate even without strong adhesions especially in confined conditions ([21], [22]). Conversely, we and others have shown that synthetic activation of the Rho family GTPase Rac1, an upstream activator of the actin cytoskeleton, induces directional cell movements toward the site of stimulation ([23], [24]). However, it is still unclear if Rac1-mediated actin polymerization alone is sufficient for cell motility in cellular settings, as Rac1 crosstalks with other Rho small GTPases such as RhoA and triggers other cytoskeletal activities including actomyosin contraction ([25]).

In a different context, molecules necessary for force generation by actin polymerization have been identified based on the motility reconstitution assay ([26]–[36]). These reports demonstrated that a mixture of purified cytoskeletal proteins, such as actin, Arp2/3, profilin, cofilin, and capping protein, can reconstitute the motility of *Listeria monocytogenes* bacteria using microspheres coated with actin nucleation-promoting factors (NPFs). Furthermore, a number of studies were conducted to reconstitute actin ([37], [38]) and other cytoskeletons ([39]–[41]) inside cell-sized membrane-bound liposomes. These studies demonstrated that actin polymerization generates membrane deformations ([42]–[45]), revealed the roles of actin-binding proteins ([46]–[52]), and further showed that internal reactions can be regulated by light and external environmental factors ([52]–[55]). However, unidirectional motion by the cytoskeletal polymerization force has remained challenging. Moreover, non-invasive and non-constitutive—thus more physiological—induction of actin polymerization has yet to be achieved, particularly within cell-sized compartments that maintain proper membrane configuration. Building on these findings, we developed a minimal membrane-bound system in which actin polymerization can be locally

77 controlled and tested whether such an artificial cytoskeletal activity is sufficient to induce cell-
78 like motility.
79

Results

Reversible and asymmetric light control of protein localization inside GUVs

To achieve *in vitro* reconstitution of cell motility, we previously developed active protocells where
chemically inducible dimerization (CID) tools were used to polymerize actin based on a bacterial
NPF inside giant unilamellar vesicles (GUVs) (*56*). While the previous system efficiently produced
actin polymerization, outward membrane extension necessary for driving movement was not
observed. We reasoned that this was due to a lack of fast turnover of NPF recruitment to the
membranes. To aim for a physiologically relevant generation of protrusive force exerted against
membranes, we adopted a light-inducible protein dimerization tool for controlling the
localization of proteins of interest. Among optogenetic dimerization tools (*57–59*), we chose iLID-
SspB (*60*) as its second-scale reversible binding kinetics match the turnover of leading-edge
polarization (Supplementary Text 1), its blue-light activation is compatible with standard
microscope setups, and its relatively small molecular size facilitates expression and purification
of recombinant proteins.
95

We first set out to test light-induced dimerization of purified iLID and SspB in GUVs. In our
construct, iLID was membrane-tethered, and blue light exposure triggered the Jα helix to extend,
revealing the binding site for SspB. In the dark, iLID returned to its closed state, releasing SspB
(Fig. 1A). We visualized the localization and the light responsiveness of the proteins by fusing iLID
with YFP and SspB with mCherry (fig. S1A). To anchor iLID to the GUV membrane, we tested four
different protein-lipid interaction strategies to mediate light-induced translocation, including
coordination bonding, electrostatic, strong non-covalent, and covalent interactions (Fig. 1A, fig.
S2, Supplementary Text 2) (*61*, *62*). Among these, the SNAP-tag/benzylguanine (BG)-conjugated
lipid pair provided efficient iLID anchoring with minimal nonspecific membrane binding of SspB,
and was therefore used for subsequent experiments.
106

When GUVs were made with 2% BG-conjugated lipid in the inner leaflet, iLID localized on the
membrane, while SspB remained in the lumen in the dark (Fig. 1B). Upon blue light illumination
of the entire GUV, SspB translocated to the membrane within 4 seconds (Fig. 1B, C). After
switching back to the dark, SspB gradually dissociated from the membrane with a half-time of
61.2 ± 13.7 seconds. The time scale is consistent with AsLOV2-based optogenetic tools measured
in cells (*58*). These data confirmed that iLID-SspB enables reversible control of protein localization
in GUVs.

Next, we tested the potential for spatially asymmetric control of iLID-SspB interactions. When
light was applied locally to one side of the GUV, SspB accumulated at the illuminated site (Fig. 1D,
E). In the dark, dissociation proceeded with a half-time of 41.8 ± 7.1 seconds (fig. S3A). Despite
faster diffusion on artificial lipid membranes compared to cell membranes (*63–65*), localized light
stimulus created an asymmetric pattern of SspB in cell-sized vesicles (Fig. 1E, F, see
Supplementary Text 1). Furthermore, SspB precisely and repeatedly responded to the directional
change of blue light illumination (Figs. 1G, S3B, C, and Movie S1). Collectively, these results
demonstrated that the iLID-SspB system can direct asymmetric protein distribution at regions of
interest on the GUV membranes.

Light-guided control of ActA
Next, we sought to manipulate the actin cytoskeleton with the iLID-SspB system. Previous studies
have established that the rate of actin polymerization depends on the surface density of NPFs
(*26, 30, 35*) and that clustering of NPFs accelerates actin polymerization (*66, 67*), as further
demonstrated by a reconstituted optogenetic VCA system (*68*). We have recently shown that
chemically-induced membrane recruitment of ActA, an N-WASP homolog NPF derived from *L.*
monocytogenes, induces actin polymerization on the membrane of GUVs, leading to symmetry
breaking (*56, 69*). Thus, we first tested ActA with the iLID-SspB system. To identify an optimal
optogenetic configuration, we tested four ActA-SspB constructs with different SspB variants and
fusion orders (Supplementary Text 3). These constructs were evaluated based on actin
polymerization activity and light-dependent membrane recruitment, leading us to select ActA-
SspB$_{nano}$-mCherry (fig. S4). When supplemented with actin and Arp2/3, light-induced global
membrane recruitment of ActA-SspB led to the emergence of actin patches on the membrane,
consistent with our previous findings with a chemical input (*56*) (fig. S5). With local light, the ActA
distribution became asymmetric, albeit the directionality of actin pattern could not be
maintained. In most cases, actin patches diffused randomly out of the illuminated region and the
GUV lost actin directionality over time (fig. S6 and Movie S2).

Reversible control of pVCA and light-guided actin polymerization
For robust maintenance of actin asymmetry, we aimed to rapidly depolymerize actin filaments in
the non-illuminated area while strengthening actin polymerization in the illuminated region. To
promote depolymerization and turnover, we incorporated factors that facilitate the actin
turnover process, namely profilin, cofilin, and capping protein (*13, 33, 70, 71*). In
ultracentrifugation assays, the combined addition of profilin, cofilin, and capping protein
increased the pool of G-actin, supporting the notion that these factors enhance actin turnover
(fig. S8A, B). To enhance polymerization in the illuminated regions within GUVs, we optimized
the polymerization module. Among several NPF constructs evaluated, GST-tagged N-WASP pVCA

(proline-rich region + VCA domain) showed the fastest actin polymerization in a bulk pyrene assay,
which monitors actin polymerization through the fluorescence increase of pyrene-labeled actin.
Notably, GST-pVCA maintained its activity when fused to SspB-mCherry, making GST-pVCA-SspB-
mCherry (pVCA-SspB-mCherry) a suitable light-inducible activator for our GUV system (Fig. 2A,
S1, S8C, D).
157
We then encapsulated this light-inducible pVCA-SspB-mCherry with actin (10% Alexa-647
labeled), Arp2/3, profilin, cofilin, and capping protein in GUVs (Fig. 2B). Upon illumination, pVCA-
SspB-mCherry rapidly translocated to the membrane, followed by the increase in membrane
actin signal. Notably, the actin signal increased gradually over time with a delay of several
minutes after pVCA translocation, consistent with light-induced actin polymerization (Fig. 2C, D,
Movie S3). In contrast, in the absence of either pVCA or light stimulation, no significant change
in membrane actin intensity was observed (Fig. 2D, fig. S9). When Arp2/3 was omitted, the
membrane actin signal showed a slight increase over time. This suggested that the G-actin
binding property of pVCA (*72*, *73*) increased the local concentration of actin at the periphery of
the membrane, albeit this effect was not as intense as Arp2/3-dependent actin assembly (Fig. 2D,
fig. S9). These data showed that light-dependent membrane accumulation of pVCA drives
Arp2/3-dependent actin enrichment on the GUV membrane.
170
Next, we employed pharmacological perturbations to probe the dynamics of membrane-
associated actin. Specifically, Latrunculin A sequesters G-actin to inhibit polymerization, whereas
Jasplakinolide stabilizes F-actin and suppresses depolymerization. To decouple actin dynamics
from optogenetic kinetics, iLID-SNAP was replaced with SsrA-SNAP, in which SsrA constitutively
binds SspB-fused pVCA independently of light (fig. S10A). Under these conditions, actin was
readily observed on the membrane (Fig. 2E, fig. S10B, C). Upon Latrunculin A treatment, the
membrane actin signal markedly decreased, confirming that the membrane actin signal derives
from F-actin. Conversely, Jasplakinolide increased membrane actin fluorescence, revealing
additional capacity for membrane actin accumulation beyond the steady-state level observed
under DMSO conditions (Fig. 2E, F, fig. S10D–F). Together, these results suggest that the
membrane-associated actin is maintained by a balance between ongoing actin assembly and
disassembly.
183
To further characterize the dynamics of pVCA and actin on the GUV membrane, we performed
fluorescence recovery after photobleaching (FRAP) analysis. pVCA-SspB-mCherry exhibited rapid
and near-complete fluorescence recovery, indicating that membrane-associated pVCA remains
highly mobile even during actin enrichment (Fig. 2G, fig. S11). In contrast, FRAP analysis of Alexa
Fluor 647–labeled actin showed slower and partial fluorescence recovery, suggesting the
presence of a less mobile F-actin fraction at the membrane (Fig. 2G, fig. S11). Notably, actin

fluorescence recovery was significantly reduced in the absence of regulatory factors, profilin,

cofilin, and capping protein (fig. S11A, E), indicating that membrane-associated actin exists in a

dynamic steady state regulated by these factors.

To examine the ultrastructure of the actin network, we performed negative staining electron

microscopy (EM). For this analysis, small unilamellar vesicles (SUVs; 50–150 nm in diameter)

containing 2% benzylguanine-modified lipids (fig. S12A) were used to recruit pVCA-SspB-mCherry

to their outer surface via SsrA-SNAP. EM images revealed dense actin filament networks (Fig. 2H,

fig. S12B–D) with frequent Y-junctions, characteristic of Arp2/3-mediated branching (fig. S12B),

providing structural evidence for Arp2/3-dependent actin polymerization.

Next, we tested the reversible actin regulation by pVCA-SspB-mCherry. As shown in Fig. 2I and J,

both pVCA-SspB-mCherry and actin displayed reversible and repetitive changes in membrane

intensity in response to blue light switching (Movie S4). Notably, once switched to the dark,

pVCA-SspB-mCherry quickly dissociated from the membrane, whereas actin decreased its

membrane intensity at a relatively slower rate (fig. S13). Taken together, these results suggested

that the polymerization state is reversible in our system and that the balance between

polymerization and depolymerization can be controlled by light.

Asymmetric and spatially controlled actin polymerization

The asymmetric distribution of NPFs and actin polymerization are fundamental features of

directed cell migration (*9*, *74*, *75*). Thus, we next tested whether locally targeted pVCA could

establish asymmetric actin polymerization (Fig. 3A). As shown in Fig. 3B, C, and fig. S14A, pVCA

showed asymmetric patterning toward the localized light stimulus (Movie S5). About 5 minutes

after pVCA translocation, the actin signal also increased from the illuminated region and created

an asymmetric pattern. Line scans along the GUV perimeter indicated that pVCA responded

rapidly with a narrow peak, whereas actin accumulated more gradually and exhibited a broader

distribution. These results suggest that actin polymerization requires a finite time to initiate and

propagate, and once filaments are formed they remain associated with the membrane for a

longer duration (fig. S14B). With continuous local illumination, both pVCA and actin maintained

stable polarity for more than 100 minutes (Fig. 3B, C, S14A, B, Movie S5). Notably, at later time

points, the actin signal appeared as two lines in the latter time points in the kymograph. This

likely reflects reduced lateral diffusion of membrane-associated actin filaments, combined with

gradual photobleaching of Alexa Fluor 647-labeled actin under repeated blue-light excitation. We

next tested whether the direction of actin polymerization could be reversibly reprogrammed.

After establishing an asymmetric actin pattern with a local light on the right side of the vesicle,

we reversed it back to the dark mode and subsequently applied new light on the left. Both pVCA

and actin rapidly re-polarized toward the new stimulation site (Fig. 3D, E, and Movie S6). Together,

these results demonstrate that the light-inducible pVCA system enables reversible and spatially
reconfigurable polarization of actin networks.

Importantly, when locally illuminated, the front membrane of some GUVs moved toward the
illuminated direction (fig. S14C), albeit occasionally (2 out of 7 instances). We speculate that
characteristics inherent to the GUV system–such as surface-to-volume ratio of individual GUVs,
surface density of pVCA, and the ratio of Arp2/3 to capping protein (*33*, *76*), either individually
or in combination, may determine whether productive protrusive behavior emerges.

mDia1-driven actin polymerization drives membrane protrusions in cells
To enhance forward membrane protrusion, we next sought to reinforce actin polymerization. At
the leading edge of migrating cells, two major actin nucleators, namely Arp2/3 and formin,
coordinate the forward extension of the plasma membrane (*77–79*). While Arp2/3 stays at the
branch points of the actin filaments and does not engage in the incorporation of monomeric actin
into the growing ends, formin continuously binds to the growing ends and actively recruits
profilin-actin complexes to the elongating tips. Consequently, one subtype of formin, mDia1,
accelerates actin polymerization 4–5-fold (*80*, *81*) and drives motility 6–23 times faster than
Arp2/3 in *in vitro* motility assays using functionalized beads (*30*, *28*, *31*). Furthermore, a
theoretical study suggests that actin polymerization perpendicular to the membrane,
characteristic of formin-driven filopodia formation, more efficiently transduces the
polymerization force to advance the membrane (*82*). Therefore, we reasoned that the
integration of mDia1 could enhance protrusive activity.

To test whether mDia1 promotes membrane protrusion, we first used living cells and performed
a plasma membrane recruitment assay (Fig. S15A) (*83–85*). More specifically, we took FH1-FH2-
DAD domains from mDia1 as a constitutively active form (*86*) and recruited it to the plasma
membrane by the CID system. As shown in fig. S15B–H, YF-mDia1 translocation induced filopodia-
like protrusions (fig. S15B–H). These protrusions were positive for Lifeact signal and showed
coiling motions with their elongating tips labeled with YF-mDia1 (fig. S15D), consistent with
formin-dependent filopodia elongation (*87–90*). Together, these data suggest that membrane
recruitment of mDia1 generates an actin-dependent protrusive force on the membrane.

mDia1-driven actin polymerization drives forward movement of GUVs
We then constructed the light-inducible version of mDia1 (mCherry-SspB-mDia1) for *in vitro*
reconstitution (Fig. S1). Pyrene actin assays suggested that mDia1 had an effect of increasing the
initial rate of actin polymerization (Fig. S8E, F). We encapsulated mCherry-SspB-mDia1 with iLID-
YFP-SNAP, actin (10% Alexa-647 labeled), profilin, and cofilin in GUVs. Upon local recruitment of
mDia1, some vesicles showed gradual drift toward the stimulated direction, suggesting that

mDia1-generated actin polymerization can produce protrusive forces in the reconstituted lipid
vesicles (fig. S16A, B). However, the response was modest and difficult to detect.

Because mDia1 alone produced only limited protrusive activity, we added pVCA-SspB, Arp2/3,
and capping protein to better mimic actin network organization of migrating cells (Fig. 4A).
Strikingly, this full component system (pVCA + mDia1) produced robust directional movement
toward light stimulation in 18 out of 22 GUVs (Fig. 4B–D, fig. S16C, Movie S7, S9, S10). Polarized
actin accumulation at the illuminated side preceded membrane displacement, consistent with
actin polymerization driving forward protrusion (Fig. 4B, Movie S7). The vesicle continued to
move steadily for an hour until the light was turned off (Fig. 4B, D, Movie S7). Furthermore, when
we shifted the position of the blue light, the signals for mDia1, pVCA, and actin reoriented,
causing the vesicle to move in the new direction, resembling chemotactic reorientation in
migrating neutrophils (Fig. 4B, D, Movie S7). The speed of pVCA/mDia1 dual system was
approximately threefold higher than that of mDia1 GUVs (Fig. 4G, H, S16B–C). The distance the
front side membrane moved in 30 minutes was also significantly longer in pVCA + mDia1 GUVs
(Fig. 4I). Furthermore, membrane recruitment of mDia1 and pVCA occasionally showed local
protrusions, suggesting the generation of protrusive force (fig. S17, Movie S8). Together, these
results indicate that mDia1-driven actin polymerization generates protrusive force that is
substantially enhanced by cooperative activity with pVCA and Arp2/3.

When mDia1 and pVCA were replaced with an equivalent amount of mCherry-SspB, vesicles did
not move even with the local blue light illumination (fig. S18A). In addition, the kymograph
analysis revealed that the timing and direction of membrane protrusions correlated with actin
intensity and its polarity, consistent with actin-driven dynamics (Fig. 4E, F). Notably, the
kymographs also revealed that protrusions at the front often coincide with retractions at the rear.
Corroborating this, the actin sequestering compound Latrunculin A and the formin inhibitor
SMIFH2 substantially compromised the motion of vesicles (Fig. 4J). Furthermore, deletion of the
DAD domain from mDia1, which is responsible for efficient actin nucleation by mDia1 (*91*),
abolished this motion (Fig. 4J). Together, these results indicate that membrane protrusion and
motion of the vesicles are driven by actin polymerization.

The pVCA–mDia1 combination forms a fast-turnover, actively polymerizing actin network
To elucidate the mechanistic basis of the distinct actin architectures generated by pVCA, mDia1,
and the pVCA–mDia1 dual system and their contributions to vesicle motion, we combined
ultrastructural, pharmacological, and FRAP analyses of reconstituted actin networks.

To examine ultrastructure, we performed EM on actin assemblies formed on SsrA-coated SUVs.
Although assemblies deposited onto a solid support may not fully represent filament

organization within GUVs, micrometer-scale filaments were observed under all conditions, with
the highest filament density observed when pVCA was used as the sole NPF (fig. S12D, E).
However, filament density alone did not explain the observed differences in vesicle motion,
underscoring the importance of actin network dynamics.
308
To probe network dynamics, we examined the pharmacological responses of actin assemblies in
a system where NPFs were constitutively tethered to the GUV membrane via SsrA. Consistent
with EM observations, pVCA generated the highest basal level of membrane-associated actin but
showed only modest additional F-actin accumulation upon Jasplakinolide treatment (fig. S10B–
F). In contrast, mDia1-driven assemblies exhibited a progressive increase in membrane-
associated actin in response to Jasplakinolide, with the most pronounced effect observed in the
combined pVCA-mDia1 system (fig. S10E, F), consistent with the enhanced polymerization
capacity of the dual condition observed in bulk pyrene–actin assays (fig. S8F). Notably,
Jasplakinolide treatment of mDia1 and pVCA-mDia1 GUVs was occasionally accompanied by
outward membrane deformation, consistent with polymerization-generated forces acting on the
membrane (fig. S10G). Moreover, the pVCA-mDia1 dual system was least susceptible to
Latrunculin A, indicating a shift in network behavior toward sustained polymerization (fig. S10E,
F).
322
FRAP analysis further distinguished network dynamics. pVCA-driven actin networks displayed
rapid fluorescence recovery, suggesting highly dynamic actin turnover (fig. S11C, E). In contrast,
actin assemblies generated by mDia1 showed minimal fluorescence recovery of actin and mDia1
itself (fig. S11B–E). This slow recovery is consistent with processive filament elongation at the
membrane, where mDia1 remains persistently associated with growing barbed ends. Importantly,
the pVCA–mDia1 dual system—despite exhibiting strong Jasplakinolide-induced actin
accumulation—retained rapid fluorescence recovery in FRAP (fig. S11C, E), indicating that the
dual network maintains dynamic turnover while sustaining high polymerization activity.
331
332 Together, these observations indicate that the pVCA–mDia1 system exhibits high polymerization
333 capacity and rapid turnover, properties potentially well suited for efficient force generation and
334 directional vesicle movement.
335
Deformation and adhesion of GUVs
Quantitative analysis revealed that the initial velocity of each GUV was strongly correlated with
its net displacement after 30 minutes ($R = 0.89$; fig. S19A). In addition, displacement showed a
moderate positive correlation with vesicle diameter, suggesting that larger vesicles can
accommodate greater chemical and cytoskeletal resources and may also be more favorable for
generating light-induced asymmetry. In contrast, vesicles showing membrane localization of

pVCA and mDia1 prior to blue light illumination tended to exhibit smaller displacement. This
suggests that nonspecific membrane binding limits the light-dependent establishment of
asymmetry (fig. S19B, C).

We next analyzed changes in velocity and vesicle shape. The vesicles initially moved at an average
velocity of 0.43 μm/min, but their speed gradually decreased over time (Fig. 4G, H, fig. S16C).
Concomitant with this deceleration, the contact area of the bottom side of the vesicles
progressively increased during the movement, stabilizing the vesicles in a dome-like shape (Fig.
4C, K). This increase in the contact area was not seen in the GUVs when pVCA and mDia1 were
replaced with mCherry-SspB (fig. S18B). Thus, we speculate that the contact to the bottom
substrate was induced by the protrusive reaction and that the adhesion and the concomitant
deformation led to cumulative tension increase (*92*) which in turn counteracted the protrusive
force. The force exerted by actin polymerization may not be enough to detach the GUV from the
BSA-coated glass substrate. This is consistent with the observation that an initial contact site
remained adherent even in a GUV that traveled farthest (Fig. 4K, fig. S20B, C, Movie S9, S10).
Additionally, actin recycling in the system may be insufficient and the force gradually diminishes
over time (*71*). Despite these potential resisting factors, the combination of mDia1 and pVCA
induced movement consistently (Fig. 4, fig. S16C, S20). Together, these observations support the
conclusion that asymmetric actin polymerization mediated by pVCA–mDia1 is sufficient to drive
forward vesicle movement under the tested conditions.

Computational modeling of actin network dynamics underlying membrane deformation
To dissect the mechanisms underlying actin-driven membrane deformation, we constructed a
computational model of actin network dynamics in a minimal vesicle system. We hypothesized
that membrane deformation could arise from intrinsic mechanical bracing within the actin
network. Accordingly, we considered two candidate mechanisms: Arp2/3-dependent network
organization and frictional interactions between actin filaments (*93*, *94*). The model was
implemented using the actin-network simulation framework Cytosim (*95*) and coupled to a
deformable membrane (*96*, *97*) (Supplementary Text 4 and Methods).

The simulations captured distinct NPF-dependent behaviors (fig. S21A, B, Movie S11, S12). pVCA
produced short, branched filaments that frequently detached from the membrane, whereas
mDia1 drove rapid filament elongation while remaining membrane-associated. Without inter-
filament steric interactions, the pVCA + mDia1 condition yielded marginal but reproducible
forward membrane motion (fig. S21A, C, Movie S11), indicating cooperativity between mDia1
and pVCA. Incorporating inter-filament interactions resulted in robust membrane deformation
(fig. S21B, D, Movie S12), with extension increasing in the order pVCA < mDia1 < pVCA+mDia1,
consistent with experiments (Fig. 4G–I). These results support a minimal internal bracing

mechanism in which membrane-anchored polymerization, Arp2/3-dependent branching, and
inter-filament interactions collectively generate protrusive force.

The simulations also revealed substantial retrograde flow of actin filaments (fig. S21A, B, Movie
S11, S12). This provides a mechanistic explanation for why the observed vesicle motility (0.1–1.0
µm/min, Fig. 4G–J, fig. S16) is more than an order of magnitude slower than the actin
polymerization rate (1–10 µm/min) (*80*, *81*) and neutrophil migration (~10 µm/min) (*98*).
Incorporating additional backing mechanisms that provide mechanical resistance to the actin
network, such as external anchoring or confinement (*21*, *22*), is therefore predicted to enhance
protrusive efficiency and bridge minimal synthetic systems toward cellular modes of amoeboid
migration.

392   Discussion

In the present study, we developed a fully reconstituted system to achieve light-guided
directional movement in a protocell. We leveraged mDia1-mediated actin polymerization to
establish the forces that drive movement. Our findings not only corroborate recent cell biological
findings on formin's significant roles in cell migration (*87*, *98–103*) but also show that the mDia1-
mediated actin polymerization can drive the forward movement of lipid vesicles in a minimal,
reconstituted system, overcoming challenges thus far posed by genetic redundancy and cellular
complexity. Formin-mediated actin polymerization exhibits several unique features compared to
free barbed end elongation in Arp2/3-dependent polymerization, including processive
elongation (*31*, *80*), accelerated polymerization rate (*80*, *81*), rotation, and coiling motion (*89*, *90*,
104). These characteristics could contribute to the effective propulsion of the vesicle. Remarkably,
our minimal system further indicated a functional interplay between pVCA and mDia1. Although
further investigation is required, this cooperative effect likely arises from coordination between
the two nucleators: mDia1 generates mother filaments required for Arp2/3 nucleation, and, in
turn, Arp2/3-mediated branching increases the number of growing ends available for mDia1
elongation (*100*, *101*). To further elucidate the mechanical coupling to force generation and
membrane protrusion, unveiling the nanoscopic architecture of the actin network in the
migrating protocell will be crucial. Our protocell approach, which bridges the gap between
cellular and biochemical domains, opens new avenues for comprehensively interrogating
modules of cellular motility.

We achieved a GUV displacement speed in the range of 0.1–1.0 µm/min, which is comparable to
the migration rates of typical fibroblasts (*105*) and other adherent cells including those that
competed in the World Cell Race (*106*). How could we make the movement even faster? *L.*
monocytogenes and NPF-coated beads in purified cytoskeletal protein solution typically move in

the range of 1–15 µm/min (*27*, *28*, *30–33*). Some adherent cells can migrate at 5 µm/min even

when actomyosin contractility is inhibited (*19*). We propose three possible ways to make artificial

cell migration faster. First, increasing actin concentration (*81*, *107*) and integrating additional

factors, such as VASP (*108*), Fascin (*109*), and NPFs (*67*, *110*), could accelerate the rate of actin

polymerization. Secondly, optimizing membrane tension could enhance the deformability of the

lipid membrane, akin to the malleable structures observed in motile cells, thereby potentially

promoting more efficient protrusion and motility (*50*, *111*). Finally, fine-tuning adhesion could

enhance migration efficiency as many cells exhibit a preference for an optimal adhesion range,

where both insufficient and excessive adhesion can hinder motility (*112*, *113*).

Interestingly, the phenotypes of mDia1 membrane recruitment differed between cells and GUVs.

In cells, mDia1 produced filopodia-like thin protrusions (fig. S12), whereas in GUVs, it pushed the

lipid membrane as a continuous plane structure on a much broader scale than filopodia (Fig. 4,

fig. S13–16). Since living cells tend to have higher membrane tension than liposomes (*114*), it

may appear counterintuitive that the cell membrane was more susceptible to mDia1-induced

local deformation. One possible explanation is that, in cells, endogenous proteins might work in

concert with mDia1 to amplify local membrane deformation and form long filopodia-like

protrusions (*100*, *115*). Alternatively, localized defects in membrane tension could predispose

certain areas to form thin protrusions (*116*). Additionally, cytoskeleton-bound transmembrane

proteins impede tension propagation in cells (*117*). Furthermore, the nuclear accumulation of

mDia1 in cells (fig. S12) could reduce its effective concentration and cause a heterogeneous

distribution on the cell membrane, resulting in focused protrusive force in regions with high

mDia1 concentration, similar to how pVCA-driven finger-like protrusions are formed in liposomes,

as demonstrated by Gat *et al.* (*118*). Since our light-inducible actuators are genetically encodable,

the same molecular toolkit can be deployed in both GUVs and living cells—providing a unified

platform to dissect cellular morphodynamics.

We have recently shown that chemically-induced membrane recruitment of ActA, an NPF derived

from *L. monocytogenes*, and subsequent global actin polymerization lead to eccentricity in the

shape of GUVs (*56*). We noted a difference in membrane deformation between the CID and LID

systems, as the CID could induce a clear membrane deformation given a global, non-directional

input. This difference could be attributed to the variations in cytoskeletal proteins and input

signals between the two systems. In the present study, we have incorporated profilin, cofilin, and

capping proteins. Since these proteins accelerate actin depolymerization (fig. S8B), they may

render the actin network more plastic rather than exerting force to deform the membrane.

Consistent with this, stabilization of F-actin by Jasplakinolide occasionally resulted in outward

membrane deformation in GUVs containing mDia1 or the pVCA–mDia1 combination (fig. S10G).

Moreover, the binding kinetics of iLID-SspB (k_{on} =1.2 × 10^3 M^{-1}s^{-1}, k_{off} =1.1 × 10^{-3} s^{-1}, K_d = 0.8 µM)

substantially differ from those of FKBP-FRB (k_{on} =1.9 × 10^6 M^{-1}s^{-1}, k_{off} =2.2 × 10^{-2} s^{-1}, K_d = 12 nM)

(*60*, *119*, *120*). This may result in different reaction dynamics between actin cytoskeleton and the

lipid membrane.

Our finding that internal cytoskeletal force can drive the unidirectional motion of lipid vesicles

offers avenues to reverse-engineer cell motility. Although we used light as an input, asymmetric

actin polymerization may be achievable with chemical inputs with consideration of binding

affinity (*83*, *121*), transmembrane signal transduction mechanisms (*122*), and self-organized

pattern formation (*123–127*). For sustained motility, it will be important to incorporate other

migratory forces—reconstructing myosin contractility at the rear (*128*) and regulating

adhesion/de-adhesion turnover (*119*, *129*) in concert with actin polymerization. Furthermore,

integrating synthetically engineered molecules (*81*, *130*, *131*) could enhance motility even

beyond the limits of natural systems. Achieving synthetic motility in GUVs could be a step toward

future developments in active therapeutic agents, self-organizing artificial tissues, and synthetic

neural circuits.

Materials and Methods

Reagents

ATP was purchased from Gold Biology (A-081) or Nacalai Tesque (01072-82), prepared as 100 mM solution in 100 mM Tris-HCl (pH 7.4 at RT), and stored at -20°C. Creatine phosphate was purchased from Gold Biotechnology (C-323-5) or Oriental Yeast (45180000), prepared as 1 M solution in PBS (Gibco 10010-023), stored at -20°C. Creatine kinase was purchased from Sigma-Aldrich (10127566001) or Oriental Yeast (46430003), prepared as 40 µM solution in buffer V (50 mM HEPES-NaOH (pH 7.5), 150 mM NaCl, 7 mM β-mercaptoethanol, 10% glycerol), snap-frozen, and stored at -20°C. Flavin mononucleotide (FMN) was purchased from TCI (R0023) and Wako (06500171), freshly prepared as 600 µM solution in Milli-Q water before iLID expression. Alexa Fluor 647 C2 Maleimide (A20347) and Pierce™ Glutathione Agarose (16100) were purchased from Thermo Fisher Scientific. Hexadecane (H6703), Silicone oil (378348), and SMIFH2 (S4826) were purchased from Sigma-Aldrich. Latrunculin A was purchased from Wako (125-04363). Jasplakinolide was purchased from Cayman Chemical Company (11705). Benzylguanine-PEG2000-DSPE was prepared and purified as previously described (*61*). POPC (1-palmitoyl-2-oleoyl-glycero-3-phosphocholine, 850457), POPS (1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine, 840034), 18:1 DGS-NTA-Ni (1,2-dioleoyl-sn-glycero-3-[(N-(5-amino-1-carboxypentyl)iminodiacetic acid)succinyl] (nickel salt), 790404), 18:1 PE-PEG2000-benzylguanine (1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-[benzylguanine(polyethylene glycol)-2000], 880137), and 18:1 Biotinyl Cap PE (1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-(cap biotinyl) (sodium salt), 870273C) were purchased from Avanti Polar Lipids as chloroform solution. Actin (AKL99), pyrene labeled actin (AP05), Arp2/3 (RP01P), Profilin (PR02), Cofilin (CF01), were purchased from Cytoskeleton. Strep-Tactin XT (2-4010-010, discontinued) resin, Strep-Tactin XT 4Flow (2-5010-010) resin, and purified Strep-Tactin protein were purchased from IBA. Ni-NTA agarose (30210) was purchased from Qiagen. Amylose resin (E8021S) was purchased from New England BioLabs.

Plasmid construction

pQE80L-6×His-MBP-TEVprorease(S219V)

Ser219Val mutation (*132*) was introduced by PCR with the template plasmid gifted from Dr. Yubin Zhou and the primer set of fwd: 5'-TGGTGAAACCTGAAGAACCTTTT-3' and rev: 5'-TTCACCATGAAAACTTTATGGCC-3'. TEV prorease (S219V) was then PCR amplified with the primer set of fwd: 5'-AATTGAGCTCGATGAGCGGCCTGGTGC-3' and rev: 5'-AATTGGATCCTTATTGCGAGTACACCAATTCATTCATG-3' and inserted between SalI and BamHI sites of pQE-80L MBP-SspB Nano plasmid (Addgene #60409) by using restriction digestion and T4 ligase ligation process.

pQE80L-6×His-MBP-TEVsite-mCherry-SspB(micro)

EYFP-2×SAGG linker was PCR amplified from the template plasmid YFP-FKBP (*133*)with the primer set fwd: 5'- GTATTTTCAGGGATCGCCGCTAGCGCTACCGG-3' and rev: 5'- TCGGGGAGCTGGATCCTCCGCCAGCGCTGC-3' and inserted into BamHI site of pQE-80L MBP-SspB Nano plasmid (Addgene #60409) by using Gibson assembly. EYFP part was then replaced with mCherry from pmCherry-C1 (Clontech) by restriction digestion and T4 ligase ligation process with AgeI and BsrGI. Arg73 of SspB was mutated to Gln by PCR with the primer set fwd: 5'- AACGCCCagTTTAAGGGCGTGTCTCGT-3' and rev: 5'- CTTAAActGGGCGTTGAACTGGATAA-3'.

pGEX-2T-GST-3Csite-TEVsite-6×His

Two sets of oligo DNA pairs were designed to introduce 3C protease cleavage site, TEV protease cleavage site, and 6×His tag (1st set, fwd: 5'- <u>GGTTCCGCGTGGATC</u>TGGTCTTGAGGTGCTCTTTCAGGGACCCGGCAGTCTCGAGGGTCTG<u>TACAAGCG</u> <u>AATTCAG</u>-3', rev: 5'- <u>CTGAATTCGCTTGTA</u>CAGACCCTCGAGACTGCCGGGTCCCTGAAAGAGCACCTCAAGACCA<u>GATCCACG</u> <u>CGGAACC</u>-3'. 2nd set, fwd: 5'- <u>TACAAGCGAATTCAG</u>GAGAACCTCTACTTTCAAAGCGATCATCATCATCATCACTAAA<u>AATTCATCGT</u> <u>GACTG</u>-3' and rev: 5'- <u>CAGTCACGATGAATT</u>TTTAGTGATGATGATGATGATGATCGCTTTGAAAGTAGAGGTTCTC<u>CTGAATTCG</u> <u>CTTGTA</u>-3' (underlines indicate overhang sequences)). To anneal oligo DNAs individually, 0.5 µM each DNAs were mixed in 1× T4 Ligase buffer (NEB, B0202S) and heated at 85°C for 10 min and cooled 5°C per 2 min down to 40°C. pGEX-2T, which was kindly gifted from Dr. Miho Iijima, was digested by BamHI and EcoRI and assembled with the two annealed DNA fragments by Gibson assembly.

pGEX2T-GST-pVCA-3Csite-SspB(micro)-mCherry-TEVsite-6×His

To first construct pGEX2T-GST-pVCA-3Csite-TEVsite-6×His, pVCA (proline rich region and VCA domain) of human N-WASP was PCR amplified with the primer set fwd: 5'- CAGTCTCGAGGGAGGTGTTGAAGCTGTTAAAAATGA-3' and rev: 5'- TCCTGAATTCCGTCTTCCCACTCATCATCATCCTC-3' and inserted between XhoI and EcoRI site of pGEX-2T-GST-3Csite-TEVsite-6×His by using restriction digestion and T4 ligase ligation. Then, SspB(micro)-mCherry was PCR amplified with the primer set fwd: 5'- TGAGTGGGAAGACGGAATTGGGCAGTCGACGGTACCGCG-3' and rev: 5'- GTAGAGGTTCTCCTGAATTCCCTTGTACAGCTCGTCCATGCC-3' and inserted in the EcoRI site of pGEX2T-GST-pVCA-3Csite-TEVsite-6×His by Gibson assembly.

pET28-2×Strep-iLID-EYFP-MARCKS

First, pET28-2×Strep-iLID-6×His was constructed as follows. iLID sequence was cut out from the
pLL7.0: Venus-iLID-Mito (Addgene #60413) by NheI and EcoRI and inserted between NheI and
EcoRI sites of pET28-2×Strep-tag gifted from Dr. Kanemaki by using restriction digestion and T4
ligase ligation. Then, to construct pET28-2×Strep-iLID-EYFP-6×His, EYFP was PCR amplified from
pEYFP-C1 (Clontech) with the primer set fwd: 5'-AATTGTCGACATGGTGAGCAAGGGCGAG-3' and
rev: 5'-AATTGCGGCCGCCTTGTACAGCTCGTCCATGC-3' and inserted between SalI and NotI sites of
pET28-2×Strep-iLID-6×His by restriction digestion and T4 ligation. MARCKS-ED fragment was
constructed   by   oligo   annealing   (1st   set   fwd: 5'-
GTACAAGGGAAGTGCTGGTGGTAAAAGAAAAAGAAGCGCTTTTCCTTC-3'   and   rev: 5'-
TTCTTGAAGGAAAAGCGCTTCTTTTTCTTTTTACCACCAGCACTTCCCTT-3',   2nd   set   fwd: 5'-
AAGAAGTCTTTCAAGCTGAGCGGCTTCTCCTTCAAGAAGAACAAGAAGTA-3'   and   rev: 5'-
GTACTACTTCTTGTTCTTCTTGAAGGAGAAGCCGCTCAGCTTGAAAGAC-3') and inserted into BsrGI
site of pET28-2×Strep-iLID-EYFP-6×His.
558
pET28-2×Strep-iLID-EYFP-SNAP
SNAP tag sequence was PCR amplified from the template Phage-ubc-nls-ha-tdMCP-SNAP, a kind
gift   from   Dr.   Bin   Wu,   with   the   primer   set   fwd: 5'-
AATTTGTACAAGTCTGCTGGCGGAAGCGCTGGAGGCAGCATGGACAAAGACTGCGAAATGAAGC-3'
and rev: 5'-AATTGCGGCCGCTTAACCCAGCCCAGGCTTG-3' and inserted between BsrGI and NotI of
pET28-2×Strep-iLID-EYFP-MARCKS by restriction digestion and T4 ligation.
565
pET28-2×Strep-SsrA-SNAP
The plasmid was constructed by replacing the iLID-EYFP part of pET28-2×Strep-iLID-EYFP-SNAP
with the sequence containing SsrA peptide (AANDENY). A set of oligo DNA (5'-
AGCGGGTGCCGCTAGC<u>GCAGCGAACGACGAAAATTAC</u>GGACCGGTCGGTCTGTACAAGTCTGCTGG-3'
and   5'-
CCAGCAGACTTGTACAGACCGACCGGTCC<u>GTAATTTTCGTCGTTCGCTGC</u>GCTAGCGGCACCCGCT-3',
(underlines indicate SsrA peptide region)) was annealed and inserted between NheI and BsrGI
site of pET28-2×Strep-iLID-EYFP-SNAP by Gibson assembly.
574
mCherry-SspB(micro) (Clontech C1)
SspB(micro) sequence was PCR amplified from pQE80L-6×His-MBP-TEVsite-mCherry-SspB(micro)
with   the   primer   set   fwd: 5'-GAACAGTACGAACGCGCC-3'   and   rev: 5'-
AATTCTCGAGGACCACCAGCACTACCACCAGCACTACCACCAGCACTACCACCAGCACTACCACCAGCAC
TACCAATATTCAGCTCGTCATAGATT-3' and inserted between BsrGI and XhoI sites by restriction
digestion and T4 ligation. A silent mutation was introduced to BamHI site upstream of SspB by
inverse PCR with the primer set fwd: 5'-GAGGATCtAGCTCCCCGAAACGCCCT-3' and rev: 5'-
GGGAGCTaGATCCTCCGCCAGCGCTG-3'

pQE80L-MBP-TEVsite-mCherry-MCS

mCherry sequence was PCR amplified from pmCherry-C1 (Clontech) with the primer set fwd: 5'-

GTATTTTCAGGGATCGCTAGCGCTACCGGTC-3'   and   rev: 5'-

TCAGCTAATTAAGCTATCAGTTATCTAGATCCGGTGGATC-3' and inserted between BamHI and

HindIII sites of pQE-80L MBP-SspB Nano by Gibson assembly.

2×Strep-mCherry-MCS

mCherry sequence was PCR amplified from pmCherry-C1 (Clontech) by the primer set fwd: 5'-

AGCGGGTGCCGCTAGTCATATGGGTACGCTAGCGCTACCGGTCG-3'   and   rev: 5'-

GGTGGTGGTGCTCGATCAGTTATCTAGATCCGGTGGATCC-3' and inserted between NheI and XhoI

sites of pET28-2×Strep-tag by Gibson assembly.

2×Strep-MBP-TEVsite-mCherry-MCS

MBP-TEVsite-mCherry-MCS was PCR amplified from the template pQE80L-MBP-TEVsite-

mCherry-MCS with the primer set fwd 5'-CAAATGGGTCGGATCGACGGATCTAAAATCGAAGAAGG-

3' and rev: 5'-GGTGGTGGTGCTCGATCAGTTATCTAGATCCGGTGGA-3' and inserted between

BamHI and XhoI sites of pET28-2×Strep-tag by Gibson assembly.

pCold-6×His-2×Strep-MBP-TEVsite-mCherry-SspB(micro)-mDia1(FH1-FH2-DAD)

A DNA fragment 2×Strep-MBP-TEVsite-mCherry was PCR amplified from 2×Strep-MBP-mCherry-

MCS with the primer set fwd: 5'-TCGAAGGTAGGCATATGGGCTGGTCTCACCC-3' and rev: 5'-

ACAGCTCGTCCATGCCG-3'. The other DNA fragment SspB(micro) was PCR amplified from

mCherry-SspB(micro) (Clontech C1) with the primer sets fwd: 5'-GCATGGACGAGCTGTACAAG-3'

and rev: 5'-TCATTCTTGGCCATAGCTTGAGCTCGAGGACC-3'. pCold-mDia1, which was a kind gift

from Dr. Roberto Dominguez, was digested by NdeI and assembled with the two DNA fragments

by Gibson assembly.

pET28-2×Strep-mCherry-SspB(micro)-mDia1(FH1-FH2-DAD)

mCherry-SspB(micro)-mDia1(FH1-FH2-DAD) was cut out from pCold-6×His-2×Strep-MBP-

TEVsite-mCherry-SspB(micro)-mDia1(FH1-FH2-DAD) by AgeI and BamHI and inserted between

AgeI and BamHI sites of 2×Strep-mCherry-MCS by restriction digestion and T4 ligation.

pET28-2×Strep-mCherry-SspB(micro)-TEVsite-6×His

A DNA fragment mCherry-SspB(micro) was PCR amplified from mCherry-SspB(micro) (Clontech

C1) with the primer set fwd: 5'-AGCGGGTGCCGCTAGCGGACCGGTCGCCACCATG-3' and rev: 5'-

AAAGTAGAGGTTCTCAGATCCGGTGGATCCCGG-3'. The other fragment containing TEV cleavage

site   was   made   by   oligo   annealing   of   a   set   of   DNA   5'-

GAGAACCTCTACTTTCAAagcgatgTCGAGCACCACCACC-3'   and   5'-
GGTGGTGGTGCTCGAcatcgctTTGAAAGTAGAGGTTCTC-3'. These fragments were inserted
between NheI and sites of pET100 by Gibson assembly.

pET28-2×Strep-mCherry-SspB(micro)-mDia1(FH1-FH2-DAD)-TEVsite-6×His
A DNA fragment mDia1(FH1-FH2-DAD) was PCR amplified from pCold-mDia1 with the primer set
fwd: 5'-GGTCCTCGAGCCAAGAATGAAATGGCTTCTCTCTC-3'   and   rev: 5'-
CGGTGGATCCGCTTGCACGGCCAACCAG-3' and inserted between XhoI and BamHI sites of pET28-
2×Strep-mCherry-SspB(micro)-TEVsite-6×His by Gibson assembly.

pET28-2×Strep-mCherry-SspB(micro)-mDia1ΔDAD(FH1-FH2)-TEVsite-6×His
A DNA fragment mDia1(FH1-FH2) was PCR amplified from pCold-mDia1 with the primer set
fwd: 5'-GGTCCTCGAGCCAAGAATGAAATGGCTTCTCTCTC-3'   and   rev: 5'-
CGGTGGATCCCTTCTCCTTGGCTAATTTTGCTCTCCG-3' and inserted between XhoI and BamHI
sites of pET28-2×Strep-mCherry-SspB(micro)-TEVsite-6×His by Gibson assembly.

EYFP-FKBP-mDia1(FH1-FH2-DAD)
mDia1(FH1-FH2-DAD) was cut out from pCold-6×His-2×Strep-MBP-TEVsite-mCherry-
SspB(micro)-mDia1(FH1-FH2-DAD) by XhoI and BamHI and inserted between XhoI and BamHI of
EYFP-FKBP by restriction digestion and T4 ligation.

pET28-6×His-2×Strep-TEVsite-Capβ2(mouse)-Capα1(mouse)
First, pET28-6×His-2×Strep was constructed as follows. 6×His sequence was synthesized by oligo
annealing of fwd: 5'-AGGAGATATACCATGGGCCATCACCATCACCATCACATGGGCTGGTCTCA-3' and
rev: 5'- TGAGACCAGCCCATGTGATGGTGATGGTGATGGCCCATGGTATATCTCCT-3' and inserted into
NcoI site of pET28-2×Strep-tag. Capβ2(mouse)-Capα1(mouse) was PCR amplified from the
template of pET3d-Capβ2(mouse)-Capα1(mouse), gifted from Dr. Julie Plastino and Dr. Cecile
Sykes,   with   the   primer   set   fwd: 5'-
CTGGAGCGGGTGCCGCTAGCGAGAACCTCTACTTTCAAAGCGATATGAGCGATCAGCAGCTGG-3' and
rev: 5'- TGTCGACGGAGCTCGAATTCTTAAGCATTCTGCATTTCTTTGCCAATC-3' and inserted
between NheI and EcoRI of pET28-6×His-2×Strep by Gibson assembly. TEV cleavage site was
designed in the forward primer and introduced between 2×Strep tag and Capβ2 ORF.

Actin preparation
Actin for GUV experiments was reconstituted from lyophilized powder as described in
manufacturer's instructions, aliquoted, snap-frozen in liquid nitrogen, and stored at -80°C.
For actin polymerization assays, actin was dissolved in G-buffer [2 mM Tris-HCl (pH 7.5 at RT), 0.1
mM $CaCl_2$, 0.2 mM ATP, 0.5 mM DTT, 1 mM NaN_3] at concentration less than 65 μM and dialyzed

against G-buffer for 3 days with daily buffer exchange. Dialyzed actin was then purified by a
Superdex 200 increase 10/300 GL column (Cytiva) to separate monomer fraction from occasional
larger molecular size (smaller elution volume) fraction. The purified actin was stored at 4°C with
dialysis against G-buffer. The dialysis buffer for actin storage was exchanged twice a week.
Alexa 647 labeling of actin was performed following the previously reported method (*134*). First,
rabbit skeletal muscle actin (Cytoskeleton Inc., #AKL99) was dissolved in G*-buffer [2 mM Tris-
HCl (pH 7.5 at RT), 0.1 mM $CaCl_2$, 0.2 mM ATP] at concentration less than 65 µM. The solution
was dialyzed against G*-buffer overnight, followed by an additional 3 hour dialysis with fresh
buffer. Dialyzed actin was collected and mixed with 4 molar excess Alexa Fluor 647 C2 maleimide
(ThermoFisher, A20347, solubilized to 10 mM in DMSO) and incubated at 4°C overnight with
rotation (10 min labeling as in the original protocol did not yield efficient labeling in the case of
Alexa Fluor 647). The reaction was quenched by adding DTT to 10 mM. Aggregated proteins and
insoluble dyes were removed by ultracentrifugation at 350,000×g (90,000 rpm, TLA-100.1 rotor)
for 12 min. The supernatant was collected and mixed 1/10 volume of 10×KMEI buffer (500 mM
KCl, 10 mM $MgCl_2$, 10 mM EGTA, 100 mM Imidazole, pH 7.0) and final 1 mM ATP to polymerize
actin at RT for 2–3 hours. Labeled filamentous actin was collected by ultracentrifugation at
195,000×g (67,000 rpm, TLA-100.1 rotor) for 30 min and resuspended with G-buffer. The labeled
actin was dialyzed against G-buffer for 3 days with daily buffer exchange and purified through
size exclusion chromatography with a Superdex 200 increase 10/300. The protein was snap-
frozen in liquid nitrogen and stored at -80°C.
679
680   **Protein purification**
681   Protein expressions were carried out in BL21-CodonPlus (DE3)-RIL (Agilent Technologies) and cell
682   lysis was performed by microfluidizer (Microfluidics, Model M-110Y), unless otherwise specified.
683
6×His-MBP-TEV protease (S219V)
The bacteria transformed with pQE80L-6×His-MBP-TEVprotease (S219V) were cultured overnight
in LB broth supplemented with 100 µg/mL Ampicillin and 25 µg/mL Chloramphenicol
(LB/Amp/Cam). Pre-culture was inoculated to LB/Amp/Cam and cultured at 37°C until OD_{600}
reached around 0.4. Protein expression was induced by adding 300 µM IPTG and allowed to
proceed for 17 hours at 18°C. The cells were re-suspended with 50 mM Tris-HCl (pH 7.5 at RT),
250 mM NaCl, 5 mM $MgCl_2$, 7 mM β-mercaptoethanol and lysed by microfluidizer. The lysate was
centrifuged at 20,000×g at 4°C for 30 min to remove cell debris. The protein was purified by Ni-
NTA, eluted around 60–100 mM imidazole, and then dialyzed against 25 mM Tris-HCl (pH 7.5 at
RT), 200 mM NaCl, 7 mM β-mercaptoethanol. The elution was concentrated by Amicon Ultra (10
kDa cutoff) and further dialyzed against 25 mM Tris-HCl (pH 7.5 at RT), 200 mM NaCl, 7 mM β-
mercaptoethanol, 50% glycerol. The protein was stored at -20 °C.
696

mCherry-SspB

The bacteria transformed with pQE80L-6×His-MBP-TEVsite-mCherry-SspB were cultured

overnight in LB broth supplemented with 100 μg/mL Ampicillin and 25 μg/mL Chloramphenicol

(LB/Amp/Cam). Pre-culture was inoculated to LB/Amp/Cam and cultured at 37°C until OD_{600}

reached 0.4–0.6. Protein expression was induced by adding 300 μM IPTG and allowed to proceed

for 18 hours at 22°C. The cells were re-suspended with 50 mM Tris-HCl (pH 7.5 at RT), 250 mM

NaCl, 7 mM β-mercaptoethanol, 20 mM Imidazole supplemented with cOmplete EDTA-free

protease Inhibitor Cocktail (MilliporeSigma) and lysed by microfluidizer. The lysate was

centrifuged at 20,000×g at 4°C for 30 min to remove cell debris. The protein was first purified by

Ni-NTA, eluted around 50 mM imidazole, and then dialyzed against 25 mM Tris-HCl (pH 7.5 at RT),

250 mM NaCl, 7 mM β-mercaptoethanol. 6×His-MBP tag was cleaved by 6×His-MBP-TEV protease

(S219V) (purified in house) during the dialysis. The cleaved 6×His-MBP tag and TEV protease were

removed through the second round of Ni-NTA column. Flow through and wash fractions were

dialyzed against buffer S [25 mM Tris-HCl (pH 7.5 at RT), 250 mM NaCl, 7 mM β-mercaptoethanol]

and concentrated by Amicon Ultra (10 kDa cutoff). The protein was snap-frozen in liquid nitrogen

in storage buffer [25 mM Tris-HCl (pH 7.5 at RT), 250 mM NaCl, 7 mM β-mercaptoethanol, 20%

glycerol] and stored at -80 °C.

iLID-YFP-MARCKS

The bacteria transformed with pET28-2×Strep-iLID-EYFP-MARCKS were cultured overnight in LB

broth supplemented with 50 μg/mL Kanamycin and 25 μg/mL Chloramphenicol (LB/Kan/Cam).

Pre-culture was inoculated to LB/Kan/Cam and cultured at 37°C until OD600 reached around 0.6.

Protein expression was induced by adding 300 μM IPTG and allowed to proceed for 17 hours at

23°C in the presence of 6 μM FMN. The cells were resuspended with buffer L1 [50 mM Tris-HCl

(pH 7.5 at RT), 200 mM NaCl, 1 mM EDTA, 7 mM β-mercaptoethanol] and lysed by microfluidizer.

The lysate was centrifuged at 20,000×g at 4°C for 30 min to remove cell debris. The supernatant

was applied to Strep-Tactin XT (IBA Lifesciences) equilibrated with buffer L1 and the protein was

eluted with 50 mM biotin in buffer L1. The elution was dialyzed against buffer L2 [25 mM Tris-

HCl (pH 7.5 at RT), 200 mM NaCl, 7 mM β-mercaptoethanol] and concentrated by Amicon Ultra

(10 kDa cutoff). The protein was snap-frozen in liquid nitrogen in storage buffer [25 mM Tris-HCl

(pH 7.5 at RT), 200 mM NaCl, 7 mM β-mercaptoethanol, 30% glycerol] and stored at -80 °C.

iLID-YFP-SNAP

The bacteria transformed with pET28-2×Strep-iLID-EYFP-SNAP were cultured overnight in LB

broth supplemented with 50 μg/mL Kanamycin and 25 μg/mL Chloramphenicol (LB/Kan/Cam).

Pre-culture was inoculated to LB/Kan/Cam and cultured at 37°C until OD_{600} reached around 0.5.

Protein expression was induced by adding 300 μM IPTG and allowed to proceed for 20 hours at

23°C in the presence of 6 μM FMN. The cells were resuspended with buffer L1 (50 mM Tris-HCl

(pH 7.5 at RT), 200 mM NaCl, 1 mM EDTA, 7 mM β-mercaptoethanol) and lysed by microfluidizer.
The lysate was centrifuged at 20,000×g at 4°C for 30 min to remove cell debris. The supernatant
was applied to Strep-Tactin XT (IBA Lifesciences) equilibrated with buffer L and the protein was
eluted with 50 mM biotin in buffer L1. The elution was further purified by Superdex 200 increase
10/300 GL column in the buffer L2 [25 mM Tris-HCl (pH 7.5 at RT), 200 mM NaCl, 7 mM β-
mercaptoethanol] and concentrated by Amicon Ultra (10 kDa cutoff). The protein was snap-
frozen in liquid nitrogen in storage buffer [25 mM Tris-HCl (pH 7.5 at RT), 200 mM NaCl, 7 mM β-
mercaptoethanol, 30% glycerol] and stored at -80 °C.

SsrA-SNAP
BL21(DE3) RP transformed with pET28-2×Strep-SsrA-SNAP were cultured overnight in LB broth
supplemented with 50 μg/mL Kanamycin and 25 μg/mL Chloramphenicol (LB/Kan/Cam). Pre-
culture was inoculated to LB/Kan/Cam and cultured at 37°C until OD_{600} reached around 0.5.
Protein expression was induced by adding 300 μM IPTG and allowed to proceed for 20 hours at
23°C. The cells were resuspended with lysis buffer [50 mM Tris-HCl (pH 7.6 at RT), 200 mM NaCl,
1 mM $MgCl_2$, 7 mM β-mercaptoethanol, 1 mM PMSF, 1× protease inhibitor cocktail V (Wako 168-
26033)] supplemented with house-purified benzonase and sonicated by Misonix Ultrasonic
Liquid Processor Sonicator S4000 (amplitude 30, 2 seconds ON, 10 seconds OFF, total 3 minutes
ON). The lysate was centrifuged at 13,420×g at 4°C for 45 min to remove cell debris. The
supernatant was applied to Strep-Tactin XT High Capacity (IBA Lifesciences) equilibrated with
buffer L3 (25 mM Tris-HCl (pH 7.6 at RT), 200 mM NaCl, 1 mM $MgCl_2$, 7 mM β-mercaptoethanol,
1 mM PMSF) and the protein was eluted with 100 mM biotin in wash buffer. The elution was
dialyzed against buffer L3 and further purified by Superdex 200 increase 10/300 GL column in
buffer L3. Peak fractions were collected and concentrated by Viva Spin Turbo (10 kDa cutoff). The
protein was snap-frozen in liquid nitrogen in storage buffer (25 mM Tris-HCl (pH 7.6 at RT), 200
mM NaCl, 7 mM β-mercaptoethanol, 40% glycerol) and stored at -80 °C.

GST-pVCA-SspB-mCherry
The bacteria transformed with pGEX2T-GST-pVCA-3Csite-SspB(micro)-mCherry-TEVsite-6×His or
pGEX2T-GST-pVCA-3Csite-SspB(nano)-mCherry-TEVsite-6×His were cultured overnight in LB
broth supplemented with 100 μg/mL Ampicillin and 25 μg/mL Chloramphenicol (LB/Amp/Cam).
Pre-culture was inoculated to LB/Amp/Cam and cultured at 37°C until OD_{600} reached around 0.5–
0.6. After cooling the culture on ice, protein expression was induced by adding 300 μM IPTG and
allowed to proceed for 24–26 hours at 18°C. The cells were resuspended with buffer V (50 mM
HEPES-NaOH (pH 7.5), 150 mM NaCl, 10% glycerol, 14 mM β-mercaptoethanol, 1 mM EDTA, 1
mM PMSF, 100 μg/mL DNase, supplemented with cOmplete EDTA-free protease inhibitor
cocktail) and lysed by microfluidizer. The lysate was centrifuged at 20,000×g at 4°C for 30 min to
remove cell debris. To enhance resin binding, final 5 mM DTT was added to the lysate. The lysate

was applied to glutathione sepharose equilibrated with buffer V [50 mM HEPES-NaOH (pH 7.5)

150 mM NaCl, 10% glycerol, 7 mM β-mercaptoethanol, and 1 mM PMSF], and the protein was

eluted with 20 mM glutathione in buffer V (pH was adjusted between 7.0–8.0 by NaOH). The

elution was then purified by Ni-NTA and eluted with 50–500 mM imidazole. After concentrated

by Amicon Ultra (10 kDa cutoff), the protein was further purified by Superdex 200 increase

10/300 GL column in buffer V and concentrated again by Amicon Ultra (10 kDa cutoff). The

protein was snap-frozen in liquid nitrogen and stored at -80 °C.

mCherry-SspB-mDia1

The bacteria transformed with pET28-2×Strep-mCherry-SspB(micro)-mDia1(FH1-FH2-DAD) were

cultured overnight in LB broth supplemented with 50 μg/mL Kanamycin and 25 μg/mL

Chloramphenicol (LB/Kan/Cam). Pre-culture was inoculated to LB/Kan/Cam and cultured at 37°C

until OD_{600} reached around 0.5–0.6. After cooling the culture on ice, protein expression was

induced by adding 300 μM IPTG and allowed to proceed for 24 hours at 16°C. The cells were

resuspended with buffer D [50 mM Tris-HCl (pH 7.5), 500 mM NaCl, 10% glycerol, 1 mM DTT, 1

mM EDTA, 1 mM PMSF, 100 μg/mL DNase, supplemented with cOmplete EDTA-free protease

inhibitor cocktail] and lysed by microfluidizer. The lysate was centrifuged at 20,000×g at 4°C for

30 min to remove cell debris. The supernatant was applied to Strep-Tactin XT (IBA Lifesciences)

equilibrated with buffer D1 (50 mM Tris-HCl (pH 7.5), 500 mM NaCl, 10% glycerol, 1 mM DTT, 1

mM EDTA, 1 mM PMSF) and the protein was eluted with 50 mM biotin in buffer D1. The elution

was concentrated by Amicon Ultra (10 kDa cutoff) and dialyzed against buffer D2 [25 mM HEPES-

NaOH (pH 7.5), 150 mM NaCl, 7 mM β-mercaptoethanol, 10% glycerol]. The protein was further

purified by Superdex 200 increase 10/300 GL column in buffer D2 and concentrated again by

Amicon Ultra (10 kDa cutoff). The protein was snap-frozen in liquid nitrogen and stored at -80 °C.

This protein was used as mDia1 in most figures unless otherwise indicated.

mCherry-SspB-mDia1-6×His and mCherry-SspB-mDia1ΔDAD-6×His

The bacteria transformed with pET28-2×Strep-mCherry-SspB(micro)-mDia1(FH1-FH2-DAD)-

TEVsite-6×His   or   pET28-2×Strep-mCherry-SspB(micro)-mDia1ΔDAD(FH1-FH2)-TEVsite-6×His

were cultured overnight in LB broth supplemented with 50 μg/mL Kanamycin and 25 μg/mL

Chloramphenicol (LB/Kan/Cam). Pre-culture was inoculated to LB/Kan/Cam and cultured at 37°C

until OD_{600} reached around 0.5–0.6. After cooling the culture on ice, protein expression was

induced by adding 300 μM IPTG and allowed to proceed for 24 hours at 16°C. The cells were

resuspended with buffer D3 [50 mM Tris-HCl (pH 8.0), 500 mM NaCl, 10% glycerol, 1 mM DTT,

1 mM PMSF] supplemented with 1× protease inhibitor cocktail V (Wako 168-26033)] and house-

purified benzonase, and sonicated by Misonix Ultrasonic Liquid Processor Sonicator S4000

(amplitude 30, 2 seconds ON, 10 seconds OFF, total 3 minutes ON). Unbroken cells were pelleted

by centrifugation at 13,420×g for 5 min at 4°C, resuspended in buffer D1, and subjected to
sonication under the same conditions. This procedure was repeated twice. The lysate was then
clarified by centrifugation at 13,420×g for 45 min at 4°C. The supernatant was applied to Strep-
Tactin XT (IBA Lifesciences) equilibrated with buffer D3 and the protein was eluted with 100 mM
biotin in buffer D3. The elution was then purified by Ni-NTA equilibrated with 50 mM Tris-HCl (pH
8.0), 150 mM NaCl, 10% glycerol, 7 mM β-mercaptoethanol, 1 mM PMSF and eluted stepwise
with 20–500 mM imidazole. Peak fractions were collected and diluted 3-fold with 50 mM Tris-
HCl (pH 8.0), 10% glycerol, 7 mM β-mercaptoethanol, 1 mM PMSF. The protein was further
purified by HiTrapQ column equilibrated with 50 mM HEPES-NaOH (pH 7.5), 50 mM NaCl, 10%
glycerol, 7 mM β-mercaptoethanol, 1 mM PMSF, and eluted with linear NaCl gradients of 50–500
mM NaCl. The protein was concentrated by Viva Spin Turbo (10 kDa cutoff), snap-frozen in liquid
nitrogen and stored at -80 °C. mCherry-SspB-mDia1-6×His was used in Figure 2E, F, H, S10, S12.
mCherry-SspB-mDia1ΔDAD-6×His was used in Figure 4J.

Capping protein
The bacteria transformed with pET28-6×His-2×Strep-TEVsite-Capβ2(mouse)-Capα1(mouse) were
cultured overnight in LB/Kan/Cam. Pre-culture was inoculated to LB/Kan/Cam and cultured at
37°C until OD_{600} reached around 0.7. Protein expression was induced by adding 300 µM IPTG and
allowed to proceed for 24 hours at 18°C. The cells were re-suspended with 50 mM Tris-HCl (pH
7.5 at RT), 150 mM NaCl, 14 mM β-mercaptoethanol, 1 mM EDTA, 1 mM PMSF, 100 µg/mL
DNase and lysed by microfluidizer. The lysate was centrifuged at 20,000×g at 4°C for 30 min to
remove cell debris. The supernatant was applied to Strep-Tactin XT 4Flow (IBA Lifesciences)
equilibrated with buffer C1 [50 mM Tris-HCl (pH 7.5 at RT), 150 mM NaCl, 7 mM β-
mercaptoethanol] and the protein was eluted with 50 mM biotin in buffer C1. The elution was
dialyzed against buffer C2 [25 mM Tris-HCl (pH 7.5 at RT), 150 mM NaCl, 7 mM β-
mercaptoethanol]. 6×His-2×Strep tag was cleaved by 6×His-MBP-TEV protease (S219V) (purified
in house) during the dialysis. The cleaved 6×His-2×Strep tag and TEV protease were removed
through Ni-NTA column. Flow through and wash fractions were dialyzed against buffer and
concentrated by Amicon Ultra (10 kDa cutoff). The protein was snap-frozen in liquid nitrogen in
buffer C2 and stored at -80 °C.

Pyrene-actin polymerization assay
For pyrene actin assay, actin was purified by size exclusion column. One milligram of actin
(Cytoskeleton Inc. AKL99-B) was dissolved in 400 µL G-buffer [2 mM Tris-HCl (pH 7.5 at RT), 0.1
mM $CaCl_2$, 0.2 mM ATP, 0.5 mM DTT, 1 mM NaN_3] and dialyzed against the same buffer. After 3
days dialysis with daily buffer exchange, the actin was purified by Superdex 200 Increase 10/300
GL column (GE Healthcare) to separate monomer fraction from a fraction occasionally appear at

larger size molecular size (smaller elution volume). The purified actin was stored at 4°C in a
dialyzed manner against G-Buffer with twice a week buffer exchange.
Pyrene-actin polymerization assay was performed with FluoroMax 3 and DataMax software.
Pyrene fluorescence and its kinetics were measured by 365 nm excitation (1 nm bandwidth) and
407 nm emission (5 nm bandwidth). The reaction was prepared by following the previously
reported method (*135*), except that the reaction was 50 μL in quartz cuvettes (Hellma, 105-251-
15-40).
855

Ultracentrifugation assay
Reaction solutions were prepared by mixing 4.5 μL of actin in G-buffer and 4.5 μL of other
proteins solution. At the final concentration, reaction solution contained 7.5 μM actin (10% Alexa
488 labeled), 1 μM GST-pVCA-SspB(micro)-mCherry, 1 μM mCherry-SspB(micro)-mDia1, 150 nM
Arp2/3, 50 nM capping protein, 3 μM profilin, 2 μM cofilin, 1 μM creatin kinase, 1 mM ATP, 25
mM creatin phosphate, 240 mM sucrose, and 1×KMEI buffer [10 mM imidazole pH 7.0, 50 mM
KCl, 1 mM $MgCl_2$, 1 mM EGTA]. Polymerization reaction for 30 min at RT. The solutions were
ultracentrifuged at 100,000×g (90,000 rpm, S55A2 rotor) for 1 hour at RT and analyzed by SDS-
PAGE.
865

GUV preparation by emulsion transfer
Lipid-in-oil solution was prepared as follows. Chloroform solutions of lipids were mixed in 5 mL
glass vial, dried at 70°C with N_2 gas flow, and left in vacuum 3 hours to overnight. The amounts
of lipids were typically 6.5 μmol (1.3 mM, 1 mg/mL at final concentration in oil) for 40 mol%
POPS/60 mol% POPC mixture (MARCKS peptide anchoring) and 1.5 μmol (0.3 mM, 0.23 mg/mL
at final concentration in oil) for 2 mol% Benzylguanine-conjugated lipid/98 mol% POPC and 100%
POPC (SNAP-tag anchoring). Lipid was dissolved in 5 mL of oil mixture (90 volume% hexadecane,
10 volume% silicone oil) by heating at 80°C for 1–2 hours, then brought to RT.
For iLID-SspB translocation experiments, inner solution contained 4 μM iLID-YFP-SNAP (or iLID-
EYFP-MARCKS) and 1 μM mCherry-SspB(micro) in buffer (22.5 mM Tris-HCl (pH 7.5 at RT), 122.5
mM NaCl, 2 mM $MgCl_2$, 240 mM sucrose, 0.7 mM β-mercaptoethanol, and 3% glycerol). Outer
solution was prepared by replacing protein fractions of inner solution with their buffers and
sucrose with glucose.
For actin polymerization experiments, inner solution was prepared by mixing 10 μL of actin (10%
Alexa 647 labeled) in G-buffer and 10 μL of other proteins solution similarly to pyrene actin
polymerization assay. At the final concentration, inner solution typically contained 11.75 μM
iLID-EYFP-SNAP, 1 μM GST-pVCA-SspB(micro)-mCherry, 1 μM mCherry-SspB(micro)-mDia1, 7.5
μM actin, 150 nM Arp2/3, 50 nM capping protein, 3 μM profilin, 2 μM cofilin, 1 μM creatine
kinase in the buffer (5.9 mM Tris-HCl (pH 7.5 at RT), 1.3 mM HEPES-NaOH (pH 7.5 at RT), 7.7 mM
Imidazole (pH 8.0 at RT), 27 μM KH_2PO_4, 74 μM Na_2HPO_4, 39 mM KCl, 33 mM NaCl, 0.8 mM $MgCl_2$,

43 μM CaCl$_2$, 1.1 mM EGTA, 1.2 mM β-mercaptoethanol, 0.21 mM DTT, 0.43 mM NaN$_3$, 1 mM
ATP, 25 mM creatine phosphate, 240 mM sucrose and 2.8% glycerol). Outer solution was
prepared by replacing protein fractions of inner solution with their buffers and sucrose with
glucose. Protein and buffer conditions are summarized in Supplementary table 1.
In experiments using pVCA alone, SspB(nano) was employed to ensure sufficiently robust light-
induced membrane recruitment and actin polymerization, and pVCA was used at 4–4.2 μM.
When mDia1 was included, SspB(micro) was used to avoid excessive basal membrane association
of mDia1 prior to illumination. To maintain comparable recruitment behavior within the
combined system, pVCA was also fused to SspB(micro) and used at 1 μM, matching the
concentration of mDia1.
GUVs were prepared as previously reported (*56*, *136*). Briefly, oil-buffer interface was created by
layering 250 μL lipid-in-oil solution on top of 250 μL outer buffer in 1.5 mL tube. Water-in-oil
emulsion was made by vigorously pipetting 20 μL of inner solution in 250 μL of lipid-in-oil solution
and then gently added to the top of the oil-buffer layers. The emulsions were transferred through
the oil-buffer interface by centrifugation at 2500×g for 2 min at RT. GUVs were collected by
pipetting.
902
Image acquisition and analysis
For imaging GUVs, 8-well glass chamber slides (ThermoFisher Scientific, 154534) were pretreated
with 10 mg/mL BSA in PBS for 10 minutes and washed with MilliQ water twice. LSM780 confocal
microscope (Zeiss) equipped with Plan-Apochromat 63X/1.40na oil immersion DIC objective lens
(Zeiss 420782-9900) was used for imaging and light stimulation of GUVs. YFP, mCherry, and Alexa
Fluor 647 were imaged with 514, 561, and 633 nm excitation lasers, respectively. Blue light
illumination for iLID stimulation was performed by using bleaching function with 458 nm laser at
intensity 1.0–5.0%, scan speed 6 (pixel dwell time = 6.3 μsec). Main beam splitters MBS458/561,
MBS 488/561/633, and MBS 458/514 were used for 458 and 561, 633, and 514 nm excitation,
respectively. The imaging was performed at room temperature. Data were analyzed by Fiji
software (*137*).
914
FRAP analysis on GUV membranes
A small region of interest (ROI) on the GUV membrane was selectively photobleached for 1 s
using high-intensity laser illumination. Fluorescence recovery within the bleached ROI was
measured. To correct for photobleaching caused by repeated imaging, fluorescence intensity in
the bleached ROI was normalized to that of a non-bleached reference region on the same GUV
membrane. To capture rapid fluorescence recovery with short imaging intervals, mCherry- and
Alexa Fluor 647–labeled signals were analyzed on different individual GUVs.
922
923   **Small unilamellar vesicles (SUVs) preparation**

SUVs used for EM experiments were prepared as follows. Chloroform solutions of lipids were mixed in 5 mL glass vial, dried at 70°C under a flow of N_2 gas, and left under vacuum overnight. The total amount of lipid was 0.5 µmol (corresponding to 5 mM, or 4 mg/mL, at the final concentration in buffer). Lipid composition was 2 mol% benzylguanine-conjugated lipid/98 mol% POPC. The lipid film was hydrated with 100 µL of SUV buffer (25 mM HEPES-KOH (pH 7.6), 50 mM KCl, 1 mM $MgCl_2$, 1 mM TCEP), incubated at 37°C for 10 min, and sonicated using Misonix Ultrasonic Liquid Processor Sonicator S4000.

Electron microscopy (EM)

The reaction for EM imaging was prepared by mixing 2.75 µL of a solution containing all components except actin (solution I) with 2.25 µL of actin in G-Mg buffer [2 mM Tris-HCl (pH 7.5), 0.2 mM ATP, 0.1 mM $MgCl_2$, 0.5 mM DTT, 1 mM NaN_3] (solution II). Solution I was first placed to carbon-coated collodion film on a Cu grid (Nisshin EM, Tokyo, Japan) and incubated for several seconds to allow SUVs to settle onto the grid surface. Solution II was then added, and actin polymerization was allowed to proceed for 3 min at room temperature (24–26°C). At final concentrations, the reaction mixture contained 0.3 mg/mL lipid SUVs, 7.5 µM actin (10% Alexa Fluor 647–labeled), 1 µM GST–pVCA–SspB(micro)–mCherry, 1 µM mCherry–SspB(micro)–mDia1(FH1–FH2–DAD)–6×His, 150 nM Arp2/3 complex, 50 nM capping protein, 3 µM profilin, 2 µM cofilin, 1 µM creatine kinase, 1 mM ATP, 25 mM creatine phosphate, and 1× KMEI buffer (10 mM imidazole, pH 7.0, 50 mM KCl, 1 mM $MgCl_2$, 1 mM EGTA). After the reaction, excess liquid on the grid was carefully removed, followed by negative staining with 2% phosphotungstic acid (pH 7.0). The samples were observed with a JEM1010 EM (JEOL, Akishima, Japan) at 80 kV equipped with a FastScan-F214(T) charge-coupled device (CCD) camera (TVIPS, Gauting, Germany).

Live-cell CID experiments

DNA constructs: Four plasmids were used which encoded Lyn-CFP-FRB (*138*), YFP-FKBP (*138*), YFP-FKBP-Tiam1 (*133*), YFP-FKBP-pVCA, and YFP-FKBP-mDia1.

Cell culture: HEK293T cells (ATCC) were cultured in DMEM (Corning, 10-013-CV) supplemented with 10% fetal bovine serum (Corning, 35-010-CV) and 1% penicillin-streptomycin (ThermoFisher, 15140163).

Transient transfection: HEK293T cells were seeded at a density of 9,000-12,000 cells in an 8-well chamber slide. Per well, a solution of 0.3 µg of DNA, 1 µL of FuGENE HD, and 30 µL of Opti-MEM was added for transfection. FRB:FKBP pairs were cotransfected in a 1:1 or 1:2 ratio. Cells were incubated at 37°C with 5% CO_2 and 95% humidity for 24–48 h before imaging.

Microscopes and imaging: Live cell imaging of actin NPF recruitment to the cell membrane was performed using an Eclipse Ti inverted fluorescence microscope (Nikon) with a 100x Oil objective lens and Andor Zyla 4.2 plus sCMOS camera, or with a 60x Oil, TIRF 60x Oil, or TIRF 100x Oil

objective lens and ORCA-Fusion Digital CMOS camera. Images were captured every 1 min for 30-

120 min, where within 5 min of starting, 0.1 μM rapamycin was added. Nikon microscopes were

driven by NIS-Elements software (Nikon). All live-cell imaging was performed at 37 °C with 5%

CO_2 and humidity control by a stage top incubator. Image processing and analysis were

performed using Fiji software.

Data analysis: Per transfection sample, a count was made based on the phenotype of cells falling

into three categories: no change, lamellipodia, and filopodia. Cells with lamellipodia or filopodia

pre-rapamycin were not counted unless new membrane protrusions formed after rapamycin

addition. If the cell under observation did not produce any new lamellipodia or filopodia after

addition of rapamycin, then it was counted as no change. Only healthy cells fully in the field of

view with clear expression of both FRB and FKBP constructs were counted. If there was any doubt

in the phenotype of the cell, then it was not included in the count.

Statistics

1. The cell counts for each category were summed across all experiments, and this sum was

divided by the total number of cells to get a percentage of cells belonging to each category when

a certain actin NPF was recruited to the membrane.

2. For each image set collected from a transfection sample, the categorized cell counts were

divided by the total cell count to get percentages. For each actin NPF, the percentages from all

transfection samples were averaged. These averages were analyzed by two-tailed Student's t-

test assuming equal variance (determined by F-test) in order to establish the significance of

lamellipodia or filopodia detection between transfection samples.

Statistics and reproducibility

Quantitative data with error bars were obtained from three or more independent experiments,

except for fig. S10 and S11, which were obtained from two independent experiments. Where

indicated, 95% confidence intervals of the mean were calculated in Microsoft Excel using the t-

distribution (T.INV.2T) and the standard error (SD/√n). For statistical comparison of multiple

groups, the Steel–Dwass test was performed using the pSDCFlig function (asymptotic option) in

the NSM3 package in R. Pearson's R and p-values in fig. S19 was calculated by cor.test function

(pearson option).

Theoretical model of actin filament-membrane interactions

We employed Cytosim (*95*), a stochastic agent-based cytoskeletal modeling framework that

incorporates Brownian dynamics, actin filament dynamics, and the action of actin-binding

proteins. To Cytosim, we added a deformable membrane model (*96*) along with terms for the

reciprocal actin-membrane interactions (*97*). In the following text we elaborate on the details

and implementation for each model component. We further refer the reader to Table 1, for a

complete list of parameters and their corresponding references from where they were obtained.

The use of these physically relevant parameters constraint our simulations against reality.

1001

1002   To model the membrane, we utilize a 2-D enclosed polygon with resistance to bending and

1003   stretching. The energy of the membrane is given by discrete 2-D Helfrich Hamiltonian,

$$E = \frac{\kappa}{4} \sum_{i=1}^{N_{edges}} k_i^2 l_i^2 + \sum_{i=1}^{N_{edges}} \sigma l_i^2,$$

1005   where κ is the bending rigidity of the membrane and σ is the membrane surface tension. k_i is the

1006   local curvature at vertex i and l_i is the length of the edge connecting vertices i and $i+1$. The force

1007   on a vertex i is given by

$$\frac{\partial E}{\partial r_i} = \frac{\kappa}{4} \sum_{i=1}^{N_{edges}} \left[l_i^2 \frac{\partial}{\partial r_i} (k_i^2) + 2 l_i k_i^2 \frac{\partial}{\partial r_i} l_i \right] + \sum_{i=1}^{N_{edges}} 2 \sigma l_i \frac{\partial}{\partial r_i} l_i.$$

1009   There are several established ways to define curvature in discrete settings (*139, 140*). For this

1010   model, we choose a definition of curvature based on the turning angle at each vertex.

$$k_i = \frac{\tan\left(\frac{\alpha_i}{2}\right) + \tan\left(\frac{\alpha_{i+1}}{2}\right)}{l_i},$$

1012   where the turning angle α_i is given by the sum of the internal angles,

$$\alpha_i = \tan^{-1}\left(\frac{l_{i,y}}{l_{i,x}}\right) + \tan^{-1}\left(\frac{l_{i+1,y}}{l_{i+1,x}}\right),$$

1014   and we can express $\partial l_i / \partial r_a$ and $\partial \alpha_i / \partial r_a$ as

$$\frac{\partial l_i}{\partial r_a} = -\frac{e_i}{|e_i|},$$

$$\frac{\partial \alpha_i}{\partial r_a} = \frac{\eta_i}{|\eta|^2},$$

respectively, as the unit vector along the edge i and the unit normal to the edge i.

Filament description

The actin filaments are modeled as described in the original Cytosim reference (*95*). In brief, actin
filaments are modeled as semiflexible, inextensible fibers composed of linear segments with
hinge points that allow for bending. The bending rigidity of the filament, k_{bend}, is chosen so that
the persistence length of an isolated filament is 17 μm.

1023

1024 **Filament-membrane interactions**
1025 We model filament-membrane interactions using a purely repulsive harmonic potential between
1026 points on the filament and the edges of the enclosed polygon. Given a position **p** on a filament *i*
1027 and an edge *j* on the enclosed membrane, the force is given by

1028

1029 $F^{excl} = K^{excl}(\boldsymbol{q} - \boldsymbol{p})$,
1030 where $\boldsymbol{q}$ is the point closest to the point $\boldsymbol{p}$ on the edge *j* and K^{excl} is the stiffness of the steric
1031 potential.

1032

1033 F^{excl} confines the actin filament within the enclosed membrane and applies a force from the
1034 actin filament to the membrane. The steric force to points on the membrane edge c_j and c_{j+1}
1035 is calculated using the lever rule,

$$F(c_{j+1}) = F^{excl}\left(\frac{|q - c_j|}{|c_{j+1} - c_j|}\right),$$

1036

$$F(c_j) = F^{excl} - F(c_{j+1}).$$

1037

1038

1039 **Inter-filament interactions**
1040 They are implemented as a repulsive harmonic potential between filaments, mimicking the
1041 physical constraints and forces arising from their proximity and entanglement.

1042

1043 **Dynamics of filament growth**
1044 We assume that filaments polymerize at their barbed ends and do not depolymerize at the
1045 pointed ends. We assume a constant polymerization rate dependent on which actin-binding
1046 protein is bound to the filament.

1047

Actin-binding proteins and nucleation promoting factors
We considered three proteins: mDia1 (formin) and Arp2/3. To model the spatial control of mDia1
and pVCA, we assume that iLID-SspB is already attached to pVCA and mDia1 at the membrane
surface. To capture the actin nucleation and polymerization dynamics of the experiment, we
modeled mDia1 and pVCA as beads that can nucleate and elongate actin filaments at prescribed
rates. For reasons of computational cost, we assume that pVCA recruitment of Arp2/3 is fast. It
has been shown experimentally that pVCA unbinds Arp2/3 after nucleation (*141*). In our model,
pVCA unbinds the mother filament after nucleation. After initial nucleation by pVCA-Arp2/3

complex, free Arp2/3 can bind to mother filaments and nucleate daughter filaments. Arp2/3
stochastically bind mother filaments with a rate k_{bind} and unbind with a rate k_{unbind} in a force-
dependent manner given by

1059
$$k_{unbind} = k_{unbind,0} \exp\left(\frac{F}{F_d}\right),$$

1060 where $k_{unbind,0}$ is the unbinding rate in the absence of force, **F** is the force experienced by the
1061 actin-binding protein and F_d is a characteristic detachment force.
1062

1063 **Dynamics of network evolution**
1064 Unbound actin-binding proteins, filament motion, and membrane deformations are governed by
1065 Langevin dynamics. In each time interval Δt, we update the positions of all particles in the system
1066 using

1067
$$x(t + \Delta t) = x(t) + \mu F(t)\Delta t + B(t)\Delta t.$$

Here, μ is the viscosity of the medium. **F**(t) is the sum total of all forces acting on the particle at
time t. **B**(t) is the Brownian noise term, which is a random variable drawn from a normal
distribution with a mean of 0 and a variance of $2\mu k_B T\Delta t$. k_B is the Boltzmann constant and T is
the temperature.
The forces on the membrane are composed of forces from membrane surface tension, bending
rigidity, and point forces from the actin filaments on the membrane. In our formulation, the
speed at which membrane deformation occurs is governed by a membrane deformation
parameter, $\mu_{mobility}$ (not to be confused with the viscosity of the medium), which acts as a
dampener to slow down the deformation of the membrane over time.
1077
1078

Model Assumptions
To capture the length control effect of capping protein in the experiments, we cap filament
growth to a particular filament length. Our current model does not account for filament turnover,
a feature we plan to incorporate in future simulations. Because we are simulating the 2-D
projection of the membrane, we assume that the membrane is in a low-tension regime and any
stretching of the membrane during deformation results from overcoming entropic and enthalpic
cost of unfolding of wrinkles. Membrane surface tension is set to 0.001 mN/m in line with
experimentally reported values (*142*).
For computational cost and technical challenges of performing simulations on the timescales of
the experiments, we assumed faster nucleation rates for mDia1 and pVCA-bound Arp2/3 and
simulated for shorter timescales.
1090

Simulation set-up

Mimicking the instance of asymmetric surface accumulation after illumination, we randomly
placed surface-bound nucleators at the right edge of the model GUV and simulated the system
for a period of 5 s. We examined three cases: one with pVCA and Arp2/3, one with mDia1 only
and one with both NPFs and Arp2/3. For each case, we simulated conditions with and without
inter-filament interactions, giving a total of 6 simulation conditions. We performed 20
independent trajectories per condition.
1098
1099 Table 1. List of Parameters used in the simulations

Parameter	Units	Value(s)	References
r_{eff}	μm	0.035	This study
t_{final}	s	5	This study
Boundary Repulsion	pN/μm	10	This study
r_{cutoff}	μm	0.0035	Based on the radius of an actin filament
Brownian ratchet force for polymerization	pN	3	Footer et al. 2007 (*143*)
Growth rate	μm/s	0.033 (Arp23), 0.167 (Formin)	Kovar 2004 (*144*)
pVCA-Arp2/3 binding rate	s⁻¹	1	Beltzner and Pollard 2008 (*141*)
pVCA-Arp2/3 unbinding rate	s⁻¹	0.001	Beltzner and Pollard 2008 (*141*)
pVCA-Arp2/3 nucleation rate	s⁻¹	5	This study
mDia1 nucleation rate	s⁻¹	10	This study
Final filament length	μm	0.75	This study
Steric strength	pN/μm	0, 10	This study
Membrane surface tension	mN/m	10⁻³	Rawicz et al. 2000 (*142*)
Membrane bending modulus	$k_B T$	20	Drabik et al. 2020 (*145*)
Filament Bending rigidity	pNμm	0.068	Akenuwa and Abel 2023 (*146*)
Number of mDia1, Arp2/3, pVCA		50	This study
Medium viscosity	Pa·s	0.001	

Membrane deformation parameter	Pa·s	1.0	This study

1100

1101

1102 **Quantification of membrane deformation in simulations**

1103 To quantify the extent of actin-induced membrane deformation, we calculated the average
1104 difference between the radius of the GUV and the distance from each node in the leading edge
1105 of the membrane to the center of the membrane at the final timepoint.

$$\Delta L = \frac{1}{M} \sum_{i=1}^{M} (|\vec{r_i}| - r_{GUV}) \ \forall M \in N(leading\ edge)$$

1107

1108

Acknowledgement

We thank Yubin Zhou for TEV protease plasmid; Masato Kanemaki for pET28-2×Strep-tag
plasmid; Miho Iijima for pGEX-2T plasmid; Bin Wu for Phage-ubc-nls-ha-tdMCP-SNAP plasmid;
Roberto Dominguez for pCold-mDia1 plasmid, Saki Takayanagi for constructing GST-pVCA-FKBP-
EYFP plasmid. We also thank Yusuke Sato, Taro Toyota, Pushpendra Singh, Tetsuya Kotani, Takeya
Masubuchi for technical advice on the experiments. We appreciate Erin D. Goley for technical
support of pyrene assay and insightful comments on actin preparation. We also appreciate Julie
Plastino and Cecile Sykes for pET3d-Capβ2(mouse)-Capα1(mouse) plasmid and insightful advice
on protein purification. Our appreciation extends to Pablo A. Iglesias for fruitful discussion. We
thank our lab members, including Allen Kim, Abhijit Deb Roy, Helen D. Wu, Yuta Nihongaki for
constructive discussion. We thank Kyoko Chiba for technical support on protein purification. We
also thank Robert DeRose for proofreading the manuscript and experimental support. FRIS CoRE
(a shared research environment in Tohoku University) is also acknowledged. This study was
supported by the PRESTO program (JPMJPR12A5 to TI, JPMJPR20KA to HTM), CREST program
(JPMJCR19S5 to MM), and FOREST program (JPMJFR2416 to HTM) of the Japan Science and
Technology Agency, MEXT/JSPS KAKENHI Grants (23K14150, 23H04397, 25K02234, 25H02347 to
HTM), National Institutes of Health (5R01GM123130, R01GM136858, R35GM149329 to TI, and
R35GM128786 to BC), ExCELLS project research (25EXC604-6 to HTM) and the National Science
Foundation (000819255 to TI, JL, BC). TI was also supported by Nitto Denko Corp. HTM was
supported by Postdoctoral Fellowships from the Japan Society for the Promotion of Science. SR
was supported by the National Science Foundation and the Japan Society for the Promotion of
Science (JSPS) East Asia and Pacific Summer Institutes and Johns Hopkins Lucille Elizabeth Hay
graduate fellowships. Computer simulations were run on hardware hosted by the Triton Shared

Computing Cluster (*147*). OHA and CTL further acknowledge high performance computing
support from the UCSD Physics Computing Facility.
1134
1135

Author contributions

HTM, SR, and TI conceived the project. HTM performed most of the experiments and analyzed
the data with contributions from SR, HN, BC, SM, SMN and TI. SR established GUV preparation
method. WR performed in-cell CID experiment and analyzed the data. RY performed biochemical
experiments. YOT performed EM under supervision by MM. OHA and CTL built and analyzed
computational models. DN, DAK, and BC provided cytoskeleton proteins. TM provided
benzylguanine-conjugated lipid. HTM wrote the manuscript in consultation with TI. HTM, SR, and
TI edited the manuscript. All the authors contributed to the final version of the manuscript.
1144

Data Availability

1146   The datasets generated during this study are available from the corresponding author upon
1147   request.
1148
1149

Figure Legends

1151

Fig. 1. Reversible and asymmetric control of protein localization with iLID-SspB within GUVs.
(A) Schematic representation of iLID-SspB reaction with global light stimulation. iLID was tethered
on the membrane via the interaction between SNAP-tag and benzylguanine-conjugated lipids.
Upon blue light illumination, Jα helix of iLID extends to expose the binding site for SspB, whereas
in the dark, iLID reverses to its closed state and releases SspB. (B) Representative images of iLID-
YFP-SNAP and mCherry-SspB. Scale bar, 10 μm. Blue bar indicates the period of blue light
illumination. (C) Time course of membrane/lumen ratio of mCherry-SspB signal (n=5). Error bars
indicate 95% CI. Blue area indicates the period of blue light stimulation. (D) Schematic
representation of iLID-SspB reaction with local light stimulation. (E) Representative images of
mCherry-SspB. Scale bar, 10 μm. Yellow box indicates the area of blue light illumination. (F)
Membrane/lumen ratio of mCherry-SspB for the representative image shown. The distance is
measured from the center of blue light illumination along the perimeter of the GUV. (G)
Representative image and kymograph of repetitive and reversible SspB translocation. Kymograph
shows membrane signal of mCherry-SspB.
1166

Fig. 2. Reversible light control of actin cytoskeleton within GUVs. (A) Bulk pyrene actin polymerization assay with 1 µM actin (5% pyrene-labeled), 10 nM Arp2/3, and 100 nM NPFs. (-NPF) represents actin and Arp2/3 only condition. (B) Schematic representation of light-inducible actin polymerization. Translocation of pVCA-SspB-mCherry increases local concentration of pVCA on the membrane and enhances actin polymerization. (C) Representative images of pVCA-SspB-mCherry and Alexa 647 actin. Blue bar indicates the period of blue light illumination. (D) Time course of membrane/lumen ratio of mCherry and Alexa 647 actin in (C) and fig. S9. pVCA +Arp2/3+Light: full set of components (n=4). -Light: no blue light stimulation (n=3). -pVCA: pVCA-SspB-mCherry was replaced with mCherry-SspB (n=4). -Arp2/3: Arp2/3 was omitted (n=4). In (E)– (H), pVCA-SspB-mCherry was constitutively anchored to the membrane via SsrA-SNAP. (E) Pharmacological perturbation of membrane-associated actin. DMSO (n=8), Jasplakinolide (Jasp, n=10), Latrunculin A (LatA, n=9). (F) Time course of normalized membrane/lumen ratio of Alexa-647 actin in (E). The values were normalized by time=0. (G) FRAP analysis of membrane fluorescence of mCherry and Alexa 647 actin in fig. S11. The values were normalized by time=0. Sample sizes are listed in fig. S11A. (H) Electron microscopy (EM) images of actin filaments in the presence and absence of pVCA. Scale bar indicates 500 nm. (I) Representative images of reversible actin polymerization. Blue bars indicate the period of blue light illumination. (J) Time course of membrane/lumen ratio of pVCA-SspB-mCherry and Alexa 647-labeled actin in (F). (D, F, G) Error bars indicate 95% CI.

Fig. 3. Asymmetric light control of actin cytoskeleton within GUVs. (A) Schematic representation of iLID-SspB reaction with local light stimulation. (B, D) Representative images of light-inducible asymmetric actin polymerization. Yellow boxes indicate the area of blue light illumination. (C, E) Kymograph of membrane pVCA-SspB and actin signals. Blue bars indicate the period of blue light illumination. Triangles indicate timepoints represented in (B, D).

Fig. 4. mDia1-mediated actin polymerization drives protrusive movement of GUVs. (A) Schematic representation of light-inducible membrane protrusion driven by pVCA and mDia1-mediated actin polymerization. (B) Representative images of pVCA-mDia1-mediated movement of GUVs. Yellow boxes indicate the area of blue light stimulation. Dotted line shows the initial position of the GUV. (C) Representative images of iLID-EYFP-SNAP showing the increase in adhesive area and vesicle deformation into dome-like shape. X-y images show the confocal plane right above the bottom substrate. Z-x and y-z images are reconstituted from z-stack. Dotted line shows the adhesive area of the previous images. (D) Color-coded boundaries of GUVs. The data correspond as follows. Left: Fig. 4B. Middle: fig. S16A. Right: fig. S16B. (E, F) Kymographs of membrane signal of mCherry and actin Alexa-647 and membrane protrusions. Triangles in (E) indicate the timepoints represented in (B, C). Data of (E) and (F) correspond to Fig. 4 (B) and fig. S16A, respectively. (G) Distance the front (illuminated) side of GUV membrane moved forward.

pVCA+mDia1 (1 μM GST-pVCA-SspB-mCherry + 1 μM mCherry-SspB-mDia1, n=14 vesicles),
mDia1 (1 μM mCherry-SspB-mDia1, n=12 vesicles), pVCA (1 μM GST-pVCA-SspB-mCherry, n=12
vesicles). Error bars indicate 95% CI (H, I) Quantification of the initial velocity and the distance
the front side of GUV membrane moved forward. P values of Steel-Dwass test (two sided) are
indicated. pVCA + mDia1: n=22 vesicles. mDia1: n=11 vesicles. pVCA: n=12 vesicles. Box whisker
plots represent median, 1st, 3rd quartiles and 1.5×inter-quartile range. (J̶H̶) Distance the front
(illuminated) side of GUV membrane moved forward. DMSO (n=9 vesicles), 30 μM LatA
(Latrunculin A) (n=10 vesicles). DMSO, ΔDAD: mCherry-SspB-mDia1 (FH1-FH2-DAD) was replaced
with mCherry-SspB-mDia1(FH1-FH2) (n=13 vesicles). 30 μM SMIFH2 (n=6 vesicles). Error bars
indicate 95% CI. (H, I) Quantification of the initial velocity and the distance the front side of GUV
membrane moved forward. P values of Steel-Dwass test (two sided) are indicated. pVCA + mDia1:
n=22 vesicles. mDia1: n=11 vesicles. pVCA: n=12 vesicles. Box whisker plots represent median,
1st, 3rd quartiles and 1.5×inter-quartile range. (K) A GUV deformed into dome-like shape after
movement. Upper images: before light stimulation. Lower images: after light stimulation. The
data corresponds to fig. S16C.
1220
1221 **fig. S1. Proteins purified in this study.** (A) SDS-PAGE results of purified proteins. Blue: Coomassie
1222 brilliant blue staining. Gray: SYPRO Ruby staining. Most mCherry-tagged proteins show two extra
1223 bands due to a known chromophore cleavage reaction during the boiling process (*148*, *149*). (B)
1224 Domain structures of NPFs used in this study.
1225
fig. S2. Optimization of membrane anchoring of iLID. (A),(D),(G),(I): Schematic representations
of membrane anchoring of iLID. (B),(C),(E),(F),(H),(J): Confocal images of iLID variants and
mCherry-SspB. (B) His-tag Ni-NTA pair caused non-specific membrane recruitment of SspB. This
is possibly caused by the presence of 2×Strep-iLID-YFP-6×His because mCherry-SspB itself did not
show non-specific binding to the membrane of 5%Ni-NTA DGS/95%POPC (C). (E),(F) MARCKS-ED
peptide requires high percentage of PS and PEGylated-lipids to reduce non-specific binding of
SspB. (H) Biotin-StrepTactin-Streptag also caused non-specific membrane binding of SspB. (J) The
data correspond with Fig. 1B.
1234
fig. S3. Asymmetric control of protein localization with iLID-SspB within GUVs. (A) Time course
of membrane/lumen ratio of mCherry-SspB signal in the illuminated area (n=6). Error bars
indicate 95% CI. Blue area indicates the time window of blue light stimulation. The data
corresponds to Fig. 1E and F. (B, C) Confocal images and time course of membrane/lumen ratio
of mCherry-SspB signal responding to repetitive local light illuminations. The data corresponds
to Fig. 1G.
1241

fig. S4. Optimization of ActA-SspB. (A) Pyrene actin polymerization assay. The reaction was performed with 1 µM actin (5% pyrene labeled), 10 nM Arp2/3, and 100 nM ActA variants or GST-VCA. (B) Fluorescence intensity/time in 0–6 min of (A) was plotted as bar chart. (C–E) Light-inducible local membrane recruitment of ActA (C) Confocal images of ActA translocation. Yellow boxes indicate the area of blue light illumination. (D) Time course of membrane/lumen ratio of ActA-SspB variants (n≥5 for each variant). Error bars indicate 95% CI. Blue bar indicates the time window of blue light stimulation. (E) Average membrane/lumen ratio of mCherry signal. In each condition, left and right bars indicate before and after light stimulation, respectively. The numbers on the bar chart indicate the fold change after stimulation.

fig. S5. Light-inducible actin polymerization with global ActA membrane recruitment. iLID-EYFP-MARCKS (5 µM), ActA-SspBnano-mCherry (2 µM), actin (1.5 µM, 10% Alexa-647 labeled), and Arp2/3 (150 nM) were encapsulated in GUVs containing 60 mol% POPC/40 mol% POPS. Blue light was applied to the entire GUV. (A) Representative images of ActA(1–183)-SspB(nano)-mCherry and actin Alexa-647. All components (n=4): w/o ActA: ActA(1–183)-SspB(nano)-mCherry was replaced with mCherry-SspB(micro) (n=3), w/o Arp2/3: Arp2/3 was omitted (n=3). No Light: no blue light stimulation (n=3). (B, C) Time course of membrane/lumen ratio of mCherry and actin Alexa 647. (D) Bar chart of membrane/lumen ratio of actin Alexa 647 before (0 min) and after (40 min) light stimulation. Error bars indicate 95% CI.

fig. S6. Asymmetric ActA membrane recruitment fails to induce directed actin polarization toward local light illumination. iLID-EYFP-MARCKS (5 µM), ActA-SspBnano-mCherry (2 µM), actin (1.5 µM, 10% Alexa-647 labeled), and Arp2/3 (150 nM), 1 mM ATP were encapsulated in GUVs containing 60 mol% POPC/40 mol% POPS. Yellow boxes indicate the area of blue light illumination. Although ActA creates asymmetric distribution, actin patches randomly distributed due to the diffusion on the membrane.

fig. S7. Global ActA membrane recruitment fails to induce actin polymerization on the membrane in the presence of profilin, cofilin, and capping protein. 15 µM iLID-YFP-MARCKS, 5 µM ActA(1-183)-SspB(nano)-mCherry, 7.5 µM actin (10% Alexa 647 labeled), 150 nM Arp2/3, 3 µM profilin, cofilin, and capping protein were encapsulated in GUVs containing 60 mol% POPC/40 mol% POPS. Blue bars indicate the period of blue light illumination.

fig. S8. Biochemical characterization of actin and actin-binding proteins used in this study. (A, B) Ultracentrifuge assay. After 30 min actin polymerization reaction at RT, the solutions were ultracentrifuged 100,000×g, 60 min, at RT. Proteins and concentrations used: 7.5 µM actin (10% Alexa-488 labeled), 3 µM profilin, 4 µM cofilin, 50 nM capping protein, 150 nM Arp2/3, 1 µM GST-pVCA-SspB-mCherry, 1 µM mCherry-SspB-mDia1. G-buffer: 2 mM Tris-HCl (pH 7.5), 0.2 mM

ATP, 0.1 mM $CaCl_2$, 0.5 mM DTT, 1 mM NaN_3. KMEI-S buffer: .10mM imidazole (pH 7.0), 50mM

KCl, 1mM $MgCl_2$, 1mM EGTA, 240 mM sucrose (sucrose was added to mimic the buffer conditions

in GUV experiments). (C–F) Pyrene actin polymerization assay. The reactions were performed

with 2 μM actin (5% pyrene labeled) at RT. (C, D) 10 nM Arp2/3, and 100 nM NPFs. (-) represents

actin and Arp2/3 only condition. (E) The reactions were performed with the indicated factors. (F)

50 nM capping protein, 150 nM Arp2/3, 3 μM profilin, 1 μM mCherry-SspB-mDia1, 1 μM GST-

pVCA-SspB-mCherry. (-) represents actin only condition.

1287

fig. S9. Light control of actin cytoskeleton with pVCA-SspB-mCherry. The data correspond to Fig.

2C–E. (+) condition contains 15 μM iLID-YFP-MARCKS, 4.2 μM pVCA-SspB(nano)-mCherry, 7.5 μM

actin (10% Alexa 647 labeled), 150 nM Arp2/3, 100 nM capping protein, 3 μM profilin, 4 μM cofilin,

and 1 mM ATP. In (-light) and (-Arp2/3) conditions, blue light illumination or Arp2/3 was omitted

from (+pVCA +Arp2/3 +Light) condition, respectively. In (-pVCA) condition, pVCA-SspB(nano)-

mCherry was replaced with mCherry-SspB(micro).

1294

fig. S10. Pharmacological perturbation of membrane-associated actin assemblies. (A)

Schematic representation of constitutive tethering of NPFs. SspB-fused NPFs (pVCA and/or

mDia1) were tethered to the GUV membrane via the SsrA–SNAP/BG-lipid system. (B)

Representative confocal images of membrane-associated actin assemblies formed by pVCA,

mDia1, or their combination. Scale bars, 5 μm. (C) Quantification of the initial membrane/lumen

fluorescence intensity ratio for mCherry and actin Alexa-647. pVCA: n=27 vesicles. mDia1: n=23

vesicles. pVCA + mDia1: n=27 vesicles. Box whisker plots represent median, 1st, 3rd quartiles and

1.5×inter-quartile range. (D) Representative images before (0 min) and after (24 min) treatment

with DMSO, 50 μM Latrunculin A, or 50 μM Jasplakinolide. Scale bars, 5 μm. (E) Time courses of

membrane/lumen ratio and normalized membrane/lumen ratio of actin Alexa-647. (F) Time

courses of normalized membrane/lumen ratio of actin Alexa-647 for each NPF condition. The y-

axis is shown on a logarithmic scale. (E, F) Error bars indicate 95% CI. (G) Representative images

showing membrane deformation after Jasplakinolide treatment. Scale bars, 10 μm. mCherry-

SspB-mDia1-6×His was used as mDia1 in this figure.

1309

fig. S11. Fluorescence recovery after photobleaching (FRAP) analysis of membrane-associated

NPFs and actin. (A) Summary table of proteins included in each conditions. In FRAP analysis, SspB-

fused NPFs (pVCA and/or mDia1) were tethered to the GUV membrane via the SsrA–SNAP/BG-

lipid system. (B, C) Representative confocal images of mCherry-tagged NPFs and actin Alexa-647.

The white dashed box indicates the photobleached region. Scale bars, 5 μm. (D, E) Time course

of mCherry and Alexa-647 fluorescence recovery at the bleached region. Fluorescence intensities

were normalized to those of a non-bleached region to compensate for photobleaching during

image acquisition. The data from Fig. 1G are reproduced to facilitate direct comparison with the
other conditions presented in this figure.

1319

fig. S12. Electron microscopy (EM) analysis of actin filaments. Ultrastructure of small unilamellar
vesicles (SUVs) and actin filaments were analyzed by negative staining electron microscopy. (A)
Representative EM images of SUVs alone. (B) Representative images of Y-junctions formed in the
presence of pVCA, Arp2/3, capping protein (CP), profilin, and cofilin. (C) Negative control sample
lacking pVCA (Arp2/3, CP, profilin, and cofilin only). (B, C) To reduce filament overlap, the
polymerization reactions were diluted 10-fold prior to deposition onto EM grids. (D)
Representative EM images of actin assemblies formed with pVCA or the corresponding negative
control lacking pVCA. (E) Representative EM images of actin assemblies formed with pVCA +
mDia1, mDia1 alone, or the corresponding negative control lacking mDia1. All samples were
deposited onto EM grids and negatively stained prior to imaging. Scale bars are shown in each
panel. mCherry-SspB-mDia1-6×His was used as mDia1 in this figure.

1331

1332   **fig. S13. Representative images of reversible actin polymerization.** Confocal images and time
1333   course of membrane/lumen ratio of pVCA-SspB-mCherry and Alexa 647-labeld actin. Blue bars
1334   on images and blue regions in graphs indicate timepoints with blue light illumination. This figure
1335   corresponds with Fig. 2F, G.

1336

fig. S14. Asymmetric actin polymerization induced by pVCA. (A) Confocal images of local pVCA
recruitment and actin polymerization. Yellow boxes indicate areas of blue light illumination. (B)
Membrane/lumen ratio of mCherry-tagged pVCA and actin Alexa-647 for the images in (A). The
distance is measured from the center of blue light illumination along the perimeter of the GUV.
(A, B) The data corresponds to Fig. 3B and C. (C) Confocal images of local pVCA recruitment
accompanied by vesicle movement toward light stimulation. Gray scaled images were created by
averaging pVCA and actin images. Dotted circles indicate positions of vesicles in the images of
one above.

1345

fig. S15. Quantitative analysis of mDia1-induced filopodia formation in Cos7 cells. (A) Schematic
representation of chemically-inducible mDia1 translocation and membrane protrusions. (B, C)
Confocal images of plasma membrane recruitment of mDia1 and subsequent filopodia formation.
Cos7 cells were transfected with Lyn-ECFP-FRB, Lifeact-mCherry, and either of YFP-FKBP or YFP-
FKBP-mDia1. At time 0, 100 nM rapamycin was added. Green: YFP-FKBP or YFP-FKBP-mDia1.
Magenta: Lifeact-mCherry. (B) YFP-FKBP control. (C) YFP-FKBP-mDia1. (D) Zoomed-up view of
filopodia extension. Images were acquired every 1.5 minutes. The bottom is merged image. (E)
Quantification of mDia1-induced filopodia. Number of filopodia whose length is longer than 15
µm was counted. Box whisker plots represent median, 1st, 3rd quartiles and 1.5×inter-quartile

range. P-values: ****: < 0.0001. Wilcoxon rank sum test. YF (YFP-FKBP): n=36 cells. YF-mDia1 (YFP-FKBP-mDia1): n=33 cells. (F–H) Quantification of the phenotypes observed after plasma membrane translocation of mDia1 (n=159 cells), pVCA (n=142 cells), Tiam1 (Rac1 GEF) (n=67 cells), and control protein YFP-FKBP (n=44 cells).

fig. S16. mDia1-mediated GUV movement and the comparison between mDia1 and pVCA + mDia1 system. (A) Representative images of mDia1-mediated GUV movement. Yellow boxes indicate the area of blue light stimulation. Dotted lines indicate the initial position of the GUV. (B, C) Time course of the distance the front (illuminated) side of GUV membrane moved forward. Each line represents each GUV. Insets show the initial response.

fig. S17. Local protrusions in pVCA + mDia1 GUVs. Representative images of local protrusions seen in pVCA + mDia1 GUVs. Yellow boxes indicate the area of blue light stimulation. Dotted lines indicate the initial position of the GUV. White arrows indicate where local protrusions arose.

fig. S18. Lack of vesicle movement and shape change with mCherry-SspB. With pVCA-SspB-mCherry and mCherry-SspB-mDia1 being replaced with mCherry-SspB, GUVs neither moved nor changed their shape after blue light illumination. (A) Time course of the distance the front (illuminated) side of GUV membrane moved forward. pVCA + mDia1 GUVs: n=6 vesicles. SspB GUVs: n=4 vesicles. Error bars indicate 95% CI. (B) 3D reconstitution of iLID-YFP-SNAP images before and after blue light illumination.

Fig. S19. Relationship between vesicle motion and biophysical parameters. (A) Relationship between initial velocity (average velocity over the first 5 minutes) and net displacement after 30 minutes of GUV forward movement. (B, C) Biophysical parameters are plotted against initial velocity and displacement are plotted in (B) and (C), respectively. R and p indicate Pearson's correlation coefficient and its p-value.

fig. S20. Representative images of pVCA-mDia1-mediated movement of GUVs. (A, B) Representative images of pVCA-mDia1-mediated movement of GUVs. Yellow boxes indicate the area of blue light stimulation. Dotted lines indicate the initial position of the GUV. White arrows indicate where local protrusions arose. (C) Distance the front side of GUV membrane moved forward. Data was quantified from the images of (B). Blue areas indicate the periods of blue light stimulation.

Fig. S21. Computational modeling of actin network dynamics underlying membrane deformation. (A, B) Snapshots showing actin filaments at the leading edge of the membrane after 5 s of simulation. Actin filaments are shown in green, pVCA-Arp2/3 is shown in red, mDia1 is

shown in orange, and Arp2/3 is shown in magenta. The membrane location at t = 0 is shown with
the cyan dotted line to highlight the extent of membrane deformation. Simulations without and
with inter-filament steric interactions are shown in A and B, respectively. (C, D) Box-whisker plots
showing distributions of the membrane extension for the different actin-binding proteins. Results
for networks without and with inter-filament steric interactions are shown in A and B,
respectively. Each box plot is constructed using 20 independent simulation trajectories. Error
bars represent standard deviation about the mean.
1400
1401
1402 **Movie S1**
1403 Representative movie of repetitive and reversible SspB translocation. The data corresponds to
1404 Fig. 1G. Time in min:sec. Yellow regions indicate the area of blue light stimulations.
1405
1406 **Movie S2**
1407 A rare example where ActA induced actin polymerization in the direction of light stimulus. The
1408 data corresponds the leftmost images in Fig. S6. Red: ActA-SspB(nano)-mCherry. Magenta: Alexa
1409 647-labeled actin. Time in hr:min:sec. Yellow regions indicate the area of blue light stimulation.
1410
1411 **Movie S3**
1412 Representative movie of global membrane recruitment of pVCA-SspB-mCherry. The data
1413 corresponds to Fig. 2C and "pVCA +Arp2/3 +Light" in Fig. S9. Red: pVCA-SspB-mCherry. Magenta:
1414 Alexa 647-labeled actin. Time in hr:min:sec. The label "Light" indicates when blue light
1415 stimulation is applied to the entire GUV.
1416
1417 **Movie S4**
1418 Representative movie of global and reversible membrane recruitment of pVCA-SspB-mCherry.
1419 The data corresponds to Fig. 2IF and S13. Red: pVCA-SspB-mCherry. Magenta: Alexa 647-labeled
1420 actin. Time in hr:min:sec. The label "Light" indicates when blue light stimulation is applied to the
1421 entire GUV.
1422
1423 **Movie S5**
1424 Representative movie of asymmetric membrane recruitment of pVCA-SspB-mCherry. The data
1425 corresponds to Fig. 3B and C. Red: pVCA-SspB-mCherry. Magenta: Alexa 647-labeled actin. Time
1426 in hr:min:sec. Yellow regions indicate the area of blue light stimulation.
1427
1428 **Movie S6**

1429   Representative movie of asymmetric and reversible membrane recruitment of pVCA-SspB-
1430   mCherry. The data corresponds to Fig. 3D and E. Red: pVCA-SspB-mCherry. Magenta: Alexa 647-
1431   labeled actin. Time in hr:min:sec. Yellow regions indicate the area of blue light stimulation.
1432

1433   **Movie S7**

1434   Representative movie of pVCA-mDia1-mediated movement of GUV. The data corresponds to Fig.
1435   4B and E. Red: pVCA-SspB-mCherry and mCherry-SspB-mDia1. Magenta: Alexa 647-labeled actin.
1436   Time in hr:min:sec. Yellow regions indicate the area of blue light stimulation.
1437

1438   **Movie S8**

1439   Representative movie of local protrusions seen in pVCA + mDia1 GUVs. The data corresponds to
1440   Fig. S17. Red: pVCA-SspB-mCherry and mCherry-SspB-mDia1. Magenta: Alexa 647-labeled actin.
1441   Time in hr:min:sec. Yellow regions indicate the area of blue light stimulation. In the zoomed-in
1442   view, the relative movement of the GUV was subtracted to better highlight the local protrusions.
1443

1444   **Movie S9**

1445   Representative movie of pVCA-mDia1-mediated movement of GUV. The data corresponds to Fig.
1446   4F and S20A. Red: pVCA-SspB-mCherry and mCherry-SspB-mDia1. Magenta: Alexa 647-labeled
1447   actin. Time in hr:min:sec. Yellow regions indicate the area of blue light stimulation.
1448

1449   **Movie S10**

1450   Representative movie of pVCA-mDia1-mediated movement of GUV. The data corresponds to Fig.
1451   S20B. Red: pVCA-SspB-mCherry and mCherry-SspB-mDia1. Magenta: Alexa 647-labeled actin.
1452   Time in hr:min:sec. Yellow regions indicate the area of blue light stimulation.
1453

1454   **Movie S11**

1455   Representative simulation movies in the absence of inter-filament steric interactions. Actin
1456   filaments are shown in green, pVCA–Arp2/3 in red, mDia1 in orange, and Arp2/3 in magenta. The
1457   membrane position at $t = 0$ is marked by a cyan line to indicate the extent of membrane
1458   deformation.
1459

1460   **Movie S12**

1461   Representative simulation movies in the presence of inter-filament steric interactions. Actin
1462   filaments are shown in green, pVCA–Arp2/3 in red, mDia1 in orange, and Arp2/3 in magenta. The
1463   membrane position at $t = 0$ is marked by a cyan line to indicate the extent of membrane
1464   deformation.

References

1465

1. M. Miyata, R. C. Robinson, T. Q. P. Uyeda, Y. Fukumori, S. ichi Fukushima, S. Haruta, M.
Homma, K. Inaba, M. Ito, C. Kaito, K. Kato, T. Kenri, Y. Kinosita, S. Kojima, T. Minamino, H.
Mori, S. Nakamura, D. Nakane, K. Nakayama, M. Nishiyama, S. Shibata, K. Shimabukuro, M.
Tamakoshi, A. Taoka, Y. Tashiro, I. Tulum, H. Wada, K. ichi Wakabayashi, Tree of motility – A
proposed history of motility systems in the tree of life. *Genes to Cells* **25**, 6–21 (2020).

2. K. Y. Wan, G. Jékely, Origins of eukaryotic excitability. *Phil. Trans. R. Soc. B* **376**, 20190758
(2021).

3. T. Froese, N. Virgo, T. Ikegami, Motility at the Origin of Life: Its Characterization and a
Model. *Artificial Life* **20**, 55–76 (2014).

4. P. N. Devreotes, S. Bhattacharya, M. Edwards, P. A. Iglesias, T. Lampert, Y. Miao, Excitable
Signal Transduction Networks in Directed Cell Migration. *Annu. Rev. Cell Dev. Biol.* **33**, 103–
125 (2017).

5. L. K. Fritz-Laylin, The evolution of animal cell motility. *Current Biology* **30**, R477–R482
(2020).

6. S. SenGupta, C. A. Parent, J. E. Bear, The principles of directed cell migration. *Nat Rev Mol
Cell Biol* **22**, 529–547 (2021).

7. Effects of Cytochalasins on Mammalian Cells | Nature.
https://www.nature.com/articles/213261a0.

8. T. D. Pollard, S. Ito, CYTOPLASMIC FILAMENTS OF *AMOEBA PROTEUS*. *The Journal of Cell
Biology* **46**, 267–289 (1970).

9. J. V. Small, G. Isenberg, J. E. Celis, Polarity of actin at the leading edge of cultured cells.
Nature **272**, 638–639 (1978).

10. L. G. Tilney, S. Inoué, Acrosomal reaction of Thyone sperm. II. The kinetics and possible
mechanism of acrosomal process elongation. *J Cell Biol* **93**, 820–827 (1982).

11. J. A. Theriot, The Polymerization Motor. *Traffic* **1**, 19–28 (2000).

12. T. D. Pollard, Landmarks in the discovery of a role for actin in cell locomotion. *Molecular
Biology of the Cell*, 4 (2022).

13. P. Lappalainen, T. Kotila, A. Jégou, G. Romet-Lemonne, Biochemical and mechanical
regulation of actin dynamics. *Nat Rev Mol Cell Biol*, doi: 10.1038/s41580-022-00508-4
(2022).

14. T. Lämmermann, M. Sixt, Mechanical modes of 'amoeboid' cell migration. *Current Opinion*
in Cell Biology **21**, 636–644 (2009).

15. P. Friedl, K. Wolf, Plasticity of cell migration: a multiscale tuning model. *Journal of Cell*
Biology **188**, 11–19 (2010).

16. M. Garrido-Casado, G. Asensio-Juárez, M. Vicente-Manzanares, Nonmuscle Myosin II
Regulation Directs Its Multiple Roles in Cell Migration and Division. *Annu Rev Cell Dev Biol*
37, 285–310 (2021).

17. D. Knecht, W. Loomis, Antisense RNA inactivation of myosin heavy chain gene expression in
Dictyostelium discoideum. *Science* **236**, 1081–1086 (1987).

18. A. De Lozanne, J. A. Spudich, Disruption of the Dictyostelium myosin heavy chain gene by
homologous recombination. *Science* **236**, 1086–1091 (1987).

19. A. B. Verkhovsky, T. M. Svitkina, G. G. Borisy, Self-polarization and directional motility of
cytoplasm. *Current Biology* **9**, 11-S1 (1999).

20. R. J. Eddy, L. M. Pierini, F. Matsumura, F. R. Maxfield, Ca2+-dependent myosin II activation
is required for uropod retraction during neutrophil migration. *J Cell Sci* **113 (Pt 7)**, 1287–
1298 (2000).

21. T. Lämmermann, B. L. Bader, S. J. Monkley, T. Worbs, R. Wedlich-Söldner, K. Hirsch, M.
Keller, R. Förster, D. R. Critchley, R. Fässler, M. Sixt, Rapid leukocyte migration by integrin-
independent flowing and squeezing. *Nature* **453**, 51–55 (2008).

22. A. Reversat, F. Gaertner, J. Merrin, J. Stopp, S. Tasciyan, J. Aguilera, I. de Vries, R. Hauschild,
M. Hons, M. Piel, A. Callan-Jones, R. Voituriez, M. Sixt, Cellular locomotion using
environmental topography. *Nature* **582**, 582–585 (2020).

23. Y. I. Wu, D. Frey, O. I. Lungu, A. Jaehrig, I. Schlichting, B. Kuhlman, K. M. Hahn, A genetically
encoded photoactivatable Rac controls the motility of living cells. *Nature* **461**, 104–108
(2009).

24. B. Lin, W. R. Holmes, C. J. Wang, T. Ueno, A. Harwell, L. Edelstein-Keshet, T. Inoue, A.
Levchenko, Synthetic spatially graded Rac activation drives cell polarization and movement.
Proceedings of the National Academy of Sciences of the United States of America **109**
(2012).

25. S. Nanda, A. Calderon, A. Sachan, T.-T. Duong, J. Koch, X. Xin, D. Solouk-Stahlberg, Y.-W. Wu,
P. Nalbant, L. Dehmelt, Rho GTPase activity crosstalk mediated by Arhgef11 and Arhgef12
coordinates cell protrusion-retraction cycles. *Nat Commun* **14**, 8356 (2023).

26. L. A. Cameron, M. J. Footer, A. Van Oudenaarden, J. A. Theriot, Motility of ActA protein-
coated microspheres driven by actin polymerization. *Proceedings of the National Academy*
of Sciences of the United States of America **96**, 4908–4913 (1999).

27. T. P. Loisel, R. Boujemaa, D. Pantaloni, M. Carlier, Reconstitution of actin-based motility of
Listeria and Shigella using pure proteins. **401**, 613–616 (1999).

28. A. Bernheim-Groswasser, S. Wiesner, R. M. Golsteyn, M.-F. Carlier, C. Sykes, The dynamics
of actin-based motility depend on surface parameters. *Nature* **417**, 308–311 (2002).

29. A. Mogilner, G. Oster, Force generation by actin polymerization II: The elastic ratchet and
tethered filaments. *Biophysical Journal* **84**, 1591–1605 (2003).

30. S. Wiesner, E. Helfer, D. Didry, G. Ducouret, F. Lafuma, M. F. Carlier, D. Pantaloni, A
biomimetic motility assay provides insight into the mechanism of actin-based motility.
Journal of Cell Biology **160**, 387–398 (2003).

31. S. Romero, C. Le Clainche, D. Didry, C. Egile, D. Pantaloni, M. F. Carlier, Formin is a
processive motor that requires profilin to accelerate actin assembly and associated ATP
hydrolysis. *Cell* **119**, 419–429 (2004).

32. C. Co, D. T. Wong, S. Gierke, V. Chang, J. Taunton, Mechanism of Actin Network Attachment
to Moving Membranes: Barbed End Capture by N-WASP WH2 Domains. *Cell* **128**, 901–913
(2007).

33. O. Akin, R. D. Mullins, Capping Protein Increases the Rate of Actin-BasedMotility by
Promoting Filament Nucleation by the Arp2/3 Complex. *Cell* **133**, 841–851 (2008).

34. J. Prost, J.-F. Joanny, P. Lenz, C. Sykes, "The Physics Of Listeria Propulsion" in *Cell Motility*
(Springer New York, New York, NY, 2008; https://doi.org/10.1007/978-0-387-73050-9_1),
pp. 1–30.

35. P. Bieling, S. D. Hansen, O. Akin, T.-D. Li, C. C. Hayden, D. A. Fletcher, R. D. Mullins, WH2 and
proline-rich domains of WASP-family proteins collaborate to accelerate actin filament
elongation. *EMBO J* **37**, 102–121 (2018).

36. C. Guérin, A.-B. N'Diaye, L. Gressin, A. Mogilner, M. Théry, L. Blanchoin, A. Colin, Balancing
limited resources in actin network competition. *Current Biology* **35**, 500-513.e5 (2025).

37. S. Razavi, T. Inoue, "Chapter 22 - Reconstitution of membrane symmetry breaking" in
Plasma Membrane Shaping, S. Suetsugu, Ed. (Academic Press, 2023;
https://www.sciencedirect.com/science/article/pii/B9780323899116000273), pp. 333–353.

38. R. Lopes Dos Santos, C. Campillo, Studying actin-induced cell shape changes using Giant
Unilamellar Vesicles and reconstituted actin networks. *Biochemical Society Transactions* **50**,
1527–1539 (2022).

39. F. C. Keber, E. Loiseau, T. Sanchez, S. J. DeCamp, L. Giomi, M. J. Bowick, M. C. Marchetti, Z.
Dogic, A. R. Bausch, Topology and dynamics of active nematic vesicles. *Science* **345**, 1135–
1139 (2014).

40. Y. Sato, Y. Hiratsuka, I. Kawamata, S. Murata, S. M. Nomura, Micrometer-sized molecular
robot changes its shape in response to signal molecules. *Science Robotics* **2**, eaal3735
(2017).

41. K. Gavriljuk, B. Scocozza, F. Ghasemalizadeh, H. Seidel, A. P. Nandan, M. Campos-Medina,
M. Schmick, A. Koseska, P. I. H. Bastiaens, A self-organized synthetic morphogenic liposome
responds with shape changes to local light cues. *Nature Communications* **12**, 1548 (2021).

42. J. D. Cortese, B. Schwab, C. Frieden, E. L. Elson, Actin polymerization induces a shape
change in actin-containing vesicles. *Proceedings of the National Academy of Sciences of the
United States of America* **86**, 5773–5777 (1989).

43. H. Miyata, H. Hotani, Morphological changes in liposomes caused by polymerization of
encapsulated actin and spontaneous formation of actin bundles. *Proceedings of the
National Academy of Sciences of the United States of America* **89**, 11547–11551 (1992).

44. H. Miyata, S. Nishiyama, K. -i. Akashi, K. Kinosita, Protrusive growth from giant liposomes
driven by actin polymerization. *Proceedings of the National Academy of Sciences* **96**, 2048–
2053 (1999).

45. M. Honda, K. Takiguchi, S. Ishikawa, H. Hotani, Morphogenesis of liposomes encapsulating
actin depends on the type of actin-crosslinking. *Journal of Molecular Biology* **287**, 293–300
(1999).

46. L. L. Pontani, J. Van Der Gucht, G. Salbreux, J. Heuvingh, J. F. Joanny, C. Sykes,
Reconstitution of an actin cortex inside a liposome. *Biophysical Journal* **96**, 192–198 (2009).

47. K. Dürre, F. C. Keber, P. Bleicher, F. Brauns, C. J. Cyron, J. Faix, A. R. Bausch, Capping protein-
controlled actin polymerization shapes lipid membranes. *Nature Communications*, doi:
10.1038/s41467-018-03918-1 (2018).

48. Y. Bashirzadeh, S. A. Redford, C. Lorpaiboon, A. Groaz, H. Moghimianavval, T. Litschel, P.
Schwille, G. M. Hocky, A. R. Dinner, A. P. Liu, Actin crosslinker competition and sorting drive
emergent GUV size-dependent actin network architecture. *Commun Biol* **4**, 1136 (2021).

49. N. H. Wubshet, Y. Bashirzadeh, A. P. Liu, Fascin-induced actin protrusions are suppressed by
dendritic networks in giant unilamellar vesicles. *MBoC* **32**, 1634–1640 (2021).

50. T. Litschel, C. F. Kelley, D. Holz, M. Adeli Koudehi, S. K. Vogel, L. Burbaum, N. Mizuno, D.
Vavylonis, P. Schwille, Reconstitution of contractile actomyosin rings in vesicles. *Nature
Communications* **12**, 2254 (2021).

51. R. Sakamoto, M. P. Murrell, Composite branched and linear F-actin maximize myosin-
induced membrane shape changes in a biomimetic cell model. *Commun Biol* **7**, 1–12 (2024).

52. R. Sakamoto, D. S. Banerjee, V. Yadav, S. Chen, M. L. Gardel, C. Sykes, S. Banerjee, M. P.
Murrell, Membrane tension induces F-actin reorganization and flow in a biomimetic model
cortex. *Commun Biol* **6**, 325 (2023).

53. K. Y. Lee, S. J. Park, K. A. Lee, S. H. Kim, H. Kim, Y. Meroz, L. Mahadevan, K. H. Jung, T. K.
Ahn, K. K. Parker, K. Shin, Photosynthetic artificial organelles sustain and control ATP-
dependent reactions in a protocellular system. *Nature Biotechnology* **36**, 530–535 (2018).

54. S. Tanaka, K. Takiguchi, M. Hayashi, Repetitive stretching of giant liposomes utilizing the
nematic alignment of confined actin. *Communications Physics* **1** (2018).

55. A. Fink, C. R. Doll, A. Yagüe Relimpio, Y. Dreher, J. P. Spatz, K. Göpfrich, E. A. Cavalcanti-
Adam, Extracellular Cues Govern Shape and Cytoskeletal Organization in Giant Unilamellar
Lipid Vesicles. *ACS Synth. Biol.*, acssynbio.2c00516 (2023).

56. S. Razavi, F. Wong, B. Abubaker-Sharif, H. T. Matsubayashi, H. Nakamura, N. T. H. Nguyen,
D. N. Robinson, B. Chen, P. A. Iglesias, T. Inoue, Synthetic control of actin polymerization
and symmetry breaking in active protocells. *Sci Adv* **10**, eadk9731 (2024).

57. K. Zhang, B. Cui, Optogenetic control of intracellular signaling pathways. *Trends in
Biotechnology* **33**, 92–100 (2015).

58. A. Pudasaini, K. K. El-Arab, B. D. Zoltowski, LOV-based optogenetic devices: Light-driven
modules to impart photoregulated control of cellular signaling. *Frontiers in Molecular
Biosciences* **2**, 1–15 (2015).

59. J. Niu, M. Ben Johny, I. E. Dick, T. Inoue, Following Optogenetic Dimerizers and Quantitative
Prospects. *Biophysical Journal* **111**, 1132–1140 (2016).

60. G. Guntas, R. A. Hallett, S. P. Zimmerman, T. Williams, H. Yumerefendi, J. E. Bear, B.
Kuhlman, Engineering an improved light-induced dimer (iLID) for controlling the localization
and activity of signaling proteins. *Proceedings of the National Academy of Sciences of the
United States of America* **112**, 112–117 (2015).

61. A. Uyeda, T. Watanabe, T. Hohsaka, T. Matsuura, Different protein localizations on the inner
and outer leaflet of cell-sized liposomes using cell-free protein synthesis. *Synthetic Biology*
3, ysy007 (2018).

62. A. K. Rudd, J. M. Valls Cuevas, N. K. Devaraj, SNAP-Tag-Reactive Lipid Anchors Enable
Targeted and Spatiotemporally Controlled Localization of Proteins to Phospholipid
Membranes. *J. Am. Chem. Soc.* **137**, 4884–4887 (2015).

63. D. Scherfeld, N. Kahya, P. Schwille, Lipid dynamics and domain formation in model
membranes composed of ternary mixtures of unsaturated and saturated
phosphatidylcholines and cholesterol. *Biophysical journal* **85**, 3758–3768 (2003).

64. F. A. Thomas, I. Visco, Z. Petrášek, F. Heinemann, P. Schwille, Diffusion coefficients and
dissociation constants of enhanced green fluorescent protein binding to free standing
membranes. *Data in Brief* **5**, 537–541 (2015).

65. M. Frick, K. Schmidt, B. J. Nichols, Modulation of Lateral Diffusion in the Plasma Membrane
by Protein Density. *Current Biology* **17**, 462–467 (2007).

66. R. D. Mullins, P. Bieling, D. A. Fletcher, From solution to surface to filament: actin flux into
branched networks. *Biophys Rev* **10**, 1537–1551 (2018).

67. L. B. Case, X. Zhang, J. A. Ditlev, M. K. Rosen, Stoichiometry controls activity of phase-
separated clusters of actin signaling proteins. *Science* **363**, 1093–1097 (2019).

68. K. Yamamoto, M. Miyazaki, Optogenetic actin network assembly on lipid bilayer uncovers
the network density-dependent functions of actin-binding proteins. [Preprint] (2024).
https://doi.org/10.1101/2024.12.25.630233.

69. H. Nakamura, E. Rho, C. T. Lee, K. Itoh, D. Deng, S. Watanabe, S. Razavi, H. T. Matsubayashi,
C. Zhu, E. Jung, P. Rangamani, S. Watanabe, T. Inoue, ActuAtor, a Listeria-inspired molecular
tool for physical manipulation of intracellular organizations through de novo actin
polymerization. *Cell Reports* **42**, 113089 (2023).

70. S. Shekhar, M.-F. Carlier, Enhanced Depolymerization of Actin Filaments by ADF/Cofilin and
Monomer Funneling by Capping Protein Cooperate to Accelerate Barbed-End Growth.
Current Biology **27**, 1990-1998.e5 (2017).

71. A. Colin, T. Kotila, C. Guérin, M. Orhant-Prioux, B. Vianay, A. Mogilner, P. Lappalainen, M.
Théry, L. Blanchoin, Recycling of the actin monomer pool limits the lifetime of network
turnover. *The EMBO Journal* **42**, e112717 (2023).

72. H. Miki, K. Miura, T. Takenawa, N-WASP, a novel actin-depolymerizing protein, regulates
the cortical cytoskeletal rearrangement in a PIP2-dependent manner downstream of
tyrosine kinases. *The EMBO Journal* **15**, 5326–5335 (1996).

73. R. Rohatgi, L. Ma, H. Miki, M. Lopez, T. Kirchhausen, T. Takenawa, M. W. Kirschner, The
Interaction between N-WASP and the Arp2/3 Complex Links Cdc42-Dependent Signals to
Actin Assembly. *Cell* **97**, 221–231 (1999).

74. O. D. Weiner, G. Servant, M. D. Welch, T. J. Mitchison, J. W. Sedat, H. R. Bourne, Spatial
control of actin polymerization during neutrophil chemotaxis. *Nature Cell Biology* **1**, 75–81
(1999).

75. O. D. Weiner, M. C. Rentel, A. Ott, G. E. Brown, M. Jedrychowski, M. B. Yaffe, S. P. Gygi, L. C.
Cantley, H. R. Bourne, M. W. Kirschner, Hem-1 Complexes Are Essential for Rac Activation,
Actin Polymerization, and Myosin Regulation during Neutrophil Chemotaxis. *PLoS Biol* **4**,
e38 (2006).

76. A. Kawska, K. Carvalho, J. Manzi, R. Boujemaa-Paterski, L. Blanchoin, J.-L. Martiel, C. Sykes,
How actin network dynamics control the onset of actin-based motility. *Proceedings of the*
National Academy of Sciences **109**, 14440–14445 (2012).

77. K. B. Velle, L. K. Fritz-Laylin, Diversity and evolution of actin-dependent phenotypes. *Current*
Opinion in Genetics and Development **58–59**, 40–48 (2019).

78. E. D. Goley, M. D. Welch, The ARP2/3 complex: An actin nucleator comes of age. *Nature*
Reviews Molecular Cell Biology **7**, 713–726 (2006).

79. D. A. Valencia, M. E. Quinlan, Formins. *Current Biology* **31**, R517–R522 (2021).

80. D. R. Kovar, E. S. Harris, R. Mahaffy, H. N. Higgs, T. D. Pollard, Control of the Assembly of
ATP- and ADP-Actin by Formins and Profilin. *Cell* **124**, 423–435 (2006).

81. J. Funk, F. Merino, L. Venkova, L. Heydenreich, J. Kierfeld, P. Vargas, S. Raunser, M. Piel, P.
Bieling, Profilin and formin constitute a pacemaker system for robust actin filament growth.
eLife **8**, e50963 (2019).

82. S. Dmitrieff, F. Nédélec, Amplification of actin polymerization forces. *Journal of Cell Biology*
212, 763–766 (2016).

83. R. DeRose, T. Miyamoto, T. Inoue, Manipulating signaling at will: chemically-inducible
dimerization (CID) techniques resolve problems in cell biology. *Pflugers Archiv : European*
journal of physiology **465**, 409–417 (2013).

84. W. Luo, Z. Z. Lieu, E. Manser, A. D. Bershadsky, M. P. Sheetz, Formin DAAM1 organizes actin
filaments in the cytoplasmic nodal actin network. *PLoS ONE* **11**, 1–22 (2016).

85. K. M. C. Chan, A. L. Arthur, J. Morstein, M. Jin, A. Bhat, D. Schlesinger, S. Son, D. A. Stevens,
D. G. Drubin, D. A. Fletcher, Evolutionarily related small viral fusogens hijack distinct but
modular actin nucleation pathways to drive cell-cell fusion. *Proc Natl Acad Sci USA* **118**,
e2007526118 (2021).

86. A. Drazic, H. Aksnes, M. Marie, M. Boczkowska, S. Varland, E. Timmerman, H. Foyn, N.
Glomnes, G. Rebowski, F. Impens, K. Gevaert, R. Dominguez, T. Arnesen, NAA80 is actin's N-
terminal acetyltransferase and regulates cytoskeleton assembly and cell motility.
Proceedings of the National Academy of Sciences **115**, 4399–4404 (2018).

87. J. Block, D. Breitsprecher, S. Kühn, M. Winterhoff, F. Kage, R. Geffers, P. Duwe, J. L. Rohn, B.
Baum, C. Brakebusch, M. Geyer, T. E. B. Stradal, J. Faix, K. Rottner, FMNL2 Drives Actin-

Based Protrusion and Migration Downstream of Cdc42. *Current Biology* **22**, 1005–1012
(2012).

88. A. Schirenbeck, R. Arasada, T. Bretschneider, M. Schleicher, J. Faix, Formins and VASPs may
co-operate in the formation of filopodia. *Biochemical Society Transactions* **33**, 4 (2005).

89. H. Mizuno, C. Higashida, Y. Yuan, T. Ishizaki, S. Narumiya, N. Watanabe, Rotational
movement of the formin mDia1 along the double helical strand of an actin filament. *Science*
331, 80–83 (2011).

90. X. Naj, A. K. Hoffmann, M. Himmel, S. Linder, The formins FMNL1 and mDia1 regulate
coiling phagocytosis of borrelia burgdorferi by primary human macrophages. *Infection and*
Immunity **81**, 1683–1695 (2013).

91. C. J. Gould, S. Maiti, A. Michelot, B. R. Graziano, L. Blanchoin, B. L. Goode, The formin DAD
domain plays dual roles in autoinhibition and actin nucleation. *Current Biology* **21**, 384–390
(2011).

92. M. P. Murrell, R. Voituriez, J.-F. Joanny, P. Nassoy, C. Sykes, M. L. Gardel, Liposome
adhesion generates traction stress. *Nature Phys* **10**, 163–169 (2014).

93. A. Ward, F. Hilitski, W. Schwenger, D. Welch, A. W. C. Lau, V. Vitelli, L. Mahadevan, Z. Dogic,
Solid friction between soft filaments. *Nature Mater* **14**, 583–588 (2015).

94. M. A. Dichtl, E. Sackmann, Microrheometry of semiflexible actin networks through enforced
single-filament reptation: Frictional coupling and heterogeneities in entangled networks.
Proc. Natl. Acad. Sci. U.S.A. **99**, 6533–6538 (2002).

95. F. Nedelec, D. Foethke, Collective Langevin dynamics of flexible cytoskeletal fibers. *New J.*
Phys. **9**, 427–427 (2007).

96. C. Zhu, C. T. Lee, P. Rangamani, Mem3DG: Modeling membrane mechanochemical
dynamics in 3D using discrete differential geometry. *Biophysical Reports* **2**, 100062 (2022).

97. S. Dmitrieff, A. Alsina, A. Mathur, F. J. Nédélec, Balance of microtubule stiffness and cortical
tension determines the size of blood cells with marginal band across species. *Proc. Natl.*
Acad. Sci. U.S.A. **114**, 4418–4423 (2017).

98. Y. Shi, J. Zhang, M. Mullin, B. Dong, A. S. Alberts, K. A. Siminovitch, The mDial Formin Is
Required for Neutrophil Polarization, Migration, and Activation of the LARG/RhoA/ROCK
Signaling Axis during Chemotaxis. *The Journal of Immunology* **182**, 3837–3845 (2009).

99. D. Sakata, H. Taniguchi, S. Yasuda, A. Adachi-Morishima, Y. Hamazaki, R. Nakayama, T. Miki,
N. Minato, S. Narumiya, Impaired T lymphocyte trafficking in mice deficient in an actin-
nucleating protein, mDia1. *Journal of Experimental Medicine* **204**, 2031–2038 (2007).

100.   C. Yang, L. Czech, S. Gerboth, S. Kojima, G. Scita, T. Svitkina, Novel Roles of Formin
mDia2 in Lamellipodia and Filopodia Formation in Motile Cells. *PLoS Biol* **5**, e317 (2007).

101.   T. Isogai, R. van der Kammen, D. Leyton-Puig, K. M. Kedziora, K. Jalink, M. Innocenti,
Initiation of lamellipodia and ruffles involves cooperation between mDia1 and the Arp2/3
complex. *Journal of Cell Science*, jcs.176768 (2015).

102.   P. Monzo, Y. K. Chong, C. Guetta-Terrier, A. Krishnasamy, S. R. Sathe, E. K. F. Yim, W. H.
Ng, B. T. Ang, C. Tang, B. Ladoux, N. C. Gauthier, M. P. Sheetz, Mechanical confinement
triggers glioma linear migration dependent on formin FHOD3. *MBoC* **27**, 1246–1261 (2016).

103.   F. Kage, M. Winterhoff, V. Dimchev, J. Mueller, T. Thalheim, A. Freise, S. Brühmann, J.
Kollasser, J. Block, G. Dimchev, M. Geyer, H.-J. Schnittler, C. Brakebusch, T. E. B. Stradal, M.-
F. Carlier, M. Sixt, J. Käs, J. Faix, K. Rottner, FMNL formins boost lamellipodial force
generation. *Nat Commun* **8**, 14832 (2017).

104.   Y. Yamazaki, Y. Miyata, K. Morigaki, M. Miyazaki, Controlling Physical and Biochemical
Parameters of Actin Nucleation Using a Patterned Model Lipid Membrane. *Nano Lett.* **24**,
1825–1834 (2024).

105.   M. Abercrombie, J. E. M. Heaysman, S. M. Pegrum, The locomotion of fibroblasts in
culture I. Movements of the leading edge. *Experimental Cell Research* **59**, 393–398 (1970).

106.   P. Maiuri, E. Terriac, P. Paul-Gilloteaux, T. Vignaud, K. McNally, J. Onuffer, K. Thorn, P. A.
Nguyen, N. Georgoulia, D. Soong, A. Jayo, N. Beil, J. Beneke, J. C. Hong Lim, C. Pei-Ying Sim,
Y.-S. Chu, A. Jiménez-Dalmaroni, J.-F. Joanny, J.-P. Thiery, H. Erfle, M. Parsons, T. J.
Mitchison, W. A. Lim, A.-M. Lennon-Duménil, M. Piel, M. Théry, The first World Cell Race.
Current Biology **22**, R673–R675 (2012).

107.   S. A. Koestler, K. Rottner, F. Lai, J. Block, M. Vinzenz, J. V. Small, F- and G-Actin
Concentrations in Lamellipodia of Moving Cells. *PLoS ONE* **4**, e4810 (2009).

108.   S. Samarin, S. Romero, C. Kocks, D. Didry, D. Pantaloni, M. F. Carlier, How VASP
enhances actin-based motility. *Journal of Cell Biology* **163**, 131–142 (2003).

109.   W. M. Brieher, M. Coughlin, T. J. Mitchison, Fascin-mediated propulsion of Listeria
monocytogenes independent of frequent nucleation by the Arp2/3 complex. *Journal of Cell
Biology* **165**, 233–242 (2004).

110.   P. Li, S. Banjade, H. C. Cheng, S. Kim, B. Chen, L. Guo, M. Llaguno, J. V. Hollingsworth, D.
S. King, S. F. Banani, P. S. Russo, Q. X. Jiang, B. T. Nixon, M. K. Rosen, Phase transitions in the
assembly of multivalent signalling proteins. *Nature* **483**, 336–340 (2012).

111.   C. Simon, R. Kusters, V. Caorsi, A. Allard, M. Abou-Ghali, J. Manzi, A. Di Cicco, D. Lévy, M.
Lenz, J.-F. Joanny, C. Campillo, J. Plastino, P. Sens, C. Sykes, Actin dynamics drive cell-like
membrane deformation. *Nature Physics* **15**, 602–609 (2019).

112.   P. DiMilla, J. Stone, J. Quinn, S. Albelda, D. Lauffenburger, Maximal migration of human
smooth muscle cells on fibronectin and type IV collagen occurs at an intermediate
attachment strength. *Journal of Cell Biology* **122**, 729–737 (1993).

113.   C. Schreiber, B. Amiri, J. C. J. Heyn, J. O. Rädler, M. Falcke, On the adhesion–velocity
relation and length adaptation of motile cells on stepped fibronectin lanes. *Proc. Natl. Acad.*
Sci. U.S.A. **118**, e2009959118 (2021).

114.   V. Caorsi, J. Lemière, C. Campillo, M. Bussonnier, J. Manzi, T. Betz, J. Plastino, K.
Carvalho, C. Sykes, Cell-sized liposome doublets reveal active tension build-up driven by
acto-myosin dynamics. *Soft Matter* **12**, 6223–6231 (2016).

115.   W. I. Goh, K. B. Lim, T. Sudhaharan, K. P. Sem, W. Bu, A. M. Chou, S. Ahmed, mDia1 and
WAVE2 Proteins Interact Directly with IRSp53 in Filopodia and Are Involved in Filopodium
Formation. *Journal of Biological Chemistry* **287**, 4702–4714 (2012).

116.   A. Bisaria, A. Hayer, D. Garbett, D. Cohen, T. Meyer, Membrane-proximal F-actin
restricts local membrane protrusions and directs cell migration. *Science* **368**, 1205–1210
(2020).

117.   Z. Shi, Z. T. Graber, T. Baumgart, H. A. Stone, A. E. Cohen, Cell Membranes Resist Flow.
Cell **175**, 1769-1779.e13 (2018).

118.   S. Gat, C. Simon, C. Campillo, A. Bernheim-Groswasser, C. Sykes, Finger-like membrane
protrusions are favored by heterogeneities in the actin network. *Soft Matter* **16**, 7222–7230
(2020).

119.   S. M. Bartelt, J. Steinkühler, R. Dimova, S. V. Wegner, Light-Guided Motility of a Minimal
Synthetic Cell. *Nano Lett.* **18**, 7268–7274 (2018).

120.   L. A. Banaszynski, C. W. Liu, T. J. Wandless, Characterization of the FKBP·Rapamycin·FRB
Ternary Complex. *J. Am. Chem. Soc.* **127**, 4715–4721 (2005).

121.   E. Rihtar, T. Lebar, D. Lainšček, K. Kores, S. Lešnik, U. Bren, R. Jerala, Chemically inducible
split protein regulators for mammalian cells. *Nat Chem Biol*, doi: 10.1038/s41589-022-
01136-x (2022).

122.   K. Yoshida, K. Abe, Y. Sato, I. Kawamata, R. J. Archer, H. T. Matsubayashi, S. Hamada, S.
Murata, S. Nomura, Transmembrane DNA Sequence Signaling via Hybridization. ChemRxiv
[Preprint] (2024). https://doi.org/10.26434/chemrxiv-2024-571kp.

123.   S. L. Veatch, S. L. Keller, Separation of Liquid Phases in Giant Vesicles of Ternary
Mixtures of Phospholipids and Cholesterol. *Biophysical Journal* **85**, 3074–3083 (2003).

124.   T. Litschel, B. Ramm, R. Maas, M. Heymann, P. Schwille, Beating Vesicles: Encapsulated
1798   Protein Oscillations Cause Dynamic Membrane Deformations. *Angewandte Chemie*
1799   *International Edition* **57**, 16286–16290 (2018).

1800   125.   S. Takada, N. Yoshinaga, N. Doi, K. Fujiwara, Controlling the Periodicity of a Reaction–
1801   Diffusion Wave in Artificial Cells by a Two-Way Energy Supplier. *ACS Nano* **16**, 16853–16861
1802   (2022).

1803   126.   S. D. Hansen, W. Y. C. Huang, Y. K. Lee, P. Bieling, S. M. Christensen, J. T. Groves,
1804   Stochastic geometry sensing and polarization in a lipid kinase–phosphatase competitive
1805   reaction. *Proceedings of the National Academy of Sciences of the United States of America*
1806   **116**, 15013–15022 (2019).

1807   127.   Y. Sato, M. Takinoue, Capsule-like DNA Hydrogels with Patterns Formed by Lateral Phase
1808   Separation of DNA Nanostructures. *JACS Au* **2**, 159–168 (2022).

1809   128.   R. Sakamoto, Z. Izri, Y. Shimamoto, M. Miyazaki, Y. T. Maeda, Geometric trade-off
1810   between contractile force and viscous drag determines the actomyosin-based motility of a
1811   cell-sized droplet. *Proc. Natl. Acad. Sci. U.S.A.* **119**, e2121147119 (2022).

1812   129.   L. T. Benk, A. S. Benk, R. B. Lira, E. A. Cavalcanti-Adam, R. Dimova, R. Lipowsky, B. Geiger,
1813   J. P. Spatz, Integrin α_{IIb} β_3 Activation and Clustering in Minimal Synthetic Cells. *Advanced*
1814   *NanoBiomed Research* **2**, 2100094 (2022).

1815   130.   S. Brühmann, D. S. Ushakov, M. Winterhoff, R. B. Dickinson, U. Curth, J. Faix, Distinct
1816   VASP tetramers synergize in the processive elongation of individual actin filaments from
1817   clustered arrays. *Proc Natl Acad Sci USA* **114**, E5815–E5824 (2017).

1818   131.   Y. Liang, Y. Takaki, H. Inaba, K. Matsuura, Local Deformation in Phase-Separated Giant
1819   Liposomes via Photo-Induced Peptide Nanofiber Formation. *Langmuir* **41**, 15173–15181
1820   (2025).

1821   132.   J. E. Tropea, S. Cherry, D. S. Waugh, "Expression and Purification of Soluble His6-Tagged
1822   TEV Protease" in *High Throughput Protein Expression and Purification: Methods and*
1823   *Protocols*, S. A. Doyle, Ed. (Humana Press, Totowa, NJ, 2009; https://doi.org/10.1007/978-1-
1824   59745-196-3_19)*Methods in Molecular Biology*, pp. 297–307.

1825   133.   T. Inoue, W. D. Heo, J. S. Grimley, T. J. Wandless, T. Meyer, An inducible translocation
1826   strategy to rapidly activate and inhibit small GTPase signaling pathways. *Nature Methods* **2**,
1827   415–418 (2005).

1828   134.   Cytoplasmic Actin: Purification and Single Molecule Assembly Assays | SpringerLink.
1829   https://link.springer.com/protocol/10.1007/978-1-62703-538-5_9.

1830   135.   L. K. Doolittle, M. K. Rosen, S. B. Padrick, Measurement and analysis of in vitro actin
1831   polymerization. *Methods in Molecular Biology*, doi: 10.1007/978-1-62703-538-5_16 (2013).

136.   S. Razavi, T. Inoue, T. LUO, D. Robinson, Methods for making giant vesicles and their use
(2024). https://patents.google.com/patent/US11951210B2/en.

137.   J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch,
C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D. J. White, V. Hartenstein, K. Eliceiri, P.
Tomancak, A. Cardona, Fiji: an open-source platform for biological-image analysis. *Nature*
methods **9**, 676–682 (2012).

138.   H. T. Matsubayashi, J. Mountain, N. Takahashi, A. Deb Roy, T. Yao, A. F. Peterson, C. Saez
Gonzalez, I. Kawamata, T. Inoue, Non-catalytic role of phosphoinositide 3-kinase in
mesenchymal cell migration through non-canonical induction of p85β/AP2-mediated
endocytosis. *Nat Commun* **15**, 2612 (2024).

139.   K. Crane, DISCRETE DIFFERENTIAL GEOMETRY: AN APPLIED INTRODUCTION.

140.   M. Meyer, M. Desbrun, P. Schröder, A. H. Barr, "Discrete Differential-Geometry
Operators for Triangulated 2-Manifolds" in *Visualization and Mathematics III*, H.-C. Hege, K.
Polthier, Eds. (Springer, Berlin, Heidelberg, 2003), pp. 35–57.

141.   C. C. Beltzner, T. D. Pollard, Pathway of Actin Filament Branch Formation by Arp2/3
Complex*. *Journal of Biological Chemistry* **283**, 7135–7144 (2008).

142.   W. Rawicz, K. C. Olbrich, T. McIntosh, D. Needham, E. Evans, Effect of Chain Length and
Unsaturation on Elasticity of Lipid Bilayers. *Biophysical Journal* **79**, 328–339 (2000).

143.   M. J. Footer, J. W. J. Kerssemakers, J. A. Theriot, M. Dogterom, Direct measurement of
force generation by actin filament polymerization using an optical trap. *Proc. Natl. Acad.*
Sci. U.S.A. **104**, 2181–2186 (2007).

144.   D. R. Kovar, T. D. Pollard, Insertional assembly of actin filament barbed ends in
association with formins produces piconewton forces. *Proc. Natl. Acad. Sci. U.S.A.* **101**,
14725–14730 (2004).

145.   D. Drabik, M. Gavutis, R. Valiokas, A. Ulčinas, Determination of the Mechanical
Properties of Model Lipid Bilayers Using Atomic Force Microscopy Indentation. *Langmuir*
36, 13251–13262 (2020).

146.   O. H. Akenuwa, S. M. Abel, Organization and dynamics of cross-linked actin filaments in
confined environments. *Biophysical Journal* **122**, 30–42 (2023).

147.   San Diego Supercomputer Center (2022): Triton Shared Computing Cluster. University of
California, San Diego. Service. https://doi.org/10.57873/T34W2R

148.   L. A. Gross, G. S. Baird, R. C. Hoffman, K. K. Baldridge, R. Y. Tsien, The structure of the
chromophore within DsRed, a red fluorescent protein from coral. *Proc Natl Acad Sci U S A*
97, 11990–11995 (2000).

149.   V. I. Martynov, A. P. Savitsky, N. Y. Martynova, P. A. Savitsky, K. A. Lukyanov, S. A.
Lukyanov, Alternative Cyclization in GFP-like Proteins Family. *Journal of Biological Chemistry*
276, 21012–21016 (2001).

1869

Figure 1

Fig. 1. Reversible and asymmetric control of protein localization with iLID-SspB within GUVs. (A) Schematic representation of iLID-SspB reaction with global light stimulation. iLID was tethered on the membrane via the interaction between SNAP-tag and bendylguanine-conjugated lipids. Upon blue light illumination, Jα helix of iLID extends to expose the binding site for SspB, whereas in the dark, iLID reverses to its closed state and releases SspB. (B) Representative images of iLID-YFP-SNAP and mCherry-SspB. Scale bar, 10 μm. Blue bar indicates the period of blue light illumination. (C) Time course of membrane/lumen ratio of mCherry-SspB signal (n=5). Error bars indicate 95% CI. Blue area indicates the period of blue light stimulation. (D) Schematic representation of iLID-SspB reaction with local light stimulation. (E) Representative images of mCherry-SspB. Scale bar, 10 μm. Yellow box indicates the area of blue light illumination. (F) Membrane/lumen ratio of mCherry-SspB for the representative image shown. The distance is measured from the center of blue light illumination along the perimeter of the GUV. (G) Representative image and kymograph of repetitive and reversible SspB translocation. Kymograph shows membrane signal of mCherry-SspB.

Figure 2

Fig. 2. Reversible light control of actin cytoskeleton within GUVs. (A) Bulk pyrene actin polymerization assay with 1 µM actin (5% pyrene-labeled), 10 nM Arp2/3, and 100 nM NPFs. (-NPF) represents actin and Arp2/3 only condition. (B) Schematic representation of light-inducible actin polymerization. Translocation of pVCA-SspB-mCherry increases local concentration of pVCA on the membrane and enhances actin polymerization. (C) Representative images of pVCA-SspB-mCherry and Alexa 647 actin. Blue bar indicates the period of blue light illumination. (D) Time course of membrane/lumen ratio of mCherry and Alexa 647 actin in (C) and fig. S9. pVCA +Arp2/3+Light: full set of components (n=4). -Light: no blue light stimulation (n=3). -pVCA: pVCA-SspB-mCherry was replaced with mCherry-SspB (n=4). -Arp2/3: Arp2/3 was omitted (n=4). In (E)–(H), pVCA-SspB-mCherry was constitutively anchored to the membrane via SsrA-SNAP. (E) Pharmacological perturbation of membrane-associated actin. DMSO (n=8), Jasplakinolide (Jasp, n=10), Latrunculin A (LatA, n=9). (F) Time course of normalized membrane/lumen ratio of Alexa-647 actin in (E). The values were normalized by time=0. (G) FRAP analysis of membrane fluorescence of mCherry and Alexa 647 actin in fig. S11. The values were normalized by time=0. Sample sizes are listed in fig. S11A. (H) Electron microscopy (EM) images of actin filaments in the presence and absence of pVCA. Scale bar indicates 500 nm. (I) Representative images of reversible actin polymerization. Blue bars indicate the period of blue light illumination. (J) Time course of membrane/lumen ratio of pVCA-SspB-mCherry and Alexa 647-labeled actin in (F). (D, F, G) Error bars indicate 95% CI.

Figure 3

Fig. 3. Asymmetric light control of actin cytoskeleton within GUVs. (A) Schematic representation of iLID-SspB reaction with local light stimulation. (B, D) Representative images of light-inducible asymmetric actin polymerization. Yellow boxes indicate the area of blue light illumination. (C, E) Kymograph of membrane pVCA-SspB and actin signals. Blue bars indicate the period of blue light illumination. Triangles indicate timepoints represented in (B, D).

Figure 4

Fig. 4. mDia1-mediated actin polymerization drives protrusive movement of GUVs. (A) Schematic representation of light-inducible membrane protrusion driven by pVCA and mDia1-mediated actin polymerization. (B) Representative images of pVCA-mDia1-mediated movement of GUVs. Yellow boxes indicate the area of blue light stimulation. Dotted line shows the initial position of the GUV. (C) Representative images of iLID-EYFP-SNAP showing the increase in adhesive area and vesicle deformation into dome-like shape. X-y images show the confocal plane right above the bottom substrate. Z-x and y-z images are reconstituted from z-stack. Dotted line shows the adhesive area of the previous images. (D) Color-coded boundaries of GUVs. The data correspond as follows. Left: Fig. 4B. Middle: fig. S16A. Right: fig. S16B. (E, F) Kymographs of membrane signal of mCherry and actin Alexa-647 and membrane protrusions. Triangles in (E) indicate the timepoints represented in (B, C). Data of (E) and (F) correspond to Fig. 4 (B) and fig. S16A, respectively. (G) Distance the front (illuminated) side of GUV membrane moved forward. pVCA+mDia1 (1 μM GST-pVCA-SspB-mCherry + 1 μM mCherry-SspB-mDia1, n=14 vesicles), mDia1 (1 μM mCherry-SspB-mDia1, n=12 vesicles), pVCA (1 μM GST-pVCA-SspB-mCherry, n=12 vesicles). Error bars indicate 95% CI. (H, I) Quantification of the initial velocity and the distance the front side of GUV membrane moved forward. P values of Steel-Dwass test (two sided) are indicated. pVCA + mDia1: n=22 vesicles. mDia1: n=11 vesicles. pVCA: n=12 vesicles. Box whisker plots represent median, 1st, 3rd quartiles and 1.5×inter-quartile range. (J) Distance the front (illuminated) side of GUV membrane moved forward. DMSO (n=9 vesicles), 30 μM LatA (Latrunculin A) (n=10 vesicles). DMSO, ΔDAD: mCherry-SspB-mDia1 (FH1-FH2-DAD) was replaced with mCherry-SspB-mDia1(FH1-FH2) (n=13 vesicles). 30 μM SMIFH2 (n=6 vesicles). Error bars indicate 95% CI. (H, I) Quantification of the initial velocity and the distance the front side of GUV membrane moved forward. P values of Steel-Dwass test (two sided) are indicated. pVCA + mDia1: n=22 vesicles. mDia1: n=11 vesicles. pVCA: n=12 vesicles. Box whisker plots represent median, 1st, 3rd quartiles and 1.5×inter-quartile range. (K) A GUV deformed into dome-like shape after movement. Upper images: before light stimulation. Lower images: after light stimulation. The data corresponds to fig. S16C.

Figure S1

Fig. S1. Proteins purified in this study. (A) SDS-PAGE results of purified proteins. Blue: Coomassie brilliant blue staining. Gray: SYPRO Ruby staining. Most mCherry-tagged proteins show two extra bands due to a known chromophore cleavage reaction during the boiling process (145, 146). (B) Domain structures of NPFs used in this study.

Figure S2

A

Ni-NTA   6xHis tag
iLID

B   5 mol% NI-NTA-DGS/0–2 mol% PEG2000-PE/ 93-95 mol% POPC

2xStrep-iLID-YFP-6xHis (4 µM)   mCherry-SspB (1 µM)

PEG2000-PE
0 mol%

0.5 mol%

1.0 mol%

2.0 mol%

C   5 mol% Ni-NTA-DGS/95 mol% POPC

mCherry-SspB (1 µM)

D

POPS   MARCKS-ED
iLID

E   2xStrep-iLID-YFP-MARCKS (4 µM)   mCherry-SspB (1 µM)

10 mol% POPS/ 90 mol% POPC

20 mol% POPS/ 80 mol% POPC

40 mol% POPS/ 60 mol% POPC

F   40 mol% POPS/58–60 mol% POPC

2xStrep-iLID-YFP-MARCKS (4 µM)   mCherry-SspB (1 µM)

PEG2000-PE
0.5 mol%

1.0 mol%

2.0 mol%

Figure S2

Fig. S2. Optimization of membrane anchoring of iLID. (A),(D),(G),(I): Schematic representations of membrane anchoring of iLID. (B),(C), (E),(F),(H),(J): Confocal images of iLID variants and mCherry-SspB. (B) His-tag Ni-NTA pair caused non-specific membrane recruitment of SspB. This is possibly caused by the presence of 2×Strep-iLID-YFP-6×His because mCherry-SspB itself did not show non-specific binding to the membrane of 5%Ni-NTA DGS/95%POPC (C). (E),(F) MARCKS-ED peptide requires high percentage of PS and PEGylated-lipids to reduce non-specific binding of SspB. (H) Biotin-StrepTactin-Streptag also caused non-specific membrane binding of SspB. (J) The data correspond with Fig. 1B.

Figure S3

Fig. S3. Asymmetric control of protein localization with iLID-SspB within GUVs. (A) Time course of membrane/lumen ratio of mCherry-SspB signal in the illuminated area (n=6). Error bars indicate 95% CI. Blue area indicates the time window of blue light stimulation. The data corresponds to Fig. 1E and F. (B, C) Confocal images and time course of membrane/lumen ratio of mCherry-SspB signal responding to repetitive local light illuminations. The data corresponds to Fig. 1G.

Figure S4

Fig. S4. Optimization of ActA-SspB. (A) Pyrene actin polymerization assay. The reaction was performed with 1 μM actin (5% pyrene labeled), 10 nM Arp2/3, and 100 nM ActA variants or GST-VCA. (B) Fluorescence intensity/time in 0–6 min of (A) was plotted as bar chart. (C–E) Light-inducible local membrane recruitment of ActA (C) Confocal images of ActA translocation. Yellow boxes indicate the area of blue light illumination. (D) Time course of membrane/lumen ratio of ActA-SspB variants (n≥5 for each variant). Error bars indicate 95% CI. Blue bar indicates the time window of blue light stimulation. (E) Average membrane/lumen ratio of mCherry signal. In each condition, left and right bars indicate before and after light stimulation, respectively. The numbers on the bar chart indicate the fold change after stimulation.

Figure S5

Figure S5

Fig S5. Light-inducible actin polymerization with global ActA membrane recruitment. iLID-EYFP-MARCKS (5 µM), ActA-SspBnano-mCherry (2 µM), actin (1.5 µM, 10% Alexa-647 labeled), and Arp2/3 (150 nM) were encapsulated in GUVs containing 60 mol% POPC/40 mol% POPS. Blue light was applied to the entire GUV. (A) Representative images of ActA(1–183)-SspB(nano)-mCherry and actin Alexa-647. All components (n=4): w/o ActA: ActA(1–183)-SspB(nano)-mCherry was replaced with mCherry-SspB(micro) (n=3), w/o Arp2/3: Arp2/3 was omitted (n=3). No Light: no blue light stimulation (n=3). (B, C) Time course of membrane/lumen ratio of mCherry and actin Alexa 647. (D) Bar chart of membrane/lumen ratio of actin Alexa 647 before (0 min) and after (40 min) light stimulation. Error bars indicate 95% CI.

Figure S6

Fig S6. Asymmetric ActA membrane recruitment fails to induce directed actin polarization toward local light illumination. iLID-EYFP-MARCKS (5 μM), ActA-SspBnano-mCherry (2 μM), actin (1.5 μM, 10% Alexa-647 labeled), and Arp2/3 (150 nM), 1 mM ATP were encapsulated in GUVs containing 60 mol% POPC/40 mol% POPS. Yellow boxes indicate the area of blue light illumination. Although ActA creates asymmetric distribution, actin patches randomly distributed due to the diffusion on the membrane.

Figure S7

Fig. S7. Global ActA membrane recruitment fails to induce actin polymerization on the membrane in the presence of profilin, cofilin, and capping protein. 15 µM iLID-YFP-MARCKS, 5 µM ActA(1-183)-SspB(nano)-mCherry, 7.5 µM actin (10% Alexa 647 labeled), 150 nM Arp2/3, 3 µM profilin, cofilin, and capping protein were encapsulated in GUVs containing 60 mol% POPC/40 mol% POPS. Blue bars indicate the period of blue light illumination.

Figure S8

fig. S8. Biochemical characterization of actin and actin-binding proteins used in this study. (A, B) Ultracentrifuge assay. After 30 min actin polymerization reaction at RT, the solutions were ultracentrifuged 100,000×g, 60 min, at RT. Proteins and concentrations used: 7.5 μM actin (10% Alexa-488 labeled), 3 μM profilin, 4 μM cofilin, 50 nM capping protein, 150 nM Arp2/3, 1 μM GST-pVCA-SspB-mCherry, 1 μM mCherry-SspB-mDia1. G-buffer: 2 mM Tris-HCl (pH 7.5), 0.2 mM ATP, 0.1 mM CaCl2, 0.5 mM DTT, 1 mM NaN3. KMEI-S buffer: .10mM imidazole (pH 7.0), 50mM KCl, 1mM MgCl2, 1mM EGTA, 240 mM sucrose (sucrose was added to mimic the buffer conditions in GUV experiments). (C–F) Pyrene actin polymerization assay. The reactions were performed with 2 μM actin (5% pyrene labeled) at RT. (C, D) 10 nM Arp2/3, and 100 nM NPFs. (-) represents actin and Arp2/3 only condition. (E) The reactions were performed with the indicated factors. (F) 50 nM capping protein, 150 nM Arp2/3, 3 μM profilin, 1 μM mCherry-SspB-mDia1, 1 μM GST-pVCA-SspB-mCherry. (-) represents actin only condition.

Figure S9

Fig. S9. Light control of actin cytoskeleton with pVCA-SspB-mCherry. The data correspond to Fig. 2C–E. (+) condition contains 15 µM iLID-YFP-MARCKS, 4.2 µM pVCA-SspB(nano)-mCherry, 7.5 µM actin (10% Alexa 647 labelled), 150 nM Arp2/3, 100 nM capping protein, 3 µM profilin, 4 µM cofilin, and 1 mM ATP. In (-light) and (-Arp2/3) conditions, blue light illumination or Arp2/3 was omitted from (+pVCA +Arp2/3 +Light) condition, respectively. In (-pVCA) condition, pVCA-SspB(nano)-mCherry was replaced with mCherry-SspB(micro).

Figure S10

fig. S10. Pharmacological perturbation of membrane-associated actin assemblies. (A) Schematic representation of constitutive tethering of NPFs. SspB-fused NPFs (pVCA and/or mDia1) were tethered to the GUV membrane via the SsrA–SNAP/BG-lipid system. (B) Representative confocal images of membrane-associated actin assemblies formed by pVCA, mDia1, or their combination. Scale bars, 5 μm. (C) Quantification of the initial membrane/lumen fluorescence intensity ratio for mCherry and actin Alexa-647. pVCA: n=27 vesicles. mDia1: n=23 vesicles. pVCA + mDia1: n=27 vesicles. Box whisker plots represent median, 1st, 3rd quartiles and 1.5×inter-quartile range. (D) Representative images before (0 min) and after (24 min) treatment with DMSO, 50 μM Latrunculin A, or 50 μM Jasplakinolide. Scale bars, 5 μm. (E) Time courses of membrane/lumen ratio and normalized membrane/lumen ratio of actin Alexa-647. (F) Time courses of normalized membrane/lumen ratio of actin Alexa-647 for each NPF condition. The y-axis is shown on a logarithmic scale. (E, F) Error bars indicate 95% CI. (G) Representative images showing membrane deformation after Jasplakinolide treatment. Scale bars, 10 μm. mCherry-SspB-mDia1-6×His was used as mDia1 in this figure.

Figure S11

A

	pVCA	Arp2/3	mDia1	actin	profilin	cofilin	capping protein	SsrA-SNAP		mCherry FRAP	Alexa 647 FRAP
pVCA/Arp2/3 + prof/cof/CP	+	+			+	+	+	+		n=9	-
pVCA/Arp2/3 + prof/cof/CP + actin	+	+		+	+	+	+	+		n=10	n=13
pVCA/Arp2/3 + actin	+	+		+				+		n=7	n=8
mDia1 + prof/cof			+		+	+		+		n=9	-
mDia1 + prof/cof + actin			+	+	+	+		+		n=6	n=8
pVCA/Arp2/3 + mDia1 + prof/cof/CP + actin	+	+	+	+	+	+	+	+		n=6	n=10

fig. S11. fluorescence recovery after photobleaching (FRAP) analysis of membrane-associated NPFs and actin. (A) Summary table of proteins included in each conditions. In FRAP analysis, SspB-fused NPFs (pVCA and/or mDia1) were tethered to the GUV membrane via the SsrA–SNAP/BG-lipid system. (B, C) Representative confocal images of mCherry-tagged NPFs and actin Alexa-647. The white dashed box indicates the photobleached region. Scale bars, 5 μm. (D, E) Time course of mCherry and Alexa-647 fluorescence recovery at the bleached region. Fluorescence intensities were normalized to those of a non-bleached region to compensate for photobleaching during image acquisition. A subset of data also presented in Fig. G.

Figure S12

A **SUVs only**
(98 mol% POPC, 2 mol% 18:1 PE-PEG2000-benzylguanine)

B **pVCA** (pVCA, Arp2/3, CP, profilin, cofilin) ×10 dilution

C **pVCA negative control** (Arp2/3, CP, profilin, cofilin) ×10 dilution

fig. S12. Electron microscopy (EM) analysis of actin filaments. Ultrastructure of small unilamellar vesicles (SUVs) and actin filaments were analyzed by negative staining electron microscopy. (A) Representative EM images of SUVs alone. (B) Representative images of Y-junctions formed in the presence of pVCA, Arp2/3, capping protein (CP), profilin, and cofilin. (C) Negative control sample lacking pVCA (Arp2/3, CP, profilin, and cofilin only). (B, C) To reduce filament overlap, the polymerization reactions were diluted 10-fold prior to deposition onto EM grids. (D) Representative EM images of actin assemblies formed with pVCA or the corresponding negative control lacking pVCA. (E) Representative EM images of actin assemblies formed with pVCA + mDia1, mDia1 alone, or the corresponding negative control lacking mDia1. All samples were deposited onto EM grids and negatively stained prior to imaging. Scale bars are shown in each panel. mCherry-SspB-mDia1-6×His was used as mDia1 in this figure.

D **pVCA** (pVCA, Arp2/3, CP, profilin, cofilin)

pVCA negative control (Arp2/3, CP, profilin, cofilin)

pVCA + mDia1 (pVCA, mDia1, Arp2/3, CP, profilin, cofilin)

E

mDia1 (mDia1, profilin, cofilin)

mDia1 negative control (profilin, cofilin)

Figure S13

Figure S13

Fig. S13. Representative images of reversible actin polymerization. Confocal images and time course of membrane/lumen ratio of pVCA-SspB-mCherry and Alexa 647-labeld actin. Blue bars on images and blue regions in graphs indicate timepoints with blue light illumination. This figure corresponds with Fig. 2F, G.

Figure S14

A

B

C

pVCA-SspB-mCherry   **actin-Alexa647**

fig. S14. Asymmetric actin polymerization induced by pVCA. (A) Confocal images of local pVCA recruitment and actin polymerization. Yellow boxes indicate areas of blue light illumination. (B) Membrane/lumen ratio of mCherry-tagged pVCA and actin Alexa-647 for the images in (A). The distance is measured from the center of blue light illumination along the perimeter of the GUV. (A, B) The data corresponds to Fig. 3B and C. (C) Confocal images of local pVCA recruitment accompanied by vesicle movement toward light stimulation. Gray scaled images were created by averaging pVCA and actin images. Dotted circles indicate positions of vesicles in the images of one above.

Figure S15

Fig. S15. Quantitative analysis of mDia1-induced filopodia formation in Cos7 cells. (A) Schematic representation of chemically-inducible mDia1 translocation and membrane protrusions. (B, C) Confocal images of plasma membrane recruitment of mDia1 and subsequent filopodia formation. Cos7 cells were transfected with Lyn-ECFP-FRB, Lifeact-mCherry, and either of YFP-FKBP or YFP-FKBP-mDia1. At time 0, 100 nM rapamycin was added. Green: YFP-FKBP or YFP-FKBP-mDia1. Magenta: Lifeact-mCherry. (B) YFP-FKBP control. (C) YFP-FKBP-mDia1. (D) Zoomed-up view of filopodia extension. Images were acquired every 1.5 minutes. The bottom is merged image. (E) Quantification of mDia1-induced filopodia. Number of filopodia whose length is longer than 15 μm was counted. Box whisker plots represent median, 1st, 3rd quartiles and 1.5×inter-quartile range. P-values: ****: < 0.0001. Wilcoxon rank sum test. YF (YFP-FKBP): n=36 cells. YF-mDia1 (YFP-FKBP-mDia1): n=33 cells. (F–H) Quantification of the phenotypes observed after plasma membrane translocation of mDia1 (n=159 cells), pVCA (n=142 cells), Tiam1 (Rac1 GEF) (n=67 cells), and control protein YFP-FKBP (n=44 cells).

Figure S16

Fig. S16. mDia1-mediated GUV movement and the comparison between mDia1 and pVCA + mDia1 system. (A) Representative images of mDia1-mediated GUV movement. Yellow boxes indicate the area of blue light stimulation. Dotted lines indicate the initial position of the GUV. (B, C) Time course of the distance the front (illuminated) side of GUV membrane moved forward. Each line represents each GUVs. Insets show the initial response.

Figure S17

Fig. S17. Local protrusions in pVCA + mDia1 GUVs. Representative images of local protrusions seen in pVCA + mDia1 GUVs. Yellow boxes indicate the area of blue light stimulation. Dotted lines indicate the initial position of the GUV. White arrows indicate where local protrusions arose.

Figure S18

A

B

SspB

0 min   51 min

Fig. S18. Lack of vesicle movement and shape change with mCherry-SspB. With pVCA-SspB-mCherry and mCherry-SspB-mDia1 being replaced with mCherry-SspB, GUVs neither moved nor changed their shape after blue light illumination. (A) Time course of the distance the front (illuminated) side of GUV membrane moved forward. pVCA + mDia1 GUVs: n=6 vesicle. SspB GUVs: n=4 vesicles. Error bars indicate 95% CI. (B) 3D reconstitution of iLID-YFP-SNAP images before and after blue light illumination.

Figure S19

Fig. S19. Relationship between vesicle motion and biophysical parameters. (A) Relationship between initial velocity (average velocity over the first 5 minutes) and net displacement after 30 minutes of GUV forward movement. (B, C) Biophysical parameters are plotted against initial velocity and displacement are plotted in (B) and (C), respectively. R and p indicate Pearson's correlation coefficient and its p-value.

Figure S20

Fig. S20. Representative images of pVCA-mDia1-mediated movement of GUVs. (A, B) Representative images of pVCA-mDia1-mediated movement of GUVs. Yellow boxes indicate the area of blue light stimulation. Dotted lines indicate the initial position of the GUV. White arrows indicate where local protrusions arose. (C) Distance the front side of GUV membrane moved forward. Data was quantified from the images of (B). Blue areas indicate the periods of blue light stimulation.

Figure S21

A pVCA   mDia1   pVCA + mDia1

-Sterics

1 µm

B

+Sterics

1 µm

C   −Sterics

Membrane extension (µm)

0.03

0.02

0.01

0.00

pVCA   mDia1   pVCA
                  +
                  mDia1

D   +Sterics

pVCA   mDia1   pVCA
                  +
                  mDia1

Fig. S21. Computational modeling of actin network dynamics underlying membrane deformation. (A, B) Snapshots showing actin filaments at the leading edge of the membrane after 5 s of simulation. Actin filaments are shown in green, pVCA-Arp2/3 is shown in red, mDia1 is shown in orange, and Arp2/3 is shown in magenta. The membrane location at t = 0 is shown with the cyan dotted line to highlight the extent of membrane deformation. Simulations without and with inter-filament steric interactions are shown in A and B, respectively. (C, D) Box-whisker plots showing distributions of the membrane extension for the different actin binding proteins. Results for networks without and with inter-filament steric interactions are shown in A and B, respectively. Each box plot is constructed using 20 independent simulation trajectories. Error bars represent standard deviation about the mean.