Dexrazoxane as a viable microsporidia control agent in *Anopheles gambiae*

Tiago G. Zeferino [1] [¶]

Luís M. Silva [1,2] [*] [¶]

[1] Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland

[2] Department of Zoology, University of British Columbia, Vancouver, Canada

[¶] Both authors contributed equally

* Corresponding author

E-mail: luis.silva@ubc.ca (LMS)

ORCID

TGZ: 0009-0004-0767-5200

LMS: 0000-0002-8203-4006

Preprinted version in bioRxiv:

https://doi.org/10.1101/2024.10.14.617874

Keywords

dexrazoxane, infection, microsporidia, control agent, *Vavraia culicis*

Abstract

Microsporidia have long been proposed as biological agents for controlling disease vectors and the parasites they transmit. However, their study in vector biology has been constrained due to challenges in manipulating microsporidia within hosts. In this study, we investigated the effect of Dexrazoxane, a candidate drug against microsporidiosis, on the establishment and development of *Vavraia culicis* infection in its natural host, the mosquito *Anopheles gambiae*, the main malaria vector. Our findings show that Dexrazoxane significantly reduces spore load, particularly in mosquitoes reared individually, without affecting the overall infection success of the parasite. This result aligns with studies in *Caenorhabditis elegans*, where Dexrazoxane inhibited new spore production without hindering initial spore integration into the host gut cells. Dexrazoxane's DNA topoisomerase II inhibitor mechanism likely explains its impact on mosquito development, as larvae exposed to the drug failed to emerge as adults. These findings highlight Dexrazoxane's potential as a viable tool for controlling microsporidia in adult mosquitoes and hope to enhance the study of mosquito-microsporidia interactions. Further research is required to explore its broader application in vector-borne disease control, including malaria.

Introduction

Microsporidia are common obligate parasites of many animals, and their prevalence has been increasingly reported over the past decades across a wide range of taxa, sparking a surge in microsporidia studies (Becnel and Weiss, 2014; Bojko et al., 2022; Franzen, 2008; Han et al., 2020; Seatamanoch et al., 2022; Wadi and Reinke, 2020; Weidner and Overstreet, 2021). While research in human-infecting microsporidia mainly focuses on clearing them (i.e., microsporidiosis) (Han et al., 2021), in invertebrates such as insects, their study is more nuanced, as evidence suggests microsporidia may serve as biological control agents for certain pests (Bjørnson and Oi, 2014) and vector species, and the diseases they transmit (Becnel and Andreadis, 2014; Bukhari et al., 2022; Koella et al., 2009).

Nearly half of the microsporidia species that infect insects have specialised in targeting dipterans, such as flies and mosquitoes, which are vectors for several human diseases. Among those infecting mosquitoes, *Vavraia culicis* (Silva et al., 2025; Vavra and Becnel, 2007) is one of the most well-studied microsporidia species (Agnew et al., 1999; Biron et al., 2005; Desjardins et al., 2015; Kelly et al., 1981; Rivero et al., 2007; Silva and Koella, 2024a, 2024b; Sy et al., 2014; Vyas-Patel, 2023; Zeferino and Koella, 2024). This generalist intracellular parasite is known to infect several mosquito genera, including *Anopheles gambiae*, the major vector of malaria (Zeferino and Koella, 2024). Infection typically occurs during the larval stage when mosquito larvae ingest spores along with food in water bodies. Once the spores penetrate the host's gut wall, they develop and produce new spores.

The limited immune response of *An. gambiae* to this particular microsporidian species (Silva, 2024)

allows for a steady progression of the infection, characterised by continuous consumption of host

resources and accumulation of spores in the body cavity, leading to costs in host longevity (Agnew et

al., 2004; Zeferino and Koella, 2024) and fecundity (Silva and Koella, 2024a).

Although *An. gambiae* incurs significant costs from *V. culicis* infection, it comes with a benefit in

protection against co-infection with *Plasmodium spp.* by negatively affecting oocyst development

(Bargielowski and Koella, 2009; Lorenz and Koella, 2011). These findings led the community to

hypothesise that microsporidian parasites, such as the one in this study, could be used as biological

control agents for mosquitoes and their parasites by reducing the mosquito lifespan and parasite

transmission to mammal hosts, respectively (Bargielowski and Koella, 2009; Bukhari et al., 2022;

Lorenz and Koella, 2011). However, the mechanisms underlying this interaction remain unclear,

partly due to the technical challenges of manipulating microsporidian infections within mosquito

hosts.

Hence, to facilitate the control and study of microsporidians within the mosquito host, a candidate

chemical, Dexrazoxane, recently shown to have anti-microsporidian activity in *Caenorhabditis*

elegans (Murareanu et al., 2022) was tested. Due to its antimitotic and redox activity, Dexrazoxane

has been extensively used as a protector agent against the cardiotoxicity of chemotherapy in humans

(Vejpongsa and Yeh, 2014). Its mechanism of action is similar to ethylenediaminetetraacetic acid

(EDTA) but diffuses into cells more readily (Hasinoff et al., 1995), where it binds to iron and lowers

oxidative stress formation (Eneh and Lekkala, 2020). However, its role against microsporidia has only

recently been studied. For instance, in *C. elegans*, Dexrazoxane is known to inhibit *Nematocida*

parisii development after it has established itself in the gut cells (Murareanu et al., 2022). While it

does not prevent initial infection, it suppresses the production of new spores, thereby hindering

disease progression and transmission. Given its promising effects on microsporidia infection in *C.*

elegans, we aimed to test whether these effects extend to microsporidian infections in mosquitoes,

using *An. gambiae* and *V. culicis* as the experimental system. Our findings suggest that Dexrazoxane's

effect might be applicable to more model systems than previously thought, extending to mosquito-

microsporidia interactions. This result may enhance our understanding of microsporidia interactions

with their host and other parasites and could potentially inform and fuel future vector-borne disease

control strategies.

Figure 1. Illustration of the experimental design. Dexrazoxane's effect on spore load and mortality was tested on both larval **(a)** and adult stages **(b)**. In the first experiment, one of three doses (0 µM, 30 µM, or 60 µM) was administered at one of three time-points (4-, 5-, or 6-days post eclosion (DPE)) to infected and uninfected larvae, in a fully combinatorial manner. In the second experiment, we tested for the effect of exposure to dexrazoxane during the adult stage. For this, we supplemented their sugar diet with 60 µM of Dexrazoxane, or not, and tested for its impact on the spore load when reared individually or collectively, and infected or not. Values between brackets represent mortality for that treatment. For more details, see Materials and Methods (section 2).

Materials and Methods

1. Experimental system

We used the *An. gambiae s.s* Kisumu strain (Vulule et al., 1994), which had been maintained at our standard laboratory conditions (about 600 individuals per cage, constant access to 6% sucrose

solution, 26 ± 1ºC, 70 ± 5% relative humidity and 12 h light/dark) for many years before the
experiments.

The microsporidian *V. culicis floridensis* was provided by J.J Becnel (USDA, Gainesville, FL, USA).
Originally discovered in *Ae. albopictus,* this parasite was later found to parasitise several mosquito
genera, including *Aedes*, *Culex*, *Anopheles*, *Culiseta*, *Ochlerotatus* and *Orthopodomyia* (Andreadis,
2007; Fukuda et al., 1997; Kelly et al., 1981). We maintained it by alternately infecting *Aedes aegypti*
and *An. gambiae* to ensure its status as a generalist parasite, as described previously (Zeferino and
Koella, 2024).

V. culicis is an obligatory, intracellular parasite. Mosquitoes are infected when, as larvae, they ingest
the spores along with their food. *V. culicis* culture and experiments described below used a dose of
10,000 spores per larva, as previous preliminary experiments have shown that it infects 100% of the
mosquitoes at this dose. After several rounds of replication, the parasite produces new infectious
spores that spread to the gut and fat body cells. The spores are released to larval sites when infected
larvae die, adults die on the surface of water, or eggs covered with spores are laid onto the water's
surface.

2. Rearing and maintenance of experimental mosquitoes

Freshly hatched (0-3 hours old) *An. gambiae* larvae were individually placed into 12-well culture
plates, each containing 3 ml of deionized water. The larvae received Tetramin Baby® fish food every
day according to their age (0.04, 0.06, 0.08, 0.16, 0.32 and 0.6 mg/larva respectively on days 0, 1, 2,
3, 4 and 5 or older (Kulma et al., 2013)).

2.1 Larvae exposure to Dexrazoxane

In the first experiment **(Fig. 1)**, we assessed the impact of Dexrazoxane (Cat# 05587, Sigma-Aldrich,
St. Louis, MO, USA) on larval development and spore growth. Since the mosquito larvae were reared
in deionized water with Dexrazoxane, we hypothesized that a long exposure time might differently
affect the mosquito fitness traits. Therefore, we not only tested three different doses (0 µM, 30 µM,
and 60 µM), but also tested for the impact of one administration at one of three time points (four, five,
or six days after egg hatching) for infected and uninfected larvae, in a fully combinatorial design. The
larvae's survival and pupation were tracked daily. Since mosquitoes only have detectable sporulation
in the adult stage (Silva and Koella, 2024a), spores were uncountable for this experiment.

2.2 Adult exposure to Dexrazoxane

In a second experiment **(Fig. 1)**, we assessed the impact of Dexrazoxane on: i) spore growth at the

adult stage, and ii) if being reared individually (Individual) as opposed to collectively could influence

this growth (Collective). The latter is to test if Dexrazoxane needs to be administered individually or

if it has the same effect when provided to a group of mosquitoes (with higher stress due to higher

density). Two-day-old larvae were exposed to either 0 or 10,000 spores of *V. culicis*. Pupae were

transferred to individual 50 ml falcon tubes with approximately 10 ml of deionised water (Dao et al.,

2010). Female adults were then transferred to different containers based on their rearing treatment:

those reared individually were placed into separate 150 ml cups (5.5 cm Ø x 10 cm) covered with a

net, while those from the collective rearing treatment were housed together in a cage (21 cm x 21 cm

x 21 cm). The cups contained a 50 mm Ø petri dish (to prevent mosquitoes from drowning) floating

on the surface of 50 ml deionised water and a 10 x 7 cm rectangular filter paper to maintain humidity.

The mosquitoes were provided with cotton balls soaked in a 6% sucrose solution or a 6% sucrose

solution supplemented with 60 µM of Dexrazoxane, which was replaced every other day. We used

only one dose due to technical constraints. Unlike the previous experiment on larvae, we allowed

treated adult mosquitoes to feed on the Dexrazoxane-sugar-soaked cotton for eight days. The

reasoning behind this decision was that while at the larval stage, they are constantly ingesting and

exposed to Dexrazoxane, in the adult stage, the exposure is not constant but dependent on the

mosquito's motivation to feed and, therefore, extending the exposure time will maximize the ingestion

of Dexrazoxane by the adult mosquitoes.

3. Statistical analysis

All analyses were conducted with the R software (Team, 2020) version 4.4.1, using the packages

DHARMa (Hartig and Hartig, 2017), car (Fox et al., 2012), glmmTMB (Brooks et al., 2017),

emmeans (Lenth et al., 2019) and multcomp (Hothorn et al., 2016). Significance was assessed with

the "Anova" function of the "car" package (Fox et al., 2012). We used a type III ANOVA in the case

of a significant interaction and a type II ANOVA otherwise. When relevant, we performed post-hoc

multiple comparisons with the package "emmeans", using the default Tukey adjustment.

The presence of spores was coded as a binary response (presence or absence of detectable spores) and

analysed with a generalised linear model with a binomial distribution of errors, where the explanatory

variables were diet (with or without Dexrazoxane), type or rearing (individual or collective) and their

interaction. Because we counted the number of spores in a haemocytometer containing 0.1 µl of the

sample (i.e. 1/1000 of the total volume), the detection threshold was estimated to be 1000 spores.

Spore load of mosquitoes with positive spore counts was analysed with a generalised linear model

with a negative binomial distribution of errors. The explanatory variables were diet (with or without

Dexrazoxane), type or rearing (individual or collective), and their interaction.

Results

Larval exposure to Dexrazoxane delayed larval development and blocked pupation, independently of
the dose, with none of the treated larvae surviving past pupation (as opposed to Dexrazoxane-
unexposed larvae). That led us to discard the use of this chemical during the larval stage.

The proportion of individuals harbouring spores was lower for the individual treatment than for the
collective (0.65 vs 0.85, χ^2 = 12.25, df = 1, p < 0.001, **Fig. 2a**) but was not affected by the
consumption of Dexrazoxane (0.75 vs 0.75, χ^2 = 0.00, df = 1, p = 0.945) or their interaction (χ^2 = 0.28,
df = 1, p = 0.595). In contrast, the spore load of mosquitoes with positive counts was reduced by
approximately 59% in individuals consuming Dexrazoxane, but only when reared individually (4.9 x
10^4 vs 1.2 x 10^5 in individual rearing, and 9.5 x 10^4 vs 6.7 x 10^4 in collective rearing, interaction type
of rearing * diet: χ^2 = 10.35, df = 1, p = 0.001, **Fig. 2b**). For further details on the statistical analysis
see **Table 1**.

Table 1. Effect of exposure to Dexrazoxane on infection. Statistical models carried on to test for
effect of exposure to Dexrazoxane, rearing condition (individual in cup, or collective in cage) and
their interaction on the presence of spores (Model 1a) and number of spores (Model 1b). See
Materials and methods for more details on the experimental design and statistical tests. Values in bold
are statistically significant.

Tested effect			
Model 1a – Presence of spores	df	χ^2	*p*
Exposure to Dexrazoxane	1	0.00	0.945
Rearing (Individual/Collective)	1	12.25	**<0.001**
Exposure to Dexrazoxane: Rearing (Individual/Collective)	1	0.28	0.595
Model 1b – Spore load	df	χ^2	*p*
Exposure to Dexrazoxane	1	2.13	0.144
Rearing (Individual/Collective)	1	4.68	**0.030**
Exposure to Dexrazoxane: Rearing (Individual/Collective)	1	10.35	**0.001**

Figure 2. Parasite load of infected mosquitoes after consuming a specific diet for seven days. The proportion of mosquitoes reared individually or collectively that had detectable spores for **(a)** each Dexrazoxane diet and their spore load **(b)**. The sample sizes were 70, 60, 60 and 60 for **(a)**, and 60, 50, 38 and 40 for **(b)**. Error bars show the 95% confidence intervals of the means. Letters indicate

197 statistically significant differences from multiple comparisons. See Table 1 for further details on the
198 statistical analyses.

Discussion

Our study investigates the effect of Dexrazoxane on the infection by the microsporidian *V. culicis* in
the mosquito host *An. gambiae*. The findings extend the current knowledge of microsporidia-
mosquito interactions and suggest potential strategies for controlling microsporidia infections and
mosquito-borne diseases like malaria.

The primary finding from this study is that Dexrazoxane significantly reduces the spore load of
infected mosquitoes but only under certain rearing conditions, mainly when mosquitoes are raised
individually during the adult stage **(Fig. 2b)**. Moreover, although the total number of spores produced
decreased, the number of successful infections remained unchanged compared to unexposed
individuals **(Fig. 2a)**. This finding is consistent with previous research in *C. elegans* (Murareanu et
al., 2022), which reported that Dexrazoxane inhibits microsporidian development primarily by
reducing spore production. However, further studies are needed to determine whether Dexrazoxane
acts by limiting the initial integration of spores into the host gut wall or by suppressing subsequent
spore proliferation, as both mechanisms could account for the observed reduction in spore load.
Importantly, this decline in spore numbers did not come at the cost of host survival—100% of the
mosquitoes exposed to Dexrazoxane survived the entire duration of the experiment and were
sacrificed only at its conclusion for spore quantification.

For future experimental purposes, we tested for the effect of the Dexrazoxane when administrated
individually or collectively. Collective administration of Dexrazoxane has two main advantages: it
requires less Dexrazoxane (as administration still happens through supplemented-sugar-cottons) and
allows for experiments to be conducted in cages, which often can ease their feasibility. Our results
show that Dexrazoxane works at these concentrations but is only administered individually.
Nevertheless, we cannot discard the fact that (for instance) higher concentrations of Dexrazoxane
administered collectively could lead to similar outcomes.

The reduced spore load in mosquitoes raised individually in cups highlights a potential indirect effect
of Dexrazoxane on the mosquito immune response or microsporidian metabolism. While any
hypothesis cannot be ruled out, this inhibition is likely due to the drug's mechanism of action. For
human purposes, Dexrazoxane primarily functions through iron chelation, where it blocks circulating
iron from being used to fuel oxidative stress and tissue damage. Although this mechanism has been
shown to affect *Plasmodium falciparium* development *in vitro* (Loyevsky et al., 1999), the same
cannot be said for microsporidia. In fact, *C. elegans* infection with the microsporidia *N. parisii* can be

blocked using Dexrazoxane but cannot be reverted through iron supplementation (Murareanu et al.,
2022). This result suggests inhibition in microsporidian infections might not happen via iron chelation
but instead through another mechanism of action. Interestingly, Dexrazoxane also acts as a DNA
topoisomerase II inhibitor, which likely explains why larvae exposed to the drug failed to emerge as
adults and had their development arrested at the pupa stage. Given the mechanism of action of
Dexrazoxane is fair to assume that this DNA topoisomerase II might be particularly relevant for
metamorphosis and transition to the adult stage. In agreement with this, a study on *Ae. aegypti*, larvae
treated with etoposide (another DNA topoisomerase II inhibitor) exhibited morphological defects and
low survival during both larval and pupal stages (Santos et al., 2021), potentially pointing to a similar
result. Therefore, it is plausible that *V. culicis* relies on the same protein or a structurally similar one,
and inhibiting this protein during the parasite growth phase might block spore production and
infection progression. Although intuitively, this positions Dexrazoxane as a possible larvicide agent,
there are a couple of reasons why it might not be the best solution to control mosquito populations.
First, our study and others have demonstrated that Dexrazoxane might act at the most basic cellular
mechanisms that most animals share. Therefore, there is a long-term potential for environmental
damage if this drug is used to target the mosquito larval stage. The use of Dexrazoxane in water
bodies where mosquitoes live is likely to impede the correct development of other arthropods or even
fish that share the environment. Second, it is crucial to consider that in these experiments,
Dexrazoxane was used at relatively high concentrations but in small volumes of water. In the wild,
water bodies would have much higher volumes of water, which would raise the cost of the control
strategy.

It is important to note that the context of mosquito hosts introduces additional complexity compared
to previous studies performed *in vitro* or with *C. elegans* (Loyevsky et al., 1999; Murareanu et al.,
2022). Dexrazoxane's reduction in spore load, especially in isolated conditions, suggests that other
factors, such as social interactions and spore transmission, may influence its efficacy. These factors
should be carefully controlled and studied in future research, along with determining the optimal
dosage. Despite these complexities, the results of this study open the door for further investigation
into microsporidia as potential biological control agents, particularly for *V. culicis*. Moreover, the
strategic timing of the suppression of the infection progression, without completely clearing it, offers
several advantages for studying interactions with vector-borne parasites, such as *Plasmodium*. We
believe this approach will allow researchers to investigate the mechanisms behind microsporidia-
driven inhibition, for instance, whether it is driven by collateral immunity or resource competition. A
further understanding of these mechanisms will allow us to better prepare and design vector control
strategies, which are essential today.

References

Agnew, P., Bedhomme, S., Haussy, C., Michalakis, Y., 1999. Age and size at maturity of the mosquito Culex pipiens infected by the microsporidian parasite Vavraia culicis. Proc. R. Soc. London. Ser. B Biol. Sci. 266, 947–952. https://doi.org/https://doi.org/10.1098/rspb.1999.0728

Agnew, P., Berticat, C., Bedhomme, S., Sidobre, C., Michalakis, Y., 2004. Parasitism increases and decreases the costs of insecticide resistance in mosquitoes. Evolution (N. Y). 58, 579–586. https://doi.org/https://doi.org/10.1111/j.0014-3820.2004.tb01680.x

Andreadis, T.G., 2007. Microsporidian parasites of mosquitoes. J. Am. Mosq. Control Assoc. 23, 3–29. https://doi.org/https://doi.org/10.2987/8756-971X(2007)23[3:MPOM]2.0.CO;2

Bargielowski, I., Koella, J.C., 2009. A possible mechanism for the suppression of Plasmodium berghei development in the mosquito Anopheles gambiae by the microsporidian Vavraia culicis. PLoS One 4, e4676. https://doi.org/https://doi.org/10.1371/journal.pone.0004676

Becnel, J.J., Andreadis, T.G., 2014. Microsporidia in insects. Microsporidia Pathog. Oppor. 521–570. https://doi.org/https://doi.org/10.1002/9781118395264

Becnel, J.J., Weiss, L.M., 2014. Microsporidia: pathogens of opportunity. John Wiley & Sons Incorporated. https://doi.org/https://doi.org/10.1002/9781118395264

Biron, D.G., Agnew, P., Marche, L., Renault, L., Sidobre, C., Michalakis, Y., 2005. Proteome of Aedes aegypti larvae in response to infection by the intracellular parasite Vavraia culicis. Int. J. Parasitol. 35, 1385–1397. https://doi.org/https://doi.org/10.1016/j.ijpara.2005.05.015

Bjørnson, S., Oi, D., 2014. Microsporidia biological control agents and pathogens of beneficial insects. Microsporidia Pathog. Oppor. 635–670. https://doi.org/https://doi.org/10.1002/9781118395264

Bojko, J., Reinke, A.W., Stentiford, G.D., Williams, B., Rogers, M.S.J., Bass, D., 2022. Microsporidia: a new taxonomic, evolutionary, and ecological synthesis. Trends Parasitol. 38, 642–659. https://doi.org/https://doi.org/10.1016/j.pt.2022.05.007

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A., Skaug, H.J., Machler, M., Bolker, B.M., 2017. glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. R J. 9, 378–400. https://doi.org/https://doi.org/10.32614/RJ-2017-066

Bukhari, T., Pevsner, R., Herren, J.K., 2022. Microsporidia: a promising vector control tool for residual malaria transmission. Front. Trop. Dis. 3, 957109. https://doi.org/https://doi.org/10.3389/fitd.2022.957109

Dao, A., Kassogue, Y., Adamou, A., Diallo, M., Yaro, A.S., Traore, S.F., Lehmann, T., 2010.
Reproduction-longevity trade-off in Anopheles gambiae (Diptera: Culicidae). J. Med. Entomol.
47, 769–777.

Deng, S., Yan, T., Jendrny, C., Nemecek, A., Vincetic, M., Gödtel-Armbrust, U., Wojnowski, L.,
2014. Dexrazoxane may prevent doxorubicin-induced DNA damage via depleting both
topoisomerase II isoforms. BMC Cancer 14, 1–11.

Desjardins, C.A., Sanscrainte, N.D., Goldberg, J.M., Heiman, D., Young, S., Zeng, Q., Madhani,
H.D., Becnel, J.J., Cuomo, C.A., 2015. Contrasting host–pathogen interactions and genome
evolution in two generalist and specialist microsporidian pathogens of mosquitoes. Nat.
Commun. 6, 7121. https://doi.org/https://doi.org/10.1038/ncomms8121

Eneh, C., Lekkala, M.R., 2020. Dexrazoxane.

Fox, J., Weisberg, S., Adler, D., Bates, D., Baud-Bovy, G., Ellison, S., Firth, D., Friendly, M.,
Gorjanc, G., Graves, S., 2012. Package 'car.' Vienna R Found. Stat. Comput. 16.
https://doi.org/https://doi.org/10.32614/CRAN.package.car

Franzen, C., 2008. Microsporidia: a review of 150 years of research. Open Parasitol. J. 2.
https://doi.org/https://doi.org/10.2174/1874421400802010001

Fukuda, T., Willis, O.R., Barnard, D.R., 1997. Parasites of the Asian tiger mosquito and other
container-inhabiting mosquitoes (Diptera: Culicidae) in northcentral Florida. J. Med. Entomol.
34, 226–233. https://doi.org/https://doi.org/10.1093/jmedent/34.2.226

Han, B., Pan, G., Weiss, L.M., 2021. Microsporidiosis in humans. Clin. Microbiol. Rev. 34, e00010-
20. https://doi.org/https://doi.org/10.1128/CMR.00010-20

Han, B., Takvorian, P.M., Weiss, L.M., 2020. Invasion of host cells by microsporidia. Front.
Microbiol. 11, 172. https://doi.org/https://doi.org/10.3389/fmicb.2020.00172

Hartig, F., Hartig, M.F., 2017. Package 'DHARMa.' R Packag.
https://doi.org/https://doi.org/10.32614/CRAN.package.DHARMa

Hasinoff, B.B., Kuschak, T.I., Yalowich, J.C., Creighton, A.M., 1995. A QSAR study comparing the
cytotoxicity and DNA topoisomerase II inhibitory effects of bisdioxopiperazine analogs of
ICRF-187 (dexrazoxane). Biochem. Pharmacol. 50, 953–958.
https://doi.org/https://doi.org/10.1016/0006-2952(95)00218-O

Hothorn, T., Bretz, F., Westfall, P., Heiberger, R.M., Schuetzenmeister, A., Scheibe, S., Hothorn,
M.T., 2016. Package 'multcomp.' Simultaneous inference Gen. Parametr. Model. Proj. Stat.
Comput. Vienna, Austria. https://doi.org/https://doi.org/10.32614/CRAN.package.multcomp

Kelly, J.F., Anthony, D.W., Dillard, C.R., 1981. A laboratory evaluation of the microsporidian

Vavraia culicis as an agent for mosquito control. J. Invertebr. Pathol. 37, 117–122.

https://doi.org/https://doi.org/10.1016/0022-2011(81)90064-1

Koella, J.C., Lorenz, L., Bargielowski, I., 2009. Microsporidians as evolution☐proof agents of

malaria control? Adv. Parasitol. 68, 315–327. https://doi.org/https://doi.org/10.1016/S0065-

308X(08)00612-X

Kulma, K., Saddler, A., Koella, J.C., 2013. Effects of age and larval nutrition on phenotypic

expression of insecticide-resistance in Anopheles mosquitoes. PLoS One 8, e58322.

https://doi.org/https://doi.org/10.1371/journal.pone.0058322

Lenth, R., Singmann, H., Love, J., Buerkner, P., Herve, M., 2019. Package 'emmeans.' R Packag.

version 1. https://doi.org/https://doi.org/10.32614/CRAN.package.emmeans

Lorenz, L.M., Koella, J.C., 2011. The microsporidian parasite Vavraia culicis as a potential late life–

acting control agent of malaria. Evol. Appl. 4, 783–790.

https://doi.org/https://doi.org/10.1111/j.1752-4571.2011.00199.x

Loyevsky, M., Sacci Jr, J.B., Boehme, P., Weglicki, W., John, C., Gordeuk, V.R., 1999. Plasmodium

falciparum and Plasmodium yoelii: Effect of the Iron Chelation Prodrug Dexrazoxane on in

Vitro Cultures. Exp. Parasitol. 91, 105–114.

https://doi.org/https://doi.org/10.1006/expr.1998.4371

Murareanu, B.M., Antao, N. V, Zhao, W., Dubuffet, A., El Alaoui, H., Knox, J., Ekiert, D.C., Bhabha,

G., Roy, P.J., Reinke, A.W., 2022. High-throughput small molecule screen identifies inhibitors

of microsporidia invasion and proliferation in C. elegans. Nat. Commun. 13, 5653.

https://doi.org/https://doi.org/10.1038/s41467-022-33400-y

Rivero, A., Agnew, P., Bedhomme, S., Sidobre, C., Michalakis, Y., 2007. Resource depletion in

Aedes aegypti mosquitoes infected by the microsporidia Vavraia culicis. Parasitology 134,

1355–1362. https://doi.org/https://doi.org/10.1017/S0031182007002703

Santos, D., Gomes, H.F., Ribeiro, L., Farias, A.B., Romeiro, N.C., da Fonseca, R.N., Nepomuceno-

Silva, J.L., Moraes, J., 2021. Inhibition of Aedes aegypti DNA topoisomerase II by etoposide:

Impact on survival and morphology of larvae and pupae. Comp. Biochem. Physiol. Part C

Toxicol. Pharmacol. 247, 109066. https://doi.org/https://doi.org/10.1016/j.cbpc.2021.109066

Seatamanoch, N., Kongdachalert, S., Sunantaraporn, S., Siriyasatien, P., Brownell, N., 2022.

Microsporidia, a highly adaptive organism and its host expansion to humans. Front. Cell. Infect.

Microbiol. 12, 924007. https://doi.org/https://doi.org/10.3389/fcimb.2022.924007

Silva, L.M., 2024. The transcriptome of the mosquito host Anopheles gambiae upon infection by
different selected lines of the microsporidian parasite Vavraia culicis. bioRxiv 2009–2024.
https://doi.org/https://doi.org/10.1101/2024.09.22.613703

Silva, L.M., Koella, J.C., 2024a. Complex interactions in the life cycle of a simple parasite shape the
evolution of virulence. bioRxiv 2024.01.28.577571. https://doi.org/10.1101/2024.01.28.577571

Silva, L.M., Koella, J.C., 2024b. Virulence evolution: thinking outside of the host. bioRxiv 2005–
2024. https://doi.org/https://doi.org/10.1101/2024.05.23.595559

Silva, L.M., Zeferino, T.G., Koella, J.C., 2025. Vavraia culicis. Trends Parasitol.
https://doi.org/https://doi.org/10.1016/j.pt.2025.02.011

Sy, V.E., Agnew, P., Sidobre, C., Michalakis, Y., 2014. Reduced survival and reproductive success
generates selection pressure for the dengue mosquito Aedes aegypti to evolve resistance against
infection by the microsporidian parasite Vavraia culicis. Evol. Appl. 7, 468–479.
https://doi.org/https://doi.org/10.1111/eva.12144

Team, R.C., 2020. RA language and environment for statistical computing, R Foundation for
Statistical. Computing.

Vavra, J., Becnel, J.J., 2007. Vavraia culicis (Weiser, 1947) Weiser, 1977 revisited: cytological
characterisation of a Vavraia culicis-like microsporidium isolated from mosquitoes in Florida
and the establishment of Vavraia culicis floridensis subsp. n. Folia Parasitol. (Praha). 54, 259.
https://doi.org/https://doi.org/10.14411/fp.2007.034

Vejpongsa, P., Yeh, E.T.H., 2014. Prevention of anthracycline-induced cardiotoxicity: challenges and
opportunities. J. Am. Coll. Cardiol. 64, 938–945.
https://doi.org/https://doi.org/10.1016/j.jacc.2014.06.1167

Vulule, J.M., Beach, R.F., Atieli, F.K., Roberts, J.M., Mount, D.L., Mwangi, R.W., 1994. Reduced
susceptibility of Anopheles gambiae to permethrin associated with the use of
permethrin□impregnated bednets and curtains in Kenya. Med. Vet. Entomol. 8, 71–75.
https://doi.org/https://doi.org/10.1111/j.1365-2915.1994.tb00389.x

Vyas-Patel, N., 2023. The Suppression of Plasmodium berghei in Anopheles coluzzii infected later
with Vavraia culicis. bioRxiv 2002–2023.
https://doi.org/https://doi.org/10.1101/2023.02.05.527158

Wadi, L., Reinke, A.W., 2020. Evolution of microsporidia: An extremely successful group of
eukaryotic intracellular parasites. PLoS Pathog. 16, e1008276.
https://doi.org/https://doi.org/10.1371/journal.ppat.1008276

Weidner, E., Overstreet, R., 2021. Microsporidia have a peculiar outer membrane with exterior

cytoplasmic proteins. Acta Parasitol. 1–6. https://doi.org/https://doi.org/10.1007/s11686-021-

00446-9

Zeferino, T.G., Koella, J.C., 2024. Host-specific effects of a generalist parasite of mosquitoes. Sci.

Rep. 14, 18365. https://doi.org/https://doi.org/10.1038/s41598-024-69475-4

Acknowledgements

400 We thank Gwendoline Acerbi for her technical support and Jacob C. Koella for early feedback on the

401 project. LMS and TGZ were supported by SNF grant 310030_192786.

Author contributions

403 LMS and TGZ designed and conducted the experiments, analysed the data and wrote the manuscript.

Data availability

405 All data generated or analysed during this study are included as Supplementary Information files.

Additional Information

407 The authors declare no competing interests.

Figure legends

Figure 1. Illustration of the experimental design. Dexrazoxane's effect on spore load and mortality

was tested on both larval **(a)** and adult stages **(b)**. In the first experiment, one of three doses (0 µM, 30

µM, or 60 µM) was administered at one of three time-points (4-, 5-, or 6-days post eclosion (DPE)) to

infected and uninfected larvae, in a fully combinatorial manner. In the second experiment, we tested

for the effect of exposure to dexrazoxane during the adult stage. For this, we supplemented their sugar

diet with 60 µM of Dexrazoxane, or not, and tested for its impact on the spore load when reared

individually or collectively, and infected or not. Values between brackets represent mortality for that

treatment. For more details, see Materials and Methods (section 2).

Figure 2. Parasite load of infected mosquitoes after consuming a specific diet for 7 days. The

proportion of mosquitoes reared individually or collectively that had detectable spores for **(a)** each

Dexrazoxane diet, and their spore load **(b)**. The sample sizes were 70, 60, 60 and 60 for **(a)**, and 60,

50, 38 and 40 for **(b)**. Error bars show the 95% confidence intervals of the means. Letters indicate

statistically significant differences from multiple comparisons. See Table 1 for further details on the
statistical analyses.

Table 1. Effect of exposure to Dexrazoxane on infection. Statistical models carried on to test for
effect of exposure to Dexrazoxane, rearing condition (individual in cup, or collective in cage) and
their interaction on the presence of spores (Model 1a) and number of spores (Model 1b). See
Materials and Methods for more details on the experimental design and statistical tests. Values in bold
are statistically significant.

	Larvae *(Exp. 1)*			**Adults** *(Exp. 2)*	
	Administration time (DPE)			Exposure for 8 days (rearing)	
Concentration (µM)	4	5	6	Individual	Collective
0	(0%) (0%)	(0%) (0%)	(0%) (0%)	(0%) (0%)	(0%) (0%)
30	(100%) (100%)	(100%) (100%)	(100%) (100%)		
60	(100%) (100%)	(100%) (100%)	(100%) (100%)	(0%) (0%)	(0%) (0%)

 Uninfected

 Infected