Structure of the turnip yellows virus particles: How is the readthrough domain exposed on the capsid surface?

Joséphine Lai-Kee-Him [a], Stefano Trapani [b,*], Sylvaine Boissinot [c,1], Catherine Reinbold [d], Chloé Fallet [e], Aurélie Ancelin [f], François Lecorre [g], François Hoh [h], Véronique Ziegler-Graff [i], Véronique Brault [j,*] and Patrick Bron [k,*]

[a] Centre de Biologie Structurale (CBS), Univ Montpellier, CNRS, INSERM, Montpellier, France; josephine.laikeehim@cbs.cnrs.fr

[b] Centre de Biologie Structurale (CBS), Univ Montpellier, CNRS, INSERM, Montpellier, France; stefano.trapani@umontpellier.fr

[c] INRAE, Université de Strasbourg, UMR SVQV, Colmar France; sylvaine.boissinot@inrae.fr

[d] INRAE, Université de Strasbourg, UMR SVQV, Colmar France; catherine.reinbold@inrae.fr

[e] Centre de Biologie Structurale (CBS), Univ Montpellier, CNRS, INSERM, Montpellier, France.

[f] Centre de Biologie Structurale (CBS), Univ Montpellier, CNRS, INSERM, Montpellier, France; aurelie.ancelin@cbs.cnrs.fr

[g] Centre de Biologie Structurale (CBS), Univ Montpellier, CNRS, INSERM, Montpellier, France; francoislecorre.ar@gmail.com

[h] Centre de Biologie Structurale (CBS), Univ Montpellier, CNRS, INSERM, Montpellier, France; francois.hoh@cnrs.fr

[i] Institut de biologie moléculaire des plantes, CNRS, Université de Strasbourg, Strasbourg, France; veronique.ziegler-graff@ibmp-cnrs.unistra.fr

[j] INRAE, Université de Strasbourg, UMR SVQV, Colmar France; veronique.brault@inrae.fr

[k] Centre de Biologie Structurale (CBS), Univ Montpellier, CNRS, INSERM, Montpellier, France; patrick.bron@cbs.cnrs.fr

[1] Current address: INRAE, BFP UMR 1332, Villenave d'Ornon, France

[*] Corresponding authors

Abstract

Turnip yellows virus (TuYV) is a plant virus infecting important crops such as oilseed rape. TuYV is phloem-restricted and transmitted by aphids. The capsid contains two subunit types: the major capsid protein (CP) and a minor component (RTP*) which arises from the C-terminal cleavage of a readthrough product (RTP). RTP* contains the CP sequence fused with a structured domain, denoted ^{N}RTD, which is a key determinant of virus transmission. Though both CP and RTP* are involved in virus movement and aphid transmission, how RTP* is incorporated into the capsid is poorly understood. We present here the structural characterisation, by immunogold labelling and 3D cryo-EM, of the wild-type TuYV and a mutant whose capsid contains the CP only. We show that incorporation of RTP* does not impair the capsid structure, and the ^{N}RTD does not adopt well-defined positions at the capsid surface. The number of incorporated RTP*s suggests a random insertion.

Keywords: polerovirus, readthrough domain, capsid, elevated dodecahedron, structure, electron cryo-microscopy, immunogold labelling

Introduction

Turnip yellows virus (TuYV), formerly beet western yellows virus, is a member of the genus *Polerovirus*. Members of the genera *Polerovirus* and *Enamovirus* (family *Solemoviridae*) and *Luteovirus* (family *Tombusviridae*), hereafter referred to as P/E/L viruses (Schiltz et al., 2022), are important plant pathogens that affect many major crops (Heck and Brault, 2018). Their economic importance extends worldwide to small grains, potato, brassicas, cotton, cucurbits, pepper, sugarcane and sorghum. P/E/L viruses are phloem-restricted and transmitted exclusively by aphids in nature (Gildow, 1999). The P/E/L genome is a linear single-stranded positive-sense RNA with an intricate organisation of the open reading frames (ORFs) (Boissinot et al., 2020; Smirnova et al., 2015; Sõmera et al., 2021) (Fig. 1a). The genome is encapsidated into a non-enveloped particle of ~30 nm in diameter composed of 180 subunits arranged according to a *T*=3 icosahedral symmetry. Most of the 180 subunits are the major capsid protein (CP) of about 22-25 kDa encoded by ORF3. However, some of the subunits (10 to 25%) contain a C-terminal extension which protrudes at the virion surface (Brault et al., 1995; Filichkin et al., 1994). This minor capsid protein, hereafter called RTP* (~55 kDa), arises from stochastic translational readthrough of the CP stop codon, whereby a non-structural readthrough protein (RTP) of about 74 kDa is produced (Fig. 1b). The RTP contains the CP sequence on the N-ter side, followed by a proline-rich flexible linker and a readthrough domain (RTD). This is composed of a structured sub-domain, denoted [N]RTD (Schiltz et al., 2022), and a C-ter sub-domain, denoted [C]RTD, which is predicted to be intrinsically disordered (LaTourrette et al., 2021). Post-translational cleavage of the [C]RTD stretch from the RTP results in the structural RTP* which is incorporated in the capsids (Fig. 1b).

The role of RTP, RTP* and CP in P/E/L virus movement and transmission by aphids has been extensively addressed, based mainly on reverse genetics on TuYV, potato leafroll virus (PLRV, genus *Polerovirus*), barley yellow dwarf virus (BYDV, genera *Luteovirus* and *Polerovirus* depending on the species) and cucurbit aphid-borne yellows virus (CABYV, genus *Polerovirus*). The full-length non-structural RTP is involved in virus systemic invasion of the plant (Boissinot et al., 2014; Brault et al., 1995; Hipper et al., 2014; Kaplan et al., 2007; Lee et al., 2005; Mutterer et al., 1999; Peter et al., 2008; Rodriguez-Medina et al., 2015; Wang et al., 1995; Xu et al., 2018) and the terminal [C]RTD plays a role in restricting PLRV to the phloem (Peter et al., 2009) and in symptom development (Bruyère et al., 1997; Rodriguez-Medina et al., 2015). The RTP* is not required for capsid assembly (Bruyère et al., 1997; Reutenauer et al., 1993), but the presence of both CP and RTP* in intact virions is required for the virus to move long distance within host plants and for transmission by aphid vectors (Brault et al., 1995; Chay et al., 1996; Hipper et al., 2014; Lee et al., 2005). It has been shown that both TuYV structural proteins interact with the ephrin receptor, an aphid protein involved in virus uptake and transmission (Mulot et al., 2018). Also, the [N]RTD of RTP* has been identified as a key determinant of transmission as mutant viruses devoid of the whole RTD are not aphid transmissible and retain the ability to infect plants only at a reduced titre (Brault et al., 1995; Chay et al., 1996; Liang et al., 2004; Reutenauer et al., 1993). Conversely, mutants lacking just the [C]RTD assemble efficiently into virions (Boissinot et al., 2014; Bruyère et al., 1997), retain the ability to interact with aphid proteins (Mulot et al., 2018) and can be

transmitted to new hosts (Wang et al., 1995). *In vitro* and *in vivo* binding assays also revealed that TuYV and PLRV virions bind to plant components (Bencharki et al., 2010; DeBlasio et al., 2016, 2015a, 2015b).

The structure of P/E/L capsids and how they interact with aphid vectors have been the object of investigation for several years. Difficulties in producing enough purified viruses had long limited structural studies to homology modelling coupled to mutational and cross-linking experiments (Alexander et al., 2017; Brault et al., 2003; Chavez et al., 2012; Kaplan et al., 2007; Terradot et al., 2001; Torrance, 1992). Significant advances were made after the production of BYDV and PLRV virus-like particles (VLPs) containing the recombinant CP, but not the RTP*, and the establishment of their atomic structure by electron cryogenic microscopy (cryo-EM) (Byrne et al., 2019). Also, the crystal structure of the [N]RTD, absent from the VLPs, was solved using recombinant forms of the isolated domains from PLRV and TuYV (Schiltz et al., 2022). These studies revealed that P/E/L viruses are structurally related to picorna-like viruses, corrected errors in previously proposed structural models, and rationalised the observations from previous mutagenesis studies. The [N]RTD structure revealed a dimeric architecture in which the sequence of each monomer goes back and forth from a jell-roll domain to a cap domain. The [N]RTD jelly roll was found to be structurally related to the constitutively expressed capsid protrusions (P domains) of tombusviruses, while the cap domain is absent from those viruses. A model, based on the tombusvirus structure, of how the [N]RTD might be presented on the surface of P/E/L viruses was proposed (Schiltz et al., 2022).

To understand if and how the presence of the RTP* affects the P/E/L capsid structure and how the [N]RTD is presented on the capsid surface, we have purified and studied two forms of TuYV particles: the wild-type virus (TuYV-WT, GenBank accession NC_003743.1, Veidt et al. (1988)), whose viral capsids contain both CP and RTP*, and a mutant (TuYV-ΔRTP) whose capsids contain the CP only. The two forms have been observed and characterised by electron microscopy (EM) techniques: negative staining, immunogold labelling and 3D cryo-EM. In this paper we present and discuss the results of our observations.

Materials and Methods

Virus production and purification

Purified virions were prepared from 320 g of frozen leaf material of *N. benthamiana* infiltrated with TuYV-WT full-length infectious clone (Leiser et al., 1992) using agroinfiltration (English et al., 1997) and from 335 g of *Montia perfoliata* similarly infiltrated with TuYV-ΔRTP infectious clone (referred to as BW6.4 in Brault et al (1995)). Infiltrated leaves were collected and frozen at –80 °C 6 to 8 days post-inoculation. The purification was performed according to van den Heuvel et al. (1991) and completed by a sucrose density gradient centrifugation followed by ultracentrifugation at 45 000 rpm during 2 h 30 min at 17 °C to obtain a highly concentrated suspension of virions. The final pellet was suspended in citrate buffer 0.1 M pH 6.0 and stored at –80 °C. The final concentration reached 1.32 µg/µL and 3.5 µg/µL for TuYV-WT and TuYV-ΔRTP respectively.

Immunogold labelling of viruses

Formvar/carbon coated 300 mesh nickel grids were allowed to float for 30 min on drops of purified TuYV-WT or TuYV-ΔRTP at about 100 ng/μL in citrate buffer 0.1 M (pH 6.0). Grids were then incubated in blocking buffer (PBS + 2% BSA) for 30 min and transferred to drops containing purified immunoglobulins at 0.8 mg/mL directed against the TuYV RTD, dilution 1:40 in PBS (Reutenauer et al., 1993). Incubation was conducted overnight at 4 °C. Grids were rinsed in blocking buffer (30 min) and incubated in goat anti-rabbit IgG conjugated with 10 nm gold beads for 3 h 30 min, 1:20 in PBS 0.1% BSA (BBInternational, Cardiff, UK). Grids were rinsed with PBS then with water and finally stained with 3% uranyl acetate. Grids were observed with a Philips EM 208 transmission electron microscope operating at 80 kV.

SDS-PAGE of purified virus

Purified TuYV-WT from *M. perfoliata* (2 μg denatured in Laemmli buffer) was loaded on a 10% SDS-PAGE which was further stained with colloidal Coomassie blue. After staining, the bands were quantified using ImageJ (see Web references). The net electrostatic charge of each structural protein was calculated using Prot Pi (see Web references).

Negative stain electron microscopy

Negative stain EM grids were prepared by applying 3 μL of virus solution previously diluted at 0.07 mg/ml to glow-discharged collodion-carbon-coated copper grids for 2 min. The excess of liquid was blotted with Whatman N4 filter paper and the grids were washed twice in 1% w/v uranyl acetate for 1 min. The excess liquid was removed, and grids were air-dried. The grids were observed using a JEOL 2200 FS electron microscope operating at 200 kV in zero-energy-loss mode with a slit width of 20 eV and equipped with a 4k × 4k slow-scan CDD camera (Gatan inc.)

Cryo-EM

Preliminary images were recorded using a JEOL 2200 FS electron microscope operating at 200 kV in zero-energy-loss mode with a slit width of 20 eV and equipped with a 4k × 4k slow-scan CDD camera (Gatan inc.). We initially prepared frozen-hydrated viruses deposited onto Quantifoil R2/2 or R1.2/1.3 grids (Quantifoil Micro tools GmbH, Germany) or glow discharged Lacey grid (Ted Pella Inc., USA), blotted for 3 s and then flash frozen in liquid ethane using the semi-automated plunge freezing device Vitrobot IV at 100% relative humidity. Observations revealed that even at high nominal concentrations, up to 3.5 mg/mL, the final concentration of viruses within the grid holes was not enough for an automatic image acquisition. Indeed, most of the virus particles interacted with the carbon film and TuYV-ΔRTP also formed nanocrystals. To overcome this problem, we used Quantifoil R2/2 grids covered with a 10-15 nm carbon film, which led to virus concentrations suitable for a single-particle cryo-EM analysis. No deformation of viral particles was observed, indicating that they were apparently not impaired by the interaction with the carbon film.

For high-resolution data collection, 3 μL of purified viral particles at ~1 mg/mL were applied during 2 min to glow-discharged Quantifoil R 2/2 grids covered with a thin carbon film (Quantifoil Micro tools GmbH, Germany), blotted for 3 s and then flash frozen in liquid ethane using the semi-automated plunge freezing device Vitrobot IV at 100% relative humidity. A Polara transmission electron microscope (Thermo Fisher Scientific, Eindhoven, Netherlands) (IBS, Grenoble, France) was used operating at 300 kV with defocus ranging from –0.8 to –2.5 μm. The pixel size estimated at the specimen level was 0.97 Å. A total of 151 and 179 movies were recorded for TuYV-WT and TuYV-ΔRTP respectively, using the Gatan Latitude single particle acquisition software (Gatan inc.) on a K2 camera in counting mode. The total exposure time was 5 s corresponding to a dose of 50 $e^-/Å^2$. Forty individual frames were collected with an electron dose of 1.0 $e^-/Å^2$ per frame.

Image processing and 3D reconstructions

Image processing was carried out using RELION 3.1 (Scheres, 2012; Zivanov et al., 2018) for icosahedral reconstructions, RELION 4.0 (Kimanius et al., 2021) for focussed asymmetric classifications, Gctf (Zhang, 2016) for contrast transfer function (CTF) calculations, Gautomatch (developed by Kai Zhang at the MRC Laboratory of Molecular Biology, UK) for automatic particle picking and RELION 3.1 for local resolution estimation. EMAN2 (Tang et al., 2007) was used to generate radial density plots.

The frames of each movie were computationally corrected for drift and beam-induced movement (Zivanov et al., 2018) and the CTF calculated. 51 930 TuYV-WT and 26 519 TuYV-ΔRTP particles were automatically picked, extracted from the movies and rescaled (pixel size: 2.91 Å, box size: 160 pixels) and submitted to reference-free 2D classification with regularisation parameter τ=2 and mask diameter=377 Å. 40 780 TuYV-WT and 18 433 TuYV-ΔRTP particles were selected from the best 2D classes and used to compute *de novo* 3D models at 15 Å resolution. Icosahedral symmetry was imposed during the model computation. Further particles were discarded based on the results of 3D classifications (pixel size: 1.22526 Å, box size: 380 pixels, τ: 4, mask diameter: 375 Å, number of classes: 4) and 3D alignment refinements with icosahedral symmetry constraints. The sets of particles producing the best reconstructions were used for per-particle defocus refinement and Bayesian polishing. Ultimately, 17 316 TuYV-WT and 3 003 TuYV-ΔRTP selected particles yielded icosahedral reconstructions at 4.1 Å and 3.5 Å resolution respectively. The resolution was estimated based on a 0.143 gold-standard FSC threshold on masked maps (Chen et al., 2013; Scheres and Chen, 2012). The refined maps were post-processed through model masking and *B*-factor sharpening, with $B = -137$ Å^2 and –60.5 Å^2 for TuYV-WT and TuYV-ΔRTP respectively.

To reconstruct the ^{N}RTD from the TuYV-WT images, several asymmetric, focussed 3D classifications were run. Regions surrounding an I_2, a Q_2 and a Q_3 axis of the CP-shell were independently targeted (see Results for the definition of these symbols). For each targeted axis, three spherical masks were used possessing a diameter of 80 Å, 120 Å and 180 Å, corresponding respectively to 1, 1.5 and 2 times the size of the ^{N}RTD (estimated based on PDB 7uln). These masks were positioned either assuming the ^{N}RTD in contact with the capsid surface or at a distance, estimated by assuming extended CP-^{N}RTD linkers (21 residues) in polyproline helix conformation parallel to the CP symmetry axis. An

elongated mask, consisting of approximately one half of an ellipsoid (half-axis lengths 50 Å, 50 Å and 180 Å), was also used. Finally, an approximately hemispherical mask (radius 180 Å) covering a large region around the targeted axis was used to account for the possibility of ^{N}RTD locations far from the symmetry axis (bending of the linkers). A 19.6 Å (16 pixels) soft border was added to all masks. The set of 17 316 aligned images of the TuYV-WT particles was symmetry-expanded using either all 60 icosahedral-symmetry operators (for 3D classifications focussed around the Q_2 and Q_3 axes) or a 30 operator subset (for 3D classifications focussed around the I_2 axis). Projections of the TuYV-WT reconstructed capsid were subtracted from the images of the expanded set, except for a small region of the protein shell corresponding approximately to the volume of four CP monomers surrounding the targeted I_2 or Q_2 axis, or three CP monomers surrounding the targeted Q_3 axis. The subtracted images were recentred at the centre of the spherical masks or at the base of the hemispherical or hemi-ellipsoidal masks and re-boxed (box size: 216 pixels for spherical masks, 312 pixels otherwise; pixel size: 1.22526 Å). With each mask, extensive 3D focussed (fixed particles), asymmetric classifications were carried out (parameters: τ=4,10,20,40,60; number of classes=4; mask diameter=300-360 Å). No preestablished atomic model of the ^{N}RTD was used as a reference for classification. All attempts to identify the ^{N}RTD in the 3D classes were unsuccessful.

Atomic model building, refinement and analysis

The TuYV-ΔRTP capsid structure was built and refined starting from an AlphaFold-3 (Abramson et al., 2024) prediction of the CP domain (Fig. S1). To save computation time and memory and, at the same time, correctly avoid steric clashes between symmetry-related neighbouring chains, only a part of the whole T=3 icosahedral capsid density was considered, which consisted of eleven subunits: the three chains of an asymmetric unit plus the eight closest neighbouring chains. Eleven copies of the AlphaFold-3 CP model (from which the N-terminal Arg-rich stretch was excluded) were fitted and iteratively refined against the experimental density map – following the general guidelines described by Afonine et al. (2018a) – by alternated steps of automatic refinement under stereochemical restraints with PHENIX.real_space_refine (Afonine et al., 2018b), visual model inspection and correction with *Coot* and model geometry validation with MolProbity (Williams et al., 2018). Non-crystallographic symmetry (NCS) restraints were applied during the first refinement run and subsequently released. Standard stereochemical restraints and interactively-checked secondary-structure restraints were applied during all refinement steps. Atomic isotropic displacement parameters (one parameter per residue) were applied and refined in the last refinement run. Finally, the refined model was pruned back to one asymmetric unit (three chains) before deposition with the PDB. The TuYV-WT atomic structure was similarly refined using the TuYV-ΔRTP structure as a starting point and no NCS restraint. Subunit-subunit interfaces were analysed using PISA (Krissinel and Henrick, 2007). The directions and positions of the local symmetry axes in the final structures were calculated by least-squares optimisation, under matrix-orthogonality constraints, of the group of matrices associated to an exact n-fold axis with respect to the transformations obtained by pair-wise alignment of the relevant subunits. The geometrical parameters describing the ideal virus architecture (see Results) were calculated based on Fig. S2, basic trigonometry and the aid of formulas for regular polyhedra (icosahedron, dodecahedron and pentagonal

pyramid) available from https://mathworld.wolfram.com (see Web references). Results of cumbersome trigonometric expressions were checked with the aid of Wolfram|Alpha (see Web references). The expected volumes of the unmodelled portions of the structure at the capsid interior (CP N-ter stretches and genomic RNA) were estimated using NucProt (Voss and Gerstein, 2005).

Artwork

Figures were prepared using: UCSF Chimera (Pettersen et al., 2004), developed by the Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco, with support from NIH P41-GM103311; UCSF ChimeraX (Meng et al., 2023), developed by the Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco, with support from National Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases; PyMOL open source, version 2.5.0 (Schrödinger, LLC, 2024).

Results and Discussion

Characterisation of purified viral particles by negative staining, immunogold labelling and SDS-PAGE

TuYV virions were purified by sucrose density centrifugation from agroinfiltrated leaves of plants inoculated with the wild-type (TuYV-WT) or a mutant clone (TuYV-ΔRTP) devoid of RTP* (Fig. 1). The mutant (called BWYV-6.4 in Brault et al. (1995)) is affected in plant systemic infection and is not transmissible by aphids (Brault et al., 1995). TuYV-ΔRTP virions are solely composed of CP subunits whereas the wild-type virions contain both CP and RTP* (Brault et al., 1995) (Fig. 1).

Observation of negatively stained TuYV-WT virions by transmission EM revealed circular particles with a diameter of approximately 30 nm (Fig. 2a). In the TuYV-ΔRTP samples two particle shapes are reproducibly observed: circular, sized as the wild-type virions (Fig. 2b), and slightly elongated (Fig. 2c). This latter shape has never been observed in preparations of purified TuYV-WT. TuYV-ΔRTP particles were rather smooth (Fig. 2b), while TuYV-WT virions displayed protruding densities at their surface (4 to 6 protrusions on each particle), indicated by black arrows in Fig. 2a, likely the [N]RTD from the RTP*.

Immunogold labelling experiments using antibodies directed against the RTD (Reutenauer et al., 1993) confirm exposition of the [N]RTD at the surface of the TuYV-WT virions (Fig. 2d-g) as previously shown on virions partially purified from TuYV-infected plant protoplasts (Brault et al., 1995). No gold labelling was observed on TuYV-ΔRTP virions (Figure 2f). We estimated that the number of gold beads per viral particles in the EM images (Table S1) varies from 0 to 4. This is consistent with the presence of the protrusions observed in negatively stained samples.

The number of incorporated RTP*s per TuYV-WT particle, estimated by electrophoretic analysis, was higher than the average number of beads observed by immunogold labelling and capsid protrusions observed by negative staining. Based on colloidal-Coomassie-blue SDS-PAGE band intensity (Fig. S3), the CP/ RTP* ratio was

estimated to be 13.5, corresponding to 12.4 incorporated RPT*s per virion. Note, however, that estimations from stained gels are approximate (Tal et al., 1985).

Schiltz et al. (2022) showed that, *in vitro*, the [N]RTD forms dimers. The discrepancy we observed between the number of incorporated RTP*s estimated by negative staining, immunogold labelling and electrophoretic analysis on denatured proteins is in line with the hypothesis that the observed capsid protrusions contain [N]RTD dimers. The varying number of beads per virion observed after immunogold labelling suggests a heterogeneous population of viruses incorporating a different number of RTP* subunits. One cannot exclude, though, the lack of accessibility of some [N]RTDs, or the presence of different RTP* conformations, some of which would not be recognised by the antibodies.

We found that TuYV-ΔRTP can form large 2D-arrays of particles in negatively stained samples (Fig. 3a) and also small 3D crystals in frozen hydrated samples at a concentration of 3.5 mg/mL (Fig. 3b) as already seen in aphids fed on purified TuYV-ΔRTP (Reinbold et al., 2001). The regularity of these assemblies seems to be interrupted by the incorporation of elongated particles, indicated by arrows in Fig. 3a. This extensive regular organisation has never been observed for the wild-type virions neither by negative stain nor by cryo-EM. Only some assemblies of 2 or 4 virions were observed in negatively stained samples (Fig. 3c). This different behaviour suggests that the presence of RTP* in TuYV-WT strongly impairs the interplay between viral particles.

Cryo-EM 3D reconstruction of TuYV-WT and TuYV-ΔRTP

Frozen-hydrated samples of TuYV-WT and TuYV-ΔRTP were prepared on Quantifoil R2/2 holey-carbon grids covered with a carbon film to avoid preferential aggregation at the hole borders and formation of nanocrystals (Fig. 3b). Transmission EM images of the samples were collected using a Polara microscope equipped with a K2 camera. In most of the images, uniform distributions of mostly undistorted, RNA-filled spherical virions were observed (Fig. 4a). The 3D scattering densities of the TuYV-WT and TuYV-ΔRTP virions were reconstructed from selected sets of, respectively, 17 316 and 3 003 homogeneous, RNA-filled particles (Fig. 4b-c). Statistics for the cryo-EM single-particle analysis are summarised in Table S2. The overall resolution of the reconstructed maps, refined by imposing icosahedral symmetry constraints, was respectively estimated to be 4.1 Å and 3.5 Å for TuYV-WT and TuYV-ΔRTP (0.143 gold-standard FSC criterion). Both the protein shell, which was well resolved in the sharpened maps (Fig. 4b) and the RNA shell obscured by icosahedral averaging, are visible in the unsharpened, unmasked maps (Fig. 4c). Local resolution in the protein shell varies in the range 3.95–4.5 Å and 3.3–4.5 Å for TuYV-WT and TuYV-ΔRTP respectively (Fig. S4). Atomic models of the capsids, comprising more than two thirds of the length of the CP domains (residues 62/64 to 202), have been built and refined against the reconstructed densities (Fig. 4d-h, Fig. S5, Table S3). No interpretable density could be attributed to the long arginine-rich N-terminal stretches (Fig. 4h), which are predicted to be disordered and likely buried in the nucleic acid at the interior of the virions. Expectedly, due to symmetry averaging during the reconstruction procedure, no extra density indicative of the presence of the [N]RTD shows up in the TuYV-WT reconstruction (asymmetric reconstructions are discussed later). The atomic models of the CP assemblies do not reveal any significant difference between the TuYV-WT and TuYV-ΔRTP protein shells (Fig. 4e). Direct comparison of the trimers in

the icosahedral asymmetric unit, without rigid-body adjustments of their positions, shows a low root mean square deviation (RMSD) between main-chain atoms (0.39 Å). Rigid body fit of individual subunit pairs results in slight improvements of the RMSD (0.32-0.34 Å), but with negligible movements of the subunits (0.2-0.6° rotations).

The CP architeture

If not otherwise stated, the terms "chain" and "subunit" in this subsection will refer to the visible (modelled) part of the CP domains in the TuYV capsid structures (unframed sequences in Fig. 4h). Though we report calculated values (e.g. RMSD, rotation angles) for the TuYV-ΔRTP structure only, all the considerations that follow are valid for both TuYV-WT and TuYV-ΔRTP.

The TuYV capsid is composed by 180 copies of the CP domain arranged according to icosahedral rotational symmetry with triangulation number T=3 (Caspar and Klug, 1962). One icosahedral asymmetric unit comprises three quasi-equivalent subunits (chains A, B and C in Fig. 5a) related by an almost exact, local threefold axis, denoted Q_3 (Fig. 5a-b). Each subunit folds into a classical jelly roll (Fig. 4g), commonly found in viruses, composed of two packed four-stranded anti-parallel β-sheets. In TuYV, one of the β-sheets faces the interior of the capsid while the other faces the exterior. Two intervening helices, most of the loops, the N- and C-ter β-strands and a few residues from other strands of the jelly roll provide interactions between subunits (Fig. 4g).

The geometry of the overall assembly can be described in terms of an equilateral elevated dodecahedron (EED, Fig. 5c). This is a non-convex polyhedron composed of sixty equilateral triangular faces (Janner, 2006a), where each face represents, with respect to the icosahedral symmetry of the assembly, one asymmetric unit. The sixty faces of the EED, grouped by five around the twelve fivefold axes, can be thought of as the faces of pentagonal pyramids whose bases sit on the faces of an underlying regular dodecahedron (outlined by blue edges in Fig. 5c). The dihedral angles between triangular faces sharing a common edge are convex towards either the capsid interior or the capsid exterior, so that two dihedral angle values, named respectively *endo* and *exo*, and two classes of edges (named accordingly) can be defined (Fig. 5c). The actual TuYV geometry is quite close to that of an ideal EED. The values of the *endo* and *exo* angles (141° and 174° respectively), evaluated by comparing the directions of the TuYV Q_3 axes, are close to the ideal values (138.19° and 168.68° respectively) (Fig. 5f-g). The virtual hinge movement that would transform an *exo* into an *endo* interface corresponds to a rotation of ~45° in TuYV as compared to 53.13° in an ideal EED.

Besides the global icosahedral symmetry, the EED arrangement implies a set of local symmetry axes. Fig. 5d (see also Fig. S2) illustrates how the EED global and local symmetry can be derived by virtually folding an ideal 2D hexagonal lattice according to the *endo* and *exo* angles. We will use the symbols I_n ($n = 2, 3, 5$) and Q_n ($n = 2, 3, 5, 6$) to denote, respectively, global (icosahedral) and local (quasi) n-fold symmetry. The approximate positions of the I_n and Q_n axes are indicated in Fig. 5e. The TuYV capsid can be formally described as an assembly of pentamers and hexamers centred on the I_5 and I_3 icosahedral axes respectively. Fig. 5e shows a set of two hexamers, chains $BC_2B_3C_4B_5C_6$ and $CB_2C_3B_4C_5B_6$, and one pentamer, chains $AA_2A_3A_4A_5$. This portion of the capsid encompasses one asymmetric unit plus all its adjacent subunits. Pentamers obey strict

fivefold symmetry. Hexamers obey threefold but not sixfold symmetry and are composed of two sets of interleaved subunits, blue and green subunits respectively in Fig. 5h, not related by icosahedral symmetry. Large deviations from ideal sixfold symmetry are observed around the I_3 axes. Superposition of adjacent chains within a hexamer requires either a quasi-sixfold (Q_6) rotation of 60.2° or a quasi-fivefold (Q_5) rotation of 67.3° around axes that are not collinear with the I_3 axis. The Q_6-I_3 and Q_5-I_3 axes misalignments are 23.0° and 20.7° respectively (Fig. 5i). Note that Q_5 rotations give rise to an arrangement of adjacent subunits (ex. subunits CB_2 in Fig. 5e,h,i) similar to that observed within pentamers (ex. subunits AA_2 in Fig. 5e), hence the preferred symbol Q_5 instead of Q_6. Also, non-equivalent triplets of subunits can be found (e.g. ABA_2 and CAB_2) which are arranged essentially in the same way and whose local symmetry axes would be compatible with those of a T=1 icosahedron (Fig. 5j, Fig. 6). This is a consequence of pentamers and hexamers sharing *endo* angles (Fig. 5e), a structural constraint described as "*endo* angle propagation" by Mannige & Brooks (2010, 2009).

A total of eight independent —*i.e.* not related by icosahedral symmetry— interfaces between subunits exist. We have labelled the interfaces according to the rotation I_n or Q_n relating the contacting subunits (Table 1). The portion of capsid illustrated in Fig. 5e contains at least one representative of the eight interfaces. Interacting residues at the subunit interfaces are mainly of a polar or charged nature (Table 2, Fig. 6). No major differences are observed among the three Q_3 interfaces, though these are not strictly equivalent in principle, nor between the Q_5 and I_5 interfaces. (Table 2, Fig. 6). On the other hand, significant differences are found at the I_2, when compared to Q_2, and Q_6, when compared to Q_5 or I_5 interfaces. These differences are related to the existence, in an EED, of different dihedral angles: the Q_2, Q_5 and I_5 interfaces are found on *endo* edges, while the I_2 and Q_6 interfaces are found on *exo* edges. Indeed, most residues involved in Q_5, I_5, Q_6, Q_2 and I_2 interfaces line the edges of the EED (Fig. 6). The virtual hinge movement transforming an *endo* interface into an *exo* interface (Fig. 5g) would occur around a line of atomic contacts between helices h1 of neighbouring subunits. This transformation maintains the two helices in close contact but disrupts interactions and separate the underlying N-ter and C-ter β-sheets in adjacent chains (Fig. 6).

The structure of the TuYV capsid shows no significant differences with the previously published structures of polerovirus VLPs (Byrne et al., 2019) (RMSD ~1.0 Å in comparison with PLRV and BYDV VLPs). More generally, the T=3 subunit arrangement of TuYV follows the schematic initially described by Rossmann et al. (1983a, 1983b) for southern bean mosaic virus (SBMV, species *Sobemovirus SBMV*), in which the quasi-equivalence of contacts (Caspar and Klug, 1962) is assured essentially by different relative orientations of the subunits at the Q_2 and I_2 interfaces (dimeric interfaces) where the h1-helices touch each other. Differently to what happens in SBMV, the cryo-EM maps of TuYV indicate that the increased space available under the h1-helices at the I_2 interfaces is not occupied by an additionally ordered stretch of polypeptide chain.

We have described the TuYV assembly in terms of an equilateral elevated dodecahedron (EED), within the framework of Caspar and Klug's theory on the principles governing regular virus structures (Caspar and Klug, 1962) (see also Brinkmann et al. (2017), Janner (2006b), Keef et al. (2013), Konevtsova et al. (2024), Lorman & Rochal (2008), Mannige & Brooks (2008, 2009, 2010), Rochal et al. (2017), Twarock (2006) and

Twarock & Luque (2019) for different or more general approaches). The EED geometrical model can be applied, to different degrees of approximation, to many but not all viruses with T=3 icosahedral symmetry. TuYV shows small deviations from this ideal geometry (Fig. 5g,i). We have analysed all virus structures classified as T=3 in the VIPERdb (Montiel-Garcia et al., 2021) and found that, among those that can be reconducted to the EED geometry, a very limited number strictly adhere to that model, for example panicum mosaic virus (species *Panicovirus panici*), while most of the others, like tomato bushy stunt virus (TBSV, species *Tombusvirus lycopersici*) and SBMV, deviate to some extent (imperfect Q_3 rotations within the asymmetric unit). Other geometrical descriptions have been used in the past for some of these viruses. TBSV and SBMV have been described as rhombic triacontahedra (Johnson and Olson, 2021; Winkler et al., 1977), polyhedra composed by non-equilateral triangles forming, in pairs, flat rhombic faces (*exo* angle = 180°). The polerovirus PLRV has also been described as possessing flat *exo* angles (Adams et al., 2022), but by comparing it with TuYV we propose that PLRV also has non-flat *exo* interfaces. Other T=3 icosahedral viruses, like brome mosaic virus (species *Bromovirus BMV*) cannot be described by the EED model. Interestingly, none of the known T=3 structures adheres to the geometry of a regular icosahedron (Fig. S2), *i.e.* that obtained by folding a 2D hexagonal lattice along *endo* edges *only* as described by Caspar and Klug (1962) in their seminal paper. We could also point out that, while the EED geometry was not explicitly cited by Caspar and Klug, they indeed included a papercraft picture of an EED, and not an icosahedron, in Fig. 8c of their paper to illustrate the T=3 architecture.

The RNA structure

The genomic RNA shows up in the unmasked, unsharpened maps as an amorphous layer of averaged density just underneath the CP shell (Fig. 7). The approximate volume of the residual density layer appears to be adequate to accommodate the whole genomic RNA plus the unmodelled N-ter stretches of the CP subunits (Fig. S6). Highest values of the residual density are found around the I_5 and Q_3 axes and underneath the BC and AC interfaces (Fig. 7b-e). CP residues in proximity of the residual density are highlighted in Fig. 7f-h.

The unreconstructed ^{N}RTD is likely to float at the capsid exterior

Inspection of some selected cryo-EM images confirmed the presence of the ^{N}RTD (four to six protrusions of the capsid) in the purified TuYV-WT samples (Fig. 8a), but its scattering density was not visible in the cryo-EM map of the capsid due to the symmetry averaging. Also, we did not see differences between the radially averaged densities of the TuYV-WT and TuYV-ΔRTP particles that could be related to the ^{N}RTD (Fig. 7a). We tried to reconstruct the ^{N}RTD domain by applying asymmetric classification and reconstruction procedures. We chose to focus the 3D classifications on regions around the I_2, Q_2 and Q_3 axes (Fig. 8b) based on the observation that, *in vitro*, the ^{N}RTD forms symmetric dimers (Schiltz et al., 2022), and that the homologous dimeric domains in tomato bushy stunt virus (TBSV, PDB 2tbv) lie on the capsid surface in strict contact with it and aligned with the capsid I_2 or Q_2 axes. Also, in almost all other known T=3 virus structures possessing protruding domains, those domains are in a dimeric arrangement around the capsid symmetry axes and in contact with the capsid surface. This has been observed in some

nodavirids (PDB 6ab6) and calicivirids (PDB 3m8l, 2gh8, 7bjp; 7dod; 9cvg; 6otf; 3j1p), except for Orsay virus (PDB 4nwv), where the protruding domains are better described as trimers. We also targeted regions further away from the capsid surface to account for the length of the CP-^{N}RTD linker (Fig. 8b). Despite our extensive attempts (see Materials and Methods), we could not obtain any convincing result.

The ^{N}RTD dimers described by Schiltz et al. (2022) are structurally related to the dimeric P domains that protrude along the I_2 and Q_2 axes of the capsids of TBSV and other tombusviruses. Based on the TuYV-ΔRTP structure described in this paper (released by the PDB in July 2022), and by comparison with TBSV, Schiltz et al. (2022) showed that, in principle, it is structurally possible for the ^{N}RTD dimers to be accommodated (like in TBSV) in tight contact with the TuYV CP layer and even occupy the complete set of 90 twofold sites on the capsid surface. Failure of our extensive attempts to reconstruct the ^{N}RTD around the twofold sites suggests, however, that the ^{N}RTD domain does not occupy (in the conditions of our purified samples) a well-defined position on the capsid surface. Indeed, the proline-rich linker (21 residues) connecting the CP and ^{N}RTD domain in TuYV is much longer than the stretch (7 residues) joining the TBSV P domain to the capsid shell, and this may allow the ^{N}RTD to float around, at least in absence of a specific interacting partner, in proximity of the capsid surface.

We have seen no significant differences between the cryo-EM maps of TuYV-WT and TuYV-ΔRTP, indicating that the presence of the ^{N}RTD domains did not influence the final CP fold and assembly organisation (of the selected particles). One cannot exclude, however, the possibility of ^{N}RTD playing a role in RNA packaging and capsid assembly (formation of elongated particles has been observed in TuYV-ΔRTP but not in TuYV-WT samples).

Conclusions

We presented here the structural characterisation by immunogold labelling and 3D cryo-EM of the wild-type TuYV and a mutant whose capsid contains the CP only. We showed that incorporation of the RTP* does not impair the capsid structure, and the ^{N}RTD does not adopt well-defined positions at the capsid surface. The varying number of beads per virion observed after immunogold labelling suggests a heterogeneous population of viruses incorporating a different number of RTP* subunits, though one cannot exclude the lack of accessibility of some ^{N}RTDs, or the presence of different RTP* conformations, some of which would not be recognised by the antibodies. This raises the question whether different forms exist of the ^{N}RTD at the surface of the capsid that may have different cellular partners and fulfil different functions in the virus cycle. Answers to these questions will require further investigation.

Deposited data references

The cryo-EM maps of TuYV-WT and TuYV-ΔRTP have been deposited with the Electron Microscopy Data Bank and released under the accession IDs EMD-10003 and EMD-10001 respectively. The model atomic coordinates have been deposited with the Protein Data Bank and released under the PDB IDs 9fhp and 9q8j respectively.

Funding sources

This work was funded by the French Infrastructure for Integrated Structural Biology (FRISBI), grant ANR-10-INSB-05. A CC-BY 4.0 public copyright license (https://creativecommons.org/licenses/by/4.0/) has been applied by the authors to the submitted manuscript and to all subsequent versions of the document up to the Author Accepted Manuscript arising from the submission, in accordance with the grant's open access conditions.

Credits

Joséphine Lai-Kee-Him: Investigation, Resources. **Stefano Trapani**: Conceptualisation, Data Curation, Formal analysis, Investigation, Software, Validation, Visualisation, Writing - Original Draft, Writing - Review & Editing. **Sylvaine Boissinot**: Investigation, Resources, Writing - Review & Editing. **Catherine Reinbold**: Investigation, Resources, Visualisation, Writing - Review & Editing. **Chloé Fallet**: Investigation. **Aurélie Ancelin**: Investigation. **François Lecorre**: Investigation. **François Hoh**: Investigation. **Véronique Ziegler-Graff**: Conceptualisation, Writing - Review & Editing. **Véronique Brault**: Conceptualisation, Investigation, Project administration, Writing - Original Draft, Writing - Review & Editing. **Patrick Bron**: Conceptualisation, Funding acquisition, Investigation, Project administration, Supervision, Writing - Original Draft, Writing - Review & Editing.

Supplementary material

Table S1. Estimation of the number of immunogold-labelled TuYV particles.

Table S2. Cryo-EM data collection and single-particle processing statistics.

Table S3. Atomic model refinement statistics.

Fig. S1. AlphaFold-3 model of the TuYV RTP* monomer.

Fig. S2. Folding of a hexagonal lattice into either an icosahedron or an elevated dodecahedron.

Fig. S3. SDS-PAGE analysis and colloidal Coomassie blue staining of 2 µg of TuYV-WT viral particles purified from *M. perfoliata* and denaturated in Laemmli buffer.

Fig. S4. Local resolution of the cryo-EM reconstructions (sharpened maps).

Fig. S5. Some details of the TuYV-ΔRTP cryo-EM map and the corresponding refined atomic model.

Fig. S6. Estimation of the residual density volumes at the capsid interior.

References

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A.J., Bambrick, J., Bodenstein, S.W., Evans, D.A., Hung, C.-C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A., Arvaniti, E., Beattie, C., Bertolli, O., Bridgland, A., Cherepanov, A., Congreve, M., Cowen-Rivers, A.I., Cowie, A., Figurnov, M., Fuchs, F.B., Gladman, H., Jain, R., Khan, Y.A., Low, C.M.R., Perlin, K., Potapenko, A., Savy, P., Singh, S., Stecula, A., Thillaisundaram, A., Tong, C., Yakneen, S., Zhong, E.D., Zielinski, M., Žídek, A., Bapst, V., Kohli, P., Jaderberg, M., Hassabis, D., Jumper, J.M., 2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3. Nature 630, 493–500. https://doi.org/10.1038/s41586-024-07487-w

Adams, M.C., Schiltz, C.J., Heck, M.L., Chappie, J.S., 2022. Crystal structure of the potato leafroll virus coat protein and implications for viral assembly. Journal of Structural Biology 214, 107811. https://doi.org/10.1016/j.jsb.2021.107811

Afonine, P.V., Klaholz, B.P., Moriarty, N.W., Poon, B.K., Sobolev, O.V., Terwilliger, T.C., Adams, P.D., Urzhumtsev, A., 2018a. New tools for the analysis and validation of cryo-EM maps and atomic models. Acta Cryst D 74, 814–840. https://doi.org/10.1107/S2059798318009324

Afonine, P.V., Poon, B.K., Read, R.J., Sobolev, O.V., Terwilliger, T.C., Urzhumtsev, A., Adams, P.D., 2018b. Real-space refinement in PHENIX for cryo-EM and crystallography. Acta Cryst D 74, 531–544. https://doi.org/10.1107/S2059798318006551

Alexander, M.M., Mohr, J.P., DeBlasio, S.L., Chavez, J.D., Ziegler-Graff, V., Brault, V., Bruce, J.E., Heck, M. (Cilia), 2017. Insights in luteovirid structural biology guided by chemical cross-linking and high resolution mass spectrometry. Virus Research, Plant Virus Epidemiology 241, 42–52. https://doi.org/10.1016/j.virusres.2017.05.005

Bencharki, B., Boissinot, S., Revollon, S., Ziegler-Graff, V., Erdinger, M., Wiss, L., Dinant, S., Renard, D., Beuve, M., Lemaitre-Guillier, C., Brault, V., 2010. Phloem Protein Partners of Cucurbit aphid borne yellows virus: Possible Involvement of Phloem Proteins in Virus Transmission by Aphids. MPMI 23, 799–810. https://doi.org/10.1094/MPMI-23-6-0799

Boissinot, S., Brault, V., Herrbach, E., 2020. Luteovirids, in: Encyclopedia of Life Sciences. Wiley, pp. 327–339. https://doi.org/10.1002/9780470015902.a0029206

Boissinot, S., Erdinger, M., Monsion, B., Ziegler-Graff, V., Brault, V., 2014. Both Structural and Non-Structural Forms of the Readthrough Protein of Cucurbit aphid-borne yellows virus Are Essential for Efficient Systemic Infection of Plants. PLOS ONE 9, e93448. https://doi.org/10.1371/journal.pone.0093448

Brault, V., Bergdoll, M., Mutterer, J., Prasad, V., Pfeffer, S., Erdinger, M., Richards, K.E., Ziegler-Graff, V., 2003. Effects of Point Mutations in the Major Capsid Protein of Beet Western Yellows Virus on Capsid Formation, Virus Accumulation, and Aphid Transmission. J Virol 77, 3247–3256. https://doi.org/10.1128/JVI.77.5.3247-3256.2003

Brault, V., van den Heuvel, J.F., Verbeek, M., Ziegler-Graff, V., Reutenauer, A., Herrbach, E., Garaud, J.C., Guilley, H., Richards, K., Jonard, G., 1995. Aphid transmission of beet western yellows luteovirus requires the minor capsid read-through protein P74. The EMBO Journal 14, 650–659. https://doi.org/10.1002/j.1460-2075.1995.tb07043.x

Brinkmann, G., Goetschalckx, P., Schein, S., 2017. Comparing the constructions of Goldberg, Fuller, Caspar, Klug and Coxeter, and a general approach to local symmetry-preserving operations. Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences 473, 20170267. https://doi.org/10.1098/rspa.2017.0267

Bruyère, A., Brault, V., Ziegler-Graff, V., Simonis, M.-T., Van den Heuvel, J.F.J.M., Richards, K., Guilley, H., Jonard, G., Herrbach, E., 1997. Effects of Mutations in the Beet Western Yellows Virus Readthrough Protein on Its Expression and Packaging and on Virus Accumulation, Symptoms, and Aphid Transmission. Virology 230, 323–334. https://doi.org/10.1006/viro.1997.8476

Byrne, M.J., Steele, J.F.C., Hesketh, E.L., Walden, M., Thompson, R.F., Lomonossoff, G.P., Ranson, N.A., 2019. Combining Transient Expression and Cryo-EM to Obtain High-Resolution Structures of Luteovirid Particles. Structure 27, 1761-1770.e3. https://doi.org/10.1016/j.str.2019.09.010

Caspar, D.L., Klug, A., 1962. Physical principles in the construction of regular viruses. Cold Spring Harb. Symp. Quant. Biol. 27, 1–24.

Chavez, J.D., Cilia, M., Weisbrod, C.R., Ju, H.-J., Eng, J.K., Gray, S.M., Bruce, J.E., 2012. Cross-linking Measurements of the Potato leafroll virus Reveal Protein Interaction Topologies Required for Virion Stability, Aphid Transmission, and Virus–Plant Interactions. J. Proteome Res. 11, 2968–2981. https://doi.org/10.1021/pr300041t

Chay, C.A., Gunasinge, U.B., Dinesh-kumar, S.P., Miller, W.A., Gray, S.M., 1996. Aphid Transmission and Systemic Plant Infection Determinants of Barley Yellow Dwarf Luteovirus-PAV are Contained in the Coat Protein Readthrough Domain and 17-kDa Protein, Respectively. Virology 219, 57–65. https://doi.org/10.1006/viro.1996.0222

Chen, S., McMullan, G., Faruqi, A.R., Murshudov, G.N., Short, J.M., Scheres, S.H.W., Henderson, R., 2013. High-resolution noise substitution to measure overfitting and validate resolution in 3D structure determination by single particle electron cryomicroscopy. Ultramicroscopy 135, 24–35. https://doi.org/10.1016/j.ultramic.2013.06.004

DeBlasio, S.L., Chavez, J.D., Alexander, M.M., Ramsey, J., Eng, J.K., Mahoney, J., Gray, S.M., Bruce, J.E., Cilia, M., 2016. Visualization of Host-Polerovirus Interaction Topologies Using Protein Interaction Reporter Technology. Journal of Virology 90, 1973–1987. https://doi.org/10.1128/jvi.01706-15

DeBlasio, S.L., Johnson, R., Mahoney, J., Karasev, A., Gray, S.M., MacCoss, M.J., Cilia, M., 2015a. Insights Into the Polerovirus–Plant Interactome Revealed by Coimmunoprecipitation and Mass Spectrometry. MPMI 28, 467–481. https://doi.org/10.1094/MPMI-11-14-0363-R

DeBlasio, S.L., Johnson, R., Sweeney, M.M., Karasev, A., Gray, S.M., MacCoss, M.J., Cilia, M., 2015b. Potato leafroll virus structural proteins manipulate overlapping, yet distinct protein interaction networks during infection. PROTEOMICS 15, 2098–2112. https://doi.org/10.1002/pmic.201400594

English, J.J., Davenport, G.F., Elmayan, T., Vaucheret, H., Baulcombe, D.C., 1997. Requirement of sense transcription for homology-dependent virus resistance and trans-inactivation. The Plant Journal 12, 597–603. https://doi.org/10.1046/j.1365-313X.1997.00597.x

Filichkin, S.A., Lister, R.M., McGrath, P.F., Young, M.J., 1994. *In Vivo* Expression and Mutational Analysis of the Barley Yellow Dwarf Virus Readthrough Gene. Virology 205, 290–299. https://doi.org/10.1006/viro.1994.1645

Gildow, F.E., 1999. Luteovirus transmission and mechanisms regulating vector specificity, in: The Luteoviridae. CABI publishing, Wallingford, pp. 88–113.

Heck, M., Brault, V., 2018. Targeted disruption of aphid transmission: a vision for the management of crop diseases caused by Luteoviridae members. Current Opinion in Virology 33, 24–32. https://doi.org/10.1016/j.coviro.2018.07.007

Hipper, C., Monsion, B., Bortolamiol-Bécet, D., Ziegler-Graff, V., Brault, V., 2014. Formation of virions is strictly required for turnip yellows virus long-distance movement in plants. Journal of General Virology 95, 496–505. https://doi.org/10.1099/vir.0.058867-0

Janner, A., 2006a. Crystallographic structural organization of human rhinovirus serotype 16, 14, 3, 2 and 1A. Acta Crystallographica: Section A (Wiley-Blackwell) 62, 270–286. https://doi.org/10.1107/S010876730601748X

Janner, A., 2006b. Towards a classification of icosahedral viruses in terms of indexed polyhedra. Acta Crystallogr A Found Crystallogr 62, 319–330. https://doi.org/10.1107/S0108767306022227

Johnson, J.E., Olson, A.J., 2021. Icosahedral virus structures and the protein data bank. Journal of Biological Chemistry 296. https://doi.org/10.1016/j.jbc.2021.100554

Kaplan, I.B., Lee, L., Ripoll, D.R., Palukaitis, P., Gildow, F., Gray, S.M., 2007. Point mutations in the potato leafroll virus major capsid protein alter virion stability and aphid transmission. Journal of General Virology 88, 1821–1830. https://doi.org/10.1099/vir.0.82837-0

Keef, T., Wardman, J.P., Ranson, N.A., Stockley, P.G., Twarock, R., 2013. Structural constraints on the three-dimensional geometry of simple viruses: case studies of a new predictive tool. Acta Cryst. A 69, 140–150. https://doi.org/10.1107/S0108767312047150

Kimanius, D., Dong, L., Sharov, G., Nakane, T., Scheres, S.H.W., 2021. New tools for automated cryo-EM single-particle analysis in RELION-4.0. Biochemical Journal 478, 4169–4185. https://doi.org/10.1042/BCJ20210708

Konevtsova, O.V., Chalin, D.V., Rochal, S.B., 2024. Theory of density waves and organization of proteins in icosahedral virus capsids. Phys. Chem. Chem. Phys. 26, 569–580. https://doi.org/10.1039/D3CP05384A

Krissinel, E., Henrick, K., 2007. Inference of Macromolecular Assemblies from Crystalline State. Journal of
Molecular Biology 372, 774–797. https://doi.org/10.1016/j.jmb.2007.05.022

LaTourrette, K., Holste, N.M., Garcia-Ruiz, H., 2021. Polerovirus genomic variation. Virus Evolution 7,
veab102. https://doi.org/10.1093/ve/veab102

Lee, L., Kaplan, I.B., Ripoll, D.R., Liang, D., Palukaitis, P., Gray, S.M., 2005. A Surface Loop of the Potato
Leafroll Virus Coat Protein Is Involved in Virion Assembly, Systemic Movement, and Aphid
Transmission. Journal of Virology 79, 1207–1214. https://doi.org/10.1128/jvi.79.2.1207-1214.2005

Leiser, R.M., Ziegler-Graff, V., Reutenauer, A., Herrbach, E., Lemaire, O., Guilley, H., Richards, K., Jonard,
G., 1992. Agroinfection as an alternative to insects for infecting plants with beet western yellows
luteovirus. Proceedings of the National Academy of Sciences 89, 9136–9140.
https://doi.org/10.1073/pnas.89.19.9136

Liang, D., Gray, S.M., Kaplan, I., Palukaitis, P., 2004. Site-Directed Mutagenesis and Generation of Chimeric
Viruses by Homologous Recombination in Yeast to Facilitate Analysis of Plant-Virus Interactions. MPMI
17, 571–576. https://doi.org/10.1094/MPMI.2004.17.6.571

Lorman, V.L., Rochal, S.B., 2008. Landau theory of crystallization and the capsid structures of small
icosahedral viruses. Phys. Rev. B 77, 224109. https://doi.org/10.1103/PhysRevB.77.224109

Mannige, R.V., Brooks, C.L., 2010. Periodic Table of Virus Capsids: Implications for Natural Selection and
Design. PLOS ONE 5, e9423. https://doi.org/10.1371/journal.pone.0009423

Mannige, R.V., Brooks, C.L., 2009. Geometric considerations in virus capsid size specificity, auxiliary
requirements, and buckling. Proceedings of the National Academy of Sciences 106, 8531–8536.
https://doi.org/10.1073/pnas.0811517106

Mannige, R.V., Brooks, C.L., 2008. Tilable nature of virus capsids and the role of topological constraints in
natural capsid design. Phys. Rev. E 77, 051902. https://doi.org/10.1103/PhysRevE.77.051902

Meng, E.C., Goddard, T.D., Pettersen, E.F., Couch, G.S., Pearson, Z.J., Morris, J.H., Ferrin, T.E., 2023.
UCSF CHIMERAX : Tools for structure building and analysis. Protein Science 32, e4792.
https://doi.org/10.1002/pro.4792

Montiel-Garcia, D., Santoyo-Rivera, N., Ho, P., Carrillo-Tripp, M., III, C.L.B., Johnson, J.E., Reddy, V.S.,
2021. VIPERdb v3.0: a structure-based data analytics platform for viral capsids. Nucleic Acids Research
49, D809–D816. https://doi.org/10.1093/nar/gkaa1096

Mulot, M., Monsion, B., Boissinot, S., Rastegar, M., Meyer, S., Bochet, N., Brault, V., 2018. Transmission of
Turnip yellows virus by Myzus persicae Is Reduced by Feeding Aphids on Double-Stranded RNA
Targeting the Ephrin Receptor Protein. Front. Microbiol. 9. https://doi.org/10.3389/fmicb.2018.00457

Mutterer, J.D., Stussi-Garaud, C., Michler, P., Richards, K.E., Jonard, G., Ziegler-Graff, V., 1999. Role of the
beet western yellows virus readthrough protein in virus movement in Nicotiana clevelandii. Journal of
General Virology 80, 2771–2778. https://doi.org/10.1099/0022-1317-80-10-2771

Peter, K.A., Gildow, F., Palukaitis, P., Gray, S.M., 2009. The C Terminus of the Polerovirus P5 Readthrough
Domain Limits Virus Infection to the Phloem. Journal of Virology 83, 5419–5429.
https://doi.org/10.1128/jvi.02312-08

Peter, K.A., Liang, D., Palukaitis, P., Gray, S.M., 2008. Small deletions in the potato leafroll virus
readthrough protein affect particle morphology, aphid transmission, virus movement and accumulation.
Journal of General Virology 89, 2037–2045. https://doi.org/10.1099/vir.0.83625-0

Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., Ferrin, T.E., 2004.
UCSF Chimera—A visualization system for exploratory research and analysis. Journal of Computational
Chemistry 25, 1605–1612. https://doi.org/10.1002/jcc.20084

Reinbold, C., Gildow, F.E., Herrbach, E., Ziegler-Graff, V., Gonçalves, M.C., van den Heuvel, J.F.J.M.,
Brault, V., 2001. Studies on the role of the minor capsid protein in transport of Beet western yellows virus
through Myzus persicae. Journal of General Virology 82, 1995–2007.
https://doi.org/10.1099/0022-1317-82-8-1995

Reutenauer, A., Ziegler-Graff, V., Lot, H., Scheidecker, D., Guilley, H., Richards, K., Jonard, G., 1993.
Identification of Beet Western Yellows Luteovirus Genes Implicated in Viral Replication and Particle
Morphogenesis. Virology 195, 692–699. https://doi.org/10.1006/viro.1993.1420

Rochal, S.B., Konevtsova, O.V., Lorman, V.L., 2017. Static and dynamic hidden symmetries of icosahedral viral capsids. Nanoscale 9, 12449–12460. https://doi.org/10.1039/C7NR04020B

Rodriguez-Medina, C., Boissinot, S., Chapuis, S., Gereige, D., Rastegar, M., Erdinger, M., Revers, F., Ziegler-Graff, V., Brault, V., 2015. A protein kinase binds the C-terminal domain of the readthrough protein of *Turnip yellows virus* and regulates virus accumulation. Virology 486, 44–53. https://doi.org/10.1016/j.virol.2015.08.031

Rossmann, M.G., Abad-Zapatero, C., Hermodson, M.A., Erickson, J.W., Huber, R., 1983a. Subunit interactions in southern bean mosaic virus. Journal of Molecular Biology 166, 37–73. https://doi.org/10.1016/S0022-2836(83)80049-7

Rossmann, M.G., Abad-Zapatero, C., Murthy, M.R.N., Liljas, L., Jones, T.A., Strandberg, B., 1983b. Structural comparisons of some small spherical plant viruses. Journal of Molecular Biology 165, 711–736. https://doi.org/10.1016/S0022-2836(83)80276-9

Scheres, S.H.W., 2012. RELION: Implementation of a Bayesian approach to cryo-EM structure determination. Journal of Structural Biology 180, 519–530. https://doi.org/10.1016/j.jsb.2012.09.006

Scheres, S.H.W., Chen, S., 2012. Prevention of overfitting in cryo-EM structure determination. Nature Methods 9, 853–854. https://doi.org/10.1038/nmeth.2115

Schiltz, C.J., Wilson, J.R., Hosford, C.J., Adams, M.C., Preising, S.E., DeBlasio, S.L., MacLeod, H.J., Van Eck, J., Heck, M.L., Chappie, J.S., 2022. Polerovirus N-terminal readthrough domain structures reveal molecular strategies for mitigating virus transmission by aphids. Nat Commun 13, 6368. https://doi.org/10.1038/s41467-022-33979-2

Schrödinger, LLC, 2024. The PyMOL Molecular Graphics System, Version 2.5.0.

Smirnova, E., Firth, A.E., Miller, W.A., Scheidecker, D., Brault, V., Reinbold, C., Rakotondrafara, A.M., Chung, B.Y.-W., Ziegler-Graff, V., 2015. Discovery of a Small Non-AUG-Initiated ORF in Poleroviruses and Luteoviruses That Is Required for Long-Distance Movement. PLOS Pathogens 11, e1004868. https://doi.org/10.1371/journal.ppat.1004868

Sõmera, M., Fargette, D., Hébrard, E., Sarmiento, C., ICTV Report Consortium, 2021. ICTV Virus Taxonomy Profile: Solemoviridae 2021. Journal of General Virology 102, 001707. https://doi.org/10.1099/jgv.0.001707

Tal, M., Silberstein, A., Nusser, E., 1985. Why does Coomassie Brilliant Blue R interact differently with different proteins? A partial answer. Journal of Biological Chemistry 260, 9976–9980. https://doi.org/10.1016/S0021-9258(17)39198-6

Tang, G., Peng, L., Baldwin, P.R., Mann, D.S., Jiang, W., Rees, I., Ludtke, S.J., 2007. EMAN2: An extensible image processing suite for electron microscopy. Journal of Structural Biology, Software tools for macromolecular microscopy 157, 38–46. https://doi.org/10.1016/j.jsb.2006.05.009

Terradot, L., Souchet, M., Tran, V., Giblot Ducray-Bourdin, D., 2001. Analysis of a Three-Dimensional Structure of *Potato leafroll virus* Coat Protein Obtained by Homology Modeling. Virology 286, 72–82. https://doi.org/10.1006/viro.2001.0900

Torrance, L., 1992. Analysis of epitopes on potato leafroll virus capsid protein. Virology 191, 485–489. https://doi.org/10.1016/0042-6822(92)90216-C

Twarock, R., 2006. Mathematical virology: a novel approach to the structure and assembly of viruses. Phil. Trans. R. Soc. A. 364, 3357–3373. https://doi.org/10.1098/rsta.2006.1900

Twarock, R., Luque, A., 2019. Structural puzzles in virology solved with an overarching icosahedral design principle. Nat Commun 10, 4414. https://doi.org/10.1038/s41467-019-12367-3

van den Heuvel, J.F.J.M., 1991. Transmission of Potato Leafroll Virus from Plants and Artificial Diets by *Myzus persicae*. Phytopathology 81, 150. https://doi.org/10.1094/Phyto-81-150

Veidt, I., Lot, H., Leiser, M., Scheidecker, D., Guilley, H., Richards, K., Jonard, G., 1988. Nucleotide sequence of beet western yellows virus RNA. Nucleic Acids Research 16, 9917–9932. https://doi.org/10.1093/nar/16.21.9917

Voss, N.R., Gerstein, M., 2005. Calculation of Standard Atomic Volumes for RNA and Comparison with Proteins: RNA is Packed More Tightly. Journal of Molecular Biology 346, 477–492. https://doi.org/10.1016/j.jmb.2004.11.072

Wang, J.Y., Chay, C., Gildow, F.E., Gray, S.M., 1995. Readthrough Protein Associated with Virions of Barley Yellow Dwarf Luteovirus and Its Potential Role in Regulating the Efficiency of Aphid Transmission. Virology 206, 954–962. https://doi.org/10.1006/viro.1995.1018

Williams, C.J., Headd, J.J., Moriarty, N.W., Prisant, M.G., Videau, L.L., Deis, L.N., Verma, V., Keedy, D.A., Hintze, B.J., Chen, V.B., Jain, S., Lewis, S.M., Arendall, W.B., Snoeyink, J., Adams, P.D., Lovell, S.C., Richardson, J.S., Richardson, D.C., 2018. MolProbity: More and better reference data for improved all-atom structure validation. Protein Science 27, 293–315. https://doi.org/10.1002/pro.3330

Winkler, F.K., Schutt, C.E., Harrison, S.C., Bricogne, G., 1977. Tomato bushy stunt virus at 5.5-Å resolution. Nature 265, 509–513. https://doi.org/10.1038/265509a0

Xu, Y., Silva, W.L.D., Qian, Y., Gray, S.M., 2018. An aromatic amino acid and associated helix in the C-terminus of the potato leafroll virus minor capsid protein regulate systemic infection and symptom expression. PLOS Pathogens 14, e1007451. https://doi.org/10.1371/journal.ppat.1007451

Zhang, K., 2016. Gctf: Real-time CTF determination and correction. Journal of Structural Biology 193, 1–12. https://doi.org/10.1016/j.jsb.2015.11.003

Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E., Scheres, S.H., 2018. New tools for automated high-resolution cryo-EM structure determination in RELION-3. eLife 7, e42166. https://doi.org/10.7554/eLife.42166

Web references

Prot Pi, https://www.protpi.ch/ (last access Jan 13, 2025).

Rasband, W.S., ImageJ, U. S. National Institutes of Health, Bethesda, Maryland, USA, https://imagej.net/ij/, 1997-2018 (last access Jan 13, 2025)

Weisstein, Eric W. "Regular Icosahedron." From MathWorld--A Wolfram Web Resource. https://mathworld.wolfram.com/RegularIcosahedron.html (last access Jan 9, 2025)

Weisstein, Eric W. "Regular Dodecahedron." From MathWorld--A Wolfram Web Resource. https://mathworld.wolfram.com/RegularDodecahedron.html (last access Jan 9, 2025)

Weisstein, Eric W. "Pentagonal Pyramid." From MathWorld--A Wolfram Web Resource. https://mathworld.wolfram.com/PentagonalPyramid.html (last access Jan 9, 2025)

Weisstein, Eric W. "Augmentation." From MathWorld--A Wolfram Web Resource. https://mathworld.wolfram.com/Augmentation.html (access Jan 9, 2025)

Wolfram Alpha LLC. "Wolfram|Alpha." http://www.wolframalpha.com/input/ (last access Jan 9, 2025)

Figure captions

Fig. 1. Encoding and organisation of the structural proteins in TuYV.
(**a**) TuYV genome organisation (5641 nucleotides). The numbered rectangles represent ORFs, whose functions in the virus cycle are described in Boissinot et al. (2020). The VPg capping is indicated at the genome 5' end. (**b**) Schematic representation of a wild-type virion (TuYV-WT). The capsid is composed of a mixture of the major coat protein (CP) and the minor coat protein (RTP*). An (arbitrary) number of ^{N}RTD extensions (from the RTP*) are represented by small spheres. (**c**) Representation of the ORF3 and 5 in the wild-type virus and their expression products (RTP, RTP* and CP). The readthrough position at the ORF3 stop codon is indicated by a triangle. The amino-acid numbers delimiting the arginine-rich N-terminal stretch, the proline-rich linker, and the putative cleavage site on the RTP are shown. (**d**) Schematic representation of the TuYV-ΔRTP mutant, whose capsid is composed exclusively of the major coat protein. (**e**) as in (**c**) for the TuYV-ΔRTP mutant. The deletion of the ORF5 sequence is indicated by a dotted line.

Fig. 2. EM images of negatively stained and immunogold-labelled TuYV samples.
(**a**) Negatively stained wild-type virions. Protrusions of the capsid (probable presence of the RTP*) are indicated by arrows. (**b**) Negatively stained mutated virions. No clear protrusion of the capsid is observed. (**c**) Elongated particles observed in samples of the mutant. (**d-g**) Immunogold-labelled wild-type virions. Gold particles (darker spots) indicate the presence of the RTP*. (**h**) Mutated virions after immunogold labelling. No gold particle is observed, indicative of the absence of the RTP*. Scale bars are 50 nm.

Fig. 3. TuYV-ΔRTP assembles into nanocrystals, while TuYV-WT does not.
(**a**) Two-dimensional array of negatively stained TuYV-ΔRTP virions. Arrows indicate elongated or distorted particles. (**b**) Cryo-EM image of TuYV-ΔRTP particles arranged in small, thin 3D crystals. (**c**) Negatively stained TuYV-WT virions forming small assemblies of 2 to 4 particles.

Fig. 4. Cryo-EM reconstruction of TuYV-WT and TuYV-ΔRTP
(**a**) Motion-corrected EM images of frozen-hydrated virions deposited on Quantifoil R2/2 grids covered with a carbon film. Scale bar is 100 nm. Images are low-pass filtered at 15 Å for illustration purposes. (**b**) 3D scattering-density (sharpened maps) of the TuYV virions reconstructed by imposing icosahedral symmetry. Three protein subunits, composing one asymmetric unit of the assembly, are highlighted in colours (blue, magenta, green). The underlying geometry of the assembly (equilateral elevated dodecahedron) is represented by coloured (blue and magenta) edges. The EMDB accession codes are reported. (**c**) Sectional view of the unsharpened, unmasked maps, illustrating the outer (protein) and inner (RNA) layers. The density is coloured according to the distance from the centre of the particle. (**d**) Atomic models of the capsid protein layers. (**e**) Comparison of the TuYV-WT and TuYV-ΔRTP atomic models. The icosahedral asymmetric units (trimers) are shown superposed on to each other and coloured by per-residue RMSD. (**f**) Ribbon representation of the atomic model of one of the CP subunits fitted in the cryo-EM map. (**g**) Two views of the atomic model of one of the CP subunits. Secondary structure elements are labelled. Residues involved in inter-subunit contacts are coloured orange. (**h**) Sequence of the three quasi-equivalent CP domains. Unmodelled regions (no interpretable density) are within red frames. Secondary structure elements are highlighted and labelled as in (g).

Fig. 5 Architecture of the TuYV capsid.

(**a**) Top view from the capsid exterior (left) and side view (right) of one asymmetric unit of the icosahedral assembly. The three subunits (chains A, B and C) are pairwise related by almost exact threefold axes (yellow rods). An ideal threefold axis (calculated by a least-squares fitting procedure) is indicated by a blue vertical rod and labelled Q_3. (**b**) Two views of the three subunits superposed onto each other and oriented as chain C in (a). The ideal threefold axis (Q_3) is shown in blue. The results of pairwise superpositions of the asymmetric unit chains are reported. (**c**) Representation of the equilateral elevated dodecahedron (EED) used for describing the TuYV capsid. The three subunits in one asymmetric unit are also shown. Two pairs of triangular faces, one in *endo* and the other in *exo* configuration, are highlighted in green and orange respectively. *Endo* and *exo* edges are coloured magenta and blue respectively. (**d**) Symmetry diagram of a virtual hexagonal lattice (top) and the resulting EED lattice (bottom) upon folding along the *endo* (magenta) and *exo* (blue) edges. The colours of the symmetry symbols in the bottom picture indicate how the nature of the symmetry axes changes upon folding: black means transformation into an icosahedral axis of the same order as in the hexagonal lattice, violet means transformation into an icosahedral axis of a different order, and orange means transformation into a local symmetry axis. (**e**) Arrangement of subunits around one asymmetric unit viewed from the capsid exterior. Subunits related by an icosahedral rotation are coloured the same and labelled with the same letter, while quasi equivalent subunits not related by icosahedral rotations are coloured differently and labelled with different letters. The approximate positions of icosahedral n-fold (I_n, n = 2, 3, 5) and local-symmetry quasi n-fold (Q_n, n = 2, 3, 5, 6) axes are indicated by the yellow labels. The triangle delimits one icosahedral asymmetric unit. *Endo* edges in magenta, *exo* edge in blue. (**f**) Two pairs of asymmetric units forming an *exo* (top) or *endo* edge (bottom). On the left: the triangular faces of and ideal EED are also shown. The leftmost faces are orthogonal to the view direction. On the right: the directions perpendicular to the faces of an ideal EED are indicated by the orange or green rods, while the actual local threefold axes of TuYV are indicated by the blue rods. (**g**) An *endo* and an *exo* pairs of asymmetric units represented superposed. The view direction is along the axis of the virtual hinge movement that would transforms an *endo* angle into an *exo* angle. (**h**) Hexameric arrangement of TuYV subunits around an I_3 axis of the ideal EED. Only subunits coloured the same are related by the icosahedral threefold symmetry. EED edges coloured as in (c). (**i**) A portion of a hexamer (three adjacent subunits) and the rotation axes relating pairs of these subunits. The thicker rods, labelled Q_6, I_3 and Q_5, indicate the actual axis directions in the TuYV capsid. The two thinner rods close to the Q_6 and Q_5 axes represent the directions of the corresponding axes in an ideal EED. For reference, the six EED faces underlying a complete hexamer are partly visible. (**j**) Two quasi-equivalent triplets of subunits, showing that the quasi-symmetry axes relating their chains would be compatible with those of a T=1 icosahedral assembly.

Fig. 6. The eight non-equivalent subunit interfaces.

In each panel a triangular asymmetric unit of the capsid (chains A, B and C) and, eventually, an additional neighbouring chain are depicted. Subunits are labelled as in Fig. 5. The *endo* and *exo* edges delimiting the asymmetric unit are coloured magenta and blue respectively. The viewpoints in panels b+e+h and and c+f differ from the viewpoint in panels a+d+g by a clockwise in-plane rotation of 120° and 240° respectively. In each panel, the interface between two subunits (yellow labels) is highlighted. Interfacial residues are represented in coloured stick mode. N_at: number of interfacial atoms. N_res: number of interfacial residues. N_hb: number of interfacial hydrogen bonds. N_sb: number of interfacial salt bridges.

Fig. 7. The residual amorphous density.

(**a**) Radially averaged density in the unsharpened, unmasked cryo-EM maps. (**b-e**) The residual density (yellow surfaces) after masking out the regions occupied by the CP subunits (shown as cartoon models). A spherical mask of radius 3 Å has been applied to each modelled CP atom. The densities have been contoured at a level corresponding to the peak of the CP-shell in the radially averaged density (the second peak in (**a**)). (**f-h**) Residues of the TuYV-ΔRTP atomic model found in proximity (distance ≤ 5 Å) of the contoured residual density. Similar results (not shown) have been found for TuYV-WT.

Fig. 8. Observation of capsid protrusions derived from ^{N}RTD on TuYV-WT virions by cryo-EM.

(**a**) Top: cryo-EM images of purified TuYV-WT virions in which some ^{N}RTDs, indicated by arrows, are visible in the proximity of the capsids. Bottom: for comparison, some images of the TuYV-ΔRTP in similar orientations. (**b**) The masks that have been used for asymmetric reconstructions of the ^{N}RTD focussed around an I_2 axis (vertical direction). Masks of the same shapes and sizes, aligned with the closest Q_2 and Q_3 axes have also been used. For reference, the putative farthest position of the ^{N}RTD from the capsid is shown assuming a polyproline-like helical conformation of the CP linkers parallel to the symmetry axis. Note that the ^{N}RTD model was never used as a reference during 3D classifications.

Superposed chains [a]	BC≈B_6C_2	AB	CA	C_2B≈CB_6≈C_3B_2	A_2A≈AA_5	B_2C≈ BC_6	CC_2	A_2B≈AB_2
C^α RMSD	0.43 Å	0.47 Å	0.48 Å	0.43 Å	0.00 Å	0.43 Å	0.00 Å	0.47 Å
Rotation angle	119.87°	120.30°	119.96°	60.19°	72.00°	67.34°	180.00°	178.78°
Translation amplitude	0.33 Å	0.45 Å	0.13 Å	0.87 Å	0.00 Å	0.58 Å	0.00 Å	1.28 Å
Rotation label	Q_3	Q_3	Q_3	Q_6 [b]	I_5	Q_5 [b]	I_2	Q_2

[a] Chain labels as in Fig. 5e. The "≈" symbol indicates equivalent interactions according to icosahedral symmetry.
[b] Note that Q_6 and Q_5 compose an I_3 rotation: $Q_6 \cdot Q_5 = I_3$.

Table 1. Geometric relationships between interacting subunits.

Results of pairwise rigid-body superposition of interacting subunits found within one asymmetric unit and its neighbourhoods. Chains are labelled as in Fig. 5e.

Interface type §	Q_3			Q_6	I_5	Q_5	I_2	Q_2
Subunit pairs §	BC	AB	CA	C_2B	A_2A	B_2C	CC_2	A_2B
Interface residues †	B: Glu 129 (s)	A: Glu 129 (s)	C: Glu 129 (s)		A_2: Val 68	B_2: Val 68		A_2: Ser 64 (s)
	B: Leu 130 (h)	A: Leu 130 (h)	C: Leu 130 (h)		A_2: Phe 69	B_2: Phe 69		A_2: Glu 65
	B: Asp 131	A: Asp 131	C: Asp 131	C_2: Ser 70	A_2: Ser 70 (h)	B_2: Ser 70		A_2: Thr 66
	B: Pro 132	A: Pro 132	C: Pro 132	C_2: Asp 72 (h)	A_2: Asp 72 (h)	B_2: Asp 72 (h)	C: Phe 67	A_2: Phe 67
	B: His 133 (hs)	A: His 133	C: His 133 (s)	C_2: Asn 73 (h)	A_2: Asn 73	B_2: Asn 73		A_2: Val 68
	B: Lys 135 (s)	A: Lys 135	C: Lys 135 (s)	C_2: Val 115	A_2: Val 115	B_2: Val 115		A_2: Cys 91
	B: Leu 136	A: Leu 136	C: Leu 136		A_2: Ser 116 (h)	B_2: Ser 116 (h)	C: Pro 92	A_2: Pro 92
	B: Ser 140		C: Ser 140	C_2: Glu 117 (hs)	A_2: Glu 117 (hs)	B_2: Glu 117 (hs)	C: Ala 93	A_2: Ala 93
	B: Ser 141 (h)	A: Ser 141 (h)	C: Ser 141	C_2: Ala 118	A_2: Ala 118	B_2: Ala 118	C: Asn 96	A_2: Asn 96
	B: Thr 142			C_2: Ser 119 (h)	A_2: Ser 119 (h)	B_2: Ser 119 (h)	C: Gly 97	A_2: Gly 97
	B: Ile 143	A: Ile 143	C: Ile 143	C_2: Ser 120	A_2: Ser 120 (h)	B_2: Ser 120 (h)	C: Met 98	A_2: Met 98
	B: Asn 144	A: Asn 144	C: Asn 144	C_2: Gln 121	A_2: Gln 121	B_2: Gln 121 (h)	C: Lys 100	A_2: Lys 100
		A: Phe 156	C: Phe 156	C_2: Asn 122	A_2: Asn 122	B_2: Asn 122	C: Ala 101	A_2: Ala 101
	B: Ser 159	A: Ser 159	C: Ser 159		A_2: Thr 149	B_2: Thr 149 (h)		A_2: Tyr 102
	B: Tyr 160	A: Tyr 160	C: Tyr 160		A_2: Pro 151	B_2: Pro 151		A_2: Arg 191
	B: Asn 162	A: Asn 162	C: Asn 162	C_2: Ser 185	A_2: Ser 185	B_2: Ser 185		A_2: Phe 198
	B: Thr 164 (h)	A: Thr 164 (h)	C: Thr 164 (h)	C_2: Ile 186 (h)	A_2: Ile 186	B_2: Ile 186		A_2: Pro 201
	B: Glu 165	A: Glu 165	C: Glu 165	C_2: Ser 189 (h)	A_2: Ser 189	B_2: Ser 189		A_2: Lys 202
					A_2: Arg 191	B_2: Arg 191 (h)		
	C: Gly 62							B: Ser 64 (s)
	C: His 103 (s)	B: His 103 (s)	A: His 103 (s)	B: Ser 120 (h)	A: Ser 120 (h)	C: Ser 120 (h)		B: Glu 65
	C: Glu 104 (s)	B: Glu 104	A: Glu 104 (s)	B: Gln 121	A: Gln 121 (h)	C: Gln 121 (h)		B: Thr 66
	C: Glu 165 (h)	B: Glu 165 (h)	A: Glu 165 (h)	B: Asn 122 (h)	A: Asn 122	C: Asn 122	C_2: Phe 67	B: Phe 67
	C: Trp 166	B: Trp 166	A: Trp 166	B: Ser 123 (h)	A: Ser 123	C: Ser 123		B: Val 68
	C: His 167	B: His 167	A: His 167	B: Gly 124 (h)	A: Gly 124 (h)	C: Gly 124 (h)		B: Cys 91
	C: Asp 168	B: Asp 168	A: Asp 168	B: Ser 125 (h)	A: Ser 125 (h)	C: Ser 125	C_2: Pro 92	B: Pro 92
	C: Ala 170	B: Ala 170	A: Ala 170		A: Tyr 128		C_2: Ala 93	B: Ala 93
	C: Glu 171 (s)	B: Glu 171	A: Glu 171 (s)	B: Thr 142 (h)	A: Thr 142 (h)	C: Thr 142 (h)	C_2: Asn 96	B: Asn 96
	C: His 199	B: His 199	A: His 199	B: Ile 143	A: Ile 143	C: Ile 143	C_2: Gly 97	B: Gly 97
	C: Asn 200 (h)	B: Asn 200 (h)	A: Asn 200 (h)		A: Asn 144	C: Asn 144 (h)	C_2: Met 98	B: Met 98
	C: Pro 201	B: Pro 201	A: Pro 201	B: Lys 145 (hs)	A: Lys 145 (s)	C: Lys 145 (s)	C_2: Lys 100	B: Lys 100
	C: Lys 202 (h)	B: Lys 202 (h)	A: Lys 202			C: Phe 146	C_2: Ala 101	B: Ala 101
					A: Gly 147	C: Gly 147	C_2: Tyr 102	B: Tyr 102
					A: Thr 149	C: Thr 149		B: Arg 191
					A: Lys 150 (h)	C: Lys 150 (h)		B: Phe 198
					A: Arg 154	C: Arg 154		B: Pro 201
				B: Lys 179				B: Lys 202
				B: Asn 181 (h)	A: Asn 181 (h)	C: Asn 181 (h)		
				B: Gly 182		B: Gly 182		

§ Chains and rotation axis labelled as in Fig. 5e.
† (h) and (s) indicate residues involved in interfacial hydrogen bonds and salt bridges respectively.

Table 2. Residues at the interfaces between subunits.

a

VPg

0		
1		
2	3a	
	3	5
	4	

5' 3' ssRNA(+)

b

RTP*

CP

TuYV-WT

c

readthrough ▽

| 3 | 5 |

ssRNA(+)

Arg-rich stretch 1–61 Pro-rich linker 203–223 Cleavage site ~459 ~460 670

CP domain		N_RTD	C_RTD	**RTP** (readthrough protein)
CP domain		N_RTD		**RTP*** (minor coat protein)
CP domain				**CP** (major coat protein)

d

CP

TuYV-ΔRTP

e

| 3 | - - - - - - - | ssRNA(+) |

Arg-rich stretch 1–61 202

| CP domain | **CP** (major coat protein) |

b

Superposed subunits	Cα RMSD	Nb. of superposed residues	Rotation angle	Translation
AB	0.47 Å	138	120.30°	0.45 Å
BC	0.43 Å	138	119.87°	0.33 Å
CA	0.48 Å	139	119.96°	0.13 Å

c

endo edge
360°−arccos[(−√5)/3]
=221.81°

exo edge
arccos(−8/15−1/√5)
=168.68°

T=3 equilateral elevated dodecahedron (EED)

d

hexagonal lattice
(p6 plane group)

T=3 EED

f

exo

endo

top views
(from capsid exterior)

capsid exterior
capsid interior

capsid exterior
capsid interior

side views
(along exo/endo edge)

g

~174°
ideal: ~168.68°
arccos[−8/15−1/√5]

capsid exterior

~45°
ideal: ~53.13°
arctan[4/3]

capsid interior

~141°
ideal: ~138.19°
arccos[−(√5)/3]

i

B₂→C₃
60.2° rot
(ideal 60°)
Q_6

C→C₃
120° rot
I_3

C→B₂
67.3° rot
(ideal 72°)
Q_6

23°
20.7°

j

capsid exterior

centre of an hypothetical
T=1 icosahedron

capsid interior

a

exo

C

A

B

Q_3 interface

N_at	113
N_res	30
N_hb	4
N_sb	4
Area	557 Å²

b

endo

B

C

A

Q_3 interface

N_at	112
N_res	28
N_hb	4
N_sb	1
Area	504 Å²

c

endo

A

B

C

Q_3 interface

N_at	110
N_res	29
N_hb	3
N_sb	4
Area	533 Å²

d

exo

C

A

C_2

B

Q_6 interface

N_at	81
N_res	25
N_hb	9
N_sb	1
Area	388 Å²

e

endo

B

C

A_2

A

I_5 interface

N_at	116
N_res	35
N_hb	10
N_sb	1
Area	556 Å²

f

endo

A

B

B_2

C

Q_5 interface

N_at	112
N_res	35
N_hb	9
N_sb	1
Area	535 Å²

g

exo

C

A

C_2

B

I_2 interface

N_at	54
N_res	17
N_hb	0
N_sb	0
Area	264 Å²

h

endo

B

C

A_2

A

Q_2 interface

N_at	110
N_res	36
N_hb	4
N_sb	2
Area	510 Å²

a — Plot of average density versus distance from particle centre (Å), comparing TuYV-WT (red) and TuYV-ΔRTP (blue). Labels indicate "residual density" and "CP shell".

b TUYV-WT — view from the capsid exterior

c TUYV-ΔRTP — view from the capsid exterior

d TUYV-WT — view from the capsid interior

e TUYV-ΔRTP — view from the capsid interior

f TUYV-ΔRTP — side view (-90°)

g TUYV-ΔRTP — view from the capsid interior

h TUYV-ΔRTP — side view (+90°)