Mechanistic insights into the inhibition of dengue virus NS5 methyltransferase by

herbacetin

Mandar Bhutkar, Shalja Verma, Vishakha Singh, Pravindra Kumar, Shailly Tomar *

Department of Biosciences and Bioengineering, Indian Institute of Technology Roorkee,

Uttarakhand, India.

*Corresponding author address: Department of Biosciences and Bioengineering, Indian Institute

of Technology Roorkee, Roorkee-247667, Uttarakhand, India; Email id:

shailly.tomar@bt.iitr.ac.in

Abstract: -

Herbacetin (HC) is a naturally occurring flavonoid compound with a dual antiviral mechanism. It

inhibits the polyamine pathway and targets the methyltransferase (MTase) enzyme of the dengue

(DENV3) and chikungunya virus. However, understanding the detailed inhibition mechanism of

DENV MTase by HC remains unclear. The crystal structure of DENV3 nonstructural protein 5

(NS5) MTase in complex with HC and S-adenosyl-L-homocysteine presented in this study gives

structural insights into the inhibition mechnism. Structural analysis revealed that HC binds to the

cap-0 RNA site near the GTP binding site in the DENV3 NS5 MTase and fluorescence

polarization assay demonstrated HC inhibits GTP binding with an inhibition constant K_i value of

~ 0.43 µM. Further, the *in vitro* cell-based assays demonstrated that HC effectively inhibits

replication of all the serotypes of DENV. This is the first study that identified an inhibitor that

targets the conserved RNA-binding region of NS5 MTase, suggesting its potential as a highly

effective scaffold for broad-spectrum antiviral agents against orthoflaviviruses.

Importance:-

Dengue virus (DENV) is a vector-borne disease affecting global health, and no antiviral treatment is available for Dengue fever. DENV NS5 methyltransferase (DENV NS5 MTase) is a viral RNA capping enzyme essential for the virus replication. This study demonstrated that herbacetin (HC), a natural flavonoid, blocked the binding of GTP to NS5 MTase and inhibited the enzyme. In addition to DENV3, HC effectively inhibited the replication of DENV1,2, and 4 serotypes. The crystal structure of the DENV3 NS5 MTase in complex with HC revealed that HC binds at the cap-0 RNA binding site, which is close to the GTP binding site and thereby perturbs the spatial arrangement of GTP binding site residues. HC, as the first identified inhibitor that targets the conserved RNA-binding site of DENV3 NS5 MTase, presents a potential scaffold for the development of new antiviral drugs.

Introduction: -

The *Flaviviridae* family comprises the most widespread arboviruses that can cause life-threatening human diseases. This family includes the genus *Orthoflavivirus,* which encompasses various viruses such as Dengue virus (DENV), Zika virus (ZIKV), West Nile virus (WNV), and Japanese encephalitis virus (JEV), etc (1). DENV has four (DENV 1-4) serotypes that can cause infection ranging from asymptomatic to mild symptoms such as fever, rash, joint pain, headache, etc. In severe cases, DENV infection can lead to dengue shock syndrome and dengue hemorrhagic fever (2). DENV gets transmitted by *Aedes aegypti* and *Aedes albopictus* mosquitoes. Approximately 390 million infections occur annually, with most cases reported from tropical and subtropical countries (3). DENV infections are endemic in more than 30 countries,

including India, Thailand, Sri Lanka and Myanmar (4). Although the FDA has approved the

vaccine for DENV (Dengvaxia), its use is limited to individuals living in endemic areas aged 6 to

16 years with previously laboratory-confirmed DENV infection (5). Considering the limitations

of the vaccine and the absence of antiviral drugs, there is an imminent need for the development

of effective antiviral treatments for DENV infection.

DENV is an enveloped, positive sense RNA virus having a single open reading frame, which

encodes for three structural proteins (capsid, envelope, and preM) and five nonstructural proteins

from NS1 – NS5 (6). NS5 is the largest and most highly conserved protein, comprising

approximately 900 amino acid residues among all nonstructural proteins (7). It consists of two

domains, N-terminal methyltransferase (MTase) and C-terminal RNA-dependent RNA

polymerase (RdRp) where each domain can function independently of the other (8). The viral

RNA has a type 1 cap structure (m7GpppAmp) at the 5' end, which plays a crucial role in the

translation of viral proteins as well as in evading host immune responses (9). The mechanism of

orthoflaviviral MTase comprises two steps. The first step involves the transfer of a methyl group

from s adenosine methionine (SAM) to the N7 position of the guanine base situated at the 5'

terminus of the viral RNA. The second step involves the methylation of 2 O' ribose sugar of the

initially transcribed nucleotide (9,10). In orthoflaviviruses, K61-D146-K180-E216 amino acid

arrangement is essential for the 2 O' methylation and remains conserved. It is a catalytic tetrad

where Lys180 plays a crucial role in transferring methyl group from SAM to 2 O' ribose sugar of

adenine. Additionally, Lys61, Asp146, and Glu216 help to stabilize the substrate and the

transition state during the reaction, ensuring that the 2 O' methylation occurs efficiently (11).

DENV NS5 MTase x-ray crystal structure revealed the presence of s adenosine methionine

(SAM), guanosine triphosphate (GTP), and cap 0 RNA binding sites (8,11). Since NS5 MTase

has only one SAM binding site and dual methylation function (N7 and 2 O'), the RNA substrate

must be repositioned to methylate 2 O' ribose sugar of the initially transcribed nucleotide (12).

Herbacetin (HC) is a naturally occurring flavonoid in plants, principally flaxseed, parsley, and

equisetum (Ramose scouring rush) (13). Herbacetin is structurally close to quercetagatein and

isoflavone and exerts various pharmacological activities, including antiviral, antioxidant, anti-

inflammatory, and anticancer effects (13–15). Moreover, HC is a known allosteric ornithine

carboxylase inhibitor that depletes cellular polyamine levels upon treatment (16,17).

Furthermore, HC exhibited viral MTase inhibition, leading to antiviral activity against DENV

and CHIKV with IC_{50} values in µM range (17). However, the structural aspects of inhibition of

DENV3 MTase by HC are yet to be revealed and are important to develop a high-efficacy

scaffold for the development of drugs. Also, the effect of HC interaction on the binding of GTP

in DENV3 NS5 MTase and its antiviral activity across different DENV serotypes is unknown.

To address these questions, this study presents a detailed crystal structure of the DENV NS5

MTase in complex with HC. In addition, a fluorescence polarization-based assay analyzed the

inhibitory effects of HC on GTP binding, which may underlie its broad-spectrum antiviral

activity across all dengue serotypes. Therefore, the findings of this study offer a detailed

understanding of the inhibition mechanism of HC.

Methodology: -

FP polarisation

The fluorescence polarization (FP) assay was employed to measure the displacement of FAM-

labeled GTP (F-GTP) binding to the DENV-3 NS5 methyltransferase (MTase) by HC. For K_D

determinations, reaction mixtures were prepared with FP binding buffer (10 mM KCl,50 mM

Tris, pH 7.5, 2 mM $MgCl_2$, and 2 mM DTT , increasing concentrations of F-GTP (3.9 nM to 125

nM) and 250 nM NS5 MTase. For IC_{50} determinations of HC, the reaction mixtures contained FP

binding buffer, increasing concentrations of HC (0 to 100 µM), 10 nM F-GTP, and 250 nM NS5

MTase. All samples were incubated in the dark at 25°C for 1 h before measuring total FP using a

Synergy HTX multimode plate reader (Agilent BioTek) with Gen5 software, at an excitation

wavelength of 485/20 nm and an emission wavelength of 528/20 nm. Baseline fluorescence

polarization values from control samples containing only F-GTP (without protein) were

subtracted to remove background signals. Data from three independent experiments collected

and analysed. K_D values were calculated using nonlinear regression analysis with a sigmoidal

curve (18). IC_{50} values for HC were determined through nonlinear regression analysis,

employing the log(inhibitor) versus normalized response curve (18), as analyzed by GraphPad

Prism 8.0. The inhibition constant (Ki) for HC was calculated using the Cheng-Prusoff equation:

$K_i = IC_{50}/1 + ([F\text{-}GTP]/K_D)$ (18–20).

Antiviral activity assay

The DENV serotypes 1 (Hawaii – NR82), 2 (New Guinea C – NR51427), and 4 (H241 (TC) –

NR86) from BEI Resources were propagated according to the protocol detailed by Bhutkar et

al.2022 (17). The propagated viruses were then aliquoted and stored at -80°C until further use.

Vero cells were plated in a 24-well plate at a density of 1.0 x 10☐ cells per well. HC 200 µM

concentration that maintained cell viability above 90% were used for the treatment (17). Cells

were pre-treated with HC 200 µM 12 h prior to infection. The cell monolayer was then infected

with the virus at a multiplicity of infection (MOI) of 0.1, with gentle shaking every 15 minutes

for 2 hours. Following this, the inoculum was removed, and the cells were washed twice to

minimize the risk of secondary infection. HC 200 µM was added to the post-infection (pi)

medium containing 2% DMEM, and the cells were incubated for 24 h. After 24 hpi, the medium

containing the HC compound was replaced with fresh 2% DMEM, which continued for an

additional 4-5 days (17).

Next, the supernatant was collected to assess the virus titer using a plaque-forming assay, and

plaques were visualized through an immunostaining protocol as previously described by Bhutkar

et al. (2022) (17,21). Data from three independent experiments were collected, and graphs were

created using GraphPad Prism 8.0.

Quantitative Real-Time PCR (qRT-PCR)

The treatment with the HC compound and DENV infection followed the procedure outlined in

the antiviral activity assessment. After completing the assay, RNA extraction was performed

using the TRIzol protocol as recommended by the manufacturer. The extracted RNA was then

quantified, and cDNA was synthesized using the PrimeScript 1st strand cDNA Synthesis Kit

(Takara), starting with 400 ng of RNA. Quantitative real-time PCR (qRT-PCR) was carried out

using the KAPA SYBR Fast Universal qPCR Kit, with the analysis performed on the

QuantStudio™ 5 System (Applied Biosystems, USA). Amplification involved the forward

(mD1) and reverse (D2) primers (22), with β-actin as the internal control (17). Data from three

independent experiments were collected, and relative quantification was performed using the

ΔΔCt method as described in (23). Graphs were prepared using GraphPad Prism 8.0.

Immunofluorescence Assay

Vero cells were plated in a 24-well plate at a density of $1.0 \times 10^\square$ cells per well. As previously

stated, cells were treated with HC 200 µM in an antiviral activity assay. Afterward, the cells were

washed three times with PBS and fixed in 3.7% formaldehyde for 30 min at room temperature

(RT). They were then permeabilized with 0.1% Triton-X-100 for 10 min. Following another

wash with 1X PBS, the cells were incubated with a DENV-specific antibody (BEI Resources

NR-50327, monoclonal anti-flavivirus group antigen, clone d1-4g2-4-15, diluted 1:1500) for 1

hour. The cells were then rinsed with 0.1% Tween-20 in 1X PBS (PBS-T). Finally, a fluorescein

(FITC)-conjugated secondary anti-mouse antibody (1:250, Sigma) was added, and the cells were

incubated in the dark at RT for 1 h. The cells were then rinsed with PBST and counter-stained

with 4′,6-diamidino-2-phenylindole (DAPI, Sigma) for 15 min in the dark. Finally, the images

were captured using a fluorescence microscope (EVOS FL AUTO, Thermo Fisher) (24).

Crystallization of NS5 MTase

DENV3 NS5 MTase was expressed and purified using recombinant plasmid pET28c(+)-NS5

MTase in *E. coli* BL21 cells, as described in the previous study (17). DENV3 NS5-MTase was

crystallized employing the sitting drop vapor diffusion method in 96-well plates at 20°C,

following the procedure outlined by Coutard et al., 2014 (17,25,26). To obtain the complex of

DENV3 NS5-MTase with HC, soaking was done in cryoprotectant solution (22.5% glycerol)

containing ~ 1 mM HC for 30 min. The crystals were then flash-frozen in liquid nitrogen at

100K. The data collection of DENV3 NS5-MTase crystals in complex with HC was performed

using Rigaku Micromax 007HF diffractometer integrated with Hypix 600C detector at the

Macromolecular crystallography unit, IIT Roorkee. Data reduction and scaling were performed

using the CrysAlis Pro PX software (Rigaku Inc.). The structure was solved by molecular

replacement method using the MOLREP program of CCP4i2, giving DENV3 NS5 MTase (PDB

code 4R8R) as a search model (27). Iterative rounds of model-building in COOT and the

refinements were performed using the REFMAC5 (28) program in CCP4i2 (29). Non-

crystallographic symmetry (NCS) and jelly-body restraints were applied during the entire data

refinement process. Eventually, omit maps were generated in the Polder omit map program in

Phenix software (30). Table 1 provides a summary of the data collection and refinement

statistics. All figures of protein and ligand structures were prepared using PyMOL (31).

Results: -

Inhibition of GTP binding to DENV3 NS5 MTase:

A previous study demonstrated that HC binds to the recombinantly purified DENV3 NS5 MTase

in the micromolar range K_D value. This binding was subsequently shown to exhibit anti-NS5

MTase activity. *In silico* work predicted that HC may bind within the GTP binding site (17). To

investigate whether HC can displace GTP binding to DENV3 NS5 MTase, a fluorescence

polarization (FP) assay was performed. To determine the dissociation constant (K_D) of F-GTP,

increasing concentrations of F-GTP were incubated with NS5 MTase. The binding curve

revealed a sigmoidal relationship between F-GTP concentration and fluorescence polarization,

resulting in a calculated K_D of ~13.12 ± 1.02 nM (Figure 1A). This value indicates a high

binding affinity of F-GTP for DENV3 NS5 MTase. A competition assay was conducted further to

explore the inhibitory effect of HC on F-GTP binding. The inhibition curve demonstrated that

HC effectively displaced F-GTP from the NS5 MTase, with a half-maximal inhibitory

concentration (IC_{50}) of 0.77 ± 0.01 µM (Figure 1B). Using the Cheng-Prusoff equation, the

inhibition constant (K_i) of HC was calculated to be approximately 0.43 ± 0.01 µM, suggesting

that HC may inhibit GTP binding to DENV3 NS5 MTase.

Antiviral efficacy against all the serotypes of dengue:

The antiviral activity of HC at 200 µM was assessed by following the methodology described by

Bhutkar et al. (2022) to ensure consistency in results. Earlier, HC was reported to show antiviral

activity against the DENV3 serotype (17). Here, treatment with HC in Vero cells has

demonstrated a reduced virus titer in theDENV1,2,4 individually-infected cells as compared to

respective untreated virus control (VC) (Figure 2D, 2E, 2F). Likewise, qRT-PCR was employed

to confirm the antiviral effect of HC on viral RNA levels in cells infected with DENV1, DENV2,

and DENV4. The results from qRT-PCR demonstrated a significant reduction in viral RNA levels

($p < 0.001$, $p < 0.0001$) in HC-treated cells during the antiviral assay for DENV1, DENV2, and

DENV4 (Figure 2A, 2B, 2C). Furthermore, HC exhibited approximately a 3-fold reduction in

DENV1 and DENV4 and a 10-fold reduction in DENV2, compared to the VC. The IFA results

demonstrated a decrease in DENV 1,2,4 after treatment with HC compared to VC (Figure 3).

Structural insights into DENV3 NS5 MTase with HC

The structure of the DENV3 NS5 MTase in complex with HC (PDB: 8ZMC) was determined in

P 2 2$_1$ 2$_1$ space group at 2.8 Å with an R$_{work}$ of 0.24 and an R$_{free}$ of 0.30. The structure was solved

using molecular replacement method, with the apo form of DENV3 NS5 MTase (PDB ID:

4R8R) as a search model with 99.2 % identity. The refinement and data collection statistics are

presented in Table 1. Structural analysis (Table 2) revealed the formation of hydrogen bonds and

specific hydrophobic interactions, pivotal for the cap 0 RNA and GTP binding (31,32). One HC

molecule was modeled in one chain of the DENV3 NS5 structure based on the difference Fourier

map, which matched the observed electron density with a real space correlation coefficient

(RSCC) of 0.78 (Figure 4B). The polder map contoured at 2.5 σ, indicated its precise location

within the DENV3 NS5 MTase (Figure 4A). The B-factor of HC was observed to be 34.1, which

is similar to the B-factors of the nearby amino acids. The Ramachandran outliers were 0.59%,

and none of the HC binding site residues contributed to this small fraction, thus proving the

accuracy of the determined interaction of the binding site residues with HC (Table 1). In HC, O3

forms a hydrogen bond with the NZ of Lys42 at 2.85 Å, and O6 forms a hydrogen bond with the

NE of Arg57 at a distance of 2.88 Å. HC hydrophobically interacts with Arg211, which is crucial

for interacting with the phosphate group of GTP (Table 3) (Figure 4C). Although SAH was not

added during crystallization, the electron density of SAH in the difference Fourier map was

observed in both chains, suggesting that it would have been obtained from *E. coli* during protein

expression (PDB:8ZMC) (Figure 4B). A similar observation of electron density of SAH obtained

from the host have been reported in previous studies (8,19) Additionally, SAH forms hydrogen

bonds with the residues Asp131, Gly86, Ser56, Trp87, Lys105, and Val132. Hydrophobic

interactions involve residues His110, Gly58, Gly81, Asp146, Lys130, Thr104, Gly83, Gly81,

Asp146, Cys82, Gly85, and Phe133 (Figure 4D). The surface representation of the DENV3 NS5

MTase-HC superimposed with DENV3 NS5 MTase-SAH-cap 0RNA (PDB:5DTO) structures

showed HC binds within the cap 0 RNA binding pockets (Figure 5A). The main chain

conformation of the DENV3 NS5 MTase-HC complex closely resembles that of the DENV NS5-

SAH-cap 0 RNA structure (PDB: 5DTO), with an RMSD of 0.354 Å across 233 Cα atoms

(Figure 5B). In DENV3 NS5-SAH-cap 0 RNA, NH2 of Arg57 forms hydrogen bonds with OP1

of A-G (Table 4), while in the DENV3 NS5 MTase-HC complex the O6 from HC forms

hydrogen bonds with NE of Arg57, displaying binding of HC at the cap 0 RNA binding site

(Table 3) (Figure 5D). The structural superposition of both complexes revealed that the HC

interaction with Arg57 mediated angular variation of 124° the residue (Figure 5D). The

superposition of DENV3 NS5 MTase bound to HC and GTP (PDB: 4V0R) demonstrates that HC

forms hydrophobic interactions with Arg211, whereas GTP establishes a hydrogen bond with

Arg211 (Figure 4C). These hydrophobic interactions of HC with the Arg211 residue of the GTP

binding site conveyed a slight conformational change in Arg211. Additionally, Lys42 exhibits a

notable angular shift of 51.6° in both complexes (Figure 5C).

Consequently, the presence of HC at this site may obstruct the binding of GTP and viral RNA to

the DENV3 NS5 MTase, resulting in effective structural inhibition of the enzyme.

Conserved HC binding residue predicted to make stable interactions across DENV MTases

of all serotypes

DENV 1-4, ZIKV, West Nile Virus (WNV), Yellow Fever Virus (YFV), and Japanese

Encephalitis Virus (JEV) are orthoflaviviruses that cause infections in humans. Multiple

sequence alignment (MSA) showed that the HC interacting residues Arg57, Lys42, and Arg211

remain conserved across all DENV serotypes (Supplementary figure 1). Also, in other

orthoflaviviruses such as ZIKV, JEV, and YFV, Lys42 is replaced by another positively charged

amino acid Arg42, having similar chemical properties (Supplementary figure 1). As the

interacting residues (Arg57, Lys42, and Arg211) are conserved across all dengue serotypes, it is

reasonable to hypothesize that HC would likely engage in similar key interactions with these

residues across different serotypes. However, the non-interacting residues that vary between the

serotypes could potentially influence the overall stability, orientation, or even the interaction

dynamics of HC within the binding pocket. Leveraging this sequence conservation, HC

coordinates from the DENV3 complex were superimposed onto the MTases of DENV1, 2, and 4.

This superimposition was performed to determine the residues occupying structurally equivalent

position and interaction with the active site, as observed in the crystallized DENV3 MTase-HC

complex, applied to all DENV serotypes (33).

Further, MD simulation studies were conducted to evaluate the interaction stability of MTase-HC

complexes, with DENV3-HC MTase as the control for comparisons with other serotypes. The

RMSD plots for DENV1, DENV2, DENV3, and DENV4 serotypes in complex with HC over the

100 ns MD simulation suggest stable binding interactions. The average RMSD value for DENV1

is 0.12 nm, DENV2 is 0.13 nm, DENV3 is 0.10 nm, and DENV4 is 0.12 nm (Supplementary

figure 2A). The root mean square fluctuation (RMSF) data suggested that in MTase-HC

complexes, the HC binding residues showed minimal fluctuations, indicating rigid and stable

interactions with the ligand (Supplementary figure 2 B). The radius of gyration (Rg) values for

all four DENV serotypes exhibit a similar trend, indicating a stable conformation with average

Rg value for DENV1 is 0.186 nm, DENV2 is 0.187 nm, DENV3 is 0.185 nm and DENV4 is

0.186 nm (Supplementary figure 2C). The MTase-HC complexes maintained stable hydrogen

bonds (H-bonds) throughout the simulation with an average number of H-bonds for DENV1 is

1.30, DENV2 is 1.25, DENV3 is 1.77, and DENV4 is 0.73 (Supplementary figure 2D). The

solvent accessible surface area (SASA) is key to understanding protein stability and folding

DENV1, DENV3, and DENV4 complexes show comparable SASA values, whereas DENV2

exhibits slightly higher SASA values (Supplementary figure 2E). The binding energies that is

van der Waals forces and electrostatic interaction, were calculated in the terms of Lennard-Jones

short-range (LJ-SR) and Coulombic SR (COU-SR), respectively. The LJ-SR in KJ/mol DENV1

is -71.6, DENV2 is -107.9, DENV3 is -75.15, and DENV4 is -32.09 and COU-SR in KJ/mol

DENV1 is -60.75, DENV2 is -44.76, DENV3 is -51.88, and, DENV4 is -27.24 (Supplementary

figure 2F). MD simulations of DENV serotypes reveal stable conformations over 100 ns, with

subtle differences in structural dynamics and solvent exposure.

Discussion: -

Previously, HC has been reported to exhibit antiviral efficacy against DENV3 by targeting the

viral replication enzyme NS5 MTase (17). This study has determined the crystal structure of

DENV3 NS5 MTase in complex with the inhibitor HC to elucidate its inhibition mechanism.

Additionally, the FP assay demonstrates that GTP is displaced due to the binding of HC (K_i value

of ~ 0.43 µM) to the conserved RNA binding site of NS5 MTase, which perturbs the spatial

arrangement of GTP binding residues (Figure 5C). The current study has further expanded the

findings and confirmed the broad-spectrum antiviral activity of HC against all the DENV

serotypes 1, 2, and 4 (Figure 3) (17).

The crystal structure of the HC-DENV3 MTase complex demonstrates that one HC molecule

binds to the cap 0 RNA and near the GTP binding site (Figure 5). No differences were observed

in the SAH molecule interactions compared to those reported in Zhao et al. 2015 (PDB: 4V0R)

(8). GTP primarily establishes a hydrogen bonds with Lys14, Leu17, Asn18, Leu20, Arg22,

Ser151, Arg211, and Ser213, residues and alongside forms multiple hydrophobic interactions

involving Asn18, Phe25, Lys29, Ser150, and Pro152 residues (PDB:4V0R) (Table 3) (8). The

RNA bases A-G form hydrogen bonds with Ser150, Thr214, Lys180, Glu111, and Arg57, and

hydrophobic interactions with residues Leu182, Arg84, Lys61, Gly148, Ser56, and Glu216

(PDB:5DTO) (Table 4) (11). The 3-hydroxyphenyl ring of HC formed Pi-alkyl bond with Arg 57

while the trihydroxyphenyl ring formed Pi alkyl interaction with Lys42 revealing the binding of

HC in the RNA binding site. Moreover, HC engages in hydrophobic interaction with Arg211.

GTP forms two hydrogen bonds with Arg211, where one hydrogen bond is with the oxygen atom

(O1B) of the second phosphate of GTP at a distance of 3.3 Å, and the other is with the oxygen

atom (O3B), which links the second and third phosphates, at a distance of 3.2 Å (Table 3) (PDB:

4V0R). Thus, the interaction of HC with DENV3 MTase suggests its potential to display dual-

mode inhibition of RNA and GTP binding.

Similarly, HC forms a hydrogen bond with Arg57 and Lys42, where Arg57 is a crucial residue

that has been demonstrated to form hydrogen bonds with the viral RNA (PDB:5DTO) (11). The

mutation Arg57Ala in the West Nile virus (WNV) MTase reduced N7 MTase activity to less than

20%(12). The detailed role of Lys42 is not well understood, but it may play an essential role in

maintaining the structural stability of the NS5 MTase. Although this work did not investigate the

inhibition of viral RNA binding to DENV3 NS5 MTase, HC's interaction with key residues

involved in RNA binding and reported MTase inhibition may hamper the viral RNA capping

process (Table 2) (17). In MD simulations, the conserved HC binding residue is predicted to

make stable interactions across the MTases of all DENV serotypes.

In conclusion, HC binds to the cap 0 RNA binding site, at the vicinity of the GTP binding site,

thereby interfering with GTP binding and inhibiting all dengue serotypes. Further, *in vitro* and *in*

vivo research is warranted to evaluate its efficacy across other orthoflaviviruses such as JEV,

YFV, ZIKV, etc. Moreover, structural information and antiviral activity are anticipated to provide

crucial insights for developing novel broad-spectrum inhibitors.

Acknowledgments

The authors thank the Department of Biosciences and Bioengineering, IIT Roorkee for central

facilities and the translational and structural bioinformatics center. The authors thank the

Macromolecular Crystallographic Facility (MCU) at IIC, Indian Institute of Technology,

Roorkee. ST and PK thank the Department of Biotechnology, Govt of India, for supporting

Bioinformatics Center at IIT Roorkee ref number BT/PR40141/BTIS/137/16/2021. The authors

thank the Ashok Soota Molecular Medicine facility, IIT Roorkee. ST and PK also would like to

thank Department of Biotechnology, Govt of India "National Network Project of Department of

Biotechnology, Indian Institute of Technology, Roorkee" Project no.

BT/PR40142/BTIS/137/72/2023. The authors would like to thank the Council of Scientific and

Industrial Research (CSIR), Indian Council of Medical Research (ICMR), and Ministry of

Human Resource Development (MHRD) for providing financial support. This work was

supported by a research grant to ST from DST SERB ref no. CRG/2023/002214

321   Ethical approval: -

322   Ethical approval has been received for the work under the reference number

323   IITR/IBSC/02/13/2024.

324   Supplemental information: -

325   Supplemental information includes two supplemental experimental procedures and two Figures.

326   Authors contribution

327   MB, SV, and VS conducted the experiments. MB, SV, VS, PK, and ST planned the experiments

328   and analyzed the data. MB, PK, and ST took the lead in writing the manuscript. All authors

329   provided critical feedback and helped shape the research, analysis, and manuscript.

332   Competing interests

333   The authors declare that they have no competing interests.

References:-

1.   Postler TS, Beer M, Blitvich BJ, Bukh J, de Lamballerie X, Drexler JF, et al. Renaming of the genus Flavivirus to Orthoflavivirus and extension of binomial species names within the family Flaviviridae. Arch Virol. 2023;168(9):1–7. DOI: 10.1007/s00705-023-05835-1

2.   Dengue and severe dengue. https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue

3.   Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution and burden of dengue. 2013;496(7446):504–7. DOI: 10.1038/nature12060

4.   Dengue- Global situation. https://www.who.int/emergencies/disease-outbreak-news/item/2023-DON498

5.   Thomas SJ, Yoon IK. A review of Dengvaxia®: development to deployment. Hum Vaccin Immunother. 2019 (10):2295–314. DOI: 10.1080/21645515.2019.1658503

6.   Murugesan A, Manoharan M. Dengue Virus. Emerging and Reemerging Viral Pathogens.2020 Jan 1;281. DOI: 10.1016/B978-0-12-819400-3.00016-8

7.   Lim SP, Noble CG, Shi PY. The dengue virus NS5 protein as a target for drug discovery. Antiviral Res. 2015 ;119:57–67. DOI: 10.1016/j.antiviral.2015.04.010

8.   Zhao Y, Soh TS, Zheng J, Chan KWK, Phoo WW, Lee CC, et al. A Crystal Structure of the Dengue Virus NS5 Protein Reveals a Novel Inter-domain Interface Essential for Protein Flexibility and Virus Replication. PLoS Pathog. 2015;11(3):1–27. DOI: 10.1371/journal.ppat.1004682

9.   Furuichi Y, Shatkin AJ. Viral and cellular mRNA capping: Past and prospects. Adv Virus Res. 2000;55:135. DOI: 10.1016/s0065-3527(00)55003-9

10.   Bisaillon M, Lemay G. Viral and Cellular Enzymes Involved in Synthesis of mRNA Cap Structure. Virology. 1997 ;236(1):1–7. DOI: 10.1006/viro.1997.8698

11.   Zhao Y, Soh TS, Lim SP, Chung KY, Swaminathan K, Vasudevan SG, et al. Molecular basis for specific viral RNA recognition and 2'-O-ribose methylation by the dengue virus nonstructural protein 5(NS5). Proc Natl Acad Sci USA. 2015;112(48):14834–9. DOI: 10.1073/pnas.1514978112

12.   Dong H, Ren S, Zhang B, Zhou Y, Puig-Basagoiti F, Li H, et al. West Nile virus methyltransferase catalyzes two methylations of the viral RNA cap through a substrate-repositioning mechanism. J Virol . 2008;82(9):4295–307. DOI: 10.1128/JVI.02202-07

13.   Struijs K, Vincken JP, Doeswijk TG, Voragen AGJ, Gruppen H. The chain length of lignan macromolecule from flaxseed hulls is determined by the incorporation of coumaric acid glucosides and ferulic acid glucosides. Phytochemistry. 2009;70(2):262–9. DOI: 10.1016/j.phytochem.2008.12.015

14.   Badshah SL, Faisal S, Muhammad A, Poulson BG, Emwas AH, Jaremko M. Antiviral activities of flavonoids. Biomedicine & Pharmacotherapy. 2021 ;140:111596. DOI: 10.1016/j.biopha.2021.111596

15. Rani R, Bhutkar M, Tomar S. Therapeutic Antiviral Potential of Flavonoids. In Flavonoids as Nutraceuticals 2024 Apr 9 (pp. 57-101). Apple Academic Press.

16. Kim DJ, Roh E, Lee MH, Oi N, Lim DY, Kim MO, et al. Herbacetin Is a Novel Allosteric Inhibitor of Ornithine Decarboxylase with Antitumor Activity. DOI: 10.1158/0008-5472.CAN-15-0442

17. Bhutkar M, Ruchi R, Kothiala A, Mahajan S, Waghmode B, Kumar R, et al. Elucidation of the antiviral mechanism of natural therapeutic molecules Herbacetin and Caffeic acid phenethyl ester against Chikungunya and Dengue virus. bioRxiv. 2022. DOI:https://www.biorxiv.org/content/10.1101/2022.05.31.494145v1

18. Bullard-Feibelman KM, Fuller BP, Geiss BJ. A Sensitive and Robust High-Throughput Screening Assay for Inhibitors of the Chikungunya Virus nsP1 Capping Enzyme. 2016. DOI: 10.1371/journal.pone.0158923

19. Lim SP, Sonntag LS, Noble C, Nilar SH, Ng RH, Zou G, et al. Small Molecule Inhibitors That Selectively Block Dengue Virus Methyltransferase. Journal of Biological Chemistry. 2011 Feb 25;286(8):6233–40. DOI: 10.1074/jbc.M110.179184

20. Dhaka P, Singh A, Choudhary S, Peddinti RK, Kumar P, Sharma GK, et al. Mechanistic and thermodynamic characterization of antiviral inhibitors targeting nucleocapsid N-terminal domain of SARS-CoV-2. Arch Biochem Biophys. 2023;750:109820. DOI: 10.1016/j.abb.2023.109820

21. Mishra AC, Arankalle VA, Gadhave SA, Mahadik PH, Shrivastava S, Bhutkar M, Vaidya VM. Stratified sero-prevalence revealed overall high disease burden of dengue but suboptimal immunity in younger age groups in Pune, India. PLoS neglected tropical diseases. 2018 Aug 6;12(8):e0006657. DOI: 10.1371/journal.pntd.0006657

22. Chien LJ, Liao TL, Shu PY, Huang JH, Gubler DJ, Chang GJJ. Development of real-time reverse transcriptase PCR assays to detect and serotype dengue viruses. J Clin Microbiol. 2006 Apr ;44(4):1295–304. DOI: 10.1128/JCM.44.4.1295-1304.2006

23. Mudgal R, Mahajan S, Tomar S. Inhibition of Chikungunya virus by an adenosine analog targeting the SAM-dependent nsP1 methyltransferase. FEBS Lett. 2020;594(4):678–94. DOI: 10.1002/1873-3468.13642

24. Bhutkar M, Saha A, Tomar S. Viral methyltransferase inhibitors: berbamine, venetoclax, and ponatinib as efficacious antivirals against chikungunya virus. Arch Biochem Biophys. 2024;759:110111. DOI: 10.1016/j.abb.2024.110111

25. Coutard B, Decroly E, Li C, Sharff A, Lescar J, Bricogne G, Barral K. Assessment of Dengue virus helicase and methyltransferase as targets for fragment-based drug discovery. Antiviral research. 2014 ;106:61-70. DOI: 10.1016/j.antiviral.2014.03.013

26. Benmansour F, Trist I, Coutard B, Decroly E, Querat G, Brancale A, et al. Discovery of novel dengue virus NS5 methyltransferase non-nucleoside inhibitors by fragment-based drug design. Eur J Med Chem. 2017;125:865–80. DOI: 10.1016/j.ejmech.2016.10.007

27.   Vagin A, Teplyakov A. MOLREP: an Automated Program for Molecular Replacement.
urn:issn:0021-8898. 1997;30(6):1022–5. DOI: 10.1107/S0907444909042589

28.   Murshudov GN, Skubák P, Lebedev AA, Pannu NS, Steiner RA, Nicholls RA, et al.
REFMAC5 for the refinement of macromolecular crystal structures. Acta Crystallogr D Biol
Crystallogr. 2011 Apr;67(Pt 4):355–67. DOI: 10.1107/S0907444911001314.

29.   Project CC. The CCP4 suite: programs for protein crystallography. Acta
crystallographica. Section D, Biological crystallography. 1994 Sep 1;50(Pt 5):760-3. DOI:
10.1107/S0907444994003112

30.   Liebschner D, Afonine PV, Moriarty NW, Poon BK, Sobolev OV, Terwilliger TC, Adams
PD. Polder maps: improving OMIT maps by excluding bulk solvent. Acta Crystallographica
Section D: Structural Biology. 2017;73(2):148-57. DOI: 10.1107/S2059798316018210

31.   DeLano WL. The PyMOL Molecular Graphics System. DeLano Scientific. 2002

32.   Laskowski RA, Swindells MB. LigPlot+: multiple ligand-protein interaction diagrams for
drug discovery. J Chem Inf Model. 2011 ;51(10):2778–86. DOI: 10.1021/ci200227u.

33.   Petrey D, Fischer M, Honig B. Structural relationships among proteins with different
global topologies and their implications for function annotation strategies. Proceedings of the
National Academy of Sciences. 2009;106(41):17377-82. DOI: 10.1073/pnas.0907971106

34.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation
of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol.
2011;7. DOI: 10.1038/msb.2011.75

35.   Weng YL, Naik SR, Dingelstad N, Lugo MR, Kalyaanamoorthy S, Ganesan A. Molecular
dynamics and in silico mutagenesis on the reversible inhibitor-bound SARS-CoV-2 main
protease complexes reveal the role of lateral pocket in enhancing the ligand affinity. Scientific
Reports 2021 11:1 . 2021;11(1):1–22. DOI: 10.1038/s41598-021-86471-0

36.   Van Der Spoel D, Lindahl E, Hess B, Groenhof G, Mark AE, Berendsen HJC.
GROMACS: fast, flexible, and free. J Comput Chem. 2005 ;26(16):1701–18. DOI:
10.1002/jcc.20291

37.   Vanommeslaeghe K, Mackerell AD. CHARMM additive and polarizable force fields for
biophysics and computer-aided drug design. Biochimica et Biophysica Acta (BBA) - General
Subjects. 2015;1850(5):861–71. DOI: 10.1016/j.bbagen.2014.08.004

38.   Vanommeslaeghe K, Hatcher E, Acharya C, Kundu S, Zhong S, Shim J, et al. CHARMM
General Force Field: A Force Field for Drug-Like Molecules Compatible with the CHARMM
All-Atom Additive Biological Force Fields. J Comput Chem. 2010;31:671–90. DOI:
10.1002/jcc.21367

39.   Berendsen HJC, Postma JPM, Van Gunsteren WF, Dinola A, Haak JR. Molecular
dynamics with coupling to an external bath. J Chem Phys. 1984;81(8):3684–90.
DOI:10.1063/1.448118

Figure 1: Fluorescence polarisation-based F-GTP displacement (A) Binding curve showing the interaction between F-GTP and DENV3 NS5 MTase. (B) Inhibition curve showing the effect ofHC on the binding of F-GTP to the DENV3 NS5 MTase.

Figure 2: Evaluation of antiviral activity HC against DENV serotypes

*(A-C) Fold change in viral RNA levels for HC-treated cells and virus control (VC) samples infected with DENV 1, DENV 2, and DENV 4. Statistical significance is indicated with asterisks (****p < 0.0001, ***p < 0.001). (D-F) Plaque-forming units (PFU) per milliliter for HC-treated cells and virus control (VC) samples infected with dengue serotypes DENV 1, DENV 2, and DENV 4. PFU measurements were performed to quantify infectious virus particles.*

Figure 3: *Evaluation of the antiviral effect by immunofluorescence assay (IFA).*

Immunofluorescent staining of HC (200 µM) treated cells and virus control (VC) samples

infected with DENV 1, DENV 2, and DENV 4. Green fluorescence indicates the virus load, and

blue fluorescence indicates the nuclear staining with DAPI with a 10 X objective lens. The scale

bar is 400 µm.

Figure 4: *Structural analysis of the binding of HC and SAH to the DENV3 NS5 MTase. A)*

Polder omit map of HC at a contour level of 2.5σ. HC is displayed in green, and the polder map

is shown as a magenta mesh. B) Surface and cartoon representation of the DENV3 NS5 MTase

complex showing the binding pockets for HC and SAH. HC and SAH are depicted in stick

representation, with HC and SAH in pink. The protein is shown in a green cartoon, and the

surface is transparent grey. C) and D) 2D LigPlot representations for HC and SAH, respectively,

showing the key residues involved in their interactions. Hydrogen bonds are depicted as green

dashed lines, and hydrophobic contacts are represented by spiked red arcs.

Figure 5: _Structural analysis of HC, GTP, and cap 0 RNA interactions with DENV3 NS5 MTase._

A) Surface representation of superposition of the DENV3 NS5 MTase-HC with DENV3 NS5

MTase-SAH-cap 0 RNA (PDB:5DTO). Binding pockets with Cap 0 RNA is shown in cyan, SAH

in cyan, and HC in blue. Residues lining the cavity for cap 0 RNA are depicted in orange, SAH in

yellow, and common residues from HC and cap 0 RNA in purple. A violet sphere indicates a

bound $Mg^{2\square}$ ion. B) C-alpha chain superposition of DENV3 NS5 MTase-HC (blue) and the

structure of GTP-bound MTase from PDB ID 5DTO (red). The ligands HC and cap 0 RNA are

visualized with specific shapes and colors: cap 0 as a stick, RNA as a ribbon structure (yellow),

and HC as a stick model (pink). All interacting residues are marked, with cap 0 RNA-interacting

residues shown in blue and HC-interacting residues in green. The green sphere represents a

bound $Mg^{2\square}$ ion. (C) Structure superposition of DENV3 NS5 MTase-HC (Pink-colored cartoon)

with the structure from DENV3 NS5 MTase-GTP (PDB 4V0R) (wheat-colored cartoon). HC is

represented in pink, with key residues interacting with HC also highlighted in pink. GTP is

shown in green, residues are marked in blue (PDB 4V0R).D) Structure superposition of DENV3

NS5 MTase- A-G from cap 0 RNA (PDB 5DTO) (violet cartoon), with the DENV3 NS5 MTase-

HC (wheat-colored cartoon). Adenine (A) and guanine (G) bases from the cap 0 RNA are colored

blue, with residues highlighted in green (PDB 5DTO). HC is shown in pink, with residues

highlighted in cyan. The red arc represents the angle of interaction. Hydrogen bonds between

interacting residues are depicted as yellow dashed lines.

Parameter	Value
Resolution range	24.88–2.80 (2.95–2.80)
Space group	P 2 2$_1$ 2$_1$
Unit cell dimensions: a b c (Å)	50.87 59.96 185.04
Completeness (%)	99.9 (100)
Rmerge	0.26 (0.924)
I/σ(I)	13.8 (4.6)
CC(1/2)	0.99 (0.89)
Refinement	
Reflections used in refinement	14586
Reflections used for R-free	707
R-work	0.245
R-free	0.302
Wilson B-factor (Å)	34.1
Number of non-hydrogen atoms	4182
Macromolecules	4078
Ligands	104
Protein residues	508
RMS (bonds) (Å)	0.01
RMS (angles) (°)	2.27
Ramachandran Plot	
Favored (%)	98.23
Allowed (%)	1.18
Outliers (%)	0.59
Average B-factor (Å)	36.0
Macromolecules (Å)	36.88
Ligands (Å)	41.23

Table 1:- Data collection, processing and refinement statistics of   DENV3 NS5 MTase

complexed with HC(PDB:8ZMC). Values in parentheses are for the highest-resolution shell.

a Rmerge = $\Sigma| I - \langle I \rangle |/\Sigma I$.

b R = $\Sigma|Fobs| - |Fcalc|/\Sigma|Fobs|$. The Rfree is the R calculated on the 5% reflections excluded for

refinement.

c RMS is root mean square.

Table 2: - Detailed molecular interactions between the DENV3 NS5 MTase and HC

(PDB:8ZMC).

	Interactions			
	Hydrogen			**Hydrophobic**
Ligand	**Residues**	**Molecules**	**Bond length (Å)**	
	Arg57	O6-NE	2.88	Arg211
HC	Lys42	O3-NZ	2.85	

Table 3: - Detailed molecular interactions between the DENV3 NS5 MTase domain and GTP
(PDB:4V0R).

Ligand	Interactions			
	Hydrogen			Hydrophobic
	Residues	Molecules	Bond length (Å)	
GTP	Lys14	O2'-NZ	2.8	Phe25, Pro152,Asn18, Lys29, Ser150
		O3'-NZ	2.9	
	Leu20	N2-O	3.4	
	Asn18	N2-O	2.9	
		O2'-OD1	3.0	
	Ser151	O3'-O	3.3	
	Arg22	O6-NH1	2.6	
	Leu17	N2-O	2.8	
	Arg211	O1B-NH1	3.3	
		O3B-NH1	3.2	
	Ser213	O1B-OG	3.1	

545 **Table 4:-** Detailed molecular interactions between the DENV3 NS5 MTase domain and A-G

546 (from cap 0 RNA) (PDB:5DTO).

| Ligand | Interactions | | | Hydrophobic |
| | Hydrogen | | | |
	Residues	Molecules	Bond length (Å)	
A-G	Ser150	OG-OP1	2.88	Lys61, Leu 182, Arg84,
	Thr214	NH2-OP1	2.66	
	Lys180	NZ-O2	2.35	Ser56, Glu 216, Gly148,
	Glu111	N2- OE2	2.58	
	Arg57	NH2-OP1	3.25	

548 These color-coded residues represent various interactions in a molecular context. Violet

549 represents hydrogen bonds with A-G, and red signifies HC's hydrophobic residues that make

550 hydrogen bonds with GTP

Supplementary Experimental Procedures

Multiple Sequence Alignment (MSA)

The amino acid sequence of the NS5 MTase domain of flaviviruses was compared with the

DENV3 NS5 MTase protein. DENV3 NS5 MTase were utilized as reference points for these

alignments. The sequence alignment profile of NS5 MTase sequences was performed via Clustal

Omega tool and analyzed by a graphical colored depiction using ESPript 3.0 (34). MSA was

performed for the MTase domain of NS5 from DENV 1 (PDB:5IKM) , DENV 2 (PDB:1R6A),

DENV3 (PDB: 5EHI), DENV 4 (Gene ID:NP_740325.1), Zika virus (ZIKV) (PDB:5WZ2),

West Nile Virus (WNV) (PDB:2OY0), Yellow Fever Virus (YFV) (PDB:3EVA), and Japanese

Encephalitis Virus (JEV) (PDB:4K6M) from the flavivirus family and the analysis was carried

out.

MD simulation studies:

The HC-bound MTase models were built using the experimentally resolved 3D structure of

MTase (PDB: 8DKZ). The process involved aligning the MTase structures (PDB: 5IKM, 1R6A,

and modelled structure of DENV4) to the template structure (PDB: 8DKZ), followed by the

removal of other ligands and molecules after alignment (35). All procedures for constructing

these HC-bound MTase models were performed using PyMOL. Subsequently, the prepared

complexes underwent a 100 ns MD simulation to assess their stability and interactions. All

simulation studies were performed using GROMACS 2022.2 (36) with the CHARMM36 force

field (37) on a LINUX-based workstation. Ligand parameters and topology files were generated

using the CHARMM General Force Field (CGenFF) program (38). The system was modelled in

a cubic box, neutralized by adding ions and solvated with water molecules. Energy minimization

was conducted using the steepest descent algorithm, followed by a two-phase equilibration for

100 ps, first under constant number of particles, volume, and temperature (NVT), and then under

constant number of particles, pressure, and temperature (NPT). During NVT equilibration, the

temperature was set to 300 K using a short-range electrostatic cut-off of 1.2 nm with the

Berendsen thermostat (39). Afterward, the system transitioned to NPT equilibration, where

coordinates were generated every 1 ps. A 100 ns production run was conducted with a 2fs time

step, saving trajectories every 10 ps. The resulting trajectories were used to assess the stability

and interactions of protein-ligand complexes.

587
Supplemetnary Figure 1: *Sequence alignment of NS5 MTase domain (residues 1 to 263) from*

DENV with MTase domains of other flaviviruses. DENV 1 (PDB: 5IKM), DENV 2 (PDB:1R6A),

DENV3 (PDB: 5EHI), DENV 4 (Gene id:NP_740325.1), ZIKV (PDB:5WZ2), WNV

(PDB:2OY0), YFV (PDB:3EVA), and JEV (PDB:4K6M). Residues are represented with single

letter amino acid code, with identical residues indicated in white font and boxed in red, while

similar residues are indicated in red font. Blue triangles correspond to the HC binding site

residues. Alignments were generated via Clustal Omega and ESPript tools.

Supplementary Figure 2: The MD analysis of MTase-HC complexes for DENV1, DENV2,

DENV3, and DENV4 serotypes over a 100 ns simulation. *A) RMSD plots of all DENV*

serotypes. B) RMSF data for all DENV serotypes with the black dashed circle indicating the HC

binding residue. C) Rg values indicate conformations for all serotypes D) The hydrogen bond

analysis of all DENV serotypes. E) SASA values of all DENV serotypes F). Radar plot showing

the short-range (SR) interaction energies in kJ/mol. The red line represents the Lennard-Jones

SR interaction energies, while the blue line represents the Coulombic SR interaction energies.

Each vertex corresponds to a different serotype: DENV1, DENV2, DENV3, and DENV4.